



UNIVERSITY OF
ILLINOIS LIBRARY
AT URBANA-CHAMPAIGN




NOV 27 1979

THE UNIVERSITY OF CHICAGO PRESS
CHICAGO, ILL. 60637
U.S.A.



RECORDS OF THE U. S. DEPARTMENT OF AGRICULTURE
ABSTRACTS



99405
V.16 no.9

Biol.


112
THE LIBRARY OF THE
MAY 03 1979
UNIVERSITY OF ALABAMA
AT URBANA-CHAMBERS

CARCINOGENESIS ABSTRACTS

MAY 10 1979

BRITISH LIBRARY
LONDON

VOLUME 16,
ISSUE 9



CARCINOGENESIS ABSTRACTS

A monthly publication sponsored by the
NATIONAL CANCER INSTITUTE

EDITOR

GEORGE P. STUDZINSKI, M.D., Ph.D.
COLLEGE OF MEDICINE AND DENTISTRY
OF NEW JERSEY, NEWARK

ASSOCIATE EDITOR

JUSSI J. SAUKKONEN, M.D.
JEFFERSON MEDICAL COLLEGE
PHILADELPHIA

NCI STAFF CONSULTANTS

ELIZABETH WEISBURGER, Ph.D.
JOAN W. CHASE, M.S.

Literature Selected, Abstracted, and Indexed
by

Information Services Department, Science Information Services Organization

BRUCE H. KLEINSTEIN, Ph.D., J.D., Assistant Director, Biomedical Projects

RUTHANN E. AUCHINLECK, Managing Editor

BIOLOGY LIBRARY
101 BURRILL HALL

The Franklin Research Center
The Benjamin Franklin Parkway, Philadelphia, Pennsylvania 19103

All rights reserved. No part of this journal may be reproduced in any form, for any purpose or by any means, abstracted, or entered into any data base, electronic or otherwise, without specific permission in writing from the publisher.

Copyright © 1979 The Franklin Institute Press

CARCINOGENESIS ABSTRACTS

6699408
CHL
V16
no 9-12
cop. 2

Biol.

VOLUME 16, ISSUE 9

CONTENTS

	Cross Reference Abbreviations	Article Numbers	Page
REVIEW	(Rev)	78-4801-78-4937	1791
CHEMICAL CARCINOGENESIS	(Chem)	78-4938-78-5167	1824
PHYSICAL CARCINOGENESIS	(Phys)	78-5168-78-5186	1873
VIRAL CARCINOGENESIS	(Viral)	78-5187-78-5276	1878
IMMUNOLOGY	(Immun)	78-5277-78-5294	1899
PATHOGENESIS	(Path)	78-5295-78-5343	1903
EPIDEMIOLOGY AND BIOMETRY	(Epid-Biom)	78-5344-78-5388	1914
MISCELLANEOUS	(Misc)	78-5389-78-5400	1923
AUTHOR INDEX			1927
SUBJECT INDEX			1937
CHEMICAL ABSTRACTS SERVICE REGISTRY NUMBER INDEX			2021
WISWESSER LINE NOTATION INDEX			2025

GENERAL INFORMATION


CARCINOGENESIS ABSTRACTS makes available abstracts, annotations or citations of significant carcinogenesis articles collected from the current major biomedical sources of world literature. This service is provided by the National Cancer Institute through a contract with the Franklin Research Center for preparation of the publication, under Contract No. NOI-CP-75885 with the National Cancer Institute, U.S. Department of Health, Education and Welfare. Published and distributed by the Franklin Institute PressSM.

SUBSCRIPTIONS

For sale by The Franklin Institute Press, Box 2266, Philadelphia, PA 19103. Subscription rate for Volume 16, consisting of 12 monthly issues and a cumulative index is \$75.00, Domestic and \$90.00, Foreign. Check should be made payable to The Franklin Institute Press.

Maruzen Co., Ltd., 3-10, Nihonbashi 2-Chome, Tokyo 103, Japan, is the exclusive distributor of CARCINOGENESIS ABSTRACTS in Japan.

MAY 10 1979



GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations, or citations of significant cancer articles collected from the current major biomedical sources of the world literature. This service is provided by the National Cancer Institute through a contract with Franklin Research Center for preparation of the publication.

FREQUENCY OF PUBLICATION: *CARCINOGENESIS ABSTRACTS*, Volume 16, is published in 12 monthly issues with indexes, plus a cumulative index.

ABBREVIATIONS

JOURNAL names are abbreviated according to the *List of Journals Indexed in Index Medicus, Abbreviation Listing*. If the journal is not listed in this, abbreviations are derived from the *International List of Periodical Title Word Abbreviations*.

LANGUAGE of the article is indicated in parentheses after the title and is represented by a three-letter code. The source for these codes is *MARC Manuals Used by the Library of Congress*, pages 183-187.

ABBREVIATIONS used in abstracts:

A	angstrom(s)	mOsm	milliosmolar
ACTH	adrenocorticotrophic hormone	max	maximum
ADP	adenosine diphosphate	mEq	milliequivalent(s)
AMP	adenosine monophosphate	min	minute(s)
ATP	adenosine triphosphate	ml	milliliter(s)
approx	approximately	μl	microliter(s)
av	average	mm	millimeter(s)
BCG	bacillus Calmette-Guerin	mo	month(s)
bid	twice daily	mol wt	molecular weight
C	degree(s) centigrade	N	normal concentration
cal	calorie(s)	NAD	nicotinamide adenine dinucleotide
kcal	kilocalorie(s)	NADH	reduced nicotinamide adenine dinucleotide
cc	cubic centimeter(s)	NADP	nicotinamide adenine dinucleotidephosphate
Ci	curie(s)	NADPH	reduced nicotinamide adenine dinucleotidephosphate
mCi	millicurie(s)	NCI	National Cancer Institute
μCi	microcurie(s)	NIH	National Institutes of Health
cm	centimeter(s)	PAS	periodic acid-Schiff
CNS	central nervous system	po	orally
cpm	counts per minute	ppb	parts per billion
DNA	deoxyribonucleic acid	ppm	parts per million
ED₅₀	median effective dose	qid	four times daily
EDTA	ethylenediamine tetraacetic acid	qod	every other day
g	gram(s)	QO₂	oxygen quotient
kg	kilogram(s)	R	roentgen
mg	milligram(s)	RBC	red blood cells (erythrocytes)
μg	microgram(s)	RNA	ribonucleic acid
Hb	hemoglobin	rpm	revolutions per minute
hr	hour(s)	sc	subcutaneous
ia	intra-arterial	sec	second(s)
id	intradermal	SGOT	serum glutamic-oxaloacetic transaminase
IgA	Immunoglobulin A	SGPT	serum glutamic-pyruvic transaminase
IgB	Immunoglobulin B	soln	solution
IgG	Immunoglobulin G	TCD	tissue culture dose
IgM	Immunoglobulin M	TCD₅₀	median tissue culture dose
ILS	increased life span	tid	three times daily
im	intramuscular	UV	ultraviolet
ip	intraperitoneal	WBC	white blood cells (leukocytes)
IU	International Unit(s)	wk	week(s)
iv	intravenous	wt	weight
Km	Michaelis constant	X	times
LD	lethal dose	yr	year(s)
LD₅₀	median lethal dose		
M	molar		
μM	micromolar		



REVIEW

4-4801 **Aspects in Carcinogen Bioassay.** (Eng) Sontag, J. M. (Div. Cancer Cause and Prevention, NCI, Bethesda, MD, 20014). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1327-1338; 1977.

Factors to be considered in the design and conduct of long-term carcinogen bioassays are reviewed. These factors include the following: (1) selection of a chemical compound in a form that will reflect the danger to humans; (2) chemical quality control; (3) choice of animal for bioassay; (4) proper care of the animal and its surroundings; (5) genetic composition of the animal; (6) safety and health of laboratory personnel; (7) establishment of the max tolerated dose that can be administered during a chronic toxicity study; (8) parameters of the chronic study such as animal group size, treatment dose levels, route and mode of administration, and the detail of pathologic examination; (9) the use of positive control; and (10) method of data collection. (9 refs.)

8-4802 **Identifying and Defining Chemical Carcinogens.** (Eng) Saffiotti, U. (Experimental Pathology Branch, Carcinogenesis Program, Div. Cancer Cause and Prevention, NCI, NIH, Bethesda, MD, 20014). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1311-1326; 1977.

Changing parameters in toxicological studies are reviewed, and the application of these studies to chemical carcinogenesis is discussed. Traditionally, toxicology was concerned with substances producing early recognizable effects, the intensity of which was directly correlated with the intensity of exposure. The manifestations of the toxicity were due to altered functional products, degenerative changes, or target cell death. Carcinogenic toxicity, however, is characterized by a delay in the appearance of toxic effects; and although the frequency of tumor incidence (injury) is directly correlated with the intensity of exposure, the intensity of the induced pathology (tumor growth and spread) is independent of the intensity of exposure. Furthermore, the manifestations of toxicity are due to the proliferative response of a new altered cell population. The application of this new toxicology include epidemiologic studies of cancer prevalence in humans exposed to potential carcinogens, experimental long-term stu-

dies of cancer induction in animals exposed to test substances, and experimental short-term studies of neoplastic transformation, mutagenic activity, and chemical interactions with target macromolecules. (103 refs.)

78-4803 **Chronic Toxicity and Carcinogenicity Guidelines.** (Eng) Page, N. P. (Natl. Inst. Occupational Safety and Health, Parklawn Building, Room 8A-30, 5600 Fishers Lane, Rockville, MD, 20857). *J Environ Pathol Toxicol* 1(2): 161-182; 1977.

Protocols and guidelines for chronic toxicity and carcinogenesis tests that have been developed by national and international organizations are reviewed with respect to two factors: the potential for standardization and the assessment of issues of greatest concern. Of the major design aspects, animal models, route of exposure, dose selection, pathology examination, and reporting requirements are the most controversial. Complete standardization of protocols is not considered desirable because test objectives may differ and overly rigid protocols would be likely to stifle development of improved methodology. One of the greatest dilemmas is the extent of pathology that should be required in routine bioassays. Although a thorough microscopic examination is obviously desirable, the increased requirements for testing chemicals using the extensive pathology protocols now in existence has created a demand exceeding available scientific manpower. The past few years have seen great improvements in standard bioassays, but refinements are still needed to provide for: the use of dose level(s) that will allow for assessment of dose response and altered physiology; comparative/correlative metabolism studies; and the routine use of a defined, contaminant-free diet. (48 refs.)

78-4804 **Metabolic Activation of Carcinogens.** (Eng) Irving, C. C. (Veterans Admin. Hosp., Dept. Urology, Univ. Tennessee Center Health Sciences, Memphis, TN). In: *Carcinogens: Identification and Mechanisms of Action. Proceedings of the Thirty-First Annual Symposium on Fundamental Cancer Research Held at Houston, Texas, 1-3 March 1978*. University of Texas System Cancer Center, M. D. Anderson Hospital and Tumor Institute, NCI, American Cancer Society. (Austin, Texas). 13 pp.; 1978. (no refs)

78-4805 **When Is the Extrapolation of Animal Carcinogenesis to Man Inappropriate? (Letter to Edi-**

tor). (Eng) Gross, P. (Industrial Health Foundation, Inc., 5231 Centre Ave., Pittsburgh, PA, 15232); Braun, D. C. *J Occup Med* 20(4): 236-240; 1978.

Positive carcinogenesis test results in animals can be extrapolated to humans except when the carcinogenic effect results from physical rather than chemical induction or when the administrative route was grossly inappropriate. Experiments in which there were ip injections or pleural implantations of large fibers in rats are presented as examples of these exceptions. (13 refs)

78-4806 Inhibitors of Chemical Carcinogenesis. (Eng) Wattenberg, L. W. (Dept. Laboratory Medicine and Pathology, Univ. Minnesota, Minneapolis, MN, 55455); Lam, L. K.; Speier, J. L.; Loub, W. D.; Borchert, P. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 785-799; 1977.

The following low-toxicity inhibitors of chemical carcinogenesis are discussed: butylated hydroxyanisole, butylated hydroxytoluene, ethoxyquin, disulfiram and related compounds, benzyl isothiocyanate and related compounds, selenium compounds, cysteamine hydrochloride, and various inducers or microsomal mixed-function oxidase activity. The latter include polycyclic hydrocarbons, polychlorinated biphenyls, phenobarbital β -naphthoflavone, phenothiazine, α -benzene hexachloride, chlordane, and quercetinpentamethyl ether. These compounds have broad structures, which indicate that a substantial number of other, as yet unidentified, inhibitors probably exist. Before an inhibitor can be considered for use in diminishing the impact of chemical carcinogens on human populations, it should be shown to have trivial or no toxicity, and evidence should exist that its intake by a particular population results in a low cancer incidence. Less absolute requirements regarding a lack of toxicity might be justified for an inhibitor that is to be used in populations with elevated exposures to chemical carcinogens. (32 refs.)

78-4807 Species Differences in Carcinogenesis Testing. (Eng) Rall, D. P. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC, 27709). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1383-1390; 1977.

Problems associated with the extrapolation of carcinogenesis test data from small animals to humans are reviewed, and the ability of animal studies to predict human toxic and carcino-

genic responses is discussed. Differences between laboratory animals and man include different metabolic patterns and cell turnover time, different life spans, differences in body sizes resulting in different distribution of chemicals in animals, number of cells at risk, extrapolation from a genetically homogeneous population to a heterogeneous one and from a well-fed one to one with various nutritional statuses, and (since test animals receive basically the pure compound) the problem of synergistic toxicity. However, the only well-controlled comparative study indicated a strong correlation between the toxicity of 17 anticancer drugs in man and laboratory animals. In this experiment, the clinical situation and dosimetry in man were duplicated in the laboratory animals. It is suggested that carcinogenic testing situations in laboratory animals approximate the conditions to which humans will likely be exposed. (10 refs.)

78-4808 The Applicability of Fish to Carcinogenesis Studies. (Hun) Lapis, K. (I. sz. Korbontani Intezet, Semmelweis Orvostudományi Egyetem, Budapest, Hungary); Simon, K.; Szende, B. *Magy Onkol* 22(2): 75-86; 1978.

Although the testing of substances for carcinogenic activity is expensive and time-consuming, in vitro short-term tests cannot be substituted for in vivo animal experiments traditionally performed with rodents. Fish are more promising models for carcinogenicity studies than rodents since they require a shorter time for tumor development, and the studies are less expensive. The results obtained with both fish and rodents are comparable. Furthermore, the use of both rodents and fish would fulfill the requirement that tests be performed on more than one species. Under certain circumstances fish can also be used to screen carcinogens. The results of carcinogenesis experiments with fish in the literature are presented. (34 refs)

78-4809 The Salmonella/Microsome Mutagenicity Test: Predictive Value for Animal Carcinogenicity. (Eng) McCann, J. (Dept. Biochemistry, Univ. California, Berkeley, CA, 94720); Ames, B. N. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1431-1450; 1977.

The predictive value of the *Salmonella*/microsome mutagenicity assay for animal carcinogenicity is discussed. Testing of hundreds of chemicals has revealed greater than millionfold range of mutagenic potency of carcinogens in the assay. Attempts are being made to determine the relationship between mutagenic and carcinogenic potency, but information on the latter is sparse. The assay was validated with about 300 organic chemicals that have been tested in conven-

tional animal carcinogenicity tests. Most (157/175) of the carcinogens were mutagenic in the assay, and 94/104 of the noncarcinogens were nonmutagens. With further improvements in both the tester strains and the metabolic activation system, the assay is expected to detect at least 95% of all carcinogens. The remaining percent will probably fall into the hormone and promoter category or will be a class of mutagens unique to animal cells. False positives may be caused by bacterial activation of chemicals by enzymes that are not functional in animals, by limitations of animal carcinogenicity tests, and by mutagenic impurities. Several chemicals used in consumer products, including two hair dye components, 4-nitro-o-phenylenediamine and 2-nitro-p-phenylenediamine, and tris-(2,3-dibromopropyl)phosphate, the main flame retardant in children's polyester pajamas, were detected as mutagens by the assay and have since proved to be carcinogenic. It is suggested that damage to DNA by environmental mutagens initiates most human cancers, as well as genetic defects. The results of these studies support the theory that radiation and chemical carcinogens cause cancer through somatic mutation. (49 refs)

78-4810 Epidemiology, Multistage Models, and Short-term Mutagenicity Tests. (Eng) Peto, R. (Regius Dept. Medicine, Oxford Univ., Oxford OX2 6HE, England). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1403-1428; 1977.

The value of multistage models in calculating cancer risks is reviewed. Most multistage models indicate that the final cancer incidence rate is the arithmetic product of more than one term, each of the different terms being dependent on different causative agents or processes. If this is so, cancer incidence could be reduced by decreasing the levels of exposure to a compound (eg, an environmental pollutant) or by manipulating the other unknown factors involved and so disrupting the sequence of events. Mutagenicity studies could play a key role in identifying these factors. In reviewing multistage models, lung cancer, epidemiology, and suggestions that almost all carcinogens are mutagens, three conclusions were reached. (1) Multistage models do not necessarily require a large number (approx 6) of distinct rate-limiting cellular changes as is often assumed to take place during epithelial cell transformation. (2) Although the mechanisms of induction of most carcinomas may be similar, they probably differ fundamentally from the mechanisms of induction of leukemias, sarcomas, etc; thus much of the research into leukemia or sarcoma viruses may be irrelevant to human carcinomas. (3) Some of the extrinsic rate-determining causes of human lung cancer are not the direct- or indirect-acting agents detected by in vitro mutagenicity tests; this is probably true for other carcinomas as well. (45 refs.)

78-4811 Safeguarding the Quality of Microbiological Mutagenicity Testing of Foreign Substances in Routine Laboratory. (Ger) Norpoth, K. (Munster, W. Germany). *Staub-Reinhalt Luft* 38(6): 235-239; 1978.

General methodological and statistical questions relevant to the use of bacteria in mutagenicity tests of environmental compounds are discussed. The probability of carcinogenic effects can be assessed from the results of mutagenicity tests on microorganisms such as *Salmonella typhimurium*. The development of routine bacterial mutagenicity tests requires standardization, quality control, and statistical validation of the results. The prerequisites include positive and negative controls, continuous checks on the indicator bacteria, constant activity of the mammalian oxygenase used, species-specificity of the oxygenase, and attention to the influence of the solvent. 3,4-benzo(a)pyrene, aflatoxin B₁, vinyl chloride, diaminotoluene, known carcinogens, and bis(chloroethyl)ether, a suspected carcinogen, were found to induce mutations in microorganisms. (10 refs.)

78-4812 Identification of Carcinogens by Mutagen Testing in *Drosophila*: The Relative Reliability for the Kinds of Genetic Damage Measured. (Eng) Vogel, E. (Dept. Radiation Genetics and Chemical Mutagenesis, Sylvius Lab., State Univ. Leiden, Leiden, Netherlands). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1483-1499; 1977.

The usefulness of mutagen testing of *Drosophila* for the identification of carcinogens is discussed. Identification is largely dependent on the genetic end point used as a diagnostic criterion. Comparative experiments have shown the existence of two effect levels; ie, concentrations required to raise the number of point mutations (recessive lethals) significantly are considerably lower than those needed for the chromosome breakage effect. This explains the superior discriminating power of recessive-lethal assays over assays of chromosome breakage. *Drosophila* can execute the essential activation steps required to convert precarcinogens to highly reactive species. Intragonadal metabolism plays an important role in this activation, and particular attention must be paid to metabolically active germ-cell stages when testing precarcinogens in *Drosophila*. The larvae also have the enzymatic machinery needed to carry out the essential activation steps, and they are more sensitive to mutagens than the adult fly. Some problems have been encountered in this system, including deactivation of some compounds before the genetic target is reached, differences in mutagenicity depending on route of application, and the nonmutagenicity of the carcinogenic polycyclic hydrocarbons and aromatic amines. The latter problem may be overcome by developing mutagen-sensitive

strains. It is proposed that the *Drosophila* assay system be used as an exploratory screening phase and also in in-depth analyses of potent mutagens. (39 refs)

- 78-4813 **Estimation of "Safe Doses" in Carcinogenic Experiments.** (Eng) Hartley, H. O. (Inst. Statistics, Texas A & M Univ., College Station, TX, 77843); Sielken, R. L. *J Environ Pathol Toxicol* 1(2): 241-278; 1977.

A multiparametric model approach to estimating safe doses in carcinogen-testing experiments is described. It is comprised of an adequately large number of parameters to allow for differences in the idiosyncracies of suspected carcinogens, and the host species and, to a limited degree, for variations in the experimental protocols. The same computer program can cover the analysis of an experiment in which latent periods have been recorded as well as experiments in which only tumor incidence rates have been recorded, or mixtures of the two. The max likelihood estimation procedure has asymptotic optimality properties. It uses the latest techniques of convex programming. Although the model is multiparametric, it is of the form of the product model for age-specific hazard rates, a form that is now widely accepted. No estimates of safe doses can be made if the spontaneous incidence rate is zero, if the experimental dose levels are too small, and if no tumors have been observed. If the experimental doses are too small and the tumor incidence rates are comparable with the spontaneous rate, the estimation procedure has extremely large errors. The statistical analyses are given in detail in a series of appendices. (32 refs)

- 78-4814 **Aspects of the Hartley-Sielken Approach for Setting "Safe Doses" of Carcinogens.** (Eng) Mantel, N. (Biostatistics Center, George Washington Univ., Bethesda, MD, 20014). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1397-1401; 1977.

Various requirements imposed by the Hartley-Sielken approach to establishing safe dose levels of carcinogens for human exposure are reviewed. Since this approach depends on the fitting of high-degree polynomials into the calculations, a sufficiently large number of dose levels must be explored. Furthermore, because the data are going to be segmented, sufficiently large numbers of animals must be employed at each dose level. Finally, the resulting data must show some frequency of tumor occurrence. A comparison between the requirements of this method and a one-hit model is presented. (6 refs.)

- 78-4815 **Comparisons of Carcinogenic and Mutagenic Potency.** (Eng) Meselson, M. (Biological Labs., Harvard Univ., Cambridge, MA, 02138); Russell, K. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1473-1481; 1977.

Fourteen compounds that are both carcinogenic and mutagenic were examined to determine if there is a quantitative relationship between carcinogenic and mutagenic potency. These compounds were aflatoxin B₁, 4-aminobiphenyl, benzdine, benzo(a)pyrene, dibenz(a,h)anthracene, diethylnitrosamine, dimethylnitrosamine, di-n-butyl nitrosamine, 4,4'-methylene-bis-2-chloroaniline, methyl methanesulfonate, β -naphthylamine, N-nitrosomethylurea, 1,3-propane sultone, and sterigmatocystin. The carcinogenic potency of most of these compounds in laboratory rodents was approx equal to their mutagenic potency in *Salmonella typhimurium*, using the definitions of the two potencies given. For several N-nitroso compounds, however, carcinogenic potency was considerably greater than mutagenic potency. It is suggested that the carcinogenic potency of a mutagenic compound for humans may be roughly equal to that in rodents if the carcinogenic potencies are expressed in terms of the av normal life-spans of the respective species. With this condition, it is concluded that the Ames *Salmonella* test is sufficiently sensitive to analyze complex mixtures in order to identify mutagenic compounds that may be responsible for the induction of an important fraction of human cancers. (36 refs.)

- 78-4816 **Ultimate Chemical Carcinogens as Reactive Mutagenic Electrophiles.** (Eng) Miller, J. A. (McArdle Lab. Cancer Res., Univ. Wisconsin Medical Center, Madison, WI, 53706); Miller, E. C. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 605-627; 1977.

Available data indicate that the ultimate carcinogenic forms of most chemical carcinogens are strong electrophilic reactants that differ from the normal cellular electrophiles in that they generally attack cellular nucleophiles nonenzymatically and with relatively little imination. In this light, the metabolic activation of 2-acetylaminofluorene, N-methyl-4-aminoazobenzene, other amines and amides, the pyrrolizidine alkaloids, cycasin, and safrole is reviewed. The metabolic activation of aflatoxin B₁ (AFB₁) and the nature of the major covalent DNA and RNA adducts of AFB₁ formed in vitro and in vivo are also discussed. The potent carcinogenicity of AFB₁ 2,3-dichloride for the skin and lungs of mice and for the sc tissue of rats provides support for the conclusion that

AFB₁ 2,3-oxide is a major ultimate carcinogenic metabolite of AFB₁. The electrophilic character of ultimate chemical carcinogens provides the basis for at least three types of precursors for the detection of potential carcinogens. Thus, the electrophilic nature of the parent compound and of known or likely metabolites can serve as an early warning that these compounds may have carcinogenic activity. (90 refs.)

78-4817 Standard-Setting for Nitrites and Nitrosamines. (Eng) Lijinsky, W. (Chemical Carcinogenesis Program, Frederick Cancer Res. Center, Frederick, MD, 21501). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1807-1811; 1977.

Several methods for reducing human exposure to nitroso compounds are proposed. These methods include reduction of the excessive use of nitrosatable amines, reduction of the amount of residual nitrite allowable in cured meats, and the administration of drugs that can react with nitrites in the stomach by injection rather than po. Adherence to these suggestions may reduce cancer incidence over the next few decades. (5 refs.)

78-4818 Nitrate and the Etiology of Gastric Cancer. (Eng) Tannenbaum, S. R. (Dept. Nutrition and Food Science, Massachusetts Inst. Technology, Cambridge, MA, 02139); Archer, M. C.; Wishnok, J. S.; Correa, P.; Cuello, C.; Haenszel, W. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1609-1625; 1977.

Two models of gastric cancer are proposed based on evidence gathered in Colombia, South America, and laboratory tests of these models are described. A survey of water supplies in a high-risk area in Colombia has shown high concentrations of nitrates in wells. In this region, the disease progresses from normal mucosa to superficial gastritis, chronic atrophic gastritis, gastritis with intestinal metaplasia, and, finally, gastric cancer. By age 25, > 75% of the population in the high-risk area suffers from gastritis, whereas in the low-risk area the population appears to be free of gastric pathology. Two models emerge from these observations: (1) an agent in the environment, possibly an N-nitroso compound, leads to gastritis, which naturally progresses to intestinal metaplasia and then to cancer; (2) gastritis leads to elevation of the stomach pH and, concomitantly, to establishment of a microbial flora capable of nitrate reduction. This process, in turn, leads to a steady formation of carcinogenic N-nitroso compounds, which leads to gastric cancer. At present, there is not enough

evidence to decide between these two models. Laboratory studies involving saliva and stomach acid indicate that bacteria in the stomach of an individual with gastritis or intestinal metaplasia play the dual roles of nitrite former and nitrosation catalyst. Other studies have disclosed the formation of cyanamides from primary and secondary amines in human saliva. Mixing of these compounds with the stomach contents would result in the formation of highly mutagenic and carcinogenic N-nitrosocyanamides. (46 refs)

78-4819 Nitrosamines and Their Precursors in Food. (Eng) Lijinsky, W. (Chemical Carcinogenesis Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Taylor, H. W. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1579-1590; 1977.

The carcinogenicity of N-nitroso compounds and their formation from food components are discussed. More than 120 N-nitroso compounds have been synthesized and tested by long-term administration of small doses to rats. Three-quarters of these compounds were found to be carcinogenic. Two compounds induced tumors in > 20 animal species. It is improbable that any species is refractory to N-nitroso compounds. These compounds are formed easily from nitrites and amines and they probably exist in the human environment, particularly in foods that contain nitrite; eg, cured meats. Nitrite is also formed by reduction of the nitrate in vegetables, eg, spinach and beets, and water. Preformed nitrosamines found in food are usually present at low levels (10 ppb). A more important source of nitrosamines is from ingested amines and nitrite. A majority of drugs and medicines are nitrosatable amines, as are agricultural chemicals, food constituents, and food additives. In tests in which secondary and tertiary amines plus nitrite were given to rats in the drinking water, several amines, including aminopyrine and Methapyrene produced a significant incidence of malignant tumors. (18 refs)

78-4820 Evidence for the Formation of Electrophilic Metabolites from N-nitroso Compounds. (Eng) Magee, P. N. (Fels Res. Inst., Temple Univ. Sch. Medicine, Philadelphia, PA, 19140). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 629-637; 1977.

Evidence that administration of N-nitroso carcinogens to animals gives rise to the same type of electrophilic alkylating intermediate in the body is presented. There appears to be

no species among those tested so far, including mammals, birds, amphibia, and fish, that has proved resistant to carcinogenesis by N-nitroso compounds. Furthermore, there appear to be sufficient similarities in the patterns of tumor induction by these carcinogens to warrant the conclusion that their molecular mechanisms are probably similar. In contrast to the nitrosamides, the nitrosamines are essentially inactive as mutagens unless a metabolic activating system is available. Studies of the metabolism of dimethylnitrosamine indicate that the methylating intermediate must pass through the vascular walls of the liver and other organs capable of metabolizing the nitrosamine and methylate the RBC. The available evidence is consistent with the hypothesis that the carcinogenic N-nitroso compounds, symmetrical dialkylhydrazines, azo- and azoxyalkanes, and derivatives of methylazoxymethanol produce their biological effects by the reaction of cellular constituents with electrophilic alkylating intermediates. (45 refs.)

78-4821 N-Nitrosodimethylamine. (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso Compounds*. Magee, P. N., Chairman (International Agency for Research on Cancer: Switzerland): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 17, 365 pp.; 125-175; 1978.

Experimental findings on the carcinogenicity of N-nitrosodimethylamine (NDMA) are reviewed. Until April 1976, NDMA was used as an intermediate in the production of rocket fuel; currently, it is used solely for research purposes. This compound is carcinogenic in mice, rats, Syrian golden hamsters, Chinese hamsters, European hamsters, guinea pigs, rabbits, ducks, mastomys, various fish, newts, and frogs. It produces benign and malignant tumors, mainly of the liver, kidney, and respiratory tract, following administration by various routes. NDMA is mutagenic in various strains of bacteria and in *Drosophila*, and it induces chromosome aberrations in Chinese hamster cells. Although no human case reports or epidemiological studies were available for analysis, there is sufficient experimental evidence of the carcinogenicity of NDMA. It is suggested that NDMA should be regarded as if it were carcinogenic to humans. (274 refs.)

78-4822 N-Nitrosodiethylamine. (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso Compounds*. Magee, P. N., Chairman (International Agency for Research on Cancer: Switzerland): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 17, 365 pp.; 83-124; 1978.

Biological data relevant to an evaluation of carcinogenic risk to humans are presented for N-nitrosodiethylamine (NDEA). The compound has been reported to occur in the air (one area of Baltimore), tobacco smoke, well-water, the waste water

from 2/19 chemical plants, various types of food, animal feed, and alcoholic beverages. NDEA is carcinogenic in all animal species tested: mice, rats, rabbits, dogs, gerbils, pigs, monkeys, hedgehogs, various fish, frogs, birds, and Syrian golden, Chinese, and European hamsters. It induces benign and malignant tumors after administration by various routes, including ingestion, parenteral injection, inhalation, and rectal instillation. The major target organs are the liver, respiratory and upper digestive tracts, and the kidney. NDEA is also carcinogenic following administration prenatally and in single doses. Dose-response relationships were established in several studies. A metabolite, N-nitroso-N-ethyl-N-(2-hydroxyethyl)amine, produces mainly liver tumors after po administration to rats. NDEA was found to be mutagenic in several types of assays. The general population may be exposed to low levels of NDEA; however, no exposed group suitable for an epidemiological investigation has been identified. No human data are available. Because of the evidence of its carcinogenicity in many animal species, NDEA should be regarded as if it were carcinogenic to humans. (203 refs.)

78-4823 N-Nitrosodiethanolamine. (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso Compounds*. Magee, P. N., Chairman (International Agency for Research on Cancer: Switzerland): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 17, 365 pp.; 77-82; 1978.

Chemical and physical data; production, use, occurrence, and analysis; and biological data relevant to the evaluation of carcinogenic risk to humans are presented for N-nitrosodiethanolamine (NDELA). This compound is found in tobacco, pesticides, cosmetics, and, in much higher concentrations (0.02%-3%), in cutting fluids. NDELA is carcinogenic in BD rats after po administration (600-1,000 mg/kg/day, total dose 150-300 g/kg) and in Syrian golden hamsters after sc injection (2,260 mg/kg 2x/wk x 7 or 565 mg/kg x 27 over 45 wk; total doses approx 41 g/kg). It produces hepatocellular carcinomas and renal adenomas in the rats and adenocarcinomas of the nasal cavity, papillary tumors of the trachea, hepatocellular adenomas, and local fibrosarcomas in the hamsters. Although no case reports or epidemiological data for humans are available, the experimental data indicate that NDELA should be regarded as if it were carcinogenic to humans. (11 refs.)

78-4824 N-Nitrosodi-N-Propylamine. (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso Compounds*. Magee, P. N., Chairman (International Agency for Research on Cancer: Switzerland): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 17, 365 pp.; 177-189; 1978.

Chemical and physical data and the results of carcinogenicity

and related studies are given for N-nitrosodi-n-propylamine (NDPA), a compound that has been found in the wastewater from 3/28 chemical plants, cheese, alcoholic beverages, and in two pesticides [Treflan (trifluralin) and isopropalin]. NDPA is carcinogenic in rats after po administration and in rats and hamsters after sc injection, producing benign and malignant tumors of the liver, kidney, esophagus, and respiratory tract. The metabolite N-nitroso-N-(2-hydroxy-n-propyl)-n-propylamine produces benign and malignant tumors of the respiratory tract and liver upon sc injection into rats and hamsters. NDPA was mutagenic in *Salmonella typhimurium*, *Escherichia coli*, and Chinese hamster V79 cells. The general population may be exposed sporadically to low levels of NDPA, but there are not human data available. Reports of relatively high levels in Treflan (154,000 and 190,000 µg/liter) may permit the identification of exposed groups. There is sufficient evidence of a carcinogenic effect of NDPA in two animal species; therefore, the compound should be regarded as if it were carcinogenic to humans. (47 refs.)

78-4825 N-Nitrosodi-N-butylamine. (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso Compounds*. Magee, P. N., Chairman (International Agency for Research on Cancer: Switzerland): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 17, 365 pp.; 51-75; 1978.

The carcinogenic risk of N-nitrosodi-n-butylamine (NDBA) to humans was evaluated by reviewing the current data. NDBA has no known commercial uses. It has been found in tobacco smoke, in the effluent from a coke plant, in food (soybean oil, cheese, various pork products, canned and powdered soups, and canned meals), experimental animal feed, and in a commercial corrosion-resistant film. NDBA is carcinogenic in all animal species tested (mice, rats, rabbits, guinea pigs, and Syrian golden, Chinese, and European hamsters) by the po, sc, ip, or iv routes. It produces leukemia and benign malignant tumors in the urinary bladder, esophagus, liver, respiratory tract, stomach, and intestine; it is particularly effective as a bladder carcinogen. Administration prenatally and in single doses also results in malignancy. The two metabolites, N-nitroso-n-butyl-N-(4-hydroxybutyl)amine and N-nitroso-n-butyl-N-(3-carboxypropyl)amine, are also carcinogenic, the first in mice, rats, hamsters, and dogs and the second in rats. NDBA was found to be mutagenic in several types of assays. The general population may be exposed sporadically to low levels of NDBA, but no human data on its carcinogenicity are available. However, the animal data suggest that NDBA should be regarded as if it were carcinogenic in humans. (93 refs.)

78-4826 N-Nitroso-N-methylurea. (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso*

Compounds. Magee, P. N., Chairman (International Agency for Research on Cancer: Switzerland): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 17, 365 pp.; 227-255; 1978.

Biological data on the carcinogenicity of N-nitroso-N-methylurea (NMU) were evaluated relative to the carcinogenic risk to humans. NMU has been studied for use as a cancer chemotherapy agent and for its mutagenic effects on various plants. NMU is carcinogenic in mice, rats, guinea pigs, rabbits, gerbils, pigs, dogs, monkeys and Syrian golden, Chinese, and European hamsters. It induces benign and malignant tumors following administration by most routes, including ingestion. Tumor induction occurs in the nervous tissue, stomach, esophagus, pancreas, respiratory tract, intestine, lymphoreticular tissues, skin, and kidney. It is carcinogenic following administration prenatally and in single doses. The mutagenicity of NMU has been demonstrated in various systems. There is sufficient evidence of a carcinogenic effect of NMU in several animal species. Although no human data were available, NMU should be regarded as if it were carcinogenic to humans. (122 refs)

78-4827 Nervous-system-specific Carcinogenesis by Ethylnitrosourea in the Rat: Molecular and Cellular Aspects. (Eng) Rajewsky, M. F. (Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Augenlicht, L. H.; Biessmann, H.; Goth, R.; Hulser, D. F.; Laerum, O. D.; Lomakina, L. Y. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 709-728; 1977.

Molecular and cellular aspects of the induction of nervous system (NS) tumors in the rat by ethylnitrosourea (ENU) are reviewed. The extranuclear oxygen-6 atom of guanine in the DNA is a major site of ethylation by ENU, and O⁶-ethylation of guanine in DNA is not restricted to a situation in which the extranuclear oxygen atom is unprotected by hydrogen bonding (ie, replicating DNA). Selective persistence of O⁶-methylguanine in rat brain DNA was also demonstrated after application of the predominantly NS-specific carcinogen methylnitrosourea. Dissociated fetal rat brain cells (FBC) treated transplacentally with ENU became tumorigenic after 200 days. This time interval is of the same order as the median time until death with neuroectodermal tumors in the offspring of rats injected with the same dose of ENU at the same stage of gestation. Solid, transplantable tumors developed upon sc reimplantation of the cultured tumor cells. These tumors appeared histologically as neurinoma-, glioma-, or glioblastomalike and frequently pleiomorphic neoplasms. Although subpopulations of probable high-risk FBC have not been defined, proliferative glial precursors are potential candidates. (71 refs.)

78-4828 Animal Toxicology. (Eng) Kimbrough, R. (Toxicology Branch, Clinical Chemistry Div., Bureau Labs., Center Disease Control, Dept. Health, Education and Welfare, Public Health Service, Atlanta, GA, 30333); Buckley, J.; Fishbein, L.; Flamm, G.; Kasza, L.; Marcus, W.; Shibko, S.; Teske, R. *Environ Health Perspect* 24: 173-185; 1978.

Recently published information on the toxic effects of polychlorinated biphenyls (PCB) in mammals and the results of experimental studies in birds are reviewed. Specific topics include the acute toxicity of PCB and polybrominated biphenyls (PBB); the subacute and reproductive effects of PCB, PBB, and chlorinated dibenzofurans; and the mutagenicity and teratogenicity of PCB and PBB. Hepatomas and hyperplastic nodules were produced in the livers of BALB c/J mice fed Aroclor 1254 (300 ppm), hepatocellular carcinomas and nodular hyperplasia in male dd mice fed Kanechlor 500, (100-500 ppm), liver tumors in mice fed 250 ppm α -hexachlorocyclohexane (α -HCH) or in mice fed 250 ppm PCB and 100 or 50 ppm α -HCH or 250 or 100 ppm β -HCH, and liver tumors in rats fed Aroclor 1260, Kanechlor 400, and Aroclor 1254. The induction of liver tumors in Sprague-Dawley rats by known carcinogens such as 3'-methyl-4-dimethylaminoazobenzene, N-2-fluorenylacetamide, and diethylnitrosamine was inhibited by Kanechlor 500. PCB have also been implicated as immunosuppressants. The degree of toxicity of chlorinated naphthalenes increases with the degree of chlorination; the higher chlorinated naphthalenes have been associated with chloracne and liver damage in humans. (96 refs.)

78-4829 Carcinogenic Potential of the Polychlorinated Biphenyls. (Eng) Allen, J. R. (Dept. Pathology, Univ. Wisconsin Medical Sch., Madison, WI, 53706); Norback, D. H. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 173-186; 1977.

The general toxic effects and carcinogenic action of polychlorinated biphenyls (PCBs) are reviewed. Marked hypertrophy of the liver and stimulation of the mixed-function oxidases of the liver occur in experimental animals exposed to PCB's, and hypertrophic, hyperplastic, dysplastic, metaplastic, and carcinogenic changes have been recorded in chronically exposed rodents. Chronic feeding studies in rats and mice suggest that the more highly chlorinated PCB congeners, which are more slowly metabolized and persist in the tissues indefinitely, are the likely precarcinogens. Biochemical data also suggest that PCB's might be carcinogenic. However, only preliminary observations have been made on the association of PCB exposure and tumor development in humans. Until the potential danger of low-level, long-term exposure of the human population to PCB's is better under-

stood, it must be assumed that any level of exposure may be injurious to human health. (51 refs.)

78-4830 Metabolism of Chlorinated Alkenes and Alkanes as Related to Toxicity. (Eng) Henschler, D. (Institut für Pharmakologie und Toxikologie, Univ. Würzburg, D-87 Würzburg, W. Germany). *J Environ Pathol Toxicol* 1(2): 125-133; 1977.

Chlorine substitution in aliphatic compounds results in a destabilization in alkanes and a stabilization in alkenes. With alkanes, therefore, the main pathways of metabolic transformation to reactive intermediates are radical formation by a C-C break and dechlorination or dehydrochlorination. In the chlorinated ethylenes, the first metabolic transformation step is oxidation to electrophilic oxiranes. These compounds may be hydrolyzed enzymatically or nonenzymatically, react with cellular nucleophiles, or rearrange to either chlorinated aldehydes or acyl chlorides. With tetrachloroethylene, 1,2-cis-dichloroethylene, trans-dichloroethylene, 1,1-dichloroethylene, and vinyl chloride, the metabolites identified in in vivo experiments are identical to the thermal rearrangement products of the respective oxiranes. An exception is trichloroethylene, whose thermal rearrangement product is dichloroacetyl chloride; its metabolites in vivo, however, are entirely derived from trichloroacetaldehyde. Mutagenic and carcinogenic activities in the chlorinated ethylenes are determined by the stability of their oxiranes. This stability is higher in symmetrical than in asymmetrical chlorine substitution compounds: the relatively unstable and asymmetric oxiranes of trichloroethylene, 1,1-dichloroethylene, and vinyl chloride are mutagenic in the Ames test, but the more stable symmetric oxiranes of tetra-, 1,2-cis-, and trans-dichloroethylenes are inactive. (20 refs)

78-4831 Recent Findings on the Carcinogenicity of Chlorinated Olefins. (Eng) Maltoni, C. (Inst. Oncology and Tumor Center, Bologna, Italy). *Environ Health Perspect* 21: 1-5; 1977.

Major factors affecting the carcinogenicity of vinyl chloride (VC) and vinylidene chloride (VDC) are discussed. Although the two compounds have very similar molecular structures, they have widely different biological effects. VS is a multipotential carcinogen, but VDC has produced tumors only in the murine kidney. Dose, concentration, length of treatment, route of administration, and animal species, strain, sex, and age also significantly affect the neoplastic response. These factors may alter the metabolic pathway of the test compounds. The carcinogenicity of VC and VDC is due to their active metabolites, probably epoxy derivatives. Animal species, strain, and sex greatly affect the production of active metabolites of VDC. In Swiss mice and Sprague-Dawley rats, there is a parallelism between the toxic and carcinogenic effects of VDC in relation to species and sex. Confirmation of

this parallelism in other strains would indicate new routes for establishing experimental animal models, and for understanding the mechanisms of action of many organic carcinogens. (5 refs)

78-4832 **Vinyl Chloride: A Case for the Use of Laboratory Bioassay in the Regulatory Control Procedure.** (Eng) Wagoner, J. K. (Industry-wide Studies Branch, Div. Surveillance, Hazards Evaluations and Field Studies, Natl. Inst. Occupational Safety and Health, Cincinnati, OH, 45202); Infante, P. F. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1977-1805; 1977.

The need for laboratory bioassays in the regulatory control of chemicals is illustrated using vinyl chloride (VC) as an example. The induction of skin, lung, and bone tumors in rats exposed by inhalation to high VC levels (levels not infrequently approached in the industrial setting) was first reported in 1971, 40 yr after VC's commercial introduction but 3 yr before three cases of liver angiosarcoma were reported among workers at a VC polymerization facility. A follow-up study, initiated as a result of these initial angiosarcoma cases, showed a significant excess of deaths from malignant neoplasms among VC workers, compared with US white male death rates. Excess cancer mortality was found for four organ systems: brain and CNS, respiratory system, hepatic system, and lymphatic and hematopoietic system. The cancer risk increased with increasing years of exposure. In the follow-up study, 11/14 confirmed cases of biliary or liver cancer were hepatic angiosarcoma. Among 8 bronchogenic carcinomas, 5 were large-cell undifferentiated and 3 were adenocarcinomas, which is not consistent with distributions previously reported for inhaled carcinogens. A recent bioassay was predictive not only for the carcinogenicity of VC but also for several of the target organs. VC is mutagenic in microbial test systems, and VC metabolites have induced mutations in mammalian cells. Wives of VC-exposed men have a significantly high risk of fetal loss. Thus, both mutagenic and carcinogenic assays predicted the potential hazards of VC. (34 refs.)

78-4833 **Pharmacokinetics of Vinyl Chloride.** (Eng) Bolt, H. M. (Inst. Toxicology, Univ. Tübingen, Wilhelmstrasse 56, D-7400 Tübingen 1, W. Germany). *Gen Pharmacol* 9(2): 91-95; 1978.

Data on the pharmacokinetic behavior of vinyl chloride (VC), all of which have been obtained from studies with rats, are reviewed. Consistent evidence shows that, above a saturation concentration, which on inhalation exposure is reached at 250 ppm VC, nonlinear (dose-dependent) pharmacokinetics

apply. Two different models describing the pharmacokinetics of VC have been published. They show that VC can leave the body very rapidly via expiration by the lungs. However, special conditions apply if rats are exposed to atmospheric VC: the compound equilibrates with the organism within 15-30 min and is removed from this equilibrium metabolically, probably by formation of the reactive epoxide. The rapid elimination of VC and its major metabolites from the organism agrees with the current theory that a reactive, short-lived metabolite, which occurs in low concentrations only, may be responsible for the toxic effects of VC. (25 refs)

78-4834 **Chloroprene: Observations of Carcinogenesis of Mutagenesis.** (Eng) Infante, P. F. (Industry-wide Studies Branch, Div. Surveillance, Hazard Evaluation and Field Studies, Natl. Inst. Occupational Safety and Health, Cincinnati, OH, 45202); Wagoner, J. K.; Young, R. J. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 205-217; 1977.

Observations of excessive lung and skin cancer associated with chloroprene production were reported in the Russian literature in 1972. However, it was not until 1974, when an epidemic of liver, brain, and lung cancer among vinyl chloride (VC) polymerization workers was identified, was attention in the US focused on the carcinogenic potential of chloroprene. Chloroprene is mutagenic in bacteria, causes recessive lethality in *Drosophila*, causes dominant lethality and chromosome aberrations in rat bone marrow cells, and has been associated with sterility in mice and rats. A significant excess of chromosome aberrations and a reduction in the numbers and motility of sperm have been reported in chloroprene workers, and a threefold excess of miscarriages has been reported among their wives. In the absence of adequate data, a final evaluation of the carcinogenicity of chloroprene cannot be made. However, given its other effects, the possible carcinogenicity of chloroprene may be only of academic interest from a public health point of view. The data for VC and chloroprene may serve as the basis for a more aggressive role by genetic toxicologists in the assay of industrial chemicals. (35 refs.)

78-4835 **Pulmonary Fibrosis and Lung Cancer Caused by Exposure to Pesticides.** (Ger) Barthel, E. (Sitzungsberichte der Akademie der Wissenschaften der DDR, Berlin, E. Germany). *Akademie der Wissenschaften* (2): 33-45; 1978.

Experimental, clinical, and epidemiological data on the association between exposure to pesticides, especially that of

farm workers, and the development of pulmonary fibrosis and lung cancer are reviewed. (48 refs)

- 78-4836 **Underestimation of Chronic Toxicities of Food Additives and Chemicals: The Bias of a Phantom Rule.** (Eng) Kon, S. H. (Henry Dazian Dept. Chemistry, Mount Sinai Hosp., New York, NY, 10029). *Med Hypotheses* 4: 324-339; 1978.

Although it has been assumed that dose-response curves of toxic compounds must never reverse, a literature review of the curves for several food additives revealed toxicity peaks at low doses, often at the lowest dose tested. Similar relationships of max effect at low doses abound in nature. It is suggested that compounds regulated solely on the basis of a progressive dose-response curve may constitute a public health hazard. (71 refs.)

- 78-4837 **Nontoxicity of Mold Cultures Used in Cheese Making.** (Ger) Sieber, R. (Forschungsanstalt für Milchwirtschaft, CH-3097 Liebefeld-Bern, Switzerland). *Z Ernährungswiss* 17(2): 112-123; 1978.

The only carcinogenic mycotoxin discovered in cheese to date has been aflatoxin M₁, which may have originated from the use of contaminated milk. Although feeding pure *Penicillium camemberti* var. *candidum* culture to animals produced tumors attributable to mycotoxins, further studies with various mold culture strains have not confirmed these findings. It is concluded that the use of mold cultures for cheese making does not involve human health risks. (55 refs.)

- 78-4838 **Review of Methods for the Determination of Aflatoxin B, G, and M in Foodstuffs and Feed.** (Ger) Luthy, J. (Institut für Toxikologie der Eidgenössischen Technischen Hochschule, Universität Zürich, Schorenstrasse 16, CH-8603 Schwerzenbach, Switzerland). *Mitt Geb Lebensmittellunters Hyg* 69(2): 200-219; 1978.

The most important methods for analyzing aflatoxin B, G, and M in foodstuffs and feed are reviewed. The topics discussed include the physical-chemical properties of aflatoxins, the analytical methods, collaborative studies with these methods, the identification and analysis of aflatoxin M, and the application of new methods, high-pressure liquid chromatography and radioimmunoassay, for aflatoxin detection. (106 refs.)

- 78-4839 **Field Studies on Liver Cell Cancer.** (Eng) Linsell, C. A. (International Agency Res. Cancer,

69008 Lyon, France); Peers, F. G. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp. 549-556; 1977.

Field studies relating human exposure to aflatoxins and viral hepatitis to an increased risk of liver cell cancer are reviewed. Studies in several African countries and Thailand showed a correlation between aflatoxins in ground nuts, market food samples, and foods ready for ingestion and increased liver cell cancer. The major problem in all these studies was the registration of cancer patients, which made interpretation of the data difficult. The studies suggest a possible dose-response relationship and justify seeking more definitive proof by an intervention program. Hepatitis B has also been suggested as an etiologic agent for liver cancer, the carrier state definitely being associated with this. There is a preponderance of liver cancer in men. It is possible that men metabolize the ultimate carcinogen of aflatoxin less well than women and that men are more susceptible than women to the acute toxicity and chronic carcinogenic effect of aflatoxin. Hepatitis B affects more men than women and the carrier state is more common in men. Liver cancer appears to be an excellent model for study of the interaction between two well-defined factors, one chemical and the other viral. (26 refs.)

- 78-4840 **Mode of Action and Human Health Aspects of Aflatoxin Carcinogenesis.** (Eng) Butler, W. H. (Dept. Histopathology, Saint George's Hosp. Medical Sch., Blackshaw Road, London SW17 0QT, England); Neal, G. E. *Ann Nutr Aliment* 31(4/5/6): 949-956; 1978.

Studies of the mode of action of aflatoxin (AF) carcinogenesis are reviewed. A high incidence of hepatic carcinoma is induced in rats by feeding them 5 ppm AF for 6 wk followed by return to a normal diet. At the end of the 6-wk treatment period, the liver contains multiple foci of cells that are characterized by the presence of starvation-resistant glycogen and a loss of glucose-6-phosphatase activity. These findings are similar to those reported in nitrosamine carcinogenesis. However, the foci in livers of rats fed AF are present throughout the liver, and it is difficult to determine which, if any, of them are related to the final neoplasm or the fate of those that do not develop into carcinoma. If the estimated daily dietary intake of AF is given to rats as a single ip injection, there is a marked inhibition of both RNA and DNA synthesis, with the inhibition of DNA synthesis being considerably greater than that of RNA. This parallels the changes observed during the early stages of the 6-wk feeding cycle. In all studies of experimental hepatocarcinogenesis using AF as the inducing agent, it is apparent that the neoplasm usually arises in a liver that is free of cirrhosis. The experience in humans is that in Europe and North America, most hepatocarcinomas arise in cirrhotic patients. However, in areas where hepatocarcinoma

is a relatively common disease, many such tumors arise in livers showing no cirrhosis. It is possible that in areas with a high incidence of hepatic neoplasia, multiple factors influence the development of the carcinoma. It has been shown, for example, that animals fed low-lipotrope diets have an increased susceptibility to hepatocarcinomas. (23 refs)

78-4841 Occupational and Environmental Standards for Asbestos and Their Relation to Human Disease.

(Eng) Nicholson, W. J. (Mount Sinai Sch. Medicine, City Univ. New York, New York, NY, 10029). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1785-1796; 1977.

The effectiveness of occupational and environmental asbestos standards is considered. The current occupational standard for asbestos is a time-weighted av exposure limit of 2 fibers/ml. Additional provisions for controlling exposures based on work practices are included in this standard, but they are mandatory only if the asbestos air concentration exceeds the established value. Extrapolation from studies on the relationship between lung cancer and exposure of workers in asbestos plants indicates that a continuous 20-yr fiber exposure of 0.4 (less than the proposed Department of Labor standard of 0.5 fibers/ml) could cause a fourfold increase in lung cancer. The ambient-air standard for asbestos simply states that there can be no visible emissions from any facility producing or using asbestos materials. Exposure of the general population occurs from the fireproofing of office buildings by spraying with material containing up to 15% asbestos, a practice no longer permitted. However, the demolition of such buildings, the erosion of fibers from previously applied asbestos, and damage to asbestos during repair work pose serious exposure problems now and in the future. No standards exist for asbestos levels in water supplies, which range from <0.1 to >200 µg/liter. Although no direct epidemiologic evidence links ingestion of asbestos to gastrointestinal cancer, an excess of such tumors has been found among asbestos workers. There are also no appropriate standards for the use and disposal of mining and quarrying material contaminated with asbestos fiber or for the use of asbestos in consumer products. (17 refs.)

78-4842 Cancer Risk of Asbestos Exposure. (Eng) Selikoff, I. J. (Environmental Sciences Lab., Mount Sinai Sch. Medicine, City Univ. New York, New York, NY, 10029). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp., 1765-1784; 1977.

The association between asbestos exposure and cancer is reviewed, and the scope of the problem is considered. Asbestos has been confirmed as a cause of asbestosis, lung cancer, pleural and peritoneal mesothelioma, and gastrointestinal cancer; it may also be related to cancer of the buccal cavity, pharynx, and larynx. Approx 1/5 asbestos workers who were exposed under conditions that existed in the past die of lung cancer. Peritoneal mesotheliomas occur more often than pleural mesotheliomas in asbestos workers, but pleural mesotheliomas are more common in cases of environmental, family contact, or neighborhood exposures. The long clinical latency period in asbestos-related disease has made it difficult to evaluate the quantitative relationship between the amount of exposure and the subsequent risk. However, there appears to be a dose-disease-response gradient for asbestos, with disease much more likely to occur as a result of heavier exposure. Based on studies of the time of death from lung cancer and pleural and peritoneal mesotheliomas in relation to the elapsed time from onset of first exposure, serious risk begins > 20 yr after onset of first exposure. Cigarette smoking increases the risk of lung cancer in asbestos workers, but it is not associated with pleural nor peritoneal mesothelioma. Cigarette-smoking asbestos workers are also at risk for esophageal cancer but not gastrointestinal cancer. From the public health perspective, low-level environmental exposure and end-product contamination are more widespread problems than occupational exposure. (45 refs)

78-4843 Possible Effects of Drugs on the Occurrence of Malignant Neoplasms. (Cro) Cicek, J. (Nazorova 37a, 41000 Zagreb, Yugoslavia). *Libri Oncol* 6(3/4): 37-45; 1977.

Radioisotopes, antilymphocyte serum, antimetabolites, chloronaphazine, melphalan, cyclophosphamide, synthetic estrogens, androgens, arsenic compounds, phenacetin, tar-base ointments, and, possibly, diphenylhydantoin, chloramphenicol, amphetamine, and reserpine are cancer-causing drugs in humans. (59 refs)

78-4844 Metronidazole, Isoniazid, and the Threat of Human Cancer. (Eng) Goldman, P. (Clinical Pharmacology Div., Dept. Pharmacology, Harvard Medical Sch., Boston, MA, 02115); Ingelfinger, J. A.; Friedman, P. A. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 465-474; 1977.

The benefits and cancer-causing potentials of metronidazole, an antibiotic drug used to treat trichomonal vaginitis, are

discussed. The effectiveness of the drug depends on the dose given, the dose currently approved by the Food and Drug Administration being 250 mg tid for 10 days. This dose, or even a dose of 250 mg tid for 7 days, may be unnecessarily high, since total doses as low as 2 g have been successful. There have been no direct estimates of human cancer risk associated with metronidazole ingestion, but the drug causes an increased incidence of pulmonary adenomas in mice. Metronidazole also has chemical and biological properties that are often associated with known carcinogens. On the basis of the laboratory evidence, there is no reason to suspect that metronidazole is a more potent carcinogen than isoniazid. The latter has been tested directly in humans and has not been associated with a large increase in cancer within the 15 yr following treatment. However, in the absence of a reliable method for extrapolating from laboratory data to human risk, it would be unwise to conclude that metronidazole is no more risky than isoniazid. Thus, metronidazole should be given only in the necessary dose to patients with symptomatic trichomonal vaginitis. (32 refs)

- 78-4845 Hycanthone and Other Antischistosomal Drugs: Lack of Obligatory Association Between Chemotherapeutic Effects and Mutagenic Activity.** (Eng) Bueding, E. (Dept. Pathobiology, Sch. Hygiene and Public Health, Johns Hopkins Univ., Baltimore, MD, 21205); Batzinger, R. P. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 445-463; 1977.

The association between the schistosomicidal and mutagenic activities of antischistosomal drugs is discussed. Hycanthone (HCT) is a potent mutagen for bacteria, yeast, and mammalian cells and it is carcinogenic in mice, particularly in those infected with *Schistosoma mansoni*. The nitroheterocyclic schistosomicide niridazole is also mutagenic and carcinogenic, and the mutagenic activity of furapromidium is of the same order of magnitude as that of niridazole (its carcinogenicity has not been tested). The mutagenic activities of two other drugs, metrifonate and oxamniquine, are several orders of magnitude lower than that of HCT. The antischistosomal potencies of HCT and of a chloroindazole analog of HCT are equal, but the mutagenic and carcinogenic potencies of the analog are much lower. N-oxidation of the analog results in the potent schistosomicide IA-4 N-oxide, whose mutagenic potency is even weaker and whose LD₅₀ is 12 times higher than that of HCT. A new antischistosomal compound, 4-isothiocyanato-4'-nitrodiphenylamine, is mutagenic in host-mediated assays, but if it is given after the host's intestinal bacterial flora has been reduced by antibacterial agents, it is nonmutagenic. Thus, schistosomicidal and mutagenic activities can be completely dissociated from each other. (83 refs)

- 78-4846 Schistosomiasis and Neoplasia.** (Eng) Cheever, A. W. (Lab. Parasitic Diseases, Natl. Inst. Allergy and Infectious Diseases, NIH, Public Health Service, U.S. Dept. Health, Education, and Welfare, Bethesda, MD, 20014). *J Natl Cancer Inst* 61(1): 13-18; 1978.

The association between schistosomiasis and cancer of the urinary bladder, hepatocellular carcinoma, lymphoma and leukemia, and carcinoma of the colon is considered. Rigorous statistical proof of the association between *Schistosoma haematobium* infection and bladder cancer has not been obtained, but most published data tend to confirm the association. Etiologic factors in schistosomal bladder cancer may be chronic inflammation of the bladder and urinary retention, carcinogenic substances in the schistosome in the schistosome eggs, effects of schistosome infection on the immune system, the interaction of bacteria in the urine with excreted substances, and the strain of schistosome. Hepatomas were equally frequent in uninfected and schistosome-infected patients in autopsy series from several countries. In Japan, both hepatoma and circulating hepatitis B surface antigen were significantly more frequent in autopsied patients with *S. japonicum* than in those without, but most autopsied patients with schistosomiasis and hepatoma also had posthepatitis cirrhosis. Equivocal increases in the frequency of hepatoma have been reported in animals infected with *S. mansoni*. An association between giant follicular lymphoma of the spleen and hepatosplenic schistosomiasis was found in Brazil. Several autopsy series suggest an association of either lymphoma, leukemia, or both with schistosomiasis. Cancers of the colon, ovaries, uterus, and other organs have been attributed to schistosomiasis, but no evidence of more than a coincidental association has been provided. (81 refs)

- 78-4847 Carcinogenic Hazards of Organic Chemicals in Drinking Water.** (Eng) Harris, R. H. (Environmental Defense Fund, Washington, DC, 20036); Page, T.; Reiches, N. A. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 309-330; 1977.

Studies dealing with elevated cancer mortality rates resulting from the presence of carcinogens in the drinking water are reviewed. Five main topics are discussed: (1) organic carcinogens in drinking water, (2) approaches to assessing the potential health consequences of these chemicals, (3) recent epidemiologic studies, (4) statistical problems in relating exposure to water pollutants to cancer mortality, and (5) the application of the results of these epidemiologic studies to assessment of the risk of low-level carcinogenic exposure. Studies indicate that although a wide variety of chemical carcinogens are present in the drinking water, their concentrations are low. However, epidemiologic studies in Louisiana,

Ohio, and New Jersey support the hypothesis that carcinogens in drinking water are related to human cancer. Since these findings reflect exposure to carcinogen levels present 10-30 yr ago and present-day levels are assumed to be higher, more noticeable effects may be expected in the future. (38 refs.)

78-4848 Potentially Carcinogenic Species Emitted to the Atmosphere by Fossil-fueled Power Plants.

(Eng) Natusch, D. F. (Dept. Chemistry, Colorado State Univ., Fort Collins, CO, 80523). *Environ Health Perspect* 22: 9-90; 1978.

The identities and physicochemical characteristics of potential carcinogens emitted by fossil-fueled power plants are presented. Many carcinogens are preferentially concentrated on the surface of respirable fly ash particles, enabling them to come in contact with lung tissue when inhaled. There is little information on the identities of particulate polycyclic organic compounds emitted from the plants. The chemical changes that these compounds may undergo following emission but before inhalation are important in determining their potential carcinogenicity. (34 refs)

78-4849 Trends in Urban Air Pollution in the United Kingdom in Relation to Lung Cancer Mortality.

(Eng) Lawther, P. J. (MRC Environmental Hazards Unit, St. Bartholomew's Hosp. Medical Coll., Charterhouse Square, London EC1M 6BQ, England); Waller, R. E. *Environ Health Perspect* 22: 71-73; 1978.

The possible role of atmospheric carcinogens in lung cancer is considered in view of the substantial reduction in air pollution, particularly in components such as benzo(a)pyrene (BP), in urban areas of the United Kingdom during the past few decades. Although overall trends in lung cancer mortality have been influenced largely by changes in smoking habits since the late 1930's, the marked contrasts that once existed between lung cancer death rates in urban and rural areas have gradually diminished. Thus, air pollution may have contributed to the urban-rural differences in lung cancer before the Clean Air Act of 1956. In London, BP concentrations are now one-tenth of those in the period 1949-1951. (5 refs)

78-4850 Are Chemicals the Main Cause of Cancer? Fateful Change Processes in the Body--Mainz Scientists Examine the Role of Benzpyrene from Waste Gases and Smoke. (Eng) Zimmermann, D. (No affiliation given). *Dtsch Forschung* 17(4): 1-3; 1978.

Studies of the carcinogenicity of benzo(a)pyrene (BP) suggest

that the compound is fat-soluble and, therefore, accumulates in the body where it is transformed by certain enzymes into a highly reactive epoxide that can disrupt cellular processes and bind to nucleic acids. It is possible that an antienzyme, which can prevent the formation of this epoxide, may be able to suppress the carcinogenicity of BP. (no refs.)

78-4851 Some Recent Findings Concerning Cigarette Smoking. (Eng) Hammond, E. C. (Dept.

Epidemiology and Statistical Res., American Cancer Society, New York, NY, 10017); Garfinkel, L.; Seidman, H.; Lew, E. A. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 101-112; 1977.

Some recent findings concerning the effects of cigarette smoking are discussed. Death rates are higher in smokers than in nonsmokers and they increase with the number of cigarettes smoked per day and with the number of years spent smoking. A prospective study of > 1,000,000 subjects was carried out between 1959 and 1965, and the 897,825 subjects who were still alive in 1965 were studied until 1971 or 1972. For both men and women for both study periods, the adjusted number of deaths and the adjusted number of deaths from lung cancer were lowest among smokers of low tar and nicotine (T/N) cigarettes and highest among high T/N smokers. The adjusted number of lung cancer deaths was greater in subjects who smoked 20-39 low T/N cigarettes/day than in subjects who smoked 1-19 high T/N cigarettes/day. The overall death rate and the lung cancer death rate were far higher in subjects who smoked low T/N cigarettes than in subjects who never smoked regularly. Thus, reduction of the T/N content of cigarette smoke did not make cigarette smoking safe, although switching from high T/N to low T/N cigarettes would appear to be a small step in the right direction. (7 refs)

78-4852 The Genetic Expression of Aryl Hydrocarbon Hydroxylase Activity: Evidence for a Receptor Mutation in Nonresponsive Mice. (Eng)

Poland, A. (Dept. Pharmacology and Toxicology, Univ. Rochester Sch. Medicine and Dentistry, Univ. Rochester, Rochester, NY, 14642); Kende, A. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 847-867; 1977.

The nature of the aryl hydrocarbon (*Ah*) locus in the mouse was determined. Previous studies have shown that the *Ah* locus determines the expression of aryl hydrocarbon hydroxylase (AHH) activity and cytochrome P₁-450; they also implicate the *Ah* locus in mediating the mutagenesis and car-

cinogenesis produced by certain polycyclic aromatic hydrocarbons (PAH). 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) was used to determine the nature of the *Ah* locus and how PAH stimulate AHH activity. TCDD was a potent inducer of hepatic AHH activity and cytochrome P₁-450 in rats and in nonresponsive mice, which were not responsive to the inductive activity of 3-methylcholanthrene. Experimental evidence indicates that the hepatic cytosol from C57BL/6J mice contains a binding species with a high affinity for TCDD, that the binding properties of this cytosol species correspond extremely well to the properties predicted for the induction receptor, and that the mutation in nonresponsive strains of mice appears to result in a receptor with a diminished affinity for inducing ligands. It is postulated that the *Ah* locus codes for the cytosol binding protein, which serves as a cellular recognition site for PAH, TCDD, and similar compounds, and that the binding of these inducers to the receptor in some manner serves as a master switch to initiate the expression of numerous noncontiguous structural genes. (43 refs.)

78-4853 Aryl Hydrocarbon Hydroxylase and Bronchogenic Carcinomas Associated with Smoking (Letter to Editor). (Eng) Korsgaard, R. (Res. Dept. I, Univ. Hosp., Wallenberg Lab., Univ. Lund, S-220 07 Lund, Sweden); Trell, E. *Lancet* 1(8073): 1103-1104; 1978.

Studies have confirmed the finding that aryl hydrocarbon hydroxylase (AHH) is the major activator of polycyclic aromatic hydrocarbon carcinogens in patients with squamous cell carcinoma of the lung. Variations in the assessment of AHH inducibility can be minimized by using T- and B-cell mitogens, equal numbers of stimulated lymphocytes for test and control studies, and by excluding any enzyme-inducing capacity from the media. (10 refs)

78-4854 Properties of the Liver Microsomal Monooxygenase System and Epoxide Hydrase: Factors Influencing the Metabolism and Mutagenicity of Benzo[a]pyrene. (Eng) Levin, W. (Dept. Biochemistry and Drug Metabolism, Hoffman-La Roche, Inc., Nutley, NJ, 07110); Lu, A. Y.; Ryan, D.; Wood, A. W.; Kapitulnik, J.; West, S.; Huang, M. T.; Conney, A. H.; Thakker, D. R.; Holder, G.; Yagi, H.; Jerina, D. M. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 659-682; 1977.

The metabolism of benzo(a)pyrene (BP) by the microsomal monooxygenase systems and epoxide hydrases of animal and human tissue is reviewed. The liver microsomal metabolism of BP varied with different enzyme inducers, animal species,

and strains of mice. In addition, the metabolism of BP by lung microsomes from 3-methylcholanthrene-treated mice differed from that by liver microsomes from the same animals. Purified cytochrome P-450 from different animal species treated with different inducers also metabolized BP at significantly different rates. In studies of the metabolism of BP arene oxides by microsomal epoxide hydrase, BP 11,12-oxide differed from the other arene oxides of BP (4,5-, 7,8-, and 9,10-oxides) in being a poor substrate for the enzyme. The dihydrodiols and phenolic metabolites of BP shared arene oxides as common precursors. Epoxide hydrase was shown to participate in both the metabolic activation and detoxification of BP by the monooxygenase system. Data indicated that the human liver contains either a single epoxide hydrase with broad substrate specificity or several epoxide hydrases under similar regulatory control. There were differences between mice and humans in the regulatory systems that control the metabolism of BP, 7-ethoxycoumarin, and 2-acetylaminofluorene. (48 refs.)

78-4855 Bay-region Epoxides of Dihydrodiols: A Concept Explaining the Mutagenic and Carcinogenic Activity of Benzo[a]pyrene and Benzo[a]anthracene. (Eng) Jerina, D. M. (Section on Oxidation Mechanisms, Natl. Inst. Arthritis, Metabolism and Digestive Diseases, NIH, Bethesda, MD, 20014); Lehr, R.; Schaefer-Ridder, M.; Yagi, H.; Karle, J. M.; Thakker, D. R.; Wood, A. W.; Lu, A. Y.; Ryan, D.; West, S.; Levin, W.; Conney, A. H. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 639-658; 1977.

In order to generalize the concept of diol epoxides as ultimate mutagenic and carcinogenic metabolites, the ability of a reconstituted monooxygenase system containing cytochrome P-448 to activate dihydrodiols of benzo(a)pyrene (BP) and benz(a)anthracene (BA) to mutagenic forms in the presence of a histidine-dependent strain of *Salmonella typhimurium* (TA98) was studied. The results of these studies support the bay-region theory; ie, that bay-region epoxides of dihydrodiols on an angular benzo ring would be good candidates for the ultimate carcinogenic forms of polycyclic aromatic hydrocarbons. The BA 3,4-diol-1,2-epoxides, which contain a bay-region epoxide, were more mutagenic than the other diol epoxides of BA. However, naphthalene, anthracene, and phenanthrene and their trans dihydrodiols could not be activated significantly to mutagens, and the naphthalene 1,2-diol-3,4-epoxides were virtually inactive. Perturbational molecular orbital calculations predict that benzylic carbonium ions at benzylic positions of saturated benzo rings of these small aromatic hydrocarbons are much less easily formed than those from larger hydrocarbons such as BP. The carcinogenicity of BP 7,8-oxide and 7,8-dihydrodiol, the high mutagenicity, upon metabolic activation, of the 7,8-dihy-

diol compared with other BP derivatives, and the high mutagenicity of the 7,8-diol-9,10-epoxides indicate the importance of the bay region for the biological activity of BP. However, other examples of bay-region epoxides of benzo[a]anthracene dihydrodiols will have to be examined to establish whether the bay-region theory has general validity. (71 refs.)

78-4856 The Workplace as a Cause of Cancer--Part I. (Eng) Schottenfeld, D. (Epidemiology and Preventive Medicine Service, Dept. Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY); Haas, J. F. *Ann Intern Med* 8(2): 54-60; 1978.

Epidemiologic methods used to determine the risks of cancer among workers exposed to suspect industrial chemicals are reviewed, and a historical overview of occupational carcinogenesis is presented. (38 refs.)

78-4857 Is There a Health Hazard in Anaesthetic Practice? (Eng) Spence, A. A. (Dept. Anaesthesia, Western Infirmary, Glasgow G11 6NT, Scotland); Snell-Jones, R. P. *Br J Anaesth* 50(7): 713-719; 1978.

A review of health hazards associated with anesthesia practice indicate that women have an increased risk of spontaneous abortion and that both sexes have an increased risk of hepatitis, but there is no evidence that operating room workers are at risk for any other disease. However, it is suggested that the incidence of cancer, particularly that of leukemia and lymphoma, be monitored in women. (59 refs.)

78-4858 Trace Concentrations of Anesthetic Gases: A Critical Review of Their Disease Potential. (Eng) Ferstandig, L. L. (Halocarbon Labs., Inc., 82 Burlew's Court, Hackensack, NJ, 07601). *Anesth Analg (Cleve)* 57(3): 328-345; 1978.

A review of the histotoxic effects of anesthetics indicates that only high exposure levels produce harmful effects in cells and animals. None of the common halogenated anesthetics produce cancer in animals, and epidemiological studies show no association between exposure and cancer in men and only a dubious association in women. Studies of the effect of anesthetics on reproductive function and disease are inconclusive. (126 refs)

78-4859 Cosmetic Science--1977. (Eng) Rieger, M. M. (Warner Lambert Co., Morris Plains, NJ). *Cosmet Toiletries* 93(5): 25-52; 1978.

Progress in surfactant chemistry during 1977 is reviewed along with recent findings on the carcinogenicity of chemicals used in cosmetics. Recent studies have contradicted earlier

reports of an association between breast cancer and exposure of hairdressers to hair dyes. Animal studies indicate that neoplasms could result from the intravesicular implantation of foreign bodies alone, suggesting that the results of animal carcinogenicity tests should be subject to careful scrutiny. (190 refs)

78-4860 Prevention of Epithelial Cancer by Vitamin A and Its Synthetic Analogs (Retinoids). (Eng) Sporn, M. B. (Lung Cancer Branch, NCI, Bethesda, MD, 20014). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 801-807; 1977.

There are two general approaches to the prevention of epithelial cancer: inhibition of initiation and inhibition of progression. The use of vitamin A and its synthetic analogs (collectively called retinoids) in the latter context is discussed. Retinoids have been shown to prevent the development of cancer in the skin, respiratory tract, mammary gland, and urinary bladder of experimental animals. There is an excellent rationale for using these agents to prevent cancer by the enhancement of intrinsic epithelial defense mechanisms. The application of these agents for the prevention of cancer in high-risk human populations is thus worthy of consideration. Future developments in the synthetic chemistry of retinoids should make new structures available that will be even more specific, more potent, and less toxic for practical cancer prevention. (26 refs.)

78-4861 Kinetics of Tumor Formation and Low Risk Assessment. (Eng) Scherer, E. (Div. Chemical Carcinogenesis, Antoni Van Leeuwenhoek-Huis, Netherlands Cancer Inst., Amsterdam, Netherlands); Emmelot, P. In: *Carcinogens: Identification and Mechanisms of Action. Proceedings of the Thirty-First Annual Symposium on Fundamental Cancer Research Held at Houston, Texas, 1-3 March 1978*. University of Texas System Cancer Center, M. D. Anderson Hospital and Tumor Institute, NCI, American Cancer Society. (Austin, Texas). 22-23 pp.; 1978. (no refs)

78-4862 The Role of Diet in Human Carcinogenesis with Special Reference to Endometrial Cancer. (Eng) Armstrong, B. K. (Dept. Medicine, Univ. Western Australia, Nedlands, Western Australia 6009, Australia). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 557-565; 1977.

Possible mechanisms for the role of diet in human carcino-

genesis are: (1) ingestion of carcinogens or carcinogen precursors, as exemplified by the positive relationship between aflatoxin ingestion and hepatocellular cancer; (2) effects on the production, activation, or inactivation of carcinogens by the intestinal flora; (3) effects on the endogenous production, activation, or inactivation of carcinogens, as exemplified by the possible modifying effect of dietary protein on the carcinogenicity of ingested nitrosamines; and (4) alterations of tissue susceptibility to carcinogenesis. The latter mechanism may be exemplified by the relationship between the incidence of endometrial cancer and dietary fat or total dietary calories. The following factors are associated with a high risk of endometrial cancer: obesity, early menarche, late menopause, diabetes mellitus, hypertension, infertility, Stein-Leventhal syndrome, previous breast or ovarian cancer, radiation exposure, and estrogen therapy. The first five may stem from a dietary excess of calories or fat, resulting in the excessive endogenous production of estrogens. Although the precise role of estrogens in the genesis of endometrial cancer is uncertain, recent studies indicate that they are probably involved. (53 refs)

- 78-4863 **The Estrogen-Cancer Hypothesis with Special Emphasis on DES.** (Eng) Hertz, R. (No affiliation given). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory: Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1665-1673; 1977.

The genesis of the estrogen-cancer hypothesis, which postulates that both exogenous and endogenous estrogens and their metabolites play a significant role in the pathogenesis of cancer in estrogen-responsive tissues of humans and animals, is reviewed. Numerous studies indicate that estrogen administration is followed by routinely reproducible tumors of the breast, cervix, endometrium, ovary, pituitary, testicle, kidney, and bone marrow in mice, rats, rabbits, hamsters, or dogs. Malignant uterine mesotheliomas have been induced in squirrel monkeys by estrogen administration. A definitive association has been established between maternal exposure to estrogen (diethylstilbestrol) during pregnancy and the development of vaginal adenocarcinomas in their daughters. There is also impressive evidence that endogenous estrogen is related to the pathogenesis of breast and endometrial cancers. The association of endometrial cancer with excessive endogenous estrogen, as in the case of women with Stein-Leventhal disease, suggests a chronic stimulatory effect on this responsive tissue. The effectiveness of progesterone, which directly antagonizes the mitosis-stimulating action of estrogen, in inducing regression of both endometrial carcinoma and endometrial adenomatous hyperplasia also implicates endogenous estrogen as a factor in endometrial cancer. Evaluation of epidemiological data is hampered by a lack of precision in the pathological diagnosis of endometrial cancer, so caution in interpretation is indicated. (55 refs)

- 78-4864 **Hepatic Tumors and Oral Contraceptives.** (Eng) Fechner, R. E. (No affiliation given). *Pathology Annual* 12(part 2): 293-310; 1977.

Clinical and experimental data linking ingestion of oral contraceptives (OC) with hepatic tumors are reviewed. Comparison of the incidence of focal nodular hyperplasia before and after 1960 (the year OC received Federal Drug Administration approval) disclosed nothing to suggest that focal nodular hyperplasia has increased in frequency or is associated with OC. However, evidence suggests that these lesions may be more prone to rupture in women taking OC. Liver cell adenomas, on the other hand, appear to have increased in frequency since 1960, and the tumors in women taking OC have a tendency to be slightly larger. These tumors have occurred almost exclusively in young women, most of whom had been taking OC for at least 3 yr. A high percentage of the latter used a preparation containing mestranol. The results of ultrastructural and experimental studies are reviewed, along with the occurrence of vascular alterations in women taking OC. (116 refs.)

- 78-4865 **Liver Adenomas and Oral Contraceptives.** (Eng) Pike, M. C. (Dept. Community Medicine, Univ. Southern California, Los Angeles, CA, 90033); Edmondson, H. A.; Benton, B.; Henderson, B. E. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory: Origins of Human Cancer. Vol. 4, 602 pp.; 423-427; 1977.

The association between liver adenomas and oral contraceptives (OC) was investigated in a case-control study of 34 adenoma patients. The frequency of OC use was similar in patients (29/34) and controls (26/34). However, the risk of liver adenomas was clearly associated with duration of use. Of the 29 users, 26 were taking them at the time of diagnosis and 2 others had stopped taking them <2 mo earlier. Twenty-four used mestranol-containing pills only, vs 15 of the controls. (9 refs)

- 78-4866 **Effects of Oral Contraceptives on Carcinogenesis in the Cervical Epithelium.** (Ger) Soost, H. J. (Institut für Klinische Zytologie der Technischen Universität München, Prinzregentenplatz 14, D-8000 Munich 80, W. Germany); Joswig-Priewe, H. *Arch Geschwulstforsch* 48(4): 345-355; 1978.

Data on the hormonal action of oral contraceptives (OC's) are reviewed. Cytophotometric investigations showed that the DNA levels in uterine cervix cells from women with a long history of OC use were normal. Statistical comparisons, based on cytological and histochemical examinations, among

the women according to type of OC formulation used indicated that some formulations produced high rates of cervical atypia, others low rates of atypia. Only a few women showed a higher rate of dysplasia and carcinoma in situ of the cervix than that in controls. Pinpointing OC's as a cause of cervical carcinogenesis is difficult because of the influence of other factors, especially a change in sexual behavior, on tumor development. At present, OC use cannot be directly related to the increased occurrence of cervical atypia. (72 refs.)

- 78-4867 **Exogenous Estrogens and the Incidence of Neoplasms in Tissues of Mullerian Origin.** (Eng) Weiss, N. S. (Dept. Epidemiology, Sch. Public Health and Community Medicine, Univ. Washington, Seattle, WA, 98195). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 413-421; 1977.

Trends in the incidence rates of cancers of mullerian-derived tissues are discussed. In the 1970's, there were sharp increases in endometrial cancer incidence in a number of areas in the US. These increases closely followed the beginning of a period of rapidly rising consumption of estrogen preparations by menopausal women. Endometrial cancer is more prevalent among women with high circulating levels of endogenous estrogens, and recent studies have shown increased rates among menopausal women using exogenous estrogens and women receiving stilbestrol. The nature of many of the lesions being produced by estrogen use remains unknown, but it is likely that menopausal estrogens play an etiologic role in endometrial cancer. Relative frequencies and incidence rates of endometrioid-clear cell tumors in three areas of the US increased between 1969 and 1974; these increases also followed increased estrogen consumption. The incidence of other epithelial ovarian tumors has not changed appreciably since the late 1940's. The relationship of contraceptive estrogens to endometrial cancer or any of the estrogens to the occurrence of endometrioid and clear cell ovarian tumors has not been established. (27 refs)

- 78-4868 **How Safe Is Nuclear Power (Letter to Editor)?** (Eng) Diesendorf, M. (20 Earle St., Lyneham, Australian Capital Territory 2602, Australia). *Med J Aust* 1(3): 162-163; 1978.

Comparisons of the health hazards of plutonium and radium are meaningful only for bone cancer, and bone should not be assumed to be the critical organ for Pu. ^{239}Pu produced lung cancers in dogs exposed to 0.2-1.0 μCi . In bone cancer experiments, ^{239}Pu was 5-10 times more toxic than ^{226}Ra . Ra is deposited uniformly throughout the bone, but Pu accumulates on the bone surface, where cells at highest risk for tumors are located. Human epidemiological investigations have

also implicated Pu in several cancers. Some human health hazard exists at every stage of Pu processing. (11 refs)

- 78-4869 **The Real Death Rays.** (Eng) Saldinger, E. (Calhoun Coll.). *Yale Scientific* 52(4): 16-21; 1978.

The various biological effects of different types of radiation are surveyed. The long-term effects of radiation were studied in Japanese survivors of the Hiroshima and Nagasaki bombings. The Hiroshima group showed a high incidence of leukemia that peaked at 30 times the av incidence. It usually appeared about 10 yr postirradiation. In addition, the incidence of cancer of the stomach, lung, breast, cervix, and ovary in these subjects was two to four times the av rate. Background radiation, ie, UV rays and cosmic rays, is associated with an increased incidence of skin cancer. Therapeutic radiation can be damaging to fetuses and to patients. Leukemia and thyroid cancer have occurred in people irradiated as children for enlarged thymus glands. Moderate amounts of radiation may cause premature aging and a shortened life-span. Ion production and the amount of energy absorbed by different tissues, six types of ionizing radiation with potential to destroy living tissue, and the molecular effects of radiation are also reviewed, along with the effects of radiation at the cellular level. Lymphocytes show high sensitivity to radiation, and in Hiroshima and Nagasaki the lymphocyte level in the population only returned to normal 3-4 yr after the bombing. (no refs)

- 78-4870 **Carcinogenic Effects of Medical Irradiation.** (Eng) Hutchison, G. B. (Dept. Epidemiology, Harvard Sch. Public Health, Boston, MA, 02115). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 501-509; 1977.

The carcinogenic risk to human populations associated with medical irradiation is discussed. Epidemiologic studies of several types of cancer that have developed after medical irradiation are reviewed. Ionizing radiation at any but the highest levels of exposure appears to be capable of inducing malignant transformation. Empiric approaches cannot identify an absolutely safe dosage level, although the hazard at arbitrarily small doses may be expressed in terms of probability concepts. Simple linear models of radiation effects give satisfactory fits to most human data, but linear interpolation on dose may systematically overestimate effects at doses ≥ 10 rads. Most observations of radiation effects in human populations involve high doses, high dose rates, and low-linear energy transfer (LET) radiation; and with this type of exposure, departure from linearity is expected. Less extensive data are available on high-LET exposure at high dose levels, and lin-

ear interpolation on these data may give adequate estimates of low dose effects at high LET. Generally, it will not be satisfactory to determine the relative effects of high- and low-LET radiations at high dose rates and employ this relation in estimating effects of low-LET radiation at low doses. Because of the nonlinearity of effect with low-LET radiation, this procedure will systematically overestimate the effect. (25 refs.)

- 78-4871 **The Risks for Radiation Workers.** (Eng) Rotblat, J. (Dept. Physics, Univ. London, London, England). *Bull At Sci* 34(7): 41-46; 1978.

The fact that, in 1977, the International Commission on Radiological Protection did not find it necessary to reduce the max permissible dose limit for radiation workers [5 roentgens-equivalent-man (rem)/yr] is discussed. The survivors of the Hiroshima and Nagasaki bombings are generally assumed to be the most reliable material for determining a reasonable estimate of the risk factor for radiation-induced cancer. However, for all types of cancer except leukemia, the risk factors calculated from the Japanese survivors are significantly lower than those from other populations. The survivors may not be suitable, therefore, as a basis for calculating radiation risk factors for other populations. There appears to be a need to revise upward the risk factors for cancer induction, and it is suggested that the present dose limit for radiation workers be reduced by a factor of 5. Furthermore, a worker who receives more than 1 rem/yr should be eligible for compensation, should he incur a disease that might be attributable to his exposure to radiation. (10 refs.)

- 78-4872 **Cancer and Low Level Ionizing Radiation.** (Eng) Morgan, K. Z. (Sch. Engineering, Georgia Inst. Technology, Atlanta, GA). *Bull At Sci* 34(7): 30-41; 1978.

The risk of cancer from low-level exposure to ionizing radiation is discussed. Since 1960, an overwhelming amount of data have shown that there is no safe level of exposure and that there is no dose of radiation so low that the risk of malignancy is zero. The risk of cancer induction from radiation exposure increases more or less with the increase or accumulation of radiation exposure. Because of many individual characteristics (eg, genetic inheritance, sex, disease, and eating habits), certain members of the general population have a higher risk of radiation-induced malignancies than others. The risk of cancer induction at low doses of radiation may be as great or greater for the human race than genetic risk. Several reasons why, in many cases, use of the linear hypothesis to estimate risk at low doses is not conservative are given. The medical profession and not the nuclear industry is the real culprit with respect to unnecessary exposure of the population to radiation. It is suggested that the max

permissible exposure should be reduced by a factor of 2 and, perhaps later, by a factor of 10 if it can be shown that all unnecessary exposure (especially medical) can be reduced and that there will be a net benefit to mankind by such action (27 refs.)

- 78-4873 **Radiation Effects.** (Eng) Upton, A. C. (Dept. Pathology, State Univ. New York, Stony Brook, NY, 11594). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 477-500; 1977.

An association between radiation and neoplasia has been detected in human populations at lower levels of exposure than heretofore suspected. However, the dose-incidence relationship cannot be specified precisely for any neoplasm, and the relation may vary depending on factors such as the type of neoplasm and age at irradiation. Susceptibility varies widely among different organs and under the influence of genetic and physiological factors. The interaction of constitutional, environmental, and radiologic variables influences the dose-incidence relationship in laboratory animals. Although the pathogenesis of radiation-induced neoplasms is poorly understood, a variety of mechanisms has been postulated, including somatic mutation, virus activation, and disturbances in homeostasis. A succession of changes requiring a relatively long latent period seems to be involved in many cases. Because quantitative assessment of the risk of radiation-induced cancer in humans must be based primarily on human dose-incidence data, epidemiologic follow-up study of irradiated human populations remains essential. (88 refs.)

- 78-4874 **Epidemiology of Radiation-induced Malignant Human Tumors.** (Rus) Strel'tsova, V. N. (No affiliation given). *Med Radiol (Mosk)* 23(8): 83-90; 1978.

Epidemiological data on malignant human tumors induced by ionizing radiation are reviewed. The bone marrow, thyroid gland, lymph nodes, reticular tissue, pharynx, bronchi, pancreas, and stomach are most susceptible to the carcinogenic effects of such radiation. (23 refs.)

- 78-4875 **Radiation Carcinogenesis In Vitro: Implications for Mechanisms.** (Eng) Little, J. B. (Dept. Physiology, Harvard Univ. Sch. Public Health, Boston, MA, 02115). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Har-

or: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 923-939; 1977.

Data on the induction of malignant transformation in C3H mouse embryo fibroblast cells by x-rays and UV light are reviewed. X-ray transformation can be divided into two steps: (1) fixation of the transformed state as a heritable cellular property, and (2) expression of this damage in terms of morphologically altered cells. The transformation frequency rises rapidly with increasing radiation dose in the range below 400 rads; at this point it reaches a plateau that remains constant up to 1,500 rads. The dose-response curve for transformation induction by UV light is similar to that for x-rays, with an inflection point at 75 ergs/mm². If transformation frequency is plotted as a function of time after irradiation, when the cultures are dispersed and seeded at low cell density, the transformation frequency per viable cell rises rapidly, reaching a max with a repair time of 3 hr. With longer repair intervals, the transformation frequency falls and is very low or not detectable if the cells are not stimulated to divide within 48 hr of irradiation. Expression of the transformed state requires about 12 additional rounds of cell division. Exposure during this expression period to a number of noncarcinogenic agents may either greatly enhance (12-O-tetradecanoylphorbol-13-acetate or interferon) or markedly suppress (Amphotericin-B) transformation. It is concluded that appropriate stimuli and capacity for cell proliferation in irradiated cell populations are major determinants in the fixation and expression of transformational damage and that the ultimate carcinogenic effect of radiation depends largely on factors secondary to the initial DNA damage. (32 refs)

78-4876 **Melanoma and Solar Radiation.** (Ger) Herrmann, W. P. (Universitäts-Hautklinik, Joseph-Stelzmann-Strasse 9, 5000 Köln 41, W.Germany). *Dtsch Med Wochenschr* 103(29): 1155-1156; 1978.

Epidemiological studies not only suggest a relationship between exposure to sunlight and the incidence of skin melanoma, but show a genetic predisposition as well. Ultraviolet radiation is a possible cofactor in the genesis of malignant melanoma. (14 refs.)

78-4877 **Ultraviolet Radiation as a Cause of Cancer.** (Eng) Scott, E. L. (Statistical Lab., Univ. California, Berkeley, CA, 94720); Straf, M. L. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 529-546; 1977.

The role of UV radiation in the production of human skin cancer is reviewed. There is a strong association between UV

flux and the incidence and mortality rates of skin cancer (both melanoma and nonmelanoma) for both sexes and all age groups. The carcinogenesis of skin cancer is uncertain, however, and it may be complicated by a complex UV repair mechanism. The dose-response relationship is not known. In principle, it can be estimated from the data on skin cancer by sex and age for different localities where the UV flux is known, but available observations are unreliable and incomplete. Estimates of the expected rate of skin cancer corresponding to a 20% increase in UV flux or a 10% depletion of stratospheric ozone indicated that there is a strong effect of UV flux and that it persists even when other possible predictors are taken into account. The numerical value of the estimated increase in melanoma was frequently greater than that for nonmelanoma. Although both temperature and pigmentation play a role in the etiology of skin cancer, UV flux is the most important factor. Corresponding calculations for cancer of the eye and lip are less reliable due to a sparsity of data. (37 refs.)

78-4878 **Radiation Induced Leukemia and Leukopenia in the Lizard *Uromastix hardwickii* (Preliminary Information).** (Eng) Ahmad, M. (Dept. Physiology, Univ. Karachi, Karachi-32, Pakistan); Taqawi, I. H. *Radiobiol Radiother (Berl)* 19(3): 353-360; 1978.

Spiny-tailed lizards (*Uromastix hardwickii*) were exposed to 0, 3,000, 3,500, or 4,000 R of ⁶⁰Co-γ irradiation, and their WBC count was monitored for 6 wk. Lizards with no exposure maintained a WBC count of almost 17,600/mm³ during the entire experiment. In lizards exposed to 3,000 R, the mean WBC count was 14,000/mm³ at the end of wk 1 and 20,600/mm³ in wk 6. In the group exposed to 3,500 R, the mean WBC count at the end of wk 1 was 13,800/mm³, but it decreased to 4,400/mm³ in wk 6. In animals exposed to 4,000 R, the mean WBC count was 8,600/mm³ at the end of wk 1 and 1,600/mm³ by wk 6. These observations, combined with those of previous studies, indicate that the WBC elevation in the group exposed to 3,000 R, after the initial decrease during the wk 1, was due to abnormal cell morphology. Thus, the 3,000 R dose induced leukemia and the higher doses induced leukopenia. (20 refs.)

78-4879 **An Analysis of Radiation-induced Malignancy Based on Somatic Mutation.** (Eng) Leenhouts, H. P. (Assoc. Euratom-ITAL, Postbus 48, Wageningen, Netherlands); Chadwick, K. H. *Int J Radiat Biol* 33(4): 357-370; 1978.

A mathematical equation is derived to provide a general description of the dose response for radiation-induced malignancy. The equation is based on the molecular theory of cell survival and on the proposal that a somatic mutation in the normal dominant homologous gene of a diploid cell, which

carries the malignant factor, may allow expression of a recessive malignant factor. The equation assumes that a radiation-induced DNA double-stranded break can cause a somatic mutation or a chromosome aberration that may leave the cell in a precancerous state. Data on the radiation-induced transformation of diploid cells and on the transformation of tetraploid cells are analyzed. The results indicate that the dose relationship for cell transformation is strictly nonlinear, increases linearly-quadratically to a max, and decreases at higher doses. Therefore, any type of extrapolation from high dose measurements to levels of interest for radiological protection should be treated with caution. (72 refs)

- 78-4880 Expression of RNA Tumor Viruses at Translation and Transcription Levels.** (Eng) Fan, H. (Tumor Virology Lab., Salk Inst., P.O. Box 1809, San Diego, CA, 92112). *Curr Top Microbiol Immunol* 79: 1-41; 1978.

Data on the translation and transcription of RNA tumor virus-specific RNA, especially that of avian and murine C-type RNA tumor viruses, in infected and uninfected cells are reviewed. Most of the virus-specific RNA in cells is identical in sequence to virion RNA, with only small amounts complementary to virion RNA being present. Intracellular virus-specific RNA is 35S and 20S-30S in size. Virus-specific RNA is synthesized in the nucleus by nuclear RNA polymerase II. The 35S molecule is the major virus-specific RNA synthesized; no large precursor molecules are evident, although larger size virus-specific RNA does exist. Cytoplasmic virus-specific RNA serves as a messenger RNA (mRNA) for viral proteins, and it is packaged into virus particles; however, only a minority of this RNA serves as mRNA. The major mRNA molecule has a size of 35S, and evidence indicates that the internal structural proteins of the virus are synthesized on polyribosomes containing 35S virus-specific mRNA. RNA coded by endogenous C-type virus genes is expressed at low levels in uninfected cells, and the level of this RNA is frequently correlated with the expression of viral proteins. Furthermore, partial expression of a particular endogenous virus may occur. DNA sequences that correspond to sarcomagenic portions of avian and murine sarcoma viruses are present in uninfected cell DNA. In some cases, transcription of the DNA into RNA is correlated with the growth state of the cell. (135 refs.)

- 78-4881 Temperature-sensitive Mutants of Avian RNA Tumor Viruses: A Review.** (Eng) Friis, R. R. (Institut für Virologie, Justus-Liebig-Universität Giessen, Fachbereich Humanmedizin, Frankfurter Strasse 107, D-6300 Giessen, W. Germany). *Curr Top Microbiol Immunol* 79: 261-293; 1978.

A review of temperature-sensitive mutants of avian RNA tumor viruses is presented. To date, at least one tempera-

ture-sensitive mutant has been identified for each genetic element of the avian sarcoma viruses. In this review, the mutants have been grouped into two categories: T mutants, which have defects exclusively in transformation, and R mutants, which are defective for virus replication, even if the function affected is also essential for transformation. Concerning T mutants, specific topics discussed include the uncertainties about the nature of the transforming gene *src*, parameters for measuring transformation, transformation-defective and partially transformation-defective mutants, and speculations about the function(s) of the *src* gene product. Specific topics discussed concerning R-mutants include mutants defective for *env*, *pol* mutants, and a mutant defective for *gag*. (156 refs.)

- 78-4882 The Function and Origin of the Transforming Gene of Avian Sarcoma Virus.** (Eng) Varmus, H. E. (Dept. Microbiology, Univ. California, San Francisco, CA, 94143); Spector, D. H.; Deng, C. T.; Padgett, T.; Bishop, J. M.; Stehelin, D.; Stubblefield, E. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1159-1168; 1977.

Studies on the origin and function of the transforming gene (*src* gene) of avian sarcoma virus (ASV) using DNA complementary (cDNA) to a portion of the genome that includes some or all of the *src* gene (cDNA-*sarc*) are reviewed. In ASV-infected avian cells (which are permissive for virus replication), large amounts of RNA transcripts containing the *src* gene are detected readily. cDNA-*sarc* anneals to both 35S and 24S RNA in these cells, but it is not known which of these species serves as a messenger in the translation of the *src* gene product. Reversion of ASV-transformed baby hamster kidney (BHK) cells to normal behavior appears to be related to relatively small differences in expression of the *src* gene, as revealed by cDNA-*sarc* hybridization experiments. Annealing of cDNA-*sarc* to nucleic acids from normal (uninfected) cells established that DNA related, but not identical, to the *src* gene of ASV is present in the genomes of all vertebrates. RNA's extracted from a large variety of avian tissues and cells have been tested for their capacity to hybridize with cDNA-*sarc*; in every case, the results indicate the presence of 1-8 "sarc" RNA molecules per cell. When the distribution and size of sarc-containing RNA were compared in normal Japanese quail embryo fibroblasts and in cells derived from a fibrosarcoma induced in quail with methylcholanthrene, virtually identical concentrations of sarc RNA were found in the nuclear, cytoplasmic, and polyribosomal fractions. The function of the sarc sequence remains unknown. (33 refs)

- 78-4883 Type C Viruses of Wild Mice: Characterization and Natural History of Amphrotropic, Ecotrop-**

c, and Xenotropic MuLV. (Eng) Gardner, M. B. (Dept. Pathology, Univ. Southern California Sch. Medicine, Los Angeles, CA, 90033). *Curr Top Microbiol Immunol* 79: 215-259; 1978.

Experiments dealing with the discovery of low- and high-virus-expressor wild mice, the characterization of amphotropic (A-tropic), ecotropic (E-tropic), and xenotropic (X-tropic) C-type murine leukemia viruses (MuLV) in wild mice, and the natural history of C-type viruses in wild mice are reviewed. C-type viruses exist in most wild mice in a strongly suppressed state; virus activation is found primarily in association with lymphomagenesis in old age. In some mice, however, latent C-type virus is genetically less suppressed and is manifest as a congenitally transmitted, persistent infection with specific immune tolerance, associated with an increased incidence of non-T-cell lymphoma and hind limb paralysis. The A-tropic viruses are the most prevalent and they are unique to wild mice. The E-tropic viruses are isolated mainly from paralyzed and some lymphoma-bearing wild mice. X-tropic virus has been isolated from two thymodexyridine-treated embryo culture. The A-tropic virus probably represents the ancestral virus of these classes. Experimental transmission in newborn laboratory mice indicates that A- and E-tropic viruses are lymphomagenic, but only the latter are paralytogenic. Autogenous humoral immunity to indigenous MuLV of all three virus classes is apparently lacking in both high- and low-expressor wild mice. However, the role of autogenous humoral and cellular immunity in these mice is still uncertain. The virus characteristics, natural history, and associated diseases in high-expressor LC wild mice are different in several major aspects from those in high-expressor lymphoma-prone AKR laboratory mice. Lymphoma and paralysis can be controlled in LC wild mice by suppression of E- and A-tropic viruses, using specific genetic and immunologic measures. (143 refs.)

78-4884 **The Development of Vaccines Against Feline Leukemia.** (Eng) Jarrett, W. F. (Dept. Veterinary Pathology, Univ. Glasgow Veterinary Sch., Bearsden, Glasgow G61 1QH, Scotland). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1215-1222; 1977.

Vaccines have been developed and successfully tested against feline leukemia. This disease lends itself to these approaches because feline leukemia virus (FeLV) is transmitted horizontally, cats are usually infected until they are young adults (5 mo), and a natural immunosurveillance system exists. Cats inoculated with 2×10^8 infected cultured cells developed high antibody titers that persisted for >1 yr. These cats were viremic, and virus was found in the bone marrow at necropsy. In contrast, cats inoculated with 4×10^7 cells produced a good feline oncornavirus-associated

membrane antigen (FOCMA) antibody response, and no virus was found in tissues taken at necropsy. In other experiments, cats were inoculated with FL74 cells (having a high density of antigen on the surface and producing virus of low infectivity) and challenged with 10^7 IU of FeLV 7 mo later. No abnormality was found in the cats, and no virus was found in any tissues. The most promising vaccine to date has been prepared from FL74 cells inactivated by paraformaldehyde. Animals inoculated with this vaccine (7×10^7 treated cells) have shown complete resistance to challenge. It is concluded that the pathogenesis and epidemiology of FeLV infection make vaccination practicable and that the feasibility of large-scale field vaccination is now essentially a technological exercise. (8 refs)

78-4885 **Leukemia, Lymphoma, and Fibrosarcoma of Cats as Models for Similar Diseases of Man.** (Eng) Essex, M. (Dept. Microbiology, Harvard Univ. Sch. Public Health, Boston, MA, 02115); Cotter, S. M.; Stephenson, J. R.; Aaronson, S. A.; Hardy, W. D. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1197-1214; 1977.

The use of spontaneous feline leukemia, lymphoma, and fibrosarcoma as models for studying analogous human tumors is reviewed. There is overwhelming evidence from epidemiologic and serologic studies that spontaneous leukemia and lymphoma of cats are caused by horizontally transmitted viruses [feline leukemia virus (FeLV)]. In studies with feline fibrosarcoma, a correlation was found between the presence of antibodies to feline oncornavirus-associated cell-membrane antigen (FOCMA) and tumor progression. In subsequent studies of laboratory and household cats, animals that did not develop malignancies following exposure to FeLV or feline sarcoma virus had elevated titers of FOCMA antibodies. A number of independent studies provide evidence that FOCMA is distinct from the major virus proteins, gp70 and p30. It is concluded that immunosurveillance works, at least for one group of tumors in one mammalian species. Therefore, attempts to determine if immunosurveillance ever occurs in humans should not be abandoned. (47 refs)

78-4886 **Progress in Studies on the Etiology and Serologic Diagnosis of Enzootic Bovine Leukosis.** (Eng) Mussgay, M. (Federal Res. Inst. Animal Virus Diseases, Paul Ehrlich Strasse 28, D-7400 Tübingen, W. Germany); Kaaden, O. R. *Curr Top Microbiol Immunol* 79: 43-72; 1978.

Advances made since 1969 concerning the properties of bovine C-type particles and their use in the serologic diagnosis of enzootic bovine leukosis are reviewed. This disease is characterized by neoplasia of the lymphocyte-forming tissue and

by its occurrence as multiple cases within a herd. It may develop in two phases: a persistent lymphocytosis followed, in approx 10%-30% of the animals, by the formation of leukotic tumors. According to the literature, the incubation time varies between 200 days and 7 yr. Etiologic studies dealing with epizootiological aspects of the disease, experimental transmission, and the production of bovine C-type particles in cell culture are reviewed. Attention is also focused on the biochemical and biophysical properties of the virus particles, their morphology, genetic relatedness to host DNA, host spectrum, and their relationship to other retroviridae. Serologic diagnoses by agar gel immunodiffusion, complement fixation, immunofluorescence, radioimmunoassay, and the indirect immunoperoxidase test and the detection of both the virus and its antigen are summarized. (98 refs.)

78-4887 Current Assessment of Human Health Hazards Associated with Bovine Leukemia Virus. (Eng)

Van Der Maaten, M. J. (U.S. Dept. Agriculture, Agricultural Res. Service, North Central Region, Natl. Animal Disease Center, Ames, IA, 50010); Miller, J. M. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1223-1234; 1977.

The properties of bovine leukemia virus (BLV) are reviewed, along with the potential hazards of this virus to human health. Seroepidemiologic studies have provided ample evidence of the association between virus infection and tumor development. Morphologic, biochemical, and immunologic studies have indicated that BLV is a unique oncornavirus that is not endogenous to the bovine species. The presence of infectious BLV in milk was established by studies in which sheep, inoculated with milk from infected cows, subsequently developed active BLV infections. Circumstantial evidence that exposure to milk or cattle may result in an increased incidence of human neoplasms has come from the occurrence of an unusual number of cases of adult nonlymphocytic leukemia in a community with a commercial milk-drying plant. Several cases of lymphoid leukemia have been reported in people living near a dairy herd with numerous cases of lymphosarcoma. Other studies, however, have not supported this correlation. Studies with chimpanzees inoculated with large doses of cell-culture-grown BLV have provided direct proof of primate susceptibility. Tests of sera from cancer patients, high-incidence cancer families, and individuals with max exposure to infected cattle or to BLV in the laboratory have failed to identify even one individual with antibodies to BLV. Thus, the risk to human health associated BLV would seem to be minimal. (57 refs)

78-4888 Herpes Simplex Viruses and Human Malignancies. (Eng) Rawls, W. E. (Dept. Pathology,

McMaster Univ., Hamilton, Ontario L8S 4J9, Canada
Adam, E. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1133-1155; 1977.

Results of a seroepidemiologic case-control study of women with cervical cancer (CC) are presented and compared with published data relating herpes simplex virus type 2 (HSV-2) to CC. Sera were collected from 285 white women (108 with invasive carcinoma, 71 with carcinoma in situ, and 106 controls with no cervical disease). Forty-five percent of the cancer patients and 17% of the control women had antibodies to HSV-2. Forty-four percent of the invasive cancer cases, 42% of the carcinoma in situ cases, and 30% of the controls had multiple sex partners. These figures are comparable to those from other studies. Antibodies to HSV-2 were more common among women with multiple sex partners than among women with one sex partner. Data from several studies using serologic assays to determine past infections with HSV-2 indicate that few, if any, women with CC have not been infected with at least one of the herpes simplex viruses. Studies of the interrelations among sexual promiscuity, CC, and HSV-2 suggest that the association between HSV-2 and CC does not simply represent one of covariability to sexual promiscuity. In addition to neutralizing antibodies, antibodies to three other virus-induced antigens (AG-4, HSV tumor-associated antigen, and VP134) have been detected in sera from women with CC. HSV-2 DNA and proteins have not been easily and reproducibly demonstrated in CC cells. It is not yet possible to establish an unequivocal etiologic relationship between HSV-2 and CC. (81 refs)

78-4889 Papillomaviruses: Possible Role in Human Cancer. (Eng) Orth, G. (Unite de Recherche sur l'Etiologie Virale des Cancers Humains, Laboratoire Associe au CNRS no. 147, Unite 140 de l'INSERM, Institut Gustave-Roussy, 94800 Villejuif, France); Breitburd, F.; Favre, M.; Croissant, O. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1043-1068; 1977.

Data on the biological properties of animal and human papillomaviruses (PV) are reviewed to show that a role of PV in the origin of human cancer must be considered. At least 13 different PV have been characterized in benign skin or mucosa tumors of one bird species and nine mammalian species. Three of these animal PV have oncogenic potential for epithelial or connective tissues in both natural and experimental conditions. When oncogenic, PV induce premalignant lesions that convert to malignancy under the influence of multiple factors, such as host and perhaps virus genetic constitution, deficient host immune response, and environmental cocar-

ogenic factors. The formation of a papilloma results from an abortive transforming infection of germinal cells. Human virus-induced conditions include skin warts, epidermodysplasia verruciformis, condylomata acuminata, oral papillomas, and juvenile laryngeal papillomatosis. Lesions are usually benign and they usually regress, but malignant transformation may occur as a rare event. It has long been considered that the different human lesions are due to the same virus and that the clinical types are determined by local conditions, but evidence is now accumulating that several different human PV may be involved. (124 refs)

78-4890 **Human Papovaviruses.** (Eng) Takemoto, K. K. (Lab. Viral Diseases, Natl. Inst. Allergy and Infectious Diseases, NIH, Bethesda, MD, 20014). *Int Rev Exp Med* (18): 281-301; 1978.

Current knowledge of the biology, biochemistry, and oncogenicity of human papovaviruses (PVs) is reviewed. Of the three types of related but immunologically distinct PVs isolated from humans, only BK virus and JC virus are true human viruses. The third type [represented by 2 isolates from progressive multifocal leukoencephalopathy (PML) case and 1 from melanoma tumor tissue] can be classified as simian virus 40 (SV40) variants that are rare and of unknown origin. JCV is probably the causative agent of PML. It has been isolated from or identified in diseased brain tissue of 21 cases. BKV has been found in the urine of immunosuppressed patients. There are at least 10 isolates from the urine of renal transplant patients and 3 from Wiskott-Aldrich cases. There is no known disease associated with BKV. PVs, their antigens, or viral DNA sequences have been detected in some human tumor cells, but there is no conclusive evidence that these viruses play any significant role in human tumor cells. BKV, JCV, and SV40 are distantly related primate PVs based on the following observations: (1) their capsid antigens show some serologic cross-reactivity in several different types of tests, and they induce similar or identical T antigens; and (2) the DNAs of the three viruses show 10%-25% homology. (105 refs)

78-4891 **The Human Papovaviruses and Their Potential Role in Human Diseases.** (Eng) Khoury, G. (Lab. DNA Tumor Viruses, NCI, NIH, Bethesda, MD, 20014); Lai, C. J.; Solomon, D.; Israel, M.; Howley, P. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 971-988; 1977.

Studies of simian virus 40 (SV40) and human papovaviruses (PVV) are reviewed. Two PVV have been isolated from humans: (1) JC, from the brain of a patient with progressive

multifocal leukoencephalopathy (PML); and (2) BK, from the urine of an immunosuppressed renal transplant patient. Although antibodies to SV40 are uncommon in humans (3%), antibodies to both BK and JC viruses are prevalent. The human PVV induce the synthesis of a tumor (T) antigen early in the lytic cycle and in transformed cells. The BK and JC T antigens cross-react strongly with the SV40 T antigen. Tumor-specific transplantation antigens (TSTA) of PVV have been localized on the surfaces of transformed and tumor cells. Structural or virus antigens can be detected late in the lytic cycle and are absent in transformed cells in which only the early events of PVV-cell interactions are expressed. Both JC and BK viruses induced tumors in inoculated newborn hamsters, and BK virus has malignantly transformed animal cells in tissue culture. Papovaviral particles were detected by electron microscopy or immunoreactivity with SV40 T antisera in human meningiomas and a metastasis from a malignant melanoma. The association of JC virus with PML seems firmly established. PVV particles have been detected in the brains of almost every patient with this disease and have been specifically localized in the areas of demyelination. All but two isolates from these lesions have been JC virus. A degenerative neurologic disease, pathologically identical to PML in humans, has been described in macaques. Virus isolates from two cases were identified as SV40. Thus, SV40 and JC virus appear to be associated with rare but pathologically similar diseases in their natural hosts. (96 refs)

78-4892 **The Role of the Papovavirus Gene A in Oncogenic Transformation.** (Eng) Tegtmeyer, P. (Dept. Microbiology, State Univ. New York, Stony Brook, NY, 11594); Rundell, K. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 957-969; 1977.

Studies of the complementation group A mutants of simian virus 40 (SV40) and polyoma virus, which are defective in both productive and transforming infections, are reviewed. Studies of SV40 infection have identified the A gene as necessary for the initiation and perhaps the maintenance of neoplastic transformation in model animal systems. The localization of the A-protein binding site to a region of viral DNA that is functionally important for both DNA replication and transcription suggests that the A protein may regulate these processes in a direct and coordinated way. Cross-reacting antigens have been detected in cells productively infected or transformed by SV40 or by human strains of papovaviruses. Immunoprecipitation confirmed that the antigen induced by the human virus is closely related to the A protein of SV40. The A proteins specified by SV40 and by RFV strains of human virus in transformed hamster cells are similar but show distinct differences in their peptide patterns, indicating that the A protein is coded for by the virus rather than the

cell. The close functional and structural relationship of the A proteins of SV40 and human viruses suggests that events known to occur in SV40 infection might also be common events in humans during infection by papovaviruses. (74 refs)

- 78-4893 **BK, A Human Polyoma Virus -- Its Biology, Biochemistry, and Association with Human Tumors.** (Eng) Major, E. O. (Dept. Microbiology, Stritch Sch. Medicine, Maywood, IL, 60153); Fiori, M.; di Mayorca, G.; Zacharias, C. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 989-1007; 1977.

Initial investigations of BK virus, a human papovavirus, from its isolation to its association with human disease are reported. BK virus was isolated from the urine of a renal allograft recipient. Antibodies to BK (and to another human papovavirus, JC virus) have been found in 80% of the world population from age 5 to adult. Hemagglutination inhibition (HAI) and plaque neutralization (PN) assays indicate no relationship between BK virus and polyoma virus and only marginal cross-reactivity between BK and SV40 or JC virus. Virion protein analysis by electrophoresis on polyacrylamide gels indicates that BK virus is an independent human isolate and not a simian variant. Molecular hybridization data from experiments in which BK DNA was hybridized with SV40 and JC viral DNA show 10%-40% homology, which exists between sequences located in the late region of the genome. BK virus transforms baby hamster kidney (BHK) cells, and these cells can induce tumors in adult hamsters. BK tumor (T) antigen cross-reacts with other papovavirus T antigens but retains a BK-unique antigenic site. Under single-cycle conditions, the growth cycle of BK virus in the most permissive host, human embryonic kidney cells, takes 48 hr. There is evidence that BK virus remains encapsidated in the nucleus of infected cells for at least 24 hr, so that some host factors must control the timing and extent of virus multiplication. Reassociation kinetics indicate that the number of genome equivalents varies from < 1 to > 3 in the different clones of BK-transformed hamster cell DNA tested. (46 refs)

- 78-4894 **The Molecular Basis of Transformation by Simian Virus 40.** (Eng) Martin, R. G. (Lab. Molecular Biology, Natl. Inst. Arthritis, Metabolism, and Digestive Diseases, NIH, Bethesda, MD, 20014); Persico-DiLauro, M.; Edwards, C. A.; Oppenheim, A. *Miami Winter Symp* 14: 87-102; 1977.

A model for the molecular basis of cell transformation by simian virus 40 (SV40) is presented. The model assumes that the SV40 tumor (T) antigen is the only viral product required for transformation and that the T antigen is an initiator of viral DNA synthesis. The model proposes that: (1) the T

antigen acting at the end of G₁ drives cells into DNA synthesis under suboptimal growth conditions; (2) the host partly regulates the expression of T antigen; (3) T antigen initiates DNA synthesis in the host either by virtue of its high- or low-affinity binding to DNA; and (4) membrane changes are largely secondary to the inability of transformed cells to enter the resting state. (65 refs)

- 78-4895 **Viruses as Causes of Some Human Tumors? Results and Prospectives of the Epidemiologic Approach.** (Eng) de-The, G. (Unit Biological Carcinogenesis, International Agency Res. Cancer, 69008 Lyon, France). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1113-1131; 1977.

Epidemiologic approaches to the study of viruses as possible causes of human cancer are reviewed, and evidence linking Epstein-Barr virus (EBV) to Burkitt's lymphoma (BL) and nasopharyngeal carcinoma (NPC) is presented. Epidemiologic characteristics of human tumors that would favor the possibility of a viral infection are: (1) horizontal transmission of tumor risk; (2) clustering of tumors in time and/or space; (3) an association between benign viral lesions and specific tumors; and (4) the association between hereditary or genetic factors with the risk for certain tumors. A prospective serological study of BL in 42,000 Ugandan children, 12 of whom developed BL, indicates the probability of EBV infection at an early age in children who develop BL and that antibodies to EBV 7-20 mo prior to clinical onset do not protect against BL. This study plus other observations suggests that a series of causal factors is involved in BL development in tropical areas, including neonatal EBV infection, genetic translocation in chromosome 14 in cells earmarked for malignant transformation, and hyperendemic malaria. A fourth unknown factor that represents the triggering event leading to the clinical onset of the disease is postulated. Patients with NPC exhibit similar and even higher EBV serologic reactivities than BL patients. Of even more critical etiologic importance is the presence of EBV markers in epithelial tumor cells. Human tumors that may be virus-associated and would be good candidates for epidemiologic studies include genital cancers (herpes simplex virus type 2, cytomegalovirus, and papillomaviruses); leukemias; lymphomas; and sarcomas; breast carcinoma; and liver carcinoma. (72 refs)

- 78-4896 **The Biology of EB Virus (Meeting Abstract).** (Eng) Pope, J. H. (Queensland Inst. Medical Res., Australia). *Pathology* 10(2): 190; 1978. (no refs)

78-4897 **Viral Etiology of Cancer.** (Spa) de Pasqualini, C. D. (Seccion Leucemia Experimental, Instituto de Investigaciones Hematologicas, Academia Nacional de Medicina, Buenos Aires, Argentina). *An Acad Bras Cienc Exactas Fis Nat* Buenos Aires 29: 229-251; 1977.

Studies of the viral etiology of cancer are reviewed. Certain RNA and DNA viruses induce cancer in many animal species, including primates. There is indirect evidence of a viral etiology for many human tumors. Leukemia with oncornavirus replication was induced in rats and primates by inoculation of these animals with human neoplastic material. In humans, C particles were found in association with leukemia, and replicating B particles were found in breast carcinoma cell lines. Oncornavirus was detected in human cancers by molecular hybridization. Human lymphoid cells were found to contain herpesvirus-Epstein-Barr virus. C-type oncornavirus was detected in normal human and primate tissues, especially in the placenta and fibroblasts. Carcinogenesis may be the result of the activation or repression of oncogenic virus by cocarcinogens, mutagenic factors such as radiation and chemicals, immunosuppressive factors such as malaria and viruses, including herpesviruses. (137 refs)

78-4898 **Type D Primate Retroviruses: A Review.** (Eng) Fine, D. (Viral Oncology Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Schochetman, S. *Cancer Res* 38(10): 3123-3139; 1978.

The unique characteristics of the D-type retroviruses, their distribution in the primate population, and their relevance to human cancer are reviewed. The prototype D-type retrovirus is the Mason-Pfizer monkey virus (MPMV), originally isolated from a breast carcinoma of a female rhesus (Old World) monkey. This is the only virus ever isolated from a primate mammary tumor, and it has in vitro transforming potential for primate cells. Viruses morphologically and immunologically indistinguishable from MPMV have been isolated from normal placenta and lactating mammary glands of other rhesus monkeys in captivity. Recently, viruses morphologically resembling MPMV have been isolated from a langur (Old World) and a squirrel (New World) monkey. Nucleic acid hybridization studies indicated that the latter viruses are endogenous in the species of origin, while MPMV appears to be a horizontally transmitted virus containing gene sequences partially related to the langur monkey isolate. Studies on the immunological relatedness of the D-type retroviruses have demonstrated interspecies cross-reactivities between the major internal and external viral proteins. Furthermore, these viruses also share cross-reactivity of their major external glycoproteins with those of the C-type baboon endogenous virus. These interspecies reactivities can also be demonstrated in natural sera from both imported and laboratory-bred monkeys. Conflicting reports on the association between MPMV and human neoplasia are reviewed. A broad

interspecies assay developed to detect immunological cross-reactivities among D-type viruses may aid in the detection of human retroviruses. (109 refs)

78-4899 **The Development of the Concept of Viral Etiology of Leukemia and Related Neoplastic Diseases. Present Status and Prospects for the Future.** (Eng) Gross, L. (Veterans Admin. Hosp., Bronx, NY, 10468). In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 1-10; 1978.

The concept of a viral etiology of leukemia and related neoplastic diseases is discussed. Using C58 and AK mice, which have a high incidence of leukemia in middle age, it was demonstrated that leukemia and lymphosarcomas are caused by a filterable virus and that this virus is transmitted from one generation to another directly through the embryos. Additional experimental data indicated that leukemia, lymphosarcomas, and related neoplastic diseases are caused by transmissible oncogenic viruses in most, possibly all, animal species. In most instances, these viruses remain latent and harmless to their carrier hosts; when activated, however, they may cause fatal disease. These oncogenic viruses appear to be widely prevalent in many species, particularly within certain families, and they may be transmitted horizontally within the same generation as well as vertically from one generation to another. Both RNA and DNA viruses may be oncogenic. Leukemia and allied neoplastic diseases have fundamental features in common with "slow virus diseases" such as scrapie, kuru, and Creutzfeldt-Jakob disease, although the oncogenic viruses are much more susceptible to inactivation by heat and other factors (physical and chemical) than the agents responsible for the slow virus diseases, and they do not readily cross species. (20 refs)

78-4900 **RNA Tumor Virus Genes (Virogenes) and the Transforming Genes (Oncogenes): Genetic Transmission, Infectious Spread, and Modes of Expression.** (Eng) Todaro, G. J. (Lab. Viral Carcinogenesis, NCI, Bethesda, MD, 20014). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation.* Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1169-1196; 1977.

The transmission of RNA tumor virus genes, the relationship between C-type RNA viruses and human neoplasia, normal functions of C-type viruses, and recombination between C-type viruses are reviewed. RNA tumor virus genes may be transmitted genetically (parent to offspring) or horizontally (animal to animal). C-type viruses, under natural conditions,

have also apparently been transferred between species that are only remotely related. In all but one case, the viral genes have become incorporated into the genome of the recipient species. The exception is the infectious primate virus group that has produced leukemias and sarcomas in primates and has been suggested to be involved in human leukemia. Several studies have indicated that this group of infectious C-type viruses, horizontally transmitted among primates, originated by transspecies infection of certain primates (gibbon, woolly monkey) by an endogenous C-type mouse virus. An infectious C-type virus closely related, if not identical, to woolly monkey virus, simian sarcoma-associated virus (SSAV), has been isolated from a cell culture derived from a woman with acute myelogenous leukemia. Endogenous primate virogenes may also be partially expressed in humans and other primates. C-type viruses may perform normal functions such as control of development, protection from infection, and transmission of genetic information. One protein that satisfies the criteria for a possible oncogene production is epidermal growth factor (EGF). Aberrant production of a growth hormone like EGF, due to introduction of sarcoma virus genes or activation of endogenous genes, may be one mechanism of transformation. (137 refs)

- 78-4901 Some Ideas on the Origin of Leukemia in Man and Recent Evidence for the Presence of Type-C Viral Related Information.** (Eng) Gallo, R. C. (Lab. Tumor Cell Biology, NCI, NIH, Bethesda, MD, 20014); Saxinger, W. C.; Gallagher, R. E.; Gillespie, D. H.; Aulakh, G. S.; Wong-Staal, F.; Ruscetti, F.; Reitz, M. S. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1253-1285; 1977.

Evidence for the presence of C-type viral related information in patients suffering from leukemia is reviewed, and five possible mechanisms of viral leukemogenesis are presented. Intracytoplasmic viruslike particles containing high-mol-wt RNA and reverse transcriptase activity have been identified in human leukemic blood and bone marrow cells. Infectious, fully mature C-type virus, containing components related to baboon endogenous virus and simian sarcoma virus, has been isolated from a patient with acute myelogenous leukemia. Proviral sequences closely related to those of baboon endogenous virus and distantly related to those of murine leukemia virus have also been found in DNA from leukemic patients. The first model of leukemogenesis proposes a classical mechanism of infection of the stem cell by a C-type virus, resulting in complete integration of the provirus, partial expression of the virus genome, and transformation of the stem cell. In the second model, only a fragment of the provirus is integrated, but it is sufficient to cause transformation. The third model involves transformation of a regulator cell. The fourth proposes that the stem cell is infected and contains a complete

provirus, but is not transformed. After a number of divisions of the stem cells, some of the progeny cells lose some or all of the provirus and adjacent cellular genes responsible for cell differentiation. These cells are blocked in normal maturation and thus have a proliferative advantage. The fifth model proposes that a C-type virus infects a target cell other than WBC. The infected cell releases DNA fragments or produces virus that secondarily infects a WBC precursor, leading to transformation. The fourth and fifth models are the most consistent with available data. (81 refs)

- 78-4902 Transformation In Vitro by Adenovirus Type 2--A Model System for Studying Mechanisms of Oncogenicity.** (Eng) McDougall, J. K. (Cold Spring Harbor Lab., Cold Spring Harbor, NY, 11524); Chen, L. B.; Galimore, P. H. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1013-1025; 1977.

The characteristics of rat cells transformed in vitro by adenovirustype 2 (Ad2) and the in vivo behavior of these cells are reviewed. Ad2-transformed cells consistently contain a minimum segment of viral DNA representing 14% of the left-hand end of the viral genome, indicating that the transformed phenotype can be induced and maintained by this region of the viral genome. With some Ad2-transformed cell lines, immunosuppression of the newborn rat results in tumor induction, suggesting that surface antigenicity is an important factor in the in vivo growth characteristics of these cell lines. There is a correlation between a decrease in one of the surface proteins, the large, external transformation-sensitive (LETS) protein, and an increase in tumorigenicity. Oncogenicity of Ad2-transformed cells is also correlated with the amount of infectious virus used to infect the monolayer cultures in which transformed foci develop. There is evidence that cells transformed at higher multiplicities of infection retain more of the virus genome. In studies with somatic-cell hybrids, there is no indication that Ad2 sequences integrate at specific loci. Rat and hamster cells transformed by adenoviruses, simian virus 40, and herpesviruses show some similar marker chromosomes, and two of the autosomes regularly involved in rearrangements in adenovirus-transformed cells are also preferentially involved in marker formation in chemical transformants. This could suggest involvement of specific loci on the chromosomes that carry genes important in the regulation of cell growth and malignancy. (33 refs)

- 78-4903 Viruses in Cancer. Evaluation of Current Evidence of Possible Role of Viruses in Human Cancer (Meeting Abstract).** (Eng) Sabin, A. B. (Medical Univ. South Carolina, Charleston, SC). *Pathology* 10(2): 187-189; 1978. (1 ref)

78-4904 **Aspects of Virus Persistence and Ubiquity in Relation to Environmental and Co-Factors (Meeting Abstract).** (Eng) Stanley, N. F. (Dept. Microbiology, Univ. Western Australia, Australia). *Pathology* 10(2): 189-190; 1978. (no refs)

78-4905 **Cancer, Virus Infections and the Immune Response (Meeting Abstract).** (Eng) Ada, G. L. (Dept. Microbiology, John Curtin Sch. Medical Res., Australian National Univ., Australia). *Pathology* 10(2): 189; 1978. (no refs)

78-4906 **Ageing, Metabolic Immunodepression and Carcinogenesis.** (Eng) Dilman, V. M. (N.N. Petrov Inst. Oncology, Leningrad, USSR). *Mech Ageing Dev* 3(3): 153-173; 1978.

Data suggesting a relationship between metabolic immunodepression and susceptibility to neoplasia are reviewed. A correlation has been suggested between increases in blood levels of free fatty acids (FFA), insulin, cholesterol, and low-density lipoproteins and immunodepressions. The same metabolic factors that inhibit cellular immunity stimulate the proliferation of somatic nonlymphoid cells, thereby increasing the possibility of malignant transformation. States such as aging, stress, pregnancy and cancer, all involving a deterioration of cell-mediated immunity, are characterized by a similar hormone-metabolic pattern. The opposing influences of the same metabolic factors on somatic, nonlymphoid cells vs lymphocytes promote the development of carcinogenesis-promoting metabolic syndrome (CPMS). Many chemical carcinogens alter the biogenic amine levels in the hypothalamus; this in turn, leads to an elevation of the hypothalamic threshold of sensitivity to feedback control, a decrease in tolerance to glucose, and an increase in blood insulin level, changes that constitute the CPMS and aging during carcinogenesis. Thus, both normal and cancer-related aging involve metabolic immunodepression and the CPMP due to implementation of the Law of deviation of Homeostasis during natural aging or exposure to chemical carcinogens, respectively. (99 refs)

78-4907 **Immunological Surveillance.** (Eng) Stutman, O. (Dept. Cellular Immunobiology, Memorial Sloan-Kettering Cancer Center, New York, NY, 10021). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. (Book B), 699 pp.; 729-750; 1977.

The experimental and clinical evidence that has been used to support the immune surveillance theory is reviewed. A reflection

analysis of the incidence of tumors in patients with primary and secondary immunodeficiencies does not support the generality of immunological surveillance as an operative mechanism, but does support its role in the development of lymphoreticular tumors. The clinical data from transplant patients also fail to support the orthodox view of immune surveillance. The clinical behavior of the more conventional tumors in immunodepressed transplant patients is comparable to that of tumors in nonimmunodepressed patients. The generality, efficiency, and even existence of a thymus-dependent immunosurveillance mechanism capable of destroying in situ tumors are questionable. Although a surveillance-like mechanism may be operative under certain special conditions (ie, polyoma oncogenesis and that of other DNA viruses, feline leukemia), its thymus dependency is also questionable. (79 refs.)

78-4908 **World-wide Variations in Cancer Incidence as Clues to Cancer Origins.** (Eng) Berg, J. W. (Dept. Preventive Medicine, Univ. Iowa, Iowa City, IA, 52242). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 15-19; 1977.

Generalizations are presented concerning the distribution of cancers that occur at particularly high rates (40 or more cases/yr/1,000 persons aged 35-64) in studied populations. International cancer risk patterns are reviewed briefly, as are problems associated with the pathogenesis of esophageal cancer and of other cancers (stomach, bowel, breast, and prostate). (14 refs)

78-4909 **Cancer Epidemic Raises Doubts on Mineral Fibres.** (Eng) McGinty, L. (No affiliation given). *New Sci* 18(1103): 428; 1978.

The results of epidemiological investigations of the cause of the high mesothelioma rate in Karain, Turkey, are reviewed. Between 1970 and 1974, almost half the people who died in Karain died of mesothelioma. It was initially thought that asbestiform fibers in the drinking water were responsible for the cancer, but recent studies suggest that fibers of volcanic silicates called zeolites are responsible. These fibers are found in the volcanic rock that the residents use for building blocks and road-making materials. Furthermore, these fibers have the same dimensions ($>0.2 \mu\text{m}$ in diameter and 3-15 μm long) as the forms of asbestos (amosite, crocidolite) most active in causing mesothelioma. If the link with zeolite is confirmed, it would be the first human evidence that nonasbestos silicate fibers cause mesothelioma. Such a finding would throw doubt on the safety of manmade mineral fibers suggested as substitutes for asbestos. (1 ref.)

- 78-4910 Epidemiologic Methods for Human Risk Assessment.** (Eng) Mack, T. M. (Dept. Community Medicine and Public Health, Univ. Southern California Sch. Medicine, Los Angeles, CA, 90033); Pike, M. C.; Casagrande, J. T. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1749-1763; 1977.

Three functions of the cancer epidemiologist are discussed. Analysis of population-based incidence data for the purpose of descriptive epidemiology may retrospectively confirm an association between a particular cancer and a known exposure, but it can not predict serious risks. Descriptive epidemiology, however, can identify patterns or changes in patterns of a particular cancer that could result in modifications of current hypotheses. The second function of epidemiology is to describe the association between a putative causal exposure and a disease under circumstances that permit the best possible discrimination between causal and alternative hypotheses. However, sample-size requirements may be prohibitive. The problems involved in designing an epidemiological study to test various hypotheses is illustrated by a study of endometrial cancer, which resulted in the identification of exogenous estrogens as a cause of endometrial cancer. The third function of epidemiology involves the reduction of pertinent biologic information into a comprehensible summary for the purpose of policy formulation. This function is illustrated by analysis of the risks and benefits from the addition of mammography to breast cancer screening programs. The analysis indicates that the benefits of mammograph in women under 50 are likely to come at the price of an increased risk of radiogenic breast cancer later in life. (15 refs)

- 78-4911 Burkitt's Lymphoma in Tropical Countries: Epidemiology and Relationship with Malaria.** (Fre) Charmot, G. (Consultation des Maladies Tropicales, Hopital Claude-Bernard, 10, av. de la Porte d'Aubervilliers, F 75019 Paris, France); Rodhain, F.; Roze, J. M. *Nouv Presse Med* 7(4): 265-269; 1977.

Epidemiological data on Burkitt's lymphoma (BL) are reviewed, and the relationship of the tumor to malaria is discussed. BL is endemic throughout rural regions of Black Africa and New Guinea, with an incidence of 3-7/100,000 children (0-15 yr of age). Characteristically, the region has a fairly dense rural population with a low standard of living. The climate is always warm (above 18 C) and humid, conditions conducive to the permanent or prolonged transmission of malaria. Infection by Epstein-Barr virus (EBV), as determined by the level of EBV antibodies, occurs before the age of 3 yr and precedes the appearance of the tumor by 6 mo to 2-3 yr. Experiments with mice show that the malaria induced by *Paludium berghei* or *P. yoelii* facilitates the development of spontaneous or virus-induced lymphomas and the growth of BL grafts that would otherwise be rejected. In certain regions of East Africa where malaria has been controlled,

the incidence of BL has decreased. It is hypothesized that a soluble mitogen produced by *P. falciparum* and found in blood of Nigerian children may have a stimulatory effect on B lymphocytes containing EBV, which may lead to growth of the lymphoma. Genetic and immunological factors may also play a role in individual susceptibility to BL. (21 refs)

- 78-4912 On the Etiology and Metabolic Epidemiology of the Main Human Cancers.** (Eng) Weisburger, J. H. (Naylor Dana Inst Disease Prevention, American Health Foundation, Valhalla, NY, 10595); Cohen, L. A.; Wynder, E. L. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 567-602; 1977.

Concepts maintaining that the etiologies of the main human cancers stem largely from human life-styles are reviewed. Most cancers of the respiratory tract are due to excessive smoking, as are cancers of the pancreas, kidneys, and bladder. Prevention of these diseases hinges on education of the public and collaboration of industry and government to produce and give preferential marketing to lower tar cigarettes. Cancers of the breast, prostate, ovary, endometrium, and colon are associated with a high fat content in the diet. Cancer of the pancreas is also associated in part with diet. Prevention of these cancers hinges on lowering fat consumption from the prevailing level of 45% of the caloric intake to 25%-33%. This would be aided by cooperation of industry with agricultural and governmental agencies to produce and market foodstuffs that would permit the consumer to adopt the lower fat diet more readily or automatically. Cancer of the stomach might be prevented by increasing the consumption of vitamin C, low-temperature food storage to prevent nitrite formation, and controlling the use of nitrite as a preservative. (155 refs)

- 78-4913 The Malignant Lymphomas: Epidemiology and Related Aspects.** (Eng) Vianna, N. J. (Bureau Occupational Health and Chronic Diseases and Lymphoma Studies, Cancer Control Bureau, Dept. Health, Albany, NY). *Pathobiol Annu* 7: 231-255; 1977.

Three issues of importance to malignant lymphoma are reviewed: the heterogeneity between and within many of the lymphoma groups, the probable importance of environmental and genetic factors in the pathogenesis of malignant lymphoma, and evidence suggesting that an infectious agent might have an etiological role in certain disorders (Hodgkin's disease and Burkitt's lymphoma). (193 refs.)

- 78-4914 Epidemiology of Esophageal Carcinoma (Letter to Editor).** (Eng) Mizroch, S. (Kaiser Foundation Hosp., San Francisco, CA). *JAMA* 239(22): 2340-2341; 1978.

age (older), sex (male), race (nonwhite), and geography exert substantial influence on the increased incidence of esophageal carcinoma. Cigarette smoking, alcohol consumption, and thermal irritation from imbibing hot foods are also predisposing factors. There are links between carcinoma and various types of obstructive esophageal lesions. An increased incidence of carcinoma in achalasia, Barrett's esophagus, lye strictures, and Plummer-Vinson syndrome should encourage the high index of suspicion incumbent to early diagnosis. (4 refs)

78-4915 **The Occurrence of Oesophageal Cancer - A World-wide Survey (Meeting Abstract).** (Eng) Muir, C. S. (International Agency Res. on Cancer, 150, Cours Albert Thomas, 69372 Lyon Cedex 2, France); Necoux, J. In: *Third International Symposium on Oncology Held at Taj Pahlavi Cancer Institute in Tehran, 4-8 March, 1978.* Taj Pahlavi Cancer Institute, Pahlavi Medical School, Tehran University. (Tehran, Iran): 115 pp.; 103-104; 1978. (no refs)

78-4916 **Pulmonary Tuberculosis and Bronchial Carcinoma (2 Letters to Editor).** (Eng) Healy, T. M. Dept. Pathology, Univ. Coll., Dublin 2, Ireland; MacGregor, G. A. *Lancet* 2(8085): 372; 1978.

The use of the word "opportunism" as applied to tuberculosis complicating bronchial carcinoma can be validated by a review of the literature. However, in another opinion, it is stated that the association may be more than fortuitous. Since mycobacterial infections appear to be more common in cancer patients than in the general population, it is suggested that the chronic immune irritation generated by tuberculosis and sarcoidosis is an initiating factor in bronchial carcinoma. (11 refs.)

78-4917 **Changing Patterns of Cancer in Japan with Special Reference to the Decrease in Stomach Cancer Mortality.** (Eng) Hirayama, T. (Epidemiology Div., Natl. Cancer Center Res. Inst., Tsukiji 5, Chuo-ku, Tokyo, Japan). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation.* Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4. 602 pp.; 55-75; 1977.

The steady downward trend in age-specific death rates for stomach cancer in Japan is discussed. Almost no increase in mortality for cancer of all sites was observed in men during the last decade and a slight decrease was observed among women. The greatest decreases occurred in cervical cancer rates and in the rates of stomach cancer in both men and women. Sharp increases were observed for leukemia and for cancer of the lung, pancreas, prostate, urinary tract, colon-rectum, liver, breast, and ovary. The risk of stomach cancer was significantly increased in daily cigarette smokers and

significantly reduced in those who ate green-yellow vegetables daily and those who drank two glasses of milk daily. The decline in stomach cancer mortality appears to be due to dietary improvements and to the effect of early detection. The mortality rate for colon cancer and the colon cancer-rectal cancer ratio were highest in persons who consumed a small amount of rice daily and in daily beer drinkers. Increases in the breast cancer mortality rate may be due to a decline in the birth rate or to the rapidly increasing intake of dietary fat. Mortality from breast cancer was highly correlated with pork intake and, to a lesser extent, with animal fat intake. Daily cigarette smoking was the most important cause of lung cancer. Causes for changing patterns in the rates of other cancers are also discussed. (13 refs.)

78-4918 **The Origin of Malignant Tumors of the Upper-Airways, the Digestive Tract and the Ear. Organ-Resistance and Organ-Disposition (Organotropy) by the Origin of Malignant Primary Tumors.** (Ger) Leicher, H. (Langenbechs. 1, 6500, Univ.-HNO-Klinik, Mainz, W. Germany). *Laryngol Rhinol Otol (Stuttg)* 57(3): 162-176; 1978.

General biological mechanisms in the genesis of malignant tumors of the upper respiratory and digestive tracts and of the ear are reviewed. Carcinomas of the nose, trachea, mouth, esophagus, and middle ear are significantly less frequent than those of the bronchi and larynx, a finding that cannot be explained by differences in the degree of exposure of these organs to carcinogenic substances. It can be explained by differences in their resistance to malignant transformation and in their disposition (organotropy). Variations in resistance and disposition are due to differences in cell regeneration rate, ability to repair nucleic acids; concentration of tissue-specific proteins (chaperones) that inhibit cell proliferation and tumor growth, increase in metabolism before and during malignant transformation, nonspecific mesenchymal reaction, immune response to tumor cells, and affinity of tissues to chemical carcinogens. In most cases, there is a positive correlation between the cell regeneration rate and the occurrence of malignant epithelial tumors. The low incidence of malignant tumors of the gingiva, floor of the mouth, tongue, and buccal mucosa, in spite of the high cell regeneration rate in these tissues, is due to the steady release of large numbers of granulocytes from the gingiva. (83 refs)

78-4919 **The Role of Unsaturated Bile Acids in the Etiology of Large Bowel Cancer.** (Eng) Hill, M. J. (Central Public Health Lab., London NW9 5DX, England). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation.* Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1627-1640; 1977.

The role of bile acids in large bowel cancer is reviewed. When populations in countries having different rates of large bowel cancer or populations in the same country but from different

socioeconomic levels were compared, the risk of large bowel cancer was found to correlate with the concentration of bile acids. The bile acids were more extensively degraded in stools from high-incidence populations than in those from groups with a lower incidence. The hypothesis that certain clostridia (referred to as NDC) are able to dehydrogenate the bile acid nucleus to produce unsaturated bile acids and that the combination of high concentrations of bile acid substrate and of NDC confers a high risk of colon cancer was tested and confirmed in a case-comparison study. A high fecal bile acid concentration alone characterized 83% of the large bowel cancer patients compared with only 19% of the comparison patients. Of the four combinations of fecal bile acid concentration and presence or absence of NDC, the combination of high fecal bile acids and NDC conferred the highest risk. Other studies have indicated that the number of organisms in stools producing cholanoil 7 α -dehydroxylase is correlated with the incidence of large bowel cancer and that 3-oxochola-4,6-dien-24-oic acid is the key etiological agent. (42 refs)

78-4920 Experimental Approaches to Benign Prostatic Hyperplasia. (Ger) Jacobi, G. H. (Urologische Universitätsklinik, Langenbeckstrasse 1, 6500 Mainz, W. Germany). *Aktuel Urol* 9(2): 73-80; 1978.

The normal and pathologic growth of the prostate gland is stimulated by testicular hormones. The concentration of dihydrotestosterone, the principal androgen in androgenic target tissues, is increased in benign prostatic hypertrophy (BPH) as a result of pathologic growth. The decrease in Leydig cell function with advancing age results in quantitative changes of the peripheral hormone pool (decrease in free testosterone, testosterone glucuronide dihydrotestosterone, and adrenal androgen level, and increase in testosterone-binding globulin and estradiol levels and in estradiol/testosterone and testosterone/dihydrotestosterone ratios). Thus, qualitative intracellular hormone alterations are most likely to be responsible for the development of BPH. Prostatic hypertrophy occurs in only two species, man and dog, which suggests the use of the latter as a model for this condition. Androstenediol induces unregulated adenomatous growth of the prostate in the dog. This hormone is a normal product of testicular secretion and prostatic metabolism. The increased formation of androstenediol in the prostate is paralleled by an increased growth of the gland. Cellular androstenediol synthesis is increased significantly in men with prostatic adenomas, compared with normal men. The findings suggest the direct role of androstenediol in the pathogenesis of BPH. (50 refs)

78-4921 Results of Research on Precancerous Changes in the Endometrium. (Rus) Zheleznev, B. I. (Moscow, USSR). *Akush Ginekol (Mosk)* (3): 10-17; 1978.

According to this review article, "precancer of the endometrium" (PCE) should include atypical hyperplasia of the func-

tional and/or basal layer of the endometrium, focal adenomatosis in glandular and basal hyperplasia, polyps, and adenomatous endometrial polyps. During the last 16 yr, PCE was diagnosed in 202 patients, other forms of hyperplasia in 1,006, and endometrial polyps in 2,031. Malignant transformation of atypical hyperplasia and focal adenomatosis was observed in 25/202 PCE patients. (no refs)

78-4922 Comparison of Neoplastic Hepatic Lesions in Man and Experimental Animals. (Eng) Popper, H. (Stratton Lab. Study of Liver Diseases, Mount Sinai Sch. Medicine, City Univ. New York, New York, NY, 10029); Maltoni, C.; Squire, R. A.; Thomas, L. B. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1359-1382; 1977.

The induction of hepatic tumors in animals and clinical observations of hepatic tumors in humans are compared. The progression from fibrotic lesions to angiosarcoma in man following exposure to vinyl chloride (VC), inorganic arsenicals, and Thorotrast is almost identical to changes observed in rodents treated with VC. This similarity in the evolution of hepatic lesions supports the predictive nature of animal experimental studies for human carcinogenesis. The major difference between the species was the greater fibroblastic reaction in man. The development of hepatocellular carcinoma in human and rodent nodules supports the concept of multiple cell populations in hepatocarcinogenesis. In the VC-associated lesions, the simultaneous proliferation of hepatocytes and sinusoidal cells, the latter potentially terminating in angiosarcoma, is stressed. The metabolic processes underlying this evolution remain to be established. (77 refs.)

78-4923 Tumor Heterogeneity and the Biology of Cancer Invasion and Metastasis. (Eng) Fidler, I. J. (Cancer Biology Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501). *Cancer Res* 38(9): 2651-2660; 1978.

The spread of tumor tissue through the lymphatic and hematogenous compartments of the circulatory system is discussed on the basis of experiments with the B16 melanoma in C57BL/6 mice. The development of a metastasis is dependent on an interplay between host factors and the intrinsic characteristics of malignant tumor cells. The process of metastasis is highly selective, and the metastatic lesion represents the end point of many destructive events that only a few cells can survive. Neoplasms, which are predominantly heterogeneous in nature, contain a variety of subpopulations of cells with differing metastatic potential. Furthermore, metastatic cell variants have been shown to preexist in murine neoplasms of old and recent origin. The possible existence of highly metastatic variant cells within a primary tumor suggests that

neoplasm should no longer be considered a uniform entity. The design of effective therapeutic agents should be directed toward the few but lethal metastatic subpopulations of cells. (3 refs.)

78-4924 **Progression and Genesis of Human Tumors** (Ita) Prodi, G. (Istituto Cancerologia, Bologna, Italy). *Recent Prog Med (Roma)* 63(6): 669-687; 1977.

Studies of the monoclonal nature of tumors and their preneoplastic stages, growth, differentiation and progression are reviewed. Tumor progression is irreversible, discontinuous, focal, and, in each individual case, unpredictable. (5 refs)

78-4925 **Introduction.** (Eng) Doll, R. (Univ. Oxford, Oxford OX2 6PS, England). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 1977.

The main clues to the origin of cancer that have been provided by epidemiological studies are reviewed. Included in the discussion are environmental agents, industrial pollution, diet, the interaction of agents, viruses, systemic factors, and genetic factors. (25 refs)

78-4926 **Genetic Predisposition to Cancer.** (Eng) Knudson, A. G. (Medical Genetics Center, Graduate School of Biomedical Sciences, Univ. Texas Health Science Center, Houston, TX, 77030). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 45-52; 1977.

In a review of the role of heredity and environment in the development of cancer, four groups of individuals can be recognized. The first group has a genetic predisposition to cancer even in the absence of environmental variation. The second group has a predisposition imposed by environmental variation in the absence of genetic variation. The third has a predisposition imposed by both genetic and environmental factors. The fourth group has neither a genetic nor environmental predisposition. Group 1, hereditary cancers, comprises only 1%-2% of all cancers but may comprise a large percentage of a particular cancer (eg, 40% for retinoblastoma). Several multiple mutation models have been proposed to explain these cancers, and most propose that there is an initial germinal mutation followed by a somatic mutation. In some cancers (ie, epithelial carcinomas of adulthood), a third event, perhaps a chromosome abnormality, may be required for the development of malignancy. The predisposition in the third group is represented by conditions such as xeroderma

pigmentosum and ataxia telangiectasia. Cancers of the second and third groups make up 70%-80% of all cancers in industrialized nations. The fourth group, comprising approx 25% of the total cancer incidence, results from spontaneous or background somatic mutations and represents the nonpreventable cancers for which diagnosis and therapy will always be important. (35 refs)

78-4927 **Genetic Regulation of Susceptibility to Polycyclic-hydrocarbon-induced Tumors in the Mouse.**

(Eng) Kouri, R. E. (Dept. Biochemical Oncology, Microbiological Associates, Bethesda, MD, 20016); Nebert, D. W. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 811-835; 1977.

The membrane-bound monooxygenase systems are reviewed, and the relationship between the murine aryl hydrocarbon (*Ah*) locus and increased susceptibility to chemical tumorigenesis, drug toxicity, and mutagenesis is discussed. A single allelic difference or a very small number of genes may have a profound influence on an individual's increased susceptibility to chemical carcinogenesis, drug toxicity, and congenital malformations. However, genes other than the *Ah* locus appear to influence an individual's risk for tumorigenesis. DNA repair, viral expression, and immunological tolerance may also be modifying factors in tumorigenesis. The interplay between the *Ah* locus and these factors, as well as the role of the factors themselves, in susceptibility to cancer can be determined hypothetically using an experimental approach outlined. (69 refs.)

78-4928 **Hereditary Factors in Human Cancer.** (Eng) Kellermann, G. (Dept. Human Oncology, Univ.

Wisconsin, Madison, WI, 53706). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 837-845; 1977.

The influence of genetic factors on human cancer is discussed. Epidemiologic data can be used to determine whether genetic factors are involved in certain common cancers. These data suggest that as much as 80% of all human cancers have some environmental element as their critical etiologic factor, but that the susceptibility to a particular cancer is inherited. Thus, the majority of carcinogens may be so-called precarcinogens that require enzymatic transformation to become real carcinogens. Interindividual variation of some of the activating enzymes might explain why some individuals are sensitive or resistant to the action of environmental carcinogens. For example, in vivo and in vitro studies demonstrate that the human population can be divided into slow and fast metabolizers of certain drugs and of the carcinogen

benzo(a)pyrene. Such studies have indicated that heavy smokers with a genetically high aryl hydrocarbon hydroxylase (AHH) activity are at greater risk for developing lung cancer than heavy smokers with low AHH activity, because the former produce more of the carcinogenic epoxides than the latter. A genetically high AHH activity may also be a factor in the etiology of laryngeal cancer. (22 refs.)

- 78-4929 Constitutional and Hereditary Factors in the Genesis of Human Tumors.** (Ita) Prodi, G. (Istituto Cancerologia, Bologna, Italy); Grilli, S. *Recent Prog Med (Roma)* 63(6): 622-640; 1977.

Studies of hereditary human tumors are reviewed. Nephroblastoma, retinoblastoma, neuroblastoma, neurofibromatosis, pheochromocytoma, colonic polyposis, and hereditary adenocarcinomatosis are the principal hereditary tumors. (37 refs)

- 78-4930 Prenatal Factors.** (Eng) Miller, R. W. (Clinical Epidemiology Branch, NCI, NIH, Bethesda, MD, 20014). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 381-388; 1977.

The association between human cancer and congenital malformations is discussed. Included in the discussion are xeroderma pigmentosum and ataxia telangiectasia, hereditary retinoblastoma with osteosarcoma, common features in persons at high risk of specific cancers, environmental causes of cancers and malformations, and the overlap at mutagenesis, carcinogenesis, and teratogenesis. (52 refs)

- 78-4931 Chromosomal Abnormalities in Human Neoplasia.** (Ger) Bartram, C. R. (I. Medizinische Klinik, Universität Hamburg, Martinistrasse 52, D-2000 Hamburg 20, W. Germany); Rudiger, H. W. *Klin Wochenschr* 56(15): 733-741; 1978.

Current knowledge on the association between chromosome aberrations and malignant tumors is reviewed. Specific topics covered include cancer-prone chromosome abnormalities, chromosome abnormalities related to tumors, and chromosome abnormalities related to carcinogens. Observations in humans are discussed in connection with in vitro experimental data. (92 refs.)

- 78-4932 Cellular Transformation and the 'Morphologic Phenotype' of Transformed Cells.** (Eng) Pastan, I. (NCI, Bethesda, MD, 20014); Willingham, M. *Nature* 274(5672): 645-650; 1978.

Changes in the adhesion, appearance, and surface properties of cells undergoing transformation are reviewed, with emphasis on the role of cyclic AMP (cAMP) and cell-surface protein (CSP) in this process. In most cases, these changes are linked to a diminished adhesion of the cell to the substratum. Treatment of transformed cells with cAMP has been reported to induce the flattened appearance characteristic of normal fibroblastic cells, increase their adhesion to a substratum, decrease their agglutinability by plant lectins, decrease the number of surface microvilli, and decrease growth rate; but it does not restore contact inhibition of growth. However, not all transformed cells respond to cAMP. The response of the parental untransformed cell usually indicates whether the transformed derivative will respond to cAMP. cAMP levels are reduced and adenylate cyclase levels are altered in some transformed cells. CSP is decreased in many types of transformed cells. In Rous sarcoma virus-transformed chick embryo cells, the decrease is primarily due to a decrease in the rate of synthesis of CSP, and it is accompanied by a reduction in the amount of translatable messenger RNA for CSP. The addition of highly purified CSP to cultures of transformed cells increased cell-substratum adhesion and produced cell elongation, but it did not restore growth control. Expression of the product of the transforming (*src*) gene of RNA tumor viruses promotes growth and usually produces the morphologic phenotype of transformation in cultured fibroblasts. The changes closely resemble those that occur when quiescent cells are stimulated to grow by addition of an appropriate peptide hormone, such as insulin. (82 refs.)

- 78-4933 Control of Normal Cell Differentiation and the Phenotypic Reversion of Malignancy in Myeloid Leukaemia.** (Eng) Sachs, L. (Dept. Genetics, Weizmann Inst. Science, Rehovot, Israel). *Nature* 274(5671): 535-539; 1978.

Cultured myeloid cells can be used to determine and dissect the controls that regulate the growth and differentiation of macrophages (M) and granulocytes (G) and the development of myeloid leukemia. Normal myeloid precursors can be induced to grow and differentiate to mature M and G by a protein M and G inducer (MGI) that is secreted by various types of cells, including fibroblasts and macrophages. One type of myeloid leukemia cell in both humans and mice can be induced to differentiate normally by MGI. In mice, this differentiation has also been demonstrated in vivo. This type of leukemia (MGI+D+, D+ for differentiation to mature cells) from different strains of mouse has been cloned in culture. Mature cells induced from MGI+D+ are no longer malignant in vivo and no longer multiply in vitro. MGI+D+ are viable and can grow in the absence of MGI, whereas normal cells require MGI for viability and growth. It is proposed that an inadequate supply of MGI can occur in vivo, possibly because of a decrease in MGI synthesis or the production of MGI inhibitors, that would limit the growth of normal cells without affecting the growth of leukemic cells. MGI+D+ cells do not have a normal diploid

karyotype, and the change in requirement for MGI for cell viability and growth was associated with a chromosome abnormality both in vitro and in vivo, suggesting that the origin of leukemia due to loss in requirement for MGI for viability and growth is genetic and due to a chromosomal change. Cell competence for induction of normal differentiation by MGI was associated with certain membrane qualities and also with the production and inducibility of C-type RNA virus. It is suggested that the mouse chromosomes (2 and 12) carrying the genes that control cell competence also carry genes that control the malignancy of the cells. (84 refs.)

78-4934 Incidence of Cancer in Population with Exceptional Exposure to Metals. (Eng) Hernberg, S. (Dept. Epidemiology and Biometry, Inst. Occupational Health, Helsinki 29, Finland). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory: Origins of Human Cancer. Vol. 4, 602 pp.; 147-157; 1977.

Experimental and human epidemiologic data on the carcinogenicity of arsenic, beryllium, cadmium, chromium, iron, nickel, and lead are reviewed. As causes skin cancer in humans and likely induces lung cancer as well. Lymphatic cancers and hemangiosarcoma of the liver also appear to be connected with As exposure. Experimental data indicate that iv Be salts cause osteogenic sarcomas in rabbits and mice and Be aerosols cause pulmonary cancer in rats and monkeys. Epidemiological studies of humans, however, are inconclusive. Cd appears to increase the risk of prostate, renal, and respiratory cancer in humans but more studies are necessary before studies are necessary before definite conclusions can be drawn. Studies with hexavalent Cr have shown an excess of respiratory cancer in chromate workers, but the exact identity of the compound or compounds responsible for the carcinogenesis is unknown. Furthermore, there are no conclusive negative epidemiologic studies of workers exposed to trivalent chromium. Data on the carcinogenicity of Fe in humans is inconclusive, even though Fe-polysaccharide complexes have induced local sarcomas in experimental animals. There is strong epidemiologic evidence supporting the carcinogenicity of Ni compounds, but the exact identity of the compounds involved is not certain. Studies of the carcinogenicity of Pb have not shown any positive results. (89 refs)

78-4935 Effects of Urethane Administration on the Sedimentation Patterns of Mouse and Rat Liver DNA in Alkaline Sucrose Gradients. (Eng) Floom, B. G. (Chemical Carcinogenesis Div., Antoni van Leeuwenhoek-Huis, Netherlands Cancer Inst., Amsterdam, Netherlands); Philippus, E. J.; Scherer, E.; Den Engelse, L. *Chem Biol Interact* 21(2/3): 331-336; 1978.

The effect of urethane on the alkaline sucrose sedimentation pattern of mouse and rat liver DNA was determined. C3Hf/A male mice and Sprague-Dawley female rats were

two-thirds hepatectomized and treated 3 wk later (24 hr before sacrifice) with 1 mg/g urethane ip. No significant changes in the DNA sedimentation pattern were observed. Thus, the reported formation of alkali-labile ethyl phosphotriesters in DNA by urethane is at most a minor process of alkylation. (10 refs.)

78-4936 Significance of Repair of Human DNA: Evidence from Studies of Xeroderma Pigmentosum.

(Eng) Robbins, J. H. (Dermatology Branch, NCI, NIH, Public Health Service, US Dept. Health, Education, and Welfare, Bethesda, MD, 20014). *J Natl Cancer Inst* 61(3): 645-656; 1978.

Recent studies relating certain clinical manifestations of xeroderma pigmentosum (XP) to the patient's defective DNA repair are reviewed. XP patients develop excessive freckling on sun-exposed skin at an early age (usually before 2 yr) and this frequently leads to tumor development. The tumors occur at the same frequency as, and are indistinguishable from, tumors in patients without XP, but XP patients can develop hundreds of these tumors by the age of 30. Cultured dermal fibroblasts from all excision repair-deficient XP cells show a reduced rate of UV-induced unscheduled DNA synthesis. Furthermore, an increased frequency of UV-induced sister chromatid exchanges exists in these cells, possibly as a result of unrepaired DNA damage. The primary inherited DNA repair defect has not been established in any form of XP. Neurologic damage accompanying XP appears to be the result of UV-induced DNA damage. It is hypothesized that UV radiation-induced damage in XP cells requires the same process(es) for repair as the currently unknown damage postulated to occur in the DNA of the CNS of XP and normal individuals during embryonic and postnatal life. The XP neurons are not repaired as efficiently as normal neurons and, although the neuronal DNA does not replicate after the second postnatal year, proper functioning of this DNA is necessary for neuron survival. Cell death eventually results from the UV-induced damage. The acute sun sensitivity of some XP patients appears to be the result of excessive killing or damage of skin cells as a result of inadequate DNA repair. An analysis of the various XP complementation groups and of the biochemical and physicochemical abnormalities in XP cells is also presented. (120 refs)

78-4937 Role of Cyclic Adenosine-3,5-monophosphate in Tumorigenesis. (Rus) Simkhovich, B. Z. (Inst. Organic Synthesis, Riga, USSR); Lidak, M. Iu.; Gilev, A. P. *Khim Farm Zh* 12(6): 17-31; 1978.

Literature on the role of cyclic AMP (cAMP) in cell metabolism and differentiation is reviewed with special attention to alterations in the cAMP system during tumorigenesis. The characteristic feature of a series of malignant tumors was an extremely low level of cAMP. The cAMP concentration was also decreased after exposure of tissue to carcinogens. Recent studies show that cAMP can inhibit tumor growth. (142 refs.)

CHEMICAL CARCINOGENESIS

78-4938 Carcinogenicity and Bacterial Mutagenicity of 47 Structurally Related Heterocyclic Compounds (Meeting Abstract). (Eng) Schwind, H. (Pharmakologisches Institut der Universität, D-6500 Mainz, W. Germany); Oesch, F.; Zajdela, F.; Jacquignon, P.; Croisy, A.; Perin, F.; Glatt, H. *Experientia* 34(7): 918; 1978. (no refs)

78-4939 Sensitivity to Five Mutagens in Fanconi's Anemia as Measured by the Micronucleus Method. (Eng) Heddle, J. A. (Dept. Natural Science, York Univ., Toronto, Ontario M3J 1P3, Canada); Lue, C. B.; Saunders, E. F.; Benz, R. D. *Cancer Res* 38(9): 2983-2988; 1978.

The micronucleus technique was used to measure the sensitivity of Fanconi's anemia (FA) patients to mitomycin C (MMC), ethyl methanesulfonate, methyl methanesulfonate, x-rays, and 5-bromodeoxyuridine. Blood samples from four groups of patients were cultured: 4 patients with FA, 1 patient with aplastic anemia of uncertain etiology, 6 carriers of the FA gene (parents of the patients), and 6 normal controls. MMC treatment of cultures from the FA patients resulted in an increase in micronuclei formation that was readily distinguishable from the response of the other groups. This corresponded to a tenfold increased sensitivity to chromosomal breakage by the cross-linking agent. The micronucleus technique was also successful at detecting the increased spontaneous rate of chromosome breakage in FA. None of the other groups differed significantly in response to MMC or to the other mutagens. FA cells were no more sensitive to the other agents than were cells of normal controls, except for a 50% increase in sensitivity to x-rays. These findings indicate that the micronucleus technique can be used to diagnose cases of FA and to extend the information available on the sensitivity of various genotypes to chromosomal damage by various agents. (32 refs.)

78-4940 Mammalian Cell Transformation In Vitro. (Eng) Styles, J. A. (Imperial Chemical Industries Ltd., Central Toxicology Lab., Alderley Park, Cheshire, England). *Br J Cancer* 37(6): 931-936; 1978.

The use of in vitro transformation as a rapid test to detect carcinogenic chemicals was evaluated. WI-38 (human lung), Chang (human liver), and BHK-21 (baby Syrian hamster kidney) cells were used to test 120 organic chemicals. The assay was 94% accurate in determining carcino-

genic activity using two cell lines. The predictive figure was only slightly reduced when one cell type was used (88% for WI-38 and Chang, and 91% for BHK-21). Only 11 compounds were detected with the use of one cell type. Thus, species or organ of origin of the tester cells is of no significance in this assay, and one cell type offers no great predictive advantage over another. Comparison of the data revealed that 106 compounds were correctly identified by both the mammalian cell transformation and bacterial-mutation tests. Of the remaining 14 compounds, 7 were correctly identified only by the transformation test and 5 were identified only by the bacterial-mutation test. Few carcinogens transformed the human cells without auxiliary metabolism by a rat liver homogenate. (40 refs)

78-4941 The Usefulness of Mutagenicity Tests as Pre-screening Cancer Tests: Comparing "Host-Mediated Assay" and "Mammalian Spot Test". (Ger) Fahrig, R. (Freiburg, W. Germany). *Staub-Reinhalt Luft* 38(6): 232-235; 1978.

The usefulness of two mutagenicity test systems for pre-screening chemical carcinogens was compared: (1) the host-mediated assay (yeast/rat) for detecting the organo-specificity of carcinogens and (2) the in vivo mammalian spot test for detecting genetic alterations, especially point mutations in murine somatic cells. Parallel testing of carcinogens demonstrated that the spot test was not only capable of identifying carcinogens and their target organs, but also required a lower carcinogen dose than the host-mediated assay. In the host-mediated assay, yeast cells were injected into rat testicles and tail veins. Within min, the tail-vein injected yeast cells were filtered from the blood and transported to organs such as the liver, lungs, and spleen. Subsequently, the yeast cells were retrieved from testicles, liver, and lungs and examined for genetic alterations. Of the 15 mutagens or carcinogens tested, 6 were most active in the lungs and 9 in the liver. Six mutagens did not appear in the testicles and 2 did not appear in either the testicles or liver. Not all mutagens reacted in this assay, but in the spot test, all compounds reacted, even those that induced genetic mutations exclusively. (45 refs.)

78-4942 Carcinogenic Activity of *Symphytum officinale*. (Eng) Hirono, I. (Dept. Carcinogenesis and Cancer Susceptibility, Inst. Medical Science, Univ. Tokyo, Shirokanedai, Minato-ku, Tokyo 108, Japan); Mori, H.; Haga, M. *J Natl Cancer Inst* 61(3): 865-869; 1978.

The carcinogenicity of *Symphytum officinale* L., or Russian comfrey, which is used as a green vegetable or a tonic in Japan, was studied in inbred ACI rats. Three groups of 19-28 rats were fed comfrey leaves (8%, 16%, or 33% of diet) for 80-600 days; four additional groups of 15-24 rats were fed comfrey roots (0.5%-8% of diet) for varying lengths of time. A control group was given a normal diet. Hepatocellular adenomas were induced in all experimental groups that received the diets containing comfrey roots and leaves. Tumor incidence ranged from 4% (group fed leaves at 8% of diet for 600 days) to 86.3% (group fed roots at 8% of diet until death). Hemangioendothelial sarcomas of the liver were also induced at a low incidence. The results show that comfrey roots were much more toxic than the leaves. They strongly suggest that the pathologic changes in the liver were induced by the pyrrolizidine alkaloids present in comfrey. (13 refs.)

78-4943 **The Implant Test.** (Eng) Longstaff, E. (No affiliation given). *Br J Cancer* 37(6): 954-958; 1978.

A technique is described whereby the histological changes observable after the sc implantation of filter disks containing test material are used to predict the carcinogenic potential of the material. An irritant (a Millipore filter of specific pore size) was included in the implant, and the test compound was suspended in gelatin to induce the production of proteolytic enzymes. The disks were implanted into the dorsolumbar region of 2-mo-old Alderley Park Swiss mice. Control implants contained dimethyl sulfoxide but no added test material. The carcinogenic potential of 76/111 compounds was predicted correctly. Only 19/52 carcinogens were positively identified, but there were only 2 false-positive results among the 59 non-carcinogenic compounds tested (3% false-positive rate). Because the implant test proved insensitive and required a protracted duration of study for little return, it would be inadequate as a routine test for identifying new carcinogens. However, it shows promise as a model for the regular production of sc sarcomas and carcinomas. (8 refs)

78-4944 **Use of the Fluctuation Test to Detect Genetic Activity of Environmental Chemicals in *Aspergillus nidulans*** (Meeting Abstract). (Eng) Kappas, A. (Dept. Biology, Nuclear Res. Center, Democritus, Athens, Greece); Demopoulos, N. *Mutat Res* 53(2): 208; 1978. (1 ref)

78-4945 **Intravenous Host-mediated Assay for Detection of Mutagenesis with *Saccharomyces cerevisiae*** (Meeting Abstract). (Eng) Frezza, D. (Lab. Environmental Mutagenesis, NIEHS, Research Triangle Park, NC, 27709); Leiger, E. *Environ Health Perspect* 20: 235; 1977. (no refs)

78-4946 **Description and Calibration of a Forward Mutation System in Salmonella** (Meeting Abstract). (Eng) Pueyo, C. (Lab. Environmental Mutagenesis, NIEHS, Research Triangle Park, NC, 27709). *Environ Health Perspect* 20: 235; 1977. (no refs)

78-4947 **Effect of Excision Repair-Deficient Mutation *uvr2* on the Mutagenic and Killing Activities of 7 Chemical Mutagens in Heterokaryons of *Neurospora crassa*** (Meeting Abstract). (Eng) Brockman, H. E. (Illinois State Univ., Normal, IL, 61561); Hung, C. Y.; Ong, T. M.; de Serres, F. J. *Mutat Res* 53(2): 156-157; 1978. (3 refs)

78-4948 **Chemical Mutagenesis in UV-sensitive Strains of *Neurospora crassa*** (Meeting Abstract). (Eng) Inoue, H. (NIEHS, Research Triangle Park, NC, 27709); Ong, T. M.; De Serres, F. J. *Environ Health Perspect* 20: 235; 1977. (no refs)

78-4949 **Test for Repair Replication of DNA In Vivo in Liver of Animals Treated with Carcinogens** (Meeting Abstract). (Eng) Craddock, V. M. (MRC Toxicology Unit, Woodmansterne Road, Carshalton, Surrey, England). *Br J Cancer* 38(1): 170; 1978. (1 ref)

78-4950 **The Effect of Bacterial Plasmids on Mutagenesis** (Meeting Abstract). (Eng) Monti-Bragadin, C. (Inst. Microbiology, Univ. Trieste, Trieste, Italy); Babudri, C. N.; Tamaro, M.; Cinco, M.; Molina, A. M. *Mutat Res* 53(2): 231-232; 1978. (no refs)

78-4951 **Mutation Induction by Indirectly Acting Mutagens in *Drosophila* Larvae** (Meeting Abstract). (Eng) Kortselius, M. J. (Dept. Radiation Genetics and Chemical Mutagenesis, State Univ. Leiden, Leiden, Netherlands); Vogel, E. *Mutat Res* 53(2): 212; 1978. (no refs)

78-4952 **Mutagenicity of Intestinal Bacteria.** (Eng) Suzuki, K. (Inst. Physical and Chemical Res., Wako-shi, Saitama 351, Japan); Mitsuoka, T. *Mutat Res* 50(2): 295-299; 1978.

The mutagenicity of 45 intestinal bacterial strains was tested by the rec-assay, which measures the lethal effects on a recombination-deficient *Bacillus subtilis* strain (M45 rec-). Of the 16 strains that showed lethal activity, 8 were Bifidobacteria strains. Positive strains that are predominant species in

the human intestine included *Bifidobacterium adolescentis*, *Bifidobacterium infantis*, *Bifidobacterium longum*, *Bacteroides fragilis* ss. *fragilis*, *Bacteroides multiacidus*, and *Eubacterium aerofaciens*. *Bacteroides multiacidus*, *Peptostreptococcus parvulus*, *Streptococcus faecium*, and *Veillonella parvula* were mutagenic for *Salmonella typhimurium* strain TA1535, indicating that they induce base-pair substitution mutations. (13 refs)

- 78-4953 **Mutagenicity Tests with Styrene Oxide in Mammals.** (Eng) Fabry, L. (Mammalian Genetics Lab., Dept. Radiobiology, CEN-SCK, B-2400 Mol, Belgium); Leonard, A.; Roberfroid, M. *Mutat Res* 51(3): 377-381; 1978.

The capability of styrene oxide to induce chromosomal damage was determined using four mammalian test systems in vivo and one in vitro. Adult male BALB/c mice were given a single ip injection of 250 mg/kg styrene oxide, and the induction of chromosomal aberrations in the bone marrow cells, the production of micronuclei in polychromatic RBC, and the effects on the spermatocytes and postmeiotic germ cells were assayed. Short-term cultures of human lymphocytes from a single donor were treated in vitro with 0.1 or 0.5 mM styrene oxide, and the effects on the chromosomes were determined. There was a significant increase in the total number of anomalies after treatment of the lymphocytes with 0.5 mM styrene oxide; this was due mainly to an increase in chromatid gaps. All tests on somatic and germ cells in vivo yielded negative results. Thus, styrene oxide appears to be capable of producing genetic damage in cells in vitro, but it could be less effective in vivo. (14 refs.)

- 78-4954 **The Induction of Tumours of the Digestive Tract and of Certain Other Organs in Rats Given T-2 Toxin, a Secondary Metabolite of *Fusarium sporotrichioides*** (Meeting Abstract). (Eng) Schoental, R. (Royal Veterinary Coll., London, England); Joffe, A. Z.; Yagen, B. *Br J Cancer* 38(1): 171; 1978. (1 ref)

- 78-4955 **Carcinogenicity of Betel Quid Ingredients Singly or in Combination Using Different Experimental Models** (Meeting Abstract). (Eng) Gothoskar, S. V. (Biology Div., Cancer Res. Inst., Tata Memorial Centre, Bombay 400012, India); Ranadive, S. N.; Shivapurkar, N. M.; Ranadive, K. J. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 66-67; 1977. (no refs)

- 78-4956 **Fungal Contamination and Mycotoxin Detection of Powdered Herbal Drugs.** (Eng) Hitokoto, H. (Dept. Microbiology, Tokyo Metropolitan Res. Lab. Public Health, Shinjuku-ku, Tokyo, 160, Japan); Morozumi, S.; Wauke, T.; Sakai, S.; Kurata, H. *Appl Environ Microbiol* 36(2): 252-256; 1978.

A total of 49 powdered herbal drugs were analyzed for their mold profile and for the potential presence of aflatoxin, sterigmatocystin, and ochratoxin A. Mycotoxins were not detected in any samples. Of 69 strains of fungus isolated from the drugs, only one strain of *Aspergillus flavus* was capable of producing aflatoxin (type B₁, 0.15 µg/ml) on laboratory media. The drugs contained predominantly *Aspergillus* and *Penicillium* fungi. (14 refs.)

- 78-4957 **Differential Effect of Aflatoxin B₁ on Ribosomal RNA Synthesis of Rat Liver Depending on the Carcinogenic or Toxic Activity of the Compound** (Meeting Abstract). (Eng) Novi, A. M. (Pathologisches Institut, Universitat Dusseldorf, Moorenstrasse 5, D-4000 Dusseldorf, W. Germany); Wagner, G. *Hoppe Seylers Z Physiol Chem* 359(9): 1128; 1978. (2 refs)

- 78-4958 **Mycotoxins. Thin Layer Chromatographic Determination of Aflatoxin B₁ in Eggs: Collaborative Study.** (Eng) Nesheim, S. (Div. of Chemistry and Physics, Food and Drug Admin., Washington, DC, 20204); Trucksess, M. W. *J Assoc Off Anal Chem* 61(3): 569-573; 1978.

The effectiveness of two-dimensional thin-layer chromatography for determining aflatoxin B₁ (AFB₁) levels in hens eggs was tested collaboratively. Nine collaborators analyzed three known practice samples and nine unknown samples containing 0, 0.05, 0.10, or 0.30 nanogram (ng)/g AFB₁. Recoveries for the three positive levels were: 0-0.13 [av 98%, coefficient of variation (CV) 83%], 0.05-0.18 (av 102%, CV 36%), and 0.11-0.42 ng/g (av 93%, CV 31%), respectively. (6 refs)

- 78-4959 **Use of High Performance Liquid Chromatography Combined with Fluorescence Detection for the Identification and Estimation of Aflatoxins and Ochratoxin in Food.** (Eng) Hunt, D. C. (Dept. Industry, Lab. Government Chemist, Cornwall House, Stamford St., London SE1 9NQ, England); Bourdon, A. T.; Wild, P. J.; Crosby, N. T. *J Sci Food Agric* 29(3): 234-238; 1978.

A method was developed for the separation, identification, and estimation of aflatoxin B₁ (AFB₁), AFB₂, AFG₁, AFG₂, AFM₁, and ochratoxin A in foods at the µg/kg level. The food is extracted with aqueous acetonitrile, washed with

2,2,4-trimethylpentane, and any mycotoxins are reextracted into chloroform. The extract is purified by thin-layer chromatography. Any fluorescent bands are then removed and examined by high-pressure liquid chromatography. Recoveries at twice the detection limit varied from 40% to 95%. (8 refs)

78-4960 Effect of Hypophysectomy and Aflatoxin B₁ on Rat Liver. (Eng) Neal, G. E. (Toxicology Unit, MRC Labs., Woodmansterne Road, Carshalton, Surrey SM5 4EF, England); Judah, D. J. *Cancer Res* 38(10): 3460-3467; 1978.

The effect of hypophysectomy (HPS) on aflatoxin B₁ (AFB₁)-induced liver carcinogenesis was studied in male Fischer rats. Four groups of rats were used: weanlings, small and large hypophysectomized (HPS-x) rats, and adult intact rats. Ingestion of 4 ppm AFB for 4-5 wk resulted in minimal changes in the livers of weanlings and HPS-x rats, but caused massive necrosis in the livers of the adult intact animals. This effect was not related to the amount of diet consumed. The liver necrosis in the adult intact animals may have been due to the presence of a large tetraploid hepatocyte population. The absence of these cells could account for the resistance of the weanlings. The considerable tetraploid hepatocyte population present in the large HPS-x rats was not susceptible to the necrogenic action of AFB₁ either fed to the rats or given as ip injections (2.0 mg/kg). Hepatic nucleic acid synthesis was reduced by HPS or the administration of AFB. In both cases, the effect was more marked in the tetraploid than in the diploid hepatocytes. Nucleic acid synthesis in the HPS-x rats was more resistant to inhibition by AFB than that in intact rats. This appeared to be due to the lower basal level of nucleic acid synthesis in the former, the final inhibited levels being similar in the two groups. The ability of the microsomal systems obtained from HPS-x rats to activate AFB did not appear to be significantly less than the ability of microsomes isolated from control rats. There may be a close correlation between the uptakes of AFB and steroid hormones into the liver, with the uptake mechanisms possibly being located in the tetraploid hepatocytes. HPS could protect against the effects of AFB by removing the steroid uptake mechanisms by which AFB enters the subcellular structures. (15 refs.)

78-4961 Aflatoxin Contamination of Grains in Flooded Areas of Mathura, Uttar Pradesh (Letter to Editor). (Eng) Mishra, R. S. (P.G. Dept. Botany, Bhagalpur Univ., Bhagalpur 812 007, Bihar, India); Singh, R. S. *Curr Sci (Bangalore)* 47(11): 396-397; 1978.

Thirty-six grain samples from fields that had been inundated by heavy rains were studied for mycotoxin contamination. *Aspergillus*, *Penicillium*, and *Rhizopus* were the main fungi associated with the samples. Aflatoxins were found in 18 samples. Aflatoxin B₁ was the most prevalent toxin (6-200

ppb); few samples contained aflatoxins B₂, G₁, and G₂, and the levels were < 10 ppb. (8 refs.)

78-4962 Identification of the Principal Aflatoxin B₁-DNA Adduct Formed In Vivo in Rat Liver. (Eng) Croy, R. G. (Dept. Nutrition and Food Science, Massachusetts Inst. Technology, Cambridge, MA, 02139); Essigmann, J. M.; Reinhold, V. N.; Wogan, G. N. *Proc Natl Acad Sci USA* 75(4): 1745-1749; 1978.

The products of in vivo covalent binding of activated aflatoxin B₁ (AFB₁) to DNA were investigated in male Fischer rats. The principal covalent product formed in liver DNA of rats treated with AFB₁ (2.0 mg/kg, ip) was identified as 2,3-dihydro-2-(N⁷-guanyl)-3-hydroxyaflatoxin B₁. This compound was isolated in sufficient quantity for characterization by physicochemical techniques, including field-desorption mass spectrometry. This information, together with results of chemical methylation of the compound, proved that the major adduct formed between DNA and AFB₁ in vivo is identical to that produced in vitro when AFB₁ is incubated with DNA in the presence of a rat liver microsome activation system. Quantitative studies of the formation of this compound revealed a dose-dependent relationship between the level of its occurrence in liver DNA and AFB₁ doses over the range 0.125-1.0 mg/kg. (31 refs)

78-4963 Effect of Vitamin A Deficiency and Excess on Aflatoxin Metabolism in the Rabbit. (Eng) Bas-sir, O. (Dept. Biochemistry, Univ. Ibadan, Ibadan, Nigeria); Adekunle, A. A.; Okoye, Z. S. *Biochem Pharmacol* 27(5): 833-838; 1978.

The effect of a diet depleted of or supplemented with vitamin A on the in vitro metabolism of aflatoxin was investigated using liver fractions from three groups of weanling male rabbits. Group 1 was fed a vitamin A-deficient diet, and Groups 2 (controls) and 3 were fed a diet supplemented with 5,000 and 60,000 IU retinylpalmitate/kg diet, respectively. The animals were killed after the 28-day feeding period, and the liver fractions were isolated. During the feeding period, the liver vitamin A content of Group 1 decreased to 45% of that in Group 2, but the vitamin A content in the livers of Group 3 animals increased to 340% of that in Group 2. Only aflatoxin B₁ demethylase activity was significantly reduced in the marginally deficient rabbits. However, for both hydroxylation and demethylation of aflatoxin, the value for Group 3 tended to be less than that of Group 2, and aflatoxin B₁ hydroxylase activity tended to be higher in Group 1 than in Group 2. It appears that a gross vitamin A deficiency is not necessary to reduce aflatoxin B₁ demethylation in the rabbit. (29 refs)

78-4964 The Estimation of Aflatoxin M₁ in Milk Using a Two-dimensional Thin-layer Chromatographic Method Suitable for Survey Work. (Eng) Patterson, D. S. (Central Veterinary Lab., Weybridge, Surrey KT15 3NB, England); Glancy, E. M.; Roberts, B. A. *Food Cosmet Toxicol* 16(1): 49-50; 1978.

An existing method for the analysis of aflatoxin M₁ in cow's milk was modified by the introduction of two-dimensional thin-layer chromatography (TLC) and certain confirmatory tests. The smallest quantity of aflatoxin M₁ that is easily visible on two-dimensional TLC is 0.2 nanogram when it is applied in 5 μ l chloroform. This corresponds to a quantitative analytical limit of 0.052 μ g aflatoxin M₁/liter milk, but even smaller amounts can be detected if they are not quantitated, so that the detection limit is 0.03 μ g/liter. This procedure is sufficiently sensitive to detect any aflatoxin M₁ carried over into milk from aflatoxin B₁ present in the cattle feed at the very low max level permitted by current regulations in the United Kingdom. (5 refs.)

78-4965 Determination of Aflatoxins in Peanut Products Using Reverse Phase HPLC. (Eng) Hurst, W. J. (Hershey Foods Res. Labs., P.O. Box 54, Hershey, PA, 17033); Toomey, P. B. *J Chromatogr Sci* 16(8): 372-376; 1978.

High-performance liquid chromatography was used to determine aflatoxin B₁, B₂, G₁, and G₂ in peanut products. This procedure can be applied to roasted and unroasted peanuts and to sweetened peanut matrixes with no interferences in the chromatography. It allowed the quantitative detection of all four aflatoxins at the 5- to 7-nanogram/kg level in <1.5 hr. (32 refs.)

78-4966 Desmofibrinogenic Activity of Transamidases and Thiol Groups in Plasma and Tissues after Poisoning with Aflatoxin B₁. (Pol) Zuch, A. (Zaklad Patologii Ogoinej i Doswiadczalnej AM, Instytut Fizjologii i Biochemii, ul. Mickiewicza 2c, 15-230 Bialystok, Poland); Januszko, T.; Poplawska, T. *Pol Arch Med Wewn* 59(2): 171-181; 1978.

Ip injection of aflatoxin B₁ (total dose: 0.56 mg/kg) into guinea pigs decreased the concentration and activity of plasma transamidases (FSF, coagulation factor XIII). A parallel transient decrease in desmofibrinogenic activity and in the concentration of lung and liver tissue transamidases was also observed. The increase in fibrinolytic activity and decrease in plasma levels of free thiol groups lasted longer (up to 10 days) than the changes in FSF. (18 refs)

78-4967 Microsomal Activation of Aflatoxin B₁ and G₁ Metabolites Mutagenic in *Neurospora crassa* (Meeting Abstract). (Eng) Ong, T. M. (Lab. Environmental Mutagenesis, NIEHS, Research Triangle Park, NC 27709). *Environ Health Perspect* 20: 234; 1977. (no refs)

78-4968 Aflatoxin Observed by Fluorescent Microscopy in Primary Liver Cancer Patients from Zaire. (Fre) Stora, C. (I.N.S.E.R.M., F.R.A. No. 12, Faculte de Medecine de Nice, Chemin de Vallombrose, 06034 Nice Cedex, France). *C R Acad Sci [D] (Paris)* 286(11): 917-920; 1978.

Biopsy samples from inhabitants of Zaire with hepatomas or cholangiocarcinomas of the liver were examined by fluorescence microscopy for the presence of aflatoxin B or G. The hepatoma tissue showed the blue fluorescence characteristic of aflatoxin B. Both the cytoplasm and nuclei of the tumor cells were highly fluorescent, with the most intense fluorescence being exhibited by the chromosomes. The cholangiocarcinoma cells were only slightly fluorescent; the fluorescence was observed in newly formed vessels and in the cytoplasm of the macrophages, which were abundant in zones of necrosis. Aflatoxin G was noted by an intense yellow-green fluorescence in the connective tissue surrounding the tumor nodes. No difference in fluorescence was observed between aflatoxins B₁ and B₂ or between G₁ and G₂ and their metabolites. (9 refs)

78-4969 Hepatic Transformation of Aflatoxin and Its Carcinogenicity. (Eng) Hsieh, D. P. (Dept. Environmental Toxicology, Coll. Agricultural and Environmental Sciences, Univ. California, Davis, CA, 95616); Wong, J. J.; Wong, Z. A.; Michas, C.; Ruebner, B. H. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conference on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp; 697-707; 1977.

The relationship between the carcinogenicity and mutagenicity of aflatoxins (AF's) is reviewed. The Ames mutagenicity assay was performed on several AF's and their metabolites. All the AF's tested required the presence of a rat liver post-mitochondrial supernatant (S-9) for mutagenic activity, indicating that none of them is the active form and that all are precarcinogens. Aflatoxin B₁ (AFB₁) had the highest mutagenic activity, followed by AFL, AFG₁, AFM₁, and other derivatives. There was a good correlation between the in vitro bacterial mutagenicity of these compounds and their in vivo animal carcinogenicity. Hepatic S-9 systems for in vitro assays of AFB₁ bioactivation were obtained from duck, rat, monkey, mouse, and human livers. The in vitro bioactivation

activity of the liver from a particular species for AFB₁ correlated with the known sensitivity of that species to the carcinogenic effect of AFB₁. This was evident from the consistent ranking of the relative mutagenic activities of AFB₁ generated by the hepatic S-9 enzymes of the different species (duck > rat > monkey > mouse > human) and the relative susceptibilities of the species to carcinogenic aflatoxicosis. The data show that humans are less susceptible to carcinogenic aflatoxicosis than the relatively resistant monkey and mouse. (31 refs.)

78-4970 Alpha Benzene Hexachloride Inhibition of Aflatoxin B₁-induced Hepatocellular Carcinoma. A Preliminary Report. (Eng) Angsubhakorn, S. (Dept. Pathobiology, Faculty Science, Ramathibodi Hosp., Mahidol Univ., Bangkok 4, Thailand); Bhamarapavati, N.; Romruen, K.; Sahaphong, S.; Thamavit, W. *Experientia* 34(8): 1069-1070; 1978.

The effect of α -benzene hexachloride (BHC) on the induction of hepatocellular carcinoma by aflatoxin B₁ was studied in male albino Fisher rats. Rats were fed a diet containing 1 ppm AFB and/or 0.05% BHC for 10 or 20 wk. Rats were sacrificed at 10 or 65 wk. In animals sacrificed after 10 wk, the only histological change noted was hypertrophy of the hepatocytes in the centrilobular region in the BHC and BHC + AFB groups. The liver wt:body wt ratio in the BHC and BHC + AFB groups was significantly higher than that in the AFB and normal diet groups. In the long-term experiment, tumors were noted in all three AFB group rats that survived until 65 wk, as well as in two rats that died during weeks 56 and 59. Tumors were not detected in any of the BHC, BHC + AFB, or control rats that survived the experiment. Rats from these three groups that died during the experiment had no signs of preneoplastic change in the liver. The liver wt:body wt ratio of the AFB-fed rats was greater than that of rats fed BHC, BHC + AFB, or a control diet. These results suggest that BHC prevent the induction of liver tumors by AFB. (8 refs.)

78-4971 Relationship Between Lipids and Scleroproteins in Asbestosis and Anthracosis. (Rus) Belobragina, G. V. (Inst. Vocational Hygiene and Occupational Diseases, Sverdlovsk, USSR); El'nichnykh, L. N. *Gig Tr Prof Zabol* (3): 31-34; 1978.

In an attempt to evaluate the possible role of phospholipids in the pathogenesis of asbestosis and anthracosis, asbestos and anthracite dusts were administered intratracheally (50 mg) to albino rats. Rats were sacrificed within 7 days-17 mo of exposure, and the lung tissue was examined. Single administration of asbestos and anthracite dusts induced mild perinodular fibrosis and the gradual development of diffuse fibrosis. The total lipid level in the lung tissue of exposed rats

was significantly greater than that in controls; the accumulation of total lipids and especially, phospholipids exceeded the accumulation of fibroproteins. These findings indicate that the relationships between total lipids and fibroproteins in asbestosis and anthracosis are similar to those in the development of presenile fibrosis. (21 refs)

78-4972 Carcinogens and Analogs: Trace Analysis of Thirteen Compounds in Admixture in Wastewater and Human Urine. (Eng) Nony, C. R. (Dept. Health, Education and Welfare, Food and Drug Admin., Natl. Center Toxicological Res., Jefferson, AR, 72079); Bowman, M. C. *Int J Environ Anal Chem* 5(3): 203-220; 1978.

Traces of 13 carcinogens and related compounds (aromatic amines and estrogens) in admixture in wastewater and human urine were determined by gas chromatography. Residues were detectable at the 0.1- and 1-ppb levels in the two liquids, respectively. Partition values for all pentafluoropropionyl derivatives in five solvent systems, structure verification by mass spectrometry, and adaptation of the method to work surfaces and air are provided. (18 refs)

78-4973 Occupationally Related Incidence of Double Cancers: Four Cases of Transitional Cell Carcinoma of the Urinary Bladder with Other Cancers. (Jpn) Matsushima, M. (Dept. Urology, Sch. Medicine, Toho Univ., Toho, Japan); Fukazawa, K.; Yagishita, T.; Hirose, K.; Miura, K.; Matsumoto, H.; Ando, K. *Acta Urol Jpn* 24(5): 409-416; 1978.

Four of 20 male dyestuff workers with transitional cell carcinoma (TCC) of the urinary bladder also had another occupationally related cancer. The 20 cases of TCC (including 1 of asynchronous bilateral ureteral tumors) were found among 90 workers who had been exposed to aromatic amines and who were subjected to periodic urine cytology tests. The four patients with metachronous double cancers were a 56-yr-old with TCC and an adenocarcinoma of the left lung, a 56-yr-old with TCC and a hepatoma, a 43-yr-old with TCC and a bile duct adenocarcinoma, and a 61-yr-old with TCC and a non-hormonal adrenocortical cancer. (28 refs.)

78-4974 Enzymatic Changes in Peripheral Blood Leukocytes in Rats in Subacute Benzene Vapours Poisoning. Acid Phosphatase Activity in Lymphocytes. (Eng) Moszczynski, P. (Internal Diseases Dept., ZOZ, 32-800 Brzesko, Poland); Starek, A. *Pol J Pharmacol Pharm* 30(1): 21-27; 1978.

Acid phosphatase (AcP) activity, its intracellular distribution in peripheral blood lymphocytes, and the activity of the lysosomal isoenzyme of acid phosphatase (AcP-L) were investigated in the serum of rats exposed to benzene vapor (27,000 mg/m³, 6 hr daily, for 10 consecutive days). There was a statistically significant fall in total lymphocytes, AcP+ and AcP- lymphocytes, and granular lymphocytes and an increase in granular-diffuse and diffuse lymphocytes. Serum AcP-L was also increased. Benzene may increase the permeability of the lysosomal membranes. (24 refs)

- 78-4975 Binding of Labeled Benzene Metabolites to Mouse Liver and Bone Marrow.** (Eng) Snyder, R. (Dept. Pharmacology, Thomas Jefferson Univ., Philadelphia, PA, 19107); Lee, E. W.; Kocsis, J. J. *Res Commun Chem Pathol Pharmacol* 29(1): 191-194; 1978.

The binding of benzene metabolites to liver and bone marrow was investigated in mice given twice daily doses of 440 or 880 mg/kg ³H-benzene sc for 10 days, or 880 mg/kg/day sc for 10 days. Irreversible binding of the radioactivity to liver and bone marrow was related to the dose of ³H-benzene administered SC over a 3 day period. Administration of 880 mg/kg twice daily definitely increased binding to liver and possibly to bone marrow over that observed with a single 880-mg/kg dose. Dividing the 880 mg dose into two daily doses of 440 mg/kg did not increase binding in either of these organs. Twice daily doses of 440 mg/kg resulted in increased binding throughout treatment in the liver, but in the bone marrow, the binding increased for days 1 to 6 and decreased between days 6 and 10. It is suggested that the increase in the liver was related to a marked reduction in the urinary metabolites of benzene between days 6 and 10, probably because of liver damage. By day 10, the marrow had lost its characteristic red color, indicating that the functional tissue had probably been replaced by fat. It is suggested that benzene binding to bone marrow is important in leukemogenesis. (5 refs)

- 78-4976 Biological Behavior of N-[4-(5-Nitro-2-furyl)-2-thiazolyl]formamide Induced Carcinoma of the Transitional Epithelium in the Rat.** (Eng) Kiel, H. (Dept. Lab. Animals, Chemie Grunenthal GmbH, Zweifaller Strasse, D-5190 Stolberg, W. Germany); Adolphs, H. D.; Thiele, J.; Steffens, L. *Z Krebsforsch* 91(3): 267-277; 1978.

The incidence and selectivity of tumor induction by N-[4-(5-nitro-2-furyl)-2-thiazolyl]formamide (FANFT) were determined in 45 female Wistar rats fed 0.188% FANFT for 8 mo, for an av weekly dose of 0.195 g/animal. At 12 mo, multiple papillomatous bladder tumors were found in all 30 rats available for assessment, but there were no distant metastases. Histologically, the tumors were papillary transitional cell carcinomas, and all bladder specimens showed infiltrative tumor growth. The tumors were Stages II-IV and Grades 2-3.

In addition, there were considerable dysplastic changes in the urothelium with focal areas of intraepithelial carcinomas. There was no significant statistical relationship between tumor stage and grade or between grade and wt, but a correlation was established between tumor stages and wt ($p < 0.05$), ie, tumor wt increased with enhanced local tumor infiltration. After 12 mo, 36/60 kidneys were normal; 20/60 had dysplasia of the renal pelvis epithelium, in part with papillary folding; 2/60 had transitional cell carcinomas of the renal pelvis; and 4/60 had hydronephrosis. It is concluded that the FANFT-induced bladder tumor of the rat provides a suitable model for studying human bladder tumors. (31 refs)

- 78-4977 The Initiation, Progress, and Diagnosis of Dog Bladder Cancer Induced by 4-Aminobiphenyl.** (Eng) Block, N. L. (Dept. Urology, Univ. Miami Sch. Medicine, PO Box 520875, Miami, FL); Sigel, M. M.; Lynne, A. B.; Ng, A. B.; Grosberg, R. A. *Invest Urol* 16(1): 50-54; 1978.

Beginning at age 4 mo, 24 female beagles were fed 4-aminobiphenyl 5 days/wk for 3.5 yr. Two dogs failed to develop detectable tumors, but 20 dogs developed tumors that progressed to transitional cell carcinomas Grades 2 and 3. Three basic patterns of response to the carcinogen were observed: (1) negative or minimal disease (4 dogs); (2) slow tumor progression with minor fluctuations in development (11 dogs); and (3) rapid tumor progression (9 dogs). The first positive diagnosis of neoplasia was made 15-34 mo after the beginning of treatment. Eleven dogs underwent surgery ranging from tumor sampling to cystectomy plus ureterosigmoidostomy. The tumors closely paralleled human bladder cancer in several characteristics, including histologic types and grades and patterns of metastasis. (16 refs.)

- 78-4978 Metabolism of Bromobiphenyls.** (Eng) Kohli, J. (Guelph Waterloo Centre for Graduate Work in Chemistry, Dept. Chemistry, Univ. Guelph, Guelph, Ontario, Canada); Wyndham, C.; Smylie, M.; Safe, S. *Biochem Pharmacol* 27(8): 1245-1249; 1978.

The metabolism in vitro of 2-bromobiphenyls, 3-bromobiphenyl, and 4-bromobiphenyl was studied in male albino rabbits, and the mutagenicity of 4-bromobiphenyl in *Salmonella typhimurium* TA 1538 was investigated. The bromobiphenyl (PBB) substrates were given ip and the urine and feces collected for 7 days. A series of mono- and dihydroxylated metabolic products were isolated by preparative thin-layer chromatography. The nuclear magnetic resonance spectra of the two major 4-bromobiphenyl metabolites confirmed their structures as 4'-bromo-4-biphenolol and 4'-bromo-3,4-biphenyldiol. The metabolism of 4'-(³H)4-bromobiphenyl gave the corresponding monohydroxy compound that retained 61% of the deuterium; the catechol retained only 19%. The deuterium shift is consistent with metabolism via an arene oxide that rearranges to the phenol and undergoes hydra-

on/dehydrogenation to give the catechol. The retention of 9% of the deuterium in the catechol also indicates that some of the diol is formed via direct hydroxylation, as observed in the metabolism of 4-chlorobiphenyl. Incubation of (³H)-4-bromobiphenyl with rabbit liver microsomes gave two major degradation products, the chromatographic behavior of which was identical to that of the metabolites in vivo. 4-Bromobiphenyl was highly mutagenic to *S. typhimurium* TA 1538 in the presence of a liver microsomal enzyme preparation. The results indicate that metabolic degradation of PBB is an important pathway for the detoxication of these chemicals via urinary excretion. (23 refs.)

78-4979 Hepatocarcinogenicity of Polychlorinated Terphenyl (PCT) in ICR Mice and Its Enhancement by Hexachlorobenzene (HCB). (Eng) Shirai, T. (First Dept. Pathology, Nagoya City Univ. Medical Sch., 1 Kawasumi, Mizuho-cho, Mizuho-ku, Nagoya 467, Japan); Miyata, Y.; Nakanishi, K.; Murasaki, G.; Ito, N. *Cancer Lett* (5): 271-275; 1978.

The carcinogenic effects of hexachlorobenzene (HCB) and polychlorinated terphenyl (PCT) were studied singly and in combination when given po to ICR mice for 24 wk. The mice were fed a basal diet supplemented with 10 or 50 ppm HCB, 50 or 500 ppm PCT, or 250 ppm PCT plus 50 ppm HCB. PCT alone induced liver tumors, nodular hyperplasias, and hepatocellular carcinomas. HCB alone had no effect, but it enhanced the induction of liver tumors by PCT. It is known that PCT accumulates in mouse liver, reaching a concentration of 4-10 times that in the diet, and this may be a reason why the liver is the target organ for PCT carcinogenesis. The tumors induced by PCT were histologically similar to those induced by hepatocarcinogens such as α -benzene hexachloride, polychlorinated biphenyl, and N-2-fluorenylacetylamide. (10 refs.)

78-4980 Distribution and Excretion of Halogenated Xenobiotics (PCB, PBB, and Kepone) (Meeting Abstract). (Eng) Matthews, H. B. (Pharmacokinetics Section, Lab. Pharmacology, NIEHS, Research Triangle Park, NC, 27709); Morales, N. M.; Kato, S.; McKinney, J. D.; Tuey, D. B. *Environ Health Perspect* 20: 233; 1977. (no refs)

78-4981 Reaction Products of the Carcinogen N-Hydroxy-4-acetyl-4'-fluorobiphenyl with DNA in Liver and Kidney of the Rat. (Eng) Kriek, E. (Chemical Carcinogenesis Div., Antoni Van Leeuwenhoek-Huis, Netherlands Cancer Inst., Amsterdam, Netherlands); Hengeveld, G. M. *Chem Biol Interact* 21(2/3): 179-201; 1978.

The in vivo binding of 4-acetyl-4'-fluorobiphenyl (AABP4'F) and 4-amino-4'-fluorobiphenyl (ABP4'F) residues to rat liver and kidney DNA was studied at various

times after inoculation of male albino rats (inbred R-Amsterdam strain) with N-[G-³H]hydroxy-AABP4'F (5 or 25 mg/kg). DNA preparations from both organs were hydrolyzed enzymatically at pH 8-9 with mixtures of DNase, snake venom phosphodiesterase and alkaline phosphatase from *Escherichia coli*. The enzymatic digests were analyzed by Sephadex LH-20 chromatography using synthetic N-([G-³H]deoxyguanosin-8-yl)-ABP4'F as marker. Elution with 30% ethanol gave three major peaks of ³H activity, one that consisted largely of N-(deoxyguanosin-8-yl)-ABP4'F decomposition products, a second that had chromatographic and chemical properties similar to those of 3-(deoxyguanosin-N²-yl)-AAF and was persistent in both liver and kidneys, and a third that cochromatographed with the marker compound. Kinetic studies revealed that the third product was removed rapidly from liver and kidney DNA at equal rates (half time = 2 days). Approx 80% of the total radioactivity bound to DNA consisted of deacetylated material, which was removed at a much slower rate (half time = 10 days) in both organs. The initial rapid removal of all products in the kidney during the first 7 days (half time = 3.3 days) with the 25-mg/kg dose was probably due to toxic effects on the kidneys because this phenomenon was not observed with the 5-mg/kg dose. It was concluded that the overall removal of arylamine adduct in DNA proceeds much more slowly than the removal of N-(guanine-8-yl)arylamide moieties. Comparisons were made between these results and those obtained with acetylaminofluorene. (40 refs.)

78-4982 Non-specific Tritiation of Some Carcinogenic Aromatic Amines. (Eng) Breeman, G. E. (Chemical Carcinogenesis Div., Antoni van Leeuwenhoek-Huis, Netherlands Cancer Inst., Amsterdam, Netherlands); Kaspersen, F. M.; Westra, J. G. *J Labelled Compd Radiopharm* 14(5): 741-750; 1978.

2-Amino-4-fluorene, 4-amino-3-methylbiphenyl, 4-aminobiphenyl, and 4-amino-4'-fluorobiphenyl were tritiated by acid-catalyzed exchange of the corresponding nitro compounds followed by catalytic reduction. The exchange reactions were carried out by heating the nitro compounds in ³H-trifluoroacetic acid and adding a catalytic amount of trifluoromethanesulfonic acid. No loss of tritium could be detected during conversion of the tritiated nitro compounds into the corresponding amines by catalytic hydrogenation. Incorporation into the ortho position was < 4%. During the metabolic activation and binding of ³H-N-acetyl-2-aminofluorene to rat liver DNA in vivo, no tritium exchange occurred. (7 refs.)

78-4983 Effect of α -Benzene Hexachloride on 2-Fluorenylacetylamide Carcinogenesis in Rats. (Eng) Hiasa, Y. (First Dept. Pathology, Nara Medical Univ., 840 Shijo-cho Kashihara, Nara-ken 634, Nara, Japan); Ohshima, M.; Ohmori, T.; Murata, Y. *Gann* 69(3): 423-426; 1978.

The effect of α -benzene hexachloride (BHC) on the induction of hepatomas in male Wistar rats treated simultaneously with 2-fluorenylacetylamide (2-FAA) was studied. Diets containing 0.025% 2-FAA (Group 1), 2-FAA + 0.06% BHC (Group 2), BHC (Group 3), or no additions (Group 4) were fed to the animals for 5 mo, followed by a stock diet for 4 wk. Hepatocellular carcinomas were observed only in 7/8 Group 1 rats; the livers of all these rats contained a few small cysts, and there were multiple hyperplastic nodules and moderate oval cell infiltration. Adenocarcinomas developed in 4/11 Group 2 rats, and the livers of all rats had multiple cysts with islands of parenchymal cells between them. There were a few hyperplastic nodules, and oval cell infiltration was slight. The adenocarcinomas, which were not found in any other groups, were small (max of 2 mm in diameter) and there were no metastases. The adenocarcinomas may originate from the bile ducts. Histopathological findings in Groups 3 and 4 were negative. (13 refs.)

78-4984 Covalent Intercalative Binding to DNA in Relation to the Mutagenicity of Hydrocarbon Epoxides and N-Acetoxy-2-acetylaminofluorene. (Eng) Drinkwater, N. R. (McArdle Lab. Cancer Res., Univ. Wisconsin Medical Center, Madison, WI, 53706); Miller, J. A.; Miller, E. C.; Yang, N. C. *Cancer Res* 38(10): 3247-3255; 1978.

The abilities of N-acetoxy-2-acetylaminofluorene (N-AAF) and a series of hydrocarbon epoxides to cause the unwinding of supercoiled simian virus 40 (SV40) DNA by covalent binding were measured by incubating the DNA with the compound at pH 7.4 and 37 C and then determining the superhelicity of the modified DNA by gel electrophoresis. The concentrations of the electrophiles required to fully unwind the supercoiled DNA ranged from 0.023 to > 3 mM in this ascending order: ($\pm$)-7 β ,8 α -dihydroxy-9 α ,10 α -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene, 1-pyrenyloxirane, 9-methyl-10-anthryloxirane, 6-chrysenyloxirane, 9-anthryloxirane, N-AAF, 9-phenanthryloxirane, 7-benzanthryloxirane, 2-naphthyloxirane, and styrene oxide (phenyloxirane). The frameshift mutagenicities of these compounds toward *Salmonella typhimurium* TA98 ranged from 540 to 0.001 revertants/nanomole, and they were well correlated with their abilities to cause DNA unwinding. The abilities of the compounds to induce missense mutations in *S. typhimurium* TA100 did not correlate as well with their activities in the DNA-unwinding assay. The hydrolysis rates of the compounds under the incubation conditions used for the DNA-unwinding assay were not correlated with their mutagenicities or their activities in causing DNA unwinding. Studies with ^{3} H-labeled N-AAF and ($\pm$)-[7- ^{14} C]7 β ,8 α -dihydroxy-9 α ,10 α -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene indicated that complete unwinding of the supercoiled DNA required binding of about 0.05 hydrocarbon residue/nucleotide. The changes in superhelicity of the SV40 DNA upon treatment with these two electrophiles were confirmed by electron microscopy; no denaturing of the modified DNA

was observed. These data suggest a model of covalent intercalative binding for the interaction of these compounds with DNA. (49 refs.)

78-4985 Microsomal Metabolism of Arylamides by the Rat and Guinea Pig--I. Oxidation of N-3-Fluorenylacetylamide at Carbon-Atom-9--A Major Metabolic Reaction. (Eng) Kaplan, E. (Lab. Cancer Res., Veterans Admin. Hosp., Minneapolis, MN, 55417); Gutmann, H. R.; Emory, T. H. *Biochem Pharmacol* 27(11): 1581-1589; 1978.

The N-hydroxylation of N-3-fluorenylacetylamide (3-FAA) by hepatic microsomes of untreated and 3-methylcholanthrene (3-MC; 20 mg/kg ip)-treated male Duncan-Hartley guinea pigs and male Sprague-Dawley rats was investigated. The N-hydroxylation by microsomes of both species was of the same low order. Both guinea pig and rat hepatic microsomes converted 3-FAA to N-(9-hydroxy)-3-FAA and to N-(9-oxo)-3-FAA. The former was the major product of 3-FAA hydroxylation by hepatic microsomes of both species, and it even exceeded the formation of N-(7-hydroxy)-3-FAA, the principal phenolic metabolite of 3-FAA. The amounts of N-(9-oxo)-3-FAA formed were about one-third those of N-(9-hydroxy)-3-FAA. In contrast to the formation of phenolic metabolites, the hydroxylation of 3-FAA to N-(9-hydroxy)-3-FAA was not increased by pretreatment of guinea pigs or rats with 3-MC. Similarly, pretreatment of rats with phenobarbital (80 mg/kg ip) did not enhance the yield of N-(9-hydroxy)-3-FAA. CO inhibited the formation of N-(9-hydroxy)-3-FAA by 80%. It is concluded that the formation of N-(9-hydroxy)-3-FAA is catalyzed by a microsomal hemoprotein not identical with cytochrome P₄₅₀ or P-450. In contrast, experiments in rats revealed that N-2-fluorenylacetylamide (2-FAA) was a poor substrate for hydroxylation to N-(9-hydroxy)-2-FAA. The susceptibility of 3-FAA to hydroxylation at carbon atom 9 of the fluorene moiety may be rationalized by resonance structures in which carbon atom 9 is positively charged and acts as an electrophilic center. Similar resonance structures cannot be written for 2-FAA. (29 refs.)

78-4986 Antibodies to DNA Modified by the Carcinogen N-Acetoxy-N-2-acetylaminofluorene. (Eng) Leng, M. (Centre de Biophysique Moléculaire, 1A, avenue de la Recherche Scientifique, 45045 Orleans Cedex, France); Sage, E.; Fuchs, R. P.; Daune, M. P. *FEBS Lett* 92(2): 207-210; 1978.

The immunogenicity of native DNA modified by reaction with N-acetoxy-N-2-acetylaminofluorene (AAAF) was investigated. In rabbits, this DNA induced the synthesis of antibodies that specifically recognized AAAF-substituted DNA. A method for purifying the antibodies is described, and association constants for the binding of the antibodies and several ligands are reported. (17 refs.)

78-4987 **Mutagenesis of Chinese Hamster Cells In Vitro by Combination Treatments with Methyl Methanesulfonate and N-Acetoxy-2-acetylaminofluorene.** (Eng) Myhr, B. C. (Somatic Cell Genetics Section, Biology Branch, NCI, Bethesda, MD, 20014); DiPaolo, J. A. *Cancer Res* 38(8): 2539-2543; 1978.

The lethal and mutagenic consequences of exposing Chinese hamster V79 cells to modify toxic doses of methyl methanesulfonate (MMS) followed at different time intervals by a strongly mutagenic dose of N-acetoxy-2-acetylaminofluorene (AcAAF) are described. Treatment with 500 μ M MMS resulted in 5% survival of cloning ability and approx four azaguanine-resistant mutants per 10^5 survivors. Treatment with 7 μ M AcAAF produced 10 times as many mutants, and survival was 7%. However, when V79 cells were treated with μ M AcAAF after MMS (250-1,000 μ M), a lethal synergism occurred that increased the lethality of the MMS pretreatment; the resulting lethality was unaffected by increasing the time interval between treatments from 1 to 48 hr. Lethal synergism was also observed at a constant dose of MMS (500 μ M) and increasing doses of AcAAF (0.5-2.5 μ g/ml). In contrast, no significant change in mutant frequency from that induced by AcAAF alone was found for treatment intervals of 1-63 hr. This result contrasts with a previously observed sixfold enhancement of AcAAF-induced transformation of Syrian hamster embryo cells similarly exposed, with a 48-hr interval between treatments. The difference in the response of these two cell types illustrates the difficulties in attempting to extrapolate the known correlation between individual mutagen and carcinogen treatments to combination treatments when different cell types are used for the two cellular responses. Thus, conclusions about a mutagenic mechanism for in vitro transformation must be derived from mutagenesis and transformation studies of the same cell type. (14 refs)

78-4988 **Culture of Hamster Hepatocytes Capable of High Rates of Carcinogen Metabolism (Meeting Abstract).** (Eng) Pooley, J. A. (Chemical Carcinogenesis Section, Frederick Cancer Res. Center, Frederick, MD, 21501); Raineri, R.; Cavanaugh, D.; Schuman, R.; Pienta, R. J. *In Vitro* 14(4): 336-337; 1978. (no refs)

78-4989 **Interaction of Aliphatic N-Hydroxylamines with Microsomal Cytochrome P450: Nature of the Different Derived Complexes and Inhibitory Effects on Monooxygenases Activities.** (Eng) Mansuy, D. (Laboratoire de Chimie, l'Ecole Normale Supérieure, 24 rue Lhomond, 75231 Paris Cedex 05, France); Rouer, E.; Bacot, C.; Gans, P.; Chottard, J. C.; Leroux, J. P. *Biochem Pharmacol* 27(8): 1229-1237; 1978.

The interactions of aliphatic primary N-hydroxylamines with rat liver microsomal cytochrome P450 was studied under

conditions of oxidative metabolism. The microsomes were obtained from male Sprague-Dawley rats treated with sodium phenobarbital (SPB; 80 mg/kg/day ip for 3 days) or 3-methylcholanthrene (MCA; 20 mg/kg/day ip for 2 days). Primary N-hydroxylamines, RR'R''CNHOH, produced difference spectra of liver microsomes with peaks around 420 nanometers (nm) and troughs around 392 nm, possibly as the result of the hydroxylamines binding to cytochromes P450-Fe(III) through their oxygen atoms. The hydroxylamines interacted with dithionite- or NADPH-anaerobically reduced microsomes, giving peaks around 423 nm. In the presence of NADPH, oxygen, and microsome, all type RR' CHNHOH hydroxylamines formed a 455 nm spectrum corresponding to cytochrome P450-Fe(II)-RR'CHNO complexes. These hydroxylamines strongly inhibited aniline hydroxylase, p-nitroanisole -O- demethylase, and 7-ethoxycoumarin -O- deoxylase activities of SPB-pretreated rat liver microsomes. These inhibitory effects were related to the ability of the hydroxylamines to form 455-nm absorbing complexes whose nitrosoalkane ligands were not displaced by oxygen. The general formation of nitrosoalkane-cytochrome P450 complexes and their inhibitory effects on monooxygenases should be taken into account in pharmacology, especially in the case of drug associations including an amino compound susceptible to metabolic oxidation to a primary or secondary nitrosoderivative. (47 refs)

78-4990 **Xenobiotic Metabolism in Marine Species Exposed to Hydrocarbons (Meeting Abstract).** (Eng) James, M. O. (Marine Pharmacology and Biomedicine Section, Lab. Pharmacology, NIEHS Lab. at C.V. Whitney Marine Lab., St. Augustine, FL, 32084); Bowen, E. R.; Weatherby, R. P.; Bend, J. R. *Environ Health Perspect* 20: 228; 1977. (no refs)

78-4991 **Histofluorescence of 3-Methylcholanthrene Metabolites in the Rat Pancreas.** (Eng) Black, O. (Gastroenterology Res. Labs., Forest Hills Div., Veterans Admin. Hosp., Augusta, GA, 30904). *Gastroenterology* 75(3): 438-444; 1978.

The pancreases of male Long-Evans rats were treated in vivo and in vitro with 3-methylcholanthrene (MC) and examined by histofluorescence techniques to determine the cell of origin of induced pancreatic tumors. For the in vitro studies, frozen sections were mounted on glass slides and dipped for 2 sec into a soln of 2 mg/ml MC dissolved in acetone:ether. The slides were then incubated for 60 min in a soln containing NADH, $MgCl_2$, and KCl. These sections revealed that acinar and ductal epithelium had equal fluorescence, which suggested equal metabolic capability. For the in vivo experiments, 10 mg MC was injected ip and the animals were killed 60 min later. Two additional rats were provided with biliary canulas to divert bile from the pancreatobiliary ducts. Frozen sections were prepared and examined. Fluorescence was con-

centrated in the epithelium and lumen of the ducts. Animals with diversion of bile from the pancreatobiliary ducts had a similar intensity and distribution of fluorescence. These studies indicated that ductal epithelium was exposed to greater concentrations of carcinogen than nonductal epithelium, and they offer a possible explanation as to why ductal epithelium is more likely to undergo neoplastic transformation. (24 refs.)

- 78-4992 **Experimental Production of Brain Tumours in the Albino Mice (Meeting Abstract).** (Eng) Sarasa Bharati, R. (Madras Medical Coll., Madras, India); Arumugam, S. *J Neuropathol Exp Neurol* 37(5): 590; 1978. (no refs)

- 78-4993 **Separation and Identification of 3-Methylcholanthrene-related Compounds by Gas Chromatography-Mass Spectrometry.** (Eng) Takahashi, G. (Dept. Pathology, Chest Disease Res. Inst., Kyoto Univ., 53 Shogoin Kawahara-cho, Sakyo-ku, Kyoto 606, Japan). *Gann* 69(3): 437-439; 1978.

Eleven synthesized derivatives of 3-methylcholanthrene (3-MC) were rapidly separated, identified, and quantified by gas chromatography-mass spectrometry (GS-MS). Hydroxyl derivatives were silylated, and all chemicals were applied to columns of OV-1 and Dexsil-300 at a constant temperature of 260 and 290 C, respectively. A strong relative intensity of the mass spectra of all derivatives was obtained at an ionization voltage of 20 electron volts, and most of the molecular ions gave base peaks. Although separation was incomplete by GC, mass fragmentography plotted against molecular ions satisfactorily separated all the standard derivatives examined. (5 refs)

- 78-4994 **Measurement of H-2 Antigen and Immunogenicity of Methylcholanthrene-induced Murine Sarcomas.** (Eng) Pearson, T. (MRC Lab. Molecular Biology, Hills Road, Cambridge CB2 2QH, England); Sikora, K.; Lennox, E. *Br J Cancer* 37(4): 530-535; 1978.

The possible relationship between amounts of tumor-specific transplantation antigen (TSTA) and H-2 antigen was investigated using tumors induced in B10 or B10.BR mice by sc injection of 0.5 mg 3-methylcholanthrene. The strength of the TSTA was measured by a transplantation assay, the amount of H-2 antigen by absorption of a specific antiserum. Cell suspensions of tumors taken directly from the animal were shown to be contaminated by host cells that made a substantial contribution to the H-2 assay. Since this contamination was lost after several in vitro passages, the amount of H-2b on the tumor cells was assayed after three in vitro passages. A plot of the antigenic ratios vs the relative amount of H-2b

indicated that the tumors varied greatly in both their degree of immunogenicity and their expression of H-2b. For most of the tumors, there was no obvious correlation between the two. However, the two most strongly immunogenic tumors expressed relatively little measurable H-2, and the most weakly immunogenic tumor expressed a fair amount. (13 refs)

- 78-4995 **Metabolic Activation of 3-Methylcholanthrene and Its Metabolites to Products Mutagenic to Bacterial and Mammalian Cells.** (Eng) Wood, A. W. (Dept. Biochemistry and Drug Metabolism, Hoffman-La Roche Inc., Nutley, NJ, 07110); Chang, R. L.; Levin, W.; Thomas, P. E.; Ryan, D.; Stoming, T. A.; Conney, A. H.; Thakker, D. R.; Jerina, D. M. *Cancer Res* 38(10): 3398-3404; 1978.

3-Methylcholanthrene (3-MC) and six of its metabolites were tested for their ability to be metabolized by either hepatic microsomes from Aroclor 1254-pretreated rats or a purified hepatic cytochrome P-450-dependent monooxygenase system to products mutagenic to bacterial and mammalian cells. With *Salmonella typhimurium* strain TA98 as a target for mutagens and microsomes as the source of metabolizing enzyme, 1-hydroxy-3-MC was metabolically activated to a 10-fold greater extent than 3-MC and was the most active compound tested. 1,9,10-Trihydroxy-9,10-dihydro-3-MC (TD-3-MC) and 3MC-2-one, the second and third most active compounds, respectively, were also activated to a greater extent than 3-MC; 2-hydroxy-3-MC was equipotent to the parent hydrocarbon. 3-MC-1-one was less active than 3-MC, and 11,12-dihydroxy-11,12-dihydro-3MC was inactive. With the highly purified and reconstituted monooxygenase system as the enzyme source, TD-3-MC was activated to a 20-fold greater extent than 3-MC and was the most active compound. 3-MC-2-one and 1-hydroxy-3MC were the second and third most active compounds. 3-MC, 3-MC-1-one, and 2-hydroxy-3-MC were poorly activated. When *S. typhimurium* TA100 was used to detect mutations, TD-3-MC was metabolically activated to the greatest extent by both hepatic microsomes and the purified monooxygenase system. The metabolites of this compound were 15 and 100 times more mutagenic than those formed from 3-MC when the microsomes and the purified monooxygenase system, respectively, were the enzyme source. 3-MC-2-one was the second most active substrate with either enzyme source. Although 3-MC-1-one was the third most active substrate in the presence of microsomes, it was the least active with the purified enzyme system. 1-Hydroxy-3-MC, the third most active substrate with the enzyme system, was the least active when activated by microsomes. When hepatic microsomes from Aroclor 1254-pretreated rats were the source of the monooxygenase activity, the compounds had the same relative mutagenic activities in Chinese hamster V79 cells as they did in *S. typhimurium* TA100. Thus, the mutagenic activity of 3-MC metabolites is dependent on both the metabolic activating system and the target cell used to detect mutations. (2 refs.)

78-4996 Identification of the Proximate Form of the Carcinogen 15,16-Dihydro-11-methylcyclopenta[a]phenanthren-17-one (Meeting Abstract). (Eng) Coombs, M. M. (Chemistry Dept., Imperial Cancer Res. Fund, London, England); Bhatt, T. S.; Kissonerghis, A. M. *Br J Cancer* 38(1): 192; 1978. (1 ref)

78-4997 Quantitative Mutation Studies with a Series of Polycyclic Hydrocarbon Derivatives (Meeting Abstract). (Eng) Newbold, R. F. (Chemical Carcinogenesis Div., Pollards Wood Res. Station, Nightingales Lane, Chalfont St. Giles Bucks, England); Amos, J.; Brookes, P. *Br J Cancer* 38(1): 170; 1978. (no refs)

78-4998 Comparative Studies on the Covalent Binding of the Carcinogen Benzo(a)pyrene (BP) to DNA in Various Model Systems (Meeting Abstract). (Eng) Jaggi, W. (Inst. Toxicology, ETH/Univ. Zurich, CH-8603 Schwerzenbach, Switzerland); Lutz, W. K.; Schlatter, C. *Experientia* 34(7): 925; 1978. (no refs)

78-4999 Viral Antigen Induction and Mutability of Different Genetic Loci by Metabolically Activated Carcinogenic Polycyclic Hydrocarbons in Cultured Mamalian Cells. (Eng) Huberman, E. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory: Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1521-1535; 1977.

A possible relationship between the degree of carcinogenicity of a series of polycyclic hydrocarbons and their ability to exhibit gene derepression representing an epigenetic process and/or to induce somatic mutations at different genetic loci was investigated. The induction of polyoma virus capsid antigen (PVCA) (which was detected by immunofluorescence) in an inducible PV-transformed rat embryo cell line was used to test for gene derepression. Somatic mutations were determined in Chinese hamster cells by genetic markers that affect the surface membrane, nucleic acid synthesis, and protein synthesis. The percentage of cells that synthesized PVCA was not significantly increased by treatment with noncarcinogenic hydrocarbons. Dibenz(a,h)anthracene, a weak carcinogen in vitro increased the proportion of cells that synthesized PVCA eightfold. The more potent carcinogens, 3-methylcholanthrene (3-MC), benzo(a)pyrene (BP), and 7,12-dimethylbenz(a)anthracene, increased the proportion of cells producing PVCA by 120-, 150-, and 180-fold, respectively. These results demonstrate a relationship between the carcinogenicity of polycyclic hydrocarbons and their ability to induce PVCA synthesis. There was also a correlation between

mutagenicity and carcinogenicity. The noncarcinogens did not induce mutation, but 3-MC and BP enhanced the mutation frequency 60-280 times for the three genetic markers tested. The major mutagenic metabolite of BP was identified as ($\pm$)-7 α ,8 β -dihydroxy-9 β ,10 β -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene. The results suggest that both somatic mutations and epigenetic processes such as gene derepression may be involved in tumor formation by chemical carcinogens. (41 refs)

78-5000 Morphological Transformation of C3H/10T1/2 Cells Subcultured at Low Cell Densities. (Eng) Haber, D. A. (Genetic Toxicology Group., Dept. Nutrition and Food Science, Massachusetts Inst. Technology, Cambridge, MA, 02139); Thilly, W. G. *Life Sci* 22(18): 1663-1674; 1978.

Morphological transformation of C3H/10T1/2 cells treated with various concentrations of benzo(a)pyrene (BP) was measured following subculture of the cells at low cell densities. Subconfluent cultures exposed to 0.2-40 μ M BP were allowed to grow to confluence, trypsinized, and reseeded at cell densities ranging from 5 to 2,300 surviving cells/cm². These secondary cultures were examined for morphological transformation after 8 to 9 wk incubation. Treated cells grown in virtual isolation (approx 5 surviving cells/cm²) exhibited a transformation frequency of 0.2%-14.5%. At these initial cell densities, transformation frequency was not significantly dependent on BP concentration. However, BP-treated cells reseeded at 11-2,300 surviving cells/cm² showed both density-dependent transformation frequencies and BP concentration dependence of transformation. The subculturing of treated cells did not affect the BP concentration dependence of focus formation. Cell density-dependent suppression of morphological transformation appears to be associated with colony interactions. BP at low concentrations appears to induce transformation potential via a nonmutational mechanism; at the higher concentrations used in this study, BP appears to result in a decline in contact-mediated inhibition phenomena. (16 refs.)

78-5001 Toxic and Transforming Effects of Polycyclic Hydrocarbons in Fetal Hamster Lung-Cell Cultures. I. Benzo(a)pyrene. (Eng) Emura, M. (Abteilung für Experimentelle Pathologie, Medizinische Hochschule Hannover, Karl-Wiechert-Allee 9, 3000 Hannover 61, W. Germany); Richter-Reichhelm, H. B.; Mohr, U. *Cancer Lett* 4(6): 343-348; 1978.

The feasibility of using fetal Syrian golden hamster lung cell cultures of mixed population at early passages for the quantification of the toxic and carcinogenic effects of benzo(a)pyrene (BP) was determined. BP was dissolved in dimethyl sulfoxide (DMSO) and the stock soln diluted to final concentrations of 1.0, 0.75, 0.5, 0.25, 0.1, or 0.05 μ g/ml. The

plating efficiency of the fetal lung cells at two to four passages varied between 1% and 4%, but this value could be held fairly constant at approx 4% by the addition of 0.025-0.2 IU/ml insulin. The colony-forming ability of the cells could be reduced by increasing the insulin concentration at constant BP concentrations. Thus, the toxic effects of BP were enhanced by the presence of insulin. The results of previous studies suggest that insulin may stimulate the cellular intake of BP or activate membrane-bound BP-metabolizing enzymes. Morphologically, only fibroblastic cell colonies were transformed. In additional studies, the insulin concentration was then held at 0.2 IU/ml and the cells were treated with various doses of BP and DMSO. At BP concentrations of 0.05 and 0.25 $\mu\text{g/ml}$, transformation rates were about 1%; however, total colony survival decreased with increasing BP concentrations. BP appeared to be more toxic to endothelial or epitheliallike cells than fibroblastic ones, but DMSO showed the opposite effect. DMSO was more toxic in the presence of 0.2 IU insulin/ml medium than without insulin or at insulin concentrations of 0.025 or 0.1 IU/ml. (9 refs.)

- 78-5002 A Model System for Evaluating the Role of Dietary Fiber in Chemical Carcinogenesis.** (Eng) Clinton, S. K. (Sch. Basic Medical Sciences, Univ. Illinois, Urbana, IL, 61801); Truex, C. R.; Visek, W. J. *Biochem Pharmacol* 27(9): 1393-1396; 1978.

A model system for investigating the role of dietary fiber in intestinal cancer is presented. Female Sprague-Dawley rats were maintained on diets with or without benzo(a)pyrene (BP) and with or without wheat bran. Wheat bran prevented the induction of aryl hydrocarbon hydroxylase by BP in the exposed intestinal epithelium. This model can be used to examine the effects of this and other fibers on other segments of the intestinal tract. (14 refs.)

- 78-5003 Photodynamic Effect of 3,4-Benzopyrene on Thermo-sensitive Mutants of *Escherichia coli* B.** (Eng) Molina, A. M. (Istituto di Microbiologia dell'Università, Via Laterina, 8, 53100 Siena, Italy); Calegari, L.; Calendi, E. *Boll Ist Sieroter Milan* 57(2): 134-139; 1978.

The sensitivity of temperature-sensitive (ts) mutants of *Escherichia coli* B to the lethal photodynamic effect of benzo(a)pyrene (BP) was studied. These mutants were chosen for their inability to form colonies at 40 C. For comparison, two mutants that retained the ability to grow at 40 C were also tested. Cultures of the mutants were incubated with 10^{-5} M BP for 45 min at 30 C and then irradiated, in a glass water-bath at 20 C, with a 500-W quartz lamp 40 cm away. Mutants were removed at various times over a period of 25 min. The ts mutants had a significantly increased sensitivity to the photodynamic effect of BP; this effect was not observed in the two mutants able to grow at 40 C. It appeared unlikely that an alteration of the DNA was responsible for the increased sen-

sitivity. The results indicate that irradiated BP can exert its action on proteins in vivo and that the presence of a protein with an impaired structure may be responsible not only for temperature sensitivity but also for increased susceptibility to the irradiated BP. However, temperature sensitivity and sensitivity to the photodynamic effect were not always correlated. This may be due to the type of amino acid substitution in the altered protein or to the position of the altered protein in the cell. (20 refs.)

- 78-5004 Carcinogenic Activation of Benzo(a)pyrene by Iodine and Ferric Chloride in the Respiratory Tract of Syrian Golden Hamsters.** (Eng) Stenback, F. (Dept. Pathology, Univ. Oulu, Kajaanintie 52 D, SF-Oulu 22, Finland); Rowland, J. *Experientia* 34(8): 1065-1066; 1978.

The effect of iodine (I_2) and ferric chloride (FeCl_3) dusts on benzo(a)pyrene (BP)-induced respiratory tract carcinogenesis was studied in six groups of 48 Syrian golden hamsters. All chemicals were administered by intratracheal instillation weekly for 20 wk. Group 1 received 0.5 mg BP; Group 2 received 0.20 mg I_2 ; Group 3 received BP + I_2 ; Group 4 received 0.16 mg FeCl_3 ; Group 5 received BP + FeCl_3 ; and Group 6 received 0.20 ml saline. Three of seven tumor-bearing animals (TBA) in Group 1 had respiratory tract tumors (RTT): a laryngeal papilloma, a tracheal papilloma, and an accumulation of squamous cells arranged in a glandular fashion and a tracheal papilloma. There were no RTT in the five Group 2 TBA. Seventeen RTT were noted in 19 Group 3 TBA: there were 2 laryngeal papillomas, 19 tracheal papillomas, 2 tracheal squamous cell carcinomas, 6 adenosquamous lesions, 9 peripheral adenomatoid lesions, and 2 each of bronchial and lung papillomas, squamous cell carcinomas, and adenomas. No RTT were noted in two Group 4 TBA. Six RTT were noted in 12 Group 5 TBA: there were 6 tracheal papillomas and 6 peripheral adenomatoid lesions. No RTT tumors were noted in the one Group 6 TBA. The specific chemical binding of hydrocarbons with biological molecules requires that the compounds be activated by cellular hydroxylating enzymes. Several hydroxylases require transitional metals for enzymic activity, and this suggests that metals may be of prime importance in hydroxylation. Covalent binding of BP and similar compounds to nucleic acids is also produced by I_2 . (11 refs.)

- 78-5005 Increase of Urinary Putrescine in 3,4-Benzopyrene Carcinogenesis and Its Inhibition by Putrescine.** (Eng) Fujita, K. (Inst. Comprehensive Medical Science, Fujita-Gakuen Univ. Sch. Medicine, Toyoake, Aichi 470-11, Japan); Nagatsu, T.; Shinpo, K.; Maruta, K.; Takahashi, H.; Sekiya, A. *Cancer Res* 38(10): 3509-3511; 1978.

The effect of putrescine on benzo(a)pyrene (BP)-induced carcinogenesis was determined in female BALB/c mice. A sig-

nificant increase in putrescine was noted in the urine of mice with tumors induced by a single sc injection of 2.52 mg BP in 0.5 ml tricaprilyn soln. However, when 10 mg putrescine was added to the BP soln, the development of tumors was completely inhibited and the increase in urinary putrescine was simultaneously suppressed. Animal wt data for a control group receiving only putrescine indicated that the inhibitory effect of putrescine was not due to its toxicity. (7 refs.)

78-5006 Role of Talc and Benzo(a)pyrene in Respiratory Tumor Formation. An Experimental Study.

(Eng) Stenback, F. (Dept. Pathology, Univ. Oulu, 90220 Oulu 22, Finland); Rowland, J. *Scand J Respir Dis* 59(3): 130-140; 1978.

Morphologic alterations induced by 18 weekly intratracheal instillations of talc (3 mg/0.2 ml saline) or talc + benzo(a)pyrene (BP: 3 mg of each/0.2 ml saline) were studied in Syrian golden hamsters. Only three benign lung lesions were found in 48 hamsters treated with talc alone, and there were no detectable dust deposits, granulomas, or mesothelial proliferations. In contrast, 123 respiratory tumors were found in the 48 hamsters treated with talc + BP, 84 of which were malignant. Tumors were found in the larynx [6 papillomas, 3 squamous cell carcinomas (SCC)], trachea and main bronchi (6 papillomas, 8 SCC, 1 adenocarcinoma, 4 undifferentiated carcinomas), and peripheral bronchi and lungs (27 adenocarcinomas, 19 SCC, 8 undifferentiated carcinomas, 1 hemangioma, 1 adenoma). It appears that, in the hamster, talc has no carcinogenic activity itself but it does influence the activity of BP. (32 refs.)

78-5007 Excisability and Persistence of Benzo(a)pyrene DNA Adducts in Epithelioid Human Lung Cells.

(Eng) Feldman, G. (Dept. Biochemistry and Molecular Biology, Univ. Florida, J. H. Miller Health Center, Gainesville, FL, 32610); Remsen, J.; Shinohara, K.; Cerutti, P. *Nature* 274(5673): 796-798; 1978.

The excisability and persistence of the covalent DNA adducts produced in the epithelioid human alveolar tumor cell A549 by benzo(a)pyrene (BP) were studied by high-pressure liquid chromatography. Chromatograms of total DNA and of micrococcal nuclease-resistant core DNA were made of digests from A549 cells that had been exposed to ^{14}C -thymidine and H-BP (1.3 μM) and incubated for 0 to 72 hr. Peak I contained exclusively (7R) N^2 -(10-[7 β ,8 α ,9 α -trihydroxy-7,8,9,10-tetrahydrobenzo(a)pyrene]yl)deoxyguanosine (BP diol-epoxide I-dG), and peak II contained approx 80% (7R or 7S) N^2 -(10-[7 β ,8 α ,9 β -trihydroxy-7,8,9,10-tetrahydrobenzo(a)pyrene]yl)deoxyguanosine (BP-diol epoxide II-dG). Ratios of $^3\text{H}/^{14}\text{C}$ were used to relate the lesion concentration to the amount of DNA present during BP treatment. They were comparable at 0 hr posttreatment incubation for total DNA (peak I, 0.302; peak II, 0.090) and for nucleosomal core

DNA (0.296; 0.070), and they remained comparable throughout the incubation period, decreasing with time. Excision essentially stopped by 72 hr, leaving a persistent fraction of lesions of 0.55 for BP diol-epoxide I-dG and of 0.40 for BP diol-epoxide II-dG. Thus, A549 cells possessed only a limited capacity to remove covalent DNA-BP adducts from their DNA. The poor repairability by excision of these and related lesions in human lung cells may represent a factor in lung carcinogenesis by polycyclic aromatic hydrocarbons. (13 refs.)

78-5008 Modulation of the In Vivo Binding of the Carcinogen Benzo(a)pyrene (BP) to Rat Liver DNA by Selective Induction of Microsomal or Nuclear Aryl Hydrocarbon Hydroxylase Activity (AHH) (Meeting Abstract).

(Eng) Viviani, A. (Inst. Toxicology, ETH/Univ. Zurich, CH-8603 Schwerzenbach, Switzerland); Lutz, W. K. *Experientia* 34(7): 931; 1978. (no refs)

78-5009 Repair of DNA Damage Induced by Benzo(a)pyrene Diol-Epoxides I and II in Human Alveolar Tumor Cells.

(Eng) Cerutti, P. A. (Dept. Biochemistry and Molecular Biology, J. Hillis Miller Health Center, Univ. Florida, Box J-245, Gainesville, FL, 32610); Sessions, F.; Hariharan, P. V.; Lusby, A. *Cancer Res* 38(7): 2118-2124; 1978.

Repair of DNA damage induced by the proximate benzo(a)pyrene (BP) metabolites, ($\pm$)-7 β ,8 α -dihydroxy-9 α ,10 α -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene (diol-epoxide I) and ($\pm$)-7 β ,8 α -dihydroxy-9 β ,10 β -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene (diol-epoxide 2) was studied in the human alveolar tumor line A549. The fragmentation and subsequent reconstitution of DNA parent strands upon treatment of the A549 cells with the diol-epoxides were followed by an alkaline elution procedure. The effect of this treatment on the lengths of the newly synthesized daughter strands and on the total amounts of DNA synthesized was determined as a function of posttreatment incubation. Parental DNA strands were fragmented within the first 3 hr following treatment with either of the diol-epoxides and then slowly elongated to the length of untreated control samples during 30 hr of incubation. The treatment also increased the fraction of short daughter strands within the first 3 hr of incubation. This fraction returned to control levels after 15 hr of incubation. Both diol-epoxides exerted similar effects on the integrity of parental DNA and on daughter strand synthesis. The total amount of DNA synthesized within a 30-min period decreased to 10%-25% of that synthesized by untreated control cultures during 30 hr of incubation. The results illustrate the repairability of BP-induced DNA damage in human lung cells that are related to the target tissues of polycyclic aromatic hydrocarbon carcinogenesis in humans. (32 refs)

- 78-5010 Binding of Benzo(a)pyrene Metabolites to Cellular DNA in Perfused Rat Lungs.** (Eng) Deckers-Schmelzle, B. (Pharmakologisches Institut, Universität Mainz, Obere Zahlbacher Strasse 67, D-6500 Mainz, W. Germany); Klaus, E.; Kahl, R.; Kahl, G. F. *Naunyn Schmiedeberg Arch Pharmacol* 303(3): 303-307; 1978.

The influence of pretreatment with the monooxygenase inducers β -naphthoflavone (NF) and phenobarbital (PB) on total irreversible binding of metabolically activated ^3H -benzo(a)pyrene (BP) to cellular DNA and the formation of DNA nucleoside-BP adducts after cytochrome P-448 induction was studied in isolated perfused lungs of male Sprague-Dawley rats. Pretreatment with PB (80 mg/kg $\times$ 3, ip) increased binding only slightly, but pretreatment with NF (80 mg/kg ip) increased binding by a factor of 23 [7 picomoles (pmol) of BP equivalents/lung bound to DNA vs 0.3 pmol in controls]. In the lungs of NF-induced rats, 0.45, 2.46, and 7.61 pmol of BP equivalents/mg were bound in the DNA, RNA, and protein fractions, respectively. The corresponding control values were 0.02, 0.12, and 0.50 pmol. The two major nucleoside adducts were partially identified by Sephadex LH-20 chromatography of hydrolyzed DNA of lungs from NF-treated rats. These adducts were ascribed to modification of the DNA with a BP 7,8-dihydrodiol-9,10-epoxide and with secondary metabolites of BP phenols. (31 refs.)

- 78-5011 Ellipticines as Potent Inhibitors of Aryl Hydrocarbon Hydroxylase: Their Binding to Microsomal Cytochromes P450 and Protective Effect Against Benzo(a)pyrene Mutagenicity.** (Eng) Lesca, P. (Laboratoire de Pharmacologie et Toxicologie Fondamentales, 205 route de Narbonne, 21078 Toulouse Cedex, France); Leconte, P.; Paoletti, C.; Mansuy, D. *Biochem Pharmacol* 27(8): 1203-1209; 1978.

The interaction of ellipticine (4,11-dimethyl-6-H-pyrido(4,3b)carbazole) and several of its derivatives with rat liver cytochromes P450 was studied. Wistar rats were inoculated ip with phenobarbital (PB: 80 mg/kg/day for 3 days), benzo(a)pyrene (BP: 20 mg/kg/day for 2 days), ellipticine (20 mg/kg/day for 2 days), or Aroclor 1254 (500 mg/kg, one dose 5 days before sacrifice). All the ellipticine derivatives produced the same type of difference spectrum with a peak of 428 nanometers (nm) using oxidized hepatic microsomes and at 445 nm using dithionite-reduced microsomes (spectral dissociation constants of 10^{-6} and 10^{-5} M, respectively). Ellipticine was a better inhibitor of microsomal BP hydroxylase than 7,8-benzoflavone or metyrapone, regardless of the type of induction. This is attributed to the high affinity of ellipticine for microsomal cytochrome P448. Very low ellipticine concentrations (2.3×10^{-6} M- 7.5×10^{-6} M) inhibited the covalent binding of BP to DNA by 50%. The efficiency of the inhibitor was high whatever the source of the microsomes, the smallest effect being obtained with PB-induced microsomes. Ellipticine and 9-hydroxyellip-

tine inhibited the mutagenicity of BP for *Salmonella typhimurium* TA100 by 60% and 78%, respectively, when BP and ellipticine concentrations were equal. Ellipticine (and, to a lesser extent, 9-hydroxyellipticine) was mutagenic by itself. The hypothesis that ellipticines might inhibit the microsome-dependent mutagenicity of many chemicals is under investigation. (34 refs)

- 78-5012 Glutathione S-Transferase Activities in Rat and Mouse Sperm and Human Semen.** (Eng) Mukhtar, H. (Lab. Pharmacology, Natl. Inst. Environmental Health Sciences, PO Box 12233, Research Triangle Park, NC, 27709); Lee, I. P.; Bend, J. R. *Biochem Biophys Res Commun* 83(3): 1093-1098; 1978.

Glutathione S-transferase activity was found in the sperm of Sprague-Dawley rats and DBA/2J and C57BL/6J mice and in human semen with the use of benzo(a)pyrene 4,5-oxide (4,5-BPO), styrene 7,8-oxide (SO), or 1-chloro-2,4-dinitrobenzene (CDNB) as substrates. Specific rat sperm activities with 4,5-BPO, SO, and CDNB were 0.88, 1.07, and 26.6 nanomoles (nmol)/min/mg protein, respectively. These values represent 1.9%, 0.4%, and 1.4% of the specific hepatic cytosol activities in the rat. The specific Δ^1 -3-ketosteroid isomerase activity in sonicated rat sperm was 2.46 nmol/min/mg protein, representing 4.1% of the hepatic cytosol activity. Mouse sperm activities were similar to those in rat with 4,5-BPO as substrate and slightly higher with CDNB as substrate. Semen from four healthy individuals had significant glutathione S-transferase activity, ranging from 47 to 112 nmol/min/ml ejaculate and from 1.0 to 3.6 nmol/min/ml ejaculate, respectively, with CDNB and 4,5-BPO as substrates. The presence of glutathione S-transferase activity in human semen suggests that it may assist in the detoxification of electrophilic metabolites of environmental chemicals. (23 refs.)

- 78-5013 Epoxide Hydrase Activity in Liver Nuclei: Hydroxylation of Benzo(a)pyrene-4,5-oxide and Styrene Oxide.** (Eng) Bornstein, W. A. (Dept. Cell and Molecular Biology, Medical Coll. Georgia, Augusta, GA); Chuang, H.; Bresnick, E.; Mukhtar, H.; Bend, J. R. *Chem Biol Interact* 21(2/3): 343-346; 1978.

The presence of epoxide hydrase activity in purified nuclei isolated from the livers of male Sprague-Dawley rats was determined by high-pressure liquid chromatography (HPLC) and by a radioactive assay with the use, at least in one case, of two different substrates. For the HPLC assay, the incubation system (total volume, 0.8 ml) consisted of nuclei (up to 10 mg protein in 0.25 M sucrose), Na phosphate buffer [pH 8.0, 60 micromoles (μmol)], and benzo(a)pyrene 4,5-oxide [BP 4,5-oxide, 185 nanomoles (nmol)]. The radioactive assay was conducted in a 0.2-ml incubation mixture of nuclei in HEPES buffer (pH 9.0, 25 μmol), $7\text{-}^3\text{H}$ -styrene oxide (200

ol) or ^3H -BP 4,5-oxide (9 nmol), and water. In the HPLC assay, enzyme activity was directly proportional to nuclear protein (up to 10 mg) and to time (to 20 min). In the radioacetylation assay, linearity was observed for both styrene oxide (up to 3 mg nuclear protein and 10 min of incubation) and BP 4,5-oxide (up to 1 mg protein and 15 min of incubation). With BP 4,5-oxide, the specific activity by either assay was 0.33-0.35 nmol/min/mg protein; with styrene oxide, the specific activity was 0.41 nmol/min/mg. Under identical conditions and with BP 4,5-oxide as substrate, the enzyme activity in liver microsomes was approx 25 times this level. It is suggested that the nucleus becomes not only a prime target for the action of polycyclic hydrocarbons, but that it may also represent the means by which these substances are metabolically activated for mutagenesis, teratogenesis, or carcinogenesis. (6 refs.)

78-5014 Changes in Peroxidation During Benzo(a)pyrene-induced Carcinogenesis. (Rus) Lankin, V. (Inst. Chemical Physics, USSR Academy of Sciences, Moscow, USSR); Gerasimov, A. M.; Poliakov, V. M.; Moroleva, L. A.; Antonenkov, V. D.; Gurevich, S. M. *Biokhimiia* 43(2): 235-239; 1978.

The effect of benzo(a)pyrene (BP) on the intensity of peroxide metabolism in the liver was studied in random-bred albino rats. The animals received a single injection of BP (5 mg, sc) and, 3-3.5 mo later, when they developed tumors (pleomorphic cell sarcoma), they were sacrificed. The amount of total lipids in the liver tissue of the tumor-bearers was 15.2 mg/g, compared with 43.7 mg/g in the liver of control animals. Tumorigenesis resulted in a significant decrease of catalase and urate oxidase activities (2x and 1.5x, respectively), a 29% decrease of glutathione reductase activity, a 142% increase of glutathione peroxidase activity, and a 12% increase of the activity of the NADPH-dependent system of microsomal phospholipids peroxidation. The amount of lipid peroxides in the liver of the tumor-bearers was 23.4 nanomoles/mg, compared with 11.2 nanomoles/mg in controls. (23 refs)

78-5015 Co-Oxygenation of Chemicals (Benzopyrene) During Prostaglandin Biosynthesis (Meeting Abstract). (Eng) Sivarajah, K. (Pharmacokinetics Section, Lab. Pharmacology, NIEHS, Research Triangle Park, NC, 27709); Anderson, M.; Eling, T. *Environ Health Perspect* 20: 233-234; 1977. (no refs)

78-5016 Effect of Dietary Phenobarbital, 3,4-Benzo(a)pyrene and 3-Methylcholanthrene on Hepatic, Intestinal, and Renal Glutathione S-Transferase Activities in the Rat. (Eng) Clifton, G. (Clinical Investigation Center, Alameda Regional Medical Center, Oakland, CA, 94627); Kawanishi, N. *Biochem Pharmacol* 27(8): 1284-1287; 1978.

The effect of dietary phenobarbital (PB), 3,4-benzo(a)pyrene (BP), and 3-methylcholanthrene (MCA) (40 mg/kg/day for 14 days) on hepatic, intestinal, and renal glutathione S-transferase activities was studied in male Sprague-Dawley rats. Significant increases in hepatic transferase activities for the following substrates were observed in the PB-treated animals vs controls: 3,4-dichloronitrobenzene (DCNB; 102%), 1-chloro-2,4-dinitrobenzene (CDNB; 145%), p-nitrobenzyl chloride (NBC; 111%), 1,2-epoxy-3-(p-nitrophenoxy)propane (31%), and ethacrynic acid (EA; 51%). BP increased hepatic activity for all substrates except CDNB; MCA had no effect on activity of the epoxide. In the intestine, PB increased transferase activity of DCNB, CDNB and NBC; BP increased activity for EA and the epoxide; and MCA increased activity for CDNB only. In the kidney, MCA increased transferase activity for DCNB, CDNB, NBC and EA. Dietary PB and BP had no inductive effects in the kidney. The PB induction in the liver and small intestine was significantly greater for activity with DCNB, CDNB, and NBC, as opposed to activity with the epoxide or EA. This difference in degree of induction was not seen with BP or MCA and suggests a greater induction of certain transferases by PB. These results differ from those of a study in which the substances were administered parenterally, which indicates that the administration route plays an important role in glutathione S-transferases induction. (24 refs.)

78-5017 Synthesis of the Non-K-region and K-Region trans-Dihydrodiols of Benzo(e)pyrene. (Eng) Lehr, R. E. (Dept. Chemistry, Univ. Oklahoma, Normal, OK, 73019); Taylor, C. W.; Kumar, S.; Mah, H. D.; Jerina, D. M. *J Org Chem* 43(18): 3462-3466; 1978.

Experimental procedures for the synthesis of trans-9,10-dihydroxy-9,10-dihydrobenzo(e)pyrene and trans-4,5-dihydroxy-4,5-dihydrobenzo(e)pyrene are described. The former was synthesized with an overall yield of 50% to 60%, and the latter was obtained in a quantitative yield. (22 refs.)

78-5018 Metabolism of Benzo[a]pyrene and Benzo[a]pyrene 4,5-Oxide by the Isolated Perfused Rabbit Lung (Meeting Abstract). (Eng) Smith, B. R. (Pharmacology and Biomedicine Section, Lab. Pharmacology, NIEHS, Research Triangle Park, NC, 27709); Bend, J. R. *Environ Health Perspect* 20: 230; 1977. (no refs)

78-5019 Inability of Vitamin A Deficiency to Alter Benzo(a)pyrene Metabolism in Syrian Hamsters. (Eng) Bornstein, W. A. (Dept. Biochemistry, Univ. Vermont Coll. Medicine, Burlington, VT, 05401); Lamden, M. P.; Chuang, A. H.; Gross, R. L.; Newberne, P. M.; Bresnick, E. *Cancer Res* 38(6): 1497-1501; 1978.

The effect of vitamin A deficiency on the *in vitro* metabolism of benzo(a)pyrene (BP) was investigated. Male Syrian golden hamsters were placed either on a control or vitamin A-deficient diet, and when the serum vitamin A levels of the latter were reduced to <10% of the former, the hamsters were decapitated and lung aryl hydrocarbon hydroxylase activity and BP metabolism were measured. Slightly greater aryl hydrocarbon hydroxylase activity was found in hepatic microsomes from vitamin A-deficient animals, but this difference was not significant. The BP metabolic profile of pulmonary microsomes was similar for control and vitamin A-deficient systems, and only a few quantitative differences were noted. The addition of a vitamin A analog, β -retinyl acetate, to the *in vitro* incubations did not substantially affect BP metabolism. (47 refs)

78-5020 Human Pulmonary Alveolar Macrophages Metabolise Benzo(a)pyrene to Proximate and Ultimate Mutagens. (Eng) Harris, C. C. (Human Tissue Studies Section, Experimental Pathology Branch, Carcinogenesis Res. Program, NCI, Bethesda, MD, 20014); Hsu, I. C.; Stoner, G. D.; Trump, B. F.; Selkirk, J. K. *Nature* 272(5654): 633-634; 1978.

The metabolic activation of benzo(a)pyrene (BP) by human pulmonary alveolar macrophages (PAM) into a promutagen, ($\pm$)7,8-dihydroxy-7,8-dihydrobenzo(a)pyrene (7,8-diol; 50% +), and an ultimate mutagen was investigated in a cell-mediated mutagenesis system using Chinese hamster V-79 cells as the indicator cells. The mutation frequency for ouabain resistance was directly dependent on the number of PAM added to the V-79 cells in medium containing either BP or 7,8-diol. Benzo(e)pyrene [B(e)P], the noncarcinogenic analog of BP, was not mutagenic. Cocultivation of V-79 cells with PAM but without BP, B(e)P, or 7,8-diol did not alter the spontaneous mutation frequency. An increase in concentration of either BP or 7,8-diol enhanced mutation frequencies. Compared with BP, 7,8-diol was a more potent promutagen. PAM also metabolized BP to 7,8-diol, which was released by PAM into the culture medium. These data suggest that PAM can metabolize BP to proximate and ultimate mutagens, which are then released into the extracellular space. The intimate contact between PAM and bronchial epithelium during clearance by the mucociliary transport system of particulates containing chemical carcinogens within PAM suggests the possibility that PAM and bronchial epithelium may act in concert to metabolically activate chemical carcinogens during bronchogenic carcinogenesis. (13 refs.)

78-5021 Binding of the Carcinogens Benzo(a)pyrene (BP) and 7,12-Dimethylbenz(a)anthracene (DMBA) to Salmonella DNA as Compared to the Corresponding Mutagenicity (Meeting Abstract). (Eng) Manthey, B. (Inst. Toxicology, ETH/Univ. Zurich, CH-8603 Schwerzenbach, Switzerland); Lutz, W. K.; L'Eplattenier, E.; Schlatter, C.; Wurgler, F. *Experientia* 34(7): 927; 1978. (no refs)

78-5022 Evidence That Benzo(a)anthracene 3,4-Diol-1,2-Epoxy Is an Ultimate Carcinogen on Mouse Skin. (Eng) Levin, W. (Dept. Biochemistry and Drug Metabolism, Hoffmann-LaRoche Inc., Nutley, NJ 07110); Thakker, D. R.; Wood, A. W.; Chang, R. L.; Lel, R. E.; Jerina, D. M.; Conney, A. H. *Cancer Res* 38(6): 1705-1710; 1978.

Benz(a)anthracene (BA) and several benzo ring derivative of BA were examined for their tumor-initiating activity following topical applications of 0.1-2.0 micromoles to the skin of female CD-1 mice, followed 7 days later by twice weekly applications of 12-O-tetradecanoylphorbol-13-acetate for 20 wk. Comparisons of the percentage of mice with papillomas and the number of papillomas per mouse indicated that both diastereomeric 3,4-dihydroxy-1,2-epoxy-1,2,3,4-tetrahydroBA's (DETBA's), in which the epoxide oxygen is either cis (diol-epoxide 1) or trans (diol-epoxide 2) to the benzylic 4-hydroxyl group, were 10 to 40 times more tumorigenic than BA. Diol-epoxide 2 was two- to threefold more tumorigenic than diol-epoxide 1 after 20 wk of promotion. The immediate metabolic precursor of the DETBAs, trans-3,4-dihydroxy-3,4-dihydroBA, had intermediate activity compared to diol-epoxides 1 and 2 and was approx 20 times more tumorigenic than BA. Resolution of the racemic trans-3,4-dihydroxy-3,4-dihydroBA into the (-)-/and (+)-enantiomers and assignment of their absolute stereochemistry revealed that the (-)-(3R,4R) isomer was at least fivefold more active as a tumor initiator than the (+)-(3S,4S) isomer. Testing of 1,2-dihydrobenz(a)anthracene and 3,4-dihydrobenz(a)anthracene for carcinogenicity revealed the latter to be the most potent compound used in this study; the former had activity comparable to that of BA. These findings suggest that a diol-epoxide in which the epoxide on a saturated, angular benzene ring forms part of a 'bay region' of the hydrocarbon is a prime candidate for an ultimate carcinogenic metabolite of the hydrocarbon. For BA, this metabolite is either or both of the diastereomeric DETBA's. (39 refs)

78-5023 Formation of Glucuronic Acid Conjugates of 7,12-Dimethylbenz(a)anthracene Phenols in First- and Second-Passage Syrian Hamster Embryo Cells Treated with 0.1-1.0 nanomole/ml ³H-DMBA. (Eng) Baird, W. M. (Wistar Inst. Anatomy and Biology, 36th St. at Spruce, Philadelphia, PA, 19104); Chemerys, R.; Chern, C. J.; Diamond, L. *Cancer Res* 38(10): 3432-3437; 1978.

The formation of glucuronic acid conjugates of 7,12-dimethylbenz(a)anthracene (DMBA) phenols in first- or second-passage Syrian hamster embryo cells treated with 0.1-1.0 nanomole/ml ³H-DMBA was studied. The cultures metabolized >90% of the DMBA within 48 hr. Samples of the medium were extracted with chloroform, methanol, and water. The chloroform phases contained about one-third of the DMBA metabolites; the major chloroform-extractable metabolite was 8,9-dihydro-8,9-dihydroxy-DMBA. Glucuronidase treatment of the aqueous methanol-soluble

metabolites converted almost 50% of them to chloroform-soluble metabolites, of which > 80% were identified as phenolic derivatives of DMBA. Similar metabolite profiles were obtained by treating the medium with β -glucuronidase after chloroform extraction. Separation of the methyl and hydroxylated derivatives of DMBA from the phenolic derivatives was accomplished by high-pressure liquid chromatography. Small amounts of hydroxymethyl derivatives were detected only in the chloroform-extractable material, but DMBA phenols were the major component of β -glucuronidase-released material. These results indicate the major pathway of DMBA metabolism in hamster erythrocytes is oxidation of the aromatic rings and not oxidation of the methyl groups. (45 refs.)

78-5024 In Vitro Development of Oncogenicity in Cell Lines Established from Tracheal Epithelium Exposed In Vivo to 7,12-Dimethylbenz(a)anthracene.

(Eng) Marchok, A. C. (Biology Div., Oak Ridge Natl. Lab., Box Y, Oak Ridge, TN, 37830); Rhoton, J. C.; Nettey, P. *Cancer Res* 38(7): 2030-2037; 1978.

Transplanted rat tracheas were exposed to 150 or 650 μ g 7,12-dimethylbenz(a)anthracene (DMBA) over a 2-wk period and epithelial cell lines derived from these tracheas were examined periodically in vivo for development of oncogenicity for phenotypic changes associated with neoplastic transformation in vitro. Among three cell lines obtained from tracheas preexposed to 150 μ g DMBA, tumorigenicity was demonstrated only in the late passages of two cell lines maintained to 400 days or more in vitro. With these cells, in vivo latency was 115-255 days. Eight of nine cell lines derived from tracheas preexposed to 640 μ g DMBA became tumorigenic. Several were oncogenic after approx 200 days in vitro, and they yielded palpable tumors within 9-80 days of inoculation. In several instances, inoculation of cells at an early passage into immunosuppressed isogenic recipients resulted in the development of keratinic cysts or tumors that regressed. Inoculation of later passages of the cell lines resulted in the development of invasive tumors. Loss of anchorage dependence of growth was, in most cases, a reliable phenotypic marker for the development of oncogenicity, as was colony-forming efficiency on plastic. All cell lines were epithelial, as indicated by ultrastructural characteristics, the production of keratin in vitro, and/or the induction of squamous cell carcinomas upon inoculation in vivo. The cell lines had stable morphological characteristics distinguishing each cell line from any other by growth pattern, cell shape, and degree of keratinization. The data indicate that the carcinogenesis initiated in epithelial cells in vivo continues in culture. Thus, it should be possible to study in vitro both the evolution of neoplasia and factors affecting tumor progression. (32 refs)

78-5025 Biological Influence of Fat Intake on Mammary Cancer and Mammary Tissue: Experimental

Correlates. (Eng) Nishizuka, Y. (Lab. Experimental Pathology, Aichi Cancer Center Res. Inst., Chikusa-ku, Nagoya 464, Japan). *Prev Med* 7(2): 218-224; 1978.

The effect of dietary fat on 7,12-dimethylbenz(a)anthracene (DMBA)-induced tumors was investigated in Sprague-Dawley rats. Following two iv injections of DMBA at 8 wk of age, the rats were given similar amounts of four different semisynthetic, isocaloric fat diets (8% and 20% coconut oil and corn oil diets) for 8 wk starting at 12 wk of age. At the end of this period, they were placed on the control diet. The induced mammary carcinoma and mammary fat pads were examined at 2- to 3-wk intervals for 13 wk after return to the control diet. Fatty acid compositions in both tissues reflected those of the diets ingested; this finding was more obvious in the high-fat groups. In the cancer tissues, polyunsaturated fatty acids were detected in larger amounts. The uptake of tritiated thymidine appeared to be increased in both tissues of rats fed the high-fat diet. The fatty acid patterns in neoplastic tissue, especially that of lauric acid in the coconut oil group, remained for a longer period of time after return to the control diet. These findings suggest that the promoting influence of a high-fat diet on chemical-induced mammary carcinomas is related to the retention of dietary fats in the adipose tissue and their slower turnover in cancer tissues. The literature data on the association between diet and mammary carcinogenesis are reviewed. (21 refs)

78-5026 Studies on the Mechanism of Neuro-Oncogenesis by Polycyclic Aromatic Hydrocarbons (Meeting Abstract). (Eng) Kleihues, P. (Abteilung Neuropathologie, Pathologisches Institut der Universität Freiburg, 78 Freiburg, W. Germany). *J Neuropathol Exp Neurol* 37(5): 643; 1978. (no refs)

78-5027 Indirect Chromosomal Fluorescence with Antibody to 7,12-Dimethylbenz(a)anthracene. (Eng) Balick, R. (The Ben May Lab. Cancer Research, Univ. Chicago, Chicago, IL, 60637); Pataki, J. *Biochem Biophys Res Commun* 82(1): 81-84; 1978.

An indirect immunofluorescence technique was applied to acid-alcohol fixed metaphase chromosome preparations from rats treated with 7,12-dimethylbenz(a)anthracene in order to study the mechanism of polycyclic aromatic hydrocarbon (PAH) carcinogenesis. Numerous binding modalities were observed reflecting the extent of chromosomal modification. Isolation of chromatin derived from specific chromosomes would further facilitate research concerning the chemical nature of PAH carcinogenesis. (10 refs)

78-5028 Neoplasia Following Cessation of DMBA Application to Hamster Buccal Pouch. (Eng) Eisenberg, E. (Dept. Oral Medicine and Oral Pathology, Har-

vard Sch. Dental Medicine, Boston, MA, 02115). *J Dent Res* 56(11): 1430; 1977.

9,10-Dimethyl-1,2-benzanthracene (DMBA: 0.5% soln) was applied 3x/wk for 8-10 wk to the buccal pouch of Syrian golden hamsters. In 100% of the animals, the dysplasia that was induced resulted in carcinoma, without the continuous application of DMBA after week 10. This progression of dysplasia to frank neoplasia is predictable, and it obviates the need for continuous application of carcinogen. (no refs)

78-5029 Restraint-induced Inhibition of 7,12-Dimethylbenz[a]anthracene-induced Mammary Tumors: Relation to Stages of Tumor Development. (Eng) Newberry, B. H. (Dept. Psychology, Kent State Univ., Kent, OH, 44242). *J Natl Cancer Inst* 61(3): 725-729; 1978.

Experiments were conducted on the use of chronically applied restraint to reduce the number of mammary tumors developing in response to treatment of Sprague-Dawley rats with 7,12-dimethylbenz[a]anthracene (5 mg/g iv at 50, 53, and 56 days of age; total dose, 6 mg). Restraint was applied by inserting animals in hardware cloth tubes that did not allow reversal of the facing direction. The restraint duration was 12 or 14 hr/day in the light portion of the light-dark cycle. The results indicated that the inhibitory effect was largely due to restraint applied after the induction period; preinduction restraint and induction period restraint had no significant effect on tumor development. Therefore, to a considerable extent at least, the restraint effect seems due to an interference with the proliferation of neoplastic cells. After termination of the restraint treatment, the rate at which new tumors appeared first increased and then decreased. Restraint did not affect the proportion of tumors regressing after ovariectomy. (19 refs.)

78-5030 Bacillus of Calmette and Guérin and 7,12-Dimethylbenz(a)anthracene Carcinogenesis in Mice. (Eng) Scelsi, R. (Istituto di Anatomia e Istologia Patologica, Via Forlanini 14-18, 27100 Pavia, Italy); Scelsi, M.; Poggi, P.; Mosca, L. *Gastroenterol Clin Biol* 64(4): 353-360; 1978.

The effects of single and repeated treatments with BCG on 7,12-dimethylbenz(a)anthracene (DMBA)-induced tumorigenesis were studied in Swiss mice. A total of 500 mice were given a single sc injection of DMBA (100 µg) within 16 hr of birth. Of this number, 100 animals received a single sc injection of BCG (1×10^7 cells suspended in 0.2 ml saline) at 20 days and 400 received multiple injections (up to 5) at intervals from 20 to 140 days after birth. All animals were sacrificed at 250 days of age. Among the animals treated with DMBA alone, there were 14 lung adenomas, 11 thymic lymphomas, 3 sc sarcomas, 1 liver hemangioma, 1 mammary adenocarcinoma, and 1 ovarian cystoma. Animals that received two or more doses of BCG had a lower tumor inci-

dence than untreated animals or those given only one dose. Reduction in tumor incidence was correlated with length of BCG treatment. Malignant thymic lymphomas were particularly sensitive to the action of BCG, none developed in animals receiving multiple injection. BCG should be able to alter the action of carcinogens by its effect on the reticuloendothelial system and enzyme levels. (26 refs.)

78-5031 The Preparation of Dihydrodiols from 7-Methylbenz(a)anthracene. (Eng) Tierney, B. (Chemter Beatty Res. Inst., Inst. Cancer Res., Royal Cancer Hospital, Fulham Road, London SW3 6JB, England); Abercrombie, B.; Walsh, C.; Hewer, A.; Grover, P. L.; Sims, P. *Chem Bio Interact* 21(2/3): 289-298; 1978.

7-Methylbenz(a)anthracene (MBA) was oxidized by an ascorbic acid-ferrous sulfate mixture, and 3/5 trans-dihydrodiols (DHD) formed were isolated, purified, and characterized. MBA (1 g) dissolved in 3 liters of acetone was divided among four flasks, each containing ascorbic acid (40 g), EDTA (2.0 g), and ferrous sulfate (5.0 g) dissolved in 750 ml water, and the reaction was allowed to proceed for 18 hr. The three non-K region DHD (trans-1,2-dihydro-1,2-dihydroxy-MBA, trans-3,4-dihydro-3,4-dihydroxy-MBA, and trans-8,9-dihydro-8,9-dihydroxy-MBA) were obtained in acceptable quantities, but only trace amounts of the 5,6- and 10,11-DHD were formed. The non-K region DHD were purified by thin-layer chromatography followed by preparative high-pressure liquid chromatography. The UV spectra of the non-K region DHD were similar to those of the corresponding non-K region DHD derived from benz(a)anthracene, except that, in the former, the peaks were shifted by 20 nanometers to longer wavelengths. The methyl group adjacent to the aromatic system probably caused the shift. The mass spectra of the three DHD confirmed the proposed mol wt and showed a strong parent ion peak at m/e 276. Both the 1,2- and 8,9-DHD had similar nuclear magnetic resonance (NMR) spectra, except that the chemical shift for H(1) of the 1,2-DHD was downfield of values typical of benzylic-type carbons, protons and was indicative of a bay region DHD group. In the 8,9-DHD, the hydroxyl groups preferred to be in a quasi-diaxial conformation, giving rise to a low coupling constant. Additional evidence for the trans-configuration of the hydroxyl groups of 8,9-DHD was provided by the identity of the NMR spectrum of the compound prepared by oxidation to that of the compound prepared by metabolism of MBA in a rat liver homogenate. The NMR spectrum of 3,4-DHD gave a large coupling constant between H(3) and H(4), typical of a trans-DHD in which a quasi-diequatorial conformation is preferred. (26 refs.)

78-5032 Carcinogenic Polycyclic Aromatic Hydrocarbons and Their Metabolites as Possible Emission Components. (Ger) Herlan, A. (Karlsruhe, W. Germany). *Reinhalt Luft* 38(7): 276-278; 1978.

oils, highway dusts, and oil furnace emissions were sampled for carcinogenic aromatic polycyclic hydrocarbons. The samples contained 200-500 aromatic compounds. Coronene, dibenz[a,h]anthracene epoxide, phenol, cresol, pyrene, anthranthol, epoxyanthracene, epoxymethanophenanthrene, epoxyphenanthrene, and benzo[a]pyrene were among the carcinogenic hydrocarbons identified. (no refs.)

5033 Binding of Dibenzo(a,e)fluoranthene, a Carcinogenic, Polycyclic Hydrocarbon Without K-Region, to Nucleic Acids in a Subcellular Microsomal System. (Eng) Perin-Roussel, O. (Laboratoire de Chimie des Substances Naturelles, C.N.R.S., 91190 Gif-sur-Yvette, France); Perin, B.; Zajdela, F.; Jacquignon, P. *Cancer Res* 38(10): 3539-3544; 1978.

Initial conditions were determined for the binding of dibenzo(a,e)fluoranthene (DBF), a carcinogenic polycyclic hydrocarbon without an apparent K-region, to nucleic acids when incubated with hepatic microsomal fractions. Methoxyanthrene-pretreated mouse liver microsomes induced a higher level of binding of DBF to DNA than did similarly treated rat liver microsomes. 7,8-Benzoflavone strongly inhibited the binding of this polycyclic aromatic hydrocarbon to DNA, but cyclohexene oxide and trichloropropene oxide had an enhancing effect when used in the presence of rat liver microsomes. An inhibitory effect was also observed with cyclohexene oxide in mouse liver microsome-enriched medium. Labeled DBF bound twice as much to denatured as to native DNA. Incubation of DBF in the presence of liver microsomes, polyribonucleotides (polyadenylate, polyuridylylate, polyguanylate, and polyinosinate) indicated that binding occurred mainly with guanine. Binding of DBF to DNA of various origins was directly proportional to the amount of guanine-cytosine pairs. Preliminary results indicate that there is a covalent bond between DBF and nucleic acids. (45 refs.)

5034 A Novel and Convenient Synthesis of Dibenzo[a,c]anthracene. (Eng) Harvey, R. G. (Ben May Research Laboratories, Cancer Res., Univ. Chicago, Chicago, IL, 60637); Harvey, C.; Konieczny, M.; Fu, P. P.; Sukumaran, K. B. *J Org Chem* 43(17): 3423-3425; 1978.

A simple method for the synthesis of dibenz(a,c)anthracene (DBA) is presented. The method involves direct reductive cyclization of 2-(9'-phenanthroyl)benzoic acid to DBA with stannous chloride in refluxing acetic acid. There was a 90% yield of DBA. (9 refs.)

5035 Synthesis of Aryloxiranes. (Eng) Yang, N. C. (Dept. Chemistry, Univ. Chicago, Chicago, IL, 60637); Chiang, W.; Leonov, D.; Leonov, E.; Bilyk, I.; Kim, J. *Org Chem* 43(17): 3425-3427; 1978.

Three experimental procedures for the synthesis of 1-pyrenyl-, 9-anthryl-, 1-naphthyl-, 2-naphthyl-, 7-benzanthryl-, 6-chrysenyl-, 10-methyl-9-anthryl-, 9-phenanthryl-, and 9,10-dimethyl-2-anthryloxirane and 9,10-bis(oxiranyl)anthracene are presented. The yields of these compounds ranged from 69% to 94%. The compounds may be used as models for the activated forms of polynuclear hydrocarbons. (14 refs.)

78-5036 Tumor Persistence and Regression in Skin Carcinogenesis. An Experimental Study. (Eng) Stenback, F. (Dept. Pathology, Univ. Oulu, Kajaanintie 52 D, SF 90220 Oulu 22, Finland). *Z Krebsforsch* 91(3): 249-259; 1978.

The incidence of skin tumors and factors in the regression and progression of these tumors were determined in 600 female Swiss mice treated topically with 7,12-dimethylbenzo(a)anthracene (DMBA) and/or 12-O-tetradecanoylphorbol-13-acetate (TPA). The animals were divided into five groups: Group 1 received a single application of 300 µg DMBA; Group 2, DMBA followed by 3.2 µg TPA 2x/wk for 15 wk; Group 3, 30 µg DMBA and TPA; Group 4, 5 µg DMBA 1x/wk for life; Group 5, untreated controls. Groups 1-3 developed a large number of skin tumors, many of which regressed spontaneously. Most of the tumors were papillomas, although keratoacanthomas and squamous cell carcinomas were observed in small numbers. Max tumor incidence and tumor regression in these groups occurred early in the experiment, with the number of developing and regressing tumors following a cyclical pattern. In Group 4 mice, tumors occurred in almost all animals and the incidence of regression was low; in contrast to Groups 1-3, it occurred mainly in tumors lasting > 3 wk. There was a definite correlation between type of treatment and regression: in Group 4, only 1/37 tumors lasting ≤ 3 wk regressed but in Group 3, only 1/16 tumors lasting ≤ 3 wk persisted. Regression was associated to a limited extent with tumor size, tumor duration, and total number of tumors. Histological disorganization, ie, disturbed stratification and polarization, cytological irregularities, nuclear hyperchromatism, and cellular enlargement, was consistent with prolonged tumor duration. The results indicate that neoplastic transformation occurs in multiple stages (hyperplastic lesions, benign regressing tumors, persistent tumors, and frankly malignant tumors), with the incidence and biological behavior of the tumors depending on treatment. (24 refs)

78-5037 The Role of the Induction of Ornithine Decarboxylase in Tumor Promotion. (Eng) Boutwell, R. K. (McArdle Lab. Cancer Res., Univ. Wisconsin Medical Center, Madison, WI, 53706). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor

Laboratory): *Origins of Human Cancer*. Vol. 4 (Book B), 699 pp.; 773-783; 1977.

Evidence supporting the induction of ornithine decarboxylase (OD) as a phenotypic change specific and essential for skin tumor formation is presented. The tumor-promoting activity of phorbol esters has been linked to the induction of OD and S-adenosylmethionine decarboxylase (SAMD). The latter, but not the former, is also induced by nonpromoting irritants such as acetic acid and cantharidin. Promoting activity is achieved only when the carcinogens are applied in excess of certain threshold levels; below these levels, promoting activity is lost and only initiation occurs. Skin tumors contain increased levels of OD but not of SAMD. It is suggested that normal tissue possesses an efficient system for the inactivation of OD and that treatment with a carcinogen results in a loss of this capacity, a loss that becomes permanent after several treatments with promoter levels of the compound. Data suggest that neither cyclic AMP nor cyclic guanosine monophosphate acts as a second messenger in the induction of OD by the tumor promoter 12-O-tetradecanoylphorbol-13-acetate (TPA). Colchicine-sensitive structures may serve in the transmission into the interior of the cell of signals resulting from TPA contact with the cell membrane. Mice pretreated with retinoic acid, an inhibitor of tumor formation, showed only a small increase in OD in response to TPA treatment; the induction of SAMD activity was not altered in these animals. (22 refs.)

- 78-5038 **Metabolism of Tritium-labeled 12-O-Tetradecanoylphorbol-13-acetate by Cells in Culture.** (Eng) O'Brien, T. G. (Wistar Inst. Anatomy and Biology, Philadelphia, PA, 19104); Diamond, L. *Cancer Res* 38(8): 2562-2566; 1978.

The metabolism of tritium-labeled 12-O-tetradecanoylphorbol-13-acetate (^3H -TPA) by Syrian hamster embryo fibroblasts (HEF) and by human fibroblasts (line HC-4, from normal conjunctiva of an infant with retinoblastoma) was studied to determine whether the tumor promoter itself or a metabolite is the biologically effective agent. Within 2-3 days after its addition to growing or confluent cultures of HEF, no unchanged ^3H -TPA remained in the medium, as determined by thin-layer chromatography of the chloroform phase obtained by extraction of the medium with chloroform:methanol:water. In contrast, little or no metabolism of ^3H -TPA occurred in cultures of human fibroblasts. Over 7 days, the radioactivity in the chloroform phase decreased from 99% to only 93% (similar to control flasks containing medium only). The major metabolite formed in the HEF cultures was ^3H -phorbol-13-acetate. With both cell types, 99% of the cell-associated radioactivity was chloroform-extractable, and virtually all was unchanged ^3H -TPA, although a minor peak of radioactivity did migrate with the solvent front of the chromatogram of the chloroform extracts from human cells. It appears that, once metabolism of ^3H -TPA

occurs, most of the metabolites produced are found in the medium. (23 refs)

- 78-5039 **Establishment of Epithelial Cell Lines Following Exposure of Cultured Tracheal Epithelium to 12-O-Tetradecanoyl-phorbol-13-acetate.** (Eng) Steel, V. E. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Marchok, A. C.; Nettesheim, P. *Cancer Res* 38(10): 3563-3565; 1978.

Permanent epithelial cell lines were established from Fischer 344 rat tracheal epithelium following its exposure in vitro to 12-O-tetradecanoyl-phorbol-13-acetate (TPA). Organ cultures were prepared from the tracheas of 12-wk-old rats. Beginning on day 9 of culture, 2 x 3-mm explants were exposed for 1 hr every 6 days to 0, 0.01, or 1.0 μg TPA/ml medium for a total of four or eight exposures. Primary cell cultures derived from 18/20 TPA-exposed explants demonstrated morphologically altered cells after approx 150-180 days. These cells had a markedly higher nuclear to cytoplasmic ratio, and they grew more rapidly across the surface of the dish than did the remainder of the primary or control cells. These cells could be repeatedly subcultured in > 95% of the cases. The exposure mode that gave the greatest number of cell lines per explant was four exposures to 1 μg /ml TPA. Fourteen of the cell lines were inoculated into isogenic animals immunosuppressed by thymectomy and x-radiation; all were nontumorigenic. (19 refs.)

- 78-5040 **The Action of Tumor-promoting Agents in Cell Culture.** (Eng) Weinstein, I. B. (Inst. Cancer Res., Columbia Univ., Coll. Physicians and Surgeons, New York, NY, 10032); Wigler, M.; Pietropaolo, C. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory: Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 751-772; 1977.

The actions of tumor-promoting agents in cell cultures were studied. Extremely low concentrations (10^{-10}M) of 12-O-tetradecanoylphorbol-13-acetate (TPA) and certain other phorbol esters caused a rapid and marked induction of the synthesis of plasminogen activator (PA), a tumor-associated proteolytic enzyme, in human cell and normal chick embryo fibroblast (CEF) cultures. The induction appeared to operate at the level of transcription. Morphologic and biochemical changes mimicking those seen upon transformation with Rous sarcoma virus (RSV) were also seen in TPA-treated CEF, and TPA further potentiated the expression of the transformed phenotype in RSV-infected CEF. However, phorbol ester-treated CEF did not display a complete phenotype of RSV-transformed CEF. Structure-function studies with a variety of phorbol esters and structurally related diterpene plant toxins showed a good correlation between structure

al requirements for PA induction in CEF and tumor promotion in mouse skin. The data suggest the involvement of a common cellular receptor in both processes. (43 refs.)

78-5041 Interactions Between Adenovirus, a Tumor Promoter, and Chemical Carcinogens in Transformation of Rat Embryo Cell Cultures. (Eng) Fisher, P. B. *Inst. Cancer Res., Columbia Univ. Coll. Physicians and Surgeons, New York, NY, 10032*; Weinstein, I. B.; Eisenberg, R.; Ginsberg, H. S. *Proc Natl Acad Sci USA* 75(5): 2311-2314; 1978.

The effect of 12-O-tetradecanoylphorbol-13-acetate (TPA) on the in vitro transformation of cells isolated from Sprague-Dawley rat embryos that had been transformed by a temperature-sensitive mutant of adenovirus type 5 (H5ts125) or treated with polycyclic aromatic hydrocarbons prior to transformation was determined. TPA (0.1 $\mu\text{g}/\text{ml}$) enhanced the transformation of secondary embryo cell cultures by H5ts125 two- to threefold. In addition, transformed foci appeared earlier and were larger in cultures grown in the presence of TPA. When TPA was added as long as 7 days after viral infection, enhancement still resulted. Exposure of the cells to benzo(a)pyrene (0.05-0.5 $\mu\text{g}/\text{ml}$) or 7,12-dimethylbenz(a)anthracene (0.0125-0.1 $\mu\text{g}/\text{ml}$) 18 hr prior to H5ts125 infection also enhanced transformation (2 to 4 times). This enhancement was further augmented two- to threefold when cells were grown in TPA after virus infection. Although TPA did not enhance the cloning efficiency of normal rat embryo cells, it did enhance the cloning efficiency of isolated colonies of adenovirus-transformed cells when these cells were grown alone or cocultured with a 100-fold excess of normal rat embryo cells. Transformation could be enhanced with as little as 30 nanograms/ml of TPA. This action of TPA appeared to be due to its ability to facilitate expression of the transformed state rather than to enhance virus uptake or integration. (29 refs)

78-5042 Stimulation of $^{86}\text{Rb}^+$ and ^{32}P Movements in 3T3 Cells by Prostaglandins and Phorbol Esters. (Eng) Moroney, J. (New York State Dept. Health, Roswell Park Memorial Inst., Buffalo, NY, 14263); Smith, A.; Tomei, L. D.; Wenner, C. E. *J Cell Physiol* 95(3): 287-294; 1978.

The relationship between tetradecanoyl phorbol acetate (TPA)-mediated changes in ion transport and alterations in ion movements induced by structurally similar prostaglandins was investigated with the use of the indicators $^{86}\text{Rb}^+$ as its chloride salt and inorganic ^{32}P (^{32}P) as phosphoric acid. Among the earliest changes induced by TPA was a significant increase in ^{32}P incorporation within 15 min incubation of the post-confluent fibroblasts in 10^{-8} to 10^{-6} M TPA. Similarly, the active phorbol ester homolog 4- β -OH phorbol didecanoate but not the inactive stereoisomeric 4- α -OH phorbol didecanoate stimulated ^{32}P incorporation. TPA also stimu-

lated $^{86}\text{Rb}^+$ influx shortly after administration. Both fluxes were ouabain-sensitive, in accord with the idea that an early effect of TPA is to alter ($\text{Na}^+ + \text{K}^+$)-ATPase activity. Prostaglandin E_1 (10^{-7} to 10^{-6} M) and prostaglandin $\text{F}-2\alpha$ (3×10^{-9} to 10^{-7} M) caused a similar stimulation of $^{86}\text{Rb}^+$ and ^{32}P uptake. The finding that water-soluble prostaglandin $\text{F}-2\alpha$ also exhibited stimulator effects indicates that the hormone-induced responses were not mediated by solvent interactions. The similar responses of phorbol esters and prostaglandin derivatives suggest that phorbol esters and prostaglandin derivatives may act at common membrane sites. Since stimulatory effects were observed at discrete times in the logarithmic phase of growth, activation of membrane receptors may be cell-cycle-dependent. (20 refs)

78-5043 A Tobacco Smoke Inhalation Exposure Device for Rodents. (Eng) Maddox, W. L. (Oak Ridge Gaseous Diffusion Plant, Oak Ridge, TN, 37830); Creasia, D. A.; Dalbey, W. E.; Guerin, M. R.; Stokely, J. R.; Creasia, D. A.; Kendrick, J. *Arch Environ Health* 33(2): 64-71; 1978.

A system that utilizes a piston pump to generate cigarette smoke under standard conditions was used to expose rodents to diluted smoke (via inhalation) for controlled periods of time. The number of puffs and the total particulate matter and nicotine levels per cigarette differed only slightly from predicted values determined with analytical smoking machines. Chamber concentrations of nicotine, CO, and CO_2 were also similar. (8 refs)

78-5044 Effect of Smoking on Excretion of Tryptophan Metabolite in Urine--3-Hydroxy Anthranilic Acid. (Eng) Sulochana, G. (Dept. Biochemistry, Medical Coll., Thanjavur 613 004, India); Arunagiri, R. *Indian J Exp Biol* 16(4): 507-508; 1978.

Twenty smokers smoked five cigarettes within a 45-min period, and changes in their urinary excretion of 3-hydroxyanthranilic acid were determined. The smokers had fasted on the morning of the test and had not smoked for at least 12 hr prior to testing. Compared with pretest values, the excretion of this tryptophan metabolite was significantly increased in all 20 subjects 1 hr after the start of the test. (3 refs.)

78-5045 Pulmonary Carcinogenesis in Rats Given Implants of Cigarette Smoke Condensate in Beeswax Pellets. (Eng) Dagle, G. E. (Biology Dept., Battelle Pacific Northwest Labs., Richland, WA, 99352); McDonald, K. E.; Smith, L. G.; Stevens, D. L. *J Natl Cancer Inst* 61(3): 905-910; 1978.

Lungs of inbred OM/NCR and outbred Sprague-Dawley rats were given implants, through a thoracotomy, of pellets of

cigarette smoke condensate (CSC) suspended in a beeswax-tricaprylin vehicle. The pellets contained 1-67 mg CSC. The pellets slowly released material into the surrounding parenchyma, which resulted in a dose-related increased incidence of lung cancer, predominantly invasive and metastasizing epidermoid carcinoma. There was a 42% incidence of pulmonary carcinoma in the highest dosage group, in which approx 1.65 cm² bronchiolar epithelium was exposed. This dose (67 mg) is approx equal to only three high-tar cigarettes. Pulmonary carcinoma was also produced with 1 mg CSC (1.87% incidence). Squamous metaplasia associated with the implanted site preceded the appearance of the carcinomas, and it increased in severity with the larger pellets that contained more concentrated CSC. There was no difference in pulmonary carcinoma incidence with the use of CSC containing high or low concentrations of nicotine. The potential values of this bioassay system are discussed. (34 refs.)

- 78-5046 Lung-Cancer Risks in the Smoking of Different Tobaccos (Meeting Abstract).** (Eng) Betts, T. E. (Res. Labs., Brompton Hosp., London, England); Elson, L. A. *Br J Cancer* 38(1): 193; 1978. (no refs)

- 78-5047 Effect of Glycerol on Local and Systemic Carcinogenicity of Topically Applied Tobacco Condensate.** (Eng) Wilson, J. (Central Toxicology Lab., Imperial Chemical Industries Ltd., Alderley Park, Macclesfield, Cheshire SK10 4TJ, England); Clapp, M. J.; Conning, D. M. *Br J Cancer* 38(2): 250-257; 1978.

The effect of glycerol on the topical carcinogenicity of tobacco condensate was studied in female Alderley Park albino mice. Doses of condensate used were 189 and 94.5 mg/mouse/wk, each alone or with 17.5% or 35% glycerol, the balance being made up from solvent (acetone/water, 80/20, volume/volume). The test solns were applied in a volume of 0.3 ml 3 days/wk to the shaved back skin of the mice. Age-standardized results show that in high-condensate-dose animals, glycerol reduced the incidence of benign and malignant tumors and that of multiple tumors. The high dose of glycerol reduced the incidence of hyperplasia in animals not developing skin tumors. In effect, glycerol appears to modify the incidence of malignant tumors, benign tumors, hyperplasia, and unaffected skin. This suggests a sequential relationship between epithelial hyperplasia, benign tumors, and malignant tumors of the skin in response to topical carcinogens, with glycerol impeding the conversion of hyperplasia to neoplasia and the transition of benign to malignant tumors and, at high dose, inhibiting skin hyperplasia. (9 refs)

- 78-5048 Mutagen-Carcinogens in Food, with Special Reference to Highly Mutagenic Pyrolytic Pro-**

ducts in Broiled Foods. (Eng) Sugimura, T. (Natl. Cancer Center Res. Inst., Tsukiji 5-1-1, Chuoku, Tokyo, Japan); Nagao, M.; Kawachi, T.; Honda, M.; Matsumoto, H.; Yahagi, T.; Seino, Y.; Sato, S.; Matsukura, N.; Matsushima, T.; Shirai, A.; Sawamura, M. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1561-1577; 1977.

The mutagenic activity (MA) of a large number of chemicals and of common food components was tested by a modified *Salmonella typhimurium*/liver microsome assay. Of 400 chemicals tested, 140 were reported to be carcinogenic, and 121/140 (87%) were mutagenic to *Salmonella*. Smoke condensate from broiled fish was mutagenic to *Salmonella* with or without metabolic activation. Dimethyl sulfoxide (DMSO) extracts of the charred surface of broiled fish and beef steak were mutagenic to *Salmonella* with metabolic activation. Smoke condensate from broiled steak showed very low mutagenicity. When MA was expressed in terms of benzo(a)-pyrene (BP) equivalents, the smoke condensate from broiled fish was calculated to contain 10,000 times the MA of its BP content (9 nanograms BP in smoke condensate from 100 g fish). Smoke produced by heating protein of high L-tryptophan content and L-tryptophan itself possessed strong MA. Smoke from heating carbohydrate and vegetable oil was not mutagenic. The MA of low-, medium, and high-tar cigarettes was assayed and expressed in BP equivalents. The BP equivalents/mg tar were similar, indicating that the equivalents are a direct function of the amount of tar. Of 35 mycotoxins tested, 10 were shown to be mutagenic for the first time. Since these compounds are likely to be food contaminants, their carcinogenicity in animals should be examined fully. (40 refs)

- 78-5049 Ingested Mutagens from Opium and Tobacco Pyrolysis Products and Cancer of the Oesophagus.** (Eng) Hower, T. (Univ. Bristol, Bristol, England); Rose, E.; Ghadirian, P.; Castegnaro, M.; Bartsch, H.; Malaveille, C.; Day, N. *Lancet* 2(8088): 494-496; 1978.

The mutagenicities of crude opium, opium dross (the residue from the opium pipe), shireh (refined opium extract), injonga (the tobacco residue extracted from the pipe stem), and isixaxa (the tobacco residues in the pipe) were determined by plate incorporation assays with *Salmonella typhimurium* strains TA98 and TA100. These substances are commonly sucked or chewed in northeast Iran and the Transkei, South Africa, where the incidence of esophageal cancer is high. The opium dross was mutagenic in both tester strains in the presence of rat liver microsomes, whereas the crude opium and shireh were not. Therefore, it is likely that mutagenic pyrolysis products are formed during opium smoking. Both injonga and isixaxa showed a concentration-dependent increase in mutagenicity in the presence of rat liver microsomes. The data indicate that opium and tobacco pyrolysis products

uld be involved in the etiology of esophageal cancer in the
anskei and northeast Iran. (14 refs.)

5050 **Effects of LSD on Human Chromosomes.** (Eng)
Muneer, R. S. (Dept. Biochemistry and Molecu-
Biology, Univ. Oklahoma Health Sciences Center, Okla-
ma City, OK, 73190). *Mutat Res* 51(3): 403-410; 1978.

Effects of lysergic acid diethylamide-25 (LSD-25) and
D tartrate (1.0 or 10 μ g/ml) on human chromosomes were
died using phytohemagglutinin-stimulated heparinized
od lymphocytes from six normal individuals. Control lym-
ocyte cultures were left untreated or were treated with
omycin C (0.1 μ g/ml). There was no significant difference
chromosome aberrations between control cells and those
posed to either concentration of LSD-25 or LSD tartrate
24 hr or either concentration of LSD-25 for 72 hr. Howev-
the aberration frequency was significantly higher in cells
ated with LSD tartrate for 72 hr compared with controls.
ost of the aberrations represented the results of single break
nts, and complex aberrations such as dicentric and quad-
radials were not seen. Exposure to mitomycin C induced
gnificantly elevated incidence of severe chromosome aber-
ions. LSD-25 at 10 μ g/ml had no significant effect on the
wth rate, viability, mutation, or recombination frequency
Escherichia coli; the growth rate or rate of macromolecular
thesis in mouse L cells; or the growth of human fibro-
s. LSD-25 at 100 μ g/ml had a toxic effect on mouse and
nan cells. The data permit no conclusions concerning pos-
deleterious effects of LSD-25 at reasonable doses on hu-
n somatic cells. (16 refs.)

5051 **Gas-Liquid Chromatographic Determination of
Nitrate and Nitrite in Cheese, Ham, Fish Sau-
e, Cod Roes, and Salmon Roes.** (Eng) Toyoda, M. (Hoen-
a, Higashi-ku, Natl. Inst. Hygienic Sciences, Osaka
nch, Hoenzaka, Higashi-ku, Osaka 540, Japan); Suzuki,
Ito, Y.; Iwaida, M. *J Assoc Off Anal Chem* 61(3): 508-512;
78.

gas-liquid chromatographic method based on extraction
the reaction product of nitrate and 2,4-xyleneol has been
veloped for determining nitrates and nitrites in foods. After
tillation, the nitrite residue is oxidized to nitrate with
assium permanganate and then chromatographed in the
ne manner as nitrate. Recoveries from several foods were
%-100% for nitrate and 80%-100% for nitrite; the limit
ensitivity was 0.1 ppm for both. (8 refs)

5052 **Nitrite and Nitrate Are Formed by Endogenous
Synthesis in the Human Intestine.** (Eng) Tan-
baum, S. R. (Dept. Nutrition and Food Science, Massa-
achusetts Inst. Technology, Cambridge, MA, 02139); Fett, D.;

Young, V. R.; Land, P. D.; Bruce, W. R. *Science* 200(4349):
1487-1489; 1978.

Nitrate balance studies were performed in healthy Caucasian
men to assess the extent of nitrate intake in the general popu-
lation. Six 68- to 72-yr-old men were given a diet of their own
choosing for 1 wk, a protein-free diet for 10 days, and a diet
containing 0.8 g/kg egg protein for 10 days. Large day-to-day
fluctuations in urinary nitrate were noted with both the pro-
tein-free and 0.8-g protein diets; there was no tendency to
equilibrium. For each subject, the excretion of nitrate greatly
exceeded the dietary intake of nitrate. In a second study, eight
25- to 40-yr-old men submitted single fecal samples for anal-
ysis, and six 25- to 30-yr-old men submitted samples of ileos-
tomy effluent. Great variations were found among individu-
als, but all 14 samples contained nitrite. Five of the fecal
samples also contained nitrate. In a series of fecal samples
from two individuals, nitrite was always present and nitrate
was sometimes present. Substances in the diet that react with
nitrite (eg, ascorbic acid) did not appear to have a significant
effect on nitrite concentration at pH > 5.5. It is suggested
that heterotrophic nitrification of ammonia or organic nitro-
gen compounds takes place in the upper, aerobic portion of
the intestine. As material passes through the intestine, the
environment becomes more anaerobic, leading to the ac-
cumulation of nitrate as well as nitrite. Thus, human expo-
sure to nitrite is much greater than previously recognized.
Absorbed nitrite is rapidly destroyed in the blood, but nitrate
is recycled into the saliva, stomach, and bladder; in this way,
intestinal nitrification may contribute to the etiology of hu-
man cancer. (21 refs)

78-5053 **Reversible Interconversion of N-Nitroso(2-me-
thylamino)acetonitrile and 3-Methyl-5-
amino-1,2,3-oxadiazolium Chloride and Related
Reactions.** (Eng) Vohra, S. K. (Dept. Chemistry, Temple
Univ., Philadelphia, PA, 19122); Harrington, G. W.; Swern,
D. *J Org Chem* 43(9): 1671-1673; 1978.

As part of an ongoing study of the anchimeric effects of the
N-nitroso group in conjunction with studies of structure-bio-
logical activity relationships in N-nitrosamines, N-ni-
troso(2-methylamino)acetonitrile (I) and N-nitroso(3-me-
thylamino)propionitrile (II) were reacted with anhydrous
HCl in nonaqueous solvents. Compound I quickly yielded
3-methyl-5-amino-1,2,3-oxadiazolium chloride III, virtually
quantitatively, suggesting anchimeric assistance by the ni-
troso group. Treatment with aqueous base converted III back
to I. Compound II did not yield a cyclic product. (8 refs)

78-5054 **Chromosomal Proteins in Hepatocarcinogene-
sis.** (Eng) Tsanev, R. (Inst. Biochemistry, Bul-
garian Acad. Sciences, Sofia, Bulgaria); Hadjiolov, D. Z
Krebsforsch 91(3): 237-247; 1978.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis was used to study the chromosomal proteins of rat liver during nitrosomorpholine (NM)-induced hepatocarcinogenesis and in the resulting primary hepatomas and metastases. Male Wistar rats were fed NM in the drinking water (10 mg/kg, 6 days/wk, for 5 mo). Experimental and control rats were killed 30, 90 and 220 days after the start of the experiment. Chromatin was isolated from liver nuclei in the absence and presence of phenylmethylsulfonylfluoride (PMSF), a powerful inhibitor of serine proteases. Trabecular- and vesicular-type hepatocellular carcinomas plus two cases of metastases to the omentum lymph nodes were observed. During carcinogenesis, the pattern of nonhistone chromosomal proteins (NHCP) from liver chromatin isolated in the absence of PMSF was altered strikingly. Many high-mol-wt NHCP disappeared. In the presence of PMSF, however, no qualitative alterations could be detected. Thus, the liver homogenates of NM-treated animals exhibited an increased proteolytic activity. Contractile protein and histone fractions were selectively preserved. Only slight quantitative changes were noted in the hepatomas: an increase in the 43,000 (43K)- and 63K-dalton protein bands and a decrease in the fast-moving histone subfraction H1^o. Apparently, primary hepatomas have an overall NHCP pattern similar to that of normal liver. Quantitative and qualitative differences were found in the metastases: a significant decrease in protein bands corresponding to the contractile proteins α - and β -tubulin and actin, a significant increase in the 63K-dalton protein, the appearance of new proteins of approx 60K, 90K, and 120K daltons, and the complete disappearance of H1^o. It is suggested that the hepatoma cells that migrate to form metastases have undergone further malignant progression associated with changes in their chromosomal proteins. (53 refs)

78-5055 The Effect of Estradiol-17-phenylpropionate and Estradiol Benzoate on N-Nitrosomorpholine-induced Liver Carcinogenesis in Ovariectomized Female Rats. (Eng) Taper, H. S. (Dept. General Pathology and Neuropathology, Univ. Louvain, Av. Mounier 52, ANPG 52.60, B-1200 Brussels, Belgium). *Cancer* 42(2): 462-467; 1978.

The effects of synthetic estrogens on N-nitrosomorpholine (NNM)-induced liver carcinogenesis in ovariectomized Wistar R rats were determined. Approx 1 mo after ovariectomy, the animals were divided into four groups. Group 1 received NNM (2.6 mg/100 g, 5x/wk for 3 wk, in aqueous soln, by intragastric intubation). Group 2 received NNM plus a single dose of 645 μ g/100 g Dimenformon Prolongatum (500 μ g estradiol-17-phenylpropionate and 145 μ g estradiolbenzoate/100 g) at the beginning of carcinogen treatment. Group 3 received NNM and, after a 2-wk interval, Dimenformon Prolongatum at 3-wk intervals until the end of the experiment. Group 4 received estrogen treatment only, as in Group 3. Animals were sacrificed approx 300 days after the beginning of NNM treatment. Twenty-nine animals were lost as a result of ovariectomy complications and NNM toxicity and

they were excluded from the calculations. Liver foci, nodules or tumors were noted in 25/31 Group 1, 19/21 Group 2, 29/31 Group 3, and 0/27 Group 4 animals. Hepatocellular tumors with varying degrees of differentiation were noted in 3 Group 1, 2 Group 2, and 9 Group 3 rats. Enlarged lymph nodes were found in all animals in Groups 3 and 4 in which autopsy of the skull was performed; no such enlargements were noted in the other groups. Since no Group 4 animals developed foci, nodules, or hepatocellular carcinomas, it was suggested that the estrogens were acting as tumor promoters and not as initiators of liver carcinogenesis. (24 refs.)

78-5056 Nitrosomorpholine-induced Tumors in Fish (Bul) Khudolei, V. V. (N. N. Petrov Res. Inst. Oncology, Leningrad, USSR); Balanski, R. M. *Onkologiya* 14(4): 168-173; 1977.

The carcinogenic effect of nitrosomorpholine (NM) was studied in guppies (*Lebistes reticulatus*) and *Brachydan rerio*. Twenty specimens of *Br. rerio* were exposed to NM for 28 wk; the NM concentration in the water was gradually increased from 100 to 320 ppm. Thirteen animals were alive at the time the first tumor was detected. Seven tumors were found in four animals and the av latent time was 16 wk. The tumors included 4 hepatocellular carcinomas, 2 intestinal adenocarcinomas, and 1 mesenchymoma of the abdominal cavity. Thirty guppies were exposed to NM for 23 wk; the concentration was increased from 75 to 250 ppm. Nine animals were alive at the time the first tumor was detected. Eight tumors (4 hepatocellular carcinomas, 3 hepatic adenomas, and 1 cystocholangiolar cyst) were found in eight animals. The av latent period was 30 wk. Forty specimens of *Br. rerio* were exposed to 300 ppm NM for 11 wk; 11 were alive at the time the first tumor was detected. Five tumors (2 intestinal adenocarcinomas and 3 hepatocellular carcinomas) were found in four animals. The av latent period was 22 wk. No tumors were found in 90 controls of both species. The findings indicate the potent carcinogenic effect of NM in fish. (15 refs)

78-5057 Detection and Determination of N-Nitrosamine Acids by Thin-Layer Chromatography Using Fluorescamine. (Eng) Young, J. C. (Chemistry and Biology Res. Inst., Agriculture Canada, Ottawa, Ontario K1A 0C6, Canada). *J Chromatogr* 151(2): 215-221; 1978.

A procedure for the detection and determination of N-nitrosoamino acids on activated silica gel thin-layer chromatographic plates is described. Using this technique, N-nitrososarcosine, N-nitrosoproline, and N-nitroso-4-hydroxyproline could be detected as fluorophors at the 2 to 30-nanogram level. An application of this method to the detection of these acid compounds in uncooked bacon is described. (58 refs)

78-5058 Occurrence and Determination of N-Nitroso-3-hydroxypyrrolidine in Cured Meat Products.

(Eng) Janzowski, C. (Inst. Toxicology and Chemotherapy, German Cancer Res. Center, Im Neuenheimer Feld 280, D-6900 Heidelberg, W. Germany); Eisenbrand, G.; Preussmann, R. *J Chromatogr* 150(1): 216-220; 1978.

An analytical method for the isolation and determination of N-nitroso-3-hydroxypyrrolidine, a mutagen in the *Salmonella typhimurium* assay, in foods is presented. The amount of this compound in bacon, boiled ham, and bologna sausage was found to be 1-7 ppb. (8 refs)

78-5059 Crypt Hyperplasia Occuring During Dimethylhydrazine Carcinogenesis in the Jejunal Mucosa of Rats (Meeting Abstract). (Eng) Sunter, J. P. (Dept. Pathology, Univ. Newcastle-upon-Tyne, Newcastle-upon-Tyne, England); Appleton, D. R.; Watson, A. J.; Wright, N. A. *J Anat* 126(2): 437; 1978. (no refs)**78-5060 Glycoprotein Metabolism in Normal Proximal and Distal Rat Colon and Changes Associated with 1,2-Dimethylhydrazine-induced Colonic Neoplasia.**

(Eng) Freeman, H. J. (Gastrointestinal Res. Lab., Veterans Admin. Hosp., 4150 Clement St., San Francisco, CA, 94121); Kim, Y.; Kim, Y. S. *Cancer Res* 38(10): 3385-3390; 1978.

The mucosal sugar content as well as the activities of several glycosyltransferases and some glycosidases were determined in the proximal and distal colon of normal male Wistar rats and in rats undergoing 1,2-dimethylhydrazine (DMH)-induced carcinogenesis. In normal rats, differences in total hexose content as well as the activities of several glycosyltransferases and some glycosidases were noted. This indicated that glycoprotein content and metabolism are different at these two sites. In rats receiving DMH (weekly injections of 25 mg/kg from age 6 to 24 wk, sacrifice at 52 wk), significant alterations in hexose content and two galactosyltransferase activities were observed in uninvolved mucosa following DMH administration. Furthermore, significant reductions in sugar content, three of four glycosyltransferase activities, and two hexosaminidase activities were observed in colonic tumors. These data indicate that significant alterations in glycoproteins occur in association with chemical carcinogenesis in both colonic tumors and uninvolved rat colonic mucosa. Furthermore, these studies emphasize the importance of considering the regional differences in quantity and quality of glycoproteins in normal colonic mucosa when studying pathological states. (37 refs.)

78-5061 Effects of Diet on Colon Carcinogenesis and the Immune System in Rats Treated with 1,2-Dime-

thylhydrazine. (Eng) Bansal, B. R. (Dept. Surgery, Alma Dea Morani Lab. Surgical Immunobiology, Medical Coll. Pennsylvania, Philadelphia, PA, 19129); Rhoads, J. E.; Bansal, S. C. *Cancer Res* 38(10): 3293-3303; 1978.

The effects of different diets on colon carcinogenesis by 1,2-dimethylhydrazine (DMH) and on the immune system were determined in male W-Fu rats. Six different diets were used: standard diet (1), semisynthetic diet (2), high-carbohydrate diet (3), high-fat diet (4), high-carbohydrate diet with lard (5), and high-fat diet with lard (6). In the first experiment, diets 1-4 were initiated upon weaning and DMH (15 mg/kg weekly in two divided doses, sc) was administered for 190 days beginning when the rats were 35 days old. In the second experiment, diets 1, 5, and 6 were initiated upon weaning and DMH was given for 133 days beginning at 45 days of age. The rats were followed by sequential laparotomies until death for the development of gastrointestinal (GI) tumors. Tumors appeared earlier and the total number of GI tumors, particularly those of the colon, was higher in rats fed diets 5 and 6. These GI tumors metastasized more frequently and the survival of the rats, after the appearance of the first GI tumor, was significantly shortened. Diet 3 reduced the number of GI tumors and delayed their appearance. Diet 2 accelerated the appearance of colon tumors without increasing the total number of GI tumors over the life-span of the animals. Serum cholesterol levels evaluated during carcinogenesis suggested a correlation between these levels and the increased frequency of colon tumors. The alterations in serum IgG levels, lymphocyte counts, and surface immunoglobulin-bearing lymphocytes, evaluated at different times during carcinogenesis, suggested a biphasic (M-type) immune response. Serum IgG levels were significantly depressed in rats fed diets 2, 4, 5, and 6 compared with levels in rats fed diet 1. However, the pattern of the immune response was similar in the groups of rats fed different types of diets. (39 refs.)

78-5062 Influence of Surgical Anastomosis on DMH Carcinogenesis in the Rat Bowel (Meeting Abstract). (Eng) Teague, C. A. (Dept. Pathology, Univ. Auckland, Auckland, New Zealand); Gavin, J. B.; Whitehouse, S. *NZ Med J* 86(595): 246; 1977. (no refs)**78-5063 Carcinogenicity of 1,2-Dimethylhydrazine Dihydrochloride (DMH) in BALB/c Mice: Influence of the Route and Dose of Administration (Meeting Abstract).** (Eng) Izumi, K. (Second Dept. Pathology, Sch. Medicine, Tokushima Univ., Tokushima, 770, Japan); Furuya, K.; Otsuka, H. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine

Cancer Society. (Manila, Philippines): 184 pp.; 110-111; 1977. (no refs)

78-5064 Inhibition of DNA Synthesis by 1,2-Dimethylhydrazine and Methylazoxymethanol Acetate in Rabbit Colon Mucosa in Organ Culture. (Eng) Mak, K. M. (Dept. Anatomy, Mount Sinai Sch. Medicine, New York, NY, 10029); Chang, W. W. *J Natl Cancer Inst* 61(3): 799-805; 1978.

Colon mucosa explants from New Zealand White rabbits were cultured and used to study the effect of 1,2-dimethylhydrazine (DMH) and methylazoxymethanol acetate (MAM acetate) on DNA synthesis in the epithelial cells. When maintained in organ culture, the mucosa showed near-normal morphology and a linear incorporation of ³H-thymidine into mucosal DNA up to 36 hr of incubation. Explants from the descending colon had a higher DNA synthetic activity than did other segments of the large bowel. Inhibition of DNA synthesis in the explants by DMH and MAM acetate was dose-dependent. After 18 hr of incubation, MAM acetate inhibited DNA synthesis at concentrations of 50, 100, 150, and 200 µg/ml. Little or no inhibition was observed with the same concentrations of DMH. At a concentration of 200 µg/ml and after only 3 or 6 hr of incubation, however, both carcinogens significantly inhibited DNA synthesis. With a longer incubation time, the inhibitory effect of DMH appeared to be reversible, but DNA synthesis was continuously inhibited by MAM acetate up to 18 hr of incubation. Neither DMH nor MAM acetate altered the uptake of ³H-thymidine during 18 hr of incubation. No morphologic changes were seen in explants treated with 200 µg/ml MAM acetate for 18 hr. Physostigmine sulfate, a cholinesterase inhibitor, had no effect on the MAM acetate-induced inhibition of DNA synthesis. These findings indicate that DMH and MAM acetate have a direct effect on the colon mucosa, and they suggest that colon epithelial cells contain enzymes capable of activating DMH and MAM acetate to their alkylating carcinogens. (33 refs.)

78-5065 Carcinogenic Action of N-Nitroso Compounds. 7. Methyl-, Trideuteromethyl-, Ethyl-, n-Propyl-, n-Butyl-, and Acetoxymethylnitrosamine and Methylbutyroxymethylnitrosamine. (Ger) Wiessler, M. (Institut für Toxikologie und Chemotherapie, Deutschen Krebsforschungszentrum, Im Neuenheimer Feld 280, D-6900 Heidelberg, W. Germany); Habs, M.; Schmahl, D. *Z Krebsforsch* 91(3): 317-322; 1978.

Methyl-, trideuteromethyl-, ethyl-, n-propyl-, n-butyl-, and acetoxymethylnitrosamine and methylbutyroxymethylnitrosamine were examined for carcinogenicity in SD rats. All compounds were carcinogenic, inducing forestomach tumors with approx the same latent period. The total doses necessary

for tumor induction were related to the length of the alkyl chain and, hence, to water solubility. (18 refs.)

78-5066 Synthesis of ¹⁴C-Labelled Nitrosamines. IV. Synthesis of ¹⁴C-Methyl(1-acetoxy)methyl Nitrosamine and 1-¹⁴C-Ethyl(acetoxy)ethyl Nitrosamine. (Eng) Braun, H. (Inst. Toxicology and Chemotherapy, German Cancer Res. Center, Im Neuenheimer Feld 280, 6900 Heidelberg, W. Germany); Wiessler, M. *J Labelled Compd Radiopharm* 14(6): 887-892; 1978.

Experimental procedures for the synthesis of ¹⁴C-methyl(1-acetoxy)ethylnitrosamine and 1-¹⁴C-ethyl(1-acetoxy)ethylnitrosamine are presented. The respective yields of the two compounds were 16.3% and 33.3%. (6 refs.)

78-5067 Possible Neuropharmacological Regulation of Diethylnitrosamine-induced Chemical Carcinogenesis. (Rus) Gurkalo, V. K. (Lab. Chemical Carcinogenesis, N. N. Petrov Res. Inst. Oncology, Leningrad, USSR); Zabezhinskii, M. A. *Farmakol Toksikol* 41(3): 338-342; 1978.

The effect of neurophilic agents on diethylnitrosamine (DENA)-induced carcinogenesis was evaluated in four groups of random-bred rats who received DENA with drinking water (100 mg/liter) plus (Group 1) sc injections of atropine (12 mg/kg, 3x/wk), (Group 2) sc injections of proserin (50 µg/kg 3x/wk), (Group 3) sc injections of an adrenergic alpha blocking agent (a 3-phenylpyrrolidine derivative: 2 mg/kg, 3x/wk), or (Group 4) no further treatment. The rats were sacrificed at 6 mo, and the activity of monoamine oxidase (MAO) in liver mitochondria was assessed. The incidence of hepatic and esophageal tumors was increased significantly in Group 1, being 12/15 and 7/15, respectively compared with 7/15 and 3/15 Group 4. The adrenergic alpha blocking agent did not affect the incidence of hepatic and esophageal tumors (6/15 and 2/15), but proserin decreased both the incidence (2/15 and 1/15) and multiplicity of tumor nodes. All three agents significantly increased MAO activity (from 2.0 IU in Group 4 to 3.20 in Group 1, 3.4 in Group 2, and 2.65 in Group 3). (13 refs)

78-5068 Ultrastructure of N-Diethylnitrosamine Induced Tumours in the Nasal Olfactory Region of the Syrian Golden Hamster. (Eng) Reznik-Schuller, H. (Frederick Cancer Res. Center, P.O. Box B, Frederick, MD 21501). *J Pathol* 124(3): 161-164; 1978.

Tumors of the nasal olfactory region induced in five Syrian golden hamsters after sc treatment with N-diethylnitrosamine (DEN: 17.8 mg/kg 1x/wk for 20 wk) were examined electron microscopically. The neoplasms consisted of two cell

types, one of which contained membrane-bound, dense-cored, neurosecretory-type granules. These dense-cored granules were frequently found closely associated with the Golgi apparatus, and a few were identified in cells undergoing mitosis. The other cell type, characterized by its content of abundant ribosomes and polyribosomes, occasionally also exhibited mitosis. The findings demonstrate that the DEN-induced neoplasms originated from the poorly differentiated precursors of olfactory sensory cells (neuroblasts) and from precursors of Sertoli cells (basal cells). (10 refs)

78-5069 Nucleic Acids from Subcellular Fractions of N-Nitrosodiethylamine-induced Hepatoma in Mice. II. Changes in Gene Expression During Tumor Progression. (Eng) Shaposhnikov, J. D. (Petrov Res. Inst. Oncology, USSR Ministry Public Health, Leningrad 188646, USSR); Ratovitski, E. A. *Neoplasma* 25(2): 145-152; 1978.

The hybridization properties of in vivo-labeled nuclear and mitochondrial RNA's of an N-nitrosodiethylamine-induced transplantable hepatoma were investigated in C3HA mice at early (5 and 6) and late (60) passages. During tumor progression, the repression of the nuclear genome noted at early stages was replaced by derepression at later stages. As a result, the percentage of nuclear RNA that hybridized with nuclear DNA increased, and new classes of hybridizable sequences appeared. These results are related only to the repetitive nucleotide sequences. The hybridizability of mitochondrial RNA with nuclear DNA was not changed. (26 refs)

78-5070 The Determination of N-Nitrosodiethanolamine in Cutting Fluids. (Eng) Williams, D. T. (Environmental Health Directorate, Health Protection Branch, Tunney's Pasture, Ottawa, Ontario K1A 0L2, Canada); Benoit, F.; Muzika, K. *Bull Environ Contam Toxicol* 20(2): 206-211; 1978.

Twenty-four samples of cutting fluid concentrates were analyzed by thin layer chromatography for the presence of N-nitrosodiethanolamine (NDETA). Nine of these samples contained nitrite at levels up to 9.8%; NDETA (0.4-4.2 mg/g) was only found in the nitrite-containing samples. (8 refs.)

78-5071 DNA Phosphotriesters as Indicators of Cumulative Carcinogen-induced Damage. (Eng) Shooter, K. V. (Div. Carcinogenesis, Inst. Cancer Res., Pollards Wood Res. Station, Nightingales Lane, Chalfont St., Giles, Bucks, England). *Nature* 274(5671): 612-614; 1978.

Female Chester Beatty hooded rats were fed dimethylnitrosamine (DMN) or diethylnitrosamine (DEN) in the drinking

water (0.1-5 ppm, mean rate of consumption, 2-75 µg/rat/day) continuously for 6-54 wk or they were inoculated ip with single doses of 2-20 mg/kg DMN, and the distribution of phosphotriesters (PTE) in the DNA of lung, liver, and kidney was compared. The concentration of PTE was determined by the differential alkaline hydrolysis method. In the long-term feeding experiment, all doses of DMN induced significant amounts of methyl PTE in the liver DNA. At the highest dose, significant and increasing amounts of PTE were observed in the lung DNA; however, at lower doses, the data on lung DNA were inconclusive. In kidney DNA, significant positive results were observed in only 3/18 experiments, which agrees with observations that long-term feeding of this compound does not tend to induce kidney tumors. After continuous ingestion of DEN, significant concentrations of ethyl PTE were found in the DNA of liver, kidney, and lung, even at low dose rates. After 1 yr, significant amounts of specific lesions could be detected in tissue DNA, even in rats receiving only 2 µg/day of either carcinogen. With a single dose of DMN, most of the methyl PTE were detected in the liver, but an appreciable fraction of DMN was metabolized in the kidney and lung. Since this assay for PTE involves simply the determination of the number of chain breaks produced in DNA by alkaline hydrolysis, it should have a fairly general applicability, the nature of the chemical moiety reacting with the phosphate group being, in general, not important. (13 refs.)

78-5072 Hepatic RNA Resistance to Enzymatic Degradation During Hepatocarcinogenesis in Rats. (Rus) Voronova, L. A. (N. N. Petrov Res. Inst. Oncology, USSR Ministry Health, Leningrad, USSR); Ivanov, S. D. *Vopr Onkol* 24(5): 30-35; 1978.

The resistance of hepatic RNA to degradation by serum RNase was investigated during hepatocarcinogenesis induced in rats by a 6-mo feeding of diethylnitrosamine (DENA). Resistance to RNase degradation was used as a criterion of changes in the spatial configuration of the polynucleotides. In the precancerous state and within the pH range studied (5.1, 7.6, and 8.5), the resistance of total, cytoplasmic, ribosomal, and transfer RNA to enzymatic degradation increased considerably and then fell. With tumor development, RNA resistance to enzymatic cleavage increased again and, in most cases, exceeded control levels. A comparison of these findings with those obtained after treatment of rats with CCl₄ indicated that the carcinogenic effect of DENA on rat liver is accompanied by changes in the spatial configuration of cytoplasmic RNA molecules. (16 refs)

78-5073 Studies on the Gastrointestinal Absorption of N-Nitrosamines: Effect of Dietary Constituents. (Eng) Agrelo, C. (British Industrial Biological Res. As-

soc., Woodmansterne Road, Carshalton Beeches, Surrey SM4 4DS, England); Lake, B. G.; Phillips, J. C.; Gangolli, S. D.; Longland, R. C. *Toxicology* 10(2): 159-167; 1978.

The effects of fat, protein, and carbohydrate on the rates of disappearance of ^{14}C -labeled N-nitrosodimethylamine (DMN) and other nitrosamines from the small intestine of male Sprague-Dawley rats were investigated. Following overnight starvation, the rats were given, by po intubation, ^{14}C -DMN (5 mg/kg) dissolved in isotonic saline soln, corn oil, or casein soln (5% wt/volume). Intestinal absorption studies showed that the presence of fat markedly reduced the rate of disappearance of DMN at concentrations of 4-200 $\mu\text{g}/\text{ml}$. Protein and carbohydrate had little effect on absorption rate. Investigations of the absorption of N-nitrosodiethylamine, N-nitropyrrolidine, and N-nitrosopiperidine from isotonic saline and lipid emulsion showed that lipids also retarded the disappearance of these compounds from the small intestine. In vitro absorption studies, using everted intestinal sac preparations, showed that the small intestinal tract of the rat can degrade DMN. Factors that prolong the retention time of nitrosamines in the gut should facilitate the degradation of these compounds by the metabolic activity of intestinal cells and, possibly, by gut flora and thereby reduce the effective concentration of nitrosamine reaching target organs of the body. (19 refs.)

78-5074 Experimentally Induced Hemangioendotheliomas in the Liver of Mastomys Using Dimethylnitrosamine (Meeting Abstract). (Ger) Wayss, K. (Heidelberg, W. Germany); Bannasch, P.; Mattern, J.; Volm, M. *Zentralbl Allg Pathol* 122(3): 289; 1978. (no refs)

78-5075 Stimulation of Microsomal Dimethylnitrosamine-N-demethylase by Pretreatment of Mice with Acetone. (Eng) Sipes, I. G. (Dept. Anesthesiology, Univ. Arizona, Tucson, AZ, 85724); Slocumb, M. L.; Holtzman, G. *Chem Biol Interact* 21(2/3): 155-166; 1978.

Male C57BL/6J mice were pretreated with acetone (Ac, 15-60 millimoles/kg in normal saline, ip) or Aroclor 1254 (Ar, 500 mg/kg in sesame oil, ip); and the relationship between in vivo microsomal enzyme modifiers and in vitro dimethylnitrosamine (DMN) metabolism was investigated. Ac and Ar are known to influence DMN-N-demethylase activity. Microsomes were isolated from the livers of mice sacrificed 2-48 hr after Ac injection, 4 days after Ar injection. They were then incubated with 0.1-200 mM DMN. Ac enhanced the in vitro microsomal activity of DMN-N-demethylase, as measured by formaldehyde production from DMN. The enhancement was dose- and time-dependent, with max activity occurring 16 hr after Ac injection. There was a concomitant increase in the covalent binding of ^{14}C -DMN to RNA, protein, and DNA. Four distinct K_m values that were dependent on DMN concentration were noted for the

N-demethylase in microsomes from control mice. Only one K_m value was observed for the N-demethylase in microsomes from Ac-treated animals, but it was significantly lower (0.09 mM) than the lowest control value (0.2 mM). Compared with control microsomes, Ac significantly increased DMN-N-demethylation at 1 and 10 mM DMN, but not at 100 mM. DMN-N-demethylase activity in microsomes from Ar-treated mice was significantly decreased at 1 mM DMN and significantly enhanced at 100 mM. It is suggested that there may be multiple forms of DMN-N-demethylase that are dependent on DMN concentration and that respond differently to modifiers of the microsomal drug-metabolizing enzymes. (31 refs.)

78-5076 Nitrosamine Formation in Bladder Infections and Its Role in the Etiology of Bladder Cancer. (Eng) Radomski, J. L. (Dept. Pharmacology, Univ. Miami Sch. Medicine, Miami, FL); Greenwald, D.; Hearn, W. L.; Block, N. L.; Woods, F. M. *J Urol* 120(1): 48-50; 1978.

To evaluate the role of nitrosamine formation in the etiology of bladder cancer, dimethylnitrosamine (DMN) formation was determined in urine specimens incubated in vitro with sodium nitrite and aminopyrine or tetracycline and in urine specimens from untreated and tetracycline-treated patients with urinary tract infections. DMN was formed during the incubation of aminopyrine and, to a lesser extent, tetracycline in urine containing sodium nitrite. Small but significant amounts of DMN were also formed in control urine specimens. DMN was formed during incubations of nitrite-containing urine with diphenhydramine, but not with dimethylaminoethanol, sulfisoxazole, or chloroquine. DMN was found in the urine of 3/10 patients with urinary tract infections before therapy, the concentration being correlated with the presence of nitrite in the urine. The highest concentrations of nitrite and DMN were produced by *Proteus mirabilis*. Therapy with tetracycline rapidly decreased the concentrations of DMN, thus exonerating the drug from any involvement in the induction of bladder cancer by this mechanism. (11 refs.)

78-5077 Kinetic Studies on the Formation of Nitrosamines. I. Formation of Dimethylnitrosamine in Aqueous Solution of Perchloric Acid. (Eng) Cachaza, J. M. (Departamento de Quimica Fisica, Instituto de Investigaciones Quimicas, C.S.I.C., Universidad de Santiago, E-Santiago de Compostela, Spain); Casado, J.; Castro, A.; Lopez Quintela, M. A. *Z Krebsforsch* 91(3): 279-290; 1978.

The kinetics of nitrosation of dimethylamine (DMA) in aqueous perchloric acid soln were determined by differential spectrophotometry. Absorbance was measured at 323 nanometers, and the reaction was performed at a temperature of 310 K. The rate of the reaction was independent of amine concen-

ration over the range tested (0.25-2.0 M). The initial rate of nitrosation was proportional to the square of the initial stoichiometric concentration of nitrite ($3-7 \times 10^2$ M) in the overall range of pH's studied (1.5-4.0). The degree of nitrosation by NO_2H_2^+ (nitrosacidium ion) had practically no influence on the reaction rate. pH values were affected by the reaction media (ionic strength, neutral salts), and it was difficult to determine their effect on the reaction rate. This rate was affected, however, by sodium perchlorate concentration. Nitrosation was inhibited with increasing ionic strength (0.5-2.5 M) as a result of a decrease in the concentration of NO_2H due to a change in pK-nitrite. The rate constant of the reaction was $4.8 \pm 0.5 \cdot 10^{-3} \text{ M}^{-2} \text{ s}^{-1}$ (extrapolated to 298 K for comparison with previous rates obtained at that temperature). Nitrogen trioxide was the only nitrosating carrier that was effective under the simplified conditions used. (30 refs)

78-5078 Hormonal Regulation of Carcinogen Metabolism in Mouse Kidney. (Eng) Bakshi, K. (Dept. Genetics, Litton Bionetics, Inc., Kensington, MD, 20795); Brusick, D.; Bullock, L.; Bardin, C. W. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 199 pp.; 683-695; 1977.

The effects of sex steroid hormones on the metabolism of dimethylnitrosamine (DMN) and N-2-fluorenylacetylacetamide (AAF) by the mouse kidney were studied. Kidney microsomes from androgen-insensitive male (*tfm/y*) mice activated DMN to the same small extent as those from normal males and much less than those from normal females. In contrast, liver microsomes from all animals markedly activated DMN. Castration reduced kidney microsomal metabolism in males. Testosterone activated DMN in normal females and castrated males and enhanced carcinogen metabolism in normal males. Microsomes from untreated and testosterone-treated *tfm/y* mice produced no response; they had little ability to form mutagenic metabolites. Thus, androgens regulate DMN metabolism by modifying renal enzyme activity. The action of medroxyprogesterone acetate (MPA) on this microsomal enzyme system was also mediated via the androgen receptor. Kidney microsomes from normal female and *tfm/y* mice did not significantly activate AAF to a mutagen, whereas those from normal males were quite active. MPA increased the ability of kidney microsomes from female, but not *tfm/y*, mice to activate AAF. The data suggest that the sex-related differences in AAF-induced bladder tumors in mice are correlated with the differential metabolic activating capacities of male vs female kidneys. (27 refs.)

78-5079 Diisopropanolnitrosamine-Induced Thyroid Tumors in Sprague-Dawley Rats (Meeting Abstract). (Ger) Reznik, G. (Hannover, W. Germany). *Zentralbl Allg Pathol* 122(1/2): 146; 1978. (no refs)

78-5080 Synthesis of Deuterium-labeled Nitrosomethylethylamines. (Eng) Mangino, M. M. (Chemical Carcinogenesis Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Lijinsky, W.; Singer, G. M. *J Labelled Compd Radiopharm* 14(5): 697-703; 1978.

Ethyl isocyanate, N-methylacetamide, N-methylacetamide- d_3 , and N-methylacetamide- d_6 were reduced with lithium aluminum deuteride and then nitrosated with sodium nitrite to yield, respectively, nitrosomethyl(- d_3)ethylamine, nitrosomethylethyl(α - d_3)amine, nitrosomethylethyl(- d_3)amine, and nitrosomethyl(- d_3)ethyl(d_3)amine. All products were characterized by nuclear magnetic resonance spectrometry, UV spectrophotometry, and mass spectrometry. The mass spectra of nitrosomethylethylamine and the four deuterium labeled analogs showed patterns that confirmed previous conclusions about the fragmentation of the undeuterated compound. The chemical and isotopic purity of the products was deemed satisfactory for the biological testing of their relative carcinogenicity. (6 refs.)

78-5081 Selective Induction of Pancreatic Ductular Tumors by Single Doses of N-Nitrosobis(2-oxopropyl)amine in Syrian Golden Hamsters. (Eng) Pour, P. M. (Eppley Inst. Res. Cancer, Univ. Nebraska Medical Center, 42nd and Dewey Ave., Omaha, NB, 68105); Salmasi, S. Z.; Runge, R. G. *Cancer Lett* 4(6): 317-323; 1978.

The effect of single injections of various N-nitrosobis(2-oxopropyl)amine (BOP) doses on the incidence of pancreatic neoplasms in Syrian golden hamsters was determined. Male and female animals received an sc injection of either 40 (Group 1), 20 (Group 2), 10 (Group 3), 5 (Group 4), or 2.5 mg/kg (Group 5); the animals were then observed for life and sacrificed when moribund. Pancreatic tumor incidence was dose-dependent, with Group 1 animals having the highest number of benign and malignant tumors and Group 5 animals having only a few benign neoplasms. Most of the tumors developed in the splenic lobe of the pancreas, which indicates that the carcinogen or its metabolites reach the pancreas systemically. Furthermore, most tumors were found along the main pancreatic ducts, suggesting that the neoplasms arise primarily from islet cell precursors, which are derivatives of ductular cells and germinative cells of the pancreas. Almost all tumors had ductular patterns, and they appeared to rise primarily from the intra- and periinsular ductules. A few tumors were also noted in the lungs, liver, gallbladder, and kidneys of some pancreatic tumor-bearing animals; animals without pancreatic tumors had no other tumors that could be related to treatment. The finding that pancreatic tumors were induced with 2.5 mg/kg BOP (2.5% of LD_{50}) demonstrates the high carcinogenic potency of this compound in the hamster pancreas. (15 refs.)

78-5082 Synthesis of ^{14}C -Labelled Compounds. III. Synthesis of N-Nitroso-O,N- ^{14}C -dimethylhydrox-

ylamine. (Eng) Arjungi, K. N. (Cancer Res. Inst., Tata Memorial Center, Parel, Bombay 400026, India); Kruger, F. W.; Wiessler, M. *J Labelled Compd Radiopharm* 14(6): 913-918; 1978.

An experimental procedure for the synthesis of N-nitroso-O,N-¹⁴C-dimethylhydroxylamine from ¹⁴C-labeled dimethyl sulfate and N-hydroxyurethane is reported. The procedure resulted in a 49% yield of the nitrosamine. (9 refs.)

78-5083 Distribution of 7-Methylguanine in Liver Chromatin DNA from Rats Treated with N,N-Di[¹⁴C]Methylnitrosamine (Meeting Abstract). (Eng) Galbraith, A. I. (Paterson Labs., Christie Hosp., Manchester, England); Itzhaki, R. F. *Br J Cancer* 38(1): 170-171; 1978. (1 ref)

78-5084 Initiation of Chemical Carcinogenesis Requires Cell Proliferation. (Eng) Cayama, E. (Dept. Pathology, Univ. Toronto, Medical Sciences Building, Toronto, Ontario M5S 1A8, Canada); Tsuda, H.; Sarma, D. S.; Farber, E. *Nature* 275(5675): 60-62; 1978.

A quantitative assay for the initiation of liver cancer was used to investigate whether cell replication exerts its first effect on initiation or on some later step in the carcinogenic process. The assay is based on the hypothesis that an early cell is resistant to the inhibitory effect of carcinogens on cell proliferation. It consists of measuring the number of foci of carcinogen-altered hepatocytes that are capable of responding to a stimulus for cell proliferation in the presence of a strong inhibitor of normal cell proliferation. Adult male Fischer 344 rats received a single ip dose of 60 mg/kg N-methyl-N-nitrosourea (MNU) followed 3 hr later by 67% partial hepatectomy (PH). Two weeks later, the rats received a diet containing 0.02% 2-acetylaminofluorene (2-AAF) and they were force-fed the mitogen α -hexachlorocyclohexane (HCH; 150 mg/kg) on days 7 and 9 after commencing 2-AAF. Foci of resistant phenotypically altered hepatocytes, positive for γ -glutamyltranspeptidase, were seen only in rats treated with MNU and PH. Very few foci were seen in rats receiving MNU + sham operation, vehicle alone + PH, or no treatment. Similar results were obtained with male Wistar rats. Furthermore, when CCl₄ (0.25 ml/100 g) replaced HCH as a stimulus for cell proliferation, significant numbers of γ -glutamyltranspeptidase-positive foci were again seen only in rats receiving MNU + PH. Recent results suggested that MNU is a much more efficient inducer of putative initiated cells when administered 18 hr after PH. Thus, initiation of carcinogenesis by MNU is intimately related to some change in the liver that occurs after two-thirds PH. Cell proliferation may stimulate the induction of putative preneoplastic hepatocytes by replication of carcinogen-damaged DNA before repair. (22 refs.)

78-5085 Changes in Trace Element Levels During the Induction of Peripheral Nervous System Tumors. (Rus) Andronikashvili, E. L. (Inst. Physics, Tbilisi, USSR); Sharashidze, L. K.; Mosulishvili, L. M.; Beniashvili, D. S.; Kuchava, N. E.; Maisuradze, S. D. *Soobshch Akad Nauk Gruz SSR* 89(2): 473-476; 1978.

Changes in the amounts of various trace elements during the development and growth of induced peripheral nervous system (PNS) tumors were evaluated in random-bred albino rats. The rats were subjected to weekly iv injections of methyl nitrosourea (MNU: 10 mg/kg). Every 30 days, some of the animals were sacrificed, and the amounts of Fe, Zn, Rb, Cr, Se, Co, Ba, Hg, Ag, and Sb were assessed in the blood, sciatic nerve, and, after 3-5 mo, in the PNS tumors (benign reticular neurofibromas). The levels of the elements did not change within the first 2 mo of the experiment. From day 90, however, amounts of Zn, Se, Cr, Fe, and Rb increased markedly. The rate of increase was max between days 60 and 120, ie, during the period when the tumors could not be yet detected morphologically. (5 refs)

78-5086 Chemical Carcinogenesis and Immunity: Immunologic Status of Rats Treated with Methyl Nitrosourea. (Eng) Zwilling, B. S. (Dept. Microbiology, Columbus, OH, 43210); Filippi, J. A.; Chorpennig, F. W.; Koestner, A.; Rheins, M. S. *J Natl Cancer Inst* 61(3): 731-735; 1978.

During the treatment of outbred Sprague-Dawley rats with methyl nitrosourea (MNU) or the noncarcinogenic analog diphenylnitrosamine, antibody levels to teichoic acid as well as several parameters of lymphocyte and macrophage function were assessed in animals not overtly stimulated with antigen. Groups of animals received 4.5, 1.5, or 0.5 mg/kg MNU or gastric intubation every other week for 36 wk or similarly 4.5 mg/kg diphenylnitrosamine. MNU treatment did not appear to alter most immunologic parameters studied. Some alterations occurred in natural antibody levels, in spleen weight, and in peripheral blood differentials of rats that had received the highest carcinogen dose. These alterations, however, appeared to coincide with the development of tumors and were not observed in animals with premalignant lesions. Levels of chemical carcinogens that were capable of inducing tumors did not necessarily suppress the immune response. (23 refs)

78-5087 Genetics of Susceptibility in the Platyfish/Swordtail Tumor System to Development of Fibrosarcoma and Rhabdomyosarcoma Following Treatment with N-Methyl-N-nitrosourea (MNU). (Eng) Schwab, M. (Genetisches Institut, Justus-Liebig-Universität Giessen, Heinrich-Buff-Ring 58-62, D-6300 Giessen, W. Germany); Abdo, S.; Ahuja, M. R.; Kollinger, G.; Anders, A.; Ander, F.; Frese, K. *Z Krebsforsch* 91(3): 301-315; 1978.

The susceptibility (S) of 65 genotypes of xiphophorine fish (platy and swordtail fish) to the development of mesenchymal tissue neoplasms [fibrosarcomas (Fi) and rhabdomyosarcomas (Rh)] was tested following treatment of 7,000 fish with N-methyl-N-nitrosourea (MNU). The fish were given four 1-hr treatments with 10^{-3} M MNU at 2-wk intervals. The degree of S to the development of Fi and Rh varied considerably, ranging from zero in certain genotypes to almost 9% in others. This suggests that the degree of S is determined by the genetic constitution of the animal. An S that was high and reproducible was obtained only in backcross generations resulting from *Platyopocilus variatus*/*Xiphophorus helleri* hybrids and *X. helleri* as the recurrent parent. Purebred species of *P. variatus* and *X. helleri* and their F_1 progeny were not susceptible. The differential S may be a function of the type of control of a tumor gene (*Tu-Fi-Rh*) endogenous to *P. variatus* and involved in the development of Fi and Rh. In *P. variatus* and in the F_1 offspring, the *Tu-Fi-Rh* gene is controlled by repressing genes (*R* genes) linked and not linked to *Tu-Fi-Rh*. Because simultaneous mutation of both *R* genes after MNU treatment is extremely unlikely, these genotypes have an extremely low S. In the backcross hybrids, however, the unlinked *R* gene is eliminated and only the linked *R* gene remains. This condition confers a high degree of S, because one single mutation may lead to impairment of this *R* gene and to the *Tu-Fi-Rh*-mediated formation of Fi and Rh. It is concluded that in xiphophorine fish, S to the development of MNU-induced Fi and Rh is related directly to the genotype. (60 refs)

78-5088 Enhanced Development of Mammary Tumors in Rats Following Transplacental and Neonatal Exposure to Ethylnitrosourea. (Eng) Mandybur, T. I. (Dept. Pathology, Univ. Cincinnati Medical Coll., Medical Sciences Building, 231 Bethesda Ave., Cincinnati, OH, 45267); Ormsby, I.; Buncher, C. R. *Cancer Res* 38(10): 3182-3185; 1978.

The transplacental and neonatal induction of mammary tumors (MT) in Sprague-Dawley rats by ethylnitrosourea (ENU) was investigated. Six experimental groups were used: (1) untreated controls, (2) rats treated transplacentally (T) on days 16 or 17 of gestation with 1 ml/kg trioctanoin; (3) rats treated T on day 16 or 20 of gestation with 10 mg/kg ENU ip; (4) rats treated on day 4, 5, or 7 of life with 30 mg/kg ENU ip; (5) rats treated T with 60 mg/kg ENU; and (6) rats treated T with 120 mg/kg ENU. Tumors developed in 10/35 Group 1, 9/21 Group 2, 4/48 Group 3, 14/25 Group 4, 25/63 Group 5, and 4/9 Group 6 animals. The median tumor induction time decreased steadily from 17.5 mo in Group 1 to 4.5 mo in Group 6. Several tumors from each group were examined histologically; the numbers of benign (B) and malignant (M) tumors in each group were as follows: Group 1 5B, 3M; Group 2 6B, 3M; Group 3 2B, 2M; Group 4 1B, 8M; Group 5 6B, 17M; Group 6, 1B 6M. Sarcomas were observed only in Groups 1 and 3; carcinomas predominated in the high-dose ENU groups. Tumors developed mostly in females. Approx 88% of the MT-bearing rats in Groups 4,

5, and 6 MT also had neurogenic tumors. It is concluded that transplacental and neonatal exposure to ENU induces mammary carcinogenesis in susceptible animals. (19 refs.)

78-5089 Scanning Electron Microscopy of Cloned Astrocytic Lines Derived from Ethylnitrosourea-induced Rat Gliomas. (Eng) Spence, A. M. (Div. Neurology, RG-20, Univ. Washington, Seattle, WA, 98195); Coates, P. W. *Virchows Arch Cell Pathol* 28(1): 77-85; 1978.

Six cloned astrocytoma cell lines derived from four ethylnitrosourea-induced F-344 rat gliomas (36B, spinal malignant astrocytoma; 40A, cerebral malignant astrocytoma; 44B, spinomedullary astrocytoma; and 48A, cerebral glioblastoma multiforme) were examined in vitro by scanning electron microscopy (SEM). Two of these lines were further examined in vivo after intracerebral transplantation into syngeneic, sex-matched rats. All clones consisted of stellate cells that were reasonably homogeneous within individual glioma lines. The surface features of the lines tended to be rough. They were marked by the presence of microvilli mainly on cell bodies, rounded elevations, miniridges, ruffles, occasional blebs, and filopodia projecting from cell processes. The degree of cell membrane roughness varied among the lines. Line 40A-2 showed an extreme degree of microvillous formation and ruffling that correlated with a short cell-doubling time. Line 36B-10 was intermediate between line 40A-2 and the other lines, which showed fewer microvilli, ruffles, and filopodia. These features of lines 40A-2 and 36B-10 were maintained after transplantation into brain. The distinct variations in the cell membrane topography in these tumors are thought to be unrelated to their species of origin or mode of genesis (spontaneous, virally, or chemically induced). (20 refs.)

78-5090 Selective Induction of Melanomas in Gerbils (*Meriones unguiculatus*) Following Postnatal Administration of N-Ethyl-N-nitrosourea. (Eng) Kleihues, P. (Pathologisches Institut, Klinikum der Albert-Ludwigs-Universität, 7800 Freiburg, W. Germany); Bucheler, J.; Riede, U. N. *J Natl Cancer Inst* 61(3): 859-863; 1978.

Following a single ip injection (50 mg/kg) of N-ethyl-N-nitrosourea (ENU) on postnatal day 7 or 20, 9/21 gerbils developed a total of 15 cutaneous melanomas. The mean survival time of tumor-bearing animals was 909 ± 212 days. Neoplasms were preferentially located at sites with little hair and with a high content of pigment cells (feet, ears, tail, or snout). Histologic studies of newborn gerbils indicated that ENU-induced malignant transformation occurred in differentiated melanocytes rather than in precursor cells of the neural crest. (21 refs.)

78-5091 Changes in Surface Morphology Related to Ethylnitrosourea-induced Malignant Transforma-

tion of Cultured Rat Brain Cells (Meeting Abstract). (Eng) Roscoe, J. P. (Sch. Pathology, Middlesex Hosp. Medical Sch., London, England); Winslow, D. P.; Rowles, P. *Br J Cancer* 38(1): 192; 1978. (no refs)

78-5092 In Vitro Malignant Transformation of Fetal Brain Cells Exposed In Uterus to Ethylnitrosourea (ENU) (Meeting Abstract). (Eng) Yoshida, J. (New York Univ. Medical Center, New York, NY); Cravito, H.; Ransohoff, J. *J Neuropathol Exp Neurol* 37(5): 711; 1978. (no refs)

78-5093 Effect of Dibutyl Cyclic AMP on Rat Tumorigenesis by Ethylnitrosourea (Meeting Abstract). (Eng) Nagai, M. (Dept. Neurosurgery, Dokkyo Univ. Sch. Medicine, Mibu, Tochigi, Japan); Arai, T.; Awa, H. *J Neuropathol Exp Neurol* 37(5): 664; 1978. (no refs)

78-5094 Induction of Malignant Tumors in Wistar and Sprague-Dawley Rats by Single Doses of n-Butyl-nitrosourea in Perinatal and Juvenile Phases of Development. (Ger) Zeller, W. J. (Deutsches Krebsforschungszentrum, Institut für Toxikologie und Chemotherapie, Im Neuenheimer Feld 280, D-69 Heidelberg 1, W. Germany); Ivankovic, S.; Zeller, J. *Arch Geschwulstforsch* 48(1): 9-16; 1978.

n-Butylnitrosourea (BNU: 120 mg/kg po, ip, or sc) was given to pregnant Wistar rats on day 22 postconception (pc); the same dose was injected in the interscapular region of fetuses delivered surgically on day 22 pc, 1- and 2-day-old Wistar rats, and 10- and 30-day-old Sprague-Dawley rats. In 50% of the progeny, tumors (mainly neurogenic) developed both after diaplacental and direct application of BNU on day 22 after conception. After neonatal application, tumors (almost exclusively neurogenic) were found in >85% of the treated animals. Comparison of survival time and tumor rates among the various groups revealed that the nervous tissue of the rat is less sensitive to BNU prenatally than in the neonatal phase of development. After application of BNU to 10-day-old rats, >95% of the animals developed neurogenic tumors almost exclusively, and there was a high percentage of brain tumors. There was a significant increase in survival time and a decrease in total tumor yield in rats treated at age 30 days. There was also a decrease in the rate of neurogenic tumors and an increase in tumors outside the nervous system. Single doses of BNU were not leukemogenic in the late prenatal and neonatal phase: only 2/154 animals developed leukemia. Following BNU treatment of 30-day-old rats, leukemia was diagnosed in 7/40. However, this increased incidence in 30-day-old rats over that seen in younger rats was not considered significant. (21 refs.)

78-5095 Involvement of Chromosome #2 in Chromosome Changes in Primary Leukemia Induced in Rats by N-Nitroso-N-butylurea. (Eng) Uenaka, H. (Dept. Pathology, Kobe Univ. Sch. Medicine, Ikuta-ku, Kobe 650 Japan); Ueda, N.; Maeda, S.; Sugiyama, T. *J Natl Cancer Inst* 60(6): 1399-1404; 1978.

Cytogenetic studies were made of 34 primary leukemias induced in outbred Long-Evans rats by N-nitroso-N-butylurea (NBU: 0.04% in drinking water for 125 days, beginning at age 7 wk). The results revealed leukemia cells with No. 2 trisomy and a long No. 2 chromosome, changes that are characteristic of the leukemias induced by polycyclic hydrocarbon carcinogens such as 7,12-dimethylbenz(a)anthracene and 7,8,12-trimethylbenz(a)anthracene. These findings suggest a common mode of action of different carcinogenic chemicals at the chromosome level, although the lower incidence of these chromosome changes and of erythroblastic leukemias with NBU implies subtle differences in their actions. (26 refs.)

78-5096 Studies on N,N'-Dimethylnitrosourea-induced Brain Tumors in Rats--Especially, on Microtumors. (Eng) Jinnai, H. (Dept. Pathology, Okayama Univ. Medical Sch., Okayama 700, Japan). *Acta Med Okayama* 32(2): 119-137; 1978.

A study was made of the incidence, distribution, and histopathology of tumors induced by N,N'-dimethylnitrosourea (DMNU) in the brains of Sprague-Dawley rats. The rats were inoculated sc, in the interscapular region, with 40 mg/kg/wt of DMNU; rats were sacrificed 24 hr after 1, 5, or 10 injections or after 150-246 days (total, 17-21 injections). Animals sacrificed after 1-10 injections showed no pathological changes in any organs. Between 157 and 246 days, however, 122 gliomas developed in 38/55 rats. Sixty-six microtumors (diameter <1 mm) were detected in 28/38 rats by light microscopic observation of serial sections. The microtumors were of three types: oligodendrogliomas (55 tumors, 30 of which developed in the corpus callosum), astrocytomas (8 tumors, 5 of which were in the cerebral cortex), and mixed gliomas (3 tumors). Anaplasia occurred as tumors became larger, and cellular atypism was observed in 11/56 enlarged tumors. The target cells of DMNU were well-differentiated glial cells that had migrated from the matrix layer. (25 refs.)

78-5097 The Relationship Between Mutagenicity and Carcinogenicity of Some Nitrosamines. (Eng) Andrews, A. W. (Chemical Carcinogenesis Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Thibault, L. H.; Lijinsky, W. *Mutat Res* 51(3): 319-326; 1978.

The mutagenicity of 4 nitrosotrialkylureas, 8 aliphatic nitrosamines, and 14 cyclic nitrosamines was determined using the Ames *Salmonella typhimurium* assay, and the results

were compared with existing carcinogenicity data. There was agreement between the mutagenicity and carcinogenicity data for 80% of the compounds and a lack of agreement for 20%. The Ames test failed to detect 3/22 carcinogens among the 26 nitrosamines tested. Until this or some other test demonstrates a quantitative and qualitative correlation between mutagenic potency and carcinogenic potency, it will be difficult to establish a hierarchy of carcinogenic risk among a group of compounds such as the nitrosamines. (16 refs.)

- 78-5098 Neoplastic Transformation of Human Diploid Fibroblast Cells by Chemical Carcinogens.** (Eng) Kakunaga, T. (Chemistry Branch, NCI, Bethesda, MD, 20014). *Proc Natl Acad Sci USA* 75(3): 1334-1338; 1978.

The ability of 4-nitroquinoline 1-oxide (4NQO) to transform KD fibroblasts derived from a cutaneous punch biopsy of a normal human adult was investigated. Alterations of the cell growth pattern such as higher density and piling up of cells were noticed in some fractions of cultures that were successively subcultured after 4NQO treatment (0.1 µg/ml). Morphologically altered cells retained this growth pattern and became established lines of transformed cells without showing the limited life-span characteristic of normal cells in culture. The transformed cells showed a higher saturation density and the ability to grow in soft agar. It is doubtful that selection of preexisting transformed human cells was the mechanism of transformation, because clones of these normal cells could also be used to assess the transforming effect of 4NQO. Preliminary studies suggested that numerous cell divisions were required for the development of transformation. Female nude mice were inoculated sc with 0.2 or 2.0 x 10⁶ transformed cells, and solid tumors appeared at the injection site after an av latent period of 6 wk. Treatment of cells with 0.1 µg/ml N-methyl-N'-nitro-N-nitrosoguanidine induced transformation in a manner similar to that following 4NQO treatment. However, transformation was not induced with 4-aminoquinoline 1-oxide, 3-methylcholanthrene (a carcinogen that cannot be metabolically activated by the target cells employed), or dimethyl sulfoxide (the solvent used in the experiments). (20 refs)

- 78-5099 Carcinogenicity of Quinoline and Its Derivatives in Rat (Meeting Abstract).** (Eng) Shirai, T. (Nagoya City Univ. Medical Sch., 4-35 Mizoho-cho, Miguho-ku, Nagaya-shi, Aichi-ku, Japan); Fukushima, S.; Murasaki, G.; Ogiso, T.; Takahashi, M.; Ito, N. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 109-110; 1977. (no refs)

- 78-5100 Binding of Polynucleotides with 4-Nitrosoquinoline 1-Oxide Producing the Same Free Radical as That Produced from 4-Hydroxyaminoquinoline 1-Oxide.** (Eng) Kodama, M. (Biophysics Div., Natl. Cancer Center Res. Inst., Tsukiji 5-1-1, Chuo-ku, Tokyo 104, Japan); Nagata, C.; Mochizuki, M.; Okada, M. *Gann* 69(2): 285-286; 1978.

Evidence is presented that 4-nitrosoquinoline 1-oxide (4-NOQO) produces the same free radical as that produced from 4-hydroxyaminoquinoline 1-oxide (4-HAQO) and that 4-NOQO binds it covalently with polynucleotides. 4-NOQO was soluble in dioxan, ethanol, and dimethyl sulfoxide, and it gave an electron spin resonance (ESR) signal with a fine structure of six lines. After the dioxan soln of 4-NOQO it was degassed, the signal was further resolved into hyperfine structure of 52 lines, which was the same signal as that observed after a dioxan soln of 4-HAQO was degassed. The free radical thus formed showed no decay, even after standing at room temperature for 3 days under anaerobic conditions. In contrast, the ESR signal from 4-HAQO decayed progressively. Thus, the free radical produced from 4-HAQO was formed by oxidation, but that formed from 4-NOQO was produced by reduction. The addition of a reducing agent such as sodium borohydride to the dioxan soln of 4-NOQO resulted in a marked increase in the ESR signal, whereas the same treatment decreased the signal in the 4-HAQO soln. Both 4-NOQO and 4-HAQO were sufficiently active to bind poly(G) without activation, the former being more active than the latter. The emission spectra of poly(G)-bound 4-NOQO and 4-HAQO showed max peaks at 450 nanometers; these peaks coincided with the emission spectrum of DNA isolated from 4-HAQO-treated cells. (7 refs)

- 78-5101 Carcinogenic Effect of N-Methyl-N'-nitro-N-nitrosoguanidine (MNNG) in European Hamsters.** (Eng) Ketkar, M. (Abteilung fur Experimentelle Pathologie, Medizinische Hochschule Hannover, Karl-Weichert-Allee 9, 3000 Hannover 61, W. Germany); Reznik, G.; Green, U. *Cancer Lett* 4(4): 241-244; 1978.

The effects of chronic administration of N-methyl-N'-nitro-N-nitrosoguanidine (MNNG) on European hamsters of both sexes were determined. The LD₅₀ was 1,378 mg/kg for males and 1,070 mg/kg for females. One-tenth of the LD₅₀ was administered in water by stomach tube once weekly for 20 wk to 10 animals of each sex. Av survival for males and females was 21 and 18 wk, respectively. Papillary hyperplasia and hyperkeratosis of the forestomach occurred in nearly all hamsters. The epithelium of the upper digestive tract was often thickened, and hemorrhages were noted in the intestine. Controls did not show neoplastic alterations. The earliest time to tumor appearance was 13 wk for males and 20 wk for females; av tumor latency was 25 wk. Tumor incidence was 80% in males, 30% in females. Eight males and three females developed multiple squamous cell papillomas of the forestomach with various degrees of keratinization. Invasive

squamous cell carcinomas of the forestomach were noted in three males and one female; invasion extended to the duodenum, small intestine, spleen, liver, and pancreas. In addition, fibrosarcomas of the forestomach (1 male) and glandular stomach (1 male) glandular stomach (1 male) were found. In males, tumors at other sites included 2 squamous cell papillomas of the pharynx, 2 squamous cell papillomas of the nasal cavity, 1 squamous cell carcinoma invading the nasal septum, and 1 squamous cell papilloma of the trachea. Females hamsters had no neoplastic growths in the nasal cavities, but they occasionally had hyper- and metaplastic epithelial changes. Two females did, however, develop invasive squamous cell papillomas of the larynx. (9 refs)

- 78-5102 Pathways of Mutagenesis and Repair in *Escherichia coli* Exposed to Low Levels of Simple Alkylating Agents.** (Eng) Schendel, P. F. (Imperial Cancer Res. Fund, Mill Hill Labs., London NW7 1AD, England); Defais, M.; Jeggo, P.; Samson, L.; Cairns, J. *J Bacteriol* 135(2): 466-475; 1978.

Repair processes induced in *Escherichia coli* strains following exposure to ethyl methane sulfonate, methyl methane sulfonate, and N-methyl-N'-nitro-N-nitrosoguanidine were studied to determine whether error-prone (*lexA*-dependent) or *lex*-independent repair occurs. Bacterial strains were incubated with various concentrations of the mutagens. Error-prone repair was responsible for the majority of mutants formed after a large dose of alkylating agent, but the results suggest that this repair probably does not contribute significantly to mutagenesis at low concentrations of these chemicals. The mutagenicity at low doses was reduced by a repair system constitutively present in *lexA* cells but absent in *lexA* mutants. This system reduced mutagenesis until a second error-free system, called the adaptive response, could be induced. The adaptive response was capable of dealing with a much larger amount of alkylation damage than the constitutive system, and it appeared to reduce mutagenesis by both decreasing the number of sites available for mutagenesis and delaying the induction of error-prone repair enzymes. Based on these observations, a model for chemically induced mutagenesis is discussed that maintains that the observed mutation frequency is dependent on a competition between these two error-free systems and the two mutagenic pathways. (29 refs.)

- 78-5103 A Possible Process of Conversion of Food Components to Gastric Carcinogens.** (Eng) Endo, H. (Cancer Res. Inst., Faculty Medicine, Kyushu Univ., Fukuoka 812, Japan); Ishizawa, M.; Endo, T.; Takahashi, K.; Utsunomiya, T.; Kinoshita, N.; Hidaka, K.; Baba, T. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring

Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1591-1607; 1977.

A possible process for the conversion of food components to gastric carcinogens was tested in order to identify causal factors for human gastric cancer. N-Methyl-N'-nitro-N-nitrosoguanidines (MNNG) was selected as a model carcinogen and MNNG-like compounds among food were component surveyed. Fifteen naturally occurring guanidine derivatives were selected; these compounds are similar in structure to N-methyl-N'-nitrosoguanidines (MNG), which yields MNNG by nitrosation under strong acidic conditions. Only methylguanidine (MG) showed mutagenicity in the *Salmonella typhimurium* assay after nitrosation in gastric juice (pH 1.2). This compound was not mutagenic when nitrosated at pH > 3.4. Methylnitrosocyanamide (MNC), which was identified as the mutagenic principle of nitrosated MG, was 10 times more mutagenic than MNNG; however, the mutagenic quality of both compounds was similar. The in vitro yield of MNC formed from MG and nitrite in gastric juice (pH 1.2) was 0.3%-0.5%; the in vivo yield was 6%-10% of the in vitro yield. MNC was much more labile in gastric juice than MNG, and it lost its activity quickly. When given po to rats with the drinking water MNC was strongly carcinogenic to the forestomach and esophagus and less carcinogenic in the glandular stomach. The production of nitrite from nitrate by the oral microflora has been estimated to be 10 mg/day. Since the daily amount of saliva swallowed is 1.5 liters, saliva may be the most important source of nitrite. (53 refs)

- 78-5104 Reticulum Cell Sarcoma Production in Mammals by Oral Administration of N-Methyl-N'-Nitro-N-Nitrosoguanidine (MNNG) (Meeting Abstract)** (Eng) Kamano, T. (Gastroenterology, Dept. Surgery, Juntendo Univ. Hosp., Hongo, Bunkyo, Tokyo, Japan); Kawasaki, M.; Kodama, T.; Kishino, H.; Kidokoro, T.; Kurihara, M. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 75-76; 1977. (no refs)

- 78-5105 The Transformation of Human Diploid Cells by Chemical Carcinogens.** (Eng) Kakunaga, T. (Chemistry Branch, NCI, NIH, Bethesda, MD, 20014). In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1537-1548; 1977.

The in vitro transformation of human diploid cells by chemical carcinogens is reported. The cells were derived from a skin

biopsy of a normal woman who did not show any sign of genetic disease. They were exposed to 4-nitroquinoline-1-oxide (4NQO), 4-aminoquinoline-1-oxide (4AQO), N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), or 3-methylcholanthrene (3-MC). Before treatment, the cells exhibited the typical morphology of fibroblast cells with no evidence of an epitheliallike morphology. Assays in which cells were (1) treated with 0.01-1.0 $\mu\text{g}/\text{ml}$ 4NQO or MNNG and then plated to form individual colonies; (2) maintained at confluence after treatment; or (3) treated, allowed to divide for four or five generations, and plated in agar failed to reveal any transformation. However, cells that were treated with 4NQO or MNNG for 30 min and then subcultured at 1:4 dilutions whenever they became confluent showed transformed foci after 5-10 passages. Morphologically altered cells were found in a small area of a very small number of dishes. The main alteration was the formation of an area with a high density of cells showing frequent mitosis. Transformed foci were not observed in cultures treated with 3-MC or 4AQO. Apparently, an av of at least 13 cell divisions is necessary for the transformation of human diploid cells. The transformed cells had an increased saturation density and an increased ability to grow in soft agar, and they grew exponentially. Treated, untransformed cells did not exhibit these characteristics. Transplantation of the transformed cells into nude mice resulted in tumors at the site of injection within 2-21 wk, depending on the cell line and the number of cells inoculated. Spontaneous tumor regression was observed in only one case. No tumors were produced by injection of morphologically unaltered cells. (17 refs.)

78-5106 A Radioimmunoassay for O⁶-Alkyl Guanine Derivatives (Meeting Abstract). (Eng) Kyrtopoulos, S. A. (Courtauld Inst. Biochemistry, Middlesex Hosp. Medical Sch., Middlesex, England); Swann, P. F. *Br J Cancer* 38(1): 170; 1978. (1 ref)

78-5107 Interaction of 9-Aminoacridine with 7-Methylguanosine and 1,N⁶-Ethenoadenosine Monophosphate. (Eng) Kubota, Y. (Dept. Chemistry, Yamaguchi Univ., Yamaguchi 753, Japan); Nakamura, H.; Morishita, M.; Fujisaki, Y. *Photochem Photobiol* 27(4): 279-481; 1978.

The fluorescence properties of 9-aminoacridine (9AA) in aqueous solns of 7-methylguanosine (7MeG) and 1,N⁶-ethenoadenosine monophosphate (eAMP) were investigated to obtain information on the interaction between 9AA and DNA bases. The fluorescence of 9AA was quenched by 7MeG and eAMP as well as by unmodified bases such as guanosine monophosphate and AMP. Both dynamic and static quenching processes were involved. (14 refs)

78-5108 Phosphate Diester Formation Following Reaction of β -Propiolactone with Thymidine-5'-

monophosphoric Acid. (Eng) Segal, A. (Lab. Organic Chemistry and Carcinogenesis, Inst. Environmental Medicine, New York Univ. Sch. Medicine, New York, NY, 10016); Mate, U.; Wörtman, M. *Chem Biol Interact* 21(2/3): 249-261; 1978.

The carcinogen β -propiolactone (BPL) was reacted with thymidine-5'-monophosphoric acid (dThd5'P, or TMPA) at pH 1.8 and at 24-76 C to determine whether BPL forms phosphodiester with TMPA. Acidic conditions were chosen to try to avoid complicating ring alkylations. At 37, 50, 63, and 76 C, the reactions produced unreacted TMPA, which represented 90% of the UV-absorbing material (UVAM) on paper chromatograms, and a product designated compound F2, which represented 10% of the UVAM; at 24 C, F2 represented 5% of the material. The identical UV spectra of TMPA and F2 at pH 1 and 13 ruled against ring alkylation; and, on the basis of chemical tests, alkaline phosphatase and snake venom phosphodiesterase (SVPD) treatment at both pH 7.2 and 9.0, and incubation at pH 1, 7, 9, 11, and 13, F2 was assigned the structure 5'-O-(2-carboxyethyl)phosphonothymidine. BPL-TMPA reaction at pH 6.6-7.0 produced two additional compounds, F3 and F4. The UV spectra of TMPA, F3, and F4 were identical at pH 1 and 13. TMPA represented 36% of the UVAM, the remainder being F2 (36%), F3 (24%), and F4 (4%). F3 contained BPL and it was a phosphodiester, since treatment of F3 with SVPD at pH 7.2 or 9.0 resulted in its complete conversion to TMPA. A tentative assignment of a structure containing a six-membered ring phosphate ester-acyl phosphate derivative was assigned to F3. Results of BPL-TMPA reactions at pH 7.0-7.5, 7.5-8.0, and 8.0-8.5 were similar to those at pH 6.6-7.0. No ring alkylation was detected. If BPL does form acyl phosphates, a phosphate ester-acyl phosphate cross-link may help to explain the association of nucleic acids and cellular membranes induced by BPL, the intermolecular linking of DNA, and the DNA-protein complexes produced by BPL. These phenomena could be responsible for abnormal gene expression leading to neoplasia. (49 refs.)

78-5109 Oral Toxicity of 1,1-Dichloroethylene in the Rat: Effects of Sex, Age, and Fasting. (Eng) Andersen, M. E. (Toxicology Detachment, Naval Medical Res. Inst., Wright-Patterson Air Force Base, OH, 45433); Jenkins, L. J. *Environ Health Perspect* 21: 157-163; 1977.

The acute po toxicity of 1,1-dichloroethylene (1,1-DCE; vinylidene chloride) was investigated in male and female Holtzman HOT:(SD)BR rats of various sizes. When fasted and fed large, mature, male rats were given 400 mg 1,1-DCE, the hepatotoxic response was four to five times greater in the fasted rats. Determinations of the po LD₅₀ of male rats indicated that for large animals (395 g), mortality increased monotonically from 0% to 100% as the dose increased from 800 to 2,000 mg/kg; the estimated acute LD₅₀ was similar to that reported previously, 1,550 mg/kg. With males weighing 224 g, mortality varied between 10% and 33% following

doses of 50-800 mg/kg 1,1-DCE. With rats weighing 73 g, 100% mortality was noted at 300 mg/kg; percent mortality then decreased as the dose was increased up to 800 mg/kg. Maxima and extended plateaus were seen in the survival curves of all males between doses of 100 and 700 mg/kg, making it impossible to calculate an exact LD₅₀. Following a dose of 50 mg/kg, mortality and hepatotoxicity were greatest in rats weighing 100-150 g; smaller and larger males were less susceptible. Female rats did not show this variation in susceptibility with size, and they were less susceptible to the toxic effects of 1,1-DCE: the threshold of toxicity in male rats of 100-150 g occurred near 50 mg/kg, while for females it was closer to 100 mg/kg. These findings suggest that 1,1-DCE is metabolized to a toxic intermediate by some saturable pathway. Based on the effects of pretreatment with microsomal enzyme inhibitors and activators on 1,1-DCE toxicity in rats of various sizes, it appears that there are at least two microsomal reactions involved in 1,1-DCE metabolism. (25 refs)

78-5110 Hepatic Macromolecular Binding Following Exposure to Vinyl Chloride. (Eng) Watanabe, P. G. (Toxicology Res. Lab., Health and Environmental Res., 1803 Building, Dow Chemical Co., Midland, MI, 48640); Zempel, J. A.; Pegg, D. G.; Gehring, P. J. *Toxicol Appl Pharmacol* 44(3): 571-579; 1978.

Male Sprague-Dawley rats were exposed by inhalation to 1, 25, 50, 250, 1,000, or 5,000 ppm ¹⁴C-labeled vinyl chloride (VC) for 6 hr, and covalent binding of VC to hepatic macromolecules and nucleic acids was studied to determine if VC-induced carcinogenesis might be related to electrophilic alkylation of macromolecules in vivo. The total amount of VC metabolized and hepatic glutathione content (GSH) content were also measured. The total amount of radioactivity bound to hepatic macromolecules did not increase proportionately to the increase in the exposure concentration of VC, but it was related directly to the total amount of VC metabolized. At exposure concentrations > 50 ppm, the total metabolism of VC and covalent binding appeared to correlate with the induction of hepatic angiocarcinomas in the rats. Isolation of RNA and DNA by a nondigestive procedure from the liver of rats exposed to 1, 100, 250, and 1,000 ppm VC failed to reveal any detectable radioactivity. Hepatic GSH was depressed significantly only at ≥ 100 ppm, suggesting that VC carcinogenicity is related to a decreased ability to detoxify the reactive metabolites of VC. The results do not associate the carcinogenic effect of VC with a disproportionate increase in binding of electrophilic metabolites of VC to hepatic macromolecules as the exposure concentration is increased. Moreover, the lack of preferential binding of the metabolites to the hepatocyte nucleic acids suggests that the carcinogenicity of VC may not be associated directly with this commonly accepted mechanism of carcinogenesis. (27 refs.)

78-5111 Biochemical Effects of Acrylonitrile on the Rat Liver, As Influenced by Various Pretreatments

of the Animals. (Eng) Duverger-van Bogaert, M. (Lab. Biotoxicology, Univ. Louvain, Sch. Pharmacy, UCL-73-69, B-1200 Brussels, Belgium); Lambotte-Vandepaer, M.; Noel, G.; Roberfroid, M.; Mercier, M. *Biochem Biophys Res Commun* 83(3): 1117-1124; 1978.

The effects of acrylonitrile (AN) on the liver enzymatic systems of adult male Wistar rats were studied. The activities of two marker enzymes of the endoplasmic reticulum, glucose-6-phosphatase and NADPH:cytochrome c reductase, were nonsignificantly lowered by treatment of the rats with 10 mg/kg AN ip. AN did not affect the integrity of the liver mitochondria or lysosomes or the levels of SGOT, SGPT, or lipid peroxidation. A dose-dependent increase in lactate dehydrogenase (LDH) and sorbitol dehydrogenase (SDH) levels induced by AN was enhanced by pretreatment with diethyl maleate (DEM) and suppressed by pretreatment with phenobarbital (PB), 3-methylcholanthrene (3-MC), Aroclor 1254, or L-cysteine. Surprisingly, however, AN inhibited 3-MC-induced increases in LDH and SDH. A non-dose-dependent decrease in cytochrome P-450 levels and in the activity of aldrin oxide synthetase was unaffected by DEM and inhibited by PB, 3-MC, Aroclor 1254, and L-cysteine. AN did not significantly decrease the K_m of aldrin oxide synthetase. Propionitrile (10 mg/kg) also increased the levels of SDH and LDH, but it had no effect on cytochrome P-450 or aldrin oxide synthetase. (25 refs.)

78-5112 Intermediate Metabolites of Thiocarbamides, Thioureylenes and Thioamides: Mechanism of Formation and Reactivity. (Eng) Ziegler, D. M. (Clayton Foundation Biochemical Inst., Univ. Texas at Austin, Austin, TX, 78712). *Biochem Soc Trans* 6(1): 94-96; 1978.

The formation of the intermediate metabolites of thiocarbamides, thioureylenes, and thioamides was investigated in vitro. Studies with microsomal amine oxidases and the thio compounds indicated that only micromolar concentrations of the latter were required to half-saturate the oxidase; at saturation, all compounds were oxidized at rates equal to those observed with tertiary amine substrates. However, unlike the tertiary amines, 2 moles each of oxygen and NADPH were consumed per mole of substrate oxidized, which indicates that the oxidase catalyzes the dioxygenation of sulfur-containing compounds. Dioxygenations catalyzed by a monooxygenase can occur by sequential enzymic oxidation of the initial substrate through a monooxygenated intermediate or by disproportion of the intermediate. Although in vitro experiments demonstrated that thiocarbamides and thioureylenes can be oxidized to sulfenic and sulfinic acids, it is unlikely that either metabolite could be formed in vivo, because the acids are rapidly reduced to the parent compound by glutathione. The reduction can lead to a cycle that would operate until the NADPH or oxygen became limiting. In vivo hamster experiments indicated that at toxic doses, the cycle is terminated by the loss of glutathione: glutathione content

in the liver, lung, and bone marrow of methimazole-treated animals decreased in a time- and dose-dependent manner. It is suggested that the toxicity of thiocarbamides and thioureylenes may be partly due to their ability to oxidize glutathione. (11 refs)

- 78-5113 Enhancement of DL-Ethionine-induced Liver Carcinogenesis in Rats Fed a Choline-devoid Diet.** (Eng) Shinozuka, H. (Dept. Pathology, Univ. Pittsburgh Sch. Medicine, Pittsburgh, PA, 15261); Lombardi, B.; Sell, S.; Iammarino, R. M. *J Natl Cancer Inst* 61(3): 813-817; 1978.

The effects of a choline-devoid (CD) or a choline-supplemented (CS) diet on the induction of liver tumors in male Sprague-Dawley rats by DL-ethionine were determined. Groups of rats were fed a plain CD or a plain CS diet, or the same diets containing 0.05% DL-ethionine. Rats were killed after 14, 22, or 30 wk and the livers were examined. In rats fed the CD plus ethionine diet, hepatocellular carcinomas developed in 3/6 rats examined at 14 wk, 13/16 seen at 22 wk; and 6/8 seen at 30 wk. No liver tumors were noted in any of the other groups. The absolute food and ethionine intake of the rats receiving the CS plus ethionine diet was approx twice that of those fed ethionine plus the CD diet. Thus, the latter developed tumors despite the ingestion of less ethionine and a lower caloric intake. These findings suggest that a CD diet alters the response of rat liver to ethionine and leads to an early and enhanced induction of hepatocellular carcinoma. (28 refs.)

- 78-5114 Synthesis, DNA-Binding and Biological Activity of a Double Intercalating Analog of Ethidium Bromide.** (Eng) Kuhlmann, K. F. (Bio-Organic Chemistry Dept., Life Sciences Div., SRI International, Menlo Park, CA, 94025); Charbeneau, N. J.; Mosher, C. W. *Nucleic Acids Res* 5(7): 2629-2641; 1978.

The synthesis and DNA-binding characteristics of a bis[6-(3-substituted phenyl)] phenanthridinium salt and its monomeric analog are reported. Evidence provided by UV and circular dichroism spectra, thermal denaturation profiles, and equilibrium dialysis of the drug-DNA complex led to the conclusion that both phenanthridine moieties intercalate in the helix. The double intercalator appears to be less potent than ethidium chloride as an inhibitor of nucleic acid synthesis in cultured L1210 cells, although it is more potent than a monomeric analog. The low potency may be due to a low cell influx rate. (30 refs.)

- 78-5115 The Mutagenicity and Cytotoxicity of Selenite, "Activated" Selenite and Selenate for Normal**

and DNA Repair-deficient Human Fibroblasts. (Eng) Lo, L. W. (Cancer Res. Centre, Univ. British Columbia, Vancouver, Canada); Koropatnick, J.; Stich, H. F. *Mutat Res* 49(3): 305-312; 1978.

At doses of 8×10^{-5} to 3×10^{-3} M, sodium selenite (Na_2SeO_3) induced DNA fragmentation, DNA-repair synthesis, chromosome aberrations, and a mitotic inhibition in cultured human fibroblasts. The response of DNA repair-deficient xeroderma pigmentosum (XP) fibroblasts to selenite was comparable to that of control cells. Incubation with mouse liver S-9 microsomal fraction enhanced the lethal effect of selenite and increased its ability to induce chromosome aberrations and DNA-repair synthesis. XP cells behaved as control cells when treated with activated selenite. Sodium selenite (Na_2SeO_3) at doses of 8×10^{-5} to 3×10^{-3} M could not be activated by incubation with an S-9 preparation. Selenate induced a small but significant DNA-repair synthesis. (24 refs)

- 78-5116 Tumours and Disorders of Porphyrin Metabolism in Rats with Chronic Experimental Lead Poisoning (Meeting Abstract).** (Eng) Zawirska, B. (Inst. Pathomorphology, Medical Sch., Wroclaw, Poland). *J Clin Chem Clin Biochem* 16(1): 56; 1978. (4 refs)

- 78-5117 In Vitro Neoplastic Transformation of Syrian Hamster Cells by Lead Acetate and Its Relevance to Environmental Carcinogenesis.** (Eng) Dipaolo, J. A. (Biology Branch, NCI, Bethesda, MD, 20014); Nelson, R. L.; Casto, B. C. *Br J Cancer* 38(3): 452-455; 1978.

Secondary Syrian hamster embryo cell cultures were treated with 0, 1.0 or 2.5 $\mu\text{g}/\text{ml}$ lead acetate for 8 days and observed for transformation. A total of 18 transformed colonies were noted in 9 petri dishes treated with 2.5 $\mu\text{g}/\text{ml}$, 9 colonies were noted in the 10 dishes treated with 1.0 $\mu\text{g}/\text{ml}$, and none were noted in the 11 controls. Cells derived from these transformed colonies produced fibrosarcomas when injected sc into Syrian hamsters or nude mice. Primary hamster embryo cultures were treated for 18 hr with varying dilutions of lead acetate, inoculated with 200 focus-forming units of simian adenovirus 7 (SA7), and assayed for enhancement of viral transformation. Cell lethality from the lead acetate treatment was less in this assay, probably because of the short time of incubation and the large number of cells used. Statistically significant enhancement of SA7 transformation was observed with lead acetate doses of 100 and 200 $\mu\text{g}/\text{ml}$. These findings suggest that lead salts may be potential carcinogens for humans. (13 refs.)

- 78-5118 Influence of Low Cadmium-Dosages in Sub-acute Experiments on the Drug Metabolism**

on Mice (Meeting Abstract). (Eng) Abel, J. (Dept. Toxicology, Univ. Duesseldorf, Moorenstrasse 5, D-4000 Duesseldorf, W. Germany); Dieter, H.; Menzel, H.; Ohnesorge, F. K. *Naunyn Schmiedebergs Arch Pharmacol* 302(Suppl): R16; 1978. (no refs)

78-5119 Carcinogenicity of Nickel Subsulfide for Respiratory Tract Mucosa. (Eng) Yarita, T. (Dept. Surgery, Inst. Pulmonary Cancer Res., Sch. Medicine, Chiba Univ., Chiba 280, Japan); Nettesheim, P. *Cancer Res* 38(10): 3140-3145; 1978.

The carcinogenicity of nickel subsulfide (Ni_3S_2) was investigated in female Fischer 344 rats bearing sc tracheal transplants from isogenic recipients. Chemical determinations in tracheas treated with 1 and 3 mg Ni_3S_2 indicated that the compound persisted in the trachea for 7 to 9 mo. It showed a marked toxicity for mucociliary epithelium, resulting in widespread atrophy and focal epithelial necrosis during the first 2 mo of exposure. The submucosa showed mononuclear infiltration and signs of fibroblastic and capillary proliferation. In carcinogenesis studies in which 1 mg Ni_3S_2 was implanted in the trachea, tumors were noted in 9/60 tracheas; the first tumor occurred at 11 mo. Of the nine, three were sarcomas and six were carcinomas. The former showed extensive invasion into the host tissues but no distant metastases. Tumors developed in 45/64 tracheas treated with 3 mg Ni_3S_2 ; the first tumor was observed at 6 mo. Only one of these tumors was a carcinoma, the rest were sarcomas. These findings suggest that, compared with some carcinogenic polycyclic hydrocarbons, Ni_3S_2 may not be a strong carcinogen for the epithelium of conducting airways. However, the carcinoma incidence in the 3 mg group may have been artificially low because of early and high sarcoma incidence; furthermore, the high toxicity of the compound might outweigh its carcinogenicity and result in a misleadingly low carcinoma incidence. (28 refs.)

78-5120 Gastro-intestinal Tumors in Mice of Three Strains Maintained on Fat-enriched Diets. (Eng) Szepsenwol, J. (Dept. Biological Sciences, Florida International Univ., Tamiami Trail, Miami, FL, 33144). *Oncology* 35(4): 143-152; 1978.

The effect of fat-enriched diets on the development of primary gastrointestinal tumors was determined in T.M., BALB/c, and C57 Br. mice. Six groups of T.M. mice were treated as follows: (1) controls, regular diet only; (2)-(6) regular diet supplemented with refined corn oil, crude corn oil, refined corn oil + up to 1.5% oleic and linoleic acids, monolein, and monostearin, respectively. There were two groups of BALB/c and C57 Br. mice, one a control group (7 and 9, respectively) and one fed a diet supplemented with raw egg yolk (8 and 10, respectively). Forestomach tumors (papil-

omas and squamous cell carcinomas) were found in all mice on fat-supplemented diets, and the incidence varied from group to group: (2) 7/209, (3) 55/196, (4) 97/328, (5) 37/166, (6) 2/155, (8) 10/365, (10) 87/177. The incidence in control groups was very low: (1) 3/195, (7) 0/218, (9) 1/165. In addition, none of the control groups developed pyloric tumors. In the test animals, the incidence of pyloric tumors was highest in Group 10 (27/177) and lowest in Group 2 (2/209) and Group 6 (0/115). None of the control groups and none of the T.M. mice developed intestinal tumors. These tumors were mostly polyps (2/365 in Group 8, 14/177 in Group 10) except for three adenocarcinomas also found in Group 10. The fact that BALB/c and C57 Br. mice developed intestinal tumors but T.M. mice did not may be due to a difference in susceptibility and/or a difference in their diets. It is suggested that cholesterol, levels of which are high in egg yolk, may play an important role in intestinal and gastric tumorigenesis. (33 refs)

78-5121 A Mutagen in the Feces of Normal Humans. (Eng) Bruce, W. R. (Ontario Cancer Inst., Toronto, M4X 1K9, Ontario, Canada); Varghese, A. J.; Furrer, R.; Land, P. C. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1641-1646; 1977.

Feces collected from four normal men (25-45 yr old) on full Western diets were assayed for mutagens using the *Salmonella* tester strains TA98, TA100, TA1535, and TA1537 with and without liver microsomes. An agent in the ether extract, E, of many feces was directly active on strains TA100, TA1535, and, to a lesser degree, TA98. The highest activities were usually seen in fraction EBE, a sample derived from E by washing with a basic aqueous solution. Tests with other fractions indicated that there were factors in the water-soluble feces fraction, EBA, that inhibit the expression of mutations. Thus, the actual mutagenicity of a feces sample can be determined only after chemical separation of the active material from the inhibitor. The fecal mutagen was further purified and identified as an N-nitroso compound. Mutagen and nitroso compound levels in the feces of the four donors were reduced by > 60% when their normal diets were supplemented with ascorbic acid. (5 refs)

78-5122 Studies on Natural Desmutagens: Screening for Vegetable and Fruit Factors Active in Inactivation of Mutagenic Pyrolysis Products from Amino Acids. (Eng) Morita, K. (Dept. Induced Mutation, Natl. Inst. Genetics, Misima, Sizuoka-ken 411, Japan); Hara, M.; Kada, T. *Agric Biol Chem* 42(6): 1235-1238; 1978.

Juices from 59 fruits and vegetables were screened for their desmutagenicity to the mutagenic pyrolysis products of

-citrulline, L-lysine, L-ornithine, L-arginine, and L-tryptophan. Mutagenicity assays were carried out with the use of *Salmonella typhimurium* strain TA98 and a rat liver homogenate activating system. Ten samples, those of cabbage, broccoli, green pepper, eggplant, apple, burdock, shallot, ginger, pineapple, and mint leaf, were remarkably effective in inactivating the mutagenicity of L-tryptophan pyrolysis products. Eight others (radish, sweet potato, grape, Japanese ginger, cauliflower, beefsteak plant, enoki mushroom, and shimeji mushroom) were moderately effective. The other 41 samples were inactive. The 10 effective samples also showed varying degrees of desmutagenicity toward the pyrolysis products of the other amino acids. Three, eggplant, burdock, and broccoli, showed a high degree of desmutagenicity toward all pyrolysis products. The cabbage desmutagenic factor was not affected by ultracentrifugation, dialysis, or treatment with DNase and RNase. It was sensitive to heat (100 C) and to pronase treatment, which suggests that it is a protein. (5 refs.)

78-5123 **Influence of Diets High and Low in Animal Fat on Bowel Habit, Gastrointestinal Transit Time, Fecal Microflora, Bile Acid, and Fat Excretion.** (Eng) Cummings, J. H. (Medical Research Council, Dunn Nutrition Unit, Cambridge, England); Wiggins, H. S.; Jenkins, D. J.; Houston, H.; Jivraj, T.; Drasar, B. S.; Hill, M. J. *J Clin Invest* 61(4): 953-963; 1978.

The effect of diets containing either 62 or 152 g/day of fat (mainly animal) on the bowel habits, gastrointestinal transit time, fecal microflora pattern, and fecal bile acids and fat excretion of six healthy men was observed over a 4-wk period. The only significant change was a higher total fecal bile acid and fatty acid excretion in subjects on the high-fat diet. These results would support a theory linking colonic cancer with increased fecal bile acid excretion due to increased dietary fat intake. (73 refs)

78-5124 **Analysis of Mutagen-Antimutagen Reactions in Food and Food Additives by the Rec-Assay and Reversion-Assay Procedures (Meeting Abstract).** (Eng) Kada, T. (Natl. Inst. Genetics, Misima, Sizuoka-ken, Japan); Sadaie, Y.; Hara, M. *Mutat Res* 53(2): 206-207; 1978. (no refs)

78-5125 **CMOS: Environmental and Human Safety.** (Eng) Singer, E. J. (Res. Center, Lever Brothers Co., Edgewater, NJ); Mones, E.; Rothenstein, A. S.; Weaver, J. M. *Soap/Cosmetics/Chemical Specialties* 54(8): 38-44; 1978.

The results of environmental and biological studies of CMOS (a soluble nonphosphate detergent builder) are reported.

CMOS was readily biodegradable, and no accumulation is expected in the food chain. None of the CMOS salts tested showed any potential to induce cancer or mutations or to affect the reproductive performance or development of offspring. It was concluded that this compound will have no adverse environmental or health effects. (6 refs)

78-5126 **The Goitrogenic Effect of 4,4'-Oxydianiline in Rats and Mice.** (Eng) Hayden, D. W. (Dept. Veterinary Pathology, Coll. Veterinary Medicine, Univ. Minnesota, St. Paul, MN, 55108); Wade, G. G.; Handler, A. H. *Vet Pathol* 15(5): 649-662; 1978.

The carcinogenicity of 4,4'-oxydianiline (OD) was studied in Fischer 344 rats and B6C3F₁ mice maintained for 13 wk on diets containing 0.03%, 0.06%, 0.1%, or 0.2% of the compound. Stunted growth and morbidity were apparent in animals given 0.06% or more OD, and lethargy was observed in the 0.1% and 0.2% groups. Alopecia, dyspnea, and cyanosis were also seen in rats given the two highest doses. Mortality was observed only among the rats, and it was two times higher among females than in males: 4/10 females in the 0.1% group and 7/10 females in the 0.2% group died. Pathologic findings in the rats included bilateral thyroid enlargement that included the isthmus, presence of clear or red-tinged ascitic fluid, accentuation of the lobular pattern of the liver, atrophy of the male gonads and seminal vesicles, and embedment of the pancreas in gelatinous tissue. Slight to moderate thyroid enlargement was noted in some mice. Microscopic lesions in the thyroid, pituitary, and testes of both species and the prostate, seminal vesicles, kidneys, and bone marrow of rats were related to the chemical and its concentration in the diet. The data showed that OD has a goitrogenic effect and can alter thyroid function to initiate pituitary hyperplasia. (16 refs.)

78-5127 **Biphasic Change of Proton Magnetic Relaxation Times During Azo-Dye Hepatocarcinogenesis.** (Eng) Kodama, M. (Biophysics Div., Natl. Cancer Center Res. Inst., Tsukiji 5-1-1, Chuo-ku, Tokyo 104, Japan); Ohki, T.; Saito, H.; Nagata, C.; Tagashira, Y. *Br J Cancer* 37(2): 233-236; 1978.

The correlation between neoplastic change and change in the proton longitudinal relaxation times (T_1) in the liver, kidney, and blood serum of male Sprague-Dawley rats was studied for periods ≥ 150 days. This is the first report of such studies covering the whole process of azo-dye hepatocarcinogenesis. Rats were fed a diet containing 0.06% 3-methyl-4-dimethylaminoazobenzene for 87 days. The T_1 and water content of liver, kidney, and blood serum were measured. The T_1 of the liver exhibited two maxima, on days 60 and 160. The change of T_1 correlated well with the change in water content, which also showed two maxima on days 60 and 160. The formation of hepatic nodules was observed macroscopically

on day 60. Histological examinations revealed that the first peak of T_1 coincided, not with the disappearance of original hepatocytes, but with the proliferation of renewed hepatocytes. The second peak coincided with the development of hepatomas. Normal hepatic regions adjoining the tumors showed low T_1 values; thus, the change of T_1 accompanying the primary hepatoma was localized. (12 refs)

- 78-5128 Multigeneration Reproduction Studies with Carmoisine in Rats.** (Eng) Holmes, P. A. (Illinois Inst. Technology Res. Inst., Chicago, IL); Pritchard, A. B.; Kirschman, J. C. *Toxicology* 10(2): 169-183; 1978.

The effects of carmoisine (CM: a water-soluble red azo dye that is the disodium salt of 2-(4-sulfo-1-naphthylazo)-1-naphthol-4-sulfonic acid) on overall reproductive capacity were studied in three generations of Sprague-Dawley rats. The rats were exposed to 0%, 0.35%, 0.8%, or 2.0% CM in the diet for various time periods up to 1 yr. No significant effect was seen in fertility and copulation indices, birth statistics, offspring viability, or wt of pups. These results indicate a max no-effect level of 2% in the diet. A shorter term study involving the feeding of CM at very high levels (0%, 2%, 4%, 6%, or 8% for 9 wk) was made to establish the highest no-effect level possible. The rats showed normal wt gain with up to 4% CM in the diet. The wt loss observed at 6% and 8% CM may be due to a caloric deficit because of the high percentage of a nonnutritive substance (the CM) in the diet rather than a direct toxic effect of CM itself. These studies indicate that the max no-effect level for CM may be > 2%, since 4% was tolerated without evidence of overt toxicity. (4 refs.)

- 78-5129 A One Year Feeding Study with Carmoisine in Rats.** (Eng) Holmes, P. A. (Illinois Inst. Technology Res. Inst., Chicago, IL); Pritchard, A. B.; Kirschman, J. C. *Toxicology* 10(2): 185-193; 1978.

The effects of long-term exposure to carmoisine (CM: a water-soluble red azo dye that is the disodium salt of 2-(4-sulfo-1-naphthylazo)-1-naphthol-4-sulfonic acid) were studied in the F-2b generation of Sprague-Dawley rats fed CM in a multigeneration reproduction study. The F-2b rats were exposed to 0.35%, 0.8%, or 2.0% CM in the diet for 52 wk and then sacrificed. Body wts, hematological values, urinalyses, and relative organ wts were not significantly different from those of controls. No malignant tumors were diagnosed. The incidence of nonneoplastic lesions was the same in both treated and control females. In males receiving 2.0% CM, a significantly higher incidence of minimal bronchitis (6 cases/13 tissues examined), tracheal inflammation (6/13), and lymph node congestion (7/13) was noted. It is suggested that the max no-effect level after 1 yr of CM in the diet is 0.8%, or 400 mg/kg/day. However, since the deleterious effects seen at the 2.0% level were considered minimal, the no-effect level may actually be much higher than 0.8%. (3 refs.)

- 78-5130 Hepatic Tumors after Oral Contraceptives (Meeting Abstract).** (Fre) Begue, J. Y. (No affiliation given); Leclerc, J.; Bigard, M. A.; Laugros, A.; Gaucher, P.; Rauber, G. *Gastroenterol Clin Biol* 2(5): 558; 1978. (no refs)

- 78-5131 Benign Tumors of the Liver and Oral Contraceptives.** (Fre) Letoublon, C. (Service des Urgences Chirurgicales, Clinique Obstetricale et Gynecologique du C.H.U. de Grenoble, Hopital Nord, 38700 La Tronche, France); Champetier, J.; Benbassa, A.; Durand, A.; Laborde, Y.; Pasquier, D. *Lyon Chir* 74(2): 121-124; 1978.

The case reports of two women (aged 36 and 33 yr), each with a hepatic hamartoma, are presented. The women had taken oral contraceptives for 4 and 7 yr, respectively. The histology of one of the tumors, which had been attached to the sigmoid colon, is presented. A review of 172 cases of benign hepatic tumor associated with oral contraceptive use indicated that the risk of developing a benign hepatic tumor is probably small. Hemorrhage with tumor rupture was the most severe complication: it occurred in > 25% of the cases, mostly in adenomas. (15 refs)

- 78-5132 Mammary Nodules in Beagle Dogs Administered Investigational Oral Contraceptive Steroids.** (Eng) Giles, R. C. (Dept. Veterinary Pathology, Armed Forces Inst. Pathology, Washington, DC, 20306); Kwapien, R. P.; Geil, R. G.; Casey, H. W. *J Natl Cancer Inst* 60(6): 1351-1364; 1978.

Of 172 beagle dogs given oral contraceptive steroids for 5-7 yr, 114 developed 1,156 nodules in the mammary gland region. Most of these nodules arose 2.5-3.5 yr after initiation of treatment. Approx 16% of the nodules disappeared spontaneously. A total of 925 nodules were present in 99 dogs at the time of death or necropsy. These nodules were classified as benign mammary dysplasias (7.0%), lobular or intraductal hyperplasias (31.4%), simple adenomas (20.8%), complex adenomas (25.4%), benign mixed tumors (5.3%), malignant tumors (3.6%), or nonmammary lesions (6.5%). Histologically, the mammary nodules were representative primarily of the hyperplasias and tumors that occur spontaneously in canine mammary glands. The major exception was the presence of 82 simple adenomas that had basaloid features. Most of the contraceptive-related mammary nodules developed in dogs receiving the combination of progestin and mestranol at 10 or 25 times the proposed human dose. Control dogs and dogs receiving mestranol alone had few mammary nodules. Combinations of anagestone acetate and mestranol and chloerthynyl norgestrel (WY-4355) and mestranol produced large numbers of nodules at 10 and 25 times the proposed human dose; ethynone plus mestranol produced large numbers of nodules only at 25 times the proposed dose. Ethynone, when given alone at 25 times the proposed dose, was associated with fewer mammary nodules. Malignant neo-

isms were seen in dogs given 10 and 25 times the proposed human dose of anagestone acetate plus mestranol and 25 times the proposed dose of WY-4355 plus mestranol and cyproterone plus mestranol. This study strongly associates certain combinations of progestin and mestranol with mammary neoplasia in dogs. (39 refs)

78-5133 Oral-Contraceptive-induced Benign Liver Tumors--The Magnitude of the Problem (Letter to the Editor). (Eng) Jick, H. (Boston Collaborative Drug Surveillance Program, Boston Univ. Medical Center, Boston, MA); Herman, R. *JAMA* 240(9): 828-829; 1978.

The annual number of benign liver tumors induced by oral contraceptives (OC) in the US estimated made based on (1) information obtained between 1970 and 1976 by the Commission on Professional and Hospital Activities (covering 40% of patients discharged from US hospitals) and (2) information on such tumors obtained in a study of 477 US hospitals during 1970-1975. The number was estimated to be roughly 100 and 140/yr, respectively, or one benign tumor for approximately 1,000 users. (3 refs.)

78-5134 Estrogen Induction of Growth Factors Specific for Hormone-responsive Mammary, Pituitary, and Kidney Tumor Cells. (Eng) Sirbasku, D. A. (Dept. Biochemistry and Molecular Biology, Univ. Texas Medical Sch., Houston, P.O. Box 20708, Houston, TX, 77025). *Proc Natl Acad Sci USA* 75(8): 3786-3790; 1978.

The problem of estrogen-promoted tumor cell growth was studied extensively in an attempt to establish the direct mitogenic role of steroid hormones. Permanent cell lines were developed from three estrogen-responsive rodent tumors: TW9/PL estrogen-, thyroid hormone-, and prolactin-responsive rat mammary tumor cells; GH3/C14 estrogen- and thyroid hormone-responsive rat pituitary tumor cells; and 301 estrogen-responsive hamster kidney tumor cells. All three cell lines required estrogens from optimal tumor formation in host animals, but they did not respond to estrogens or prolactin with increased cell growth rate, as measured by increased cell number, in culture. Extracts of rodent uterus, kidney, and liver were found to contain growth-promoting activity for all three tumor lines. Estrogen treatment increased the tissue content of these activities, and the estrogen-induced activities were specific for the estrogen-responsive cells. The data suggest that estrogen-responsive tumor growth in vivo involves estrogen stimulation of uterus, kidney, or liver to produce specific growth factors that, in turn, stimulate estrogen-responsive tumor cells. (37 refs.)

78-5135 Sex Hormones and Intra-Cranial Gliomata (Meeting Abstract). (Eng) Hopewell, J. W. (Res.

Inst., Churchill Hosp., Oxford, England). *Br J Cancer* 38(1): 171; 1978. (no refs)

78-5136 Colchicine-like Effect of Diethylstilbestrol (DES) on Mammalian Cells In Vitro. (Eng) Sawada, M. (Dept. Chemical Pathology, Natl. Inst. Hygienic Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158, Japan); Ishidate, M. *Mutat Res* 57(2): 175-182; 1978.

The cytological effects of diethylstilbestrol (DES) on mammalian cells in culture were examined in the following cell lines: CHT, derived from the thymus of a young male Chinese hamster; DL, derived from a Japanese Donryu rat liver; EDEN-1/TC, derived from a rat erythroblastic leukemia; and HeLa-S3. Metaphase arrest was induced 3 hr after treatment with 15 μ g/ml DES, and polyploid or polynucleated cells were prominent > 24 hr after treatment. The arrest was reversible when the agent was removed from the medium. Tetraploid karyotypes induced by DES in CHT cells consisted of double sets of diploid chromosomes except for one marker chromosome, which may have been formed by fusion or translocation of two No. 4 chromosomes. However, this marker may not be DES-specific, as it was not detected in other lines after DES treatment. By clonal selection, several hypotetraploid sublines were successfully isolated from a CHT cell population after it was treated with 15 μ g/ml DES for 48 hr. These studies demonstrate that DES has a direct effect on cell mitosis in vitro, probably on the formation of the spindle apparatus, similar to the effect induced by colchicine. (21 refs)

78-5137 Cytogenetic Studies with Diethyl Stilboestrol. (Eng) Bishun, N. P. (Tissue Culture and Cytogenetics Unit, Marie Curie Memorial Foundation, The Chart, Oxted, Surrey, England); Eddie, H.; Williams, D. C. *Clin Oncol* 4(2): 159-165; 1978.

Normal human peripheral blood WBC, two human carcinoma cell lines, and one established rat cancer cell line were exposed in culture to diethylstilbestrol at levels of 20, 15, 10, or 5 μ g/ml and examined for chromosome aberrations. No significant chromosome damage was noted at any of the concentrations used. The discrepancy between these findings and previous results is discussed. (6 refs.)

78-5138 Oxidative Metabolites of Diethylstilbestrol in the Fetal, Neonatal and Adult Mouse. (Eng) Metzler, M. (Inst. Toxicology, Univ. Wurzburg, D-8700 Wurzburg, W. Germany); McLachlan, J. A. *Biochem Pharmacol* 27(7): 1087-1094; 1978.

Urinary metabolites of ^{14} C-labeled diethylstilbestrol (DES) in adult female mice were identified by radio gas chromatogra-

phy and gas chromatography-mass spectrometry as β -dienestrol and the ω -hydroxy derivatives of DES, ψ -DES, and dienestrol. β -Dienestrol did not appear to be formed through a DES epoxide. It is suggested that DES might be oxidized to a p-quinone, which could easily be tautomerized to β -dienestrol. Most of the metabolites found in adult mice were also detected by mass fragmentography in the whole-body homogenates obtained from 1-, 4- and 8-day-old mice, indicating that oxidative metabolism of DES is operative in neonatal mice. Extensive conjugation of DES and its metabolites to glucuronic acid was found in adult and neonatal mice. Part of the administered radioactivity was unextractable from tissue macromolecules in neonatal mice. The biotransformation of DES and the binding of its metabolites might be associated with its teratogenic and carcinogenic potential for neonatal and fetal mice. (34 refs)

78-5139 Diethylstilbestrol Binding to Estrogen Receptor in Mammary Tissue and Uteri of C3H/HeJ Mice. (Eng) Okey, A. B. (Dept. Biology, Univ. Windsor, Windsor, Ontario, Canada N9B 3P4); Bondy, G. P. *J Natl Cancer Inst* 61(3): 719-724; 1978.

Diethylstilbestrol (DES) binding to components in mammary cytosol from C3H/HeJ mice was assessed directly (with ^3H -DES) and indirectly [by competition between DES and ^3H -17 β -estradiol (E_2)]. On sucrose density gradients, the qualitative properties of cytosol estrogen receptor were similar whether ^3H -DES was used as the ligand or whether the receptor was labeled with ^3H - E_2 . Binding of either ^3H -DES or ^3H - E_2 was extinguished by incubation in the presence of a 100-fold excess of unlabeled DES or E_2 . ^3H -DES also sedimented with a component in the 4S-5S region; this component bound much larger quantities of ^3H -DES than ^3H - E_2 and was not eliminated by incubation with a 100-fold excess of unlabeled estrogen. Both ^3H - E_2 - and ^3H -DES-labeled receptors dissociated to a 4S-5S form when centrifuged on gradients containing 0.4 M KCl. DES also induced receptor transformation and nuclear uptake, as evidenced by recovery of a 5S-region receptor from mammary nuclei of mice given injections in vivo with ^3H -DES. Dextran-coated charcoal assays/Scatchard plot analyses with ^3H -DES indicated a DES-receptor interaction with a dissociation constant of 2.6×10^{-10} M compared with 6.3×10^{-11} M for ^3H - E_2 . By competition studies, the apparent affinity of receptor for DES was essentially the same as that for E_2 . The apparent concentration of high-affinity binding sites in mammary cytosol was higher for ^3H -DES than for ^3H - E_2 ; this probably was due to ^3H -DES binding to nonreceptor proteins, possibly serum albumin. Overall, intracellular (receptor) binding of DES did not appear to be fundamentally different from binding of natural estrogens. (24 refs.)

78-5140 Acute Myeloblastic Leukemia Following Chlorambucil Treatment. Two New Cases. (Fre)

Witz, F. (Clinique Medical A, C.H.U., Nancy-Brabois, F 54500 Vandoeuvre, France); Lederlin, P.; Aymard, J. P.; Thibaut, G.; Guerci, O. *Nouv Presse Med* 7(27): 2392; 1978.

Two women, aged 47 and 52 yr, developed acute myeloblastic leukemia following long-term treatment with chlorambucil for multiple sclerosis. Each patient received 6 mg/day for a total dose of 13.3 g in 75 mo and 9.9 g in 55 mo, respectively. Both women died. (3 refs.)

78-5141 Comparative Study of Mutagenic Activity of a Series of Hypotensive Drugs in Mice. (Rus) Zolotareva, G. N. (Res. Inst. Biological Trials of Chemical Compounds, Moscow, USSR); Akaeva, E. A.; Iskhakova, E. N.; Oblapenko, N. G.; Kotlianskaia, A. E.; Epshtein, S. I. *Khim Farm Zh* 12(3): 19-22; 1978.

The potential mutagenic activity of a series of hypotensive drugs [reserpine, its commercial forms raucedil and raunatine, and clopheline, or 2,2,6-dichlorophenylamino-2-imidazoline hydrochloride] was tested in CBA x C57BL/6 mice. The drugs were given ip and po, at levels ranging from 0.5/LD₅₀ to therapeutic dosages. Mutagenicity was measured by the incidence of chromosome aberrations in bone marrow cells and by the induction of dominant lethals in germ cells. Raucedil at (10 and 5 mg/kg) and raunatine at (15 and 10 mg/kg) increased the incidence of single fragments, but therapeutic doses of these drugs lacked mutagenic activity. Clopheline at all doses tested showed no mutagenic activity. Exposure of male mice to reserpine decreased the number of pregnant females and increased the preimplantation death of the zygotes. (5 refs)

78-5142 Thyroid Tumours in Rats and Hepatomas in Mice after Griseofulvin Treatment. (Eng) Rustia, M. (Eppley Inst. Res. Cancer, Univ. Nebraska Medical Center, Omaha, NB, 68105); Shubik, P. *Br J Cancer* 38(2): 237-249; 1978.

The carcinogenicity of the antibiotic, griseofulvin, was studied in Swiss mice, MRC-Wistar rats, and Syrian hamsters. Three groups of mice and rats were given the drug in a powdered diet in alternating 5-wk periods for life, at dose levels of 3.0%, 1.5%, and 0.3% (mice) and 2.0%, 1.0%, and 0.2% (rats). A fourth group of mice received continuous daily treatment for life with 0.1% griseofulvin, and three groups of hamsters were similarly treated at levels of 3.0%, 1.5%, and 0.3%. In mice, the two highest doses produced a significant incidence of hepatic tumors in both males and females. At the 3.0% level, females had a higher percentage of hepatomas than males (87% vs 83.3%); at the 1.5% level, males had a higher percentage than females (68% vs 53.6%). Histologically, the liver tumors were either well-differentiated and solid, with a structure similar to that of normal liver; less-differentiated and trabecular; or (occasionally) glandular, in

which case the liver structure was completely obliterated. Contrary to previous findings, statistically significant rates (≤ 0.001 and/or $P \leq 0.02$) of thyroid tumors were recorded in male rats at all dose levels and in females at the two higher levels. A classic dose-response relationship was observed. Histologically, the thyroid tumors were predominant adenomas, consisting of follicles (macro- or microfollicular types) and/or papillary-cystic structures. In hamsters, no neoplasms developed in response to treatment at any level. (1 refs.)

78-5143 Accelerated Growth of Mammary Tumor Cells in Normal and Athymic Mice after Treatment In Vitro with Dexamethasone. (Eng) Gillette, R. W. (Cancer Center Hawaii, Honolulu, HI, 96822); Wunderlich, D. A. *Cancer Res* 38(10): 3146-3149; 1978.

The tumorigenicity of dexamethasone (DM)-treated C3HMT mouse mammary adenocarcinoma cells was compared with that of untreated C3HMT cells. Cultures of tumor cells were treated with $0.1 \mu\text{g/ml}$ DM and harvested after 4 days of steroid treatment. DM increased RNA-directed DNA polymerase activity in the cultures, which indicated increased murine mammary tumor virus expression. Both treated and untreated cells induced tumors in 20/20 C3H mice inoculated with 10^6 cells. At doses of 10^5 , 10^4 , and 10^3 cells however, tumors were noted in 19/19, 14/20, and 8/20 mice receiving DM-treated cells, respectively, compared with 15/20, 2/20, and 0/20 in mice receiving the untreated cells. Further studies suggested that the DM treatment did not measurably change the immunogenicity of the C3HMT cells. Upon injection of 10^5 treated or untreated cells into athymic nude mice, all animals developed tumors, but the DM-treated tumors progressed more rapidly. Thus, immunostimulation by T cells was not a factor in the accelerated growth of the DM-induced tumor cells. DM treatment (1 mg) of the host at the time of tumor cell inoculation did not influence tumor growth, but thrice weekly injections of 1 mg DM for 3 wk after tumor inoculation had a negative effect on C3HMT progression. The possibility of DM immunostimulation of other kinds of lymphoid cells is discussed. (12 refs.)

78-5144 Oncogenic Transformation In Vitro by the Hypoxic Cell Sensitizer Misonidazole. (Eng) Miller, R. C. (Radiological Res. Lab., Coll. Physicians and Surgeons Columbia Univ., New York, NY, 10032); Hall, E. *Br J Cancer* 38(3): 411-417; 1978.

The effect of the hypoxic cell sensitizer Misonidazole (MIS; 0-07-0582) on aerated C3H/10T1/2 cells was determined. Cultured cells were exposed to varying doses of radiation and/or to 0.4-8.0 mM MIS for 3 days or 0.5-2.0 mM MIS for 6 days. Doses as low as 0.4 mM resulted in a significant transformation rate. This rate clearly increased with dose. However, the period during which MIS was in contact with

the cells had little influence on the number of cells transformed. A concentration of 0.4 mM for 3 or 6 days resulted in a transformation rate comparable to that observed following an x-ray dose of 1 Gray. A concentration of 6.0 mM was equivalent to 4 Gray. When cells were exposed to 1 mM MIS for 3 days and, 24 hr after the start of MIS exposure, to 1, 2, or 4 Gray of x-rays, there was a significant increase in the frequency of transformation over that seen with either the drug or the radiation alone. It is not known whether the action between the two agents was additive or synergistic. (18 refs.)

78-5145 Caffeine Inhibition of Prereplication Repair of Mitomycin C-induced DNA Damage in Human Peripheral Lymphocytes. (Eng) Ishii, Y. (Dept. Fundamental Radiology, Faculty Medicine, Osaka Univ., Kita-ku, Osaka 530, Japan); Bender, M. A. *Mutat Res* 51(3): 419-425; 1978.

The effect of caffeine ($0-300 \mu\text{g/ml}$) on the prereplication repair of mitomycin C (MMC)-induced DNA damage in human peripheral lymphocytes was studied. Caffeine alone induced some sister chromatid exchanges (SCEs) and also interacted synergistically to produce large increases in the yields of MMC-induced SCE's. This enhancement decreased when the MMC-treated cells were held in medium before phytohemagglutinin (PHA) stimulation or when caffeine was added to the cultures some time after PHA stimulation but prior to DNA synthesis. The data confirm the existence of a caffeine-sensitive prereplicative DNA repair mechanism in human peripheral lymphocytes. It appears that this repair system becomes saturated and is only capable of repairing a fraction of those MMC-induced DNA lesions capable of inducing SCE's. (45 refs.)

78-5146 Cancers among Users of Preparations Containing Vitamin A. A Case-Control Investigation. (Eng) Smith, P. G. (DHSS Cancer Epidemiology and Clinical Trials Unit, Univ. Oxford, 9 Keble Road, Oxford, England); Jick, H. *Cancer* 42(2): 808-811; 1978.

The regular use of vitamin A preparations was determined among 800 newly diagnosed cancer patients and 3,433 patients with certain nonmalignant conditions. The age range of both groups was 20-75 yr. Regular usage was defined as at least once a week for the 3-mo period or more prior to hospital admission. Overall, there was no convincing evidence that regular consumption of such preparations protected against the development of cancer (risk ratio 0.85). However, there was a suggestion of a negative association among men (risk ratio 0.54) but not among women (risk ratio 1.11). The negative association among men did not increase with the period that the preparations had been regularly taken and was not greater for squamous cell tumors. There is experimental evidence from animal experiments and in vitro labora-

tory investigations indicating that vitamin A deficiency predisposes to squamous metaplasia and that vitamin A protects against the development of some chemically induced neoplasms, particularly squamous carcinomas. (8 refs.)

- 78-5147 **Aplasia and Leukaemia Following Chloroquine Therapy.** (Eng) Nagaratnam, N. (General Hosp., Colombo, Sri Lanka); Chetiyawardana, A. D.; Rajiyah, S. *Postgrad Med J* 54(628): 108-112; 1978.

Case reports are presented for three patients who developed aplastic anemia following chloroquine (CQ) therapy. A 69-yr-old man, who subsequently developed acute myeloblastic leukemia, had taken large doses for several months for the treatment of discoid lupus erythematosus. A 29-yr-old woman had received CQ for malaria on several occasions and had been on prophylactic CQ for several months. In a 22-yr-old man, anemia with neutropenia occurred after a 3-wk course of CQ for malaria. Although parallels can be drawn with chloramphenicol-induced blood dyscrasias, the relationship between CQ and bone marrow injury is unclear. (12 refs)

- 78-5148 **Carcinogenic and Other Adverse Effects of Procarbazine in Nonhuman Primates.** (Eng) Sieber, S. M. (Lab. Chemical Pharmacology, NCI, NIH, Building 37, Room 5A13, Bethesda, MD, 20014); Correa, P.; Dalgard, D. W.; Adamson, R. H. *Cancer Res* 38(7): 2125-2134; 1978.

The carcinogenic and toxic effects of procarbazine (PRO) were investigated in 55 monkeys who received the drug sc, po, or ip for periods of up to 12 yr. Thirteen monkeys are alive and without evidence of tumor. Among the 42 animals that were necropsied, 11 have had malignant neoplasms. The malignancies included 6 cases of acute leukemia (5 myelogenous and 1 undifferentiated), 2 of osteogenic sarcomas, 2 hemangiosarcomas, and 1 lymphocytic lymphoma. In contrast, only 2/211 control and breeder monkeys maintained during the same time period developed malignant tumors. The av total dose of PRO received by the monkeys developing cancers was 36.0 g (range, 2.69-101.65 g), and the av duration of PRO treatment was 75 mo. Two of the 11 monkeys with malignant neoplasms received PRO by ip injections, 7 were given the drug both sc and po, and 2 were treated both ip and po. The toxic effects of PRO included vomiting, myelosuppression, and testicular atrophy, with complete aplasia of the germinal epithelium. (51 refs)

- 78-5149 **Induction of Sister Chromatid Exchanges In Vivo in Bone Marrow Cells of AKR Mice.** (Fre) Renault, G. (Institut de Recherches scientifiques sur le Cancer [C.N.R.S.], B.P. no. 8, 94800 Villejuif, France); Pot-Deprun, J.; Chouroulinkov, I. *C R Acad Sci [D] (Paris)* 286(11): 887-890; 1978.

Four drugs known for their mutagenic activity were tested in vivo in 3-to 5-mo-old AKR mice for their ability to induce sister chromatid exchanges (SCE). Mice were treated with 5-bromodeoxyuridine for 7 days (40 mg/kg, ip), the test drug was given ip on day 9, colchicine (0.05 mg ip) was injected on day 18, and mice were killed on day 21. SCE were counted in samples of bone marrow obtained from the femurs. In order of decreasing activity, the drugs tested were cyclophosphamide (5-25 mg/kg), procarbazine (25-200 mg/kg), methylmethane sulfonate (25-100 mg/kg), and diethylnitrosamine (10-100 mg/kg). It is concluded that induction of SCE in vivo is an objective measure of the action of chemical agents on the animal. (8 refs)

- 78-5150 **Potentially Carcinogenic Analogues of the Carcinogen Hexamethylphosphoramide: Evaluation In Vitro.** (Eng) Ashby, J. (Central Toxicology Lab., Imperial Chemical Industries Ltd., Alderley Park, Macclesfield, Cheshire, England); Styles, J. A.; Paton, D. *Br J Cancer* 38(3): 418-427; 1978.

Several analogs of hexamethylphosphoramide were assayed for carcinogenicity using baby hamster kidney cells (BHK 21/C13) and the cell transformation assay of Styles under two metabolic activation conditions: an S-9 microsomal mix at a standard ratio of S-9 fraction to cofactors (1:9 10 μ /tube) or the mix at a modified ratio (1:1; 50 μ l). Each experiment was repeated twice. Hexamethylphosphoramide, hexamethylphosphorus triamide, hexamethylphosphorothioic triamide, tripiperidinophosphine oxide, phosphorothioic trimorpholide, and diethoxymorpholinophosphine oxide gave positive responses in this assay. Different transformation frequencies were obtained with the two S-9 mixes, suggesting that a small but widely separated range of S-9 mixes be used in experimental situations in which an initially derived negative response is not compatible with what is already known about, or expected from, a chemical or class of chemicals. It appears that the broad structural requirements for carcinogenicity observed for nitrosamine carcinogens also apply to phosphoramidate carcinogens. (18 refs.)

- 78-5151 **Acute Myelogenous Leukemia Following Immunosuppressive Therapy for Rheumatoid Arthritis.** (Eng) Kapadia, S. B. (Dept. Pathology, Presbyterian-Univ. Hosp., Pittsburgh, PA, 15213); Kaplan, S. S. *Am J Clin Pathol* 70(2): 301-302; 1978.

The development of acute myelogenous leukemia in a 51-yr-old woman following long-term cyclophosphamide therapy for rheumatoid arthritis is reported. During her 4-yr history of cyclophosphamide ingestion, an av daily dose of 25 mg had been administered. The development of pancytopenia resulted in a discontinuation of the drug; however, 4 mo later the pancytopenia had progressed to acute myelogenous leukemia. Cytogenetic study of the marrow cells re-

realed a missing B-group chromosome in all 20 cells studied and one to three extra C-group chromosomes in 12/20 cells. These findings suggest a possible leukemogenic role of cyclophosphamide in this case; caution should be exercised in treating nonfatal diseases with antitumor drugs. (19 refs.)

78-5152 Use of Human-Liver Microsomes from Kidney-Transplant Donors for the Induction of Chromatid Aberrations and Sister-Chromatid Exchanges by Means of Pre-carcinogens in Chinese Hamster Cells In Vitro. (Eng) Thust, R. (Res. Group Preventive Oncology, Inst. Pathology, Medical Acad. Erfurt, Nordhauser Strasse 74, D-99089 Erfurt, E. Germany); Warzok, R.; Grund, E.; Menzel, J. *Mutat Res* 51(3): 397-402; 1978.

The induction of chromatid aberrations and sister chromatid exchanges (SCE's) in Chinese hamster fibroblasts by cyclophosphamide (CP) and dimethylnitrosamine (DMN) metabolized by human liver microsomes in combination with an ADPH-generating system was studied. The liver microsomes were obtained from two kidney transplant donors. The liver microsomes from one subject, who also suffered from Gaucher's disease, activated CP and DMN to induce chromatid aberrations and a slight increase in SCE's. The microsomes from this subject metabolized DMN at the same level as uninduced rat liver microsomes. The microsomes from the second subject failed to activate CP, but they did activate DMN, although to a somewhat lesser extent than rat-liver microsomes or the microsomes from the first subject. Based on absolute concentration, DMN showed slight SCE induction compared with CP. The use of tissues from renal transplant donors can provide a practicable means for studying the metabolism of carcinogens and for detecting active metabolites by means of genetic and cytogenetic assays. (16 refs.)

78-5153 Acute Leukemia Following Immunosuppressive Therapy for Disseminated Sclerosis and for Behcet's Syndrome. (Fre) Mougeot-Martin, M. (Clinique Medicale "A", Hopital Saint-Antoine, 184, rue du Faubourg Saint-Antoine, F 75571 Paris Cedex 12, France); Krulik, M.; Larousseau, J. L.; Audebert, A. A.; Chaouat, Y.; Debray, J. *Ann Med Interne (Paris)* 129(3): 175-180; 1978.

The development of acute myelocytic leukemia (AML) in two patients treated with immunosuppressive drugs is reported, and the literature is reviewed. Evidence of AML was noted in the first patient, a 52-yr-old woman, after 49 mo of treatment with cyclophosphamide (100 mg/day, then 50 mg/day; total dose 81 g) for Behcet's syndrome. The second patient, a 47-yr-old woman, developed anemia and leukemia with a high proportion of myeloblasts in the bone marrow after 2 yr of therapy with chlorambucil (6 mg/day) for multiple sclerosis. The latent period between the end of therapy and the appearance of AML was 2 yr. A total of 33 cases of acute leukemia following immunosuppressive therapy with

chlorambucil, cyclophosphamide, azathioprine, melphalan, methotrexate, and busulphan are summarized. There were 16 men and 17 women aged 21-71 yr (av age 50.8 yr). The patients were treated for a variety of diseases, but rheumatoid arthritis was the most frequent (13/33). Except for two patients treated 4.5 and 6 mo, respectively, duration of treatment was 12 mo or more (av 37.9 mo). The time between the end of treatment and the leukemia was 14.3 mo on the av. There was only one lymphocytic leukemia; three were undifferentiated, and the rest were myelocytic. Possible mechanisms of action of leukemia induction by immunosuppressive therapy are discussed. (41 refs)

78-5154 Sister Chromatid Exchanges Induced by Cyclophosphamide in V79 Cells Cultured in Diffusion Chambers in Mice. (Eng) Furukawa, M. (Dept. Experimental Biology, Roswell Park Memorial Inst., 666 Elm St., Buffalo, NY, 14263); Huang, C. C. *Mutat Res* 57(2): 233-239; 1978.

Line V79 Chinese hamster cells, cultured in diffusion chambers (DC) and implanted into mice, were tested as a potential assay system for mutagenicity. An exponential growth was observed from days 2-4 after implantation. Max growth was reached on day 6. After that, cell growth and viable cell counts decreased. Three days after implantation of DC with V79 cells, the hosts received six hourly injections of 0.2 ml 5-bromodeoxyuridine (BUdR) soln at concentrations of 0.125 to 1.0×10^{-2} M. DC were removed and analyzed for chromosome and sister chromatid exchanges (SCE) 24 hr after the first BUdR injection. The frequency of metaphases with differentially stained chromatids and with aberrations and the number of SCE per cell increased with BUdR dose. The frequency of metaphases with differentially stained chromatids was also positively correlated with the duration of BUdR exposure or the number of hourly injections of BUdR. The effects of cyclophosphamide (CY) were also studied. Injections of 2.5, 5, 10, and 15 μ g/g into hosts carrying V79 cells in DC increased in the number of SCE per cell in a linear manner. There was no increase of chromosome aberrations compared with controls. This study indicates that V79 cells cultured in DC in mice may prove to be a sensitive test system for screening potential mutagens or carcinogens, especially those that require metabolic activation. (26 refs)

78-5155 Cytogenetics in Short-Term Tests (Meeting Abstract). (Eng) Hughes, L. O. (Pharmaceuticals Div., Imperial Chemical Industries, Ltd., Macclesfield, Cheshire, England); Topham, J. C. *Mutat Res* 53(2): 203; 1978. (no refs)

78-5156 The Nucleus Anomaly Test as a Sensitive Method of Detecting Mutagenic Effects on Somatic Cells (Meeting Abstract). (Eng) Langauer, M. (Res. Dept.,

Pharmaceuticals Div., CIBA-GEIGY Ltd., Basel, Switzerland); Muller, D. *Mutat Res* 53(2): 216-217; 1978. (2 refs)

78-5157 Mutagenic Effects of Different Mutagens on Human Chromosomes In Vitro and In Vivo as Detected by Conventional and 'Harlequin' Methods (Meeting Abstract). (Eng) Kucerovala, M. (Pediatric Dept., Postgraduate Medical Inst., Prague, Czechoslovakia); Polivkova, Z. *Mutat Res* 53(2): 215; 1978. (no refs)

78-5158 Comparative Mutagenicity of Alkylsulfate and Alkanesulfonate Derivatives in Chinese Hamster Ovary Cells. (Eng) Couch, D. B. (Chemical Industry Inst. Toxicology, Research Triangle Park, NC, 27709); Forbes, N. L.; Hsie, A. W. *Mutat Res* 57(2): 217-224; 1978.

The mutagenicity and cytotoxicity of several alkyl sulfates and alkane sulfonates were quantitated using the Chinese hamster ovary/hypoxanthine-guanine phosphoribosyltransferase system. Dose-response relationships were examined for two alkyl sulfates, dimethyl sulfate (DMS) and diethyl sulfate (DES), and three alkyl alkanesulfonates, methyl methanesulfonate (MMS) ethyl methanesulfonate (EMS), and isopropyl methanesulfonate (iPMS). Cytotoxicity decreased with the size of the alkyl group. DMS was more toxic than DES, and MMS was more toxic than EMS and iPMS. All agents gave a linear dose response of mutation induction: DMS was more mutagenic than DES and MMS was more mutagenic than EMS and iPMS, based on mutants induced per unit of mutagen concentration. When comparisons were made at 10% survival, however, the following relative mutagenic potency was observed; DES > DMS; EMS > MMS > iPMS. It does not appear that mutagenic potency, as determined in this study, is clearly associated with S(n)1 (unimolecular nucleophilic substitution) reactivity. iPMS has been reported to have the greatest S(n)1 reactivity, but it was less mutagenic than DES or EMS. (21 refs)

78-5159 Promotion of Azoxymethane-induced Colonic Neoplasia by Resection of the Proximal Small Bowel. (Eng) Williamson, R. C. (Dept. Surgery, Bristol Royal Infirmary, Bristol, England); Bauer, F. L.; Oscarson, J. E.; Ross, J. S.; Malt, R. A. *Cancer Res* 38(10): 3212-3217; 1978.

Potential enhancement of intestinal neoplasia by compensatory mucosal hyperplasia was tested in male Fischer rats subjected to 50% proximal small bowel resection (PSBR) 10 days after the last of 16 weekly injections of 10 mg azoxymethane (AZM). AZM alone increased the jejunal contents of RNA and DNA each by 26% at 17-18 wk before there was macroscopic evidence of neoplasia. Three months after PSBR alone, ileal hyperplasia was characterized by increased amounts of RNA (42%-76%) and DNA (68%-95%), taller

villi, deeper crypts, and luminal dilation; however, the colon showed only patchy hyperplasia. When the combined effects of AZM and PSBR were observed 26-30 wk after the first injection, rats with PSBR had an increased number of colonic tumors/animal (2.9 vs 1.6 for controls). Despite the intense ileal hyperplasia produced by PSBR, ileal neoplasia did not occur. Thus, enhanced colonic carcinogenesis was only observed following sequential exposure of the mucosa to the carcinogen and the promoting factors. (43 refs.)

78-5160 A Study of the Mutagenicity of Anesthetics and Their Metabolites. (Eng) Waskell, L. (Dept. Biochemistry, Univ. California, Berkeley, CA, 94720). *Mutat Res* 57(2): 141-153; 1978.

The commonly used volatile anesthetics, several of their metabolites, and drugs frequently used by the anesthesiologist were screened for mutagenicity in the Ames *Salmonella typhimurium*/rat liver microsomal assay system. Chloral hydrate was weakly mutagenic. Other compounds tested, including halothane (1,1,1-trifluoro-2-bromo-2-chloroethane), isoflurane (1-chloro-2,2,2-trifluoroethyl difluoromethyl ether), methoxyflurane (2,2-dichloro-1,1-difluoroethyl methyl ether), diazepam (7-chloro-1,3-dihydro-1,1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one), and chlordiazepoxide (7-chloro-2-methylamino-5-phenyl-3H-1,4-benzodiazepine) were not mutagenic. Non volatile compounds were tested for their ability to inhibit growth of bacterial strains with decreased capacity to repair damaged DNA. None of the compounds tested inhibited the growth of DNA repair-deficient strains relative to a strain with normal DNA repair. Halothane and trileone (1,1,2-trichloroethylene) were tested for direct interaction with DNA with negative results. These results support the hypothesis that the inhalation anesthetics, which are chemically stable molecules of low toxicity, are unlikely to react covalently directly with DNA. (49 refs)

78-5161 Metabolism of Methylated Analogs of Aldrin and Dieldrin by Liver Preparations from Male and Female Rat. (Eng) Hassall, K. A. (Dept. Physiology and Biochemistry, Univ. Reading, Reading, England); Dionyssiou-Asteriou, A.; Manning, D. *Pestic Biochem Physiol* 8(3): 287-301; 1978.

Aldrin and dieldrin analogs whose molecules lack a methylene bridge but have two methyl groups attached to olefinic or oxirane ring carbon atoms were synthesized, and their metabolism was studied using NADPH-supplemented 12,000 g x 30-min supernates of Wistar and CD strain rat livers. The dimethylated olefin was not metabolized to its epoxide but was converted by rat liver supernates of both sexes to a labile hydroxylated metabolite. Exogenous NADPH increased the rate at which the olefin was metabolized, but concentrations of NADPH that increased the yield of the labile metabolite

When added to female rat liver preparations decreased the yield when added to those of males. This metabolite had a mass spectrum that was consistent with its being a hydroxymethyl derivative. The dimethylated epoxide was metabolized more slowly than the olefin, and a notable sex difference in the route of metabolism was evident. With supernate from the livers of female rats, the only metabolite detected by gas-liquid chromatography had a mass spectrum consistent with its being a ring-hydroxylated epoxide. Male rat liver supernate formed small quantities of the same compound, but the major metabolite had a mass spectrum indicative of a hydroxymethyl epoxide. No hydrolysis of the oxirane ring of the epoxide was detected. (17 refs)

78-5162 Study of the Mutagenic Potential of Captan by the Heritable Translocation Test in Mice (Feeding Abstract). (Eng) Jorgenson, T. A. (Stanford Res. Inst., Menlo Park, CA); Rushbrook, C. J.; Newell, G. W.; Waters, M. *Mutat Res* 53(2): 206; 1978. (no refs)

78-5163 Effect of Fungicide Folpet on Growth and Chromosomes of Human Lymphoid Cell Lines. (Eng) Sirianni, S. R. (Dept. Experimental Biology, Roswell Park Memorial Inst., Buffalo, NY, 14263); Huang, C. C. *Can Genet Cytol* 20(2): 193-197; 1978.

The cytotoxic and cytogenetic effects of Folpet (N-tri-chloromethylthiophthalimide; a fungicide used in paints) on normal (Jeff) and Burkitt (Raji) human lymphoid cell lines were determined. Jeff cells were continuously treated with Folpet concentrations of 0.5, 1, 2, and 4 $\mu\text{g}/\text{ml}$ media and Raji cells were pulse-treated for 15 min with 2.5, 5, 10, and 15 $\mu\text{g}/\text{ml}$. Various times after treatment, cell viability counts and chromosome analyses were made. Severe cell growth inhibition and an increase in chromosome aberrations were observed in both cell lines. Two types of chromosome aberrations were induced. Type-B aberrations appeared in cultures treated with lower doses, and they were characterized by chromosome or chromatid gaps and breaks; dicentric and exchanges were also noted infrequently. In the later stages of the experiment, the cell growth rate and number of aberrations returned to control levels. Type-B chromosome aberrations appeared in early samples of cultures treated with higher doses of the compound. The aberrations included chromosome stickiness, despiralization, and pulverization. The cultures died off within 2 days. Pulse treatment of Jeff cultures with the same doses used to treat Raji cells indicated that the former were more sensitive to Folpet than the latter. Based on the clastogenic effect of Folpet in this experiment and the close correlation between clastogens and mutagens, it can be inferred that the compound is mutagenic. The environmental risk of Folpet as a paint additive needs further study. (13 refs.)

78-5164 Determination of Carcinogenicity of O,O-Dimethyl(1-hydroxy-2,2,2-trichloroethyl)phosphonate (Trichlorfon) in Mice by Oral, Intraperitoneal, and Dermal Application. (Ger) Teichmann, B. (Zentralinstitut für Krebsforschung, Akademie der Wissenschaften der DDR, Lindenberger Weg 80, DDR-1115 Berlin-Buch, E. Germany); Hauschild, F. *Arch Geschwulstforsch* 48(4): 301-307; 1978.

The carcinogenicity of O,O-dimethyl(1-hydroxy-2,2,2-trichloroethyl)phosphonate (Trichlorfon; TCP) was investigated in male and female AB/Jena mice. The mice received TCP by po (esophageal-gastric intubation), ip, or dermal administration. In po experiments, male mice received a max of 157.5 mg and females 154 mg over 73 wk. With ip administration, males received a max of 160.8 mg and females 149.7 mg over 73 wk. With dermal application, both sexes received a max of 375 mg TCP over 75 wk. The experiment was terminated at 80 wk. There was no statistically significant difference between the total tumor incidence in TCP-treated animals and controls. In addition, no significant difference in tumor incidence or type was observed among the mice receiving TCP by the different routes. (9 refs.)

78-5165 Effects of Feeding Lindane to Dogs for Periods of up to 2 Years. (Eng) Rivett, K. F. (Huntingdon Res. Centre, Huntingdon, England). *Toxicology* 9(3): 273-289; 1978.

The effect of chronic exposure to lindane on dogs were studied by feeding the pesticide to male and female beagles at levels of 25, 50, or 100 ppm for 104 wk or 200 ppm for 32 wk. In a preliminary study in which the same levels were administered for 7 wk, the only changes were decreases in body wt at 100 and 200 ppm and increased liver wt at 200 ppm. In animals given 100 ppm lindane for 104 wk, the only change in the electroencephalogram was a superimposition of slightly irregular slow waves on the normal sleep pattern in two animals. Body wt gain, appetite, urinalysis, ophthalmological and hematological parameters, and organ wt were unaffected. The only biochemical change was a significant elevation in serum alkaline phosphatase (SAP) levels in this group; the livers in this group were also unusually dark. In the 200-ppm group, high-voltage slow-wave activity, which was possibly indicative of a nonspecific neuronal irritation, was observed. SAP levels were also elevated in this group, and the livers were dark, friable, and slightly enlarged. The negative findings in the 25- and 50-ppm groups are consistent with a "no-effect level" of 1.25 mg/kg in dogs that is comparable with that in rats and consistent with the acceptable daily intake of 0.0125 mg/kg proposed for humans. (23 refs)

78-5166 Children Absorb Tris-BP Flame Retardant from Sleepwear: Urine Contains the Mutagenic Metabolite, 2,3-Dibromopropanol. (Eng) Blum, A. (Dept. Bi-

ochemistry, Univ. California, Berkeley, CA, 94720); Gold, M. D.; Ames, B. N.; Kenyon, C.; Jones, F. R.; Hett, E. A.; Dougherty, R. C.; Horning, E. C.; Dzidic, I.; Carroll, D. I.; Stillwell, R. N.; Thenot, J. P. *Science* 201(4360): 1020-1023; 1978.

Urine samples from children who were wearing or who had worn sleepwear treated with the flame retardant tris(2,3-dibromopropyl)phosphate (tris-BP) were examined for 2,3-dibromopropanol (DBP), a mutagenic metabolite of tris-BP. Tris-BP is also a mutagen and it causes sterility and cancer in animals. Preliminary studies were made of morning urine samples from a 7-yr-old girl who wore repeatedly washed sleepwear that may have been treated with tris-BP on days 1 and 2 and 8-12 and new tris-BP-treated pajamas on days 3-7. On days 1 and 2, her urine contained 0.4 nanogram (ng)/ml DBP; considerably more DBP (up to 29 ng/ml on day 4) was found in her urine both during and after the wearing of the new pajamas. Urine samples were collected from 10 other children and 1 adult about 5 mo after the sale of new tris-BP-treated pajamas was banned. No DBP was found in the urine of the one child and one adult who had never worn flame-retardant sleepwear. Eight children were wearing well-washed tris-BP-treated sleepwear: seven had urinary DBP levels of approx 0.5 ng/ml and one child had a level of 5 ng/ml. A trace of DBP was found in the urine of one child who reportedly stopped wearing tris-BP-treated sleepwear 6 mo previously. It was estimated that the amount of tris-BP absorbed could be as much as 15 times as high as the amount of DBP found in urine. Absorption of tris-BP from old treated sleepware could pose a continuing problem, since half the children's sleepware sold between July 1973 and April 1977 was treated with this chemical. (25 refs.)

78-5167 Mouse Skin Carcinogenicity Tests of the Flame Retardants Tris(2,3-dibromopropyl)phosphate, Tetrakis(hydroxymethyl)phosphonium Chloride, and Polyvinyl Bromide. (Eng) Van Duuren, B. L. (Lab. Organic Chemistry and Carcinogenesis, Inst. Environmental Medicine, New York, NY, 10016); Loewengart, G.; Seidman, I.; Smith, A. C.; Melchionne, S. *Cancer Res* 38(10): 3236-3240; 1978.

The flame retardants tris(2,3-dibromopropyl)phosphate (tris), tetrakis(hydroxymethyl)phosphonium chloride (THPC), and polyvinyl bromide (PVB) were tested for carcinogenic activity by thrice weekly skin application in female Icr/Ha Swiss mice for 420-496 days. Tris, at doses of 30 or 10 mg/application (30 mice/group), induced benign and malignant tumors of the skin, forestomach, and oral cavity (tongue and gingiva) in a statistically significant number of mice ($p < 0.0005$). A statistically significant incidence of papillary tumors of the lung was observed at both doses ($p < 0.0005$). One carcinoma of the liver was observed at both doses, and the higher dose also resulted in a tubular adenocarcinoma of the kidney in one mouse. THPC (2 mg/application, 60 mice) and PVB (0.1 ml latex suspension/application, 30 mice) were inactive. PVB was also injected sc (approx 23 mg) into the mice once weekly for 48 wk; the mice were observed for an additional 12 wk. Liposarcomas were induced in 19/30 mice; the tumors were ascribed to physical carcinogenesis because the aqueous suspension contained 90% solids. The internal tumors observed in the PVB-treated animals were not significantly different from those observed in controls. (28 refs.)

PHYSICAL CARCINOGENESIS

5168 **Host-Cell Reactivation of UV-irradiated and Chemically-treated Herpes Simplex Virus-1 by Xeroderma Pigmentosum, XP Heterozygotes and Normal Human Fibroblasts.** (Eng) Selsky, C. A. (Lab. Radiobiology, Dept. Physiology, Harvard Sch. Public Health, Boston, MA, 02115); Greer, S. *Mutat Res* 50(3): 395-405; 1978.

to further characterize the xeroderma pigmentosum (XP) phenotype, the survival of UV-irradiated herpes simplex virus type 1 (HSV-1) strain MP was compared with the survival of N-acetoxy-2-acetylaminofluorene (NAO-AAF)-treated virus in normal human skin fibroblasts, XP cells from the different genetic complementation groups, an XP variant, and two presumed XP heterozygotes. In addition, the survival of NAO-AAF-treated HSV-1 MP virus was compared with that of nitrogen mustard-treated virus in complementation group D XP cells. The relative ability of XP cells from the different complementation groups to reactivate UV-irradiated virus was in the order $D < C < B$, and the relative reactivation ability of these cells for NAO-AAF-treated virus was in the order $D \approx \text{equal} < A < B < C$. The survival of UV- and NAO-AAF-treated HSV-1 MP in the XP variant was similar to that of treated virus in normal fibroblasts. Caffeine in the postinfection assay medium had no effect on the results. Relative to normal skin fibroblast strains, groups A and B XP cell strains showed efficient reactivation of virus treated with up to 100 μg NAO-AAF/ml. The inactivation kinetics of the treated virus in the XP heterozygotes were the same as those in normal cell strains. UV and NAO-AAF treatment resulted in reduced virus survival in the XP cells relative to normal cells, whereas the survival of nitrogen mustard-treated virus was similar in group D XP cells and normal fibroblasts. The data indicate that UV- and NAO-AAF-treated virus is repaired by the same pathway or by different pathways that share a common intermediate step and that the excision defect of XP plays little if any role in the reactivation of nitrogen mustard-treated virus. (24 refs)

5169 **Xeroderma Pigmentosum: Damage to DNA Is Involved in Carcinogenesis.** (Eng) Setlow, R. B. (Biology Dept., Brookhaven Natl. Lab., Upton, NY, 11973); Friedberg, F. E.; Grist, E. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. Cold Spring Harbor: Cold Spring Harbor Laboratory; Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 889-902; 1977.

The role of UV in skin cancer was studied with the use of human fibroblasts from normal individuals, and from patients with xeroderma pigmentosum (XP). Cells were treated with acetoxyacetylaminofluorene (AAAF) and/or UV. DNA was isolated and then treated with an endonuclease from *Micrococcus luteus*, which recognizes UV- but not AAAF-induced DNA damage. AAAF treatment did not inhibit the removal of sites sensitive to *M. luteus* endonuclease, as would be expected if AAAF damage truly mimicked UV damage. It is concluded that the enzymatic steps involved in the repair of the two types of damage are not identical and that the defect in XP cells controls more than one function. The defect may be at the level of formation or stability of the presumptive repair complex involved in excision repair. In a study of postreplication repair, cells were preilluminated with an inducing exposure of 2.5 joules (J)/ m^2 of 254-nanometer UV radiation. After 2.8 or 14 hr, the cells were treated with a second exposure of 5.0 J/m^2 and, 15 min later, the DNA was pulse-labeled with ^3H -thymidine. Postreplication repair was enhanced in all the cells tested, but it was much larger in XP variant cells than in normal or conventional XP cells. In cells from XP-variant individuals, excision is normal, but postreplication repair is slow. It is inferred that in these cells the probability of a carcinogenic event is high not because many dimers remain, but because replication past a dimer is more error-prone than the process in normal cells. (41 refs)

78-5170 **A Mutagen Screening Test for UV-Mimetic Chemicals Based on Differential Killing of Human Cell Lines (Meeting Abstract).** (Eng) Burrell, A. D. (General Products Div., IBM Corporation, San Jose, CA, 95193); Anderson, J. J. *Mutat Res* 53(2): 161; 1978. (2 refs)

78-5171 **Mutational Synergism of Nitrous Acid and Ultraviolet Light (Meeting Abstract).** (Eng) Kubitschek, H. E. (Div. Biological and Medical Res., Argonne Natl. Lab., Argonne, IL). *Mutat Res* 53(2): 214-215; 1978. (no refs)

78-5172 **Excision Repair of Ultraviolet Damage in Mammalian Cells: Evidence for Two Steps in the Ex-**

cision of Pyrimidine Dimers. (Eng) Williams, J. I. (Dept. Pathology, Stanford Univ. Medical Center, Stanford, CA, 94305); Cleaver, J. E. *Biophys J* 22(2): 265-279; 1978.

The post-UV irradiation incidence of pyrimidine dimer formation and the kinetics of DNA repair in African green monkey kidney CV-1 cells were studied by measuring survival; T4 endonuclease V-sensitive sites; the fraction of pyrimidine dimers in acid-insoluble DNA, as determined by thin layer chromatography (TLC); and repair replication. The CV-1 cells demonstrated a survival curve with extrapolation number $n = 7.8$ and D_0 (the single dose of radiation necessary to kill 63% of the cells using the straight part of the cell survival curve) of 2.5 joules/m². Pyrimidine dimers were lost from acid-insoluble DNA more slowly than endonuclease-sensitive sites were lost from or new bases were incorporated into high-mol-wt DNA during the course of repair. Addition of ³H-thymidine to the cultures of x-irradiation (2 or 10 kilorads) 24 hr before UV irradiation had no effect on repair replication induced by UV (25 joules/m²). These findings suggest that pyrimidine dimer excision measurements by TLC are probably not affected by radiation from high levels of incorporated radionuclides. The endonuclease-sensitive site and TLC measurements can be reconciled by the assumption that pyrimidine dimers are excised from high mol wt DNA in acid-insoluble oligonucleotides that are slowly degraded to acid-soluble fragments. Since the CV-1 cell line is often used as a host for simian virus 40 or herpesvirus studies, these findings provide biochemical information valuable in the study of host cell reactivation and viral mutagenesis. (57 refs)

78-5173 Mechanisms of Depressed Reactivity to Dinitrochlorobenzene and Ultraviolet-induced Tumors During Ultraviolet Carcinogenesis in BALB/c Mice. (Eng) Jessup, J. M. (Dept. Surgery, Univ. Texas Medical Sch. at Houston, Houston, TX, 77025); Hanna, N.; Palaszynski, E.; Kripke, M. L. *Cell Immunol* 38(1): 105-115; 1978.

Experiments were conducted to determine (1) whether UV radiation renders BALB/cAnN female mice susceptible to challenge with a highly antigenic, syngeneic, UV-induced fibrosarcoma (2C16) and (2) whether chronic UV radiation induces a transient, reversible depression of dinitrochlorobenzene (DNCB) hypersensitivity in these mice. Mice were irradiated (av dose rate, 2.8 joules/m²/sec) for 1 hr, 3x/wk, for 1-4 mo. Upon transplantation, fibrosarcoma 2C16 grew progressively in 9/10 mice irradiated for 1 mo, 7/10 irradiated for 2 mo, and 10/10 irradiated for 3 mo. The lack of tumor rejection in the irradiated mice could be transferred with the spleen cells to lethally x-irradiated syngeneic recipients. Groups of UV-treated and untreated mice were sensitized with DNCB at monthly intervals after the initial UV treatment, and their responses were compared with those of unsensitized mice in a footpad assay. UV-treated mice failed to exhibit a primary immune response to DNCB for the first 2 mo of irradiation. However, mice irradiated for 3 or 4 mo

could be sensitized, and they exhibited the same levels of footpad reactivity as sensitized controls. The two immunologic dysfunctions demonstrated are mediated by different mechanisms, even though both are induced by UV radiation. The inability of UV-treated mice to reject UV-induced tumors seems to be due, at least in part, to the development of specific suppressor cells, probably, T lymphocytes. The temporary inability of UV-treated mice to respond to DNCB seems to be due to a block in antigen processing or presentation, since DNCB-reactive lymphocytes can be demonstrated in these animals by passive transfer studies. (14 refs)

78-5174 Ultraviolet Carcinogenesis in T-Cell-depleted Mice. (Eng) Norbury, K. C. (Merck Inst. Therapeutic Res., West Point, PA, 19486); Kripke, M. L. *J Natl Cancer Inst* 61(3): 917-921; 1978.

The induction of skin tumors by chronic UV irradiation was compared in normal C3H/HeN(MTV-) mice, T-cell-depleted mice, and T-cell-depleted mice reconstituted with thymus grafts. The T-cell-depleted mice were thymectomized young adults, lethally x-irradiated, and repopulated with neonatal liver cells. All mice were irradiated (dose rate, 2 joules/m²/sec, wavelength range 280-320 nanometers) 60 min 3x/wk for 3 wk, then 60 min 3x/wk until tumor development. Skin tumors appeared earlier in the immunosuppressed T-cell-depleted mice (50% tumor incidence, 30 wk) than in the normal mice (35 wk). However, tumors developed more rapidly in the group that was T-cell-depleted and immunologically restored with thymus grafts (24 wk). In the latter group, most of the tumors were squamous cell carcinomas, whereas in the normal mice, fibrosarcomas predominated. These experiments illustrate the limitations of this approach to studying the role of the immune response in carcinogenesis. (26 refs.)

78-5175 Ultraviolet Radiation-induced Chromosomal Abnormalities in Fetal Fibroblasts from New Zealand Black Mice. (Eng) Reddy, A. L. (Medical Genetics Section, Medical Service, Veterans Admin. Hosp., 4435 Bonifant Ave. S., Seattle, WA, 98108); Fialkow, P. J.; Salo, J. *Science* 201(4359): 920-922; 1978.

Cultured fetal fibroblasts from New Zealand Black (NZB) and BALB/c mice were exposed to 8, 16, or 32 joules (J)/m² of UV radiation, and the incidence of chromosome (chr) breaks and rearrangements was determined. Chr abnormalities were found more often in untreated NZB cells than in BALB/c cells, but the difference was not significant. After irradiation with 8 and 16 J/mm², the number of aberrations did not increase in the BALB/c fibroblasts; in contrast, the numbers of chromatid and chr breaks and interchanges increased significantly in the NZB fibroblasts. There was a direct relation between the frequency of abnormal cells and radiation dose: at the lower doses, chr abnormalities were

increased only in NZB cells, but at 32 J/mm², both NZB and ALB/c cells exhibited a high incidence of abnormalities. At this dose, however, significantly more abnormal cells were found in the former (57%) than in the latter (26%). The number of abnormal cells was max at 24 hr after radiation and then decreased to untreated levels at 72 hr. This temporal decline may be due to reproductive death of the abnormal cells consequent to chr damage. Thus, fibroblasts from NZB mice manifest increased sensitivity to UV radiation. This instability is similar to that reported for cells from xeroderma pigmentosum patients, and it may be related to the aneuploid clones and lymphoreticular malignancy characteristic of adult NZB mice. Chr instability may favor the development of specific chr changes that, in turn, play a role in the evolution of neoplasia. (13 refs.)

78-5176 Fluorescent Light-induced DNA Crosslinkage and Chromatid Breaks in Mouse Cells in Culture. (Eng) Gantt, R. (Lab. RNA Tumor Viruses, NCI, Bethesda, MD, 20014); Parshad, R.; Ewig, R. A.; Sanford, K. L.; Jones, G. M.; Tarone, R. E.; Kohn, K. W. *Proc Natl Acad Sci USA* 75(8): 3809-3812; 1978.

The effect of cool-white fluorescent light (20 hr exposure to 5 W/m²) on the DNA and chromosomes of cultured mouse embryo and lung cells was studied. Compared with shielded cells, the exposed cultures showed a significant increase in the percentage of cells with chromatid breaks and in the number of chromatid breaks per cell. The breaks appeared to be density-dependent. The number of minute chromosomes and the number of chromatid exchanges per cell were significantly increased in the exposed cultures. The fluorescent light had no apparent effect on the frequency of metacentric chromosomes. In Kohn's alkaline elution assay, the DNA from exposed cells eluted more slowly than that from shielded cells, suggesting that the DNA in the exposed cells was larger mol wt and cross-linked. The estimated frequency of crosslinks was sufficient to account for the observed number of chromatid breaks. The data suggest that the DNA cross-linking and chromosome aberrations produced by fluorescent light may be related. (19 refs.)

78-5177 Effect of Radiation Dosage on the Synergism Between Radiation and Estrogen in the Production of Mammary Cancer in the Rat. (Eng) Segaloff, A. (Div. 5, Alton Ochsner Medical Foundation, 1516 Jefferson Highway, New Orleans, LA, 70121); Pettigrew, H. M. *Cancer Res* 38(10): 3445-3452; 1978.

The effect of radiation dosage on the synergism between radiation and estrogen in the production of mammary cancer was studied in A x C rats. Rats that had previously been oophorectomized were exposed to estrogen by implantation of 20-mg cholesterol pellets containing 25% diethylstilbestrol (DES). The left mammary chain of the rats was then exposed

to 50, 150, or 450 rads of x-radiation while the rats were anesthetized with urethan or sodium pentobarbital. Control animals were implanted with cholesterol pellets containing no DES or they were not implanted at all. Some of the animals died in the 24 hr following irradiation; these deaths were independent of the anesthetic used, but there was a positive dose-response curve for radiation. In terms of median survival time, the highest radiation doses produced the shortest survival times, and rats receiving estrogen also had the shortest survival times. Overall growth was also inhibited by both of the agents used. With radiation alone, there was an increase in benign tumors with a dose-response curve, but the increase in mammary cancers was not significantly proportionate to dose. There was no significant difference in tumor incidence in any of the glands in each mammary chain. The cholesterol carrier did not affect the synergism between estrogen and radiation for mammary carcinogenesis; however, synergism between radiation and DES was apparent. Malignant mammary tumors were relatively rare in groups that did not receive DES. The optimal expression of synergism was dependent on the amount of radiation given. (23 refs.)

78-5178 Second Malignancies of the Irradiated Area in Patients Treated for Uterine Cervix Cancer. (Eng) Czesnin, K. (Dept. Gynecologic Oncology, Inst. Oncology in Warsaw, P.O. Box 47, Warsaw 02-034, Poland); Wronkowski, Z. *Gynecol Oncol* 6(4): 309-315; 1978.

The incidence of second malignancies in the irradiated area was analyzed in 8,043 patients with carcinoma of the uterine cervix treated primarily by irradiation between 1948 and 1966 at the Institute of Oncology in Warsaw, Poland. Radiotherapy had been administered by the modified Paris technique (dose to point A = 5,500 rads, point B 1,500 rads) and irradiation from external fields (deep field dose of x-rays = 3,000 R; ⁶⁰Co = 3,500-4,500 R). Follow-up time ranged from 9 to 27 yr; 4,041 patients survived 5 yr. Second malignancies developed in the irradiated area of 28 women 18 mo-22 yr after primary treatment. The most frequent type of second neoplasm was cancer of the uterine corpus, followed by ovarian cancer, cancer of the vulva, cancer of the large bowel and rectum, and malignant neoplasms of the integument. Uterine corpus cancers occurred after the longest period of time (13 yr after therapy); cancer of the vulva and bladder and sarcoma of the uterine corpus occurred in a shorter period of time (6 yr after therapy). The incidence rates for uterine sarcoma and soft tissue malignant neoplasms in the irradiated area were significantly higher in the examined group than the rates expected for the whole population. No significant increase in incidence rates for other pelvic neoplasms was demonstrated. (22 refs.)

78-5179 Sebaceous Carcinoma of the Eyelid Following Radiation Therapy. (Eng) Lemos, L. B.

(Memorial Sloan-Kettering Cancer Center, New York, NY); Santa Cruz, D. J.; Baba, N. *Am J Surg Pathol* 2(3): 305-311; 1978.

A 12-yr-old boy developed a metastatic sebaceous carcinoma of the eyelid 11 yr after curative radiation therapy for bilateral retinoblastoma. The right eye had been enucleated at the age of 15 mo; the right orbit was then treated with 2,100 rads and the left eye with 1,475 rads. The gross and ultrastructural pathologic findings are presented. A review of the literature revealed only five reports of sebaceous carcinoma associated with previous irradiation. Three of the patients had been treated for benign disorders, and two for bilateral retinoblastoma. This small series of six cases suggests a cause-and-effect relationship between irradiation and sebaceous carcinoma. However, the possibility that genetic factors may predispose patients with bilateral retinoblastoma to develop sebaceous carcinoma cannot be excluded. (31 refs.)

78-5180 Postirradiation Fibrosarcoma Following Radical Mastectomy. (Eng) Iwasaki, K. (Dept. Surgery I, Kyushu Univ. Faculty Medicine, Fukuoka, Japan); Nagamitsu, S.; Tsuneyoshi, M. *Jpn J Surg* 8(1): 73-77; 1978.

A 52-yr-old woman underwent radical mastectomy for Stage I infiltrating breast carcinoma followed by radiation therapy (2,500 rads 20 days later). Over the next 11 yr, five fibrosarcomas arose from the mastectomy scar that were considered to be a result of the radiation therapy. (8 refs)

78-5181 Radioactive Fallout Effects of the Nagasaki Atomic Bomb. (Eng) Okajima, S. (Dept. Radiation Biophysics, Atomic Disease Inst., Nagasaki Univ. Sch. Medicine, Nagasaki, Japan); Takeshita, K.; Antoku, S.; Shiomoto, T.; Russell, W. J.; Fujita, S.; Yoshinaga, H.; Neriishi, S.; Kawamoto, S.; Norimura, T. *Health Phys* 34(6): 621-633; 1978.

Whole-body counting techniques, physical and laboratory examinations, and chromosome studies were used to determine retained radionuclide activity and late radiation effects in Nishiyama residents exposed to fallout from the Nagasaki atomic bomb. Soil, reservoir sediments, and crops from Nishiyama and comparison areas were examined for ^{137}Cs content by scintillation counting. Body ^{137}Cs concentrations were significantly higher among those exposed to the fallout than among those who were not exposed. However, the av annual exposure dose was low compared with the dose limit recommended by the ICRP, and no deleterious effects were evident. The body burden of ^{137}Cs in the Nishiyama residents was maintained by constant replenishment via ingestion of contaminated farm products. The ^{137}Cs content of the drinking water used by Nishiyama residents was essentially the same as that of the Nagasaki City and the Nishiyama and Urakami reservoirs. (32 refs.)

78-5182 X-ray Sensitivity of Diploid Fibroblasts from Patients with Hereditary or Sporadic Retinoblastoma. (Eng) Weichselbaum, R. R. (Laboratory Radiobiology, Dept. Physiology, Harvard Univ., Sch. Public Health, Boston, MA, 02115); Nove, J.; Little, J. B. *Proc Natl Acad Sci USA* 75(8): 3962-3964; 1978.

The x-ray sensitivity, expressed D_0 values, of diploid fibroblasts from six patients with hereditary retinoblastoma, seven patients with sporadic retinoblastoma, a retinoblastoma patient with a D-deletion of chromosome 13, and four normal controls was studied. Cultured fibroblast cells were exposed to an x-ray dose of 80/rads/min and fixed and stained 15 days later; colonies composed of more than 50 cells were scored as survivors. The mean D_0 (inverse of the slope) of the x-ray survival curves was significantly lower for the hereditary group than for the sporadic group. Cells from a patient belonging to a family with an extremely high incidence of malignant tumors appeared to be considerably more x-ray sensitive than those from other hereditary patients. Only one hereditary strain was more resistant than the most sensitive sporadic strain, the sensitivity of most of the sporadic strains being well within the normal range. The fibroblasts from the patient with the D-deletion were not significantly x-ray sensitive. The increased radiosensitivity of hereditary retinoblastoma fibroblasts could result from a defect in a molecular DNA repair process. (17 refs.)

78-5183 Liver Cell Proliferation and Failure of X Radiation to Produce Hepatomas in Rats. (Eng) Christov, K. (Acad. Medicine, Natl. Inst. Oncology, Sofia-56, Darvenitza, Bulgaria). *Radiat Res* 74(2): 378-381; 1978.

X-ray doses that are carcinogenic in mice failed to induce hepatomas in intact or partially hepatectomized male Wistar rats. In contrast, hepatoma incidence was increased in rats treated with dimethylnitrosamine during late G_1 . This difference could result from different mechanisms of chemical and radiation carcinogenesis in the liver. (11 refs)

78-5184 A Compartment Model of Plutonium Dynamics in a Deciduous Forest Ecosystem. (Eng) Garman, C. T. (Environmental Sciences Div., Oak Ridge National Lab., Oak Ridge, TN, 37830); Gardner, R. H.; Dahlman, C. *Health Phys* 34(6): 611-619; 1978.

An ecosystem model of plutonium (^{239}Pu and ^{240}Pu) behavior in a contaminated deciduous forest is described. The model was formulated as a linear compartment model with donor-controlled flows between compartments. It employs time-variant transfer coefficients. At steady states predicted by the models, the amount of Pu in the biota is $<0.25\%$ of the existing inventory in the soil. The soil would be the major

reservoir of Pu in the forest, and reciprocal exchanges of Pu between soil and litter or soil and tree roots would be the dominant factors affecting the Pu ecosystem distribution. However, the average predicted amount of Pu in each forest ecosystem, given a uniform soil contamination, could vary depending on site conditions and forest species composition. Present attempts to model Pu dynamics in terrestrial ecosystems are hampered by uncertainties in predictions arising from problems with system identification, quality of available data, and a lack of validation. The ability to develop reliable models of Pu transport in ecosystems will improve with more precise data from natural environments and a better understanding of the sources of variation in Pu data. (24 refs.)

78-5185 Late Effects of Inhaled $\text{Pu}(\text{NO}_3)_4$ in Rats (Meeting Abstract). (Eng) Ballou, J. E. (Battelle, Pacific Northwest Labs., Richland, WA, 99352); Dagle, G. E.; McDonald, K. E. *Health Phys* 33(6): 665-666; 1977. (no refs)

78-5186 Trauma and Brain Tumors: An Experimental Study (Meeting Abstract). (Eng) Morantz, R. A. (Univ. Kansas Medical Center, Kansas City, KS); Shain, W. *J Neuropathol Exp Neurol* 37(5): 662; 1978. (no refs)

VIRAL CARCINOGENESIS

- 78-5187 Genetic Analysis of the Defectiveness in Strain MC29 Avian Leukosis Virus.** (Eng) Bister, K. (Institut für Virologie, Zentrum für Hygiene, Universität Freiburg, 78 Freiburg, W. Germany); Vogt, P. K. *Virology* 88(2): 213-221; 1978.

The nature of the replication defect of avian myelocytomatosis virus (MC29) was analyzed in complementation experiments, in which the replicative genes (*gag*, *pol*, *env*) were tested genetically. Continuous nonproducer (NP) Japanese quail cell lines transformed by MC29 were superinfected with various helper viruses. Infectious MC29 pseudotypes were formed only with viruses related to the avian leukosis-sarcoma complex (Rous-associated virus type 1 or ring-necked pheasant virus), which demonstrated that its replication defect is not restricted to a lack of envelope glycoprotein. Reticuloendotheliosis virus, golden pheasant virus, or an amphotrophic murine leukemia virus (MuLV-1313) could not rescue MC29 from the NP cells, but they complemented the envelope-defective Bryan high-titer strain of Rous sarcoma virus (RSV-B). Analysis of the host range and viral interference patterns of MC29 rescued from NP clones indicated that both are determined by the envelope specificity of the activating helper virus. Co-cultivation or Sendai virus-induced fusion of MC29 NP cells with quail or White Leghorn C/BE chicken cells transformed by RSV-B did not result in complementation of the defects in either virus: fluorescent antibody staining did not detect virus-specific antigens at the surface of the MC29 NP cells. Temperature-sensitive mutants of RSV with lesions in the genes coding for RNA-directed DNA polymerase (*pol* gene) or nonglycosylated structural proteins (*gag* gene) did not acquire wild-type characteristics when grown in the NP cells. These experiments indicate that MC29 is defective in all three genes essential for retrovirus replication; ie, *gag*, *pol*, and *env*. These defects may be significant for the origin of the retroviruses and for their oncogenic activities. (48 refs.)

- 78-5188 In Vitro Synthesis of a Functional Avian Sarcoma Virus Transforming-Gene Product.** (Eng) Erikson, E. (Dept. Pathology, Univ. Colorado Medical Center, Denver, CO, 80262); Collett, M. S.; Erikson, R. L. *Nature* 274(5674): 919-921; 1978.

The in vitro product of the Schmidt-Ruppin avian sarcoma virus (ASV, subgroup D) *src* gene was tested for phosphotransferase activity. The 70S RNA was extracted from nondefective (nd) and transformation-defective (td) virions and then purified and fractionated. The RNA in fractions 12-15 was used to program messenger RNA-dependent reticulocyte lysates. The ³⁵S-methionine-labeled polypeptide

products were analyzed by gel electrophoresis, and the results indicated that p60src was translated only from nd RNA. No p60src was translated from td RNA, although this RNA stimulated protein synthesis 9-fold above the background synthesis without RNA, compared with 10-fold stimulation with nd RNA. Serum from rabbits bearing ASV-induced tumors (TBR serum) precipitated p60src and a small amount of Pr76 (product of ASV *gag* gene) from the products synthesized in response to nd RNA; in addition, a small amount of Pr76 was found in the immunoprecipitate of the products synthesized in response to td RNA. Only the products of translation of the RNA that immunoprecipitated with TBR serum yielded phosphorylated IgG. The translation of viral RNA in vitro resulting in the synthesis of an enzymatically functional p60src strongly indicates that the product of the *src* gene is a protein kinase. These and other results show that p60src is a product of the ASV transformation gene in vitro and in vivo and that its function is likely to result in the phosphorylation of cellular proteins. (12 refs.)

- 78-5189 Inhibition of Reverse Transcription of 70S and 35S Avian Myeloblastosis RNAs by Nonprimer tRNA's.** (Eng) Yamaura, I. (Sloan-Kettering Inst. Cancer Res., Walker Lab., Rye, NY, 10580); Cavalieri, L. F. *J. Virol.* 27(2): 300-306; 1978.

The kinetics of the reverse transcription of 70S and 35S RNAs of avian myeloblastosis virus (AMV) was studied in the presence and absence of various transfer RNA's (tRNA's). All tRNA's inhibited synthesis due to the binding of tRNA's to the polymerase. The reverse transcription kinetics of both 70S and 35S RNAs showed a complex inhibition pattern in the presence of nonprimer tRNA's obtained from 70S RNAs. The nonprimer tRNA's that bound to the reverse transcriptase decreased the affinity of the viral RNA for the enzyme. Thus, the binding of 70S or 35S RNA was decreased. In contrast, tRNA's from *Escherichia coli* and yeast showed classical noncompetitive inhibition: they bound reversibly and randomly and did not alter the affinity of viral RNA for the polymerase. The max rate of synthesis with 70S RNA as the template was less than that with 35S RNA, presumably because the former contains nonprimer tRNA's that can interact with the polymerase. (18 refs.)

- 78-5190 The Defectiveness of Mill Hill 2, a Carcinoma-inducing Avian Oncovirus.** (Eng) Hu, S. (Dept. Microbiology, Univ. Southern California, School of Medicine, 2011 Zonal Ave., Los Angeles, CA, 90033); Moricovici, C.; Vogt, P. K. *Virology* 89(1): 162-178; 1978.

The results of characterization experiments on the avian carcinoma virus Mill Hill 2 (MH2) are presented. MH2 transformed avian fibroblast and macrophage cultures derived from 10-day chick embryos, and it also transformed secondary cultures of macrophages from chick or Japanese quail embryo yolk sacs. It was defective in replication and dependent on a helper virus, MH2 associated-virus (MH2 AV), for the production of infectious progeny. MH2 AV contained both subgroup C and A envelope determinants; its helper functions were required by MH2 in the *env*, *pol*, and probably the *gag* genes. The defects of MH2 could be complemented only by helper viruses of the chicken leukosis group, further suggesting that MH2 is defective in *gag* or *pol*, or both. MH2 transformed-nonproducer cells synthesized an aberrant viral glycoprotein of 100,000 daltons. This MH2p100 carried antigenic determinants of the *gag* protein p27, but not of the *env* protein gp85. MH2p100 was not cleaved into smaller, functional proteins and was not glycosylated. (43 refs.)

78-5191 **Evidence for Splicing of Avian Sarcoma Virus 5'-terminal Genomic Sequences onto Viral-specific RNA in Infected Cells.** (Eng) Krzyzek, R. A. (Dept. Microbiology, Univ. Minnesota Medical Sch., Minneapolis, MN, 55455); Collett, M. S.; Lau, A. F.; Perdue, M. L.; Leis, P.; Faras, A. J. *Proc Natl Acad Sci USA* 75(3): 1284-1288; 1978.

The 5'-terminal nucleotide sequences of the avian sarcoma virus (ASV) genome are transcribed by reverse transcriptase in vitro into a DNA transcript that represents the entire distance (approx 100 nucleotides) between the transfer RNA^{Trp} (tRNA^{Trp}) primer molecule and the 5' terminus. These tRNA^{Trp} transcripts were used in hybridization reactions with SV-specific RNA from infected avian cells. Nucleotide sequences complementary to these transcripts were found on all the various size classes of viral messenger RNA (mRNA) identified. Similar hybridization results were obtained with specific DNA transcript complementary to viral genomic nucleotide sequences between the tRNA^{Trp} primer molecule and up to, but not including, the terminal redundant sequences (DNA₇₀), indicating that the observed hybridization of DNA₁₀₀ to all size classes of viral RNA in infected cells did not reflect hybridization of DNA₁₀₀ to the terminal redundant sequences at the 3' end of the viral genome. *Escherichia coli* RNase H hydrolysis of RNA-DNA hybrids consisting of genomic 35S RNA obtained from virus and DNA₁₀₀ transcripts indicated that viral genomic sequences complementary to these DNA transcripts were not present at sites distal to the ends of the RNA genome and, therefore, not adjacent to the corresponding gene sequences representing the various species of viral mRNA from infected cells. These findings suggest that the 5'-terminal genomic nucleotide sequences, or a portion thereof, are somehow added or "spliced" onto each ASV-specific mRNA species in infected cells either during or after transcription of proviral DNA. (35 refs)

78-5192 **The Effect of Infection and Transformation by Rous Sarcoma Virus upon Lysophosphoglyceride Biosynthesis in Chicken Embryo Fibroblasts.** (Eng) Edwards, R. G. (Div. Biochemical Virology and Membrane Res., Scheie Eye Inst., Univ. Pennsylvania Sch. Medicine, Philadelphia, PA, 19104); Blough, H. A. *Virology* 89(1): 314-317; 1978.

Lysophosphoglyceride biosynthesis was studied in secondary chicken embryo fibroblasts infected with the Schmidt-Ruppin strain (subgroup A) of Rous sarcoma virus (RSV) or its transformation-defective mutant (td₁₀₅) that lacks the *src* gene. The cells were infected with 5×10^6 focus-forming units of RSV or 5×10^5 to 5×10^6 infectious units of td₁₀₅ and harvested 10 days later. De novo biosynthesis lysophosphoglycerides was greatly depressed in cells transformed by the wild-type virus but not in those infected with the mutant. This depression may be due to a lack of the appropriate enzyme, rather than to an inhibitory factor, and thus represent an effect of expression of the *src* gene on lipid biosynthesis. (9 refs.)

78-5193 **Rous Sarcoma Viral Protein p15 Induces Intracellular Processing of Viral Precursor p76** (Meeting Abstract). (Eng) von der Helm, K. (ISREC, CH-1066 Epalinges, Switzerland); Rungger, D.; Wille, W.; Willecke, K. *Experientia* 34(7): 942; 1978. (1 ref)

78-5194 **A Transformation Defective Mutant of Rous Sarcoma Virus Inducing Chick Embryo Neuroretinal Cell Proliferation.** (Eng) Calothy, G. (Fondation Curie, Institut du Radium, 91405 Orsay, France); Poirier, F.; Dambrine, G.; Pessac, B. *Virology* 89(1): 75-84; 1978.

A transformation-defective (td) Rous sarcoma virus (RSV) mutant was isolated using induction of chick embryo neuroretinal (NR) cell multiplication as a genetic marker. The mutant was derived from a variant of Schmidt-Ruppin RSV subgroup D (SR-RSV-D) that was rescued by Sendai-mediated fusion transformed chick embryo fibroblasts (CEF) with simian virus 40-transformed Lesch-Nyhan cells infected with SR-RSV-D. The mutant, designated td PA-101, induced sustained multiplication of NR cells from 7-day-old chick embryos. However, the mitogenic and transforming properties of the mutant were permanently dissociated in NR cells. tdPA-101 was defective in transforming NR cells and CEF, and it did not produce sarcomas upon inoculation into white Leghorn chicks. Thus, this mutant represents a novel class of nonconditional RSV mutants that differ from other td viruses in retaining fully the capacity to induce NR cell proliferation. (31 refs.)

- 78-5195 Differential Expression of Rous Sarcoma Virus-specific Transformation Parameters in Enucleated Cells.** (Eng) Beug, H. (Biologisch-Medizinische Abteilung, Max-Planck-Institut für Virusforschung, 7400 Tübingen, W. Germany); Claviez, M.; Jockusch, B. M.; Graf, T. *Cell* 14(4): 843-856; 1978.

To obtain information about the primary site of action of the Rous sarcoma virus (RSV) *src* gene product(s), chicken embryo fibroblasts transformed with the Ta and ts68 mutants of RSV were enucleated and studied for their capacity to express reversibly the transformed phenotype in response to temperature changes. After shift to the permissive temperature (35 C), the cytoplasts acquired a transformed morphology and displayed characteristic ruffles and microvilli at their surface. Immunofluorescence revealed that they also lost their actin filament cables and exhibited characteristic changes in the pattern of cell-surface structures containing the large external transformation-sensitive (LETS) protein. Expression of all these transformation parameters was reversible after shiftback to the nonpermissive temperature (41 C). These results indicate that a whole set of changes characteristic of the transformed phenotype can be expressed independently of the cell nucleus. In contrast, ts mutant-infected cytoplasts were no longer able to respond to temperature shifts with changes in their hexose transport rate. Cytoplasts prepared from cells grown at 41 C retained their low rate of hexose uptake after shift to 35 C, but cytoplasts from cells grown at 35 C exhibited a high rate of hexose transport even after 10 hr of shift to 41 C. These findings agree with the hypothesis that the product of the RSV *src* gene represents a multifunctional protein that acts independently on nuclear and extranuclear sites. (51 refs.)

- 78-5196 Suppression and Enhancement of Mitogen Response in Chickens Infected with Marek's Disease Virus and the Herpesvirus of Turkeys.** (Eng) Lee, L. F. (Regional Poultry Res. Lab., Science and Education Admin.-Federal Res., U.S. Dept. Agriculture, East Lansing, MI, 48823); Sharma, J. M.; Nazerian, K.; Witter, R. L. *Infect Immun* 21(2): 474-479; 1978.

The kinetics of the phytohemagglutinin (PHA) response of peripheral blood lymphocytes from chickens infected with oncogenic Marek's disease virus (MDV) or nononcogenic herpesvirus of turkeys (HVT) was studied using a whole-blood microassay. Approx 7 days after inoculation, the PHA response was depressed in MDV-inoculated resistant line N or susceptible line 7₂ chickens and in HVT-inoculated line 7₂ chickens. All chickens initially regained their PHA responsiveness. Susceptible chickens that died of MD or developed MD lymphoma in later stages of virus infection showed a second severe depression in the PHA response. No depression was observed in HVT-vaccinated chickens when challenged with MDV. The PHA response of MDV-inoculated chickens that survived MD, HVT-inoculated chickens,

and HVT-vaccinated MDV-challenged chickens showed evidence of enhancement. The depression of the PHA response was attributed to the suppressive effect of macrophages on T-cell response. This finding is consistent with the results of previous studies on MDV suppression of the PHA response. (27 refs.)

- 78-5197 Identification of an Abelson Murine Leukemia Virus-encoded Protein Present in Transformed Fibroblast and Lymphoid Cells.** (Eng) Witte, O. N. (Dept. Biology, Massachusetts Inst. Technology, Cambridge, MA 02139); Rosenberg, N.; Paskind, M.; Shields, A.; Baltimore, D. *Proc Natl Acad Sci USA* 75(5): 2488-2492; 1978.

A search for gene products in Abelson murine leukemia virus (A-MuLV)-transformed cells that could play a role in transformation revealed that extracts from A-MuLV-transformed lymphoid and fibroblast cell lines contained a protein with a mol wt of 120,000 daltons (P120). Immunoprecipitation with specific sera indicated that P120 contained regions homologous to the 5'-terminal segment of the MuLV *gag* gene complex (p15, p12, and at least part of p30), but it lacked detectable determinants of p10, reverse transcriptase, and the envelope glycoprotein. P120 was phosphorylated and had an intracellular half-life of 3-6 hr. In vitro translation of viral RNA from A-MuLV, with Moloney MuLV as helper, yielded a 120,000-dalton product with serological reactivity similar to that of cellular P120. Translation of the RNA from the helper produced no P120. P120 was expressed in A-MuLV-transformed lymphoid and fibroblastic lines tested, but it was not detectable in a lymphoid line in which the A-MuLV genome was established by infection but was not responsible for the transformation. Expression of P120 was selectively retained in clones of A-MuLV-transformed lymphocytes that converted to a nonproducer state after loss of expression of helper MuLV intracellular precursors. These findings suggest that the P120 product of the A-MuLV genome may be responsible for maintenance of the transformed phenotype of lymphoid and fibroblast cells transformed by the virus. (28 refs)

- 78-5198 Abelson Virus-induced Lymphomagenesis in Mice.** (Eng) Risser, R. (McArdle Lab. Cancer Res., Univ. Wisconsin, Madison, WI, 53706); Potter, M. Rowe, W. P. *J Exp Med* 148(3): 714-726; 1978.

The age- and virus-dose-dependence of Abelson murine leukemia virus (MuLV-A) lymphomagenesis and the pattern of susceptibility of various mouse strains to MuLV-A were examined. MuLV-A lymphoma induction was age-dependent, virus-dose-dependent, and under the control of host genes unrelated to other genes known to control murine leukemia (*Fv-1* or *Fv-2*). The most common sites of lymphoma were

the vertebral regions and skull, frequently resulting in the characteristic caput previously described for Abelson disease. Of 16 mouse strains tested, only BALB/c and some of its derivative strains showed high sensitivity to MuLV-A infection. Studies with CXB (BALB x B6) recombinant inbred strains and hybrids between them suggest that BALB/c carries dominant sensitivity alleles at two loci, tentatively designated *Av-1* and *Av-2*, which confer on these mice partial susceptibility to MuLV-A lymphoma induction. In addition, the *Av-2* gene may play a minor role in determining the susceptibility of mice to MuLV-A, its effect being seen only in mice homozygous for resistance at both *Av-1* and *Av-2*. Virologic studies indicate that the resistance of adult B6 mice is not related to restriction in the helper virus replication, but is specific for the effective transforming virus genome. (25 refs)

78-5199 Cells Nonproductively Transformed by Abelson Murine Leukemia Virus Express a High Molecular Weight Polyprotein Containing Structural and Nonstructural Components. (Eng) Reynolds, F. H. (NCI, Frederick Cancer Research Center, Frederick, MD, 20014); Sacks, T. L.; Deobagkar, D. N.; Stephenson, J. R. *Proc Natl Acad Sci USA* 75(8): 3974-3978; 1978.

Cell clones nonproductively transformed by the replication-defective Abelson strain of murine leukemia virus (A-MuLV) were analyzed for C-type viral antigen expression by competition immunoassay. AbLV-transformed mink non-producer lines expressed a 110,000- to 130,000-mol-wt polyprotein containing MuLV gag proteins p15 and p12 covalently linked to nonstructural A-MuLV-coded component(s) of approx 80,000-100,000-mol-wt. This polyprotein lacked detectable antigenic cross-reactivity with other virion-coded gag gene proteins such as p30, p10, viral reverse transcriptase, and gp70. The results suggest that a portion of this polyprotein might represent the A-MuLV *src* gene product and that in translation, it is initially linked in precursor form to gag structural proteins. Superinfection of mink cells nonproductively transformed by A-MuLV with a wild mouse amphotropic type C virus isolate, 4070-A, or the endogenous cat virus RD114 led to the production of pseudotype virus containing high concentrations of the A-MuLV-coded precursor polyprotein. (41 refs.)

78-5200 Abelson Antigen: A Viral Tumor Antigen That Is Also a Differentiation Antigen of BALB/c Mice. (Eng) Risser, R. (McArdle Lab. Cancer Res., Univ. Wisconsin, Madison, WI, 53706); Stockert, E.; Old, L. J. *Proc Natl Acad Sci USA* 75(8): 3918-3922; 1978.

The hypothesis that Abelson murine leukemia virus (A-MuLV) induces a cell surface antigen cross-reactive with one normally present only in uninfected BALB/c mice was tested. Serum from C57BL/6 mice hyperimmunized with

syngeneic A-MuLV lymphoma cells was cytotoxic for the immunizing cells. Using this reaction as a serologic test system for recognition of A-MuLV antigens, two serologic specificities were detected by absorption analysis with 40 tumors and 21 cell lines; (1) Friend-Moloney-Rauscher antigen(s) related to the Moloney MuLV helper and (2) an antigen expressed on all cells transformed by A-MuLV. The A-MuLV-specific antigen was also present on uninfected cells from BALB/c bone marrow, spleen, and fetal liver, but not cells from adult liver, thymus, lymph nodes, or peripheral blood. Abelson antigen was not expressed on bone marrow or spleen cells of 12 other murine strains. The Abelson antigen may be an antigenic determinant specified by a gene of BALB/c mice that was incorporated by recombination into the Moloney MuLV genome. (34 refs.)

78-5201 Murine Leukemia Virus (T-8)-transformed Cells: Identification of a Precursor Polyprotein Containing gag Gene-coded Proteins (p15 and p12) and a Nonstructural Component. (Eng) Sacks, T. L. (Lab. RNA Tumor Viruses, NCI, Bethesda, MD, 20014); Reynolds, F. H.; Deobagkar, D. N.; Stephenson, J. R. *J Virol* 27(3): 809-814; 1978.

Mink cells nonproductively transformed by the T-8 strain of mink cell focus-inducing virus, Abelson leukemia virus (AbLV), Moloney murine sarcoma virus (M-MSV), or the S+L- isolate of MSV were analyzed for high-mol-wt polyproteins. Cells transformed by T-8 or AbLV expressed the two amino terminal murine leukemia virus (MuLV) gag gene-coded proteins p15 and p12, but not the remaining gag gene-coded proteins p30 and p10, the *env* gene product gp70, or the reverse transcriptase. Virus-coded polyproteins of around 90,000 to 110,000 and 110,000 to 130,000 mol wt were demonstrated in T-8- and AbLV-transformed mink cells, respectively, by gel filtration and by immunoprecipitation followed by gel electrophoresis. In both cases, these proteins were precipitated by sera against Rauscher MuLV or M-MuLV, but not by control sera or by high-titered sera against MuLV gp70. To date, identification of high-mol-wt virus-coded precursor polyproteins containing structural and nonstructural components has been limited to those transforming viruses coding only for p15 and p12. (27 refs)

78-5202 Cell-free Synthesis of Rauscher Murine Leukemia Virus "gag" and "gag-pol" Precursor Polyproteins from Virion 35 S RNA in a mRNA-dependent Translation System Derived from Mouse Tissue Culture Cells. (Eng) Murphy, E. C. (Dept. Biology, Univ. Texas System Cancer Center, M.D. Anderson Hosp. and Tumor Inst., Houston, TX, 77030); Arlinghaus, R. B. *Virology* 86(2): 329-343; 1978.

The capacity of 35S genomic RNA from Rauscher leukemia

virus (RLV) to code for the synthesis of viral proteins was examined using a messenger RNA (mRNA)-dependent cell-free protein synthesis system derived from mouse tissue culture cell extracts by treatment with micrococcal nuclease. Analysis of the polypeptide product by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) indicated that RLV 35S RNA directed the synthesis of RLV-specific polypeptides of 55,000, 65,000, 75,000 and 200,000 apparent mol wt, identical in size to the known intracellular precursors of RLV mature polypeptides. None of the individual mature virus proteins appeared to be synthesized. When canavanine, an arginine analog, was substituted for arginine under conditions for cell-free protein synthesis, a RLV-specific polypeptide with a mol wt of approx 80,000 was synthesized at the expense of the 65,000- and 75,000-mol wt polypeptides. Monospecific antisera against p30, p15, p12, and p10 recognized polypeptides of mol wt 65,000, 75,000, 80,000 and 200,000, indicating that each shared antigenic determinants with all of these gag proteins. Using the appropriate authentic RLV precursor polypeptides as standards, comparative tryptic maps were performed with the ³H-tyrosine or ³⁵S-methionine-labeled in vitro-synthesized 65,000- and 200,000-mol wt polypeptides. The former was found to be identical to Pr65-gag, a 65,000-mol wt RLV gag protein precursor obtained from infected cells by immunoprecipitation and gel electrophoresis. The second polypeptide was found to contain methionine-labeled tryptic peptides characteristic of RLV reverse transcriptase (pol) and p30. (44 refs)

- 78-5203 Reverse Transcriptase-associated RNase H Activity: II. Inhibition by Natural and Synthetic RNA.** (Eng) Marcus, S. L. (Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Smith, S. W.; Modak, M. J. *J Virol* 27(3): 576-581; 1978.

The effect of exogenously added RNA molecules of natural and synthetic origin on the RNase H activities associated with purified avian myeloblastosis virus (AMV) and Rauscher murine leukemia virus (R-MuLV) DNA polymerases as well as on polymerase-independent RNase H activities was investigated. None of the polymerase-independent RNase H activities was inhibited by the addition of exogenous RNA, whereas those activities associated with R-MuLV and AMV DNA polymerases were strongly inhibited by RNA addition. It was also shown that the formation of duplex RNA completely abolished inhibitory activity and that various homopolymeric DNA's gave similar, if not identical, patterns of inhibition with R-MuLV DNA polymerase-associated RNase. However, polyguanylic acid was the only homopolymeric RNA that effectively inhibited this enzyme activity. An examination of RNA inhibition of the avian and murine oncoviral DNA polymerase-associated RNase H activities showed that they differed. Firstly, the relative efficiency of inhibition by various homopolymeric RNA molecules varied. Secondly, the RNase H activity associated with the AMV enzyme could be completely inhibited by increasing concen-

trations of naturally occurring RNA's, but that association with the R-MuLV DNA polymerase was unaffected by the presence of such molecules at similar concentrations. Lastly, R-MuLV DNA polymerase-associated RNase H activity could be inhibited by the addition of homopolymeric RNA to an ongoing reaction, whereas similar additions did not affect the ongoing activity of the AMV polymerase-associated RNase H. (20 refs.)

- 78-5204 A Combined Assay for the Rapid Preliminary Detection of Structural Retroviruses.** (Eng) Lick, B. J. (David H. David Labs., Cancer Inst., Hahnemann Medical Coll. and Hosp., 230 N. Broad St., Philadelphia, PA, 19102); Fuscaldo, A. A.; West, W.; Fuscaldo, K.; Brodsky, I. *Cancer Lett* 4(3): 163-169; 1978.

Equilibrium gradients, RNA-dependent DNA polymerase assays, and electron microscopy (EM) were used in a combined assay for the rapid preliminary detection of intact retroviruses in crude tissue preparations. The use of critical point drying as a preparative technique for EM is an integral part of this assay because it combines many of the advantages of thin sectioning and negative staining, but few of their limitations. Positive combined assays of platelets from preleukemic patients corresponded with their karyotypic abnormalities. Reconstruction experiments with Rauscher leukemia virus added to buffer or disrupted mouse spleen demonstrated the ease of detecting $\geq 10^3$ particles/g crude tissue; however, 10^4 particles or less were not detectable. The presence of added protein or other components of spleen tissue appeared to enhance viral stability or affinity to the EM grid. In unpublished results, platelets from patients with chronic myelogenous leukemia demonstrated the presence of reverse transcriptase without visible particles. These findings indicate that intact particles may be present in some preleukemic states but only viral information and/or subunits remain in most leukemic states. (18 refs)

- 78-5205 Type-specific Radioimmunoassays for the gp70 of Mink Cell Focus-inducing Murine Leukemia Viruses: Expression of a Cross-reacting Antigen in Cells Infected with the Friend Strain of the Spleen Focus-forming Virus.** (Eng) Ruscetti, S. (Lab. Tumor Virus Genetics, NCI, Bethesda, MD, 20014); Linemeyer, D.; Feild, J.; Troxler, L.; Scolnick, E. *J Exp Med* 148(3): 654-663; 1978.

The gp70 of a helper-independent strain of a Friend mink cell focus-inducing (MCF) virus, Fr-MCF-1, was isolated and used in a radioimmunoassay to detect any immunologically determinant(s) of the gp70s of MFC viruses derived from Friend murine leukemia virus (F-MuLV), Moloney MuLV and AKR-MuLV. Previous biochemical and biological studies showed that Fr-MCF-1 is an envelope gene recombinant between F-MuLV and a mouse xenotropic virus. The immunoassay utilized ¹²⁵I-Fr-MCF-1 gp70 and antiserum prepared

pared against an MCF strain of Moloney C-type virus. The MCF determinant(s) was not detected in the ecotropic parents of each of these MCF viruses, nor in helper-independent murine xenotropic viruses derived from Swiss or BALB/c mice. A protein partially cross-reactive with the MCF gp70 determinant(s) was detected in a replicating xenotropic virus derived from New Zealand Black mice. The MCF gp70-specific immunoassay could also detect a cross-reacting gene product coded for by the Friend strain of spleen focus-forming virus (SFFV) in rat fibroblasts nonproductively infected with SFFV. These findings support earlier molecular hybridization studies, which indicated that the genome of SFFV contains genetic information derived from both F-MuLV and xenotropic virus and that the xenotropic-related sequences in SFFV are highly related to those found in MCF murine C-type viruses. (14 refs.)

78-5206 Helper-independent Mink Cell Focus-inducing Strains of Friend Murine Type-C Virus: Potential Relationship to the Origin of Replication-defective Spleen Focus-forming Virus. (Eng) Troxler, D. H. (Lab. Tumor Virus Genetics, NCI, Bethesda, MD, 20014); Yuan, E.; Linemeyer, D.; Ruscetti, S.; Scolnick, E. M. *J Exp Med* 148(3): 639-653; 1978.

Cloned ecotropic Friend murine leukemia virus (F-MuLV) was inoculated into newborn NIH Swiss mice and the MuLV isolated from the preleukemic and leukemic spleens was used to study the relationship between the Friend erythroleukemia system and the AKR thymic leukemia system. Two helper-independent mink cell focus-inducing (MCF) strains of F-MuLV were isolated and shown by host range, interference pattern, and tryptic peptide analysis of the MuLV gp70s to be *env* gene recombinants between an ecotropic virus (F-MuLV) and a mouse xenotropic virus. Furthermore, RNA from the Friend MCF viruses hybridized completely to complementary DNA of spleen focus-forming virus (SFFV), a probe that detects that portion of SFFV not derived from F-MuLV. The ability to isolate replicating MCF viruses derived from F-MuLV strengthens the parallels between the Friend erythroleukemia system and the AKR thymic leukemia system. The potential relationship of helper-independent *env* gene recombinants between F-MuLV and xenotropic virus to the highly leukemogenic SFFV is discussed. (31 refs.)

78-5207 An Analysis of the Role of Lymphatic Leukemia Virus in the Immunodepression Exerted by Friend Complex in Leukaemia-resistant C57BL/6 Mice. (Eng) Bendinelli, M. (Inst. Microbiology, Univ. Pisa, 56100 Pisa, Italy); Campa, M.; Toniolo, A.; Garzelli, C. *J Gen Virol* 39(2): 243-251; 1978.

The transient depression in the ability of leukemia-resistant C57BL/6 mice to develop IgM and IgG antibody-producing

cells to sheep RBC following inoculation with the Friend leukemia complex (FLC) was investigated. Analysis of the lymphatic leukemia virus (LLV) component of FLC indicated that it was not capable of producing the same amount of immunodepression as the entire FLC. Mice inoculated with the Rowson-Parr isolate of LLV (RP-LLV) at various times before immunization constantly developed normal plaque-forming cell responses to sheep RBC. Following inoculation with the anemia-inducing strain of FLC from which RP-LLV was originally obtained, they exhibited a suppression of this response. A different isolate of LLV also failed to depress the plaque-forming cell responsiveness of C57BL/6 mice. Thus, the immunodepression cannot be attributed solely to the LLV component of FLC. This conclusion was also supported by evidence obtained from examining the replication of FLC and LLV, the antibody response to *Escherichia coli* lipopolysaccharide, and the ability of syngeneic macrophages administered together with the antigen to restore the response to sheep RBC. The mechanisms whereby FLC and its LLV component immunodepress are therefore partially different. (29 refs)

78-5208 Analysis of Translational Products of Friend Strain of Spleen Focus-forming Virus. (Eng) Barbacid, M. (Lab. Cellular and Molecular Biology, NCI, Bethesda, MD, 20014); Troxler, D. H.; Scolnick, E. M.; Aaronson, S. A. *J Virol* 27(3): 826-830; 1978.

Normal rat kidney (NRK) cells nonproductively infected with the Friend strain of spleen focus-forming virus (SFFV) were analyzed for the presence of murine leukemia virus (MuLV)-specific C-type viral proteins. SFFV was found to code for the p15 and p12 proteins of Friend-MuLV, as determined by immunological typing of their antigenic determinants. Molecular size analysis of the p15 and p12 proteins in SFFV nonproductively infected NRK cells indicated that these proteins are translated as parts of polypeptide molecules. The apparent mol wts of the polypeptides bearing p12 antigenic determinants revealed the presence of translational products of the SFFV genome that could not be accounted for by known C-type virus helper structural proteins. (29 refs)

78-5209 Elimination of Infectious MuLV and Related Disease in LC Wild Mice by Selective Breeding. (Eng.) Gardner, M. B. (Dept. Pathology, Univ. Southern California Sch. Medicine, Los Angeles, CA, 90032); Dougherty, M. F.; Casagrande, J.; Estes, J. D. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 88-89; 1978.

Infectious murine leukemia virus was eliminated from lym-

phoma/paralysis-prone LC wild mice by the selective breeding of nonviremic LC mothers. When the LC mothers were viremic, regardless of the serum virus status of the LC fathers, 92/115 of the F₁ progeny were viremic at 2 mo of age. When the LC mothers were nonviremic, only 2/172 of the progeny were viremic; all 114 of these progeny tested remained nonviremic to 9 mo of age. (6 refs)

78-5210 Nearest-Neighbor Interactions of the Major RNA Tumor Virus Glycoprotein on Murine Cell Surfaces. (Eng) Takemoto, L. J. (Dept. Microbiology, Univ. California, Los Angeles, CA, 90024); Fox, C. F.; Jensen, F. C.; Elder, J. H.; Lerner, R. A. *Proc Natl Acad Sci USA* 75(8): 3644-3648; 1978.

Chemical cross-linking was used in conjunction with immunoadsorption to probe for evidence of nearest-neighbor interactions of gp70 molecules, the major envelope constituent of C-type RNA viruses, on the surfaces of non-virus-producing [endogenous murine leukemia virus (MuLV)] murine cells. Radiolabeled gp70 molecules in LM fibroblast cells, B6 lymphocytes, and EL-4 lymphoma cells were linked to a protein of approx 15,000 daltons via native disulfide bonding. Prior treatment of the cells with the reversible, bifunctional, cross-linking reagent dimethyl-3,3'-dithiobispropionimidate, followed by immunoadsorption and two-dimensional diagonal electrophoresis, revealed apparent homodimers and homotrimers of the 85,000-dalton complex. Identical treatment of purified type C RNA tumor virus from murine cells also revealed homodimeric and homotrimeric species, demonstrating similar self-associating tendencies of this glycoprotein in both intact virus and the plasma membranes of nonproducing murine cells. One cross-linked product consistently detected on the surfaces of murine cells was not present after cross-linking of a representative strain of MuLV. (49 refs.)

78-5211 Distinguishable Biologic Effects of RNA Type-C Murine Leukemia and Sarcoma Viruses in Long-Term Bone Marrow Culture (Meeting Abstract). (Eng) Greenberger, J. (Joint Center Radiation, Sidney Farber Cancer Inst., Boston, MA); Davisson, P.; Gans, P. *J Reticuloendothel Soc* 23(6): 36i; 1978. (no refs)

78-5212 Induction of Endogenous Mouse Type C Virus (Meeting Abstract). (Eng) Hellman, K. B. (George Washington Univ., Washington, DC, 20006). *Diss Abstr Int B* 39(1): 68B-69B; 1978. (no refs)

78-5213 Two Different Murine Sarcoma Virus Isolates Have Homologous Genome Sequences: Implica-

tion for Their Origin. (Eng) Deng, C. (Div. Chemistry and Chemical Engineering, California Inst. Technology, Pasadena, CA, 91125); Wimmer, E. *Virology* 89(1): 309-313; 1978.

The RNase T1-resistant oligonucleotide fingerprints of the Moloney murine sarcoma virus clone 124 (MSV-214) and MSV clone 1712 were compared. Fingerprint comparison indicated that the RNA genomes of MSV-124 and MSV-1712 shared at least 14 large oligonucleotides, suggesting that the viruses are closely related. In view of this relationship, the hypothesis that MSV-1712 arose by recombination/deletion events of M4-MLV with mouse cellular DNA should be re-considered. (16 refs.)

78-5214 Heteroduplex Analysis of the Nonhomology Region Between Moloney MuLV and the Dual Host Range Derivative HIX Virus. (Eng) Donoghue, D. J. (Dept. Biology, Massachusetts Inst. Technology, Cambridge, MA, 02139); Rothenberg, E.; Hopkins, N.; Baltimore, D.; Sharp, P. A. *Cell* 14(4): 959-970; 1978.

The location of the glycoprotein coding sequences of the dual host range virus HIX was investigated by heteroduplex analysis. This virus was previously characterized as an envelope gene recombinant between Moloney murine leukemia virus (Mo-MuLV) and an unidentified xenotropic murine leukemia virus. Using long reverse transcripts of Mo-MuLV, a region of nonhomology was mapped by electron microscopic analysis of heteroduplexes formed with HIX 35S virion RNA. In this nonhomology region, the Mo-MuLV complementary DNA (cDNA) strand measured approx 900 nucleotides, mapping between 1.6 and 2.5 kilobases from the 3' end. In a previous study, hybridization of Mo-MuLV 21S RNA with Mo-MuLV cDNA resulted in the formation of different heteroduplex structures diagnostic of a noncontiguously coded leader sequence at the 5' end of the 21S RNA. Following hybridization of poly(A)+ HIX 21S RNA with 8.2-kilobase Mo-MuLV cDNA, analogous heteroduplex structures were observed exhibiting the Mo-MuLV:HIX substitution loop in the DNA:RNA segment of the molecules. This analysis permitted more precise mapping of the nonhomology region with respect to the splice point in the 21S presumptive glycoprotein messenger RNA. The mapping of this nonhomology region in HIX virus provides an internal visual marker for the 3' end of the genome. (33 refs.)

78-5215 High Frequency of Aberrant Expression of Moloney Murine Leukemia Virus in Clonal Infections. (Eng) Shields, A. (Dept. Biology, Massachusetts Inst. Technology, Cambridge, MA, 02139); Witte, O. N.; Rothenberg, E.; Baltimore, D. *Cell* 14(3): 601-609; 1978.

The clonal progeny of a culture of NIH/3T3 cells infected with Moloney murine leukemia virus (M-MuLV) were studied for production of infectious virus, antigens of the virus,

and noninfectious particles. There was approx one defective infected clone for every two normal productive infected clones. One clone produced virus with altered plaque morphology, but others failed to produce particles able to make plaques on XC cells. Two clones made particles lacking reverse transcriptase, and they did not synthesize the reverse transcriptase precursor Pr180-gag-pol. One clone (M23) lacked both detectable glycoprotein and reverse transcriptase. Despite these defects, all clones released particles with C-type morphology, suggesting that gag gene function alone may be sufficient for particle production. The monomer RNA's resulting from the denaturation of 70S RNA from all clones but M23 were electrophoretically indistinguishable from the RNA isolated from the producer line. All particles contained viral RNA of 60S-70S that was composed of the normal 35S size subunits except for M23, which had a deletion in the viral genome of approx 1,000-1,500 nucleotides. A variety of defective clones were also isolated from normal rat kidney cells infected with M-MuLV. Apparently, the M-MuLV genome is often mutated by spontaneous processes generating a wide range of phenotypes. (36 refs.)

8-5216 **A Competition Immunoassay for Characterizing the Reverse Transcriptase of Mammalian RNA Tumor Viruses.** (Eng) Panet, A. (Dept. Virology, Hebrew Univ., Hadassah Medical Sch., Jerusalem, Israel); Kra-oz, Z. *Virology* 89(1): 95-101; 1978.

A simple and rapid competition enzyme immunoassay for the reverse transcriptase (RT) of mammalian RNA tumor viruses is described. The assay is based on competition between an enzymatically active RT probe and a sample tested for RT antigen on a limited amount of monospecific antibody. The extent of competition is determined from the residual DNA polymerase activity left in the probe. Any RT activity in the test sample that could interfere with the enzymatic assay is inhibited irreversibly by pretreatment with N-ethylmaleimide. Thus, increasing amounts of test antigen neutralize more antibodies and restore more of the RT probe activity. The type, group, and interspecies antigenic determinants in the RT of mammalian RNA tumor viruses were resolved using the competition assay. The assay was also useful for quantitating the amount of RT present in virions. Application of this procedure to Moloney murine leukemia virus revealed that RT comprised 2.5% of the total viral proteins. (21 refs.)

8-5217 **Genome Organization of RNA Tumor Viruses. I. In Vitro Synthesis of Full-Genome-Length Single-stranded and Double-stranded Viral DNA Transcripts.** (Eng) Verma, I. M. (Tumor Virology Lab., Salk Inst., San Diego, CA, 92112). *J Virol* 26(3): 615-629; 1978.

The synthesis of large complementary DNA (cDNA) transcripts from avian myeloblastosis virus (AMV), a cloned iso-

late of Moloney murine leukemia virus (M-MuLV), and Moloney murine sarcoma virus (M-MSV) were studied. The longest cDNA transcripts synthesized using AMV, M-MuLV, and M-MSV (clone 124) were 7, 9, and 6.5 kilobases (kb), respectively. Double-stranded DNA could also be synthesized in vitro, but the second strand had an av size of only 0.8 to 1.2 kb. Two methods were used to synthesize double-stranded M-MuLV: (1) the 3' end of the minus strand was extended by the addition of 50 to 100 deoxyAMP residues using terminal deoxynucleotidyl transferase, and the plus strands were synthesized using the (dA)n-tailed cDNA transcripts as templates, an oligomer of deoxythymidine (dT) as primer, and *Escherichia coli* DNA polymerase; and (2) DNase-digested calf thymus DNA was used to prime the synthesis of plus strands on long cDNA with *E. coli* DNA polymerase I. Regardless of the size of the parent cDNA transcripts, both methods were capable of synthesizing double-stranded viral DNA. The plus strand of DNA synthesized by oligo(dT) primer sedimented heterogeneously with an av size of 5 to 6 kb, whereas the plus strands synthesized by calf thymus DNA primer appeared to be no longer than 0.3 to 0.4 kb. It was possible to synthesize double-stranded DNA without primer, but the plus strand of DNA was covalently linked to the parent strand. (60 refs)

78-5218 **Generation of New Mouse Sarcoma Viruses in Cell Culture.** (Eng) Rapp, U. R. (Lab. Viral Carcinogenesis, NCI, NIH, Bethesda, MD, 20014); Todaro, G. J. *Science* 201(4358): 821-824; 1978.

Endogenous non-tumor-producing C-type viruses from C3H mice that replicated in transformed mouse fibroblast cell lines generated variants that could directly transform cells in culture. The variants also induced rapid formation of undifferentiated sarcomas in 100% of newborn NIH Swiss mice inoculated ip with 50 µl of culture fluid. The foci that were induced by the new variants in cultured cells initially showed signs of only minimal transformation when compared with foci induced by standard laboratory strains of murine sarcoma virus. Also, the morphological effects that accompanied the adaptation to high-titer growth in a particular cell line were transient; immediate selection of virus released from these areas of transient transformation was needed in order to isolate the more stable, solid-tumor-inducing variant. The data show that variants of endogenous C-type virus that contain transforming genes (oncogenes) can arise during spread of the endogenous virus in fibroblast lines in vitro as well as in susceptible tissues in vivo. (26 refs.)

78-5219 **'Nonproductive' Mouse Sarcoma Virus Transformed Gorilla Cells which Contain Noninfectious Type C Virus Particles.** (Eng) Rhim, J. S. (Microbiological Associates, Inc., Bethesda, MD, 20016); Park, D. K.; Vernon, M. L. *Proc Soc Exp Biol Med* 158(3): 304-309; 1978.

Gorilla spleen cells were readily transformed morphologically in vitro by Kirsten murine sarcoma virus (Ki-MSV) and the MSV pseudotype of baboon placenta virus (MSV-BP). The Ki-MSV stock used was a rat cell-grown virus. MSV-BP was produced in human sarcoma-positive, leukemia-negative cells by cocultivation with a baboon virus-releasing human osteosarcoma line. The morphology of the Ki-MSV- and MSV-BP-transformed foci was similar. However, cells transformed by MSV-BP were producers, but the Ki-MSV-transformed cells were nonproducers (NP). These NP cells were induced by simple infection of gorilla cells with Ki-MSV. The NP cells did not release any infectious virus or murine leukemia virus group-specific antigen, but they did contain a rescuable MSV genome and noninfectious C-type particles with a buoyant density of 1.16 g/ml. Evidence is presented to show that the NP gorilla cells represent a new class of transformants; that is, gorilla NP cells containing noninfectious C-type virus without viral antigen. Attempts to rescue the genome of the focus-forming virus from gorilla NP cells by 5-bromodeoxyuridine yielded negative results. (23 refs.)

- 78-5220 Growth Factors from Murine Sarcoma Virus-transformed Cells.** (Eng) De Larco, J. E. (Lab. Viral Carcinogenesis, NCI, NIH, Bethesda, MD, 20014); Todaro, G. J. *Proc Natl Acad Sci USA* 75(8): 4001-4005; 1978.

The partial purification of a family of growth-stimulating polypeptides, sarcoma growth factors, from the supernatant fluids of Moloney murine sarcoma virus (MuSV)-transformed mouse 3T3 cells is described. These factors stimulate cells to divide in monolayer cultures and also to form colonies that grow progressively in soft agar. Three major peaks of activity were seen with apparent mol wts of 25,000, 12,000, and 7,000 daltons. The sarcoma growth factors were heat-stable, trypsin-sensitive, and active in nanogram quantities when tested for growth stimulation of untransformed rat and mouse fibroblasts. All three molecular species were also capable of competing for membrane epidermal growth factor (EGF) receptors when tested with ^{125} I-labeled EGF. However, they differed from mouse EGF in their mol wt, their inability to react with anti-EGF antibodies, and in their ability to convert cells to anchorage-independent (agar) growth. It is concluded that these sarcoma growth factors are a new class of polypeptide tropic factors that confer properties associated with the transformed phenotype on fibroblasts in vitro. (20 refs.)

- 78-5221 Particles from Mouse Melanoma B16 Containing Reverse Transcriptase and 70 S RNA Related to Human Melanoma Cytoplasmic RNA.** (Eng) Hellmann, R. (Medizinische Poliklinik, Ludwig-Maximilians-Universität, Pettenkoferstrasse 8a, D-8000 Munich 2, W. Germany); Balda, B. R.; Spiegelman, S. *Arch Dermatol Res* 262(2): 157-165; 1978.

The viruslike particles in B16 mouse melanomas were characterized. Material from a spontaneous melanoma was density-fractionated, and the viral-enriched fractions (density 1.1 g/cm³) were used for an endogenous reverse transcriptase reaction followed by glycerol sedimentation analysis of deproteinized nucleic acids. A portion of the ³H-DNA sedimented in discrete 70S peaks in the gradients. After RNase digestion of the 70S-³H-DNA, all the radioactivity was found in the low-mol-wt region of the gradient, showing that ³H-DNA was complexed to a high-mol-wt DNA. Thus, particles in the B16 melanoma contained a high-mol-wt 70S RNA associated with reverse transcriptase and possessed density characteristics of RNA tumor viruses. The ³H-DNA synthesized by the particles hybridized back to its template RNA; no hybrids were observed with mouse mammary tumor virus or avian myeloblastosis virus 70S RNA. The ³H-DNA also showed homology to RNA isolated from hamster melanoma and from three human malignant melanomas: > 7% hybridized to hamster melanoma RNA and to 5.7% hybridized to the human melanoma RNA. (30 refs.)

- 78-5222 Mammary Tumor Virus DNA: A Glucocorticoid-responsive Transposable Element.** (Eng) Yamamoto, K. R. (Dept. Biochemistry and Biophysics, Univ. California, San Francisco, CA, 94143); Stallcup, M. L.; Ring, J.; Ringold, G. M. *Cold Spring Harbor Symp Quant Biol* 42(part 2): 625-638; 1978.

Studies of the effects of dexamethasone on the expression of chromosomal sequences coding for mammary tumor virus (MTV) are described, along with evidence consistent with the idea that the MTV DNA sequence is a glucocorticoid-responsive transposable genetic element. When viral DNA was present in the genomes of a variety of mammalian cell types and was transcribed at a basal level, dexamethasone appeared to stimulate viral RNA production selectively. Evidence is presented indicating that the stimulation of viral RNA production is the result of an immediate shift to a new rate of transcription of the MTV genes is reviewed. Preliminary results indicate that RNA synthesis under cell-free conditions appears to reflect faithfully these different efficiencies of specific gene transcription. In experiments with MTV-infected HTC cells, the extent of viral gene expression and hormone responsiveness varied from clone to clone. Furthermore, infected clones were altered in the expression of a few specific genes. Unintegrated viral DNA was produced in these cells, perhaps via reverse transcription, but it probably was not extensively transcribed into RNA. (93 refs.)

- 78-5223 Genetic Transmission of Mammary Tumor Virus in the DBA/2 Mouse Strain.** (Eng) Versteeg, A. A. (Div. Genetics, Antoni van Leeuwenhoekhuis, Netherlands Cancer Inst., Amsterdam, Netherlands); van Nie, R. *Int J Cancer* 21(4): 473-475; 1978.

Mouse mammary tumor virus (MTV) antigens were demonstrable by radioimmunoassay (RIA) in milk samples from individual DBA/f mice and in samples from (female BALB/c x male DBA/f)_{F1} mice. Some samples collected during the first lactation periods were virus-negative but all samples of later lactation periods were virus-positive. Milk samples from 42/75 mice of the [female BALB/c x male (female BALB/c x male DBA/f)]Bc I population were virus-positive. In the [BALB/c x (BALB/c x DBA/f) x BALB/c]Bc II population two groups were distinguished. In the first group, the progeny of virus-positive Bc I mothers, 37/62 mice had detectable levels of viral antigen in the milk. None of the 43 samples from mice of the second group, derived from MTV-negative Bc I females, were virus-positive. Statistical analysis of the Bc I and Bc II data revealed that the segregation ratios did not deviate significantly from 1:1 ratios, which suggests that one dominant gene of DBA/f mice controls the expression of viral antigens in the milk. DBA/f mice have the non-agouti/brown/dilute coat color phenotype (a, b, d) and BALB/c mice the albino phenotype (alleles b, c). When the RIA data were grouped according to the coat color phenotype, a linkage relationship was observed between the absence of viral antigens in the milk and the c allele of the albino locus on chromosome 7 (recombination percentage: 20%). (17 refs.)

78-5224 Dependent and Autonomous Phases During Leukemogenesis. (Eng) Haran-Ghera, N. (Dept. Chemical Immunology, Weizmann Inst. Science, Rehovot, Israel). *Leuk Res* 2(1): 11-21; 1978.

The induction of leukemia in C57BL/6 mice by intrathymic injection of radiation leukemia virus variants that induce a high leukemia incidence (A-RadLV, 80%-100% incidence) or a low incidence (D-RadLV, 0%-20%) was investigated. Some mice were treated with D-RadLV plus, 3 days later, 400 R of whole-body x-radiation, which also yields a high leukemia incidence (70%-100%). The occurrence of preleukemic (PL) cells in the thymus (Th) and bone marrow (BM) was analyzed by cell transfer-genotype analysis. Samples of BM, Th, and whole blood, collected after 10-80 days from the treated mice, were injected iv into normal or irradiated syngeneic or (BALB/c x C57BL/6)_{F1} hybrids. The earliest (10 days) and highest incidence of PL cells was found in the BM of mice treated with D-RadLV alone. In contrast, in mice treated with A-RadLV or D-RadLV + radiation, PL cells were identified approx 30 and 40 days after injection, respectively. The latent period of leukemia development following transfer of BM cells from D-RadLV-treated donors was always delayed (100-125 days), irrespective of the time between virus injection and BM sampling. The ability of D-RadLV to induce a high incidence of PL cells indicates that a leukemogenic agent might exert a transformation effect without necessarily providing the proliferative capacity that would ultimately contribute to autonomous tumor development. An intact Th was required for the initiation of PL cell proliferation, which consisted of two phases, dependent and autonomous. In the former, the preferential growth of PL

cells in pretreated hosts depended on specific host environments (presence of Th, host irradiation, availability of specific lymphocyte populations). Ultimately, following sustained proliferation, the dependent PL cells underwent changes, with the emergence of autonomous clones whose proliferation was independent of host environment. The high or low potential of leukemogenic agents might be related to their capacity to induce dependent or autonomous transformed cells. Possible mechanisms involved in the transition from dependence to autonomy are given. (16 refs.)

78-5225 Characterization by Molecular Hybridization of Two Viral Populations Derived from a Radiation Leukemia Virus. (Eng) Janowski, M. (Departement de Radiobiologie, Centre d'Etude de l'Energie Nucleaire, B-2400 Mol, Belgium); Astier, T.; Guillemain, B.; Baugnet-Mahieu, L.; Duplan, J. F.; Maisin, J. R. *Cancer Res* 38(10): 3487-3493; 1978.

Two leukemogenic viral populations were derived from a radiation leukemia virus of the C57BL mouse and characterized. One (FB), in which only B-tropic virus could be detected, was obtained in vivo by serial passage of a cell-free extract in newborn rats. The second (3C), a complex containing at least B-tropic and xenotropic viruses, was produced in vitro by a permanent cell line (13-3C) established from the spleen of a virus-infected C57BL mouse. In molecular hybridization experiments, the 70S RNA of Gross leukemia virus hybridized 96% and 78% of FB and 3C radioactive complementary DNA's, respectively, with the duplexes formed having a relatively high thermal stability. In contrast, the 70S RNA of Rauscher leukemia virus hybridized 23% and 20% of the FB and 3C DNA probes, respectively, and the duplexes had low thermal stability. The rat-grown FB virions exhibited 50% genome homology with the viruses produced in vitro on the 13-3C cells. Finally, hybridizing the FB and 3C probes with normal or leukemic mouse spleen DNA's resulted in 89% to 100% homology. The rat-grown virions did not appear to contain detectable rat cellular DNA sequences, but about 20 complete copies of their nucleotide sequences were detected in covalent linkage with FB-infected rat spleen DNA. These findings strongly support the endogenous murine origin of the investigated virions. (27 refs.)

78-5226 Double-stranded DNA Synthesized by RD-114 Virion DNA Polymerase. (Eng) Yamazaki, H. (Dept. Molecular Biology, Cancer Res. Inst., Sapporo Medical Coll., Sapporo 060, Japan); Fujinaga, K. *Tumor Res (Sapporo)* 12: 29-36; 1977.

The double-stranded DNA (dsDNA) of RD-114 virus (an endogenous C-type feline virus) was synthesized by RD-114 virion DNA polymerase and used as a probe to investigate RD-114 virus-related DNA sequences in human cells. According to the reassociation kinetics of purified ³H-dsDNA,

the dsDNA was heterogeneous. DNA-DNA reassociation was used to separate the dsDNA into three fractions: a rapidly reassociating fraction (RR), and two slowly reassociating fractions (SR-1 and SR-2). The nucleotide complex of the RR fraction weighed 1.32×10^6 daltons (13.1% of the viral genome), and that of the SR-1 fraction weighed 3.34×10^6 daltons (33.1% of the viral genome). The size of nucleotide complex of SR-2 fraction could not be determined because of the heterogeneity of this fraction. Addition of human cell DNA (from a Burkitt's lymphoma line P3HR-1, peripheral blood of patients with acute myelogenous leukemia, and a rhabdomyosarcoma line RD) did not significantly accelerate the reassociation of the RR fraction. However, RD cell DNA appreciably accelerated the reassociation of SR-1 and SR-2, which suggests that RD-114 virus-related sequences were present in RD cells. Hybridization experiments with single-stranded DNA (ssDNA), which represented most of the viral genome sequences, and human DNA showed that only a small percentage of the ssDNA hybridized with human DNA. In addition, the thermal denaturation curves were broad. These findings indicate that few sequences specifically related to RD-114 DNA were present in human cells. (14 refs.)

- 78-5227 **Potentiation of the Neutralization of Feline Leukaemia Virus in Hypotonic Medium.** (Eng) Russell, P. H. (ICRF Tumour Immunology Unit, Dept. Zoology, Univ. Coll. London, Gower St., London WC1E 6BT, England). *J Gen Virol* 40(2): 373-378; 1978.

The neutralization of feline leukemia virus was found to be enhanced by performing the reaction in hypotonic medium. Absorption of neutralizing antibody by virus-specific antigens was also more sensitive in this medium. In one experiment, the serum titer of the virus was increased 30 times and the infectivity was decreased fivefold. (10 refs.)

- 78-5228 **Production of Antibodies against Feline Leukemia and Sarcoma Viruses and the Feline Oncovirus-associated Cell Membrane Antigen in Strain 2 Guinea Pigs.** (Eng) Log, T. (NCI, NIH, Bethesda, MD, 20014); Chang, S. S.; Sarma, P. S. *Virology* 89(1): 300-305; 1978.

Cloned cell lines of a guinea pig tumor induced by the Snyder-Theilen strain of feline sarcoma virus with feline leukemia virus subgroup C as helper [ST-FeSV (FeLV-C)] were established to study the host-virus relationship in an inbred laboratory animal strain. The cell line released high ST-FeSV titers (FeLV-C) (10^3 - 10^4 focus-forming units/0.2 ml) and 1 log excess of FeLV-C. In vivo studies with one cell line, GPTC C1, indicated that as little as 100 cells could induce tumors in weanling strain-2 guinea pigs following sc administration. The tumors grew progressively and killed the animals within 23 days. Sera from tumor-bearing guinea pigs contained antibodies to FeLV group-specific antigen, FeLV-C

virus envelope antigen, and feline oncovirus-associated cell membrane antigen. Absorption tests with the appropriate antigens suggested that the three antibodies were distinct. (2 refs.)

- 78-5229 **Studies on a Possible Viral Aetiology of Canine Lymphosarcoma and Mammary Tumours.** (Meeting Abstract). (Eng) Armstrong, S. J. (Dept. Clinical Veterinary Medicine, New Addenbrooks Hosp., Cambridge, England); Nunes de Souza, P.; Wreghitt, T. G.; Nagington, J.; Mahy, B. W.; Owen, L. N. *Br J Cancer* 38(1): 175; 1978. (1 ref)

- 78-5230 **Distribution of Bovine Leukemia Virus Proviral DNA Sequences in Tissues of Animals with Endemic Zootic Bovine Leukosis.** (Eng) Kettmann, R. (Departement de Biologie Moleculaire, Universite Libre de Bruxelles, Brussels, Belgium); Burny, A.; Cleuter, Y.; Ghysdael, J.; Mammerickx, M. *Leuk Res* 2(1): 23-32; 1978.

Normal and malignant tissues of bovine, ovine, porcine, and human origin were investigated for the presence of bovine leukemia virus (BLV) proviral DNA sequences using a BLV ^{3} H-thymidine-labeled, single-stranded, complementary DNA (cDNA) probe synthesized on purified BLV 60S-70S RNA as template by avian myeloblastosis virus. The malignant tissues were obtained from cattle and sheep experimentally infected with BLV, field cases of enzootic bovine leukosis (EBL, sporadic cases), a lymphoid tumor from a leukemic pig, and human leukemias and sarcomas. After recycling against normal bovine DNA, the probe hybridized with EBL tumor DNA in sheep and cattle but not with normal bovine DNA. The highest hybridization levels were obtained with tumorous lymph nodes (35%-38%). These results indicate that BLV is an exogenous RNA oncogenic virus and confirm that EBL is an infectious disease. Proviral BLV sequences were also detected in buffy coat cells of animals with persistent lymphocytosis (PL) and carrying antibodies to BLV but without tumors (30%-45% hybridization). This result clearly establishes that PL (without tumor formation) is most of the time, if not always, a subclinical form of EBL. In tumor-bearers, the proviral sequences were found in buffy coat cells, in solid tumors (lymphosarcomas) and in organs infiltrated with tumoral lymphoid cells but not in apparently normal organs. No hybridization above background was observed with DNA's extracted from buffy coat cells and tumors from the sporadic forms of bovine leukosis, pig leukemia, and the human cancers. Thus, BLV is probably unrelated to human cancers. (27 refs)

- 78-5231 **A Gene (Bevi) on Human Chromosome 6 Is an Integration Site for Baboon Type C DNA Provirus in Human Cells.** (Eng) Lemons, R. S. (Lab. Viral Car-

ogenesis, NCI, Bethesda, MD, 20014); Nash, W. G.; O'Brien, S. J.; Benveniste, R. E.; Sherr, C. J. *Cell* 14(4): 95-1005; 1978.

An attempt was made to map genetically a putative integration site for baboon endogenous C-type virus in humans using the technique of somatic cell hybridization. Human VA-2 cells infected with baboon C-type virus were cloned and fused to Syrian hamster cells, and 33 primary hybrid colonies were obtained. These cells segregated human chromosomes and retained the complete hamster genome. Assays for C-type viral p30 antigen and reverse transcriptase were performed in conjunction with analyses of 30 gene-enzyme systems representing 22 different human chromosomes. The results indicated that a gene, *Bevi*, previously assigned to human chromosome 6, dominantly controls baboon C-type virus expression in hybrid cells. Representative hybrid clones were studied by nucleic acid hybridization techniques for the presence of integrated proviral DNA using complementary H-DNA transcripts of the baboon viral RNA genome. For each of 12 clones examined, there was a concordance between the presence of human chromosome 6, the presence of baboon C-type proviral DNA sequences, and virus expression. Clones that segregated chromosome 6 (according to isozyme and karyological analyses) lost detectable proviral DNA sequences and failed to produce virus. There was no syntenic association between the replication of baboon virus and the presence of 21 other human chromosomes. It is concluded that *Bevi* is a preferred integration site for the baboon C-type provirus in the human genome. (47 refs.)

8-5232 **Susceptibility of Human Cultured Cells to Mason-Pfizer Monkey Virus.** (Eng) Ogura, H. Dept. Virology, Cancer Inst., Okayama Univ. Medical Sch., Okayama, Japan); Ocho, M.; Tanaka, T.; Oda, T. *Gann* 9(3): 413-415; 1978.

Cultures of the following seven types of human cells were inoculated with Mason-Pfizer monkey virus (M-PMV) to ascertain which types are susceptible to this virus: fibroblasts from whole embryo (HEC) and embryonic lung (HEL) cells, Detroit-551 embryonic skin cells, RSb cells (embryonic cells usually transformed by Rous sarcoma virus and simian virus 40), glial (KC) cells, and epithelioid carcinoma cells (HeLa and HEP-2). The infected cultures were maintained routinely with periodic transfers for 25 days and then examined electron microscopically and by the KC syncytium assay. All of the cells produced M-PMV-like particles approx 100 nanometers in diameter. Budding, extracellular, and intracytoplasmic particles resembling A-type particles were seen. None of the infected cells showed morphological changes. Mock-infected control cultures did not contain viruslike particles. Syncytium formation was observed at the second transfer of M-PMV-infected HEC, HEL, and Detroit-551 cells cocultivated with KC cells, but not in HeLa, HEP-2, RSb, and KC cells, even at the fifth transfer. However, these syncytium-negative cells showed M-PMV under the electron micro-

scope and in an infectivity assay of KC syncytium formation. The titers were 10^{-4} syncytium-forming unit/0.2 ml for HEC, HEL, and Detroit-551 cells; 10^{-3} for RSb and KC cells; and 10^{-1} for HeLa and HEP-2 cells. Thus, M-PMV propagated well in human fibroblastic cells and poorly in human epithelioid cells. (17 refs.)

78-5233 **Inhibition of Adenovirus Replication by 5,6-Dichloro-1- β -D-ribofuranosylbenzimidazole.** (Eng) Brotz, M. (Inst. Genetics, Univ. Cologne, Cologne, W. Germany); Doerfler, W.; Tamm, I. *Virology* 86(2): 516-529; 1978.

The halogenated benzimidazole derivative 5,6-dichloro-2- β -D-ribofuranosylbenzimidazole (DRB) reversibly inhibited the replication of human adenovirus type 2 (Ad2) and viral DNA in human KB cells. At 60 μ M, DRB reversibly inhibited Ad2 replication in KB cells by > 99.9% when it was added at the time of infection and present throughout infection. Viral DNA replication was almost completely inhibited when the drug was added < 4 hr postinfection in concentrations between 50 and 150 μ M. This inhibition affected all four size classes of viral DNA (> 100S, 50S-90S, 34S, and < 20S). The replication block was reversible upon withdrawal of the inhibitor. When DRB was administered at the time of max viral DNA replication, 16-18 hr postinfection, the inhibitor had no apparent effect on viral DNA synthesis. In the presence of 150 μ M DRB, the synthesis of early virus-specific RNA in the nucleus was reduced by approx 90% and the appearance of virus-specific RNA sequences in the cytoplasm by > 95%, as demonstrated by DNA-RNA filter hybridization. Thus, the block in viral DNA replication is best explained by the inhibition of early virus-specific RNA synthesis. (36 refs)

78-5234 **Alterations in Surface Glycoproteins and Level of Sialyltransferase Activity of Human Embryo Kidney Cells Infected with Oncogenic Adenovirus Type 12.** (Eng) Aoi, Y. (Inst. Medical Science, Univ. Tokyo, Tokyo 108, Japan); Yokota, M. *Tohoku J Exp Med* 125(2): 177-183; 1978.

A comparative study of plasma membrane glycoproteins and microsomal sialyltransferase activity was conducted in cultures of human embryo kidney (HEK) cells and in HEK cells infected with wild-type human adenovirus type 12 (H12) or *Cyt* mutants of H12. The H12 mutants used were *Cyt66* (highly tumorigenic) and *Cyt31* (less tumorigenic). The sodium dodecyl sulfate-polyacrylamide gel electrophoretic patterns of HEK cells demonstrated major glycoproteins in fractions 10, 18, and 19. Cells infected with H12 synthesized fraction 10 glycoprotein but not that of fractions 18 and 19. Cells infected with the *Cyt* mutants did not produce fraction 10 glycoprotein. Investigations of microsomal sialyltransferase indicated that the enzyme activity in H12-infected cells was increased by 230% over that seen in uninfected cells. The

level of sialyltransferase in *Cy66*-infected cells was 1.23 times higher than that of *Cy31*-infected cells. Thus, there was a positive correlation between sialyltransferase activity in the microsomes and tumorigenicity. (27 refs)

- 78-5235 **Are Oncogenic Human Adenoviruses Associated with Human Cancer? Analysis of Human Tumors for Adenovirus Transforming Gene Sequences.** (Eng) Green, M. (Inst. Molecular Virology, St. Louis Univ. Sch. Medicine, St. Louis, MO, 63110); Mackey, J. K. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1027-1042; 1977.

An adenovirus (Ad) etiology of human cancer was tested using sensitive Ad DNA probes to detect Ad DNA sequences in human tumors. Ad12 DNA and the Ad12 transforming restriction fragments (*Eco*RI-C fragment) were labeled in vitro to 10^7 to 4×10^8 cpm/ μ g by nick translation. Using Ad12- ^{3} H-DNA as a probe, DNA or RNA from 18 normal gastrointestinal (GI) tissues and 34 tumors, including cancers of the colon, rectum, small intestine, and stomach, was analyzed under conditions that would detect 0.1 copy of the Ad12 genome per tumor cell. No hybridization was detected with DNA or RNA extracted from 57 GI tumors, indicating that Ad12 is not a major cause of human GI cancer. The *Eco*RI-C fragment hybridized 50%-60% with other Ad12 isolates and with Ad18 and Ad31 (members of oncogenic group A), but not with 28 other human Ad serotypes. Thus, other group A members probably are also not involved in human GI cancer. No Ad12 DNA sequences were detected in 22 lung tumors, and no Ad12 *Eco*RI-C sequences were detected in 57 lung tumor DNA or RNA specimens. Similar experiments in which DNA or RNA from 55 GI tumors and 53 lung tumors was analyzed for the presence of sequences homologous to the transforming Group C Ad2 *Hind*III-G restriction fragment excluded the presence of less than one copy of 1% of the Ad2 genome in these tumors. Since Ad2 *Hind*III-G sequences are also present in all group C Ad's, these data indicate that group C Ad's do not cause 98% of human GI and lung tumors. (24 refs)

- 78-5236 **Target Cells of Adenovirus Type 12 in Brain Cell Cultures (Meeting Abstract).** (Eng) Ogawa, K. (Okayama Univ. Medical Sch., Okayama, Japan); Sonobe, H. *J Neuropathol Exp Neurol* 37(5): 667; 1978. (no refs)

- 78-5237 **Patterns of Integration of Viral DNA Sequences in the Genomes of Adenovirus Type 12-Transformed Hamster Cells.** (Eng) Sutter, D. (Inst. Genetics, Univ. Cologne, Cologne, W. Germany); Westphal, M.; Doerfler, W. *Cell* 14(3): 569-585; 1978.

The patterns of integration of the viral genome were analyzed in four hamster cell lines (HA12/7, T637, A2497-2, and A2497-3) transformed by adenovirus type 12 (Ad12). For each cell line, a distinct and specific pattern of integrated viral DNA sequences was observed for each restriction endonuclease (*Bam* HI, *Bgl* II, *Eco* RI, *Hind* III, *Hpa* II, and *Sma* I) used. The distribution patterns of the viral sequences complementary to the isolated *Eco* RI restriction endonuclease fragments were also specific for each cell line. Some of the viral DNA molecules were fragmented prior to or during integration. The *Eco* RI B, D, and E fragments, comprising a contiguous segment from 0.17 to 0.62 fractional length units of the viral DNA, remained intact during integration in a portion of the viral DNA molecules. Although each cell line carried multiple copies of Ad12 DNA, the viral DNA sequences were concentrated in a small number of distinct size classes of fragments. In the three cell lines tested (T637, A2497-2, and HA12/7), the integration pattern was stable over many generations and was identical in a number of randomly isolated clones. The data support the hypothesis that at least a portion of the viral DNA sequences is integrated into repetitive sequences or else that the integrated viral sequences are amplified after integration. (33 refs.)

- 78-5238 **Transformation-specific Antigen Induced by Oncogenic Human Adenovirus.** (Eng) Ledinko, N. (Dept. Biology, Univ. Akron, Akron, OH, 44325). *Nature* 274(5673): 812-813; 1978.

The human adenovirus type 12 (H12) transformation-specific protein(s) in H12-transformed cell lines was studied. A 60,000-mol wt transformation-specific antigen in H12-transformed rat 3Y1 cells was identified by immunoprecipitation of ^{35} S-methionine-labeled polypeptides with serum from H12-tumor-bearing hamsters. The expression of this antigen was temperature-dependent in 3Y1 cells transformed by the temperature-sensitive H12 mutant *ts401*. The altered behavior of the 60,000-mol wt protein observed in mutant infection strongly suggests that it is the product of the 401 gene. Alternatively, the protein may be a cell-coded protein, induced by the 401 function, and its reduction under restrictive conditions may reflect host cell alterations associated with a return to the normal phenotype. (25 refs.)

- 78-5239 **Transformation of Marmoset Lymphocytes In Vitro with *Herpesvirus ateles*.** (Eng) Falk, I. (Dept. Microbiology, Rush-Presbyterian-St. Luke's Medical Center, Chicago, IL, 60614); Johnson, D.; Deinhardt, F. *Int J Cancer* 21(5): 652-657; 1978.

Continuous lymphoblastoid cell lines with T-lymphocyte cell-surface markers were established from marmoset circulating lymphocytes transformed in vitro by *Herpesvirus ateles* (HVA). Blood samples were obtained from wild-caught cotton-topped (*Saguinus oedipus*), white-lipped (*S. fuscicollis*)

S. nigricollis), and common (*Callithrix jacchus*) marmosets. Transformation was effected by cocultivating the lymphocytes with lethally x-irradiated HVA-producing lymphoblastoid cells (I) or by infecting them with cell-free virus (II). A total of 10/17 cotton-topped, 13/16 white-lipped, and 19/39 common marmoset lines were obtained 17-35 days after transformation method I was initiated; 2/4 cotton-topped marmoset lines were established 51-53 days after the start of method II. Attempts to transform squirrel monkey lymphocytes by method I were not successful. The association of HVA with each transformed culture was demonstrated by staining of antigen-positive cells in indirect fluorescent antibody tests and recovery of HVA after cocultivation of the transformed cells with permissive monolayer cells. Cells of most of the transformed cultures formed E (erythrocyte) rosettes, reacted with a specific anti-marmoset T-lymphocyte serum, and lacked surface immunoglobulin, properties identifying them as T cells. (22 refs)

78-5240 Replication of Herpesviruses in Human Cells Transformed by Cytomegalovirus. (Eng) Figueroa, M. E. (Dept. Microbiology, Milton S. Hershey Medical Center, Pennsylvania State Univ. Coll. Medicine, Hershey, PA, 17033); Geder, L.; Rapp, F. *J Gen Virol* 40(2): 391-398; 1978.

The replication of human cytomegalovirus (CMV) and herpes simplex virus (HSV) was studied in three human embryo cell lines (CMV-Mj-HEL-1, CMV-Mj-HEL-2, and CMV-Mj-HEL-2,T-1) transformed in vitro by human CMV. Growth studies revealed that these cells were completely resistant to infection by CMV strains AD169 and Mj and partially resistant to HSV types 1 and 2. Neither virus DNA nor virus proteins were synthesized in the transformed cells infected with CMV AD169. HSV production in CMV-transformed human embryo lung (HEL) cells was delayed when compared with the virus production in normal HEL cells, and spread of the cytopathic effect of HSV was slower in the transformed cells. The treatment of normal HEL cells with a crude extract of CMV-transformed HEL cells also resulted in inhibition of the spread of the cytopathic effect of HSV types 1 and 2. The inhibitory effect was not due to interferon, since vesicular stomatitis virus replication was not affected and several experiments showed that it was not due to mycoplasma. It is postulated that there are virus inhibitor molecules in CMV-transformed cells and that they are absent in normal HEL cells. (16 refs.)

78-5241 Detection of Epstein-Barr Virus DNA in Anaplastic Carcinoma of the Nasopharynx and Other Tumors. (Ger.) Wolf, H. (Max von Pettenkofer-

Institut Hygiene, Munich, W. Germany). *Fortschr Med* 96(15): 832; 1978.

Epstein-Barr virus (EBV) DNA was found in anaplastic nasopharyngeal carcinoma cells by in situ nucleic acid hybridization. An etiological relationship was found between EBV and Burkitt's lymphoma, but not between EBV and Hodgkin's disease or Kaposi's sarcoma. (no refs)

78-5242 Relation of Herpesvirus Hominis Type II to Carcinoma of the Cervix: An Animal Model for the Induction of Long-Term Latency of Herpesvirus Hominis Type II. (Eng) Tobin, S. M. (Dept. Res., Mount Sinai Hosp., 600 University Ave., Toronto, Ontario M5G 1X5, Canada); Wilson, W. D.; Fish, E. N.; Papsin, F. R. *Obstet Gynecol* 51(6): 707-712; 1978.

An animal model was developed to demonstrate the long-term persistence of acute herpesvirus hominis type II (HVH-II) vaginocervical infection, one of the conditions that must be proved if HVH-II is to be considered an etiologic factor in carcinoma of the cervix. Female Wistar rats and female Balb/c mice were infected with HVH-II. Acute herpetic infection of the cervix was verified by several techniques: cytopathic effects (CPE), serum antibody titers to HVH-II, cytologic examination of cervical squamous epithelial cells, and indirect double-antibody immunofluorescence staining. Rats and mice that gave a positive result in these tests were maintained for 30 and 18 mo, respectively, after which they were sacrificed and the pelvic sensory ganglia examined for persistence of HVH-II. Of the ganglia from infected rats and mice, 91.5% and 80.0%, respectively, showed persistence of the virus as determined by CPE and 51.5% and 40.0%, respectively, showed persistence of the virus as determined by immunofluorescent staining. These results indicate that following a proved acute cervicovaginal infection with HVH-II, the virus persists in the local pelvic sensory ganglia of a significant number of animals for the equivalent of 60 human yr. (25 refs)

78-5243 An Investigation into the Relationship of the Immunological Status of Herpes Simplex Virus Type-2 (HSV-2) Infected Animals and the Tumor-enhancing Property of this Infection. (Eng) Reiss-Gutfreund, R. J. (Inst. Cancer Res., Univ. Vienna, Vienna, Austria); Dostal, V.; Fanta, D. *Oesterr Z Onkol* 4(2/3): 35-39; 1978.

The immunological status of herpes simplex virus 2 (HSV-2)-infected animals was compared with that of controls by three in vivo tests of cell-mediated immunity plus an in vitro test involving stimulation of DBF, mouse splenocytes with phytohemagglutinin (PHA). In the skin-grafting test, C3H mice received tail skin grafts from C57Bl mice and, 48

hr later, were infected with 2.5×10^3 plaque-forming units (PFU) of HSV-2 sc. In a footpad test, Swiss mice were infected similarly and inoculated with 50 μ g PHA 12 days later. In the tuberculin test, Hartley strain II guinea pigs were inoculated with 10^5 PFU HSV-2 into both hind footpads and sensitized simultaneously with BCG. Thirteen days later, the guinea pigs were inoculated with tuberculin. The skin-grafting and PHA-footpad tests did not reveal any differences in cell-mediated immunity between the test groups and similarly treated, but uninfected, control groups. However, in the tuberculin test, the test animals had a slightly impaired reactivity compared with controls. In addition, in the *in vitro* PHA assay there was a transient (approx 1 day) reduction of the stimulation index in cultures of splenocytes from virus-infected mice approx 10 days postinfection. This reduction was the result of a low rate of ^{3} H-thymidine incorporation by the PHA-stimulated cultures and a relatively high rate of uptake of ^{3} H-thymidine by the control (unstimulated) cultures. The short-lived immunodeficiency of HSV-2-infected mice might be associated with the tumor-enhancing property of HSV-2, as the time of impairment of cell-mediated immunity was directly correlated with the time of tumor initiation found in previous animal experiments. (26 refs.)

- 78-5244 Molecular Genetics of Herpes Simplex Virus: Demonstration of Regions of Obligatory and Nonobligatory Identity Within Diploid Regions of the Genome by Sequence Replacement and Insertion.** (Eng) Knipe, D. M. (Marjorie B. Kovler Viral Oncology Labs., Univ. Chicago, 910 East 58th St., Chicago, IL, 60637); Ruyechan, W. T.; Roizman, B.; Halliburton, I. W. *Proc Natl Acad Sci USA* 75(8): 3896-3900; 1978.

The structure of the reiterated regions of the S component of the DNA's of herpes simplex viruses 1 and 2 (HSV-1, HSV-2) were probed in marker rescue experiments involving transfection of cells with mixtures of intact HSV-1 mutant DNA and individual DNA fragments generated by restriction endonuclease digestion of wild-type HSV-1 or HSV-2 DNA. DNA fragments from both reiterated regions of the S component of HSV-1 DNA rescued temperature-sensitive (ts) mutants tsLB2 and tsD. However, only one end of the S component of wild-type HSV-1 rescued tsD, although both ends rescued tsLB2. Recombinants produced with the heterotypic ends of the S complement specified corresponding polypeptides typical of HSV-1 and HSV-2. Several recombinant clones produced by rescue with HSV-2 DNA contained identical HSV-2 DNA insertions within both reiterated regions of the HSV-1 S component. Thus, the reiterated S sequences may contain a region of obligatory identity. HSV DNA could be expended by at least 5×10^6 by means of insertion in the S component, suggesting that it could be a vehicle for exogenous DNA. (14 refs.)

- 78-5245 Cytoskeletal Proteins in Normal, Infected and Virus-transformed Cells (Meeting Abstract).**

(Eng) Gabbiani, G. (Dept. Pathology and Molecular Biology, Univ. Geneva, CH-1211 Geneva, Switzerland); Borgia, R.; Chaponnier, C.; Rungger, E.; Weil, R. *Experientia* 34(7): 940; 1978. (no refs)

- 78-5246 Forms of Intracellular SV40 DNA-Protein Complexes (Meeting Abstract).** (Eng) Fanning, E. (Fachbereich Biologie, Universität Konstanz, Postfach 733, D-7750 Konstanz, W. Germany); Baumgartner, I.; Klempnauer, K. H.; Richter, A.; Otto, B. *Hoppe Seylers Z Physiol Chem* 359(9): 1081; 1978. (4 refs)

- 78-5247 Selective Extraction of Replicating Simian Virus 40 Nucleoprotein Complexes from BSC-1 Monkey Cells (Meeting Abstract).** (Eng) Adams, R. L. (Dept. Biochemistry, Univ. Glasgow, Glasgow G12 8QQ, Scotland); Eason, R.; Digges, I.; Lindsay, J. G.; Wolstencroft, L. *Hoppe Seylers Z Physiol Chem* 359(9): 1051-1171; 1978. (4 refs)

- 78-5248 Phenotypic Complementation of the SV40 tsA Mutant Defect in Viral DNA Synthesis Following Microinjection of SV40 T Antigen.** (Eng) Kriegler, M. P. (Div. Medical Oncology, Sidney Farber Cancer Inst., Boston, MA, 02115); Griffin, J. D.; Livingston, D. M. *Cell* 14(4): 983-994; 1978.

To determine whether the simian virus 40 (SV40) A gene product, the large tumor (T) antigen, is involved in the initiation of viral DNA replication, African green monkey CV-1 cells were microinjected with highly purified SV40 T antigen using protein-labeled RBC ghosts and polyethylene glycol as fusagen. The microinjected cells were then infected with a temperature-sensitive SV40 mutant (tsA209) that is defective in the initiation of viral DNA synthesis. *In situ* hybridization assays indicated that microinjection of both partially and highly purified T antigen increased the amount of viral DNA sequences in the monolayer. Furthermore, ^{3} H-thymidine-labeled and unlabeled Hirt supernatant from microinjected, tsA209-infected cells contained significantly more SV40 DNA than comparable extracts from sham-injected tsA209-infected or uninfected cells. Thus, the introduction of highly purified large SV40 T antigen led to phenotypic complementation of the tsA defect in viral DNA synthesis. (35 refs.)

- 78-5249 Complete Nucleotide Sequence of SV40 DNA** (Eng) Fiers, W. (Lab. Molecular Biology, Univ. Ghent, Ghent, Belgium); Contreras, R.; Haegeman, G.; Rogiers, R.; Van de Voorde, A.; Van Heuverswyn, H.; Van Herreweghe, J.; Volckaert, G.; Ysebaert, M. *Nature* 273(5658): 113-120; 1978.

the entire nucleotide sequence of simian virus 40 (SV40) DNA (strain 776), a supercoiled circular duplex of 5,224 base pairs, was determined. It is estimated that at least 15.2% of the genome is not translated into polypeptides. DNA replication initiates bidirectionally from approx position 0.67 and terminates on the other side of the circle, where the two replication forks meet. The half-circle from 0.67 to 0.17, replicated and transcribed counterclockwise, corresponds almost exactly to the early region, but the segment from 0.67 to 0.17, replicated and transcribed clockwise, corresponds to the late region. The early t and T antigens are initiated at the same position, and the T antigen is coded by two noncontiguous regions of the genome; the T antigen messenger RNA (TmRNA) is spliced in the coding region. In the late region, the gene for the major protein VP₁ overlaps those for VP₂ and VP₃ by over 122 nucleotides, but it is read in a different frame. It is thought that the mRNA's for these three proteins are spliced out of a common primary RNA transcript. The use of degenerate codons is nonrandom, but it is similar in both the early and late regions. Codons such as NUC, NCG, and UGA are absent or very rare. (99 refs)

78-5250 Recently Replicated Simian Virus 40 DNA Is a Preferential Template for Transcription and Replication. (Eng) Green, M. H. (Dept. Biology, Univ. California at San Diego, La Jolla, CA, 92093); Brooks, T. L. *J Virol* 26(2): 325-334; 1978.

The template activities for replication and transcription were compared for recently replicated (new) and uniformly labeled (old) simian virus 40 (SV40) DNA in TC7 infected monkey cells. New SV40 DNA (pulse-labeled for 1 hr) served as a template for a second round of replication with a relatively high probability (8% of the DNA replicated per hour) for a period of 5 hr; after this time, the template activity rapidly decreased severalfold. Old SV40 DNA (labeled for 24 hr) functioned as a template for replication with a constant probability (1.8% of the DNA replicated per hour) for at least 24 hr. The proportion of RNA polymerase associated with newly replicated and with recently replicated (bromodeoxyuridine-substituted) viral DNA was determined by an assay that used the Triton-soluble SV40 transcription complex. The proportion of RNA polymerase associated with nonreplicating SV40 DNA decreased very slowly (to 50% in 6 hr); this suggested that replicating viral genomes are not required as templates for the initiation of late transcription. This suggestion was supported by the finding that the RNA synthesized *in vitro* was associated with covalently closed circular SV40 DNA. Furthermore, after 9 hr in bromodeoxyuridine, the recently replicated viral DNA had nearly three times more RNA polymerase per unit of DNA than did the nonreplicating DNA. It is concluded that recently replicated SV40 DNA is preferentially utilized as a template for transcription and replication. (36 refs)

78-5251 An Immunochemical Investigation of SV40 T Antigens. 1. Production Properties and

Specificity of a Rabbit Antibody to Purified Simian Virus 40 Large-T Antigen. (Eng) Lane, D. P. (Dept. Zoology, Imperial Coll., London SW7 2AZ, England); Robbins, A. K. *Virology* 87(1): 182-193; 1978.

The specificity of a rabbit antiserum to purified large tumor (T) antigen, a polypeptide of simian virus 40 (SV40), was compared with that of hamster antitumor serum to gain insight into the specificity of the hamster immune response to an SV40-transformed cell-induced tumor (H65). (T antigen has an apparent mol wt of 84,000 or 94,000 daltons, depending on the extraction method.) Immune complexes of SV40-infected monkey kidney cell extracts and hamster antitumor sera were isolated and analyzed on sodium dodecyl sulfate-polyacrylamide gels. A band of 84,000 apparent mol wt that was strongly stained by Coomassie blue was obtained. The rabbit antiserum bound specifically to a 94,000-mol wt polypeptide present in the transformed H65 cell line. The hamster antitumor serum also bound to these components in extracts of transformed or lytically infected cells. Both rabbit anti-T serum and hamster anti-tumor serum specifically bound up to 80% of a purified 94,000-mol wt T antigen that had been isolated by prior immunoprecipitation with a hamster antitumor serum. Native T antigen was heat-stable and resistant to denaturation. The rabbit anti-T serum specifically precipitated an additional low-mol wt (19,000 daltons) polypeptide (small T antigen). This same component was precipitated by the hamster antitumor serum, and it was found during the course of an SV40 lytic infection and in SV40-transformed cells. The data indicated that t and T antigens share common antigenic determinants. A third species of 60,000 mol wt was also specifically precipitated by both the rabbit anti-T serum and hamster antitumor serum; its relationship to t and T is not known. (37 refs)

78-5252 New Region of the Simian Virus 40 Genome Required for Efficient Viral Transformation. (Eng) Bouck, N. (Dept. Microbiology, Univ. Illinois at Medical Center, Chicago, IL, 60680); Beales, N.; Shenk, T.; Berg, P.; di Mayorca, G. *Proc Natl Acad Sci USA* 75(5): 2473-2477; 1978.

The transformation of normal nonpermissive rat cells (F111) to anchorage independence was measured after they were infected with viable mutants of simian virus 40 (SV40) that had deletions in three regions of the virus genome: map coordinates 0.21-0.17, 0.59-0.54, and 0.67-0.74. Only those mutants whose deletions occurred between 0.59 and 0.55 in the proximal part of the early region were defective in transforming ability. The most severely defective mutants transformed with less than one-hundredth the efficiency of wild-type virus. These mutants retained their defect when tested in Chinese hamster lung cells and when infection was initiated with viral DNA instead of intact virions. Complementation for transformation was observed between these transformation-defective deletions and an SV40 temperature-sensitive A mutant. (31 refs)

- 78-5253 Replication of Simian Virus 40 in Simian Virus 40-transformed Hamster Kidney Cells Induced by Mitomycin C or ^{60}Co γ Irradiation.** (Eng) Rakusanova, T. (Dept. Medicine, Massachusetts General Hosp., Boston, MA, 02114); Smales, W. P.; Kaplan, J. C.; Black, P. H. *Virology* 88(2): 300-313; 1978.

The kinetics of simian virus 40 (SV40) DNA replication and late virus protein synthesis were examined in various clones of induced SV40-transformed Syrian golden hamster kidney (SV-HK) cells and compared with those in lytically infected, permissive, monkey (SV-PM) cells. Virus yields in the SV-HK cells obtained after induction with ^{60}Co γ -radiation (250 rads/sec) or mitomycin C (0.5 $\mu\text{g}/\text{ml}$) were low, and the clones were heterogeneous for infectious virus induction. To investigate the reason(s) for the heterogeneity and low virus yields, the extent of SV40 replication and production of virion (V) antigen and infectious virus after induction were examined. Cells replicating SV40 DNA or synthesizing V antigen were determined by in situ hybridization and immunofluorescence, respectively. Only some cells in induced cultures produced SV40, and those that did were less efficient than SV-PM cells. The inducers acted by inducing more cells to replicate virus rather than by increasing the amount of SV40 released from individual cells. In all clones studied, a greater proportion of cells could be induced to replicate SV40 DNA than to synthesize V antigen. Furthermore, SV40 DNA replication could be induced by lower doses of γ -radiation than could the production of V antigen or infectious virus, which suggests that, in induced cells, synthesis of late virus protein is more restricted than SV40 DNA replication. Thus, one of the effects of induction of SV-HK cells was an enhancement of the cells' capacity to support SV40 replication. (19 refs.)

- 78-5254 Biochemical and Ultrastructural Analysis of SV40 Chromatin.** (Eng) Keller, W. (Cold Spring Harbor Lab., Cold Spring Harbor, NY, 11524); Muller, U.; Eicken, I.; Wendel, I.; Zentgraf, H. *Cold Spring Harbor Symp Quant Biol* 42(part 2): 227-244; 1978.

Simian virus 40 (SV40) chromatin containing a full complement of histone H1 in addition to the intranucleosomal histones H2A, H2B, H3, and H4 was isolated from SV40-infected African green monkey kidney CV1 cells and characterized. SV40 chromatin isolated at moderate ionic strength ($\mu = 0.1-0.3$) contained all five histones and appeared highly condensed under the electron microscope: it was composed of a few closely connected large spheres with an av diameter of 190 Å (range 160-220 Å). At high ionic strength ($\mu = 0.6-0.8$) or after prolonged exposure to very low ionic strengths ($\mu < 0.02$), the compact form unfolded and the chromatin exhibited a typical nucleosomal morphology. Native SV40 DNA-protein complexes contained a median number of 24 nucleosomes. The number of superhelical turns did not differ in DNA obtained from the compact and the unfolded forms of chromatin. DNA-relaxing enzyme

was found to be associated with SV40 chromatin, and it was capable of acting both on extraneously added circular DNA and on its own DNA in the nucleoprotein complex. Purified DNA-relaxing enzyme formed transiently nicked DNA intermediates, and the enzyme could be found covalently attached to the site of the nick in the DNA. Transcriptionally active SV40 complexes underwent the same ionic strength-dependent structural transition as that of bulk SV40 chromatin and, therefore, it may also have a compact configuration at physiological salt concentrations. (71 refs.)

- 78-5255 A Novel Sequence Arrangement of SV40 Late RNA.** (Eng) Hsu, M. T. (Rockefeller Univ., New York, NY, 10021); Ford, J. *Cold Spring Harbor Symp Quant Biol* 42(part 2): 571-576; 1978.

The 16S and 19S late messenger RNA's (mRNA's) of simian virus 40 (SV40) were mapped on the SV40 genome. The simple technique used is as follows: SV40 form-I DNA was cleaved with *EcoRI* endonuclease to generate full-length linear molecules. Subsequent treatment of these linear DNA molecules with exonuclease III exposed SV40 DNA sequences complementary to the 5' end of the mRNA molecules. Hybrids between the mRNA and exonuclease-treated DNA could be observed by electron microscope or by gel electrophoresis after enzymatic digestion of the hybrid. The size of the RNA-DNA hybrid defined the distance between the *EcoRI* site and the 5' ends of mRNA. Both the 16s and 19s late mRNA's had a 5' spliced sequence derived from a DNA region not immediately adjacent to the main coding region of the RNA. Within the precision of the technique, the 5' leader of both mRNA species is concluded to be derived from the region of the SV40 genome from 0.70 to 0.75 map unit. (13 refs.)

- 78-5256 Assembly of SV40 and Polyoma Minichromosomes During Replication.** (Eng) Cremisi, C. (Dept. Molecular Biology, Institut Pasteur, 75015 Paris, France); Chestier, A.; Yaniv, M. *Cold Spring Harbor Symp Quant Biol* 42(part 2): 409-416; 1978.

The assembly of simian virus 40 (SV40) and polyoma virus minichromosomes during replication was studied using African green monkey kidney CV1 cells and 3T6 mouse fibroblasts infected with SV40 and polyoma virus. The replicating intermediates of the viral DNA were associated with histones in the form of nucleosomes in both the parental and daughter double-helical DNA. Newly synthesized histones were preferentially associated with these replicating intermediates and did not exchange with the majority of the histones present on the mature complexes. Polyoma viral DNA that replicated in the absence of protein synthesis contained an av of 1 nucleosomes; this suggested that parental histones are redistributed between the two newly replicated viral DNA molecules. Their repartition on the newly replicated viral DNA appeared to be partially cooperative. (37 refs.)

78-5257 **Nucleosome Structure II: Structure of the SV40 Minichromosome and Electron Microscopic Evidence for Reversible Transitions of the Nucleosome Structure.** (Eng) Oudet, P. (Laboratoire de Genetique Moleculaire des Eucaryotes du CNRS, U.44 de l'INSERM, Institut de Chimie Biologique, Faculte de Medecine, 67085 Strasbourg Cedex, France); Spadafora, C.; Chambon, P. *Cold Spring Harbor Symp Quant Biol* 42(part 2): 301-312; 1978.

Simian virus 40 (SV40) minichromosomes extracted from CV1 cells were used as a model for chromatin in studies to determine whether the compact nucleosomal form of chromatin could be opened and extended in a reversible manner that would not involve the complete disruption of the histone-DNA bonds. Electron microscopic evidence indicates that at very low ionic strength, nucleosomes can open into half-nucleosomes and more extended, morphologically disorganized, structures. However, these structural transitions were readily reversible, and the basic repeat structure of the chromatin was preserved upon micrococcal nuclease digestion, even when the highly ordered nucleosomal appearance had disappeared. Recent findings relating to the nucleosomal structure of the SV40 minichromosome are summarized. (35 refs.)

78-5258 **Stimulation of Ribosomal RNA Synthesis in Isolated Nuclei and Nucleoli by Partially Purified Preparations of SV40 T Antigen.** (Eng) Baserga, R. (Fels Res. Inst., Temple Univ. Sch. Medicine, Philadelphia, PA, 19140); Ide, T.; Whelly, S. *Cold Spring Harbor Symp Quant Biol* 42(part 2): 685-691; 1978.

The ability of partially purified simian virus 40 (SV40) tumor (T)-antigen preparations to stimulate RNA synthesis in the isolated nuclei and nucleoli of resting rat liver cells was determined. Addition of T-antigen preparations isolated from SV40-transformed cells to the nuclei and nucleoli resulted in the stimulation of ribosomal RNA (rRNA) synthesis; most of the stimulation was due to an increased synthesis of rRNA precursors. Isolated nuclei had to be preincubated with T-antigen preparations for at least 20 min to obtain a good stimulation, but a 5-min preincubation was sufficient to produce a max stimulatory effect in nucleoli. These findings combined with those in the literature make SV40 T antigen the virally directed counterpart of the hypothetical protein that is induced by serum stimulation and that activates rRNA synthesis in density-inhibited normal cells stimulated to proliferate. These findings could also explain, at least partly, the mechanism of viral transformation. T antigen is expressed in SV40-transformed cells, and it is hypothesized that its presence continually stimulates the nucleolus of the transformed cells. This continuous interaction would keep the transformed cells from entering G_0 . The cells would then continue to grow until the nutritional conditions became such that they could not sustain growth and cell death would result. (66 refs.)

78-5259 **Density-Dependent Regulation of Amino Acid Transport in SV40-Transformed 3T3 Cells (Meeting Abstract).** (Eng) Borghetti, A. F. (Istituto di Patologia Generale, Universita di Parma, Via Gramsci 14, I-43100 Parma, Italy); Piedimonte, G.; Tramacere, M.; Guidotti, G. G. *Hoppe Seylers Z Physiol Chem* 359(9): 1061-1062; 1978. (11 refs)

78-5260 **Synthesis and Degradation of Simian Virus 40 RNA Sequences Not Found in the mRNA Population.** (Eng) Fried, A. H. (Institut fur Virusforschung, Deutsches Krebsforschungszentrum, Im Neuenheimer Feld 280, 6900 Heidelberg 1, W. Germany). *Virology* 88(2): 286-299; 1978.

The synthesis and degradation of a class of RNA sequences, referred to as dRNA sequences, which are transcribed from the late DNA strand of simian virus 40 (SV40) along with late messenger RNA (mRNA), were analyzed. The RNA isolated from SV40-infected CV-1 monkey cells was pulse-chase labeled with ^3H -uridine, the total cellular RNA extracted, and the RNA hybridized against the late strands of restriction endonuclease fragments of SV40 DNA that specifically hybridize to dRNA or mRNA sequences. Minimum estimates of the relative rate of synthesis of the dRNA sequences, compared to that of the mRNA sequences, were obtained from the relative amounts of radioactivity in these sequences following 10 min of labeling. These minimum estimates ranged from 0.32 to 0.55 and depended both on the individual labeling experiment and on which DNA fragments were used to assay for the dRNA and mRNA sequences. With longer chase periods, the ratio of labeled dRNA to that of labeled mRNA decreased, suggesting that the former are less stable than the latter. Further studies indicated that the dRNA was being degraded. In contrast to a previous report, sedimentation analysis of pulse-labeled RNA in dimethyl sulfoxide did not reveal more than minor amounts of labeled multilength transcripts of the late DNA strand. (35 refs)

78-5261 **Experimental (JC Virus-Induced) Intraocular and Extraocular Orbital Tumors in the Syrian Hamster (Meeting Abstract).** (Eng) Ohashi, T. (Univ. Wisconsin, Madison, WI); ZuRhein, G. M.; Varakis, J. N.; Padgett, B. L.; Walker, D. L. *J Neuropathol Exp Neurol* 37(5): 667; 1978. (1 ref)

78-5262 **Minichromosome from BK Virus as a Template for Transcription In Vitro.** (Eng) Meneguzzi, G. (Faculte des Sciences, Centre de Biochimie, Nice, France); Pignatti, P. F.; Barbanti-Brodano, G.; Milanesi, G. *Proc Natl Acad Sci USA* 75(3): 1126-1130; 1978.

BK virus (BKV) DNA was isolated from infected Vero cells

as a nucleoprotein complex containing about 20 nucleosomes and studied. This "minichromosome" had an *av* sedimentation coefficient of 55S-60S; no naked DNA was seen in any of the gradient fractions. No endogenous RNA synthesis activity was detected in the absence of added RNA polymerase. Addition of this polymerase indicated that both initiation and elongation of RNA chains are reduced by the presence of nucleosomes. Digestion of BKV DNA by the restriction endonuclease *Hind*III into four segments and hybridization analysis of the RNA indicated that with superhelical DNA as the template, all regions of the BKV genome were equally transcribed. With the minichromosome as the transcription template, however, transcription occurred preferentially on the largest fragment. No increase in strand selection was observed with respect to the transcription of purified superhelical BKV DNA. (39 refs)

78-5263 Ependymomas, Malignant Tumors of Pancreatic Islets, and Osteosarcomas Induced in Hamsters by BK Virus, a Human Papovavirus. (Eng) Corallini, A. (Inst. Microbiology, Univ. Ferrara, Via Luigi Borsari 46, I-44100 Ferrara, Italy); Altavilla, G.; Cecchetti, M. G.; Fabris, G.; Grossi, M. P.; Balboni, P. G.; Lanza, G.; Barbanati-Brodano, G. *J Natl Cancer Inst* 61(3): 875-883; 1978.

BK virus (BKV), a human papovavirus, was injected iv into 73 3-wk-old Syrian golden hamsters. Between 2.5 and 9 mo after inoculation, 60 of the animals developed tumors. The induced neoplasms were ependymoma (47 animals), carcinoma of the pancreatic islets (8), osteosarcoma (6), adenocarcinoma (1), angiosarcoma (1), angioma (1), lymphoma (1), and seminoma (1). Some of the animals had multiple tumors. Hypersecretion of insulin, glucagon, C peptide, and calcitonin was detected in the pancreatic islet tumors. A BKV etiology of the tumors was supported by the following evidence: (1) No tumors with BKV-specific markers appeared in animals given injections of buffer, in animals inoculated with BKV neutralized by anti-BKV-specific serum, or in uninoculated controls; (2) BKV tumor (T) antigen was detected by immunofluorescence and complement fixation tests in tumors of animals inoculated with infectious BKV and in transplanted tumors; (3) antibodies to BKV T antigen were detected in sera of animals bearing primary or transplanted tumors; (4) BKV could be activated by Sendai virus-mediated fusion of neoplastic cells with susceptible Vero cells; and (5) no endogenous hamster oncornaviruses were found in the tumors. (38 refs.)

78-5264 Polyoma and Other Virus Infections Following Bone Marrow Transplantation for Aplastic Anaemia. (Eng) Henry, K. (Bone Marrow Transplant Team, Westminster Hosps., London, England). *Pathol Biol (Paris)* 26(1): 55-56; 1978.

The case of a 16-yr-old girl with severe aplastic anemia whose course was complicated by graft-vs-host reaction (GVHR)

and fluctuating mental symptoms after a bone marrow transplant from a compatible sister is presented. Polyoma virion and oncornaviruslike particles were discovered via negative stain of the urine sediment. The polyoma virus was of the BF strain. Comprehensive virological screening is suggested both donor and recipient. (3 refs)

78-5265 A Mouse Embryo Cell Line Carrying an Infectible, Temperature-sensitive, Polyoma Virus Genome. (Eng) Bourgaux, P. (Departement de Microbiologie, Faculte de Medicine, Centre Hospitalier Universitaire de Sherbrooke, Sherbrooke, Quebec, Canada J1H 5N4); Delbecchi, L.; Yu, K. K.; Herring, E. *Virology* 88(2): 348-360; 1978.

Transformed (Cyp) cells were obtained by exposing secondary cultures from whole albino Swiss mouse embryos to temperature-sensitive (ts) mutant of polyoma (Py) virus (ts-P155, 1 plaque-forming unit/cell) at the restrictive temperature (38 C). At this temperature, ts-P155 transforms but replicates poorly. The cultures were then passaged at 39 C in the presence of Py virus antiserum until sufficient cell degeneration occurred (approx 4 mo later) to allow detection of foci of cells exhibiting lack of contact inhibition. One such focus gave rise to the Cyp strain, which was cloned after passages in vitro. No significant differences in phenotype were observed among any of the clones or between cloned and uncloned cells. At 39 C, they displayed several properties of transformed cells and, as indicated by indirect immunofluorescence, were negative for virus antigen and clearly positive for tumor (T) antigen. They did not produce detectable amounts of viral DNA and they produced only low amounts of viral hemagglutinin, although they were still fully permissive for Py virus replication. Exposure of Cyp cells to nonrestrictive temperature (33 C) resulted in an extensive cytopathic effect within 30-34 hr that was concurrent with the production of large amounts of infectious, ts, Py virus. T-antigen synthesis was even more pronounced. Covalently closed viral DNA molecules sedimenting at 53S were synthesized; max levels were reached 32-34 hr after transfer to 33 C. Cleavage by *Hpa*II yielded fragments characteristic of parent virus strain of ts-P155. Py virus antiserum neither curtailed the cytopathic effect nor depressed viral DNA synthesis, although it suppressed virus production. The highly efficient induction of virus replication by the Cyp cells at 33 C and the temperature sensitivity of the rescued virus suggest that Cyp cells are not carrying a revertant of the ts-P155 genome. (49 refs)

78-5266 Neoplastic Transformation of Hamster Glial Cells by Polyoma Virus (Meeting Abstract). (Eng) Hassoun, J. (Laboratoire de Neuropathologie, Faculte de Medecine, Marseille, France); de Micco, C.; Tripiet, F.; Toga, M. *J Neuropathol Exp Neurol* 37(5): 624; 1978. (4 refs)

78-5267 **The Frequencies of Transcription from the E- and L-Strands of Polyoma DNA.** (Eng) Ar-Shavit, R. (Dept. Genetics, Weizmann Inst. Science, Rehovot, Israel); Laub, O.; Aloni, Y. *J Gen Virol* 39(2): 37-360; 1978.

Exhaustion-type hybridization was used to measure the amount of nuclear virus RNA complementary to the early (E) and late (L) polyoma virus DNA strands. 3T-6 cells were infected with 2 ml of stock virus (10^6 plaque-forming units/ml), and the infected cells were labeled at 30 hr postinfection with ^3H -uridine. At 36 hr postinfection, 2.5%-7.3% of the newly synthesized virus RNA was complementary to the E-strand (-strand) and 92.7%-97.5% was complementary to the L-strand (+strand). This proportion was independent of labeling time, indicating similar accumulation of the E-RNA and L-RNA transcripts in the nucleus. These findings indicate that the transcription of polyoma virus DNA is regulated in a manner similar to that of simian virus 40 (SV40). Thus, at least two control mechanisms are at work. One is at the transcriptional level and it determines the frequencies of transcription from each strand, and the second is at the posttranscriptional level and it determines the stability of the primary transcripts and the choice of specific fragments transported to the cytoplasm. (13 refs)

78-5268 **Cellular Alterations Dependent upon the Polyoma Virus Hr-t Function: Separation of Mitogenic from Transforming Capacities.** (Eng) Schlegel, R. (Dept. Pathology, Harvard Medical Sch., Boston, MA, 02115); Benjamin, T. L. *Cell* 14(3): 587-599; 1978.

The role of the hr-t polyoma virus gene in inducing properties of transformation was explored by studying the early phase of nonproductive infection in cells infected by wild-type virus, hr-t mutants, and ts-a mutants. Within the first 24 hr after infection by wild-type or ts-a mutant virus, a transformed phenotype consisting of stellate cell shape, loss of organized cytoplasmic actin cables, nuclear and nucleolar enlargement, and behavioral changes such as cellular underlapping became evident. Cells infected by hr-t mutants underwent none of these changes and remained morphologically indistinguishable from mock-infected cultures. Wild-type virus and hr-t mutants induced cellular DNA synthesis in confluent monolayers of rat cells beginning 12-14 hr postinfection. There was a virus-mediated transit of cells from the G1 to the S and G2 phases of the cell cycle and an increase in the proportion of cells with an octaploid DNA content. Infection by wild-type virus gave rise to nearly 50% polyploid clones, most of which were not stably transformed. Less than 5% of the clones derived from the mock-infected and hr-t mutant-infected cultures were polyploid, and no transformants were found. The hr-t mutants promoted only a single round of cell division, whereas the wild-type virus and ts-a mutants promoted multiple rounds. Thus, loss of the hr-t function in polyoma virus results in a residual viral mitogenic activity

but prevents the virus from effecting the morphological transformation of cells. (46 refs.)

78-5269 **Detection of Airborne Polyoma Virus.** (Eng) McGarrity, G. J. (Inst. Medical Res., Camden, NJ); Dion, A. S. *J Hyg (Camb)* 81(1): 9-13; 1978.

The detection of airborne polyoma virus in an animal laboratory housing mice infected with the virus is described. Air samples were obtained by a high-volume air sampler, and the samples were concentrated by high-speed centrifugation. Polyoma virus was assayed by mouse antibody production tests, and it was found in 4/6 samples. (8 refs.)

78-5270 **Absence of C-Type Virus Production in Human Leukemic B Cell, T Cell and Null Cell Lines.** (Eng) Ogura, H. (Dept. Virology, Cancer Inst., Okayama Univ. Medical Sch., Okayama 700, Japan); Hiraki, S.; Omura, S.; Miyoshi, I.; Oda, T. *Acta Med Okayama* 32(2): 169-172; 1978.

Electron microscopy and enzymatic and biological techniques were used to determine whether C-type RNA virus is produced in human leukemic B-cell (BALL-1), T-cell (TALL-1), and null cell (NALL-1) lines. The results of the electron microscope studies and of an assay for reverse transcriptase activity in the culture supernatants were negative. Next, an attempt was made to amplify any possible human leukemia virus in the cells by cocultivation of the three lines with a bat lung cell line (BAT/88), which propagates primate C-type virus well. After 3 wk of cocultivation, the culture supernatants were again examined for reverse transcriptase activity and the cells were tested for syncytia formation in XC, KC, and RSb cells. All of these tests, except for those using a positive control (simian sarcoma virus-simian sarcoma-associated virus), were negative. It is concluded that the three human leukemic cell lines do not produce C-type RNA virus, although it is possible that the virus is present in a masked form in these lines. (16 refs.)

78-5271 **Analysis of RNA Tumor Virus Genomes.** (Eng) Haseltine, W. A. (Sidney Farber Cancer Inst., Boston, MA, 02115); Kleid, D. G.; Pedersen, F. S. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 437-442; 1977.

RNA tumor viruses were divided into four groups according

to the length of the runoff product (strong-stop DNA) produced by transcription of 70S viral RNA's with avian myeloblastosis virus reverse transcriptase. Group 1 comprised murine, woolly gibbon, and feline viruses; Group 2, endogenous baboon viruses and endogenous feline viruses; Group 3, chicken viruses; and Group 4, murine mammary tumor virus. On the basis of the oligopyrimidine composition of the strong stop DNA, the murine, woolly gibbon, baboon, and feline RD114 viruses were then divided into subgroups that show the extent of sequence divergence within the main groups. Preliminary data are also presented on the T₁-resistant oligonucleotides shared by viruses isolated from preleukemic or leukemic AKR mice. The viruses are AKV-1, AKV-2, AKR-247 CN-tropic xenotropic virus from a preleukemic mouse, and MB-34, a similar isolate from a mouse partially congenic for the AKV-Z genomes. (9 refs)

- 78-5272 **Abolition of Lymphocyte Blastogenesis by On-cornaviral Components (Meeting Abstract).** (Eng) Hellman, A. (VOP, NCI, Frederick, MD); Fowler, A, K.; Twardzik, D. R.; Weislow, O. S. *J Reticuloendothel Soc* 23(6): 37i; 1978. (no refs)

- 78-5273 **Relationship Between A-Type and C-Type Particles in Ehrlich Ascites Tumor Cells.** (Eng) Chew, E. C. (Dept. Anatomy, Univ. Hong Kong, Hong Kong). *Experientia* 34(8): 1086-1088; 1978.

Electron microscopy of Ehrlich ascites tumor cells obtained from the peritoneal fluid and im implants of mice carrying the tumor revealed intracisternal A-type particles in the cytoplasm. On rare occasions, intracytoplasmic A particles were also detected. In addition, mature and immature C-type particles were observed in some tumor cells. Many mature C-type particles were found only in the intercellular space. (14 refs.)

- 78-5274 **Search for Infective Mammalian Type-C Virus-related Genes in the DNA of Human Sarcomas and Leukemias.** (Eng) Nicolson, M. O. (Div. Infectious Diseases and Virology, Children's Hosp., Los Angeles, CA); Gilden, R. V.; Charman, H.; Rice, N.; Heberling, R.; McAllister, R. M. *Int J Cancer* 21(6): 700-706; 1978.

Transfection techniques were used to study the infectivity of proviral DNA's of C-type murine, gibbon, and baboon viruses in human and dog cells. DNA's from four gibbon tissues (liver, lung, skin-muscle, and kidney) and from baboon embryo fibroblast strains BEF No. 1 and BEF No. 2 did not transfect TE-32 human sarcoma cells or D-17 dog cells. DNA's from two baboon placentas and from BEF No. 3, containing 100 viral genome equivalents per haploid cellu-

lar genome, transfected D-17 but not TE-32 cells. TE-32 and D-17 cells experimentally infected with baboon virus contained only five genome equivalents per haploid cellular genome, but they released high titers of infectious virus. The DNA's of these cells contained baboon proviruses infectious for both TE-32 and D-17 cells. Proviruses were not detectable in any of the DNA's of human sarcoma or leukemia cells tested. (46 refs)

- 78-5275 **DNA Viruses in Urine after Renal Transplantation.** (Eng) Lecatsas, G. (Dept. Microbiology, Inst. Pathology, Univ. Pretoria, P.O. Box 2034, Pretoria 0001, Republic South Africa); Van Wyk, J. A. *S Afr Med J* 53(20): 787-788; 1978.

Over a 5-yr period, approx 2,500 urine specimens from 96 renal transplants were examined for virus particles. Electron microscopy of negatively stained preparations showed that only three virus groups were present: polyoma virus (36 patients), herpesvirus (10), and adenovirus (11). The failure to detect RNA viruses suggests that they are absent or may require different means of activation. (13 refs)

- 78-5276 **Localization of Hepatitis B Surface Antigen in Neoplastic Cells of Primary Hepatocellular Carcinoma (Meeting Abstract).** (Eng) Murugasu, R. (Royal Newcastle Hosp., Newcastle, England). *Pathology* 10(2): 196-197; 1978. (no refs)

5277 **Sequence Variation among Heavy Chains from Inulin-binding Myeloma Proteins.** (Eng) Vrana, J. (Lab. Cell Biology, NCI, NIH, Bethesda, MD, 20014); Udikoff, S.; Potter, M. *Proc Natl Acad Sci USA* 75(4): 1957-1961; 1978.

The entire sequences of the variable region of three heavy chains from BALB/c mouse inulin-binding myeloma proteins were determined by cyanogen bromide cleavage and Sephadex G-100 column chromatography. (The variable region sequence of the fourth inulin-binding protein has been described previously). Among the four proteins there are six amino acid differences, all of which occur in the framework portion of the variable region. All of the six amino acid substitutions can be explained by single base mutations at the DNA level. Unlike the previously described group of heavy chains from phosphorylcholine-binding myeloma proteins, which (with the exception of 2 that are identical) have size and sequence differences in their complementarity regions, the inulin-binding heavy chains all have identical complementarity regions, with H3 being extremely short. The pattern of variation observed in the antiinulin heavy chains appears to be most easily explained by a somatic mutation mechanism. However, because none of the substitutions occur in complementarity-determining regions, they presumably could have no selective advantage and would not alter binding specificity. Five of the six sequence differences occur at positions that are internal in the molecule and thus do not account for previously observed idiotypic differences. The results suggest that most of the idiotypic cross-reactivities are due to differences in the light chains of the antiinulin proteins. (35 refs)

5278 **Shedding of Anti-IgM from Antibody-coated Human Lymphoma Lines: Differences Between Epstein-Barr Virus-negative Lines and Their Virus-converted Sublines.** (Eng) Yefenof, E. (Dept. Tumor Biology, Karolinska Institutet, S-104 01 Stockholm 60, Sweden); Lundin, M.; Klein, G. *Eur J Immunol* 8(3): 190-193; 1978.

Surface IgM-positive, Epstein-Barr virus (EBV)-negative cell lines and their virus-converted sublines were compared with respect to their ability to shed membrane-bound anti-IgM antibodies, in order to investigate the membrane changes that accompany EBV conversion. The in vitro converted EBV genome-carrying sublines showed a reduced ability to shed surface-bound antibodies, compared with their EBV-free parental B lymphoma lines. On the other hand, antibody degradation was higher in the EBV-positive cells. Analysis of antibody-coated single cells indicated that the cells that failed to

redistribute or cap their ligand-bound surface components at 37 C were also capable of shedding bound antibody. This finding suggests that capping and shedding are essentially independent phenomena. However, both may be influenced by the same virally induced change in the lateral mobility of membrane constituents. (33 refs)

5279 **Further Characterization of the Suppressor Cells, Activated by Goat Anti-Th-B Antibody Reagent and Responsible for Enhanced Growth of Sarcoma 180 in AKR Mice.** (Eng) Kakimoto, K. (Dept. Immunology Res., Roswell Park Memorial Inst., Buffalo, NY, 14263); Fuji, H.; Grossberg, A. L.; Pressman, D. *Cell Immunol* 37(1): 61-76; 1978.

The suppressor cells that enhance the growth of allogeneic sarcoma 180 (S180) in AKR mice previously inoculated with goat anti-Th-B antibody reagent (prepared from antiserum raised in goats against BALB/c myeloma MOPC 104E cells and purified) were characterized. These suppressor cells were T cells, since they carried the Thy-1 antigen on their surface. Treatment of the cells in vitro with anti-Thy-1 plus complement eliminated their ability to enhance tumor growth. The T cells responsible for tumor enhancement did not carry the Th-B determinant. Treating T cells in vitro with goat anti-Th-B antibody reagent plus complement did not abrogate their tumor-enhancing activity. This suggests that the suppressor T cells involved in tumor enhancement are generated by the interaction of anti-Th-B antibodies with precursor suppressor cells that do carry Th-B. Once generated, the active suppressor cells lose the Th-B antigen; this contention is supported by the finding that the thymic precursors of concanavalin A (Con A)-inducible suppressor cells bear Th-B, since they are killed by anti-Th-B plus complement, but active suppressor cells induced by Con A do not carry Th-B, since they are not killed by anti-Th-B plus complement. Evidence also indicated that nonthymic suppressor cell precursors could be activated by anti-Th-B, since spleen cells from anti-Th-B-treated thymectomized mice bearing S180 can transfer the tumor-enhancing effect. It is concluded that mature suppressor cells cause tumor enhancement by suppressing the immune mechanisms responsible for allogeneic tumor graft rejection. (25 refs)

5280 **Preleukemic AKR Mice are Selectively IgA-deficient (Meeting Abstract).** (Eng) Mink, J. G. (Dept. Cell Biology and Genetics, Erasmus Univ., Rotterdam, Netherlands); van Muiswinkel, W. B.; de Laat, A.

M.; Benner, R. *Z Immunitaetsforsch* 54(4): 339; 1978. (3 refs)

78-5281 Demonstration of Humoral Antibodies Against Autologous Glioma Cells in Experimental Rabbit Gliomas (Meeting Abstract). (Eng) Stavrou, D. (Inst. Veterinary Pathology, Univ. Munich, Munich, W. Germany); Elling, H.; Haglid, K. G. *J Neuropathol Exp Neurol* 37(5): 694; 1978. (no refs)

78-5282 Specific Suppression of Antigen-reactive Cells in Neonatal Transplantation Tolerance. (Eng) Holan, V. (Inst. Molecular Genetics, Czechoslovak Acad. Sciences, Flemingovo nam. 2, 166 10 Prague 5, Czechoslovakia); Chutna, J.; Hasek, M. *Nature* 274(5674): 895-897; 1978.

Immunological reactivity was analyzed in two experimental systems of neonatal tolerance in mice involving a disparity in either the entire *H-2* complex or the *D* region only. Tolerance to different strains was induced by inoculation of B10.A newborn mice with spleen cells from B10 or B10.A(2R) mice within 20 hr of birth. The B10.A mice receiving the neonatal injections were grafted with allogeneic skin at 8-9 wk of age, and those that did not reject the graft were used in the study. In each case, cells from the tolerant mice could not be sensitized in vitro to the tolerated antigens. These cells, which were nonreactive in the in vitro cytotoxicity test, could be separated into nonadherent and nylon-adherent cells according to adherence to nylon wool. The former cells, when sensitized in vitro, were cytotoxic, as were the sensitized cells from normal animals. The latter remained nonreactive even after sensitization and, in addition, specifically inhibited the in vitro sensitization of cells from normal animals. (10 refs.)

78-5283 B and T Lymphocyte Activation by Murine Leukemia Virus Infection (Meeting Abstract). (Eng) Bendinelli, M. (Inst. Microbiology, Pisa, Italy); Friedman, H. *J Reticuloendothel Soc* 23(6): 28i; 1978. (no refs)

78-5284 Alveolar Macrophage-Splenic Lymphocyte Interactions Following Chronic Asbestos Inhalation in the Rat (Meeting Abstract). (Eng) Kagan, E. (Dept. Pathology, Georgetown Univ. Medical Center, Washington, DC); Miller, K. *J Reticuloendothel Soc* 23(6): 24i; 1978. (no refs)

78-5285 Ultraviolet Light Induced Murine Suppressor Lymphocytes Dictate Specificity of Anti-ul-

traviolet Tumor Immune Responses. (Eng) Spellman, C. V. (Dept. Pathology, Univ. Utah Medical Center, Salt Lake City, UT, 84132); Daynes, R. A. *Cell Immunol* 38(1): 25-31; 1978.

Suppressor lymphocyte activity was investigated in UV-irradiated female C3Hf/HeN mice. When normal mice were hyperimmunized with the regressor UV tumor RD 87 and then challenged with progressor tumors, they exhibited cross-protective immunity. However, the adoptive transfer of spleen cells from tumor immune mice into UV-irradiated hosts resulted in tumor-specific immunity. Transfer of 3.5 x 10⁷ nylon-nonadherent spleen cells from UV-treated mice into RD 87-hyperimmune animals did not inhibit the antitumor responses. However, when the same number of nylon-nonadherent UV spleen cells were transferred into normal mice, the animals were unable to reject RD 87. It was then determined whether mice hyperimmunized with a regressor tumor were capable of rejecting a secondary UV regressor tumor following x-radiation-induced block of primary immune responses. The 450-R dose selected rendered normal mice tumor-susceptible, but it did not inhibit the antitumor responses of hyperimmunized mice to challenge with either of two regressor tumors. Cross-reactive immunity was unaffected in the immune animals. The suppression of antitumor responses in hyperimmune mice could be effected by adoptive transfer of suppressor cells from UV-treated mice. These results are consistent with the suggestion that UV-induced suppressor cells inhibit antitumor responses directed against the common cross-protective tumor-associated antigens. Although many immune interactions may be involved in the antitumor response, ultimately, effective tumor rejection may be mediated by cells subject to suppressor control. In vivo, suppression may dominate and cause potential effector cells directed against the common antigens remain as committed preeffector cells. In vitro, suppression may be broken, and cross-reactivity may occur as a result of the survival and expression of effector cells directed against the common antigenic determinants. (44 refs)

78-5286 Use of Stimulating Capacity of Mixed Lymphocyte Reaction (MLR-S) as a Possible Marker for the Cell-Origin of Null-Cell Acute Lymphoblastic Leukaemia. (Eng) Han, T. (Roswell Park Memorial Inst. New York State Dept. Health, 666 Elm St., Buffalo, NY 14263); Minowada, J. *Immunology* 35(2): 333-339; 1978

The stimulating capacity of the mixed lymphocyte reaction (MLR-S) of fresh leukemic cells from eight patients with null cell acute lymphoblastic leukemia (ALL) and cultured leukemic null cells from five different cell lines was investigated. A one-way MLR conducted with a 1:1 ratio of responder cells to stimulating cells revealed that the cells of five of the patients were good stimulators but those of the other three were poor. No stimulating capacity of leukemic cells was observed in one of the poor stimulators, even though high numbers of stimulating cells were employed. Three of

ltured leukemic cell lines consistently exerted a very strong stimulation on allogeneic lymphocytes. Another line exhibited a lesser but significant stimulation of one-way MLR. No significant MLR-S of the fifth cell line was observed, even with the use of higher numbers of cells. These findings suggest that leukemic null cells possessing and not possessing a stimulating capacity may represent early B- and T-cell precursors, respectively. (28 refs.)

78-5287 **Functional Heterogeneity and T-Cell Dependent Activation of Macrophages from MSV-induced Tumors (Meeting Abstract).** (Eng) Holden, H. T. (Lab. Immunodiagnosis, NCI, Bethesda, MD, 20014); Varesio, L.; Imaniyama, T. *J Reticuloendothel Soc* 23(6): 39i; 1978. (no refs)

78-5288 **Enhancement of the Growth of a Chemically-induced Fibrosarcoma by Peritoneal Macrophages of Normal and Tumor-bearing Mice (Meeting Abstract).** (Eng) Gabizon, A. (Dept. Cell Biology, Weizmann Inst. Science, Rehovot, Israel); Trainin, N. *J Reticuloendothel Soc* 23(6): 38i; 1978. (no refs)

78-5289 **Idiotypic-specific Transplantation Resistance to MOPC-315: Abrogation by Post-immunization Thymectomy.** (Eng) Daley, M. J. (Immunobiology Labs., Univ. New Mexico, Sch. Medicine, Albuquerque, NM, 87131); Lynch, R. G.; Gebel, H. M. *J Immunol* 120(5): 1620-1624; 1978.

Studies were undertaken to determine if the thymus plays a role in idiotype-specific myeloma graft resistance. The idiotype-specific myeloma transplantation resistance induced in ALB/c mice by immunization with the DNP-binding IgA protein produced by plasmacytoma MOPC-315 was abrogated by postimmunization thymectomy. Sham thymectomy had no effect. The ablative effect of thymectomy was observed in mice challenged sc with MOPC-315 cells either 3 or 44 days after thymectomy. These observations imply that short-lived thymic effector contributes to idiotype-specific myeloma graft resistance. (34 refs)

78-5290 **Suppression of Host Immunity by Dietary Hypercholesterolemia (Meeting Abstract).** (Eng) Iorio, R. M. (Dept. Microbiology, Medical Coll. Virginia, Virginia Commonwealth Univ., Richmond, VA); Kos, W. L.; Kaplan, A. M. *J Reticuloendothel Soc* 23(6): 49i; 1978. (no refs)

78-5291 **Immunologic Basis of Tumor Dormancy (Meeting Abstract).** (Eng) Wheelock, E. F. (Dept. Microbiology, Thomas Jefferson Univ., Philadelphia, PA). *J Reticuloendothel Soc* 23(6): 38i; 1978. (no refs)

78-5292 **Role of Immunocompetence, Immunosuppression and Social Stress in Susceptibility of Chickens to Rous Sarcoma (Meeting Abstract).** (Eng) Misra, L. K. (Texas A&M Univ., College Station, TX, 77843). *Diss Abstr Int B* 39(1): 7B-8B; 1978. (no refs)

78-5293 **Nonimmune and Immune Surveillance. II. Effect of Recipient's Age, Tumor Immunogenicity, and Neonatal Thymectomy on Tumor Growth Inhibition.** (Eng) Zinzar, S. N. (Lab. Virology, Cancer Res. Center, Acad. Medical Sciences USSR, Moscow 115478, USSR); Svet-Moldavsky, G. J.; Karmanova, N. V. *J Natl Cancer Inst* 61(3): 737-745; 1978.

Suspensions of homogenates of Lewis lung adenocarcinoma, a mammary gland adenocarcinoma, rectal sarcoma SR-1-75, hepatoma H-2-73, and ovarian carcinoma OC-1-75 were implanted sc into newborn and adult (CBA x C57BL/6J)F₁ and BALB/c mice. The growth of the lung and the mammary tumor was inhibited in newborn mice, but the rectal sarcoma grew at the same rate in newborn and adult recipients. Neonatal thymectomy stimulated the growth of the hepatoma in newborns. The degree of tumor growth inhibition was age-dependent: max inhibition was observed in 1- to 6-day-old recipients, but later it gradually decreased. The hepatoma and the ovarian carcinoma, whose growth was inhibited in newborns, and the rectal sarcoma, whose growth was uninhibited, had equally low immunogenicity. The data suggest that newborns possess factors that inhibit tumor growth but that these factors disappear with the increased age of the recipients. (27 refs.)

78-5294 **Inhibition of Lymphoma Growth in the Spleen and Liver of Lethally Irradiated Mice.** (Eng) Iorio, A. (Inst. Hygiene, Chair Virology, Inst. Pharmacology, Univ. Perugia, 06100 Perugia, Italy); Campanile, F.; Neri, M.; Spreafico, F.; Goldin, A.; Bonmassar, E. *J Immunol* 120(5): 1679-1685; 1978.

The growth of murine lymphomas carrying the *H-2d* (LSTRA, P388), the *H-2K* (6C3HED), and the *H-2a* (LAF-17) haplotypes was studied in the spleen and liver of lethally irradiated (850-1,000 R) compatible or allogeneic mice by the ¹²⁵I-5-iodo-2'-deoxyuridine uptake method. Ac-

According to the genetic pattern of the *Hh* (hemopoietic-histocompatibility) system, recipient hosts were either susceptible or resistant to bone marrow grafts carrying the same haplotypes as those of the lymphomas studied. Tumor cell growth was impaired in the spleen of irradiated *Hh*-incompatible resistant hosts (eg, P388 into B10, B10.BR, and CBA), suggesting that lymphoma and bone marrow cells could share common *Hh* antigens. However, inhibition of lymphoma growth occurred also in the spleen of irradiated *Hh*-susceptible mice, either *Hh*-compatible (eg, P388 into B10.D2; 6C3HED into AKR or CBA) or *Hh*-incompatible, but genetically unable to respond against bone marrow cells (eg, LSTRA or P388 into C3H, or 6C3HED into BALB/c or SJL). In almost all cases, growth inhibition was absent or weak in the liver. These results suggest that lymphoma cells can express antigens not detectable on normal bone marrow cells that are capable of eliciting radioresistant inhibition of

tumors predominantly in the spleens of mice susceptible to marrow grafts of the same genetic origin as that of the lymphomas tested. (38 refs)

PATHOGENESIS

5295 **Characteristics of Preleukemia Cells Induced in Mice.** (Eng) Haran-Ghera, N. (Dept. Chemical Immunology, Weizmann Inst. Science, Rehovot, Israel); Boio, N.; Leef, F.; Goldstein, G. *Cell Immunol* 37(2): 314; 1978.

Studies were conducted to determine whether preleukemia cells produced by inoculation of mice with radiation leukemia virus (D-RadLV) are prothymocytes and whether virus-induced preleukemia marrow cells could immunize against transplanted leukemia cells induced by the same virus. A high proportion of irradiated C57BL/6 mice inoculated with D-RadLV developed overt T cell leukemias originating in the thymus; the incidence in unirradiated hosts was much lower. As early as 10 days after injection of D-RadLV, the bone marrow contained preleukemia cells that, although not frankly leukemic, would develop into leukemia cells upon transfer to a specially pretreated recipient mouse. The leukemia cells expressed the T-cell surface component Thy-1 (Thy-1+), but the preleukemia cells did not. However, the leukemia cells could be induced in vitro by thymopoietin and ubiquitin to become Thy-1+, suggesting that they are prothymocytes. Unlike leukemia cells, the injection of preleukemia cells into normal recipients immunized them against transplants of leukemias induced by the same D-RadLV. It is suggested that D-RadLV induces a critical change in prothymocytes that in a later (Thy-1+) phase of differentiation manifested in overt leukemia transformation. (18 refs)

5296 **Trisomy of 1q in Preleukaemia with Progression to Acute Leukaemia.** (Eng) Najfeld, V. (Medical Genetics, Veterans Admin. Hosp., 4435 Beacon Ave. S., Seattle, WA, 98105); Singer, J. V.; James, M. C.; Fialkow, P. J. *Scand J Haematol* 21(1): 24-28; 1978.

Chromosome studies were performed on the marrow cells of a 48-yr-old man with refractory megaloblastic anemia prior to treatment. The cells had trisomy for the long arm of chromosome 1, which was translocated to the long arm of chromosome 11 [46,XY,-11,+t(1:11)(q11 or q12; q25)]. The same abnormality was found in metaphases from eight individual granulocytic colonies. When the disease progressed into acute myelomonocytic leukemia 5 mo later, trisomy of chromosome 8 and an extra chromosome number 19 with partial deletion were found in addition to the basic chromosome abnormality. Fibroblasts grown from marrow biopsy material showed a normal diploid complement, 46,XY. (11 refs.)

78-5297 **Chronic Lymphocytic Leukaemia: Clonal Origin in a Committed B-Lymphocyte Progenitor.** (Eng) Fialkow, P. J. (Medical Service, Veterans Admin. Hosp., 4435 Beacon Ave. S., Seattle, WA, 98108); Najfeld, V.; Reddy, A. L.; Singer, J.; Steinmann, L. *Lancet* 2(8087): 444-446; 1978.

Determination was made of the glucose-6-phosphate dehydrogenase (G-6-PD) types of isolated blood cell populations and normal skin in two patients (patient 1, 47 yr old, patient 2, 72 yr old) with chronic lymphocytic leukemia (CLL) who were heterozygous at the G-6-PD locus. Approx equal amounts of B and A isoenzyme activities were found in skin specimens. In contrast, only type A G-6-PD was found in the B lymphocytes isolated from patient 1. The homogeneity of this preparation of B lymphocytes was confirmed by failure of the cells to stain with peroxidase or to form erythrocyte (E) rosettes, by the presence of monoclonal immunoglobulin (μ and λ) on the cell surfaces, and by the ability of the cells to form rosettes with mouse RBC. The B-cell preparation from patient 2 showed 5% type B and 95% type A G-6-PD. Peroxidase stains were positive in 3% of the cells, indicating some admixture with granulocytes or monocytes, which may have accounted for the small amount of B isoenzyme. These observations indicate that at the time of diagnosis, all CLL cells were descended from one progenitor; ie, the disease has a clonal origin. Approx equal amounts of B and A isoenzymes were found in granulocytes, RBC, platelets, and T lymphocytes. The findings indicate that CLL involves committed B-lymphocyte progenitors and that the disease differs from chronic myelocytic leukemia and other myeloproliferative syndromes, all of which involve multipotent hemopoietic stem cells. (23 refs.)

78-5298 **Acute Myelogenous Leukemia Subsequent to Therapy for a Different Neoplasm: Clinical Features and Response to Therapy.** (Eng) Preisler, H. D. (Dept. Medicine A, Roswell Park Memorial Inst., 666 Elm St., Buffalo, NY, 14263); Lyman, G. H. *Am J Hematol* 3: 209-218; 1977.

The case records of 10 patients with acute myelogenous leukemia (AML) who had initially been treated for other malignancies are reviewed. These patients differed from those with "spontaneous" AML in that, at the time of diagnosis, they had lower WBC counts, and significantly less cellular bone marrow, and a lower proportion of leukemic cells in the mar-

row. Eight of these patients initially had advanced primary hematologic malignancies with bone marrow involvement. Most had been exposed to extensive radiation therapy and combination therapy (including alkylating agents) during treatment for their primary malignancy. Extensive chromosome changes were observed in 9/10 patients, compared with only about 40% of previously reported spontaneous AML patients. Only 1/9 patients who received chemotherapy attained remission status, and the mean and median survivals from the initiation of chemotherapy were 2.7 and 1.0 mo, respectively. Nine of the 10 patients died as a result of infection. The refractoriness of this form of AML to chemotherapy was borne out by a review of the literature, which revealed only 2 remissions in 32 treated patients. Cytotoxic chemotherapy should be delayed as long as possible in patients whose AML developed subsequent to therapy for another malignancy. (28 refs)

- 78-5299 Non-Hodgkin's Lymphoma Terminating in Acute Myelogenous Leukemia.** (Eng) Cavalli, F. (Div. Oncology, Univ. Hosp., Inselspital, CH-3010 Bern, Switzerland); Sonntag, R. W.; Zimmermann, A.; Deubelbeiss, K.; Ryssel, H. J. *Acta Haematol (Basel)* 60(4): 250-256; 1978.

The case report of a 77-yr-old man who developed acute myelogenous leukemia (AML) 5.25 yr after a diagnosis of diffuse histiocytic lymphoma is presented. The disease was initially treated by radiotherapy; later, upon progression of the disease, moderately intensive chemotherapy was instituted. This therapy was given for almost 5 yr. In Hodgkin's disease, radiotherapy followed by chemotherapy has been shown to carry a high risk for developing a second neoplasia. At present, it is not known why AML is less frequently seen in patients with non-Hodgkin's lymphoma (NHL) than in those with Hodgkin's disease or multiple myeloma. However, since the survival of patients with disseminated NHL is much poorer than of those with Hodgkin's disease, the disease may not have time to develop. It is possible that the radiotherapy and chemotherapy of the NHL contributed to the development of AML in this patient. (31 refs.)

- 78-5300 Immunological Family Studies in Hodgkin's Disease. Is the Immunodeficiency Horizontally Transmitted?** (Eng) Bjorkholm, M. (Dept. Medicine, Serafi-merlasarettet, S-112 83 Stockholm, Sweden); Holm, G.; Mellstedt, H. *Scand J Haematol* 20(4): 297-305; 1978.

Lymphocyte reactivity was evaluated in 27 consanguineous and nonconsanguineous relatives of 12 patients with Hodgkin's disease (HD). The lymphocyte DNA synthesis induced by concanavalin (Con A) was depressed in six patients. Seven of 15 relatives (3 first-degree relatives and 4 spouses) showed a profound impairment in lymphocyte response to Con A

stimulation. None of 12 relatives of patients with a normal Con A response was defective in this respect. In relatives of Con A-hyporesponsive patients, there was a small but statistically significant decrease in lymphocyte response to pokeweed mitogen and purified protein derivative. No differences in lymphocyte subpopulations was seen. The findings suggest that an exogenous factor (probably a virus) depressed the mitogen response in relatives of immunodeficient HD patients. A possible link between the immunodeficiency and the pathogenesis of HD is discussed. (35 refs)

- 78-5301 Lymphoproliferative Disorders in Four Patients Receiving Chronic Diphenylhydantoin Therapy. Etiologic Correlation or Chance Association?** (Eng) Matzner, Y. (Dept. Hematology, Hadassah Univ. Hosp., Jerusalem, Israel); Polliack, A. *Isr J Med Sci* 14(8): 865-869; 1978.

The occurrence of angioimmunoblastic lymphadenopathy (AL), Hodgkin's disease, and IgG- λ multiple myeloma in four epileptic patients who had been receiving diphenylhydantoin (DPH) therapy is described, and current knowledge of the association between DPH therapy and the development of lymphoproliferative disorders is reviewed. The four patients were the only examples of lymphoplasma-cytic disorders encountered among 300 epileptics treated with DPH at one institution. Case 1, a 26-yr-old woman, was treated for 6 mo with DPH before she developed AL. Cessation of DPH therapy led to remission. Case 2, a 61-yr-old man had received DPH for several years before developing AL. He recovered slowly after cessation of the therapy. Case 3, a 22-yr-old man, developed Hodgkin's disease after treatment with DPH for 5 yr. Discontinuation of DPH and total nodal irradiation led to complete remission 1 yr later. Case 4, a 54-yr-old man treated with DPH for centrencephalic epilepsy for > 20 yr, developed IgG- λ multiple myeloma. Serum and urine immunoelectrophoresis showed the presence of an IgG λ paraprotein associated with combined IgA and IgM deficiency. Although the relation between DPH therapy and the development of immunosuppression and lymphoma is well-documented, the role of the drug in the etiology of these disorders is still controversial. A chance association between ingestion of the drug and the development of lymphoma cannot be excluded in these cases. (27 refs.)

- 78-5302 Multiple Myeloma Terminating in Lymphocytic Leukemia with B-Lymphocyte Membrane Markers.** (Eng) Polliack, A. (Dept. Hematology, Kirya Hadassah, Hadassah Univ. Hosp., Jerusalem, Israel); Napavsky, E.; Eldor, A.; Zvi, A. B.; Bassat, H. B.; Zlotnik, A. *Am J Hematol* 3: 153-163; 1977.

The development of an acute, rapidly fatal lymphocytic leukemia in a 64-yr-old woman with kappa light-chain multiple myeloma is reported. The leukemia developed after a relative

short period of Alkeran therapy and it was fulminant, with a rapidly fatal course associated with a bleeding diathesis. Most of the leukemic cells lacked the ultrastructural features of plasma cells but bore readily detectable B-cell markers and resembled lymphocytes at various stages of maturation under light, transmission, and scanning electron microscopes. This case might best be regarded as one of myeloma developing as a variant of plasma cell leukemia in which the leukemic cells were lymphocytic in type, suggesting a degree of dedifferentiation of the original malignancy. It appears to represent the first case of myeloma terminating as lymphocytic leukemia in which the leukemic cells were not obviously of plasmacytic type. (28 refs.)

5303 Chronic Lymphocytic Leukemia (CLL) Terminating in Multiple Myeloma: Report of Two Cases. (Eng) Kough, R. H. (Dept. Hematology-Oncology, Hershey Medical Center, Danville, PA, 17821); Makary, A. *Blood* 52(3): 532-536; 1978.

The late development of multiple myeloma (MM) in two elderly patients with chronic lymphocytic leukemia (CLL) is reported. In the first case, an 81-yr-old woman developed symptoms of MM 80 mo after the onset of CLL, and a diagnosis of IgA κ MM was made 95 mo after the diagnosis of CLL. In the second case, a 74-yr-old man developed symptoms of MM 169 mo after the onset of CLL, and a diagnosis of κ light-chain MM was made 192 mo after the diagnosis of CLL. Both cases of CLL had responded well to intermittent low-dose chlorambucil therapy, but the MM failed to respond to cyclic melphalan-prednisone therapy. The finding of 2 cases of MM late in the course of CLL among 105 cases of CLL reviewed remains unexplained. However, the suggestion that there is an increased prevalence of MM in CLL is an attractive one, because both are B-cell neoplasms and because of the increased frequency of asymptomatic monoclonal gammopathies in CLL found by others. (21 refs.)

5304 Lodgement and Extravasation of Tumour Cells in Blood-Borne Metastasis: An Electron Microscopic Study. (Eng) Kinjo, M. (Dept. Pathology, Faculty of Medicine, Kyushu Univ., 3-1-1 Maidashi, Higashi-ku, Fukuoka 812, Japan). *Br J Cancer* 38(2): 293-301; 1978.

The lodgement and extravasation of ascites hepatoma cells with high (line AH 130) and low [line AH 130 F(N)] thromboplastic (TP) activities were examined by electron microscopy, and the role of TP activity in blood-borne metastasis was determined. Donryu rats were inoculated with 10^7 cells/0.2 ml cell suspension of either line and sacrificed immediately or at times up to 72 hr postinjection. Soon after injection, three types of tumor cell emboli were found in the arterioles and capillaries of the lung. Type I, which had marked aggregation of platelets and deposition of fibrin, was

mainly seen when AH 130 cells were injected. Detachment and fragmentation of endothelial (ET) cells resulted 1-3 hr later. Type II had loosely aggregated platelets, and type III had no aggregation of platelets or fibrin deposition. Types II and III were mainly seen with AH 130 F(N) cells, and they did not accompany severe structural changes in the ET cells. Tumor cell-platelet complexes appeared to be induced by TP released from tumor cells rather than that from the detached ET cells. At 1-6 hr after infection of AH 130, tumor cells were found beneath the detached ET cells in areas with microthrombi. In places, tumor cells closely adhered to and extended cytoplasmic processes to the ET cells, forming indentations. Some of these processes penetrated through the defects to the subendothelial space, with desquamation of the ET cells. At 3 days, the number of metastatic foci was higher in rats inoculated with AH 130. The results indicate that TP activity is important to microthrombi formation in the lodgement phase and that it might be one of the factors facilitating blood-borne metastasis. When tumor cells had low TP activity, there were no severe structural changes of the ET cells in the entire process from lodgement to extravasation. (20 refs.)

78-5305 Cutaneous Angiosarcoma of the Hip. (Eng) Zdrojewski, J. F. (Beaver Medical Clinic, Redlands, CA); Hodge, S. J.; Scheen, S. R.; Owen, L. G.; Smith, S. Z. *South Med J* 71(8): 914-916; 1978.

The case of a 63-yr-old woman who developed a fatal angiosarcoma of the hip approx 10 yr after surgery and intracavitary irradiation (8,000 rads) for a uterine malignancy is reported. The patient was treated by lymph node dissection and cobalt irradiation (5,000 rads) for right inguinal metastases 4 yr after the initial treatment. She then developed aseptic necrosis of both femoral heads, resulting in a right hip replacement procedure with an Austin-Moore prosthesis. The latter was removed after 5 yr because of continuing purulent drainage from the hip incision. Mild chronic lymphedema of the involved lower extremity was noticed after insertion of the prosthesis, and the patient eventually developed a cutaneous angiosarcoma in the affected area. Any or all of these factors might bear a causative relationship to the angiosarcoma, but none can be proved. (27 refs.)

78-5306 Leiomyomatosis Peritonealis Disseminata. An Ultrastructural Study. (Eng) Nogales, F. F. (Dept. Pathology, Univ. Seville Medical Sch., Seville, Spain); Matilla, A.; Carrascal, E. *Am J Clin Pathol* 69(4): 452-457; 1978.

A histopathologic study was made of the eighth documented case of leiomyomatosis peritonealis disseminata (LPD), a disease that consists of widespread peritoneal nodules of a leiomyomatous nature that involve the parietal peritoneum and the visceral peritoneum of abdominal organs. Six of the pa-

tients were black women. LPD was discovered upon cesarean section in an asymptomatic 34-yr-old Caucasian woman admitted for delivery. The main proliferative cell type was identified by electron microscopy as a mature leiomyocyte with all its associated ultrastructural features. Intimately intermixed with the leiomyocytes were hybrid cells (myofibroblasts) in which fibroblastic features such as abundant rough endoplasmic reticulum, prominent Golgi complexes, and a resemblance to stromal collagen fibers coexisted with variable amounts of cytoplasmic fibrils and electron-dense bodies. The histogenesis of LPD is considered to be a multicentric metaplastic change determined by an abnormal tissue response to elevated hormone levels associated with pregnancy or ovarian functioning tumors. The cell of origin is suggested to be the multipotent subcelomic mesenchymal cell, which retains the capacity to differentiate into several tissues, among them smooth muscle. The myofibroblasts may represent a transitional morphologic stage between the mesenchymal cell and the smooth muscle proliferating cell. (17 refs)

78-5307 Malignant Fibrous Histiocytoma. Clinicopathologic and Ultrastructural Study of 12 Cases.

(Eng) Limacher, J. (Dept. Pathology, Hotel-Dieu de Quebec, Quebec City, Quebec G1R 2J6, Canada); Delage, C.; Lagace, R. *Am J Surg Pathol* 2(3): 265-274; 1978.

The light and electron microscopy findings for 12 specimens of malignant fibrous histiocytoma of the soft tissues are presented. The specimens were removed from eight men and four women aged 29-88 yr. All primary lesions were located in the lower extremities except for one in the arm, one in the thoracic wall, and one in the jaw. Nine patients had recurrences and three had metastases. The essential histologic feature of this tumor was its biphasic composition, with both histiocytic and fibroblastic components, the latter showing a characteristic storiform pattern. Furthermore, a variable number of xanthomatous cells and multinucleated giant cells were observed. Ultrastructural studies confirmed the presence of these cells and also revealed the existence of a primitive mesenchymal cell. It is suggested that this mesenchymal cell is a primitive histiocyte from which the two main cellular components of the tumor arise. The precise significance of the xanthomatous cells and the multinucleated giant cells is unknown. The former may represent some sort of terminal state of either main cell line. (23 refs.)

78-5308 Familial Pheochromocytoma With von Hippel's Disease in One Sister and Ectopic Adrenal Cortex in the Kidney in Another. (Eng) Mendelow, A. D.

(Medical Sch., Hospital St., Johannesburg 2001, S. Africa); Kay, S.; Levin, E.; Mieny, C. J. *Br J Surg* 65(2): 138-140; 1978.

Two sisters, aged 12 and 20 yr at presentation, developed

bilateral adrenal pheochromocytomas. The former had ectopic adrenal cortical tissue and the latter had von Hippel's disease. This is the first known association of the former with familial pheochromocytoma. (7 refs)

78-5309 Aniridia, Cataracts, and Wilms' Tumor in Monozygous Twins. (Eng) Cotlier, E. (Dept.

Ophthalmology and Visual Science, Yale Univ. Sch. Medicine, 333 Cedar St., New Haven, CT, 06510); Rose, M.; Moe, S. A. *Am J Ophthalmol* 86(1): 129-132; 1978.

The first instance of aniridia-Wilms' tumor syndrome in mentally retarded identical twin girls is reported. Both had congenital bilateral aniridia, cataracts, and glaucoma, but only one developed a Wilms' tumor. It is proposed that a combined gonadal-somatic cell mutation occurred in the twin with Wilms' tumor and a gonadal mutation occurred in the other twin. A gonadal somatic mutation probably occurred in a Wilms' tumor allele and a pure gonadal mosaicism affected the aniridia allele. A two-allele gonadal and a somatic mutation could explain the manifestation of the tumor in only one twin. The aniridia-Wilms' syndrome in twins further documents the relationship of teratogenic malformations and neoplasms. (21 refs.)

78-5310 Chemodectomas of the Carotid Body and Glomus Tympanicum. Sequential Occurrence in Sisters. (Eng) Schmidt, F. E. (Dept. Surgery, Tulane Univ. Sch. Medicine, 1430 Tulane Ave., New Orleans, LA, 70112)

Lewis, M. L.; Kahle, H. R. *JAMA* 240(12): 1266-1267; 1978.

Two sisters, aged 47 and 44 yr, developed chemodectomas of the carotid body 10 and 1.5 yr, respectively, after they were diagnosed with tumors of the glomus tympanicum. No recurrences have been noted in either patient since excision of the chemodectomas, and neither patient has had evidence of contralateral tumors at either site. (3 refs.)

78-5311 Extensive Metastases in Peutz-Jeghers Syndrome. (Eng) Ryo, U. Y. (Div. Nuclear Medicine, Michael Reese Hosp. and Medical Center, 2839 S. Ellis Ave., Chicago, IL, 60616); Roh, S. K.; Balkin, R. B.; Sieband, L. B.; Sobel, G. W.; Pinsky, S. M. *JAMA* 239(21): 2268-2269; 1978.

A 27-yr-old woman with Peutz-Jeghers (PJ) syndrome since age 11 yr was hospitalized with a sudden onset of weakness of the right extremity, expressive aphasia, and a 3 mo history of back pain. Liver and whole-body scintigrams demonstrated multiple metastatic disease. Postmortem examination showed extensive metastases in bone, liver, brain, lung, and

left ovary. The source of the metastases was a malignant change from PJ polyps. The incidence of malignant transformation resulting from PJ polyps is 2%-3%. (4 refs)

- 78-5312 **Familial Cushing's Syndrome.** (Eng) Arce, B. (Inst. Endocrinology and Metabolic Diseases, Capata y D, Havana 4, Cuba); Licea, M.; Hung, S.; Padron, A. *Acta Endocrinol (Kbh)* 87(1): 139-147; 1978.

The case reports of four siblings, two women and two men, with Cushing's syndrome are presented. Their ages at diagnosis were 15, 15, 13, and 32 yr, respectively. All four patients were short in stature and they had a disturbed carbohydrate metabolism. Adenomatous hyperplasia of the adrenal glands was demonstrated in three patients, and a virilizing adrenal carcinoma was noted in the fourth. A four-generation genealogical study of both branches of the family revealed no other case of adrenal hyperfunction. It is suggested that these tumors could have arisen from anomalous ACTH hyperproduction. These cases indicate a possible familial pathogenic origin of Cushing's syndrome. (3 refs)

- 78-5313 **Ultrastructural Study of the Normal Mucosa-Adenoma-Cancer Sequence in the Development of Familial Polyposis Coli.** (Eng) Mughal, S. (Dept. Histopathology, Westminster Medical Sch., Univ. London, Udall Street Labs., Udall St., London SW1P 2PP, England); Filipe, J. I. *J Natl Cancer Inst* 60(4): 753-768; 1978.

The ultrastructural changes that occur in the mucosae of patients with familial polyposis coli (FPC) were studied. The epithelial atypia were graded on the basis of vesiculation and increasing numbers of electron-dense bodies (EDB) and lysosomes in the mature and immature absorptive cells (IAC), the presence of immature and undifferentiated cells, variations in the goblet cells and the appearance of atypical secretory cells (ASC), and nuclear changes. With Grade 0 atypia, few EDB were noted in the apical region of the mature absorptive cells. With Grades 1 and 2, vesiculation and increased EDB were noted in mature cells. Goblet cells were smaller and sometimes distended with mucus; this distension may lead to disruption of the theca and leakage of secretory material into the adjacent cytoplasm. In Grades 3 and 4, immature and undifferentiated cells appeared at a higher than normal level in the crypt, and IAC and ASC were common. In Grade 5, the population consisted mainly of immature cells with numerous EDB, ASC were prominent, and nuclei showed marked pseudostratification. Grades 6 and 7 consisted mainly of undifferentiated cells, and there was marked nuclear irregularity. Carcinomas contained Grade 7 epithelia and Grades 5 and 6 glandular structures. Specimens from normal mucosa of FPC patients showed ultrastructural changes in 56%. The incidence of both atypia and adenomatous polyps was higher on the left side of the colon than on

the right side. These findings illustrate the adenoma-carcinoma sequence and support its occurrence. (27 refs)

- 78-5314 **Polyposis Coli. The Clinical Spectrum in Adults.** (Eng) Cooke, S. A. (Dept. Surgery, Univ. of the Witwatersrand, Johannesburg, S. Africa). *S Afr Med J* 53(12): 454-457; 1978.

The case reports of four patients with polyposis coli are presented to illustrate the unusual features of different types of this disease. The first patient is a 42-yr-old man with no family history of colonic disease who developed metaplastic polyps. The second is a 41-yr-old man with the multiple form of juvenile polyposis coli. The third is a 57-yr-old woman with adenomatous polyposis coli that is thought to have arisen by a spontaneous mutation. The fourth is an 18-yr-old girl who underwent total colectomy; she had had adenomatous polyposis coli at the age of 10, was a member of a family with a known predisposition to polyposis, and had multiple osteomas in the mandible. The types of lesions represented by these cases are discussed. A comparison of the fourth patient with Gardner's syndrome patients suggests that the differences between familial polyposis and Gardner's syndrome may be due only to a variation in the expressivity of the different extracolonic manifestations associated with the polyposis. Furthermore, juvenile polyposis, familial adenomatous polyposis, and carcinoma of the colon may be interrelated because they have occasionally been diagnosed in the same family. (15 refs)

- 78-5315 **Thymomas in Patients with Myasthenia Gravis.** (Eng) Slater, G. (Dept. Surgery, Mount Sinai Sch. Medicine, Fifth Ave. and 100th St., New York, NY, 10029); Papatestas, A. E.; Jenkins, G.; Kornfeld, P.; Horowitz, S. H.; Bender, A. *Ann Surg* 188(2): 171-174; 1978.

Thymoma occurred among 10% of all myasthenia gravis patients seen at one hospital. The histological classification was 72% lymphoepithelial, 24% lymphocytic, and 4% epithelial. Forty-three percent of the thymomas were invasive. (13 refs)

- 78-5316 **The Formal Pathogenesis of Primary Squamous Epithelial Carcinomas of the Thyroid (Meeting Abstract).** (Ger) Bocker, W. (Hamburg, W. Germany). *Zentralbl Allg Pathol* 122(3): 292; 1978. (no refs)

- 78-5317 **Multiple Endocrine Adenomatosis: In Support of the Common Origin Theories.** (Eng) Janson, K. L. (Section of Urology, Dept. Anatomy, Tulane Univ.

Sch. Medicine, New Orleans, LA); Roberts, J. A.; Varela, M. *J Urol* 119(2): 161-165; 1978.

A family with features of multiple endocrine adenomatosis types 1 and 2 was studied, and evidence in support of a common etiologic origin of the tumors associated with this syndrome is presented. A 9-yr-old girl (case 1) was found to have a right pheochromocytoma; chromosome study revealed a marker on the 15th chromosomal pair. Her 31-yr-old mother (case 2) had bilateral adrenal pheochromocytomas and an accessory pheochromocytoma in a paraaortic position, a right renal cystadenoma with renal cysts and a left renal adenoma, and three islet cell tumors of the pancreas. Chromosome studies indicated a marker on the 15th chromosomal pair identical to that of her daughter and the same histocompatibility haplotype as her daughter. The mother of case 2 (case 3) had had a pituitary tumor removed at the age of 22 and was later diagnosed as having bilateral pheochromocytomas. Chromosomal analysis of the mother of case 3 (case 4) revealed no markers. A 50-yr-old sister of case 3 has a marker on the 15th chromosomal pair but no adenomatosis. A younger sister and brother of case 1 have this marker but no evidence of disease. These findings suggest a common genetic origin of multiple endocrine adenomatosis but that not all persons with this genetic information develop tumors. (20 refs)

- 78-5318 Association of Acromegaly and Meningiomas.** (Eng) Bunick, E. M. (Div. Endocrinology and Metabolism, Hahnemann Medical Coll. and Hosp., 230 N. Broad St., Philadelphia, PA, 19102); Mills, L. C.; Rose, L. I. *JAMA* 240(12): 1267-1268; 1978.

The coexistence of a growth hormone-producing pituitary tumor and a subfrontal meningioma at the time of initial diagnosis in a 57-yr-old man is reported. There are only six similar cases in the literature. In two of these patients the pituitary tumor and the meningioma were present before the initiation of therapy. The other four patients underwent irradiation of the pituitary fossa for treatment of acromegaly before the meningioma developed. The present patient had both tumors without any history of head trauma, infection, or irradiation. There is no way to determine which tumor developed first, and the exact nature of the relationship between acromegaly and meningiomas is not known. (3 refs.)

- 78-5319 The Role of the Subependymal Plate in the Origin of Experimental Gliomas: A Sequential Electron Microscope Study (Meeting Abstract).** (Eng) Lantos, P. L. (Middlesex Hosp. Medical Sch., London, England). *J Neuropathol Exp Neurol* 37(5): 647; 1978. (no refs)

- 78-5320 Human Neuroblastoma Serially Transplanted in Nude Mice and Metastases.** (Eng) Hata, J. I. (Dept. Pathology, Tokai Univ., Sch. Medicine, Bohseidai, Isehara, Kanagawa, 259-11, Japan); Ueyama, Y.; Tamaoki,

N.; Furukawa, T.; Morita, K. *Cancer* 42(2): 468-473; 1978.

The direct transplantation of a neuroblastoma from a 3-yr-old girl into nude BALB/c mice is reported. A metastatic tumor of the paraaortic lymph node obtained at surgery was minced, and the fragments were transplanted sc into the mice; serial transplantation was then established. The histology of the transplanted tumor was almost the same as that of the original tumor. During serial transplantation, a high frequency of metastasis to distant sites such as the ovaries, lymph nodes, and cerebrum was observed; the metastatic rate to the ovaries reached 66%. The distribution of the organs affected by metastasis was similar to that in humans with neuroblastoma. This transplantable human neuroblastoma may be useful for studies of the morphological and biological nature of these tumors. (7 refs.)

- 78-5321 Massive Peripheral Giant-Cell Repara- Granuloma of the Jaw: A Pregnancy Dependent Tumor.** (Eng) Caillouette, J. C. (50 Bellefontaine St., Pasadena, CA, 91105); Mattar, N. *Am J Obstet Gynecol* 131(2): 176-179; 1978.

A 21-yr-old pregnant woman had a large giant-cell granuloma that may have been enhanced by the diminished immune response and hormonal changes associated with pregnancy. The high levels of estrogens and gonadotropic hormones during pregnancy have a positive trophic effect on the buccal mucosa and augment glycogen deposition in the epithelium. Peripheral giant-cell reparative granuloma should not be mistaken for true giant-cell tumor, which is rare. (19 refs)

- 78-5322 Malignant Transformation of Esophageal Columnar Epithelium.** (Eng) Berenson, M. (Div. Gastroenterology, Univ. Medical Center, Salt Lake City, UT, 84132); Riddell, R. H.; Skinner, D. B.; Friesen, J. W. *Cancer* 41(2): 554-561; 1978.

The cases of two men (aged 47 and 69 yr) with esophageal columnar epithelium (Barrett's esophagus) in which microinvasive adenocarcinoma of the esophagus developed are reported. The former had multiple foci of in situ carcinoma contiguous with epithelium resembling that of the stomach and intestine. The latter had signet-ring type adenocarcinoma. The adenocarcinoma appeared to develop directly from the dysplastic epithelium, providing direct evidence that the dysplastic process is neoplastic. The origin of esophageal columnar epithelium is not known, but it may originate from reflux-induced ulceration and subsequent regeneration by mature gastric cells that undergo metaplastic change. Patients with columnar esophageal epithelium should be routinely checked for malignant transformation. (28 refs)

- 78-5323 Cylindrical Metaplasia of the Lower Esophagus (Meeting Abstract).** (Fre) Ozzello, L. (Institute of Pathology, Universite de Lausanne, Lausanne, Switzerland).

d); Savary, M.; Roethlisberger, B. *Schweiz Med Wochenschr* 108(13): 506; 1978. (no refs)

5324 **The Aryl Hydrocarbon Hydroxylase Polymorphism and Respiratory Cancer (Meeting Abstract).** (Eng) Paigen, B. (Roswell Park Memorial Inst., Buffalo, NY, 14263); Ward, E.; Gurtoo, H. L.; Minowada, J.; Vincent, R.; Parker, N. B.; Paigen, K. *Proc Am Assoc Cancer Res* 19: 100; 1978. (no refs)

5325 **Histochemical Observations on Nodules Induced in the Mouse Liver by Phenobarbitone.** (Eng) Butler, W. H. (Pharmaceuticals Div., ICI Ltd., Macclesfield, Cheshire SK10 4TG, England); Hemsall, V. *J Pathol* 125(3): 155-161; 1978.

Meaning C3H mice were fed phenobarbitone in the diet (500 ppm for 1 wk, followed by 1,000 ppm for the duration of the experiment, 2 yr), and their livers were examined histochemically to determine which features are associated with malignant neoplasms. A total of 13 livers from phenobarbitone-fed mice and 3 livers from control animals, all with macroscopic nodules, were examined. The morphology of the lesions in the treated animals depended on the preparative techniques used. On examination of hematoxylin and eosin-stained sections, nodules from 11 treated livers showed a simple trabecular pattern of cells with abundant eosinophilic cytoplasm; nodules from the other 2 livers had thickened and irregular trabeculae with wide vascular spaces. Examination of paraffin sections indicated that six nodules contained areas showing a dilated vascular pattern and irregular trabeculae. On serial sectioning, vascular emboli of hepatic cells were observed in the lungs of the two animals with the thickened trabeculae. A varying pattern of histochemical reaction was noted. The aniline hydroxylase reaction was dense in the simple nodules and centrilobular areas of the treated mice, but was weak in the abnormal trabecular lesions. Tissues from 10 of the nodules in treated mice were transplanted sc into leanling male C3H mice; two successful transplants were recorded. These findings suggest that it is possible to predict behavioral characteristics of mouse hepatic nodules on the basis of certain morphological characteristics. (12 refs.)

5326 **Conglomerative Siderosis and Carcinoma of the Lung.** (Fre) Lemenager, J. (Clinique medicale, C.H.U., 14033 Caen Cedex, France); Wyplaz, J.; Mandel, J. C.; Chasles, J.; Le Bouffant, L.; Caillard, J. F.; Benard, J. *Rev Fr Mal Respir* 6(4): 431-433; 1978.

51-yr-old worker in the metallurgy industry presented with an opaque focus in each lung. Thoracotomy revealed a focus of conglomerative siderosis in the left lung and an adenocarcinoma in the right lung. Histological and mineralogical examination of the siderosis revealed a giant cell granuloma, high iron content, and particles of titanium oxide and silver. Aggregates of different minerals were found in the fibrosis

surrounding the siderosis. The adenocarcinoma, which was moderately differentiated, contained numerous amphibole asbestos fibers as well as iron oxides, quartz, and traces of silver and feldspar. (3 refs.)

78-5327 **A Histochemical and Ultrastructural Study of Adenocarcinoma of the Lung.** (Eng) Kimura, Y. (Dept. Pathology, Tokyo Metropolitan Inst. Gerontology, 35-2, Sakaecho, Itabashiku, Tokyo 173, Japan). *Am J Surg Pathol* 2(3): 253-264; 1978.

Twenty-five cases of solitary nodular-type adenocarcinoma of the human lung were studied ultrastructurally in an attempt to elucidate the histogenesis of adenocarcinoma at an early and developing stage. Twenty-one cases were type D adenocarcinoma (resembling nonciliated bronchiolar cells), two were type B (resembling the mucous cell of the bronchial gland), one was type A (resembling the bronchial goblet cell), and one was type C (resembling the type II alveolar lining cell). According to criteria of the World Health Organization, 15 lesions were adenocarcinomas and 10 were bronchioloalveolar carcinomas; there were no histochemical or ultrastructural differences between the two. These findings suggest that pulmonary adenocarcinoma may arise from more than one cell type but that the predominant type is apparently from the terminal bronchiolar epithelium. (18 refs.)

78-5328 **Special Features of Intestinal Metaplasia and Its Relation to Early Gastric Carcinoma in Man: Observation by a Method in Which Leucine Amino-peptidase Activity Is Used.** (Eng) Nakahara, K. (Dept. Surgery, Natl. Kyushu Cancer Center, 595 Minami-Ku, Notame, Fukuoka-City 815, Japan). *J Natl Cancer Inst* 61(3): 693-702; 1978.

A leucine amino-peptidase (LAP) staining procedure was used to determine the degree of intestinal metaplasia and its macroscopic distribution in gastric mucosa in order to investigate the relationship between intestinal metaplasia and gastric carcinoma. LAP is a specific enzyme of intestinal metaplasia, and the area showing a positive reaction for this enzyme corresponded strictly with the zone of intestinal metaplasia examined microscopically. This method was used to examine specimens of gastric carcinoma confined to the mucosa from 40 Japanese patients. Gastric mucosa containing differentiated tubular-type carcinoma (24 cases) was associated with a high degree of metaplasia, but a low degree of metaplasia was found in the 16 cases of poorly differentiated carcinoma. The more differentiated a tubular-type carcinoma was, the more closely it was related to intestinal metaplasia. In contrast, all carcinomas arising from the mucosa without intestinal metaplasia (18%) were poorly differentiated. Therefore, gastric carcinomas occurring in Japanese patients are frequently and closely related to intestinal metaplasia. (32 refs.)

- 78-5329 Cancer of the Stomach after Resection for Peptic Ulcer.** (Rus) Patiutko, Iu. I. (Dept. Oncology, I. M. Sechenov First Medical Inst., Moscow, USSR). *Khirurgiia (Mosk)* (3): 27-31; 1978.

To evaluate whether resection of the stomach for peptic ulcer increases the relative risk of cancer in the remaining part of the stomach, the case histories of 942 patients with gastric ulcer treated surgically between 1946 and 1952 were subjected to retrospective analysis. Nineteen patients had developed cancer of the gastric stump: 1 within 6-10 yr of surgery, 3 within 11-15 yr, 3 within 16-20 yr, 7 within 21-25 yr, and 5 > 26 yr postsurgery. Calculation of the expected age-predicted incidence of cancer of the proximal region of the stomach (patients had undergone resection of distal region) showed that observed and expected values were in good agreement only within the first 10 yr after resection; 21-25 yr later the observed incidence was 3.5 times greater and > 25 yr later it was 12.5 times greater than expected. (11 refs)

- 78-5330 Pathogenesis of Gastric Cancer in Pernicious Anemia.** (Eng) Ruddell, W. S. (Dept. Gastroenterology, Central Middlesex Hosp., London NW10 7NS, England); Bone, E. S.; Hill, M. J.; Walters, C. L. *Lancet* 1(8063): 521-523; 1978.

Thirteen patients with pernicious anemia (PA) and 13 age-matched controls were studied to determine if nitrite concentrations in the PA patients could be involved in the pathogenesis of gastric cancer. The gastric juice of the PA patients had significantly higher nitrite concentrations, total bacterial counts, and counts of bacteria showing nitrate-reductase activity than the gastric juice of the controls. The results of an in vitro nitrosation experiment indicated that appreciable quantities of nitrosamines were detected in extracts of gastric juice from the PA patients. Nitrosamines were not detected in any of the extracts from the control experiments without added nitrite. If similar nitrosation occurs in vivo, the intragastric production of carcinogenic N-nitroso compounds could explain the high incidence of gastric cancer in patients with PA. (17 refs)

- 78-5331 Squamous Cell Carcinoma of the Stomach.** (Eng) Won, O. H. (1221 Mowry St., Old Forge, PA, 18518); Farman, J.; Krishnan, M. N.; Iyer, S. K.; Vuletin, J. C. *Am J Gastroenterol* 69(5): 594-598; 1978.

The case report of a 55-yr-old woman with a pure squamous cell carcinoma of the stomach is given. The tumor arose in an area lined by squamous mucosa continuous with that of the esophagus, but located well within the stomach. It is thought that this squamous mucosa was present congenitally and that it progressed through dysplasia to neoplasia and invasive cancer. (21 refs.)

- 78-5332 Frequency of Primary Liver Carcinoma Associated by Cirrhosis of the Liver at Cipto Mangunkusumo Central Hospital, Jakarta; Assessed by Peritoneoscopy (Meeting Abstract).** (Eng) Akbar, N. (Dept. Hepatology Medical Faculty, Univ. Indonesia, Jakarta, Indonesia); Lesmana, L.; Sulaiman, A.; Noer, A. M.; Sukato, U. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 35-36; 1977. (no refs)

- 78-5333 Familial Liver-Cell Adenomas and Diabetes Mellitus.** (Eng) Foster, J. H. (Dept. Surgery, Hartford Hosp., Hartford, CT, 06115); Donohue, T. A.; Berman, M. M. *N Engl J Med* 299(5): 239-241; 1978.

A family was encountered in which four members, a 43-yr-old mother, two daughters aged 17 and 18 yr, and a 15-yr-old son, had liver cell adenomas. All were diabetic, and the two daughters also had sclerocystic ovaries, suggestive of a change in their internal hormonal milieu. Liver-function tests gave normal results in all, except for elevated enzyme in one daughter (presumably related to an acute episode of hemorrhagic necrosis), and a twice-normal elevation of alkaline phosphatase in the son. A fourth child, a 12-yr-old nondiabetic girl, was not affected. The mother's father, also diabetic, died with diffuse pulmonary metastases from an autopsy-confirmed hepatocellular carcinoma occurring in a cirrhotic liver. His nondiabetic father had died with biopsy-confirmed hepatocellular carcinoma in a cirrhotic liver. There are no reports of an association of diabetes mellitus (DM) with benign or malignant liver tumors. The mild nature and strong family penetrance of DM in this family suggest that the diabetic phenotype is that of maturity-onset diabetes of the young as opposed to classic juvenile-onset diabetes. This apparent autosomal dominant inheritance of DM may correlate with a higher incidence of liver cell adenomas in other families with similar diabetic phenotypes. It is hypothesized that a change in the hormonal milieu caused by endogenous or exogenous sex steroids, masks a fundamental, perhaps hereditary, defect in carbohydrate metabolism, resulting in nodule formation in the liver. (10 refs)

- 78-5334 Evaluation of Hepatocellular Neoplasms in Mice.** (Eng) Ward, J. M. (Tumor Pathology Branch, Carcinogenesis Testing Program, NCI, NIH, Public Health Service, US Dept. Health, Education, and Welfare, Bethesda, MD, 20014); Vlahakis, G. *J Natl Cancer Inst* 61: 807-811; 1978.

The genesis of hepatocellular neoplasms was studied in C3H-AvylB male mice, a strain with a high incidence of neoplasms.

y occurring hepatocellular tumors. At 3, 6, 9, 12, and 15 of age, preselected groups of six mice were killed and a complete necropsy was performed. The number of mice with tumors in these age groups was 0%, 33.3%, 50%, 66.7% and 100%, respectively. As the mice aged, the number of tumors increased and the mean diameter of the tumors increased. Histologically, the smallest tumors were composed of sheets of cells and were better differentiated than the largest tumors. All foci of hepatocytes in prominent trabeculae were seen in a few small tumors, but prominent trabecular formation was a common characteristic of large tumors. Metastatic tumors in these mice plus those in C3H/c, C3H, C3H/fb, C3H/fb x YBR/F₁, and C57BL/6 mice were studied morphologically. The pulmonary metastases were composed of a well-differentiated sheet of hepatocytes that resembled those of the solid liver tumors. The smallest hepatocellular neoplasms appeared to represent early stages in the development of metastatic hepatocellular carcinoma. (12 refs.)

5335 Vaginal Metastasis from Renal Cell Carcinoma: Routes of Metastasis. (Jpn) Masuda, F. (Dept. Urology, Jikei Univ. Sch. Medicine, Tokyo, Japan); Hinuma, H.; Sasaki, T.; Arai, Y.; Shoji, R.; Teramoto, K.; Nagisawa, M.; Chen, Z.; Machida, T. *Jpn J Urol* 69(4): 479; 1978.

Left renal cell carcinoma was diagnosed in a 22-yr-old woman. Renal venography revealed a tumor embolism from the renal vein to the inferior vena cava, and a chest x-ray indicated metastasis to the bilateral lymph nodes. Five months later, the patient developed vaginal bleeding. One small tumor was found near the vaginal introitus and another at the left vaginal wall adjacent to the uterine neck. Histological examination of the vaginal lesions showed a clear cell adenocarcinoma of the capillaries, confirming the presence of metastases from renal lesion. The possible routes of metastasis were studied in adult mongrel dogs by following the circulation of the ovarian vein before and after ligation of the left renal vein. The results suggest that renal cell carcinomas metastasize to the vagina as follows: in a left renal cell carcinoma, the tumor cells flow from the renal vein, directly or through the ureteral vein, to the left ovarian vein, the ovarian plexus, the parovaginal plexus, and the vagina. In the case of a right renal cell carcinoma, the tumor cells flow from the renal capillary veins or the ureteral vein to the vagina through the right ovarian vein. (13 refs)

5336 Functional and Morphological Studies of Human Renal Cell Carcinoma Using a Nude Mouse Cell Culture System. (Jpn) Katsuoka, Y. (Dept. Urology, Sch. Med., Keio Univ., Keio, Japan). *Jpn J Urol* 69(3): 303; 1978.

Thirty-six surgical specimens of human renal cell carcinoma

obtained over a 5-yr period were cultured, transplanted into nude mice, and analyzed with respect to their functional and morphological characteristics. Eleven primary cultures, 2 continuous cultures, and 6/12 tumor transplants were successful, and 4/6 transplanted tumors are still growing. Among the latter is a tumor capable of producing erythropoietin (EP) and derived from metastatic foci in a lung. Histopathologically, the transplanted tumors, the continuous cultures, and the original tumors were basically identical. Indirect culture through a nude mouse produced a cell line (KU-2) whose histopathological characteristics remained unchanged even after it was transplanted back into a nude mouse. There were brush border-type structures in the transplanted tumors and radially arranged microvilli on the culture cells (as shown by electron microscopy), proof that renal cell carcinoma originates in a proximal tubular cell. The transfer of erythropoietin to the tumor-bearing mice and the determination of EP activity in the tumor tissue raise the possibility that the tumor per se is capable of producing EP. The development of hepatomegaly in the tumor bearers, plus the other findings, may explain the unique functional characteristics of these tumors. (82 refs)

78-5337 Cancer Risk with Different Types of Gallstones. (Ger) Grosse, H. (Bezirkskrankenhaus-Prosektur, Billrothstrasse 22, DDR-23 Stralsund, E. Germany). *Arch Geschwulstforsch* 47(8): 710-718; 1977.

The association between gallstones and different types of cancer was investigated in 22,868 autopsied adults. Of these, 7,411 had cholelithiasis or had undergone cholecystectomy. In cancerous gallbladders, multiple faceted stones and solitary combined stones were seen most frequently; pigmented stones, pure crystalline cholesterol stones, and lobulated stones were rarely seen. As the partial volume of gallstones increased, so did the number of cases of cholecystitis and carcinoma. In men with all forms of gallstones, carcinomas of the lung, stomach, colon, rectum, and prostate gland were more common than gallbladder carcinoma. In women with multiple faceted stones, the most frequent cancer was that of the gallbladder. The area of the gallbladder affected by the stones was more important to the localization of a later carcinoma in women than in men. The percentage of unoperated patients with gallstones who later developed any type of cancer was not higher than that of patients with a previous cystectomy. (42 refs)

78-5338 A New Aspect of the Urinary Excretion of Tryptophan Metabolites in Patients with Cancer of the Bladder. (Eng) Teulings, F. A. (Dept. Biochemistry, Rotterdamsch Radio-Therapeutisch Instituut, P.O. Box 5201, 3075 EA Rotterdam, Netherlands); Peters, H. A.; Hop,

W. C.; Fokkens, W.; Haije, W. G.; Portengen, H.; van der Werf-Messing, B. *Int J Cancer* 21(2): 140-146; 1978.

Excretion of tryptophan metabolites was examined in 165 urine samples obtained from bladder cancer patients (67), renal cancer patients (19), patients cured of bladder cancer (37), and control subjects (40). The metabolites kynurenine, kynurenic acid, 3-hydroxykynurenine, xanthurenic acid, and 3-hydroxyanthranilic acid were separated by ion-exchange chromatography followed by thin-layer chromatography on cellulose. Their concentration was determined by measuring the fluorescent intensity of the spots on the cellulose plates. Thirty-six percent of the bladder cancer patients and 67% of the renal carcinoma patients excreted abnormally high amounts of at least one of the metabolites, most frequently 3-hydroxyanthranilic acid, 3-hydroxykynurenine, and kynurenine. In bladder cancer an elevated excretion of metabolites occurred principally in patients with obstruction to the outflow of urine from the kidneys. Ureteral obstruction, probably a direct consequence of the presence of tumor in the urinary tract, causes an alteration in renal function that is reflected in urea and creatinine retention. The high incidence of abnormal excretion of tryptophan metabolites in renal cancer also points to alterations in renal function caused by the tumor. These observations do not support the hypothesis that tryptophan metabolites play a part in the etiology of bladder cancer, since raised basal urinary levels of these metabolites are a result, rather than a cause of the disease. (19 refs)

78-5339 Blue Nevus of the Endocervix: Report of Two Cases and Ultrastructure. (Eng) Diaz de Molnar, A. M. (Dept. Pathology, Jewish General Hosp., 3755 Cote St. Catherine Road, Montreal, Quebec H3T 1E2, Canada); Guralnick, M.; Ferenczy, A. *Gynecol Oncol* 6(4): 373-382; 1978.

Examination of the surgical specimens from two women, aged 44 and 58 yr, who had undergone hysterectomy for menorrhagia revealed blue nevus lesions in each. The younger woman had a solitary 4- x 4-mm lesion in the posterior one-third of the endocervical canal. In the older women, multiple nevi, the largest measuring 2 x 1.5 cm in diameter, were noted in the upper two-thirds of the endocervical canal. Electron microscopy of specimens removed from the older woman showed that the cells were engaged in melanogenesis similar in all respects to that observed in blue nevus cells of the dermis. Their ultrastructural resemblance and close location to perineurial fibroblasts of nonmyelinated nerve bundles suggest that at least in some instances, the perineurial fibroblasts may be the most immediate neural crest precursors of blue nevus cells of the endocervix. The light microscopic appearance of the lesions is also outlined. (19 refs.)

78-5340 The Histogenetic Significance of Dysplastic Lesions of the Cervix. (Fre) Hoang-Ngoc Minh

(Service Central d'Anatomie et Cytologie Pathologiques C.H.U. F 80000 Amiens, France); Smadja, A.; Orcel, L. *Gynecol Obstet Biol Reprod (Paris)* 7(3): 387-398; 1978.

The results of morphological studies of dysplastic lesions of the uterine cervix are described. These lesions appear to develop from the vestigial cells that are situated at the endocervical junction. The dysplasias consist of cells with a basophilic cytoplasm and nondistinctive edges. The cells form streams that run from the external os of the cervix toward the endocervix, where they replace and disturb the columnar epithelium. These lesions may take one of two forms, regular or irregular, according to the degree of nuclear-cytoplasmic disharmony. The slow and insidious evolution of these lesions must be distinguished from dystrophic endocervical lesions that result from a disturbance in tissue nutrition and from exocervical lesions that can undergo atypical scarring. (19 refs.)

78-5341 Genital Infections and Cervical Carcinoma (Letter to Editor). (Eng) Astedt, B. (Dept. Obstetrics and Gynecology, Malmo Allmanna Sjukhus, Malmo, Sweden); Furgyik, S.; Hansson, H. *Br Med J* 2(6131): 201-202; 1978.

A link between genital infection with neisseria and cervical cancer is proposed. Of 162 patients infected with neisseria, 34 had cervical carcinoma (20% incidence, vs 5% age-matched controls). The results agree with the opinion that genital infections predispose to cervical neoplasia. (19 refs)

78-5342 Endometrial Carcinoma in Gonadal Dysgenesis with and Without Estrogen Therapy. (Eng) Cator, A. G. (Dept. Pathology, Royal Women's Hosp., 7 Swanston St., Carlton, Victoria, 3053, Australia); Fortum, D. W.; Evans, J. H.; Kneale, B. L. *Gynecol Oncol* 6(4): 316-327; 1978.

The case reports of two women, aged 29 and 21 yr, who had gonadal dysgenesis and who subsequently developed endometrial carcinoma are presented. The former was initially seen because of menorrhagia; both initial and recurrent episodes were treated by curettage. One of these curettages revealed Grade 1 adenocarcinoma. Chromosomal analysis revealed that 80% of the cells had a 45,X constitution and 20% had a 47,XXX constitution. She had never received estrogen therapy. The second patient was also initially seen because of menorrhagia, and she was treated by curettage and later by estrogen therapy. One year later, regular withdrawal bleeding was produced by cyclic progestogen therapy only. During the next 2 yr, she was treated with an estrogen-progestogen preparation; the total amount of estrogen she received was estimated to be 15 mg. A curettage during this estrogen therapy period revealed well-differentiated

grade 1 adenocarcinoma. Chromosome analysis revealed that 56% of the cells had a 45,X constitution, 44% a 46,XXr constitution. Autoradiography revealed that the ring chromosome was a late replicating X chromosome. Both patients were surgically treated and are alive 13 and 7 yr, respectively, after surgery. It is suggested that the carcinomas developed because of unopposed endogenous estrogen secretion resulting from anovulation, but a genetic disposition to cancer in gonadal dysgenesis patients cannot be excluded. (24 refs.)

8-5343 **Carcinoma in a Nipple Preserved by Heterotopic Auto-implantation.** (Eng) Allison, A. B. (Oxford-Lafayette County Hosp., Oxford, MS); Howorth, M. B. *Engl J Med* 298(20): 1132; 1978.

An infiltrating lobular breast carcinoma, with metastasis identified in two lymph nodes, was removed from a 40-yr-old woman by modified radical mastectomy, and the nipple was transplanted to her groin to preserve it for breast reconstruction. Carcinoma was present in sections taken from beneath the nipple. Upon removal of the nipple at the patient's request

1 yr later, pathological examination revealed carcinoma in the nipple. It is not known whether the cancer was already present or developed after transplantation. (2 refs)

EPIDEMIOLOGY AND BIOMETRY

78-5344 Cancer by County: Etiologic Implications.

(Eng) Blot, W. J. (Environmental Epidemiology Branch, NCI, Bethesda, MD, 20014); Mason, T. J.; Hoover, R.; Fraumeni, J. F. In: *Incidence of Cancer in Humans, Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 21-32; 1977.

Lung, colon, and oral and pharyngeal cancer incidence was determined in various counties throughout the US, and the results were correlated with those of analytical field studies. In white men, lung cancer mortality was elevated in the Northeast and around large metropolitan areas, but the highest rates were clustered in contiguous counties along the Gulf of Mexico and the Southeast Atlantic coast. Rates among women were high in urban areas. In general, rates were lowest in farming areas; mortality was inversely correlated with county education levels. Mortality among white men was elevated in counties with paper, chemical, and petroleum manufacturing industries. Geographic patterns for colon cancer mortality were identical for men and women. For both sexes and among races, high rates were present throughout the Northeast and low rates were present throughout the South. High mortality rates were associated with urbanization, income, and ethnicity (Irish, German, Czechoslovakian). Colon cancer did not appear to be associated with dietary habits. High mortality from oral cancer among white men predominated in the Northeast, while high rates in white women were clustered in the Southeast. The northern excess among men was attributed to urbanization, tobacco smoking, and alcohol consumption. It is suspected that the poor use of snuff among southern women accounts for their excess of oral cancer. Among men, oral cancer mortality was elevated in counties with leather industries, particularly in the central US. Other findings include high skin cancer rates in the South, high bladder cancer rates in men in industrial areas, high stomach cancer rates in Northern European and Spanish American ethnic groups, and rectal cancer rates paralleling those of colon cancer. Studies of cancer mortality in counties grouped according to specific environmental variables are reviewed. (31 refs)

78-5345 Minimum Incidence Rate Based on Microscopically Diagnosed Cancer in the Semarang City Population. 1970-1974 (Meeting Abstract). (Eng) Tirtosugondo (Dept. Pathology, Kariadi Hosp., Semarang, Indonesia); Soetoto, H. J.; Brotosaputro, B.; Widagdo, L. In: *Summary Program and Abstracts of Papers. Proceedings of*

the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 101-103; 1977. (no refs)

78-5346 Cancer Statistics of Sabah (Meeting Abstract) (Eng) Kulkarni, M. G. (Queen Elizabeth Hospital, Kota Kinabalu, Sabah, Malaysia).

In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 83-84; 1977. (no refs)

78-5347 Cancer Pattern Among Filipino Population in Hawaii (Meeting Abstract). (Eng) Hirohata, T. (Biomedical Sciences D-102, Univ. Hawaii, Honolulu, HI 96822); Kolonel, L.; Nomura, A.; Burch, T. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 80-81; 1977. (no refs)

78-5348 Recent Changes in Estimated Cancer Mortality in Hawaii (Meeting Abstract). (Eng) Chiazzese, L. (Div. Biostatistics, Epidemiology, Dept. Community Medicine and International Health, Georgetown Univ. Sch. Medicine, Washington, DC, 20007); Levin, D.L.; Silverman, D. T. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 33-44; 1977.

Statistical problems involved in calculating recent changes in estimated cancer mortality are outlined, and appropriate methods for examining these changes are discussed. A comparison of cancer mortality data for 1974 with the most recent provisional 12-mo data for 1975 reveals that there was a 2.3% increase in the crude mortality rate and a 0.7% increase in the mortality rate adjusted to the 1940 population.

to the 1940 population. This 0.7% increase contrasts with the 2% av annual increase observed in the final data from 1968 to 1974. The appropriateness of extrapolating from short-term monthly trends to determine recent changes in cancer mortality is questioned. However, there appears to be some increase in cancer mortality from 1974 to 1975, although not of the magnitude originally reported. Most of the increase appears to be due to respiratory cancer, particularly among white women. Lung cancer mortality rates have been increasing for several years, but mortality rates for cervical and stomach cancer have been declining at a significant rate. The use of age-, race-, sex-, and site-specific final mortality data for determining changes in cancer mortality removes some of the confusion in discerning observed changes and may provide clues to the etiologies of these changes. (9 refs)

78-5349 **The Relationship of Serum Vitamin A and Serum Cholesterol to the Incidence of Cancer in Evans County, Georgia.** (Eng) Kark, J. D. (Univ. North Carolina, Chapel Hill, NC, 27514). *Diss Abstr Int B* 39(1): 78B; 1978. (no refs)

78-5350 **Environment and Cancer Epidemiology.** (Ger) Friedl, H. P. (Institut für Krebsforschung, Universität Wien, Borschkegasse 8a, A-1090 Vienna, Austria). *Oesterr Z Onkol* 4(2-3): 46-52; 1977.

Cancer mortality rates in Austria were analyzed in view of the contention that environmental factors influence the development of at least 80% of all cancers. The mortality rates for the most common malignancies were similar to those in other western industrialized countries. A relationship between cancer risk and the size of the Austrian communities was demonstrated. Compared with other common cancer sites, cancer of the stomach was most common in small communities with mainly an agricultural population. Etiological hypotheses about the origin of stomach cancer are discussed with regard to epidemiological findings and the dietary habits of the population. (13 refs.)

78-5351 **Correlation of Sites.** (Eng) Clemmesen, J. (Danish Cancer Registry, Finsen Inst., Copenhagen 2100, Denmark). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 87-100; 1977.

Patterns in the occurrence of malignant neoplasms in Denmark during the period 1943 through 1972 (data obtained from the Danish Cancer Registry) were analyzed. Although there was a considerable regularity in the patterns over the

years, there were some outstanding changes: morbidity rates for cancers of the male digestive organs, particularly the stomach, have decreased, as has that for esophageal cancer; cancers of the pancreas, respiratory tract and lung, genitals (especially in men), breast (women only), urinary tract, skin, and hemopoietic system have increased. A socioeconomic grading for lung cancer among men and for cervical uterine cancer is unfavorable to the poor. Age-adjusted mortality rates showed a definite correlation between cigarette consumption and lung cancer and a less pronounced correlation between cigarette consumption and bladder cancer. The ratio of lung to bladder cancer in Denmark, Sweden, Norway, and the US was approx 3:1, whereas the ratios for Britain and Finland were considerably higher. There was no negative correlation between the morbidity rates for cervical and breast cancers in several countries, but reverse ratios appeared in Israel and Japan. (14 refs.)

78-5352 **Regularly Used Drugs and Cancer.** (Eng) Jick, H. (Dept. Medicine, Boston Univ. Sch. Medicine, Boston, MA, 02118); Smith, P. G. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 389-398; 1977.

The possible role of the aggregate of all drugs taken on a regular basis on the induction of cancer was studied based on 800 cancer patients and 3,443 comparison patients whose reasons for entering the hospital were thought to be such that their use of drugs on a regular basis would be similar to that of the cancer patients. In the 3 mo prior to hospitalization, 444 of the cancer patients and 1,710 of the comparison patients reported using one or more drugs at least once a week. When examined by age, the proportion of regular users of one or more drugs was similar in both patient groups. When determined for each age-sex stratum separately, the risk ratio estimates associated with regular use of one or more drugs were also similar for the cancer and comparison patients. The upper limit on the proportion of all cancers attributable to regular drug use, in both regular drug users and nonusers, was estimated to be about 1%. Among long-term (5 yr or more) users, the proportion of all cancers that may be associated with long-term drug use was about 2%. Thus, regular usage of drugs accounts for, at most, only a small fraction of all new cases of cancer diagnosed between the ages of 20 and 75 yr. (11 refs.)

78-5353 **Effects of Drugs--Immunosuppression.** (Eng) Hoover, R. (NCI, Bethesda, MD, 20014). In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold

Spring Harbor Laboratory): *Origins of Human Cancer*. Vol. 4, 602 pp.; 369-379; 1977.

The risks of various types of malignancy in renal transplant recipients were studied based on data from 16,290 patients who received transplants between 1951 and 1975. Expected risks for the general population were established from the Connecticut Cancer Registry. Eleven patients for which there was evidence that tumors had been transplanted from the donor along with the kidney were excluded. The risk of lymphoma among transplant recipients was 1.5/1,000 patients/yr, or about 19 times that in the general population. This excess was due almost exclusively to a risk of reticulum cell sarcoma, which was approx 150 times greater than expected. Fifty percent of the lymphomas were localized to the brain. The risk of other cancers was about twice that expected, the excess being attributable to increases in soft-tissue sarcomas, adenocarcinomas of the lung, primary liver and bile duct tumors, lower urinary tract cancers, thyroid cancers, and, perhaps, cancers of the uterine cervix. No excesses were noted for cancers of the stomach, colon and rectum, or breast. The relative risk of lymphoma was increased to 16-fold within the year following transplant and thereafter, whereas the increased risk of other cancers appeared 2 yr after transplant and increased progressively thereafter. The risk of cancer seemed to be particularly excessive among those aged <20 yr at transplantation. The implications of these data for several hypothesized mechanisms of carcinogenesis are discussed. (21 refs.)

- 78-5354 **Estimates of the Cancer Risk Due to Nuclear Electric Power Generation.** (Eng) Ellett, W. H. (Office Radiation Programs, Environmental Protection Agency, Washington, DC, 20460); Richardson, A. C. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): *Origins of Human Cancer*. Vol. 4, 602 pp.; 511-527; 1977.

The risk of fatal cancer associated with the annual production of one gigawatt (1,000 megawatts) of nuclear electric power was estimated. The estimated number of cancer fatalities due to the extraction and fabrication of uranium fuel was about 0.5/gigawatt-year [GW(e)-y, the amount of energy produced annually by a single nuclear reactor of the type and size currently being constructed]. The estimated number of cancer fatalities due to the reprocessing of spent fuel was about 0.3/GW(e)-y, and the estimated number of fatal cancers due to nuclear reactor effluents was about 0.1/GW(e)-y. The estimated number due to occupational exposure was 0.2/GW(e)-y, and that due to the accidental release of nuclear material was about 0.1/GW(e)-y. The latter estimate was uncertain because fatalities related to non-core-related accidents, fuel reprocessing, and waste storage could not be estimated. In total, about 1.2 cancer deaths could be associat-

ed with each gigawatt-year of electrical energy produced. (3 refs.)

- 78-5355 **Estrogen Use and Cancer of the Uterine Corpus in Alberta.** (Eng) Wigle, D. T. (Bureau Epidemiology, Room 230A, Lab. Centre Disease Control, Ottawa, Ontario K1A 0L2, Canada); Grace, M.; Smith, E. S. *Can Med Assoc J* 118(10): 1279-1280; 1978.

Persons first attending cancer clinics in Alberta, Canada, between 1971 and 1973 were interviewed to determine whether the risk of cancer of the uterine corpus (CUC) was increased among women with a history of estrogen therapy. Two hundred and two women aged 55 to 74 yr with histologically confirmed CUC were compared with 1,243 55- to 74-yr-old women with primary cancers other than cancer of the breast, cervix, uterus, ovary, or other female genital organs. Estrogen use was reported by 47.2% of the women in the case group and by 26.3% of the comparison subjects. Use of estrogen for at least 5 yr was reported by 23.9% of the users in the case group and 5.5% of the users in the comparison group. Women who had used estrogens at some time were at a significantly higher relative risk of CUC than were women who had never used estrogens, and the risk was higher for current users than for former users. The risk increased as the duration of hormone use increased, and it was especially high among women who reported use for ≥ 5 yr. About 24% of the CUC cases may have been due to current or past use of estrogen. The rate of increased incidence over time among women in Alberta and Saskatchewan was substantially greater for the age group 55 to 74 yr than for the age group 35 to 54 yr. The data suggest that the use of estrogens for the treatment of menopause-related symptoms should be reconsidered. (11 refs)

- 78-5356 **Adenocarcinoma of Cervix Uteri (Meeting Abstract).** (Eng) Boonyanit, S. (Dept. Obstetrics and Gynecology, Mahidol Univ., Siriraj Hosp., Bangkok, Thailand). In: *Summary Program and Abstracts of Paper Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control, Manila, 26-30 September, 1977*. Philippine Cancer Society (Manila, Philippines): 184 pp.; 79-80; 1977. (no refs)

- 78-5357 **Epidemiologic Features of Canine Ovarian Neoplasms.** (Eng) Hayes, H. M. (Environmental Epidemiology Branch, NCI, Bethesda, MD, 20014); Young, J. L. *Gynecol Oncol* 6(4): 348-353; 1978.

The epidemiologic features of histologically confirmed ova-

neoplasms diagnosed in 94 dogs seen by 13 North American veterinary university hospitals and clinics in March 1974 through June 1976 are described. They are classified by cell type or origin, epithelial tumors presented 60% of the case series and granulosa-theca tumors, 27%. Malignant tumors predominated 66%, excluding those of mesenchymal and unknown origin. The risk of epithelial ovarian neoplasms increased strikingly with age, but the risk of granulosa-theca tumors was constant through middle age. Pointer breed dogs had a statistically significant risk of epithelial ovarian tumors, and English Bulldogs had a significant risk of granulosa-theca cell tumors. Mongrel dogs were significantly freed from malignant tumors, but they had about the same risk as purebreeds for benign ovarian tumors. The epidemiologic features of ovarian neoplasia in dogs appeared similar to features noted in women. (21 refs.)

78-5358 **Faecal Bile Acids and Clostridia in the Aetiology of Colorectal and Breast Cancer (Meeting Abstract).** (Eng) Blackwood, A. (Dept. Clinical Oncology, Western Infirmary, Glasgow, Scotland); Murray, W. R.; McKay, C.; Calman, K. C. *Br J Cancer* 38(1): 175; 1978. (2 refs.)

78-5359 **Breast Cancer in Thailand (Meeting Abstract).** (Eng) Sa-Ngobwarchar, P. (National Cancer Institute, Bangkok, Thailand); Tangthatswasd, K.; Sontipong, S. *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 25-26; 1977. (no refs)

78-5360 **Case-Control Study of Genetic Admixture and Breast Cancer in American Black Women (Meeting Abstract).** (Eng) Petrakis, N. L. (G. W. Hooper Foundation, Univ. California, San Francisco, CA, 94143); Spuy, M. E.; Wong, R.; King, M. C. *In: Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 24; 1977. (no refs)

78-5361 **An Epidemiological Investigation of Cancer of Cervix in Thailand (Meeting Abstract).** (Eng) Panyanit, S. (Dept. Obstetrics and Gynecology, Siriraj

Hosp., Mahidol Univ., Bangkok, Thailand). *In: Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 28; 1977. (no refs)

78-5362 **The Replacement Hormone Use and Breast Cancer (Meeting Abstract).** (Eng) Choi, N. W. (Univ. Manitoba, Winnipeg R3E OW3, Canada); Nelson, N. A.; Shalev, C.; Abu Zeid, H. A. *Am J Epidemiol* 108(3): 230; 1978. (no refs)

78-5363 **Evidence Against the Association of Breast Cancer and DES Exposure (Meeting Abstract).** (Eng) Brian, D. (Mayo Clinic, Rochester, MN, 55901); Tilley, B.; Labarthe, D.; O'Fallon, W. M.; Noller, K. *Am J Epidemiol* 108(3): 230; 1978. (no refs)

78-5364 **Genetic Epidemiology of Families with High Incidence of Breast Cancer (Meeting Abstract).** (Eng) King, M. C. (Univ. California, Berkeley, CA, 94720); Elston, R. C.; Lynch, H. T. *Am J Epidemiol* 108(3): 230-231; 1978. (no refs)

78-5365 **Breast Cancer in Young Women: The Effect of Prior Pregnancy (Meeting Abstract).** (Eng) Janerich, D. T. (New York State Dept. Health, Albany, NY, 12237); Flink, E. M.; Glebatis, D. *Am J Epidemiol* 108(3): 235; 1978. (no refs)

78-5366 **Hypothesis on Relationship Between Breast Carcinoma and Diet. Analysis of Secondary Statistical Material.** (Ger) Vutuc, C. (Hygiene-Institut, Universitat Wien, Kinderspitalgasse 15, A-1095 Vienna, Austria); Kunze, M. *Wien Med Wochenschr* 128(14): 428-430; 1978.

Possible relationships between breast cancer mortality in women and diet were studied in Austria. In this country, breast cancer mortality has increased by 52% from 1957 to 1974. Concomitantly, the consumption of fat and protein of animal origin has increased by 25.4% and 27%, respectively. These data show a significant positive correlation between increased breast cancer mortality and fat and protein consumption. The findings are in agreement with those observed

in other countries. Among the other factors examined, young age at the first childbirth had no significant influence on breast cancer mortality. (27 refs.)

- 78-5367 Nutrition and Carcinoma of the Breast: Preliminary French Investigation.** (Fre) Le, M. (INSERM, Unite de Recherches statistiques, 16 bis, avenue P. Vaillant-Couturier, F 948000 Villejuif, France); Flamant, R. *Bull Cancer (Paris)* 65(1): 65-69; 1978.

A French survey is being conducted to determine if diet is a risk factor in the development of breast cancer in postmenopausal women. A preliminary evaluation of the first 654 questionnaires showed no differences between women with breast cancer and a population of age matched controls with regard to wt, height, or various indices of obesity. (16 refs)

- 78-5368 Bilateral Breast Cancer and Familial Aggregations.** (Eng) Sakamoto, G. (Dept. Pathology, Cancer Inst., Tokyo 170, Japan); Sugano, H.; Kasumi, F. *Prev Med* 7(2): 225-229; 1978.

A study was made of the incidence of primary bilateral breast cancer and familial aggregations of this condition among 3,365 Japanese breast cancer patients who underwent mastectomy between 1946 and 1975. The findings were compared with those in Caucasian women. Of the 3,365 patients, 92 had primary bilateral breast cancer; 23 of these cases were simultaneous. The crude incidence of bilateral cancer was 2.7%, compared with 3.0%-6.8% among American women. The true annual incidence rate of breast cancer after surgery for unilateral breast cancer was 3.4/1,000 Japanese patients at risk. Values for American women range from 5.8 to 7.1/1,000. Of the 92 patients with bilateral breast cancer, 13 had a family history of breast cancer among first- to third-degree relatives. In contrast, the familial incidence was only 4.1% among unilateral breast cancer patients. Three of the 13 patients showed multiple familial occurrences, with more than three women being affected within a given family. As many as 46.7% of the bilateral breast cancer patients were nulliparous compared with 18.4% of the unilateral breast cancer patients. Fibrocystic disease was found in association with the cancer in 60% of those in the bilateral group compared with 23% of those in the unilateral group. (8 refs)

- 78-5369 Mammographic Patterns and Risk of Breast Cancer.** (Eng) Hainline, S. (Duke Comprehensive Cancer Center, Durham, NC, 27710); Myers, L.; McLeland, R.; Newell, J.; Grufferman, S.; Shingleton, W. *Am J Roentgenol* 130(6): 1157-1158; 1978.

The mammograms 171 women with breast cancer and a and race-matched asymptomatic controls were evaluated to check a previous claim that breast cancer risk depends on mammographic classifications N1 (normal breast), P1 (lesser and greater degrees of ductal prominence), and DY (dense breast extensively involved with dysplasia). For each subject, the mammogram of only one breast was examined for all but two of the cancer patients, the mammograms of the negative breast were used. The distribution of the four categories among the breast cancer patients was consistent with the distribution noted previously. Age-standardized ratio estimates of the categories, relative to N1, were 1.5 for P1, 2.7 for P2, and 7.2 for DY. The cases and controls differed significantly with respect to the distributions of the four categories. It is concluded that ductal prominence and dysplasia, as determined by mammography, are significantly related to breast cancer risk. (6 refs)

- 78-5370 Abnormal Development of the Human Genital Tract following Prenatal Exposure to Diethylstilbestrol.** (Eng) Herbst, A. L. (Dept. Obstetrics & Gynecology, Univ. Chicago, Chicago, IL, 60637); Scully, E.; Robboy, S. J.; Welch, W. R.; Cole, P. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 6 pp.; 399-412; 1977.

Data from the Registry of Clear Cell Adenocarcinoma of the Genital Tract in Young Females were used to study diethylstilbestrol (DES)-related abnormalities of the female genital tract. Among 244 registry cases with clear cell carcinoma, 63% had been exposed prenatally to DES or similar preparations and 10% had been exposed to an unknown medication given because of high-risk pregnancy. Most had been exposed before the 18th week of gestation, and the total amount of drug taken during pregnancy ranged from 135 to 18,200 mg. The first case was diagnosed in 1961; the number of cases began to rise strikingly in the late 1960's, and the greatest number occurred in 1972. There was a sharp rise in the age-specific rate beginning at age 14; it peaked at age 19 and then declined sharply through age 22. The rise in incidence at age 14 suggests that stimuli associated with the onset of puberty may be related to the development of many of the tumors. The range of the absolute risk of development of adenocarcinoma in a DES-exposed female is 0.14 to 1.4/1,000. These tumors are considered curable. Nonneoplastic abnormalities commonly found in DES-exposed females are vaginal adenosis, usually associated with inflammation and squamous metaplasia, and cervical ectropion. These abnormalities can be explained by a disturbance in the development of the müllerian epithelium and/or vaginal plate. It is likely that endogenous estrogens play a role in the development of tumors in DES-exposed females. (28 refs.)

5371 Multiple Primary Cancers and Prolonged Survival: Familial Colonic and Endometrial Cancers. (Eng) Lynch, H. T. (Creighton Univ. Sch. Medicine, 400 California St., Omaha, NB, 68178); Bardawil, W. A.; Morris, R. E.; Lynch, P. M.; Guirgis, H. A.; Lynch, J. F. *Dis Colon Rectum* 21(3): 165-168; 1978.

The case histories of a woman and her mother, each of whom developed multiple primary cancers, and who showed prolonged survival are presented. The proband was treated at the age of 36 for adenocarcinoma of the rectum. Both ovaries and fallopian tubes were removed at the time of colostomy; in one ovary there was a polypoid nodule that was classified as a small pedunculated fibroma. At age 41, she was treated for well-differentiated endometrial carcinoma. Her mother had been treated at age 46 for bilateral papillary adenocarcinoma of the ovaries and adenocarcinoma of the uterus, at age 59 for colonic malignancy, at age 63 for a poorly differentiated mucinous adenocarcinoma of the sigmoid colon, at age 67 for adenocarcinoma of the transverse colon, at age 70 for a malignancy of the transverse colon, at age 73 for recurrent adenocarcinoma of the large bowel, and at age 75 for chronic myelogenous leukemia. She died at age 79. Analysis of the proband's family pedigree indicated that seven persons with cancer (including the proband and her mother) were direct descendants of a maternal grandmother who had died of colonic cancer at age 58. Another of the seven had developed multiple primary tumors of the colon at ages 27 and 48. The spectrum of tumors occurring at an early age among relatives of this kindred supports a genetic etiology for the tumors. (13 refs)

5372 Diet in the Epidemiology of Cancer of the Colon and Rectum. (Eng) Graham, S. (Dept. Social and Preventive Medicine, State Univ. New York at Buffalo, 100 Main Building, Buffalo, NY, 14261); Dayal, H.; Swanson, M.; Mittelman, A.; Wilkinson, J. *Natl Cancer Inst* 61(3): 709-714; 1978.

The diets of 256 male patients with cancer of colon (CC) and 100 male patients with cancer of the rectum (CR) were examined and compared with those of control patients with non-neoplastic, nondigestive diseases. There were 783 controls for the CC patients and 628 controls for the CR patients, and their age distributions were similar to those of the respective cancer patient group. No increase in risk for CC or CR was observed regardless of the amounts of beef or other meats consumed. However, an increase in the risk of CC was associated with a decrease in the frequency with which vegetables were eaten. A study of 214 women with CC and 182 women with CR yielded similar results. The decrease in CC risk associated with frequent ingestion of vegetables, especially cabbage, Brussels sprouts, and broccoli, is consistent with the decreased numbers of tumors observed in animals challenged with carcinogens and fed compounds found in these vegetables. Further studies to determine the effects of the possible

contamination of peanut butter by aflatoxins indicated an increased relative risk of CR in men who were currently eating peanut butter, and the risk appeared to increase with years of use. Ingestion of foods salted both during and after cooking was associated with an increased relative risk of colon cancer in men. Several other characteristics of diet were examined, but only slightly suggestive differences were found. (30 refs)

78-5373 Chronic Anacid Gastritis and Cancer of the Stomach. (Rus) Druzova, T. A. (Dept. Epidemiology of Malignant Tumors, Cancer Res. Center, Moscow, USSR). *Vopr Onkol* 24(7): 103; 1978.

The results of a retrospective study of the role of chronic anacid gastritis in the etiology of stomach cancer are presented. Of 5,000 patients with anacid gastritis, 47 developed stomach cancer. Most (65%) of the tumors were located in the corpus; in contrast, 90% of patients with a history of hyperacid gastritis develop cancer in the antral region of the stomach. (no refs.)

78-5374 Comparative Study on the Bowel Transit Times in Populations with Different Gastric Cancer Mortalities. (Jpn) Shimada, A. (Dept. Hygiene, Akita Univ. Sch. Medicine, Akita, Japan); Kamiyama, S. *Jpn J Hyg* 32(6): 712-716; 1978.

Bowel transit times (BTT) and other bowel behaviors were studied at Chokai, Akita Prefecture, and Ohasama, Iwate Prefecture. The male and female gastric cancer mortality of the Chokai inhabitants was 6 and 1.6 times that of the inhabitants of Ohasama, respectively. Transit times of at least 80% excretion of marker pellets were used as an indicator of BTT. BTT of the Akita inhabitants was 34.09 hr, but that of the Iwate inhabitants was 43.89 hr. Frequency of excretion showed the same trend as BTT, with 1.79 (Akita) and 2.20 (Iwate) movements required for excretion of at least 80% of the pellets. About 65% of the stool markers were eliminated at the first movement of the Akita inhabitants, compared with only 50% for the Iwate inhabitants. No differences in frequency of bowel movement were observed between the two regions, but wide individual differences were observed. Stool wt per excretion and stool wt per day were greater for the Akita inhabitants, and these people had a significant correlation between BTT and stool wt. A positive correlation between stool wt per day and 5-day frequency of excretion was noted only in Akita. (7 refs)

78-5375 Epidemiology of Renal and Ureteral Cancer in Rochester, Minnesota, 1950-1974, with Special

Reference to Clinical and Pathologic Features. (Eng) Resseguie, L. J. (Dept. Medical Statistics, Mayo Clinic, Rochester, MN); Nobrega, F. T.; Farrow, G. M.; Timmons, J. W.; Worobec, T. G. *Mayo Clin Proc* 53(8): 503-510; 1978.

An analysis was made of the epidemiology of renal and ureteral cancer in Rochester, Minnesota, during 1950-1974 and of the survival of these patients using the thorough medical records available for this population. Ninety cases were identified, 75 renal cell carcinomas (Group 1) and 11 pelvic and 6 ureteral carcinomas (Group 2). For women in Group 1, the incidence of renal carcinoma increased from 1/100,000 in 1950-1954 to 12/100,000 in 1970-1974; no clear trend was seen among men. In Group 1, 27 cases were diagnosed at autopsy (listed as primary cause of death in only 7), and about half of all 73 cases lacked specific symptoms that would focus attention on the urinary tract. Gastrointestinal symptoms were the major complaint in 18/46 cases diagnosed during life. In this group, no patient in whom metastasis was detected at diagnosis survived as long as 5 yr, but patients with carcinoma localized to the kidney cortex (Stage I) had 5- and 10-yr survival rates approx 87% of that expected from the age and sex distribution at diagnosis. Metastases were found in 7/27 cases diagnosed at autopsy and, at last follow-up, in 25/46 cases diagnosed during life. In Group 2, the survival rate was 41% of that expected at 5 yr and 56% at 10 yr. Sixteen patients had symptoms that drew attention directly to the tumor, and all 17 were diagnosed during life. Primary bladder malignancies were found in eight of these patients, but none were found in Group 1. (28 refs.)

78-5376 The Pattern of Malignant Disease of the Pancreas in Sri Lanka (Meeting Abstract). (Eng) Attygalle, D. (Faculty Medicine, Sri Lanka Cancer Society, Kynsey R. D. Colombo 8, Sri Lanka). In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 80; 1977. (no refs)

78-5377 Pleural and Peritoneal Mesotheliomas. (Ger) Weicksel, P. (Facharzt für Lungen- und Bronchialheilkunde, Kaiserstrasse 26, D-8700 Würzburg, W. Germany). *Med Klin* 72(47): 2005-2011; 1977.

A total of 1,687 cases of pleural and peritoneal mesothelioma were registered in different parts of the world during 1963-1976. Of this number, 986 cases were attributed to asbestos exposure, including 744 cases of occupational exposure. (32 refs.)

78-5378 Lung Cancer Epidemiology (Letter to Editor). (Dan) Wilson, K. (Strandby, Denmark). *Ugeskr Laeg* 140(11): 634-636; 1978.

Lung cancer was diagnosed in 15 men who had lived in Strandby, Denmark, (population approx 1,000) for $\geq$ 10 yr. All cases were clustered in a small area (population 200) about 500 m from an asphalt factory built 15 yr previously (no refs.)

78-5379 Morbidity due to Cancer of the Nasal Cavity and Maxillary Sinuses in Altai Oblast (USSR) During 1961-1970. (Rus) Zavarzin, A. A. (Altai Regional Clinical Hosp., Barnaul, USSR). *Zh Ushn Nos Gorl Bolezni* 1: 51-54; 1978.

The incidence of cancer of the nasal cavity and maxillary sinus was studied in Altai Oblast, USSR. From 1961 to 1970 there were 164 primary cases of these cancers, which comprised 0.3% of the total cancer morbidity in the district. 17.3% of the otorhinolaryngological cancer morbidity. The group included 75 men and 89 women. Max morbidity was detected among men $>$ 50-yr-old and among women $>$ 60-yr-old. Of the 125 patients studied histologically, 76 carcinomas, 47 sarcomas, 1 melanoma, and 1 plasmacytoma. Stage I was detected in 9 patients, Stage II in 13, Stage III in 80, and Stage IV in 13. (17 refs)

78-5380 Nasopharyngeal Carcinoma in Alaskan Natives (Eskimo, Indian, Aleut) 1966-1976 (Meeting Abstract). (Eng) Lanier, A. (Alaska Natives Medical Center, P.O. Box 7-741, Anchorage, AK 99510). In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 170 pp.; 70-71; 1977. (no refs)

78-5381 Oesophageal Carcinoma in Sri-Lanka-Lanka Deficiency as a Possible Cause of Its High Incidence and Its Unusually Common Occurrence in Young Females (Meeting Abstract). (Eng) Ganeshananthan, N. (Dharmapala Maravite, Ratnapura, Sri-Lanka). In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 168 pp.; 68-69; 1977. (no refs)

78-5382 **Association of Various Chewing and Smoking Habits with Carcinoma of the Oesophagus (Meeting Abstract).** (Eng) Jafarey, N. A. (Jinnah Postgraduate Medical Center, Karachi, Pakistan); Zaidi, S. H. In: *Third International Symposium on Oncology Held by Taj Pahlavi Cancer Institute in Tehran, 4-8 March, 1978*. Taj Pahlavi Cancer Institute, Pahlavi Medical School, Tehran University. Tehran, Iran): 115 pp.; 105; 1978. (no refs)

78-5383 **The Frequency of Thyroid Gland Carcinoma in Tehran. A Comparison of Surgically Removed Thyroid Gland and Autopsy Findings (Meeting Abstract).** (Eng) Bahadori, M. (Dept. Pathology, Tehran Univ., Tehran, Iran); Bahadori N. P. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 88-89; 1977. (no refs)

78-5384 **Malignant Neoplasms Following Cardiac Transplantation.** (Eng) Krikorian, J. G. (Div. Oncology, Stanford Univ. Sch. Medicine, Stanford, CA, 94305); Anderson, J. L.; Bieber, C. P.; Penn, I.; Stinson, E. B. *JAMA* 240(7): 639-643; 1978.

Malignant neoplasms developed in 7/124 patients receiving cardiac allografts between January 1968 and April 1977 at Stanford University Medical Center. The mean and median periods of follow-up were 18.3 and 9.7 mo, respectively. There were 3 lymphoproliferative neoplasms, 2 skin cancers, 1 acute leukemia, and 1 colon carcinoma. Of the 35 deaths occurring > 3 mo after transplantation, 4 were due to malignant neoplasms. Comparison of the patients with and without malignant neoplasms showed no significant differences with respect to sex, underlying disease, or blood type. Malignant neoplasms occurred with increased frequency in patients bearing histocompatibility antigens A1, B8, and A1 + B8. The acute myelogenous leukemia may have been due to immunosuppression or therapy with cyclophosphamide. The patients with lymphoproliferative diseases were younger (av age, 20 yr) than the other patients (av age, 50 yr), and their tumors occurred earlier (av time, 33 mo after transplantation) than those in the other patients (av time, 53 mo). The incidence and spectrum of cancers were similar to those observed in recipients of renal homografts. (12 refs)

78-5385 **Mesothelioma in Relation to Asbestos Fibre Exposure: A Review of 70 Serial Cases.** (Eng) Cochrane, J. C. (Natl. Res. Inst. Occupational Diseases, S.

African Medical Res. Council, Johannesburg, S. Africa); Webster, I. *S Afr Med J* 54(7): 279-281; 1978.

The relationship of pleural mesothelioma to a history of asbestos exposure was established in 69/70 cases. In one case, it was not possible to elicit a history of significant exposure to asbestos dust. Malignant mesothelioma was confirmed by open thoracotomy biopsy specimens. Detailed environmental and occupational histories were obtained from the patients. Most of the patients were exposed occupationally (asbestos mines, locomotive workshops); the 13 patients with nonoccupational exposure lived on asbestos mine property or in mining areas, where fibers were more or less constantly in the air. The interval from exposure to tumor occurrence ranged from 9 to 40 yr. A minimum of 4 mo of constant exposure to an atmosphere of visible floating asbestos fibers or 4 yr in room contact with loose fibers not visible in the atmosphere was required for adequate and significant exposure. For cases of environmental exposure, a minimum of 3 yr residence in a mining area was required. (9 refs)

78-5386 **Cancer Family Syndrome. A Case Report and Literature Review.** (Eng) Boland, C. R. (US Public Health Service Hosp., 15 and Lake Sts., San Francisco, CA, 94118). *Am J Dig Dis* 23(5, Suppl): 25s-27s; 1978.

A 49-yr-old man, who had survived a right hemicolectomy for adenocarcinoma of the cecum 23 yr earlier, presented with three simultaneous colonic carcinomas in the transverse colon, sigmoid colon, and rectum. Pedigree analysis revealed that he was a member of a family with cancer family syndrome, an autosomal dominant genetic disease in which the affected members are at risk at early age for carcinoma, particularly of the colon and female genital tract. He was a member of a sibship of 13, 5 of whom had had colonic carcinoma at early ages. Endometrial or ovarian carcinoma had been documented in three sisters; one lung cancer and one brain tumor had also occurred among the siblings. Three sibs had solitary adenomatous colonic polyps detected during annual radiographic examinations. The proband's father underwent removal of carcinoma of the cecum at age 27, and he died of carcinoma of the rectum at age 45. The proband's paternal uncle had a colonic carcinoma at age 64, and a paternal aunt died of breast cancer at age 76. The proband's grandfather died of colonic cancer at age 41. No polyps or neoplasms have been documented among the descendants of this sibship. However, in the genetic lines at risk, all members are still under the age of 30. Carcinoembryonic antigen levels in all living members, determined by the Hansen method, were < 4.0 nanograms/ml. (16 refs)

78-5387 **Malignant Melanoma in Spouses.** (Eng) Mintzis, M. J. (Dept. Dermatology, New York Univ.

Sch. Medicine-Bellevue Medical Center, New York, NY); Berger, A. P.; Greenwald, E.; Greenwald, E.; Golomb, F. *Cancer* 42(2): 804-807; 1978.

A study of 743 white patients > 14 yr of age entered in the New York University-Bellevue melanoma registry between 1950 and 1976 revealed three married couples in which both spouses developed malignant melanoma. Statistical calculations based on data supplied by the Third National Cancer Survey and on the 1970 age distribution of married men and women in the US indicated that at most, 0.44 cases of melanoma would be expected to occur among the spouses of the patients. The difference was statistically significant. Various explanations for this occurrence of connubial melanoma are proposed. (1) One explanation is based on the association of melanoma with fair skin and with sun exposure. People tend to marry within their ethnic group, often of a distinctive skin type. All six members of the doubly affected couples had blond hair and blue eyes, features associated with fair skin. (2) Genetic factors or (3) a viral etiology could also be involved. However, the registry records had no information pertaining to any of these issues. (14 refs.)

78-5388 Actual Volume Doubling Time Values for Pulmonary Metastases from Soft Tissue Sarcomas.

(Eng) Plesnicar, S. (Inst. Oncology, Vrazov trg 4, 61000 Ljubljana, Yugoslavia); Klanjscek, G.; Modic, S. *Cancer Lett* 4(6): 311-316; 1978.

The doubling times of pulmonary metastases in 24 patients with histologically proved soft tissue sarcomas were determined. Among these patients there were 15 cases of fibrosarcoma, 5 of rhabdomyosarcoma, and 1 each of leiomyosar-

coma, liposarcoma, hemangiopericytoma, and angiosarcoma. Sixty metastases were measured by chest radiograms on consecutive days, and 116 doubling times were calculated. The mean doubling time was 69.9 days. Twenty-two doubling times were eliminated because they were > 100 days long and the mean for the 941 doubling time values was 37.7 days. Metastases with diameters ≤ 14.9 mm had a mean doubling time of 29.7 days; those with diameters of 15.0-111.0 mm had a mean of 43.4 days. Apparently, the slowing of growth rates of these metastases is size dependent. Of doubling times > 100 days, the mean was 207.6 days, with a range of 103.0-370.6 days. When these times were eliminated, the mean doubling time of the tumors was similar to the previously reported for pulmonary metastases of osteosarcoma. It is suggested that similar mechanisms govern growth rates of metastases from both types of sarcoma. (refs.)

MISCELLANEOUS

5389 **Infantile Histiocytosis X (Letterer-Siwe Disease).** Investigations with Enzyme-histochemical and Sheep-Erythrocyte Rosetting Techniques. (Eng) Elema, D. (Dept. Pathology, Univ. Groningen, Groningen, Netherlands); Poppema, S. *Cancer* 42(2): 555-565; 1978.

In biopsies and one lymph node specimen from three infants with histiocytosis-X (H-X, Letterer-Siwe disease) were analyzed enzyme histochemically for acid phosphatase (AP), adenosine triphosphatase (ATP), leucyl- β -naphthylamidase (LNA), α -naphthylacetate esterase and peroxidase and by sheep erythrocyte rosetting on frozen sections for complement C₃ and for the Fc fragment of IgG. Light microscopy showed a dense dermal and scattered epidermal infiltration of histiocytic cells. Electron microscopy showed the well-known features of H-X cells with often numerous Langerhans' cell granules. Most of the cells comprising the specimens had weak AP and weak or absent nonspecific esterase activity but considerable LNA activity. Although its biological significance is unknown, the uniform presence of LNA could seem to make it a useful marker for H-X cells. The ATP reaction was variable in intensity, ranging from negative to strongly positive, and the peroxidase reaction was uniformly negative in the infiltrating cells but was found sporadically in the eosinophils. Rosetting techniques indicated the presence of receptors for the Fc fragment of IgG but not for C₃. Cells with the same enzyme histochemical characteristics could be found in T-cell areas of normal spleen, normal and reactive lymph nodes, and thymic medulla but not in B-cell areas or in the thymic cortex. The enzyme analyses provide additional support for the close relation of H-X cells and Langerhans' cells. It is suggested that H-X cells belong to the mononuclear phagocyte system and that they are closely linked to the functioning of T-cell-mediated immunity. (44 refs)

8-5390 **Pregnancy and Sarcoma.** (Eng) Jafari, K. (Dept. Obstetrics and Gynecology, Cook County Hospital, 1825 W. Harrison St., Chicago, IL, 60612); Lash, A.; Webster, A. *Acta Obstet Gynecol Scand* 57(3): 265-271; 1978.

The case reports of six women (aged 15-39) whose pregnancies were complicated by sarcoma are presented. The types and location of the sarcomas were: pleomorphic sarcoma of the right leg with pulmonary metastases, osteogenic sarcoma, lymphosarcoma, malignant mixed mesodermal tumor of the vulva compatible with sarcoma botryoides, giant cell tumor

of the left leg, and liposarcoma of the thigh treated 2 yr previously--metastasis to the lungs appeared during pregnancy. All six women delivered viable infants. The first three women died of their disease within 90 days of delivery, the fourth was lost to follow-up, the fifth is alive with no evidence of disease, and the sixth is in remission. A review of the literature on similar cases is presented. Most authors believe that pregnancy has no deleterious effect on the course of cancer, but this is not supported by adequate data. Fetal or placental metastases are rare. Of 29 reports of maternal cancer metastasizing to the placenta or fetus, 14 were cases of malignant melanoma. (15 refs)

78-5391 **Neoplasia and Dysplasia of the Cervix Uteri and Contraception: A Possible Protective Effect of the Diaphragm.** (Eng) Wright, N. H. (Dept. Community Medicine, Rutgers Medical Sch., Rutgers, NJ); Vessey, M. P.; Kenward, B.; McPherson, K.; Doll, R. *Br J Cancer* 38(2): 273-279; 1978.

The hypothesis that use of a diaphragm is protective against cervical neoplasia (CN) was analyzed in a cohort of women (aged 25-39 yr) entered in a contraceptive study. Among 17,032 women, 65 were found with histologically proved neoplasia or dysplasia diagnosed after cone biopsy prior to September 1, 1977. Two had frankly invasive cancer, 4 microinvasive cancer, 33 carcinoma in situ, 26 dysplasia. The incidence rates of CN in diaphragm users, oral contraceptive users, and intrauterine device users were, respectively, 0.17, 0.95, and 0.87/1,000 woman-yr of observation. The negative relationship between diaphragm use and CN could not be explained in terms of the confounding variables of age, social class, smoking habits, age at first marriage, and age at first pregnancy. It also could not be attributed to a lower frequency of cervical smearing among diaphragm users. In more detailed study of 52 of the women with CN and 139 matched controls, information concerning age at first intercourse, numbers of sexual partners, and frequency of cervical smearing outside the clinics was obtained. In comparison with users of other methods of contraception, diaphragm users were less likely to have had coitus at an early age and they had had fewer partners. After adjusting for the effects of these variables, however, the risk of CN was still only about one-quarter that in the users of the other two methods. In this study, smoking emerged as a major risk factor for CN. Smoking habit may reflect some important aspect of sexual behavior relevant to the production of CN that the investigators have been unable to measure. (10 refs)

- 78-5392 Effects of Insulin, Human Placental Lactogen and Human Growth Hormone on DNA Synthesis in Organ Cultures of Benign Human Breast Tumours.** (Eng) Welsch, C. W. (Dept. Anatomy, Michigan State Univ., E. Lansing, MI, 48824); Patten, S. E.; McManus, M. J. *Br J Cancer* 38(2): 258-262; 1978.

The effects of bovine insulin (5.0 $\mu\text{g/ml}$), human placental lactogen (HPL: 10.0 $\mu\text{g/ml}$), and human growth hormone (HGH: 10.0 $\mu\text{g/ml}$) on ^3H -thymidine incorporation into DNA were studied in organ cultures of 19 benign human breast tumors from 19 premenopausal patients. An attempt was made to determine if HGH has mitogenic activities toward these tumors, as has been reported for HPL. Most of the tumors were classified histologically as fibrocystic disease with varying degrees of adenosis and ductal hyperplasia. There was no correlation between the histopathological diagnosis and response to the hormones. Insulin and HPL significantly ($P < 0.01$) increased the mean incorporation of ^3H -thymidine into DNA, but HGH was ineffective. The marked difference in the ability of HPL and HGH to stimulate DNA synthesis in benign human breast tumors, despite their close chemical and physiological properties, may provide insight into the structural entity of HPL required for human mammotrophic activities. The results provide evidence that HPL and insulin, but not HGH, may be important factors in the etiology of benign disease of the human breast. (11 refs)

- 78-5393 Migration and Morphological Changes Induced in Normal Cells Treated with Purified Extracts of Conditioned Medium from a Transformed Cell Line (Meeting Abstract).** (Eng) Stocker, M. (Friedrich-Miescher-Institut, P.O. Box 273, CH-4002 Basel, Switzerland); Burk, R. R. *Experientia* 34(7): 958; 1978. (no refs)

- 78-5394 Establishment of a Human Plasma Cell Line In Vitro.** (Eng) Burk, K. H. (Dept. Lab. Med., Univ. Texas System Cancer Center, M. D. Anderson Hosp. and Tumor Inst., Houston, TX, 77030); Drewinko, B.; Trujillo, J. M.; Ahearn, M. J. *Cancer Res* 38(8): 2508-2513; 1978.

The morphological, cytogenetic, ultrastructural, and cell kinetic parameters of a cell line (ARH-77) derived from the peripheral blood of a 33-yr-old woman with IgG plasma cell leukemia were evaluated. The line has been propagated in culture for 2 yr. Morphologically, the ARH-77 cells showed identifiable markers of plasma cells. This was confirmed by electron microscopy, which revealed the presence of substantial (although atypical) endoplasmic reticulum and numerous polyribosomes. An av of 40% cells were positive for IgG by direct immunofluorescence, while IgG-specific radioim-

munoassay revealed the production of 1.21×10^4 nanograms/ 10^6 plasma cells. The karyotype was aneuploid, with a modal chromosome number of 45-46 and no marked chromosome. Growth kinetic characteristics were: doubling time, 110.4 hr; generation time, 56.4 hr; $G_1 + G_2$ -phase transit time, 45.5; S-phase transit time, 10.9 hr; growth fraction, 74%; mitotic index 1.5%; labeling index, 14.3%; and cell loss, 31.0%. Some of the growth kinetic characteristics were markedly similar to those displayed in vivo by plasma cells of multiple myeloma patients. The ARH-77 cell line might be a useful in vitro model for the study of human myeloma. (15 refs)

- 78-5395 Separation of Cells from a Murine Fibrosarcoma on the Basis of Size. I. Relationship Between Cell Size and Age as Modified by Growth In Vivo In Vitro.** (Eng) Grdina, D. J. (Section Experimental Radiotherapy, Univ. Texas System Cancer Center, M. D. Anderson Hosp. and Tumor Inst., 6723 Bertner Ave., Houston, TX, 77030); Peters, L. J.; Jones, S.; Chan, E. *J Natl Cancer Inst* 61(1): 209-214; 1978.

Cells from a 3-methylcholanthrene-induced murine fibrosarcoma were separated and synchronized on the basis of cell size by centrifugal elutriation. The tumor cells were elutriated immediately after disaggregation of a solid tumor or after 4 hr of incubation in vitro. Cell recovery was $> 85\%$, and cell viability was $> 95\%$. No difference in recovery was observed between cells separated directly from the tumor or following in vitro incubation. Av cell volume, along with sedimentation rate, increased with increasing fraction number, indicating that separation was being achieved according to differences in cell size. The largest size variation was observed among cells incubated in vitro: modal volumes ranged from 740 to 2,400 μ^3 , and the modal volume of the whole population was 1,620 μ^3 . In contrast, cells separated without prior incubation varied in volume from 60 to 1,400 μ^3 , and the modal volume of the population was 980 μ^3 . The incubated tumor cells were more readily synchronized than cells separated directly from the tumor. The reduced synchrony of the latter reflects the heterogeneity of the cell sizes in the same phase of the division cycle present in the tumor growing in vivo. In contrast, tumor cells incubated in vitro are exposed to more uniform growth conditions, which minimizes physical differences among the various cell classes and makes the relationship between cell size and age more apparent. (14 refs)

- 78-5396 Highly Reiterated Sequences of SIMIAN SIMIAN SIMIAN SIMIAN SIMIAN.** (Eng) Rosenberg, H. (Israel Inst. Biological Res., Prime Minister's Office, Ness Ziona P.O.B. 19, Israel); Singer, M.; Rosenberg, M. *Science* 200(4340): 394-402; 1978.

72-base pair (bp) segment of DNA that is repeated several million times in the genome of the African green monkey was characterized. The DNA was digested with restriction endonuclease Hind III, and the products were fractionated by electrophoresis on polyacrylamide gels. A unique nucleotide sequence representing the most abundant residue at each of 172 positions, but not necessarily the structure of any particular member of the set, is described. The sequence is internally repetitive. The isolated fragments represent a mixture of molecules of identical chain length that differ by the presence of a nucleotide other than the most abundant one at one or more positions. The divergence from the determined sequence within the members of the set is not random. Direct quantitation and hybridization kinetics indicate that the set occurs several million times within the genome, but the reiteration frequency of particular members of the set is unknown. This uncloned segment was compared with a cloned segment of the same DNA that was recombined into the genome of a defective, substituted variant of simian virus during permissive infection. The cloned DNA preparation showed no evidence of divergence. The sequences of the cloned and uncloned DNA fragments were identical except at one position. Thus, the particular DNA may be derived from one of the more infrequent members of the set comprising the monkey fragments. (35 refs)

78-5397 **Cell Surface Markers on Pre-leukemic Cells. I. Soluble Proteins.** (Eng) Hollinshead, A. C. Dept. Medicine, George Washington Univ. Medical Center, Washington, DC, 20037; White, C. S.; Ladoulis, C. T.; Ellis, D.; Gilcher, R. O.; Lessin, L. S. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 256-259; 1978.

Profiles of soluble membrane proteins from WBC, RBC, and platelets of preleukemic patients with hemopoietic dysplasia (D) or myeloid metaplasia were compared with those of leukemic and normal individuals. Changes in these profiles were examined in relation to those seen during blast crisis and conversion to leukemia. (8 refs)

78-5398 **Theoretical Mechanisms for Synthesis of Carcinogen-induced Embryonic Proteins. II. Perturbations of Post-translational Nuclear Protein Modification.** (Eng) Hancock, R. L. (High River Inst. Theoretical Cancer Study, High River, Alberta, Canada). *Med Hypotheses* 4(3): 290-301; 1978.

It is hypothesized that carcinogenic chemicals initially induce

a derepression, typically by an alkylation reaction on specific acidic protein repressors, of structural genes of enzymes that modify the posttranslational structure of nuclear proteins. This causes a cascade of events leading to an array of phenotypic changes. Although massive genic changes in activity may result because of the large number of differentiated cells, only when these changes occur in a stemlike cell does the possibility of a neoplastic array of genic expression occur by virtue of its repressors being more vulnerable to conformational modification. It is also theorized that there are three major classes of repressors, that may be manipulated via the posttranslational modifying enzymes to become more or less weak, giving rise to a specific differentiated state. It is the repressors synthesized during embryogenesis that are particularly susceptible to derepression. Quasi-differentiated liver stem cells, eg, become relatively easily altered to express many embryonic genes and, if a particular required array of derepression occurs, the overall result would be a large variety of neoplastic clones. However, an organism would have only a limited number of stem cells capable of producing such clones. This would explain the small number of hepatomas produced during chemical induction, although a large number of cells are undoubtedly affected. (39 refs)

78-5399 **Growth Characteristics and Drug Responses of a Murine Lung Carcinoma In Vitro and In Vivo.** (Eng) Kaneko, T. (McEachern Lab., Cancer Res. Unit, Univ. Alberta, Edmonton, Alberta, Canada T6G 2H7); LePage, G. A. *Cancer Res* 38(7): 2084-2090; 1978.

The growth characteristics of a recently established mouse lung tumor line were studied, and several drugs in common clinical use for the treatment of lung cancers were tested in this model system. Nettesheim carcinomas were initially transplanted sc into male DBA/2 mice, and the resulting growths were cultured in vitro. Three cultures slowly developed monolayers of epithelial-like cells that produced tumors when inoculated into DBA/2 mice. The cells were subcultured 15 times, and cloning was carried out with one of the lines. A clone designated KLN205 was selected for further work because it grew relatively rapidly in vitro. This cell line was able to produce lung colonies in male DBA/2 mice when injected into a tail vein, and it could be transplanted sc in male DBA/2 and BD2F₁ mice. The sc tumors were composed of poorly differentiated epidermoid carcinoma cells with numerous mitotic figures. The tumors showed evidence of keratinization and frequent extensive areas of necrosis, with a narrow discontinuous zone of lymphocytic infiltration and fibrosis at the periphery. The antimetabolites arabinosyl-6-mercaptopurine and 6-selenoguanosine (SeGR) were used to treat KLN205 cells in vitro. The former was only marginally effective against the lung carcinoma, but the latter was highly effective. 5-Fluorouracil, thioguanine, methotrexate, and adriamycin also had some inhibitory effect in vitro, but methylcyclohexylnitrosourea had no effect. In vivo, only SeGR, thioguanine, and hexamethylmelamine

were significantly effective. This cell line is proposed as a potential model for testing other chemotherapeutic agents. (8 refs)

78-5400 Proteins Conferred by the Virulence-specifying Plasmid of *Agrobacterium tumefaciens* C-58. (Eng) Sonoki, S. (Dept. Plant Pathology, Univ. California, Davis, CA, 95616); Kado, C. I. *Proc Natl Acad Sci USA* 75(8): 3796-3800; 1978.

Membrane-associated and periplasmic proteins of *Agrobacterium tumefaciens* C-58 were compared with those from avirulent, nontumorigenic derivative strains by slab and

two-dimensional gel electrophoresis. Two proteins (Per-1 and Per-2) with mol wt of 37,500 and 37,000, respectively, were detected in the supernatant fraction of C-58 cells treated with EDTA and lysozyme. The same proteins were missing from an avirulent C-58 derivative that did not contain the 117-megadalton plasmid found in the virulent strain. When the derivative was mated with C-58, the resulting virulent transconjugants regained the large C-58 plasmid and expressed the Per-1 and Per-2 proteins. When the plasmid was removed from the transconjugants, they became avirulent and did not express Per-1 and Per-2. The functional roles of these proteins are unknown, but they are associated with the outer membrane and periplasmic fraction of the *Agrobacterium* cells. The Per-1 and Per-2 proteins appear to be plasmid-coded gene products that may be involved in, but not the sole determinants of, tumorigenesis. (23 refs.)

- Aaronson, S. A., 78-4885, 78-5208
 Abdo, S., 78-5087
 Abel, J., 78-5118
 Abercrombie, B., 78-5031
 Abu Zeid, H. A., 78-5362
 Ada, G. L., 78-4905
 Adam, E., 78-4888
 Adams, R. L., 78-5247
 Adamson, R. H., 78-5148
 Adekunle, A. A., 78-4963
 Adolphs, H. D., 78-4976
 Agrelo, C., 78-5073
 Ahearn, M. J., 78-5394
 Ahmad, M., 78-4878
 Ahmed, F. E., 78-5169
 Ahuja, M. R., 78-5087
 Akaeva, E. A., 78-5141
 Akbar, N., 78-5332
 Allen, J. R., 78-4829
 Allison, A. B., 78-5343
 Aloni, Y., 78-5267
 Altavilla, G., 78-5263
 Ames, B. N., 78-4809, 78-5166
 Amos, J., 78-4997
 Anders, A., 78-5087
 Anders, F., 78-5087
 Andersen, M. E., 78-5109
 Anderson, J. J., 78-5170
 Anderson, J. L., 78-5384
 Anderson, M., 78-5015
 Ando, K., 78-4973
 Andrews, A. W., 78-5097
 Andronikashvili, E. L., 78-5085
 Angsubhakorn, S., 78-4970
 Antoku, S., 78-5181
 Antonenkov, V. D., 78-5014
 Aoi, Y., 78-5234
 Appleton, D. R., 78-5059
 Arai, T., 78-5093
 Arai, Y., 78-5335
 Arce, B., 78-5312
 Archer, M. C., 78-4818
 Arjungi, K. N., 78-5082
 Arlinghaus, R. B., 78-5202
 Armstrong, B. K., 78-4862
 Armstrong, S. J., 78-5229
 Arumugam, S., 78-4992
 Arunagiri, R., 78-5044
 Ashby, J., 78-5150
 Astedt, B., 78-5341
 Astier, T., 78-5225
 Attygalle, D., 78-5376
 Audebert, A. A., 78-5153
 Augenlicht, L. H., 78-4827
 Aulakh, G. S., 78-4901
 Awa, H., 78-5093
 Aymard, J. P., 78-5140
 Baba, N., 78-5179
 Baba, T., 78-5103
 Babudri, C. N., 78-4950
 Bacot, C., 78-4989
 Bahadori, M., 78-5383
 Bahadori N. P., 78-5383
 Baird, W. M., 78-5023
 Bakshi, K., 78-5078
 Balanski, R. M., 78-5056
 Balboni, P. G., 78-5263
 Balda, B. R., 78-5221
 Balick, R., 78-5027
 Balkin, R. B., 78-5311
 Ballou, J. E., 78-5185
 Baltimore, D., 78-5197, 78-5214
 78-5215
 Bannasch, P., 78-5074
 Bansal, B. R., 78-5061
 Bansal, S. C., 78-5061
 Bar-Shavit, R., 78-5267
 Barbacid, M., 78-5208
 Barbanti-Brodano, G., 78-5262, 78-5263
 Bardawil, W. A., 78-5371
 Bardin, C. W., 78-5078
 Barthel, E., 78-4835
 Bartram, C. R., 78-4931
 Bartsch, H., 78-5049
 Baserga, R., 78-5258
 Bassir, O., 78-4963
 Batzinger, R. P., 78-4845
 Bauer, F. L., 78-5159
 Baugnet-Mahieu, L., 78-5225
 Baumgartner, I., 78-5246
 Beales, N., 78-5252
 Begue, J. Y., 78-5130
 Belobragina, G. V., 78-4971
 Ben Bassat, H., 78-5302
 Ben Zvi, A., 78-5302
 Benard, Y., 78-5326
 Benbassa, A., 78-5131
 Bend, J. R., 78-4990, 78-5012, 78-5013
 78-5018
 Bender, A., 78-5315
 Bender, M. A., 78-5145
 Bendinelli, M., 78-5207, 78-5283
 Beniashvili, D. S., 78-5085
 Benjamin, T. L., 78-5268
 Benner, R., 78-5280
 Benoit, F., 78-5070
 Benton, B., 78-4865
 Benveniste, R. E., 78-5231
 Benz, R. D., 78-4939
 Berenson, M. M., 78-5322
 Berg, J. W., 78-4908
 Berg, P., 78-5252
 Berger, A. P., 78-5387
 Berman, M. M., 78-5333
 Betts, T. E., 78-5046
 Beug, H., 78-5195
 Bhamarapravati, N., 78-4970
 Bhatt, T. S., 78-4996
 Bieber, C. P., 78-5384
 Biessmann, H., 78-4827
 Bigard, M. A., 78-5130
 Bilyk, I., 78-5035
 Bishop, J. M., 78-4882
 Bishun, N. P., 78-5137
 Bister, K., 78-5187
 Bjorkholm, M., 78-5300
 Black, O., 78-4991
 Black, P. H., 78-5253
 Blackwood, A., 78-5358
 Block, N. L., 78-4977, 78-5076
 Blot, W. J., 78-5344
 Blough, H. A., 78-5192
 Blum, A., 78-5166
 Bocker, W., 78-5316
 Boland, C. R., 78-5386
 Bolt, H. M., 78-4833
 Bondy, G. P., 78-5139
 Bone, E. S., 78-5330
 Bonmassar, E., 78-5294
 Boonyanit, S., 78-5356, 78-5361
 Borchert, P., 78-4806
 Borghetti, A. F., 78-5259
 Borgia, R., 78-5245
 Bornstein, W. A., 78-5013, 78-5019
 Bouck, N., 78-5252
 Bourdon, A. T., 78-4959
 Bourgaux, P., 78-5265
 Boutwell, R. K., 78-5037
 Bowen, E. R., 78-4990
 Bowman, M. C., 78-4972
 Braun, D. C., 78-4805
 Braun, H., 78-5066
 Breeman, G. E., 78-4982
 Breitburd, F., 78-4889
 Bresnick, E., 78-5013, 78-5019
 Brian, D., 78-5363
 Brockman, H. E., 78-4947
 Brodsky, I., 78-5204
 Brookes, P., 78-4997
 Brooks, T. L., 78-5250
 Brotosaputro, B., 78-5345
 Brotz, M., 78-5233
 Bruce, W. R., 78-5052, 78-5121
 Brusick, D., 78-5078
 Bucheler, J., 78-5090
 Buckley, J., 78-4828
 Bueding, E., 78-4845
 Bullock, L., 78-5078
 Buncher, C. R., 78-5088
 Bunick, E. M., 78-5318
 Burch, T., 78-5347
 Burk, K. H., 78-5394
 Burk, R. R., 78-5393
 Burny, A., 78-5230
 Burrell, A. D., 78-5170
 Butler, W. H., 78-4840, 78-5325
 Cachaza, J. M., 78-5077
 Caillard, J. F., 78-5326
 Caillouette, J. C., 78-5321
 Cairns, J., 78-5102
 Calegari, L., 78-5003
 Calendi, E., 78-5003
 Calman, K. C., 78-5358
 Calothy, G., 78-5194
 Campa, M., 78-5207
 Campanile, F., 78-5294
 Carrascal, E., 78-5306
 Carroll, D. I., 78-5166
 Casado, J., 78-5077

- Casagrande, J., 78-5209
 Casagrande, J. T., 78-4910
 Casey, H. W., 78-5132
 Castegnaro, M., 78-5049
 Casto, B. C., 78-5117
 Castro, A., 78-5077
 Cavallieri, L. F., 78-5189
 Cavalli, F., 78-5299
 Cavanaugh, D., 78-4988
 Cayama, E., 78-5084
 Cecchetti, M. G., 78-5263
 Cerutti, P., 78-5007
 Cerutti, P. A., 78-5009
 Chadwick, K. H., 78-4879
 Chambon, P., 78-5257
 Champetier, J., 78-5131
 Chan, E., 78-5395
 Chang, R. L., 78-4995, 78-5022
 Chang, S. S., 78-5228
 Chang, W. W., 78-5064
 Chaouat, Y., 78-5153
 Chaponnier, C., 78-5245
 Charbeneau, N. J., 78-5114
 Charman, H., 78-5274
 Charmot, G., 78-4911
 Chasles, J., 78-5326
 Cheever, A. W., 78-4846
 Chemerys, R., 78-5023
 Chen, L. B., 78-4902
 Chen, Z., 78-5335
 Chern, C. J., 78-5023
 Chestier, A., 78-5256
 Chetiyawardana, A. D., 78-5147
 Chew, E. C., 78-5273
 Chiang, W., 78-5035
 Chiazze, L., 78-5348
 Choi, N. W., 78-5362
 Chorpenning, F. W., 78-5086
 Chottard, J. C., 78-4989
 Chouroulinkov, I., 78-5149
 Christov, K., 78-5183
 Chuang, A. H., 78-5019
 Chuang, H., 78-5013
 Chutna, J., 78-5282
 Cicek, J., 78-4843
 Cinco, M., 78-4950
 Clapp, M. J., 78-5047
 Claviez, M., 78-5195
 Cleaver, J. E., 78-5172
 Clemmesen, J., 78-5351
 Cleuter, Y., 78-5230
 Clifton, G., 78-5016
 Clinton, S. K., 78-5002
 Coates, P. W., 78-5089
 Cochrane, J. C., 78-5385
 Cohen, L. A., 78-4912
 Cole, P., 78-5370
 Collett, M. S., 78-5188, 78-5191
 Conney, A. H., 78-4854, 78-4855
 78-4995, 78-5022
 Conning, D. M., 78-5047
 Contreras, R., 78-5249
 Cooke, S. A., 78-5314
 Coombs, M. M., 78-4996
 Corallini, A., 78-5263
 Correa, P., 78-4818, 78-5148
 Cotlier, E., 78-5309
 Cotter, S. M., 78-4885
 Couch, D. B., 78-5158
 Craddock, V. M., 78-4949
 Cravioto, H., 78-5092
 Creasia, D. A., 78-5043
 Cremisi, C., 78-5256
 Croissant, O., 78-4889
 Croisy, A., 78-4938
 Crosby, N. T., 78-4959
 Croy, R. G., 78-4962
 Cuello, C., 78-4818
 Cummings, J. H., 78-5123
 Czesnin, K., 78-5178
 Dagle, G. E., 78-5045, 78-5185
 Dahlman, R. C., 78-5184
 Dalbey, W. E., 78-5043
 Daley, M. J., 78-5289
 Dalgard, D. W., 78-5148
 Dambrine, G., 78-5194
 Daune, M. P., 78-4986
 Davisson, P., 78-5211
 Day, N., 78-5049
 Dayal, H., 78-5372
 Daynes, R. A., 78-5285
 de Laat, A. M., 78-5280
 De Larco, J. E., 78-5220
 de Micco, C., 78-5266
 de Pasqualini, C. D., 78-4897
 de Serres, F. J., 78-4947, 78-4948
 de-The, G., 78-4895
 Debray, J., 78-5153
 Deckers-Schmelzle, B., 78-5010
 Defais, M., 78-5102
 Deinhardt, F., 78-5239
 Delage, C., 78-5307
 Delbecchi, L., 78-5265
 Demopoulos, N., 78-4944
 Den Engelse, L., 78-4935
 Deng, C., 78-5213
 Deng, C. T., 78-4882
 Deobagkar, D. N., 78-5199, 78-5201
 Deubelbeiss, K., 78-5299
 di Mayorca, G., 78-4893, 78-5252
 Diamond, L., 78-5023, 78-5038
 Diaz de Molnar, A. M., 78-5339
 Diesendorf, M., 78-4868
 Dieter, H., 78-5118
 Digges, I., 78-5247
 Dilman, V. M., 78-4906
 Dion, A. S., 78-5269
 Dionyssiou-Asteriou, A., 78-5161
 DiPaolo, J. A., 78-4987, 78-5117
 Doerfler, W., 78-5233, 78-5237
 Doll, R., 78-4925, 78-5391
 Donoghue, D. J., 78-5214
 Donohue, T. A., 78-5333
 Dore, J. F., 78-5064
 Dostal, V., 78-5243
 Dougherty, M. F., 78-5209
 Dougherty, R. C., 78-5166
 Drasar, B. S., 78-5123
 Drewinko, B., 78-5394
 Drinkwater, N. R., 78-4984
 Druzova, T. A., 78-5373
 Duplan, J. F., 78-5225
 Dupuy, M. E., 78-5360
 Durand, A., 78-5131
 Duverger-van Bogaert, M., 78-5111
 Dzidic, I., 78-5166
 Eason, R., 78-5247
 Eddie, H., 78-5137
 Edmondson, H. A., 78-4865
 Edwards, C. A., 78-4894
 Edwards, R. G., 78-5192
 Eicken, I., 78-5254
 Eisenberg, D., 78-5041
 Eisenberg, E., 78-5028
 Eisenbrand, G., 78-5058
 Ekert, B., 78-5033
 El'nichnykh, L. N., 78-4971
 Elder, J. H., 78-5210
 Eldor, A., 78-5302
 Elema, J. D., 78-5389
 Eling, T., 78-5015
 Ellett, W. H., 78-5354
 Elling, H., 78-5281
 Ellis, L. D., 78-5397
 Elson, L. A., 78-5046
 Elston, R. C., 78-5364
 Emmelot, P., 78-4861
 Emory, T. H., 78-4985
 Emura, M., 78-5001
 Endo, H., 78-5103
 Endo, T., 78-5103
 Epshtein, S. I., 78-5141
 Erikson, E., 78-5188
 Erikson, R. L., 78-5188
 Erlick, B. J., 78-5204
 Essex, M., 78-4885
 Essigmann, J. M., 78-4962
 Estes, J. D., 78-5209
 Evans, J. H., 78-5342
 Ewig, R. A., 78-5176
 Fabris, G., 78-5263
 Fabry, L., 78-4953
 Fahrig, R., 78-4941
 Falk, L., 78-5239
 Fan, H., 78-4880
 Fanning, E., 78-5246
 Fanta, D., 78-5243
 Faras, A. J., 78-5191
 Farber, E., 78-5084
 Farman, J., 78-5331
 Farrow, G. M., 78-5375
 Favre, M., 78-4889
 Fechner, R. E., 78-4864
 Feild, J., 78-5205
 Feldman, G., 78-5007
 Ferenczy, A., 78-5339
 Ferstandig, L. L., 78-4858
 Fett, D., 78-5052
 Fialkow, P. J., 78-5175, 78-5296
 78-5297
 Fidler, I. J., 78-4923
 Fiers, W., 78-5249
 Figueroa, M. E., 78-5240
 Filipe, M. I., 78-5313
 Filppi, J. A., 78-5086
 Fine, D., 78-4898
 Fiori, M., 78-4893
 Fish, E. N., 78-5242

- Fishbein, L., 78-4828
 Fisher, P. B., 78-5041
 Flamant, R., 78-5367
 Flamm, G., 78-4828
 Flink, E. M., 78-5365
 Floot, B. G., 78-4935
 Fokkens, W., 78-5338
 Forbes, N. L., 78-5158
 Ford, J., 78-5255
 Fortune, D. W., 78-5342
 Foster, J. H., 78-5333
 Fowler, A. K., 78-5272
 Fox, C. F., 78-5210
 Fraumeni, J. F., 78-5344
 Freeman, H. J., 78-5060
 Frese, K., 78-5087
 Freston, J. W., 78-5322
 Frezza, D., 78-4945
 Fried, A. H., 78-5260
 Friedl, H. P., 78-5350
 Friedman, H., 78-5283
 Friedman, P. A., 78-4844
 Friis, R. R., 78-4881
 Fu, P. P., 78-5034
 Fuchs, R. P., 78-4986
 Fuji, H., 78-5279
 Fujinaga, K., 78-5226
 Fujisaki, Y., 78-5107
 Fujita, K., 78-5005
 Fujita, S., 78-5181
 Fukazawa, K., 78-4973
 Fukushima, S., 78-5099
 Furgyk, S., 78-5341
 Furrer, R., 78-5121
 Furukawa, M., 78-5154
 Furukawa, T., 78-5320
 Furuya, K., 78-5063
 Fuscaldo, A. A., 78-5204
 Fuscaldo, K., 78-5204
 Gabbiani, G., 78-5245
 Gabizon, A., 78-5288
 Galbraith, A. I., 78-5083
 Gallagher, R. E., 78-4901
 Gallimore, P. H., 78-4902
 Gallo, R. C., 78-4901
 Ganeshananthan, N., 78-5381
 Gangolli, S. D., 78-5073
 Gans, P., 78-4989, 78-5211
 Gantt, R., 78-5176
 Gardner, M. B., 78-4883, 78-5209
 Gardner, R. H., 78-5184
 Garfinkel, L., 78-4851
 Garten, C. T., 78-5184
 Garzelli, C., 78-5207
 Gaucher, P., 78-5130
 Gavin, J. B., 78-5062
 Gebel, H. M., 78-5289
 Geder, L., 78-5240
 Gehring, P. J., 78-5110
 Geil, R. G., 78-5132
 Jenkins, G., 78-5315
 Gerasimov, A. M., 78-5014
 Ghadirian, P., 78-5049
 Ghysdael, J., 78-5230
 Gilcher, R. O., 78-5397
 Gilden, R. V., 78-5274
 Giles, R. C., 78-5132
 Gilev, A. P., 78-4937
 Gillespie, D. H., 78-4901
 Gillette, R. W., 78-5143
 Ginsberg, H. S., 78-5041
 Glancy, E. M., 78-4964
 Glatt, H., 78-4938
 Glebatis, D., 78-5365
 Gold, M. D., 78-5166
 Goldin, A., 78-5294
 Goldman, P., 78-4844
 Goldstein, G., 78-5295
 Golomb, F., 78-5387
 Goth, R., 78-4827
 Gothoskar, S. V., 78-4955
 Grace, M., 78-5355
 Graf, T., 78-5195
 Graham, S., 78-5372
 Grdina, D. J., 78-5395
 Green, M., 78-5235
 Green, M. H., 78-5250
 Green, U., 78-5101
 Greenberger, J., 78-5211
 Greenwald, D., 78-5076
 Greenwald, E., 78-5387, 78-5387
 Greer, S., 78-5168
 Griffin, J. D., 78-5248
 Grilli, S., 78-4929
 Grist, E., 78-5169
 Grosberg, R. A., 78-4977
 Gross, L., 78-4899
 Gross, P., 78-4805
 Gross, R. L., 78-5019
 Grossberg, A. L., 78-5279
 Grosse, H., 78-5337
 Grossi, M. P., 78-5263
 Grover, P. L., 78-5031
 Grufferman, S., 78-5369
 Grund, E., 78-5152
 Guerci, O., 78-5140
 Guerin, M. R., 78-5043
 Guidotti, G. G., 78-5259
 Guillemain, B., 78-5225
 Guirgis, H. A., 78-5371
 Guralnick, M., 78-5339
 Gurevich, S. M., 78-5014
 Gurkalo, V. K., 78-5067
 Gurtoo, H. L., 78-5324
 Gutmann, H. R., 78-4985
 Haas, J. F., 78-4856
 Haber, D. A., 78-5000
 Habs, M., 78-5065
 Hadjiolov, D., 78-5054
 Haegeman, G., 78-5249
 Haenszel, W., 78-4818
 Haga, M., 78-4942
 Haglid, K. G., 78-5281
 Haije, W. G., 78-5338
 Hainline, S., 78-5369
 Hall, E. J., 78-5144
 Halliburton, I. W., 78-5244
 Hammond, E. C., 78-4851
 Han, T., 78-5286
 Hancock, R. L., 78-5398
 Handler, A. H., 78-5126
 Hanna, N., 78-5173
 Hansson, H., 78-5341
 Hara, M., 78-5122, 78-5124
 Haran-Ghera, N., 78-5224, 78-5295
 Hardy, W. D., 78-4885
 Hariharan, P. V., 78-5009
 Harousseau, J. L., 78-5153
 Harrington, G. W., 78-5053
 Harris, C. C., 78-5020
 Harris, R. E., 78-5371
 Harris, R. H., 78-4847
 Hartley, H. O., 78-4813
 Harvey, R. G., 78-5034
 Hasek, M., 78-5282
 Haseltine, W. A., 78-5271
 Hassall, K. A., 78-5161
 Hassoun, J., 78-5266
 Hata, J. I., 78-5320
 Hauschild, F., 78-5164
 Hayden, D. W., 78-5126
 Hayes, H. M., 78-5357
 Healy, T. M., 78-4916
 Hearn, W. L., 78-5076
 Heberling, R., 78-5274
 Heddle, J. A., 78-4939
 Hehlmann, R., 78-5221
 Hellman, A., 78-5272
 Hellman, K. B., 78-5212
 Hemsall, V., 78-5325
 Henderson, B. E., 78-4865
 Hengeveld, G. M., 78-4981
 Henry, K., 78-5264
 Henschler, D., 78-4830
 Herbst, A. L., 78-5370
 Herlan, A., 78-5032
 Herman, R., 78-5133
 Hernberg, S., 78-4934
 Herring, E., 78-5265
 Herrmann, W. P., 78-4876
 Hertz, R., 78-4863
 Hett, E. A., 78-5166
 Hewer, A., 78-5031
 Hewer, T., 78-5049
 Hiasa, Y., 78-4983
 Hidaka, K., 78-5103
 Hill, M. J., 78-4919, 78-5123
 Hiraki, S., 78-5270
 Hirayama, T., 78-4917
 Hirohata, T., 78-5347
 Hirono, I., 78-4942
 Hirose, K., 78-4973
 Hishinuma, H., 78-5335
 Hitokoto, H., 78-4956
 Hoang-Ngoc Minh, 78-5340
 Hodge, S. J., 78-5305
 Holan, V., 78-5282
 Holden, H. T., 78-5287
 Holder, G., 78-4854
 Hollinshead, A. C., 78-5397
 Holm, G., 78-5300
 Holmes, P. A., 78-5128, 78-5129
 Holtzman, G., 78-5075
 Honda, M., 78-5048
 Hoover, R., 78-5344, 78-5353
 Hop, W. C., 78-5338
 Hopewell, J. W., 78-5135
 Hopkins, N., 78-5214
 Horning, E. C., 78-5166

- Horowitz, S. H., 78-5315
 Houston, H., 78-5123
 Howley, P., 78-4891
 Howorth, M. B., 78-5343
 Hsie, A. W., 78-5158
 Hsieh, D. P., 78-4969
 Hsu, I. C., 78-5020
 Hsu, M. T., 78-5255
 Hu, S. S., 78-5190
 Huang, C. C., 78-5154, 78-5163
 Huang, M. T., 78-4854
 Huberman, E., 78-4999
 Hughes, L. O., 78-5155
 Hulser, D. F., 78-4827
 Hung, C. Y., 78-4947
 Hung, S., 78-5312
 Hunt, D. C., 78-4959
 Hurst, W. J., 78-4965
 Hutchison, G. B., 78-4870
 Iammarino, R. M., 78-5113
 IARC Working Group, 78-4821
 78-4822, 78-4823, 78-4824, 78-4825
 78-4826
 Ide, T., 78-5258
 Infante, P. F., 78-4832, 78-4834
 Ingelfinger, J. A., 78-4844
 Inoue, H., 78-4948
 Iorio, A., 78-5294
 Irving, C. C., 78-4804
 Ishidate, M., 78-5136
 Ishii, Y., 78-5145
 Ishizawa, M., 78-5103
 Iskhakova, E. N., 78-5141
 Israel, M., 78-4891
 Ito, N., 78-4979, 78-5099
 Ito, Y., 78-5051
 Itzhaki, R. F., 78-5083
 Ivankovic, S., 78-5094
 Ivanov, S. D., 78-5072
 Iwaida, M., 78-5051
 Iwasaki, K., 78-5180
 Iyer, S. K., 78-5331
 Izumi, K., 78-5063
 Jacobi, G. H., 78-4920
 Jacquignon, P., 78-4938, 78-5033
 Jafarey, N. A., 78-5382
 Jafari, K., 78-5390
 Jaggi, W., 78-4998
 James, M. C., 78-5296
 James, M. O., 78-4990
 Janerich, D. T., 78-5365
 Janowski, M., 78-5225
 Janson, K. L., 78-5317
 Januszko, T., 78-4966
 Janzowski, C., 78-5058
 Jarrett, W. F., 78-4884
 Jeggo, P., 78-5102
 Jenkins, D. J., 78-5123
 Jenkins, L. J., 78-5109
 Jensen, F. C., 78-5210
 Jerina, D. M., 78-4854, 78-4855
 78-4995, 78-5017, 78-5022
 Jessup, J. M., 78-5173
 Jick, H., 78-5133, 78-5146, 78-5352
 Jinnai, H., 78-5096
 Jivraj, T., 78-5123
 Jockusch, B. M., 78-5195
 Joffe, A. Z., 78-4954
 Johnson, D., 78-5239
 Jones, F. R., 78-5166
 Jones, G. M., 78-5176
 Jones, S., 78-5395
 Jorgenson, T. A., 78-5162
 Joswig-Priewe, H., 78-4866
 Judah, D. J., 78-4960
 Kaaden, O. R., 78-4886
 Kada, T., 78-5122, 78-5124
 Kado, C. I., 78-5400
 Kagan, E., 78-5284
 Kahl, G. F., 78-5010
 Kahl, R., 78-5010
 Kahle, H. R., 78-5310
 Kakimoto, K., 78-5279
 Kakunaga, T., 78-5098, 78-5105
 Kamano, T., 78-5104
 Kamiyama, S., 78-5374
 Kaneko, T., 78-5399
 Kapadia, S. B., 78-5151
 Kapitulnik, J., 78-4854
 Kaplan, A. M., 78-5290
 Kaplan, E., 78-4985
 Kaplan, J. C., 78-5253
 Kaplan, S. S., 78-5151
 Kaplowitz, N., 78-5016
 Kappas, A., 78-4944
 Kark, J. D., 78-5349
 Karle, J. M., 78-4855
 Karmanova, N. V., 78-5293
 Kaspersen, F. M., 78-4982
 Kasumi, F., 78-5368
 Kasza, L., 78-4828
 Kato, S., 78-4980
 Katsuoka, Y., 78-5336
 Kawachi, T., 78-5048
 Kawamoto, S., 78-5181
 Kawasaki, M., 78-5104
 Kay, S., 78-5308
 Keller, W., 78-5254
 Kellermann, G., 78-4928
 Kende, A., 78-4852
 Kendrick, J., 78-5043
 Kenward, B., 78-5391
 Kenyon, C., 78-5166
 Ketkar, M., 78-5101
 Kettmann, R., 78-5230
 Khoury, G., 78-4891
 Khudolei, V. V., 78-5056
 Kidokoro, T., 78-5104
 Kiel, H., 78-4976
 Kim, B., 78-5035
 Kim, Y., 78-5060
 Kim, Y. S., 78-5060
 Kimbrough, R., 78-4828
 Kimula, Y., 78-5327
 King, M. C., 78-5360, 78-5364
 Kinjo, M., 78-5304
 Kinoshita, N., 78-5103
 Kirschman, J. C., 78-5128, 78-5129
 Kishino, H., 78-5104
 Kissonerghis, A. M., 78-4996
 Klanjscek, G., 78-5388
 Klaus, E., 78-5010
 Kleid, D. G., 78-5271
 Kleihues, P., 78-5026, 78-5090
 Klein, G., 78-5278
 Klempnauer, K. H., 78-5246
 Kneale, B. L., 78-5342
 Knill-Jones, R. P., 78-4857
 Knipe, D. M., 78-5244
 Knudson, A. G., 78-4926
 Kocsis, J. J., 78-4975
 Kodama, M., 78-5100, 78-5127
 Kodama, T., 78-5104
 Koestner, A., 78-5086
 Kohli, J., 78-4978
 Kohn, K. W., 78-5176
 Kollinger, G., 78-5087
 Kolonel, L., 78-5347
 Kon, S. H., 78-4836
 Konieczny, M., 78-5034
 Kornfeld, P., 78-5315
 Koroleva, L. A., 78-5014
 Koropatnick, J., 78-5115
 Korsgaard, R., 78-4853
 Kortselius, M. J., 78-4951
 Kos, W. L., 78-5290
 Kotelianskaia, A. E., 78-5141
 Kough, R. H., 78-5303
 Kouri, R. E., 78-4927
 Kra-oz, Z., 78-5216
 Kriegler, M. P., 78-5248
 Kriek, E., 78-4981
 Krikorian, J. G., 78-5384
 Kripke, M. L., 78-5173, 78-5174
 Krishnan, M. N., 78-5331
 Kruger, F. W., 78-5082
 Krulik, M., 78-5153
 Krzyzek, R. A., 78-5191
 Kubitschek, H. E., 78-5171
 Kubota, Y., 78-5107
 Kucerova, M., 78-5157
 Kuchava, N. E., 78-5085
 Kuhlmann, K. F., 78-5114
 Kulkarni, M. G., 78-5346
 Kumar, S., 78-5017
 Kunze, M., 78-5366
 Kurata, H., 78-4956
 Kurihara, M., 78-5104
 Kwapien, R. P., 78-5132
 Kyrtopoulos, S. A., 78-5106
 L'Eplattenier, E., 78-5021
 Labarthe, D., 78-5363
 Laborde, Y., 78-5131
 Ladoulis, C. T., 78-5397
 Laerum, O. D., 78-4827
 Lagace, R., 78-5307
 Lai, C. J., 78-4891
 Lake, B. G., 78-5073
 Lam, L. K., 78-4806
 Lambotte-Vandepaer, M., 78-5111
 Lamden, M. P., 78-5019
 Land, P. C., 78-5121
 Land, P. D., 78-5052
 Lane, D. P., 78-5251
 Langauer, M., 78-5156
 Lanier, A., 78-5380
 Lankin, V. Z., 78-5014
 Lantos, P. L., 78-5319

nza, G., 78-5263
 pis, K., 78-4808
 sh, A. F., 78-5390
 u, A. F., 78-5191
 ub, O., 78-5267
 ugros, A., 78-5130
 wther, P. J., 78-4849
 Bouffant, L., 78-5326
 , M., 78-5367
 catsas, G., 78-5275
 clere, J., 78-5130
 cointe, P., 78-5011
 derlin, P., 78-5140
 dinko, N., 78-5238
 e, E. W., 78-4975
 e, I. P., 78-5012
 e, L. F., 78-5196
 ef, F., 78-5295
 enhouts, H. P., 78-4879
 hr, R., 78-4855
 hr, R. E., 78-5017, 78-5022
 ichter, H., 78-4918
 is, J. P., 78-5191
 menager, J., 78-5326
 mons, R. S., 78-5231
 mos, L. B., 78-5179
 ng, M., 78-4986
 ennox, E., 78-4994
 onard, A., 78-4953
 onov, D., 78-5035
 onov, E., 78-5035
 ePage, G. A., 78-5399
 erner, R. A., 78-5210
 eroux, J. P., 78-4989
 esca, P., 78-5011
 esmana, L., 78-5332
 assin, L. S., 78-5397
 etoublon, C., 78-5131
 evin, D. L., 78-5348
 evin, E., 78-5308
 evin, W., 78-4854, 78-4855, 78-4995
 78-5022
 ew, E. A., 78-4851
 ewis, M. L., 78-5310
 eyba, C., 78-5034
 icea, M., 78-5312
 idak, M. Iu., 78-4937
 ijinsky, W., 78-4817, 78-4819, 78-5080
 78-5097
 imacher, J., 78-5307
 indsay, J. G., 78-5247
 inemeyer, D., 78-5205, 78-5206
 insell, C. A., 78-4839
 ittle, J. B., 78-4875, 78-5182
 ivingston, D. M., 78-5248
 o, L. W., 78-5115
 ewengart, G., 78-5167
 og, T., 78-5228
 omakina, L. Y., 78-4827
 ombardi, B., 78-5113
 onglan, R. C., 78-5073
 ongstaff, E., 78-4943
 opez Quintela, M. A., 78-5077
 oria, R. M., 78-5290
 oub, W. D., 78-4806
 u, A. Y., 78-4854, 78-4855

Lue, C. B., 78-4939
 Lundin, L., 78-5278
 Lusby, A., 78-5009
 Luthy, J., 78-4838
 Lutz, W. K., 78-4998, 78-5008, 78-5021
 Lyman, G. H., 78-5298
 Lynch, H. T., 78-5364, 78-5371
 Lynch, J. F., 78-5371
 Lynch, P. M., 78-5371
 Lynch, R. G., 78-5289
 Lynne, A. B., 78-4977
 MacGregor, G. A., 78-4916
 Machida, T., 78-5335
 Mack, T. M., 78-4910
 Mackay, C., 78-5358
 Mackey, J. K., 78-5235
 Maddox, W. L., 78-5043
 Maeda, S., 78-5095
 Magee, P. N., 78-4820
 Mah, H. D., 78-5017
 Mahy, B. W., 78-5229
 Maisin, J. R., 78-5225
 Maisuradze, S. D., 78-5085
 Major, E. O., 78-4893
 Mak, K. M., 78-5064
 Makary, A. Z., 78-5303
 Malaveille, C., 78-5049
 Malt, R. A., 78-5159
 Maltoni, C., 78-4831, 78-4922
 Mammerickx, M., 78-5230
 Mandard, J. C., 78-5326
 Mandybur, T. I., 78-5088
 Mangino, M. M., 78-5080
 Manning, D., 78-5161
 Mansuy, D., 78-4989, 78-5011
 Mantel, N., 78-4814
 Manthey, B., 78-5021
 Marchok, A. C., 78-5024, 78-5039
 Marcus, S. L., 78-5203
 Marcus, W., 78-4828
 Martin, R. G., 78-4894
 Maruta, K., 78-5005
 Mason, T. J., 78-5344
 Masuda, F., 78-5335
 Mate, U., 78-5108
 Matilla, A., 78-5306
 Matsukura, N., 78-5048
 Matsumoto, H., 78-4973, 78-5048
 Matsushima, M., 78-4973
 Matsushima, T., 78-5048
 Mattar, N., 78-5321
 Mattern, J., 78-5074
 Matthews, H. B., 78-4980
 Matzner, Y., 78-5301
 McAllister, R. M., 78-5274
 McCann, J., 78-4809
 McDonald, K. E., 78-5045, 78-5185
 McDougall, J. K., 78-4902
 McGarrity, G. J., 78-5269
 McGinty, L., 78-4909
 McKinney, J. D., 78-4980
 McLachlan, J. A., 78-5138
 McLelland, R., 78-5369
 McManus, M. J., 78-5392
 McPherson, K., 78-5391
 Melchionne, S., 78-5167

Mellstedt, H., 78-5300
 Mendel, J., 78-5152
 Mendelow, A. D., 78-5308
 Meneguzzi, G., 78-5262
 Menzel, H., 78-5118
 Mercier, M., 78-5111
 Meselson, M., 78-4815
 Metzler, M., 78-5138
 Michas, C., 78-4969
 Mieny, C. J., 78-5308
 Milanesi, G., 78-5262
 Miller, E. C., 78-4816, 78-4984
 Miller, J. A., 78-4816, 78-4984
 Miller, J. M., 78-4887
 Miller, K., 78-5284
 Miller, R. C., 78-5144
 Miller, R. W., 78-4930
 Mills, L. C., 78-5318
 Mink, J. G., 78-5280
 Minowada, J., 78-5286, 78-5324
 Mintzis, M. J., 78-5387
 Mishra, R. S., 78-4961
 Misra, L. K., 78-5292
 Mitsuoka, T., 78-4952
 Mittelman, A., 78-5372
 Miura, K., 78-4973
 Miyata, Y., 78-4979
 Miyoshi, I., 78-5270
 Mizroch, S., 78-4914
 Mochizuki, M., 78-5100
 Modak, M. J., 78-5203
 Modic, S., 78-5388
 Moel, S. A., 78-5309
 Mohr, U., 78-5001
 Molina, A. M., 78-4950, 78-5003
 Mones, E., 78-5125
 Monti-Bragadin, C., 78-4950
 Morales, N. M., 78-4980
 Morantz, R. A., 78-5186
 Morgan, K. Z., 78-4872
 Mori, H., 78-4942
 Morishita, M., 78-5107
 Morita, K., 78-5122, 78-5320
 Moroney, J., 78-5042
 Morozumi, S., 78-4956
 Mosca, L., 78-5030
 Moscovici, C., 78-5190
 Mosher, C. W., 78-5114
 Mosulishvili, L. M., 78-5085
 Moszczynski, P., 78-4974
 Mougeot-Martin, M., 78-5153
 Mughal, S., 78-5313
 Muir, C. S., 78-4915
 Mukhtar, H., 78-5012, 78-5013
 Muller, D., 78-5156
 Muller, U., 78-5254
 Muneer, R. S., 78-5050
 Murasaki, G., 78-4979, 78-5099
 Murata, Y., 78-4983
 Murphy, E. C., 78-5202
 Murray, W. R., 78-5358
 Murugasu, R., 78-5276
 Mussgay, M., 78-4886
 Muzika, K., 78-5070
 Myers, L., 78-5369
 Myhr, B. C., 78-4987

- Nagai, M., 78-5093
 Nagamitsu, S., 78-5180
 Nagao, M., 78-5048
 Nagaratnam, N., 78-5147
 Nagata, C., 78-5100, 78-5127
 Nagatsu, T., 78-5005
 Nagington, J., 78-5229
 Najfeld, V., 78-5296, 78-5297
 Nakahara, K., 78-5328
 Nakamura, H., 78-5107
 Nakanishi, K., 78-4979
 Naparstek, E., 78-5302
 Nash, W. G., 78-5231
 Natusch, D. F., 78-4848
 Nazerian, K., 78-5196
 Neal, G. E., 78-4840, 78-4960
 Nebert, D. W., 78-4927
 Nectoux, J., 78-4915
 Nelson, N. A., 78-5362
 Nelson, R. L., 78-5117
 Neri, M., 78-5294
 Neriishi, S., 78-5181
 Nesheim, S., 78-4958
 Nettesheim, P., 78-5024, 78-5039
 78-5119
 Newberne, P. M., 78-5019
 Newberry, B. H., 78-5029
 Newbold, R. F., 78-4997
 Newell, G. W., 78-5162
 Newell, J., 78-5369
 Ng, A. B., 78-4977
 Nicholson, W. J., 78-4841
 Nicolson, M. O., 78-5274
 Nishizuka, Y., 78-5025
 Nobrega, F. T., 78-5375
 Noel, G., 78-5111
 Noer, H. M., 78-5332
 Nogales, F. F., 78-5306
 Noller, K., 78-5363
 Nomura, A., 78-5347
 Nony, C. R., 78-4972
 Norback, D. H., 78-4829
 Norbury, K. C., 78-5174
 Norimura, T., 78-5181
 Norpoth, K., 78-4811, 78-4811
 Nove, J., 78-5182
 Novi, A. M., 78-4957
 Nunes de Souza, P., 78-5229
 O'Brien, S. J., 78-5231
 O'Brien, T. G., 78-5038
 O'Fallon, W. M., 78-5363
 Oblapenko, N. G., 78-5141
 Ocho, M., 78-5232
 Oda, T., 78-5232, 78-5270
 Oesch, F., 78-4938
 Ogawa, K., 78-5236
 Ogiso, T., 78-5099
 Ogura, H., 78-5232, 78-5270
 Ohashi, T., 78-5261
 Ohki, T., 78-5127
 Ohmori, T., 78-4983
 Ohnesorge, F. K., 78-5118
 Ohshima, M., 78-4983
 Okada, M., 78-5100
 Okajima, S., 78-5181
 Okey, A. B., 78-5139
 Okoye, Z. S., 78-4963
 Old, L. J., 78-5200
 Omura, S., 78-5270
 Ong, T. M., 78-4947, 78-4948, 78-4967
 Oppenheim, A., 78-4894
 Orcel, L., 78-5340
 Ormsby, I., 78-5088
 Orth, G., 78-4889
 Oscarson, J. E., 78-5159
 Ostor, A. G., 78-5342
 Otsuka, H., 78-5063
 Otto, B., 78-5246
 Oudet, P., 78-5257
 Owen, L. G., 78-5305
 Owen, L. N., 78-5229
 Ozzello, L., 78-5323
 Padgett, B. L., 78-5261
 Padgett, T., 78-4882
 Padron, R., 78-5312
 Page, N. P., 78-4803
 Page, T., 78-4847
 Paigen, B., 78-5324
 Paigen, K., 78-5324
 Palaszynski, E., 78-5173
 Panet, A., 78-5216
 Paoletti, C., 78-5011
 Papatestas, A. E., 78-5315
 Papsin, F. R., 78-5242
 Park, D. K., 78-5219
 Parker, N. B., 78-5324
 Parshad, R., 78-5176
 Paskind, M., 78-5197
 Pasquier, D., 78-5131
 Pastan, I., 78-4932
 Pataki, J., 78-5027
 Patiutko, Iu. I., 78-5329
 Paton, D., 78-5150
 Patten, S. E., 78-5392
 Patterson, D. S., 78-4964
 Pearson, T., 78-4994
 Pedersen, F. S., 78-5271
 Peers, F. G., 78-4839
 Pegg, D. G., 78-5110
 Penn, I., 78-5384
 Perdue, M. L., 78-5191
 Perin, F., 78-4938
 Perin-Roussel, O., 78-5033
 Persico-DiLauro, M., 78-4894
 Pessac, B., 78-5194
 Peters, H. A., 78-5338
 Peters, L. J., 78-5395
 Peto, R., 78-4810
 Petrakis, N. L., 78-5360
 Pettigrew, H. M., 78-5177
 Philippus, E. J., 78-4935
 Phillips, J. C., 78-5073
 Piedimonte, G., 78-5259
 Pienta, R. J., 78-4988
 Pietropaolo, C., 78-5040
 Pignatti, P. F., 78-5262
 Pike, M. C., 78-4865, 78-4910
 Pinsky, S. M., 78-5311
 Plesnicar, S., 78-5388
 Poggi, P., 78-5030
 Pooley, J. A., 78-4988
 Poirier, F., 78-5194
 Poland, A., 78-4852
 Poliakov, V. M., 78-5014
 Polivkova, Z., 78-5157
 Polliack, A., 78-5301, 78-5302
 Pope, J. H., 78-4896
 Poplawska, T., 78-4966
 Poppema, S., 78-5389
 Popper, H., 78-4922
 Portengen, H., 78-5338
 Pot-Deprun, J., 78-5149
 Potter, M., 78-5198, 78-5277
 Pour, P. M., 78-5081
 Preisler, H. D., 78-5298
 Pressman, D., 78-5279
 Preussmann, R., 78-5058
 Pritchard, A. B., 78-5128, 78-5129
 Prodi, G., 78-4924, 78-4929
 Pueyo, C., 78-4946
 Radomski, J. L., 78-5076
 Raineri, R., 78-4988
 Rajewsky, M. F., 78-4827
 Rajiyah, S., 78-5147
 Rakusanova, T., 78-5253
 Rall, D. P., 78-4807
 Ranadive, K. J., 78-4955
 Ranadive, S. N., 78-4955
 Ransohoff, J., 78-5092
 Rapp, F., 78-5240
 Rapp, U. R., 78-5218
 Ratovitski, E. A., 78-5069
 Rauber, G., 78-5130
 Rawls, W. E., 78-4888
 Reddy, A. L., 78-5175, 78-5297
 Reiches, N. A., 78-4847
 Reinhold, V. N., 78-4962
 Reiss-Gutfreund, R. J., 78-5243
 Reitz, M. S., 78-4901
 Remsen, J., 78-5007
 Renault, G., 78-5149
 Resseguie, L. J., 78-5375
 Reynolds, F. H., 78-5199, 78-5201
 Reznik, G., 78-5079, 78-5101
 Reznik-Schuller, H., 78-5068
 Rheins, M. S., 78-5086
 Rhim, J. S., 78-5219
 Rhoads, J. E., 78-5061
 Rhoton, J. C., 78-5024
 Rice, N., 78-5274
 Richardson, A. C., 78-5354
 Richter, A., 78-5246
 Richter-Reichhelm, H. B., 78-5001
 Riddell, R. H., 78-5322
 Riede, U. N., 78-5090
 Rieger, M. M., 78-4859
 Ring, J., 78-5222
 Ringold, G. M., 78-5222
 Risser, R., 78-5198, 78-5200
 Rivett, K. F., 78-5165
 Robbins, A. K., 78-5251
 Robbins, J. H., 78-4936
 Robboy, S. J., 78-5370
 Roberfroid, M., 78-4953, 78-5111
 Roberts, B. A., 78-4964
 Roberts, J. A., 78-5317
 Rodhain, F., 78-4911
 Roethlisberger, B., 78-5323


- Rogiers, R., 78-5249
 Roh, S. K., 78-5311
 Roizman, B., 78-5244
 Romruen, K., 78-4970
 Roscoe, J. P., 78-5091
 Rose, E., 78-5049
 Rose, L. I., 78-5318
 Rose, M., 78-5309
 Rosenberg, H., 78-5396
 Rosenberg, M., 78-5396
 Rosenberg, N., 78-5197
 Ross, J. S., 78-5159
 Rothblat, J., 78-4871
 Rothenberg, E., 78-5214, 78-5215
 Rothenstein, A. S., 78-5125
 Rouer, E., 78-4989
 Rowe, W. P., 78-5198
 Rowland, J., 78-5004, 78-5006
 Rowles, P., 78-5091
 Roze, J. M., 78-4911
 Rubio, N., 78-5295
 Ruddell, W. S., 78-5330
 Rudiger, H. W., 78-4931
 Rudikoff, S., 78-5277
 Ruebner, B. H., 78-4969
 Rundell, K., 78-4892
 Runge, R. G., 78-5081
 Rungger, D., 78-5193
 Rungger, E., 78-5245
 Ruscetti, F., 78-4901
 Ruscetti, S., 78-5205, 78-5206
 Rushbrook, C. J., 78-5162
 Russell, K., 78-4815
 Russell, P. H., 78-5227
 Russell, W. J., 78-5181
 Rustia, M., 78-5142
 Ruyechan, W. T., 78-5244
 Ryan, D., 78-4854, 78-4855, 78-4995
 Ryo, U. Y., 78-5311
 Ryssel, H. J., 78-5299
 Sa-Ngobwarchar, P., 78-5359
 Sabin, A. B., 78-4903
 Sachs, L., 78-4933
 Sacks, T. L., 78-5199, 78-5201
 Sadaie, Y., 78-5124
 Safe, S., 78-4978
 Saffiotti, U., 78-4802
 Sage, E., 78-4986
 Sahaphong, S., 78-4970
 Saito, H., 78-5127
 Sakai, S., 78-4956
 Sakamoto, G., 78-5368
 Saldinger, E., 78-4869
 Salmasi, S. Z., 78-5081
 Salo, A., 78-5175
 Samson, L., 78-5102
 Sanford, K. K., 78-5176
 Santa Cruz, D. J., 78-5179
 Sarasa Bharati, R., 78-4992
 Sarma, D. S., 78-5084
 Sarma, P. S., 78-5228
 Sasaki, T., 78-5335
 Sato, S., 78-5048
 Saunders, E. F., 78-4939
 Savary, M., 78-5323
 Sawada, M., 78-5136
 Sawamura, M., 78-5048
 Saxinger, W. C., 78-4901
 Scelsi, M., 78-5030
 Scelsi, R., 78-5030
 Schaefer-Ridder, M., 78-4855
 Scheen, S. R., 78-5305
 Schendel, P. F., 78-5102
 Scherer, E., 78-4861, 78-4935
 Schlatter, C., 78-4998, 78-5021
 Schlegel, R., 78-5268
 Schmahl, D., 78-5065
 Schmidt, F. E., 78-5310
 Schochetman, G., 78-4898
 Schoental, R., 78-4954
 Schottenfeld, D., 78-4856
 Schuman, R., 78-4988
 Schwab, M., 78-5087
 Schwind, H., 78-4938
 Scolnick, E., 78-5205
 Scolnick, E. M., 78-5206, 78-5208
 Scott, E. L., 78-4877
 Scully, R. E., 78-5370
 Segal, A., 78-5108
 Segaloff, A., 78-5177
 Seidman, H., 78-4851
 Seidman, I., 78-5167
 Seino, Y., 78-5048
 Sekiya, A., 78-5005
 Selikoff, I. J., 78-4842
 Selkirk, J. K., 78-5020
 Sell, S., 78-5113
 Selsky, C. A., 78-5168
 Sessions, F., 78-5009
 Setlow, R. B., 78-5169
 Shain, W., 78-5186
 Shalev, C., 78-5362
 Shaposhnikov, J. D., 78-5069
 Sharashidze, L. K., 78-5085
 Sharma, J. M., 78-5196
 Sharp, P. A., 78-5214
 Shenk, T., 78-5252
 Sherr, C. J., 78-5231
 Shibko, S., 78-4828
 Shields, A., 78-5197, 78-5215
 Shimada, A., 78-5374
 Shingleton, W., 78-5369
 Shinohara, K., 78-5007
 Shinozuka, H., 78-5113
 Shinpo, K., 78-5005
 Shiomi, T., 78-5181
 Shirai, A., 78-5048
 Shirai, T., 78-4979, 78-5099
 Shivapurkar, N. M., 78-4955
 Shoji, R., 78-5335
 Shooter, K. V., 78-5071
 Shubik, P., 78-5142
 Sieber, R., 78-4837
 Sieber, S. M., 78-5148
 Siedband, L. B., 78-5311
 Sielken, R. L., 78-4813
 Sigel, M. M., 78-4977
 Sikora, K., 78-4994
 Silverman, D. T., 78-5348
 Simkhovich, B. Z., 78-4937
 Simon, K., 78-4808
 Sims, P., 78-5031
 Singer, E. J., 78-5125
 Singer, G. M., 78-5080
 Singer, J., 78-5297
 Singer, J. V., 78-5296
 Singer, M., 78-5396
 Singh, R. S., 78-4961
 Sipes, I. G., 78-5075
 Sirbasku, D. A., 78-5134
 Sirianni, S. R., 78-5163
 Sivarajah, K., 78-5015
 Skinner, D. B., 78-5322
 Slater, G., 78-5315
 Slocumb, M. L., 78-5075
 Smadja, A., 78-5340
 Smales, W. P., 78-5253
 Smith, A., 78-5042
 Smith, A. C., 78-5167
 Smith, B. R., 78-5018
 Smith, E. S., 78-5355
 Smith, L. G., 78-5045
 Smith, P. G., 78-5146, 78-5352
 Smith, S. W., 78-5203
 Smith, S. Z., 78-5305
 Smylie, M., 78-4978
 Snyder, R., 78-4975
 Sobel, G. W., 78-5311
 Soetoto, H. J., 78-5345
 Solomon, D., 78-4891
 Sonntag, R. W., 78-5299
 Sonobe, H., 78-5236
 Sonoki, S., 78-5400
 Sontag, J. M., 78-4801
 Sontipong, S., 78-5359
 Soost, H. J., 78-4866
 Spadafora, C., 78-5257
 Spector, D. H., 78-4882
 Speier, J. L., 78-4806
 Spellman, C. W., 78-5285
 Spence, A. A., 78-4857
 Spence, A. M., 78-5089
 Spiegelman, S., 78-5221
 Sporn, M. B., 78-4860
 Spreafico, F., 78-5294
 Squire, R. A., 78-4922
 Stallcup, M. R., 78-5222
 Stanley, N. F., 78-4904
 Starek, A., 78-4974
 Stavrou, D., 78-5281
 Steel, V. E., 78-5039
 Steffens, L., 78-4976
 Stehelin, D., 78-4882
 Steinmann, L., 78-5297
 Stenback, F., 78-5004, 78-5006, 78-5036
 Stephenson, J. R., 78-4885, 78-5199
 78-5201
 Stevens, D. L., 78-5045
 Stich, H. F., 78-5115
 Stillwell, R. N., 78-5166
 Stinson, E. B., 78-5384
 Stocker, M., 78-5393
 Stockert, E., 78-5200
 Stokely, J. R., 78-5043
 Stoming, T. A., 78-4995
 Stoner, G. D., 78-5020
 Stora, C., 78-4968
 Straf, M. L., 78-4877

- Strel'tsova, V. N., 78-4874
 Stubblefield, E., 78-4882
 Stutman, O., 78-4907
 Styles, J. A., 78-4940, 78-5150
 Sugano, H., 78-5368
 Sugimura, T., 78-5048
 Sugiyama, T., 78-5095
 Sukaton, U., 78-5332
 Sukumaran, K. B., 78-5034
 Sulaiman, A., 78-5332
 Sulochana, G., 78-5044
 Sunter, J. P., 78-5059
 Sutter, D., 78-5237
 Suzuki, H., 78-5051
 Suzuki, K., 78-4952
 Svet-Moldavsky, G. J., 78-5293
 Swann, P. F., 78-5106
 Swanson, M., 78-5372
 Swern, D., 78-5053
 Szende, B., 78-4808
 Szepsenwol, J., 78-5120
 Tagashira, Y., 78-5127
 Takahashi, G., 78-4993
 Takahashi, H., 78-5005
 Takahashi, K., 78-5103
 Takahashi, M., 78-5099
 Takemoto, K. K., 78-4890
 Takemoto, L. J., 78-5210
 Takeshita, K., 78-5181
 Tamaoki, N., 78-5320
 Tamaro, M., 78-4950
 Tamm, I., 78-5233
 Tanaka, T., 78-5232
 Tangthatswasd, K., 78-5359
 Taniyama, T., 78-5287
 Tannenbaum, S. R., 78-4818, 78-5052
 Taper, H. S., 78-5055
 Taqawi, I. H., 78-4878
 Tarone, R. E., 78-5176
 Taylor, C. W., 78-5017
 Taylor, H. W., 78-4819
 Teague, C. A., 78-5062
 Tegtmeier, P., 78-4892
 Teichmann, B., 78-5164
 Teramoto, K., 78-5335
 Teske, R., 78-4828
 Teulings, F. A., 78-5338
 Thakker, D. R., 78-4854, 78-4855
 78-4995, 78-5022
 Thamavit, W., 78-4970
 Thenot, J. P., 78-5166
 Thibault, L. H., 78-5097
 Thibaut, G., 78-5140
 Thiele, J., 78-4976
 Thilly, W. G., 78-5000
 Thomas, L. B., 78-4922
 Thomas, P. E., 78-4995
 Thust, R., 78-5152
 Tierney, B., 78-5031
 Tilley, B., 78-5363
 Timmons, J. W., 78-5375
 Tirtosugondo, 78-5345
 Tobin, S. M., 78-5242
 Todaro, G. J., 78-4900, 78-5218
 78-5220
 Toga, M., 78-5266
 Tomei, L. D., 78-5042
 Toniolo, A., 78-5207
 Toomey, P. B., 78-4965
 Topham, J. C., 78-5155
 Toyoda, M., 78-5051
 Trainin, N., 78-5288
 Tramacere, M., 78-5259
 Trell, E., 78-4853
 Tripier, M. F., 78-5266
 Troxler, D., 78-5205
 Troxler, D. H., 78-5206, 78-5208
 Trucksess, M. W., 78-4958
 Truex, C. R., 78-5002
 Trujillo, J. M., 78-5394
 Trump, B. F., 78-5020
 Tsanev, R., 78-5054
 Tsuda, H., 78-5084
 Tsuneyoshi, M., 78-5180
 Tuey, D. B., 78-4980
 Twardzik, D. R., 78-5272
 Ueda, N., 78-5095
 Uenaka, H., 78-5095
 Ueyama, Y., 78-5320
 Upton, A. C., 78-4873
 Utsunomiya, T., 78-5103
 Van de Voorde, A., 78-5249
 van der Werf-Messing, B., 78-5338
 Van Duuren, B. L., 78-5167
 Van Herreweghe, J., 78-5249
 Van Heuverswyn, H., 78-5249
 van Muiswinkel, W. B., 78-5280
 van Nie, R., 78-5223
 Van Wyk, J. A., 78-5275
 Van Der Maaten, M. J., 78-4887
 Varakis, J. N., 78-5261
 Varela, M., 78-5317
 Varesio, L., 78-5287
 Varghese, A. J., 78-5121
 Varmus, H. E., 78-4882
 Verma, I. M., 78-5217
 Vernon, M. L., 78-5219
 Verstraeten, A. A., 78-5223
 Vessey, M. P., 78-5391
 Vianna, N. J., 78-4913
 Vincent, R., 78-5324
 Visek, W. J., 78-5002
 Viviani, A., 78-5008
 Vlahakis, G., 78-5334
 Vogel, E., 78-4812, 78-4951
 Vogt, P. K., 78-5187, 78-5190
 Vohra, S. K., 78-5053
 Volckaert, G., 78-5249
 Volm, M., 78-5074
 von der Helm, K., 78-5193
 Voronova, L. A., 78-5072
 Vrana, M., 78-5277
 Vuletin, J. C., 78-5331
 Vutuc, C., 78-5366
 Wade, G. G., 78-5126
 Wagner, G., 78-4957
 Wagoner, J. K., 78-4832, 78-4834
 Walker, D. L., 78-5261
 Waller, R. E., 78-4849
 Walsh, C., 78-5031
 Walters, C. L., 78-5330
 Ward, E., 78-5324
 Ward, J. M., 78-5334
 Warzok, R., 78-5152
 Waskell, L., 78-5160
 Watanabe, P. G., 78-5110
 Waters, M., 78-5162
 Watson, A. J., 78-5059
 Wattenberg, L. W., 78-4806
 Wauke, T., 78-4956
 Wayss, K., 78-5074
 Weatherby, R. P., 78-4990
 Weaver, J. M., 78-5125
 Webster, A., 78-5390
 Webster, M., 78-5385
 Weichselbaum, R. R., 78-5182
 Weicksel, P., 78-5377
 Weil, R., 78-5245
 Weinstein, I. B., 78-5040, 78-5041
 Weisburger, J. H., 78-4912
 Weislow, O. S., 78-5272
 Weiss, N. S., 78-4867
 Welch, W. R., 78-5370
 Welsch, C. W., 78-5392
 Wendel, I., 78-5254
 Wenner, C. E., 78-5042
 West, S., 78-4854, 78-4855
 West, W., 78-5204
 Westphal, M., 78-5237
 Westra, J. G., 78-4982
 Wheelock, E. F., 78-5291
 Whelly, S., 78-5258
 White, C. S., 78-5397
 Whitehouse, S., 78-5062
 Widagdo, L., 78-5345
 Wiessler, M., 78-5065, 78-5066
 78-5082
 Wiggins, H. S., 78-5123
 Wigle, D. T., 78-5355
 Wigler, M., 78-5040
 Wild, P. J., 78-4959
 Wilkinson, G., 78-5372
 Wille, W., 78-5193
 Willecke, K., 78-5193
 Williams, D. C., 78-5137
 Williams, D. T., 78-5070
 Williams, J. I., 78-5172
 Williamson, R. C., 78-5159
 Willingham, M., 78-4932
 Wilson, J., 78-5047
 Wilson, K., 78-5378
 Wilson, W. D., 78-5242
 Wimmer, E., 78-5213
 Winslow, D. P., 78-5091
 Wishnok, J. S., 78-4818
 Witte, O. N., 78-5197, 78-5215
 Witter, R. L., 78-5196
 Witz, F., 78-5140
 Wogan, G. N., 78-4962
 Wolf, H., 78-5241
 Wolstencroft, L., 78-5247
 Won, O. H., 78-5331
 Wong, J. J., 78-4969
 Wong, R., 78-5360
 Wong-Staal, F., 78-4901
 Wong, Z. A., 78-4969
 Wood, A. W., 78-4854, 78-4855
 78-4995, 78-5022

Woods, F. M., 78-5076
Worobec, T. G., 78-5375
Wortman, M., 78-5108
Wreghitt, T. G., 78-5229
Wright, N. A., 78-5059
Wright, N. H., 78-5391
Wronkowski, Z., 78-5178
Wunderlich, D. A., 78-5143
Wurgler, F., 78-5021
Wynder, E. L., 78-4912
Wyndham, C., 78-4978
Wyplatz, J., 78-5326
Yagen, B., 78-4954
Yagi, H., 78-4854, 78-4855
Yagishita, T., 78-4973
Yahagi, T., 78-5048
Yamamoto, K. R., 78-5222
Yamaura, I., 78-5189
Yamazaki, H., 78-5226
Yanagisawa, M., 78-5335

Yang, N. C., 78-4984, 78-5035
Yaniv, M., 78-5256
Yarita, T., 78-5119
Yefenof, E., 78-5278
Yokota, M., 78-5234
Yoshida, J., 78-5092
Yoshinaga, H., 78-5181
Young, J. C., 78-5057
Young, J. L., 78-5357
Young, R. J., 78-4834
Young, V. R., 78-5052
Ysebaert, M., 78-5249
Yu, K. K., 78-5265
Yuan, E., 78-5206
Zabzhinskii, M. A., 78-5067
Zacharias, C., 78-4893
Zaidi, S. H., 78-5382
Zajdela, F., 78-4938, 78-5033
Zavarzin, A. A., 78-5379
Zawirska, B., 78-5116

Zdrojewski, J. F., 78-5305
Zeiger, E., 78-4945
Zeller, J., 78-5094
Zeller, W. J., 78-5094
Zempel, J. A., 78-5110
Zentgraf, H., 78-5254
Zheleznov, B. I., 78-4921
Zieqler, D. M., 78-5112
Zimmermann, A., 78-5299
Zimmermann, D., 78-4850
Zinzar, S. N., 78-5293
Zlotnik, A. , 78-5302
Zolotareva, G. N., 78-5141
Zuch, A., 78-4966
ZuRhein, G. M., 78-5261
Zwilling, B. S., 78-5086



Subject Index

malities

anesthetics
Occupational Hazard, 78-4858
aryl Hydrocarbon Hydroxylases
Genetics, Review, 78-4927
nide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
ames Test
Dose-Response Study, 78-4984
azaguanine Resistance
Hamster V79 Cells, 78-4987
DNA
Antibody Formation, 78-4986
DNA Repair
Endonucleases, 78-5169
fibroblasts
DNA Repair, 78-5169
methanesulfonic Acid, Methyl Ester
Lethal/Mutagenic Activity, 78-4987
ultraviolet Rays
DNA Repair, 78-5169
Xeroderma Pigmentosum, 78-5169
virus, SV40
DNA, Binding, 78-4984
Xeroderma Pigmentosum
DNA Repair, 78-5169
Virus, Herpes Simplex 1, 78-5168
nide, *N*-Fluoren-2-yl-
ceto hydroxamic Acid, *N*-Fluoren-2-yl-
Carcinogenic Metabolite, 78-4988
ames Test
Microsomes, Kidney, 78-5078
aryl Hydrocarbon Hydroxylases
Carcinogenic Metabolite, 78-4988
cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -isomer
Hepatocarcinogenicity, Rat, 78-4983
lutamyl Transpeptidase
Liver, Rat, 78-5084
epatoma
Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -isomer
78-4983
anechlor 500
Hepatocarcinogenicity, Review, 78-4828
iver Neoplasms
Adenocarcinoma, 78-4983
etabolic Activation
Carcinogenic Metabolite, Review, 78-4816
icrosomes, Liver
Hydroxylation, Carbon Atom 9, 78-4985
xygenases
Metabolism, Review, 78-4854
regn-4-ene-3,20-dione, 17-(Acetyloxy)-6 α -methyl-
Metabolism, Kidney, 78-5078
ex Hormones
Metabolism, Kidney, 78-5078
estosterone
Metabolism, Kidney, 78-5078
ritium
DNA, Binding, 78-4982
rea, Methyl Nitroso-
Hepatocarcinogenicity, 78-5084
nide, *N*-Fluoren-3-yl-
arbituric Acid, 5-Ethyl-5-phenyl-

Acetamide, *N*-Fluoren-3-yl- (cont'd)
Metabolism, Rat, Guinea Pig, 78-4985
Cholanthrene, 3-Methyl-
Metabolism, Rat, Guinea Pig, 78-4985
Microsomes, Liver
Hydroxylation, Carbon Atom 9, 78-4985
Metabolism, Rat, Guinea Pig, 78-4985
Acetamide, *N*-Methyl-
Ethylamine, *N*-Methyl-*N*-nitroso-
Nitrous Acid, Sodium Salt, 78-5080
Acetamide, Thio-
DNA Repair
Liver, 78-4949
Acetanilide, 4'-(*p*-Fluorophenyl)-
Acetohydroxamic Acid, *N*-4-(4'-Fluorobiphenyl)-
DNA, Binding, 78-4981
Acetic Acid
Benzo(b)triphenylene
Benzoic Acid, 2-(9'-Phenanthroyl)-, 78-5034
Carboxy-Lyases
Enzymatic Activity, 78-5037
Acetic Acid, Butylnitrosaminomethyl Ester
Stomach Neoplasms
Carcinogenic Activity, Rat, 78-5065
Acetic Acid, (2,3-Dichloro-4-(2-methylenebutyryl)phenoxy)
Barbituric Acid, 5-Ethyl-5-phenyl-
Transferases, 78-5016
Benzo(a)pyrene
Transferases, 78-5016
Cholanthrene, 3-Methyl-
Transferases, 78-5016
Acetic Acid, (Ethylenedinitrilo)tetra-
Agrobacterium tumefaciens
Proteins, 78-5400
Acetic Acid, Ethylnitrosaminoethyl Ester
Ethylamine Hydrochloride
Synthesis, 78-5066
Acetic Acid, Ethylnitrosaminomethyl Ester
Stomach Neoplasms
Carcinogenic Activity, Rat, 78-5065
Acetic Acid, Lead Salt
Cell Transformation, Neoplastic
Cells, Cultured, 78-5117
Fibrosarcoma
Cells, Cultured, 78-5117
Kidney Neoplasms
Porphyrins, 78-5116
Porphyrins
Metabolism, Rat, 78-5116
Virus, Adeno 7
Cell Transformation, Neoplastic, 78-5117
Acetic Acid, Methylnitrosaminomethyl Ester
Methylamine Hydrochloride
Synthesis, 78-5066
Stomach Neoplasms
Carcinogenic Activity, Rat, 78-5065
Acetic Acid, Propylnitrosaminomethyl Ester
Stomach Neoplasms

Acetic Acid, Propylnitrosaminomethyl Ester (cont'd)
 Carcinogenic Activity, Rat, 78-5065

Acetic Acid, Trideuteromethylnitrosaminomethyl Ester
 Stomach Neoplasms
 Carcinogenic Activity, Rat, 78-5065

Acetohydroxamic Acid, *N*-Fluoren-2-yl-
 Acetamide, *N*-Fluoren-2-yl-
 Carcinogenic Metabolite, 78-4988
 Hepatocytes
 Carcinogenic Metabolite, 78-4988

Acetohydroxamic Acid, *N*-Fluoren-3-yl-
 Microsomes, Liver
 Hydroxylation, Carbon Atom 9, 78-4985
 Metabolism, Rat, Guinea Pig, 78-4985

Acetohydroxamic Acid, *N*-4-(4'-Fluorobiphenyl)-
 Acetanilide, 4'-(*p*-Fluorophenyl)-
 DNA, Binding, 78-4981
 4-Biphenylamine, 4'-Fluoro-
 DNA, Binding, 78-4981
 Carcinogenic Metabolite
 Liver, Kidney, Rat, 78-4981

Acetone
 Dimethylamine, *N*-Nitroso-
 Oxidoreductases, *N*-Demethylating, 78-5075

Acetonitrile, 2-(*N*-Methyl-*N*-nitroso)amino-
 Hydrochloric Acid
 Reaction Product, 78-5053
 1,2,3-Oxadiazolium, 3-Methyl-5-amino-, Chloride
 Reversible Interconversion, 78-5053

***p*-Acetophenetidine**
 Carcinogenic Potential
 Review, 78-4843

Acetoxymethyl Methylnitrosamine
 see Acetic Acid, Methylnitrosaminomethyl Ester

Acid Phosphatase
 Benzene
 Enzymatic Activity, 78-4974
 Inhalation Study, Rat, 78-4974
 Lymphocytes, 78-4974
 Letterer-Siwe Disease
 Enzymatic Activity, 78-5389

Acridine, 9-Amino-
 Adenosine Monophosphate
 Fluorescence Quenching, 78-5107
 Guanosine, 7-Methyl-
 Fluorescence Quenching, 78-5107
 Guanosine Monophosphate
 Fluorescence Quenching, 78-5107

Acromegaly
 Meningioma
 Case Report, 78-5318

Acrylonitrile
 Aldrin
 Enzymatic Activity, 78-5111
 Cytochrome Reductases
 Liver, Rat, 78-5111
 Glucosephosphatase
 Liver, Rat, 78-5111
 Glutamyl Transpeptidase
 Enzymatic Activity, 78-5111
 Lactate Dehydrogenase

Acrylonitrile (cont'd)

Aroclor 1254, 78-5111
 Barbituric Acid, 5-Ethyl-5-phenyl, 78-5111
 Cholanthrene, 3-Methyl-, 78-5111
 Cysteine, 78-5111
 Liver, Rat
 Enzymatic Activity, 78-5111
 Maleic Acid, Diethyl Ester
 Lactate Dehydrogenase, 78-5111
 Oxidoreductases, 78-5111

Actin

Cell Transformation, Neoplastic
 Cell Cycle Kinetics, 78-5245
 Hepatoma
 Neoplasm Metastasis, 78-5054
 Virus, Polyoma
 Cell Transformation, Neoplastic, 78-5245
 Virus, Rous Sarcoma
 Temperature Sensitive Mutants, 78-5195
 Virus, SV40
 Cell Transformation, Neoplastic, 78-5245

Actinomycin D

Neurospora crassa
 Lethal/Mutagenic Activity, 78-4947

Adenocarcinoma

Bile Duct Neoplasms
 Dyes, 78-4973
 Cervix Neoplasms
 Epidemiology, Thailand, 78-5356, 78-5361
 Colonic Neoplasms
 Genetics, 78-5386
 Esophageal Neoplasms
 Case Report, Barrett's Syndrome, 78-5322
 Gastrointestinal Neoplasms
 Cholesterol, 78-5120
 Gynecologic Neoplasms
 4,4'-Stilbenediol, α,α' -Diethyl-, 78-5370
 Hereditary Diseases
 Review, 78-4929
 Intestinal Neoplasms
 Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063
 Morpholine, *N*-Nitroso-, 78-5056
 Kidney Neoplasms
 Neoplasm Metastasis, 78-5335
 1-Propanol, 2,3-Dibromo-, Phosphate, 78-5167
 Liver Neoplasms
 Acetamide, *N*-Fluoren-2-yl-, 78-4983
 Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -isomer
 78-4983
 Lung Neoplasms
 Asbestos, 78-5326
 Dyes, 78-4973
 Histochemical Study, 78-5327
 Iron Oxide, 78-5326
 Kidney Transplantation, 78-5353
 Quartz, 78-5326
 Siderosis, 78-5326
 Silver, 78-5326
 Transplantation Immunology, 78-5293
 Ultrastructural Study, 78-5327
 Mammary Neoplasms, Experimental
 Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
 Dexamethasone, 78-5143
 Transplantation Immunology, 78-5293
 Nose Neoplasms
 Ethanol, *N*-Nitrosoiminodi-, 78-4823

ocarcinoma (cont'd)**Ovarian Neoplasms**

Epidemiology, Dog, 78-5357

Pancreatic Neoplasms

Animal Model, Rat, 78-4991

Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081

Epidemiology, Sri Lanka, 78-5376

Peutz-Jeghers Syndrome

Neoplasm Metastasis, 78-5311

Respiratory Tract Neoplasms

Benzo(a)pyrene, 78-5004, 78-5006

4,4'-Stilbenediol, α,α' -Diethyl-

Carcinogenic Potential, Review, 78-4863

Thyroid Neoplasms

2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5079

Uterine Neoplasms

Epidemiology, Review, 78-4867

Radiation Effects, 78-5305

Turner's Syndrome, 78-5342

Virus, Polyoma, BK

Carcinogenic Activity, Hamster, 78-5263

oma**Colonic Neoplasms**

Polyps, 78-5313

Intestinal Neoplasms

Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063

Kidney Neoplasms

Ethanol, *N*-Nitrosoiminodi-, 78-4823

Liver Neoplasms

Contraceptives, Oral, 78-5130

Epidemiology, Review, 78-4865

Morpholine, *N*-Nitroso-, 78-5056

Lung Neoplasms

Benz(a)anthracene, 7,12-Dimethyl-, 78-5030

Imidazole-1-ethanol, 2-Methyl-5-nitro-, 78-4844

Mammary Neoplasms, Experimental

Contraceptives, Oral, 78-5132

Pancreatic Neoplasms

Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081

Respiratory Tract Neoplasms

Benzo(a)pyrene, 78-5004, 78-5006

Thyroid Neoplasms

Epidemiology, Iran, 78-5383

2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5079

oma, Liver Cell

see Hepatoma

omatosis, Familial Endocrine**Chromosome Abnormalities**

Chromosomes, Human, 13-15, 78-5317

Pheochromocytoma

Case Report, 78-5317

Chromosomes, Human, 13-15, 78-5317

Histocompatibility Antigens, 78-5317

osine Cyclic 3',5' Monophosphate**Adenyl Cyclase**

Enzymatic Activity, Review, 78-4932

Carcinogen, Environmental

Concentration Levels, Review, 78-4937

Cell Transformation, Neoplastic

Cell Adhesion, Review, 78-4932

Neoplasms, Experimental

Concentration Levels, Review, 78-4937

osine Monophosphate**Acridine, 9-Amino-**

Fluorescence Quenching, 78-5107

Adenyl Cyclase

Adenosine Cyclic 3',5' Monophosphate

Enzymatic Activity, Review, 78-4932

Adrenal Gland Neoplasms**Carcinoma**

Cushing's Syndrome, 78-5312

Cushing's Syndrome

Case Report, 78-5312

Genetics, 78-5312

Hyperplasia

Cushing's Syndrome, 78-5312

Neoplasms, Multiple Primary

Dyes, 78-4973

Occupational Hazard, 78-4973

Pheochromocytoma

Genetics, 78-5308

Adrenergic Alpha Receptor Blockaders**Esophageal Neoplasms**

Dimethylamine, *N*-Nitroso-, 78-5067

Liver Neoplasms

Dimethylamine, *N*-Nitroso-, 78-5067

Adriamycin**Lung Neoplasms**

Carcinoma, Epidermoid, 78-5399

Aflatoxin B1**Aflatoxin B1 2,3-Dichloride**

Carcinogenic Metabolite, Review, 78-4816

Ames Test

Carcinogenic Potential, Review, 78-4815

Mutagenic Activity, Review, 78-4811

Aspergillus flavus

Drug Contamination, 78-4956

Cholangioma

Ultrastructural Study, 78-4968

DNA

Binding, Liver, 78-4962

Carcinogenic Metabolite, 78-4962

Enzyme Activation

Species Difference, 78-4969

Food Contamination

Carcinogenic Metabolite, 78-4964

Quantitation Method, 78-4958, 78-4959, 78-4965

Quantitation Method, Review, 78-4838

Rain, 78-4961

Hepatoma

Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
78-4970

Ultrastructural Study, 78-4968

Hypophysectomy

Enzyme Activation, 78-4960

Hepatotoxicity, Rat, 78-4960

Nucleic Acids, 78-4960

RNA Polymerase, 78-4960

Liver Neoplasms

Epidemiology, Zaire, 78-4968

Metabolic Activation

Carcinogenic Metabolite, Review, 78-4816

Microsomes, Liver

Ames Test, 78-4969

Mutagenic Activity, 78-4969

Mutagenic Activity

Dose-Response Study, 78-4966

Neurospora crassa

Lethal/Mutagenic Activity, 78-4947

Mutagenic Activity, 78-4967

Aflatoxin B1 (cont'd)

- Nucleic Acids
 - Carcinogenic Metabolite, Review, 78-4816
 - Liver, Rat, 78-4960
- Retinol Palmitate
 - Hydroxylation and Demethylation, 78-4963
 - Metabolism, Rabbit, 78-4963
- RNA Replication
 - Liver, Rat, 78-4957
- RNA, Ribosomal
 - RNA Replication, 78-4957
- Species Difference
 - Hepatocarcinogenicity, 78-4969

Aflatoxin B2

- Food Contamination
 - Quantitation Method, 78-4959, 78-4965

Aflatoxin B1 2,3-Dichloride

- Aflatoxin B1
 - Carcinogenic Metabolite, Review, 78-4816

Aflatoxin G1

- Cholangioma
 - Ultrastructural Study, 78-4968
- Food Contamination
 - Quantitation Method, 78-4959, 78-4965
 - Quantitation Method, Review, 78-4838
- Hepatoma
 - Ultrastructural Study, 78-4968
- Liver Neoplasms
 - Epidemiology, Zaire, 78-4968
- Microsomes, Liver
 - Ames Test, 78-4969
 - Mutagenic Activity, 78-4969

Aflatoxin G2

- Food Contamination
 - Quantitation Method, 78-4959, 78-4965

Aflatoxin M1

- Food Contamination
 - Quantitation Method, 78-4964
 - Quantitation Method, Review, 78-4838
 - Review, 78-4837
- Microsomes, Liver
 - Mutagenic Activity, 78-4969

Aging

- Carcinogen, Chemical
 - Immunity, Cellular, Review, 78-4906
- Homeostasis
 - Cell Transformation, Neoplastic, Review, 78-4906
- Immunosuppression
 - Cell Transformation, Neoplastic, Review, 78-4906
- Insulin
 - Immunity, Cellular, Review, 78-4906
- T-Lymphocytes
 - Immunity, Cellular, Review, 78-4906

Agrobacterium tumefaciens

- Acetic Acid, (Ethylenedinitrilo)tetra-
 - Proteins, 78-5400
- Muramidase
 - Proteins, 78-5400
- Proteins
 - Isolation and Characterization, 78-5400
 - Plasmids, 78-5400
 - Tumorigenicity, 78-5400

Air Pollutants

- Power Plants, Fossil-Fueled
 - Carcinogenic Potential, Review, 78-4848
 - Physicochemical Properties, Review, 78-4848

Air Pollution

- Asbestos
 - Epidemiology, Review, 78-4841, 78-4842
- Benzene, (Epoxyethyl)
 - Fossil-Fuel Combustion Products, 78-5032
- Coronene
 - Fossil-Fuel Combustion Products, 78-5032
- m*-Cresol
 - Fossil-Fuel Combustion Products, 78-5032
- Lung Neoplasms
 - Asphalt Plant, 78-5378
 - Epidemiology, 78-5378
 - Epidemiology, Great Britain, Review, 78-4849
- Phenol
 - Fossil-Fuel Combustion Products, 78-5032
- Plutonium
 - Carcinogenic Potential, Review, 78-4868
- Pyrene
 - Fossil-Fuel Combustion Products, 78-5032
- Radium
 - Carcinogenic Potential, Review, 78-4868

Alanine, 3-Anthraniloyl-

- Bladder Neoplasms
 - Tryptophan, 78-5338

Alanine, 3-(*p*-(Bis(2-chloroethyl)amino)phenyl)-

- Carcinogenic Potential
 - Review, 78-4843
- Leukemia, Lymphoblastic
 - Drug Therapy, 78-5302

Alanine, 3-(3-Hydroxyanthraniloyl)-

- Bladder Neoplasms
 - Tryptophan, 78-5338

Aldrin

- Acrylonitrile
 - Enzymatic Activity, 78-5111
- Hydroxymethyl Derivatives
 - Metabolism, Rat, 78-5161
- Liver
 - Metabolism, Rat, 78-5161
- NADP
 - Metabolism, Rat, 78-5161

Alexan

- see* Thrombin

Alkaline Phosphatase

- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer
 - Enzymatic Activity, 78-5165

Alkeran

- see* Alanine, 3-(*p*-(Bis(2-chloroethyl)amino)phenyl)-

Ames Test

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Dose-Response Study, 78-4984
- Acetamide, *N*-Fluoren-2-yl-
 - Microsomes, Kidney, 78-5078
- Aflatoxin B1
 - Carcinogenic Potential, Review, 78-4815
 - Microsomes, Liver, 78-4969
 - Mutagenic Activity, Review, 78-4811
- Aflatoxin G1
 - Microsomes, Liver, 78-4969

nes Test (cont'd)

- Anthracene
 - Dihydrodiols, 78-4855
- Arginine
 - Pyrolysis Products, 78-5122
- Benz(a)anthracene
 - Carcinogenic Metabolite, Review, 78-4855
- Benz(a)anthracene, 3,4-Dihydroxy-1,2-oxy-1,2,3,4-tetrahydro-
 - Carcinogenic Metabolite, Review, 78-4855
- Benz(a)anthracene, 7,12-Dimethyl-
 - DNA, Binding, 78-5021
- Benzenamine, 4,4'-Methylenebis(2-chloro)-
 - Carcinogenic Potential, Review, 78-4815
- Benzene, (Epoxyethyl)-
 - Dose-Response Study, 78-4984
- Benzidine
 - Carcinogenic Potential, Review, 78-4815
- Benzo(a)pyrene
 - Carcinogenic Metabolite, Review, 78-4855
 - Carcinogenic Potential, Review, 78-4815
 - DNA, Binding, 78-5021
 - Ellipticine, 78-5011
 - Mutagenic Activity, Review, 78-4811
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Carcinogenic Metabolite, Review, 78-4855
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
 - Carcinogenic Metabolite, Review, 78-4855
 - Dose-Response Study, 78-4984
- Benzo(a)pyrene 7,8-Oxide
 - Carcinogenic Metabolite, Review, 78-4855
- Biphenyl, 2-Bromo-
 - Microsomes, Liver, 78-4978
- Biphenyl, 3-Bromo-
 - Microsomes, Liver, 78-4978
- Biphenyl, 4-Bromo-
 - Microsomes, Liver, 78-4978
- 4-Biphenylamine
 - Carcinogenic Potential, Review, 78-4815
- 1-Butanamine, *N*-Butyl-*N*-nitroso-
 - Carcinogenic Potential, Review, 78-4815
- Carcinogen, Chemical
 - Aroclor 1254, 78-4938
 - Microsomes, Liver, 78-4940
 - Riboflavine, 78-5048
 - Somatic Mutation, Review, 78-4809
- Chloral Hydrate
 - Mutagenic Activity, 78-5160
- Chlordiazepoxide
 - Mutagenic Activity, 78-5160
- Cholanthren-1-ol, 3-Methyl-
 - Microsomes, Liver, 78-4995
- Cholanthren-2-ol, 3-Methyl-
 - Microsomes, Liver, 78-4995
- Cholanthren-1-one, 3-Methyl-
 - Microsomes, Liver, 78-4995
- Cholanthren-2-one, 3-Methyl-
 - Microsomes, Liver, 78-4995
- Cholanthrene, 9,10-Dihydro-3-methyl-1,9,10-trihydroxy-
 - Microsomes, Liver, 78-4995
- Cholanthrene, 3-Methyl-
 - Microsomes, Liver, 78-4995
- Citrulline
 - Pyrolysis Products, 78-5122
- Diazepam
 - Mutagenic Activity, 78-5160

Ames Test (cont'd)

- Dibenz(a,h)anthracene
 - Carcinogenic Potential, Review, 78-4815
- Diethylamine, *N*-Nitroso-
 - Carcinogenic Potential, Review, 78-4815
- Dimethylamine, *N*-Nitroso-
 - Carcinogenic Potential, Review, 78-4815
- Ellipticine
 - Hydroxy Derivative, 78-5011
- Ethane, 2-Bromo-2-chloro-1,1,1-trifluoro-
 - Mutagenic Activity, 78-5160
- Ether, Bis(2-chloroethyl)-
 - Mutagenic Activity, Review, 78-4811
- Ether, 1-Chloro-2,2,2-trifluoroethyl Difluoromethyl
 - Mutagenic Activity, 78-5160
- Ether, 2,2-Dichloro-1,1-difluoroethyl Methyl
 - Mutagenic Activity, 78-5160
- Ethylene, Chloro-
 - Mutagenic Activity, Review, 78-4811
- Guanidine, Methyl-
 - Gastric Juice, 78-5103
 - Nitrous Acid, 78-5103
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Nitrosated Derivatives, 78-5103
- Lysine
 - Pyrolysis Products, 78-5122
- Methanesulfonic Acid, Methyl Ester
 - Carcinogenic Potential, Review, 78-4815
- Plasmids, 78-4950
- Mutagens
 - Feces, 78-5121
- Mycotoxins
 - Food Contamination, 78-5048
- Naphthalene
 - Dihydrodiols, 78-4855
- 2-Naphthylamine
 - Carcinogenic Potential, Review, 78-4815
- Nitrosamines
 - Aroclor 1254, 78-5097
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5097
 - Carcinogenic Activity, Correlation, 78-5097
- Opium
 - Microsomes, Liver, 78-5049
- Ornithine
 - Pyrolysis Products, 78-5122
- 1,2-Oxathiolane, 2,2-Dioxide
 - Carcinogenic Potential, Review, 78-4815
- Oxirane, 9-Anthryl-
 - Dose-Response Study, 78-4984
- Oxirane, 7-Benzanthryl-
 - Dose-Response Study, 78-4984
- Oxirane, 6-Chrysenyl-
 - Dose-Response Study, 78-4984
- Oxirane, 9-Methyl-10-anthryl-
 - Dose-Response Study, 78-4984
- Oxirane, 2-Naphthyl-
 - Dose-Response Study, 78-4984
- Oxirane, 9-Phenanthryl-
 - Dose-Response Study, 78-4984
- Oxirane, 1-Pyrenyl-
 - Dose-Response Study, 78-4984
- Phenanthrene
 - Dihydrodiols, 78-4855
- o*-Phenylenediamine, 4-Nitro-
 - Carcinogenic Potential, Review, 78-4809
- p*-Phenylenediamine, 2-Nitro-
 - Carcinogenic Potential, Review, 78-4809

Ames Test (cont'd)

- Polycyclic Hydrocarbons
 - Aroclor 1254, 78-4938
- 1-Propanol, 2,3-Dibromo-, Phosphate
 - Carcinogenic Potential, Review, 78-4809
- Smoke Condensate
 - Benzo(a)pyrene, 78-5048
 - Broiled Fish, Meat, 78-5048
 - Tryptophan, 78-5048
- Sterigmatocystin
 - Carcinogenic Potential, Review, 78-4815
- Tobacco
 - Pyrolysis Products, 78-5049
- Toluene-2,4-diamine
 - Mutagenic Activity, Review, 78-4811
- Tryptophan
 - Pyrolysis Products, 78-5122
- Urea, Methyl Nitroso-
 - Carcinogenic Potential, Review, 78-4815

Amino Acid Naphthylamidases

- Letterer-Siwe Disease
 - Enzymatic Activity, 78-5389

Amino Acids

- Myeloma Proteins
 - Immunoglobulins, Heavy Chain, 78-5277
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-5259
 - Contact Inhibition, 78-5259
 - Isobutyric Acid, α -Amino-, 78-5259
 - Membrane Transport, 78-5259
 - Proline, 78-5259

Aminopyrine

- see Antipyrine, 4-(Dimethylamino)-

Anagestone Acetate

- Mammary Neoplasms, Experimental
 - Hyperplasia, 78-5132

5 α -Andostan-3-one, 17 β -Hydroxy-

- Prostatic Neoplasms
 - Hyperplasia, 78-4920

Androgens

- Prostatic Neoplasms
 - Hyperplasia, 78-4920

Androstanediol

- Prostatic Neoplasms
 - Hyperplasia, 78-4920

Anemia, Aplastic

- Leukemia, Myeloblastic
 - Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-, 78-5147
- Methanesulfonic Acid, Ethyl Ester
 - Chromosome Aberrations, 78-4939
- Methanesulfonic Acid, Methyl Ester
 - Chromosome Aberrations, 78-4939
- Micronucleus Test
 - Chromosome Aberrations, 78-4939
- Mitomycin C
 - Chromosome Aberrations, 78-4939
 - Micronucleus Test, 78-4939
- Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-
 - Case Report, 78-5147
- Radiation, Ionizing
 - Chromosome Aberrations, 78-4939

Anemia, Aplastic (cont'd)

- Micronucleus Test, 78-4939
- Uridine, 5-Bromo-2'-deoxy-
 - Chromosome Aberrations, 78-4939
- Virus, Polyoma, BK
 - Bone Marrow Transplantation, 78-5264

Anemia, Macrocytic

- Chromosome Aberrations
 - Chromosomes, Human, 1-3, 78-5296
- Leukemia, Myeloblastic
 - Chromosome Aberrations, 78-5296

Anemia, Pernicious

- Nitrosamines
 - Gastric Juice, 78-5330
- Nitrous Acid
 - Gastric Juice, 78-5330
- Stomach Neoplasms
 - Nitrous Acid, 78-5330
 - Precancerous Conditions, 78-5330

Anesthetics

- Hematopoietic Stem Cells
 - Immune Response, Review, 78-4858
- Hepatitis
 - Epidemiology, Review, 78-4857
- Leukemia
 - Epidemiology, Review, 78-4857
- Lymphoma
 - Epidemiology, Review, 78-4857
- Occupational Hazard
 - Abnormalities, 78-4858
 - Carcinogenic Potential, Review, 78-4858
 - Epidemiology, Review, 78-4857, 78-4858

Angiosarcoma

- Arsenic
 - Epidemiology, Review, 78-4934
- Ethylene, Chloro-
 - Epidemiology, 78-4832
- Lung
 - Neoplasm Metastasis, 78-5388
- Prosthesis
 - Case Report, Hip Tumor, 78-5305
- Radiation, Ionizing
 - Case Report, Hip Tumor, 78-5305
 - Radiotherapy, 78-5305
- p*-Tolamide, *N*-Isopropyl- α -(2-methylhydrazino)-
 - Carcinogenic Activity, Monkey, 78-5148
- Virus, Polyoma, BK
 - Carcinogenic Activity, Hamster, 78-5263

Aniline, 2,6-Dinitro-*N,N*-dipropyl-*p*-isopropyl-

- Dipropylamine, *N*-Nitroso-
 - Pesticides, Review, 78-4824

Aniline, *N*-Methyl-*p*-(phenylazo)-

- Metabolic Activation
 - Carcinogenic Metabolite, Review, 78-4816

Aniline, 4,4'-Oxydi-

- Bone Neoplasms
 - Precancerous Conditions, 78-5126
- Hyperplasia
 - Dose-Response Study, 78-5126
 - Pituitary Gland, 78-5126
 - Thyroid Gland, 78-5126
- Hyperthyroidism
 - Dose-Response Study, 78-5126
 - Goitrogenic Effect, Mouse, Rat, 78-5126

ne, 4,4'-Oxydi- (cont'd)

- Pituitary Neoplasms
 - Precancerous Conditions, 78-5126
- Thyroid Neoplasms
 - Precancerous Conditions, 78-5126
- Urogenital Neoplasms
 - Precancerous Conditions, 78-5126

thracenamide

- Xeroderma Pigmentosum
 - DNA Repair, 78-5170

racene

- Ames Test
 - Dihydrodiols, 78-4855
- Carcinogenic Metabolite
 - Fossil-Fuel Combustion Products, 78-5032

racene, 9,10-Bis(oxiranyl)-

- Carcinogenic Metabolite
 - Synthesis, 78-5035

racosilicosis

- Phospholipids
 - Scleroproteins, 78-4971

ranilic Acid, 3-Hydroxy-

- see Benzoic Acid, 2-Amino-3-hydroxy-

Antibodies

- Sarcoma 180, Crocker
 - T-Lymphocytes, 78-5279
- Virus, Epstein-Barr
 - Binding, Cell Membrane, 78-5278
 - IgM, 78-5278

odies, Neoplasm

- Brain Neoplasms
 - Urea, Methyl Nitroso-, 78-5281
- Glioma
 - Urea, Methyl Nitroso-, 78-5281

odies, Viral

- Virus, Bovine Leukemia
 - Seroepidemiology, Review, 78-4887
- Virus, Feline Leukemia
 - Immune Response, Review, 78-4885
- Virus, Feline Sarcoma
 - Immune Response, Review, 78-4885

ody Formation

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - DNA, 78-4986
- Diphenylamine, *N*-Nitroso-
 - Antigens, Bacterial, 78-5086
 - Immune Response, Rat, 78-5086
- Urea, Methyl Nitroso-
 - Antigens, Bacterial, 78-5086
 - Immune Response, Rat, 78-5086
- Virus, Coxsackie B
 - Hypercholesteremia, 78-5290
- Virus, Friend Murine Leukemia
 - Immunosuppression, 78-5207
- Virus, Murine Leukemia
 - Immune Response, Wild Mouse, Review, 78-4883
- Virus, Rous Sarcoma
 - Genetics, 78-5292
 - Radiation, Ionizing, 78-5292

ody Specificity

- Lymphoma
 - Virus, Abelson Murine Leukemia, 78-5200

Antibody Specificity (cont'd)

- Virus, Abelson Murine Leukemia
 - Antigens, Viral, 78-5200
- Virus, SV40
 - Antigens, Viral, 78-5251

Antifungal Agents

- Aspergillus nidulans*
 - Methionine Requirement, Fluctuation Test, 78-4944

Antigen-Antibody Reactions

- Ependymoma
 - Virus, Polyoma, BK, 78-5263
- Virus, Bovine Leukemia
 - Diagnosis, Review, 78-4886
- Virus, Feline Leukemia
 - Guinea Pig, 78-5228
 - Hypotonic Medium, 78-5227
- Virus, Feline Sarcoma
 - Guinea Pig, 78-5228
- Virus, Friend Murine Leukemia
 - Cells, Cultured, 78-5283
 - Lipopolysaccharides, 78-5207
 - B-Lymphocytes, 78-5283
 - Macrophages, 78-5207

Antigenic Determinants

- Ultraviolet Rays
 - Transplantation Immunology, 78-5285
- Virus, Abelson Murine Leukemia
 - Cell Transformation, Neoplastic, 78-5199
 - Viral Proteins, 78-5199
- Virus, Friend Murine Leukemia, 78-5200
- Virus, Moloney Murine Leukemia, 78-5200
- Virus, Rauscher Murine Leukemia, 78-5200
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5238
- Virus, Avian Carcinoma
 - Viral Proteins, 78-5190
- Virus, Friend Murine Leukemia
 - Virus, Mink Cell Focus-Inducing, 78-5205
- Virus, Friend Spleen Focus-Forming
 - Radioimmunoassay, 78-5205
- Virus, Friend Murine Leukemia, 78-5208
- Virus, Mink Cell Focus-Inducing, 78-5205
- Virus, Moloney Murine Leukemia
 - Virus, Mink Cell Focus-Inducing, 78-5205
- Virus, Murine Leukemia
 - Virus, Mink Cell Focus-Inducing, 78-5205
- Virus, Polyoma, BK
 - Isolation and Characterization, Review, 78-4893
- Virus, Rauscher Murine Leukemia
 - Viral Proteins, 78-5202
- Virus, RNA Tumor
 - Reverse Transcriptase, 78-5216
- Virus, Rous Sarcoma
 - Cell-Free Synthesis, 78-5193
 - Viral Proteins, 78-5193
- Virus, SV40
 - Antigens, Viral, 78-5251

Antigens, Bacterial

- Diphenylamine, *N*-Nitroso-
 - Antibody Formation, 78-5086
- Urea, Methyl Nitroso-
 - Antibody Formation, 78-5086

Antigens, Neoplasm

- Leukemia, Lymphoblastic
 - Lymphocyte Culture Test, Mixed, 78-5286

Antigens, Neoplasm (cont'd)

- Leukemia, Myeloblastic
Blood Platelets, 78-5397
Precancerous Conditions, 78-5397

Antigens, Viral

- Benz(a)anthracene, 7,12-Dimethyl-
Virus, Polyoma, 78-4999
- Benzo(a)pyrene
Virus, Polyoma, 78-4999
- Cervix Neoplasms
Virus, Herpes Simplex 2, 78-5242
- Cholanthrene, 3-Methyl-
Virus, Polyoma, 78-4999
- Dibenz(a,h)anthracene
Virus, Polyoma, 78-4999
- Lymphoma
Cholanthrene, 3-Methyl-, 78-5291
Virus, Abelson Murine Leukemia, 78-5200
- Virus, Abelson Murine Leukemia
Antibody Specificity, 78-5200
Cell Differentiation, 78-5200
Cell Transformation, Neoplastic, 78-5200
- Virus, Adeno
Immune Response, Review, 78-4905
- Virus, Adeno 12
Isolation and Characterization, 78-5238
Temperature Sensitive Mutants, 78-5238
- Virus, Baboon
Hybrid Cells, 78-5231
- Virus, Feline Leukemia
Immune Response, Review, 78-4884
Viral Vaccines, Review, 78-4884
- Virus, Herpes Simplex 2
Precancerous Conditions, 78-5242
- Virus, Murine Mammary Tumor
Radioimmunoassay, 78-5223
- Virus, Polyoma
Temperature Sensitive Mutants, 78-5265
- Virus, SV40
Antibody Specificity, 78-5251
Antigenic Determinants, 78-5251
Carcinogenic Activity, Review, 78-4894
Cell Cycle Kinetics, 78-5258
DNA Replication, 78-5248, 78-5253
DNA Replication, Review, 78-4894
Immune Response, Hamster, 78-5251
Immune Response, Review, 78-4905
Immune Serums, 78-5251
Phenotype Complementation, 78-5248
RNA, Messenger, 78-5249
RNA Replication, 78-5258
RNA, Ribosomal, 78-5258
Temperature Sensitive Mutants, 78-5248

Antilymphocyte Serum

- Carcinogenic Potential
Review, 78-4843
- Virus, Herpes Ateles
T-Lymphocytes, 78-5239

Antineoplastic Agents

- Neoplasms, Experimental
Carcinogen, Chemical, 78-4807

Antipyrine, 4-(Dimethylamino)-

- Dimethylamine, *N*-Nitroso-
Nitrous Acid, Sodium Salt, 78-5076
- Nitrous Acid

Antipyrine, 4-(Dimethylamino)- (cont'd)

- Carcinogenic Activity, Review, 78-4819

Areca

- Esophageal Neoplasms
Epidemiology, Pakistan, 78-5382
Epidemiology, Sri Lanka, 78-5381
- Mouth Neoplasms
Carcinogenic Activity, Mouse, Rat, 78-4955
- Sarcoma
Carcinogenic Activity, Mouse, Rat, 78-4955
- Tobacco
Carcinogenic Activity, Mouse, Rat, 78-4955

Arginine

- Ames Test
Pyrolysis Products, 78-5122
- Mutagens
Inactivation, Fruit/Vegetable Juices, 78-5122
Pyrolysis Products, 78-5122

Aroclor 1254

- Acrylonitrile
Lactate Dehydrogenase, 78-5111
- Carcinogen, Chemical
Ames Test, 78-4938
- Cholanthrene, 3-Methyl-
Mutagenic Activity, 78-4995
- Dimethylamine, *N*-Nitroso-
Nucleic Acids, Binding, 78-5075
Oxidoreductases, *N*-Demethylating, 78-5075
- Ellipticine
Aryl Hydrocarbon Hydroxylases, 78-5011
- Hepatoma
Carcinogenic Activity, Review, 78-4828
- Nitrosamines
Ames Test, 78-5097
- Polycyclic Hydrocarbons
Ames Test, 78-4938

Aroclor 1260

- Hepatoma
Carcinogenic Activity, Review, 78-4828

Arsenic

- Angiosarcoma
Epidemiology, Review, 78-4934
- Carcinogenic Potential
Review, 78-4843
- Liver Neoplasms
Precancerous Conditions, Review, 78-4922
- Lymphoma
Epidemiology, Review, 78-4934
- Occupational Hazard
Epidemiology, Review, 78-4934
- Skin Neoplasms
Epidemiology, Review, 78-4934

Arthritis, Rheumatoid

- Leukemia, Myeloblastic
Drug Therapy, 78-5151
Immunosuppressive Agents, 78-5153

Aryl Hydrocarbon Hydroxylases

- Abnormalities
Genetics, Review, 78-4927
- Acetamide, *N*-Fluoren-2-yl-
Carcinogenic Metabolite, 78-4988
- Aroclor 1254
Ellipticine, 78-5011
- Barbituric Acid, 5-Ethyl-5-phenyl-

aryl Hydrocarbon Hydroxylases (cont'd)

- Ellipticine, 78-5011
- Benzo(a)pyrene**
 - Dietary Fiber, 78-5002
 - DNA, Binding, 78-5008
 - Ellipticine, 78-5011
 - Genetics, Review, 78-4928
 - Metabolism, Liver, 78-5008
 - Retinol, 78-5019
- Benzo(b)triphenylene**
 - Fish, 78-4990
- 7,8-Benzoflavone**
 - Microsomes, Liver, 78-5011
- Carcinogen, Chemical**
 - Genetics, Review, 78-4927
- Cell Transformation, Neoplastic**
 - Genetics, Review, 78-4927
- Cholanthrene, 3-Methyl-**
 - Fish, 78-4990
- Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-**
 - Liver, Mouse, Review, 78-4852
- DNA Repair**
 - Genetics, Review, 78-4927
- Ellipticine**
 - Microsomes, Liver, 78-5011
- Laryngeal Neoplasms**
 - Genetics, Review, 78-4928
- Lung Neoplasms**
 - Genetics, Review, 78-4928
- Oncogenic Viruses**
 - Genetics, Review, 78-4927
- Polycyclic Hydrocarbons**
 - Genetics, Review, 78-4927
- 1-Propanone, 2-Methyl-1,2-di-3-pyridyl-, Tartrate**
 - Microsomes, Liver, 78-5011
- Respiratory Tract Neoplasms**
 - Genetics, 78-5324
 - Lymphocytes, 78-5324
- Retinol**
 - Lung, Hamster, 78-5019
- Smoking**
 - Genetics, Review, 78-4928

Asbestos

- Air Pollution**
 - Epidemiology, Review, 78-4841, 78-4842
- Cysteine**
 - Macrophages, 78-5284
- Digestive System Neoplasms**
 - Epidemiology, Review, 78-4842
- Gastrointestinal Neoplasms**
 - Epidemiology, Review, 78-4841, 78-4842
- Lung Neoplasms**
 - Adenocarcinoma, 78-5326
 - Epidemiology, Review, 78-4841, 78-4842
- T-Lymphocytes**
 - Immune Response, Rat, 78-5284
- Macrophages**
 - Immune Response, Rat, 78-5284
- Mesothelioma**
 - Epidemiology, 78-5377, 78-5385
 - Epidemiology, Review, 78-4842
 - Epidemiology, Turkey, 78-4909
- Neoplasms, Experimental**
 - Trans-Species Extrapolation, 78-4805
- Occupational Hazard**
 - Epidemiology, 78-5377
 - Epidemiology, Review, 78-4842

Asbestos (cont'd)

- Periodic Acid, Sodium Salt
 - Macrophages, 78-5284
- Peritoneal Neoplasms
 - Epidemiology, Review, 78-4842
- Phospholipids
 - Lung, Rat, 78-4971
- Pleural Neoplasms
 - Epidemiology, Review, 78-4842
- Sclerosis
 - Lung, Rat, 78-4971
- Smoking
 - Co-carcinogenic Activity, Review, 78-4842

Asbestosis

- Phospholipids
- Scleroproteins, 78-4971

Ascorbic Acid

- Benz(a)anthracene, 7-Methyl-Dihydrodiols, 78-5031
- Mutagens
 - Feces, 78-5121
- Nitrous Acid
 - Metabolism, Intestines, 78-5052
- Stomach Neoplasms
 - Epidemiology, Review, 78-4912

Aspergillus flavus

- Aflatoxin B1
 - Drug Contamination, 78-4956

Aspergillus nidulans

- Antifungal Agents
 - Methionine Requirement, Fluctuation Test, 78-4944
- Mutagens
 - Methionine Requirement, Fluctuation Test, 78-4944

Astrocytoma

- Brain Neoplasms
 - Urea, 1,3-Dimethyl-1-nitroso-, 78-5096
- Neoplasm Transplantation
 - Ultrastructural Study, 78-5089
- Urea, Ethyl Nitroso-
 - Ultrastructural Study, 78-5089

Ataxia Telangiectasia

- Genetics
 - Epidemiology, Review, 78-4926
- Skin Neoplasms
 - Epidemiology, Review, 78-4930

Atropine

- Esophageal Neoplasms
 - Dimethylamine, *N*-Nitroso-, 78-5067
- Liver Neoplasms
 - Dimethylamine, *N*-Nitroso-, 78-5067

Australia Antigen

- Hepatoma
 - Cytoplasm, 78-5276
 - Epidemiology, Review, 78-4846
 - Liver Cirrhosis, 78-5276
- Schistosomiasis
 - Epidemiology, Review, 78-4846

Autoimmune Diseases

- IgA
 - Mouse, AKR, 78-5280
- Virus, Gross Murine Leukemia
 - Precancerous Conditions, 78-5280

- Azathioprine**
Lymphocytes
Chromosome Aberrations, 78-5157
- Bacillus subtilis**
Intestinal Bacteria
Mutagenic Activity, 78-4952
- Bacteria**
Carcinogen, Chemical
Mutagenic Activity, Review, 78-4811
Colonic Neoplasms
Carcinogen, Chemical, 78-5123
Dietary Fats, 78-5123
Diethylamine, *N*-Nitroso-
Mutagenic Activity, Review, 78-4822
Intestines
Mutagenic Activity, 78-4952
- Barbituric Acid, 5-Ethyl-5-phenyl-**
Acetamide, *N*-Fluoren-3-yl-
Metabolism, Rat, Guinea Pig, 78-4985
Acetic Acid, (2,3-Dichloro-4-(2-methylenebutyl)phenoxy)
Transferases, 78-5016
Acrylonitrile
Lactate Dehydrogenase, 78-5111
Benzene, 1-Chloro-2,4-dinitro-
Transferases, 78-5016
Benzene, 1-(Chloromethyl)-4-nitro-
Transferases, 78-5016
Benzene, 1,2-Dichloro-4-nitro-
Transferases, 78-5016
Benzo(a)pyrene
DNA, Binding, 78-5010
Metabolism, Liver, 78-5008
Cadmium
Oxidoreductases, *N*-Demethylating, 78-5118
Carcinogen, Chemical
Inhibitors, Review, 78-4806
Ellipticine
Aryl Hydrocarbon Hydroxylases, 78-5011
Liver Neoplasms
Histochemical/Histological Study, Mouse, 78-5325
Precancerous Conditions, 78-5325
Nitrosamines
Ames Test, 78-5097
Propane, 1,2-Epoxy-3-(*p*-nitrophenoxy)-
Transferases, 78-5016
Transferases
Enzymatic Activity, Rat, 78-5016
- Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt**
Hydroxylamine
Cytochrome P-450, 78-4989
- BCG**
Benz(a)anthracene, 7,12-Dimethyl-
Carcinogenic Activity, Mouse, 78-5030
Sarcoma
Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
Thymoma
Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
Virus, Herpes Simplex 2
Immunity, Cellular, 78-5243
- Behcet's Syndrome**
Leukemia, Myeloblastic
Cyclophosphamide, 78-5153
- Benz(a)anthracene**
Ames Test
Carcinogenic Metabolite, Review, 78-4855
Benz(a)anthracene, 3,4-Dihydroxy-1,2-oxy-1,2,3,4-tetrahydro-
Carcinogenic Metabolite, 78-5022
Skin Neoplasms
Papilloma, 78-5022
12-*O*-Tetradecanoylphorbol-13-acetate, 78-5022
- Benz(a)anthracene, 1,2-Dihydro-**
Skin Neoplasms
Carcinogenic Potential, 78-5022
- Benz(a)anthracene, 3,4-Dihydro-**
Skin Neoplasms
Carcinogenic Potential, 78-5022
- Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-**
Skin Neoplasms
Carcinogenic Potential, 78-5022
- Benz(a)anthracene, 8,9-Dihydro-8,9-dihydroxy-7,12-dimethyl-**
Chloroform
Water-Soluble Metabolite, 78-5023
- Benz(a)anthracene, 1,2-Dihydro-1,2-dihydroxy-7-methyl-**
Benz(a)anthracene, 7-Methyl-
Spectrum Analysis, 78-5031
- Benz(a)anthracene, 3,4-Dihydroxy-1,2-oxy-1,2,3,4-tetrahydro-**
Ames Test
Carcinogenic Metabolite, Review, 78-4855
Benz(a)anthracene
Carcinogenic Metabolite, 78-5022
- Benz(a)anthracene, 7,12-Dimethyl-**
Ames Test
DNA, Binding, 78-5021
- BCG**
Carcinogenic Activity, Mouse, 78-5030
Carcinoma, Epidermoid
Cells, Cultured, 78-5024
12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
Cell Transformation, Neoplastic
Cells, Cultured, 78-5024
Trachea, Rat, 78-5024
Chloroform
Metabolism, Hamster Embryo Cells, 78-5023
Chromosomes
DNA, Binding, 78-5027
DNA
Carcinogenic Metabolite, 78-5026
DNA, Binding
Indirect Immunofluorescence, 78-5027
Glucuronidase
Metabolism, Hamster Embryo Cells, 78-5023
Phenol Glucuronides, 78-5023
Keratoacanthoma
12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
Liver Neoplasms
Hemangioma, 78-5030
Lung Neoplasms
Adenoma, 78-5030
Mammary Neoplasms, Experimental
Adenocarcinoma, 78-5030
Dietary Fats, 78-5025
Stress, 78-5029
Mouth Neoplasms
Histological Study, Hamster, 78-5028
Precancerous Conditions, 78-5028

12(a)anthracene, 7,12-Dimethyl- (cont'd)

- Mutagenic Activity
 - Hamster V79 Cells, 78-4999
- Nervous System Neoplasms
 - Carcinogenic Metabolite, 78-5026
 - Transplacental Carcinogenesis, 78-5026
- Ovarian Neoplasms
 - Cystadenoma, 78-5030
- Papilloma
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
- Sarcoma
 - BCG, 78-5030
- Skin Neoplasms
 - Carcinoma, Epidermoid, 78-5036
 - Keratoacanthoma, 78-5036
 - Papilloma, 78-5036
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Neoplasm Regression, Spontaneous, 78-5036
- Thymoma
 - BCG, 78-5030
- Virus, Adeno 5
 - Cell Transformation, Neoplastic, 78-5041
- Virus, Polyoma
 - Antigens, Viral, 78-4999

12(a)anthracene, 7-Methyl-

- Ascorbic Acid
 - Dihydrodiols, 78-5031
- Benz(a)anthracene, 1,2-Dihydro-1,2-dihydroxy-7-methyl-
 - Spectrum Analysis, 78-5031
- Sulfuric Acid, Iron Salt
 - Dihydrodiols, 78-5031

12-N-aminamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-

- Hepatoma
 - Precancerous Conditions, 78-5127
 - Proton Magnetic Relaxation Times, 78-5127
- Kanechlor 500
 - Hepatocarcinogenicity, Review, 78-4828

12-N-aminamine, 4,4'-Methylenebis(2-chloro)-

- Ames Test
 - Carcinogenic Potential, Review, 78-4815

12-N-azene

- Acid Phosphatase
 - Enzymatic Activity, 78-4974
 - Inhalation Study, Rat, 78-4974
- Binding, Covalent
 - Bone Marrow, Liver, Mouse, 78-4975
- Lymphocytes
 - Acid Phosphatase, 78-4974

12-N-azene, 4-Allyl-1,2-(Methylenedioxy)-

- Metabolic Activation
 - Carcinogenic Metabolite, Review, 78-4816

12-N-azene, 1-Chloro-2,4-dinitro-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Transferases, 78-5016
- Benzo(a)pyrene
 - Transferases, 78-5016
- Cholanthrene, 3-Methyl-
 - Transferases, 78-5016
- Transferases
 - Spermatozoa, 78-5012
- Ultraviolet Rays
 - Hypersensitivity, Delayed, 78-5173

Benzene, 1-(Chloromethyl)-4-nitro-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Transferases, 78-5016

Benzo(a)pyrene

- Transferases, 78-5016

Cholanthrene, 3-Methyl-

- Transferases, 78-5016

Benzene, 1,2-Dichloro-4-nitro-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Transferases, 78-5016

Benzo(a)pyrene

- Transferases, 78-5016

Cholanthrene, 3-Methyl-

- Transferases, 78-5016

Benzene, (Epoxyethyl)-

- Air Pollution
 - Fossil-Fuel Combustion Products, 78-5032

Ames Test

- Dose-Response Study, 78-4984

Chromosome Aberrations

- Bone Marrow Cells, 78-4953

Lymphocytes, 78-4953

Germ Cells

- Mutagenic Activity, 78-4953

Hydro-Lyases

- Liver, Rat, 78-5013

Micronucleus Test

- Mutagenic Activity, 78-4953

Transferases

- Spermatozoa, 78-5012

Virus, SV40

- DNA, Binding, 78-4984

α -Benzene Hexachloride

- see Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer

γ -Benzene Hexachloride

- see Cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer

Benzene, Hexachloro-

- Hepatoma
 - Kanechlor C, 78-4979

Kanechlor C

- Co-carcinogenic Activity, 78-4979

Benzene, (2-Isocyanatomethyl)-

- Carcinogen, Chemical
 - Inhibitors, Review, 78-4806

Benzidine

- Ames Test
 - Carcinogenic Potential, Review, 78-4815

Benzimidazole, 5,6-Dichloro-1- β -D-ribofuranosyl-

- Virus, Adeno 2
 - DNA Replication, 78-5233
 - RNA, Viral, 78-5233

Benzo(a)pyrene

- Acetic Acid, (2,3-Dichloro-4-(2-methylenebutyl)phenoxy)
 - Transferases, 78-5016
- Ames Test
 - Carcinogenic Metabolite, Review, 78-4855
 - Carcinogenic Potential, Review, 78-4815
 - DNA, Binding, 78-5021
 - Mutagenic Activity, Review, 78-4811
- Arene Oxides
 - Enzyme Activation, Review, 78-4854
- Aryl Hydrocarbon Hydroxylases

Benzo(a)pyrene (cont'd)

- Dietary Fiber, 78-5002
- DNA, Binding, 78-5008
- Genetics, Review, 78-4928
- Metabolism, Liver, 78-5008
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - DNA, Binding, 78-5010
 - Metabolism, Liver, 78-5008
- Benzene, 1-Chloro-2,4-dinitro-
 - Transferases, 78-5016
- Benzene, 1-(Chloromethyl)-4-nitro-
 - Transferases, 78-5016
- Benzene, 1,2-Dichloro-4-nitro-
 - Transferases, 78-5016
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Metabolism, 78-5020
- 5,6-Benzoflavone
 - DNA, Binding, 78-5010
- Carcinogenic Metabolite
 - DNA, Binding, 78-5010
 - Fossil-Fuel Combustion Products, 78-5032
- Carcinoma, Bronchiolar
 - Diol Epoxides, 78-5007
- Cell Transformation, Neoplastic
 - Cell Density, 78-5000
 - Cells, Cultured, 78-5000
 - Fetal Lung Cells, Hamster, 78-5001
- Cells, Cultured
 - Metabolism, 78-5020
 - Mutagenic Activity, 78-5020
- Cholanthrene, 3-Methyl-
 - Microsomes, Liver, 78-4854
- Cytochromes
 - DNA, Binding, 78-5010
- Dieldrin
 - Metabolism, Liver, 78-5008
- DNA
 - Binding, 78-4998
 - Diol Epoxides, 78-5007
 - Photodynamic Effects, 78-5003
- DNA, Binding
 - Carcinogenic Metabolite, 78-4997
 - Lung, Rat, 78-5010
 - Mutagenic Activity, 78-4997
- Ellipticine
 - Ames Test, 78-5011
 - Aryl Hydrocarbon Hydroxylases, 78-5011
 - DNA, Binding, 78-5011
- Enzyme Activation
 - Carcinogenic Metabolite, Review, 78-4855
- Escherichia coli*
 - Temperature Sensitive Mutants, 78-5003
- Ferric Chloride
 - Co-carcinogenic Activity, Hamster, 78-5004
- Glioma
 - Sex Hormones, 78-5135
- Hydro-Lyases
 - Carcinogenic Activity, Review, 78-4850
 - Enzyme Activation, Review, 78-4854
- Insulin
 - Cell Transformation, Neoplastic, 78-5001
- Intestinal Neoplasms
 - Dietary Fiber, 78-5002
- Iodine
 - Co-carcinogenic Activity, Hamster, 78-5004
- Lipids
 - Liver, Rat, 78-5014

Benzo(a)pyrene (cont'd)

- Liver Neoplasms
 - Mesenchymoma, 78-5014
- Lung
 - Metabolism, 78-5015
- Lung Neoplasms
 - Air Pollution, Review, 78-4849
- Lung, Rabbit
 - Metabolism, 78-5018
- Macrophages
 - Metabolism, 78-5020
- Mesenchymoma
 - Catalase, 78-5014
 - Glutathione Peroxidase, 78-5014
 - Glutathione Reductase, 78-5014
 - Urate Oxidase, 78-5014
- Methane, Sulfinylbis-
 - Cell Transformation, Neoplastic, 78-5001
- Microsomes, Liver
 - Metabolism, Review, 78-4854
- Mutagenic Activity
 - Hamster V79 Cells, 78-4999
 - Ouabain Resistance, 78-5020
- Nucleic Acids
 - Binding, Review, 78-4850
- Oxygenases
 - Enzyme Activation, Review, 78-4854
- Peroxides
 - Liver, Rat, 78-5014
- Propane, 1,2-Epoxy-3-(*p*-nitrophenoxy)-
 - Transferases, 78-5016
- Putrescine
 - Carcinogenic Activity, 78-5005
- Radiation
 - Cell Survival, 78-5003
 - Escherichia coli*, 78-5003
- Respiratory Tract Neoplasms
 - Adenocarcinoma, 78-5004, 78-5006
 - Adenoma, 78-5004, 78-5006
 - Carcinoma, 78-5006
 - Carcinoma, Epidermoid, 78-5004, 78-5006
 - Ferric Chloride, 78-5004
 - Hemangioma, 78-5006
 - Histological Study, Rat, 78-5006
 - Iodine, 78-5004
 - Papilloma, 78-5004, 78-5006
 - Talc, 78-5006
- Retinol
 - Aryl Hydrocarbon Hydroxylases, 78-5019
 - Metabolism, 78-5019
- Retinol Acetate
 - Metabolism, 78-5019
- Seminal Vesicles
 - Metabolism, 78-5015
- Skin Neoplasms
 - Putrescine, 78-5005
- Smoke Condensate
 - Ames Test, 78-5048
- Talc
 - Co-carcinogenic Activity, Rat, 78-5006
- Thymidine
 - Diol Epoxides, 78-5007
- Transferases
 - Enzymatic Activity, Rat, 78-5016
- Virus, Adeno 5
 - Cell Transformation, Neoplastic, 78-5041
- Virus, Polyoma

- benzo(a)pyrene (cont'd)**
Antigens, Viral, 78-4999
- benzo(a)pyrene, 4,5-Dihydro-4,5-dihydroxy-**
Carcinogenic Metabolite
Nuclear Magnetic Resonance Spectra, 78-5017
Synthesis, 78-5017
- benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-**
Ames Test
Carcinogenic Metabolite, Review, 78-4855
- Benzo(a)pyrene**
Metabolism, 78-5020
- Cells, Cultured**
Metabolism, 78-5020
Mutagenic Activity, 78-5020
- Macrophages**
Metabolism, 78-5020
- Mutagenic Activity**
Ouabain Resistance, 78-5020
- benzo(a)pyrene, 9,10-Dihydro-9,10-dihydroxy-**
Carcinogenic Metabolite
Nuclear Magnetic Resonance Spectra, 78-5017
Synthesis, 78-5017
- benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-**
Ames Test
Carcinogenic Metabolite, Review, 78-4855
Dose-Response Study, 78-4984
- Carcinogenic Metabolite**
Lung, Rat, 78-5010
- DNA Repair**
Cells, Cultured, 78-5009
- Lung Neoplasms**
DNA Repair, 78-5009
- Mutagenic Activity**
Hamster V79 Cells, 78-4999
Ouabain Resistance, 78-4999
- Virus, SV40**
DNA, Binding, 78-4984
- benzo(a)pyrene 4,5-Oxide**
Hydro-Lyases
Liver, Rat, 78-5013
- Transferases**
Spermatozoa, 78-5012
- benzo(a)pyrene 7,8-Oxide**
Ames Test
Carcinogenic Metabolite, Review, 78-4855
- benzo(b)triphenylene**
Acetic Acid
Benzoic Acid, 2-(9'-Phenanthroyl)-, 78-5034
- Aryl Hydrocarbon Hydroxylases**
Fish, 78-4990
- Benzoic Acid, 2-(9'-Phenanthroyl)-**
Synthesis, 78-5034
- Benzo(a)pyrene**
DNA, Binding, 78-5010
- Carcinogen, Chemical**
Inhibitors, Review, 78-4806
- Benzo(a)pyrene**
Dibenz(a,e)aceanthrylene
DNA, Binding, 78-5033
- Microsomes, Liver**
Aryl Hydrocarbon Hydroxylases, 78-5011
- Benzoic Acid, 2-Amino-3-hydroxy-**
Bladder Neoplasms
Tryptophan, 78-5338
- Smoking**
Metabolism, 78-5044
- Benzoic Acid, 2-(9'-Phenanthroyl)-**
Benzo(b)triphenylene
Acetic Acid, 78-5034
Synthesis, 78-5034
- Benzyl Isothiocyanate**
see Benzene, (2-Isocyanatomethyl)-
- Beryllium**
Occupational Hazard
Epidemiology, Review, 78-4934
Respiratory Tract Neoplasms
Carcinogenic Potential, Review, 78-4934
Sarcoma, Osteogenic
Carcinogenic Potential, Review, 78-4934
- Betel**
see Areca
- Bile Acids and Salts**
Breast Neoplasms
Feces, 78-5358
- Clostridium**
Carcinogenic Activity, Review, 78-4919
- Colonic Neoplasms**
Carcinogen, Chemical, 78-5123
- Intestinal Neoplasms**
Epidemiology, Review, 78-4919
Feces, 78-5358
- Bile Duct Neoplasms**
Adenocarcinoma
Dyes, 78-4973
- Cysts**
Morpholine, *N*-Nitroso-, 78-5056
Neoplasms, Multiple Primary
Occupational Hazard, 78-4973
- 3,4-Biphenyldiol, 4'-Bromo-**
Biphenyl, 4-Bromo-
Metabolism, Rat, 78-4978
- Biphenyl, 2-Bromo-**
Microsomes, Liver
Ames Test, 78-4978
- Biphenyl, 3-Bromo-**
Microsomes, Liver
Ames Test, 78-4978
- Biphenyl, 4-Bromo-**
3,4-Biphenyldiol, 4'-Bromo-
Metabolism, Rat, 78-4978
- 4-Biphenylol, 4'-Bromo-**
Metabolism, Rat, 78-4978
- Microsomes, Liver**
Ames Test, 78-4978
- 4-Biphenylamine**
Ames Test
Carcinogenic Potential, Review, 78-4815
Bladder Neoplasms
Carcinoma, Transitional Cell, 78-4977
Histological Study, Dog, 78-4977
- Tritium**
Catalytic Hydrogenation, 78-4982

4-Biphenylamine, 4'-Fluoro-
 Acetohydroxamic Acid, *N*-4-(4'-Fluorobiphenyl)-
 DNA, Binding, 78-4981
 Tritium
 Catalytic Hydrogenation, 78-4982

4-Biphenylamine, 3-Methyl-
 Tritium
 Catalytic Hydrogenation, 78-4982

4-Biphenylol, 4'-Bromo-
 Biphenyl, 4-Bromo-
 Metabolism, Rat, 78-4978

Bladder Neoplasms
 4-Biphenylamine
 Histological Study, Dog, 78-4977
 1-Butanamine, *N*-Butyl-*N*-nitroso-
 Carcinogenic Activity, Review, 78-4825
 Carcinoma
 Epidemiology, 78-5375
 Precancerous Conditions, 78-5375
 Carcinoma, Transitional Cell
 Animal Model, Rat, 78-4976
 4-Biphenylamine, 78-4977
 Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
 78-4976
 Occupational Hazard, 78-4973
 Dyes
 Occupational Hazard, 78-4973
 Epidemiology
 Geographic Factors, 78-5344
 Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
 Histological Study, Rat, 78-4976
 Neoplasms, Multiple Primary
 Occupational Hazard, 78-4973
 Retinol
 Antineoplastic Activity, Review, 78-4860
 Schistosomiasis
 Epidemiology, Review, 78-4846
 Smoking
 Epidemiology, 78-5351
 Epidemiology, Review, 78-4912
 Tryptophan
 Alanine, 3-Anthraniloyl-, 78-5338
 Alanine, 3-(3-Hydroxyanthraniloyl)-, 78-5338
 Benzoic Acid, 2-Amino-3-hydroxy-, 78-5338
 Metabolism, 78-5338
 Quinaldic Acid, 4,8-Dihydroxy-, 78-5338
 Quinaldic Acid, 4-Hydroxy-, 78-5338

Blood Platelets
 Leukemia, Myeloblastic
 Antigens, Neoplasm, 78-5397
 Precancerous Conditions, 78-5397
 Leukemia, Myelocytic
 Reverse Transcriptase, 78-5204
 Precancerous Conditions
 Reverse Transcriptase, 78-5204

Bone Marrow
 Virus, Radiation Leukemia
 Cell Differentiation, 78-5295

Bone Marrow Cells
 Benzene, (Epoxyethyl)-
 Chromosome Aberrations, 78-4953
 Cyclophosphamide
 Chromosome Aberrations, 78-5155
 Sirotidesmin A

Bone Marrow Cells (cont'd)
 Chromosome Aberrations, 78-5155
 Virus, Kirsten Murine Sarcoma
 Cell Transformation, Neoplastic, 78-5211
 Virus, Radiation Leukemia
 Precancerous Conditions, 78-5224

Bone Neoplasms
 Aniline, 4,4'-Oxydi-
 Precancerous Conditions, 78-5126
 Estrogens
 Carcinogenic Potential, Review, 78-0486
 Plutonium
 Carcinogenic Potential, Review, 78-4868
 Radiation, Ionizing
 Epidemiology, Review, 78-4874
 Radium
 Carcinogenic Potential, Review, 78-4868

Brain Neoplasms
 Astrocytoma
 Urea, 1,3-Dimethyl-1-nitroso-, 78-5096
 Cholanthrene, 3-Methyl-
 Carcinogenic Activity, Mouse, 78-4992
 Ultrastructural Study, Mouse, 78-4992
Fusarium sporotrichioides
 Toxins, 78-4954
 Glioma
 Cholanthrene, 3-Methyl-, 78-4992
 Urea, 1,3-Dimethyl-1-nitroso-, 78-5096
 Urea, Ethyl Nitroso-, 78-5186, 78-5319
 Urea, Methyl Nitroso-, 78-5281
 Kidney Transplantation
 Epidemiology, 78-5353
 Oligodendroglioma
 Urea, 1,3-Dimethyl-1-nitroso-, 78-5096
 Urea, 1,3-Dimethyl-1-nitroso-
 Ultrastructural Study, Rat, 78-5096
 Urea, Ethyl Nitroso-
 Precancerous Conditions, 78-5091
 Ultrastructural Study, Rat, 78-5091
 Wounds and Injuries, 78-5186
 Urea, Methyl Nitroso-
 Antibodies, Neoplasm, 78-5281
 Virus, Polyoma
 Clone Cells, 78-5266

Breast Neoplasms
 Age Factors
 Epidemiology, Thailand, 78-5359
 Bile Acids and Salts
 Feces, 78-5358
 Carcinoma
 Nipples, 78-5343
Clostridium
 Epidemiology, 78-5358
 Diet
 Epidemiology, 78-5367
 Epidemiology, Japan, 78-4917
 Epidemiology, Review, 78-4912
 Dietary Fats
 Epidemiology, 78-5358
 Epidemiology, Austria, 78-5366
 Dietary Proteins
 Epidemiology, Austria, 78-5366
 DNA Replication
 Insulin, 78-5392
 Placental Lactogen, 78-5392
 Precancerous Conditions, 78-5392

east Neoplasms (cont'd)

- Somatotropin, 78-5392
 - Epidemiology
 - Indonesia, 78-5345
 - Review, 78-4908
 - Estrogens
 - Carcinogenic Potential, Review, 78-0486
 - Epidemiology, Canada, 78-5362
 - Genetics
 - Epidemiology, 78-5364
 - Epidemiology, Blacks, 78-5360
 - Epidemiology, Japan, United States, 78-5368
 - Hair Dyes
 - Epidemiology, Review, 78-4859
 - Insulin
 - Precancerous Conditions, 78-5392
 - Isoantigens
 - Epidemiology, Blacks, 78-5360
 - Mammographic Classification
 - Risk Factors, 78-5369
 - Mammography
 - Epidemiology, Review, 78-4910
 - Neoplasms, Multiple Primary
 - Genetics, 78-5368
 - Nipples
 - Case Report, Autotransplantation, 78-5343
 - Obesity
 - Epidemiology, 78-5367
 - Placental Lactogen
 - Precancerous Conditions, 78-5392
 - Precancerous Conditions
 - Mammographic Classification, 78-5369
 - Pregnancy
 - Epidemiology, 78-5365
 - Radiation, Ionizing
 - Dose-Response Study, Review, 78-4871
 - Socioeconomic Factors
 - Epidemiology, 78-5351
 - 4,4'-Stilbenediol, α,α' -Diethyl-
 - Epidemiology, 78-5363
- Murkitt's Lymphoma**
- Malaria
 - Epidemiology, Review, 78-4911
 - Parasitic Diseases
 - Epidemiology, Review, 78-4913
 - Virus, Epstein-Barr
 - DNA-RNA Hybridization, 78-5241
 - Epidemiology, Review, 78-4895, 78-4911, 78-4913
 - Immune Response, Review, 78-4913
 - Seroepidemiology, Review, 78-4896
 - Virus, RD-114
 - DNA-RNA Hybridization, 78-5226

3-Butadiene, 2-Chloro-

- Chromosome Aberrations
 - Mutagenic Activity, Review, 78-4834
- Lung Neoplasms
 - Epidemiology, Review, 78-4834
- Occupational Hazard
 - Epidemiology, Review, 78-4834
- Skin Neoplasms
 - Epidemiology, Review, 78-4834

1-Butanamine, *N*-Butyl-*N*-nitroso-

- Ames Test
 - Carcinogenic Potential, Review, 78-4815
- Bladder Neoplasms
 - Carcinogenic Activity, Review, 78-4825

1-Butanamine, *N*-Butyl-*N*-nitroso- (cont'd)

- 1-Butanol, 4-(Butylnitrosamino)-
 - Carcinogenic Metabolite, Review, 78-4825
- Butyric Acid, 4-Butylamino-*N*-nitroso-
 - Carcinogenic Metabolite, Review, 78-4825
- Digestive System Neoplasms
 - Carcinogenic Activity, Review, 78-4825
- Respiratory Tract Neoplasms
 - Carcinogenic Activity, Review, 78-4825

1-Butanol, 4-(Butylnitrosamino)-

- 1-Butanamine, *N*-Butyl-*N*-nitroso-
 - Carcinogenic Metabolite, Review, 78-4825

Butyric Acid, 2-Amino-4-(ethylthio)-

- DNA Repair
 - Liver, 78-4949
- Hepatoma
 - Choline, 78-5113
 - Histological Study, Rat, 78-5113
- Protein Kinase
 - Liver, Rat, 78-5398
- RNA, Transfer, Methyltransferases
 - Liver, Rat, 78-5398

Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-

- Leukemia, Myeloblastic
 - Case Report, 78-5140, 78-5153
 - Multiple Sclerosis, 78-5153

Butyric Acid, 4-Butylamino-*N*-nitroso-

- 1-Butanamine, *N*-Butyl-*N*-nitroso-
 - Carcinogenic Metabolite, Review, 78-4825

Butyric Acid, Methylnitrosaminomethyl Ester

- Stomach Neoplasms
 - Carcinogenic Activity, Rat, 78-5065

Cadmium

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Oxidoreductases, *N*-Demethylating, 78-5118
- Carcinogenic Potential
 - Liver, Rat, 78-5118
- Hydro-Lyases
 - Enzymatic Activity, 78-5118
- Occupational Hazard
 - Epidemiology, Review, 78-4934
- Oxidoreductases
 - Enzymatic Activity, 78-5118
- Oxidoreductases, *N*-Demethylating
 - Enzymatic Activity, 78-5118
- Prostatic Neoplasms
 - Epidemiology, Review, 78-4934
- Respiratory Tract Neoplasms
 - Epidemiology, Review, 78-4934

Caffeine

- Chromatids
 - Lymphocytes, 78-5145
- DNA Repair
 - Lymphocytes, 78-5145
- Mitomycin C
 - Chromatids, 78-5145
 - DNA Repair, 78-5145
- Virus, Murine Sarcoma
 - Virus, Helper, 78-5212
- Xeroderma Pigmentosum
 - Virus, Herpes Simplex 1, 78-5168

Calcitonin

- Islet Cell Tumor

- Calcitonin (cont'd)**
Virus, Polyoma, BK, 78-5263
- Canavanine**
Virus, Rauscher Murine Leukemia
Viral Proteins, 78-5202
- Cantharidin**
Carboxy-Lyases
Enzymatic Activity, 78-5037
- Carbamic Acid, Ethyl Ester**
DNA
Alkaline Sucrose Sedimentation, 78-4935
Liver, Mouse, Rat, 78-4935
- Carbamic Acid, Hydroxy-, Ethyl Ester**
Hydroxylamine, *O,N*-Dimethyl-*N*-nitroso-
Synthesis, 78-5082
- Carbon Dioxide**
Smoking
Inhalation Study, Hamster, 78-5043
- Carbon Monoxide**
Smoking
Inhalation Study, Hamster, 78-5043
- Carbon Tetrachloride**
DNA Repair
Liver, 78-4949
Urea, Methyl Nitroso-
Hepatocarcinogenicity, 78-5084
- Carboxy-Lyases**
Acetic Acid
Enzymatic Activity, 78-5037
Cantharidin
Enzymatic Activity, 78-5037
Skin Neoplasms
12-*O*-Tetradecanoylphorbol-13-acetate, 78-5037
12-*O*-Tetradecanoylphorbol-13-acetate
Colchicine, 78-5037
Nucleotides, Cyclic, 78-5037
Retinoic Acid, 78-5037
- Carboxymethylloxysuccinate**
see Succinic Acid, Carboxymethyloxy-
- Carcinoembryonic Antigen**
Carcinogen, Chemical
Hematopoietic Stem Cells, 78-5398
Hepatoma
Carcinogen, Chemical, 78-5398
- Carcinogen, Chemical**
Aging
Immunity, Cellular, Review, 78-4906
Ames Test
Somatic Mutation, Review, 78-4809
Aroclor 1254
Ames Test, 78-4938
Aryl Hydrocarbon Hydroxylases
Genetics, Review, 78-4927
Bacteria
Mutagenic Activity, Review, 78-4811
Barbituric Acid, 5-Ethyl-5-phenyl-
Inhibitors, Review, 78-4806
Benzene, (2-Isocyanatomethyl)-
Inhibitors, Review, 78-4806
5,6-Benzoflavone
Inhibitors, Review, 78-4806
Bioassay
- Carcinogen, Chemical (cont'd)**
Carcinogenic Activity, Review, 78-4803
Toxicity, Review, 78-4803
Biogenic Amines
Cell Transformation, Neoplastic, Review, 78-4906
Carcinoembryonic Antigen
Hematopoietic Stem Cells, 78-5398
Carcinogenic Activity
Fish, Review, 78-4808
Cells, Cultured
Cell Transformation, Neoplastic, 78-4940
Toxicology, Review, 78-4802
Chlordane
Inhibitors, Review, 78-4806
Colonic Neoplasms
Bacteria, 78-5123
Bile Acids and Salts, 78-5123
Cosmetics
Mutagenic Activity, Review, 78-4859
p-Cresol, 2,6-Di-*tert*-butyl-
Inhibitors, Review, 78-4806
Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
Inhibitors, Review, 78-4806
Diet
Enzyme Activation, Review, 78-4862
Disulfide, Bis(diethylthiocarbamoyl)-
Inhibitors, Review, 78-4806
Dose-Response Study
Carcinogenic Activity, Review, 78-4813
Hartley-Sielken Model, Review, 78-4814
Drinking Water
Epidemiology, Review, 78-4847
Drosophila
Mutagenic Activity, Review, 78-4812
Epidemiology
Species Difference, Review, 78-4807
Ethanethiol, 2-Amino-, Hydrochloride
Inhibitors, Review, 78-4806
Fibrosarcoma
Macrophages, 78-5288
Flavone, 3,3',4',5,7-Pentahydroxy-
Inhibitors, Review, 78-4806
Hair Dyes
Mutagenic Activity, Review, 78-4859
Hepatoma
Carcinoembryonic Antigen, 78-5398
RNA, Transfer, Methyltransferases, 78-5398
T-Lymphocytes
Immune Response, Review, 78-4907
Microsomes, Liver
Ames Test, 78-4940
Mutagenic Activity
Host-Mediated Assay, 78-4941
Spot Test, 78-4941
Neoplasms, Experimental
Antineoplastic Agents, 78-4807
Bioassay Criteria, Review, 78-4801
Organ Culture, 78-4802
Species Difference, Review, 78-4807
Nucleic Acids
Metabolism, Review, 78-4804
Occupational Hazard
Epidemiology, Review, 78-4856
Organ Culture
Toxicology, Review, 78-4802
Organ Specificity
Lung, Liver, Rat, 78-4941

Carcinogen, Chemical (cont'd)

- Testicles, Rat, 78-4941
- Peptide Hydrolases
 - Filter Implant Test, 78-4943
- Phenol, (1,1-Dimethylethyl)-4-methoxy-
 - Inhibitors, Review, 78-4806
- Phenothiazine
 - Inhibitors, Review, 78-4806
- Polychlorobiphenyl Compounds
 - Inhibitors, Review, 78-4806
- Polycyclic Hydrocarbons
 - Inhibitors, Review, 78-4806
- Quinoline, 6-Ethoxy-1,2-Dihydro-2,2,4-trimethyl-
 - Inhibitors, Review, 78-4806
- Riboflavine
 - Ames Test, 78-5048
- Sarcoma
 - Filter Implant Test, 78-4943
- Selenium
 - Inhibitors, Review, 78-4806
- Soft Tissue Neoplasms
 - Filter Implant Test, 78-4943
- Threshold Levels
 - Hartley-Sielken Model, Review, 78-4814
- Toxicity
 - Dose-Response Study, Review, 78-4836
- Urine
 - Quantitation Method, 78-4972
- Water Pollution
 - Epidemiology, Review, 78-4847
 - Quantitation Method, 78-4972

Carcinogen, Environmental

- Adenosine Cyclic 3',5' Monophosphate
 - Concentration Levels, Review, 78-4937
- Cell Transformation, Neoplastic
 - Cell Cycle Kinetics, 78-4861
 - Fibroblasts, 78-5098
- Chromosome Aberrations
 - Induction, Review, 78-4931
- Co-carcinogenic Activity
 - Risk Evaluation, 78-4861
- Diet
 - Epidemiology, Review, 78-4925
- Epidemiology
 - Austria, 78-5350
- Genetics
 - Epidemiology, Review, 78-4926
 - Teratogenic/Mutagenic Activity, Review, 78-4930
- Lung Neoplasms
 - Epidemiology, Review, 78-4810
 - Multistate Model, Review, 78-4810
 - Mutagenicity Assay, Review, 78-4810
- Neoplasms
 - Epidemiology, Review, 78-4925
- Occupational Hazard
 - Epidemiology, Review, 78-4925
- Risk Factors
 - Epidemiology, Review, 78-4910
- Virus, Epstein-Barr
 - Co-carcinogenic Activity, Review, 78-4904

Carcinoma

- Adrenal Gland Neoplasms
 - Cushing's Syndrome, 78-5312
- Bladder Neoplasms
 - Epidemiology, 78-5375
 - Precancerous Conditions, 78-5375

Carcinoma (cont'd)

- Breast Neoplasms
 - Nipples, 78-5343
- Colonic Neoplasms
 - Polyps, 78-5313
- Digestive System Neoplasms
 - Histological Study, 78-4918
- Ear Neoplasms
 - Histological Study, 78-4918
- Esophageal Neoplasms
 - Diagnosis, Review, 78-4914
 - Epidemiology, Review, 78-4914
- Eye Neoplasms
 - Virus, Polyoma, JC, 78-5261
- Eyelid Neoplasms
 - Radiation, Ionizing, 78-5179
- Kidney Neoplasms
 - Epidemiology, 78-5375
 - Precancerous Conditions, 78-5375
 - Transplantation, Heterologous, 78-5336
- Mammary Neoplasms, Experimental
 - Urea, Ethyl Nitroso-, 78-5088
- Nasopharyngeal Neoplasms
 - Virus, Epstein-Barr, 78-5241
- Nose Neoplasms
 - Epidemiology, USSR, 78-5379
- Otorhinolaryngologic Neoplasms
 - Epidemiology, USSR, 78-5379
- Ovarian Neoplasms
 - Transplantation Immunology, 78-5293
- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-5006
 - Histological Study, 78-4918
- Sebaceous Gland Neoplasms
 - Radiation, Ionizing, 78-5179
- Stomach Neoplasms
 - Metaplasia, 78-5328
- Thyroid Neoplasms
 - Epidemiology, Iran, 78-5383
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5079
- Tracheal Neoplasms
 - Nickel Sulfide, 78-5119

Carcinoma, Basal Cell

- Ultraviolet Rays
 - Epidemiology, Review, 78-4877

Carcinoma, Bronchiolar

- Benzo(a)pyrene
 - Diol Epoxides, 78-5007
- Lung Neoplasms
 - Ultrastructural Study, 78-5327

Carcinoma, Bronchogenic

- Epidemiology
 - Immune response, 78-4916
- Ethylene, Chloro-
 - Epidemiology, 78-4832
 - Occupational Hazard, Review, 78-4832
- Sarcoidosis
 - Immune response, 78-4916
- Tuberculosis, Pulmonary
 - Immune response, 78-4916

Carcinoma, Ductal

- Pancreatic Neoplasms
 - Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081

Carcinoma, Ehrlich Tumor

- Virus-Like Particles

- Carcinoma, Ehrlich Tumor (cont'd)**
 A- and C-Type Particles, 78-5273
 Ultrastructural Study, 78-5273
- Carcinoma, Epidermoid**
 Benz(a)anthracene, 7,12-Dimethyl-
 Cells, Cultured, 78-5024
 12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
 Cervix Neoplasms
 Epidemiology, Thailand, 78-5361
 Gastrointestinal Neoplasms
 Corn Oil, 78-5120
 Dietary Fats, 78-5120
 Intestinal Neoplasms
 Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063
 Lung Neoplasms
 Adriamycin, 78-5399
 Aryl Hydrocarbon Hydroxylases, Review, 78-4853
 Guanosine, 6-Seleno-, 78-5399
 Melamine, Hexamethyl-, 78-5399
 Methotrexate, 78-5399
 Neoplasm Transplantation, 78-5399
 Purine-6-thiol, 9- β -*D*-Arabinofuranosyl-, 78-5399
 Purine-6(1*H*)-thione, 2-Amino-, 78-5399
 Smoke Condensate, 78-5045
 Uracil, 5-Fluoro-, 78-5399
 Urea, Methylcyclohexyl Nitroso-, 78-5399
 Pancreatic Neoplasms
 Epidemiology, Sri Lanka, 78-5376
 Respiratory Tract Neoplasms
 Benzo(a)pyrene, 78-5004, 78-5006
 Retinol
 Epidemiology, 78-5146
 Skin Neoplasms
 Benz(a)anthracene, 7,12-Dimethyl-, 78-5036
 Ultraviolet Rays, 78-5174
 Stomach Neoplasms
 Case Report, 78-5331
 Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101
 Quinoline, 8-Nitro-, 78-5099
 Ultrastructural Study, 78-5331
 Thyroid Neoplasms
 Immunohistochemical Study, 78-5316
 Ultrastructural Study, 78-5316
 Ultraviolet Rays
 Epidemiology, Review, 78-4877
 Immunosuppression, 78-5174
- Carcinoma In Situ**
 Cervix Neoplasms
 Contraceptive Devices, 78-5391
 Contraceptives, Oral, 78-4866
 Epidemiology, Thailand, 78-5361
 Precancerous Conditions, 78-5391
 Virus, Herpes Simplex 2, 78-4888
- Carcinoma, Mucinous**
 Esophageal Neoplasms
 Case Report, Barrett's Syndrome, 78-5322
- Carcinoma, Transitional Cell**
 Bladder Neoplasms
 Animal Model, Rat, 78-4976
 4-Biphenylamine, 78-4977
 Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
 78-4976
 Occupational Hazard, 78-4973
 Kidney Neoplasms
 Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
- Carcinoma, Transitional Cell (cont'd)**
 Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
 78-4976
- Carotid Body**
 Paraganglioma, Nonchromaffin
 Case Report, 78-5310
- Catalase**
 Mesenchymoma
 Benzo(a)pyrene, 78-5014
- Cataract**
 Nephroblastoma
 Genetics, 78-5309
- Cell Adherence**
 Histocompatibility Antigens
 Lymphocytotoxicity, 78-5282
- Cell Differentiation**
 Virus, Abelson Murine Leukemia
 Antigens, Viral, 78-5200
 Virus, Murine Leukemia
 Hematopoietic Stem Cells, 78-5211
 Virus, Radiation Leukemia
 Bone Marrow, 78-5295
 Thymopoietins, 78-5295
 Virus Variants, 78-5224
 Virus, Rauscher Murine Leukemia
 Hematopoietic Stem Cells, 78-5211
 Virus, Rous Sarcoma
 Neuroretinal Cells, 78-5194
 Transformation Defective Mutants, 78-5194
- Cell Division**
 12-*O*-Tetradecanoylphorbol-13-acetate
 Trachea, Mouse, 78-5039
 Urea, Methyl Nitroso-
 Hepatocarcinogenicity, 78-5084
 Virus, Polyoma
 Host-Range Mutants, 78-5268
 Temperature Sensitive Mutants, 78-5268
- Cell Membrane**
 Virus, Adeno 12
 Glycoproteins, 78-5234
 Virus, C-Type RNA Tumor
 Viral Proteins, 78-5210
 Virus, Murine Leukemia
 Viral Proteins, 78-5210
 Virus, Rous Sarcoma
 Proteins, 78-5195
- Cell Transformation, Neoplastic**
 Acetic Acid, Lead Salt
 Cells, Cultured, 78-5117
 Actin
 Cell Cycle Kinetics, 78-5245
 Adenosine Cyclic 3',5' Monophosphate
 Cell Adhesion, Review, 78-4932
 Aryl Hydrocarbon Hydroxylases
 Genetics, Review, 78-4927
 Benz(a)anthracene, 7,12-Dimethyl-
 Cells, Cultured, 78-5024
 Trachea, Rat, 78-5024
 Benzo(a)pyrene
 Cell Density, 78-5000
 Cells, Cultured, 78-5000
 Fetal Lung Cells, Hamster, 78-5001
 Insulin, 78-5001

Transformation, Neoplastic (cont'd)

- Methane, Sulfinylbis-, 78-5001
- Carcinogen, Chemical
 - Cells, Cultured, 78-4940
- Carcinogen, Environmental
 - Cell Cycle Kinetics, 78-4861
 - Fibroblasts, 78-5098
- Cholanthrene, 3-Methyl-
 - Fibroblasts, 78-5098
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Diploid Cells, 78-5105
 - Fibroblasts, 78-5098
- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-
 - Cells, Cultured, 78-5144
- Immunosuppression
 - Kidney Transplantation, Review, 78-4907
- Migration Inhibitory Factor
 - Isolation and Characterization, Hamster Cells 78-5393
- Polyps
 - Peutz-Jeghers Syndrome, 78-5311
- Precancerous Conditions
 - Review, 78-4924
- Proteins
 - Cell Membrane, Review, 78-4932
- Quinoline, 4-Amino-, 1-Oxide
 - Fibroblasts, 78-5098
- Quinoline, 4-Nitro-, 1-Oxide
 - Diploid Cells, 78-5105
 - Fibroblasts, 78-5098
- Radiation, Ionizing
 - Chromosome Aberrations, 78-4879
 - DNA Repair, Review, 78-4875
 - Dose-Response Study, 78-4879
 - Dose-Response Study, Fibroblasts, 78-4875
 - 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-5144
 - Mathematical Model, 78-4879
- Tubulin
 - Cell Cycle Kinetics, 78-5245
- Ultraviolet Rays
 - DNA Repair, Review, 78-4875
 - Dose-Response Study, Fibroblasts, 78-4875
- Urea, Ethyl Nitroso-
 - Rat Brain, Fetal, 78-5092
- Virus, Abelson Murine Leukemia
 - Antigenic Determinants, 78-5199
 - Antigens, Viral, 78-5200
 - Immunoprecipitation, 78-5201
 - Viral Proteins, 78-5197
- Virus, Adeno 5
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5041
 - Benzo(a)pyrene, 78-5041
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-5041
- Virus, Adeno 7
 - Acetic Acid, Lead Salt, 78-5117
- Virus, Adeno 12
 - Antigenic Determinants, 78-5238
 - Brain Cell Cultures, 78-5236
 - DNA, Viral, 78-5237
- Virus, Avian Carcinoma
 - Fibroblasts, 78-5190
 - Macrophages, 78-5190
- Virus, Avian Sarcoma
 - Genes, Viral, Review, 78-4882
- Virus, Baboon
 - Murine Sarcoma Virus Pseudotype, 78-5219

Cell Transformation, Neoplastic (cont'd)

- Virus, Bovine Leukemia
 - DNA, Viral, 78-5230
- Virus, Cytomegalo
 - Virus Replication, 78-5240
- Virus, Feline Leukemia
 - Guinea Pig, 78-5228
- Virus, Feline Sarcoma
 - Guinea Pig, 78-5228
- Virus, Kirsten Murine Sarcoma
 - Bone Marrow Cells, 78-5211
 - C-Type Particles, 78-5219
 - Spleen Cells, Gorilla, 78-5219
 - Virus, Helper, 78-5211
- Virus, Moloney Murine Sarcoma
 - Growth Factors, Isolation and Characterization 78-5220
- Virus, Murine Sarcoma
 - Cells, Cultured, 78-5218
- Virus, Polyoma
 - Actin, 78-5245
 - Antigenic Determinants, Review, 78-4892
 - Hamster Brain Cells, 78-5266
 - Neuroglia, 78-5266
 - Protein A, Review, 78-4892
 - Tubulin, 78-5245
 - Virus Activation, 78-5265
- Virus, Polyoma, BK
 - Plaque Assay, Review, 78-4893
- Virus, Radiation Leukemia
 - T-Lymphocytes, 78-5295
 - Radiation, Ionizing, 78-5295
 - Thymopoietins, 78-5295
- Virus, Rous Sarcoma
 - Cell Adhesion, Review, 78-4932
 - Genes, Viral, 78-5188
 - Glycerides, 78-5192
 - Phospholipids, 78-5192
 - Plasminogen Activators, 78-5040
 - Protein Kinase, 78-5188
 - Temperature Sensitive Mutants, 78-5195
 - Viral Proteins, 78-5193
- Virus, SV40
 - Actin, 78-5245
 - Amino Acids, 78-5259
 - Antigenic Determinants, Review, 78-4892
 - Deletion Mutants, 78-5252
 - DNA Replication, 78-5253
 - DNA, Viral, 78-5252
 - Protein A, Review, 78-4892
 - Temperature Sensitive Mutants, 78-5252
 - Tubulin, 78-5245
- Cells, Cultured**
 - Acetic Acid, Lead Salt
 - Cell Transformation, Neoplastic, 78-5117
 - Benz(a)anthracene, 7,12-Dimethyl-
 - Cell Transformation, Neoplastic, 78-5024
 - Benzo(a)pyrene
 - Cell Transformation, Neoplastic, 78-5000
 - Metabolism, 78-5020
 - Mutagenic Activity, 78-5020
 - Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Metabolism, 78-5020
 - Mutagenic Activity, 78-5020
 - Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro
 - DNA Repair, 78-5009

Cells, Cultured (cont'd)

- Carcinogen, Chemical
 - Cell Transformation, Neoplastic, 78-4940
 - Toxicology, Review, 78-4802
- Carcinoma, Epidermoid
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5024
- Fibrosarcoma
 - Acetic Acid, Lead Salt, 78-5117
- Fluorescence
 - DNA, 78-5176
- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-
 - Cell Transformation, Neoplastic, 78-5144
- Leukemia, Plasmacytic
 - Cell Cycle Kinetics, 78-5394
 - IgG, 78-5394
 - Karyotyping, 78-5394
 - Plasma Cells, 78-5394
- Lysergic Acid Diethylamide
 - DNA Replication, 78-5050
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Chromosome Abnormalities, 78-5136
 - Mitosis, 78-5136
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Metabolism, Fibroblasts, 78-5038
- Virus, Friend Murine Leukemia
 - Antigen-Antibody Reactions, 78-5283
 - Lymphocyte Transformation, 78-5283
- Virus, Murine Sarcoma
 - Cell Transformation, Neoplastic, 78-5218
 - Fibroblasts, 78-5218
 - Virus Cultivation, 78-5218

Cervix

- Nevus, Pigmented
 - Case Report, 78-5339
 - Ultrastructural Study, 78-5339

Cervix Neoplasms

- Adenocarcinoma
 - Epidemiology, Thailand, 78-5356, 78-5361
- Carcinoma, Epidermoid
 - Epidemiology, Thailand, 78-5361
- Carcinoma In Situ
 - Contraceptive Devices, 78-5391
 - Contraceptives, Oral, 78-4866
 - Epidemiology, Thailand, 78-5361
 - Precancerous Conditions, 78-5391
 - Virus, Herpes Simplex 2, 78-4888
- Contraceptive Devices
 - Epidemiology, 78-5391
- Contraceptives, Oral
 - Epidemiology, 78-5391
 - Precancerous Conditions, Review, 78-4866
- Epidemiology
 - United States, 78-5348
- Intestinal Neoplasms
 - Radiation, Ionizing, 78-5178
- Neisseria
 - Virus, Herpes, 78-5341
- Neoplasm Metastasis
 - Virus, Herpes Simplex 2, 78-4888
- Neoplasms, Multiple Primary
 - Radiation, Ionizing, 78-5178
- Precancerous Conditions
 - Dysplasia, 78-5340
 - Histological Study, 78-5340
- Smoking
 - Epidemiology, 78-5391

Cervix Neoplasms (cont'd)

- Virus, Herpes
 - Epidemiology, 78-5341
- Virus, Herpes Simplex 2
 - Animal Model, Mouse, Rat, 78-5242
 - Antigen-Antibody Reactions, Review, 78-4888
 - Antigens, Viral, 78-5242
 - Precancerous Conditions, 78-5242
 - Seroepidemiology, Review, 78-4888

Cesium Radioisotopes

- Food Contamination
 - Body Burden, 78-5181
- Radioactive Fallout
 - Body Burden, 78-5181
 - Food Contamination, 78-5181

Chloral Hydrate

- Ames Test
- Mutagenic Activity, 78-5160

Chlorambucil

- see Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-

Chloramphenicol

- Carcinogenic Potential
- Review, 78-4843

Chlordane

- Carcinogen, Chemical
- Inhibitors, Review, 78-4806

Chlordiazepoxide

- Ames Test
- Mutagenic Activity, 78-5160

Chloroform

- Benz(a)anthracene, 8,9-Dihydro-8,9-dihydroxy-7,12-dimethyl-
 - Water-Soluble Metabolite, 78-5023
- Benz(a)anthracene, 7,12-Dimethyl-
 - Metabolism, Hamster Embryo Cells, 78-5023

Chloroprene

- see 1,3-Butadiene, 2-Chloro-

Chloroquine

- see Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)

Cholangioma

- Aflatoxin B1
 - Ultrastructural Study, 78-4968
- Aflatoxin G1
 - Ultrastructural Study, 78-4968

Cholanthren-1-ol, 3-Methyl-

- Enzyme Activation
- Mutagenic Activity, 78-4995
- Microsomes, Liver
- Ames Test, 78-4995

Cholanthren-2-ol, 3-Methyl-

- Enzyme Activation
- Mutagenic Activity, 78-4995
- Microsomes, Liver
- Ames Test, 78-4995

Cholanthren-1-one, 3-Methyl-

- Enzyme Activation
- Mutagenic Activity, 78-4995
- Microsomes, Liver
- Ames Test, 78-4995

- anthren-2-one, 3-Methyl-**
Enzyme Activation
Mutagenic Activity, 78-4995
Microsomes, Liver
Ames Test, 78-4995
- anthrene, 9,10-Dihydro-3-methyl-1,9,10-trihydroxy-**
Enzyme Activation
Mutagenic Activity, 78-4995
Microsomes, Liver
Ames Test, 78-4995
- anthrene, 3-Methyl-**
Acetamide, *N*-Fluoren-3-yl-
Metabolism, Rat, Guinea Pig, 78-4985
Acetic Acid, (2,3-Dichloro-4-(2-methylenebutyryl)phenoxy)
Transferases, 78-5016
Acrylonitrile
Lactate Dehydrogenase, 78-5111
Aroclor 1254
Mutagenic Activity, 78-4995
Aryl Hydrocarbon Hydroxylases
Fish, 78-4990
Benzene, 1-Chloro-2,4-dinitro-
Transferases, 78-5016
Benzene, 1-(Chloromethyl)-4-nitro-
Transferases, 78-5016
Benzene, 1,2-Dichloro-4-nitro-
Transferases, 78-5016
Benzo(a)pyrene
Microsomes, Liver, 78-4854
Brain Neoplasms
Carcinogenic Activity, Mouse, 78-4992
Glioma, 78-4992
Ultrastructural Study, Mouse, 78-4992
Carcinogenic Potential
Diploid Cells, 78-5105
Cell Transformation, Neoplastic
Fibroblasts, 78-5098
Dibenz(a,e)aceanthrylene
DNA, Binding, 78-5033
Enzyme Activation
Mutagenic Activity, 78-4995
Fibrosarcoma
DNA, 78-5395
Hypercholesterolemia, 78-5290
Hydroxyl Derivatives
Quantitation Method, 78-4993
Hydroxylamine
Cytochrome P-450, 78-4989
Lymphoma
Antigens, Viral, 78-5291
Tumor Dormancy, 78-5291
Microsomes, Liver
Ames Test, 78-4995
Mutagenic Activity
Hamster V79 Cells, 78-4999
Pancreas
Histofluorescence, Rat, 78-4991
Precancerous Conditions, 78-4991
Propane, 1,2-Epoxy-3-(*p*-nitrophenoxy)-
Transferases, 78-5016
Sarcoma
Histocompatibility Antigens, 78-4994
Transferases
Enzymatic Activity, Rat, 78-5016
Virus, Polyoma
- Cholanthrene, 3-Methyl- (cont'd)**
Antigens, Viral, 78-4999
- Cholecystectomy**
Gallbladder Neoplasms
Epidemiology, 78-5337
- Cholelithiasis**
Gallbladder Neoplasms
Epidemiology, 78-5337
Gastrointestinal Neoplasms
Epidemiology, 78-5337
Lung Neoplasms
Epidemiology, 78-5337
Prostatic Neoplasms
Epidemiology, 78-5337
- Cholesterol**
Colonic Neoplasms
Serum Levels, 78-5061
Gastrointestinal Neoplasms
Adenocarcinoma, 78-5120
Papilloma, 78-5120
Immunosuppression
Cell Transformation, Neoplastic, Review, 78-4906
Neoplasms
Epidemiology, 78-5349
Serum Levels, 78-5349
- Choline**
Hepatoma
Butyric Acid, 2-Amino-4-(ethylthio)-, 78-5113
- Cholinesterases**
Hydrazine, 1,2-Dimethyl-
Enzyme Activation, 78-5064
Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
Enzyme Activation, 78-5064
- Chondrosarcoma**
Virus, Bovine Leukemia
DNA-DNA Hybridization, 78-5230
- Chromatids**
Caffeine
Lymphocytes, 78-5145
Cyclophosphamide
Bone Marrow, Mouse, 78-5149
Hamster V79 Cells, 78-5154
Microsomes, Liver, 78-5152
Diethylamine, *N*-Nitroso-
Bone Marrow, Mouse, 78-5149
Dimethylamine, *N*-Nitroso-
Microsomes, Liver, 78-5152
Fluorescence
Chromosome Aberrations, 78-5176
Methanesulfonic Acid, Methyl Ester
Bone Marrow, Mouse, 78-5149
Mitomycin C
Caffeine, 78-5145
p-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
Bone Marrow, Mouse, 78-5149
Uridine, 5-Bromo-2'-deoxy-
Bone Marrow, Mouse, 78-5149
Hamster V79 Cells, 78-5154
- Chromatin**
Dimethylamine, *N*-Nitroso-
Guanine, 7-Methyl-, 78-5083
Hepatoma
Fluoride, Phenylmethylsulfonyl-, 78-5054

Chromatin (cont'd)

- Virus, SV40
 - DNA, Binding, 78-5257
 - Histones, 78-5254
 - Isolation and Characterization, CV1 Cells, 78-5254
 - Nucleosomes, 78-5257

Chromium

- Neurilemmoma
 - Urea, Methyl Nitroso-, 78-5085
- Occupational Hazard
 - Epidemiology, Review, 78-4934

Chromosome Aberrations

- Anemia, Aplastic
 - Micronucleus Test, 78-4939
- Anemia, Macrocytic
 - Chromosomes, Human, 1-3, 78-5296
- Azathioprine
 - Lymphocytes, 78-5157
- Benzene, (Epoxyethyl)-
 - Bone Marrow Cells, 78-4953
 - Lymphocytes, 78-4953
- 1,3-Butadiene, 2-Chloro-
 - Mutagenic Activity, Review, 78-4834
- Carcinogen, Environmental
 - Induction, Review, 78-4931
- Cyclophosphamide
 - Bone Marrow Cells, 78-5155
 - Host-Mediated Assay, 78-5154
 - Lymphocytes, 78-5155
 - Microsomes, Liver, 78-5152
- Dimethylamine, *N*-Nitroso-
 - Microsomes, Liver, 78-5152
 - Mutagenic Activity, Review, 78-4821
- Ethylene, Chloro-
 - Lymphocytes, 78-5157
 - Mutagenic Activity, Review, 78-4834
- Fluorescence
 - Chromatids, 78-5176
 - DNA, 78-5176
- 2-Imidazoline, 2-(2,6-Dichlorophenylamino)-, Hydrochloride
 - Mutagenic Activity, Mouse, 78-5141
- Leukemia, Myeloblastic
 - Anemia, Macrocytic, 78-5296
 - Case Report, 78-5298
 - Chromosomes, Human, 6-12, 78-5296
 - Chromosomes, Human, 19-20, 78-5296
 - Precancerous Conditions, 78-5296
- Lysergic Acid Diethylamide
 - Dose-Response Study, 78-5050
 - Lymphocytes, 78-5050
- Lysergic Acid Diethylamide, Tartrate
 - Lymphocytes, 78-5050
- Methanesulfonic Acid, Ethyl Ester
 - Anemia, Aplastic, 78-4939
- Methanesulfonic Acid, Methyl Ester
 - Anemia, Aplastic, 78-4939
- Mitomycin C
 - Anemia, Aplastic, 78-4939
 - Lymphocytes, 78-5050, 78-5157
- Neoplasms
 - Hereditary Diseases, Review, 78-4931
 - Tumor Specificity, Review, 78-4931
- Phosphine Oxide, Tris(1-aziridinyl)-
 - Lymphocytes, 78-5157
- Phthalimide, *N*-((Trichloromethyl)thio)-

Chromosome Aberrations (cont'd)

- Dose-Response Study, 78-5163
- Raji, Jeff Lymphoid Cells, 78-5163
- Radiation, Ionizing
 - Anemia, Aplastic, 78-4939
 - Cell Transformation, Neoplastic, 78-4879
- Reserpine
 - Mutagenic Activity, Mouse, 78-5141
- Selenious Acid, Disodium Salt
 - Fibroblasts, 78-5115
- Sirodesmin A
 - Bone Marrow Cells, 78-5155
 - Lymphocytes, 78-5155
- Smallpox Vaccine
 - Lymphocytes, 78-5157
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Cell line, 78-5137
 - Leukocytes, 78-5137
- Thrombin
 - Lymphocytes, 78-5157
- Uridine, 5-Bromo-2'-deoxy-
 - Anemia, Aplastic, 78-4939
 - Host-Mediated Assay, 78-5154
- Vinblastine
 - Lymphocytes, 78-5157
- Vincristine
 - Lymphocytes, 78-5157
- Xeroderma Pigmentosum
 - Selenious Acid, Disodium Salt, 78-5115

Chromosome Abnormalities

- Adenomatosis, Familial Endocrine
 - Chromosomes, Human, 13-15, 78-5317
- Leukemia
 - Precancerous Conditions, 78-5204
 - Urea, 1-Butyl-1-nitroso-, 78-5095
- Leukemia, Myeloblastic
 - Cyclophosphamide, 78-5151
- Leukemia, Myelocytic
 - Colony Stimulating Factor, Review, 78-4933
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Cells, Cultured, 78-5136
 - Fibroblasts, Hamster, 78-5136
- Ultraviolet Rays
 - Dose-Response Study, NZB Mouse, 78-5175
 - Embryo Fibroblasts, Mouse, 78-5175

Chromosomes

- Benz(a)anthracene, 7,12-Dimethyl-
 - DNA, Binding, 78-5027
- Virus, Murine Mammary Tumor
 - Dexamethasone, 78-5222
- Virus, Polyoma
 - Virus Replication, 78-5256
- Virus, SV40
 - Nucleosomes, 78-5257
 - Virus Replication, 78-5256

Chromosomes, Human, 1-3

- Anemia, Macrocytic
 - Chromosome Aberrations, 78-5296

Chromosomes, Human, 6-12

- Leukemia, Myeloblastic
 - Chromosome Aberrations, 78-5296
- Virus, Baboon
 - DNA, Viral, 78-5231
 - Hybrid Cells, 78-5231
 - Virus Replication, 78-5231

Chromosomes, Human, 13-15

- Adenomatosis, Familial Endocrine
Chromosome Abnormalities, 78-5317
- Pheochromocytoma
Adenomatosis, Familial Endocrine, 78-5317
- Virus, Epstein-Barr
Chromosome Aberrations, Review, 78-4895

Chromosomes, Human, 16-18

- Virus, Adeno 2
Chromosome Aberrations, Review, 78-4902

Chromosomes, Human, 19-20

- Leukemia, Myeloblastic
Chromosome Aberrations, 78-5296

Catritz

- Fibrosarcoma
Mastectomy, 78-5180

Crulline

- Ames Test
Pyrolysis Products, 78-5122
- Mutagens
Inactivation, Fruit/Vegetable Juices, 78-5122
Pyrolysis Products, 78-5122

Clostridium

- Bile Acids and Salts
Carcinogenic Activity, Review, 78-4919
- Breast Neoplasms
Epidemiology, 78-5358
- Intestinal Neoplasms
Epidemiology, 78-5358
Epidemiology, Review, 78-4919
Steroid Nuclear Dehydrogenation, Review, 78-4919

Dal

- Phospholipids
Lung, Rat, 78-4971
- Sclerosis
Lung, Rat, 78-4971

Dealt Radioisotopes

- Virus, SV40
DNA Replication, 78-5253

Dechicine

- 12-O-Tetradecanoylphorbol-13-acetate
Carboxy-Lyases, 78-5037

Colonc Neoplasms

- Adenocarcinoma
Genetics, 78-5386
- Adenoma
Polyps, 78-5313
- Bacteria
Carcinogen, Chemical, 78-5123
- Bile Acids and Salts
Carcinogen, Chemical, 78-5123
- Carcinoma
Polyps, 78-5313
- Cholesterol
Serum Levels, 78-5061
- Diet
Epidemiology, Japan, 78-4917
Epidemiology, Review, 78-4912
- Dietary Fats
Bacteria, 78-5123
- Epidemiology
Geographic Factors, 78-5344
- Ethnic Groups

Colonc Neoplasms (cont'd)

- Epidemiology, 78-5344
- Genetics
Case Report, 78-5386
Precancerous Conditions, 78-5313
Ultrastructural Study, 78-5313
- Glycoside Hydrolases
Hexoses, 78-5060
- Heart Transplantation
Case Report, 78-5384
- Hydrazine, 1,2-Dimethyl-
Dietary Carbohydrates, 78-5061
Dietary Fats, 78-5061
Galactosyltransferases, 78-5060
Glycoproteins, Metabolism, 78-5060
Glycoside Hydrolases, 78-5060
Immune Response, Rat, 78-5061
- Hydrazine, 1,2-Dimethyl-, Dihydrochloride
Precancerous Conditions, 78-5059
- IgG
Immune Response, Rat, 78-5061
- Methane, Azoxy-
Enterectomy, 78-5159
Hyperplasia, 78-5159
- Neoplasms, Multiple Primary
Case Report, 78-5386
Genetics, 78-5371
- Polyps
Case Report, 78-5314
Genetics, 78-5313, 78-5314
- Schistosomiasis
Epidemiology, Review, 78-4846

Concanavalin A

- Hodgkin's Disease
DNA Replication, 78-5300
Lymphocytes, 78-5300

Condylomata Acuminata

- Virus, Papilloma
Epidemiology, Review, 78-4889

Contact Inhibition

- Selenious Acid, Disodium Salt
Fibroblasts, 78-5115
- Virus, SV40
Amino Acids, 78-5259

Contraceptive Devices

- Cervix Neoplasms
Carcinoma In Situ, 78-5391
Epidemiology, 78-5391

Contraceptives, Oral

- Cervix Neoplasms
Carcinoma In Situ, 78-4866
Epidemiology, 78-5391
Precancerous Conditions, Review, 78-4866
- Hepatoma
Epidemiology, Review, 78-4864
Precancerous Conditions, Review, 78-4864
Ultrastructural Study, Review, 78-4864
- Liver Neoplasms
Adenoma, 78-5130
Case Report, 78-5130, 78-5131
Epidemiology, 78-5133
Epidemiology, Review, 78-4865
Hamartoma, 78-5131
Precancerous Conditions, 78-5133
- Mammary Neoplasms, Experimental

- Contraceptives, Oral (cont'd)**
 Adenoma, 78-5132
 Histological Study, Dog, 78-5132
- Corn Oil**
 Gastrointestinal Neoplasms
 Carcinoma, Epidermoid, 78-5120
 Papilloma, 78-5120
- Coronene**
 Air Pollution
 Fossil-Fuel Combustion Products, 78-5032
- Cosmetics**
 Carcinogen, Chemical
 Mutagenic Activity, Review, 78-4859
- Coumarin, 7-Ethoxy-**
 Oxygenases
 Metabolism, Review, 78-4854
- p*-Cresol, 2,6-Di-*tert*-butyl-**
 Carcinogen, Chemical
 Inhibitors, Review, 78-4806
- m*-Cresol**
 Air Pollution
 Fossil-Fuel Combustion Products, 78-5032
- Cushing's Syndrome**
 Adrenal Gland Neoplasms
 Carcinoma, 78-5312
 Case Report, 78-5312
 Genetics, 78-5312
 Hyperplasia, 78-5312
 Genetics
 Case Report, 78-5312
- Cyanamide, Methylnitroso-**
 Esophageal Neoplasms
 Carcinogenic Activity, Rat, 78-5103
 Guanidine, Methyl-
 Nitrosated Derivatives, 78-5103
 Stomach Neoplasms
 Carcinogenic Activity, Rat, 78-5103
- Cycasin**
 Metabolic Activation
 Carcinogenic Metabolite, Review, 78-4816
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -isomer**
 Acetamide, *N*-Fluoren-2-yl-
 Hepatocarcinogenicity, Rat, 78-4983
 Carcinogen, Chemical
 Inhibitors, Review, 78-4806
 Hepatoma
 Acetamide, *N*-Fluoren-2-yl-, 78-4983
 Aflatoxin B1, 78-4970
 Carcinogenic Activity, Review, 78-4828
 Liver Neoplasms
 Adenocarcinoma, 78-4983
 Urea, Methyl Nitroso-
 Hepatocarcinogenicity, 78-5084
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, β -Isomer**
 Hepatoma
 Carcinogenic Activity, Review, 78-4828
 Toxicology
 Animals, Review, 78-4828
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer**
 Alkaline Phosphatase
 Enzymatic Activity, 78-5165
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer (cont'd)**
 Toxicity
 Dose-Response Study, Dog, 78-5165
- 4-Cyclohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-**
 Mutagenic Activity
 Translocation Test, Mouse, 78-5162
- Cyclohexene Oxide**
 Dibenz(a,e)aceanthrylene
 DNA, Binding, 78-5033
- Cyclopenta(a)phenanthren-17-one, 15,16-Dihydro-11-methyl-**
 Carcinogenic Metabolite
 Isolation and Characterization, Epidermal DNA
 78-4996
- Cyclophosphamide**
 Bone Marrow Cells
 Chromosome Aberrations, 78-5155
 Carcinogenic Potential
 Review, 78-4843
 Chromatids
 Bone Marrow, Mouse, 78-5149
 Hamster V79 Cells, 78-5154
 Chromosome Aberrations
 Host-Mediated Assay, 78-5154
 Enzyme Activation
 Mutagenic Activity, 78-5155
 Leukemia, Myeloblastic
 Behcet's Syndrome, 78-5153
 Case Report, 78-5151, 78-5153
 Chromosome Abnormalities, 78-5151
 Lymphocytes
 Chromosome Aberrations, 78-5155
 Metabolism
 Liver S-9 Fractions, 78-5152
 Microsomes, Liver
 Chromatids, 78-5152
 Chromosome Aberrations, 78-5152
 Mutagenic Activity
 Micronucleus/Nucleus Anomaly Tests, Hamster
 78-5156
- Cystadenocarcinoma**
 Ovarian Neoplasms
 Epidemiology, Dog, 78-5357
- Cystadenoma**
 Ovarian Neoplasms
 Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
 Epidemiology, Dog, 78-5357
- Cysteine**
 Acrylonitrile
 Lactate Dehydrogenase, 78-5111
 Asbestos
 Macrophages, 78-5284
- Cysts**
 Bile Duct Neoplasms
 Morpholine, *N*-Nitroso-, 78-5056
- Cytochrome P-450**
 Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
 Hydroxylamine, 78-4989
 Cholanthrene, 3-Methyl-
 Hydroxylamine, 78-4989
 Dibenz(*p*-dioxin, 2,3,7,8-Tetrachloro-
 Liver, Mouse, Review, 78-4852
 Ethane, 1-Hydroxylamino-2-phenyl-
N- and *O*-Substituted Analogs, 78-4989

chrome P-450 (cont'd)

Hydroxylamine
Microsomes, Liver, 78-4989

chrome Reductases

Acrylonitrile
Liver, Rat, 78-5111

chromes

Benzo(a)pyrene
DNA, Binding, 78-5010
Ellipticine
DNA, Binding, 78-5011
Microsomes, Liver, 78-5011

sol

Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
Binding, Review, 78-4852
Liver, Mouse, Review, 78-4852

rgents

Succinic Acid, Carboxymethyloxy-
Biodegradability, 78-5125
Carcinogenic Potential, 78-5125
Metabolism, Rat, 78-5125

amethasone

Mammary Neoplasms, Experimental
Adenocarcinoma, 78-5143
Mouse, Nude, 78-5143
Reverse Transcriptase, 78-5143
Transplantation Immunology, 78-5143
Virus, Murine Mammary Tumor, 78-5143
Virus, Murine Mammary Tumor
Chromosomes, 78-5222
DNA, Viral, 78-5222
RNA Replication, 78-5222

etes Mellitus

Hepatoma
Case Report, 78-5333
Genetics, 78-5333

epam

Ames Test
Mutagenic Activity, 78-5160

nz(a,c)anthracene

see Benzo(b)triphenylene

nz(a,e)aceanthrylene

7,8-Benzoflavone
DNA, Binding, 78-5033
Cholanthrene, 3-Methyl-
DNA, Binding, 78-5033
Cyclohexene Oxide
DNA, Binding, 78-5033
Microsomes, Liver
DNA, Binding, 78-5033
RNA, Transfer, 78-5033
Polyribonucleotides
Microsomes, Liver, 78-5033
Propane, 1,2-Epoxy-3,3,3-trichloro-
DNA, Binding, 78-5033

nz(a,h)anthracene

Ames Test
Carcinogenic Potential, Review, 78-4815
Virus, Polyoma
Antigens, Viral, 78-4999

Dibenzo(a,e)fluoranthene

see Dibenz(a,e)aceanthrylene

Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-

Aryl Hydrocarbon Hydroxylases
Liver, Mouse, Review, 78-4852
Cytochrome P-450
Liver, Mouse, Review, 78-4852
Cytosol
Binding, Review, 78-4852
Liver, Mouse, Review, 78-4852

Dibutyl Cyclic AMP

Glioma
Urea, Ethyl Nitroso-, 78-5093
Neurilemmoma
Urea, Ethyl Nitroso-, 78-5093
Urea, Ethyl Nitroso-
Transplacental Carcinogenesis, 78-5093

Dieldrin

Benzo(a)pyrene
Metabolism, Liver, 78-5008
Hydroxymethyl Derivatives
Metabolism, Rat, 78-5161
Liver
Metabolism, Rat, 78-5161
NADP
Metabolism, Rat, 78-5161

Dienestrol

4,4'-Stilbenediol, α,α' -Diethyl-
Carcinogenic Metabolite, 78-5138

Diet

Breast Neoplasms
Epidemiology, 78-5367
Epidemiology, Japan, 78-4917
Epidemiology, Review, 78-4912
Carcinogen, Chemical
Enzyme Activation, Review, 78-4862
Carcinogen, Environmental
Epidemiology, Review, 78-4925
Colonic Neoplasms
Epidemiology, Japan, 78-4917
Epidemiology, Review, 78-4912
Diethylamine, *N*-Nitroso-
Metabolism, Rat, 78-5073
Dimethylamine, *N*-Nitroso-
Metabolism, Rat, 78-5073
Gynecologic Neoplasms
Epidemiology, Review, 78-4912
Intestinal Neoplasms
Epidemiology, 78-5372
Nitric Acid
Body Burden, 78-5052
Metabolism, Intestines, 78-5052
Nitrous Acid
Body Burden, 78-5052
Metabolism, Intestines, 78-5052
Piperidine, 1-Nitroso-
Metabolism, Rat, 78-5073
Prostatic Neoplasms
Epidemiology, Review, 78-4912
Pyrrolidine, 1-Nitroso-
Metabolism, Rat, 78-5073
Quinoline, 2-Chloro-
Carcinogenic Activity, Rat, 78-5099
Quinoline, 8-Methyl-
Carcinogenic Activity, Rat, 78-5099

Diet (cont'd)

- 8-Quinolinol, 5,7-Dibromo-
Carcinogenic Activity, Rat, 78-5099
- Stomach Neoplasms
Epidemiology, Austria, 78-5350
Epidemiology, Japan, 78-4917, 78-5374
Intestinal Transit Time, 78-5374
- Uterine Neoplasms
Epidemiology, Review, 78-4862

Dietary Carbohydrates

- Colonic Neoplasms
Hydrazine, 1,2-Dimethyl-, 78-5061

Dietary Fats

- Breast Neoplasms
Epidemiology, 78-5358
Epidemiology, Austria, 78-5366
- Colonic Neoplasms
Bacteria, 78-5123
Hydrazine, 1,2-Dimethyl-, 78-5061
- Gastrointestinal Neoplasms
Carcinoma, Epidermoid, 78-5120
Hydrazine, 1,2-Dimethyl-, 78-5061
Papilloma, 78-5120
Stearic Monoglyceride, 78-5120
Strain Difference, Mouse, 78-5120
- Intestinal Neoplasms
Epidemiology, 78-5358
- Mammary Neoplasms, Experimental
Benz(a)anthracene, 7,12-Dimethyl-, 78-5025
Co-carcinogenic Effect, Rat, 78-5025

Dietary Fiber

- Benzo(a)pyrene
Aryl Hydrocarbon Hydroxylases, 78-5002
- Intestinal Neoplasms
Benzo(a)pyrene, 78-5002

Dietary Proteins

- Breast Neoplasms
Epidemiology, Austria, 78-5366

Diethylamine, 2,2'-Dichloro-*N*-methyl-

- Xeroderma Pigmentosum
Virus, Herpes Simplex 1, 78-5168

Diethylamine, *N*-Nitroso-

- Ames Test
Carcinogenic Potential, Review, 78-4815
- Bacteria
Mutagenic Activity, Review, 78-4822
- Chromatids
Bone Marrow, Mouse, 78-5149
- Diet
Metabolism, Rat, 78-5073
- Digestive System Neoplasms
Carcinogenic Activity, Review, 78-4822
- DNA Phosphotriesters
Liver, Lung, Kidney, Rat, 78-5071
- Drosophila*
Mutagenic Activity, 78-4951
- Ethanol, 2-(Ethyl nitrosamino)-
Carcinogenic Metabolite, Review, 78-4822
- Hepatoma
DNA-RNA Hybridization, 78-5069
- Intestine, Small
Metabolism, Rat, 78-5073
- Kanechlor 500
Hepatocarcinogenicity, Review, 78-4828

Diethylamine, *N*-Nitroso- (cont'd)

- Kidney Neoplasms
Carcinogenic Activity, Review, 78-4822
- Lipids
Metabolism, Rat, 78-5073
- Liver Neoplasms
Carcinogenic Activity, Review, 78-4822
- Nose Neoplasms
Carcinogenic Activity, 78-5068
Ultrastructural Study, 78-5068
- Oxidoreductases
Mutagenic Activity, 78-4951
- Respiratory Tract Neoplasms
Carcinogenic Activity, Review, 78-4822
- Ribonuclease
RNA, Liver, 78-5072
- RNA
Hepatocarcinogenesis, 78-5072

Digestive System Neoplasms

- Asbestos
Epidemiology, Review, 78-4842
- 1-Butanamine, *N*-Butyl-*N*-nitroso-
Carcinogenic Activity, Review, 78-4825
- Carcinoma
Histological Study, 78-4918
- Diethylamine, *N*-Nitroso-
Carcinogenic Activity, Review, 78-4822
- Epidemiology
Japan, 78-4917
- Ethnic Groups
Epidemiology, Hawaii, 78-5347
- Fusarium sporotrichioides*
Toxins, 78-4954
- Socioeconomic Factors
Epidemiology, 78-5351

Dimethyl-3,3'-dithiobispropionimide Dihydrochloride
see Propionamide, Dimethyl-3,3'-dithiobis-, Dihydrochloride**Dimethylamine, 1-Acetoxy-*N*-Nitroso-**

- see Acetic Acid, Methyl nitrosaminomethyl Ester

Dimethylamine, *N*-Nitroso-

- Acetone
Oxidoreductases, *N*-Demethylating, 78-5075
- Ames Test
Carcinogenic Potential, Review, 78-4815
- Aroclor 1254
Nucleic Acids, Binding, 78-5075
Oxidoreductases, *N*-Demethylating, 78-5075
- Chromatids
Microsomes, Liver, 78-5152
- Chromosome Aberrations
Mutagenic Activity, Review, 78-4821
- Diet
Metabolism, Rat, 78-5073
- DNA Phosphotriesters
Liver, Lung, Kidney, Rat, 78-5071
- Electrophiles
Carcinogenic Metabolite, Review, 78-4820
- Esophageal Neoplasms
Adrenergic Alpha Receptor Blockaders, 78-5067
Atropine, 78-5067
Proserine, 78-5067
- Guanine, 7-Methyl-
Chromatin, 78-5083
Liver, Rat, 78-5083

- thylamine, *N*-Nitroso- (cont'd)**
Hemangioendothelioma
 Histological Study, Mouse, 78-5074
Hepatoma
 Radiation, Ionizing, 78-5183
Intestine, Small
 Metabolism, Rat, 78-5073
Kidney Neoplasms
 Carcinogenic Activity, Review, 78-4821
Lipids
 Metabolism, Rat, 78-5073
Liver Neoplasms
 Adrenergic Alpha Receptor Blockaders, 78-5067
 Atropine, 78-5067
 Carcinogenic Activity, Review, 78-4821
 Hemangioendothelioma, 78-5074
 Proserine, 78-5067
Metabolism
 Liver S-9 Fractions, 78-5152
Microsomes, Liver
 Chromosome Aberrations, 78-5152
 Metabolism, Mouse, 78-5075
Neoplasms, Experimental
 Carcinogenic Activity, Review, 78-4821
Nitrous Acid
 Nitrosation Kinetics, 78-5077
 Perchloric Acid, 78-5077
Nitrous Acid, Sodium Salt
 Antipyrine, 4-(Dimethylamino)-, 78-5076
 Ethanol, 2-(Dimethylamino)-, 78-5076
 Ethylamine, 2-(Diphenylmethoxy)-*N,N*-dimethyl-
 78-5076
 Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-, 78-5076
 Sulfanilamide, *N*-(3,4-Dimethyl-4-isoxazolyl)-
 78-5076
 Tetracycline, 78-5076
Nucleic Acids
 Methylation, Review, 78-4820
Oxidoreductases, *N*-Demethylating
 Dose-Response Study, 78-5075
Pregn-4-ene-3,20-dione, 17-(Acetyloxy)-6 α -methyl-
 Metabolism, Kidney, 78-5078
Respiratory Tract Neoplasms
 Carcinogenic Activity, Review, 78-4821
Saccharomyces cerevisiae
 Mutagenic Activity, 78-4945
Sex Hormones
 Metabolism, Kidney, 78-5078
Testosterone
 Metabolism, Kidney, 78-5078
Urinary Tract Infections
Proteus mirabilis, 78-5076
 Tetracycline, 78-5076
ethylsulfate
 see Sulfuric Acid, Dimethyl Ester
enylamine, 4-Isothiocyanato-4'-nitro-
Antischistosomal Drugs
 Mutagenic Activity, Review, 78-4845
enylamine, *N*-Nitroso-
Antibody Formation
 Immune Response, Rat, 78-5086
Antigens, Bacterial
 Antibody Formation, 78-5086
Diphenylhydramine
 see Ethylamine, 2-(Diphenylmethoxy)-*N,N*-dimethyl-
Dipropylamine, 2,2'-Dioxo-*N*-nitroso-
Pancreatic Neoplasms
 Adenocarcinoma, 78-5081
 Adenoma, 78-5081
 Carcinoma, Ductal, 78-5081
 Dose-Response Study, Hamster, 78-5081
 Neoplasms, Multiple Primary, 78-5081
Dipropylamine, *N*-Nitroso-
 Aniline, 2,6-Dinitro-*N,N*-dipropyl-*p*-isopropyl-
 Pesticides, Review, 78-4824
Esophageal Neoplasms
 Carcinogenic Activity, Review, 78-4824
Kidney Neoplasms
 Carcinogenic Activity, Review, 78-4824
Liver Neoplasms
 Carcinogenic Activity, Review, 78-4824
 Propylamine, *N*-(2-Hydroxypropyl)-*N*-nitroso-
 Carcinogenic Metabolite, Review, 78-4824
Respiratory Tract Neoplasms
 Carcinogenic Activity, Review, 78-4824
p-Toluidine, α,α,α -Trifluoro-2,6-dinitro-*N,N*-dipropyl-
 Pesticides, Review, 78-4824
Disgerminoma
 Virus, Polyoma, BK
 Carcinogenic Activity, Hamster, 78-5263
Disulfide, Bis(diethylthiocarbamoyl)-
 Carcinogen, Chemical
 Inhibitors, Review, 78-4806
DNA
 Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 Antibody Formation, 78-4986
Aflatoxin B1
 Binding, Liver, 78-4962
 Carcinogenic Metabolite, 78-4962
Benz(a)anthracene, 7,12-Dimethyl-
 Carcinogenic Metabolite, 78-5026
Benzo(a)pyrene
 Binding, 78-4998
 Diol Epoxides, 78-5007
 Photodynamic Effects, 78-5003
Carbamic Acid, Ethyl Ester
 Alkaline Sucrose Sedimentation, 78-4935
 Liver, Mouse, Rat, 78-4935
DNA, Viral
 Endonucleases, 78-5396
 Nucleotide Sequence, 78-5396
Ethane, 2-Bromo-2-chloro-1,1,1-trifluoro-
 Mutagenic Activity, 78-5160
Ethylene, Trichloro-
 Mutagenic Activity, 78-5160
Fibrosarcoma
 Cholanthrene, 3-Methyl-, 78-5395
 Flow Microfluorometry, 78-5395
Fluorescence
 Alkaline Elution Assay, 78-5176
 Cells, Cultured, 78-5176
 Chromosome Aberrations, 78-5176
Nucleotide Sequence
 Genes, Monkey, 78-5396
2-Oxetanone
 Phosphodiester, 78-5108
Radiation, Ionizing
 Mutagenic Activity, 78-4879

DNA, Neoplasm

- Leukemia
 - Virus, C-Type RNA Tumor, 78-5274
- Sarcoma
 - Virus, C-Type RNA Tumor, 78-5274
- Virus, C-Type RNA Tumor
 - DNA, Viral, 78-5274

DNA Polymerase

- Virus, Avian Myeloblastosis
 - DNA Replication, 78-5217
- Virus, Moloney Murine Leukemia
 - DNA Replication, 78-5217
- Virus, Moloney Murine Sarcoma
 - DNA Replication, 78-5217
- Virus, RD-114
 - DNA, Viral, 78-5226
- Virus, SV40
 - Nucleoprotein Complex, 78-5246, 78-5247

DNA Repair

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Endonucleases, 78-5169
 - Fibroblasts, 78-5169
 - Ultraviolet Rays, 78-5169
 - Xeroderma Pigmentosum, 78-5169
- Acetamide, Thio-
 - Liver, 78-4949
- 2-Anthracenamide
 - Xeroderma Pigmentosum, 78-5170
- Aryl Hydrocarbon Hydroxylases
 - Genetics, Review, 78-4927
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro
 - Cells, Cultured, 78-5009
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Liver, 78-4949
- Caffeine
 - Lymphocytes, 78-5145
- Carbon Tetrachloride
 - Liver, 78-4949
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Escherichia coli*, 78-5102
 - Neurospora crassa*, 78-4948
- Lung Neoplasms
 - Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro, 78-5009
- Methanesulfonic Acid, Ethyl Ester
 - Escherichia coli*, 78-5102
- Methanesulfonic Acid, Methyl Ester
 - Escherichia coli*, 78-5102
- Mitomycin C
 - Caffeine, 78-5145
 - Lymphocytes, 78-5145
- Mycotoxins
 - Liver, 78-4949
- Quinoline, 4-Nitro, 1-Oxide
 - Neurospora crassa*, 78-4948
 - UV-Mimetic Compounds, 78-5170
 - Xeroderma Pigmentosum, 78-5170
- Retinoblastoma
 - Genetics, 78-5182
 - Radiation, Ionizing, 78-5182
- Retrovirus
 - Liver, 78-4949
- Selenious Acid, Disodium Salt
 - Fibroblasts, 78-5115
- Ultraviolet Rays

DNA Repair (cont'd)

- Endonucleases, 78-5169
- Escherichia coli*, 78-5171
- Fibroblasts, 78-5169
- Monkey CV-1 Cells, 78-5172
- Nitrous Acid, 78-5171
- Pyrimidine Dimers, 78-5172
- Xeroderma Pigmentosum, 78-5169
- Xeroderma Pigmentosum
 - Selenious Acid, Disodium Salt, 78-5115
- Xeroderma Pigmentosum
 - Fibroblasts, 78-5170

DNA Replication

- Breast Neoplasms
 - Insulin, 78-5392
 - Placental Lactogen, 78-5392
 - Precancerous Conditions, 78-5392
 - Somatotropin, 78-5392
- Ethidium Bromide
 - Double Intercalating Analog, 78-5114
- Hodgkin's Disease
 - Concanavalin A, 78-5300
- Hydrazine, 1,2-Dimethyl-
 - Colon, Rat, 78-5064
 - Physostigmine, Sulfate, 78-5064
- Leukemia L1210
 - Ethidium Bromide, 78-5114
 - Ethidium Chloride, 78-5114
- Lysergic Acid Diethylamide
 - Cells, Cultured, 78-5050
- Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
 - Colon, Rat, 78-5064
 - Physostigmine, Sulfate, 78-5064
- Virus, Adeno 2
 - Benzimidazole, 5,6-Dichloro-1- β -*D*-ribofuranosyl-78-5233
- Virus, Avian Myeloblastosis
 - DNA Polymerase, 78-5217
 - DNA, Viral, 78-5217
- Virus, Moloney Murine Leukemia
 - DNA Polymerase, 78-5217
 - DNA, Viral, 78-5217
- Virus, Moloney Murine Sarcoma
 - DNA Polymerase, 78-5217
 - DNA, Viral, 78-5217
- Virus, Polyoma
 - Cell Cycle Kinetics, 78-5268
 - Host-Range Mutants, 78-5268
 - RNA, Viral, 78-5267
 - Temperature Sensitive Mutants, 78-5265
- Virus, SV40
 - Antigens, Viral, 78-5248, 78-5253
 - Cell Transformation, Neoplastic, 78-5253
 - Cobalt Radioisotopes, 78-5253
 - DNA-RNA Hybridization, 78-5253
 - Mitomycin C, 78-5253
 - Radiation, Ionizing, 78-5253
 - Temperature Sensitive Mutants, 78-5248

DNA, Viral

- DNA
 - Endonucleases, 78-5396
 - Nucleotide Sequence, 78-5396
- Leukemia
 - Virus, C-Type RNA Tumor, 78-5274
- Sarcoma
 - Virus, C-Type RNA Tumor, 78-5274

NA, Viral (cont'd)

- Virus, Adeno 2
 - Carcinogenic Activity, Review, 78-4902
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5237
 - DNA-DNA Hybridization, 78-5237
 - Endonucleases, 78-5237
 - Hamster Cell Lines, 78-5237
- Virus, Avian Myeloblastosis
 - DNA Replication, 78-5217
- Virus, Baboon
 - Chromosomes, Human, 6-12, 78-5231
- Virus, Bovine Leukemia
 - Cell Transformation, Neoplastic, 78-5230
 - DNA-DNA Hybridization, 78-5230
- Virus, C-Type RNA Tumor
 - DNA, Neoplasm, 78-5274
- Virus, Herpes Simplex 1
 - Temperature Sensitive Mutants, 78-5244
 - Virus, Recombinant, 78-5244
 - Virus Rescue, 78-5244
- Virus, Herpes Simplex 2
 - Virus Rescue, 78-5244
- Virus, Moloney Murine Leukemia
 - DNA Replication, 78-5217
- Virus, Moloney Murine Sarcoma
 - DNA Replication, 78-5217
- Virus, Murine Mammary Tumor
 - Dexamethasone, 78-5222
 - HTC Clone Cells, 78-5222
- Virus, Polyoma
 - Complementation Group A Mutants, 78-4892
 - DNA-RNA Hybridization, 78-5267
 - Endonucleases, 78-5265
 - Nucleosomes, 78-5256
- Virus, Polyoma, BK
 - Isolation and Characterization, 78-5262
 - Isolation and Characterization, Review, 78-4893
- Virus, Radiation Leukemia
 - DNA-DNA Hybridization, Spleen Cells, 78-5225
- Virus, RD-114
 - DNA Polymerase, 78-5226
 - DNA-RNA Hybridization, 78-5226
 - Isolation and Characterization, 78-5226
- Virus, RNA Tumor
 - Virus, Avian Myeloblastosis, 78-5271
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-5252
 - Complementation Group A Mutants, 78-4892
 - Nucleoprotein Complex, 78-5246, 78-5247
 - Nucleoproteins, 78-5254
 - Nucleosomes, 78-5256
 - Nucleotide Sequence, 78-5249, 78-5396
 - RNA Polymerase, 78-5250
 - RNA Replication, 78-5250, 78-5260
 - Virus Replication, 78-5250
- isophila**
 - Carcinogen, Chemical
 - Mutagenic Activity, Review, 78-4812
 - Diethylamine, *N*-Nitroso-
 - Mutagenic Activity, 78-4951
 - Mutagens
 - Test Evaluation, 78-4812
 - Polycyclic Hydrocarbons
 - Mutagenic Activity, Review, 78-4812

Drug Therapy

- Hodgkin's Disease
 - Epilepsy, 78-5301
- Imidazole-1-ethanol, 2-Methyl-5-nitro-
 - Carcinogenic Potential, Review, 78-4844
- Isonicotinic Acid Hydrazide
 - Carcinogenic Potential, Review, 78-4844
- Leukemia, Lymphoblastic
 - Alanine, 3-(*p*-(Bis(2-chloroethyl)amino)phenyl)-
78-5302
- Leukemia, Myeloblastic
 - Arthritis, Rheumatoid, 78-5151
 - Case Report, 78-5298
 - Multiple Sclerosis, 78-5140
 - Neoplasms, Multiple Primary, 78-5298
 - Sarcoma, Reticulum Cell, 78-5299
- Lymphoma
 - Epilepsy, 78-5301
- Multiple Myeloma
 - Epilepsy, 78-5301
 - Leukemia, Lymphocytic, 78-5303

Drugs

- Carcinogenic Potential
 - Epidemiology, 78-5352
- Neoplasms
 - Epidemiology, 78-5352

Dyes

- Adrenal Gland Neoplasms
 - Neoplasms, Multiple Primary, 78-4973
- Bile Duct Neoplasms
 - Adenocarcinoma, 78-4973
- Bladder Neoplasms
 - Occupational Hazard, 78-4973
- Hepatoma
 - Neoplasms, Multiple Primary, 78-4973
- Lung Neoplasms
 - Adenocarcinoma, 78-4973

Dysgammaglobulinemia

- Virus, Gross Murine Leukemia
 - IgA, 78-5280
 - Mouse, AKR, 78-5280

Ear Neoplasms

- Carcinoma
 - Histological Study, 78-4918

Ellipticine

- Ames Test
 - Hydroxy Derivative, 78-5011
- Aroclor 1254
 - Aryl Hydrocarbon Hydroxylases, 78-5011
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Aryl Hydrocarbon Hydroxylases, 78-5011
- Benzo(a)pyrene
 - Ames Test, 78-5011
 - Aryl Hydrocarbon Hydroxylases, 78-5011
 - DNA, Binding, 78-5011
- Cytochromes
 - DNA, Binding, 78-5011
- Microsomes, Liver
 - Aryl Hydrocarbon Hydroxylases, 78-5011
 - Cytochromes, 78-5011

Endonucleases

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - DNA Repair, 78-5169
- DNA

- Endonucleases (cont'd)**
 DNA, Viral, 78-5396
 Ultraviolet Rays
 DNA Repair, 78-5169
 Virus, Adeno 12
 DNA, Viral, 78-5237
 Virus, Polyoma
 DNA, Viral, 78-5265
- Environmental Hazard**
 Polychlorobiphenyl Compounds
 Epidemiology, Review, 78-4829
- Ependymoma**
 Eye Neoplasms
 Virus, Polyoma, JC, 78-5261
 Virus, Polyoma, BK
 Antigen-Antibody Reactions, 78-5263
 Carcinogenic Activity, Hamster, 78-5263
 Neoplasm Transplantation, 78-5263
- Epilepsy**
 Hodgkin's Disease
 Drug Therapy, 78-5301
 Lymphoma
 Drug Therapy, 78-5301
 Multiple Myeloma
 Drug Therapy, 78-5301
- Erythroleukemia**
 4,4'-Stilbenediol, α,α' -Diethyl-
 Mitosis, 78-5136
- Erythropoietin**
 Kidney Neoplasms
 Neoplasm Metastasis, 78-5336
 Transplantation, Heterologous, 78-5336
- Escherichia coli***
 Benzo(a)pyrene
 Radiation, 78-5003
 Temperature Sensitive Mutants, 78-5003
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 DNA Repair, 78-5102
 Mutagenic Activity, 78-5102
 Methanesulfonic Acid, Ethyl Ester
 DNA Repair, 78-5102
 Mutagenic Activity, 78-5102
 Methanesulfonic Acid, Methyl Ester
 DNA Repair, 78-5102
 Mutagenic Activity, 78-4950, 78-5102
 Nitrous Acid
 Mutagenic Activity, 78-5171
 Ultraviolet Rays
 DNA Repair, 78-5171
 Mutagenic Activity, 78-5171
- Esophageal Neoplasms**
 Adenocarcinoma
 Case Report, Barrett's Syndrome, 78-5322
 Areca
 Epidemiology, Pakistan, 78-5382
 Epidemiology, Sri Lanka, 78-5381
 Carcinoma
 Diagnosis, Review, 78-4914
 Epidemiology, Review, 78-4914
 Carcinoma, Mucinous
 Case Report, Barrett's Syndrome, 78-5322
 Cyanamide, Methylnitroso-
 Carcinogenic Activity, Rat, 78-5103
 Dimethylamine, *N*-Nitroso-
- Esophageal Neoplasms (cont'd)**
 Adrenergic Alpha Receptor Blockaders, 78-5067
 Atropine, 78-5067
 Proserine, 78-5067
 Dipropylamine, *N*-Nitroso-
 Carcinogenic Activity, Review, 78-4824
 Epidemiology
 Review, 78-4908
 Epithelium, Columnar
 Ultrastructural Study, 78-5322
 Geographical Factors
 Epidemiology, 78-4915
 Iron Deficiency
 Epidemiology, Sri Lanka, 78-5381
 Metaplasia
 Precancerous Conditions, 78-5323
 Opium
 Epidemiology, 78-5049
 Precancerous Conditions
 Esophageal Reflux, 78-5323
 Smoking
 Epidemiology, 78-5049
 Epidemiology, Pakistan, 78-5382
 Tobacco
 Chewing, 78-5382
- Esterases**
 Letterer-Siwe Disease
 Enzymatic Activity, 78-5389
- Estradiol**
 Receptors, Hormone
 Mammary/Uterine Tissue, Mouse, 78-5139
 4,4'-Stilbenediol, α,α' -Diethyl-
 Receptors, Hormone, 78-5139
- Estradiol, 3-Benzate**
 Hepatoma
 Morpholine, *N*-Nitroso-, 78-5055
 Liver Neoplasms
 Morpholine, *N*-Nitroso-, 78-5055
- Estradiol, 17-Hydrocinnamate**
 Hepatoma
 Morpholine, *N*-Nitroso-, 78-5055
 Liver Neoplasms
 Morpholine, *N*-Nitroso-, 78-5055
- Estrogens**
 Bone Neoplasms
 Carcinogenic Potential, Review, 78-0486
 Breast Neoplasms
 Carcinogenic Potential, Review, 78-0486
 Epidemiology, Canada, 78-5362
 Carcinogenic Potential
 Review, 78-4843
 Gynecologic Neoplasms
 Carcinogenic Potential, Review, 78-0486
 Kidney Neoplasms
 Growth Factors, 78-5134
 Mitosis, 78-5134
 Mammary Neoplasms, Experimental
 Growth Factors, 78-5134
 Mitosis, 78-5134
 Prolactin, 78-5134
 Thyroid Hormones, 78-5134
 Mesothelioma
 Carcinogenic Potential, Review, 78-0486
 Pituitary Neoplasms
 Growth Factors, 78-5134

- gens (cont'd)**
 Mitosis, 78-5134
 Thyroid Hormones, 78-5134
- Urine**
 Quantitation Method, 78-4972
- Urogenital Neoplasms**
 Carcinogenic Potential, Review, 78-0486
- Uterine Neoplasms**
 Epidemiology, Canada, 78-5355
 Epidemiology, Review, 78-4862, 78-4867, 78-4910
 Menopause, 78-5355
 Turner's Syndrome, 78-5342
- Water Pollution**
 Quantitation Method, 78-4972
- Acetic Acid**
 see Acetic Acid, (2,3-Dichloro-4-(2-methylenebutyryl)phenoxy)
- Acetic Acid, 2-Bromo-2-chloro-1,1,1-trifluoro-**
 Ames Test
 Mutagenic Activity, 78-5160
- DNA**
 Mutagenic Activity, 78-5160
- Acetic Acid, 1-Hydroxylamino-2-phenyl-**
 Cytochrome P-450
N- and *O*-Substituted Analogs, 78-4989
- Ethanol, 2-Amino-, Hydrochloride**
 Carcinogen, Chemical
 Inhibitors, Review, 78-4806
- Ethanol, 2-(Dimethylamino)-**
 Dimethylamine, *N*-Nitroso-
 Nitrous Acid, Sodium Salt, 78-5076
- Ethanol, 2-(Ethylnitrosamino)-**
 Diethylamine, *N*-Nitroso-
 Carcinogenic Metabolite, Review, 78-4822
- Ethanol, *N*-Nitrosoiminodi-**
 Cutting Fluids
 Quantitation Method, 78-5070
- Fibrosarcoma**
 Carcinogenic Potential, Review, 78-4823
- Hepatoma**
 Carcinogenic Potential, Review, 78-4823
- Kidney Neoplasms**
 Adenoma, 78-4823
 Carcinogenic Potential, Review, 78-4823
- Nitrous Acid**
 Cutting Fluids, 78-5070
- Nose Neoplasms**
 Adenocarcinoma, 78-4823
 Carcinogenic Potential, Review, 78-4823
- Bis(2-chloroethyl)-**
 Ames Test
 Mutagenic Activity, Review, 78-4811
- 1-Chloro-2,2,2-trifluoroethyl Difluoromethyl**
 Ames Test
 Mutagenic Activity, 78-5160
- 2,2-Dichloro-1,1-difluoroethyl Methyl**
 Ames Test
 Mutagenic Activity, 78-5160
- Ethidium Bromide**
 DNA, Binding
 Double Intercalating Analog, 78-5114
 Spectrum Analysis, 78-5114
- Ethidium Chloride**
 DNA, Binding
 Spectrum Analysis, 78-5114
 Leukemia L1210
 DNA Replication, 78-5114
- Ethionine**
 see Butyric Acid, 2-Amino-4-(ethylthio)-
 RNA, Transfer, Methyltransferases
 Liver, Rat, 78-5398
- Ethyl Alcohol**
 Pharyngeal Neoplasms
 Epidemiology, 78-5344
- Ethylamine, 2-(Diphenylmethoxy)-*N,N*-dimethyl-**
 Dimethylamine, *N*-Nitroso-
 Nitrous Acid, Sodium Salt, 78-5076
- Ethylamine Hydrochloride**
 Acetic Acid, Ethylnitrosaminoethyl Ester
 Synthesis, 78-5066
- Ethylamine, *N*-Methyl-*N*-nitroso-**
 Acetamide, *N*-Methyl-
 Nitrous Acid, Sodium Salt, 78-5080
 Isocyanic Acid, Ethyl Ester
 Nitrous Acid, Sodium Salt, 78-5080
 Nitrous Acid, Sodium Salt
 Deuterium Labeled Analogs, Synthesis, 78-5080
- Ethylene, Bromo- Polymer**
 Liposarcoma
 Carcinogenic Potential, Rat, 78-5167
- Ethylene, Chloro-**
 Ames Test
 Mutagenic Activity, Review, 78-4811
 Angiosarcoma
 Epidemiology, 78-4832
 Carcinoma, Bronchogenic
 Epidemiology, 78-4832
 Occupational Hazard, Review, 78-4832
 Chromosome Aberrations
 Mutagenic Activity, Review, 78-4834
 Dose-Response Study
 Pharmacokinetics, Review, 78-4833
 Glutathione
 Liver, Rat, 78-5110
 Leukemia
 Epidemiology, 78-4832
 Occupational Hazard, Review, 78-4832
 Liver Neoplasms
 Epidemiology, 78-4832
 Occupational Hazard, Review, 78-4832
 Precancerous Conditions, Review, 78-4922
 Lymphocytes
 Chromosome Aberrations, 78-5157
 Metabolism
 Pharmacokinetics, Review, 78-4833
 Species Difference, Review, 78-4831
 Metabolism, Liver
 Dose-Response Study, 78-5110
 Neoplasms, Experimental
 Age Factors, Review, 78-4831

Ethylene, Chloro- (cont'd)

- Carcinogenic Activity, Review, 78-4831
- Precancerous Conditions, Review, 78-4922
- Species Difference, Review, 78-4831

Nervous System Neoplasms

- Epidemiology, 78-4832
- Occupational Hazard, Review, 78-4832

Nucleic Acids

- Liver, Rat, 78-5110

Occupational Hazard

- Epidemiology, Review, 78-4834

Respiratory Tract Neoplasms

- Epidemiology, 78-4832

Structure-Activity Relationship

- Ames Test, Review, 78-4830
- Carcinogenic/Mutagenic Activity, Review, 78-4830

Ethylene, 1,1-Dichloro-**Age Factors**

- Hepatotoxicity, Rat, 78-5109
- Sex Factors, 78-5109

Metabolism

- Species Difference, Review, 78-4831

Neoplasms, Experimental

- Age Factors, Review, 78-4831
- Carcinogenic Activity, Review, 78-4831
- Species Difference, Review, 78-4831

Structure-Activity Relationship

- Ames Test, Review, 78-4830
- Carcinogenic/Mutagenic Activity, Review, 78-4830

Ethylene, 1,2-Dichloro-**Structure-Activity Relationship**

- Ames Test, Review, 78-4830
- Carcinogenic/Mutagenic Activity, Review, 78-4830

Ethylene, Tetrachloro-**Structure-Activity Relationship**

- Ames Test, Review, 78-4830
- Carcinogenic/Mutagenic Activity, Review, 78-4830

Ethylene Thiourea

see 2-Imidazolidinethione

Ethylene, Trichloro-**DNA**

- Mutagenic Activity, 78-5160

Ethynone

- Mammary Neoplasms, Experimental
- Hyperplasia, 78-5132

Eye Neoplasms**Carcinoma**

- Virus, Polyoma, JC, 78-5261

Ependymoma

- Virus, Polyoma, JC, 78-5261

Fibrosarcoma

- Virus, Polyoma, JC, 78-5261

Neurilemmoma

- Virus, Polyoma, JC, 78-5261

Retinoblastoma

- Virus, Polyoma, JC, 78-5261
- Virus, Polyoma, JC
- Carcinogenic Activity, Hamster, 78-5261

Eyelid Neoplasms**Carcinoma**

- Radiation, Ionizing, 78-5179

Radiation, Ionizing

- Case Report, 78-5179

Eyelid Neoplasms (cont'd)

- Ultrastructural Study, 78-5179

Fatty Acids**Immunosuppression**

- Cell Transformation, Neoplastic, Review, 78-4906

Ferric Chloride**Benzo(a)pyrene**

- Co-carcinogenic Activity, Hamster, 78-5004
- Respiratory Tract Neoplasms
- Benzo(a)pyrene, 78-5004

Fiber Glass**Neoplasms, Experimental**

- Trans-Species Extrapolation, 78-4805

Fibrin**Hepatoma**

- Neoplasm Circulating Cells, 78-5304

Fibroblasts

- Acetamide, *N*-(Acetyloxy)-*N*-fluorenyl-2-yl-
- DNA Repair, 78-5169

Carcinogen, Environmental

- Cell Transformation, Neoplastic, 78-5098

Cholanthrene, 3-Methyl-

- Cell Transformation, Neoplastic, 78-5098

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Cell Transformation, Neoplastic, 78-5098

Quinoline, 4-Amino-, 1-Oxide

- Cell Transformation, Neoplastic, 78-5098

Quinoline, 4-Nitro-, 1-Oxide

- Cell Transformation, Neoplastic, 78-5098

Selenious Acid, Disodium Salt

- Chromosome Aberrations, 78-5115

Contact Inhibition, 78-5115**DNA Repair, 78-5115****Ultraviolet Rays**

- DNA Repair, 78-5169

Virus, Avian Carcinoma

- Cell Transformation, Neoplastic, 78-5190

Virus, Murine Sarcoma

- Cells, Cultured, 78-5218

Virus, SV40

- Anchorage Independent Growth, 78-5252

Xeroderma Pigmentosum

- DNA Repair, 78-5170

Fibrosarcoma**Acetic Acid, Lead Salt**

- Cells, Cultured, 78-5117

Carcinogen, Chemical

- Macrophages, 78-5288

Cholanthrene, 3-Methyl-

- DNA, 78-5395
- Hypercholesteremia, 78-5290

Cicatrix

- Mastectomy, 78-5180

DNA

- Flow Microfluorometry, 78-5395

Ethanol, *N*-Nitrosoiminodi-

- Carcinogenic Potential, Review, 78-4823

Eye Neoplasms

- Virus, Polyoma, JC, 78-5261

Hypercholesteremia

- Transplantation Immunology, 78-5290

Lung

- Neoplasm Metastasis, 78-5388

Macrophages

Sarcoma (cont'd)

- Growth, 78-5288
- Winn Test, 78-5288
- Radiation, Ionizing
 - Case Report, 78-5180
 - Mastectomy, 78-5180
- Skin Neoplasms
 - Ultraviolet Rays, 78-5174
- Stomach Neoplasms
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101
- Ultraviolet Rays
 - Hypersensitivity, Delayed, 78-5173
 - Immunity, Passive, 78-5285
 - Immunosuppression, 78-5285
 - Transplantation Immunology, 78-5173, 78-5285
- Urea, Methyl Nitroso-
 - Carcinogenic Activity, Xiphophorus Fish, 78-5087
 - Genetics, Fish, 78-5087

Urea, 3,3',4',5,7-Pentahydroxy-

- Carcinogen, Chemical
- Inhibitors, Review, 78-4806

Urea-2-amine

- Tritium
 - Catalytic Hydrogenation, 78-4982

Urea, 2-Nitro-

- Salmonella typhimurium*
 - Mutagenic Activity, 78-4946

Urease

- Chromatids
 - Chromosome Aberrations, 78-5176

Urea

- Alkaline Elution Assay, 78-5176
- Cells, Cultured, 78-5176
- Chromosome Aberrations, 78-5176

Urea, Phenylmethylsulfonyl-

- Hepatoma
 - Chromatin, 78-5054
 - Chromosomal Proteins, Non-Histone, 78-5054

Urea Phthalimide, N-((Trichloromethyl)thio)-

- Additives
 - Toxicity
 - Dose-Response Study, Review, 78-4836

Contamination

- Aflatoxin B1
 - Carcinogenic Metabolite, 78-4964
 - Quantitation Method, 78-4958, 78-4959, 78-4965
 - Quantitation Method, Review, 78-4838
 - Rain, 78-4961
- Aflatoxin B2
 - Quantitation Method, 78-4959, 78-4965
- Aflatoxin G1
 - Quantitation Method, 78-4959, 78-4965
 - Quantitation Method, Review, 78-4838
- Aflatoxin G2
 - Quantitation Method, 78-4959, 78-4965
- Aflatoxin M1
 - Quantitation Method, 78-4964
 - Quantitation Method, Review, 78-4838
 - Review, 78-4837
- Cesium Radioisotopes
 - Body Burden, 78-5181
- Guanidine, 1-Methyl-3-nitro-1-nitroso-

Food Contamination (cont'd)

- Nitrosated Derivatives, 78-5103
- Intestinal Neoplasms
 - Mycotoxins, 78-5372
- Mycotoxins
 - Ames Test, 78-5048
- Nitric Acid
 - Quantitation Method, 78-5051
- Nitrosamines
 - Carcinogenic Activity, Review, 78-4819
 - Carcinogenic Potential, Review, 78-4817
- Nitrous Acid
 - Carcinogenic Potential, Review, 78-4817
 - Quantitation Method, 78-5051
- Ochratoxin A
 - Quantitation Method, 78-4959
- Proline, 4-Hydroxy-1-nitroso-
 - Quantitation Method, 78-5057
- Proline, 1-Nitroso-
 - Quantitation Method, 78-5057
- Pyrrolidine, 3-Hydroxy-1-nitroso-
 - Quantitation Method, 78-5058
- Radioactive Fallout
 - Cesium Radioisotopes, 78-5181
- Sarcosine, N-Nitroso-
 - Quantitation Method, 78-5057
- Sorbic Acid
 - Nitrous Acid, Sodium Salt, 78-5124

Formaldehyde

- Virus, Feline Leukemia
 - Viral Vaccines, Review, 78-4884

Formamide, N-(4-(5-Nitro-2-furyl)-2-thiazolyl)-

- Bladder Neoplasms
 - Carcinoma, Transitional Cell, 78-4976
 - Histological Study, Rat, 78-4976
- Kidney Neoplasms
 - Carcinoma, Transitional Cell, 78-4976

Furanacrylamide, N-Isopropyl-5-Nitro-

- Antischistosomal Drugs
 - Mutagenic Activity, Review, 78-4845

Fusarium sporotrichioides

- Brain Neoplasms
 - Toxins, 78-4954
- Digestive System Neoplasms
 - Toxins, 78-4954
- Mammary Neoplasms, Experimental
 - Toxins, 78-4954
- Pituitary Neoplasms
 - Toxins, 78-4954

Galactosyltransferases

- Colonic Neoplasms
 - Hydrazine, 1,2-Dimethyl-, 78-5060

Gallbladder Neoplasms

- Cholecystectomy
 - Epidemiology, 78-5337
- Cholelithiasis
 - Epidemiology, 78-5337

Gastrectomy

- Stomach Neoplasms
 - Epidemiology, 78-5329

Gastric Juice

- Anemia, Pernicious
 - Nitrosamines, 78-5330

Gastric Juice (cont'd)

- Nitrous Acid, 78-5330
- Guanidine, Methyl-
 - Ames Test, 78-5103
- Stomach Neoplasms
 - Nitrosation, Review, 78-4818

Gastric Mucosa

- Metaplasia
 - Precancerous Conditions, 78-5328

Gastritis

- Nitroso Compounds
 - Nitrosation, Review, 78-4818
- Stomach Neoplasms
 - Anacidity, 78-5373
 - Epidemiology, 78-5373
 - Epidemiology, Review, 78-4818

Gastrointestinal Neoplasms

- Adenocarcinoma
 - Cholesterol, 78-5120
- Asbestos
 - Epidemiology, Review, 78-4841, 78-4842
- Carcinoma, Epidermoid
 - Corn Oil, 78-5120
 - Dietary Fats, 78-5120
- Cholelithiasis
 - Epidemiology, 78-5337
- Dietary Fats
 - Stearic Monoglyceride, 78-5120
 - Strain Difference, Mouse, 78-5120
- Epidemiology
 - Review, 78-4908
- Hydrazine, 1,2-Dimethyl-
 - Dietary Fats, 78-5061
- Papilloma
 - Cholesterol, 78-5120
 - Corn Oil, 78-5120
 - Dietary Fats, 78-5120
 - Linoleic Acid, 78-5120
 - Monoolein, 78-5120
 - 9-Octadecenoic Acid, 78-5120
- Virus, Adeno 12
 - Carcinogenic Potential, 78-5235
 - DNA-DNA Hybridization, 78-5235

Genetics

- Adrenal Gland Neoplasms
 - Cushing's Syndrome, 78-5312
 - Pheochromocytoma, 78-5308
- Ataxia Telangiectasia
 - Epidemiology, Review, 78-4926
- Breast Neoplasms
 - Epidemiology, 78-5364
 - Epidemiology, Blacks, 78-5360
 - Epidemiology, Japan, United States, 78-5368
 - Neoplasms, Multiple Primary, 78-5368
- Carcinogen, Environmental
 - Epidemiology, Review, 78-4926
 - Teratogenic/Mutagenic Activity, Review, 78-4930
- Colonic Neoplasms
 - Adenocarcinoma, 78-5386
 - Case Report, 78-5386
 - Neoplasms, Multiple Primary, 78-5371
 - Polyps, 78-5313, 78-5314
 - Precancerous Conditions, 78-5313
 - Ultrastructural Study, 78-5313
- Cushing's Syndrome

Genetics (cont'd)

- Case Report, 78-5312
- Gynecologic Neoplasms
 - Case Report, 78-5386
- Hepatoma
 - Case Report, 78-5333
 - Diabetes Mellitus, 78-5333
 - Ovarian Cysts, 78-5333
- Hodgkin's Disease
 - Immune Response, Review, 78-4913
 - Immunity, Cellular, 78-5300
- Lymphoma
 - Immune Response, Review, 78-4913
- Melanoma
 - Epidemiology, Review, 78-4876
- Neoplasms
 - Epidemiology, Review, 78-4925
- Neoplasms, Multiple Primary
 - Case Report, 78-5371
- Nephroblastoma
 - Aniridia, 78-5309
 - Cataract, 78-5309
- Ovarian Neoplasms
 - Epidemiology, 78-5364
- Paraganglioma, Nonchromaffin
 - Case Report, 78-5310
 - Neoplasm Recurrence, 78-5310
- Pheochromocytoma
 - Angiomatosis, 78-5308
 - Case Report, 78-5308
 - Neoplastic Endocrine-Like Symptoms, 78-5308
- Respiratory Tract Neoplasms
 - Aryl Hydrocarbon Hydroxylases, 78-5324
- Retinoblastoma
 - DNA Repair, 78-5182
 - Epidemiology, Review, 78-4926
 - Radiation Effects, 78-5182
- Uterine Neoplasms
 - Neoplasms, Multiple Primary, 78-5371
- Virus, Epstein-Barr
 - Epidemiology, Review, 78-4904
- Virus, Rous Sarcoma
 - Antibody Formation, 78-5292
 - Graft Rejection, 78-5292
 - Immune Response, 78-5292
 - Immunity, Cellular, 78-5292
 - Immunosuppression, 78-5292
 - Stress, 78-5292
- Xeroderma Pigmentosum
 - Epidemiology, Review, 78-4926

Giant Cell Tumors

- Lung Neoplasms
 - Siderosis, 78-5326
 - Silver, 78-5326
 - Titanium Oxide, 78-5326

Glioblastoma Multiforme

- Urea, Ethyl Nitroso-
 - Carcinogenic Activity, Review, 78-4827
 - Ultrastructural Study, 78-5089

Glioma

- Benzo(a)pyrene
 - Sex Hormones, 78-5135
- Brain Neoplasms
 - Cholanthrene, 3-Methyl-, 78-4992
 - Urea, 1,3-Dimethyl-1-nitroso-, 78-5096
 - Urea, Ethyl Nitroso-, 78-5186, 78-5319

- oma (cont'd)**
 Urea, Methyl Nitroso-, 78-5281
- Sex Hormones**
 Age Factors, 78-5135
 Epidemiology, 78-5135
- Subependymal Plate**
 Ultrastructural Study, Rat, 78-5319
- Urea, Ethyl Nitroso-**
 Carcinogenic Activity, Review, 78-4827
 Dibutyryl Cyclic AMP, 78-5093
 Transplacental Carcinogenesis, 78-5093
 Ultrastructural Study, Rat, 78-5091, 78-5319
- Urea, Methyl Nitroso-**
 Antibodies, Neoplasm, 78-5281
- sarcoma**
 see Glioma
- tagon**
 Islet Cell Tumor
 Virus, Polyoma, BK, 78-5263
- osephosphatase**
 Acrylonitrile
 Liver, Rat, 78-5111
- osephosphate Dehydrogenase**
 Leukemia, Lymphocytic
 A and B Isoenzymes, 78-5297
 B-Lymphocytes, 78-5297
 T-Lymphocytes, 78-5297
- uronidase**
 Benz(a)anthracene, 7,12-Dimethyl-
 Metabolism, Hamster Embryo Cells, 78-5023
 Phenol Glucuronides, 78-5023
- amyl Transpeptidase**
 Acetamide, N-Fluorenyl-2-yl-
 Liver, Rat, 78-5084
- Acrylonitrile**
 Enzymatic Activity, 78-5111
- Urea, Methyl Nitroso-**
 Hepatectomy, 78-5084
 Liver, Rat, 78-5084
- thione**
 Ethylene, Chloro-
 Liver, Rat, 78-5110
- 2-Imidazolidinethione**
 Metabolism, 78-5112
- Isonicotinamide, 2-Ethylthio-**
 Metabolism, 78-5112
- Urea, Thio-**
 Metabolism, 78-5112
- thione Peroxidase**
 Mesenchymoma
 Benzo(a)pyrene, 78-5014
- thione Reductase**
 Mesenchymoma
 Benzo(a)pyrene, 78-5014
- rides**
 Virus, Rous Sarcoma
 Cell Transformation, Neoplastic, 78-5192
 Transformation Defective Mutants, 78-5192
- rol**
 Skin Neoplasms
 Precancerous Conditions, 78-5047
 Smoke Condensate, 78-5047
- Glycoproteins**
 Leukemia, Myeloblastic
 Precancerous Conditions, 78-5397
- Myelofibrosis**
 Precancerous Conditions, 78-5397
- Myeloid Metaplasia**
 Precancerous Conditions, 78-5397
- Virus, Adeno 12**
 Cell Membrane, 78-5234
 Cytocidal Mutants, 78-5234
- Glycoside Hydrolases**
 Colonic Neoplasms
 Hexoses, 78-5060
 Hydrazine, 1,2-Dimethyl-, 78-5060
- Goiter**
 Thyroid Neoplasms
 Epidemiology, Iran, 78-5383
- Gonadal Dysgenesis**
 see Turner's Syndrome
- Graft Rejection**
 Histocompatibility Antigens
 Mouse, Neonate, 78-5282
- Plasmacytoma**
 Myeloma Proteins, 78-5289
 Thymus Gland, 78-5289
- Virus, Rous Sarcoma**
 Genetics, 78-5292
- Granulosa Cell Tumor**
 Jaw Neoplasms
 Case Report, 78-5321
- Ovarian Neoplasms**
 Epidemiology, Dog, 78-5357
- Pregnancy**
 Case Report, 78-5321
- Griseofulvin**
 Hepatoma
 Dose-Response Study, Mouse, 78-5142
- Thyroid Neoplasms**
 Dose-Response Study, Rat, 78-5142
- Growth**
 Fibrosarcoma
 Macrophages, 78-5288
- Sarcoma 180, Crocker**
 T-Lymphocytes, 78-5279
- 12-O-Tetradecanoylphorbol-13-acetate**
 Trachea, Mouse, 78-5039
- Guanidine, Methyl-**
 Cyanamide, Methylnitroso-
 Nitrosated Derivatives, 78-5103
- Gastric Juice**
 Ames Test, 78-5103
- Nitrous Acid**
 Ames Test, 78-5103
- Guanidine, 1-Methyl-3-nitro-1-nitroso-**
 Ames Test
 Nitrosated Derivatives, 78-5103
- Carcinogenic Activity**
 Nitrosated Derivatives, 78-5103
- Cell Transformation, Neoplastic**
 Diploid Cells, 78-5105
 Fibroblasts, 78-5098
- DNA Repair**
Escherichia coli, 78-5102

Guanidine, 1-Methyl-3-nitro-1-nitroso- (cont'd)

- Escherichia coli*
 - Mutagenic Activity, 78-5102
- Food Contamination
 - Nitrosated Derivatives, 78-5103
- Intestinal Neoplasms
 - Animal Model, Mouse, 78-5104
 - Sarcoma, Reticulum Cell, 78-5104
- Laryngeal Neoplasms
 - Papilloma, 78-5101
- Neurospora crassa*
 - DNA Repair, 78-4948
 - Lethal/Mutagenic Activity, 78-4947
 - Mutagenic Activity, 78-4948
- Nitrous Acid
 - Saliva, 78-5103
- Stomach Neoplasms
 - Carcinogenic Potential, Hamster, 78-5101
 - Carcinoma, Epidermoid, 78-5101
 - Fibrosarcoma, 78-5101
 - Neoplasm Metastasis, 78-5101
 - Papilloma, 78-5101

Guanidine, Nitroso-

- Salmonella typhimurium*
 - Mutagenic Activity, 78-4946

Guanine, 7-Methyl-

- Dimethylamine, *N*-Nitroso-
 - Chromatin, 78-5083
 - Liver, Rat, 78-5083

Guanosine, 2'-Deoxy-

- DNA, Alkylation
 - Radioimmunoassay, 78-5106

Guanosine, 7-Methyl-

- Acridine, 9-Amino-
 - Fluorescence Quenching, 78-5107

Guanosine Monophosphate

- Acridine, 9-Amino-
 - Fluorescence Quenching, 78-5107

Guanosine, 6-Seleno-

- Lung Neoplasms
 - Carcinoma, Epidermoid, 78-5399

Gynecologic Neoplasms

- Adenocarcinoma
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-5370
- Diet
 - Epidemiology, Review, 78-4912
- Epidemiology
 - Sabah, Malaysia, 78-5346
- Estrogens
 - Carcinogenic Potential, Review, 78-0486
- Genetics
 - Case Report, 78-5386
- Leiomyomatosis Peritonealis Disseminata
 - Case Report, 78-5306
 - Ultrastructural Study, 78-5306
- Neoplasms, Connective Tissue
 - Radiation, Ionizing, 78-5178
- Neoplasms, Multiple Primary
 - Radiation, Ionizing, 78-5178
- Oncogenic Viruses
 - Epidemiology, Review, 78-4895
- Progesterone
 - Neoplasm Regression, Review, 78-4863
- Sarcoma

Gynecologic Neoplasms (cont'd)

- Radiation, Ionizing, 78-5178
- Schistosomiasis
 - Epidemiology, Review, 78-4846
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Carcinogenic Potential, Review, 78-0486
 - Epidemiology, 78-5370
 - Transplacental Carcinogenesis, 78-5370

Hair Dyes

- Breast Neoplasms
 - Epidemiology, Review, 78-4859
- Carcinogen, Chemical
 - Mutagenic Activity, Review, 78-4859

Halothane

- see Ethane, 2-Bromo-2-chloro-1,1,1-trifluoro-

Hamartoma

- Liver Neoplasms
 - Contraceptives, Oral, 78-5131

Heart Neoplasms

- Virus, Polyoma, JC
 - Carcinogenic Activity, Hamster, 78-5261

HeLa Cells

- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Mitosis, 78-5136
- Virus, Mason-Pfizer Monkey
 - Virus Replication, 78-5232

Hemangioendothelioma

- Dimethylamine, *N*-Nitroso-
 - Histological Study, Mouse, 78-5074
- Liver Neoplasms
 - Dimethylamine, *N*-Nitroso-, 78-5074
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063
 - Quinoline, 78-5099
 - Symphytum officinale*, 78-4942
- Quinoline
 - Carcinogenic Activity, Rat, 78-5099
- Soft Tissue Neoplasms
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063
 - Symphytum officinale*
 - Histological Study, Rat, 78-4942

Hemangioma

- Liver Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-5006

Hemangiopericytoma

- Lung
 - Neoplasm Metastasis, 78-5388

Hematopoietic Stem Cells

- Anesthetics
 - Immune Response, Review, 78-4858
- Carcinogen, Chemical
 - Carcinoembryonic Antigen, 78-5398
- Virus, C-Type RNA Tumor
 - Carcinogenic Activity, Review, 78-4901
- Virus, Murine Leukemia
 - Cell Differentiation, 78-5211
- Virus, Rauscher Murine Leukemia
 - Cell Differentiation, 78-5211

Hepatitis

- Anesthetics
 - Epidemiology, Review, 78-4857

toma
 Acetamide, *N*-Fluoren-2-yl-
 Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -isomer
 78-4983
 Actin
 Neoplasm Metastasis, 78-5054
 Aflatoxin B1
 Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
 78-4970
 Ultrastructural Study, 78-4968
 Aflatoxin G1
 Ultrastructural Study, 78-4968
 Aroclor 1254
 Carcinogenic Activity, Review, 78-4828
 Aroclor 1260
 Carcinogenic Activity, Review, 78-4828
 Australia Antigen
 Cytoplasm, 78-5276
 Epidemiology, Review, 78-4846
 Liver Cirrhosis, 78-5276
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 Precancerous Conditions, 78-5127
 Proton Magnetic Relaxation Times, 78-5127
 Butyric Acid, 2-Amino-4-(ethylthio)-
 Choline, 78-5113
 Histological Study, Rat, 78-5113
 Carcinogen, Chemical
 Carcinoembryonic Antigen, 78-5398
 RNA, Transfer, Methyltransferases, 78-5398
 Contraceptives, Oral
 Epidemiology, Review, 78-4864
 Precancerous Conditions, Review, 78-4864
 Ultrastructural Study, Review, 78-4864
 Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
 Carcinogenic Activity, Review, 78-4828
 Cyclohexane, 1,2,3,4,5,6-Hexachloro-, β -Isomer
 Carcinogenic Activity, Review, 78-4828
 Diabetes Mellitus
 Case Report, 78-5333
 Genetics, 78-5333
 Diethylamine, *N*-Nitroso-
 DNA-RNA Hybridization, 78-5069
 Ethanol, *N*-Nitrosoiminodi-
 Carcinogenic Potential, Review, 78-4823
 Fluoride, Phenylmethylsulfonyl-
 Chromatin, 78-5054
 Chromosomal Proteins, Non-Histone, 78-5054
 Genetics
 Case Report, 78-5333
 Irioseofulvin
 Dose-Response Study, Mouse, 78-5142
 Isonitrochlor 500
 Carcinogenic Activity, Review, 78-4828
 Isonitrochlor C
 Benzene, Hexachloro-, 78-4979
 Histological Study, Mouse, 78-4979
 Precancerous Conditions, 78-4979
 Liver Cirrhosis
 Epidemiology, Indonesia, 78-5332
 Mestranol
 Epidemiology, Review, 78-4864
 Nitrophenol, *N*-Nitroso-
 Carcinogenic Potential, Fish, 78-5056
 Chromosomal Proteins, Non-Histone, 78-5054
 Estradiol, 3-Benzoyl, 78-5055
 Estradiol, 17-Hydrocinnamate, 78-5055
 Neoplasm Metastasis, 78-5054

Hepatoma (cont'd)

 Ultrastructural Study, Rat, 78-5055
 Mycotoxins
 Co-carcinogenic Activity, Review, 78-4840
 Epidemiology, Review, 78-4839, 78-4840, 78-4862
 Food Contamination, Review, 78-4839
 Neoplasm Circulating Cells
 Fibrin, 78-5304
 Lung, Rat, 78-5304
 Platelet Aggregation, 78-5304
 Thromboplastin, 78-5304
 Neoplasm Metastasis
 Chromosomal Proteins, Non-Histone, 78-5054
 Lung, Mouse, 78-5334
 Ultrastructural Study, Rat, 78-5304
 Neoplasm Transplantation
 RNA, Neoplasm, 78-5069
 Neoplasms, Multiple Primary
 Dyes, 78-4973
 Occupational Hazard, 78-4973
 Ovarian Cysts
 Genetics, 78-5333
 Precancerous Conditions
 Histological Study, Mouse, 78-5334
 1-Propanol, 2,3-Dibromo-, Phosphate
 Carcinogenic Potential, Rat, 78-5167
 Quinoline
 Carcinogenic Activity, Rat, 78-5099
 Radiation, Ionizing
 Dimethylamine, *N*-Nitroso-, 78-5183
 Liver Regeneration, 78-5183
 Species Difference, 78-5183
 RNA, Neoplasm
 DNA-RNA Hybridization, 78-5069
 Schistosomiasis
 Epidemiology, Review, 78-4846
 Steroids
 Precancerous Conditions, Review, 78-4922
Symphitum officinale
 Carcinogenic Activity, Roots, Leaves, 78-4942
 Histological Study, Rat, 78-4942
 Thromboplastin
 Neoplasm Metastasis, 78-5304
 Transplantation Immunology
 Age Factors, Mouse, 78-5293
 Thymectomy, 78-5293
 Tubulin
 Neoplasm Metastasis, 78-5054
 Virus, Hepatitis
 Epidemiology, Review, 78-4839

Hereditary Diseases

Adenocarcinoma
 Review, 78-4929
 Neoplasms
 Teratogenic/Mutagenic Activity, Review, 78-4930
 Nephroblastoma
 Review, 78-4929
 Neuroblastoma
 Review, 78-4929
 Neurofibromatosis
 Review, 78-4929
 Pheochromocytoma
 Review, 78-4929
 Polyps
 Review, 78-4929
 Retinoblastoma
 Epidemiology, Review, 78-4930

Hereditary Diseases (cont'd)

- Review, 78-4929
- Sarcoma, Osteogenic
- Epidemiology, Review, 78-4930

Hexoses

- Colonic Neoplasms
- Glycoside Hydrolases, 78-5060

Histiocytes

- Letterer-Siwe Disease
- Ultrastructural Study, 78-5389

Histiocytoma

- Soft Tissue Neoplasms
- Clinopathologic Study, 78-5307
- Neoplasm Metastasis, 78-5307
- Ultrastructural Study, 78-5307

Histocompatibility Antigens

- Cell Adherence
- Lymphocytotoxicity, 78-5282
- Graft Rejection
- Mouse, Neonate, 78-5282
- Hodgkin's Disease
- Immune Response, Review, 78-4913
- Lymphoma
- Radiation, Ionizing, 78-5294
- Transplantation Immunology, 78-5294
- Pheochromocytoma
- Adenomatosis, Familial Endocrine, 78-5317
- Sarcoma
- Cholanthrene, 3-Methyl-, 78-4994
- Immunogenicity, 78-4994
- Transplantation Immunology
- Mouse, Neonate, 78-5282
- Virus, Polyoma, BK
- Isolation and Characterization, Review, 78-4891
- Virus, Polyoma, JC
- Isolation and Characterization, Review, 78-4891

Histones

- Virus, SV40
- Chromatin, 78-5254
- Nucleoprotein Complex, 78-5246

Hodgkin's Disease

- Concanavalin A
- DNA Replication, 78-5300
- Lymphocytes, 78-5300
- Epilepsy
- Drug Therapy, 78-5301
- Genetics
- Immune Response, Review, 78-4913
- Immunity, Cellular, 78-5300
- Histocompatibility Antigens
- Immune Response, Review, 78-4913
- Hydantoin, 5,5-Diphenyl-
- Case Report, 78-5301
- Parasitic Diseases
- Epidemiology, Review, 78-4913
- Virus, Epstein-Barr
- DNA-RNA Hybridization, 78-5241
- Epidemiology, Review, 78-4913

Homeostasis

- Aging
- Cell Transformation, Neoplastic, Review, 78-4906

Hybrid Cells

- Virus, Baboon

Hybrid Cells (cont'd)

- Antigens, Viral, 78-5231
- Chromosomes, Human, 6-12, 78-5231
- Reverse Transcriptase, 78-5231
- Virus Replication, 78-5231

Hycanthone

- Antischistosomal Drugs
- Mutagenic Activity, Review, 78-4845

Hydantoin, 5,5-Diphenyl-

- Carcinogenic Potential
- Review, 78-4843
- Hodgkin's Disease
- Case Report, 78-5301
- Lymphoma
- Case Report, 78-5301
- Multiple Myeloma
- Case Report, 78-5301

Hydrazine, 1,2-Dimethyl-

- Cholinesterases
- Enzyme Activation, 78-5064
- Colonic Neoplasms
- Dietary Carbohydrates, 78-5061
- Dietary Fats, 78-5061
- Galactosyltransferases, 78-5060
- Glycoproteins, Metabolism, 78-5060
- Glycoside Hydrolases, 78-5060
- Immune Response, Rat, 78-5061
- DNA Replication
- Colon, Rat, 78-5064
- Gastrointestinal Neoplasms
- Dietary Fats, 78-5061
- Physostigmine, Sulfate
- DNA Replication, 78-5064

Hydrazine, 1,2-Dimethyl-, Dihydrochloride

- Administration Route
- Carcinogenic Activity, Mouse, 78-5063
- Colonic Neoplasms
- Precancerous Conditions, 78-5059
- Intestinal Neoplasms
- Adenocarcinoma, 78-5063
- Adenoma, 78-5063
- Anastomosis, 78-5062
- Carcinoma, Epidermoid, 78-5063
- Precancerous Conditions, 78-5059
- Liver Neoplasms
- Hemangioendothelioma, 78-5063
- Soft Tissue Neoplasms
- Hemangioendothelioma, 78-5063

Hydro-Lyases

- Benzene, (Epoxyethyl)-
- Liver, Rat, 78-5013
- Benzo(a)pyrene
- Carcinogenic Activity, Review, 78-4850
- Enzyme Activation, Review, 78-4854
- Benzo(a)pyrene 4,5-Oxide
- Liver, Rat, 78-5013
- Cadmium
- Enzymatic Activity, 78-5118

Hydrochloric Acid

- Acetonitrile, 2-(*N*-Methyl-*N*-nitroso)amino-
- Reaction Product, 78-5053
- Propionitrile, 3-(*N*-Methyl-*N*-nitroso)amino-
- Reaction Product, 78-5053

droxy-3-fluorenylacetamide
 see Acetohydroxamic Acid, *N*-Fluoren-3-yl-
 oxylamine
 Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
 Cytochrome P-450, 78-4989
 Cholanthrene, 3-Methyl-
 Cytochrome P-450, 78-4989
 Microsomes, Liver
 Cytochrome P-450, 78-4989
 Enzymatic Activity, 78-4989
 Hydroxylases, 78-4989
 Oxidoreductases, 78-4989
 oxylamine, *O,N*-Dimethyl-*N*-nitroso-
 Carbamic Acid, Hydroxy-, Ethyl Ester
 Synthesis, 78-5082
 Sulfuric Acid, Dimethyl Ester
 Synthesis, 78-5082
 oxylases
 Hydroxylamine
 Microsomes, Liver, 78-4989
 droxyurethane
 see Carbamic Acid, Hydroxy-, Ethyl Ester
 cholesteremia
 Fibrosarcoma
 Cholanthrene, 3-Methyl-, 78-5290
 Transplantation Immunology, 78-5290
 Virus, Coxsackie B
 Antibody Formation, 78-5290
 Immune Response, 78-5290
 Macrophages, 78-5290
 rplasia
 Adrenal Gland Neoplasms
 Cushing's Syndrome, 78-5312
 Aniline, 4,4'-Oxydi-
 Dose-Response Study, 78-5126
 Pituitary Gland, 78-5126
 Thyroid Gland, 78-5126
 Colonic Neoplasms
 Methane, Azoxy-, 78-5159
 Mammary Neoplasms, Experimental
 Anagestone Acetate, 78-5132
 Ethinerone, 78-5132
 Mestranol, 78-5132
 Norgestrel, Chloroethynyl-, 78-5132
 Prostatic Neoplasms
 5 α -Andostan-3-one, 17 β -Hydroxy-, 78-4920
 Androgens, 78-4920
 Androstanediol, 78-4920
 Testosterone, 78-4920
 Uterine Neoplasms
 Precancerous Conditions, Review, 78-4921
 rsensitivity, Delayed
 Fibrosarcoma
 Ultraviolet Rays, 78-5173
 Ultraviolet Rays
 Benzene, 1-Chloro-2,4-dinitro-, 78-5173
 rthyroidism
 Aniline, 4,4'-Oxydi-
 Dose-Response Study, 78-5126
 Goitrogenic Effect, Mouse, Rat, 78-5126
 physectomy
 Aflatoxin B1
 Enzyme Activation, 78-4960

Hypophysectomy (cont'd)

Hepatotoxicity, Rat, 78-4960
 Nucleic Acids, 78-4960
 RNA Polymerase, 78-4960

Hypoxanthine Phosphoribosyltransferase

Methanesulfonic Acid, Ethyl Ester
 Mutagenic Activity, 78-5158
 Methanesulfonic Acid, Isopropyl Ester
 Mutagenic Activity, 78-5158
 Methanesulfonic Acid, Methyl Ester
 Mutagenic Activity, 78-5158
 Sulfuric Acid, Diethyl Ester
 Mutagenic Activity, 78-5158
 Sulfuric Acid, Dimethyl Ester
 Mutagenic Activity, 78-5158

IgA

Autoimmune Diseases
 Mouse, AKR, 78-5280
 Virus, Gross Murine Leukemia
 Dysgammaglobulinemia, 78-5280

IgG

Colonic Neoplasms
 Immune Response, Rat, 78-5061
 Letterer-Siwe Disease
 Immunoglobulins, Fc, 78-5389
 Leukemia, Plasmacytic
 Cells, Cultured, 78-5394

IgM

Virus, Epstein-Barr
 Anti-Antibodies, 78-5278

1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-

Cells, Cultured
 Cell Transformation, Neoplastic, 78-5144
 Radiation, Ionizing
 Cell Transformation, Neoplastic, 78-5144
 Co-carcinogenic Activity, 78-5144

Imidazole-1-ethanol, 2-Methyl-5-nitro-

Drug Therapy
 Carcinogenic Potential, Review, 78-4844
 Lung Neoplasms
 Adenoma, 78-4844
 Carcinogenic Potential, Review, 78-4844

2-Imidazolidinethione

Glutathione
 Metabolism, 78-5112
 Oxidoreductases
 Metabolism, 78-5112

2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-

Antischistosomal Drugs
 Mutagenic Activity, Review, 78-4845

2-Imidazoline, 2-(2,6-Dichlorophenylamino)-, Hydrochloride

Chromosome Aberrations
 Mutagenic Activity, Mouse, 78-5141

Immune Serums

Virus, Polyoma
 Virus Activation, 78-5265
 Virus, SV40
 Antigens, Viral, 78-5251

Immunity, Cellular

Hodgkin's Disease
 Genetics, 78-5300
 Virus, C-Type RNA Tumor

Immunity, Cellular (cont'd)

- Viral Proteins, 78-5272
- Virus, Herpes Simplex 2
 - BCG, 78-5243
 - Plant Agglutinins, 78-5243
 - Skin Graft, Footpad Tests, 78-5243
 - Tuberculin, 78-5243
- Virus, Rous Sarcoma
 - Genetics, 78-5292
 - Radiation, Ionizing, 78-5292

Immunity, Passive

- Fibrosarcoma
- Ultraviolet Rays, 78-5285

Immunoglobulins, Fc

- Letterer-Siwe Disease
- IgG, 78-5389

Immunoglobulins, Heavy Chain

- Myeloma Proteins
 - Amino Acids, 78-5277
 - Insulin, Binding, 78-5277

Immunoglobulins, Light Chain

- Leukemia, Lymphoblastic
- Multiple Myeloma, 78-5302
- Multiple Myeloma
 - Leukemia, Lymphocytic, 78-5303

Immunoglobulins, Surface

- Leukemia, Lymphocytic
- B-Lymphocytes, 78-5297
- Lymphoma
 - Virus, Epstein-Barr, 78-5278

Immunosuppression

- Aging
 - Cell Transformation, Neoplastic, Review, 78-4906
- Carcinoma, Epidermoid
 - Ultraviolet Rays, 78-5174
- Cell Transformation, Neoplastic
 - Kidney Transplantation, Review, 78-4907
- Cholesterol
 - Cell Transformation, Neoplastic, Review, 78-4906
- Fatty Acids
 - Cell Transformation, Neoplastic, Review, 78-4906
- Fibrosarcoma
 - Ultraviolet Rays, 78-5285
- Insulin
 - Cell Transformation, Neoplastic, Review, 78-4906
- Lipoproteins
 - Cell Transformation, Neoplastic, Review, 78-4906
- Sarcoma, Reticulum Cell
 - Kidney Transplantation, 78-5353
- Skin Neoplasms
 - Ultraviolet Rays, 78-5174
- Ultraviolet Rays
 - Lymphocytes, 78-5285
 - T-Lymphocytes, 78-5173
- Virus, Adeno 2
 - Carcinogenic Activity, Review, 78-4902
- Virus, C-Type RNA Tumor
 - T-Lymphocytes, 78-5272
- Virus, Friend Murine Leukemia
 - Antibody Formation, 78-5207
 - Lipopolysaccharides, 78-5207
- Virus, Marek's Disease Herpes
 - T-Lymphocytes, 78-5196
 - Precancerous Conditions, 78-5196

Immunosuppression (cont'd)

- Virus, Polyoma, BK
 - Isolation and Characterization, Review, 78-4890
- Virus, Rous Sarcoma
 - Genetics, 78-5292
- Virus, Turkey Herpes
 - T-Lymphocytes, 78-5196

Immunosuppressive Agents

- Leukemia, Myeloblastic
- Arthritis, Rheumatoid, 78-5153

Infectious Mononucleosis

- Virus, Epstein-Barr
- Seroepidemiology, Review, 78-4896

Insulin

- Aging
 - Immunity, Cellular, Review, 78-4906
- Benzo(a)pyrene
 - Cell Transformation, Neoplastic, 78-5001
- Breast Neoplasms
 - DNA Replication, 78-5392
 - Precancerous Conditions, 78-5392
- Immunosuppression
 - Cell Transformation, Neoplastic, Review, 78-4906
- Islet Cell Tumor
 - Virus, Polyoma, BK, 78-5263

Intestinal Neoplasms

- Adenocarcinoma
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-506
 - Morpholine, *N*-Nitroso-, 78-5056
- Adenoma
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-506
- Benzo(a)pyrene
 - Dietary Fiber, 78-5002
- Bile Acids and Salts
 - Epidemiology, Review, 78-4919
 - Feces, 78-5358
- Carcinoma, Epidermoid
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-506
- Cervix Neoplasms
 - Radiation, Ionizing, 78-5178
- Clostridium*
 - Epidemiology, 78-5358
 - Epidemiology, Review, 78-4919
 - Steroid Nuclear Dehydrogenation, Review, 78-4919
- Diet
 - Epidemiology, 78-5372
- Dietary Fats
 - Epidemiology, 78-5358
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Animal Model, Mouse, 78-5104
- Hydrazine, 1,2-Dimethyl-, Dihydrochloride
 - Anastomosis, 78-5062
 - Precancerous Conditions, 78-5059
- Morpholine, *N*-Nitroso-
 - Carcinogenic Potential, Fish, 78-5056
- Mycotoxins
 - Epidemiology, 78-5372
 - Food Contamination, 78-5372
- 3-Oxochola-4,6-dien-24-oic Acid
 - Hydroxylases, Review, 78-4919
- Sarcoma, Reticulum Cell
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5104
- Vegetables
 - Epidemiology, 78-5372

testine, Small

- Diethylamine, *N*-Nitroso-
Metabolism, Rat, 78-5073
- Dimethylamine, *N*-Nitroso-
Metabolism, Rat, 78-5073
- Piperidine, 1-Nitroso-
Metabolism, Rat, 78-5073
- Pyrrolidine, 1-Nitroso-
Metabolism, Rat, 78-5073

testines

- Bacteria
Mutagenic Activity, 78-4952

dine

- Benzo(a)pyrene
Co-carcinogenic Activity, Hamster, 78-5004
- Respiratory Tract Neoplasms
Benzo(a)pyrene, 78-5004

on

- Neurilemmoma
Urea, Methyl Nitroso-, 78-5085
- Occupational Hazard
Epidemiology, Review, 78-4934

on Oxide

- Lung Neoplasms
Adenocarcinoma, 78-5326

on Sulfate

- see Sulfuric Acid, Iron Salt

let Cell Tumor

- Pancreatic Neoplasms
Epidemiology, Sri Lanka, 78-5376
- Virus, Polyoma, BK, 78-5263
- Virus, Polyoma, BK
Calcitonin, 78-5263
- Glucagon, 78-5263
- Insulin, 78-5263

oantigens

- Breast Neoplasms
Epidemiology, Blacks, 78-5360

obutyric Acid, α -Amino-

- Virus, SV40
Amino Acids, 78-5259

ocyanic Acid, Ethyl Ester

- Ethylamine, *N*-Methyl-*N*-nitroso-
Nitrous Acid, Sodium Salt, 78-5080

oflurane

- see Ether, 1-Chloro-2,2,2-trifluoroethyl Difluoromethyl

onicotinamide, 2-Ethylthio-

- Glutathione
Metabolism, 78-5112
- Oxidoreductases
Metabolism, 78-5112

onicotinic Acid Hydrazide

- Drug Therapy
Carcinogenic Potential, Review, 78-4844

opropalin

- see Aniline, 2,6-Dinitro-*N,N*-dipropyl-*p*-isopropyl-

ow Neoplasms

- Granulosa Cell Tumor
Case Report, 78-5321

Kanechlor 500

- Acetamide, *N*-Fluoren-2-yl-
Hepatocarcinogenicity, Review, 78-4828
- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
Hepatocarcinogenicity, Review, 78-4828
- Diethylamine, *N*-Nitroso-
Hepatocarcinogenicity, Review, 78-4828
- Hepatoma
Carcinogenic Activity, Review, 78-4828

Kanechlor C

- Benzene, Hexachloro-
Co-carcinogenic Activity, 78-4979
- Hepatoma
Benzene, Hexachloro-, 78-4979
- Histological Study, Mouse, 78-4979
- Precancerous Conditions, 78-4979

Karyotyping

- Leukemia, Plasmacytic
Cells, Cultured, 78-5394
- Uterine Neoplasms
Turner's Syndrome, 78-5342

Kepone

- Metabolism
Rat, 78-4980

Keratoacanthoma

- Benz(a)anthracene, 7,12-Dimethyl-
12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
- Skin Neoplasms
Benz(a)anthracene, 7,12-Dimethyl-, 78-5036

Kidney Neoplasms

- Acetic Acid, Lead Salt
Porphyrins, 78-5116
- Adenocarcinoma
Neoplasm Metastasis, 78-5335
- 1-Propanol, 2,3-Dibromo-, Phosphate, 78-5167
- Adenoma
Ethanol, *N*-Nitrosoiminodi-, 78-4823
- Carcinoma
Epidemiology, 78-5375
- Precancerous Conditions, 78-5375
- Transplantation, Heterologous, 78-5336
- Carcinoma, Transitional Cell
Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
78-4976
- Diethylamine, *N*-Nitroso-
Carcinogenic Activity, Review, 78-4822
- Dimethylamine, *N*-Nitroso-
Carcinogenic Activity, Review, 78-4821
- Dipropylamine, *N*-Nitroso-
Carcinogenic Activity, Review, 78-4824
- Erythropoietin
Neoplasm Metastasis, 78-5336
- Estrogens
Growth Factors, 78-5134
- Mitosis, 78-5134
- Ethanol, *N*-Nitrosoiminodi-
Carcinogenic Potential, Review, 78-4823
- Hepatomegaly
Mouse, Nude, 78-5336
- Neoplasm Metastasis
Case Report, 78-5335
- Epidemiology, 78-5375
- 1-Propanol, 2,3-Dibromo-, Phosphate
Dose-Response Study, Rat, 78-5167
- Smoking

Kidney Neoplasms (cont'd)

- Epidemiology, Review, 78-4912
- Transplantation, Heterologous
- Erythropoietin, 78-5336
- Mouse, Nude, 78-5336
- Ultrastructural Study, 78-5336
- Vagina
- Neoplasm Metastasis, 78-5335

Kynurenic Acid

- see Quinaldic Acid, 4-Hydroxy-

Kynurenine

- see Alanine, 3-Anthraniloyl-

Lactate Dehydrogenase

- Acrylonitrile
- Aroclor 1254, 78-5111
- Barbituric Acid, 5-Ethyl-5-phenyl, 78-5111
- Cholanthrene, 3-Methyl-, 78-5111
- Cysteine, 78-5111
- Maleic Acid, Diethyl Ester, 78-5111
- Propionitrile
- Enzymatic Activity, 78-5111

Laryngeal Neoplasms

- Aryl Hydrocarbon Hydroxylases
- Genetics, Review, 78-4928
- Papilloma
- Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101
- Virus, Papilloma
- Epidemiology, Review, 78-4889

Lead

- Occupational Hazard
- Epidemiology, Review, 78-4934

Leiomyoma

- Peritoneal Neoplasms
- Case Report, 78-5306

Leiomyosarcoma

- Lung
- Neoplasm Metastasis, 78-5388
- Virus, Bovine Leukemia
- DNA-DNA Hybridization, 78-5230

Letterer-Siwe Disease

- Acid Phosphatase
- Enzymatic Activity, 78-5389
- Amino Acid Naphthylamidases
- Enzymatic Activity, 78-5389
- Esterases
- Enzymatic Activity, 78-5389
- Histiocytes
- Ultrastructural Study, 78-5389
- Immunoglobulins, Fc
- IgG, 78-5389
- T-Lymphocytes
- Rosette Formation, 78-5389

Leucine Aminopeptidase

- Stomach Neoplasms
- Metaplasia, 78-5328

Leukemia

- Anesthetics
- Epidemiology, Review, 78-4857
- Chromosome Abnormalities
- Precancerous Conditions, 78-5204
- Epidemiology
- Japan, 78-4917

Leukemia (cont'd)

- Ethnic Groups
- Epidemiology, Hawaii, 78-5347
- Ethylene, Chloro-
- Epidemiology, 78-4832
- Occupational Hazard, Review, 78-4832
- Oncogenic Viruses
- Epidemiology, Review, 78-4895
- Horizontal Transmission, 78-4899
- Radiation
- Japan, Review, 78-4869
- Radiation, Ionizing
- Dose-Response Study, Lizard, 78-4878
- Dose-Response Study, Review, 78-4871
- Epidemiology, Review, 78-4874
- Reverse Transcriptase
- Precancerous Conditions, 78-5204
- Schistosomiasis
- Epidemiology, Review, 78-4846
- Socioeconomic Factors
- Epidemiology, 78-5351
- Urea, 1-Butyl-1-nitroso-
- Age Factors, Rat, 78-5094
- Chromosome Abnormalities, 78-5095
- Virus, Baboon
- DNA-RNA Hybridization, Review, 78-4901
- Virus, C-Type RNA Tumor
- DNA, Neoplasm, 78-5274
- DNA, Viral, 78-5274
- Horizontal Transmission, Review, 78-4900
- B-Lymphocytes, 78-5270
- T-Lymphocytes, 78-5270
- Null Cells, 78-5270
- Reverse Transcriptase, 78-5270
- Trans-Species Infection, 78-5274
- Transformation, Genetic, 78-5274
- Ultrastructural Study, 78-5270
- Virus-Like Particles, Review, 78-4901
- Virus-Like Particles
- Dogs, 78-5229
- Precancerous Conditions, 78-5204
- Virus, Murine Leukemia
- Horizontal Transmission, 78-4899
- Virus, RNA Tumor
- Isolation and Characterization, 78-5271

Leukemia L1210

- Ethidium Bromide
- DNA Replication, 78-5114
- Ethidium Chloride
- DNA Replication, 78-5114

Leukemia, Lymphoblastic

- Alanine, 3-(*p*-Bis(2-chloroethyl)amino)phenyl)-
- Drug Therapy, 78-5302
- Antigens, Neoplasm
- Lymphocyte Culture Test, Mixed, 78-5286
- Multiple Myeloma
- Case Report, 78-5302
- Immunoglobulins, Light Chain, 78-5302
- B-Lymphocytes, 78-5302
- Null Cells
- Immune Response, 78-5286
- Lymphocyte Culture Test, Mixed, 78-5286

Leukemia, Lymphocytic

- Glucosephosphate Dehydrogenase
- A and B Isoenzymes, 78-5297
- B-Lymphocytes, 78-5297

emia, Lymphocytic (cont'd)

T-Lymphocytes, 78-5297

B-Lymphocytes

Immunoglobulins, Surface, 78-5297

Multiple Myeloma

Case Report, 78-5303

Drug Therapy, 78-5303

Immunoglobulins, Light Chain, 78-5303

Virus, Bovine Leukemia

DNA-DNA Hybridization, 78-5230

emia, Myeloblastic**Anemia, Aplastic**

Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-, 78-5147

Antigens, Neoplasm

Precancerous Conditions, 78-5397

Arthritis, Rheumatoid

Drug Therapy, 78-5151

Blood Platelets

Antigens, Neoplasm, 78-5397

Precancerous Conditions, 78-5397

Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-

Case Report, 78-5140, 78-5153

Multiple Sclerosis, 78-5153

Chromosome Aberrations

Anemia, Macrocytic, 78-5296

Case Report, 78-5298

Chromosomes, Human, 6-12, 78-5296

Chromosomes, Human, 19-20, 78-5296

Precancerous Conditions, 78-5296

Cyclophosphamide

Behcet's Syndrome, 78-5153

Case Report, 78-5151, 78-5153

Chromosome Abnormalities, 78-5151

Drug Therapy

Case Report, 78-5298

Glycoproteins

Precancerous Conditions, 78-5397

Heart Transplantation

Case Report, 78-5384

Immunosuppressive Agents

Arthritis, Rheumatoid, 78-5153

Multiple Sclerosis

Drug Therapy, 78-5140

Neoplasms, Multiple Primary

Drug Therapy, 78-5298

Radiotherapy, 78-5298

Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-

Case Report, 78-5147

Radiotherapy

Case Report, 78-5298

Sarcoma, Reticulum Cell

Case Report, 78-5299

Drug Therapy, 78-5299

Radiotherapy, 78-5299

***p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-**

Carcinogenic Activity, Monkey, 78-5148

Virus, Baboon

Virus-Like Particles, Review, 78-4901

Virus, RD-114

DNA-RNA Hybridization, 78-5226

Virus, Simian Sarcoma

Isolation and Characterization, Cell Culture
78-4900

Virus-Like Particles, Review, 78-4901

Leukemia, Myelocytic

Chromosome Abnormalities

Colony Stimulating Factor, Review, 78-4933

Leukocytes

Cell Differentiation, Review, 78-4933

Lipopolysaccharides

Leukocytes, 78-4933

Macrophages

Cell Differentiation, Review, 78-4933

Colony Stimulating Factor, Review, 78-4933

Reverse Transcriptase

Blood Platelets, 78-5204

Virus, C-Type RNA Tumor

Colony Stimulating Factor, Review, 78-4933

Leukemia, Plasmacytic

Cells, Cultured

Cell Cycle Kinetics, 78-5394

Plasma Cells, 78-5394

IgG

Cells, Cultured, 78-5394

Karyotyping

Cells, Cultured, 78-5394

Multiple Myeloma

Case Report, 78-5302

Leukocytes

Leukemia, Myelocytic

Cell Differentiation, Review, 78-4933

Lipopolysaccharides, 78-4933

4,4'-Stilbenediol, α,α' -Diethyl-

Chromosome Aberrations, 78-5137

Leukoencephalopathy, Progressive Multifocal

Virus, Polyoma, JC

Isolation and Characterization, Review, 78-4890
78-4891

Virus, SV40

Isolation and Characterization, Review, 78-4890

Leukopenia

Radiation, Ionizing

Dose-Response Study, Lizard, 78-4878

Linoleic Acid

Gastrointestinal Neoplasms

Papilloma, 78-5120

Lipids

Benzo(a)pyrene

Liver, Rat, 78-5014

Diethylamine, *N*-Nitroso-

Metabolism, Rat, 78-5073

Dimethylamine, *N*-Nitroso-

Metabolism, Rat, 78-5073

Piperidine, 1-Nitroso-

Metabolism, Rat, 78-5073

Pyrrolidine, 1-Nitroso-

Metabolism, Rat, 78-5073

Lipopolysaccharides

Leukemia, Myelocytic

Leukocytes, 78-4933

Virus, Friend Murine Leukemia

Antigen-Antibody Reactions, 78-5207

Immunosuppression, 78-5207

Lipoproteins

Immunosuppression

Cell Transformation, Neoplastic, Review, 78-4906

Liposarcoma

- Ethylene, Bromo- Polymer
Carcinogenic Potential, Rat, 78-5167
- Pregnancy
Case Report, 78-5390

Liver

- Aldrin
Metabolism, Rat, 78-5161
- Dieldrin
Metabolism, Rat, 78-5161

Liver Cirrhosis

- Hepatitis
Australia Antigen, 78-5276
Epidemiology, Indonesia, 78-5332

Liver Neoplasms

- Adenocarcinoma
Acetamide, *N*-Fluorenyl-, 78-4983
Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -isomer
78-4983
- Adenoma
Contraceptives, Oral, 78-5130
Epidemiology, Review, 78-4865
Morpholine, *N*-Nitroso-, 78-5056
- Aflatoxin B1
Epidemiology, Zaire, 78-4968
- Aflatoxin G1
Epidemiology, Zaire, 78-4968
- Arsenic
Precancerous Conditions, Review, 78-4922
- Barbituric Acid, 5-Ethyl-5-phenyl-
Histochemical/Histological Study, Mouse, 78-5325
Precancerous Conditions, 78-5325
- Contraceptives, Oral
Case Report, 78-5130, 78-5131
Epidemiology, 78-5133
Epidemiology, Review, 78-4865
Precancerous Conditions, 78-5133
- Diethylamine, *N*-Nitroso-
Carcinogenic Activity, Review, 78-4822
- Dimethylamine, *N*-Nitroso-
Adrenergic Alpha Receptor Blockers, 78-5067
Atropine, 78-5067
Carcinogenic Activity, Review, 78-4821
Proserine, 78-5067
- Dipropylamine, *N*-Nitroso-
Carcinogenic Activity, Review, 78-4824
- Epidemiology
Indonesia, 78-5345
- Ethylene, Chloro-
Epidemiology, 78-4832
Occupational Hazard, Review, 78-4832
Precancerous Conditions, Review, 78-4922
- Hamartoma
Contraceptives, Oral, 78-5131
- Hemangiopericytoma
Dimethylamine, *N*-Nitroso-, 78-5074
Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063
Quinoline, 78-5099
Symphitum officinale, 78-4942
- Hemangioma
Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
- Kidney Transplantation
Epidemiology, 78-5353
- Mesenchymoma
Benzo(a)pyrene, 78-5014
- Mestranol

Liver Neoplasms (cont'd)

- Epidemiology, Review, 78-4865
- Morpholine, *N*-Nitroso-
Carcinogenic Potential, Fish, 78-5056
Estradiol, 3-Benzoate, 78-5055
Estradiol, 17-Hydrocinnamate, 78-5055
- Mycotoxins
Precancerous Conditions, Review, 78-4840
- Polychlorobiphenyl Compounds
Carcinogenic Potential, Review, 78-4829
- Propylamine, *N*-(2-Hydroxypropyl)-*N*-nitroso-
Carcinogenic Metabolite, Review, 78-4824
- Thorium Dioxide
Precancerous Conditions, Review, 78-4922

LSD-25

see Lysergic Acid Diethylamide

Lung

- Angiosarcoma
Neoplasm Metastasis, 78-5388
- Fibrosarcoma
Neoplasm Metastasis, 78-5388
- Hemangiopericytoma
Neoplasm Metastasis, 78-5388
- Leiomyosarcoma
Neoplasm Metastasis, 78-5388
- Neoplasm Metastasis
Animal Model, B16 Tumor, 78-4923
- Rhabdomyosarcoma
Neoplasm Metastasis, 78-5388
- Sarcoma
Neoplasm Metastasis, 78-5388

Lung Neoplasms

- Adenocarcinoma
Asbestos, 78-5326
Dyes, 78-4973
Histochemical Study, 78-5327
Iron Oxide, 78-5326
Kidney Transplantation, 78-5353
Quartz, 78-5326
Siderosis, 78-5326
Silver, 78-5326
Transplantation Immunology, 78-5293
Ultrastructural Study, 78-5327
- Adenoma
Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
Imidazole-1-ethanol, 2-Methyl-5-nitro-, 78-4844
- Air Pollution
Asphalt Plant, 78-5378
Epidemiology, 78-5378
Epidemiology, Great Britain, Review, 78-4849
- Aryl Hydrocarbon Hydroxylases
Genetics, Review, 78-4928
- Asbestos
Epidemiology, Review, 78-4841, 78-4842
- Benzo(a)pyrene
Air Pollution, Review, 78-4849
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro
DNA Repair, 78-5009
- 1,3-Butadiene, 2-Chloro-
Epidemiology, Review, 78-4834
- Carcinogen, Environmental
Epidemiology, Review, 78-4810
Multistate Model, Review, 78-4810
Mutagenicity Assay, Review, 78-4810
- Carcinoma, Bronchiolar

ig Neoplasms (cont'd)

- Ultrastructural Study, 78-5327
- Carcinoma, Epidermoid
 - Adriamycin, 78-5399
 - Aryl Hydrocarbon Hydroxylases, Review, 78-4853
 - Guanosine, 6-Seleno-, 78-5399
 - Melamine, Hexamethyl-, 78-5399
 - Methotrexate, 78-5399
 - Neoplasm Transplantation, 78-5399
 - Purine-6-thiol, 9- β -D-Arabinofuranosyl-, 78-5399
 - Purine-6(1*H*)-thione, 2-Amino-, 78-5399
 - Smoke Condensate, 78-5045
 - Uracil, 5-Fluoro-, 78-5399
 - Urea, Methylcyclohexyl Nitroso-, 78-5399
- Cholelithiasis
 - Epidemiology, 78-5337
- Epidemiology
 - Geographic Factors, 78-5344
- Giant Cell Tumors
 - Siderosis, 78-5326
 - Silver, 78-5326
 - Titanium Oxide, 78-5326
- Imidazole-1-ethanol, 2-Methyl-5-nitro-
 - Carcinogenic Potential, Review, 78-4844
- Kidney Transplantation
 - Epidemiology, 78-5353
- Lymphocytes
 - Aryl Hydrocarbon Hydroxylases, Review, 78-4853
- Neoplasms, Multiple Primary
 - Occupational Hazard, 78-4973
- Nitric Acid, Plutonium Salt
 - Dog, 78-5185
- Occupational Hazard
 - Case Report, 78-5326
 - Epidemiology, 78-5344
 - Epidemiology, Review, 78-4841
- Papilloma
 - 1-Propanol, 2,3-Dibromo-, Phosphate, 78-5167
- Pesticides
 - Epidemiology, Review, 78-4835
- Phosphonium, Tetrakis(hydroxymethyl)-, Chloride
 - Carcinogenic Potential, Rat, 78-5167
- Plutonium
 - Carcinogenic Potential, Review, 78-4868
- 1-Propanol, 2,3-Dibromo-, Phosphate
 - Dose-Response Study, Rat, 78-5167
- Radiation, Ionizing
 - Dose-Response Study, Review, 78-4871
- Smoke Condensate
 - Beeswax Pellet Implantation, 78-5045
 - Neoplasm Metastasis, 78-5045
- Smoking
 - Epidemiology, Great Britain, Review, 78-4849
 - Epidemiology, Japan, 78-4917
 - Epidemiology, Review, 78-0485
 - Low Tar/Nicotine Cigarettes, Review, 78-0485
- Transplantation Immunology
 - Age Factors, Mouse, 78-5293
- Virus, Adeno 12
 - Carcinogenic Potential, 78-5235
 - DNA-DNA Hybridization, 78-5235
- Lymphocyte Depletion
 - Skin Neoplasms
 - T-Lymphocytes, 78-5174
 - Virus, Friend Murine Leukemia
 - T-Lymphocytes, 78-5283

Lymphocyte Transformation

- Virus, Friend Murine Leukemia
 - Cells, Cultured, 78-5283
- Virus, Herpes Ateles
 - T-Lymphocytes, 78-5239
 - Marmoset, 78-5239
 - Radiation, Ionizing, 78-5239
- Virus, Herpes Simplex 2
 - Plant Agglutinins, 78-5243
 - Spleen Cells, 78-5243

Lymphocytes

- Azathioprine
 - Chromosome Aberrations, 78-5157
- Benzene
 - Acid Phosphatase, 78-4974
- Benzene, (Epoxyethyl)-
 - Chromosome Aberrations, 78-4953
- Caffeine
 - Chromatids, 78-5145
 - DNA Repair, 78-5145
- Cyclophosphamide
 - Chromosome Aberrations, 78-5155
- Ethylene, Chloro-
 - Chromosome Aberrations, 78-5157
- Hodgkin's Disease
 - Concanavalin A, 78-5300
- Lung Neoplasms
 - Aryl Hydrocarbon Hydroxylases, Review, 78-4853
- Lysergic Acid Diethylamide
 - Chromosome Aberrations, 78-5050
- Lysergic Acid Diethylamide, Tartrate
 - Chromosome Aberrations, 78-5050
- Mitomycin C
 - Chromosome Aberrations, 78-5050, 78-5157
 - DNA Repair, 78-5145
- Phosphine Oxide, Tris(1-aziridinyl)-
 - Chromosome Aberrations, 78-5157
- Respiratory Tract Neoplasms
 - Aryl Hydrocarbon Hydroxylases, 78-5324
- Sirodesmin A
 - Chromosome Aberrations, 78-5155
- Smallpox Vaccine
 - Chromosome Aberrations, 78-5157
- Thrombin
 - Chromosome Aberrations, 78-5157
- Ultraviolet Rays
 - Immunosuppression, 78-5285
- Urea, Methyl Nitroso-
 - Immune Response, Rat, 78-5086
- Vinblastine
 - Chromosome Aberrations, 78-5157
- Vincristine
 - Chromosome Aberrations, 78-5157

B-Lymphocytes

- Leukemia
 - Virus, C-Type RNA Tumor, 78-5270
- Leukemia, Lymphoblastic
 - Multiple Myeloma, 78-5302
- Leukemia, Lymphocytic
 - Glucosephosphate Dehydrogenase, 78-5297
 - Immunoglobulins, Surface, 78-5297
- Virus, Friend Murine Leukemia
 - Antigen-Antibody Reactions, 78-5283

T-Lymphocytes

- Aging
 - Immunity, Cellular, Review, 78-4906

T-Lymphocytes (cont'd)

- Asbestos
 - Immune Response, Rat, 78-5284
- Carcinogen, Chemical
 - Immune Response, Review, 78-4907
- Letterer-Siwe Disease
 - Rosette Formation, 78-5389
- Leukemia
 - Virus, C-Type RNA Tumor, 78-5270
- Leukemia, Lymphocytic
 - Glucosephosphate Dehydrogenase, 78-5297
- Sarcoma
 - Macrophages, 78-5287
 - Virus, Murine Sarcoma, 78-5287
- Sarcoma 180, Crocker
 - Anti-Antibodies, 78-5279
 - Growth, 78-5279
 - Transplantation Immunology, 78-5279
- Skin Neoplasms
 - Lymphocyte Depletion, 78-5174
- Ultraviolet Rays
 - Immunosuppression, 78-5173
 - Transplantation Immunology, 78-5173
- Virus, Adeno
 - Immune Response, Review, 78-4905
- Virus, C-Type RNA Tumor
 - Immunosuppression, 78-5272
 - Suppressive Component, Isolation and Characterization, 78-5272
- Virus, DNA Tumor
 - Immune Response, Review, 78-4907
- Virus, Feline Leukemia
 - Immune Response, Review, 78-4907
- Virus, Friend Murine Leukemia
 - Lymphocyte Depletion, 78-5283
- Virus, Herpes Ateles
 - Antilymphocyte Serum, 78-5239
 - Lymphocyte Transformation, 78-5239
 - Rosette Formation, 78-5239
- Virus, Marek's Disease Herpes
 - Immunosuppression, 78-5196
 - Virus, Turkey Herpes, 78-5196
- Virus, Radiation Leukemia
 - Cell Transformation, Neoplastic, 78-5295
 - Precancerous Conditions, 78-5295
 - Transplantation Immunology, 78-5295
- Virus, SV40
 - Immune Response, Review, 78-4905
- Virus, Turkey Herpes
 - Immunosuppression, 78-5196
 - Plant Agglutinins, 78-5196

Lymphocytosis

- Virus, Bovine Leukemia
 - Epidemiology, 78-4886

Lymphoma (General and Unspecified)

- Anesthetics
 - Epidemiology, Review, 78-4857
- Arsenic
 - Epidemiology, Review, 78-4934
- Cholanthrene, 3-Methyl-
 - Antigens, Viral, 78-5291
 - Tumor Dormancy, 78-5291
- Epilepsy
 - Drug Therapy, 78-5301
- Genetics
 - Immune Response, Review, 78-4913

Lymphoma (General and Unspecified) (cont'd)

- Heart Transplantation
 - Case Report, 78-5384
- Histocompatibility Antigens
 - Transplantation Immunology, 78-5294
- Hydantoin, 5,5-Diphenyl-
 - Case Report, 78-5301
- Kidney Transplantation
 - Epidemiology, 78-5353
- Macrophages
 - Tumor Dormancy, 78-5291
- Oncogenic Viruses
 - Epidemiology, Review, 78-4895
- Parasitic Diseases
 - Epidemiology, Review, 78-4913
- Radiation, Ionizing
 - Epidemiology, Review, 78-4874
 - Growth, Spleen, Liver, 78-5294
 - Histocompatibility Antigens, 78-5294
 - Transplantation Immunology, 78-5294
- Virus, Abelson Murine Leukemia
 - Age Factors, 78-5198
 - Antibody Specificity, 78-5200
 - Antigens, Viral, 78-5200
 - Dose-Response Study, 78-5198
 - Strain Difference, 78-5198
- Virus, C-Type RNA Tumor
 - Viral Proteins, 78-5210
- Virus, Epstein-Barr
 - Immunoglobulins, Surface, 78-5278
- Virus, Gross Murine Leukemia
 - Transplantation Immunology, 78-5291
- Virus, Murine Leukemia
 - Epidemiology, Wild Mouse, Review, 78-4883
 - Vertical Transmission, 78-5209
- Virus, Polyoma, BK
 - Carcinogenic Activity, Hamster, 78-5263

Lymphoma, Giant Follicular

- Schistosomiasis
 - Epidemiology, Review, 78-4846

Lymphosarcoma

- Oncogenic Viruses
 - Horizontal Transmission, 78-4899
- Pregnancy
 - Case Report, 78-5390
- Reverse Transcriptase
 - Dogs, 78-5229
- p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
 - Carcinogenic Activity, Monkey, 78-5148
- Virus, Bovine Leukemia
 - DNA-DNA Hybridization, 78-5230
 - Epidemiology, 78-4886
 - Horizontal/Vertical Transmission, Review, 78-4886
- Virus-Like Particles
 - Dogs, 78-5229
- Virus, Murine Leukemia
 - Horizontal Transmission, 78-4899

Lysergic Acid Diethylamide

- Chromosome Aberrations
 - Dose-Response Study, 78-5050
 - Lymphocytes, 78-5050
- DNA Replication
 - Cells, Cultured, 78-5050

Lysergic Acid Diethylamide, Tartrate

- Chromosome Aberrations

Ergic Acid Diethylamide, Tartrate (cont'd)
Lymphocytes, 78-5050

Ames Test

Pyrolysis Products, 78-5122

Mutagens

Inactivation, Fruit/Vegetable Juices, 78-5122
Pyrolysis Products, 78-5122

Macrophages

Asbestos

Cysteine, 78-5284
Immune Response, Rat, 78-5284
Periodic Acid, Sodium Salt, 78-5284

Benzo(a)pyrene

Metabolism, 78-5020
Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
Metabolism, 78-5020

Fibrosarcoma

Carcinogen, Chemical, 78-5288
Growth, 78-5288
Winn Test, 78-5288

Leukemia, Myelocytic

Cell Differentiation, Review, 78-4933
Colony Stimulating Factor, Review, 78-4933

Lymphoma

Tumor Dormancy, 78-5291

Sarcoma

T-Lymphocytes, 78-5287
Migration Inhibitory Factor, 78-5287
Virus, Murine Sarcoma, 78-5287

Urea, Methyl Nitroso-

Immune Response, Rat, 78-5086

Virus, Avian Carcinoma

Cell Transformation, Neoplastic, 78-5190

Virus, Coxsackie B

Hypercholesteremia, 78-5290

Virus, Friend Murine Leukemia

Antigen-Antibody Reactions, 78-5207

Malaria

Burkitt's Lymphoma

Epidemiology, Review, 78-4911

Virus, Epstein-Barr

Epidemiology, Review, 78-4895, 78-4904

Maleic Acid, Diethyl Ester

Acrylonitrile

Lactate Dehydrogenase, 78-5111
Oxidoreductases, 78-5111

Maleimide, N-Ethyl-

see 1*H*-Pyrrole-2,5-dione, 1-Ethyl-

Mammary Neoplasms, Experimental

Adenocarcinoma

Benz(a)anthracene, 7,12-Dimethyl-, 78-5030
Dexamethasone, 78-5143
Transplantation Immunology, 78-5293

Adenoma

Contraceptives, Oral, 78-5132

Anagestone Acetate

Hyperplasia, 78-5132

Benz(a)anthracene, 7,12-Dimethyl-

Dietary Fats, 78-5025
Stress, 78-5029

Carcinoma

Urea, Ethyl Nitroso-, 78-5088

Contraceptives, Oral

Mammary Neoplasms, Experimental (cont'd)

Histological Study, Dog, 78-5132

Dexamethasone

Mouse, Nude, 78-5143
Reverse Transcriptase, 78-5143
Transplantation Immunology, 78-5143

Dietary Fats

Co-carcinogenic Effect, Rat, 78-5025

Estrogens

Growth Factors, 78-5134
Mitosis, 78-5134

Ethynone

Hyperplasia, 78-5132

Fusarium sporotrichioides

Toxins, 78-4954

Mestranol

Hyperplasia, 78-5132

Norgestrel, Chloroethynyl-

Hyperplasia, 78-5132

Prolactin

Estrogens, 78-5134

Radiation, Ionizing

Anesthetics, 78-5177
Dose-Response Study, Rat, 78-5177

Retinol

Antineoplastic Activity, Review, 78-4860

Sarcoma

Urea, Ethyl Nitroso-, 78-5088

4,4'-Stilbenediol, α,α' -Diethyl-

Radiation, Ionizing, 78-5177

Stress

Neoplasm Regression, 78-5029

Thyroid Hormones

Estrogens, 78-5134

Transplantation Immunology

Age Factors, Mouse, 78-5293
Mouse, Nude, 78-5143

Urea, Ethyl Nitroso-

Dose-Response Study, 78-5088

Neonate, Rat, 78-5088

Transplacental Carcinogenesis, 78-5088

Virus-Like Particles

Dogs, 78-5229

Virus, Murine Mammary Tumor

Dexamethasone, 78-5143

Medroxyprogesterone Acetate

see Pregn-4-ene-3,20-dione, 17-(Acetyloxy)-6 α -methyl-

Melamine, Hexamethyl-

Lung Neoplasms

Carcinoma, Epidermoid, 78-5399

Melanoma

Genetics

Epidemiology, Review, 78-4876

Marriage

Case Report, 78-5387
Epidemiology, 78-5387

Neoplasm Circulating Cells

Cell Survival, 78-4923

Neoplasm Metastasis

Animal Model, B16 Tumor, 78-4923

Nose Neoplasms

Epidemiology, USSR, 78-5379

Otorhinolaryngologic Neoplasms

Epidemiology, USSR, 78-5379

Reverse Transcriptase

Isolation and Characterization, B16 Tumor, 78-5221

- Melanoma (cont'd)**
 RNA, Viral, 78-5221
 Ultraviolet Rays
 Epidemiology, Review, 78-4876, 78-4877
 Urea, Methyl Nitroso-
 Histological Study, Gerbil, 78-5090
 Site Distribution, 78-5090
 Virus-Like Particles
 DNA-RNA Hybridization, 78-5221
 Isolation and Characterization, B16 Tumor, 78-5221
 Virus, RNA Tumor
 Virus-Like Particles, 78-5221
 Virus, SV40
 Isolation and Characterization, Review, 78-4890
 Virus-Like Particles, Review, 78-4891
- Meningioma**
 Acromegaly
 Case Report, 78-5318
 Pituitary Neoplasms
 Case Report, 78-5318
 Virus, SV40
 Virus-Like Particles, Review, 78-4891
- Mesenchymoma**
 Benzo(a)pyrene
 Catalase, 78-5014
 Glutathione Peroxidase, 78-5014
 Glutathione Reductase, 78-5014
 Urate Oxidase, 78-5014
 Liver Neoplasms
 Benzo(a)pyrene, 78-5014
 Stomach Neoplasms
 Morpholine, *N*-Nitroso-, 78-5056
- Mesothelioma**
 Asbestos
 Epidemiology, 78-5377, 78-5385
 Epidemiology, Review, 78-4842
 Epidemiology, Turkey, 78-4909
 Estrogens
 Carcinogenic Potential, Review, 78-0486
 Peritoneal Neoplasms
 Epidemiology, 78-5377
 Pleural Neoplasms
 Epidemiology, 78-5377, 78-5385
 Occupational Hazard, 78-5385
 Silicate
 Epidemiology, Turkey, 78-4909
- Mestranol**
 Hepatoma
 Epidemiology, Review, 78-4864
 Liver Neoplasms
 Epidemiology, Review, 78-4865
 Mammary Neoplasms, Experimental
 Hyperplasia, 78-5132
- Metaplasia**
 Esophageal Neoplasms
 Precancerous Conditions, 78-5323
 Gastric Mucosa
 Precancerous Conditions, 78-5328
 Precancerous Conditions
 Histological Study, 78-5323
 Stomach Neoplasms
 Carcinoma, 78-5328
 Intestines, 78-5328
 Leucine Aminopeptidase, 78-5328
 Precancerous Conditions, 78-5328
- Metaplasia (cont'd)**
 Ultrastructural Study, 78-5328
- Methane, Azoxy-**
 Colonic Neoplasms
 Enterectomy, 78-5159
 Hyperplasia, 78-5159
 Nucleic Acids
 Intestinal Mucosa, 78-5159
- Methane, Sulfinylbis-**
 Benzo(a)pyrene
 Cell Transformation, Neoplastic, 78-5001
- Methanesulfonic Acid, Ethyl Ester**
 Anemia, Aplastic
 Chromosome Aberrations, 78-4939
 Cell Survival
 Dose-Response Study, Hamster Ovary Cells
 78-5158
 DNA Repair
Escherichia coli, 78-5102
Escherichia coli
 Mutagenic Activity, 78-5102
 Hypoxanthine Phosphoribosyltransferase
 Mutagenic Activity, 78-5158
Neurospora crassa
 Lethal/Mutagenic Activity, 78-4947
- Methanesulfonic Acid, Isopropyl Ester**
 Cell Survival
 Dose-Response Study, Hamster Ovary Cells
 78-5158
 Hypoxanthine Phosphoribosyltransferase
 Mutagenic Activity, 78-5158
- Methanesulfonic Acid, Methyl Ester**
 Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 Lethal/Mutagenic Activity, 78-4987
 Ames Test
 Carcinogenic Potential, Review, 78-4815
 Anemia, Aplastic
 Chromosome Aberrations, 78-4939
 Azaguanine Resistance
 Hamster V79 Cells, 78-4987
 Cell Survival
 Dose-Response Study, Hamster Ovary Cells
 78-5158
 Chromatids
 Bone Marrow, Mouse, 78-5149
 DNA Repair
Escherichia coli, 78-5102
Escherichia coli
 Mutagenic Activity, 78-4950, 78-5102
 Hypoxanthine Phosphoribosyltransferase
 Mutagenic Activity, 78-5158
Neurospora crassa
 Lethal/Mutagenic Activity, 78-4947
 Plasmids
 Ames Test, 78-4950
 Mutagenic Activity, 78-4950
- Methanesulfonic Acid, 1-Methylethyl Ester**
 see Methanesulfonic Acid, Isopropyl Ester
- Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)**
 Cholinesterases
 Enzyme Activation, 78-5064
 DNA Replication
 Colon, Rat, 78-5064
 Physostigmine, Sulfate

thanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester) (cont'd)
DNA Replication, 78-5064

thanol, (Methyl-*ONN*-azoxy)-
Electrophiles
Carcinogenic Metabolite, Review, 78-4820
Nucleic Acids
Methylation, Review, 78-4820

thionine
Protein Kinase
Liver, Rat, 78-5398

thotrexate
Lung Neoplasms
Carcinoma, Epidermoid, 78-5399

thoxyflurane
see Ether, 2,2-Dichloro-1,1-difluoroethyl Methyl

thylamine Hydrochloride
Acetic Acid, Methylnitrosaminomethyl Ester
Synthesis, 78-5066

thylamine, *N*-Nitroso-
Purine, 2-Amino-6-methoxy-
DNA, Alkylation, 78-5106
Radioimmunoassay, 78-5106

thylazoxymethanol Acetate
see Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)

thylbutyroxymethylnitrosamine
see Butyric Acid, Methylnitrosaminomethyl Ester

trifonate
Administration Route
Carcinogenic Potential, Mouse, 78-5164
Antischistosomal Drugs
Mutagenic Activity, Review, 78-4845

icrosomes
Virus, Adeno 12
Transferases, 78-5234

icrosomes, Liver
Acetamide, *N*-Fluoren-2-yl-
Hydroxylation, Carbon Atom 9, 78-4985
Acetamide, *N*-Fluoren-3-yl-
Hydroxylation, Carbon Atom 9, 78-4985
Metabolism, Rat, Guinea Pig, 78-4985
Acetohydroxamic Acid, *N*-Fluoren-3-yl-
Hydroxylation, Carbon Atom 9, 78-4985
Metabolism, Rat, Guinea Pig, 78-4985

Aflatoxin B1
Ames Test, 78-4969
Mutagenic Activity, 78-4969

Aflatoxin G1
Ames Test, 78-4969
Mutagenic Activity, 78-4969

Aflatoxin M1
Mutagenic Activity, 78-4969

Benzo(a)pyrene
Cholanthrene, 3-Methyl-, 78-4854
Metabolism, Review, 78-4854

7,8-Benzoflavone
Aryl Hydrocarbon Hydroxylases, 78-5011

Biphenyl, 2-Bromo-
Ames Test, 78-4978

Biphenyl, 3-Bromo-
Ames Test, 78-4978

Biphenyl, 4-Bromo-
Ames Test, 78-4978

Microsomes, Liver (cont'd)

Carcinogen, Chemical
Ames Test, 78-4940
Cholanthren-1-ol, 3-Methyl-
Ames Test, 78-4995
Cholanthren-2-ol, 3-Methyl-
Ames Test, 78-4995
Cholanthren-1-one, 3-Methyl-
Ames Test, 78-4995
Cholanthren-2-one, 3-Methyl-
Ames Test, 78-4995
Cholanthrene, 9,10-Dihydro-3-methyl-1,9,10-trihydroxy-
Ames Test, 78-4995
Cholanthrene, 3-Methyl-
Ames Test, 78-4995

Cyclophosphamide
Chromatids, 78-5152
Chromosome Aberrations, 78-5152

Dibenz(a,c)aceanthrylene
DNA, Binding, 78-5033
Polyribonucleotides, 78-5033
RNA, Transfer, 78-5033

Dimethylamine, *N*-Nitroso-
Chromatids, 78-5152
Chromosome Aberrations, 78-5152
Metabolism, Mouse, 78-5075

Ellipticine
Aryl Hydrocarbon Hydroxylases, 78-5011
Cytochromes, 78-5011

Hydroxylamine
Cytochrome P-450, 78-4989
Enzymatic Activity, 78-4989
Hydroxylases, 78-4989
Oxidoreductases, 78-4989

Opium
Ames Test, 78-5049
Phosphine Oxide, Diethoxymorpholino-
Carcinogenic Activity, 78-5150
Phosphine Oxide, Tripiperidino-
Carcinogenic Activity, 78-5150
Phosphoric Triamide, Hexamethyl-
Carcinogenic Activity, 78-5150
Phosphorothioic Triamide, Hexamethyl-
Carcinogenic Activity, 78-5150
Phosphorothioic Trimorpholide
Carcinogenic Activity, 78-5150
Phosphorus Triamide, Hexamethyl-
Carcinogenic Activity, 78-5150
1-Propanone, 2-Methyl-1,2-di-3-pyridyl-, Tartrate
Aryl Hydrocarbon Hydroxylases, 78-5011
Selenic Acid, Disodium Salt
Mutagenic Activity, 78-5115
Selenious Acid, Disodium Salt
Mutagenic Activity, 78-5115

Migration Inhibitory Factor

Cell Transformation, Neoplastic
Isolation and Characterization, Hamster Cells
78-5393

Sarcoma
Macrophages, 78-5287

Misonidazole

see 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-

Mitomycin C

Anemia, Aplastic
Chromosome Aberrations, 78-4939
Micronucleus Test, 78-4939

Mitomycin C (cont'd)

- Caffeine
 - Chromatids, 78-5145
 - DNA Repair, 78-5145
- Chromosome Aberrations
 - Lymphocytes, 78-5050
- DNA Repair
 - Lymphocytes, 78-5145
- Lymphocytes
 - Chromosome Aberrations, 78-5157
- Mutagenic Activity
 - Micronucleus/Nucleus Anomaly Tests, Hamster 78-5156
- Virus, SV40
 - DNA Replication, 78-5253

Mitosis

- Erythroleukemia
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-5136
- Kidney Neoplasms
 - Estrogens, 78-5134
- Mammary Neoplasms, Experimental
 - Estrogens, 78-5134
- Pituitary Neoplasms
 - Estrogens, 78-5134
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Cells, Cultured, 78-5136
 - HeLa Cells, 78-5136

Monoolein

- Gastrointestinal Neoplasms
 - Papilloma, 78-5120

Monostearin

- see Stearic Monoglyceride

Morpholine, N-Nitroso-

- Bile Duct Neoplasms
 - Cysts, 78-5056
- Hepatoma
 - Carcinogenic Potential, Fish, 78-5056
 - Chromosomal Proteins, Non-Histone, 78-5054
 - Estradiol, 3-Benzoate, 78-5055
 - Estradiol, 17-Hydrocinnamate, 78-5055
 - Neoplasm Metastasis, 78-5054
 - Ultrastructural Study, Rat, 78-5055
- Intestinal Neoplasms
 - Adenocarcinoma, 78-5056
 - Carcinogenic Potential, Fish, 78-5056
- Liver Neoplasms
 - Adenoma, 78-5056
 - Carcinogenic Potential, Fish, 78-5056
 - Estradiol, 3-Benzoate, 78-5055
 - Estradiol, 17-Hydrocinnamate, 78-5055
- Stomach Neoplasms
 - Carcinogenic Potential, Fish, 78-5056
 - Mesenchymoma, 78-5056

Mouth Neoplasms

- Areca
 - Carcinogenic Activity, Mouse, Rat, 78-4955
- Benz(a)anthracene, 7,12-Dimethyl-
 - Histological Study, Hamster, 78-5028
 - Precancerous Conditions, 78-5028
- 1-Propanol, 2,3-Dibromo-, Phosphate
 - Carcinogenic Potential, Rat, 78-5167
- Virus, Papilloma
 - Epidemiology, Review, 78-4889

Multiple Myeloma

- Epilepsy
 - Drug Therapy, 78-5301
- Hydantoin, 5,5-Diphenyl-
 - Case Report, 78-5301
- Leukemia, Lymphoblastic
 - Case Report, 78-5302
 - Immunoglobulins, Light Chain, 78-5302
 - B-Lymphocytes, 78-5302
- Leukemia, Lymphocytic
 - Case Report, 78-5303
 - Drug Therapy, 78-5303
 - Immunoglobulins, Light Chain, 78-5303
- Leukemia, Plasmacytic
 - Case Report, 78-5302
- Plasma Cells
 - In Vitro Model, 78-5394

Multiple Sclerosis

- Leukemia, Myeloblastic
 - Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl) 78-5153
 - Drug Therapy, 78-5140

Muramidase

- Agrobacterium tumefaciens*
 - Proteins, 78-5400

Mutagens

- Ames Test
 - Feces, 78-5121
- Arginine
 - Inactivation, Fruit/Vegetable Juices, 78-5122
 - Pyrolysis Products, 78-5122
- Ascorbic Acid
 - Feces, 78-5121
- Aspergillus nidulans*
 - Methionine Requirement, Fluctuation Test, 78-4944
- Citrulline
 - Inactivation, Fruit/Vegetable Juices, 78-5122
 - Pyrolysis Products, 78-5122
- Drosophila*
 - Test Evaluation, 78-4812
- Lysine
 - Inactivation, Fruit/Vegetable Juices, 78-5122
 - Pyrolysis Products, 78-5122
- Nitroso Compounds
 - Feces, 78-5121
- Ornithine
 - Inactivation, Fruit/Vegetable Juices, 78-5122
 - Pyrolysis Products, 78-5122
- Tryptophan
 - Inactivation, Fruit/Vegetable Juices, 78-5122
 - Pyrolysis Products, 78-5122

Myasthenia Gravis

- Thymoma
 - Histological Study, 78-5315
 - Neoplasm Invasiveness, 78-5315

Mycotoxins

- DNA Repair
 - Liver, 78-4949
- Food Contamination
 - Ames Test, 78-5048
- Hepatoma
 - Co-carcinogenic Activity, Review, 78-4840
 - Epidemiology, Review, 78-4839, 78-4840, 78-4862
 - Food Contamination, Review, 78-4839
- Intestinal Neoplasms

otoxins (cont'd)

- Epidemiology, 78-5372
- Food Contamination, 78-5372

Liver Neoplasms

- Precancerous Conditions, Review, 78-4840

Penicillium camemberti

- Review, 78-4837

Virus, Hepatitis

- Co-carcinogenic Activity, Review, 78-4839

elofibrosis

- Glycoproteins
- Precancerous Conditions, 78-5397

eloid Metaplasia

- Glycoproteins
- Precancerous Conditions, 78-5397

eloma Proteins

- Immunoglobulins, Heavy Chain
- Amino Acids, 78-5277
- Insulin, Binding, 78-5277
- Plasmacytoma
- Graft Rejection, 78-5289

osarcoma

- Pregnancy
- Case Report, 78-5390

DP

- Aldrin
- Metabolism, Rat, 78-5161
- Dieldrin
- Metabolism, Rat, 78-5161

hthalene

- Ames Test
- Dihydrodiols, 78-4855

aphthalenesulfonic Acid, 4-Hydroxy-3,4'-azodi-, Disodium Salt

- Dose-Response Study
- Carcinogenic Potential, Rat, 78-5129
- Fertility, Rat, 78-5128
- Toxicity, Rat, 78-5128, 78-5129
- Respiratory Tract Diseases
- Dose-Response Study, 78-5129

aphthylamine

- Ames Test
- Carcinogenic Potential, Review, 78-4815

opharyngeal Neoplasms

- Carcinoma
- Virus, Epstein-Barr, 78-5241
- Epidemiology
- Sabah, Malaysia, 78-5346
- Ethnic Groups
- Epidemiology, Alaska, 78-5380
- Virus, Epstein-Barr
- Epidemiology, Alaska, 78-5380
- Epidemiology, Review, 78-4895
- Seroepidemiology, Review, 78-4896

sseria

- Cervix Neoplasms
- Virus, Herpes, 78-5341

plasm Circulating Cells

- Hepatoma
- Fibrin, 78-5304
- Lung, Rat, 78-5304
- Platelet Aggregation, 78-5304

Neoplasm Circulating Cells (cont'd)

- Thromboplastin, 78-5304
- Melanoma
- Cell Survival, 78-4923

Neoplasm Metastasis

- Adenocarcinoma
- Peutz-Jeghers Syndrome, 78-5311
- Angiosarcoma
- Lung, 78-5388
- Cervix Neoplasms
- Virus, Herpes Simplex 2, 78-4888
- Fibrosarcoma
- Lung, 78-5388
- Hemangiopericytoma
- Lung, 78-5388
- Hepatoma
- Actin, 78-5054
- Chromosomal Proteins, Non-Histone, 78-5054
- Lung, Mouse, 78-5334
- Morpholine, *N*-Nitroso-, 78-5054
- Thromboplastin, 78-5304
- Tubulin, 78-5054
- Ultrastructural Study, Rat, 78-5304

Kidney Neoplasms

- Adenocarcinoma, 78-5335
- Case Report, 78-5335
- Epidemiology, 78-5375
- Erythropoietin, 78-5336
- Vagina, 78-5335

Leiomyosarcoma

- Lung, 78-5388

Lung

- Animal Model, B16 Tumor, 78-4923

Lung Neoplasms

- Smoke Condensate, 78-5045

Melanoma

- Animal Model, B16 Tumor, 78-4923

Nephroblastoma

- Transplantation, Heterologous, 78-5320

Peutz-Jeghers Syndrome

- Case Report, 78-5311

Rhabdomyosarcoma

- Lung, 78-5388

Sarcoma

- Doubling Time, 78-5388
- Lung, 78-5388

Soft Tissue Neoplasms

- Histiocytoma, 78-5307

Stomach Neoplasms

- Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101

Neoplasm Regression, Spontaneous

- Benz(a)anthracene, 7,12-Dimethyl-12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036

Skin Neoplasms

- Histological Study, Mouse, 78-5036

Neoplasm Transplantation

- Astrocytoma
- Ultrastructural Study, 78-5089
- Ependymoma
- Virus, Polyoma, BK, 78-5263
- Heart Transplantation
- Epidemiology, 78-5384
- Hepatoma
- RNA, Neoplasm, 78-5069
- Lung Neoplasms
- Carcinoma, Epidermoid, 78-5399

Neoplasms (General and Unspecified)

- Carcinogen, Environmental
 - Epidemiology, Review, 78-4925
- Cholesterol
 - Epidemiology, 78-5349
 - Serum Levels, 78-5349
- Chromosome Aberrations
 - Hereditary Diseases, Review, 78-4931
 - Tumor Specificity, Review, 78-4931
- Drugs
 - Epidemiology, 78-5352
- Genetics
 - Epidemiology, Review, 78-4925
- Hereditary Diseases
 - Teratogenic/Mutagenic Activity, Review, 78-4930
- Oncogenic Viruses
 - Epidemiology, Review, 78-4925
- Radiation
 - Japan, Review, 78-4869
- Retinol
 - Antineoplastic Activity, 78-5146
 - Case-Control Study, 78-5146
 - Epidemiology, 78-5146, 78-5349
 - Serum Levels, 78-5349
- Virus, C-Type RNA Tumor
 - Isolation and Characterization, Review, 78-4897
- Virus, DNA Tumor
 - Isolation and Characterization, Review, 78-4897

Neoplasms, Connective Tissue

- Gynecologic Neoplasms
 - Radiation, Ionizing, 78-5178

Neoplasms, Experimental

- Adenosine Cyclic 3',5' Monophosphate
 - Concentration Levels, Review, 78-4937
- Antineoplastic Agents
 - Carcinogen, Chemical, 78-4807
- Asbestos
 - Trans-Species Extrapolation, 78-4805
- Carcinogen, Chemical
 - Bioassay Criteria, Review, 78-4801
 - Organ Culture, 78-4802
 - Species Difference, Review, 78-4807
- Dimethylamine, *N*-Nitroso-
 - Carcinogenic Activity, Review, 78-4821
- Ethylene, Chloro-
 - Age Factors, Review, 78-4831
 - Carcinogenic Activity, Review, 78-4831
 - Precancerous Conditions, Review, 78-4922
 - Species Difference, Review, 78-4831
- Ethylene, 1,1-Dichloro-
 - Age Factors, Review, 78-4831
 - Carcinogenic Activity, Review, 78-4831
 - Species Difference, Review, 78-4831
- Fiber Glass
 - Trans-Species Extrapolation, 78-4805
- Penicillium camemberti*
 - Review, 78-4837
- Steroids
 - Precancerous Conditions, Review, 78-4922
- Urea, Methyl Nitroso-
 - Carcinogenic Potential, Review, 78-4826

Neoplasms, Multiple Primary

- Adrenal Gland Neoplasms
 - Dyes, 78-4973
 - Occupational Hazard, 78-4973
- Bile Duct Neoplasms

Neoplasms, Multiple Primary (cont'd)

- Occupational Hazard, 78-4973
- Bladder Neoplasms
 - Occupational Hazard, 78-4973
- Breast Neoplasms
 - Genetics, 78-5368
- Cervix Neoplasms
 - Radiation, Ionizing, 78-5178
- Colonic Neoplasms
 - Case Report, 78-5386
 - Genetics, 78-5371
- Genetics
 - Case Report, 78-5371
- Gynecologic Neoplasms
 - Radiation, Ionizing, 78-5178
- Hepatoma
 - Dyes, 78-4973
 - Occupational Hazard, 78-4973
- Leukemia, Myeloblastic
 - Drug Therapy, 78-5298
 - Radiotherapy, 78-5298
- Lung Neoplasms
 - Occupational Hazard, 78-4973
- Pancreatic Neoplasms
 - Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081
- Uterine Neoplasms
 - Genetics, 78-5371

Neoplasms, Radiation-Induced

- Nuclear Reactors
 - Epidemiology, 78-5354
- Radiation, Ionizing
 - Dose-Response Study, Review, 78-4873
 - Epidemiology, Review, 78-4872, 78-4873, 78-4874
- Radiotherapy
 - Epidemiology, Review, 78-4870
- Ultraviolet Rays
 - Epidemiology, Review, 78-4873

Neoplastic Endocrine-Like Symptoms

- Pheochromocytoma
 - Genetics, 78-5308

Nephroblastoma

- Genetics
 - Aniridia, 78-5309
 - Cataract, 78-5309
- Hereditary Diseases
 - Review, 78-4929
- Transplantation, Heterologous
 - Mouse, Nude, 78-5320
 - Neoplasm Metastasis, 78-5320
- Twins
 - Case Report, 78-5309

Nervous System Neoplasms

- Benz(a)anthracene, 7,12-Dimethyl-
 - Carcinogenic Metabolite, 78-5026
 - Transplacental Carcinogenesis, 78-5026
- Ethylene, Chloro-
 - Epidemiology, 78-4832
 - Occupational Hazard, Review, 78-4832
- Neurilemmoma
 - Urea, Methyl Nitroso-, 78-5085
- Trace Elements
 - Blood Levels, 78-5085
- Urea, 1-Butyl-1-nitroso-
 - Age Factors, Rat, 78-5094
 - Transplacental Carcinogenesis, 78-5094

ous System Neoplasms (cont'd)

- Urea, Ethyl Nitroso-
Transplacental Carcinogenesis, Review, 78-4827
Ultrastructural Study, 78-5089
- Virus, Polyoma, JC
Carcinogenic Activity, Hamster, 78-5261

arilemmoma

- Eye Neoplasms
Virus, Polyoma, JC, 78-5261
- Nervous System Neoplasms
Urea, Methyl Nitroso-, 78-5085
- Urea, Ethyl Nitroso-
Carcinogenic Activity, Review, 78-4827
Dibutyl Cyclic AMP, 78-5093
Transplacental Carcinogenesis, 78-5093
- Urea, Methyl Nitroso-
Chromium, 78-5085
Iron, 78-5085
Rubidium, 78-5085
Selenium, 78-5085
Trace Elements, 78-5085
Zinc, 78-5085

roblastoma

- Hereditary Diseases
Review, 78-4929

rofibromatosis

- Hereditary Diseases
Review, 78-4929

uroglia

- Virus, Polyoma
Cell Transformation, Neoplastic, 78-5266

urospora crassa

- Actinomycin D
Lethal/Mutagenic Activity, 78-4947
- Aflatoxin B1
Lethal/Mutagenic Activity, 78-4947
Mutagenic Activity, 78-4967
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
DNA Repair, 78-4948
Lethal/Mutagenic Activity, 78-4947
Mutagenic Activity, 78-4948
- Methanesulfonic Acid, Ethyl Ester
Lethal/Mutagenic Activity, 78-4947
- Methanesulfonic Acid, Methyl Ester
Lethal/Mutagenic Activity, 78-4947
- 2-Oxetanone
Lethal/Mutagenic Activity, 78-4947
- 1*H*-Purin-2-amine
Lethal/Mutagenic Activity, 78-4947
- Quinoline, 4-Nitro, 1-Oxide
DNA Repair, 78-4948
Mutagenic Activity, 78-4948

us, Pigmented

- Cervix
Case Report, 78-5339
Ultrastructural Study, 78-5339

ikel

- Occupational Hazard
Epidemiology, Review, 78-4934

ikel Sulfide

- Tracheal Neoplasms
Carcinogenic Activity, Rat, 78-5119
Carcinoma, 78-5119

Nickel Sulfide (cont'd)

- Sarcoma, 78-5119

Nicotine

- Smoking
Inhalation Study, Hamster, 78-5043

Niridazole

- see 2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-

Nitric Acid

- Diet
Body Burden, 78-5052
Metabolism, Intestines, 78-5052
- Food Contamination
Quantitation Method, 78-5051
- Nitrogen
Metabolism, Intestines, 78-5052
- Stomach Neoplasms
Epidemiology, Review, 78-4818
Water Pollution, Review, 78-4818

Nitric Acid, Plutonium Salt

- Lung Neoplasms
Dog, 78-5185

p-Nitrobenzyl Chloride

- see Benzene, 1-(Chloromethyl)-4-nitro-

Nitrogen

- Nitric Acid
Metabolism, Intestines, 78-5052
- Nitrous Acid
Metabolism, Intestines, 78-5052

Nitrosamines

- Ames Test
Carcinogenic Activity, Correlation, 78-5097
- Anemia, Pernicious
Gastric Juice, 78-5330
- Aroclor 1254
Ames Test, 78-5097
- Barbituric Acid, 5-Ethyl-5-phenyl-
Ames Test, 78-5097
- Food Contamination
Carcinogenic Activity, Review, 78-4819
Carcinogenic Potential, Review, 78-4817
- Nitrous Acid
Carcinogenic Activity, Review, 78-4819

Nitroso Compounds

- Electrophiles
Carcinogenic Metabolite, Review, 78-4820
- Gastritis
Nitrosation, Review, 78-4818
- Mutagens
Feces, 78-5121
- Stomach Neoplasms
Nitrosation, Review, 78-4818

N-Nitrosobis(2-oxopropyl)amine

- see Dipropylamine, 2,2'-Dioxo-N-nitroso-

Nitrous Acid

- Anemia, Pernicious
Gastric Juice, 78-5330
- Antipyrine, 4-(Dimethylamino)-
Carcinogenic Activity, Review, 78-4819
- Ascorbic Acid
Metabolism, Intestines, 78-5052
- Diet
Body Burden, 78-5052

Nitrous Acid (cont'd)

- Metabolism, Intestines, 78-5052
- Dimethylamine, *N*-Nitroso-
 - Nitrosation Kinetics, 78-5077
 - Perchloric Acid, 78-5077
- Escherichia coli*
 - Mutagenic Activity, 78-5171
- Ethanol, *N*-Nitrosoiminodi-
 - Cutting Fluids, 78-5070
- Food Contamination
 - Carcinogenic Potential, Review, 78-4817
 - Quantitation Method, 78-5051
- Guanidine, Methyl-
 - Ames Test, 78-5103
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Saliva, 78-5103
- Nitrogen
 - Metabolism, Intestines, 78-5052
- Nitrosamines
 - Carcinogenic Activity, Review, 78-4819
- Pyridine, 2-(2-(Dimethylamino)ethyl)-2-thenylamino)-
 - Carcinogenic Activity, Review, 78-4819
- Stomach Neoplasms
 - Anemia, Pernicious, 78-5330
 - Bacteria, Review, 78-4818
 - Epidemiology, Review, 78-4818
- Ultraviolet Rays
 - DNA Repair, 78-5171

Nitrous Acid, Sodium Salt

- Dimethylamine, *N*-Nitroso-
 - Antipyrine, 4-(Dimethylamino)-, 78-5076
 - Ethanol, 2-(Dimethylamino)-, 78-5076
 - Ethylamine, 2-(Diphenylmethoxy)-*N,N*-dimethyl-
78-5076
 - Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-, 78-5076
 - Sulfanilamide, *N*-(3,4-Dimethyl-4-isoxazolyl)-
78-5076
 - Tetracycline, 78-5076
- Ethylamine, *N*-Methyl-*N*-nitroso-
 - Acetamide, *N*-Methyl-, 78-5080
 - Deuterium Labeled Analogs, Synthesis, 78-5080
 - Isocyanic Acid, Ethyl Ester, 78-5080
- Sorbic Acid
 - Food Contamination, 78-5124
 - Mutagenic Reaction Products, 78-5124
- Urinary Tract Infections
 - Proteus mirabilis*, 78-5076

Norgestrel, Chloroethynyl-

- Mammary Neoplasms, Experimental
 - Hyperplasia, 78-5132

Nose Neoplasms

- Adenocarcinoma
 - Ethanol, *N*-Nitrosoiminodi-, 78-4823
- Carcinoma
 - Epidemiology, USSR, 78-5379
- Diethylamine, *N*-Nitroso-
 - Carcinogenic Activity, 78-5068
 - Ultrastructural Study, 78-5068
- Ethanol, *N*-Nitrosoiminodi-
 - Carcinogenic Potential, Review, 78-4823
- Maxillary Sinus
 - Epidemiology, USSR, 78-5379
- Melanoma
 - Epidemiology, USSR, 78-5379
- Plasmacytoma

Nose Neoplasms (cont'd)

- Epidemiology, USSR, 78-5379
- Sarcoma
 - Epidemiology, USSR, 78-5379

Nucleic Acids

- Aflatoxin B1
 - Carcinogenic Metabolite, Review, 78-4816
 - Hypophysectomy, 78-4960
 - Liver, Rat, 78-4960
- Benzo(a)pyrene
 - Binding, Review, 78-4850
- Carcinogen, Chemical
 - Metabolism, Review, 78-4804
- Dimethylamine, *N*-Nitroso-
 - Methylation, Review, 78-4820
- Ethylene, Chloro-
 - Liver, Rat, 78-5110
- Methane, Azoxy-
 - Intestinal Mucosa, 78-5159
- Methanol, (Methyl-*ONN*-azoxy)-
 - Methylation, Review, 78-4820

Nucleoproteins

- Virus, SV40
 - DNA, Viral, 78-5254

Obesity

- Breast Neoplasms
 - Epidemiology, 78-5367

Occupational Hazard

- Adrenal Gland Neoplasms
 - Neoplasms, Multiple Primary, 78-4973
- Anesthetics
 - Abnormalities, 78-4858
 - Carcinogenic Potential, Review, 78-4858
 - Epidemiology, Review, 78-4857, 78-4858
- Arsenic
 - Epidemiology, Review, 78-4934
- Asbestos
 - Epidemiology, 78-5377
 - Epidemiology, Review, 78-4842
- Beryllium
 - Epidemiology, Review, 78-4934
- Bile Duct Neoplasms
 - Neoplasms, Multiple Primary, 78-4973
- Bladder Neoplasms
 - Carcinoma, Transitional Cell, 78-4973
 - Dyes, 78-4973
 - Neoplasms, Multiple Primary, 78-4973
- 1,3-Butadiene, 2-Chloro-
 - Epidemiology, Review, 78-4834
- Cadmium
 - Epidemiology, Review, 78-4934
- Carcinogen, Chemical
 - Epidemiology, Review, 78-4856
- Carcinogen, Environmental
 - Epidemiology, Review, 78-4925
- Chromium
 - Epidemiology, Review, 78-4934
- Ethylene, Chloro-
 - Epidemiology, Review, 78-4834
- Hepatoma
 - Neoplasms, Multiple Primary, 78-4973
- Iron
 - Epidemiology, Review, 78-4934
- Lead
 - Epidemiology, Review, 78-4934

Occupational Hazard (cont'd)**Lung Neoplasms**

- Case Report, 78-5326
- Epidemiology, 78-5344
- Epidemiology, Review, 78-4841
- Neoplasms, Multiple Primary, 78-4973

Nickel

- Epidemiology, Review, 78-4934

Pleural Neoplasms

- Mesothelioma, 78-5385

Polychlorobiphenyl Compounds

- Epidemiology, Review, 78-4829

Radiation, Ionizing

- Dose-Response Study, Review, 78-4871

Uranium

- Epidemiology, 78-5354

Chratoxin A**Food Contamination**

- Quantitation Method, 78-4959

Herbs

- Drug Contamination, 78-4956

Octadecenoic Acid**Gastrointestinal Neoplasms**

- Papilloma, 78-5120

oleic Acid

- see 9-Octadecenoic Acid

ligodendroglioma**Brain Neoplasms**

- Urea, 1,3-Dimethyl-1-nitroso-, 78-5096

ncogenic Viruses**Aryl Hydrocarbon Hydroxylases**

- Genetics, Review, 78-4927

Gynecologic Neoplasms

- Epidemiology, Review, 78-4895

Leukemia

- Epidemiology, Review, 78-4895
- Horizontal Transmission, 78-4899

Lymphoma

- Epidemiology, Review, 78-4895

Lymphosarcoma

- Horizontal Transmission, 78-4899

Neoplasms

- Epidemiology, Review, 78-4925

Sarcoma

- Epidemiology, Review, 78-4895

pium**Esophageal Neoplasms**

- Epidemiology, 78-5049

Microsomes, Liver

- Ames Test, 78-5049

rnithine**Ames Test**

- Pyrolysis Products, 78-5122

Mutagens

- Inactivation, Fruit/Vegetable Juices, 78-5122
- Pyrolysis Products, 78-5122

torhinolaryngologic Neoplasms**Carcinoma**

- Epidemiology, USSR, 78-5379

Melanoma

- Epidemiology, USSR, 78-5379

Plasmacytoma

- Epidemiology, USSR, 78-5379

Otorhinolaryngologic Neoplasms (cont'd)**Sarcoma**

- Epidemiology, USSR, 78-5379

Ouabain

- 12-*O*-Tetradecanoylphorbol-13-acetate
- Ion Transport, Fibroblasts, 78-5042

Ovarian Cysts**Hepatoma**

- Genetics, 78-5333

Ovarian Neoplasms**Adenocarcinoma**

- Epidemiology, Dog, 78-5357

Age Factors

- Histological Study, Dog, 78-5357

Carcinoma

- Transplantation Immunology, 78-5293

Cystadenocarcinoma

- Epidemiology, Dog, 78-5357

Cystadenoma

- Benz(a)anthracene, 7,12-Dimethyl-, 78-5030

- Epidemiology, Dog, 78-5357

Epidemiology

- Indonesia, 78-5345

Genetics

- Epidemiology, 78-5364

Granulosa Cell Tumor

- Epidemiology, Dog, 78-5357

1,2,3-Oxadiazolium, 3-Methyl-5-amino-, Chloride

- Acetonitrile, 2-(*N*-Methyl-*N*-nitroso)amino-
- Reversible Interconversion, 78-5053

Oxamniquine**Antischistosomal Drugs**

- Mutagenic Activity, Review, 78-4845

1,2-Oxathiolane, 2,2-Dioxide**Ames Test**

- Carcinogenic Potential, Review, 78-4815

2-Oxetanone**DNA**

- Phosphodiester, 78-5108

Neurospora crassa

- Lethal/Mutagenic Activity, 78-4947

Phosphodiester

- Isolation and Characterization, 78-5108

Salmonella typhimurium

- Mutagenic Activity, 78-4946

5'-Thymidylic Acid

- Phosphodiester, 78-5108

Oxidoreductases**Acrylonitrile**

- Maleic Acid, Diethyl Ester, 78-5111

Cadmium

- Enzymatic Activity, 78-5118

Diethylamine, *N*-Nitroso-

- Mutagenic Activity, 78-4951

Hydroxylamine

- Microsomes, Liver, 78-4989

2-Imidazolidinethione

- Metabolism, 78-5112

Isonicotinamide, 2-Ethylthio-

- Metabolism, 78-5112

Propionitrile

- Enzymatic Activity, 78-5111

Urea, Thio-

- Oxidoreductases (cont'd)**
Metabolism, 78-5112
- Oxidoreductases, *N*-Demethylating**
Cadmium
Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5118
Enzymatic Activity, 78-5118
Dimethylamine, *N*-Nitroso-
Acetone, 78-5075
Aroclor 1254, 78-5075
Dose-Response Study, 78-5075
- Oxirane, 9-Anthryl-**
Ames Test
Dose-Response Study, 78-4984
Carcinogenic Metabolite
Synthesis, 78-5035
Virus, SV40
DNA, Binding, 78-4984
- Oxirane, 7-Benzanthryl-**
Ames Test
Dose-Response Study, 78-4984
Carcinogenic Metabolite
Synthesis, 78-5035
Virus, SV40
DNA, Binding, 78-4984
- Oxirane, 6-Chrysenyl-**
Ames Test
Dose-Response Study, 78-4984
Carcinogenic Metabolite
Synthesis, 78-5035
Virus, SV40
DNA, Binding, 78-4984
- Oxirane, 9,10-Dimethyl-2-anthryl-**
Carcinogenic Metabolite
Synthesis, 78-5035
- Oxirane, 9-Methyl-10-anthryl-**
Ames Test
Dose-Response Study, 78-4984
Virus, SV40
DNA, Binding, 78-4984
- Oxirane, 10-Methyl-9-anthryl-**
Carcinogenic Metabolite
Synthesis, 78-5035
- Oxirane, 1-Naphthyl-**
Carcinogenic Metabolite
Synthesis, 78-5035
- Oxirane, 2-Naphthyl-**
Ames Test
Dose-Response Study, 78-4984
Carcinogenic Metabolite
Synthesis, 78-5035
Virus, SV40
DNA, Binding, 78-4984
- Oxirane, 9-Phenanthryl-**
Ames Test
Dose-Response Study, 78-4984
Carcinogenic Metabolite
Synthesis, 78-5035
Virus, SV40
DNA, Binding, 78-4984
- Oxirane, 1-Pyrenyl-**
Ames Test
Dose-Response Study, 78-4984
- Oxirane, 1-Pyrenyl- (cont'd)**
Carcinogenic Metabolite
Synthesis, 78-5035
Virus, SV40
DNA, Binding, 78-4984
- Oxygenases**
Acetamide, *N*-Fluoren-2-yl-
Metabolism, Review, 78-4854
Benzo(a)pyrene
Enzyme Activation, Review, 78-4854
Coumarin, 7-Ethoxy-
Metabolism, Review, 78-4854
- Pancreas**
Cholanthrene, 3-Methyl-
Histofluorescence, Rat, 78-4991
Precancerous Conditions, 78-4991
- Pancreatic Neoplasms**
Adenocarcinoma
Animal Model, Rat, 78-4991
Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081
Epidemiology, Sri Lanka, 78-5376
Adenoma
Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081
Carcinoma, Ductal
Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081
Carcinoma, Epidermoid
Epidemiology, Sri Lanka, 78-5376
Dipropylamine, 2,2'-Dioxo-*N*-nitroso-
Dose-Response Study, Hamster, 78-5081
Islet Cell Tumor
Epidemiology, Sri Lanka, 78-5376
Virus, Polyoma, BK, 78-5263
Neoplasms, Multiple Primary
Dipropylamine, 2,2'-Dioxo-*N*-nitroso-, 78-5081
Radiation, Ionizing
Epidemiology, Review, 78-4874
Smoking
Epidemiology, Review, 78-4912
Virus, Polyoma, BK
Carcinogenic Activity, Hamster, 78-5263
- Papilloma**
Benz(a)anthracene, 7,12-Dimethyl-
12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
Gastrointestinal Neoplasms
Cholesterol, 78-5120
Corn Oil, 78-5120
Dietary Fats, 78-5120
Linoleic Acid, 78-5120
Monoolein, 78-5120
9-Octadecenoic Acid, 78-5120
Laryngeal Neoplasms
Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101
Lung Neoplasms
1-Propanol, 2,3-Dibromo-, Phosphate, 78-5167
Respiratory Tract Neoplasms
Benzo(a)pyrene, 78-5004, 78-5006
Skin Neoplasms
Benz(a)anthracene, 78-5022
Benz(a)anthracene, 7,12-Dimethyl-, 78-5036
Stomach Neoplasms
Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101
Quinoline, 8-Nitro-, 78-5099
- Paraganglioma, Nonchromaffin**
Carotid Body
Case Report, 78-5310

ganglioma, Nonchromaffin (cont'd)

- Genetics
- Case Report, 78-5310
- Neoplasm Recurrence, 78-5310

lysis

- Virus, Murine Leukemia
- Epidemiology, Wild Mouse, Review, 78-4883

sitic Diseases

- Burkitt's Lymphoma
- Epidemiology, Review, 78-4913
- Hodgkin's Disease
- Epidemiology, Review, 78-4913
- Lymphoma
- Epidemiology, Review, 78-4913

icillium camemberti

- Mycotoxins
- Review, 78-4837
- Neoplasms, Experimental
- Review, 78-4837

ic Ulcer

- Stomach Neoplasms
- Epidemiology, 78-5329

ide Hydrolases

- Carcinogen, Chemical
- Filter Implant Test, 78-4943

ides

- Virus, Moloney Murine Sarcoma
- Growth Factors, Isolation and Characterization
- 78-5220

aloric Acid

- Dimethylamine, *N*-Nitroso-
- Nitrous Acid, 78-5077

dic Acid, Sodium Salt

- Asbestos
- Macrophages, 78-5284

oneal Neoplasms

- Asbestos
- Epidemiology, Review, 78-4842
- Leiomyoma
- Case Report, 78-5306
- Mesothelioma
- Epidemiology, 78-5377

cides

- Benzo(a)pyrene
- Liver, Rat, 78-5014

cides

- Lung Neoplasms
- Epidemiology, Review, 78-4835
- Pulmonary Fibrosis
- Epidemiology, Review, 78-4835

-Jeghers Syndrome

- Adenocarcinoma
- Neoplasm Metastasis, 78-5311
- Neoplasm Metastasis
- Case Report, 78-5311

Polyps

- Cell Transformation, Neoplastic, 78-5311

ngelial Neoplasms

- Epidemiology
- Geographic Factors, 78-5344
- Ethyl Alcohol

Pharyngeal Neoplasms (cont'd)

- Epidemiology, 78-5344
- Smoking
- Epidemiology, 78-5344

Phenanthrene

- Ames Test
- Dihydrodiols, 78-4855
- Carcinogenic Metabolite
- Fossil-Fuel Combustion Products, 78-5032

Phenethylamine, α -Methyl-

- Carcinogenic Potential
- Review, 78-4843

Phenol

- Air Pollution
- Fossil-Fuel Combustion Products, 78-5032

Phenol, (1,1-Dimethylethyl)-4-methoxy-

- Carcinogen, Chemical
- Inhibitors, Review, 78-4806

Phenothiazine

- Carcinogen, Chemical
- Inhibitors, Review, 78-4806

***o*-Phenylenediamine, 4-Nitro-**

- Ames Test
- Carcinogenic Potential, Review, 78-4809

***p*-Phenylenediamine, 2-Nitro-**

- Ames Test
- Carcinogenic Potential, Review, 78-4809

Pheochromocytoma

- Adenomatosis, Familial Endocrine
- Case Report, 78-5317
- Chromosomes, Human, 13-15, 78-5317
- Histocompatibility Antigens, 78-5317
- Adrenal Gland Neoplasms
- Genetics, 78-5308
- Angiomatosis
- Genetics, 78-5308
- Genetics
- Case Report, 78-5308
- Hereditary Diseases
- Review, 78-4929
- Neoplastic Endocrine-Like Symptoms
- Genetics, 78-5308

Phorbol 13-Acetate

- 12-*O*-Tetradecanoylphorbol-13-acetate
- Metabolism, Fibroblasts, 78-5038

4 α -Phorbolol 12,13-Didecanoate

- Rubidium
- Ion Transport, Fibroblasts, 78-5042

4 β -Phorbolol 12,13-Didecanoate

- Rubidium
- Ion Transport, Fibroblasts, 78-5042

Phosphine Oxide, Diethoxymorpholino-

- Microsomes, Liver
- Carcinogenic Activity, 78-5150

Phosphine Oxide, Tripiperidino-

- Microsomes, Liver
- Carcinogenic Activity, 78-5150

Phosphine Oxide, Tris(1-aziridinyl)-

- Lymphocytes
- Chromosome Aberrations, 78-5157

Phospholipids

- Anthracosilicosis
 - Scleroproteins, 78-4971
- Asbestos
 - Lung, Rat, 78-4971
- Asbestosis
 - Scleroproteins, 78-4971
- Coal
 - Lung, Rat, 78-4971
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-5192

Phosphonium, Tetrakis(hydroxymethyl)-, Chloride

- Lung Neoplasms
 - Carcinogenic Potential, Rat, 78-5167

Phosphoric Acid

- Prostaglandins E
 - Ion Transport, Fibroblasts, 78-5042
 - Rubidium, 78-5042
- Prostaglandins F
 - Ion Transport, Fibroblasts, 78-5042
- Rubidium
 - Ion Transport, Fibroblasts, 78-5042

Phosphoric Triamide, Hexamethyl-

- Enzyme Activation
 - Carcinogenic Analog, 78-5150
- Microsomes, Liver
 - Carcinogenic Activity, 78-5150

Phosphorothioic Triamide, Hexamethyl-

- Microsomes, Liver
 - Carcinogenic Activity, 78-5150

Phosphorothioic Trimorpholide

- Microsomes, Liver
 - Carcinogenic Activity, 78-5150

Phosphorus Triamide, Hexamethyl-

- Microsomes, Liver
 - Carcinogenic Activity, 78-5150

Phosphotransferases

- Virus, Rous Sarcoma
 - Viral Proteins, 78-5188

Phthalimide, N-((Trichloromethyl)thio)-

- Chromosome Aberrations
 - Dose-Response Study, 78-5163
 - Raji, Jeff Lymphoid Cells, 78-5163

Physostigmine, Sulfate

- Hydrazine, 1,2-Dimethyl-
 - DNA Replication, 78-5064
- Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
 - DNA Replication, 78-5064

Piperidine, 1-Nitroso-

- Diet
 - Metabolism, Rat, 78-5073
- Intestine, Small
 - Metabolism, Rat, 78-5073
- Lipids
 - Metabolism, Rat, 78-5073

Pituitary Gland

- Aniline, 4,4'-Oxydi-
 - Hyperplasia, 78-5126

Pituitary Neoplasms

- Aniline, 4,4'-Oxydi-
 - Precancerous Conditions, 78-5126

Pituitary Neoplasms (cont'd)

- Estrogens
 - Growth Factors, 78-5134
 - Mitosis, 78-5134
- Fusarium sporotrichioides*
 - Toxins, 78-4954
- Meningioma
 - Case Report, 78-5318
- Thyroid Hormones
 - Estrogens, 78-5134

Placental Lactogen

- Breast Neoplasms
 - DNA Replication, 78-5392
 - Precancerous Conditions, 78-5392

Plant Agglutinins

- Virus, Herpes Simplex 2
 - Immunity, Cellular, 78-5243
 - Lymphocyte Transformation, 78-5243
- Virus, Turkey Herpes
 - T-Lymphocytes, 78-5196

Plasmacytoma

- Myeloma Proteins
 - Graft Rejection, 78-5289
- Nose Neoplasms
 - Epidemiology, USSR, 78-5379
- Otorhinolaryngologic Neoplasms
 - Epidemiology, USSR, 78-5379
- Thymus Gland
 - Graft Rejection, 78-5289

Plasmids

- Methanesulfonic Acid, Methyl Ester
 - Ames Test, 78-4950
 - Mutagenic Activity, 78-4950

Plasminogen Activators

- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Carcinogenic Activity, 78-5040
 - Fibroblasts, Chick Embryo, 78-5040
 - Virus, Rous Sarcoma, 78-5040
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-5040

Pleural Neoplasms

- Asbestos
 - Epidemiology, Review, 78-4842
- Mesothelioma
 - Epidemiology, 78-5377, 78-5385
 - Occupational Hazard, 78-5385

Plutonium

- Air Pollution
 - Carcinogenic Potential, Review, 78-4868
- Bone Neoplasms
 - Carcinogenic Potential, Review, 78-4868
- Ecosystem Distribution
 - Mathematical Model, 78-5184
- Lung Neoplasms
 - Carcinogenic Potential, Review, 78-4868
- Soil Pollutants
 - Ecosystem Distribution, 78-5184

Polybromobiphenyl Compounds

- Toxicology
 - Animals, Review, 78-4828

Polychlorinated Terphenyl

- see* Kanechlor C

ychlorobiphenyl Compounds

- Carcinogen, Chemical
Inhibitors, Review, 78-4806
- Environmental Hazard
Epidemiology, Review, 78-4829
- Liver Neoplasms
Carcinogenic Potential, Review, 78-4829
- Metabolism
Rat, 78-4980
- Occupational Hazard
Epidemiology, Review, 78-4829
- Structure-Activity Relationship
Carcinogenic Potential, Review, 78-4829
- Toxicology
Animals, Review, 78-4828

ycyclic Hydrocarbons

- Aroclor 1254
Ames Test, 78-4938
- Aryl Hydrocarbon Hydroxylases
Genetics, Review, 78-4927
- Carcinogen, Chemical
Inhibitors, Review, 78-4806
- DNA, Binding
Mutagenic Activity, 78-4997
- Drosophila*
Mutagenic Activity, Review, 78-4812

yps

- Colonic Neoplasms
Adenoma, 78-5313
Carcinoma, 78-5313
Case Report, 78-5314
Genetics, 78-5313, 78-5314
- Hereditary Diseases
Review, 78-4929
- Peutz-Jeghers Syndrome
Cell Transformation, Neoplastic, 78-5311
- Uterine Neoplasms
Precancerous Conditions, Review, 78-4921

yribonucleotides

- Dibenz(a,e)aceanthrylene
Microsomes, Liver, 78-5033
- Quinoline, 4- (Hydroxyamino)-, 1-Oxide
Binding, 78-5100
- Quinoline, 4-Nitro-, 1-Oxide
Binding, 78-5100
- Virus, Avian Myeloblastosis
Ribonuclease, 78-5203
- Virus, Rauscher Murine Leukemia
Ribonuclease, 78-5203

phyrins

- Acetic Acid, Lead Salt
Metabolism, Rat, 78-5116
- Kidney Neoplasms
Acetic Acid, Lead Salt, 78-5116

cancerous Conditions

- Bladder Neoplasms
Carcinoma, 78-5375
- Bone Neoplasms
Aniline, 4,4'-Oxydi-, 78-5126
- Brain Neoplasms
Urea, Ethyl Nitroso-, 78-5091
- Breast Neoplasms
DNA Replication, 78-5392
Insulin, 78-5392
Mammographic Classification, 78-5369

Precancerous Conditions (cont'd)

- Placental Lactogen, 78-5392
- Cell Transformation, Neoplastic
Review, 78-4924
- Cervix Neoplasms
Carcinoma In Situ, 78-5391
Dysplasia, 78-5340
Histological Study, 78-5340
Virus, Herpes Simplex 2, 78-5242
- Cholanthrene, 3-Methyl-
Pancreas, 78-4991
- Colonic Neoplasms
Genetics, 78-5313
Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5059
- Esophageal Neoplasms
Esophageal Reflux, 78-5323
Metaplasia, 78-5323
- Gastric Mucosa
Metaplasia, 78-5328
- Hepatoma
Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-5127
Histological Study, Mouse, 78-5334
Kanechlor C, 78-4979
- Intestinal Neoplasms
Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5059
- Kidney Neoplasms
Carcinoma, 78-5375
- Leukemia
Chromosome Abnormalities, 78-5204
Reverse Transcriptase, 78-5204
Virus-Like Particles, 78-5204
- Leukemia, Myeloblastic
Antigens, Neoplasm, 78-5397
Blood Platelets, 78-5397
Chromosome Aberrations, 78-5296
Glycoproteins, 78-5397
- Liver Neoplasms
Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5325
Contraceptives, Oral, 78-5133
- Metaplasia
Histological Study, 78-5323
- Mouth Neoplasms
Benz(a)anthracene, 7,12-Dimethyl-, 78-5028
- Myelofibrosis
Glycoproteins, 78-5397
- Myeloid Metaplasia
Glycoproteins, 78-5397
- Pituitary Neoplasms
Aniline, 4,4'-Oxydi-, 78-5126
- Reverse Transcriptase
Blood Platelets, 78-5204
- Skin Neoplasms
Glycerol, 78-5047
Smoke Condensate, 78-5047
- 4,4'-Stilbenediol, α,α' -Diethyl-
Epidemiology, 78-5370
- Stomach Neoplasms
Anemia, Pernicious, 78-5330
Metaplasia, 78-5328
- Thyroid Neoplasms
Aniline, 4,4'-Oxydi-, 78-5126
- Urogenital Neoplasms
Aniline, 4,4'-Oxydi-, 78-5126
- Uterine Neoplasms
Epidemiology, Review, 78-4862
- Virus, Bovine Leukemia

- Precancerous Conditions (cont'd)**
 Epidemiology, 78-4886
 Virus, Gross Murine Leukemia
 Autoimmune Diseases, 78-5280
 Virus, Herpes Simplex 2
 Antigens, Viral, 78-5242
 Virus, Marek's Disease Herpes
 Immunosuppression, 78-5196
 Virus, Radiation Leukemia
 Bone Marrow Cells, 78-5224
 T-Lymphocytes, 78-5295
 Radiation, Ionizing, 78-5224
 Thymus Gland, 78-5224
- Pregn-4-en-20-one, 17-(Acetyloxy)-6 α -methyl-**
see Anagestone Acetate
- Pregn-4-ene-3,20-dione, 17-(Acetyloxy)-6 α -methyl-**
 Acetamide, *N*-Fluoren-2-yl
 Metabolism, Kidney, 78-5078
 Dimethylamine, *N*-Nitroso-
 Metabolism, Kidney, 78-5078
- Pregnancy**
 Breast Neoplasms
 Epidemiology, 78-5365
 Granulosa Cell Tumor
 Case Report, 78-5321
 Liposarcoma
 Case Report, 78-5390
 Lymphosarcoma
 Case Report, 78-5390
 Myosarcoma
 Case Report, 78-5390
 Rhabdomyosarcoma
 Case Report, 78-5390
 Sarcoma
 Case Report, 78-5390
 Sarcoma, Osteogenic
 Case Report, 78-5390
- Procarbazine**
see p-Toluidamide, *N*-Isopropyl- α -(2-methylhydrazino)-
- Progesterone**
 Gynecologic Neoplasms
 Neoplasm Regression, Review, 78-0486
 Uterine Neoplasms
 Turner's Syndrome, 78-5342
- Prolactin**
 Mammary Neoplasms, Experimental
 Estrogens, 78-5134
- Proline**
 Virus, SV40
 Amino Acids, 78-5259
- Proline, 4-Hydroxy-1-nitroso-**
 Food Contamination
 Quantitation Method, 78-5057
- Proline, 1-Nitroso-**
 Food Contamination
 Quantitation Method, 78-5057
- Propane, 1,2-Epoxy-3-(*p*-nitrophenoxy)-**
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Transferases, 78-5016
 Benzo(a)pyrene
 Transferases, 78-5016
 Cholanthrene, 3-Methyl-
- Propane, 1,2-Epoxy-3-(*p*-nitrophenoxy)- (cont'd)**
 Transferases, 78-5016
- Propane, 1,2-Epoxy-3,3,3-trichloro-**
 Dibenz(a,e)acanthrylene
 DNA, Binding, 78-5033
- 1-Propanol, 2,3-Dibromo-**
 1-Propanol, 2,3-Dibromo-, Phosphate
 Mutagenic Metabolite, Urine, 78-5166
- 1-Propanol, 2,3-Dibromo-, Phosphate**
 Ames Test
 Carcinogenic Potential, Review, 78-4809
 Hepatoma
 Carcinogenic Potential, Rat, 78-5167
 Kidney Neoplasms
 Adenocarcinoma, 78-5167
 Dose-Response Study, Rat, 78-5167
 Lung Neoplasms
 Dose-Response Study, Rat, 78-5167
 Papilloma, 78-5167
 Mouth Neoplasms
 Carcinogenic Potential, Rat, 78-5167
 1-Propanol, 2,3-Dibromo-
 Mutagenic Metabolite, Urine, 78-5166
 Skin Absorption
 Mutagenic Metabolite, Urine, 78-5166
 Skin Neoplasms
 Carcinogenic Potential, Rat, 78-5167
 Stomach Neoplasms
 Carcinogenic Potential, Rat, 78-5167
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-**
 Thyroid Neoplasms
 Adenocarcinoma, 78-5079
 Adenoma, 78-5079
 Carcinoma, 78-5079
 Rat, 78-5079
- 1-Propanone, 2-Methyl-1,2-di-3-pyridyl-, Tartrate**
 Microsomes, Liver
 Aryl Hydrocarbon Hydroxylases, 78-5011
- β -Propiolactone**
see 2-Oxetanone
- Propionamide, Dimethyl-3,3'-dithiobis-, Dihydrochloride**
 Virus, C-Type RNA Tumor
 Viral Proteins, 78-5210
 Virus, Murine Leukemia
 Viral Proteins, 78-5210
- Propionic Pentafluoro-, Anhydride**
 Urine
 Quantitation Method, 78-4972
 Water Pollution
 Carcinogenic Metabolite, 78-4972
- Propionitrile**
 Lactate Dehydrogenase
 Enzymatic Activity, 78-5111
 Oxidoreductases
 Enzymatic Activity, 78-5111
- Propionitrile, 3-(*N*-Methyl-*N*-nitroso)amino-**
 Hydrochloric Acid
 Reaction Product, 78-5053
- Propylamine, *N*-(2-Hydroxypropyl)-*N*-nitroso-**
 Dipropylamine, *N*-Nitroso-
 Carcinogenic Metabolite, Review, 78-4824
 Liver Neoplasms

N-Nitrosodimethylamine, N-(2-Hydroxypropyl)-N-nitroso- (cont'd)
 Carcinogenic Metabolite, Review, 78-4824

Respiratory Tract Neoplasms
 Carcinogenic Metabolite, Review, 78-4824

Serine
 Esophageal Neoplasms
 Dimethylamine, *N*-Nitroso-, 78-5067

Liver Neoplasms
 Dimethylamine, *N*-Nitroso-, 78-5067

Straglandins E
 Phosphoric Acid
 Ion Transport, Fibroblasts, 78-5042

Rubidium
 Phosphoric Acid, 78-5042

Straglandins F
 Phosphoric Acid
 Ion Transport, Fibroblasts, 78-5042

Rubidium
 Ion Transport, Fibroblasts, 78-5042

Static Neoplasms
 Cadmium
 Epidemiology, Review, 78-4934

Cholelithiasis
 Epidemiology, 78-5337

Diet
 Epidemiology, Review, 78-4912

Epidemiology
 Review, 78-4908

Hyperplasia
 5 α -Andostan-3-one, 17 β -Hydroxy-, 78-4920
 Androgens, 78-4920
 Androstenediol, 78-4920
 Testosterone, 78-4920

Protein Kinase
 Butyric Acid, 2-Amino-4-(ethylthio)-
 Liver, Rat, 78-5398

Methionine
 Liver, Rat, 78-5398

Virus, Rous Sarcoma
 Cell Transformation, Neoplastic, 78-5188
 RNA, Viral, 78-5188

Yeins
Agrobacterium tumefaciens
 Acetic Acid, (Ethylenedinitrilo)tetra-, 78-5400
 Isolation and Characterization, 78-5400
 Muramidase, 78-5400
 Plasmids, 78-5400
 Tumorigenicity, 78-5400

Cell Transformation, Neoplastic
 Cell Membrane, Review, 78-4932

Virus, Adeno 2
 Cell Membrane, Review, 78-4902

Virus, Rous Sarcoma
 Cell Adhesion, Review, 78-4932
 Cell Membrane, 78-5195
 Temperature Sensitive Mutants, 78-5195

Pseudomonas mirabilis
 Urinary Tract Infections
 Dimethylamine, *N*-Nitroso-, 78-5076
 Nitrous Acid, Sodium Salt, 78-5076

Urinary Fibrosis
 Pesticides
 Epidemiology, Review, 78-4835

1*H*-Purin-2-amine
Neurospora crassa
 Lethal/Mutagenic Activity, 78-4947

Purine, 2-Amino-6-methoxy-
 Methylamine, *N*-Nitroso-
 DNA, Alkylation, 78-5106
 Radioimmunoassay, 78-5106

Urea, Ethyl Nitroso-
 DNA, Review, 78-4827

Purine-6-thiol, 9- β -D-Arabinofuranosyl-
 Lung Neoplasms
 Carcinoma, Epidermoid, 78-5399

Purine-6(1*H*)-thione, 2-Amino-
 Lung Neoplasms
 Carcinoma, Epidermoid, 78-5399

Putrescine
 Benzo(a)pyrene
 Carcinogenic Activity, 78-5005

Skin Neoplasms
 Benzo(a)pyrene, 78-5005
 Urine, Mouse, 78-5005

Pyrene
 Air Pollution
 Fossil-Fuel Combustion Products, 78-5032

Carcinogenic Metabolite
 Fossil-Fuel Combustion Products, 78-5032

Pyridine, 2-((2-(Dimethylamino)ethyl)-2-thenylamino)-
 Nitrous Acid
 Carcinogenic Activity, Review, 78-4819

1*H*-Pyrrole-2,5-dione, 1-Ethyl-
 Virus, RNA Tumor
 Reverse Transcriptase, 78-5216

Pyrrolidine, 3-Hydroxy-1-nitroso-
 Food Contamination
 Quantitation Method, 78-5058

Pyrrolidine, 1-Nitroso-
 Diet
 Metabolism, Rat, 78-5073

Intestine, Small
 Metabolism, Rat, 78-5073

Lipids
 Metabolism, Rat, 78-5073

Quartz
 Lung Neoplasms
 Adenocarcinoma, 78-5326

Quinaldic Acid, 4,8-Dihydroxy-
 Bladder Neoplasms
 Tryptophan, 78-5338

Quinaldic Acid, 4-Hydroxy-
 Bladder Neoplasms
 Tryptophan, 78-5338

Quinoline
 Hemangioendothelioma
 Carcinogenic Activity, Rat, 78-5099

Hepatoma
 Carcinogenic Activity, Rat, 78-5099

Liver Neoplasms
 Hemangioendothelioma, 78-5099

Quinoline, 4-Amino-, 1-Oxide
 Carcinogenic Potential

Quinoline, 4-Amino-, 1-Oxide (cont'd)

- Diploid Cells, 78-5105
- Cell Transformation, Neoplastic Fibroblasts, 78-5098

**Quinoline, 2-Chloro-
Diet**

- Carcinogenic Activity, Rat, 78-5099

Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-

- Anemia, Aplastic
Case Report, 78-5147
- Dimethylamine, *N*-Nitroso-
Nitrous Acid, Sodium Salt, 78-5076
- Leukemia, Myeloblastic
Anemia, Aplastic, 78-5147
Case Report, 78-5147

Quinoline, 6-Ethoxy-1,2-Dihydro-2,2,4-trimethyl-

- Carcinogen, Chemical
Inhibitors, Review, 78-4806

Quinoline, 4-(Hydroxyamino)-, 1-Oxide

- Free Radicals
Electron Spin Resonance Study, 78-5100
- Polyribonucleotides
Binding, 78-5100

**Quinoline, 8-Methyl-
Diet**

- Carcinogenic Activity, Rat, 78-5099

Quinoline, 8-Nitro-

- Stomach Neoplasms
Carcinogenic Activity, Rat, 78-5099
Carcinoma, Epidermoid, 78-5099
Papilloma, 78-5099

Quinoline, 4-Nitro-, 1-Oxide

- Cell Transformation, Neoplastic
Diploid Cells, 78-5105
Fibroblasts, 78-5098
- DNA Repair
UV-Mimetic Compounds, 78-5170
- Free Radicals
Electron Spin Resonance Study, 78-5100
- Neurospora crassa*
DNA Repair, 78-4948
Mutagenic Activity, 78-4948
- Polyribonucleotides
Binding, 78-5100
- Xeroderma Pigmentosum
DNA Repair, 78-5170

**8-Quinolinol, 5,7-Dibromo-
Diet**

- Carcinogenic Activity, Rat, 78-5099

Radiation

- Benzo(a)pyrene
Cell Survival, 78-5003
Escherichia coli, 78-5003
- Leukemia
Japan, Review, 78-4869
- Neoplasms
Japan, Review, 78-4869

Radiation, Ionizing

- Anemia, Aplastic
Chromosome Aberrations, 78-4939
Micronucleus Test, 78-4939
- Angiosarcoma

Radiation, Ionizing (cont'd)

- Case Report, Hip Tumor, 78-5305
- Radiotherapy, 78-5305
- Bone Neoplasms
Epidemiology, Review, 78-4874
- Breast Neoplasms
Dose-Response Study, Review, 78-4871
- Carcinogenic Activity
Dose-Response Study, Review, 78-4872
- Cell Transformation, Neoplastic
DNA Repair, Review, 78-4875
Dose-Response Study, 78-4879
Dose-Response Study, Fibroblasts, 78-4875
Mathematical Model, 78-4879
- Cervix Neoplasms
Neoplasms, Multiple Primary, 78-5178
- Chromosome Aberrations
Cell Transformation, Neoplastic, 78-4879
- DNA
Mutagenic Activity, 78-4879
- Eyelid Neoplasms
Carcinoma, 78-5179
Case Report, 78-5179
Ultrastructural Study, 78-5179
- Fibrosarcoma
Case Report, 78-5180
Mastectomy, 78-5180
- Gynecologic Neoplasms
Neoplasms, Connective Tissue, 78-5178
Neoplasms, Multiple Primary, 78-5178
Sarcoma, 78-5178
- Hepatoma
Dimethylamine, *N*-Nitroso-, 78-5183
Liver Regeneration, 78-5183
Species Difference, 78-5183
- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-
Cell Transformation, Neoplastic, 78-5144
Co-carcinogenic Activity, 78-5144
- Intestinal Neoplasms
Cervix Neoplasms, 78-5178
- Leukemia
Dose-Response Study, Lizard, 78-4878
Dose-Response Study, Review, 78-4871
Epidemiology, Review, 78-4874
- Leukopenia
Dose-Response Study, Lizard, 78-4878
- Lung Neoplasms
Dose-Response Study, Review, 78-4871
- Lymphoma
Epidemiology, Review, 78-4874
Growth, Spleen, Liver, 78-5294
Histocompatibility Antigens, 78-5294
Transplantation Immunology, 78-5294
- Mammary Neoplasms, Experimental
Anesthetics, 78-5177
Dose-Response Study, Rat, 78-5177
4,4'-Stilbenediol, α,α' -Diethyl-, 78-5177
- Mutagenic Activity
Mathematical Model, 78-4879
- Neoplasms, Radiation-Induced
Dose-Response Study, Review, 78-4873
Epidemiology, Review, 78-4872, 78-4873, 78-4874
- Occupational Hazard
Dose-Response Study, Review, 78-4871
- Pancreatic Neoplasms
Epidemiology, Review, 78-4874
- Radiotherapy

ation, Ionizing (cont'd)

Epidemiology, Review, 78-4870

Respiratory Tract Neoplasms

Epidemiology, Review, 78-4874

Retinoblastoma

Cell Survival, Fibroblasts, 78-5182

DNA Repair, 78-5182

Sebaceous Gland Neoplasms

Carcinoma, 78-5179

Case Report, 78-5179

Ultrastructural Study, 78-5179

4,4'-Stilbenediol, α,α' -Diethyl-

Co-carcinogenic Activity, Rat, 78-5177

Stomach Neoplasms

Epidemiology, Review, 78-4874

Thyroid Neoplasms

Dose-Response Study, Review, 78-4871

Epidemiology, Review, 78-4874

Virus, Herpes Ateles

Lymphocyte Transformation, 78-5239

Virus, Radiation Leukemia

Cell Transformation, Neoplastic, 78-5295

Precancerous Conditions, 78-5224

Virus, Rous Sarcoma

Antibody Formation, 78-5292

Immunity, Cellular, 78-5292

Virus, SV40

DNA Replication, 78-5253

radioactive Fallout**Cesium Radioisotopes**

Body Burden, 78-5181

Food Contamination, 78-5181

radioisotopes**Carcinogenic Potential**

Review, 78-4843

12-O-Tetradecanoylphorbol-13-acetate

Rubidium, 78-5042

am**Air Pollution**

Carcinogenic Potential, Review, 78-4868

Bone Neoplasms

Carcinogenic Potential, Review, 78-4868

ptors, Hormone**Estradiol**

Mammary/Uterine Tissue, Mouse, 78-5139

4,4'-Stilbenediol, α,α' -Diethyl-

Estradiol, 78-5139

Mammary/Uterine Tissue, Mouse, 78-5139

il Neoplasms**Sarcoma**

Transplantation Immunology, 78-5293

Transplantation Immunology

Age Factors, Mouse, 78-5293

pine**Carcinogenic Potential**

Review, 78-4843

Chromosome Aberrations

Mutagenic Activity, Mouse, 78-5141

ratary Tract Diseases**1-Naphthalenesulfonic Acid, 4-Hydroxy-3,4'-azodi-,**

Disodium Salt

Dose-Response Study, 78-5129

Respiratory Tract Neoplasms**Adenocarcinoma**

Benzo(a)pyrene, 78-5004, 78-5006

Adenoma

Benzo(a)pyrene, 78-5004, 78-5006

Benzo(a)pyrene

Ferric Chloride, 78-5004

Histological Study, Rat, 78-5006

Iodine, 78-5004

Talc, 78-5006

Beryllium

Carcinogenic Potential, Review, 78-4934

1-Butanamine, *N*-Butyl-*N*-nitroso-

Carcinogenic Activity, Review, 78-4825

Cadmium

Epidemiology, Review, 78-4934

Carcinoma

Benzo(a)pyrene, 78-5006

Histological Study, 78-4918

Carcinoma, Epidermoid

Benzo(a)pyrene, 78-5004, 78-5006

Diethylamine, *N*-Nitroso-

Carcinogenic Activity, Review, 78-4822

Dimethylamine, *N*-Nitroso-

Carcinogenic Activity, Review, 78-4821

Dipropylamine, *N*-Nitroso-

Carcinogenic Activity, Review, 78-4824

Epidemiology

Indonesia, 78-5345

United States, 78-5348

Ethylene, Chloro-

Epidemiology, 78-4832

Genetics

Aryl Hydrocarbon Hydroxylases, 78-5324

Hemangioma

Benzo(a)pyrene, 78-5006

Lymphocytes

Aryl Hydrocarbon Hydroxylases, 78-5324

Papilloma

Benzo(a)pyrene, 78-5004, 78-5006

Propylamine, *N*-(2-Hydroxypropyl)-*N*-nitroso-

Carcinogenic Metabolite, Review, 78-4824

Radiation, Ionizing

Epidemiology, Review, 78-4874

Retinol

Antineoplastic Activity, Review, 78-4860

Smoking

Epidemiology, 78-5351

Epidemiology, Review, 78-4912

Socioeconomic Factors

Epidemiology, 78-5351

Talc

Histological Study, Rat, 78-5006

Retinoblastoma**Eye Neoplasms**

Virus, Polyoma, JC, 78-5261

Genetics

DNA Repair, 78-5182

Epidemiology, Review, 78-4926

Radiation Effects, 78-5182

Hereditary Diseases

Epidemiology, Review, 78-4930

Review, 78-4929

Radiation, Ionizing

Cell Survival, Fibroblasts, 78-5182

DNA Repair, 78-5182

Retinoic Acid

- 12-*O*-Tetradecanoylphorbol-13-acetate
Carboxy-Lyases, 78-5037

Retinol

- Aryl Hydrocarbon Hydroxylases
Lung, Hamster, 78-5019
- Benzo(a)pyrene
Aryl Hydrocarbon Hydroxylases, 78-5019
Metabolism, 78-5019
- Bladder Neoplasms
Antineoplastic Activity, Review, 78-4860
- Carcinoma, Epidermoid
Epidemiology, 78-5146
- Mammary Neoplasms, Experimental
Antineoplastic Activity, Review, 78-4860
- Neoplasms
Antineoplastic Activity, 78-5146
Case-Control Study, 78-5146
Epidemiology, 78-5146, 78-5349
Serum Levels, 78-5349
- Respiratory Tract Neoplasms
Antineoplastic Activity, Review, 78-4860
- Skin Neoplasms
Antineoplastic Activity, Review, 78-4860
- Smoking
Serum Levels, 78-5349

Retinol Acetate

- Benzo(a)pyrene
Metabolism, 78-5019

Retinol Palmitate

- Aflatoxin B1
Hydroxylation and Demethylation, 78-4963
Metabolism, Rabbit, 78-4963

Retrorsine

- DNA Repair
Liver, 78-4949

Reverse Transcriptase

- Leukemia
Precancerous Conditions, 78-5204
Virus, C-Type RNA Tumor, 78-5270
- Leukemia, Myelocytic
Blood Platelets, 78-5204
- Lymphosarcoma
Dogs, 78-5229
- Mammary Neoplasms, Experimental
Dexamethasone, 78-5143
- Melanoma
Isolation and Characterization, B16 Tumor, 78-5221
RNA, Viral, 78-5221
- Precancerous Conditions
Blood Platelets, 78-5204
- Virus, Avian Myeloblastosis
Ribonuclease, 78-5203
RNA, Transfer, 78-5189
- Virus, Avian Sarcoma
DNA-RNA Hybridization, 78-5191
- Virus, Baboon
Hybrid Cells, 78-5231
- Virus, Moloney Murine Leukemia
Competition Radioimmunoassay, 78-5216
Mouse 3T3 Clone Cells, 78-5215
- Virus, Rauscher Murine Leukemia
Ultrastructural Study, Crude Tissue, 78-5204
- Virus, RNA Tumor
Antigenic Determinants, 78-5216

Reverse Transcriptase (cont'd)

- Carcinogenic Potential, Review, 78-4903
- Competition Radioimmunoassay, 78-5216
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-, 78-5216
- Ultrastructural Study, Crude Tissue, 78-5204
- Virus, Avian Myeloblastosis, 78-5271

Rhabdomyosarcoma

- Lung
Neoplasm Metastasis, 78-5388
- Pregnancy
Case Report, 78-5390
- Urea, Methyl Nitroso-
Carcinogenic Activity, Xiphophorus Fish, 78-5087
Genetics, Fish, 78-5087
- Virus, Bovine Leukemia
DNA-DNA Hybridization, 78-5230
- Virus, RD-114
DNA-RNA Hybridization, 78-5226

Riboflavin

- Carcinogen, Chemical
Ames Test, 78-5048

Ribonuclease

- Diethylamine, *N*-Nitroso-
RNA, Liver, 78-5072
- Virus, Avian Myeloblastosis
Polyribonucleotides, 78-5203
Reverse Transcriptase, 78-5203
RNA, Viral, 78-5203
- Virus, Rauscher Murine Leukemia
Polyribonucleotides, 78-5203
RNA, Viral, 78-5203

RNA

- Diethylamine, *N*-Nitroso-
Hepatocarcinogenesis, 78-5072

RNA, Messenger

- Virus, Rauscher Murine Leukemia
Viral Proteins, 78-5202
- Virus, SV40
Antigens, Viral, 78-5249
DNA-RNA Hybridization, 78-5255, 78-5260
Ultrastructural Study, CV1 Cells, 78-5255

RNA, Neoplasm

- Hepatoma
DNA-RNA Hybridization, 78-5069
Neoplasm Transplantation, 78-5069

RNA Polymerase

- Aflatoxin B1
Hypophysectomy, 78-4960
- Virus, SV40
DNA, Viral, 78-5250

RNA Replication

- Aflatoxin B1
Liver, Rat, 78-4957
RNA, Ribosomal, 78-4957
- Virus, Avian Myeloblastosis
RNA, Transfer, 78-5189
- Virus, Avian Sarcoma
Genes, Viral, Review, 78-4882
- Virus, Murine Mammary Tumor
Dexamethasone, 78-5222
- Virus, SV40
Antigens, Viral, 78-5258
DNA, Viral, 78-5250, 78-5260

- Ribosomal**
 latotoxin B1
 RNA Replication, 78-4957
 rus, SV40
 Antigens, Viral, 78-5258
- Transfer**
 benz(a,e)aceanthrylene
 Microsomes, Liver, 78-5033
 rus, Avian Myeloblastosis
 Reverse Transcriptase, 78-5189
 RNA Replication, 78-5189
- Transfer, Methyltransferases**
 tyric Acid, 2-Amino-4-(ethylthio)-
 Liver, Rat, 78-5398
- Thionine**
 Liver, Rat, 78-5398
- Thymoma**
 Carcinogen, Chemical, 78-5398
- Viral**
 melanoma
 Reverse Transcriptase, 78-5221
 rus, Abelson Murine Leukemia
 Viral Proteins, 78-5197
 Virus, Helper, 78-5197
 rus, Adeno 2
 Benzimidazole, 5,6-Dichloro-1- β -D-ribofuranosyl-
 78-5233
 rus, Avian Myeloblastosis
 Ribonuclease, 78-5203
 rus, Avian Sarcoma
 DNA-RNA Hybridization, Review, 78-4880
 78-4882
 Genes, Viral, Review, 78-4882
 rus, C-Type RNA Tumor
 DNA-RNA Hybridization, Review, 78-4880
 rus, Friend Murine Leukemia
 DNA-RNA Hybridization, 78-5206
 rus, Moloney Murine Leukemia
 Heteroduplex Structures, 78-5214
 Mouse 3T3 Clone Cells, 78-5215
 rus, Moloney Murine Sarcoma
 Nucleotide Sequence, 78-5213
 rus, Murine Sarcoma
 DNA-RNA Hybridization, Review, 78-4880
 rus, Polyoma
 DNA Replication, 78-5267
 rus, Rauscher Murine Leukemia
 Ribonuclease, 78-5203
 Viral Proteins, 78-5202
 rus, Rous Sarcoma
 Protein Kinase, 78-5188
 Viral Proteins, 78-5188
 rus, SV40
 Degradation, 78-5260
 DNA-RNA Hybridization, 78-5260
- Willemoma**
 Urea, Methyl Nitroso-, 78-5085
 Phorbolol 12,13-Didecanoate
 Ion Transport, Fibroblasts, 78-5042
 Phorbolol 12,13-Didecanoate
 Ion Transport, Fibroblasts, 78-5042
 Phosphoric Acid
 Ion Transport, Fibroblasts, 78-5042
 Prostaglandins E
- Rubidium (cont'd)**
 Phosphoric Acid, 78-5042
 Prostaglandins F
 Ion Transport, Fibroblasts, 78-5042
 12-*O*-Tetradecanoylphorbol-13-acetate
 Ion Transport, Fibroblasts, 78-5042
 Radioisotopes, 78-5042
- Saccharomyces cerevisiae**
 Dimethylamine, *N*-Nitroso-
 Mutagenic Activity, 78-4945
- Safrole**
 see Benzene, 4-Allyl-1,2-(Methylenedioxy)-
- Saliva**
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Nitrous Acid, 78-5103
 Stomach Neoplasms
 Nitrosation, Review, 78-4818
- Salmonella typhimurium**
 Fluorene, 2-Nitro-
 Mutagenic Activity, 78-4946
 Guanidine, Nitroso-
 Mutagenic Activity, 78-4946
 Intestinal Bacteria
 Mutagenic Activity, 78-4952
 2-Oxetanone
 Mutagenic Activity, 78-4946
- Sarcoidosis**
 Carcinoma, Bronchogenic
 Immune response, 78-4916
- Sarcoma**
 Areca
 Carcinogenic Activity, Mouse, Rat, 78-4955
 Benz(a)anthracene, 7,12-Dimethyl-
 BCG, 78-5030
 Carcinogen, Chemical
 Filter Implant Test, 78-4943
 Cholanthrene, 3-Methyl-
 Histocompatibility Antigens, 78-4994
 Gynecologic Neoplasms
 Radiation, Ionizing, 78-5178
 Histocompatibility Antigens
 Immunogenicity, 78-4994
 Kidney Transplantation
 Epidemiology, 78-5353
 Lung
 Neoplasm Metastasis, 78-5388
 Macrophages
 T-Lymphocytes, 78-5287
 Migration Inhibitory Factor, 78-5287
 Mammary Neoplasms, Experimental
 Urea, Ethyl Nitroso-, 78-5088
 Neoplasm Metastasis
 Doubling Time, 78-5388
 Nose Neoplasms
 Epidemiology, USSR, 78-5379
 Oncogenic Viruses
 Epidemiology, Review, 78-4895
 Otorhinolaryngologic Neoplasms
 Epidemiology, USSR, 78-5379
 Pregnancy
 Case Report, 78-5390
 Rectal Neoplasms
 Transplantation Immunology, 78-5293
 Tracheal Neoplasms

Sarcoma (cont'd)

- Nickel Sulfide, 78-5119
- Virus, C-Type RNA Tumor
 - DNA, Neoplasm, 78-5274
 - DNA, Viral, 78-5274
 - Trans-Species Infection, 78-5274
 - Transformation, Genetic, 78-5274
- Virus, Murine Sarcoma
 - T-Lymphocytes, 78-5287
 - Macrophages, 78-5287

Sarcoma 180, Crocker

- T-Lymphocytes
 - Anti-Antibodies, 78-5279
 - Growth, 78-5279
 - Transplantation Immunology, 78-5279

Sarcoma, Kaposi's

- Virus, Epstein-Barr
 - DNA-RNA Hybridization, 78-5241

Sarcoma, Osteogenic

- Beryllium
 - Carcinogenic Potential, Review, 78-4934
- Hereditary Diseases
 - Epidemiology, Review, 78-4930
- Pregnancy
 - Case Report, 78-5390
- p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
 - Carcinogenic Activity, Monkey, 78-5148
- Virus, Bovine Leukemia
 - DNA-DNA Hybridization, 78-5230
- Virus, Polyoma, BK
 - Carcinogenic Activity, Hamster, 78-5263

Sarcoma, Polymorphous

see Mesenchymoma

Sarcoma, Reticulum Cell

- Heart Transplantation
 - Case Report, 78-5384
- Immunosuppression
 - Kidney Transplantation, 78-5353
- Intestinal Neoplasms
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5104
- Kidney Transplantation
 - Epidemiology, 78-5353
- Leukemia, Myeloblastic
 - Case Report, 78-5299
 - Drug Therapy, 78-5299
 - Radiotherapy, 78-5299

Sarcosine, *N*-Nitroso-

- Food Contamination
 - Quantitation Method, 78-5057

Schistosomiasis

- Australia Antigen
 - Epidemiology, Review, 78-4846
- Bladder Neoplasms
 - Epidemiology, Review, 78-4846
- Colonic Neoplasms
 - Epidemiology, Review, 78-4846
- Gynecologic Neoplasms
 - Epidemiology, Review, 78-4846
- Hepatoma
 - Epidemiology, Review, 78-4846
- Leukemia
 - Epidemiology, Review, 78-4846
- Lymphoma, Giant Follicular
 - Epidemiology, Review, 78-4846

Sclerosis

- Asbestos
 - Lung, Rat, 78-4971
- Coal
 - Lung, Rat, 78-4971

Sebaceous Gland Neoplasms

- Carcinoma
 - Radiation, Ionizing, 78-5179
- Radiation, Ionizing
 - Case Report, 78-5179
 - Ultrastructural Study, 78-5179

Selenic Acid, Disodium Salt

- Microsomes, Liver
 - Mutagenic Activity, 78-5115

Selenious Acid, Disodium Salt

- Fibroblasts
 - Chromosome Aberrations, 78-5115
 - Contact Inhibition, 78-5115
 - DNA Repair, 78-5115
- Microsomes, Liver
 - Mutagenic Activity, 78-5115
- Xeroderma Pigmentosum
 - Chromosome Aberrations, 78-5115
 - DNA Repair, 78-5115

Selenium

- Carcinogen, Chemical
 - Inhibitors, Review, 78-4806
- Neurilemmoma
 - Urea, Methyl Nitroso-, 78-5085

Sex Hormones

- Acetamide, *N*-Fluoren-2-yl-
 - Metabolism, Kidney, 78-5078
- Dimethylamine, *N*-Nitroso-
 - Metabolism, Kidney, 78-5078
- Glioma
 - Age Factors, 78-5135
 - Benzo(a)pyrene, 78-5135
 - Epidemiology, 78-5135

Siderosis

- Lung Neoplasms
 - Adenocarcinoma, 78-5326
 - Giant Cell Tumors, 78-5326

Silicate

- Mesothelioma
 - Epidemiology, Turkey, 78-4909

Silver

- Lung Neoplasms
 - Adenocarcinoma, 78-5326
 - Giant Cell Tumors, 78-5326

Sirodesmin A

- Bone Marrow Cells
 - Chromosome Aberrations, 78-5155
- Lymphocytes
 - Chromosome Aberrations, 78-5155

Skin Neoplasms

- Arsenic
 - Epidemiology, Review, 78-4934
- Ataxia Telangiectasia
 - Epidemiology, Review, 78-4930
- Benz(a)anthracene
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-5022
- Benz(a)anthracene, 1,2-Dihydro-

Neoplasms (cont'd)

- Carcinogenic Potential, 78-5022
- Benz(a)anthracene, 3,4-Dihydro-
 - Carcinogenic Potential, 78-5022
- Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-
 - Carcinogenic Potential, 78-5022
- Benz(a)anthracene, 7,12-Dimethyl-
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-5036
- Benzo(a)pyrene
 - Putrescine, 78-5005
- 1,3-Butadiene, 2-Chloro-
 - Epidemiology, Review, 78-4834
- Carcinoma, Epidermoid
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5036
 - Ultraviolet Rays, 78-5174
- Epidemiology
 - Indonesia, 78-5345
 - Sabah, Malaysia, 78-5346
- Fibrosarcoma
 - Ultraviolet Rays, 78-5174
- Glycerol
 - Precancerous Conditions, 78-5047
- Heart Transplantation
 - Case Report, 78-5384
- Keratoacanthoma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5036
- L-Lymphocytes
 - Lymphocyte Depletion, 78-5174
- Neoplasm Regression, Spontaneous
 - Histological Study, Mouse, 78-5036
- Papilloma
 - Benz(a)anthracene, 78-5022
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5036
- Propanol, 2,3-Dibromo-, Phosphate
 - Carcinogenic Potential, Rat, 78-5167
- Putrescine
 - Urine, Mouse, 78-5005
- Retinol
 - Antineoplastic Activity, Review, 78-4860
- Smoke Condensate
 - Glycerol, 78-5047
 - Precancerous Conditions, 78-5047
- Socioeconomic Factors
 - Epidemiology, 78-5351
- 2-*O*-Tetradecanoylphorbol-13-acetate
 - Carboxy-Lyases, 78-5037
- Ultraviolet Rays
 - Epidemiology, Review, 78-4877
 - Immunosuppression, 78-5174
- Virus, Papilloma
 - Precancerous Conditions, Review, 78-4889
- Xeroderma Pigmentosum
 - Epidemiology, Review, 78-4930
- Condensate
 - Ames Test
 - Broiled Fish, Meat, 78-5048
- Benzo(a)pyrene
 - Ames Test, 78-5048
- Lung Neoplasms
 - Beeswax Pellet Implantation, 78-5045
 - Carcinoma, Epidermoid, 78-5045
 - Neoplasm Metastasis, 78-5045
- Liver Neoplasms
 - Glycerol, 78-5047
 - Precancerous Conditions, 78-5047
- Pytophan
 - Ames Test, 78-5048

Smoking

- Aryl Hydrocarbon Hydroxylases
 - Genetics, Review, 78-4928
- Asbestos
 - Co-carcinogenic Activity, Review, 78-4842
- Benzoic Acid, 2-Amino-3-hydroxy-
 - Metabolism, 78-5044
- Bladder Neoplasms
 - Epidemiology, 78-5351
 - Epidemiology, Review, 78-4912
- Carbon Dioxide
 - Inhalation Study, Hamster, 78-5043
- Carbon Monoxide
 - Inhalation Study, Hamster, 78-5043
- Carcinogenic Activity
 - Respiratory Tract, Mouse, 78-5046
- Cervix Neoplasms
 - Epidemiology, 78-5391
- Esophageal Neoplasms
 - Epidemiology, 78-5049
 - Epidemiology, Pakistan, 78-5382
- Kidney Neoplasms
 - Epidemiology, Review, 78-4912
- Lung Neoplasms
 - Epidemiology, Great Britain, Review, 78-4849
 - Epidemiology, Japan, 78-4917
 - Epidemiology, Review, 78-0485
 - Low Tar/Nicotine Cigarettes, Review, 78-0485
- Nicotine
 - Inhalation Study, Hamster, 78-5043
- Pancreatic Neoplasms
 - Epidemiology, Review, 78-4912
- Pharyngeal Neoplasms
 - Epidemiology, 78-5344
- Respiratory Tract Neoplasms
 - Epidemiology, 78-5351
 - Epidemiology, Review, 78-4912
- Retinol
 - Serum Levels, 78-5349
- Stomach Neoplasms
 - Epidemiology, Japan, 78-4917
- Sodium Selenite
 - see Selenious Acid, Disodium Salt
- Soft Tissue Neoplasms
 - Carcinogen, Chemical
 - Filter Implant Test, 78-4943
 - Hemangioendothelioma
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5063
 - Histiocytoma
 - Clinopathologic Study, 78-5307
 - Neoplasm Metastasis, 78-5307
 - Ultrastructural Study, 78-5307
- Soil Pollutants
 - Plutonium
 - Ecosystem Distribution, 78-5184
- Somatotropin
 - Breast Neoplasms
 - DNA Replication, 78-5392
- Sorbic Acid
 - Nitrous Acid, Sodium Salt
 - Food Contamination, 78-5124
 - Mutagenic Reaction Products, 78-5124
- Stearic Monoglyceride
 - Gastrointestinal Neoplasms

Stearic Monoglyceride (cont'd)

Dietary Fats, 78-5120

Sterigmatocystin

Ames Test

Carcinogenic Potential, Review, 78-4815

Herbs

Drug Contamination, 78-4956

Steroids

Hepatoma

Precancerous Conditions, Review, 78-4922

Neoplasms, Experimental

Precancerous Conditions, Review, 78-4922

4,4'-Stilbenediol, α,α' -Diethyl-

Adenocarcinoma

Carcinogenic Potential, Review, 78-4863

Age Factors

Oxidative Metabolism, Mouse, 78-5138

Breast Neoplasms

Epidemiology, 78-5363

Chromosome Aberrations

Cell line, 78-5137

Leukocytes, 78-5137

Chromosome Abnormalities

Cells, Cultured, 78-5136

Fibroblasts, Hamster, 78-5136

Dienestrol

Carcinogenic Metabolite, 78-5138

Erythroleukemia

Mitosis, 78-5136

Estradiol

Receptors, Hormone, 78-5139

Gynecologic Neoplasms

Adenocarcinoma, 78-5370

Carcinogenic Potential, Review, 78-0486

Epidemiology, 78-5370

Transplacental Carcinogenesis, 78-5370

HeLa Cells

Mitosis, 78-5136

Mammary Neoplasms, Experimental

Radiation, Ionizing, 78-5177

Mitosis

Cells, Cultured, 78-5136

Precancerous Conditions

Epidemiology, 78-5370

Radiation, Ionizing

Co-carcinogenic Activity, Rat, 78-5177

Receptors, Hormone

Mammary/Uterine Tissue, Mouse, 78-5139

Urine

Carcinogenic Metabolite, 78-5138

Uterine Neoplasms

Epidemiology, Review, 78-4867

Stomach Neoplasms

Acetic Acid, Butylnitrosaminomethyl Ester

Carcinogenic Activity, Rat, 78-5065

Acetic Acid, Ethylnitrosaminomethyl Ester

Carcinogenic Activity, Rat, 78-5065

Acetic Acid, Methylnitrosaminomethyl Ester

Carcinogenic Activity, Rat, 78-5065

Acetic Acid, Propylnitrosaminomethyl Ester

Carcinogenic Activity, Rat, 78-5065

Acetic Acid, Trideuteromethylnitrosaminomethyl Ester

Carcinogenic Activity, Rat, 78-5065

Anemia, Pernicious

Nitrous Acid, 78-5330

Stomach Neoplasms (cont'd)

Precancerous Conditions, 78-5330

Ascorbic Acid

Epidemiology, Review, 78-4912

Butyric Acid, Methylnitrosaminomethyl Ester

Carcinogenic Activity, Rat, 78-5065

Carcinoma

Metaplasia, 78-5328

Carcinoma, Epidermoid

Case Report, 78-5331

Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101

Quinoline, 8-Nitro-, 78-5099

Ultrastructural Study, 78-5331

Cyanamide, Methylnitroso-

Carcinogenic Activity, Rat, 78-5103

Diet

Epidemiology, Austria, 78-5350

Epidemiology, Japan, 78-4917, 78-5374

Intestinal Transit Time, 78-5374

Epidemiology

United States, 78-5348

Ethnic Groups

Epidemiology, 78-5344

Fibrosarcoma

Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101

Gastrectomy

Epidemiology, 78-5329

Gastric Juice

Nitrosation, Review, 78-4818

Gastritis

Acidity, 78-5373

Epidemiology, 78-5373

Epidemiology, Review, 78-4818

Guanidine, 1-Methyl-3-nitro-1-nitroso-

Carcinogenic Potential, Hamster, 78-5101

Neoplasm Metastasis, 78-5101

Mesenchymoma

Morpholine, *N*-Nitroso-, 78-5056

Metaplasia

Intestines, 78-5328

Leucine Aminopeptidase, 78-5328

Precancerous Conditions, 78-5328

Ultrastructural Study, 78-5328

Morpholine, *N*-Nitroso-

Carcinogenic Potential, Fish, 78-5056

Nitric Acid

Epidemiology, Review, 78-4818

Water Pollution, Review, 78-4818

Nitroso Compounds

Nitrosation, Review, 78-4818

Nitrous Acid

Bacteria, Review, 78-4818

Epidemiology, Review, 78-4818

Papilloma

Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5101

Quinoline, 8-Nitro-, 78-5099

Peptic Ulcer

Epidemiology, 78-5329

1-Propanol, 2,3-Dibromo-, Phosphate

Carcinogenic Potential, Rat, 78-5167

Quinoline, 8-Nitro-

Carcinogenic Activity, Rat, 78-5099

Radiation, Ionizing

Epidemiology, Review, 78-4874

Saliva

Nitrosation, Review, 78-4818

Smoking

Stomach Neoplasms (cont'd)

Epidemiology, Japan, 78-4917

Urea, Methyl Nitroso-

Immune Response, Rat, 78-5086

Virus, Polyoma, JC

Carcinogenic Activity, Hamster, 78-5261

ss

Mammary Neoplasms, Experimental

Benz(a)anthracene, 7,12-Dimethyl-, 78-5029

Neoplasm Regression, 78-5029

Virus, Rous Sarcoma

Genetics, 78-5292

ene Oxide

see Benzene, (Epoxyethyl)-

cinic Acid, Carboxymethyloxy-

Detergents

Biodegradability, 78-5125

Carcinogenic Potential, 78-5125

Metabolism, Rat, 78-5125

anilamide, *N*-(3,4-Dimethyl-4-isoxazolyl)-

Dimethylamine, *N*-Nitroso-

Nitrous Acid, Sodium Salt, 78-5076

isoxazole

see Sulfanilamide, *N*-(3,4-Dimethyl-4-isoxazolyl)-

uric Acid, Diethyl Ester

Cell Survival

Dose-Response Study, Hamster Ovary Cells

78-5158

Hypoxanthine Phosphoribosyltransferase

Mutagenic Activity, 78-5158

uric Acid, Dimethyl Ester

Cell Survival

Dose-Response Study, Hamster Ovary Cells

78-5158

Hydroxylamine, *O,N*-Dimethyl-*N*-nitroso-

Synthesis, 78-5082

Hypoxanthine Phosphoribosyltransferase

Mutagenic Activity, 78-5158

uric Acid, Iron Salt

Benz(a)anthracene, 7-Methyl-

Dihydrodiols, 78-5031

phytum officinale

Hemangioendothelioma

Histological Study, Rat, 78-4942

Hepatoma

Carcinogenic Activity, Roots, Leaves, 78-4942

Histological Study, Rat, 78-4942

Liver Neoplasms

Hemangioendothelioma, 78-4942

Benzo(a)pyrene

Co-carcinogenic Activity, Rat, 78-5006

Respiratory Tract Neoplasms

Benzo(a)pyrene, 78-5006

Histological Study, Rat, 78-5006

sterone

Acetamide, *N*-Fluoren-2-yl-

Metabolism, Kidney, 78-5078

Dimethylamine, *N*-Nitroso-

Metabolism, Kidney, 78-5078

Prostatic Neoplasms

Hyperplasia, 78-4920

Tetracycline

Dimethylamine, *N*-Nitroso-

Nitrous Acid, Sodium Salt, 78-5076

Urinary Tract Infections

Dimethylamine, *N*-Nitroso-, 78-5076

12-*O*-Tetradecanoylphorbol-13-acetate

Benz(a)anthracene, 7,12-Dimethyl-

Neoplasm Regression, Spontaneous, 78-5036

Carboxy-Lyases

Colchicine, 78-5037

Nucleotides, Cyclic, 78-5037

Retinoic Acid, 78-5037

Carcinoma, Epidermoid

Benz(a)anthracene, 7,12-Dimethyl-, 78-5036

Cell Division

Trachea, Mouse, 78-5039

Cells, Cultured

Metabolism, Fibroblasts, 78-5038

Epithelial Cells

Trachea, Mouse, 78-5039

Growth

Trachea, Mouse, 78-5039

Keratoacanthoma

Benz(a)anthracene, 7,12-Dimethyl-, 78-5036

Ouabain

Ion Transport, Fibroblasts, 78-5042

Papilloma

Benz(a)anthracene, 7,12-Dimethyl-, 78-5036

Phorbol 13-Acetate

Metabolism, Fibroblasts, 78-5038

Plasminogen Activators

Carcinogenic Activity, 78-5040

Fibroblasts, Chick Embryo, 78-5040

Rubidium

Ion Transport, Fibroblasts, 78-5042

Radioisotopes, 78-5042

Skin Neoplasms

Benz(a)anthracene, 78-5022

Benz(a)anthracene, 7,12-Dimethyl-, 78-5036

Carboxy-Lyases, 78-5037

Virus, Adeno 5

Cell Transformation, Neoplastic, 78-5041

Virus, Rous Sarcoma

Plasminogen Activators, 78-5040

Thioamide

see Isonicotinamide, 2-Ethylthio-

Thioguanine

see Purine-6(1*H*)-thione, 2-Amino-

Thorium Dioxide

Liver Neoplasms

Precancerous Conditions, Review, 78-4922

Thrombin

Lymphocytes

Chromosome Aberrations, 78-5157

Thromboplastin

Hepatoma

Neoplasm Circulating Cells, 78-5304

Neoplasm Metastasis, 78-5304

Thymidine

Benzo(a)pyrene

Diol Epoxides, 78-5007

Thymidine-5'-phosphoric Acid

see 5'-Thymidylic Acid

5'-Thymidylic Acid

2-Oxetanone

Phosphodiester, 78-5108

Thymoma

Benz(a)anthracene, 7,12-Dimethyl-
BCG, 78-5030

Myasthenia Gravis

Histological Study, 78-5315

Neoplasm Invasiveness, 78-5315

Urea, Methyl Nitroso-

Immune Response, Rat, 78-5086

Thymopoietins

Virus, Radiation Leukemia

Cell Differentiation, 78-5295

Cell Transformation, Neoplastic, 78-5295

Thymus Gland

Plasmacytoma

Graft Rejection, 78-5289

Virus, Radiation Leukemia

Precancerous Conditions, 78-5224

Thyroid Gland

Aniline, 4,4'-Oxydi-

Hyperplasia, 78-5126

Thyroid Hormones

Mammary Neoplasms, Experimental

Estrogens, 78-5134

Pituitary Neoplasms

Estrogens, 78-5134

Thyroid Neoplasms

Adenocarcinoma

2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5079

Adenoma

Epidemiology, Iran, 78-5383

2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5079

Aniline, 4,4'-Oxydi-

Precancerous Conditions, 78-5126

Carcinoma

Epidemiology, Iran, 78-5383

2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5079

Carcinoma, Epidermoid

Immunohistochemical Study, 78-5316

Ultrastructural Study, 78-5316

Goiter

Epidemiology, Iran, 78-5383

Griseofulvin

Dose-Response Study, Rat, 78-5142

Kidney Transplantation

Epidemiology, 78-5353

2-Propanol, 1,1'-Iminodi-*N*-nitroso-

Rat, 78-5079

Radiation, Ionizing

Dose-Response Study, Review, 78-4871

Epidemiology, Review, 78-4874

Titanium Oxide

Lung Neoplasms

Giant Cell Tumors, 78-5326

Tobacco

Ames Test

Pyrolysis Products, 78-5049

Areca

Carcinogenic Activity, Mouse, Rat, 78-4955

Tobacco (cont'd)

Carcinogenic Activity

Flue- and Air-Curing, 78-5046

Esophageal Neoplasms

Chewing, 78-5382

***p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-**
Angiosarcoma

Carcinogenic Activity, Monkey, 78-5148

Chromatids

Bone Marrow, Mouse, 78-5149

Leukemia, Myeloblastic

Carcinogenic Activity, Monkey, 78-5148

Lymphosarcoma

Carcinogenic Activity, Monkey, 78-5148

Sarcoma, Osteogenic

Carcinogenic Activity, Monkey, 78-5148

Toluene-2,4-diamine

Ames Test

Mutagenic Activity, Review, 78-4811

***p*-Toluidine, α,α,α -Trifluoro-2,6-dinitro-*N,N*-dipropyl-**

Dipropylamine, *N*-Nitroso-

Pesticides, Review, 78-4824

Toxins

Brain Neoplasms

Fusarium sporotrichioides, 78-4954

Digestive System Neoplasms

Fusarium sporotrichioides, 78-4954

Mammary Neoplasms, Experimental

Fusarium sporotrichioides, 78-4954

Pituitary Neoplasms

Fusarium sporotrichioides, 78-4954

Trace Elements

Nervous System Neoplasms

Blood Levels, 78-5085

Neurilemmoma

Urea, Methyl Nitroso-, 78-5085

Tracheal Neoplasms

Carcinoma

Nickel Sulfide, 78-5119

Nickel Sulfide

Carcinogenic Activity, Rat, 78-5119

Sarcoma

Nickel Sulfide, 78-5119

Transfection

see Transformation, Genetic

Transferases

Acetic Acid, (2,3-Dichloro-4-(2-methylenebutyryl)phenoxy)

Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5016

Benzo(a)pyrene, 78-5016

Cholanthrene, 3-Methyl-, 78-5016

Barbituric Acid, 5-Ethyl-5-phenyl-

Enzymatic Activity, Rat, 78-5016

Benzene, 1-Chloro-2,4-dinitro-

Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5016

Benzo(a)pyrene, 78-5016

Cholanthrene, 3-Methyl-, 78-5016

Spermatozoa, 78-5012

Benzene, 1-(Chloromethyl)-4-nitro-

Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5016

Benzo(a)pyrene, 78-5016

Cholanthrene, 3-Methyl-, 78-5016

Benzene, 1,2-Dichloro-4-nitro-

- ases (cont'd)**
 Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5016
 Benzo(a)pyrene, 78-5016
 Cholanthrene, 3-Methyl-, 78-5016
 enzene, (Epoxyethyl)-
 Spermatozoa, 78-5012
 enzo(a)pyrene
 Enzymatic Activity, Rat, 78-5016
 enzo(a)pyrene 4,5-Oxide
 Spermatozoa, 78-5012
 olanthrene, 3-Methyl-
 Enzymatic Activity, Rat, 78-5016
 opane, 1,2-Epoxy-3-(*p*-nitrophenoxy)-
 Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5016
 Benzo(a)pyrene, 78-5016
 Cholanthrene, 3-Methyl-, 78-5016
 irus, Adeno 12
 Cytocidal Mutants, 78-5234
 Microsomes, 78-5234
- ormation, Genetic**
 eukemia
 Virus, C-Type RNA Tumor, 78-5274
 arcoma
 Virus, C-Type RNA Tumor, 78-5274
- lantation, Heterologous**
 idney Neoplasms
 Carcinoma, 78-5336
 Erythropoietin, 78-5336
 Mouse, Nude, 78-5336
 Ultrastructural Study, 78-5336
 ephroblastoma
 Mouse, Nude, 78-5320
 Neoplasm Metastasis, 78-5320
- lantation Immunology**
 ibrosarcoma
 Hypercholesteremia, 78-5290
 Ultraviolet Rays, 78-5173, 78-5285
 epatoma
 Age Factors, Mouse, 78-5293
 Thymectomy, 78-5293
 listocompatibility Antigens
 Mouse, Neonate, 78-5282
 ung Neoplasms
 Adenocarcinoma, 78-5293
 Age Factors, Mouse, 78-5293
 ymphoma
 Histocompatibility Antigens, 78-5294
 Radiation, Ionizing, 78-5294
 Virus, Gross Murine Leukemia, 78-5291
 mammary Neoplasms, Experimental
 Adenocarcinoma, 78-5293
 Age Factors, Mouse, 78-5293
 Dexamethasone, 78-5143
 Mouse, Nude, 78-5143
 ovarian Neoplasms
 Carcinoma, 78-5293
 rectal Neoplasms
 Age Factors, Mouse, 78-5293
 Sarcoma, 78-5293
 arcoma 180, Crocker
 T-Lymphocytes, 78-5279
 Ultraviolet Rays
 Antigenic Determinants, 78-5285
 T-Lymphocytes, 78-5173
 Virus, Radiation Leukemia
 T-Lymphocytes, 78-5295
- Treflan**
 see *p*-Toluidine, α,α,α -Trifluoro-2,6-dinitro-*N,N*-dipropy
- Trichlorfon**
 see Metrifonate
- Tris-BP**
 see 1-Propanol, 2,3-Dibromo-, Phosphate
- Tritium**
 Acetamide, *N*-Fluoren-2-yl-
 DNA, Binding, 78-4982
 4-Biphenylamine
 Catalytic Hydrogenation, 78-4982
 4-Biphenylamine, 4'-Fluoro-
 Catalytic Hydrogenation, 78-4982
 4-Biphenylamine, 3-Methyl-
 Catalytic Hydrogenation, 78-4982
 Fluoren-2-amine
 Catalytic Hydrogenation, 78-4982
- Tryptophan**
 Ames Test
 Pyrolysis Products, 78-5122
 Bladder Neoplasms
 Alanine, 3-Anthraniloyl-, 78-5338
 Alanine, 3-(3-Hydroxyanthraniloyl)-, 78-5338
 Benzoic Acid, 2-Amino-3-hydroxy-, 78-5338
 Metabolism, 78-5338
 Quinaldic Acid, 4,8-Dihydroxy-, 78-5338
 Quinaldic Acid, 4-Hydroxy-, 78-5338
 Mutagens
 Inactivation, Fruit/Vegetable Juices, 78-5122
 Pyrolysis Products, 78-5122
 Smoke Condensate
 Ames Test, 78-5048
- Tuberculin**
 Virus, Herpes Simplex 2
 Immunity, Cellular, 78-5243
- Tuberculosis, Pulmonary**
 Carcinoma, Bronchogenic
 Immune response, 78-4916
- Tubulin**
 Cell Transformation, Neoplastic
 Cell Cycle Kinetics, 78-5245
 Hepatoma
 Neoplasm Metastasis, 78-5054
 Virus, Polyoma
 Cell Transformation, Neoplastic, 78-5245
 Virus, SV40
 Cell Transformation, Neoplastic, 78-5245
- Turner's Syndrome**
 Uterine Neoplasms
 Adenocarcinoma, 78-5342
 Case Report, 78-5342
 Estrogens, 78-5342
 Karyotyping, 78-5342
 Progesterone, 78-5342
- Ultraviolet Rays**
 Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 DNA Repair, 78-5169
 Xeroderma Pigmentosum, 78-5169
 Antigenic Determinants
 Transplantation Immunology, 78-5285
 Carcinoma, Basal Cell
 Epidemiology, Review, 78-4877
 Carcinoma, Epidermoid

Ultraviolet Rays (cont'd)

- Epidemiology, Review, 78-4877
- Immunosuppression, 78-5174
- Cell Transformation, Neoplastic
 - DNA Repair, Review, 78-4875
 - Dose-Response Study, Fibroblasts, 78-4875
- Chromosome Abnormalities
 - Dose-Response Study, NZB Mouse, 78-5175
 - Embryo Fibroblasts, Mouse, 78-5175
- DNA Repair
 - Endonucleases, 78-5169
 - Monkey CV-1 Cells, 78-5172
 - Pyrimidine Dimers, 78-5172
- Escherichia coli*
 - DNA Repair, 78-5171
 - Mutagenic Activity, 78-5171
- Fibroblasts
 - DNA Repair, 78-5169
- Fibrosarcoma
 - Hypersensitivity, Delayed, 78-5173
 - Immunity, Passive, 78-5285
 - Immunosuppression, 78-5285
 - Transplantation Immunology, 78-5173, 78-5285
- Hypersensitivity, Delayed
 - Benzene, 1-Chloro-2,4-dinitro-, 78-5173
- Lymphocytes
 - Immunosuppression, 78-5285
- T-Lymphocytes
 - Immunosuppression, 78-5173
 - Transplantation Immunology, 78-5173
- Melanoma
 - Epidemiology, Review, 78-4876, 78-4877
- Neoplasms, Radiation-Induced
 - Epidemiology, Review, 78-4873
- Nitrous Acid
 - DNA Repair, 78-5171
- Skin Neoplasms
 - Carcinoma, Epidermoid, 78-5174
 - Epidemiology, Review, 78-4877
 - Fibrosarcoma, 78-5174
 - Immunosuppression, 78-5174
- Virus, C-Type RNA Tumor
 - Virus Activation, Mouse, 78-5212
- Virus, Murine Sarcoma
 - Virus, Helper, 78-5212
- Xeroderma Pigmentosum
 - DNA Repair, 78-5169
 - DNA Repair, Review, 78-4936
 - Neurologic Manifestations, Review, 78-4936
 - Virus, Herpes Simplex 1, 78-5168

Uracil, 5-Fluoro-

- Lung Neoplasms
 - Carcinoma, Epidermoid, 78-5399

Uranium

- Nuclear Reactors
 - Epidemiology, 78-5354
- Occupational Hazard
 - Epidemiology, 78-5354

Urate Oxidase

- Mesenchymoma
 - Benzo(a)pyrene, 78-5014

Urea, 1-Butyl-1-nitroso-

- Leukemia
 - Age Factors, Rat, 78-5094
 - Chromosome Abnormalities, 78-5095

Urea, 1-Butyl-1-nitroso- (cont'd)

- Nervous System Neoplasms
 - Age Factors, Rat, 78-5094
 - Transplacental Carcinogenesis, 78-5094

Urea, 1,3-Dimethyl-1-nitroso-

- Brain Neoplasms
 - Astrocytoma, 78-5096
 - Glioma, 78-5096
 - Oligodendroglioma, 78-5096
 - Ultrastructural Study, Rat, 78-5096

Urea, Ethyl Nitroso-

- Astrocytoma
 - Ultrastructural Study, 78-5089
- Brain Neoplasms
 - Glioma, 78-5186, 78-5319
 - Precancerous Conditions, 78-5091
 - Ultrastructural Study, Rat, 78-5091
 - Wounds and Injuries, 78-5186
- Cell Transformation, Neoplastic
 - Rat Brain, Fetal, 78-5092
- Dibutyl Cyclic AMP
 - Transplacental Carcinogenesis, 78-5093
- Glioblastoma Multiforme
 - Carcinogenic Activity, Review, 78-4827
 - Ultrastructural Study, 78-5089
- Glioma
 - Carcinogenic Activity, Review, 78-4827
 - Dibutyl Cyclic AMP, 78-5093
 - Transplacental Carcinogenesis, 78-5093
 - Ultrastructural Study, Rat, 78-5091, 78-5319
- Mammary Neoplasms, Experimental
 - Carcinoma, 78-5088
 - Dose-Response Study, 78-5088
 - Neonate, Rat, 78-5088
 - Sarcoma, 78-5088
 - Transplacental Carcinogenesis, 78-5088
- Nervous System Neoplasms
 - Transplacental Carcinogenesis, Review, 78-4827
 - Ultrastructural Study, 78-5089
- Neurilemmoma
 - Carcinogenic Activity, Review, 78-4827
 - Dibutyl Cyclic AMP, 78-5093
 - Transplacental Carcinogenesis, 78-5093
- Purine, 2-Amino-6-methoxy-
 - DNA, Review, 78-4827

Urea, Methyl Nitroso-

- Acetamide, *N*-Fluoren-2-yl-
 - Hepatocarcinogenicity, 78-5084
- Ames Test
 - Carcinogenic Potential, Review, 78-4815
- Antibody Formation
 - Immune Response, Rat, 78-5086
- Antigens, Bacterial
 - Antibody Formation, 78-5086
- Brain Neoplasms
 - Antibodies, Neoplasm, 78-5281
 - Glioma, 78-5281
- Carbon Tetrachloride
 - Hepatocarcinogenicity, 78-5084
- Cell Division
 - Hepatocarcinogenicity, 78-5084
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
 - Hepatocarcinogenicity, 78-5084
- Fibrosarcoma
 - Carcinogenic Activity, Xiphophorus Fish, 78-5087
 - Genetics, Fish, 78-5087

rea, Methyl Nitroso- (cont'd)

- Glioma
 - Antibodies, Neoplasm, 78-5281
- Glutamyl Transpeptidase
 - Hepatectomy, 78-5084
 - Liver, Rat, 78-5084
- Lymphocytes
 - Immune Response, Rat, 78-5086
- Macrophages
 - Immune Response, Rat, 78-5086
- Melanoma
 - Histological Study, Gerbil, 78-5090
 - Site Distribution, 78-5090
- Neoplasms, Experimental
 - Carcinogenic Potential, Review, 78-4826
- Nervous System Neoplasms
 - Neurilemmoma, 78-5085
- Neurilemmoma
 - Chromium, 78-5085
 - Iron, 78-5085
 - Rubidium, 78-5085
 - Selenium, 78-5085
 - Trace Elements, 78-5085
 - Zinc, 78-5085
- Rhabdomyosarcoma
 - Carcinogenic Activity, Xiphophorine Fish, 78-5087
 - Genetics, Fish, 78-5087
- Stomach Neoplasms
 - Immune Response, Rat, 78-5086
- Thymoma
 - Immune Response, Rat, 78-5086

rea, Methylcyclohexyl Nitroso-

- Lung Neoplasms
 - Carcinoma, Epidermoid, 78-5399

rea, Thio-

- Glutathione
 - Metabolism, 78-5112
- Oxidoreductases
 - Metabolism, 78-5112

ridine, 5-Bromo-2'-deoxy-

- Anemia, Aplastic
 - Chromosome Aberrations, 78-4939
- Chromatids
 - Bone Marrow, Mouse, 78-5149
 - Hamster V79 Cells, 78-5154
- Chromosome Aberrations
 - Host-Mediated Assay, 78-5154
- Virus, Murine Sarcoma
 - Virus, Helper, 78-5212

ridine, 2'-Deoxy-5-iodo-

- Virus, Murine Sarcoma
 - Virus, Helper, 78-5212

riary Tract Infections

- Dimethylamine, *N*-Nitroso-
 - Tetracycline, 78-5076
- Proteus mirabilis*
 - Dimethylamine, *N*-Nitroso-, 78-5076
 - Nitrous Acid, Sodium Salt, 78-5076

rine

- Virus, Adeno
 - Kidney Transplantation, 78-5275
- Virus, Herpes
 - Kidney Transplantation, 78-5275
- Virus, Polyoma

Urine (cont'd)

- Kidney Transplantation, 78-5275

Urogenital Neoplasms

- Aniline, 4,4'-Oxydi-
 - Precancerous Conditions, 78-5126
- Epidemiology
 - Japan, 78-4917
- Estrogens
 - Carcinogenic Potential, Review, 78-0486
- Ethnic Groups
 - Epidemiology, Hawaii, 78-5347
- Kidney Transplantation
 - Epidemiology, 78-5353
- Socioeconomic Factors
 - Epidemiology, 78-5351

Uterine Neoplasms

- Adenocarcinoma
 - Epidemiology, Review, 78-4867
 - Radiation Effects, 78-5305
 - Turner's Syndrome, 78-5342
- Diet
 - Epidemiology, Review, 78-4862
- Estrogens
 - Epidemiology, Canada, 78-5355
 - Epidemiology, Review, 78-4862, 78-4867, 78-4910
 - Menopause, 78-5355
- Hyperplasia
 - Precancerous Conditions, Review, 78-4921
- Neoplasms, Multiple Primary
 - Genetics, 78-5371
- Polyps
 - Precancerous Conditions, Review, 78-4921
- Precancerous Conditions
 - Epidemiology, Review, 78-4862
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Epidemiology, Review, 78-4867
- Turner's Syndrome
 - Case Report, 78-5342
 - Estrogens, 78-5342
 - Karyotyping, 78-5342
 - Progesterone, 78-5342

Vagina

- Kidney Neoplasms
 - Neoplasm Metastasis, 78-5335

Vinblastine

- Lymphocytes
 - Chromosome Aberrations, 78-5157

Vincristine

- Lymphocytes
 - Chromosome Aberrations, 78-5157

Viral Proteins

- Lymphoma
 - Virus, C-Type RNA Tumor, 78-5210
- Virus, Abelson Murine Leukemia
 - Antigenic Determinants, 78-5199
 - Cell Transformation, Neoplastic, 78-5197
 - Isolation and Characterization, 78-5197, 78-5199
 - Isolation and Characterization, Mink Cells, 78-5201
 - RNA, Viral, 78-5197
- Virus, Avian Carcinoma
 - Antigenic Determinants, 78-5190
 - Isolation and Characterization, MH2 Virus, 78-5190
- Virus, Baboon C-Type RNA Tumor
 - Antigenic Determinants, Review, 78-4898

Viral Proteins (cont'd)

- Virus, C-Type RNA Tumor
 - Cell Membrane, 78-5210
 - Cross-Linking, 78-5210
 - Fibroblasts, Mouse, 78-5210
 - Immunity, Cellular, 78-5272
 - Propionamide, Dimethyl-3,3'-dithiobis-, Dihydrochloride, 78-5210
 - RNA, Messenger, Review, 78-4880
 - Thymus Gland, Mouse, 78-5210
- Virus, Friend Spleen Focus-Forming
 - Isolation and Characterization, 78-5208
 - Virus, Helper, 78-5208
- Virus, Mason-Pfizer Monkey
 - Antigenic Determinants, Review, 78-4898
- Virus, Mink Cell Focus-Inducing
 - Isolation and Characterization, Mink Cells, 78-5201
- Virus, Moloney Murine Sarcoma
 - Immunoprecipitation, 78-5201
- Virus, Murine Leukemia
 - Cell Membrane, 78-5210
 - Cross-Linking, 78-5210
 - Propionamide, Dimethyl-3,3'-dithiobis-, Dihydrochloride, 78-5210
- Virus, Radiation Leukemia
 - Isolation and Characterization, 78-5225
- Virus, Rauscher Murine Leukemia
 - Antigenic Determinants, 78-5202
 - Canavanine, 78-5202
 - RNA, Messenger, 78-5202
 - RNA, Viral, 78-5202
- Virus, Rous Sarcoma
 - Antigenic Determinants, 78-5193
 - Cell Transformation, Neoplastic, 78-5193
 - Phosphotransferases, 78-5188
 - RNA, Viral, 78-5188

Viral Vaccines

- Virus, Marek's Disease Herpes
- Virus, Turkey Herpes, 78-5196

Virus, Abelson Murine Leukemia

- Antigenic Determinants
 - Cell Transformation, Neoplastic, 78-5199
- Antigens, Viral
 - Antibody Specificity, 78-5200
 - Cell Differentiation, 78-5200
 - Cell Transformation, Neoplastic, 78-5200
- Cell Transformation, Neoplastic
 - Immunoprecipitation, 78-5201
- Lymphoma
 - Age Factors, 78-5198
 - Antibody Specificity, 78-5200
 - Antigens, Viral, 78-5200
 - Dose-Response Study, 78-5198
 - Strain Difference, 78-5198
- Viral Proteins
 - Antigenic Determinants, 78-5199
 - Cell Transformation, Neoplastic, 78-5197
 - Isolation and Characterization, 78-5197, 78-5199
 - Isolation and Characterization, Mink Cells, 78-5201
 - RNA, Viral, 78-5197
- Virus, Friend Murine Leukemia
 - Antigenic Determinants, 78-5200
- Virus, Helper
 - RNA, Viral, 78-5197
- Virus, Mink Cell Focus-Inducing
 - Immunoprecipitation, 78-5201

Virus, Abelson Murine Leukemia (cont'd)

- Virus, Moloney Murine Leukemia
 - Antigenic Determinants, 78-5200
- Virus, Helper, 78-5197
- Virus, Rauscher Murine Leukemia
 - Antigenic Determinants, 78-5200
- Virus, RD-114
 - Pseudotype Formation, 78-5199

Virus, Adeno

- Antigens, Viral
 - Immune Response, Review, 78-4905
- T-Lymphocytes
 - Immune Response, Review, 78-4905
- Urine
 - Kidney Transplantation, 78-5275

Virus, Adeno 2

- Benzimidazole, 5,6-Dichloro-1- β -D-ribofuranosyl-
 - DNA Replication, 78-5233
 - RNA, Viral, 78-5233
- Chromosomes, Human, 16-18
 - Chromosome Aberrations, Review, 78-4902
- DNA, Viral
 - Carcinogenic Activity, Review, 78-4902
- Immunosuppression
 - Carcinogenic Activity, Review, 78-4902
- Proteins
 - Cell Membrane, Review, 78-4902

Virus, Adeno 5

- Benz(a)anthracene, 7,12-Dimethyl-
 - Cell Transformation, Neoplastic, 78-5041
- Benzo(a)pyrene
 - Cell Transformation, Neoplastic, 78-5041
- 12-O-Tetradecanoylphorbol-13-acetate
 - Cell Transformation, Neoplastic, 78-5041

Virus, Adeno 7

- Acetic Acid, Lead Salt
 - Cell Transformation, Neoplastic, 78-5117

Virus, Adeno 12

- Antigenic Determinants
 - Cell Transformation, Neoplastic, 78-5238
- Antigens, Viral
 - Isolation and Characterization, 78-5238
 - Temperature Sensitive Mutants, 78-5238
- Cell Transformation, Neoplastic
 - Brain Cell Cultures, 78-5236
- DNA, Viral
 - Cell Transformation, Neoplastic, 78-5237
 - DNA-DNA Hybridization, 78-5237
 - Endonucleases, 78-5237
 - Hamster Cell Lines, 78-5237
- Gastrointestinal Neoplasms
 - Carcinogenic Potential, 78-5235
 - DNA-DNA Hybridization, 78-5235
- Glycoproteins
 - Cell Membrane, 78-5234
 - Cytocidal Mutants, 78-5234
- Lung Neoplasms
 - Carcinogenic Potential, 78-5235
 - DNA-DNA Hybridization, 78-5235
- Transferases
 - Cytocidal Mutants, 78-5234
 - Microsomes, 78-5234

Virus, Avian Carcinoma

- Fibroblasts

- 1, Avian Carcinoma (cont'd)**
 - Cell Transformation, Neoplastic, 78-5190
- Macrophages**
 - Cell Transformation, Neoplastic, 78-5190
- Viral Proteins**
 - Antigenic Determinants, 78-5190
 - Isolation and Characterization, MH2 Virus, 78-5190
- Virus, Avian Leukosis**
 - Virus, Helper, 78-5190
- Virus, Helper**
 - Virus Replication, 78-5190
- 2, Avian Leukosis**
 - Virus, Avian Carcinoma
 - Virus, Helper, 78-5190
- 3, Avian Leukosis-Sarcoma**
 - Virus, Avian Myelocytomatosis
 - Virus, Helper, 78-5187
- 4, Avian Myeloblastosis**
 - DNA Replication
 - DNA Polymerase, 78-5217
 - DNA, Viral
 - DNA Replication, 78-5217
 - Polyribonucleotides
 - Ribonuclease, 78-5203
 - Ribonuclease
 - Reverse Transcriptase, 78-5203
 - RNA, Transfer
 - Reverse Transcriptase, 78-5189
 - RNA Replication, 78-5189
 - RNA, Viral
 - Ribonuclease, 78-5203
 - Virus, RNA Tumor
 - DNA, Viral, 78-5271
 - Reverse Transcriptase, 78-5271
- 5, Avian Myelocytomatosis**
 - Genes, Viral
 - MC29 Virus, 78-5187
 - Virus, Avian Leukosis-Sarcoma
 - Virus, Helper, 78-5187
 - Virus, Helper
 - Virus Replication, 78-5187
 - Virus Replication
 - Genes, Viral, 78-5187
 - MC29 Virus, 78-5187
 - Virus, Rous Sarcoma
 - Virus, Helper, 78-5187
- 6, Avian Sarcoma**
 - Cell Transformation, Neoplastic
 - Genes, Viral, Review, 78-4882
 - Replication Defective Mutants
 - Genes, Viral, Review, 78-4881
 - Isolation and Characterization, Review, 78-4881
 - Reverse Transcriptase
 - DNA-RNA Hybridization, 78-5191
 - RNA Replication
 - Genes, Viral, Review, 78-4882
 - RNA, Viral
 - DNA-RNA Hybridization, Review, 78-4880
 - 78-4882
 - Genes, Viral, Review, 78-4882
 - Temperature Sensitive Mutants
 - Genes, Viral, Review, 78-4881
 - Isolation and Characterization, Review, 78-4881
 - Virus, Recombinant, Review, 78-4881
- Virus, Baboon**
 - Cell Transformation, Neoplastic
 - Murine Sarcoma Virus Pseudotype, 78-5219
 - DNA, Viral
 - Chromosomes, Human, 6-12, 78-5231
 - Hybrid Cells
 - Antigens, Viral, 78-5231
 - Chromosomes, Human, 6-12, 78-5231
 - Reverse Transcriptase, 78-5231
 - Virus Replication, 78-5231
 - Leukemia
 - DNA-RNA Hybridization, Review, 78-4901
 - Leukemia, Myeloblastic
 - Virus-Like Particles, Review, 78-4901
 - Virus, Kirsten Murine Sarcoma
 - Virus, Helper, 78-5219
 - Virus Replication
 - Chromosomes, Human, 6-12, 78-5231
- Virus, Baboon C-Type RNA Tumor**
 - Viral Proteins
 - Antigenic Determinants, Review, 78-4898
 - Virus, D-Type RNA Tumor
 - Antigenic Determinants, Review, 78-4898
- Virus, Bovine Leukemia**
 - Antibodies, Viral
 - Seroepidemiology, Review, 78-4887
 - Antigen-Antibody Reactions
 - Diagnosis, Review, 78-4886
 - C-Type Particles
 - Isolation and Characterization, Review, 78-4886
 - Chondrosarcoma
 - DNA-DNA Hybridization, 78-5230
 - DNA, Viral
 - Cell Transformation, Neoplastic, 78-5230
 - DNA-DNA Hybridization, 78-5230
 - Leiomyosarcoma
 - DNA-DNA Hybridization, 78-5230
 - Leukemia, Lymphocytic
 - DNA-DNA Hybridization, 78-5230
 - Lymphocytosis
 - Epidemiology, 78-4886
 - Lymphosarcoma
 - DNA-DNA Hybridization, 78-5230
 - Epidemiology, 78-4886
 - Horizontal/Vertical Transmission, Review, 78-4886
 - Milk
 - Horizontal Transmission, Review, 78-4887
 - Precancerous Conditions
 - Epidemiology, 78-4886
 - Rhabdomyosarcoma
 - DNA-DNA Hybridization, 78-5230
 - Sarcoma, Osteogenic
 - DNA-DNA Hybridization, 78-5230
- Virus, C-Type RNA Tumor**
 - DNA, Neoplasm
 - DNA, Viral, 78-5274
 - Epidermal Growth Factor
 - Carcinogenic Activity, Review, 78-4900
 - Hematopoietic Stem Cells
 - Carcinogenic Activity, Review, 78-4901
 - Leukemia
 - DNA, Neoplasm, 78-5274
 - DNA, Viral, 78-5274
 - Horizontal Transmission, Review, 78-4900
 - B-Lymphocytes, 78-5270
 - T-Lymphocytes, 78-5270

- Virus, C-Type RNA Tumor (cont'd)**
 - Null Cells, 78-5270
 - Reverse Transcriptase, 78-5270
 - Trans-Species Infection, 78-5274
 - Transformation, Genetic, 78-5274
 - Ultrastructural Study, 78-5270
 - Virus-Like Particles, Review, 78-4901
- Leukemia, Myelocytic
 - Colony Stimulating Factor, Review, 78-4933
- T-Lymphocytes
 - Immunosuppression, 78-5272
 - Suppressive Component, Isolation and Characterization, 78-5272
- Lymphoma
 - Viral Proteins, 78-5210
- Neoplasms
 - Isolation and Characterization, Review, 78-4897
- Propionamide, Dimethyl-3,3'-dithiobis-, Dihydrochloride
 - Viral Proteins, 78-5210
- RNA, Viral
 - DNA-RNA Hybridization, Review, 78-4880
- Sarcoma
 - DNA, Neoplasm, 78-5274
 - DNA, Viral, 78-5274
 - Trans-Species Infection, 78-5274
 - Transformation, Genetic, 78-5274
- Ultraviolet Rays
 - Virus Activation, Mouse, 78-5212
- Viral Proteins
 - Cell Membrane, 78-5210
 - Cross-Linking, 78-5210
 - Fibroblasts, Mouse, 78-5210
 - Immunity, Cellular, 78-5272
 - RNA, Messenger, Review, 78-4880
 - Thymus Gland, Mouse, 78-5210
- Virus Activation
 - Carcinogenic Potential, Review, 78-4897
- Virus, Coxsackie B**
 - Hypercholesteremia
 - Antibody Formation, 78-5290
 - Immune Response, 78-5290
 - Macrophages, 78-5290
- Virus Cultivation**
 - Virus, Murine Sarcoma
 - Cells, Cultured, 78-5218
- Virus, Cytomegalo**
 - Virus, Herpes Simplex 1
 - Viral Interference, 78-5240
 - Virus Replication, 78-5240
 - Virus, Herpes Simplex 2
 - Viral Interference, 78-5240
 - Virus Replication, 78-5240
 - Virus Replication
 - Cell Transformation, Neoplastic, 78-5240
- Virus, D-Type RNA Tumor**
 - Mammae, Monkey
 - Isolation and Characterization, Review, 78-4898
 - Placenta, Monkey
 - Isolation and Characterization, Review, 78-4898
 - Virus, Baboon C-Type RNA Tumor
 - Antigenic Determinants, Review, 78-4898
 - Virus, Mason-Pfizer Monkey
 - Antigenic Determinants, Review, 78-4898
- Virus, DNA Tumor**
 - DNA-RNA Hybridization
- Virus, DNA Tumor (cont'd)**
 - Carcinogenic Potential, Review, 78-4903
- T-Lymphocytes
 - Immune Response, Review, 78-4907
- Neoplasms
 - Isolation and Characterization, Review, 78-4897
- Virus Activation
 - Carcinogenic Potential, Review, 78-4897
- Virus-Like Particles
 - Kidney Transplantation, 78-5275
- Virus, Epstein-Barr**
 - Anti-Antibodies
 - Binding, Cell Membrane, 78-5278
 - Burkitt's Lymphoma
 - DNA-RNA Hybridization, 78-5241
 - Epidemiology, Review, 78-4895, 78-4911, 78-4913
 - Immune Response, Review, 78-4913
 - Seroepidemiology, Review, 78-4896
 - Carcinogen, Environmental
 - Co-carcinogenic Activity, Review, 78-4904
 - Chromosomes, Human, 13-15
 - Chromosome Aberrations, Review, 78-4895
 - DNA-RNA Hybridization
 - Carcinogenic Potential, Review, 78-4903
 - Genetics
 - Epidemiology, Review, 78-4904
 - Hodgkin's Disease
 - DNA-RNA Hybridization, 78-5241
 - Epidemiology, Review, 78-4913
 - IgM
 - Anti-Antibodies, 78-5278
 - Infectious Mononucleosis
 - Seroepidemiology, Review, 78-4896
 - Lymphoma
 - Immunoglobulins, Surface, 78-5278
 - Malaria
 - Epidemiology, Review, 78-4895, 78-4904
 - Nasopharyngeal Neoplasms
 - Carcinoma, 78-5241
 - Epidemiology, Alaska, 78-5380
 - Epidemiology, Review, 78-4895
 - Seroepidemiology, Review, 78-4896
 - Parasites
 - Carcinogenic Activity, Review, 78-4911
 - Sarcoma, Kaposi's
 - DNA-RNA Hybridization, 78-5241
- Virus, Feline Leukemia**
 - Antibodies, Viral
 - Immune Response, Review, 78-4885
 - Antigen-Antibody Reactions
 - Guinea Pig, 78-5228
 - Hypotonic Medium, 78-5227
 - Antigens, Viral
 - Immune Response, Review, 78-4884
 - Viral Vaccines, Review, 78-4884
 - Cell Transformation, Neoplastic
 - Guinea Pig, 78-5228
 - Formaldehyde
 - Viral Vaccines, Review, 78-4884
 - Horizontal Transmission
 - Seroepidemiology, Review, 78-4885
 - T-Lymphocytes
 - Immune Response, Review, 78-4907
 - Neutralization
 - Hypotonic Medium, 78-5227

- us, Feline Sarcoma**
 - Antibodies, Viral
 - Immune Response, Review, 78-4885
 - Antigen-Antibody Reactions
 - Guinea Pig, 78-5228
 - Cell Transformation, Neoplastic
 - Guinea Pig, 78-5228
 - Horizontal Transmission
 - Seroepidemiology, Review, 78-4885
- us, Friend Murine Leukemia**
 - Antibody Formation
 - Immunosuppression, 78-5207
 - Antigen-Antibody Reactions
 - Cells, Cultured, 78-5283
 - Lipopolysaccharides
 - Antigen-Antibody Reactions, 78-5207
 - Immunosuppression, 78-5207
 - Lymphocyte Transformation
 - Cells, Cultured, 78-5283
 - B-Lymphocytes
 - Antigen-Antibody Reactions, 78-5283
 - T-Lymphocytes
 - Lymphocyte Depletion, 78-5283
 - Macrophages
 - Antigen-Antibody Reactions, 78-5207
 - RNA, Viral
 - DNA-RNA Hybridization, 78-5206
 - Virus, Abelson Murine Leukemia
 - Antigenic Determinants, 78-5200
 - Virus, Friend Spleen Focus-Forming
 - Antigenic Determinants, 78-5208
 - Virus, Recombinant, 78-5205, 78-5206
 - Virus, Mink Cell Focus-Inducing
 - Antigenic Determinants, 78-5205
 - Isolation and Characterization, 78-5206
 - Virus, Recombinant, 78-5206
- us, Friend Spleen Focus-Forming**
 - Antigenic Determinants
 - Radioimmunoassay, 78-5205
 - Viral Proteins
 - Isolation and Characterization, 78-5208
 - Virus, Friend Murine Leukemia
 - Antigenic Determinants, 78-5208
 - Virus, Recombinant, 78-5205, 78-5206
 - Virus, Helper
 - Viral Proteins, 78-5208
 - Virus, Mink Cell Focus-Inducing
 - Antigenic Determinants, 78-5205
- us, Gross Murine Leukemia**
 - Autoimmune Diseases
 - Precancerous Conditions, 78-5280
 - Dysgammaglobulinemia
 - IgA, 78-5280
 - Mouse, AKR, 78-5280
 - Lymphoma
 - Transplantation Immunology, 78-5291
 - Virus, Radiation Leukemia
 - DNA-RNA Hybridization, 78-5225
- us, Helper**
 - Virus, Abelson Murine Leukemia
 - RNA, Viral, 78-5197
 - Virus, Moloney Murine Leukemia, 78-5197
 - Virus, Avian Carcinoma
 - Virus, Avian Leukosis, 78-5190
 - Virus Replication, 78-5190
- Virus, Helper (cont'd)**
 - Virus, Avian Myelocytomatosis
 - Virus, Avian Leukosis-Sarcoma, 78-5187
 - Virus Replication, 78-5187
 - Virus, Rous Sarcoma, 78-5187
 - Virus, Friend Spleen Focus-Forming
 - Viral Proteins, 78-5208
 - Virus, Kirsten Murine Sarcoma
 - Cell Transformation, Neoplastic, 78-5211
 - Virus Activation, 78-5219
 - Virus, Baboon, 78-5219
 - Virus, Murine Sarcoma
 - Caffeine, 78-5212
 - Ultraviolet Rays, 78-5212
 - Uridine, 5-Bromo-2'-deoxy-, 78-5212
 - Uridine, 2'-Deoxy-5-iodo-, 78-5212
 - Virus Activation, Mouse, 78-5212
- Virus, Hepatitis**
 - Hepatitis
 - Epidemiology, Review, 78-4839
 - Mycotoxins
 - Co-carcinogenic Activity, Review, 78-4839
- Virus, Herpes**
 - Cervix Neoplasms
 - Epidemiology, 78-5341
 - Neisseria, 78-5341
 - Urine
 - Kidney Transplantation, 78-5275
- Virus, Herpes Ateles**
 - Lymphocyte Transformation
 - Marmoset, 78-5239
 - T-Lymphocytes
 - Antilymphocyte Serum, 78-5239
 - Lymphocyte Transformation, 78-5239
 - Rosette Formation, 78-5239
 - Radiation, Ionizing
 - Lymphocyte Transformation, 78-5239
- Virus, Herpes Simplex**
 - DNA-RNA Hybridization
 - Carcinogenic Potential, Review, 78-4903
- Virus, Herpes Simplex 1**
 - DNA, Viral
 - Temperature Sensitive Mutants, 78-5244
 - Virus, Recombinant, 78-5244
 - Virus Rescue, 78-5244
 - Virus, Cytomegalo
 - Viral Interference, 78-5240
 - Virus Replication, 78-5240
 - Virus, Herpes Simplex 2
 - Virus, Recombinant, 78-5244
 - Virus Rescue
 - Temperature Sensitive Mutants, 78-5244
 - Xeroderma Pigmentosum
 - Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-, 78-5168
 - Caffeine, 78-5168
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-5168
 - Reactivation Kinetics, 78-5168
 - Ultraviolet Rays, 78-5168
 - Virus Replication, 78-5168
- Virus, Herpes Simplex 2**
 - Antigens, Viral
 - Precancerous Conditions, 78-5242
 - Cervix Neoplasms
 - Animal Model, Mouse, Rat, 78-5242

Virus, Herpes Simplex 2 (cont'd)

- Antigen-Antibody Reactions, Review, 78-4888
- Antigens, Viral, 78-5242
- Carcinoma In Situ, 78-4888
- Neoplasm Metastasis, 78-4888
- Precancerous Conditions, 78-5242
- Seroepidemiology, Review, 78-4888

Cytopathogenic Effect, Viral

- Cervix, Mouse, Rat, 78-5242

DNA, Viral

- Virus Rescue, 78-5244

Immunity, Cellular

- BCG, 78-5243
- Plant Agglutinins, 78-5243
- Skin Graft, Footpad Tests, 78-5243
- Tuberculin, 78-5243

Lymphocyte Transformation

- Spleen Cells, 78-5243

Plant Agglutinins

- Lymphocyte Transformation, 78-5243

Sensory Ganglia, Pelvis

- Animal Model, Mouse, Rat, 78-5242

Virus, Cytomegalo

- Viral Interference, 78-5240
- Virus Replication, 78-5240

Virus, Herpes Simplex 1

- Virus, Recombinant, 78-5244

Virus, Kirsten Murine Sarcoma**Bone Marrow Cells**

- Cell Transformation, Neoplastic, 78-5211

Cell Transformation, Neoplastic

- C-Type Particles, 78-5219
- Spleen Cells, Gorilla, 78-5219

Virus, Baboon

- Virus, Helper, 78-5219

Virus, Helper

- Cell Transformation, Neoplastic, 78-5211
- Virus Activation, 78-5219

Virus-Like Particles**Carcinoma, Ehrlich Tumor**

- A- and C-Type Particles, 78-5273
- Ultrastructural Study, 78-5273

Leukemia

- Dogs, 78-5229
- Precancerous Conditions, 78-5204

Lymphosarcoma

- Dogs, 78-5229

Mammary Neoplasms, Experimental

- Dogs, 78-5229

Melanoma

- DNA-RNA Hybridization, 78-5221
- Isolation and Characterization, B16 Tumor, 78-5221
- Virus, RNA Tumor, 78-5221

Virus, DNA Tumor

- Kidney Transplantation, 78-5275

Virus, Marek's Disease Herpes**Immunosuppression**

- Precancerous Conditions, 78-5196

T-Lymphocytes

- Immunosuppression, 78-5196

Virus, Turkey Herpes

- T-Lymphocytes, 78-5196
- Viral Vaccines, 78-5196

Virus, Mason-Pfizer Monkey**HeLa Cells****Virus, Mason-Pfizer Monkey (cont'd)**

- Virus Replication, 78-5232

Viral Proteins

- Antigenic Determinants, Review, 78-4898

Virus, D-Type RNA Tumor

- Antigenic Determinants, Review, 78-4898

Virus Replication

- Epithelial Cells, 78-5232
- Fibroblasts, 78-5232

Virus, Mink Cell Focus-Inducing**Viral Proteins**

- Isolation and Characterization, Mink Cells, 78-520

Virus, Abelson Murine Leukemia

- Immunoprecipitation, 78-5201

Virus, Friend Murine Leukemia

- Antigenic Determinants, 78-5205
- Isolation and Characterization, 78-5206
- Virus, Recombinant, 78-5206

Virus, Friend Spleen Focus-Forming

- Antigenic Determinants, 78-5205

Virus, Moloney Murine Leukemia

- Antigenic Determinants, 78-5205

Virus, Murine Leukemia

- Antigenic Determinants, 78-5205
- Virus, Recombinant, 78-5206

Virus, Moloney Murine Leukemia**DNA Replication**

- DNA Polymerase, 78-5217

DNA, Viral

- DNA Replication, 78-5217

Reverse Transcriptase

- Competition Radioimmunoassay, 78-5216
- Mouse 3T3 Clone Cells, 78-5215

RNA, Viral

- Heteroduplex Structures, 78-5214
- Mouse 3T3 Clone Cells, 78-5215

Virus, Abelson Murine Leukemia

- Antigenic Determinants, 78-5200
- Virus, Helper, 78-5197

Virus, Mink Cell Focus-Inducing

- Antigenic Determinants, 78-5205

Virus, Murine Leukemia

- Virus, Recombinant, 78-5214

Virus, Recombinant

- DNA-RNA Hybridization, 78-5214
- Heteroduplex Structures, 78-5214

Virus, Moloney Murine Sarcoma**Cell Transformation, Neoplastic**

- Growth Factors, Isolation and Characterization 78-5220

DNA Replication

- DNA Polymerase, 78-5217

DNA, Viral

- DNA Replication, 78-5217

Nucleotide Sequence

- Clone Cells, 78-5213

Peptides

- Growth Factors, Isolation and Characterization 78-5220

RNA, Viral

- Nucleotide Sequence, 78-5213

Viral Proteins

- Immunoprecipitation, 78-5201

Virus, Murine Leukemia

- A-, E-, X-Tropic Viruses

is, Murine Leukemia (cont'd)

- Isolation and Characterization, Review, 78-4883
- Antibody Formation
 - Immune Response, Wild Mouse, Review, 78-4883
- Hematopoietic Stem Cells
 - Cell Differentiation, 78-5211
- Leukemia
 - Horizontal Transmission, 78-4899
- Lymphoma
 - Epidemiology, Wild Mouse, Review, 78-4883
 - Vertical Transmission, 78-5209
- Lymphosarcoma
 - Horizontal Transmission, 78-4899
- Paralysis
 - Epidemiology, Wild Mouse, Review, 78-4883
- Propionamide, Dimethyl-3,3'-dithiobis-, Dihydrochloride
 - Viral Proteins, 78-5210
- Vertical Transmission
 - Breeding, Selective, 78-5209
- Viral Proteins
 - Cell Membrane, 78-5210
 - Cross-Linking, 78-5210
- Viremia
 - Vertical Transmission, 78-5209
- Virus, Mink Cell Focus-Inducing
 - Antigenic Determinants, 78-5205
 - Virus, Recombinant, 78-5206
- Virus, Moloney Murine Leukemia
 - Virus, Recombinant, 78-5214

is, Murine Mammary Tumor

- Antigens, Viral
 - Radioimmunoassay, 78-5223
- Dexamethasone
 - Chromosomes, 78-5222
 - DNA, Viral, 78-5222
 - RNA Replication, 78-5222
- DNA, Viral
 - HTC Clone Cells, 78-5222
- Mammary Neoplasms, Experimental
 - Dexamethasone, 78-5143

is, Murine Sarcoma

- Cell Transformation, Neoplastic
 - Cells, Cultured, 78-5218
- Fibroblasts
 - Cells, Cultured, 78-5218
- RNA, Viral
 - DNA-RNA Hybridization, Review, 78-4880
- Sarcoma
 - T-Lymphocytes, 78-5287
 - Macrophages, 78-5287
- Virus Cultivation
 - Cells, Cultured, 78-5218
- Virus, Helper
 - Caffeine, 78-5212
 - Ultraviolet Rays, 78-5212
 - Uridine, 5-Bromo-2'-deoxy-, 78-5212
 - Uridine, 2'-Deoxy-5-iodo-, 78-5212
 - Virus Activation, Mouse, 78-5212

us, Papilloma

- Condylomata Acuminata
 - Epidemiology, Review, 78-4889
- Connective Tissue
 - Carcinogenic Activity, Review, 78-4889
- Epithelium
 - Carcinogenic Activity, Review, 78-4889
- Laryngeal Neoplasms

Virus, Papilloma (cont'd)

- Epidemiology, Review, 78-4889
- Mouth Neoplasms
 - Epidemiology, Review, 78-4889
- Skin Neoplasms
 - Precancerous Conditions, Review, 78-4889

Virus, Polyoma

- Actin
 - Cell Transformation, Neoplastic, 78-5245
- Airborne Virus
 - Quantitation Method, 78-5269
- Antigens, Viral
 - Temperature Sensitive Mutants, 78-5265
- Benz(a)anthracene, 7,12-Dimethyl-
 - Antigens, Viral, 78-4999
- Benzo(a)pyrene
 - Antigens, Viral, 78-4999
- Brain Neoplasms
 - Clone Cells, 78-5266
- Cell Division
 - Host-Range Mutants, 78-5268
 - Temperature Sensitive Mutants, 78-5268
- Cell Transformation, Neoplastic
 - Antigenic Determinants, Review, 78-4892
 - Hamster Brain Cells, 78-5266
 - Protein A, Review, 78-4892
 - Virus Activation, 78-5265
- Cholanthrene, 3-Methyl-
 - Antigens, Viral, 78-4999
- Chromosomes
 - Virus Replication, 78-5256
- Dibenz(a,h)anthracene
 - Antigens, Viral, 78-4999
- DNA, Binding
 - Protein A, Review, 78-4892
- DNA Replication
 - Cell Cycle Kinetics, 78-5268
 - Host-Range Mutants, 78-5268
 - Temperature Sensitive Mutants, 78-5265
- DNA, Viral
 - Complementation Group A Mutants, 78-4892
 - DNA-RNA Hybridization, 78-5267
 - Endonucleases, 78-5265
 - Nucleosomes, 78-5256
- Horizontal Transmission
 - Airborne Virus, 78-5269
- Immune Serums
 - Virus Activation, 78-5265
- Neuroglia
 - Cell Transformation, Neoplastic, 78-5266
- RNA, Viral
 - DNA Replication, 78-5267
- Tubulin
 - Cell Transformation, Neoplastic, 78-5245
- Urine
 - Kidney Transplantation, 78-5275
- Virus Replication
 - Ultrastructural Study, 78-5268
- Virus, SV40
 - Antigenic Determinants, Review, 78-4892

Virus, Polyoma, BK

- Adenocarcinoma
 - Carcinogenic Activity, Hamster, 78-5263
- Anemia, Aplastic
 - Bone Marrow Transplantation, 78-5264
- Angiosarcoma

Virus, Polyoma, BK (cont'd)

- Carcinogenic Activity, Hamster, 78-5263
- Antigenic Determinants
 - Isolation and Characterization, Review, 78-4893
- Bone Marrow Transplantation
 - Case Report, 78-5264
- Carcinogenic Activity
 - Hamster, Review, 78-4891
- Cell Transformation, Neoplastic
 - Plaque Assay, Review, 78-4893
- Disgerminoma
 - Carcinogenic Activity, Hamster, 78-5263
- DNA-RNA Hybridization
 - Nucleosomes, 78-5262
- DNA, Viral
 - Isolation and Characterization, 78-5262
 - Isolation and Characterization, Review, 78-4893
- Ependymoma
 - Antigen-Antibody Reactions, 78-5263
 - Carcinogenic Activity, Hamster, 78-5263
 - Neoplasm Transplantation, 78-5263
- Histocompatibility Antigens
 - Isolation and Characterization, Review, 78-4891
- Immunosuppression
 - Isolation and Characterization, Review, 78-4890
- Islet Cell Tumor
 - Calcitonin, 78-5263
 - Glucagon, 78-5263
 - Insulin, 78-5263
- Lymphoma
 - Carcinogenic Activity, Hamster, 78-5263
- Pancreatic Neoplasms
 - Carcinogenic Activity, Hamster, 78-5263
 - Islet Cell Tumor, 78-5263
- Sarcoma, Osteogenic
 - Carcinogenic Activity, Hamster, 78-5263
- Virus, Polyoma, JC
 - DNA-DNA Hybridization, Review, 78-4890
 - 78-4893
- Virus, SV40
 - Antigenic Determinants, Review, 78-4891
 - DNA-DNA Hybridization, Review, 78-4890
 - 78-4893

Virus, Polyoma, JC

- Carcinogenic Activity
 - Hamster, Review, 78-4891
- Eye Neoplasms
 - Carcinogenic Activity, Hamster, 78-5261
 - Carcinoma, 78-5261
 - Ependymoma, 78-5261
 - Fibrosarcoma, 78-5261
 - Neurilemmoma, 78-5261
 - Retinoblastoma, 78-5261
- Heart Neoplasms
 - Carcinogenic Activity, Hamster, 78-5261
- Histocompatibility Antigens
 - Isolation and Characterization, Review, 78-4891
- Leukoencephalopathy, Progressive Multifocal
 - Isolation and Characterization, Review, 78-4890
 - 78-4891
- Nervous System Neoplasms
 - Carcinogenic Activity, Hamster, 78-5261
- Stomach Neoplasms
 - Carcinogenic Activity, Hamster, 78-5261
- Virus, Polyoma, BK
 - DNA-DNA Hybridization, Review, 78-4890
 - 78-4893

Virus, Polyoma, JC (cont'd)

- Virus, SV40
 - Antigenic Determinants, Review, 78-4891

Virus, Radiation Leukemia

- Bone Marrow Cells
 - Precancerous Conditions, 78-5224
- Cell Differentiation
 - Bone Marrow, 78-5295
 - Virus Variants, 78-5224
- Cell Transformation, Neoplastic
 - Thymopoietins, 78-5295
- DNA, Viral
 - DNA-DNA Hybridization, Spleen Cells, 78-5224
- T-Lymphocytes
 - Cell Transformation, Neoplastic, 78-5295
 - Precancerous Conditions, 78-5295
 - Transplantation Immunology, 78-5295
- Radiation, Ionizing
 - Cell Transformation, Neoplastic, 78-5295
 - Precancerous Conditions, 78-5224
- Thymopoietins
 - Cell Differentiation, 78-5295
- Thymus Gland
 - Precancerous Conditions, 78-5224
- Viral Proteins
 - Isolation and Characterization, 78-5225
- Virus, Gross Murine Leukemia
 - DNA-RNA Hybridization, 78-5225
- Virus, Rauscher Murine Leukemia
 - DNA-RNA Hybridization, 78-5225

Virus, Rauscher Murine Leukemia

- Hematopoietic Stem Cells
 - Cell Differentiation, 78-5211
- Polyribonucleotides
 - Ribonuclease, 78-5203
- Reverse Transcriptase
 - Ultrastructural Study, Crude Tissue, 78-5204
- RNA, Viral
 - Ribonuclease, 78-5203
 - Viral Proteins, 78-5202
- Viral Proteins
 - Antigenic Determinants, 78-5202
 - Canavanine, 78-5202
 - RNA, Messenger, 78-5202
- Virus, Abelson Murine Leukemia
 - Antigenic Determinants, 78-5200
- Virus, Radiation Leukemia
 - DNA-RNA Hybridization, 78-5225

Virus, RD-114

- Burkitt's Lymphoma
 - DNA-RNA Hybridization, 78-5226
- DNA, Viral
 - DNA Polymerase, 78-5226
 - DNA-RNA Hybridization, 78-5226
 - Isolation and Characterization, 78-5226
- Leukemia, Myeloblastic
 - DNA-RNA Hybridization, 78-5226
- Rhabdomyosarcoma
 - DNA-RNA Hybridization, 78-5226
- Virus, Abelson Murine Leukemia
 - Pseudotype Formation, 78-5199

Virus Replication

- Virus, Avian Carcinoma
 - Virus, Helper, 78-5190
- Virus, Avian Myelocytomatosis

rus Replication (cont'd)

- Genes, Viral, 78-5187
- MC29 Virus, 78-5187
- Virus, Helper, 78-5187

Virus, Baboon

- Chromosomes, Human, 6-12, 78-5231
- Hybrid Cells, 78-5231

Virus, Cytomegalo

- Cell Transformation, Neoplastic, 78-5240
- Virus, Herpes Simplex 1, 78-5240
- Virus, Herpes Simplex 2, 78-5240

Virus, Mason-Pfizer Monkey

- Epithelial Cells, 78-5232
- Fibroblasts, 78-5232
- HeLa Cells, 78-5232

Virus, Polyoma

- Chromosomes, 78-5256
- Ultrastructural Study, 78-5268

Virus, SV40

- Chromosomes, 78-5256
- DNA, Viral, 78-5250
- Nucleoprotein Complex, 78-5247

Xeroderma Pigmentosum

- Virus, Herpes Simplex 1, 78-5168

irus, RNA Tumor**Leukemia**

- Isolation and Characterization, 78-5271

Melanoma

- Virus-Like Particles, 78-5221

1H-Pyrrole-2,5-dione, 1-Ethyl-

- Reverse Transcriptase, 78-5216

Reverse Transcriptase

- Antigenic Determinants, 78-5216
- Carcinogenic Potential, Review, 78-4903
- Competition Radioimmunoassay, 78-5216
- Ultrastructural Study, Crude Tissue, 78-5204

Virus, Avian Myeloblastosis

- DNA, Viral, 78-5271
- Reverse Transcriptase, 78-5271

irus, Rous Sarcoma**Actin**

- Temperature Sensitive Mutants, 78-5195

Antigenic Determinants

- Cell-Free Synthesis, 78-5193

Cell Differentiation

- Neuroretinal Cells, 78-5194
- Transformation Defective Mutants, 78-5194

Cell Transformation, Neoplastic

- Cell Adhesion, Review, 78-4932
- Genes, Viral, 78-5188
- Temperature Sensitive Mutants, 78-5195

Genes, Viral

- Fibroblasts, Enucleated, 78-5195

Genetics

- Antibody Formation, 78-5292
- Immune Response, 78-5292
- Immunity, Cellular, 78-5292
- Immunosuppression, 78-5292
- Stress, 78-5292

Glycerides

- Cell Transformation, Neoplastic, 78-5192
- Transformation Defective Mutants, 78-5192

Graft Rejection

- Genetics, 78-5292

Phospholipids

- Cell Transformation, Neoplastic, 78-5192

Virus, Rous Sarcoma (cont'd)**Plasminogen Activators**

- Cell Transformation, Neoplastic, 78-5040

Protein Kinase

- Cell Transformation, Neoplastic, 78-5188

Proteins

- Cell Adhesion, Review, 78-4932
- Cell Membrane, 78-5195
- Temperature Sensitive Mutants, 78-5195

Radiation, Ionizing

- Antibody Formation, 78-5292
- Immunity, Cellular, 78-5292

RNA, Viral

- Protein Kinase, 78-5188
- Viral Proteins, 78-5188

Surface Properties

- Fibroblasts, Enucleated, 78-5195

Temperature Sensitive Mutants

- Hexose, Metabolism, 78-5195

12-O-Tetradecanoylphorbol-13-acetate

- Plasminogen Activators, 78-5040

Transformation Defective Mutants

- Isolation and Characterization, 78-5194

Viral Proteins

- Antigenic Determinants, 78-5193
- Cell Transformation, Neoplastic, 78-5193
- Phosphotransferases, 78-5188

Virus, Avian Myelocytomatosis

- Virus, Helper, 78-5187

Virus, Simian Sarcoma**Leukemia, Myeloblastic**

- Isolation and Characterization, Cell Culture 78-4900

- Virus-Like Particles, Review, 78-4901

Virus, SV40**Acetamide, N-(Acetyloxy)-N-fluoren-2-yl-**

- DNA, Binding, 78-4984

Actin

- Cell Transformation, Neoplastic, 78-5245

Amino Acids

- Cell Transformation, Neoplastic, 78-5259
- Contact Inhibition, 78-5259
- Membrane Transport, 78-5259

Anchorage Independent Growth

- Deletion Mutants, 78-5252

Antigens, Viral

- Antibody Specificity, 78-5251
- Antigenic Determinants, 78-5251
- Carcinogenic Activity, Review, 78-4894
- Cell Cycle Kinetics, 78-5258
- DNA Replication, 78-5248
- DNA Replication, Review, 78-4894
- Immune Response, Hamster, 78-5251
- Immune Response, Review, 78-4905
- Immune Serums, 78-5251
- Phenotype Complementation, 78-5248
- RNA, Messenger, 78-5249
- RNA Replication, 78-5258
- RNA, Ribosomal, 78-5258
- Temperature Sensitive Mutants, 78-5248

Benzene, (Epoxyethyl)-

- DNA, Binding, 78-4984

Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-

- DNA, Binding, 78-4984
- Cell Transformation, Neoplastic

Virus, SV40 (cont'd)

- Antigenic Determinants, Review, 78-4892
- Deletion Mutants, 78-5252
- Protein A, Review, 78-4892
- Temperature Sensitive Mutants, 78-5252
- Chromatin
 - DNA, Binding, 78-5257
 - Histones, 78-5254
 - Isolation and Characterization, CV1 Cells, 78-5254
 - Nucleosomes, 78-5257
- Chromosomes
 - Nucleosomes, 78-5257
 - Virus Replication, 78-5256
- Cobalt Radioisotopes
 - DNA Replication, 78-5253
- Deletion Mutants
 - Agar Assay, 78-5252
- DNA, Binding
 - Protein A, Review, 78-4892
- DNA Polymerase
 - Nucleoprotein Complex, 78-5246, 78-5247
- DNA Replication
 - Antigens, Viral, 78-5253
 - Cell Transformation, Neoplastic, 78-5253
 - DNA-RNA Hybridization, 78-5253
 - Temperature Sensitive Mutants, 78-5248
- DNA, Viral
 - Cell Transformation, Neoplastic, 78-5252
 - Complementation Group A Mutants, 78-4892
 - Nucleoprotein Complex, 78-5246, 78-5247
 - Nucleoproteins, 78-5254
 - Nucleosomes, 78-5256
 - Nucleotide Sequence, 78-5249, 78-5396
 - RNA Polymerase, 78-5250
 - RNA Replication, 78-5250, 78-5260
 - Virus Replication, 78-5250
- Fibroblasts
 - Anchorage Independent Growth, 78-5252
- Histones
 - Nucleoprotein Complex, 78-5246
- Isobutyric Acid, α -Amino-
 - Amino Acids, 78-5259
- Leukoencephalopathy, Progressive Multifocal
 - Isolation and Characterization, Review, 78-4890
- T-Lymphocytes
 - Immune Response, Review, 78-4905
- Melanoma
 - Isolation and Characterization, Review, 78-4890
 - Virus-Like Particles, Review, 78-4891
- Meningioma
 - Virus-Like Particles, Review, 78-4891
- Mitomycin C
 - DNA Replication, 78-5253
- Nucleosomes
 - Ultrastructural Study, CV1 Cells, 78-5257
- Oxirane, 9-Anthryl-
 - DNA, Binding, 78-4984
- Oxirane, 7-Benzanthryl-
 - DNA, Binding, 78-4984
- Oxirane, 6-Chrysenyl-
 - DNA, Binding, 78-4984
- Oxirane, 9-Methyl-10-anthryl-
 - DNA, Binding, 78-4984
- Oxirane, 2-Naphthyl-
 - DNA, Binding, 78-4984
- Oxirane, 9-Phenanthryl-
 - DNA, Binding, 78-4984

Virus, SV40 (cont'd)

- Oxirane, 1-Pyrenyl-
 - DNA, Binding, 78-4984
- Proline
 - Amino Acids, 78-5259
- Radiation, Ionizing
 - DNA Replication, 78-5253
- RNA, Messenger
 - DNA-RNA Hybridization, 78-5255, 78-5260
 - Ultrastructural Study, CV1 Cells, 78-5255
- RNA, Viral
 - Degradation, 78-5260
 - DNA-RNA Hybridization, 78-5260
- Tubulin
 - Cell Transformation, Neoplastic, 78-5245
- Virus, Polyoma
 - Antigenic Determinants, Review, 78-4892
- Virus, Polyoma, BK
 - Antigenic Determinants, Review, 78-4891
 - DNA-DNA Hybridization, Review, 78-4890, 78-4893
- Virus, Polyoma, JC
 - Antigenic Determinants, Review, 78-4891
- Virus Replication
 - Nucleoprotein Complex, 78-5247
- Virus, Turkey Herpes**
 - T-Lymphocytes
 - Immunosuppression, 78-5196
 - Plant Agglutinins
 - T-Lymphocytes, 78-5196
 - Virus, Marek's Disease Herpes
 - T-Lymphocytes, 78-5196
 - Viral Vaccines, 78-5196
- Water Pollution**
 - Carcinogen, Chemical
 - Epidemiology, Review, 78-4847
 - Quantitation Method, 78-4972
 - Estrogens
 - Quantitation Method, 78-4972
 - Propionic Pentafluoro-, Anhydride
 - Carcinogenic Metabolite, 78-4972
- Wounds and Injuries**
 - Brain Neoplasms
 - Urea, Ethyl Nitroso-, 78-5186
- WY-4355**
 - see Norgestrel, Chloroethynyl-
- Xanthurenic Acid**
 - see Quinaldic Acid, 4,8-Dihydroxy-
- Xeroderma Pigmentosum**
 - Selenious Acid, Disodium Salt
 - Chromosome Aberrations, 78-5115
 - DNA Repair, 78-5115
- Xeroderma Pigmentosum**
 - Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - DNA Repair, 78-5169
 - Ultraviolet Rays, 78-5169
 - 2-Anthracenamide
 - DNA Repair, 78-5170
 - Complementation Group
 - DNA Repair, Review, 78-4936
 - Fibroblasts
 - DNA Repair, 78-5170
 - Genetics
 - Epidemiology, Review, 78-4926

erma Pigmentosum (cont'd)

quinoline, 4-Nitro-, 1-Oxide

DNA Repair, 78-5170

kin Neoplasms

Epidemiology, Review, 78-4930

Ultraviolet Rays

DNA Repair, 78-5169

DNA Repair, Review, 78-4936

Neurologic Manifestations, Review, 78-4936

irus, Herpes Simplex 1

Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-, 78-5168

Caffeine, 78-5168

Xeroderma Pigmentosum (cont'd)

Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-5168

Reactivation Kinetics, 78-5168


Ultraviolet Rays, 78-5168

Virus Replication, 78-5168

Zinc

Neurilemmoma

Urea, Methyl Nitroso-, 78-5085



Chemical Abstracts Service Registry Number Index

- 06-6, 78-5042
- 00-0, 78-4884
- 02-2, 78-5143, 78-5222
- 06-6, 78-4806, 78-4985, 78-5008
78-5010, 78-5011, 78-5016
78-5097, 78-5111, 78-5118
78-5325
- 07-7, 78-4939, 78-5050, 78-5145
78-5156, 78-5253
- 18-0, 78-4843, 78-5149, 78-5151
78-5152, 78-5153, 78-5154
78-5155, 78-5156
- 28-2, 78-5139
- 32-8, 78-4811, 78-4849, 78-4850
78-4854, 78-4855, 78-4929
78-4998, 78-4999, 78-5000
78-5001, 78-5002, 78-5003
78-5004, 78-5005, 78-5006
78-5007, 78-5008, 78-5010
78-5011, 78-5014, 78-5015
78-5016, 78-5018, 78-5020
78-5021, 78-5032, 78-5041
78-5048
- 37-3, 78-5050
- 50-0, 78-5055
- 55-5, 78-4843, 78-5141
- 76-0, 78-4947
- 81-7, 78-4912, 78-5031, 78-5052
78-5121
- 89-5, 78-5007
- 21-8, 78-5399
- 55-8, 78-5067
- 64-9, 78-4843
- 75-2, 78-5168
- 79-6, 78-4935
- 68-6, 78-4845, 78-5164
- 90-4, 78-5111, 78-5284
- 59-8, 78-5161
- 70-3, 78-4815, 78-4999
- 85-0, 78-5233
- 95-2, 78-4998
- 96-3, 78-4816, 78-4828, 78-4854
78-4982, 78-4983, 78-4985
78-4988, 78-5078, 78-5084
- 05-7, 78-5076, 78-5147
- 11-5, 78-5043
- 12-6, 78-5048, 78-5122, 78-5338
- 36-4, 78-5011
- 54-42-2, 78-5212
- 54-85-3, 78-4844
- 55-17-4, 78-4949, 78-5113, 78-5398
- 55-18-5, 78-4815, 78-4822, 78-4828
78-4951, 78-5068, 78-5072
78-5073, 78-5149
- 55-80-1, 78-4828, 78-5127
- 56-23-5, 78-4949, 78-5084
- 56-25-7, 78-5037
- 56-49-5, 78-4854, 78-4985, 78-4989
78-4990, 78-4991, 78-4992
78-4993, 78-4995, 78-4999
78-5016, 78-5033, 78-5098
78-5105, 78-5111, 78-5290
78-5291, 78-5395
- 56-53-1, 78-4863, 78-4867, 78-5117
78-5136, 78-5137, 78-5138
78-5139, 78-5363, 78-5370
- 56-55-3, 78-4855, 78-5022
- 56-57-5, 78-4948, 78-5098, 78-5100
78-5105, 78-5170
- 56-75-7, 78-4843
- 56-81-5, 78-5047
- 56-87-1, 78-5122
- 57-22-7, 78-5157
- 57-33-0, 78-4989
- 57-41-0, 78-4843, 78-5301
- 57-57-8, 78-4946, 78-4947, 78-5108
- 57-74-9, 78-4806
- 57-83-0, 78-4863, 78-5342
- 57-88-5, 78-4906, 78-5061, 78-5120
78-5349
- 57-97-6, 78-4999, 78-5021, 78-5023
78-5024, 78-5025, 78-5026
78-5027, 78-5028, 78-5029
78-5030, 78-5036
- 58-08-2, 78-5145, 78-5168, 78-5212
- 58-15-1, 78-4819, 78-5076
- 58-22-0, 78-4920, 78-5078
- 58-25-3, 78-5160
- 58-54-8, 78-5016
- 58-73-1, 78-5076
- 58-89-9, 78-5165
- 59-00-7, 78-5338
- 59-05-2, 78-5399
- 59-14-3, 78-4939, 78-5149, 78-5154
78-5212
- 59-51-8, 78-5398
- 59-89-2, 78-5054, 78-5055, 78-5056
- 60-00-4, 78-5400
- 60-15-1, 78-4843
- 60-33-3, 78-5120
- 60-54-8, 78-5076
- 60-57-1, 78-5008, 78-5161
- 60-92-4, 78-4932, 78-4937
- 61-19-8, 78-5107
- 61-57-4, 78-4845
- 62-44-2, 78-4843
- 62-49-7, 78-5113
- 62-50-0, 78-4939, 78-4947, 78-5102
78-5158
- 62-55-5, 78-4949
- 62-56-6, 78-5112
- 62-75-9, 78-4815, 78-4820, 78-4821
78-4945, 78-5067, 78-5071
78-5073, 78-5075, 78-5076
78-5077, 78-5078, 78-5152
78-5183
- 63-68-3, 78-5398
- 64-17-5, 78-5344
- 64-19-7, 78-5034, 78-5037
- 64-47-1, 78-5064
- 64-67-5, 78-5158
- 64-86-8, 78-5037
- 66-27-3, 78-4815, 78-4939, 78-4947
78-4950, 78-4987, 78-5102
78-5149, 78-5158
- 67-21-0, 78-4949, 78-5113, 78-5398
- 67-64-1, 78-5075
- 67-66-3, 78-5023
- 67-68-5, 78-5001
- 68-26-8, 78-4860, 78-5019, 78-5146
78-5349
- 70-18-8, 78-5110, 78-5112
- 70-25-7, 78-4947, 78-4948, 78-5098
78-5101, 78-5102, 78-5103
78-5104, 78-5105
- 70-26-8, 78-5122
- 71-43-2, 78-4974, 78-4975
- 71-58-9, 78-5078
- 72-33-3, 78-4864, 78-4865, 78-5132
- 74-79-3, 78-5122

75-01-4, 78-4811, 78-4830, 78-4831
 78-4832, 78-4833, 78-4834
 78-4922, 78-5110, 78-5157
 75-35-4, 78-4830, 78-4831, 78-5109
 76-38-0, 78-5160
 77-78-1, 78-5082, 78-5158
 79-01-6, 78-5160
 79-16-3, 78-5080
 79-81-2, 78-4963
 83-88-5, 78-5048
 84-17-3, 78-5138
 85-01-8, 78-4855, 78-5032
 85-32-5, 78-5107
 86-30-6, 78-5086
 90-45-9, 78-5107
 91-20-3, 78-4855
 91-22-5, 78-5099
 91-53-2, 78-4806
 91-59-8, 78-4815
 91-80-5, 78-4819
 92-66-0, 78-4978
 92-67-1, 78-4815
 92-84-2, 78-4806
 92-87-5, 78-4815
 94-59-7, 78-4816
 95-80-7, 78-4811
 96-09-3, 78-4953, 78-4984, 78-5012
 78-5013, 78-5037
 96-13-9, 78-5166
 96-45-7, 78-5112
 97-00-7, 78-5012, 78-5016, 78-5173
 97-77-8, 78-4806
 99-54-7, 78-5016
 99-56-9, 78-4809
 100-14-1, 78-5016
 100-75-4, 78-5073
 101-14-4, 78-4815
 101-80-4, 78-5126
 107-12-0, 78-5111
 107-13-1, 78-5111
 108-01-0, 78-5076
 108-39-4, 78-5032
 108-95-2, 78-5032
 109-90-0, 78-5080
 110-44-1, 78-5124

110-60-1, 78-5005
 111-03-5, 78-5120
 111-44-4, 78-4811
 112-80-1, 78-5120
 114-80-7, 78-5067
 117-39-5, 78-4806
 118-74-1, 78-4979
 120-12-7, 78-4855, 78-5032
 124-38-9, 78-5043
 124-64-1, 78-5167
 126-07-8, 78-5142
 126-72-7, 78-4809, 78-5166, 78-5167
 126-99-8, 78-4834
 127-18-4, 78-4830
 127-47-9, 78-5019
 127-69-5, 78-5076
 128-37-0, 78-4806
 128-53-0, 78-5216
 129-00-0, 78-5032
 133-06-2, 78-5162
 133-07-3, 78-5163
 141-05-9, 78-5111
 143-50-0, 78-4980
 144-90-1, 78-5259
 147-85-3, 78-5259
 148-82-3, 78-4843, 78-5302
 151-67-7, 78-5160
 153-78-6, 78-4982
 154-42-7, 78-5399
 156-57-0, 78-4806
 156-59-2, 78-4830
 191-07-1, 78-5032
 215-58-7, 78-4990, 78-5034
 286-20-4, 78-5033
 300-62-9, 78-4843
 301-04-2, 78-5116, 78-5117
 302-17-0, 78-5160
 302-79-4, 78-5037
 303-47-9, 78-4956
 305-03-3, 78-5140, 78-5153
 306-37-6, 78-5059, 78-5062, 78-5063
 309-00-2, 78-5111, 78-5161
 319-84-6, 78-4806, 78-4828, 78-4970
 78-4983, 78-5084
 319-85-7, 78-4828

324-93-6, 78-4981, 78-4982
 329-98-6, 78-5054
 343-65-7, 78-5338
 356-42-3, 78-4972
 362-74-3, 78-5093
 363-24-6, 78-5042
 365-07-1, 78-5108
 372-75-8, 78-5122
 398-32-3, 78-4981
 439-14-5, 78-5160
 443-48-1, 78-4844
 446-86-6, 78-5157
 452-06-2, 78-4947
 471-29-4, 78-5103
 480-54-6, 78-4949
 484-78-6, 78-5338
 492-27-3, 78-5338
 519-23-3, 78-5011
 521-18-6, 78-4920
 521-74-4, 78-5099
 536-33-4, 78-5112
 540-59-0, 78-4830
 540-73-8, 78-5060, 78-5061, 78-5064
 543-38-4, 78-5202
 545-55-1, 78-5157
 548-93-6, 78-5044, 78-5338
 551-11-1, 78-5042
 557-66-4, 78-5066
 578-76-7, 78-5083
 589-41-3, 78-5082
 590-96-5, 78-4820
 592-62-1, 78-5064
 593-51-1, 78-5066
 602-52-8, 78-5114
 604-59-1, 78-4806, 78-5011
 607-35-2, 78-5099
 607-57-8, 78-4946
 612-62-4, 78-5099
 613-13-8, 78-5170
 621-64-7, 78-4824
 621-90-9, 78-4816
 622-78-6, 78-4806
 630-08-0, 78-5043
 645-05-6, 78-5399

1-16-9, 78-5148, 78-5149
 4-81-7, 78-4946
 0-31-9, 78-5150
 4-93-5, 78-4815, 78-4826, 78-5084
 78-5085, 78-5086, 78-5087
 78-5090, 78-5281
 5-65-3, 78-5042
 9-73-9, 78-4827, 78-5088, 78-5089
 78-5091, 78-5092, 78-5093
 78-5186, 78-5319
 5-21-4, 78-5157
 9-01-2, 78-5094, 78-5095
 12-17-1, 78-4996
 12-49-9, 78-5399
 18-35-0, 78-5011
 14-16-3, 78-4815, 78-4825
 16-06-7, 78-5158
 10-55-2, 78-5073
 11-07-9, 78-5106
 116-54-7, 78-4823, 78-5070
 120-71-4, 78-4815
 162-65-8, 78-4811, 78-4815, 78-4816
 78-4947, 78-4956, 78-4957
 78-4960, 78-4962, 78-4963
 78-4966, 78-4967, 78-4968
 78-4969, 78-4970
 139-45-8, 78-5114
 109-37-1, 78-5326
 109-63-3, 78-5326
 114-20-1, 78-4922
 117-79-9, 78-5326
 119-95-5, 78-5120
 132-21-4, 78-4805, 78-4841, 78-4842
 78-4909, 78-4971, 78-5284
 78-5326, 78-5385
 185-95-1, 78-4838, 78-4959, 78-4965
 78-4968
 182-09-8, 78-4824
 146-01-6, 78-4852
 151-56-0, 78-4845
 113-57-7, 78-4978
 115-28-3, 78-5132
 508-86-3, 78-5098, 78-5105
 541-69-7, 78-5031
 1083-23-6, 78-5033
 105-97-3, 78-4845
 124-93-4, 78-5132
 137-73-3, 78-5132

3308-64-3, 78-4995
 3342-98-1, 78-4995
 3343-07-5, 78-4995
 3343-08-6, 78-4995
 3567-69-9, 78-5128, 78-5129
 3732-82-9, 78-5150
 3817-11-6, 78-4825
 4637-56-3, 78-5100
 5255-75-4, 78-5016
 5307-14-2, 78-4809
 5385-75-1, 78-5033
 6051-87-2, 78-4806, 78-5010
 6098-44-8, 78-4984, 78-4986, 78-4987
 78-5168, 78-5169
 6292-55-3, 78-4985
 6613-85-0, 78-5160
 6795-23-9, 78-4837, 78-4838, 78-4964
 78-4969
 7220-81-7, 78-4959, 78-4965
 7241-98-7, 78-4959, 78-4965
 7439-89-6, 78-4934, 78-5085
 7439-92-1, 78-4934
 7440-02-0, 78-4934
 7440-17-7, 78-5085
 7440-22-4, 78-5326
 7440-38-2, 78-4843
 7440-41-7, 78-4934
 7440-43-9, 78-4934, 78-5118
 7440-44-0, 78-4971
 7440-47-3, 78-4934, 78-5085
 7440-61-1, 78-5354
 7440-65-5, 78-5085
 7440-66-6, 78-5085
 7553-56-2, 78-5004
 7632-00-0, 78-5076, 78-5080, 78-5124
 7647-01-0, 78-5053
 7664-38-2, 78-5042
 7697-37-2, 78-4818, 78-5052
 7705-08-0, 78-5004
 7720-78-7, 78-5031
 7727-37-9, 78-5052
 7782-49-2, 78-4806, 78-5085
 7782-77-6, 78-4817, 78-4818, 78-4819
 78-5051, 78-5052, 78-5070
 78-5077, 78-5103, 78-5171
 78-5330

7790-28-5, 78-5284
 7803-49-8, 78-4989
 8001-30-7, 78-5120
 8008-60-4, 78-5049
 9001-05-2, 78-5014
 9001-45-0, 78-5023
 9001-48-3, 78-5014
 9001-60-9, 78-5111
 9001-63-2, 78-5400
 9001-77-8, 78-4974, 78-5389
 9001-78-9, 78-5165
 9001-99-4, 78-5072
 9002-04-4, 78-5157
 9002-12-4, 78-5014
 9002-62-4, 78-5134
 9002-72-6, 78-5392
 9004-10-8, 78-4906, 78-5001, 78-5263
 78-5392
 9007-12-9, 78-5263
 9012-42-4, 78-4932
 9035-50-1, 78-4852, 78-4989
 9035-58-9, 78-5304
 10028-17-8, 78-4982
 10045-97-3, 78-5181
 10048-13-2, 78-4815, 78-4956
 10102-18-8, 78-5115
 10198-40-0, 78-5253
 10595-95-6, 78-5080
 11028-71-0, 78-5300
 11085-36-2, 78-5392
 11096-82-5, 78-4828
 11097-69-1, 78-4828, 78-4938, 78-4995
 78-5075, 78-5097, 78-5111
 12001-28-4, 78-4805, 78-4841, 78-4842
 78-4909, 78-4971, 78-5284
 78-5326, 78-5385
 12001-29-5, 78-4805, 78-4841, 78-4842
 78-4909, 78-4971, 78-5284
 78-5326, 78-5385
 12035-72-2, 78-5119
 12172-73-5, 78-4326, 78-4805, 78-4841
 78-4842, 78-4909, 78-4971
 78-5284, 78-5385
 13147-25-6, 78-4822
 13256-22-9, 78-5057
 13256-32-1, 78-5096
 13410-01-0, 78-5115

13551-87-6, 78-5144	17181-37-2, 78-4909	33580-12-0, 78-5399
13823-27-3, 78-5185	17676-08-3, 78-5050	33868-17-6, 78-5103
13982-63-3, 78-4868	20244-86-4, 78-5107	36504-66-2, 78-4855
14046-83-4, 78-5023	20535-83-5, 78-4827, 78-5106	37114-16-2, 78-5057
14119-33-6, 78-5184	21738-42-1, 78-4845	37317-41-2, 78-4828
14797-73-0, 78-5077	22225-32-7, 78-4985	37574-47-3, 78-5012, 78-5013
14807-96-6, 78-5006	23214-92-8, 78-5399	38252-74-3, 78-4825
14808-60-7, 78-5326	24554-26-5, 78-4976	39386-07-7, 78-5054, 78-5195, 78-
14901-08-7, 78-4816	24909-09-9, 78-5017	39603-43-5, 78-5399
14932-53-7, 78-5042	25013-16-5, 78-4806	39603-53-7, 78-4824
15117-48-3, 78-4868, 78-5184	25126-76-5, 78-4920	56484-47-0, 78-5017
15232-63-0, 78-5050	25843-45-2, 78-5159	57303-99-8, 78-4855, 78-5020
16110-14-8, 78-5031	26443-03-8, 78-5055	59299-51-3, 78-4979
16561-29-8, 78-5022, 78-5036, 78-5037 78-5038, 78-5039, 78-5040 78-5041, 78-5042	26675-46-7, 78-5160	59963-01-8, 78-4855, 78-4984, 78- 78-5009, 78-5010
16812-54-7, 78-5119	28622-84-6, 78-5017	62641-69-4, 78-5031
16941-32-5, 78-5263	29411-74-3, 78-5399	62641-70-7, 78-5031
17068-78-9, 78-4805, 78-4841, 78-4842 78-4909, 78-4971, 78-5284 78-5326, 78-5385	30310-80-6, 78-5057	
	31005-02-4, 78-4854	
	32752-29-7, 78-5038	

Wiswesser Line Notation Index

.G, 78-5053 78-5302
 3..P-O4, 78-5042
 3, 78-5326
 3, 78-4843
 3, 78-4934
 3, 78-4971
 3, 78-4934, 78-5118
 3, 78-5253
 3, 78-4934, 78-5085
 3, 78-5181
 3, 78-4934, 78-5085
 3..G3, 78-5004
 3..S-O4, 78-5031
 3..O3, 78-5326
 3, 78-5004
 3, 78-5052
 3..I-O3-Q, 78-5284
 3..N-O-Q, 78-5076, 78-5080, 78-5124
 3..SE-O-Q2, 78-5115
 3..SE-O4, 78-5115
 3, 78-4934
 3..S2, 78-5119
 3, 78-4934
 3, 78-4868, 78-5184
 3..N-O4, 78-5185
 3, 78-4868
 3, 78-5042, 78-5085
 3, 78-4806, 78-5085
 3..O2, 78-4805, 78-5326
 3..O4, 78-4909
 3..I.O2, 78-4922
 3..O2, 78-5326
 3, 78-5354
 3, 78-5085
 3, 78-5043
 3..BR, 78-4978
 3..CR, 78-4978
 3..DR, 78-4978
 3..DR CQ DQ, 78-4978
 3..DR DQ, 78-4978
 3..YE1O 3PO, 78-5166, 78-5167
 3..DR, 78-4981

FXFFR CNW ENW DN3&3, 78-4824
 G 6-R, 78-4979
 GXGGG, 78-4949, 78-5084
 GXGGYQPO&O1&O1, 78-4845, 78-5164
 GYEXFFF, 78-5160
 GYGG, 78-5023
 GYGUYGG, 78-4830
 GYGU1, 78-4830, 78-4831, 78-5109
 GYGU1G, 78-5160
 G1U1, 78-4811, 78-4830, 78-4831, 78-4832, 78-4833, 78-4834
 78-4922, 78-5110, 78-5157
 G1U1G, 78-4830
 G2N1&2G, 78-5168
 G2O2G, 78-4811
 L B656 HHJ DMVI, 78-4985
 L B656 HHJ DNQV1, 78-4985
 L B656 HHJ EMV1, 78-4816, 78-4828, 78-4854, 78-4982
 78-4983, 78-4985, 78-4988, 78-5078, 78-5084
 L B656 HHJ ENW, 78-4946
 L B666J, 78-4855, 78-5032
 L B666J HHJ EZ, 78-4982
 L B677 MV&T&J CO1 DO1 EO1 JMV1 NO1, 78-5037
 L C555 A IUTJ AG AG BG DG EG HG IG JG, 78-4806
 L C666J, 78-4855, 78-5032
 L C666J EZ, 78-5170
 L D5 C555 A D- EU JUTJ AG AG BG IG JG KG, 78-5111
 78-5161
 L D6 B666&&&TJ OQ PQ, 78-5022
 L D6 B666J, 78-4855, 78-5022
 L D6 B666J C J, 78-4999, 78-5021, 78-5023, 78-5024
 78-5025, 78-5026, 78-5027, 78-5028, 78-5029
 78-5030, 78-5036
 L D6 B666J J, 78-5031
 L D6 B6666 2AB TJ, 78-4811, 78-4849, 78-4850, 78-4854
 78-4855, 78-4929, 78-4998, 78-4999, 78-5000
 78-5001, 78-5002, 78-5003, 78-5004, 78-5005
 78-5006, 78-5007, 78-5008, 78-5010, 78-5011
 78-5014, 78-5015, 78-5016, 78-5018, 78-5020
 78-5021, 78-5032, 78-5041, 78-5048
 L D6 B6666 2AB TJ GQ HQ, 78-4855, 78-5020
 L D6 J6 C666J, 78-4990, 78-5034
 L E5 B666 LUTJ A E FY&3Y QQ -B&AEFO, 78-4906
 78-5061, 78-5120, 78-5349
 L E5 B666 OV MUTJ A E FQ -B&AEF, 78-4920, 78-5078
 L E5 B666 OV MUTJ A1 COV1 E1 FV1 L -A&L -
 B&ACEF
 78-5078

L E5 B666 OV MUTJ A1 E1 FV1 -B&AEF, 78-4863
 78-5342
L E5 B666 OVJ A1 E1 FQ -B&AEF, 78-4920
L E5 B666TJ A1Q CQ E IQ MQ QQ F- DT5OV EHJ&
OO- BT6OTJ CQ DQ EQ F
 78-5042
L E5 B666TTT&J E FQ OQ, 78-5139
L E6 C666 BV FV CU GUTTT&J DQ EQ GVZ HQ IN1&1
MQ M RQ
 78-5076
L E6 D6656 1A T OV PHJ R, 78-4995
L E6 D6656 1A T PV OHJ R, 78-4995
L E6 D6656 1A T&&&T&J JQ KQ OQ R1, 78-4995
L E6 D6656 1A T&&&T&J OQ R, 78-4995
L E6 D6656 1A T&&&T&J PQ R, 78-4995
L E6 D6656 1A T&&&T&J R, 78-4854, 78-4985, 78-4989
 78-4990, 78-4991, 78-4992, 78-4993, 78-4995
 78-4999, 78-5016, 78-5033, 78-5098, 78-5105
 78-5111, 78-5290, 78-5291, 78-5395
L G6 D6 B666J, 78-4815, 78-4999
L545 B4 C5 D 4ABCE J DVTJ-/G 1 O, 78-4980
L6TJ AG BG CG DG EG FG *ALPHA, 78-4806, 78-4828
 78-4970, 78-4983, 78-5084
L6TJ AG BG CG DG EG FG *BETA, 78-4828
L6TJ AG BG CG DG EG FG *GAMMA, 78-5165
L6UTJ A A BIUIY&UZUIYUIVQ C -T, 78-5037
L6UTJ A BL/UIY&U2/ 2Q C C -T, 78-4860, 78-5019
 78-5146, 78-5349
L64TJ A BIUIY&U2UIY&U2OV1 C C, 78-5019
L66J, 78-4855
L66J BSWQ ENUN- CL66J BQ ESWQ &-NA- 2, 78-5128
 78-5129
L66J CZ, 78-4815
L666 B6 2AB PJ, 78-5032
NCN1&NO, 78-5103
NC1U1, 78-5111
OCN2, 78-5080
OCO, 78-5043
ONNR&R, 78-5086
ONN1&VM1, 78-5096
ONN1&1, 78-4815, 78-4820, 78-4821, 78-4945, 78-5067
 78-5071, 78-5073, 78-5075, 78-5076, 78-5077
 78-5078, 78-5152, 78-5183
ONN2&2, 78-4815, 78-4822, 78-4828, 78-4951, 78-5068
 78-5072, 78-5073, 78-5149
ONN3&3, 78-4824
ONN4&4, 78-4815, 78-4825
ON1&U&1, 78-5159
ON1&UN1OV1, 78-5064

OS1&1, 78-5001
OV1 & 2-PB-, 78-5116, 78-5117
QMVO2, 78-5082
QR, 78-5032
QR C, 78-5032
QR DY2& 2U, 78-4863, 78-4867, 78-5117, 78-5136, 78-5138, 78-5139, 78-5363, 78-5370
QVYQYQVQ & 2 &621 T6NJ C1 2/ XV/ &622, 78-5016
QVYZ1R DN2G2G -L, 78-4843
QVYZ2S1, 78-5398
QVYZ2S2 -DL, 78-4949, 78-5113, 78-5398
QV1, 78-5034, 78-5037
QVIN1&NO, 78-5057
QVIN1VQ 22, 78-5400
QV1OR BG CG DVYZ&U1, 78-5016
QV1U2U2, 78-5124
QV3R DN2G2G, 78-5140, 78-5153
QV8U9 -C, 78-5120
QYQXGGG, 78-5160
Q1 4P &G, 78-5167
Q1NUNO&1, 78-4820
Q1YE1E, 78-5166
Q1YQ1Q, 78-5047
Q1YQ1Q / YV17, 78-5120
Q2, 78-5344
Q2K &Q, 78-5113
Q2N1&1, 78-5076
Q2N2&NO, 78-4822
Q3N3&NO, 78-4824
Q4N4&NO, 78-4825
R, 78-4974, 78-4975
T B666 HKJ EX H2 IR& LZ &G &9/26, 78-5114
T B666 HKJ EZ H2 IR& LZ &E &9/26, 78-5114
T C555 A AO DVOVTJ C G, 78-5037
T C6 B5665 2AB S BX IN QN NU JH&&TTTJ FO1 IV
KVO1 KQ LOV1 M2 E- NT F6 E596 A B
LM&&TTJ NVO1 RQ R2, 78-5157
T C6 B5665 2AB S BX IN QN NU JH&&TTTTJ FO1 I
KVO1 KQ LOV1 M2 E- NT F6 E596 A B
LM&&TTJ NVO1 Q Q2-, 78-5157
T C6656 1A P GN LM CUTT&&J EVN2&2 G -B, 78-5056
T C666 BM ISJ, 78-4806
T C666 BN DNVMV INJ B1YQYQYQ1Q L M -BBB
 78-5048
T C666 BO EV INJ D FZ N G- K-/VM- OT5-16- AN FV
IVN LVO PVM SVTJ G J KY N RY 2
 78-4947


D3 B556 BN EM JV MVTJTJ GO1 HIOVZ KZ L
 78-4939, 78-5050, 78-5145, 78-5156, 78-5253
 D36 I666 B6 2AB U EOTJ&&&J, 78-5012, 78-5013
 E3 D5 C555 A D- FO KUTJ AG AG BG JG KG LG
 78-5008, 78-5161
 E3 D6 B6666 2AB U FOTT&&&J HQ IQ, 78-4855
 78-4984, 78-4999, 78-5009, 78-5010
 E6 C6 B655 DO KV PO RO SU&&&TTJ IQ MO1
 78-4815, 78-4956
 F5 C6 B655 DOV GV OO QO RUT&&TTJ LO1, 78-4811
 78-4815, 78-4816, 78-4947, 78-4956, 78-4957
 78-4960, 78-4962, 78-4963, 78-4966, 78-4967
 78-4968, 78-4969, 78-4970
 F5 C6 B655 DOV GV OO QOT&&TTJ LO1, 78-4959
 78-4965
 F6 C6 B655 DOV GVO PO ROT&&TTJ MO1, 78-4959
 78-4965
 F6 D5 C666 EM ON&&TTTJ HO1 SOVR CO1 DO1
 EO1& TO1 UVO1
 78-4843, 78-5141
 NTJ A- 3PO, 78-5157
 OTJ BR, 78-4953, 78-4984, 78-5012, 78-5013, 78-5037
 OTJ BXGGG, 78-5033
 OVTJ, 78-4946, 78-4947, 78-5108
 MVMV EHV ER& ER, 78-4843, 78-5301
 MYMTJ BUS, 78-5112
 N CNJ A2Q B1 ENW, 78-4844
 N CSJ DNW B- CT5MVTJ, 78-4845
 NNVJ A BR& DN1&1 E, 78-4819, 78-5076
 NTJ ANO, 78-5073
 NTJ ANO CQ, 78-5058
 OJ BNW E- ET5N CSJ BMVH, 78-4976
 OSWTJ, 78-4815
 OV EHV CQ DQ EYQ1Q, 78-4912, 78-5031, 78-5052
 78-5121
 VNVJ B2, 78-5216
 5-12- 1A Q EN IOVY OVO BUTJ KU2 M NQ N1Q
 78-4949
 5 A ANTJ A GOVYR&1Q, 78-5067
 5 BMJ DIYZVQ -L, 78-5048, 78-5122, 78-5338
 5 BN CM FYM INJ FUS HZ, 78-5399
 5 BN DN FNVNVJ B F H, 78-5145, 78-5168, 78-5212
 5 BO DO CHJ G2U1, 78-4816
 5 BOXVJ FO1 HO1 IG C-& DL6V DX BUTJ CO1 E
 78-5142
 5 BVNV GUTJ CSXGGG, 78-5162
 5 BVNVJ CSXGGG, 78-5163
 MPOTJ BO BN2G2G, 78-4843, 78-5149, 78-5151, 78-5152
 78-5153, 78-5154, 78-5155, 78-5156

T6MV DVN CHJ CY3 C2 FO &-NA-, 78-4989
 T6MVMVJ EF, 78-5399
 T6N CN ENJ-/N1&1 3, 78-5399
 T6N DOTJ ANO, 78-5054, 78-5055, 78-5056
 T6NJ BN2N1&1&1- BT5SJ, 78-4819
 T6NJ B2 DYZUS, 78-5112
 T6NJ C- BT5NTJ A, 78-5043
 T6NJ DVMZ, 78-4844
 T6NTJ ANO, 78-5073
 T6NVMVJ EE A- ET5OTJ B1Q CQ -A&C, 78-4939
 78-5149, 78-5154, 78-5212
 T6NVMVJ EI A- ET5OTJ B1Q CQ -A&C, 78-5212
 T6OTJ BO1NUNO&1 CQ DQ EQ F1Q-A&D -D, 78-4816
 T6MVMVJ FHJ F2 FR, 78-4806, 78-4985, 78-5008, 78-5010
 78-5011, 78-5016, 78-5097, 78-5111, 78-5118
 78-5325
 T66 BM CHJ C C E HO2, 78-4806
 T66 BM DN FN HNJ IS- ET5N ONJ DNW, 78-5157
 T66 BN DN GN JNJ CZ EZ H1N1&R DVMYVQ2VQ *L
 L DX
 78-5399
 T66 BNJ, 78-5099
 T66 BNJ BO EMQ, 78-5100
 T66 BNJ BO ENW, 78-4948, 78-5098, 78-5100, 78-5105
 78-5170
 T66 BNJ BO EZ, 78-5098, 78-5105
 T66 BNJ CG, 78-5099
 T66 BNJ CVQ EQ, 78-5338
 T66 BNJ CVQ EQ JQ, 78-5338
 T66 BNJ EMY&3N2&2 IG, 78-5076, 78-5147
 T66 BNJ JNW, 78-5099
 T66 BNJ J1, 78-5099
 T66 BN1 GE IE J1, 78-5099
 T66 BO EVJ CR CQ DQ& DQ GQ IQ, 78-4806
 T66 BVOT&J D GG IVMYVQ1R& JQ, 78-4956
 T67 GN JN IHJ CG HM1 JO KR, 78-5160
 T67 GNV JN IHJ CG G KR, 78-5160
 VHH, 78-4884
 WNMVUM&N1&NO, 78-4947, 78-4948, 78-5098, 78-5101
 78-5102, 78-5103, 78-5104, 78-5105
 WNR BG ENW, 78-5012, 78-5016, 78-5173
 WNR CG DG, 78-5016
 WNR DYQY1QM VYGG -DL, 78-4843
 WNR D1G, 78-5016
 WS1&OY, 78-5158
 WS1&O1, 78-4815, 78-4939, 78-4947, 78-4950, 78-4987
 78-5102, 78-5149, 78-5158


WS1&O2, 78-4939, 78-4947, 78-5102, 78-5158
ZQ, 78-4989
ZR BNW DZ, 78-4809
ZR BQ FVQ, 78-5044, 78-5338
ZR BQ FV1YZVQ, 78-5338
ZR BZ DNW, 78-4809
ZR CZ D, 78-4811
ZR DOR DZ, 78-5126
ZR DR, 78-4815
ZR DR C1, 78-4982
ZR DR DF, 78-4981, 78-4982
ZR DR DZ, 78-4815
ZVN1&NO, 78-4815, 78-4826, 78-5084, 78-5085, 78-5086
78-5087, 78-5090, 78-5281
ZVN2&NO, 78-4827, 78-5088, 78-5089, 78-5091, 78-5092
78-5093, 78-5186, 78-5319
ZVN4&NO, 78-5094, 78-5095
ZVO2, 78-4935
ZV4YZVQ, 78-5122
ZYUS, 78-4949
ZYZUS, 78-5112
ZY1R, 78-4843
Z1 &GH, 78-5066
Z2 &GH, 78-5066

Z2SH &GH, 78-4806
Z3Z, 78-5005
1MM1, 78-5060, 78-5061, 78-5064
1MM1 &GH &GH, 78-5059, 78-5062, 78-5063
1MR DNUNR, 78-4816
1N1& 3PO1N1& 3PO, 78-5150
1N1&R DNUNR C, 78-4828, 78-5127
1N1&2OYR&R, 78-5076
1OSWO1, 78-5082, 78-5158
1UYG1U1, 78-4834
1VN1, 78-5080
1V1, 78-5075
1VINNO&1V1, 78-5081
1X&&R BQ E CX, 78-4806
1Y&MVR DIMM1, 78-5148, 78-5149
2N2&YUS&S 2, 78-4806
2OR DMV1, 78-4843
2OSWO2, 78-5158
2OVMMVO2, 78-5111
2OV1U1VO2, 78-5111
9AA, 78-5107

مجلسه اول



1875



Handwritten text, likely bleed-through from the reverse side of the page.



405
670

8101


CARCINOGENESIS ABSTRACTS

JUN 1 1979

VOLUME 16,
ISSUE 10



THE FRANKLIN INSTITUTE PRESS™



CARCINOGENESIS ABSTRACTS

A monthly publication sponsored by the
NATIONAL CANCER INSTITUTE

EDITOR

GEORGE P. STUDZINSKI, M.D., Ph.D.
COLLEGE OF MEDICINE AND DENTISTRY
OF NEW JERSEY, NEWARK

ASSOCIATE EDITOR

JUSSI J. SAUKKONEN, M.D.
JEFFERSON MEDICAL COLLEGE
PHILADELPHIA

NCI STAFF CONSULTANTS

ELIZABETH WEISBURGER, Ph.D.
JOAN W. CHASE, M.S.

Literature Selected, Abstracted, and Indexed
by

Information Services Department, Science Information Services Organization

BRUCE H. KLEINSTEIN, Ph.D., J.D., Assistant Director, Biomedical Projects

RUTHANN E. AUCHINLECK, Managing Editor

The Franklin Research Center
The Benjamin Franklin Parkway, Philadelphia, Pennsylvania 19103

All rights reserved. No part of this journal may be reproduced in any form, for any purpose or by any means, abstracted, or entered into any data base, electronic or otherwise, without specific permission in writing from the publisher.

Copyright © 1979 The Franklin Institute Press

CARCINOGENESIS ABSTRACTS

VOLUME 16, ISSUE 10

CONTENTS


	Cross Reference Abbreviations	Article Numbers	Page
VIEW	(Rev)	78-5401-78-5463	2029
CHEMICAL CARCINOGENESIS	(Chem)	78-5464-78-5689	2040
PHYSICAL CARCINOGENESIS	(Phys)	78-5690-78-5735	2092
VIRAL CARCINOGENESIS	(Viral)	78-5736-78-5853	2100
IMMUNOLOGY	(Immun)	78-5854-78-5868	2132
PATHOGENESIS	(Path)	78-5869-78-5914	2137
EPIDEMIOLOGY AND BIOMETRY	(Epid-Biom)	78-5915-78-5977	2147
MISCELLANEOUS	(Misc)	78-5978-78-6000	2160
AUTHOR INDEX			2167
SUBJECT INDEX			2177
CHEMICAL ABSTRACTS SERVICE REGISTRY NUMBER INDEX			2265
SWESSER LINE NOTATION INDEX			2269

GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations or citations of significant carcinogenesis articles collected from the current major biomedical sources of world literature. This service is provided by the National Cancer Institute through a contract with the Franklin Research Center for preparation of the publication, under Contract No. NOI-CP-75885 with the National Cancer Institute, U.S. Department of Health, Education and Welfare. Published and distributed by the Franklin Institute PressSM.

SUBSCRIPTIONS

Subscription sale by The Franklin Institute Press, Box 2266, Philadelphia, PA 19103. Subscription rate for Volume 16, consisting of 12 monthly issues and a cumulative index is \$75.00, Domestic and \$90.00, Foreign. Check should be made payable to The Franklin Institute Press.
Nihonbashi 2-Chome, Tokyo 103, Japan, is the exclusive distributor of CARCINOGENESIS ABSTRACTS in Japan.



GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations, or citations of significant cancer articles collected from the current major biomedical sources of the world literature. This service is provided by the National Cancer Institute through a contract with Franklin Research Center for preparation of the publication.

FREQUENCY OF PUBLICATION: *CARCINOGENESIS ABSTRACTS*, Volume 16, is published in 12 monthly issues with indexes, plus a cumulative index.


ABBREVIATIONS

JOURNAL names are abbreviated according to the *List of Journals Indexed in Index Medicus, Abbreviation*. If the journal is not listed in this, abbreviations are derived from the *International List of Periodical Title and Abbreviations*.

LANGUAGE of the article is indicated in parentheses after the title and is represented by a three-letter code. The source for these codes is *MARC Manuals Used by the Library of Congress*, pages 183-187.

ABBREVIATIONS used in abstracts:

angstrom(s)	mOsm	milliosmolar
adrenocorticotrophic hormone	max	maximum
adenosine diphosphate	mEq	milliequivalent(s)
adenosine monophosphate	min	minute(s)
adenosine triphosphate	ml	milliliter(s)
approximately	μl	microliter(s)
average	mm	millimeter(s)
bacillus Calmette-Guerin	mo	month(s)
twice daily	mol wt	molecular weight
degree(s) centigrade	N	normal concentration
calorie(s)	NAD	nicotinamide adenine dinucleotide
kilocalorie(s)	NADH	reduced nicotinamide adenine dinucleotide
cubic centimeter(s)	NADP	nicotinamide adenine dinucleotidephosphate
curie(s)	NADPH	reduced nicotinamide adenine dinucleotidephosphate
millicurie(s)	NCI	National Cancer Institute
microcurie(s)	NIH	National Institutes of Health
centimeter(s)	PAS	periodic acid-Schiff
central nervous system	po	orally
counts per minute	ppb	parts per billion
deoxyribonucleic acid	ppm	parts per million
median effective dose	qid	four times daily
ethylenediamine tetraacetic acid	qod	every other day
gram(s)	QO ₂	oxygen quotient
kilogram(s)	R	roentgen
milligram(s)	RBC	red blood cells (erythrocytes)
microgram(s)	RNA	ribonucleic acid
hemoglobin	rpm	revolutions per minute
hour(s)	sc	subcutaneous
intra-arterial	sec	second(s)
intradermal	SGOT	serum glutamic-oxaloacetic transaminase
Immunoglobulin A	SGPT	serum glutamic-pyruvic transaminase
Immunoglobulin B	soln	solution
Immunoglobulin G	TCD	tissue culture dose
Immunoglobulin M	TCD ₅₀	median tissue culture dose
increased life span	tid	three times daily
intramuscular	UV	ultraviolet
intraperitoneal	WBC	white blood cells (leukocytes)
International Unit(s)	wk	week(s)
intravenous	wt	weight
Michaelis constant	X	times
lethal dose	yr	year(s)
median lethal dose		
molar		
micromolar		



REVIEW

5401 **Environmental Pathology, Target Environmental Cancer.** (Eng) Garry, V. F. (Dept. Lab. Medicine and Pathology, Univ. Minnesota, Minneapolis, MN). *Minn Med* 61(5): 328-330; 340; 1978.

Statistics show an almost logarithmic increase in overall cancer incidence per 100,000 persons over 40 yr of age. Problems screening for environmental carcinogens include the latent period between exposure and onset of disease, as well as species differences among experimental animals and man. Current screening tools are discussed. Occupational bladder cancer and other cancer clusters in Minnesota are reviewed. (23 refs.)

5402 **Environmental Carcinogenesis.** (Eng) Wynder, E. L. (Dept. Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY). *Clin Bull* 8(1): 3-9; 1978.

The role of tobacco, alcohol, and diet in cancer induction is discussed, and measures to prevent environmental carcinogenesis are outlined. (46 refs.)

5403 **Harmful Effect of Water Chlorination with Special Regard to the Cocarcinogenicity of Chlorine.** Part I: Literature Survey. (Ger) Pfeiffer, E. H. (Hygiene-Institut, Johannes-Gutenberg-Universität, D-6500 Mainz, W. Germany). *Zentralbl Bakteriologie (Orig B)* 166(2/3): 99-184; 1978.

Studies of the toxicological aspects of chlorine in water and certain foods are reviewed. In one experiment, seven generations of BD II rats received tap water containing 100 ppm chlorine ad libitum over a 9-yr period. Tumor incidence was the same (approx 10%) as that in controls, which received nonchlorinated water. (no refs.)

5404 **Nutritional and Environmental Factors in Carcinogenesis (Meeting Abstract).** (Eng) Wynder, E. L. (No affiliation given). In: *Arnold O. Beckman Conference on Biochemistry of Cancer held in San Antonio, September 8, 1978*. American Association for Clinical Chemistry (San Antonio, TX): p. 9; 1978. (3 refs.)

5405 **The Environment and Prenatal Origins of Human Disease.** (Eng) Miller, R. W. (Clinical

Epidemiology Branch, NCI, Bethesda, MD, 20014). *Excerpta Med Int Congr Ser* (432): 297-302; 1978.

Patterns of discovery of environmental effects on the embryo, monitoring for new human teratogens, teratogenic pollutants (ie, polychlorinated biphenyls, methylmercury), and the interaction of genetics and the environment are surveyed briefly. (23 refs)

78-5406 **Environmental Carcinogens: Mechanisms of Action and Occurrence.** (Ger) Preussmann, R. (Institut für Toxikologie und Chemotherapie, Deutsches Krebsforschungszentrum, Im Neuenheimer Feld 280, D-6900 Heidelberg 1, W. Germany). *Zentralbl Bakteriologie (Orig B)* 166(2/3): 144-158; 1978.

The mechanism of action of known environmental carcinogens is described. These compounds are not carcinogenic per se, but are metabolically bioactivated to electrophilic reactants by chemically reactive intermediates. The electrophilic reactants then form covalent bonds with nucleic acids. This change in the genetic code is in agreement with the mutation hypothesis of carcinogenesis. The bioactivation of aflatoxin B₁ (AFB), N-nitroso compounds, and benzo(a)pyrene (BP) is presented to illustrate this mechanism. Threshold levels (no-effect levels) determined by animal experiments are compared with data on human exposure to BP, AFB, N-nitroso compounds, and vinyl chloride. The difficulties involved in risk evaluations are discussed. (39 refs)

78-5407 **Genetic Variability and Extrapolation from Animals to Man: Some Perspectives on Susceptibility to Chemical Carcinogenesis from Aromatic Amines.** (Eng) Weber, W. W. (Dept. Pharmacology, Univ. Michigan Medical Sch., Ann Arbor, MI, 48109). *Environ Health Perspect* 22: 141-144; 1978.

The predictive value for man of data obtained in other species is illustrated using carcinogenesis arising from exposure to aromatic amines as an example. Studies with an experimental mouse genetics model indicated that the mutagenic effect of acetylaminofluorene (AAF) is related to genetically mediated increases in microsomal oxidative enzyme activity associated with cytochrome P-448. Genetic studies have shown that persons could be classified into two major genetic subgroups, rapid and slow isoniazid acetylators. A genetic polymorphism also exists in rabbits, and the rate of elimination of isoniazid by acetylation is highly correlated in both species

with the amount of isoniazid N-acetyltransferase in the liver. The rate of acetylation of amines with liver homogenates prepared from rapid and slow acetylators rabbits was determined. The results suggest that individuals that are genetically rapid acetylators of isoniazid might also be expected to transform aminofluorene and other hazardous amines to activated carcinogenic forms more rapidly than slow acetylators. The role of this genetic polymorphism as a human determinant of susceptibility to chemical carcinogenesis from aromatic amines is being assessed. (16 refs)

- 78-5408 Estimating Cancer Risks to a Population.** (Eng) Schneiderman, M. A. (Field Studies and Statistics Program, DCCP, NCI, Bethesda, MD, 20014); Brown, C. C. *Environ Health Perspect* 22: 115-124; 1978.

General principles guiding current laboratory-to-humans extrapolations of cancer risk are as follows: (1) effects in animals, properly qualified, are applicable to humans; (2) methods to establish a threshold for long-delayed effects of toxic agents such as cancer do not exist; (3) the exposure of experimental animals to toxic agents in high doses is a necessary and valid method of discovering possible carcinogenic hazards in humans; and (4) material should be assessed in terms of human risk, rather than as "safe" or "unsafe". A number of mathematical models have been developed to extrapolate from animal data to human risk of carcinogenesis and to estimate carcinogenic threshold doses. Statistical analysis of standard animal carcinogenicity experiments does not, however, resolve the threshold question, because there are too many biologically reasonable mathematical models, both implying and denying the existence of thresholds, that will fit the observed results. The search for thresholds should not be a search for one specific no-effect dose applicable to all members of the population at risk, but rather a search for many thresholds for each of the population subgroups at various degrees of risk. (20 refs.)

- 78-5409 Identification and Reduction of Carcinogens in the Respiratory Environment.** (Eng) Hoffmann, D. (Naylor Dana Inst. Disease Prevention, American Health Foundation, Valhalla, NY, 10595); Wynder, E. L. *Zentralbl Bakteriol [Orig B]* 166(2/3): 113-135; 1978.

The three major factors responsible for the overall increase in lung cancer (tobacco, air pollution, and industrial chemical exposure) are reviewed. Concerning industrial carcinogenesis, particular attention is focused on bis(chloromethyl)ether, vinyl chloride, benzene, coke oven effluents, asbestos, and nitrosamines. Sources of atmospheric carcinogens and their reduction are outlined. The tumorigenic agents in cigarette smoke are being studied so that a less harmful cigarette can be developed. (58 refs)

- 78-5410 Inhalation Carcinogenesis.** (Eng) Nelson, (Inst. Environmental Medicine, New York Univ. Medical Center, New York, NY, 10016). *Ecotox Environ Safety* 1(3): 289-295; 1977.

A limited review of the carcinogenicity of various inhaled substances is presented. Chemical hazards in the workplace include benzene, nickel carbonyl, mustard gas, vinyl chloride, chloroprene, and bis(chloromethyl)ether (BCME). To date, at least 35 cases of lung cancer in the US have been attributed to BCME. Several methods for the safe and controlled inhalation of experimental animals to carcinogenic material are outlined. (22 refs)

- 78-5411 Carcinogenesis and Aging.** (Eng) Magee, P. (Fels Res. Inst., Temple Univ. Sch. Medicine, Philadelphia, PA, 19140). *Adv Exp Med Biol* 97: 133-137; 1978.

Factors affecting carcinogenesis that might be influenced by age are reviewed. Evidence that some human cancer is caused by chemicals is derived from studies on occupational and other forms of exposure and from studies on the geographical pathology of cancer. Studies with migrant populations indicate that genetic factors do not play a major role in the geographical variations of some cancers and that these variations are the result of environmental factors. Since the efficacy of most chemical carcinogens is dependent on the balance between their metabolic activation and deactivation, changes in the activities of any of the enzyme systems involved could influence their carcinogenic effects. Some enzyme activities are known to vary with age, but it is too early to draw conclusions on the effects of these changes on tumor induction. DNA appears to be the most likely target for the activation of forms of chemical carcinogens, but there is no evidence of any decline in DNA repair with increasing age. Cancers that are influenced by hormonal balance, including mammary and prostatic cancer, may be affected by age-related changes in the endocrine system. There is substantial evidence from human and animal studies that the increasing incidence of cancer with increasing age simply reflects the longer time that the cells have been exposed to carcinogens. (53 refs)

- 78-5412 Can Permissible Levels of Carcinogenic Chemicals in the Environment Be Envisaged? Chemical Remarks.** (Eng) Truhaut, R. (Centre de Recherches Toxicologiques, Faculté des Sciences Pharmaceutiques et Biologiques, Université René Descartes, 75006 Paris, France). *Ecotox Environ Safety* 1(1): 31-37; 1978.

Arguments for and against the theory of no thresholds for carcinogens are reviewed. The data available on dose-effect relationships for most environmental carcinogens, especially at low levels, are limited. Since all chemical carcinogens p

hazard, human exposure must be reduced to a socially acceptable level, similar to the approach taken with ionizing radiation. Such an approach will avoid the imposition of a burden or of an unrealistic economic burden. (13 refs)

8-5413 **Methods of Analyzing Environmental N-Nitrosamines.** (Rus) Kostiukovskii, I. L. (Inst. Nutrition, Moscow, USSR); Melamed, D. B. *Gig Sanit* (1): 6-68; 1978.

Methods of identifying, purifying, and determining environmental N-nitrosamines (NA) are reviewed. The following protocol is recommended for isolating and purifying NA: homogenization, steam distillation, distillation, evaporation from an acid and then alkaline medium, extraction, rising and dehydration of the extract, and chromatographic purification. Gas-liquid chromatography with nitrogen-sensitive detectors, chromatography/mass spectrometry, chemiluminescence, and Raman spectroscopy can be used to detect these compounds. (51 refs)

8-5414 **An Assessment of Human Exposure to N-Nitroso Compounds.** (Eng) Fine, D. H. (Thermo Electron Res. Center, Waltham, MA). *IARC Sci Publ* (19): 67-278; 1978.

Current knowledge concerning nitrosamine (NA) impurities is reviewed, and data on known exposure routes are normalized to their relative significance in terms of NA exposure per person per day. Volatile NA's are ingested in nitrite-preserved foods; data are not available for the occurrence of nonvolatile NA's in foods. N-Nitrosodimethylamine is present in wastewater from chemical factories at levels of 0.2-5 µg/liter, but the < 0.01-µg/liter levels of NA's found in drinking water are considered insignificant. Airborne NA's are inhaled in localized areas around certain specialized industries, tobacco smoke, aqueous mists of herbicides, and cutting fluid vapors. Dermal contact is experienced by machinists who handle machine parts that are wet with cutting fluid, agricultural workers who formulate or apply herbicides, and by persons (especially actresses and models) who use facial cosmetics, hair shampoos, hand lotions, and skin creams. All of these substances contain significant amounts of NA's. Tumors have been induced in laboratory animals by po administration of precursor amines + nitrite, presumably via in vivo nitrosation. In vivo nitrosation also takes place in humans fed diphenylamine + nitrite. Possible N-nitrosation mechanisms are presented. A high pH does not necessarily minimize the reaction, and in vivo nitrosation proceeds at a rate considerably faster than the one that could be predicted by conventional kinetics. (37 refs)

78-5415 **Interference of Carcinogens with Information Structures.** (Ita) Prodi, G. (Istituto di Can-

cerologia, Universita, Bologna, Italy); Grilli, S. *Recent Prog Med (Roma)* 63(6): 549-570; 1977.

The molecular mechanisms of chemical and viral carcinogenesis and relationships between mutagenesis and carcinogenesis are reviewed. The mechanisms of action of polycyclic aromatic hydrocarbons, aromatic amines, and nitrosamines are described. (37 refs)

78-5416 **Determination of N-Nitroso Compounds by Mass Spectrometry. A Review.** (Eng) Gough, T. A. (Lab. Government Chemist, Cornwall House, Stamford St., London SE1 9NQ, England). *Analyst* 103(1229): 785-806; 1978.

The detection of N-nitroso compounds, particularly the volatile N-nitrosamines, by mass spectrometry and by gas chromatography-mass spectrometry is reviewed, and the merits of these and other procedures reported in the literature are assessed. N-nitrosamines have been found in food, polluted air, industrial and agricultural chemicals, tobacco products, and drugs. (200 refs)

78-5417 **Experiments With In-Vitro-Test Systems to Prove Mutagenic Substances.** (Ger) Greim, H. (Munich, W. Germany); Goggelmann, W.; Wolff, T. *Staub-Reinhalt Luft* 38(6): 227-232; 1978.

Some relatively simple in vitro tests use bacteria or mammalian cells to identify mutagenic chemicals; the test compounds are metabolically activated by liver microsomes or the S-9 fraction with corresponding cofactors. The most widely used bacteria test systems, the Ames test and the Liquid test, are discussed with emphasis on problems that complicate the interpretation of test results, including those related to a compound's metabolism. Methods used to identify chemical mutagens in mammalian cells are summarized. (37 refs)

78-5418 **Experimental Carcinogenesis and Mutagenesis.** (Fre) Chouroulinkov, I. (Institut de Recherches Scientifiques sur le Cancer, BP-8, 94800 Villejuif, France). *Ann Mines* 184(7/8): 51-56; 1978.

The general hypotheses of carcinogenesis (genetic and epigenetic theories, viral gene activation, and cellular selection) are reviewed with special regard to the hypothesis on somatic mutations and the relationship between the carcinogenic and mutagenic properties of chemicals. A review of the literature shows that 88% of the tested carcinogens were also found to be mutagens, and 24% of the noncarcinogens were mutagens. Inversely, 93% of the mutagens and 61% of the non-mutagens proved to be carcinogens. (45 refs)

- 78-5419 **Aspects of Carcinogenesis.** (Ger) Reusch, K. (Institut für Toxikologie und Chemotherapie, Deutsches Krebsforschungszentrum, Im Neuenheimer Feld 280, 6900 Heidelberg, W. Germany). *Roentgenpraxis* 31(9): 197-204; 1978.

The mechanisms of chemical carcinogenesis are reviewed. The metabolic activation of a substance to a reactive product (ultimate carcinogen), the interaction of this product with cellular macromolecules, and the induction of changes in the genetic information of the cell are the principal events in chemical carcinogenesis. (11 refs)

- 78-5420 **Bladder Carcinoma after Long-Term Treatment with Cyclophosphamide for Multiple Myeloma.** (Ger) Schmidt, C. G. (Innere Klinik und Poliklinik, Westdeutsches Tumorzentrum, Universität Essen, Hufelandstrasse 55, D-4300 Essen 1, W. Germany). *Urologe[A]* 17(2): 109-110; 1978.

A patient with multiple myeloma developed carcinoma of the urinary bladder after long-term treatment with cyclophosphamide (total dose 135 g po). Most tumors that occurred after cytostatic therapy were found in patients treated with cyclophosphamide. This drug should not be used in the treatment of nonneoplastic diseases. (no refs)

- 78-5421 **A Review of the Genetic Toxicology of Chlorinated Dibenzo-p-dioxins.** (Eng) Wassom, J. S. (Environmental Mutagen Information Center, Information Center Complex/Information Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Huff, J. E.; Loprieno, N. *Mutat Res* 47(3/4): 141-160; 1978.

Published and unpublished information on the genetic toxicology of chlorinated dibenzo-p-dioxins, particularly 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), is reviewed. TCDD may be formed along with other dioxin compounds during manufacture of the herbicide 2,4,5-trichlorophenoxyacetic acid, the fungicide pentachlorophenol, and the germicide hexachlorophene. Research on dioxins as potential mutagens was initiated because of their structural similarity to acridines, a class of known intercalating agents. Only four dioxins have been evaluated for mutagenicity: the di-, tetra-, and octachlorinated derivatives and the unsubstituted dibenzo-p-dioxin. The results of these evaluations are inconclusive. Experiments with mice suggest that TCDD may be a hepatocarcinogen because of its specific cytological effect on proliferating hepatocytes. However, the cytological effects produced by TCDD in the liver of animals may be due to the toxic effects of the short-term high dose levels administered and to the fact that TCDD tends to concentrate in the liver. The

positive mutagenic and cytological effects reported with a few dioxin isomers examined seem to depend on the position of chlorine substitution, with TCDD being the most active form. Preliminary results of cytogenetic analyses of maternal blood and abortive fetal tissue from persons exposed to TCDD in Seveso, Italy (accidental release from a chemical plant), are inconclusive. However, workers exposed to herbicides Klorinol and Buvinol, which may have very low levels of TCDD contamination, had a higher incidence of chromatid breaks in peripheral lymphocytes than nonexposed individuals. The conflicting results of mutagenicity tests with dioxins could be due to solubility problems, treatment protocols, sample purity, and the extremes in toxicity of these compounds. (70 refs)

- 78-5422 **Genetic Toxicology of Lysergic Acid Diethylamide (LSD-25).** (Eng) Cohen, M. M. (Dept. of Human Genetics, Hadassah-Hebrew Univ. Medical Center, Jerusalem, Israel); Shiloh, Y. *Mutat Res* 47(3/4): 183-200; 1978.

The clastogenicity, mutagenicity, teratogenicity, and carcinogenicity of the hallucinogen lysergic acid diethylamide (LSD-25) are reviewed. It is impossible to draw clear-cut conclusions regarding any of the biological effects of LSD, despite the considerable scientific effort invested. Much of the difficulty arises from the lack of similarity of the experimental systems used, and, therefore, the lack of comparability. Most *in vitro* studies of the clastogenicity of LSD indicate that there is either a suppression of mitosis or enhanced chromosomal damage. However, extrapolation of these results to the *in vivo* situation is difficult. In humans, no consensus can be made with respect to chromosome breakage, and the inconsistencies within and between studies remain inexplicable. However, several controlled investigations of the *in vivo* effect of chemically pure LSD suggest that it produces a transient increase in lymphocyte chromosome breakage. In contrast, the results of cytogenetic studies in animals are contradictory. Although human mutagenicity studies are nonexistent, in those experimental organisms tested, using accepted techniques, LSD proved to be at best a weak mutagen. Teratogenicity studies in animals are confusing because of the large number of animal species and strains, dosages, and parameters studied. With regard to humans, however, enough time has elapsed after exposure for any such effects, if it exists, to have been expressed, and it can be concluded that LSD is not teratogenic in humans. As to the drug's oncogenic potential, the three reported cases of leukemia in LSD users are probably the result of coincidence. (160 refs)

- 78-5423 **Epidemiological Studies of the Occupational Hazards of Anaesthesia--A Review.** (Eng) V

M. P. (Univ. Dept. Social and Community Medicine, Radcliffe Road, Oxford OX1 3QN, England). *Anaesthesia* 33(4): 430-438; 1978.

British, and Scandinavian studies of the possible occupational hazards of anesthesia were reviewed. The studies, published in the period 1968-1977, indicate a moderate increase in the risk of spontaneous abortion among exposed women. There is no convincing evidence of any other hazards such as malformation of offspring, stillbirth, low birth wt, or infertility. (14 refs)

78-5424 **About Comparative Sensitivity of Certain Methods of "Drugs" Mutagenicity Estimation (Meeting Abstract).** (Eng) Revazova, J. (Res. Inst. Biological Testing Chemicals, Moscow Region, USSR); Radchenko, L.; Gerasimovskiy, V.; Podolnaya, M. *Mutat Res* 53(2): 252; 1978. (no refs)

78-5425 **Mutagenicity of Hair Dyes: Some More Evidence and the Problems of its Interpretation (Meeting Abstract).** (Eng) Venitt, S. (Div. Chemical Carcinogenesis, Inst. Cancer Res., Pollards Wood Res. Station, Buntingford Lane, Chalfont St. Giles, Bucks, England). *Mutat Res* 53(2): 278-279; 1978. (no refs)

78-5426 **Review of Submammalian Systems for Detecting Mutagens and Carcinogens (Meeting Abstract).** (Eng) Sugimura, T. (Natl. Cancer Center Res. Inst., Tokyo, Japan). *Mutat Res* 53(2): 270-271; 1978. (no refs)

78-5427 **Deficiencies in the Microbial Activation System for Mutagenic-Carcinogenic Screening (Meeting Abstract).** (Eng) Rinkus, S. (Div. Environmental Toxicology and Epidemiology, Univ. Texas Medical Branch, Galveston, TX, 77550); Legator, M. *Mutat Res* 53(2): 254; 1978. (no refs)

78-5428 **In Vivo Cytogenetic Screening of Drugs for Mutagenic Potential (Meeting Abstract).** (Eng) Uchimoto, T. (Biological and Medical Res. Div., Sandoz

Ltd., Basel, Switzerland); Matter, B. E. *Mutat Res* 53(2): 277; 1978. (no refs)

78-5429 **Metal Carcinogenesis (Meeting Abstract).** (Eng) Sunderman, F. W. (No affiliation given). In: *Arnold O. Beckman Conference on Biochemistry of Cancer held in San Antonio, September 6-8, 1978*. American Association for Clinical Chemistry (San Antonio, TX): p. 9; 1978. (3 refs)

78-5430 **Pseudolymphomas and Lymphomas Induced by Hydantoins.** (Fre) Reiffers, J. (Service d'Hematologie, Hopital Henri-Mondor, Bordeaux, France). *Nouv Rev Fr Hematol Blood Cells* 20(1): 131-137; 1978.

Literature cases of patients who developed pseudolymphomas (PL) and malignant lymphomas after treatment with phenytoin or diphenylhydantoin are reviewed. More than 90 patients with PL and 19 with lymphomas have been reported. The similarity of the PL induced by hydantoins to immunoblastic lymphadenopathy and the precancerous nature of PL are discussed. (52 refs)

78-5431 **Hepatic Changes Produced by Contraceptive Steroids.** (Spa) Uribe, M. (Departamento de Gastroenterologia, Instituto Nacional de la Nutricion, San Fernando y Viaducto Tlalpan, Mexico City 22, Mexico); Ramos de Uribe, M. H.; Cario, A. A. *Rev Gastroenterol Mex* 42(3): 139-147; 1978.

The side effects of hormonal contraceptives include jaundice, cholestasis, hepatitis, hepatic adenoma, and liver cancer. Literature cases of hepatic adenoma and liver carcinoma number about 250 and 21 patients, respectively. (75 refs)

78-5432 **Estrogens and Risk of Carcinoma. Practical Consequences for Estrogen Therapy.** (Ger) Kaiser, R. (Univ.-Frauenklinik, Kerpener Strasse 34, D-5000 Cologne 41, W. Germany). *Munch Med Wochenschr* 120(37): 1185-1186; 1978.

Relationships between estrogen treatment and cancer in women are discussed. An association between estrogen treatment and precancerosis or carcinoma of the endometrium cannot be ruled out with certainty. There is no firmly established relationship between breast cancer and estrogens. (no refs)

- 78-5433 **Genetic Effects of Coumarins.** (Eng) Grigg, G. W. (Molecular and Cellular Biology Unit, CSIRO, P.O. Box 184, North Ryde 2113, New South Wales, Australia). *Mutat Res* 47(3/4): 161-181; 1978.

Coumarins (CM) that modify or cause the cell to modify the structure or properties of genetic material, except for the brightening agents (psoralens, or furocoumarins) and the aflatoxins, which have been surveyed recently, are reviewed. CM have been shown to induce C-mitotic effects, chromosome breakage, and various sorts of rearrangements and to affect DNA repair and mutation frequency in bacterial, plant, and animal cells. The hepatic toxicity to rats that showed up in routine toxicology tests caused the voluntary withdrawal of CM from the foodstuffs industry, in which it was used as a flavoring substance. Tests with rats and dogs indicated that feeding CM initiated liver damage by damaging the endoplasmic reticulum. The principal metabolite of CM in rats, O-hydroxyphenylacetic acid, is thought to be the main cause of liver toxicity to the animal. The emergence of lines of rodents resistant to the CM derivative coumadin (warfarin) indicates that there is a genetic variability in the toxic response to the drug or in its metabolism in these animals. A similar variability emerged in human studies. Patients have been reported as being very sensitive or resistant to the anticoagulant dicoumarol. Many CM are effective chromosome-breaking agents, yet others have antimutagenic activity (coumadin). In one study high levels of CM (0.5% of diet for 2 yr) were hepatocarcinogenic in rats, but in another CM suppressed 7,12-dimethylbenzanthracene-induced breast carcinomas in rats, possibly because of its ability to modify the activity of liver microsomal enzymes. Long-term adjuvant therapy with coumadin, phenprocoumon, and dicoumarol enhanced the survival of cancer patients. Experiments with mice indicated that CM enhances survival by preventing metastasis. The paradoxical effects of CM are similar to those of caffeine, which has been found to be a chromosome-breaking agent, a mutagen, and an antimutagen. The fact that both caffeine and CM bind selectively to single-stranded DNA and consequently destabilize the double helical conformation in favor of the random coil configuration may help to explain their genetic properties. (152 refs)

- 78-5434 **PUVA: A Caution.** (Eng) Segal, S. (No affiliation given); Cohen, S. N.; Freeman, J.; Hill, R. M.; Kagan, B. M.; Kauffman, R. E.; Pruitt, A. W.; Soyka, L. F.; Vickers, S. M. *Pediatrics* 62(2): 253; 1978.

Caution should be exercised in the use of a form of photochemotherapy known as PUVA (psoralen with UV light in the A range) for the treatment of severe psoriasis. PUVA produces changes in DNA that have been linked experimentally to teratogenesis and carcinogenesis. Under no circumstances should children be treated with PUVA, except under established IND (investigational new drug) protocols. (7 refs)

- 78-5435 **Toxic Effects of Vinyl Chloride.** (Ger) Veltr G. (Univ.-Hautklinik Bonn-Venusberg, D-53001 Bonn 1, W. Germany); Lange, C. E.; Stein, G. *Hautarzt* 29(4): 177-182; 1978.

The toxicological aspects of vinyl chloride (VC) are reviewed on the basis of examinations of workers occupationally exposed to VC over long periods of time. The liver histology revealed periportal, septal, and intralobular fibrosis; focal reticular collagenization of the sinusoid walls; hepatocyte regeneration; and activation and proliferation of the sinusoidal cells, with cell atypia and transition into angiosarcoma occurring in some cases. The metaphase analysis revealed no remarkable pathological changes; ie, no mutagenic effect of VC. (35 refs)

- 78-5436 **Vinyl Chloride and Vinyl Benzene (Styrene) Metabolism, Mutagenicity and Carcinogenicity.** (Eng) Vainio, H. (Dept. Industrial and Toxicology, Occupational Health, Haartmaninkatu 1, SF-00290 Helsinki 29, Finland). *Chem Biol Interact* 22(1): 117-124; 1978.

The metabolism, mutagenicity, and carcinogenicity of vinyl chloride (VC) and vinyl benzene (styrene) are reviewed briefly. VC and styrene are mutagenic in bacterial test systems, *Drosophila*, yeast, and mammalian cells. The mutagenicity of VC in bacterial test systems depends, at least partially, on metabolic activation by microsomal enzymes. Chloroethylene oxide, the primary biotransformation product of VC, is a potent mutagenic and alkylating agent. Styrene is mutagenic to *Salmonella typhimurium*, but only after metabolic activation, whereas styrene oxide, its primary biotransformation product, is mutagenic to *Salmonella* without activation. In several studies, an excess of chromosome aberrations (compared with controls) was found in the lymphocytes of workers exposed to VC monomer. In addition, an excess of fetal loss occurred among women whose husbands are heavily exposed to VC. Workers exposed to styrene also had increased chromosome aberrations in their lymphocytes. Both chloroethylene oxide and styrene oxide bind covalently to cellular macromolecules. VC is carcinogenic in rats (skin, lung, bone marrow, liver) and humans (liver, brain, lung, and lymphatic tissue). Styrene oxide was a weak carcinogen when applied to mouse skin. Styrene is currently being tested in animals. These findings raise concern about the possible genetic risks of VC exposure to humans. (55 refs)

- 78-5437 **How Much Benzene Causes Cancer?** (Eng) Thorpe, J. J. (Exxon Corp., New York, NY). *Hydrocarbon Process* 57(5): 172-174; 1978.

Data on the acute and chronic toxicity and carcinogenicity

enzene are reviewed. Although benzene has not induced leukemia in experimental animals, it appears that occupational exposure to benzene is causally related to acute myelogenous leukemia and its variants. Arguments for and against a new US permanent standard (February 1978) are presented. (10 refs)

78-5438 Antigens of Chemically Induced Sarcomas: Their Possible Origins and Interrelationships. (Eng) Blackstock, R. (Dept. Pediatrics, Univ. Oklahoma Health Sciences Center, Oklahoma City, OK, 73100). *South Med J* 71(5): 586-590; 1978.

Antigens on the surface of chemically induced tumors (ACT) are categorized, and their relationship to H-2 antigens is discussed. There are four types of ACT: tumor-specific antigens, viral or embryonic antigens, viral antigens, and normal tissue antigens. ACT have several characteristics in common with normal H-2 alloantigens of mice: both serve as targets for transplantation immunity, both have similar cell-surface location and mobility, and both show a high degree of polymorphism. (104 refs)

78-5439 Immunoproliferation and Cancer: A Common Macrophage-derived Promotor Substance. (Eng) Salmon, S. E. (Section Hematology and Oncology, Univ. Arizona Coll. Medicine, Tucson, AR, 85724); Hamberger, A. W. *Lancet* 1(8077): 1289-1290; 1978.

It is hypothesized that a common proliferative factor is required for the growth of clones of both antigen-triggered B lymphocytes and carcinogen-triggered epithelial cells and that this factor is secreted by macrophages that have originally differentiated in support of the immune response. The promoter in the two-stage model of carcinogenesis in mouse skin induces epithelial proliferation and intense cutaneous infiltration with inflammatory cells. Viable adherent splenic macrophages from mineral oil-treated BALB/c mice produce a factor that, in concert with 2-mercaptoethanol (2-ME), promotes soft-agar colony formation by human myeloma cells. Medium conditioned by the oil-primed BALB/c macrophages is able to support in vitro tumor colony growth from fresh biopsy specimens of a wide variety of cancers. Normal murine B-lymphocyte colony formation also requires 2-ME and is promoted by a soluble factor elaborated by macrophages. Thus, both immune and tumor clones may require a common and specific growth stimulus that promotes clonal proliferation of an already triggered hypermutable cell population. This factor would also incidentally stimulate the growth of clones of tumor cells that had been previously oncogenically transformed. Some clinical features of metastasis

may also be consistent with the role of the macrophage as a promoter of tumor growth. (15 refs)

78-5440 Hamster Immune Responses: Experimental Models Linking Immunogenetics, Oncogenesis and Viral Immunity. Introduction. (Eng) Streilein, J. W. (Dept. Cell Biology, Univ. Texas Health Science Center, Dallas, TX, 75235). *Fed Proc* 37(7): 2023-2024; 1978.

Subjects covered at a symposium on hamster immune responses are outlined. The topics include the normal immune response, immunogenetics and transplantation, response to infection (particularly viral), and tumor immunity, along with the high susceptibility of hamsters to oncogenic agents, their low spontaneous rate of malignant degeneration, and the development of spontaneous lymphomas in a hamster colony. (no refs)

78-5441 Epstein-Barr Virus as the Cause of a Human Cancer. (Eng) Epstein, M. A. (Dept. Pathology, Univ. Bristol, Bristol, England). *Nature* 274(5673): 740; 1978.

Although Epstein-Barr virus (EBV) appears to be the cause of Burkitt's lymphoma, a climate-dependent cofactor, possibly malaria, also appears to be involved in the disease. The fact that only B cells have receptors for EBV and that malaria is both immunosuppressive and a stimulator of B-cell proliferation support this idea. The malaria may stimulate and maintain a subclass of B cells susceptible to EBV-induced transformation. (20 refs.)

78-5442 Biological Approaches to Human Cancer Virus Detection. (Eng) Kaplan, H. S. (Cancer Biology Res. Lab., Dept. Radiology, Stanford Univ. Sch. Medicine, Stanford, CA, 94305). *Prog Exp Tumor Res* 21: 2-18; 1978.

Procedures for detecting a human cancer virus are outlined in detail. The first step is the cultivation of freshly procured human tumor tissue in vitro. Care must be taken to insure that the culture is not infected with other cell lines. Cultured cells should be identified by morphological, cytochemical, and surface marker techniques with the original population, and it must be shown that they have the biological characteristics of neoplastic cells. The cultures should then be examined for release of viruses and the presence of viral antigens and reverse transcriptase. Cocultivation of the tumor cells with indicator cell lines derived from the same and other species is a sensitive technique for the transcellular transmission and detection of candidate viruses. Since the expression

of virus-specific antigens and the replication of infectious virus may be under strict control of the host genome, the cultures should be subjected to treatment that interferes with the host cell gene regulatory mechanisms. Once virus particles are isolated, they should be tested for oncogenicity in subhuman primates and congenitally athymic nude mice. Also, normal human cells infected with the virus could be propagated in vitro and observed for distinctive changes. Another approach would be to inject the putative tumor virus into rabbits or goats, isolate tumor antiserum, and test the antiserum for specificity against the original tumor. If positive reactions are observed with virus-specific antigens, the serum could be used to screen other tumors of the same morphology from other patients. A candidate human tumor virus that is truly the cause of the tumor from which it was isolated should induce the same or similar antigens in all similar-type tumors. (77 refs)

- 78-5443 The Immunological Approach to Study of Possibly Virus-induced Human Malignancies Using the Epstein-Barr Virus as Example.** (Eng) Henle, W. (Div. Virology, Joseph Stokes, Jr. Res. Inst., Children's Hosp. Philadelphia, Conshohocken Ave., Philadelphia, PA, 19131); Henle, G. *Prog Exp Tumor Res* 21: 19-48; 1978.

Immunological evidence linking Epstein-Barr virus (EBV) with Burkitt's lymphoma (BL) and nasopharyngeal carcinoma (NPC) by serologic surveys and by detection of EBV-determined antigens in tumor cells is reviewed. Particular attention is focused on antibody titers to the various EBV antigens in BL and NPC. The role of EBV in infectious mononucleosis is also reviewed. Since BL and NPC have either immunosuppressive effects or require therapy by immunosuppressive agents, activation of the latent persistent viral carrier state in patients with high anti-EB viral capsid antigen titers appears to be the most plausible explanation for the elevated antibody titers. (121 refs)

- 78-5444 On the Toxicity of Inhaled Hot Particles with Special Reference to Plutonium.** (Eng) Muth, H. (Institut für Biophysik, Universität des Saarlandes, Boris Rajewsky-Institut, D-6650 Hamburg, W. Germany); Jacobi, W.; Feinendegen, L. E.; Hug, O.; Oberhausen, E. *Radiat Environ Biophys* 15(1): 1-11; 1978.

A review of the epidemiological literature indicated that after > 30 yr of work with ^{239}Pu in the US, no lung cancer cases could be attributed to the incorporation of Pu. Experimental studies in animals gave no indication that hot-particle inhalation leads to a larger risk of lung cancer than homogeneous irradiation. It is suggested that no reduction in the existing

limit for inhalation of insoluble ^{239}Pu compounds be made. (35 refs)

- 78-5445 Asbestos Fibre Dust and Gastro-intestinal Malignancies. Review of Literature with regard to a Cause/Effect Relationship.** (Eng) Miller, A. (NCIC Epidemiology Unit, Univ. Toronto, 121 Saint Joseph St., Toronto, Ontario, M5S 2R9, Canada). *J Chronic Dis* 31(1): 23-33; 1978.

A review of the literature on the association between exposure to asbestos dust and gastrointestinal malignancies indicated that such an association does exist, particularly cancer of the esophagus, stomach, colon, and rectum. In the absence of an explanation to the contrary, this exposure may be regarded as causal. There appears to be a latent period of ≥ 20 yr between the first exposure and the development of malignancy. A worker with ≥ 20 yr asbestos exposure has approximately a threefold greater risk of gastrointestinal cancer than an unexposed individual. There is no evidence that other factors contribute to or subtract from this association. (39 refs)

- 78-5446 Assessment of Hazards in Engineering Work.** (Eng) Kinchin, G. H. (Safety and Reliability, UKAEA, Culcheth, Warrington, England). *Proc Institution Civil Eng* 64: 431-438; 1978.

Acceptable risk criteria to be incorporated in the design of nuclear power plant facilities are outlined. (14 refs)

- 78-5447 Low-Dose Radiation: The Hanford Evidence (Letter to Editor).** (Eng) Stewart, A. (Regional Cancer Registry, Queen Elizabeth Hosp., Birmingham 2TH, England). *Lancet* 1(8072): 1048-1049; 1978.

Data on deaths and radiation doses at a plutonium plant that was monitored from 1944 to 1972 suggest that the cancer risk from the current permissible dose (5 rads/yr) could be as high as 1/150. The mean dose was higher for the 760 employees with fatal cancers than for the 2,850 employees with deaths due to other causes. Expansion of nuclear facilities should be curtailed until correct measures of the cancer hazard are made. (4 refs)

- 78-5448 Derepression of a Dormant Bacteriophage as a Test for Potential Carcinogens (Meeting Abstract).** (Eng) Devoret, R. (Section de Radiobiologie C

re, Laboratoire d'Enzymologie, CNRS, 91190
sur-Yvette, France); Moreau, P.; Bailone, A. *Mutat Res*
: 178-179; 1978. (4 refs)

5449 **Experience and Correlation of Bacterial Muta-
genic Tests (Meeting Abstract).** (Eng) Watkins,
A. (Pharmaceuticals Div., Imperial Chemical Industries
Ltd., Macclesfield, Cheshire, England). *Mutat Res* 53(2): 283;
1978. (no refs)

5450 **Fundamental Mutagenic Mechanisms (Meeting
Abstract).** (Eng) Drake, J. W. (Dept. Microbi-
ology, Univ. Illinois, Urbana, IL, 61801). *Mutat Res* 53(2):
181; 1978. (no refs)

5451 **Human Inherited Disease with Altered Mech-
anisms for DNA Repair and Mutagenesis.** (Eng)
Singer, J. E. (Lab. Radiobiology, Univ. California, San
Francisco, CA). *Excerpta Med Int Congr Ser* (432): 85-100;
1978.

A variety of human disease (ie, xeroderma pigmentosum and
ataxia telangiectasia) involving increased cancer risk and as-
sociated disorders in the CNS, in hematopoietic, immunolog-
ical, ocular, and cutaneous tissues, and in embryological de-
velopment have defects in biochemical pathways for excision
repair of damaged DNA. Excision repair has multiple
branches by which damaged nucleotides, bases, and
cross-links are excised, and it required cofactors that control
access of repair enzymes to damage in chromatin DNA.
(no refs)

5452 **Growth Factors for Tumours.** (Eng) Shields, R.
(Imperial Cancer Res. Fund Lab., London, En-
gland). *Nature* 272(5655): 670-671; 1978.

Evidence that ectopic growth factors are involved in tumor
growth and development is summarized. A human fibrosar-
coma cell line produces a growth factor similar to MSA, a
matomedinlike factor that was first described as the prod-
uct of a rat liver cell line grown in serum-free medium. Pa-
tients with nonislet cell tumors and hypoglycemia have been

reported to have elevated serum levels of MSA-like material
whose nonsuppressible insulinlike activity accounts for the
hypoglycemia. A variety of other cells also manufacture
growth factors. For example, a line of BHK cells transformed
by simian virus 40 produces a factor that stimulates the
growth and movement of 3T3 cells. Growth factors have also
been found in medium conditioned by L929 cells; these in-
clude nerve growth factor, a colony stimulating factor that
is identical to macrophage growth factor, and other unidenti-
fied growth factors. Cells that synthesize ectopic growth fac-
tors, make new hormone receptors, or induce fresh blood
supply through the synthesis of tumor angiogenesis factor
may gain a selective advantage over their normal counter-
parts. Ectopic protein synthesis could then play a vital part
in the growth and development of many types of tumors. (25
refs)

78-5453 **Developmental Potency of Normal and Neo-
plastic Cells of the Early Mouse Embryo.** (Eng)
Gardner, R. L. (Dept. Zoology, Univ. Oxford, Oxford, En-
gland). *Excerpta Med Int Congr Ser* (432): 154-166; 1978.

Recent studies of the potency of normal early mammalian
embryo cells and their neoplastic counterparts are reviewed
along with studies of X inactivation, and some of the
implications of this work are considered. Teratocarcinomas
develop from both male and female germ cells in mice, and
they can be produced by grafting early embryos ectopically
in syngeneic adult hosts. This review is restricted to studies
of the potency of the embryonal carcinoma stem cells (EC
cells) of these tumors by blastocyst injection. So far, such
transplantations have been carried out only with EC cells of
embryo-derived teratocarcinomas, both from in vivo-pass-
aged tumors and in vitro cultures. The results of these experi-
ments demonstrate that EC cells from both sources can partic-
ipate in embryogenesis and exhibit normal differentiation
in a wide variety of organs and tissues of the conceptus and
postnatal mouse. The range of normal cell types formed by
EC cells in chimeras exceeds that found in both the tumors
and in culture and has included functional spermatozoa in
one instance. Most chimeras produced by injecting groups of
cultured EC cells into blastocysts also developed tumors that
invariably contained cells of donor origin. In single cell injec-
tion experiments, the EC cell clone in one animal autopsied
at 4 wk postpartum had colonized every somatic organ and
tissue tested. Several studies show that EC cells can retain
their totipotency despite in vivo passage for several years,
with or without an additional period of months in culture.
EC Cells may correspond to cells of the germ line that re-
quire totipotency precociously as a consequence of disruption
of embryonic organization through ectopic grafting. Studies
of injecting EC cells into blastocysts also furnish an example
of reversal of neoplasia in animal cells. However, it remains
to be seen whether teratomas and teratocarcinomas represent
special cases or whether conditions can be found that will

enable normal growth and differentiation of stem cells from other types of neoplasms to occur. (84 refs)

- 78-5454 Growth of Human Tumor Cells in Established Cultures.** (Eng) Sinkovics, J. G. (Section Clinical Tumor Virology and Immunology, Dept. Medicine, M. D. Anderson Hosp. and Tumor Res. Inst., Univ. Texas System Cancer Center, Houston, TX, 77030); Gyorkey, F.; Kusyk, C.; Siciliano, M. J. *Methods Cancer Res* 14: 243-323; 1978.

Experience with the growth of human neoplastic cells in established cultures is reviewed. In 75% of the primary cultures set, groups of cells deemed to be neoplastic were observed, but only a small number of these cultures developed into established neoplastic cell lines. Sarcomas, carcinoma, and malignant melanoma cell lines are considered. The morphology and ultrastructure of the cell lines are illustrated in numerous figures, and some of the characteristics of each line (highest cell passage attained; morphology; patient's age, gender, race, and blood group; and verified tissue diagnosis of tumor-yielding cell line) are tabulated. Methods for characterizing established tissue cultures include determination of generation time and growth kinetics, cloning, karyotypic and chromosome banding, enzyme electrophoresis, and growth in immunosuppressed or immunoincompetent animals. The major aims of working with human tumor cell lines should remain (1) the representation of each human tumor category by several established cell lines for extensive laboratory studies and (2) the achievement of technical and procedural advances in the yield of established cell lines. (176 refs)

- 78-5455 Genetics of Neoplasia--Impact of Ecogenetics on Oncogenesis. A Review.** (Eng) Purtilo, D. T. (Dept. Pathology, Univ. Massachusetts Medical Sch., 55 Lake Ave. N., Worcester, MA, 01505); Paquin, L.; Gindhart, T. *Am J Pathol* 91(3): 607-688; 1978.

The basic genetic mechanisms responsible for neoplasia are reviewed to guide the clinician in the differential diagnosis of inherited neoplastic syndromes. Particular attention is focused on the role of ecogenetics, the long-term variable effects of the environment on the genome, in the development of cancer. The specific topics include the mutational theory of oncogenesis, chromosome aberrations, marker chromosomes, clastogenic syndromes, the association of mutagenesis with teratogenesis and oncogenesis, familial cancer syndromes, the X-linked lymphoproliferative syndrome as a model demonstrating the impact of ecogenesis on oncogenesis, and the diagnosis of familial cancer syndromes. The familial cancer syndromes discussed include genodermatoses, neurofibromatosis, multiple endocrine neoplasia, familial CNS tumors, familial colon carcinoma, breast carcinoma, lung carcinoma, urinary system carcinomas, and familial hematopoietic malignancies. A medical history questionnaire for lymphoproliferative study is appended. (410 refs)

- 78-5456 Highlights of Neurosurgery in Nigeria.** (Eng) Adeloye, A. (Dept. Surgery, Univ. Ibadan, Ibadan, Nigeria). *Trop Geogr Med* 29(4): 325-334; 1977.

The major types of neurological operations performed in Nigeria are reviewed. In Ibadan, the incidence of brain tumors is less than that seen among European communities. The male:female ratio is 5:1, and the patients are younger than those noted in a European series. The most common tumors are meningiomas and pituitary tumors; glioblastoma multiforme is rare, and acoustic neuroma is virtually unknown. Approx 75% of the meningiomas are found in the convexity region, the remainder in the basal region histologically, and of these tumors are of the syncytial type. The lesions drain the bulk of their blood from the internal carotid artery. The most common pituitary tumor is the chromophobe adenoma. Tuberculoma is one of the most important causes of space-occupying lesions. Burkitt's lymphoma affects the brain in about 60% of the cases and the brain, spinal cord, and peripheral nerves in about 20% of the cases. The intracranial canal tumor pattern in Nigeria does not differ markedly from that of other places. One interesting lesion found is intramedullary spinal cord schistosomal granuloma due to *Schistosoma hematobium*. Malignant trophoblastic disease involves the nervous system in about 12% of the cases. Carcinoma of the thyroid gland is the commonest source of metastases to the skull in Nigeria. Metastatic tumors of the brain carry a 4:1 female:male incidence; the most frequent source of secondary tumor is choriocarcinoma. (41 refs)

- 78-5457 Nelson's Syndrome: Chickens, Eggs, and Scapegoats. Explosions.** (Eng) Conomy, J. P. (Cleveland Clinic Foundation, 9500 Euclid Ave., Cleveland, OH 44106). *Arch Intern Med* 138(5): 691-692; 1978.

The origin of pituitary tumors in Nelson's syndrome is discussed. The sequence and tempo of the illness usually consist of hyperadrenal corticism, adrenalectomy, progressive hyperpigmentation, and pituitary tumor; the pituitary tumor may follow the adrenal extirpation by many months or years. Studies have indicated that the incidence of this syndrome in postadrenalectomy patients is 10%. Inferential clinical evidence suggests that, most frequently, removal of the adrenal tumor is followed by development of the pituitary tumor; in one series the reverse sequence is less common: < 3%. Of the 120 patients undergoing surgical treatment of hyperadrenal corticism were known to have a preexisting pituitary tumor. Of the 8% to 40% of postadrenalectomy patients in whom pituitary tumors are manifested, 50% are known to have any pituitary tumors to begin with. The pituitary tumors that do develop are prone to enlarge, explode, and compress parasellar structures. One study indicated that half the patients with postadrenalectomy pituitary tumors had an abnormal sella turcica and that 25% of these had an acute symptomatic necrotic disruption of the tumor. (8 refs)

78-5458 **Genetically Susceptible Subgroups.** (Eng) Strong, L. C. (Graduate Sch. Biomedical Sciences, Univ. Texas Health Science Center at Houston, Houston, TX, 77025). *Environ Health Perspect* 22: 139-140; 1978.

Disorders characterized by chromosomal instability or increased mutagenesis in vivo or in vitro may predispose to malignancy. These disorders include xeroderma pigmentosum, Fanconi's anemia, Bloom's syndrome, and ataxia telangiectasia. There may be a genetically determined increased risk of mutation and cancer following exposure to certain environmental agents as a result of a genetically determined metabolism of that particular agent. If malignancy is the result of sequential mutations, susceptible subgroups might include all individuals who have sustained a primary mutation through a germinal or somatic cell. Individuals who inherit the predisposing mutation are at greatest risk because each cell of the sensitive tissue carries the mutant gene, and a subsequent mutation in any cell may lead to malignancy. The sequential mutation model also accounts for the synergistic, multiplicative effects of mutagens on the risk of cancer in general. Promotional factors may be critical in the probability of cancer in individuals who acquire the first mutation in a somatic cell. Study of uniquely susceptible subgroups may contribute to an understanding of carcinogenesis in general. (no refs)

78-5459 **Genetics and Cancer in Children.** (Eng) Lynch, H. T. (Dept. Preventive Medicine and Public Health, Creighton Univ. Sch. Medicine, Omaha, NB); Lynch, P. M. *Pediatr Ann* 7(6): 71-93; 1978.

The influence of primary genetic events, radiation, congenital birth defects, environmental interactions, and cytogenetic aberrations in familial childhood cancer is discussed. The development of a national registry of cancer-prone families is recommended as an aid to the study, management, and control of high-risk children. (80 refs)

78-5460 **Breast Cancer Morbidity and Mortality in the Soviet Union.** (Rus) Merabishvili, V. M. (N. N. Petrov Res. Inst. Oncology, Leningrad, USSR); Dymarskii, A. Iu. *Vopr Onkol* 24(8): 3-9; 1978.

Indices of morbidity and mortality due to malignant breast tumors in the Soviet Union are reviewed. Breast cancer morbidity increased from $8.4/10^5$ in 1960 to $12.5/10^5$ in 1975; correspondingly, the mortality index increased from $9.9/10^5$ to $6.4/10^5$. The absolute number of patients

with breast cancer increased by 36.9% from 1970 to 1975 and it reached 177,469. (16 refs.)

78-5461 **Epidemiology of Cervical Cancer.** (Ger) Berndt, H. (Klinik für innere Medizin, Bezirkskrankenhaus, Dr. Wilhelm-Kulz Strasse, Postfach 480, DDR-20 Neubrandenburg, E. Germany); Neuser, D. *Arch Geschwulstforsch* 48(3): 250-275; 1978.

Differences in cervical cancer incidence and mortality between countries, the known risk factors in the disease, and East Germany's preventive control program are reviewed. The importance of treating preneoplastic lesions (dysplasia and carcinoma in situ) is emphasized. (262 refs)

78-5462 **HLA System and Malignant Diseases.** (Fre) Feingold, N. (Group de Recherches de Genetique epidemiologique, INSERM, U.155, Chateau de Longchamp, Bois de Boulogne, F75016 Paris, France). *Bull Cancer (Paris)* 65(1): 79-82; 1978.

Investigative work in the last 10 yr on the possible association of antigens of the major histocompatibility system (HLA) and malignant disease is reviewed. There is some evidence for the association of antigens B5, B15, B18, BW35, and A1 with Hodgkin's disease. Antigens A2 and B12 appear to be associated with acute lymphoblastic leukemia. Except for the association of Sin2 and nasopharyngeal cancer, there is no statistical support for an association of specific HLA antigens and solid tumors. Intrapopulation studies show that the incidence of some cancers may follow the incidence of certain HLA antigens. These coincident cancer and HLA patterns may indicate a factor of susceptibility or resistance to cancer in a particular population. (16 refs)

78-5463 **Epidemiological Outlook on Cancer.** (Eng) Curnen, M. G. (Sch. Public Health, Columbia Univ., 600 W. 168th St., New York, NY, 10033). *Bull NY Acad Med* 54(4): 349-365; 1978.

An epidemiological overview of cancer incidence in the US is presented. The most common cancers in men are those of the lung, colon and rectum, and prostate; the most common in women are those of the breast, colon and rectum, and lung. Survival rates of these and other major cancers indicate that only cancers of the colon and rectum, lung, and pancreas have 5-yr survivals of $<50\%$. Most cancers appear to be linked to environmental pollutants. (10 refs)

CHEMICAL CARCINOGENESIS

- 78-5464 **Experimental Studies on the Carcinogenicity of Bracken Fern (*Pteridium aquilinum*) in the Rat.** (Jpn) Kato, T. (First Dept. Pathology, Sch. Medicine, Gifu Univ., Tsukasa-cho, Gifu 500, Japan). *Gifu Daigaku Igakubu Kiyo* 26(2): 318-333; 1978.

The carcinogenicity of bracken fern (*Pteridium aquilinum*) was determined in feeding experiments in ACI rats. Three groups of rats received a diet containing 5%, 10%, and 30% bracken, respectively, for 60 days and three groups received the corresponding diets for 480 days. Most of the resulting tumors were adenomas and adenocarcinomas of the ileum, and the commonest site was the terminal 20 cm of the ileum, particularly the terminal 3-cm section nearest the ileocecal junction. Ileal sarcomas and urinary bladder carcinomas were infrequent. For the three dose groups, rats treated for 60 days had tumor incidences of 11.1%, 25.0%, and 56.2%, respectively; rats treated for 480 days had incidences of 38.4%, 93.3%, and 92.8%. Tumor multiplicity was dose-related. The intestine was much more susceptible than the urinary bladder to the carcinogenicity of bracken: a > 10% bracken diet for 480 days was necessary for the induction of urinary bladder tumors. Atypical hyperplasia of the small intestinal mucosa usually appeared 14 wk after the start of the bracken diet. After about 28 wk, the frequency of atypical hyperplasia was reduced and the adenomas and adenocarcinomas became prominent. (18 refs)

- 78-5465 **Carcinogenicity of Boiling Water Extract of Bracken, *Pteridium aquilinum*.** (Eng) Hirono, I. (Dept. Carcinogenesis and Cancer Susceptibility, Inst. Medical Science, Shiroganedai 4-6-1, Minato-ku, Tokyo 108, Japan); Ushimaru, Y.; Kato, K.; Mori, H.; Sasaoka, I. *Gann* 69(3): 383-388; 1978.

The carcinogenicity of an aqueous extract of bracken fern (*Pteridium aquilinum*) was studied in inbred ACI rats. In experiment I, Group 1 rats received a diet containing unprocessed dry bracken powder for 3 mo; Group 2 received a boiling water extract of dry bracken as drinking water for 16 mo; Group 3 received a cold water extract of dry bracken as drinking water for 16 mo; and Group 4 received a diet containing a concentrated boiling water extract of dry bracken for 16 mo. Rats in experiment II received a diet containing a concentrated boiling water extract of dry bracken, for 500 days. A control group was fed a normal diet. Most of the Group 1 animals developed multiple ileal tumors, such as adenomas and adenocarcinomas. In Group 2, 6/11 rats that survived > 7 mo developed tumors in the ileum and/or the urinary bladder. Group 3 rats had no tumors. In Group 4 and experiment II rats, all animals except one in experiment

II had urinary bladder tumors, and almost all the rats in the groups also had ileal tumors. These results show that carcinogen in bracken fern is extractable in boiling water. (refs)

- 78-5466 **Mycotoxin Mutagenesis--Mutagenic Activity of Some Species of Subarctic Grain Fungi (Meeting Abstract).** (Eng) Sorsa, M. (Dept. Genetics, Univ. Helsinki, Helsinki, Finland); Linnainmaa, K.; Ilus, T. *Mutat Res* 53(2): 267; 1978. (1 ref)

- 78-5467 **Mitotoxicity of β -Exotoxin and δ -Endotoxin Protein from *Bacillus thuringiensis* (Meeting Abstract).** (Eng) Sahu, R. K. (Cytogenetics Lab., Berhampur Univ., Berhampur 760 007, Orissa, India); Panda, B. Sharma, C. B. *Mutat Res* 53(2): 256-257; 1978. (no refs)

- 78-5468 **Induction of Renal and Hepatic Tumors in Mice by Ochratoxin A, a Mycotoxin.** (Eng) Kaniwa, M. (Dept. Pathology, Tokyo Metropolitan Inst. Gerontology, 35-2 Sakae-cho, Itabashi-ku, Tokyo 173, Japan); Suzuki, S. *Gann* 69(4): 599-600; 1978.

Ochratoxin A (Och A), a mycotoxin produced by *Aspergillus* and *Penicillium*, was found to have a carcinogenic effect on the liver and kidney of ddY mice, and this effect was enhanced synergistically by the presence of aflatoxin B₁ (AFB₁). Mice were divided into six groups and given: 1) AFB₁ (20 mg/kg by gastric intubation), 2) Och A (40 ppm of diet), 3) AFB₁ + Och A (40 ppm of diet), 4) DMSO (0.1 mg/20 g gastric intubation), 5) Och A + DMSO, 6) AFB₁ + DMSO, no treatment. Och A produced 1 hyperplastic nodule in the liver, 5 hepatic cell tumors, 4 atypical kidney tubules, 9 cystic adenomas, and 2 renal cell tumors among 20 mice. AFB₁ produced 1 hyperplastic nodule, 2 hepatic cell tumors and 1 cystic adenoma among 18 mice. Och A + AFB₁ produced 8 hyperplastic nodules, 15 hepatic cell tumors, 11 atypical tubules, 19 cystic adenomas, and 2 renal cell tumors among 20 mice. The renal cell tumor appeared to be of epithelial cell origin, corresponding to Grawitz tumors in humans; the hepatic tumors were well-differentiated trabecular adenomas. (2 refs)

- 78-5469 **Action of Dihydroaflatoxins on Rat Liver Mitochondria. I. Membrane-dependent Osmotic and**

Effects of Aflatoxins B₂ and G₂. (Eng) Obidoa, O. t. Biochemistry, Univ. Nigeria, Nsukka, Nigeria); Babani, E. A.; Bassir, O. *Toxicology* 11(1): 29-36; 1978.

many fungal-coumarin metabolites can inhibit rat liver chondria, it is suggested that aflatoxin B₂ (AFB₂) and oxin G₂ (AFG₂) possess similar activity that may partial-count for their toxicity. Therefore, the membrane-dependent osmotic and redox effects of the two dihydroaflatoxin isolated rat liver mitochondria were investigated. At thal yet carcinogenic doses (1-2 x 10⁻⁶ M). Both comds had no deleterious effect on the permeability of the chondrial membrane. They did not cause any marked ling, inhibit any ATP-induced contraction or stimulate chondrial ATPase activity. These findings are signifi- because reversible changes in the permeability of mito- drial membranes are important in the regulation of cel- metabolic activity. The toxins were capable of inhibiting ron transport terminally, between cytochrome c and en (cytochrome oxidase). The significance of these find- in relation to the differential toxicity of the aflatoxins scussed. (23 refs)

5470 Influence of Dietary Protein and Vitamin B₁₂ on the Toxicity and Carcinogenicity of Aflatoxin in Rat Liver. (Eng) Temcharoen, P. (Dept. Pathobiology, ulty Science, Ramathibodi Hosp., Mahidol Univ., Bang- , Thailand); Anukarahanonta, T.; Bhamarapravati, N. *Cancer Res* 38(7): 2185-2190; 1978.

influence of dietary protein content and dietary vitamin supplement on the hepatotoxicity and carcinogenicity of toxins (a mixture of aflatoxins B₁ and G₁, with trace ounts of B₂ and G₂) in rat liver was studied. In rats fed w-protein diet (5% casein), the aflatoxin mixture (1 ppm) he diet induced extensive toxic and carcinogenic effects. rhosis was significantly prevented by vitamin B₁₂ adminis- ion (5 µg/100 g diet), but the incidence of cholangiofibro- and hyperplastic nodules was unchanged. No toxic effect s observed in rats receiving a high-protein diet (20% ca-) with no vitamin B₁₂ supplement for the 33 wk duration study. Only one rat bearing a hepatoma was observed ong 25 rats receiving the high-protein diet with no vitamin supplement, whereas 6 hepatomas and 4 hyperplastic ules were seen among 25 rats receiving the high-protein t plus vitamin B₁₂. Cholangiofibrosis and cirrhosis were berved in the high-protein groups, regardless of vitamin administration. The above observations and those of oth- suggest that vitamin B₁₂ may potentiate some chemical cinogens in the induction of biochemical derangements d the initiation of tumor formation. (63 refs)

5471 A Diploid Rat Liver Cell Culture. IV. Malignant Transformation by Aflatoxin B₁. (Eng)

Schaeffer, W. I. (Dept. Medical Microbiology, Coll. Medi- cine, Univ. Vermont, Given Medical Building, Burlington, VT, 05401); Heintz, N. H. *In Vitro* 14(5): 418-427; 1978.

Cloned rat hepatocyte cultures (RL-PR-C) were chronically exposed to a subtoxic, sublethal dose of aflatoxin B₁ (AFB₁): 0.5 µg/ml AFB₁ was administered to the cultures at each weekly passage beginning at passage 12. Continuous exposure to AFB₁ increased the tumorigenic potential of the RL-PR-C cells, as tested by back-injection into isogenic animals. Control cultures exhibited spontaneous transformation at passage 56, about 20 passages beyond the chemically induced event. Neither AFB₁-treated nor control cultures exhibited cytopathological morphology, formation of cell foci, growth in soft agar, or irregular fibroblastlike growth patterns that could be specifically related to the onset of tumorigenic potential. Karyotypic analysis revealed no specific chromosomal aberrations associated with AFB₁ treatment; however, chromosomal instability was a property of the tumorigenic cell populations. Injection of both AFB₁-treated and control cultures at passage 56 into isogenic newborn Wistar/Lewis rats resulted in tumors indicative of both carcinoma and sarcoma, demonstrating the multipotency of these epithelial cells transformed in vitro. The microenvironment at the location of the injected cells might dictate the ability of the cells to form tumors and may affect the morphology of the resultant tumors. The results suggest that it will be necessary to examine biochemical rather than morphological differences when monitoring epithelial cells for transformational events. (19 refs)

78-5472 The Major Metabolite of Aflatoxin B₁ in the Rat Is a Glutathione Conjugate. (Eng) Degen, G. H. (Institut für Pharmakologie und Toxikologie, Universität Würzburg, Verbacher Landstrasse 9, 8700 Würzburg, W. Germany); Neumann, H. G. *Chem Biol Interact* 22(2/3): 239-255; 1978.

The water-soluble, nonextractable metabolites of aflatoxin B₁ (AFB₁) were characterized in vivo and in vitro. When ¹⁴C-AFB₁ was injected ip into female Wistar rats (0.5 or 2.5 mg/kg), 60%-65% of the dose was eliminated into the bile, mostly as polar, non-extractable metabolites. A glutathione conjugate was the major metabolite produced. This conjugate was also formed when a rat liver postmitochondrial supernatant was incubated with AFB₁ and ³H-glutathione. The conjugate was purified by ion-exchange, gel-filtration, and thin-layer chromatography. It was tentatively identified as 2,3-dihydro-2-(S-glutathionyl)-3-hydroxy aflatoxin B₁. This structure was derived mainly from amino acid analysis, UV spectra, and the enzymatic requirements for its formation in vitro. In the rat, this detoxification product of the potentially ultimate reactive AFB₁ epoxide constitutes about 10% of the administered dose and, thus, underlines the quantitative importance of this activation pathway. The toxicity of AFB₁ should largely depend on the relative rates of the epoxidation and glutathione conjugation reactions, as these rates deter-

mine the degree of exposure of tissue to the ultimate reactive form. (36 refs)

- 78-5473 Dietary Lipotropes, Hepatic Microsomal Mixed-Function Oxidase Activities, and In Vivo Covalent Binding of Aflatoxin B₁ in Rats.** (Eng) Campbell, T. C. (Div. Nutritional Sciences, Cornell Univ., Ithaca, NY, 14853); Hayes, J. R.; Newberne, P. M. *Cancer Res* 38(12): 4569-4573; 1978.

Studies were carried out to determine whether the increased susceptibility of rats to aflatoxin B₁ (AFB₁) hepatocarcinogenesis when they were fed a high-fat diet deficient in lipotropes could be related to the mixed-function oxidase (MFO) metabolism of AFB₁. Weanling male Sprague-Dawley rats were fed a nutritionally complete synthetic diet or a diet marginally deficient in choline and methionine and lacking folic acid for 97-100 days. Hepatic microsomal MFO activities, as assayed by ethylmorphine N-demethylation, ethoxycoumarin O-dealkylation, and cytochrome c reduction, were depressed to a similar extent (21%-39%) by the lipotrope-deficient (LD) diet, when expressed on a microsomal protein basis. In addition, cytochrome P-450 content was reduced by 37%, and AFB₁ metabolism was lowered slightly. The proportion of an ip dose of AFB₁ (1 mg/kg) that became covalently bonded to DNA and RNA was reduced to a similar extent by the LD diet 6 hr after injection, but the formation of AFB₁-protein adducts was not affected. The depression of DNA and RNA adduct formation in the animals that received the LD diet was probably related to the lower MFO activities and not to an alteration of glutathione levels, which remained unchanged by dietary treatment. These results suggest that the marginally LD diet does not enhance tumor formation through an increased microsomal activation of AFB₁. (37 refs)

- 78-5474 Binding of Aflatoxin to DNA as Studied by Optical Detection of Magnetic Resonance.** (Eng) Kwiram, A. L. (Dept. Chemistry, Univ. Washington, Seattle, WA, 98195); Liu, Y. C.; Farquhar, M. N.; Smuckler, E. A. *Biochem Biophys Res Commun* 83(4): 1354-1359; 1978.

The binding of aflatoxin B₁ (AFB₁) to intact DNA was studied by optical detection of magnetic resonance (ODMR) using both in vitro and in vivo samples. For the in vivo experiments, rats were injected with 10 µg/kg AFB₁ bid for 2 days, and DNA was subsequently extracted from the liver. The in vitro studies used DNA-AFB complexes formed in microsomal preparations. ODMR measurements were carried out for free AFB₁ and AFB₂, for the reaction-DNA sample, and for solutions of native DNA to which either AFB₁ (designated DNA/AFB₁) or AFB₂ (DNA/AFB₂) had been added. The wavelength-dependent data showed that there is a small but distinct difference between AFB₁ and AFB₂. The results for

DNA/AFB₁ and DNA/AFB₂ were essentially the same as those for free AFB₁ and AFB₂. However, the reaction-DNA data coincided with the data for AFB₂, which was the more for covalently bound AFB₁. The small shift in ODMR transition frequencies is consistent with the view that binding occurs at the furan ring, since derivatization in the aromatic region of the molecule or at the keto groups would have a substantially larger effect than that observed. The difference in the data for AFB₂/DNA and the reaction-DNA sample supports the view that aflatoxin is covalently bound and not merely a physically bound AFB₂ metabolite. (20 refs)

- 78-5475 The Induction of Sister-Chromatid Exchanges in Chinese Hamster Ovary Cells by Prolonged Exposure to 2-Acetylaminofluorene and S-9 Mix.** (Eng) Takehisa, S. (Lab. Radiobiology, Univ. California, San Francisco, CA, 94143); Wolff, S. *Mutat Res* 58(1): 103-106; 1978.

When the time of treatment of Chinese hamster ovary cells with 2-acetylaminofluorene plus S-9 mix (rat liver microsomes supplemented with an NADPH-generating system) was extended from 0.5 to 2.5 hr, the yield of sister chromatid exchanges was approx doubled. (19 refs)

- 78-5476 The Effects of Discontinuing Administration of High Levels of 2-Acetylaminofluorene on the Transitional Epithelium of the Mouse Urinary Bladder.** (Eng) Frith, C. H. (Pathology Services Project, Natl. Cancer Toxicological Res., Jefferson, AR, 72079); Rule, J. E. *J. Environ Pathol Toxicol* 1(5): 581-585; 1978.

The effects of discontinuing the administration of a high level of 2-acetylaminofluorene (2-AAF) on the transitional epithelium of the urinary bladder were studied in BALB/c mice. A group of 288 mice was given 500 ppm 2-AAF in the diet for 6 wk, and subgroups of 96 were killed at 6, 13, and 26 wk. Another group of 288 mice was given the same dose of 2-AAF for 13 wk, and subgroups were killed at 13, 26, and 39 wk. Hyperplasia of the transitional epithelium was present in 99% and nodular downgrowths of transitional epithelium into the lamina propria were present in 22% of the animals sacrificed at 6 wk. Seven weeks after withdrawal of the carcinogen, the incidence of both hyperplasia and nodular downgrowths had decreased to 4%. Twenty weeks after carcinogen removal, hyperplasia was present in 5% of the animals, but nodular downgrowths were absent. A similar pattern was seen in animals treated for 13 wk, but the urothelial hyperplasia regressed at a slower rate. Four mice sacrificed at 26 wk had small transitional cell carcinomas. At 39 wk, 26% of the animals and transitional cell carcinomas were found in 2%. The results suggest that exposure to a high level of carcinogen even for a relatively brief period may result

development of carcinomas after the cessation of exposure (10 refs)

5477 **DNA Repair in Primary Cultures of Rat Hepatocytes.** (Eng) Yager, J. D. (Dept. Pathology, Dartmouth Medical Sch., Hanover, NH, 03755); Miller, J. *Cancer Res* 38(12): 4385-4394; 1978.

A repair was analyzed quantitatively in cultured Wistar hepatocytes following exposure to UV light, 2-acetylaminofluorene (AAF), and its derivatives N-hydroxy-2-acetylaminofluorene (N-OH-AAF) and N-acetyl-2-acetylaminofluorene (N-Ac-AAF). The nuclear thymidine (^3H -dThd) labeling index of control cultures was less than 0.1%, but significant cytoplasmic labeling was present in autoradiographs. Of the total acid-precipitable radioactivity present in control cultures, 54% of the ^3H was bound in protein, demonstrating the ability of these nonreducing hepatocytes to catabolize ^3H -dThd and reutilize the released metabolites. UV light irradiation of the cultured hepatocytes resulted in a dose-dependent increase of ^3H -dThd incorporation into DNA. That this represented nuclear DNA repair synthesis was demonstrated by detecting nonsemiconservative DNA synthesis (repair replication) with NaI isonic centrifugation and autoradiography. Hydroxyurea (25 and 100 mM) had only a small inhibitory effect, but 1- α -arabinofuranosylcytosine (25 and 100 μM) and ethidium bromide (25 μM) dramatically inhibited the UV-induced increase in ^3H -dThd incorporation. Repair synthesis also occurred in response to treatment with AAF, demonstrating the ability of the hepatocytes to metabolize this prohepatocarcinogen to a form capable of damaging DNA. N-OH-AAF and N-Ac-AAF were more effective in inducing a repair response. These results demonstrate the usefulness of primary hepatocyte cultures for studies of carcinogenesis and as a screening system for environmental chemicals suspected of damaging DNA and causing cancer. (49 refs)

5478 **Effect of 3-Methylcholanthrene Pretreatment on 2-Acetylaminofluorene N- and Ring-Hydroxylation by Rat and Hamster Liver Microsomes.** (Eng) Siklikar, P. D. (Fels Res. Inst., Temple Univ. Med. Sch., Philadelphia, PA, 19140); Hong, Y. S.; Baldy, W. J. *Toxicol* 2(3): 135-139; 1978.

The effect of pretreatment with 3-methylcholanthrene (MC) prior to sacrifice of rats and hamsters on the ability of liver microsomal preparations to induce N- and ring-hydroxylation of 2-acetylaminofluorene (AAF) was investigated. Hydroxylation of AAF has been established as an activation step in the latter compound's carcinogenic effect, and ring-hydroxylation is considered to be an inactivation step. MC, 100 mg/kg body wt, was injected ip into male Sprague-Dawley rats and male Syrian golden hamsters 24 hr before

death. Liver microsome fractions were incubated with either 2.3 or 100 μM AAF, and results were expressed in nanomoles of hydroxy-AAF formed/30 min per nanomol of cytochrome P-450 or P-448. Rats pretreated with MC had a significant ($p < 0.05$) increase in both N- and ring-hydroxylation for low and high concentrations of AAF. Pretreatment of hamster with MC induced a specific and large increase ($p < 0.05$) in N-hydroxylation only. In rats, it has been shown that simultaneous administration of MC and AAF inhibits liver tumor induction. Liver carcinogenicity by AAF in hamsters is not inhibited by treatment with MC. The in vitro results of this study are discussed in relationship to these in vivo studies. (21 refs)

78-5479 **In Vitro Binding of 2-Acetylaminofluorene to RNA and Protein with Rat Liver Microsomes.** (Eng) Hara, E. (Biochemistry Div., Saitama Cancer Center Res. Inst., Ina-Machi, Kitaadachi-gun, Saitama-ken 362, Japan); Yonekawa, H.; Kawajiri, K.; Masuda, A.; Tagashira, Y. *Gann* 69(4): 533-538; 1978.

The binding of 2-acetylaminofluorene (AAF) with exogenous Torula yeast RNA and endogenous protein was studied in vitro in a Sprague-Dawley rat liver microsome system. The covalent binding of ^{14}C -labeled AAF with RNA and microsomal protein depended on the concentration of AAF or microsomes and increased linearly with time up to 45 min. AAF bound to protein 100 times more readily than to RNA. The enhancement of binding by methylcholanthrene treatment, its inhibition by 7,8-benzoflavone, and the requirement for NADPH suggested that mixed-function oxidases were involved in the reaction. The binding of AAF was not due to cytosol contamination in the microsomal fraction. In fact, addition of cytosol or the cofactors ATP and sulfate ion suppressed the binding to both RNA and protein. N-Hydroxy-2-acetylaminofluorene, a mixed-function metabolite of AAF, contributed only 1.74% of the binding to RNA and 0.39% of the binding to microsomal protein. A ring-hydroxylated metabolite may be responsible for most of the binding. (21 refs)

78-5480 **Use of Intact Liver Cells for Metabolic Activation in a Simplified Fluctuation Test for Detection of Carcinogens and Mutagens (Meeting Abstract).** (Eng) Hubbard, S. A. (Dept. Biochemistry, Univ. Surrey, Surrey, England); Bridges, J. W.; Green, M. H.; Bridges, B. A. *Mutat Res* 53(2): 202; 1978. (1 ref)

78-5481 **Enhancing Effect of Chemicals on Production of Hyperplastic Liver Nodules Induced by N-2-Fluorenylacetamide in Hepatectomized Rats.** (Eng) Ito,

N. (First Dept. Pathology, Nagoya City Univ. Medical Sch., 1 Kawasumi, Mizuho-cho, Mizuho-ku, Nagoya 467, Japan); Tatematsu, M.; Hirose, M.; Nakanishi, K.; Murasaki, G. *Gann* 69(1): 143-144; 1978.

A comparison was made of the effects of various inducers of drug-metabolizing enzymes on the production of hyperplastic nodules in the livers of partially hepatectomized rats that had been treated with N-2-fluorenylacetamide (2-FAA). Male Fischer rats were maintained for 2 wk on a diet containing 200 ppm 2-FAA. Some animals were then maintained for 8 wk on diets containing 1,000 ppm polychlorinated biphenyl (PCB), 500 ppm phenobarbital, 2,500 ppm 3-(3,5-dichlorophenyl)-5,5-dimethylloxazoline-2,4-dione (DDOD), or 500 ppm phenobarbital plus 500 ppm β -naphthoflavone; controls were maintained on the 2-FAA diet for these 8 wk. The incidence of hyperplastic nodules in the livers of animals partially hepatectomized during the third week was highest in the group fed PCB, decreasing sequentially in the animals fed phenobarbital, phenobarbital plus β -naphthoflavone, and DDOD. In all cases, the incidence was significantly higher than that in animals fed 2-FAA only. The hyperplastic nodules were generally < 1.5 mm in diameter in all groups. This system may prove to be a valid in vivo short-term screening test for hepatocarcinogens. (11 refs)

78-5482 Effects of 2-Acetylaminofluorene on Diurnal Oscillations in Nucleic Acid Metabolism in Two Classes of Nuclei Isolated from Rat Liver. (Eng) Bushnell, D. E. (Dept. Pathology, Veterans Admin. Hosp., 150 Muir Drive, Martinez, CA, 94553); Jeter, J. R. *Cancer Res* 38(8): 2533-2538; 1978.

The effects of 2-acetylaminofluorene (AAF) on the diurnal fluctuations in nuclear nucleic acid labeling from pyrimidine precursors were investigated. ARS/Sprague-Dawley rats were fed a 60% protein diet with or without 0.028% AAF for 5 wk, on an 8 + 16 feeding schedule (AAF was given only during the first 8 hr of a 12-hr dark period). Liver nuclei were pulse-labeled (P-L) for 2 hr with ^3H -thymidine and ^{14}C -orotic acid three times during the feeding-lighting schedule, and isotope incorporation into DNA and RNA was determined. To do this, nuclei were fractionated into two classes, N-1 and N-2. In control livers, hepatocyte nuclei were found primarily in the N-1 class, stromal nuclei in the N-2 class. P-L of control liver nuclear DNA (nDNA) revealed that labeling peaked 2-4 hr after the feeding period. In treated livers, the N-2 population increased significantly during AAF feeding, apparently as the result of a shift of hepatocyte nuclei in AAF-fed rats reached a higher level than that in controls; however, the labeling did not peak, and there was a large variation in labeling from animal to animal at individual pulse periods. In controls, P-L of nRNA peaked 4-6 hr after the start of feeding, but RNA labeling in both nuclei classes from AAF-fed animals did not peak during the times examined. In addition, there was little variation among

individual rats. The results suggest that feeding AAF to rats alters nuclear nucleic acid metabolism in most liver cells. Changes in liver enzyme composition may be related to alterations in liver nRNA metabolism in AAF-fed animals. Presumably, alteration of nuclear structure and function could explain the changes in the regulation of gene expression found in hepatomas and precancerous liver. (32 refs)

78-5483 Site of Origin of 2,7-FAA-induced Rat Leukemia. (Eng) Takayama, S. (Dept. Pathology, Faculty Medicine, Tokyo Medical and Dental Univ., Yushima 1-5-45, Bunkyo-ku, Tokyo 113, Japan); Fujiwara, M. *Acta Pathol Jpn* 28(4): 531-539; 1978.

The primary site of mature granulocytic leukemia (MGL) was compared with that of erythroblastic leukemia (EL) in Wistar and Sprague-Dawley rats, in which leukemia was induced by N,N'-2,7-fluorenylenebisacetamide (2,7-FAA) (0.025% of diet). The appearance of orthochromatic erythroblasts in the peripheral blood was used as an index of the initial stage of leukemia. Most animals were sacrificed when a large number of erythroblasts appeared. In addition, almost all animals were examined histologically. Nodular foci of solid lesions and conspicuously large foci of scattered lesions were considered to be the primary site of leukemia. Leukemia was induced in 151/356 Wistar rats and 27/74 Sprague-Dawley rats, with the main types being MGL or EL. The foci appeared mainly in the bone marrow of various bones. Most of the MGL originated in the short bones, but most of the EL originated in the long bones. Two cases of EL also originated in the spleen. The difference in primary site between MGL and EL was significant ($p < 0.001$). The short bones have abundant bony trabeculae in the shaft; the long bones have scanty bone trabeculae. Thus, the origin of MGL seems to be connected with bone tissue, but that of EL seems to have little connection with this tissue. Morphologically, the MGL resembled human chronic myelogenous leukemia. (19 refs)

78-5484 Hematogenous Development of Rat Mature Granulocytic Leukemia. Comparison of 2,7-FAA-induced and Transplanted Leukemia. (Eng) Takayama, S. (Dept. Pathology, Faculty Medicine, Tokyo Medical and Dental Univ., Yushima 1-5-45, Bunkyo-ku, Tokyo 113, Japan); Fujiwara, M. *Acta Pathol Jpn* 28(4): 523-529; 1978.

In order to prove the hypothesis that mature granulocytic leukemia (MGL) is spread by hematogenous metastasis from nodular foci in the bone marrow to various organs of the body, the degree of leukemic cell infiltration of N, N'-2,7-fluorenylenebisacetamide (2,7-FAA)-induced MGL was compared with that of transplanted MGL in Wistar rats. Transplantation of leukemia exemplified hematogenous metastasis. Group 1 rats were fed 2,7-FAA (0.025% of diet)

Group 2 rats were inoculated iv with 2×10^5 to 4×10^5 from eight Group 1 rats with MGL. MGL was induced 63/503 Group 1 rats and 67/318 Group 2 rats. (Morphologically, the MGL resembled human chronic myelogenous leukemia.) Hematologically and histologically, the leukemias in the two groups were similar. In both, the degree of leukemia cell infiltration was similar, being high in the bone marrow, medium in the spleen, and low in the liver. The similarity of infiltration between 2,7-FAA-induced and transplanted MGL suggests that the former is spread by hematogenous metastasis from the primary nodular foci of the bone marrow to various parts of the bone marrow, spleen, and other organs. (14 refs)

78-5485 Association of an Angiosarcoma of the Liver with a Hepatoma in a Vinyl Chloride Worker. (Eng) Delorme, F. (Centre Hospitalier Regional de la Mauricie, C.P. 1130, Shawinigan-Sud, Quebec, Canada). *Ann Anat Pathol (Paris)* 23(2): 105-113; 1978.

A 51-yr-old man who had been exposed occupationally to vinyl chloride vapors for 23 yr, 4 mo died in a coma with liver failure. Autopsy revealed the association of a hepatic angiosarcoma with a hepatoma. Alcohol consumption by the patient was negligible. The association of the two tumors is the first ever to be reported in the literature. (22 refs)

78-5486 Ten Cases of Angiosarcoma of the Liver in Vinyl Chloride Workers in Canada. (Fre) Delorme, F. (Centre Hospitalier Regional de la Mauricie, C.P. 1130, Shawinigan-Sud, Quebec, Canada). *Ann Anat Pathol (Paris)* 23(2): 97-104; 1978.

Ten cases of hepatic angiosarcoma were found among employees of a vinyl chloride plant in Shawinigan, Canada. All patients had been exposed to vinyl chloride vapors. The average age of the patients was 50 yr (41-61 yr); the length of occupational exposure to vinyl chloride averaged 16 yr 10 mo (range from 5 yr, 8 mo to 26 yr); and the time lapse between last exposure and diagnosis was 20.5 yr (11-18 yr). Survival after diagnosis averaged 4.4 mo (1-8 mo). The risk of hepatic angiosarcoma appears to be highest among vat cleaners, pipe fitters, and polymerization workers. (16 refs)

78-5487 Lack of Mutagenicity of Extracts from Polyethylene (PE) in Strains of *Salmonella typhimurium* (Meeting Abstract). (Eng) Fevolden, S. E. (Central Inst. Industrial Res., Blindern, Oslo 3, Norway); Moller, E. *Mutat Res* 53(2): 186-187; 1978. (no refs)

78-5488 External Chromosome Studies Undertaken on Persons and Animals with VC Illness (Meeting Abstract). (Eng) Fleig, I. (Occupational Medicine and Health Protection, BASF, Aktiengesellschaft, 6700 Ludwigshafen, W. Germany); Thiess, A. M. *Mutat Res* 53(2): 187; 1978. (no refs)

78-5489 Cytogenetic Investigations on Lymphocytes from Workers Exposed to Vinyl Chloride (Meeting Abstract). (Eng) Leonard, A. (Mammalian Genetics Lab., C.E.N.-S.C.K., B-2400 Mol, Belgium); Decat, G.; Leonard, E. D. *Mutat Res* 53(2): 219; 1978. (no refs)

78-5490 The Metabolic Activation of Vinyl Chloride In Vitro (Meeting Abstract). (Eng) Ivanetich, K. M. (Dept. Physiology and Medical Biochemistry, Univ. Cape Town Medical Sch., Cape Town, South Africa); Katz, I. D.; Aronson, I. *Mutat Res* 53(2): 204; 1978. (no refs)

78-5491 External Chromosome Investigations of a Person with an Angiosarcoma due to VC Exposure (Meeting Abstract). (Eng) Fleig, I. (Arbeitsmedizin und Gesundheitsschutz, BASF, Aktiengesellschaft, 6700 Ludwigshafen, W. Germany); Thiess, A. M. *Mutat Res* 53(2): 187-188; 1978. (no refs)

78-5492 Detection of Electrophilic Metabolites of Halogenated Olefins with 4-(4-nitrobenzyl)pyridine (NBP) or with *Salmonella typhimurium* (Meeting Abstract). (Eng) Barbin, A. (Unit Chemical Carcinogenesis, International Agency for Res. on Cancer, 150 cours Albert Thomas, 69008 Lyon, France); Planche, G.; Croisy, A.; Malaveille, C.; Bartsch, H. *Mutat Res* 53(2): 150; 1978. (no refs)

78-5493 Lung Tumorigenesis by 1,2-Diformylhydrazine in Mice. (Eng) Toth, B. (Eppley Inst. Res. Cancer, Univ. Nebraska Medical Center, Omaha, NB, 68105); Erickson, J. *Z Krebsforsch* 92(1): 11-16; 1978.

Beginning at 6 wk of age, randomly bred Swiss albino mice (50 males and 50 females) were administered 1,2-diformylhydrazine (DFH: 2% soln in drinking water) over their entire life-span in order to determine the relative carcinogenicity of

mono- and dialkylhydrazines and also their possible environmental significance as carcinogens. The av daily intake of DFH was 144 mg for females and 146 mg for males. Of the females, 48 developed 467 lung tumors: 19 had 113 adenomas and 29 had 307 adenomas and 47 adenocarcinomas. Of the males, 41 developed 376 lung tumors: 26 had 154 adenomas and 15 had 189 adenomas and 33 adenocarcinomas. The average age at death of both males and females was 66 wk. In contrast, only 15/100 female controls developed 17 lung tumors and only 22/100 male controls developed 25 lung tumors. Although a few instances of other types of tumors were observed in various groups, their low incidences suggest that their appearance cannot be attributed to the treatment. (20 refs)

78-5494 Strain Susceptibility and Resistance to 1,2-Dimethylhydrazine-induced Enteric Tumors in Germfree Rats. (Eng) Asano, T. (Lobund Lab., Dept. Microbiology, Univ. Notre Dame, Notre Dame, IN, 46556); Pollard, M. *Proc Soc Exp Biol Med* 158(1): 89-91; 1978.

Three strains of germfree rats (Sprague-Dawley, Lobund-Wistar, and Buffalo) were tested for their susceptibility to 1,2-dimethylhydrazine (DMH)-induced enteric neoplasms. At 20 wk after the start of DMH administration (30 mg/kg po 1x/wk for 10 wk), the rats were killed and examined for tumors. Sprague-Dawley rats showed the most susceptibility to DMH carcinogenesis, Buffalo rats were intermediate, and Lobund-Wistar rats were not susceptible. The results demonstrate that intestinal microflora are not essential to the induction of intestinal tumors by DMH in Sprague-Dawley and Buffalo strain rats and that there is a strain dependent differential susceptibility to DMH-induced enteric tumors. (11 refs)

78-5495 Transplantation of Uterine Sarcoma Induced by 1,2-Dimethylhydrazine in CBA Mice. (Rus) Bazzlova, L. S. (Lab. Carcinogenic Substances, Oncological Res. Center, Acad. Medical Sciences USSR, Moscow, USSR). *Vopr Onkol* 24(9): 80-82; 1978.

The transplantability of 1,2-dimethylhydrazine (DMH)-induced uterine sarcomas was studied in nine consecutive generations of CBA mice. The primary tumor was induced in the mice by sc administration of DMH (8 mg/kg/wk) for 30-35 wk, after which time the animals were sacrificed. In addition, the primary tumor tissue homogenate was transplanted sc into 2-mo-old female, castrated and old female, and male CBA mice. The tumor take rate was 100% in all nine generations. The latent period averaged 33-73 days in generation 1, 7-12 days in generation 3, and 4-5 days in generation 9. Survival from the date of transplantation averaged 63-108 days in generation 6, 23-38 days in generation 3, and 19-50 days in generation 9. The tumor growth dynamics were the same in young mice as in old and castrated females, and

in males. The transplanted tumors retained the histological characteristics of the primary tumor. The findings indicate a reduction of the latent period and survival over the consecutive generations. (3 refs)

78-5496 Ultrastructure of 1,2-Dimethylhydrazine-induced Adenocarcinomas in Rat Colon. (Eng) Barkla, D. H. (Dept. Anatomy, Monash Univ., Clayton Victoria 3168, Australia); Tutton, P. J. *J Natl Cancer Inst* 1291-1299; 1978.

The ultrastructure of 1,2-dimethylhydrazine (DMH)-induced adenocarcinomas in the SD rat colon was examined by transmission electron microscopy and compared with that of epithelial cells in the normal colon. The ultrastructure of tumor cells in the S-phase was examined using ³H-thymidine and electron microscope autoradiography. The tumors were induced in eight male SD rats by weekly sc injection of DMH (21 mg/kg) for 20 wk. Morphologically, most tumor cells resembled undifferentiated crypt cells of normal colon epithelium; a minority of tumor cells closely resembled differentiated colonic epithelial cells. Consistent abnormalities in the morphology of the basal lamina at the interface between tumor epithelium and adjacent lamina propria mucosae were observed. Tumor cells showed ultrastructural but not light microscopic evidence of degeneration. Comparisons between tumor and normal epithelial cells did not reveal a single morphological feature characteristic of the former. Autoradiography indicated that tumor cells of widely divergent morphology were capable of thymidine uptake; these tumor cells included cells showing changes usually associated with neoplasia. S-phase tumor cells lacked distinctive ultrastructural features in the autoradiographic studies. The ultrastructural morphology of tumor cells during the cell cycle may be better studied with tumor cells in culture, because of the opportunity to synchronize cell cycles. (29 refs)

78-5497 1,2-Dimethylhydrazine-induced Large Bowel Cancer in Rats. (Jpn) Miyoshi, T. (Third Dept. Surgery, Faculty Medicine, Univ. Tokyo, Bunkyo-ku, Tokyo 113, Japan); Ohhara, T.; Morioka, M. *Jpn J Gastroenterol* 75(3): 284-293; 1978.

1,2-Dimethylhydrazine (DMH)-induced large bowel cancer was studied in 40 rats autopsied 220 days after the start of DMH treatment. Histologically, four types of lesions were found upon serial sectioning of the entire colon: atypical glands, adenoma, noninvasive cancer, and invasive cancer. As many as 86% of the colonic adenocarcinomas originated de novo; the remaining 14% arose from an adenoma. Most of the tumors occurred in the left half of the colon, and there was a tendency toward tumor multiplicity. The tumors included well-differentiated adenocarcinomas, poorly

iated adenocarcinomas, and mucinous carcinomas, and was high incidence of peritoneal dissemination and abdominal lymph node metastasis. (27 refs)

98 **Acute and Chronic Effects of 1,2-Dimethylhydrazine in Mouse Large Bowel.** (Eng) Deschner, (Memorial Sloan Kettering Cancer Center, New York, 0021); Long, F. C. In: *Prevention and Detection of Cancer Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, Vol. 2(Part 1), 2404 pp.; 1319-1328; 1978.

acute and chronic effects of 1,2-dimethylhydrazine (DMH) on the colon were studied in 90 female CF₁ mice that received 20 mg/kg DMH sc 1x/wk for 6 wk. Acute effects were evaluated in 24 mice sacrificed within the first 6 wk; chronic effects were evaluated in the remaining animals, which were sacrificed at 12-86 wk. Massive cell loss was seen after the first DMH injection, followed by an elevated thymidine labeling index at 48 hr and an increased mitotic frequency at 72 hr. These changes were accompanied by decreased mucus production and deformation of the crypt structure. By days 7 and 8, the histologic and autoradiographic parameters appeared normal. Between 2 and 4 days after the first injection, increased numbers of labeled cells were seen in the upper third of the crypt. This labeling in the crypt zone appeared to occur whenever cells in the middle third of the crypt were more actively involved in DNA synthesis. The loss of regulatory controls that normally repress DNA synthesis in the upper crypt and the increased proliferative activity of the mid crypt were the earliest molecular changes observed in the DMH-treated colonic mucosa. Focal dysplasias were observed after three injections. Focal atypias, adenomas, and carcinomas developed, primarily in the distal colon. In 95% of the mice sacrificed after 20 wk. The colonic tumors were mainly sessile or pedunculated adenomas, adenocarcinomas, and squamous cell carcinomas of the anal canal. This cancer model, involving six injections rather than the previously used 26, more closely mimics the number, distribution, and developmental pattern of colonic tumors in humans. (15 refs)

99 **New Aspects of Intestinal Carcinogenesis by 1,2-Dimethylhydrazine Dihydrochloride (DMH) and the Influence of Antilymphocyte Globuline (ALG) on its Progress.** (Eng) Nomura, K. (Dept. Surgery, Univ. Tokyo Sch. Medicine, Hongo 7-3-1, Bunkyo-ku, Tokyo, Japan); Schlake, W.; Grundmann, E. *Z Krebsforsch* 53: 17-33; 1978.

induction of intestinal tumors by 1,2-dimethylhydrazine dihydrochloride (DMH) was studied in 10-wk-old male Wi-

star rats that were treated with or without antilymphocyte globulin (ALG). A 1% aqueous solution of DMH was administered sc, 1x/wk over 16 wk, at a weekly dose of 20 mg/kg; total dose was about 120 mg/rat. Some rats were also given a 1-mg/0.5-ml aqueous solution of ALG ip, 2x/wk ip over 13 wk, starting 1 wk prior to DMH. During the first week, a double dose was given; total dose was 28 mg/rat. All rats had the following changes in the intestinal mucosa, regardless of treatment: glandular hyperplasia with goblet cell depletion and elongation of the germ cell zone (5-13 wk); glandular atrophy intermingled with hyperplasia (after 13 wk); neoplastic microlesions (17 wk); macroscopic tumors (21 wk). Tumor induction was negative in 5/37 rats at the last autopsy (after 33 wk). The classification of the growth pattern of macroscopic tumors corresponded to their histology. Adenomas or well-differentiated adenocarcinomas formed pedunculated tumors; moderately differentiated adenocarcinomas formed sessile tumors, often with cyst formation; and poorly differentiated or signet ring cell carcinomas formed flat tumors. Most of the tumors were located 7-25 cm from the pylorus, and they were distributed diffusely throughout the large intestine. Tumors were often accompanied by metastases, which were first seen at 29 wk. Neither tumor induction nor tumor growth were accelerated by ALG. However, metastatic dissemination and conglomerate formation occurred more frequently in the ALG + DMH group than in rats receiving only DMH. (37 refs)

78-5500 **Effect of Hydrazine, Isonicotinic Acid Hydrazide, Hydrazine Sulfate, and Dimethylhydrazine on Guanylate Cyclase Activity.** (Eng) Vesely, D. L. (Div. Endocrinology and Metabolism, Dept. Medicine, Univ. Miami Sch. Medicine, Miami, FL, 33155); Rovere, L. E.; Levey, G. S. *Enzyme* 23(5): 289-294; 1978.

Isonicotinic acid hydrazide, hydrazine sulfate, and dimethylhydrazine, three chemical carcinogens structurally related to hydrazine [a potent stimulator of guanylate cyclase (GC)], decreased GC activity. In addition, hydrazine increased DNA synthesis but the other three compounds decreased DNA synthesis. There may be a relationship between the GC-cyclic guanosine monophosphate system and DNA synthesis and chemical carcinogenesis. (41 refs)

78-5501 **The Measurement of PAHs in Industrial Effluents.** (Eng) Jones, P. W. (Battelle, Columbus Labs., Columbus, OH, 43201). *Proc Anal Div Chem Soc* 15(5): 158-159; 1978.

Methods for measuring of polycyclic aromatic hydrocarbons in industrial emissions are outlined. Macroreticular resins are used to collect samples from stack gases and aqueous effluents. Quantitative analysis is accomplished by capillary column gas chromatography-mass spectrometry utilizing selec-

tive ion monitoring. Some of the specificities for these procedures are outlined. (no refs)

- 78-5502 Computer-generated Graphic Models of the N²-substituted Deoxyguanosine Adducts of 2-Acetylaminofluorene and Benzo(a)pyrene and the O⁶-substituted Deoxyguanosine Adduct of 1-Naphthylamine in the DNA Double Helix.** (Eng) Beland, F. A. (Dept. Health, Education, and Welfare, Food and Drug Admin., Natl. Center Toxicological Res., Jefferson, AR, 72079). *Chem Biol Interact* 22(2/3): 329-339; 1978.

Computer models of three deoxyguanosine-carcinogen adducts in double-helical DNA are presented. The 2-acetylaminofluorene adduct was situated in the minor groove, had little freedom of rotation, and caused little helical distortion. The anti-benzo(a)pyrenediol epoxide-N² adduct had a similar fit with a greater degree of steric interaction. The 1-naphthylamine derivative was in the major groove, did not perturb the helix, and had considerable freedom of movement. (31 refs)

- 78-5503 Ca²⁺-dependent ATPase Activity in Primary Cultures of Normal Rat Cells after Exposure to Carcinogens and Their Noncarcinogenic Analogs.** (Rus) Khodosova, I. A. (Lab. Tumor Cell Genetics, Inst. Cytology, Leningrad, USSR); Fridlianskaia, I. I. *Tsitologiya* 20(4): 470-472; 1978.

The effect of the carcinogens dimethylnitrosamine (DMNA), urethane (UT), 1,2,5,6-dibenz(a)anthracene (DBA), and 7,12-dimethylbenz(a)anthracene (DMBA) and their noncarcinogenic analogs on the activity of Ca²⁺-dependent ATPase was studied in primary cultures of rat explant (heterogeneous population of muscle and connective tissue cells). The cell cultures were incubated with the test agents for 20 hr. Ca²⁺-dependent ATPase activity was significantly decreased after exposure to 1,2,5,6-DBA and DMBA (to 1.44 and 1.49 IU, compared with 2.54 IU in controls and 2.75 IU in cultures exposed to the noncarcinogen 1,2,3,4-DBA). UT also inhibited the activity of the enzyme (1.95 IU), but not DMNA. (9 refs)

- 78-5504 The Covalent Binding of Polycyclic Hydrocarbons to DNA in the Skin of Mice of Different Strains.** (Eng) Phillips, D. H. (Chester Beatty Res. Inst., Inst. Cancer Res., Royal Cancer Hosp., Fulham Road, London SW3 6JB, England); Grover, P. L.; Sims, P. *Int J Cancer* 22(4): 487-494; 1978.

The binding of ³H-labeled 7,12-dimethylbenz(a)anthracene (DMBA), benzo(a)pyrene (BP), and 3-methylcholanthrene

(3-MC) to DNA in mouse skin was studied in C57BL/6, DBA/2, and Swiss mice following topical application. C57BL mice were treated with 0.025-1 micromole (g) DMBA, and the levels of hydrocarbon bound to DNA after treatment were determined; there was no evidence of a threshold dose below which no binding to DNA occurred. From 6 hr to 8 days after treatment, DMBA was bound to DNA to a similar extent in Swiss and C57BL mice and to a slightly greater extent in DBA/2 mice. The rate of disappearance of bound DMBA from DNA was similar in all three strains and 3-MC (each, 1 μmol) were bound to DNA to a similar extent at 19 and 48 hr after treatment in all three strains. 0.1 μmol) was bound to DNA in the order DBA/2 > C57BL/6 > Swiss 19 hr after treatment. The levels of binding did not correlate with the reported susceptibilities of the three strains to polycyclic hydrocarbon carcinogenesis. The rapid disappearance of bound DMBA from DNA suggests that the modified bases are detected and excised from DNA by repair enzymes. DNA repair in strains susceptible to hydrocarbon carcinogenesis may be more prone to error than repair in the more resistant strains, and carcinogenesis may result from an alteration of the DNA sequence caused by the incorrect repair of hydrocarbon-modified bases. (15 refs)

- 78-5505 Multiple Regions of Metabolic Activation of Carcinogenic Hydrocarbons.** (Eng) Harvey, G. (Ben May Lab. Cancer Res., Univ. Chicago, Chicago, IL 60637); Dunne, F. B. *Nature* 273(5663): 566-568; 1978.

The carcinogenicities of a series of methyl and fluoro derivatives of benzo(a)pyrene (BP), 7-methylbenz(a)anthracene (MBA), and 7,12-dimethylbenz(a)anthracene (DMBA) were studied in male Long-Evans rats to determine whether the introduction of methyl or fluoro groups into the ring of a polycyclic aromatic hydrocarbon (PAH) could alter the carcinogenic activity. Thus, 7,10-dimethyl-2-methyl-DMBA, 3-methyl-DMBA, 1-fluoro-DMBA, 2-fluoro-DMBA, and 4-methyl-MBA were inactive. 10-methyl-BP was an extremely weak carcinogen, 7-methyl-BP and 4-methyl-DMBA were very potent carcinogens and 4-fluoro-DMBA and 2-fluoro-MBA exhibited diminished but still significant carcinogenic activity. These results suggest that metabolic inactivation by mixed-function oxidases occurs relatively efficiently at substituted positions or, more probably, that it can occur at alternative molecular regions when reaction at preferred sites is blocked. Although clear-cut evidence concerning the structures of the active metabolites of any of these compounds is lacking, the most probable candidates are arene oxides in the substituted ring, K-region arene oxides, or non-bay region diol epoxides. (15 refs)

506 Difference in the Induction of Osteosarcoma in Rabbit Bone with Single Administration of the Kinds of Chemical Carcinogens. (Eng) Sato, K. (Dept. Surgery, Sch. Medicine, Chiba Univ., Inohana 1-8-1, Chiba 280, Japan); Nukaga, H.; Horikoshi, T.; Iwasaki, I. *Int J Cancer* 69(4): 579-583; 1978.

Osteosarcoma (OS) induction was evaluated in mixed breed rabbits treated with a single intramedullary dose (15-20 mg) of nitroquinoline 1-oxide (4-NQO), 3-methylcholanthrene (DMBA), or 7,12-dimethylbenz(a)anthracene (DMBA) in the mandible, diaphysis, or distal metaphysis of the femur. 4-NQO was the most effective carcinogen, with OS incidence in the mandible, diaphysis, and metaphysis being 6/10, 3/10, and 5/8 rabbits, respectively. In addition, there were two fibrosarcomas in the diaphysis and a chondrosarcoma in the metaphysis. Tumor incidence with DMBA was 1/10 and 2/6 rabbits in the mandible and metaphysis, respectively; and 4-NQO produced only one fibrosarcoma in a mandible. Histological examination of each site during the latent period revealed that three types of tissue reaction were common, depending on the degree of necrotic (I), fibroblastic (II), or mixed (III) tissue encountered. The development of each type was related to the dose and nature of the carcinogen. Type I resulted when the effectiveness or dosage of carcinogen was low, and it usually appeared in sites lacking cancellous bone marrow, such as the diaphysis. Type III resulted from the use of a more potent carcinogen and dose, and it usually appeared in bones containing abundant cancellous bone marrow, such as the metaphysis. Type II resulted in the formation of OS's and occasional chondrosarcomas; fibrosarcomas were seldom observed. (8 refs)

507 Transformation of Human Cells in Culture by Chemical Carcinogens. (Eng) Rhim, J. S. (Dept. Cancer Res., Microbiological Associates, Inc., Bethesda, MD, 20814). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 1-5, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2 (Part 2) 2404 pp.; 1337-1357; 1978.

The transformation of human osteosarcoma clonal cells in culture with N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), 7,12-dimethylbenz(a)anthracene (DMBA) was investigated. In cultures treated with MNNG (0.01 or 0.1 $\mu\text{g}/\text{ml}$), morphological alterations of cells and abnormal growth patterns were observed in the sixth subculture. In cultures treated with DMBA (2.5 or 1.0 $\mu\text{g}/\text{ml}$), morphological alterations were seen in the fifth subculture. Five/five mice given 5×10^6 cells sc from cultures treated with 0.01 $\mu\text{g}/\text{ml}$ MNNG developed poorly differentiated sarcomas within 2 wk at the inoculation site. Three/five mice given cells from cultures treated with 0.1 $\mu\text{g}/\text{ml}$

MNNG produced small, persistent nodules resembling osteosarcomas. Three/three mice inoculated sc with 5×10^6 cells from cultures treated with 1.0 $\mu\text{g}/\text{ml}$ DMBA developed tumors at the inoculation site within 3 wk; 0/4 animals inoculated sc with cells treated with 2.5 $\mu\text{g}/\text{ml}$ DMBA developed tumors. However, cells from cultures treated with both DMBA concentrations were tumorigenic when inoculated intracranially. Human cells resembling the original, nontransformed osteosarcoma were recovered from the brain tumors, while sc tumors resembled the DMBA-treated cells. MNNG- and Kirsten sarcoma virus-transformed cells formed larger cell aggregates than did control, normal counterpart cells within 1-3 days when suspended in the liquid growth medium above an agar base. The size and degree of cell proliferation in the aggregates correlated well with tumorigenicity in nude mice. (25 refs)

78-5508 Explant Culture of Rat Colon: A Model System for Studying Metabolism of Chemical Carcinogens. (Eng) Autrup, H. (NCI, Bethesda, MD, 20014); Stoner, G. D.; Jackson, F.; Harris, C. C.; Shamsuddin, A. K.; Barrett, L. A.; Trump, B. F. *In Vitro* 14(10): 869-877; 1978.

An explant culture system was developed for the long-term maintenance of rat colonic tissue. Colonic explants (1 cm^2) from male Charles River CD rats (4-5 wk old) or adult male Fisher F-344 rats were placed in tissue culture dishes with the epithelium facing the gas-medium interface. Two milliliters of CMRL-1066 culture medium supplemented with glucose, hydrocortisone, β -retinyl acetate, and either 2.5% bovine albumin or 5% fetal bovine serum were added to each dish. The explants were incubated at 30 C in an atmosphere of 95% O_2 and 5% CO_2 on a rocker platform (10 cycles/min) so that the explant was submerged half the time. Under these conditions, the normal appearance of the colonic epithelium was maintained for at least 63 days. After 4 days in culture, one of the following carcinogens was added to the medium to investigate the ability of the explant to metabolize carcinogens: benzo(a)pyrene (BP: 1.5 μM), 7,12-dimethylbenz(a)anthracene (DMBA: 3.2 μM), aflatoxin B₁ (AFB₁: 0.5 μM), 1,2-dimethylhydrazine (1,2-DMH: 1.3 mM), methylazoxymethanol acetate (MAM: 0.26 mM), and dimethylnitrosamine (DMN: 135 μM). After 24 hr incubation (8 hr for MAM), the highest binding level was observed with 1,2-DMH [13,526 picomoles (pmol) carcinogen bound/10 mg DNA]. The other carcinogens bound in the following order: DMH (5,000 pmol/10 mg), MAM (1,650), DMN (845), AFB₁ (125), DMBA (5.2), and BP (3.5). Thus, cultured rat colon was able to metabolize all carcinogens tested into chemical species that formed covalent bonds with cellular DNA. (27 refs)

78-5509 Stimulation by Carcinogens and Promoters of Prostaglandin Production by Dog Kidney (MDCK) Cells in Culture. (Eng) Levine, L. (Graduate Dept Biochemistry, Brandeis Univ., Waltham, MA, 02154); Ohu-

chi, K. *Cancer Res* 38(11, part 2): 4142-4146; 1978.

The effect of carcinogens and tumor promoters on prostaglandin (PG) production was studied in dog kidney cells (MDCK) in culture. The most active polycyclic aromatic hydrocarbon carcinogens, benzo(a)pyrene (BP), 7,12-dimethylbenz(a)anthracene, and 3-methylcholanthrene (each, 0.5 $\mu\text{g}/\text{ml}$), stimulated PG production and the deacylation of cellular lipids, as did aflatoxin B₁ at 0.05 and 0.02 $\mu\text{g}/\text{ml}$. Neither the aromatic amines nor the nitrosamines, at 0.5 $\mu\text{g}/\text{ml}$, affected PG production. Purified epidermal growth factor (EGF: 10^{-9} to 10^{-10} M) also stimulated PG production and cellular lipid deacylation. 12-Tetradecanoylphorbol-13-acetate (TPA) and phorbol-12,13-didecanoate (PDD), each 10^{-9} to 10^{-10} M, stimulated PG synthesis, the deacylation of cellular lipids, and morphological changes in the MDCK cells. The increased PG synthesis and morphological changes required at least 24 hr for expression. Cycloheximide inhibited the TPA- and PDD-stimulated changes but hydrocortisone did not. The PG synthesis stimulated by TPA and PDD was reversible. In terms of the two-stage model of skin carcinogenesis, some of the carcinogens, EGF, TPA, and PDD may be acting as promoters. The mechanism of action of the biologically active phorbol diesters (and that of all promoters) may be related to their antidifferentiation properties. It is possible that deacylation of cellular lipids and/or arachidonic acid metabolism via the cyclooxygenase system mediates cellular differentiation. (27 refs)

78-5510 Association and Dissociation of the Ah Locus with the Metabolism of Aflatoxin B₁ by Mouse Liver: Co-segregation of Aflatoxin B₁ Hydroxylase Induction with Aryl Hydrocarbon (Benzo(a)pyrene) Hydroxylase Induction. (Eng) Gurtoo, H. L. (Dept. Experimental Therapeutics, Grace Cancer Drug Center, Roswell Park Memorial Inst., Buffalo, NY, 14263); Dahms, R. P.; Kanter, P.; Vaught, J. B. *J Biol Chem* 253(11): 3952-3961; 1978.

The hepatic microsomal NADPH-mediated metabolism of aflatoxin B₁ (AFB) and its response to induction by phenobarbital (PB) and/or 3-methylcholanthrene (3-MC) were studied in 70 mice representing various inbred strains, recombinant inbred lines, and progenies of various crosses. The metabolism of AFB to aflatoxin B₁-2,3-oxide and to aflatoxin M₁ (AFM) was preferentially catalyzed by cytochrome P-450- and cytochrome P₁-450-dependent pathways, respectively. AFB epoxidation, which was induced by PB in mice of various inbred strains or recombinant inbred lines, was not associated with the Ah locus. 3-MC pretreatment depressed the epoxidation of AFB 10%-70% in all the strains and recombinant inbred lines tested. This pretreatment induced AFB hydroxylase activity, leading to the formation of AFM, in only some inbred strains. The strains of mice inducible for AFB hydroxylase (8/17 strains tested) were also inducible for aryl hydrocarbon hydroxylase (AHH). The induction of AFB hydroxylase and AHH by 3-MC was also examined in vari-

ous hybrid and backcross progenies and in several recombinant inbred lines. The results suggest that the induction of AFB hydroxylase is regulated by a single gene locus with two alleles, one of which is dominant over the other. The results also suggest that the regulation of the induction of AFB hydroxylase and its association with the induction of AHH occur at the level of common regulatory factors and that the same genetic factors regulate the induction of microsomal and nuclear AFB hydroxylase activities. (72 refs)

78-5511 Induction of Sister-Chromatid Exchange in Chinese Hamster Ovary Cells Treated In Vitro with Non-K-Region Dihydrodiols of 7-Methylbenz(a)anthracene and Benzo(a)pyrene. (Eng) Pal, K. (Chester Beatty Inst., Inst. Cancer Res., Royal Cancer Hosp., Fulham Rd., London SW3 6JB, England); Tierney, B.; Grover, P. L.; P. *Mutat Res* 50(3): 367-375; 1978.

Benzo(a)pyrene (BP) and 7-methylbenz(a)anthracene (MBA) and a series of related K-region and non-K-region dihydrodiols were tested for their abilities to induce sister chromatid exchanges (SCE's) in Chinese hamster ovary (CHO) cells in vitro. BP and the related 4,5- and 9,10-dihydrodiols were weakly active in inducing SCE's, the hydrocarbon being somewhat more cytotoxic than either of the diols. However, the 7,8-dihydrodiol was much more active than the other compounds, and it was highly cytotoxic. The 3,4-dihydrodiol of MBA was cytotoxic to CHO cell cultures at high concentrations and was considerably more active in inducing SCE's than the parent compound or the 1,2-, 5,6-, or 8,9-dihydrodiols. The compounds in the MBA series induced more SCE's per metaphase plate than those in the BP series. The results suggest that this system may be useful in studies aimed at determining which dihydrodiols are important in the metabolic activation of the carcinogenic polycyclic hydrocarbons. It is concluded that CHO cells possess some intrinsic ability to metabolize such compounds in the absence of exogenous activation systems. (39 refs)

78-5512 Molecular Orbital Theoretical Prediction of Isomeric Products Formed from Reaction of Arenes with Oxygen: Arene Oxides and Related Metabolites of Polycyclic Aromatic Hydrocarbons. (Eng) Fu, P. P. (Ben May Lab., Chicago, IL, 60637); Harvey, R. G.; Beland, R. *Tetrahedron* 34(7): 857-866; 1978.

The simplified perturbational molecular orbital theory of Woodward and Ruffin was used to predict the isomeric reaction products of arene oxides and related metabolites of polycyclic aromatic hydrocarbons (PAH). This technique provided a good correlation between the predicted and observed structures of the products formed during the isomerization of arene oxides to dihydrodiols, the hydration and nucleophilic addition to arene oxides, and the dehydration of arene dihydrodiols. The method is also applicable to the arene oxides and dihydrodiols of

carcinogenic and noncarcinogenic PAH. Extension of findings to the enzymatic reactions occurring during metabolism of carcinogenic hydrocarbons and to the reactivity of the biologically active arene diolepoxides and aryloxy intermediates suggests that this method may be useful in predicting metabolite structure and structural requirements for carcinogenic and mutagenic activity. (39 refs)

513 **Effect of Light on the Hydrocarbon-DNA Adducts Formed in Hamster Embryo Cells.** (Eng) S. M. W. (Wistar Inst. Anatomy and Biology, 36th St. Philadelphia, PA, 19104). *Int J Cancer* 22(3): 297; 1978.

Hamster embryo cells in culture were used as a model system to determine how exposure to light (fluorescent light, UV) affects the formation and structure of the DNA adducts of 12-dimethylbenz(a)anthracene (DMBA, highly photolabile) and benzo(a)pyrene (BP, less photosensitive). Cells were exposed to FL (Group A) or kept in darkness (Group B) during a 48-hr incubation with 0.5 μ M of hydrocarbon. DNA of the Group B cells was exposed to FL before enzymatic digestion to deoxyribonucleosides. In cells treated with DMBA, the total amount of DMBA-DNA adducts formed per milligram of DNA was less in Group A cells than in Group B cells. No appreciable amounts of hydrocarbon-DNA adducts were formed in Groups A and B cells treated with DMBA endoperoxide, a major photooxidation product of DMBA. The DMBA-deoxyribonucleoside adducts obtained from Group B cells were analyzed by chromatography on Sephadex LH20 columns. Exposure of these adducts to FL converted more than half to photoproducts. DMBA that was exposed to FL prior to degradation contained less than half as many photoproducts and had an elution profile similar to that obtained from DNA isolated from Group A cells. The BP-DNA adducts were identical for the DNA from Groups A and B cells and for the DNA exposed to FL after isolation. These results indicate that exposure to light does not alter the mechanism of binding of hydrocarbons to DNA in hamster embryo cells but that it does affect the formation of DNA adducts of DMBA in a similar manner in cells and isolated DNA. The biological effects of the light-induced formation of DMBA-DNA adducts in cells are not known. It is possible that degradation-resistant DMBA cross-links formed in the DNA, a phenomenon that would greatly increase the biological consequences of the interaction of this hydrocarbon with DNA. (31 refs)

5514 **Comparative Mutagenesis of Test Materials from the Synthetic Fuel Technologies** (Meeting Abstract). (Eng) Epler, J. L. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Larimer, F. W.; Nix, C. E.; Rao, T. K. *Mutat Res* 53(2): 185; 1978. (no refs)

78-5515 **Testing of Carcinogens and Chemotherapeutic Agents as Inducers of Bacteriophage Lambda** (Meeting Abstract). (Eng) Elespuru, R. K. (Molecular Genetics Section, Frederick Cancer Res. Center, Frederick, MD, 21501); Yarmolinsky, M. B. *Mutat Res* 53(2): 184-185; 1978. (1 ref)

78-5516 **A Biological Analysis of Chemical Hepatocarcinogenesis.** (Meeting Abstract). (Eng) Solt, D. B. (Temple Univ., Philadelphia, PA, 19122). *Diss Abstr Int* B 39(4): 1711; 1978. (no refs)

78-5517 **Induction of the Metabolic Activation System in Rat Liver by a Combined Treatment with 5,6-Benzoflavone and Phenobarbital; Comparison with Aroclor 1254, Using the Ames Test** (Meeting Abstract). (Eng) Mattern, I. E. (Medical Biological Lab. TNO, P.O. Box 45, Rijswijk, Netherlands). *Mutat Res* 53(2): 227-228; 1978. (1 ref)

78-5518 **Mutability of Different Genetic Loci in Mamalian Cells by Metabolically Activated Chemical Carcinogens** (Meeting Abstract). (Eng) Huberman, E. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN). *Mutat Res* 53(2): 202-203; 1978. (no refs)

78-5519 **Comparative Survey of Microsomal Activation Systems for Mutagenic Studies of Safrrole** (Meeting Abstract). (Eng) Dorange, J. L. (Station de Technologie des Produits Vegetaux, 21034 Dijon, France); Janiaud, P.; Delaforge, M.; Levi, P.; Padieu, P. *Mutat Res* 53(2): 179-180; 1978. (6 refs)

78-5520 **Inhibition of Chemical Carcinogenesis by *Corynebacterium parvum*.** (Eng) Woodruff, M. (MRC Clinical and Population Cytogenetics Unit, Western General Hosp., Crewe Road, Edinburgh, EH4 2XU, Scotland); Speedy, G. *Proc R Soc Lond [Biol]* 201(1143): 209-215; 1978.

The development of fibrosarcomas in female CBA mice inoculated im with 20-methylcholanthrene (20-MC: 0.5 mg in 0.1 ml tricaprillin) was inhibited significantly by *Corynebacterium parvum* (0.7 mg) administered iv 4 days before the carcinogen and then every 28 days. Similar treatment starting 4 wk after 20-MC injection did not alter the tumor incidence. Based on these results and previous studies of the effect of *C. parvum* on macrophage activity, it is postulated that a transformation event occurring shortly after 20-MC exposure

results in the emergence of a cell population that is liable to destruction by activated macrophages. However, if the cells are not destroyed, they may multiply so rapidly that subsequent treatment is unable to prevent or delay tumor appearance. (27 refs)

78-5521 Induction of Neoplasms in Hamster Tracheal Grafts with 3-Methylcholanthrene-coated Lycra Fibers. (Eng) Mossman, B. T. (Dept. Pathology, Univ. Vermont Coll. Medicine, Burlington, VT, 05401); Craighead, J. E. *Cancer Res* 38(11, part 1): 3717-3722; 1978.

A method was devised to improve the quantitation of carcinogen dosage to the tracheal epithelium of hamsters. Synthetic Lycra spandex fibers were immersed in 0.3-30 mg/ml solns of ¹⁴C-labeled or unlabeled 3-methylcholanthrene (3-MC) for 1 wk and then inserted into the lumina of 104 tracheas removed from young adult Syrian hamsters. Tracheas were grafted sc onto the backs of 22 weanling hamsters (each animal received 3-4 grafts), and the animals were examined at 3-wk intervals for tumors. Tumors appeared in the tracheal epithelium 3-21 mo after implantation, and they included small cell carcinomas, squamous cell carcinomas, noninvasive carcinomas, papillomas, fibrosarcomas, and adenocarcinomas. Most were carcinomas. The neoplasms were induced with 3-MC doses up to 100-fold lower (9-75 µg/fiber) than amounts used by other investigators. The use of tracer quantities of ¹⁴C-3-MC allowed quantitation of the dose to the tracheal mucosa before and at intervals after grafting. A dose-dependent sensitivity of the hamster tracheal epithelium to relatively small quantities of 3MC was demonstrated. (18 refs)

78-5522 Sequential Cytologic Study of the Development of Squamous Cell Carcinoma Induced in Subcutaneously Implanted Bronchial Autograft of Dog. (Eng) Kobayashi, N. (Inst. Pulmonary Cancer Res., Sch. Medicine, Chiba Univ., Chiba, Japan); Kanisawa, M.; Okamoto, T.; Okita, M.; Katsuki, H. *Acta Cytol (Baltimore)* 22(2): 99-104; 1978.

The lumina of canine bronchi implanted in sc tissue were exposed to the carcinogens 3-methylcholanthrene or benzo(a)pyrene (30-90 mg in a saline soln containing 1% carboxymethylcellulose). The exfoliative cytology and the corresponding histologic and ultrastructural changes during the development of squamous cell metaplasia and carcinoma in the autografts were followed in needle biopsy specimens obtained at intervals of 4-22 wk. The cytologic specimens showed a progression from mild atypia of squamous metaplastic cells to moderate and marked atypia before the development of invasive squamous cell carcinoma. The histologic findings corresponded to these cytologic changes. In the premalignant phase, squamous dysplasia and changes resembling carcinoma in situ were also seen. Ultrastructural

changes in the preneoplastic stages included widened cellular space, increasing tonofilaments, enlarged nuclei, deep indentations, and prominent nucleoli with microvilli. It is concluded that exfoliative cytology, reflecting histologic and ultrastructural changes, is a useful tool for comparative study of premalignant lesions in the bronchus and their early neoplastic stages. (21 refs)

78-5523 Carcinogenicity and Metabolic Profile of 3-Methylcholanthrene Oxygenated Derivatives at the 1 and 2 Positions. (Eng) Cavalieri, E. (Eppley Inst. Cancer, Univ. Nebraska Medical Center, 42nd and I Ave., Omaha, NB, 68105); Roth, R.; Althoff, J.; Grant, C.; Patil, K.; Marsh, S.; McLaughlin, D. *Chem Biol Interact* 22(1): 69-81; 1978.

To support the hypothesis that the critical mechanism of activation of 3-methylcholanthrene (3-MC) is one-electron oxidation, its carcinogenicity was compared with that of 1-hydroxy-3-methylcholanthrene (MC-1-OH), 3-methylcholanthrene-1-one (MC-1-one), 2-hydroxy-3-methylcholanthrene (MC-2-OH), 3-methylcholanthrene (MC-2-one), and 3-methylcholanthrylene (MCL) by repeated application on mouse skin. Six groups of 30 7-wk-old Swiss mice were treated with 0.2 micromole of compound in acetone 2x/wk for 20 wk. All mice tested with 3-MC, MC-2-OH, and MC-2-one developed skin neoplasms, but the latter two groups showed later tumor development and lower survival. MCL and MC-1-OH showed a weaker carcinogenicity, and MC-1-one was not carcinogenic. In the MCL group, the percentage of epidermal carcinoma metastases to the lungs was more than twice that of the other groups. The results support the hypothesis that one-electron oxidation for 3-MC, MC-2-OH, and MC-1-one might be the critical mechanism of carcinogenic activation, with C-1 being the binding site to cellular nucleophiles. The carcinogenicity of MC-1-OH could be due to the formation of an ester intermediate, which might be the ultimate intermediate in the reaction with cellular nucleophiles. Finally, the lack of carcinogenicity of MC-1-one may be due to absence of nucleophilic trapping at C-1 via the radical cation pathways as well as to the inability of mouse skin to reduce MC-1-one to MC-1-OH. The results of metabolism studies of the compounds using mouse liver homogenates are also reported. (20 refs)

78-5524 Cyclophosphamide Delays 3-Methylcholanthrene Sarcoma Induction in Mice. (Eng) Hellstrom, I. (Div. Immunology, Fred Hutchinson Cancer Center, Seattle, WA, 98104); Hellstrom, K. E. *J Natl Cancer Inst* 275(5676): 129-130; 1978.

The hypothesis that treatment of mice with cyclophosphamide (CP) during the latency period of tumor induction might prevent or at least delay the appearance of premalignant tumors was investigated. BALB/c mice were divided into

ps, all of which were inoculated im with 0.1 mg methylcholanthrene (3-MC) on day 1. On days 10, 20, 50, 80, Groups 1 and 4 were inoculated with 10^6 heavily irradiated cells from an 3-MC-induced BALB/c sarcoma, Groups 2 and 5 were inoculated with normal BALB/c kidney cells and Groups 3 and 6 received phosphate-buffered saline. A 100 mg/kg dose of CP was injected ip into Groups 4, 5, and 6 at 10-day intervals starting on day -13 and continuing throughout the experiment. Groups 4, 5, and 6 developed tumors more slowly than Groups 1, 2, and 3 (controls). At day 142, all but one of the control mice had at least one tumor, while 40% of the CP-treated mice remained tumor-free. However, the major effect of CP was probably a delay of tumor appearance rather than a prevention of tumor induction, as the appearance of new tumors in the drug-treated group was not leveled off at the time of writing (142 days). The delay in tumor appearance is consistent with the hypothesis that CP inhibits tumor growth by interfering with either a suppressor cell mechanism or an immunostimulatory mechanism. However, it is equally consistent with the nonimmunological interpretations that CP destroys neoplastic cells as they develop or it interferes with carcinogenesis. (14 refs)

78-5525 Effects of Essential Amino Acid Deficiencies on Syngeneic Tumor Immunity and Carcinogenesis in Mice. (Eng) Worthington, B. S. (Sch. Nutritional Sciences, Textiles, Univ. Washington, Seattle, WA, 98195); Syrota, J. A.; Ahmed, S. I. *J Nutr* 108(9): 1402-1411; 1978.

The long-term effect of essential amino acid (AA) deficiencies on methylcholanthrene (MC)-induced carcinogenesis and on immunity to a transplanted MC-induced tumor was studied in inbred female Balb/c AN mice. Mice were fed a standard diet or one containing 25% of the requirement of isoleucine (Ile), leucine (Leu) or phenylalanine-tyrosine (Phe-Tyr) up to 54 wk. Tumors were induced by implanting paraffin pellets containing 5% MC sc. In the Phe-Tyr-deficient groups, tumor incidence was significantly greater than that in controls; Ile or Leu deficiency had no significant effect on incidence. There was generally a deficit in the specific serum-free response corresponding to the dietary restriction. However, tumors from experimental and control mice were similar in AA composition. Mice in the immunity study were inoculated with 5×10^5 tumor cells/mouse; all tumors were excised when the largest tumor in the controls reached 12 mm, 3 days later the animal was challenged with 5×10^5 tumor cells sc. In response to the first inoculation, the Ile-deficient group had larger (20%) and smaller tumors but the Leu-deficient group had more (40%) and larger tumors than controls (32.5%). However, animals in the Phe-Tyr-deficient group had tumors (20%), but they were of the same size as those in controls. There were no significant differences between the controls and AA-deficient groups in response to tumor challenge. The body wt of experimental and control mice did not differ markedly except for the Ile-deficient group, in which there was pronounced wt loss and debilitation. It is concluded that

selected AA acid restriction was of no therapeutic value and appeared detrimental in the case of Phe-Tyr deficiency. (21 refs)

78-5526 Regenerative Proliferation of Mouse Epidermal Cells Following Application of a Carcinogenic Skin Irritant (MCA): Micro-flow Fluorometric DNA Measurements and ^3H -TdR Incorporation Studies of Isolated Basal Cells. (Eng) Clausen, O. P. (Inst. Pathology, Univ. Oslo, Rikshospitalet, Oslo 1, Norway). *Virchows Arch Cell Pathol* 27(3): 205-215; 1978.

Changes in the cell-cycle traverse of basal cells in hairless mouse epidermis were studied after a single application of 20-methylcholanthrene (20-MC: 0.025 ml of a 1% soln in benzene). At different time intervals up to 3 days after 20-MC administration, groups of animals were injected ip with 30 μCi ^3H -thymidine (^3H -TdR) 30 min before they were killed. Single cell suspensions of epidermal basal cells were prepared, and changes in the labeling index, mean counts, and DNA distribution patterns in each cell suspension were studied in microflow fluorometry. After a slight initial depression at 6 hr after 20-MC application, the labeling index reached a peak at 16 hr, followed by a trough from 24 to 36 hr and then a second peak at 40 hr. The mean grain count was below control level at all times, except at 40 hr. Of the cells entering mitosis from 36 to 44 hr after 20-MC treatment, 65%-70% were labeled with ^3H -TdR. The proportion of cells with S-phase DNA content increased gradually starting at 6 hr after 20-MC application, reaching 250% of control values at 16 hr and staying at this level until 40 hr. After a slight transient increase at 6 hr, the proportion of cells with G₂-phase DNA content decreased to below control values until 40 hr, when there was an increase. From 40 to 72 hr, however, the proportion decreased again. These data indicate that there is a marked prolongation of DNA synthesis despite a consistently increased labeling index. They support the theory that a specific injury to DNA is an initial event in the carcinogen-target cell interaction. (29 refs)

78-5527 Aprotinin and Sarcoma Induction by Methylcholanthrene in Rats. (Ita) Laurentaci, G. (Cattedra di Chirurgia Plastica e Ricostruttiva, Università degli Studi di Bari, Bari, Italy); Dioguardi, D.; Pascone, M. *Arch Sci Med (Torino)* 134(3): 275-281; 1978.

The effect of the immunosuppressive agent aprotinin (Trasylo) on the sarcoma-inducing effect of methylcholanthrene was studied in male and female Wistar rats. Ten rats were injected with 0.3 mg% methylcholanthrene (MC) into the proximal segment of one hindlimb (controls). Ten other rats also received injections of aprotinin (20,000 kallikrein-inhibiting units/kg/day) for 2 wk starting on the day of MC injection, at the same site. The latent period of tumor induction

was 103-157 days in the controls, and their survival after tumor appearance was 34-65 days. The latent period was 103-141 days in the aprotinin-treated group, and the survival time was 37-56 days. The findings indicate that there were no significant differences between the two groups; ie, immunosuppression failed to facilitate the growth of MC-induced sarcomas. (27 refs)

78-5528 Studies of Experimental Lung Tumors Induced by Chemical Carcinogens and Inorganic Substances. III. Histopathological Studies of Lung Tumors Induced in Rats by Pertracheal Infusion of 20-Methylcholanthrene Combined with Chromium and Nickel Powder. (Eng) Mukubo, K. (Second Dept. Internal Medicine, Nara Medical Univ., Nara, Japan). *J Nara Med Assoc* 29(1): 321-340; 1978.

Female rats were given a single pertracheal infusion of 20-methylcholanthrene (3-MC: 1 or 5 mg) and/or powdered chromium (10 mg) or powdered nickel (10 mg) via a vinyl tube, and histopathological studies were made 1-12 hr later. The earliest change noted was hyperplasia of the bronchioalveolar epithelium, followed by squamous metaplasia and atypical proliferation of the bronchiolar epithelium (APBE). At either dose of 3-MC, the incidence of APBE was significantly higher in the 3-MC + CR and 3-MC + NI groups than in the 3-MC alone groups. Cornifying squamous cell carcinoma developed in rats of several groups as early as 12 wk after infusion of 3-MC. The carcinoma incidence was 7/12 rats in the 3-MC (5 mg) + Cr group, 3/5 in the 3-MC (5 mg) + Ni group, 3/7 in the 3-MC (5 mg) group, 3/12 in the 3-MC (1 mg) + Cr group, 2/7 in the 3-MC (1 mg) + Ni group, and 1/8 in the 3-MC (1 mg) group. None of the tumors metastasized. In all 3-MC groups, early alveolitis reached max severity 1 wk after infusion and then subsided in subsequent weeks. This inflammatory change and the later fibrosing reactions were more apparent in the groups given 3-MC and metallic powder than in those given 3-MC alone. The results indicate that the action of 3-MC and Cr or Ni was synergistic with respect to the production of lung cancer, as APBE, which is considered a precancerous lesion, was promoted by adding the powdered metals to 3-MC. The only change noted in the Cr and Ni alone groups was hyperplasia of the bronchiolar epithelium in 3/12 and 3/7 rats, respectively. (53 refs)

78-5529 Development of Malignant Tumors after Transplantation of Organ Lung Cultures Treated with 20-Methylcholanthrene. (Bul) Vlasov, V. (Dept. Pathoanatomy, Medical-Biological Inst., Sofia, Bulgaria); Uzunov, P. *Onkologiya* 15(1): 25-29; 1978.

Lung explants derived from 630 A/Sn and BALB/cY mice (1-3 mo old) were cultured in medium 199, treated with 20-methylcholanthrene (4 mg/ml, for 7 days), and then transplanted sc into 126 mice. Of 87 surviving mice, 41 ani-

mals developed 44 malignant tumors from 3 to 12 mo after transplantation. Most of the tumors developed after 4-7 mo. Histologically, there were 11 carcinomas, 9 carcinosarcomas and 24 sarcomas. (9 refs)

78-5530 The Formation of Dihydrodiols by Chemical and Enzymic Oxidation of 3-Methylcholanthrene. (Eng) Tierney, B. (Chester Beatty Res. Inst., Inst. Cancer Res., Royal Cancer Hosp., Fulham Road, London SW3 6JJ, England); Hewer, A.; Rattle, H.; Grover, P. L.; Simmonds, D. *Chem Biol Interact* 23(1): 121-135; 1978.

The dihydrodiol metabolites of 3-methylcholanthrene (3-MC) were prepared by direct oxidation of the hydrocarbon with ascorbic acid-ferrous sulfate-EDTA. All six structurally identifiable dihydrodiols were obtained and identified as: cis-2a,3-dihydroxy-3-methylcholanthrene (A), trans-4,5-dihydro-4,5-dihydroxy-3-methylcholanthrene (B), trans-7,8-dihydro-7,8-dihydroxy-3-methylcholanthrene (C), trans-9,10-dihydro-9,10-dihydroxy-3-methylcholanthrene (D), cis-11,12-dihydro-11,12-dihydroxy-3-methylcholanthrene (E), and trans-11,12-dihydro-11,12-dihydroxy-3-methylcholanthrene (F). Examination by high-pressure liquid chromatography of the dihydrodiols formed during the metabolism of MC by rat liver microsomal preparations showed the presence of D, E, and F. Cis- and trans-1,2-dihydroxy-3-methylcholanthrene, A, and E were tentatively identified as metabolites. Microsomal preparations from the livers of rats pretreated with 5 mg MC (ip, 48 hr before killing) formed larger amounts of dihydrodiols from ³H-labeled 3-MC than did preparations from untreated rats, with the exception of metabolite A. The metabolism of MC by mouse skin microsomes in short-term organ culture yielded only D, together with cis- and trans-1,2-dihydroxy-3-methylcholanthrene (refs)

78-5531 Nature of Oncofetal Antigen Expressed on 3-Methylcholanthrene-induced Sarcoma Cells. (Jpn) Takami, T. (Dept. Pathology, Sapporo Medical Coll., Sapporo-shi, Hokkaido 060, Japan). *Sapporo Igakusha Zasshi* 47(4): 409-420; 1978.

The immunogenicity of oncofetal antigen(s), which are expressed on the surface of methylcholanthrene (MC)-induced sarcoma cells, was investigated with the use of anti-mouse sera (AES) cross-reactive between tumor and fetal cells. Sera were obtained by immunizing rats with syngeneic sarcoma cells in midgestation. Immunofluorescence studies and complement-dependent cytotoxicity tests revealed that the sarcoma cells reacted with MC-induced tumor cells (KMT-17, KMT-18 and 16 fetal cells, but it did not react with day 10 fetal cells and normal adult thymus cells. In absorption test, KMT-17 and KMT-50 cells had the same absorptive capability against AES. Multiparous sera (MPS) reacted with tumor cells, but their reactivity tended to increase according to the num-

gnancies. Absorption activity against MPS was high day 16 fetuses and weak with day 13 and 14 fetuses. No was seen with day 14 or 16 placenta. Normal sera at rats were cytotoxic against syngeneic tumor cells, but sensitive finding was obtained by indirect immunofluorescence using FITC-conjugated anti-IgG serum. tion of tumor transplantation resistance failed in immunized with syngeneic fetal cells. The results m that oncofetal antigen(s) is expressed in most ally induced tumors and that immunization with heic fetal cells can induce a humoral antibody that reacts with tumors and fetus, but no strong tumor nce such as tumor rejection is induced. (64 refs)

32 Effect of Oil-attached BCG Cell-Wall Skeleton and Thymectomy on the Incidence of Lung Cancer and Amyloidosis Induced by Chemical Carcinogens in Rats. (Eng) Hirao, F. (Third Dept. Internal Medicine, Univ. Medical School, Fukushima 1-1-50, Fukushima, Osaka 553, Japan); Nishikawa, H.; Yasaki, S.; Taniyama, T.; Azuma, I.; Yoshimoto, T.; Kito, S.; Yamamura, Y. *Gann* 69(4): 453-459; 1978.

Investigation was made of the effect of oil-attached BCG cell skeletons (BCG-CSW) and of thymectomy on orally induced lung cancer and amyloidosis in rabbits. Rats were given a mixture of 3-methylcholanthrene (40 mg/kg) and 4-nitroquinoline 1-oxide (0.4 mg) intrabronchially every 40 days. BCG-CSW was administered either once, in a single injection of 5 mg, or continuously, 5 mg the first day and then 2 mg every 30-40 days. Lung cancer developed in 13/36 rabbits treated with carcinogens alone (controls), but no cancer was seen in animals treated with multiple doses of BCG-CSW. Rabbits given the single dose of BCG-CSW had a cancer incidence (50%) similar to that of controls (47%). Rabbits thymectomized at 30 days and subsequently treated with the carcinogen mixture had a high lung cancer rate (32/40). There were no differences in the appearance of amyloidosis among thymectomized, BCG-CSW, and control rabbits. (17 refs)

33 Particulate Enhanced Membrane Uptake of 1,2-Benzanthracene Observed by Fluorescence Spectroscopy. (Eng) Lakowicz, J. R. (Freshwater Biological Dept. Biochemistry, Univ. Minnesota, P. O. Box 100, St. Paul, MN, 55392); Englund, F.; Hidmark, A. *Biochim Biophys Acta* 543(2): 202-216; 1978.

Transfer of 1,2-benzanthracene (BA) from the adsorbed state to phospholipid vesicles of dipalmitoyl-L- α -phosphatidylcholine was studied and compared with rates of transfer from crystalline and microcrystalline suspensions. The fluorescence spectral shifts that occurred upon transfer of BA from the silica surface to the phospholipid vesicles were used to quantitate the membrane

uptake of BA from particulates. The rate of BA uptake by the dipalmitoyl-L- α -phosphatidylcholine vesicles was independent of the lipid concentration, suggesting that the solubilization rate of BA in the aqueous phase was rate limiting. The rate of membrane uptake was also independent of the surface density of BA on silica, indicating that the BA molecules were uniformly dispersed on the silica surface. Adsorption of BA to silica greatly enhanced its rate of transport into the vesicles, the silica-adsorbed BA being more rapidly transported into the vesicles than was microcrystalline BA at temperatures from 8°C to 42°C. Silica apparently did not disrupt the vesicles under the conditions used. The data indicate that particulates can act as cocarcinogens by enhancing the cellular uptake of carcinogens and by facilitating the expression of preneoplastic transformation. (27 refs)

78-5534 Cyclic Nucleotide Concentrations in 7,12-Dimethylbenz(a)anthracene-induced Pancreatic Cancer in Rats. (Eng) Stevens, R. H. (Radiation Res. Lab., Dept. Radiology, Univ. Iowa, Iowa City, IA, 52242); Graves, J. M.; Meek, E. S.; Osborne, J. W.; Cheng, H. F.; Loven, D. P. *J Natl Cancer Inst* 61(5): 1281-1284; 1978.

Cyclic AMP (cAMP) and GMP (cGMP) levels were measured in pancreatic adenocarcinomas induced in noninbred male Holtzman rats by the implantation of beeswax containing 7,12-dimethylbenz(a)anthracene (DMBA) into the head of the pancreas. Within 4-6 mo after implantation, >80% of the rats developed pancreatic tumors. The cells measured for intracellular concentrations of cAMP and cGMP belonged to two distinct categories: (1) tumors and adjacent tissue that contained significantly lower amounts of the cyclic nucleotides than normal tissue; and (2) tumors and adjacent tissue that contained significantly greater amounts. The cyclic nucleotide levels detected in adjacent tissue, which appeared to be noncancerous, may reflect the presence of some neoplastic cells or may be due to effects of DMBA or its metabolites. A significantly greater cAMP/cGMP ratio was calculated for the two tumor groups, whereas no difference was found for the tissue adjacent to the tumor. These results suggest the existence of two types of tumor cells that may be under similar cyclic nucleotide control. The roles of cAMP and cGMP in the growth and differentiation of pancreatic cancer remain to be determined. (32 refs)

78-5535 Expression of RNA of an Endogenous Replication-defective Retrovirus in Rat Mammary Adenocarcinomas Induced by 7,12-Dimethylbenz(a)anthracene. (Eng) Young, H. A. (Lab. Tumor Virus Genetics, Div. Cancer Cause and Prevention, NCI, Bethesda, MD, 20014); Wenk, M. L.; Goodman, D. G.; Scolnick, E. M. *J Natl Cancer Inst* 61(5): 1329-1337; 1978.

The possible relationship between chemical carcinogen induction of rat mammary tumors and the expression of an

endogenous retroviral genome was investigated in female SD rats. At 50 days of age, the rats received a single 1.0-ml intragastric dose of a 20-mg/ml soln of 7,12-dimethylbenz(a)anthracene (DMBA). At various times, tumor-bearing rats were killed and the tumor type was identified histologically. A total of 50 tumors was obtained from 17 DMBA-treated rats. Thirty-five were mammary adenocarcinomas, 5 fibroadenomas, and the remaining 10 miscellaneous lesions. The mammary adenocarcinomas were analyzed for the RNA of a replication-defective (RD) endogenous retrovirus or the RNA of a helper-independent endogenous C-type virus. Expression of RD virus RNA was detected in mammary tumors weighing 0.2-2.0 g. Larger tumors, for which histologic examination revealed proportionally more fibroblastic tissue than epithelial cells, did not contain comparable concentrations of this viral RNA. RNA homologous to a helper-independent rat C-type retrovirus was not detected in tumors of any size. A cell line was established from a primary DMBA-induced mammary adenocarcinoma and it appeared similar to the small mammary tumors with respect to endogenous C-type viral RNA expression. The results indicate the necessity of considering the role of RD endogenous retroviruses in carcinogen-induced, radiation-induced, or naturally occurring tumors in which viral particles have not been detected. (51 refs)

- 78-5536 Reaction of 7,12-Dimethylbenz(a)anthracene with DNA of Fetal and Maternal Rat Tissues In Vivo.** (Eng) Doerjer, G. (Abteilung Neuropathologie, Pathologisches Institut der Universität Freiburg, 7800 Freiburg, W. Germany); Diessner, H.; Bucheler, J.; Kleihues, P. *Int J Cancer* 22(3): 288-291; 1978.

The binding of 7,12-dimethylbenz(a)anthracene (DMBA) to DNA of various maternal and fetal rat tissues was investigated following a single injection of ³H-DMBA (15 mg/kg, 48 mCi/kg, iv) to two pregnant BD-IX rats on day 21 of gestation. This dose is known to induce a high incidence of nervous system tumors in the offspring. The rats were sacrificed 12 hr later, and DMBA-deoxyribonucleoside products from the DNA of the tissues were separated on Sephadex LH20 columns. The extent to which the major DMBA-DNA adduct was formed in vivo varied considerably in both maternal and fetal organs. In the mother, concentrations of this adduct (disintegrations/minute per micromole of adenine) were highest in the small intestine (19,278), liver (11,576), and lung (10,711), followed by spleen (7,757), kidney (3,656), and brain (1,328). Concentrations in fetal intestine (12,703) and liver (9,681) were much lower than those in the maternal organs, but the concentration in fetal brain (3,292) was 2.5 times higher than that in maternal brain DNA, possibly due to differences in DMBA uptake through the blood-brain barrier. The results indicate that there is no significant placental barrier to DMBA or DMBA metabolites involved in DNA binding and that rat fetuses participate in the metabolic formation of the ultimate carcinogen. (26 refs)

- 78-5537 DNA Labeling Index and Structure of the Mammary Gland as Determinants of its Susceptibility to Carcinogenesis.** (Eng) Russo, J. (Experimental Pathology Lab., Dept. Biology, Michigan Cancer Foundation, 110 E. Warren Ave., Detroit, MI, 48201); Russo, I. H. *Cancer Inst* 61(6): 1451-1459; 1978.

In a study of 7,12-dimethylbenz(a)anthracene (DMBA)-induced mammary carcinomas, an attempt was made to correlate the number of mammary gland terminal end-buds (TEB), terminal ducts (TD), and alveolar buds (AB), together with their DNA labeling indexes, at the time of DMBA administration with the number and types of tumors induced by the carcinogen. Young virgin (55 days old), old virgin (180 days old), and multiparous (180 days old) Sprague-Dawley rats were inoculated iv with methyl-³H-thymidine (2.5 μ Ci) and killed 1 hr later (Group 1). The number of ducts and buds and the DNA labeling index in the epithelial cells of these structures were compared with the number of tumors developing in age-matched mice (Group 2) 22 wk after they received DMBA (10 mg intragastrically) + methyl-³H-thymidine. The number of TEB, TD, and labeled cells was highest in young virgins, and 100% of the tumors that developed in these animals were adenocarcinomas. The older virgins had fewer TEB and TD and fewer labeled cells, and these rats developed fewer adenocarcinomas (63%). Glands of multiparous rats contained fewer TEB, a small number of TD, and few labeled cells; these rats had the lowest incidence (21%) of adenocarcinomas. In the old virgins and multiparous rats, the incidence of fibroadenomas was 36% and 79%, respectively. The tumor latency period was longer for these rats than for the young virgins. The percentage of DNA-labeled AB at the time of DMBA administration was 50% in young virgins compared with 0.5% in old virgins and 0.9% in multiparous rats. These results demonstrate that susceptibility to carcinogenesis is correlated with the density of TEB and TD and with DNA labeling indexes. (69 refs)

- 78-5538 DMBA Induced Papillomas in Congenitally Athymic (Nude) and Hereditarily Asplenic (Dh/+) Mice: Contrasts and Comparisons with Immunologically Intact Littermates.** (Eng) Gershwin, M. E. (School of Rheumatology-Clinical Immunology, Dept. Medicine, University of California, Sch. Medicine, Davis, CA, 95616); Ikeda, R. *Dev Comp Immunol* 2(3): 529-537; 1978.

The induction of papillomas by dimethylbenz(a)anthracene (DMBA) and hereditarily asplenic (Dh/+) mice and in immunologically normal littermates [nude/+ and B6.CBA (+/+) respectively]. DMBA (200 μ g) was applied weekly for 20 wk to the shaved flank skin of the mice. The incidence of papillomas at the end of 20 wk was 17% among the nude mice, 78% among the nude/+ mice, 21% among the Dh/+ mice, and 75% among the B6.CBA (+/+) mice. Furthermore, the number of papillomas/mouse and the tumor growth rate were greater for the normal mice than for their immune-deficient littermates, and the tumor la-

was shorter. Nude/+ skin that was transplanted to mice and then painted with DMBA reacted in a manner similar to that of nude skin. The data suggest that the immune system has a major role in stimulating papilloma induction (22 refs)

539 Resistance to a Transplantable Rat Mammary Tumor (DMBA 8BT) after Pre-treatment with Syngeneic Fetal Cells. (Eng) Cocuzzi, E. (Dept. Clinical Biochemistry, Sunnybrook Medical Centre, Univ. Toronto, 2075 Kennedy Ave., Toronto, Ontario, Canada M4N 3M5); Kelce, J. A. *Res Commun Chem Pathol Pharmacol* 20(2): 108; 1978.

Effect of presensitization to fetal antigens on the transplantability of a chemically induced rat mammary tumor, the DMBA 8BT adenocarcinoma (induced by feeding female Fischer 344 rats with 7,12-dimethylbenz(a)anthracene), was investigated in female Fischer 344/Mia albino rats. The rats were immunized with mid-term syngeneic fetal cells (3 sections of 5×10^5 viable cells at 7-day intervals) and injected with tumor cells 11 days after the final fetal cell immunization. This immunization schedule provided complete protection against an inoculum of up to 5×10^5 tumor cells, an amount that leads to a 100% tumor take without pretreatment. No difference in tumor take, growth rate, or survival was observed between virgin and retired breeder rats. Multiplicity does not protect against this tumor model. (4 refs)

540 Cultivation of Human Adult Skin Fibroblasts & Induction of Malignant Transformation with 7,12-Dimethylbenz(a)anthracene. (Eng) Gokhale, J. A. (Biophysics Div., Cancer Res. Inst., Tata Memorial Centre, Bombay 400 012, India); Rao, K. V.; Bhat, A. V.; Bapat, C. V. *Indian J Exp Biol* 16(5): 549-551; 1978.

Effect of 7,12-dimethylbenz(a)anthracene (DMBA) on fibroblasts from the skin of a patient with mucopolysaccharidosis was studied to examine the possible susceptibility of genetically deficient human cells to chemical carcinogens. These cells exhibited a typical fibroblastic morphology and normal growth with contact inhibition. Exposure of confluent cells to DMBA ($0.1 \mu\text{g/ml}$) for 72 hr resulted in loss of cell orientation and the occurrence of small colonies with a criss-cross growth pattern. However, these cells did not induce tumors when inoculated (1×10^6 cells) into the cheek pouch of golden hamsters. In another experiment, confluent cultures were exposed to UV light (160 ergs/cm^2 at 250-260 nm) for 30 sec, followed immediately by DMBA ($10 \mu\text{g/ml}$). These fibroblasts piled up into an irregular multilayer with increased saturation density. These DMBA-treated cells produced a solid 1-cm nodule very similar to a tumor, when inoculated into the hamster cheek pouch. In this study, certain morphological, behavioral, and biochemical changes occurred in the cells that were indicative

of malignant change but did not yield a proliferative population. However, these observations suggest that cells with a genetic defect may be a feasible model for the induction of malignant transformation in vitro. (12 refs)

78-5541 Transplacental Penetration of 7,12-Dimethylbenz(a)anthracene and Its Distribution in Organs of the Rat Fetus. (Rus) Baranova, L. N. (Lab. Biophysics, N. N. Petrov Scientific Res. Inst. Oncology, Leningrad, USSR); Aleksandrov, V. A. *Biull Eksp Biol Med* 86(10): 477-479; 1978.

The distribution and elimination of 7,12-dimethylbenz(a)anthracene (DMBA) in maternal and fetal rat organs were studied 5-300 min after iv injection of a single 15-mg/kg dose to white rats on day 21 of pregnancy. The DMBA level reached its max in the liver ($45 \mu\text{g/kg}$) and in the placenta ($6.3 \mu\text{g/kg}$) at 10-15 min after injection. The DMBA concentration peaked at 1 hr after injection in the fetal tissues, and only trace amounts were found after 5 hr. The DMBA levels in the fetus 1 hr after injection were $4.7 \mu\text{g/kg}$ in the liver, $2.4 \mu\text{g/kg}$ in the kidney, $2.2 \mu\text{g/kg}$ in the lungs, $1.5 \mu\text{g/kg}$ in the brain, $1.8 \mu\text{g/kg}$ in the intestine, and $1.1 \mu\text{g/kg}$ in the carcass. The findings do not correlate with data on the preferential formation of renal and nervous system tumors following transplacental exposure to DMBA. (4 refs)

78-5542 Differentiation of Malignant Myoblasts in 7,12-Dimethylbenz(a)anthracene-induced Rhabdomyoblastomas. An Electron Microscopic Study. (Eng) Markov, D. V. (Regeneration Res. Lab., Bulgarian Acad. Sciences, 1431 Sofia, Bulgaria); Hadjiolov, D. C. *Arch Geschwulstforsch* 48(6): 521-537; 1978.

Samples of necrotic tissue and rhabdomyoblastomas induced in male Syrian hamsters by im injections of 1 mg 7,12-dimethylbenz(a)anthracene were analyzed by light and electron microscopy at various times postinjection. Early neoplastic growth was characterized by the presence of small cells (SC) and long spindle-shaped cells (LSSC). SC always possessed a single heterochromatic nucleus with a small, compact nucleolus. The cytoplasm was characterized by well-developed Golgi apparatus (GA), dense, tubular smooth endoplasmic reticulum (SER), and almost complete absence of granular endoplasmic reticulum (GER). Also present were numerous centrioles and small lysosomelike bodies. These features were indicative of a low level of biosynthetic activity in these cells; this conclusion was supported by their low degree of differentiation. SC were hypothesized to be the precursors of the LSSC, which had the characteristics of elevated biosynthetic activity. Variable numbers of large euchromatic nuclei were present with multiple nucleoli. The cytoplasm was characterized by well-developed GA and GER, varying amounts of cytofilaments, and numerous ribosomes. Occasional LSSC contained neither GA nor cytofilaments but showed GER

with a dense substance accumulated within enlarged cisternae. These LSSC had a tendency to fuse and resembled normal myoblasts morphologically and developmentally, but they did not normally form myofibrils. It is concluded that the SC formed the proliferative cell compartment of the tumor. (67 refs)

- 78-5543 Vasopressin Stimulation of Acetate Incorporation into Lipids in a Dimethylbenz(a)anthracene-induced Rat Mammary Tumor Cell Line.** (Eng) Monaco, M. E. (Medicine Branch, NCI, Bethesda, MD, 20014); Lippman, M. E.; Knazek, R.; Kidwell, W. R. *Cancer Res* 38(11, part 2): 4101-4104; 1978.

The possibility that the effects of rat prolactin B-1 (PL) on acetate incorporation into lipids are caused by vasopressin (VP) contamination was investigated in a cloned cell line (WRK-1) derived from a dimethylbenz(a)anthracene-induced rat mammary tumor. There was a twofold increase in ^{14}C -acetate incorporation with PL alone. In the presence of insulin ($5 \times 10^{-7} \text{ M}$), the effect was increased 10-fold. Concentrations of PL required for max stimulation in both instances were $1.5 \mu\text{g/ml}$, and max activity was not seen until 36 hr after addition of the hormones. Rat growth hormone B-3 elicited a response similar to that with PL. More purified rat PL 1-2, human growth hormone HS-2019G, and ovine PL P-S-12 had minimal effects on the incorporation of ^{14}C -acetate into lipids. Activity appeared to be inversely related to the purity of the preparation. Both VP (10^{-10} M ; 20 micro units/ml) and oxytocin elicited the response observed with impure PL and growth hormone, but 100 times more oxytocin was required. This suggests that VP is the physiologically relevant hormone. Following filtration of rat PL, virtually all the activity resided in the low-mol-wt filtrate. The activity observed in this system would require a minimal VP contamination of 0.02%, which is well within the concentrations detected by radioimmunoassay in the rat PL and growth hormone B-3 preparations used in this study. Thus, VP is probably responsible for some of the activity observed with both rat PL and growth hormone. (11 refs)

- 78-5544 Synthesis of Potentially Estrogenic Carcinogens: 7,12-Dimethylbenz(a)anthracene-3,9-diol.** (Eng) Morreal, C. E. (Dept. Breast Surgery, Roswell Park Memorial Inst., Buffalo, NY, 14263); Bronstein, R. E. *J Chem Eng Data* 23(4): 354-356; 1978.

The synthesis of a diphenolic derivative of 7,12-dimethylbenz(a)anthracene, 7,12-dimethylbenz(a)anthracene-3,9-diol, from 4-methoxyphthalic anhydride and 6-methoxy-2-bromonaphthalene is described. The diol has the potential to behave as an estrogen and as a carcinogen. (5 refs)

- 78-5545 The Effect of Altered Thyroid State on Prolactin Binding to Liver and 7,12-Dimethylbenz(a)anthracene-induced Mammary Tumors in Rats.** Smith, R. D. (Dept. Biochemistry, Univ. Rochester Medical Center, Rochester, NY, 14642); Hilf, R.; Senior, A. E. *Soc Exp Biol Med* 158(4): 517-520; 1978.

The effect of thyroid state on prolactin (PRL) binding to liver and on 7,12-dimethylbenz(a)anthracene (DMBA)-induced mammary tumors was studied in female Sprague-Dawley rats. Rats were given 5 mg DMBA/wk for 5 wk by gavage. Rats bearing tumors of sufficient size (2-3 cm) were made hypothyroid by administration of propylthiouracil (0.1% wt/vol in the drinking water for 21 days) or hyperthyroid by injections of L-thyroxine ($75 \mu\text{g}/100 \text{ g}$ body weight daily for 10 days). Neither hypo- nor hyperthyroidism significantly influenced the growth rates of established DMBA-induced tumors, PRL binding to these tumors, or plasma PRL in tumor-bearing rats. PRL binding to membrane particles was reduced 85% following either hypothyroid or hyperthyroid induction in tumor-bearing rats. The physiological role of the liver PRL receptors remains unknown. (9 refs)

- 78-5546 Effect of Oophorectomy on Estrogen Receptor Binding in Rat Mammary Tumors Induced by 7,12-Dimethylbenz(a)anthracene.** (Eng) Tominaga, T. (Dept. Surgery, Tokyo Metropolitan Komagome Hosp., 3-18-1 Honkomagome, Bunkyo-ku, Tokyo, Japan); Kitamura, T.; Tei, N. *Biken J* 20(3/4): 143-145; 1977.

The relationship between the response of 7,12-dimethylbenz(a)anthracene (DMBA)-induced rat mammary tumors to oophorectomy and the presence of estrogen receptors (ER) in the tumor tissue was examined. Intravenous administration of 10 mg/kg DMBA to female Sprague-Dawley rats resulted in 100% tumor incidence. Small sections of tumor were removed and tested for the presence of ER. Bilateral oophorectomy was then performed, the residual tumors were removed 7-11 days later, and tests for ER were repeated. Before oophorectomy, ER were found in 18/33 tumors. After oophorectomy, a special correlation was noted between the histological differentiation of the mammary carcinomas or the time of tumor induction after DMBA administration and the presence of ER. Oophorectomy resulted in the disappearance of ER in 17/18 tumors within several days of surgery. ER were still present in the remaining rat 7 day after oophorectomy, but one of the 18 tumors with ER decreased in size 7-11 days after oophorectomy. Of the 15 tumors without ER, 4 regressed, and 2 remained unchanged following oophorectomy; no ER were found in any of these tumors after oophorectomy. Since a tumor that decreased in size after oophorectomy lost its ER completely within 10 days and another tumor without ER increased in size after oophorectomy, it was concluded that tumor growth was not dependent on ovarian hormones alone. Further studies in which rats received estradiol-17 β suggested that tumor growth is closely related to this compound, and not with the presence of ER. (9 refs)

78-5547 Synthesis and Carcinogenic Activity of 5-Fluoro-7-(oxygenated methyl)-12-methylbenz(a)anthracenes. (Eng) Newman, M. S. (Chemistry Dept., Ohio State Univ., Columbus, OH, 43210); Fikes, L. E.; Hamann, M. M.; Kannan, R.; Sankaran, V. *J Med Chem* 21(10): 16-1078; 1978.

5-fluoro-7-(oxygenated methyl)-12-methylbenz(a)anthracenes were synthesized and their carcinogenicity was assayed in rats. Groups of 18 male Fischer rats were injected sc with 12-dimethylbenz(a)anthracene (DMBA: single injection of 2.3 mg), 5-fluoro-7-hydroxymethyl-12-methylbenz(a)anthracene (5-F-7-OH-DMBA: 2.3 mg 1x/wk for 4 wk), or 5-fluoro-7-acetoxymethyl-12-methylbenz(a)anthracene (5-F-7-OAc-DMBA: 2.7 mg 1x/wk for 4 wk). Three of the eight rats injected with DMBA had sarcomas at the injection site by 6, 9, and 16 mo after injection. By contrast, only 1/18 rats that had received 5-F-7-OH-DMBA and 0/18 that had received 5-F-7-OAc-DMBA developed sarcomas by the end of 16 mo. In another experiment, male Sprague-Dawley rats were injected sc with 5-fluoro-7,12-benz(a)anthraquinone (5-F-BAQ); 5-F-7-OAc-DMBA, or 5-F-7-OH-DMBA (0.1 mg/mole 3x/wk for 30 doses, beginning at age 30 days). One of the rats injected with 5-F-7-OH-DMBA, 3/13 injected with 5-F-7-OAc-DMBA, and 1/12 injected with 5-F-BAQ developed sarcomas at the injection site. Since neither 5-F-7-OH-DMBA nor 5-F-7-OAc-DMBA was appreciably carcinogenic, the carcinogenic metabolism of DMBA probably does not involve attack at the 7-methyl group. The methyl group merely serves to block a detoxification metabolism that is initiated at the 7 position. When the detoxification pathway is blocked, the cancer-producing mechanism can become effective. (22 refs)

78-5548 Metabolic Activation of Dibenzo(a,h)anthracene and Its Dihydrodiols to Bacterial Mutagens. (Eng) Wood, A. W. (Dept. Biochemistry and Drug Metabolism, Hoffmann-LaRoche Inc., Nutley, NJ, 07110); Levin, W.; Thomas, P. E.; Ryan, D.; Karle, J. M.; Yagi, H.; Pina, D. M.; Conney, A. H. *Cancer Res* 38(7): 1967-1973; 1978.

Dibenzo(a,h)anthracene (DBA) and the three metabolically possible trans-dihydrodiols of DBA were tested for mutagenicity toward *Salmonella typhimurium* strain TA100. The tests were performed in the presence of hepatic microsomes from Aroclor 1254-pretreated rats or in the cytochrome P-450-dependent monooxygenase system isolated and purified to near homogeneity from these microsomes. With either enzyme system, the 3,4-dihydrodiol was activated to products that were more mutagenic than the metabolites of DBA or its 1,2- and 5,6-dihydrodiols. With microsomes, but not with the purified monooxygenase system, the 5,6-dihydrodiol was also activated to products that were highly mutagenic. However, significant activation was seen only when high concentrations of the 5,6-dihydrodiol were incubated with relatively high amounts of enzyme. Saturation

of the double bond at position 1,2 of the 3,4-dihydrodiol produced a tetrahydrodiol that was poorly activated in either metabolic activation system. The high activity of the 3,4-dihydrodiol in both systems and the importance of the double bond in position 1,2 of the dihydrodiol strongly suggest that a bay-region 3,4-diol-1,2-epoxide is a biologically important active metabolite of DBA, thereby providing support for the bay region theory. (39 refs)

78-5549 Determination of Benzo(a)pyrene in a Filtered Retort Water Sample by Solid-Surface Fluorescence. (Eng) Hurtubise, R. J. (Dept. Chemistry, Univ. Wyoming, Laramie, WY, 82071); Phillip, J. D.; Skar, G. T. *Anal Chim Acta* 101(2): 333-338; 1978.

A method is described for determining benzo(a)pyrene (BP) in filtered retort water from an in situ oil shale process by liquid-liquid extraction, dry-column chromatography, thin-layer chromatography, and fluorescence spectrometry. BP was identified by detecting its fluorescence directly from chromatoplates. The accuracy and precision of the method were good; the detection limit was approx 0.08 ppb. (17 refs)

78-5550 Factors Affecting the Binding of Polycyclic Aromatic Hydrocarbons to Human Embryo Cells, and Transformable and Non-transformable Hamster Embryo Cells. (Eng) Raineri, R. (Chemical Carcinogenesis Program, NCI, Frederick Cancer Res. Center, Frederick, MD, 21501); Pooley, J. A.; Hillesund, T.; Pienta, R. J. *Chem Biol Interact* 22(2/3): 257-270; 1978.

The effects of various factors on the binding of three polycyclic aromatic hydrocarbons to human (HuEC) and Syrian golden hamster embryo cells (HaEC) were studied. The binding of 7,12-dimethylbenz(a)anthracene (DMBA) to HaEC DNA, RNA, and protein was max after 22 hr of treatment. In contrast, binding to HuEC macromolecules increased for at least 55 hr. Treatment of HaEC at 100% confluence resulted in much less binding than treatment at 70% confluence, whereas with HuEC the binding increased or remained constant following treatment at the greater confluence. The transforming frequency of HaEC decreased with increasing population doubling number (PDN): binding of DMBA to hamster embryo DNA, RNA, and protein decreased approx 100-fold between PBN's 8 and 20. In transformable cell cultures, DMBA was bound to HaEC DNA to a greater extent than to RNA or protein. The binding of DMBA to HuEC DNA was low at population doubling 30 and did not change with increasing PDN. The binding to HuEC RNA was also low, but the binding to protein increased 2.5-fold between PDN's 5 and 70, after which binding decreased sharply. The binding of DMBA to nucleic acids was much greater than binding by dibenz(a,h)anthracene or dibenz(a,c)anthracene, both of which had low binding values at all PDN's tested. Therefore, the best correlation of binding with carcinogenicity

ty and transforming activity was observed with DMBA. (16 refs)

78-5551 Fluorometric Detection of Carcinogen Binding.

(Eng) Rho, J. H. (Univ. California, Schs. Medicine and Pharmacy, Los Angeles, CA); Okigaki, T.; Bonner, J. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2 (Part 1), 2404 pp.; 1365-1374; 1978.

The results of a three-dimensional fluorometric analysis of benzo(a)pyrene (BP) metabolites bound to the chromatin components of cultured Syrian golden hamster embryonic lung cells are reported. The cells were exposed to ^3H -BP (0.5 $\mu\text{g}/\text{ml}$) for 24 hr. The chromosomal constituents were separated by centrifugation and electrophoresis into the DNA fraction, proteins tightly bound to DNA, and 4 M NaCl-soluble proteins. The three-dimensional fluorescence spectra revealed that the NaCl-soluble proteins were bound predominantly with monohydroxylated BP, the tightly bound proteins with *cis*-7,8-diol-9,10-epoxy-BP, and the DNA with BP itself. These results indicate that a BP metabolite similar to the *cis*-7,8-diol-9,10-epoxide forms a molecular complex between various nonhistone chromosomal proteins and DNA through a carcinogenic reaction of the metabolite. (20 refs)

78-5552 Glucuronidation of Benzo(a)pyrene in Hamster

Embryo Cells. (Eng) Nemoto, N. (Dept. Experimental Pathology, Cancer Inst., 1-37-1, Kami-Ikebukuro, Toshima-ku, Tokyo 170, Japan); Hirakawa, T.; Takayama, S. *Chem Biol Interact* 22(1): 1-14; 1978.

Cultures of Syrian hamster embryo cells were exposed to 3.75 or 37.5 nanomoles (nmol)/ml of ^3H -labeled benzo(a)pyrene (BP), and the formation of BP metabolites before and after treatment of the medium with β -glucuronidase (GU) was analyzed by thin-layer chromatography (TLC) and high-pressure liquid chromatography (HPLC). The ratio of radioactivity (RA) in the aqueous phase to that in the organic phase increased with time of incubation. After incubation for 48 hr with 3.75 nmol/ml BP, 92.5% of the RA was found in the aqueous phase, but with 37.5 nmol/ml BP, 53.4% of the RA remained in the organic phase. Medium containing the higher dose of BP was extracted with ethyl acetate, and BP metabolites in the organic phase were separated by HPLC. The major metabolites eluted were the 9,10-diol and 7,8-diol; the amount of 4,5-diol was negligible. After treatment of the medium with GU, however, the major oxygenated metabolites were shown by TLC to be phenols, with the amount of 3-hydroxy-BP being greater than that of 9-hy-

droxy-BP. GU also released 9,10-diol and 7,8-diol, but most of these diols were in free form in the medium. With the low BP concentration, the metabolic profile was different: diols and phenols were barely detectable, and almost nothing except intact BP was found. However, on treatment with large amounts of phenols and small amounts of diols appeared. The results show that in hamster embryo cells, BP is mainly metabolized to conjugates of phenols with glucuronic acid. (29 refs)

78-5553 Modulation of the Binding of the Carcinogen Benzo(a)pyrene to Rat Liver DNA In Vivo

Selective Induction of Microsomal and Nuclear Aryl Hydrocarbon Hydroxylase Activity. (Eng) Viviani, A. (Inst. Toxicology, Federal Inst. Technology, CH-8603 Schwerzenbach, Zurich, Switzerland); Lutz, W. K. *Cancer Res* 38(11): 4640-4644; 1978.

The influence of microsomal and nuclear aryl hydrocarbon hydroxylase (AHH) activity on the covalent binding of ^3H -benzo(a)pyrene to rat liver DNA was evaluated in vivo. Phenobarbital treatment (1 g/liter in drinking water) induced microsomal AHH in male Sprague-Dawley rats to 160% of control. This treatment also increased DNA binding to 190%, but left the nuclear activity unchanged. Nuclear AHH was induced to 150% of control by dieldrin treatment (1 mg/kg/day, ip), and binding was decreased to 75%, whereas the microsomal AHH was at control level. The increased effect of microsomal AHH induction as well as the decreased effect of nuclear AHH induction on binding was shown clearly when the data on individual rats were used to solve the following equation: binding = $a \cdot (\text{microsomal AHH}) + b \cdot (\text{nuclear AHH}) + C$. Multiple linear regression analysis with the data from 10 animals resulted in positive values for a and c, a negative value for b, and a good multiple correlation coefficient of $r = 0.974$. Pretreatment of the rats with 3-methylcholanthrene (20 mg/kg/day, ip) induced microsomal AHH to 380% of control and nuclear AHH to 590% of control, and increased binding to 175%. The binding was higher than that predicted by the formula, probably because the increasing influence of induced microsomal AHH overshadowed the decreasing effect of nuclear AHH. The results show that the binding of a foreign compound to DNA in vivo is dependent not only on microsomal enzyme activities but also on nuclear activities, even if the latter are considerably lower than those of microsomes. (21 refs)

78-5554 [^3H]Benzo(a)pyrene Metabolism in Tracheal Epithelial Microsomes and Tracheal Organ Cultures.

(Eng) Mass, M. J. (Dept. Pathology, Sch. Medicine, Univ. North Carolina at Chapel Hill, Chapel Hill, NC 27514); Kaufman, D. G. *Cancer Res* 38(11, part 2): 3861-3866; 1978.

^3H -Benzo(a)pyrene (BP) metabolism was studied in Syn-

in hamster tracheal epithelial microsomes and tracheal cell cultures. Microsomal BP metabolism was induced by treatment of tracheas with 5 μ M BP in organ culture. The apparent K_m 's for the induced and constitutive (uninduced) monooxygenase were not significantly different and were 1.5 μ M. However, the induced enzyme had a V_{max} of 86 pmol product/mg microsomal protein/min, whereas the constitutive enzyme had a value of 31 pmol/mg/min, a 2.8-fold increase. Microsomes from BP-treated and uninduced tracheas mediated the binding of 12.84 $\times 10^{-12}$ and 8.06 $\times 10^{-12}$ pmol BP/ μ g DNA/ μ g tracheal microsomal protein respectively, during incubation at 37 C for 30 min. High-pressure liquid chromatography revealed that the major metabolites produced by tracheal microsomes incubated with 2 μ M ^{3} H-BP for 30 min were diols, quinones, and phenols of BP. After 24 hr of incubation with ^{3} H-BP, the metabolites formed in organ cultures of induced tracheas were largely tetrols, triols, and the *trans*-1,2,3,4-dihydro-9,10-dihydroxybenzo(a)pyrene. The disparity between metabolites produced in these two systems may be attributed to coupled secondary metabolic reactions that occur in prolonged incubation in the intact cells of organ cultures. (38 refs)

78-5555 Comparative Studies of the Carcinogenic Polycyclic Aromatic Hydrocarbon Levels in Smoked Meat Products in Hungary and in East Germany. (Hun) Soos, J. (Urszagos Elelmezes- és Taplalkozástudományi Intézet, Budapest, Hungary); Fritz, W. *Egeszsegstudomány* 21(3): 285; 1978.

Forty-two kinds of commercial and homemade smoked meat products from Hungary and 29 kinds of smoked meat products from East Germany, as well as smoked cheese and fish samples, were analyzed for benzo(a)pyrene (BP) and other carcinogenic and cocarcinogenic polycyclic aromatic hydrocarbons (PAH) by UV spectrophotometry and spectrophotometry. The av BP level in the commercial meat products was 0.6 μ g/kg in Hungary and 0.43 μ g/kg in East Germany. The BP levels in the home-smoked meat products were somewhat lower in both countries (0.74 μ g/kg in Hungary and 0.76 μ g/kg in East Germany). The BP level was 0.6 μ g/kg in 8/32 Hungarian samples (2 commercial, 6 home-smoked) and in 4/29 samples from East Germany (1 commercial, 3 home-smoked). The total PAH level was somewhat lower in cold-smoked meat products than in those smoked at 40 or 100 C in both countries. The av BP levels were 0.95 μ g/kg in smoked cheese, 8.65 μ g/kg in smoked fish with skin, and 1.39 μ g/kg in skinned smoked fish. The findings indicate that the permissible BP level in smoked food could be set at 1 μ g/kg in both countries. (20 refs)

78-5556 Asbestos-mediated Membrane Uptake of Benzo(a)pyrene Observed by Fluorescence Spectroscopy. (Eng) Lakowicz, J. R. (Freshwater Biological Inst.,

Univ. Minnesota, County Roads 15 and 19, P.O. Box 100, Navarre, MN, 55392); Hylden, J. L. *Nature* 275(5679): 446-448; 1978.

The ability of asbestos and silica to adsorb benzo(a)pyrene (BP) and to deliver the adsorbed carcinogen to a model membrane was investigated by fluorescence spectroscopy. The surface area of the silica sample was 381 m²/g, and most particles were 1.5-15 μ m in diameter (av 3 μ m). The surface area of the asbestos sample was 5.6 m²/g, and the particle size was 1.5-10 μ m (av 2 μ m). Particulate samples containing BP were prepared by mixing the particulate with a benzene soln of BP (0.1-3.0 mg BP/g particulate), followed by evaporation of the benzene under reduced pressure. The amounts of BP adsorbed to silica and asbestos were 85% of the amount added. Although the surface area of the asbestos sample was 60 times less than that of the silica sample, asbestos was more effective than silica in transferring BP to the membrane at all BP-to-particulate wt ratios examined. Asbestos or silica, when added to preformed microcrystals of BP in buffer, did not increase the rate of BP membrane uptake, indicating that only BP that is adsorbed to the particles displays an increased rate of membrane uptake. It is concluded that increased cellular exposure to carcinogens, resulting from the presence of particulates that can adsorb and deliver these carcinogens to the cell, may be involved in the process of cocarcinogenesis. (20 refs)

78-5557 Inhibition of Benzo(a)pyrene Binding to DNA in Cultured Human Bronchi. (Eng) Harris, C. C. (Human Tissue Studies Section, NCI, Building 37, Room 3A07, Bethesda, MD, 20014); Autrup, H.; van Haaften, C.; Connor, R.; Frank, A. L.; Barrett, L. A.; McDowell, E. M.; Trump, B. F. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1359-1364; 1978.

The binding of benzo(a)pyrene (BP) to cultured human bronchial cell DNA was shown to be reduced by inhibitors of BP metabolism and inhibitors of chemical carcinogenesis. Cells from specimens of human bronchi removed at 'immediate autopsy' or during surgery from 9 patients with and from 5 patients without lung cancer were cultured. After 7 days, the cultures were exposed for 24 hr to ^{3} H-BP (25 Ci/mmol) either with or without (controls) one of the following: 7,8-benzoflavone (BF) (BF: 20 μ M), vitamin E (VE: 10 and 100 μ M), disulfiram (DS: 10 and 100 μ M) or vitamin A (VA: 3 and 30 μ M). Compared with controls, the level of bound BP was decreased by BF and, to a lesser extent, by DS (100 μ M) and VE (100 μ M). The binding level was not significantly altered by VA, DS (10 μ M), or VE (10 μ M). There were no binding differences between cultures from cancer or non-cancer patients. The results show that it may be possible to

quantify and modify the interaction of chemical carcinogens with specific human target tissues in a controlled experimental setting. (17 refs)

- 78-5558 Fluorescence Studies of Benzo(a)pyrene Uptake by Cultured Macrophages and Lymphocytes.** (Fre) Salmon, J. M. (Laboratoire de Chimie physique, Centre Universitaire, avenue de Villeneuve, 66025 Perpignan, France); Thierry, C.; Serrou, B.; Viallet, P. *C R Acad Sci D (Paris)* 287(4): 345-348; 1978.

The kinetics of benzo(a)pyrene uptake by human lymphocytes and mouse macrophages was studied in vitro by microspectrofluorometry. The uptake appeared to be a first-order reaction. A fluorescent metabolite was detected in the macrophages, but not in the lymphocytes. (8 refs)

- 78-5559 Modification of the Mutagenicity of Benzo(a)pyrene on Bacteria by Substrates of Enzymes Producing Water Soluble Conjugates.** (Eng) Nemoto, N. (Dept. Experimental Pathology, Cancer Inst., 1-37-1, Kami-Ikebukuro, Toshima-ku, Tokyo 170, Japan); Takayama, S.; Nagao, M.; Umezawa, K. *Toxicol Lett* 2(4): 205-211; 1978.

The effects of enzyme substrates that produce water-soluble conjugates on the mutagenicity of benzo(a)pyrene (BP) to *Salmonella typhimurium* strains TA98 and TA100 were studied. At a BP concentration of 2.5 $\mu\text{g}/\text{plate}$, the number of revertants decreased with increasing concentration of uridine 5'-diphosphoglucuronic acid (UDPGA) in the incubation mixture. However, at a BP concentration of 10 $\mu\text{g}/\text{plate}$, the number of revertants was slightly increased in the presence of UDPGA; this increase was greater with TA98 than with TA100. The mutagenicity of BP (5 or 10 $\mu\text{g}/\text{plate}$) was also enhanced by ATP, whereas glutathione (GSH) reduced the number of revertants. UDPGA and, to a lesser extent, ATP inhibited the apparent aryl hydrocarbon hydroxylase (AHH) activity of the S-9 liver fraction from polychlorinated biphenyl-treated rats. GSH had no significant effect on this activity. Thus, one reason why results on the mutagenicities of certain chemicals vary is that UDPGA, ATP, and GSH, present as contaminants in the S-9 fraction, influence the reactions of detoxifying enzymes in this fraction. (23 refs)

- 78-5560 The Effect of Overcrowding Stress on the Carcinogen-metabolizing Enzymes of Rats.** (Eng) Capel, I. D. (Res. Dept., Marie Curie Memorial Foundation, The Chart, Oxted, Surrey RH8 0TL, England); Williams, D. C. *ICRS Med Sci (Cancer)* 6(8): 329; 1978.

The effect of overcrowding (OC) on the hepatic microsomal enzyme system responsible for activating benzo(a)pyrene and

on the plasma disposition of antipyrine [monitored as an estimate of in vivo aryl hydrocarbon hydroxylase (AHH) activity] was determined in Sprague-Dawley rats. OC decreased the rate and amount of antipyrine absorption. The plasma cortisol level was max on day 4 of OC, when there was a fourfold increase in the in vitro AHH level. Although OC increased AHH activity, there was a gradual adaptation to this form of stress. (5 refs)

- 78-5561 Binding of Benzo(a)pyrene at the 1,3,6 Positions to Nucleic Acids In Vivo on Mouse Skin and In Vitro with Rat Liver Microsomes and Nuclei.** (Eng) Rogge, E. (Eppley Inst. Res. Cancer, Univ. Nebraska Medical Center, 42nd and Dewey Ave., Omaha, NB, 68105); Roth, J.; Katomski, P.; Benderson, J.; Cavalieri, E. *Chem Biol Interact* 22(1): 35-51; 1978.

Double-label studies of benzo(a)pyrene (BP) binding to nucleic acids were undertaken to determine if binding of BP to nucleic acids occurs at the 1, 3, and 6 positions of BP and if tritium loss from a specific position is a valid measure of binding at that site. Tritium loss from (6- ^3H , ^{14}C)BP, (1,3- ^{14}C)BP, (1,3,6- ^3H , ^{14}C)BP, (6,7- ^3H , ^{14}C)BP, and (7- ^3H , ^{14}C)BP was studied in vitro using liver nuclei and microsomes from 3-methylcholanthrene-induced Sprague-Dawley rats and in vivo on Charles River CD-1 mouse skin. The relative tritium loss from (^3H , ^{14}C)BP was measured after binding to skin DNA and RNA, to nuclear DNA, and to native and denatured calf thymus and rat liver DNA's and poly(G) by microsome activation. In skin, nuclei, and microsomes plus native DNA, virtually all BP binding occurred at positions 1, 3, and 6; with microsomes plus denatured DNA or poly(G), BP showed no binding at position 6 and little if any at positions 1 and 3. In vivo and in isolated nuclei, binding at position 6 predominated. There was little loss of tritium from position 7; this was expected, because binding at this position is not thought to occur. This confirms the interpretation of loss of tritium as an indication of binding at a given position. The results demonstrate that the use of microsomes to activate BP is not a valid model system for delineating the in vivo mechanism of BP activation, and they support previous evidence for one-electron oxidation as the mechanism of activation of hydrocarbons in binding to nucleic acids. (44 refs)

- 78-5562 Metabolism of Benzo(a)pyrene and Identification of the Major Benzo(a)pyrene-DNA Adducts in Cultured Human Colon.** (Eng) Autrup, H. (Human Tissue Studies Section, Carcinogenesis Res. Program, NCI, Building 37, Room 3A09, Bethesda, MD, 20014); Harris, C.; Trump, B. F.; Jeffrey, A. M. *Cancer Res* 38(11, part 2): 3689-3696; 1978.

The metabolism of benzo(a)pyrene (BP) in cultured non-malignant colonic tissue from patients with and without colon cancer was investigated. Also studied were the effects of lith

lic acid (LC) and taurodeoxycholic acid (TDC), second-bile acids, on the binding level of BP to cellular macromolecules and on the metabolite profile of BP. Cultured human colonic tissue exposed to 1.5 μ M BP for 24 hr produced two DNA-BP adducts: the major adduct (72%-100%) formed between the 10 position of BP diol-epoxide I and the 2-amino group of guanine; the minor adduct was formed between BP diol-epoxide II and the 2-amino group of guanine. The major metabolites of BP extracted with ethyl acetate/acetone from the tissue culture media were (7/8,9)-tetrahydroxy-7,10,8,9-tetrahydro-BP, trans-7,8-dihydroxy-7,8-dihydro-BP, and a peak containing (7/8,9)-tetrahydroxy-7,10,8,9-tetrahydro-BP, (7/8,9)-trihydroxy-7,8-dihydro-BP, and trans-9,10-dihydroxy-7,10-dihydro-BP. The metabolites formed by the cultured colon varied among individuals. There was a slight difference ($p < 0.1$) in binding levels between tumorous tissues from cancer patients (40) and those from noncancer patients (16). In addition, binding level of BP to DNA varied five fold among different sections of colon from the same patient, with levels being highest in the ascending colon. LC significantly ($p < 0.05$) enhanced the binding of BP to DNA at concentrations of 0.5 and 1.0 μ M, but higher concentrations decreased binding. TDC slightly enhanced binding levels in a dose-dependent manner ($p < 0.05$). Apparently, cultured human colon can metabolize BP by pathways similar to those observed in human bronchus and in cells of experimental animals. (67 refs)

78-5563 **Lack of Mutagenicity and Putative Carcinogenicity of Several Novel Benzopyrene Derivatives.** (Eng) Ashwood-Smith, M. J. (Laboratoire de Biologie Physico-Chimique, Universite de Dijon, Ecole Nationale Supérieure de Biologie, Appliquée à la Nutrition et à l'Alimentation, Dijon, France); Mitchell, R. H.; Kennedy, A. *Mutat Res* 57(2): 123-125; 1978.

Three benzopyrene derivatives with the same gross structure and the same electronic periphery as benzo(a)pyrene, but with some alteration in the complete electronic structure, were tested for mutagenicity in the Salmonella/microsome test using strains TA1537, TA100, and TA98. The three derivatives were trans-12b,12c-dimethyl-12b,12c-dihydrobenzo(a)pyrene, trans-12b,12c-dimethyl-12b,12c-dihydrobenzo(e)pyrene, and trans-13b,13c-dimethyl-13b,13c-dihydrodibenzo(b,def)-chrysene. These derivatives were completely without activity in any of the three tester strains with or without differing amounts of microsomes (from rats stimulated with phenobarbital or aroclor). Thus, it is not sufficient for a molecule to have the same electronic periphery as a known mutagen/carcinogen for it to induce mutagenicity (and probably carcinogenicity). (6 refs)

78-5564 **Reconstitution of Benzo(a)pyrene 4,5-Oxide Reductase Activity by Purified Cytochrome**

P-450. (Eng) Yamazoe, Y. (Dept. Pharmacology, Sch. Medicine, Keio Univ., 35 Shinanomachi, Shinjuku-ku, Tokyo 160, Japan); Sugiura, M.; Kamataki, T.; Kato, R. *FEBS Lett* 88(2): 337-340; 1978.

Requirements for the reconstitution of NADPH-dependent benzo(a)pyrene 4,5-oxide (BP 4,5-oxide) reductase activity were examined. Cytochrome P-450 was purified from phenobarbital-treated rat liver microsomes (PB P-450) and from 3-methylcholanthrene-treated rat liver microsomes (MC P-450 and MC P-448). Cytochrome P-450 (MC P-448) and NADPH-cytochrome P-450 reductase were essential requirements for the reconstitution of reductase activity. Lipid was also required for max activity. Using liver microsomes from intact and MC-treated rats as controls, MC P-448 in the reconstituted system exhibited 18- and 6-fold greater activity than intact and MC-treated microsomes, respectively. The addition of riboflavin (100 μ M) resulted in a 1.7-fold enhancement of BP 4,5-oxide reductase activity. BP 4,5-oxide was reduced by cytochrome P-450 even in the absence of NADPH and NADPH-cytochrome P-450 reductase if a reducing equivalent, such as reduced methylviologen, was present. In this system, MC P-448 showed the highest activity, followed by MC P-450 and PB P-450. The results suggest that both BP and BP 4,5-oxide bind to a common site in the cytochrome P-450 molecule. (15 refs)

78-5565 **Solubilization of 6-Oxybenzo(a)pyrene Radical by Caffeine and DNA as Studied by Magnetic Resonance. Observation of Intermolecular Charge Transfer.** (Eng) Mishra, K. P. (Dept. Chemistry, Faculty Science, Kyoto Univ., Kyoto 606, Japan); Nosaka, Y.; Akasaka, K.; Nagata, C.; Hatano, H. *Biochim Biophys Acta* 520(3): 679-687; 1978.

The reaction of 6-oxybenzo(a)pyrene (6-OBP) free radical (synthesized from 6-hydroxy-BP) with aqueous caffeine and DNA soln's was investigated using nuclear magnetic resonance (NMR) and electron spin resonance (ESR) spectroscopy. The ESR studies indicated that crystals of 6-OBP radical dissolved in the hydrophobic environment of caffeine molecules and existed as dispersed monomers. In aqueous DNA, 6-OBP radical was seen to bind chemically to the molecule. Comparison of NMR spin-lattice and spin-spin relaxation times in caffeine protons disclosed that a part of the 6-OBP unpaired electron was transferred from the radical to the caffeine molecule. This suggested that the reaction involves the formation of a π -bond. It is concluded that the intermolecular charge pathway is not a major source of stabilization energy of this complex. (24 refs)

78-5566 **Sulfate Conjugation of Benzo[a]pyrene Metabolites and Derivatives.** (Eng) Nemoto, N. (Dept. Experimental Pathology, Cancer Inst., Kami-Ikeburo, Toshima-ku, Tokyo, Japan); Takayama, S.; Gelboin, H. V. *Chem Biol Interact* 23(1): 19-30; 1978.

The sulfate conjugation of various metabolites and derivatives of benzo(a)pyrene (BP) was studied using rat liver 105,000 g supernatant as an enzyme source. On incubation of 3-OH BP with ^{35}S - Na_2SO_4 and a sulfate activating system, a new compound identified as a sulfate conjugate of 3-OH BP was formed. The reaction required MgCl_2 as well as ATP, indicating that the sulfate was activated to 3'-phosphoadenosine-5'-phosphosulfate prior to conjugation. The rate of 3-OH BP conjugation depended in part on the period of preincubation, a 20-min preincubation period giving maximum subsequent conjugation. All of the BP phenols and quinone metabolites formed conjugates with sulfate, whereas the dihydrodiol metabolites showed different reactivities. The sulfate conjugates were more polar than their parent BP derivatives and, except for the quinone conjugates, were easily extracted with ethyl acetate. The role of sulfate conjugation in BP carcinogenesis is discussed. (36 refs)

78-5567 Lack of Involvement of 6-Hydroxymethylation in Benzo(a)pyrene Skin Tumor Initiation in Mice. (Eng) Slaga, T. J. (Biology Div., Oak Ridge Natl. Lab., P.O. Box Y, Oak Ridge, TN, 37830); Bracken, W. M.; Viaje, A.; Berry, D. L.; Fischer, S. M.; Miller, D. R. *J Natl Cancer Inst* 61(2): 451-455; 1978.

The skin tumor-initiating activities of benzo(a)pyrene (BP), 6-hydroxymethylbenzo(a)pyrene (6-OH- CH_2 -BP), and 6-methylbenzo(a)pyrene (6- CH_3 -BP) were compared in female CD-1 mice. Further, the effects of 7,8-benzoflavone (7,8-BF), 1-benzylimidazole, and quercetin on these activities were determined. All mice were initiated with 400 or 800 nanomoles of one of the three BP compounds; 1 wk later, the mice were promoted twice weekly with 10 μg 12-O-tetradecanoylphorbol-13-acetate. BP was a much more effective tumor initiator than 6-OH- CH_2 -BP or 6- CH_3 -BP, the respective activities of these two compounds being 12.5% and 20% that of BP. The tumor-initiating activity of 6- CH_3 -BP was greater than 6-OH- CH_2 -BP, but only the latter exhibited a dose-response relationship. The administration of 10, 100, and 200 μg of 7,8-BF 5 min before BP resulted in a 20%, 27%, and 32% inhibition of papillomas per mouse, respectively. The number of mice with tumors was only slightly inhibited by the two higher doses of 7,8-BF. This compound had no effect on 6-OH- CH_2 -BP and only slightly inhibited 6- CH_3 -BP tumor initiation. Quercetin and 1-benzylimidazole (both, 100 μg) inhibited the number of BP-initiated papillomas per mouse by 28% and 40%, respectively. 1-Benzylimidazole + 7,8-BF 5 min before BP resulted in 40% inhibition. These results indicate that direct 6-hydroxymethylation is probably not an important pathway for the activation of BP in mouse skin. (35 refs)

78-5568 Atmospheric Reactions of Polycyclic Aromatic Hydrocarbons: Facile Formation of Mutagenic Nitro Derivatives. (Eng) Pitts, J. N. (Statewide Air Pollution

Res. Center, Dept. Chemistry, Univ. California, Riverside, CA, 92521); Van Cauwenberghe, K. A.; Grosjean, D.; Schmid, J. P.; Fitz, D. R.; Belser, W. L.; Knudson, G. B.; Hynds, P. M. *Science* 202(4367): 515-519; 1978.

In order to test the hypothesis that oxidative reactions of polycyclic aromatic hydrocarbons in the atmosphere lead to the formation of directly active mutagens, two sets of experiments were conducted: one in which the carcinogenic benzo(a)pyrene and the noncarcinogen perylene were exposed to pollutant gases under simulated atmospheric conditions (including 1 ppm NO_2 and traces of nitric acid) and another in which BP was exposed to the gases in ambient photochemical smog (ppm levels of NO_2 , O_3 , and peroxyacetyl nitrate). After exposure, the reaction products were separated and purified by thin layer chromatography and analyzed by mass spectrometry. The products were then tested for mutagenic activity with *Salmonella typhimurium* strains TA1535, TA1538, TA98, and TA100, with and without the addition of 0.5 ml rat liver S9 mix as the metabolic activator. Results of the Ames reversion assay revealed that directly active mutagenic nitro derivatives were readily formed from both perylene in stimulated atmospheres and that directly active mutagens were formed from BP in photochemical smog. The formation of direct mutagens by similar reactions on samples collected on air pollution collection filters, in exhaust effluents, and in the atmosphere is possible. (36 refs)

78-5569 Cytologic Correlates of Chemical Carcinogenesis in Respiratory Tracts of Dogs. (Eng) Muro, K. (Chiba Univ. Sch. Medicine, Inst. Pulmonary Medicine, Chiba, Japan); Cohen, A. H.; Shors, E. C.; Iversen, J. R. *J Natl Cancer Inst* 61(5): 1209-1214; 1978.

Serial cytologic studies were made of 54 dogs during respiratory carcinogenesis induced by benzo(a)pyrene (BP) and N-methyl-N-nitrosourea (NMU). Dogs received submucosal injections of either 45 mg BP coupled with ferric oxide or 45 mg NMU to total doses of 810 and 810-945 mg, respectively. Another group received 21 weekly submucosal injections of 15 mg BP followed by 27 weekly doses of 10 NMU for a total of 315 mg BP and 270 mg NMU. Abnormal cells appeared within 1 mo after the first carcinogen injection. In the early stage, BP induced changes primarily in the cytoplasm. These changes consisted of increased size, basophilia, and vacuolization. NMU induced primarily nuclear enlargement and enhanced the prominence of the nucleolus. After 6 mo, dogs exposed to BP demonstrated pleomorphism of the cytoplasm and nucleus, but cells exposed to NMU formed bizarre spindle-shaped cells. Cells were generally more scattered from one another following BP treatment than after NMU treatment. When the mucosa was exposed to BP + NMU, cytologic features became those characteristic of lesions induced by NMU from the outset. These serial cytologic studies of bronchial brushings provided insights that repetitive biopsies did not, but biological significance of the observed stereotypic patterns to cytologic morphogenesis is not clear. (35 refs)

78-5570 **Ultrastructure of Early Stages in N-Diethylnitrosamine and Benzo(a)pyrene Induced Lung Carcinogenesis.** (Eng) Reznik-Schuller, H. (Abteilung für Experimentelle Pathologie, Medizinische Hochschule Hannover, 3000 Hannover 61, W. Germany); Mohr, U. In: *Detection and Prevention of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1976. pp. 1315-1318; 1978.

The early effects of N-diethylnitrosamine (DEN) and benzo(a)pyrene (BP) in the lungs of Syrian golden hamsters were examined by electron microscopy. Forty animals were injected sc with DEN (1/10 LD₅₀, 2x/wk), and 20 were given intratracheal instillations of BP (0.63 mg/animal, 1x/wk). Animals were sacrificed at regular intervals up to 20 wk of treatment. Following DEN treatment, nonciliated cells frequently contained lamellated inclusion bodies in their cytoplasm, and proliferation of both nonciliated and APUD (amine precursor uptake and decarboxylase activity) cells was observed. In addition, squamous metaplasias developed in the par bronchi. None of these effects were observed after BP administration. The earliest BP-induced alterations occurred after 3 wk in ciliated cells, and they were either a reduction in the number of cilia per cell or a deformation of cilia. These alterations occurred late in DEN treatment. Penetration of the basement membrane, with subsequent development of adenomas, was induced by both carcinogens. These findings indicate that in bronchial epithelia, very early cellular reactions to carcinogen treatment vary according to the carcinogen and its mode of administration but that the later steps are similar. (10 refs)

78-5571 **Effects of Exposure to Furfural Vapour in Hamsters Simultaneously Treated with Benzo(a)pyrene or Diethylnitrosamine.** (Eng) Feron, V. J. (Central Inst. Nutrition and Food Res. TNO, Zeist, Netherlands); Kruysse, J. *Toxicology* 11(2): 127-144; 1978.

Syrian golden hamsters were exposed to furfural vapor (0 or 50/400 ppm for 7 hr/day) 5 days/wk for 52 wk to study the effects of long-term repeated exposures. Some animals were simultaneously given intratracheal benzo(a)pyrene (BP, 0.2 or 36.4 mg total) weekly for 52 wk or sc diethylnitrosamine (DENA, 2.1 µl total) every third wk for 52 wk. The study was terminated after 81 wk. The animals' coats became yellowish-brown during furfural treatment, and body wts decreased relative to control animals. The death rate was not affected by furfural or clearly influenced by BP or DENA. There were no significant differences in hematological or biochemical findings or organ-to-body-wt ratios between furfural-treated and control hamsters. Furfural exposure caused irritation of the nasal mucosa, atrophy and downward growth of sensory cells of the olfactory epithelium, degenerative

changes in Bowman's glands, and the occurrence of cyst-like structures in the lamina propria beneath the olfactory epithelium. Furfural did not affect the tumorigenicity of BP or DENA. It is concluded that furfural is not a significant cofactor in respiratory tract carcinogenesis. (10 refs)

78-5572 **Relative Carcinogenic Effectiveness of Derivatives of Nitrosodiethylamine in Rats.** (Eng) Lijinsky, W. (Chemical Carcinogenesis Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Taylor, H. W. *Cancer Res* 38(8): 2391-2394; 1978.

The carcinogenicity of five of nitrosodiethylamine derivatives, administered po to Sprague-Dawley rats, was compared with that of the parent compound. All were less potent than nitrosodiethylamine. When nitrosobis(2-methoxyethyl)amine and nitrosobis(2-ethoxyethyl)amine were administered at equimolar doses (0.68×10^{-3} M) in drinking water, there was a high incidence of liver tumors, but the animals survived longer than after nitrosodiethylamine treatment, which also induced esophageal tumors. Nitrosoiminodipropionitrile and nitrosobis(2,2-diethoxyethyl)amine failed to induce tumors at the same doses. Nitrosobis(2-chloroethyl)amine was administered in oil by gavage at a dose (1.15 mg 2x/wk) lower than that of nitrosodiethylamine, and produced a much weaker tumor response: 5/15 treated rats had forestomach papillomas and 1 had an olfactory adenocarcinoma. (17 refs)

78-5573 **Carcinogenesis of N-Nitrosodiethylamine (DENA) in Chickens and Domestic Cats.** (Eng) Schmahl, D. (Institut für Toxikologie und Chemotherapie, Deutsches Krebsforschungszentrum, im Neuenheimer Feld 280, 6900 Heidelberg, W. Germany); Habs, M.; Ivankovic, S. *Int J Cancer* 22(5): 552-557; 1978.

The carcinogenicity of N-nitrosodiethylamine (DENA) was investigated in two new animal models, the chicken and cat, and reviewed in 16 other animal species. Six White Leghorn chickens were inoculated im 2x/wk with 10 mg/kg DENA for 50 wk (median total dose, 1,000 mg/kg) and five domestic short-haired cats were fed a 10% DENA soln in milk 5x/wk (daily dose, 1.5-2 mg/kg; median total dose, 870 mg/kg). All test animals showed severe toxic injury to the liver. In the chickens, median survival was 518 days. Three of the six chickens developed tumors: 1 developed multiple benign adenomas of the kidney; 1 a hepatocellular carcinoma and a pheochromocytoma of the adrenal gland; and 1 a hepatocellular carcinoma and an adenocarcinoma of the kidney. In the cats, median survival was 545 days. Three cats died with hepatocellular carcinomas; one of these animals also had multiple cholangiogenic adenomas and an early cholangiocellular carcinoma, and another showed peliosis hepatis and a cavernous hemangioma. DENA is carcinogenic in all species in which it has been tested adequately, with the liver proving

to be the main target organ. It is highly probable that DENA will act as a carcinogen in humans. (49 refs)

- 78-5574 Enhancement by WY-14,643, A Hepatic Peroxisome Proliferator, of Diethylnitrosamine-Initiated Hepatic Tumorigenesis in the Rat.** (Eng) Reddy, J. K. (Dept. Pathology, Northwestern Univ. Medical Sch., 303 East Chicago Ave., Chicago, IL, 60611); Rao, M. S. *Br J Cancer* 38(4): 537-543; 1978.

The hepatic peroxisome proliferator (4-chloro-6-(2,3-xylidino)-2-pyrimidylthio)acetic acid (WY-14,643) enhanced the development of liver tumors in male F-344 Charles River rats given diethylnitrosamine (DEN: 100 parts/10⁶) in drinking water for 14 days. Beginning 1 wk after DEN treatment, rats in groups of 30 animals each were fed diets containing 0.1% WY-14,643 or 0.5% clofibrate (CFB) for 48 wk. Laparotomies at 22 wk revealed grossly visible liver nodules in only an occasional rat fed CFB, compared with approx 60% of rats fed WY-14,643. By 48 wk all 28 surviving WY-14,643 treated rats had developed multiple tumors, 2-35 mm in diameter, involving all lobes. All tumors > 10 mm were hepatocellular carcinomas; two rats also developed kidney adenomas. In the CFB group, 25/28 animals had liver tumors. The tumor incidence in this group was significantly greater than that (5/18) in rats receiving a control diet after DEN, but it was significantly less than that in the WY-14,643 group. Although the enhancement of liver tumorigenesis by WY-14,643 may be due to its hepatomegalic and peroxisome proliferative properties, the final mechanism remains to be elucidated. (33 refs)

- 78-5575 The Influence of Diazepam and Thiouracil upon the Carcinogenic Effect of Diethylnitrosamine in Gerbils.** (Eng) Green, U. (Abteilung für Experimentelle Pathologie, Medizinische Hochschule Hannover, Karl-Wiechert-Allee 9, D-3000 Hannover-Kleefeld, W. Germany); Ketkar, M. *Z Krebsforsch* 92(1): 55-62; 1978.

The influence of diazepam (DZP) and thiouracil (TU) on N-diethylnitrosamine (DEN)-induced carcinogenesis was examined in gerbils. One group of gerbils received DZP (10 mg intragastrically 1 x/wk for life) and, 30 min later, DEN (23 mg/kg sc 1 x/wk for life). Another group was fed TU daily (2,000 ppm in the diet) + DEN. Other groups received DEN, DZP, or TU alone. Animals receiving DZP + DEN or TU + DEN had significantly higher av survival times (53%-80% and 43%-90% higher, respectively) than gerbils receiving DEN alone. This longer av survival time resulted in a significantly higher total dosage of DEN. However, similarly high incidences of tumor (mainly adenocarcinomas), originating in the nasal cavities were seen in the DEN alone and combined treatment groups. Hepatic neoplasms were found as early as 15 wk after the start of treatment with DEN alone, but these tumors were first observed after 40 wk in the

DZP + DEN animals and after 44 wk in the TU + DEN group. DEN alone selectively induced well-differentiated cholangiocarcinomas, whereas cholangiomas were mainly found in animals given the combined treatments. No tumors were found in animals treated with DZP or TU alone. DEN has to be activated enzymatically to form its proximate and/or ultimate carcinogen. The results indicate that this metabolism is inhibited in gerbils by the simultaneous administration of DZP or TU. (13 refs)

- 78-5576 Glucocorticoid Binding and Response in Liver and Diethylnitrosamine (DENA)-Induced Hepatomas.** (Eng) Snart, R. S. (Dept. Zoology, The University of Sheffield, S10 2TN, England); Thorne, J. *J Steroid Biochem* 9(8): 709-715; 1978.

The binding of ³H-dexamethasone (³H-dex) in normal and DENA-treated rat liver was studied. Wistar rats were given DENA (50 mg/ml) in their drinking water beginning at age 6 wk. Liver preparations were obtained from normal and DENA-treated rats at 4, 8, 12, and 17 wk after the start of DENA treatment. The binding of ³H-dex to rat liver cytosol, plasma, and nuclei was determined, as was the binding of the ³H-dex-receptor complex to DNA. There were no differences between normal rat liver and DENA-induced hepatomas with regard to glucocorticoid binding to receptor proteins and translocation of the receptor-steroid complex into the nucleus or the ability of the complex to bind to DNA. Tyrosine amino transferase (TAT) activity in rat livers was refractory to corticosterone and dexamethasone treatments, but following adrenalectomy, a highly significant response was observed. The DENA-induced hepatomas had slightly reduced levels of TAT activity, but this enzyme was stimulated significantly by corticosterone and very well by dex. The plasma corticosterone concentrations in DENA-treated rats were significantly lower than normal concentrations. It is suggested that the resulting feedback inhibition due to unmetabolized glucocorticoid decreases ACTH release and seriously affects adrenal function. A decreased steroid metabolism in DENA-treated rat livers could also account for the increased tissue sensitivity to the hormone. (16 refs)

- 78-5577 Chemical Studies on Carcinogenic Nitrosamines. 1. Hydrolysis of α -Acetoxynitrosamines.** (Eng) Baldwin, J. E. (Dept. Chemistry, Massachusetts Institute of Technology, Cambridge, MA, 02139); Scott, A.; Branz, S.; Tannenbaum, S. R.; Green, L. *J Org Chem* 43(12): 2427-2431; 1978.

Chemical reactions of the two structural isomers of α -acetoxynitrosamines, one of which is a powerful bacterial mutagen whereas the other is inactive, were studied. Nucleophilic attack at the α carbon resulted in hydrolysis to methanolysis, but attack at the carbonyl carbon by n-propyl

e.g., yielded N-n-propylacetamide and the unstable α -acetoxy-nitrosamine. The hydrolysis and methanolysis appear to occur via an SN1 reaction, involving a resonance-stabilized benzylnitrosiminium ion. The deacylation of primary amines was strongly activated by the nitrosamino group, probably via an inductive effect, and it did not involve a cyclic intermediate. (12 refs)

578 Oxidative Decarboxylation of Nitrosoamino Acids: A Synthetic Approach to Cyclic α -Acetoxy-nitrosamines. (Eng) Saavedra, J. E. (NCI Frederick Cancer Center, Frederick, MD, 21501). *Tetrahedron Lett* 1923-1926; 1978.

Primary studies of the one-step synthesis of α -acetoxy-nitrosopyrrolidine and α -acetoxy-nitrosopiperidine by oxidative decarboxylation of the corresponding nitrosoamino acids and lead tetraacetate are reported. Since α -acetoxy-nitrosamines may be direct-acting carcinogens, they should be treated with caution. (14 refs)

579 The Fragmentation Reaction of β -Hydroxynitrosamines: Possible Environmental and Biological Significance for Nitrosamine Carcinogenicity. (Eng) Loeppky, R. N. (Dept. Chemistry, Univ. Missouri-Columbia, Columbia, MO); Christiansen, R. *IARC Sci Publ* 17(19): 117-126; 1978.

Ability of β -hydroxynitrosamines (OH-NA's) to undergo acid-catalyzed fragmentation (ie, retroaldol-type, cleavage and decarboxylation) into an NA and a carbonyl product was determined. Six OH-NA's were synthesized, and their behavior toward strong base (3 M KOH) was examined. All six compounds underwent fragmentation, three with high reactivity, and three with low reactivity. The reverse reaction (aldol-type condensation) is prevented by the occurrence of Cannizzaro-type condensation of the carbonyl product. This fact may account for the low reactivity of three of the OH-NA's, the nature of their carbonyl products prevented or made it difficult for them to enter into condensation reactions. It is suggested that fragmentation and condensation could occur while NA's are being isolated from substances such as food products. Severe alkaline conditions should be avoided in the isolation and analysis of NA's from natural sources, consideration must be given to the possibility that fragmentation may have occurred in past work. The fragmentation may have a biochemical counterpart that is significant in the biochemical mechanism of NA carcinogenesis. (17 refs)

5580 Effect of Dose on the Carcinogenic Activity of Orally Administered N-Bis(2-hydroxypropyl)nitrosoamine in Rat. (Eng) Konishi, Y. (Dept.

Oncological Pathology, Cancer Center, Nara Medical Univ., 840 Shijo-cho, Kashihara 634 Japan); Kondo, H.; Ikeda, T.; Kawabata, A.; Shoji, Y.; Denda, A. *Gann* 69(4): 573-577; 1978.

Male Wistar rats were given 100 ppm N-bis(2-hydroxypropyl)nitrosoamine (DHPN) in drinking water and sacrificed at 25 wk (Group 1) or 52 wk (Group 2), or they were given 500 ppm DHPN and sacrificed at 25 wk (Group 3) or 29-39 wk (Group 4). Two lung adenomas occurred among 12 Group 1 animals. Three hepatocellular carcinomas, 1 liver hemangioma, 3 thyroid adenomas, and 6 lung adenomas, 3 adenocarcinomas, and 1 squamous cell carcinoma of the lungs were found among 10 Group 2 rats. All 12 Group 3 rats developed lung adenomas; in addition, 2 developed a lung adenocarcinoma and 4, a squamous cell carcinoma. Nine surviving animals in Group 4 developed numerous tumors in various tissues, including 22 lung tumors, 7 liver tumors, 1 thyroid tumor, 5 esophageal tumors, 2 kidney tumors, and 1 bladder tumor. Therefore, lung tissue was affected most severely by this carcinogen; the fact that no pancreatic tumors were identified is interesting, because DHPN is a potent pancreatic carcinogen in Syrian golden hamsters. (20 refs)

78-5581 A Cell Line Derived from BBN [N-Butyl-N-(4-hydroxybutyl)nitrosamine]-induced Rat Bladder Cancer: Establishment and Scanning Electron Microscopic Cell Surface Characteristics. (Eng) Nishi, M. (Dept. Urology, Okayama Univ. Medical Sch., Okayama 700, Japan). *Acta Med Okayama* 32(3): 181-205; 1978.

The surface structure of bladder cancer cells was studied by scanning electron microscopy using synchronized cultures of lines cloned from rat bladder cancers induced by N-butyl-N-(4-hydroxybutyl)nitrosamine. Tumors were induced in male Wistar rats raised with drinking water containing 0.05% of the carcinogen. After several subcultures, the cells exhibited loss of contact inhibition and marked piling-up behavior. Because single-cell cloning was difficult, colonies of cells were cloned. The population doubling time of the parent strain and the clones was approx 30 hr. The chromosome mode was in the triploid to tetraploid range. Plating efficiency was <20%. Backtransplantation of 5×10^6 cells ip into newborn rats produced tumors in all cases. Histopathologically, some parts of these tumors resembled primary transitional cell carcinoma. However, the major tumor cell groups showed marked keratinization and resembled squamous cell carcinoma. Compared with normal bladder mucosa cells, the nucleus cytoplasm ratio and the number of nucleoli, free ribosomes, and intracytoplasmic microfibrils were increased. The ultrastructure was characterized by a profuse arrangement of microvilli, parts of which were joined to desmosomes. During mitosis, the cells became large and globular; the microvilli decreased in height and were gathered profusely over the entire surface. Cells in G₁-S developed lamellipodia and pseudopodia-like cytoplasmic processes and were polygonal. Microvilli were present in the central part

containing the nucleus, but they were somewhat decreased in number and increased in height. (37 refs)

- 78-5582 Inhibitory Effect of 13-*cis*-Retinoic Acid on Urinary Bladder Carcinogenesis Induced in C57BL/6 Mice by N-Butyl-N-(4-hydroxybutyl)nitrosamine.** (Eng) Becci, P. J. (Life Sciences Div., IIT Res. Inst., Chicago, IL, 60616); Thompson, H. J.; Grubbs, C. J.; Squire, R. A.; Brown, C. C.; Sporn, M. B.; Moon, R. C. *Cancer Res* 38(12): 4463-4466; 1978.

The effect of 13-*cis*-retinoic acid (CRA) on the induction of highly invasive urinary bladder carcinoma by N-butyl-N-(4-hydroxybutyl)nitrosamine (OH-BBN) was studied in male C57BL/6 mice. Tumors were induced by gastric intubation of OH-BBN in doses of 7.5 or 10.0 mg 2 x/wk for 6 or 7 wk, respectively. Seven days after the last carcinogen intubation, mice were fed a placebo diet or a diet supplemented with 200 mg/kg CRA. Autopsies performed 6 mo following the first carcinogen treatment revealed highly invasive squamous and transitional cell carcinomas of the urothelium. In most cases, tumor cells had penetrated the bladder muscle wall. The CRA diets reduced the incidence of carcinomas and noninvasive papillomas, as well as the extent of neoplastic development in the bladder, induced by either dose of OH-BBN. At the lower dose of OH-BBN, CRA prevented the appearance of both squamous and transitional cell carcinoma, with a reduction of incidence from 33% to 0%. These findings suggest that CRA can reduce both the severity and the incidence of chemically induced urinary bladder carcinomas. (8 refs)

- 78-5583 Effect of Protease Inhibitors in Mice Treated with the Hepatocarcinogen, Hexachlorocyclohexane (α -Isomer) and the Bladder Carcinogen, N-Butyl-N-(4-hydroxybutyl)nitrosamine.** (Eng) Arai, H. (First Dept. Pathology, Nagoya City Univ. Medical Sch., 1 Kawasumi, Mizuho-cho, Mizuho-ku, Nagoya 467, Japan); Shirai, T.; Hananouchi, M.; Hirose, M.; Murasaki, G.; Ito, N. *Gann* 69(4): 593-595; 1978.

One of the protease inhibitors pepstatin (PS), leupeptin (LP), and antipain (AP) (each, 0.1% in diet) were administered to male DDY mice that had been fed the α -isomer of 1,2,3,4,5,6-hexachlorocyclohexane (α -BHC) (500 ppm for 24 wk) or to male ICR mice that had been given N-butyl-N-(4-hydroxybutyl)nitrosamine (BBN) (0.05% of drinking water for 8 wk). Animals treated with LP + α -BHC experienced a significant increase in hepatic nodular hyperplasia (HNH: 5/12 vs 1/13 in α -BHC-only mice), and they developed the only α -BHC-induced hepatoma. Groups given α -BHC and the other two protease inhibitors did not show any significant differences in HNH from the α -BHC-only group. In all BBN + protease inhibitor groups, the urinary bladder changes were similar to those of the BBN-only group:

7 simple hyperplasias, 4 papillary or nodular hyperplasias, 1 papilloma, 1 cancer, and no hemangiomas among 13 mice. Hemangiomas did occur in the BBN + PS (3/10) BBN + LP (1/16) groups. Treatment with protease inhibitor alone in both experiments did not produce any changes. Thus, it appears that LP enhances the induction of NBN + α -BHC. (10 refs)

- 78-5584 Cell Kinetics of Mouse Urinary Bladder Epithelium. VII. Changes in Proliferation and Nuclear DNA Content after Repeated Doses of Dibutyltinamine.** (Eng) Farsund, T. (Haukeland Sykehus, N-5015 Bergen, Norway); Iversen, O. H. *Virchows Arch [Cell Pathol]* 27(2): 119-135; 1978.

The effect of repeated sc injections of dibutyltinamine (DBN; 0.01 ml in 0.2 ml propylene glycol) on the cell kinetics of the urinary bladder epithelium was investigated in hairless hr/hr Oslo strain mice. The animals were divided into two groups: Group 1 received the injections three times weekly for 3 mo; Group 2 received the injections once a week for 6 mo. A high mortality (24/72) was noted in Group 1, no Group 2 animals died during the observation period. DNA synthesis in the tetraploid cells of both groups was inhibited early in the experimental period; this inhibition was followed by a disappearance of the octaploid and, finally, tetraploid cells. In the diploid cells, however, the labeling index, the proportion of cells in the S phase, and the mitotic rate increased. Thus, repeated DBN injections led to a hyperplasia of the diploid cells due to a high rate of diploid proliferation and a block in the mechanism of repeated DNA synthesis. Malignant transformation was noted in Group 1 after 5-6 mo. These changes in urothelial cell turnover were qualitatively similar to, but more pronounced, than those induced by injections of cyclophosphamide or propylene glycol. It is suggested that the strong and early effect of DBN on diploid, basal cell DNA synthesis and on the mechanism of repeated DNA synthesis in the bladder mucosa is linked to the carcinogenicity of DBN. (22 refs)

- 78-5585 Ultrastructure of Nitrosoheptamethyleneimine-induced Lung Tumors in European Hamsters.** (Eng) Reznik-Schuller, H. (Chemical Carcinogenesis Section, Frederick Cancer Res. Center, P. O. Box B, Frederick, MD, 21501). *Am J Pathol* 93(1): 45-52; 1978.

The ultrastructure of nitrosoheptamethyleneimine (NHMI)-induced lung tumors (LT) in European hamsters (EH) is described and compared with that of tumors induced in the same species and in Syrian golden hamsters by other nitrosamines. Ten male EH were given weekly sc injections of 66 mg/kg NHMI until they showed clinical signs of toxicity. All animals developed nasal cavity tumors and multiple lung tumors. Light microscopically, the LT showed an adenomatous mixed (areas with adenoid and poorly keratinized squamous

erentiation) pattern. Electron microscopically, the tumorous neoplasms (adenomas and adenocarcinomas) in the areas of adenoid differentiation in the mixed tumors consisted predominantly of Clara cells and of cells with bellated inclusion bodies and various amounts of smooth endoplasmic reticulum in their cytoplasm. Areas of squamous differentiation contained cells in which a gradual loss of dense-cored granules coincided with an increasing occurrence of cytoplasmic filaments. These LT were similar to those induced in EH by N-dibutyl nitrosamine and to those induced in Syrian golden hamsters by diethylnitrosamine, N-butyl nitrosamine, and nitrosomorpholine. (20 refs)

5586 **Autoradiography and Organ Distribution of N-Methyl-N-nitrosobenzylamine in Rats.** (Eng) Hidaka, T. (Serology Div., Natl. Cancer Center Res. Inst., Tsukiji, 5-1-1, Chuo-ku, Tokyo 104, Japan); Ichimura, S.; Sawachi, T. *Gann* 69(4): 487-492; 1978.

The mechanism of esophageal carcinoma induction by N-methyl-N-nitrosobenzylamine (MNBA) or by N-methyl-N-nitrosamine (MBA) + sodium nitrite, which form MNBA in stomach acid, was studied in male Donryu rats. The distribution of labeled MNBA was determined by autoradiography. One group of rats was given a soln of 120-150 mg/kg MNBA + 77-100 mg/kg NaNO₂ intragastrically. Four groups were given 83 mg/kg MNBA ip, im, sc, and intragastrically, respectively. All groups of rats developed esophageal ulcers within 1-4 days. Those produced by MNBA were deeper and more severe than those produced by MBA + NaNO₂. Whole-body autoradiography showed that the radioactive label was high in the esophagus and liver of rats inoculated with labeled MNBA; the radioactivity persisted in these tissues at 24 hr. Incorporation of label was higher in the esophagus than in any other part of the gastrointestinal tract. Microautoradiography displayed a concentration of silver grains in both esophagus and liver, particularly in the mucous membrane of the esophagus. Thus, MNBA has a specific affinity for the esophagus, eventually resulting in the development of carcinoma. (10 refs)

5587 **Nitrosobenzylmethylamine, Zinc Deficiency and Oesophageal Cancer.** (Eng) Fong, L. Y. (Massachusetts Inst. Technology, Cambridge, MA); Newberne, P. M. *IARC Sci Publ* (19): 503-513; 1978.

The effects of zinc deficiency on esophageal cancer induction by nitrosobenzylmethylamine (NBMA) were investigated by feeding weanling Charles River CD strain male rats a basal zinc-deficient (Zn-) diet containing 7 ppm Zn (30 rats) or a control diet containing 60 ppm Zn ad libitum (20 rats; ad libitum-fed controls). After 12 wk (at 15 wk of age), NBMA (2 mg/kg 2 x/wk) was administered by intragastric intubation for 12 wk. In a second experiment, 40 rats received a Zn-deficient diet for 4 wk and 40 rats served as pair-fed controls. After

4 wk (at 7 wk of age), NBMA was administered as above. Zn determinations were made in the serum, hair, esophagus, testes, liver, and kidneys; Zn- rats were retarded in growth compared with ad libitum-fed controls. Furthermore, pair-fed controls, although of comparable size, did not show the hair deficiencies and dermal lesions common in Zn- rats. In the first experiment, 43% of the Zn- rats developed esophageal carcinoma vs 15% of the ad libitum-fed controls. In the second experiment, 83% and 33% of the Zn- animals (sacrificed at 58 and 93 wk, respectively) developed esophageal carcinoma vs 0% of the pair-fed controls. Tumor induction time was shorter in all Zn- rats. Zn levels in serum, hair, testis, and esophagus were depressed in all Zn- animals, although liver levels were unchanged in all groups. It is suggested that Zn deficiency enhances the induction of esophageal cancer in the rat. (17 refs)

78-5588 **Mode of Mutagenic Action of Methyl-N-nitrosocyanamide, a Potent Carcinogen.** (Eng) Hidaka, K. (Cancer Res. Inst., Faculty Medicine, Kyushu Univ., Fukuoka 812, Japan); Ishizawa, M.; Otsuji, N.; Endo, H. *Cancer Res* 38(12): 4630-4633; 1978.

To study the mode of mutagenic action of methyl-N-nitrosocyanamide (MNC), N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), and N-methyl-N-nitrosourea (MNU), revertants induced by treatment of a strain of *Escherichia coli* carrying nonsense mutations in the tryptophan (*trp*) and alkaline phosphatase (*phoA*) genes were analyzed. Trp+ revertants were purified and classified into seven categories based on their ability to support the growth of particular nonsense mutants of phage λ and on their content of alkaline phosphatase. About 90% of the trp+ revertants induced by MNC were due to mutations in suppressor genes, and 85% of the suppressor mutations occurred in the suppressor gene *supE*. Intragenic reversion cannot occur by a GC(to)AT base-substitution mutation, whereas this is the obligate mode of mutation in gene *supE*. It is concluded that MNC preferentially induces GC(to)AT transition mutations but that other base-substitution mutations are also induced at about 10% of this frequency. NMU and, particularly, MNNG, also preferentially induce GC(to)AT transition mutations. (25 refs)

78-5589 **Mutagenicity of N-Nitrosopiperazine Derivatives in *Salmonella typhimurium*.** (Eng) Rao, T. K. (Biology Div., Oak Ridge Natl. Lab., Post Office Box Y, Oak Ridge, TN, 37830); Young, J. A.; Lijinsky, W.; Epler, J. L. *Gen Pharmacol* 57(2): 127-134; 1978.

A series of nitrosopiperazine derivatives was tested for mutagenicity in the *Salmonella typhimurium*/microsomal activation system, using strains TA1535, TA100, TA1537, TA158, and TA98. Compounds tested were N-nitrosopiperazine, N-methyl-N-nitrosopiperazine, dinitrosopiperazine, 2-methyldinitrosopiperazine, 2,5-dimethyldinitrosopiperazine,

zine, 2,6-dimethyldinitrosopiperazine, and 2,3,5,6-tetramethyldinitrosopiperazine, dinitrosohomopiperazine, and 4-benzoyl-3,5-dimethylnitrosopiperazine. These compounds required metabolic activation; max activity was obtained with phenobarbital-induced rat liver microsome enzymes. There was a good correlation between a positive effect in the mutation assay and the known carcinogenic potency of the compound. These results demonstrate the predictive value of an in vitro activation system using *S. typhimurium* to detect carcinogenic compounds as mutagens. (14 refs)

- 78-5590 Metabolism of Acyclic and Cyclic N-Nitrosamines by Cultured Human Colon.** (Eng) Autrup, H. (Human Tissue Studies Section, Experimental Pathology Branch, Carcinogenesis Program, NCI, Bethesda, MD, 20014); Harris, C. C.; Trump, B. F. *Proc Soc Exp Biol Med* 159(1): 111-115; 1978.

The metabolism of several acyclic and cyclic N-nitrosamines was studied in human colon mucosa cultures derived from 11 patients, 7 with and 4 without cancer of the colon. Nitrosamine metabolism was measured by ^{14}C - CO_2 formation and reaction of the activated moieties with cellular DNA and protein. Dimethylnitrosamine (DMN) and N-nitrosopyrrolidine (NPy) were metabolized by explants from all patients studied. Diethylnitrosamine (DEN) was metabolized by explants from two patients, N',N'-dinitrosopiperazine (DNP) by five, and N-nitrosopiperidine by one. No ^{14}C - CO_2 was formed from N-nitrosornicotine. A positive correlation between binding of DMN to DNA and CO_2 formation was observed. DMN alkylated DNA in both the O-6 and N-7 positions of guanine. However, most of the radioactivity was associated with an acid-labile compound. High binding levels of DNP to protein without concomitant binding to DNA were detected. Interindividual variation in both binding level to DNA and ability to metabolize the different N-nitrosamines was observed. These findings suggest that the colon should be considered one of the organs that are likely to be susceptible to the carcinogenic action of N-nitrosamines. (36 refs)

- 78-5591 Demonstration of the Presence of Nitrosamine in Human Urine: Preliminary Observations on a Possible Etiology for Bladder Cancer in Association with Chronic Urinary Tract Infection.** (Eng) Hicks, R. M. (Sch. Pathology, Middlesex Hosp. Medical Sch., London, England); Gough, T. A.; Walters, C. L. *IARC Sci Publ* (19): 465-475; 1978.

In view of the ability of nitrate-reducing bacteria to synthesize N-nitrosodimethylamine (DMNA) from precursors, a study was undertaken to determine whether human urine contains elevated levels of nitrosamines (NA's) during bacterial cystitis of the bladder and whether there is any connection between these elevated levels and subsequent bladder cancer. Urine was collected from hemi- and paraplegic Brit-

ish men, British bladder cancer patients, Egyptians with bladder cancer and superimposed bilharzial bladder disease (BBD), Egyptians with uncomplicated BBD, British controls, and Egyptian controls. Total NA content was determined by optical density measurements, volatile NA's by gas chromatography-mass spectrometry. NA's were present in 3/10 paraplegics, all three with active bacterial infection, and in the single British cancer patient with active bacterial infection. Elevated levels of urinary NA's were detected in 4/15 Egyptian bladder cancer patients with BBD. Their urine was not examined for bacteria, but infection in these patients is commonplace. Six of 10 Egyptians with uncomplicated BBD, 7 of whom had infected urine, showed traces of NA's, but these levels were much lower than in the cancer patients with BBD. The NA levels in Egyptian bladder cancer patients were much higher than those found in the British patients, which suggests that there is an unidentified factor in the urine of the former that promotes nitrosation. The only NA positively identified was DMN, but strong evidence was obtained that other extractable, not yet identified, NA's were also present. The results suggest that NA's were only in urine in which bacterial infection was established or commonplace. It is suggested that bacterial enzymes in infected patients reduce nitrate to nitrite, which can then nitrosate available precursors to potentially carcinogenic NA's. (12 refs)

- 78-5592 Repeated Partial Hepatectomy as a Prominent Stimulus for Carcinogenic Response of Liver to Nitrosamines in Rats.** (Eng) Pound, A. W. (Dept. Pathology, Univ. Queensland, Brisbane, Australia); McGuire, L. *J Cancer* 37(4): 585-594; 1978.

The effect of partial hepatectomy (PH) on the tumorigenic response of nitrosamine-treated male Sprague-Dawley rats was investigated. Rats were given a two-thirds PH followed by 100 mg/kg diethylnitrosamine (DEN), 20 mg/kg diethylnitrosamine, or 30 mg/kg ethylmethylnitrosamine. Compared with control rats (laparotomy alone plus nitrosamines), this treatment increased both the number of liver tumors and the number of focal proliferations. In a second experiment, rats (Group I) were given a hepatonecrotic dose of 1.0 mg/kg CCl_4 followed 24 hr later by a dose of 100 mg/kg DEN. Five, 10, and 15 wk later, these rats underwent three PH's. Three additional groups received CCl_4 + 1 PH, the three PH's, or no treatment, respectively. The number of tumors in surviving animals was greatest in Group I rats, but this result was not significant. The yields of focal proliferations and cell foci could not be compared because of variation in areas of the liver scanned. The incidence of focal proliferations was significantly greater in Group I animals following the second PH (at 15-20 wk), but the increase found in necropsy specimens (at 52 wk) was not statistically significant. If the single dose of DEN is regarded as an initiating agent, the stimulus of repeated PH appears to act as a promoting agent for hepatocarcinogenesis. (35 refs)

5593 **Microsome-mediated Mutagenesis of a Chinese Hamster Cell Line by Various Chemicals (Meeting Abstract).** (Eng) Drevon, C. (International Agency for Cancer, 150 cours Albert Thomas, 69008 Lyon, France); Hiroki, T.; Montesano, R. *Mutat Res* 53(2): 181-182; 1978. (10 refs)

5594 **A Comparison of Ascorbic Acid Excretion with Other Indicators of Nitrosamine Hepatotoxicity.** (Eng) Nishie, K. (Richard B. Russell Agricultural Research Center, Science and Education Administration, U.S. Dept. of Agriculture, Athens, GA, 30604). *Res Commun Chem Pathol Pharmacol* 20(3): 549-557; 1978.

Sprague-Dawley rats given single or multiple doses of four noncarcinogenic and seven carcinogenic nitrosamines (NA: various doses in 0.5-1.0 ml olive oil/kg body wt, po), changes in urinary ascorbic acid output were not indicative of NA carcinogenicity or acute hepatotoxicity. Histologic examination of the liver with glycogen staining was the most sensitive test of the acute effects of the NA carcinogens. The pentobarbital-induced sleeping time test was intermediate, and the GOT test was the least sensitive. (12 refs)

5595 **Diffusion of Nitrosamines Through Protective Gloves.** (Eng) Gough, T. A. (Lab. Government Chemist, London, England); Webb, K. S.; McNeill, M. F. *IARC Sci Publ* (19): 531-534; 1978.

The protection afforded by rubber and polyvinyl chloride gloves against N-nitrosodimethylamine (DMNA), N-nitrosodiethylamine (DENA), N-nitrosodipropylamine (NDPA), N-nitrosodibutylamine (NDBA), N-nitrosopiperidine (NPPI), and N-nitrosopyrrolidine (NPYR) was evaluated. The gloves were turned inside out, and 1 ml of a 10 µl/liter nitrosamine soln was placed in each of the four fingers. The outer faces of the fingers were washed every 10 min, and the washings were evaluated for nitrosamines by gas chromatography-mass spectrometry. In another experiment, the rubber in the gloves themselves were analyzed for nitrosamines, but none were found. For both glove types, NPYR permeated the fingers most rapidly, especially within the first 30 min, and to the greatest extent. Rubber gloves allowed diffusion in the following hierarchy: NPYR > NPIP > DENA > DMNA > NDPA > NDBA. After 2 hr, approx 57% of NPYR and 2.5% of NDBA had diffused through the gloves. Polyvinyl chloride gloves allowed diffusion in the following order: NPYR > NDPA > NPIP > DENA > DMNA. After 1 hr, approx 45% NPYR and 10% of DMNA had diffused through the gloves. It is concluded that the use of gloves does not give adequate protection from nitrosamine contamination of the hands. (3 refs)

5596 **The Permeability of Laboratory Gloves to Selected Nitrosamines.** (Eng) Sansone, E. B. (Frederick Cancer Res. Center, Frederick, MD); Tewari, Y. B. *IARC Sci Publ* (19): 517-529; 1978.

The permeability of five varieties of laboratory gloves to 0.136 to M solns of N-nitrosodimethylamine, N-nitrosodiethylamine, N-nitrosoethylmethylamine, N-nitrosobutylmethylamine, and N-nitrosodipropylamine, N-nitrosodiisopropylamine, and N-nitrosodibutylamine in acetone, dichloromethane, ethanol, and water was determined. Disks cut from gloves made of natural rubber, polyvinyl chloride, nitrile, neoprene, and a neoprene + natural rubber mixture were used to partition a permeability apparatus into two compartments that contained the permeant soln and water, respectively. Samples were withdrawn from the water side after 0.5, 1, 2, 4, 6, and 8 hr and analyzed for nitrosamine (NA's). In nearly all cases, readily measurable amounts of NA's penetrated glove materials within 0.5 hr. In general, penetration of NA's decreased with increasing mol wt in acetone, dichloromethane, and ethanol, whereas with water as solvent, NA's penetrated more readily as their mol wt increased. The results indicate that gloves should not be assumed to provide full protection, they should be discarded immediately following overt contamination, and they should not be reused. (14 refs)

5597 **Limitations to the Protective Effect of Rubber Gloves for Handling Nitrosamines.** (Eng) Walker, E. A. (International Agency Res. Cancer, Lyon, France); Castegnaro, M.; Garren, L.; Pignatelli, B. *IARC Sci Publ* (19): 535-543; 1978.

The rate of diffusion of N-nitrosodimethylamine (DMNA) and N-nitrosodiethylamine (DENA) through various types of rubber gloves was investigated. When a glove was filled with methylene chloride and suspended in a beaker containing an equal volume of methylene chloride soln containing 10 mg/liter each of DMNA and DENA, about 25% of both nitrosamines diffused through in 15 min. When saline was substituted on the inside of the barrier to simulate human perspiration, 10% of DMNA diffused through in 15 min. Slightly lesser values were determined for DENA. The thicker household gloves tested actually provided less protection than surgical gloves, possibly because of the properties of the materials constituting the two types of gloves. DMNA showed a tendency to concentrate in the rubber of surgical gloves and then equilibrate with the saline phase. The diffusion of other nitrosamines across a surgical glove barrier from methylene chloride to saline was also investigated, and after 20 min the relative hierarchy of diffusion was DMNA > N-nitrosopyrrolidine > DENA > N-nitrosopiperidine > N-nitrosomethylpentylamine > N-nitrosodipropylamine > N-nitrosodibutylamine. A double layer of surgical gloves successfully reduced exposure. Better protection was afforded by dusting the inner pair with talc. However, max protection

was afforded by smearing the inner pair with oil-resistant barrier cream, with only 2.9% of DMNA diffusing through the material after 30 min of exposure. (no refs)

- 78-5598 **Carcinogenesis by Nitroso Derivatives of Methylcarbamate Insecticides and Other Nitrosamides in Rats and Mice.** (Eng) Lijinsky, W. (Chemical Carcinogenesis Program, Frederick Cancer Res. Center, Frederick, MD); Schmahl, D. *IARC Sci Publ* (19): 495-501; 1978.

Nitroso derivatives of methylcarbamide insecticides and other common nitrosamines were tested for carcinogenicity and mutagenicity by feeding and topical application to female BD rats. Animals were fed (by gavage) specified amounts of each derivative or nitroso-N-methylphenylcarbamate (NMPC) dissolved in corn oil 1 x/wk, or they had a 0.02 M soln of each nitroso compound in acetone painted on a shaved interscapular region 2 x/wk. All six nitrosated methylcarbamide insecticides and NMPC gave rise to forestomach tumors. Of 32 animals tested in each group, the numbers of tumor-bearers for each compound were: nitrosocarbaryl, 19; nitrosobaygon, 15; nitrosocarbofuran, 12; nitrosobux-ten, 11; NMPC, 11; nitrosolandrin, 9; and nitrosomethomyl, 4. Most of the tumors (63) were papillomas; the remainder (30) were carcinomas. Nitrosobaygon and nitrosocarbofuran each produced the greatest number of carcinomas (7). In skin-painting tests, N-methyl-N'-nitro-N-nitrosoguanidine produced 16 tumors, methyl-(d₃)-nitrosonitroguanidine 11, ethylnitrosonitroguanidine, 8, methylnitrosourea 3, and methyl-(d₃)-nitrosourea 1. None of the other nitroso compounds gave rise to more than one tumor per group of 20 animals, an unexpected finding since approx 0.8 millimole/kg of each compound was applied. (11 refs)

- 78-5599 **Distribution of 7-Methylguanine and of Replication Sites in the Different Kinetic Classes of DNA from Rats Treated with Dimethylnitrosamine.** (Eng) Galbraith, A. I. (Paterson Labs., Christie Hosp. and Holt Radium Inst., Manchester M20 9BX, England); Chapleo, M. R.; Itzhaki, R. F. *Nucleic Acids Res* 5(9): 3357-3364; 1978.

Examination of the radioactivity level in the families of repetitive and unique sequences of DNA, after treatment of male Wistar rats with ¹⁴C-dimethylnitrosamine (DMN) showed that the distribution of 7-methylguanine was random. Examination of the distribution of newly synthesized DNA after treatment of rats with DMN + ³H-thymidine indicated that there is no difference in the extent of synthesis between the two kinetic classes. (28 refs)

- 78-5600 **Formation and Subsequent Removal of O⁶-Methylguanine from Deoxyribonucleic Acid in Rat**

Liver and Kidney after Small Doses of Dimethylnitrosamine (Eng) Pegg, A. E. (Dept. Physiology, Milton S. Hershey Medical Center, 500 University Drive, Hershey, PA, 17033); Hui, G. *Biochem J* 173(3): 739-748; 1978.

The amounts of 7-methylguanine (7-MG) and O⁶-methylguanine (OMG) present in rat liver and kidney DNA were measured 4 and 24 hr after female Sprague-Dawley rats were administered low doses of dimethylnitrosamine (DMN). DMN levels were measured. Less than 15% of the expected amount of OMG (on the basis of 7-MG found) was present in liver DNA within 4 hr after ip doses of ≤0.25 mg/kg. Within 24 hr after doses of ≤1 mg/kg, >85% of the expected amount of OMG was removed. Removal was max for DMN doses of 0.25-0.5 mg/kg; however, the reaction did not appear to be saturated even at the highest dose tested (20 mg/kg). Alkylation of kidney DNA after 1 μg/kg DMN ip occurred at about one-tenth the extent of alkylation of liver DNA. Removal of OMG from kidney DNA was slower than that from the liver. The alkylation of kidney DNA was much less after po administration of DMN than after ip administration, representing only 1% of that found in the liver. Alkylation of liver and kidney DNA was readily detectable when measured 24 hr after the first injection in rats that received daily injections of 1 μg/kg ³H-DMN for 2 or 3 wk. After 3 wk, OMG levels in liver DNA were about 1% of the 7-MG levels. The amount of 7-MG in liver DNA was 10 times that in kidney DNA, but liver OMG levels were only twice those in the kidney. Extracts able to catalyze the removal of OMG from alkylated DNA in vitro were isolated from liver and kidney. These extracts did not lead to the loss of 7-MG from DNA. Although the liver OMG-removal system may protect against carcinogenesis by DMN and may be responsible for lowering the carcinogenic risk from daily doses of 5-0.5 mg/kg by more than the simple factor between the doses, its existence does not imply that there is necessarily a threshold dose at which no tumors would be expected. The findings that there is little reaction with the kidney after low po doses of DMN may explain why only liver tumors and not renal tumors are produced by prolonged feeding of low DMN doses. (70 refs)

- 78-5601 **Effect of Pretreatment with Other Dialkyl Nitrosamines on Excision from Hepatic DNA of O⁶-Methylguanine Produced by Dimethylnitrosamine.** (Eng) Pegg, A. E. (Dept. Physiology, Milton S. Hershey Medical Center, 500 Univ. Drive, Hershey, PA, 17033). *Chem Interact* 22(1): 109-116; 1978.

The effect of pretreatment of rats with higher dialkyl nitrosamines (DAN's) on the amount of O⁶-methylguanine (OMG) found in hepatic DNA after administration of 10 μg/kg ³H-dimethylnitrosamine (DMN) were determined. Female Sprague-Dawley rats were inoculated ip with DAN and, 4 hr later, with DMN, and they were sacrificed 4 or 24 hr after the latter inoculation. Pretreatment with diethylnitrosamine (DEN: 200 mg/kg) or dipropylnitrosamine (240 mg/kg) increased the amount of labeled OMG in liver DNA 4 and

after injection of DMN. Dibutyl nitrosamine (600 mg/kg) had a similar though smaller effect at 4 hr, but the effect had worn off by 24 hr. These pretreatments did not effect methylguanine levels in the DNA, showing that metabolic conversion of DMN into a methylating agent was not altered. Increases in OMG levels must be due to interference with the removal of this product from the DNA. Since the efficiency of removal of OMG from rat liver DNA has been shown to be greatly reduced by increasing the dose of DMN, the most plausible explanation of the reduced rate of excision following administration of other DAN is that they also produce altered bases in the DNA similar to OMG and thus slow the rate of repair. This hypothesis was supported by other experiments, which showed that 24 hr after administration of DEN, the amount of O⁶-ethylguanine in hepatic DNA was approximately equal to the amount of OMG found after injection of DMN and that the rate of loss of the two guanines from the DNA was comparable; ie, both are removed by the same process. This method used here may be of value for determining whether cyclic nitrosamines produce similar products. The results suggest that different nitrosamines might have additive effects in prolonging damage to DNA, a phenomenon that could be important in carcinogenesis. (30 refs)

78-5602 **Effects of Dimethylnitrosamine on Induction of Cholangiocarcinoma in *Opisthorchis viverrini*-infected Syrian Golden Hamsters.** (Eng) Thamavit, W. (Dept. Pathobiology, Faculty Science, Mahidol Univ., Rama 6 Road, Bangkok 4, Thailand); Bhamarapravati, N.; Sahapongkarn, S.; Vajrasthira, S.; Angsubhakorn, S. *Cancer Res* (12): 4634-4639; 1978.

Interactions between nitroso compounds and the presence of liver fluke (*Opisthorchis viverrini*) infections were studied in Syrian hamsters, in order to determine whether the concomitant occurrence of the endemic parasitic disease and cholangiocarcinoma in northeastern Thailand could be related to the presence of nitrosating agents, nitrosatable substances, and nitrosamine compounds commonly present in foods in this region. All 18 animals that received dimethylnitrosamine (DMN, 0.0025%) in the drinking water for 10 wk, beginning 1 wk after they were infected with metacercariae, developed cholangiocarcinoma and cholangiofibrosis. The tumor was not observed in controls receiving either DMN or parasites alone, although cholangiofibrosis was found in some animals treated with DMN alone. It is postulated that the cholangiocarcinomas arose because the DMN exerted a carcinogenic effect on the altered proliferating epithelial cells of bile ducts that had been stimulated by the parasite. These findings suggest that the combination of DMN ingestion and liver fluke infestation may play an important role in the carcinogenesis of intrahepatic duct neoplasms in humans. (28 refs)

78-5603 **Mechanism of Dimethylnitrosamine Metabolism and Activation in Rats.** (Eng) Godoy, H.

M. (Laboratorio de Quimica Biotoxicologica, CITEFA, Zufriategui y Varela, Villa Martelli, Buenos Aires, Argentina); Diaz Gomez, M. I.; Castro, J. A. *J Natl Cancer Inst* 61(5): 1285-1289; 1978.

The effect of inducers and inhibitors of hepatic microsomal metabolism on dimethylnitrosamine (DMN) metabolism and activation was studied in Sprague-Dawley rat liver preparations. DMN was metabolized in liver slices to CO₂, and reactive metabolites were formed that bound to nucleic acids. SKF 525-A [2-(diethylamino)ethyl-2,2-diphenylvalerate hydrochloride: 0.5-2 mM] inhibited CO₂ production by 27.2%-92.9% and binding by 46%-91.4%. In liver slices from rats pretreated with phenobarbital sodium (PB) for 3 days, CO₂ production was 30.8% of control values, but covalent binding was only slightly altered. In liver slices from animals pretreated with CoCl₂, the metabolism of CO₂ was 32.4% of control values and binding was 60% of control values. An NADPH-dependent, heat-sensitive, DMN-activation process leading to reactive metabolites that bound covalently to proteins was also observed in the microsomes and 9,000 x g (9K) supernatants but not in the 100,000 x g (100K) supernatants. Treatment with 1 mM SKF 525-A strongly inhibited the activation process in microsomes but only slightly inhibited this process in the 9K supernatants. Microsomes or 9K supernatants from PB-pretreated rats were less able to activate DMN than those from controls. The 9K supernatants from CoCl₂-treated animals had a markedly decreased capacity for DMN activation, but only a moderately decreased one for microsome activation. These results suggest that there are multiple pathways for DMN metabolism and activation, only some of which are dependent on cytochrome P450. (29 refs)

78-5604 **Proliferating Cell Populations in the Early Phase of Dimethylnitrosamine-induced Renal Carcinogenesis.** (Eng) Hard, G. C. (Baker Medical Res. Inst., Commercial Road, Prahran, Victoria 3181, Australia). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1267-1272; 1978.

Wistar rats were given a single 60-mg/kg ip injection of dimethylnitrosamine (DMN) and then sacrificed sequentially on successive days for 3 wk after being pulsed ip with ³H-thymidine. Kidneys were prepared for autoradiography 1 hr after sacrifice. The percentages of labeled cells in each subpopulation representing cells involved in DNA synthesis at each daily interval were correlated with the prevailing tissue changes at that time. The only cells stimulated by DMN treatment to engage in substantial waves of increased proliferative activity in the acute phase of injury (cortical fibrocytes, proximal and distal tubule lining cells) were the same as those

from which the resulting tumors (mesenchymal and cortical epithelial tumors, respectively) are thought to be derived. The greater proportion of responding cells in the mesenchymal cell population may be relevant to the subsequent high incidence of mesenchymal tumors in relation to cortical epithelial tumors induced by DMN, and it may imply a larger hit on interstitial target cells than on epithelial cells by DMN. (9 refs)

- 78-5605 Enhancement of Liver and Kidney Carcinogenesis by Dimethylnitrosamine in Thyroidectomized NZR/Gd Rats.** (Eng) Noronha, R. F. (Dept. Surgery, St. Louis Univ. Sch. Medicine, St. Louis, MO); Goodall, C. M. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1273-1282; 1978.

The effect of thyroidectomy on dimethylnitrosamine (DMN) carcinogenesis was studied in inbred male and female NZR/Gd albino rats. The rats were surgically thyroidectomized (Th-x) at age 40 days and, together with intact rats, injected ip with 20 mg/kg DMN at age 85 days. Renal tubular neoplasms were induced in 20/45 and liver tumors in 8/45 intact rats following DMN treatment. In Th-x rats, 34/49 rats developed renal epithelial tumors, 29/49 rats liver tumors. Alveogenic lung tumors occurred in 44% of the intact rats and in 47% of the Th-x rats. There was a significant sex difference in intact rats, with 70% of males and 16% of females having lung tumors. There was no sex difference in lung tumor incidence in the Th-x group. Mortality from malignant renal and liver tumors in Th-x rats was significantly greater than that in the intact group. It is suggested that tissue-carcinogen interactions are altered at the stage of oncogenic initiation in Th-x animals due to modification of the metabolism and excretion of injected DMN. It is also suggested that local metabolism of DMN in responding tissues may be of greater significance to oncogenesis than systemically distributed metabolic products arising in the liver. (24 refs)

- 78-5606 Interaction of Vitamin E and Selenium with the Hepatotoxic Agent Dimethylnitrosamine.** (Eng) Skaare, J. U. (Dept. Pharmacology and Toxicology, Veterinary Coll. Norway, Postbox 8146, Oslo Dept., Oslo 1, Norway); Nafstad, I. *Acta Pharmacol Toxicol (Kbh)* 43(2): 119-128; 1978.

The effects of vitamin E and selenium on the acute hepatotoxicity of dimethylnitrosamine (DMNA) were determined in six groups of male albino rats. Groups 1 and 4 received vitamin E (0.02% wt/wt) in the diet for 50 days + a single dose of 200 mg sc 48 hr before a single dose of DMNA (30 mg/kg

ip) or vehicle (saline), respectively. Groups 3 and 6 were given a single dose of 0.5 mg/kg sc ip 48 hr before DMNA or vehicle, respectively. Group 2 received vehicles only, and Group 5 received oil vehicle 48 hr before DMNA. Rats were killed 24 hr after the last injection. In Group 1, vitamin E ameliorated the acute hepatotoxicity of DMNA, as reflected in reduced plasma asparagine-aminotransferase (AspAT) activity. This effect was confirmed histologically. DMNA significantly increased hepatic vitamin E levels in these rats, but it had no effect on plasma levels. In Group 3, Se enhanced the acute hepatotoxicity of DMNA, as reflected in increased AspAT activity. This effect was not confirmed morphologically. DMNA did not have any effect on hepatic Se levels, however, Se decreased hepatic and plasma tocopherol levels. Vitamin E and Se may interfere with DMNA metabolism, accelerating DMNA biotransformation or by producing alternative metabolic pathways, leading to less or more toxic end products. Finally, a special function of vitamin E may be to protect membranes from DMNA damage. (51 refs)

- 78-5607 Effect of Hypophysectomy on Persistence of Methylated Purines in Rat Liver Deoxyribonucleic Acid after Administration of Dimethylnitrosamine.** (Eng) Pegg, A. E. (Dept. Physiology, Specialized Cancer Research Center, Milton S. Hershey Medical Center, 500 University Drive, Hershey, PA, 17033); Hui, G.; Rogers, K. J. *Biochim Biophys Acta* 520(3): 671-678; 1978.

The effect of hypophysectomy and subsequent growth hormone (1 mg/kg sc) treatment on the formation of methylated purines in the liver DNA of Sprague-Dawley rats treated with dimethylnitrosamine (DMNA: 1 or 10 mg/kg ip) was investigated. In control and hypophysectomized DMNA-treated rats, the major products were 7-methylguanine (7-MG) and 0⁶-methylguanine followed by 3-methyladenine (3-MA), 7-MA, 1-MA, and 3-MG. Levels of 7-MG, 3-MG, 7-MA, 3-MA, and 1-MA were slightly greater than control levels in hypophysectomized rats (expected due to enlarged livers), but levels of 0⁶-MG were significantly greater than control levels. This was due to the slower rate of loss of 0⁶-methylguanine from the DNA of the hypophysectomized animals. The rate of loss of all other purines was relatively fast and did not differ significantly between experimental and control animals. Growth hormone treatment of hypophysectomized rats decreased DNA 0⁶-methylguanine levels but had no effect on 7-MG levels, which suggested that the hormone increased the activity of the enzymatic system that removes 0⁶-MG from DNA. Although this hormone was able to restore partial activity to the 0⁶-methylguanine excision system, other factors, possibly hormonal, are apparently required for full activity. (32 refs)

- 78-5608 Formation of Dimethylnitrosamine in Polluted Environment and the Role of Photosynthetic Bacteria.** (Eng) Kobayashi, M. (Dept. Agricultural Chemistry,

oto Univ., Kyoto 606, Japan); Tchan, Y. T. *Water Res* 199-201; 1978.

ation of sludge with a fish meat medium, to simulate
uted environment, revealed that dimethylnitrosamine
N) was formed during putrefaction. However, this com-
was metabolized upon addition of the photosynthetic
ia *Rhodopseudomonas capsulata* to the medium. Rats
then given DMN in their drinking water with or without
acteria. Liver tumors were prevented in rats receiving
+ *R. capsulata*. (11 refs)

509 **Formation of Nitrosamines from Drugs, III:
Quantitative Determination of N-Nitrosodime-
amine Formed from Promethazine and Nitrite.** (Ger) Ab-
aki, A. (Pharmazeutisches Institut der Universitat, An-
mmenbourg 4, 53 Bonn-Endenich, W. Germany); Lau-
Wetjé, A. K.; Roth, H. J. *Arch Pharm (Weinheim)*
): 775-779; 1978.

formation of N-nitrosodimethylamine (NDMA) by reac-
of promethazine with sodium nitrite in HCl at 37 C was
mined quantitatively by a polarographic technique.
MA concentrations of 5×10^{-6} M could be determined.
fering depolarizing agents were eliminated by vacuum
lation below 60 C. (no refs)

610 **Possible Nitrosodimethylamine Formation in
Comparative In Vitro Nitrosation Experiments
Six Different Tetracycline Antibiotics.** (Eng) Roper, H.
itut fur Organische Chemie und Biochemie der Univer-
Hamburg, 2000 Hamburg 13, W. Germany); Heyns, K.
C Sci Publ (19): 219-237; 1978.

hydrochlorides of tetracycline (TC) and six of its deriva-
s were reacted with sodium nitrite for 2 hr at 37 C in
eous buffer solns at pH 2 and 4 under simulated stomach
ditions. Minocycline (MC), doxycycline (CD), oxytetra-
line, and anhydrochlorotetracycline (AHCTC) formed
nitrosodimethylamine (NDMA) in 0.02%-2.0% yields, as
ected by gas-liquid chromatography. NDMA formation
n MC and DC was blocked 97% by ascorbic acid. In
osation experiments with MC and a thiocyanate catalyst,
yields (0.5%-0.6%) were close to those of the noncat-
ed reaction (0.5%-0.7%). However, the possibility of an
eleration of the reaction during the initial phase cannot
discounted, as NDMA levels were measured after a reac-
time of 2 hr. The different reactivities of the tested TC's
ard nitrite in acidic solns can be explained stereochemi-
y. The failure of dealkylative nitrosation reactions to oc-
with TC, chlorotetracycline, and desmethylchlorotetra-
line is due to the formation of intramolecular hydrogen
dge linkages between epidimethylamino groups at C-4 and
l groups at C-6. An electrophilic attack of NO^+ on the
ethylamino nitrogen is hindered, and NDMA formation

is not possible. This hypothesis was proved by the formation
of NDMA from AHCTC and sodium nitrite at pH 2. (59 refs)

78-5611 **Mechanism of N-Nitrosodimethylamine Forma-
tion from Trimethylamine and Trime-
thylaminoxide.** (Eng) Ohshima, H. (Dept. Biomedical Res.
Food, Natl. Inst. Health, Tokyo, Japan); Kawabata, T. *IARC
Sci Publ* (19): 143-153; 1978.

The kinetics of the in vitro nitrosation of trimethylamine
(TMA) and trimethylaminoxide (TMAO) to produce
N-nitrosodimethylamine (NDMA) was studied. TMA or
TMAO (both, 0.5 millimole) was reacted with sodium nitrite
(2.5 millimoles) in a pH range of 1-7 and at temperatures of
25-100 C for 2 hr. The formation of NDMA was max at pH
3 and 100 C, with yields as high as 53% being produced from
TMA, 55% from TMAO. At < 75 C, the pH dependence
of the nitrosation was similar to that observed with alkyla-
mides. The initial rate of NDMA formation from TMAO at
pH 3 and 25 C was proportional to the TMAO and nitrite
concentrations, but not to the square of the nitrite concentra-
tion. In contrast, the rates of NDMA formation from TMA
and TMAO at pH 3 and 100 C were proportional to the
square and the cube, respectively, of nitrite concentration.
The mechanisms of the nitrosation of TMA and TMAO were
very complex, and the pathways of NDMA formation dif-
fered according to pH and temperature. At higher tempera-
tures NDMA formation may involve oxidative cleavage of
tertiary amines to produce secondary amines (dimethyla-
mine, DMA), which may react with nitrite to form NDMA.
At lower temperatures, however, NDMA may be formed
from TMA or TMAO by a pathway not involving DMA. (14
refs)

78-5612 **Acceleration of Nitrosamine Formation at pH
3.5 by Microorganisms.** (Eng) Archer, M. C.
(Dept. Nutrition and Food Science, Massachusetts Inst.
Technology, Cambridge, MA); Yang, H. S.; Okun, J. D.
IARC Sci Publ (19): 239-246; 1978.

The effects of microorganisms on nitrosamine formation at
acidic pH, conditions of relevance to nitrosation of amines
in the stomach, were investigated. When 20 mM dihexyla-
mine (DHA) was nitrosated at pH 3.5 in the presence of 12
mg/ml *Saccharomycopsis lipolytica*, *Saccharomyces cerevisiae*,
Escherichia coli, or *Bacillus brevis* cells, the rate of dihexyl-
nitrosamine formation was enhanced 12-49 times over that
without the bacteria or yeast cells. Rates were similar in the
presence of either boiled or unheated cells. As the alkyl chain
length of the secondary amine increased, the magnitude of
the rate enhancement also increased, with DHA having the
greatest rate acceleration. Nitrosation of the heterocyclic se-
condary amines morpholine and piperidine was not affected
by the cells. Sonic disruption of *E. coli* cells followed by cen-
trifugation showed that the particulate fraction was responsi-

ble for 80% of the acceleration effect, the soluble cell fraction for 20%. A nonenzymatic mechanism involving hydrophobic interactions of the precursor amines and cellular constituents is proposed. (15 refs)

- 78-5613 N-Nitroso Compounds from the Reaction of Primary Amines with Nitrite and Thiocyanate.** (Eng) Tannenbaum, S. R. (Dept. Nutrition and Food Science, MA Inst. Technology, Cambridge, MA); Wishnok, J. S.; Hovis, J. S.; Bishop, W. W. *IARC Sci Publ* (19): 155-159; 1978.

When primary amines are reacted with nitrite ion in the presence of thiocyanate ion, large numbers of thermal energy analyzer-positive compounds are formed, some of which are likely to be N-nitroso derivatives. Since nitrate and thiocyanate are found in human saliva and primary amines are widely distributed in food, these reactions are of significance in the context of human cancer. (9 refs)

- 78-5614 Formation In Vivo of Traces of Volatile N-Nitrosamines in the Cow after Direct Administration of Nitrate into the Rumen--Preliminary Communication.** (Eng) van Broekhoven, L. W. (Center Agrobiological Res., Wageningen, Netherlands); Stephany, R. W. *IARC Sci Publ* (19): 461-463; 1978.

N-Nitrosamine formation was investigated in the digestive tract of a quadruple-fistulated cow with a fistula in the rumen. The cow was fed a low-nitrate hay diet and then 260 mg/kg potassium nitrate, directly into the rumen. The variable amounts of N-nitrosodiethylamine, N-nitrosopyrrolidine, and N-nitrosodimethylamine detected in rumen fluid correlated with nitrite formation as a function of time after nitrate administration. (31 refs)

- 78-5615 The Intake of Nitrate, Nitrite, and Volatile N-Nitrosamines and the Occurrence of Volatile N-Nitrosamines in Human Urine and Veal Calves.** (Eng) Stephany, R. W. (Natl. Inst. Public Health, Bilthoven, Netherlands); Schuller, P. L. *IARC Sci Publ* (19): 443-460; 1978.

The occurrence of volatile N-nitrosamines (NA's) in 100 total diets, 19 cheese samples, 30 veal calves fed nitrate-rich (16,400 mg/kg KNO_3) diets, and in the urine of 14 kidney stone patients medicated with massive doses of ammonium nitrate (equivalent to up to 180 mg/kg KNO_3) was evaluated by gas chromatography-thermal energy analysis. N-Diethylnitrosamine (NDMA) was frequently found in all samples, at concentrations $< 1 \mu\text{g/kg}$ except for two diets (1.2 and 1.7 $\mu\text{g/kg}$). Other NA's were found sporadically in the same concentration range except for one diet having an N-nitrosopyrrolidine concentration of 1.9 $\mu\text{g/kg}$. The mean daily

intake of NDMA, KNO_3 , and NaNO_2 from the total was calculated to be 0.5 μg , 215 mg, and 7.7 mg, respectively. NA recoveries were mostly at the 1- $\mu\text{g/kg}$ level. There was no correlation between NDMA levels and the composition of the diets or the nitrate load of the patients or veal calves. Although the patients received KNO_3 doses that were 2 to 10 times the mean daily intake of dietary nitrate, NA levels in the urine were not extraordinary. In addition, there were no significant differences in NDMA levels between tissue samples from nondosed and dosed calves. The KNO_3 and NaNO_2 levels in 142 vegetable samples and 48 human saliva samples collected at various times after the intake of vegetable-rich meals were determined. Some vegetables had high to very high nitrite levels, but the vegetables contained only traces of nitrite. The nitrite content was highly dependent on dietary intake of nitrate and it increased up to 60 times after the consumption of vegetable-rich meals. (29 refs)

- 78-5616 New Data from Collaborative Studies on the Analysis of Nitrosamines.** (Eng) Castegnaro, M. (International Agency Res. Cancer, Lyon, France); Walker, A. *IARC Sci Publ* (19): 53-62; 1978.

The results of collaborative studies of the analysis of volatile and nonvolatile nitrosamines (NA's) are reported. In the first study, 18 laboratories analyzed heavily spiced processed meat samples spiked with N-nitrosodimethylamine (DMNA), N-nitrosodiethylamine, N-nitrosodibutylamine, and N-nitrosopyrrolidine. Preliminary mass spectrometric (MS) analysis showed that the unspiked meat contained measureable levels of DMNA and N-nitrosopiperidine. The analytical techniques used included MS, gas chromatography (GC), thermal energy analysis (TEA), and other chemiluminescence techniques. The chemiluminescence methods (including TEA) were as accurate as GC or MS for trace quantities of NA's. The results indicated that volatile NA's may be determined in spiced, processed meat samples without special difficulties by several established methods. In the second study, freeze-dried solns of three nonvolatile NA's (N-nitrososarcosine, N-nitrosoproline, and N-nitrosohydroxyproline) were analyzed by 12 laboratories. Participants were provided with a tube containing a known amount of each NA and one containing comparable but unknown amounts. The analytical methods used included high-pressure liquid chromatography or GC plus detection by UV or TEA, differential pulse polarography, and thin-layer chromatography followed by fluorimetric detection. The results indicated that a sufficient number of laboratories are able to perform this analysis and are able to be involved in a collaborative study of nonvolatile NA's in food matrix samples. (refs)

- 78-5617 Rapid Formation of N-Nitrosamines from Nitrogen Oxides under Neutral and Alkaline Conditions.** (Eng) Challis, B. C. (Chemistry Dept., Imperial Coll., London, England); Edwards, A.; Hunma, R. R.; K

los, S. A.; Outram, J. R. *IARC Sci Publ* (19): 127-142;

formation of N-nitrosamines (NA's) under a wide range of conditions is demonstrated. At pH 6-14 and at 25 C, NA's are formed very rapidly in organic solvents from N_2O_3 to N_2O_4 . The reactions are much faster than those with acidified nitrite. A 2×10^{-3} M amine gives approx 10%-50% NA in a few minutes with an excess of nitrogen oxide. NA yields are not affected by the absence of air (O_2), I_2 , or metal salts. O_2 accelerates NA formation by converting NO (via NO_2) to N_2O_3 or N_2O_4 , but I_2 and metal salts are effective under anaerobic conditions, in which reaction rates are independent of amine basicity but dependent on the nature of the reagent. The most effective substance is I_2 , which gives quantitative yields of NA in a few minutes at 25 C by forming the reactive nitrosyl iodide (NOI) reagent. Acceleration in ethanol is also observed with several metal salts, with substantial amounts of NA being produced in 30-300 min. Metal iodides intervene in the NOI reagent, as for I_2 , but other salts require a mechanism involving reaction between a metal-amine complex and itself. The relevance of NA formation to human exposure is discussed, with particular reference to urban pollution and the effect of dietary antioxidants. (34 refs)

5618 **Possible Breakdown of Volatile N-Nitrosamines by Enzymatic and/or Microbiological Agents in Swiss Raw Sausage.** (Eng) Hauser, E. (Swiss Federal Veterinary Office, Berne, Switzerland); Heiz, H. J. *IARC Sci Publ* (19): 289-296; 1978.

The possible degradation of four added nitrosamines (N-nitrosodimethylamine (DMNA), N-nitrosodiethylamine (DENA), N-nitrosopiperidine (NPPI), and N-nitrosopyrrolidine (NPYR)) in sodium nitrate-containing Italian salami was investigated. Two of the four salamis contained starting culture (micrococcus and lactobacillus) and two contained no culture. The degradation of each nitrosamine was determined by gas chromatography (GC) coupled with a thermal energy analyzer and by GC-mass spectrometry. DMNA and DENA decreased over the 29-day ripening period, and the decrease was enhanced over the first 21 days by the addition of starting culture. On day 29, however, salamis with and without starting culture had the same total degradation. NPPI and NPYR levels fell rapidly until day 7 and then they rose to their original amounts by day 21 (NPPI) and day 14 (NPYR). Thereafter, they both dropped to about half the initial amount. The use of starting cultures had no influence on total degradation. Thus, all the nitrosamines were partly degraded during the ripening period,

with the aliphatic compounds being less stable than the heterocyclic ones. (no refs)

78-5619 **Determination of Nitrosaminoalcohols and Amines by Fluorine-19 Nuclear Magnetic Resonance of Hexafluoroacetone Derivatives.** (Eng) Gaffield, W. (Western Regional Res. Center, Agricultural Res. Service, US Dept. Agriculture, Berkeley, CA); Lundin, R. E. *IARC Sci Publ* (19): 87-95; 1978.

Fluorine-19 nuclear magnetic resonance (NMR) data for several hexafluoroacetone adducts of nitrosamino alcohols and amines were obtained by Fourier transform (FT) ^{19}F NMR spectroscopy. The spectra readily differentiated among stereoisomers and positional isomers. The FT method has great potential for signal/noise enhancement, and FT NMR is approx 20 times more sensitive than continuous wave NMR for studies of a spectral range similar to that reported. (9 refs)

78-5620 **Detection and Determination of Volatile and Non-Volatile N-Nitrosamines by Thin-Layer Chromatography Using Fluorescamine.** (Eng) Young, J. C. (Chemistry and Biology Res. Inst., Agriculture Canada, Ottawa, Ontario, Canada). *IARC Sci Publ* (19): 63-73; 1978.

Volatile and nonvolatile N-nitrosamines (NA's) were analyzed quantitatively by chromatography on thin-layer plates of activated silica gel in one of three solvent systems. The NA's were cleaved to primary or, in a few cases, secondary amines and to keto compounds by UV radiation, and the amines were subsequently detected as fluorescent spots after treatment with the primary amine-specific reagent fluorescamine. Detection limits of 10-15 nanograms (approx 75 picomoles) were observed for most NA's. The method was sensitive to volatile NA's, nonvolatile N-nitrosamino acids, and the N-nitroso derivative of the herbicide glyphosate. Unfortunately, detection limits were poorest for the volatile NA's of greatest interest (dimethyl-, diethyl-, and pyrrolidinyl-). The high limits were probably due to evaporative loss of the volatile parent NA and/or the resulting primary amine. Plots of relative fluorescence vs nanomoles of NA were close to linear, so that NA concentrations could be quantitated. Applications of the method are given. (24 refs)

78-5621 **An Examination of Some Foodstuffs for Trace Amounts of Volatile Nitrosamines Using the Thermal Energy Analyzer.** (Eng) Gough, T. A. (Lab. Government Chemist, London, England). *IARC Sci Publ* (19): 297-304; 1978.

A wide range of foodstuffs was examined for trace amounts of nitrosamines (NA's) by chemiluminescence using a ther-

mal energy analyzer (TEA). Many of the 264 samples had been examined previously by mass spectrometry and shown to contain no NA's at levels $> 1 \mu\text{g/kg}$. All eight fried bacon samples contained 1-10 $\mu\text{g/kg}$ N-nitrosodimethylamine (DMNA) and N-nitrosopyrrolidine (NPYR), 0.08-0.25 $\mu\text{g/kg}$ N-nitrosoethylmethylamine (PNIP) and 0.3-0.5 $\mu\text{g/kg}$ of an unidentified NA. N-Nitrosoethylmethylamine and N-nitrosodiethylamine also occurred often in cured meats up to 0.2 $\mu\text{g/kg}$. The only NA detected frequently in the other food samples was DMNA, which was sometimes present up to 0.2 $\mu\text{g/kg}$. Other volatile NA's were rarely found, and their contribution to the total volatile NA content of food is negligible. The unidentified NA's sometimes detected by TEA do not contribute significantly to intakes, except, possibly, those in cured meats. Cured meat, fish, and cheese contained an av dialkyl-NA level about 1 order of magnitude higher than that in all the other foods examined. The only significant contribution of NPYR and NPIP to the diet is from cured meats, as, in all other samples, these NA's were detected in only a few instances in amounts $> 0.01 \mu\text{g/kg}$. (13 refs)

- 78-5622 Observations on the Use of the Thermal Energy Analyzer as a Specific Detector for Nitrosamines.** (Eng) Fiddler, W. (Eastern Regional Res. Center, Agricultural Res. Service, US Dept. Agriculture, Philadelphia, PA); Doerr, R. C.; Piotrowski, E. G. *IARC Sci Publ* (19): 33-39; 1978.

Since most nitrosamines (NA's) undergo photolytic decomposition when exposed to UV light, this property was used to develop an analytical method for NA's sensitive to picogram levels. After photolysis, NA determinations are performed on a gas-liquid chromatography interfaced with a thermal energy analyzer. This technique is intended not to replace but to supplement mass spectrometric confirmation. (14 refs)

- 78-5623 Survey of Cured Meat Products for Volatile N-Nitrosamines: Comparison of Two Analytical Methods.** (Eng) Havery, D. C. (Dept. Health, Education and Welfare, Washington, DC); Fazio, T.; Howard, J. W. *IARC Sci Publ* (19): 41-52; 1978.

A total of 106 cured meat samples were analyzed by multidetector gas liquid chromatography-mass spectrometry and the mineral oil distillation-thermal energy analyzer method for volatile N-nitrosamines. From 5 to 75 ppb N-nitrosopyrrolidine were found in fried bacon, and traces of N-nitrosodimethylamine were found in various cooked meats. The two techniques produced comparable results, especially at the 10-ppb level. (24 refs)

- 78-5624 Determination of N-Nitrosamine Levels in Meat and Some Meat Products.** (Rus) Pokrov-

skii, A. A. (Inst. Nutrition, Moscow, USSR); Kostiukskii, I. A.; Melamed, D. B.; Medvedev, F. A. *Vopr Pita* 65-72; 1978.

Gas chromatography/mass spectrometry was used to trace amounts of N-nitrosamines (NA) in food products. Sensitivity of the method was 3 nanograms. The amount of NA in various meat products ranged from 0.9 to 23.4 $\mu\text{g/kg}$ (the highest amount was found in pork sausage). NA was not found in meat. (16 refs)

- 78-5625 Electronic Properties of the Nitrosamine Function.** (Eng) Lyle, R. E. (Dept. Chemistry, Texas State Univ., Denton, TX); Fribush, H. M.; Lytle, G.; Saavedra, J. E. *IARC Sci Publ* (19): 99-107; 1978.

The interaction of nitrosamines (NA's) with an oxidizing enzyme was detected by studying the effect of chiral substituents capable of hydrogen bonding on the circular dichroism in the presence of achiral NA's. The addition of a chiral alcohol to an achiral NA causes a dissymmetric perturbation of the symmetrical chromophore, and a Cotton effect is evident in the CD curve at the wavelength of UV absorption of the NA function. Structural implications are considered. (16 refs)

- 78-5626 Chemical Models for Possible Nitrosamine Artifact Formation in Environmental Analysis.** (Eng) Angeles, R. M. (Carcinogen Metabolism and Toxicity Branch, NCI, Analytical Chemistry Section, Bethesda, MD, 20014); Keefer, L. K.; Roller, P. P.; Uhm, S. J. *IARC Sci Publ* (19): 109-115; 1978.

Two recent discoveries concerning artifact formation in the analysis of nitrosamines were made: (1) solid-phase inorganic nitrite can efficiently nitrosate amines in the presence of certain common solvents, and (2) deionized water can contain significant quantities of nitrosamines. Possible implications of both phenomena are discussed, along with measures for their control. (12 refs)

- 78-5627 Comprehensive Analytical Procedures for the Determination of Volatile and Non-volatile Polar and Non-polar N-Nitroso Compounds.** (Eng) Fan, T. (Thermo Electron Res. Center, Waltham, MA); Krull, R. D.; Ross, R. D.; Wolf, M. H.; Fine, D. H. *IARC Sci Publ* 3-17; 1978.

Comprehensive methods for the quantitative analysis of volatile and nonvolatile, polar and nonpolar N-nitroso compounds combine the use of thermal energy analysis, high-pressure liquid chromatography or standard chromatography. However, independent techniques are required to confirm the results. (25 refs)

528 Characteristics of the TEA Detector in Use with High Pressure Liquid Chromatography. (Eng)

, J. K. (Dept. Medicinal Chemistry, Sch. Pharmacy, Mississippi, University, MS); Ma, C. Y. *IARC Sci Publ* 19-32; 1978.

se of a thermal energy analyzer detector in conjunction high-pressure liquid chromatography for determining nitroso compounds is described in detail, particularly the operating parameters of the detector. The detector response is highly dependent on furnace temperature, the flow rate of gas used to purge the furnace, and the volatility of the compounds. (4 refs)

529 Non-volatile N-Nitroso Compounds in Human Feces. (Eng) Varghese, A. J. (Ontario Cancer

Toronto, Ontario, Canada); Land, P. C.; Furrer, R.; W. R. *IARC Sci Publ* (19): 257-264; 1978.

nature of the previously detected mutagens in human was investigated. Samples were collected from healthy, non-smoking men and they were frozen and extracted with ether. The extracts were dried, reconstituted in ether, passed through a 0.5- μ m millipore filter, and purified by silica gel chromatography followed by high-pressure liquid chromatography. The mutagenicity of the purified mutagen was tested in the Ames assay using *Salmonella typhimurium* TA100 without microsomal activation. The exact chemical nature of the mutagen could not be determined. However, the following evidence suggests that it could be an nitroso compound: the mutagenically active ether extracts released nitrite upon heating, these extracts contained aldehydic agents and at least four new ninhydrin-positive compounds, and signals for nitroso compounds were seen in the ether extract and in the mutagen fraction by thermal mass analysis. Additional evidence was provided by a diet experiment based on the fact that ascorbic acid inhibits nitrosation in vitro and in vivo. A volunteer was given 1 g ascorbic acid before each meal and at bedtimes for 2 wk. In feces collected 1 wk prior to the experiment, the level of nitroso compounds averaged 0.45 A/g feces and mutagenic activity was 180 revertants/plate. After 2 wk of ascorbic acid, the levels decreased to 0.2 A/g and the revertants decreased to 160/plate. Nitroso compounds in the bowel are probably formed in the stomach or small bowel, and they may be the agents responsible for cancer of the colon. (21 refs)

530 Some Effects of Phenol- and Thiol-Nitrosation Reactions of N-Nitrosamine Formation. (Eng)

es, R. (Ministry Agriculture Fisheries and Food, Food Research Institute, Colney Lane, Norwich NR4 7UA, England); Massey, M. J.; Massey, R. C.; McWeeny, D. J. *IARC Sci Publ* 183-197; 1978.

ts of recent studies showing that phenols and thiols can

lead to enhanced nitrosamine (NA) formation under conditions in which it would not normally take place are reported. C-Nitrosophenols, produced by the nitrosation of phenols, catalyze NA formation. The rate of nitrosation of pyrrolidine catalyzed by p-nitroso-o-cresol was fastest at about pH 5. The catalytic species is thought to be the quinone monoxime tautomer of the nitrosophenol. The catalysis may involve initial nitrosation of the catalyst, either in or via the quinone monoxime tautomeric form, to give an intermediate that then undergoes attack by the amine to give NA and to regenerate the catalyst. When pyrrolidine was reacted with nitrite at pH 5.0, N-nitrosopyrrolidine was further enhanced when the reaction was performed under anaerobic conditions. Nitrosocysteine (NC) HCl reacted with N-methylaniline at pH 2.65, 5.5, and 9.75 in the absence of nitrite to give N-nitroso-N-methylaniline. The yield of the latter was reduced significantly at pH 9.75 under anaerobic conditions. NC HCl also nitrosated morpholine and pyrrolidine at pH 9.75. The rate of the reaction at this pH decreased with increasing amine basicity. The implications of these results in terms of the formation of NA's in cured meats and in vivo are not clear. The complicating effects of other components and of adsorption in heterogeneous systems cannot be predicted quantitatively. (15 refs)

78-5631 Fate of Nitrate and Nitrite in Saliva and Blood of Monkey Administered Orally Sodium Nitrate Solution, and Microflora of Oral Cavity of the Monkey.

(Jpn) Hayashi, N. (Natl. Inst. Hygienic Sciences, 18-1, Kamiyoga 1-chome, Setagaya-ku, Tokyo, Japan); Watanabe, K.; Ishiwata, H.; Mizushiro, H.; Tanimura, A.; Kurata, H. *J Food Hyg Soc Jpn* 19(4): 392-400; 1978.

The fate of nitrate (NO_3) and nitrite (NO_2) in monkey saliva was investigated in five monkeys (2.5-5.0 kg) that were fed by stomach tube with 4, 20, and 40 mg/kg of sodium nitrate (NaNO_3) at 10-day intervals. The monkeys were then designated as very good (M1), good (M2 and M3), or poor converters (M4 and M5). To obtain more detailed information about the reduction of ingested NO_3 , further investigation was performed on the M1 monkey, with M4 and M5 monkeys serving as controls. All monkeys were fed a 100 mg/kg NaNO_3 soln by stomach tube. NO_3 and NO_2 concentrations in blood and saliva were calculated. Parallel with this experiment, a comparative survey of oral microflora in the monkeys was conducted. Blood NO_3 concentrations in all monkeys peaked 7 hr after ingestion (180 ppm). In the M1 monkey, about 70 ppm was maintained for 48 hr after administration. Salivary concentrations in the M1 monkey showed a max of 800 ppm after 7 hr, and the salivary NO_3 concentration was consistently higher than that in blood. The M4 and M5 monkeys, in contrast, exhibited a lower concentration of NO_3 in blood. Salivary NO_2 in the M1 monkey reached a max concentration of 30 ppm, and 10-20 ppm was maintained in saliva for 24 hr. In the M4 and M5 monkeys, no apparent correlation between ingestion of NO_3 and formation of NO_3 in saliva was observed. Counts of bacteria and molds in the oral

cavity of the M1 monkey showed a consistently greater number of microbes than in the M4 and M5 monkeys. *Streptococcaceae*, *Peptococcaceae*, and *Bacteroidaceae* predominated in the oral cavity of the M1 monkey, and *Bacteroides melaninogenicus* was the most common species among the microflora. NO_3^- reducing anaerobic bacteria, ie, *Bacteroidaceae* and *Veillonella*, seemed to be indigenous in the mouth of the M1 monkey. (26 refs)

- 78-5632 **Gastric Juice Nitrite: Its Source and Relationship to Hydrogen Ion Concentration.** (Eng) Walters, C. L. (British Food Manufacturing Industries Res. Assoc., Leatherhead, Surrey, England); Hill, M. J.; Ruddell, W. S. *IARC Sci Publ* (19): 279-288; 1978.

Nitrite concentrations were determined in the fasting gastric juice (GJ) of 17 patients undergoing routine pentagastrin tests. Nitrite was present in the GJ from all 17 subjects, the mean concentration being $4.9 \mu\text{M}$. Pentagastrin stimulation increased the volume and H^+ concentration of the GJ and decreased the gastrin thiocyanate concentration from a mean of 0.9 to $0.3 \mu\text{M}$ after 30-45 min. However, there was no significant change from the basal level of the mean nitrite concentration of $4.9 \mu\text{M}$ throughout the stimulation period, in spite of increased gastric aspirate volumes. In GJ and saliva, the thiocyanate levels of smokers were significantly greater than those of nonsmokers, but, there were no significant differences in nitrite levels between the two groups. Patients with demonstrable gastroduodenal lesions could be divided into normals and hypochlorhydric (HCH) normals on the basis of their fasting H^+ concentrations (> 15 and $< 5 \text{ mM}$, respectively). Neither mean H^+ nor nitrite concentrations in the normals were significantly different from those of duodenal ulcer patients. H^+ concentrations in the GJ of HCH normals were significantly lower and their nitrite concentrations were significantly higher than those of all groups other than the six gastric cancer patients. The mean GJ nitrite concentration of the gastric ulcer patients was significantly higher than that of normal and duodenal ulcer patients, and the H^+ concentration was significantly lower. The six gastric cancer patients had a mean nitrite concentration significantly higher and an H^+ concentration significantly lower than those of the gastric ulcer group. The difference in total bacterial counts and those of nitrate-reducing organisms among the diagnostic groups were secondary to those in H^+ concentrations. (28 refs)

- 78-5633 **Gastric-Juice Nitrite and Gastric Cancer (Letter to Editor).** (Eng) Jones, S. M. (Univ. Dept. Surgery, Bristol Royal Infirmary, Bristol BS2 8HW, England); Davies, P. W.; Savage, A. *Lancet* 1(8078): 1355; 1978.

Among 27 patients who had undergone partial gastrectomy for duodenal ulcers, 8/10 with significant dysplasia and 2/8 with intestinal metaplasia had gastric juice nitrite concentra-

tions > 10 micromoles/liter. This finding supports a association between increased nitrite in the hypochlor stomach and the development of gastric carcinoma after partial gastrectomy. (5 refs)

- 78-5634 **Possibility of Nitrosation During Food Digestion. I. Evaluation of Precursors and Affecting Factors In Vivo. II. Effect of Food Intake on Nitrosation Reaction.** (Eng) Klein, D. (Departement de Nutrition, Maladies Metaboliques, Universite de Nancy I, Nancy, France); Gaconnet, N.; Poullain, B.; Debry, G. *IARC Sci Publ* (19): 485-494; 1978.

The influence of ingested nitrate on concentrations of nitrate and nitrite was investigated in two groups of subjects, only one of which swallowed their saliva. Each subject received a 450-ml test meal containing, per 100 ml, 10 g powder, 10 g glucose, 5 g corn oil, 2.2 g polyethylene glycol and 25 mg sodium nitrate. Saliva and gastric juice samples (10 ml) were taken at 30-min intervals. Salivary nitrate increased to a max ($35 \pm 29 \text{ mg/liter}$) after 90 min and declined. Salivary nitrite followed a similar course, but in a much smaller quantity. As the meal was digested, gastric pH decreased from pH 5 at 30 min to pH 3 at 2 hr, and dilution of the alimentary bolus increased. Gastric nitrate decreased in all individuals as a result of dilution. However, the gastric nitrite concentration in subjects who swallowed their saliva rose rapidly and reached an average value of 7 mg/liter within 10 min; nonswallowers displayed continued low nitrite levels. It is suggested that salivary nitrate is converted to nitrite in the esophagus. An in vitro study showed that nitrosation is initiated at time of meal ingestion, but the effect is diminished as the meal is diluted and digested. The results show that gastric nitrite concentration depends essentially on the salivary nitrate contribution, which is related to the nitrate concentration in the alimentary bolus. It increases during digestion and reaches values $> 10 \text{ ppm}$ in a medium that changes in a direction favorable to nitrosation; ie, the pH decreases and the effect of the alimentary pool components diminishes. (12 refs)

- 78-5635 **Effects of Some Commonly Used Food Preservatives on *Vicia Faba* Root Meristems (Memorandum Abstract).** (Eng) Gopalan, H. N. (Dept. Botany, Kenyatta Univ., P.O. Box 43844, Nairobi, Kenya); Njagi, G. D. *Kenya J. Res* 53(2): 194; 1978. (no refs)

- 78-5636 **Studies Related to Nitrosamide Formation: Nitrosation in Solvent: Water and Solvent Systems. Nitrosomethylurea Formation in the Rat Stomach and Analysis of a Fish Product for Ureas.** (Eng) Mirvish, S. S. (Ep. Inst. Res. Cancer, Univ. Nebraska Medical Center, Omaha, NE); Karlowski, K.; Sams, J. P.; Arnold, S. D. *IARC Sci Publ* (19): 161-174; 1978.

experiments related to nitrosamide formation were conducted: (1) production of nitrosamides by nitrous acid in water mixtures and in methylene chloride (MeCl); (2) N-methylurea (NMU) formation in the stomach of rats fed methylurea (MU: 100 mg/kg) + sodium nitrite (4 g/kg); (3) analysis of a salt-dried bonito fish from Japan for nitrosamine before and after nitrosation-denitrosation by ion-exchange and paper chromatography. In experiment 1, nitrous acid was partly extracted from water by organic solvents, especially polar ones. Carbaryl was nitrosated 20 times faster in MeCl:water than in water alone, mainly because of its insolubility in water. For ethylurea, hexylurea, ethyl N-ethylurea, and aminopyrine, nitrosation in sodium sulfite-extracted MeCl extracts of nitrous acid at 6°C was at least complete after 5 sec. Nitrosation of N-butylacetamide in MeCl by the same extract was slower, as was nitrosation of N-methylurea in MeCl by equivalent concentrations of dried nitrous acid, dinitrogen trioxide, and dinitrogen tetroxide. In experiment 2, the NMU yield in rats killed 3 hr after feeding was 10% of the MU. When the food also contained 11.5 g/kg ascorbate, NMU formation was inhibited completely. In experiment 3, 2.4 g/liter NaNO_2 in the drinking water and MU in the food, no MNU was detected. In experiment 3, the fish contained 80 mg/kg urea but no MU. When the fish was nitrosated at pH 1 and then denitrosated at pH 0, 25 g MU was detected. The identity of MU was confirmed by mass spectrometry. (13 refs)

37 **Methylnitrosophenylurea - A Highly Carcinogenic Compound.** (Eng) Warzok, R. (Res. Preventive Oncology, Inst. General Pathology and Anatomical Anatomy, Medical Acad. Erfurt, 50 Erfurt, E. Germany); Thust, R.; Mendel, J.; Schwarz, H.; Blaufuss, E. *Cancer Lett* 4(5): 299-303; 1978.

possible carcinogenic effects of 1-methyl-1-nitrosophenylurea (MNPU), which is easily formed in water from 1-methyl-3-phenylurea (MPU) and sodium nitrite in an acid environment, were examined in male rats of inbred strain E. The rats were given MPU, MNPU, or MPU plus NaNO_2 by a stomach tube. In the group given MPU alone, no tumors were observed in rats more than 500 days after the start of the experiment. In the group given MPU plus nitrite, only four animals that before the 137th day had no tumors; the remaining 21 developed benign or malignant growths of the forestomach. In the group given MNPU, 22/25 rats developed tumors of the forestomach. The induction of forestomach tumors in the rats given MPU and nitrite indicates that MNPU is formed endogenously after administration of these two compounds. In the metabolism of the herbicide 1,1-dimethyl-3-phenylurea (fenuron), MPU results as a demethylation product. In estimating carcinogenic risk, therefore, it is not enough to investigate only the residues of methyl-substituted ureas; the secondary amides formed by their demethylation seem to be of greater importance as possible carcinogenic

factors (by virtue of their possible nitrosation) and should be decreased to a minimum. (10 refs)

78-5638 **Combined Transplacental Carcinogenic Effect of N-Nitrosomethylurea (NMU) and N-Nitrosoethylurea (NEU) in Rats.** (Rus) Alexandrov, V. A. (Lab. Experimental Tumors, N. N. Petrov Res. Inst. Oncology, Leningrad, USSR); Shraiber, D. *Vopr Onkol* 24(4): 38-43; 1978.

The transplacental teratogenic and carcinogenic effects of N-methylnitrosourea (MNU) and N-ethylnitrosourea (ENU) were studied in random-bred albino rats. Pregnant females received a single injection of MNU (5 or 20 mg/kg, ip), or ENU (20 mg/kg, iv) on day 13, 15, 17, or 21 of pregnancy. In another experiment, females were inoculated with MNU on day 15 and with ENU on day 17, or with ENU on day 17 and with MNU on day 21. Transplacental exposure to MNU alone induced tumors in 25% (at 5 mg/kg) and 51.5% (at 20 mg/kg) of the progeny. However, tumor incidence was greater after ENU administration: 91.2% and 86.4% after exposure on days 13 and 17, respectively. MNU + ENU induced tumors in 92.6%-95.4% of the progeny. The induced tumors were predominantly located in the CNS: of 135 CNS tumors, there were 62 oligodendrogliomas, 38 mixed oligoastrocytomas, 12 astrocytomas, 12 glioblastomas, 3 ependymomas, and 1 pinealoma. All 34 tumors of the peripheral nervous system had the histology of malignant neurinoma. MNU alone and in combination with ENU induced malformations of the CNS (microcephaly or hypoplasia of the cerebellum). The incidence of brain tumors against a background of microcephaly was 3 times greater after exposure to MNU and then to ENU than after exposure to ENU and then to MNU. (19 refs)

78-5639 **Alkylation of Deoxyribonucleic Acid In Vivo in Various Organs of C57BL Mice by the Carcinogens N-Methyl-N-nitrosourea, N-Ethyl-N-nitrosourea and Ethyl Methanesulphonate in Relation to Induction of Thymic Lymphoma. Some Applications of High-Pressure Liquid Chromatography.** (Eng) Frei, J. V. (Dept. Pathology, Univ. Western Ontario, London, Ontario N6A 5C1, Canada); Swenson, D. H.; Warren, W.; Lawley, P. D. *Biochem J* 174(3): 1031-1044; 1978.

The in vivo DNA alkylation (Alk) products of N-methyl-N-nitrosourea (MNU), N-ethyl-N-nitrosourea (ENU), and ethyl methanesulphonate (EMS) were measured by high-pressure liquid chromatography to determine whether there is a quantitative relationship between tumor yield and DNA Alk at specific sites. Female C57BL mice were injected ip with single doses of the carcinogens and sacrificed 0.5-122 hr later. Alk by the nitrosoureas was complete at 1 hr, but with EMS it was max at 2-4 hr. In addition to the rapid loss of 3- and 7-alkyladenines for all tissues, there was a rapid loss

of O⁶-alkylguanines from DNA of liver that was more rapid at lower doses. In brain, lung, and kidney, excision of O⁶-alkylguanine was much less marked. In thymus, bone marrow, and small bowel, all alkylated bases were lost, with half-lives of 12-24 hr at noncytotoxic Alk doses. There was no evidence for marked excision of other minor products; 1-methyladenine, O²-ethylcytosine (found in appreciable amount only with ENU), 3-methylguanine, and dTp(Alk)dt persisted in alkylated DNA, including that of liver. In a second experiment, mice were injected ip with up to 60 mg/kg MNU, up to 240 mg/kg EN, and up to 400 mg/kg EM in order to induce thymic lymphomas. Lymphoma induction was related to the extent of Alk of the target tissues, thymus and bone marrow. Tumor yields were > 90% with 60 mg/kg MNU, < 52% with 240 mg/kg ENU, but only a few percent with 400 mg/kg EMS. The carcinogenic effectiveness of these agents was positively correlated with the extent of Alk of guanine in target tissue DNA at the O-6 atom. Assuming that at doses giving equal carcinogen response these extents of Alk would be equal, the ratio of equipotent doses to that for MNU would be 5.3 for ENU and 21 for EMS. (51 refs)

78-5640 Tumor Induction in a Localized Area of the Hamster Trachea with N-Nitroso-N-methylurea: Effect of Exposure Conditions. (Jpn) Yarita, T. (Div. Surgery, Inst. Pulmonary Cancer Res. Sch. Medicine, Chiba Univ., Inohana, Chiba 280, Japan); Nettesheim, P. *Haigan* 18(2): 181-194; 1978.

The tumor-response of the hamster trachea to two concentrations of N-nitroso-N-methylurea (NMU) and the number of exposures required for tumor induction were investigated. The animals were divided into four groups and given (1) 20 or (2) 30 twice-weekly exposures to 1% NMU, or (3) 20 or (4) 30 twice-weekly exposures to 0.5% NMU. A high incidence of benign tumors was observed in Group 3. The incidence of invasive carcinoma in the four groups was 67%, 94%, 39%, and 76%, respectively. Histological types resembling those seen in human lung cancer were noted. The major tumor types in Groups 1 and 2 were epidermoid and anaplastic large cell carcinomas. The lower concentration of NMU induced a higher incidence of adenocarcinoma (43% of all the invasive carcinomas in Groups 3 and 4, vs 13% in Groups 1 and 2). The mean tumor-induction period in Groups 3 and 4 was 10-14 wk longer than that in Groups 1 and 2. With 0.5% NMU, the epithelium bordering the tumors mainly underwent hyperplastic and hypertrophic changes. With 1% NMU, the main changes in the epithelium bordering the tumors were squamous metaplasia with various degrees of atypia. (30 refs)

78-5641 Sodium Azide-mediated Recovery from N-Methyl-N-nitrosourea-induced Genetic Injury and Repair of DNA Damage in Barley. (Eng) Gichner, T. (Inst. Experimental Botany, Flemingovo n. 2, 160 00 Pra-

gue 6, Czechoslovakia); Veleminsky, J.; Pokorny, V. *J. Exp. Bot* 18(1): 27-31; 1978.

The sodium azide-mediated recovery of barley seed N-methyl-N-nitrosourea (MNU)-induced genetic injury is reported. Dry or presoaked barley seeds were treated for 5 hr with 3-6 mM MNU, washed for 1 hr, and then treated for 3 hr with 40-600 mM azide at pH 7. The sodium azide posttreatment reduced the yield of genetic injuries (seedling height reduction, M₁ sterility, and M₂ chlorophyll mutants). The number of single-stranded break (SSB) in DNA isolated from embryos immediately after the treatment of MNU and MNU + azide treatments was approximately the same. However, when the isolation of DNA from MNU-azide-treated seeds was postponed for 24 hr, a reduction in MNU-induced SSB was noted. It is suggested that sodium azide prolonged the period between MNU-induced damage and the onset of the first semiconservative DNA synthesis, allowing more time for prereplication repair to occur. (15 refs)

78-5642 Effects of Carcinogen Dose on the Characteristics of the Tracheal Tumor Response Induced by N-Nitroso-N-methylurea in Hamsters. (Eng) Yarita, T. (Dept. Surgery, Inst. Pulmonary Res., Sch. Medicine, Chiba Univ., Chiba 280, Japan); Nettesheim, P. *Int J Cancer* 29: 298-303; 1978.

The effects of carcinogen dose on the characteristics of tracheal tumors induced by N-nitroso-N-methylurea (NMU) in male Syrian golden hamsters are described. Animals were exposed, via an intratracheal catheter, to 0.5% or 1% NMU 2 x/wk for a total of 20 or 30 exposures. Data from a previous report on exposure to 1% NMU (same regimen) are included for comparison. Most of the tumors developed in the midportion of the tracheas, and virtually all developed after the end of the 10- to 15-wk exposure period. The incidence ranged from 20% to 94% with mean tumor induction times of 13-46 wk, depending on dose (ie, exposure frequency x NMU concentration; total, 6 doses). Tumor incidence, carcinoma incidence, and number of tumors per animal were directly proportional to dose, and tumor latency was inversely proportional to dose. The number of benign tumors (papillomas, polyps) was high at the three higher doses (47%-75% of all tumors) and low at the three lower doses (2%-22%). The proportion of invasive carcinomas ranged from 5% to 40% at the lower doses and 67% to 94% at the higher doses. These included epidermoid, adenocarcinoma, adenoid-adeno-, and anaplastic large- and small-cell carcinomas. A relatively high proportion of the adenocarcinomas (7/17) occurred at the three lower dose concentrations. The possible mechanisms of the dose effect of NMU are presented. (21 refs)

78-5643 Significance of Modifying Factors in the Development of Peripheral Nervous System Tumors

Rabbits. (Rus) Beniashvili, D. Sh. (Lab. Chemical Carcinogenesis, Inst. Oncology Georgian SSR Ministry Health, Tbilisi, USSR). *Vopr Onkol* 24(9): 60-65; 1978.

Influence of modifying factors on the induction of peripheral nervous system tumors by methylnitrosourea (MNU) was studied in 170 chinchilla rabbits. MNU was introduced into the ear vein in weekly doses of 10 mg/kg. Group 1 was immunized with rabbit sciatic nerve tissue homogenate (4 ml of 0.3-0.7 ml at 2-wk intervals) in complete Freund's adjuvant; in Group 2, the sciatic nerve was chronically irritated mechanically; Group 3 comprised females castrated before MNU treatment; Group 4 comprised females treated with estradiol (60 mg/kg/mo sc); Group 5 was treated with testosterone (250 mg/kg/day); Group 6 was treated with mesalyl (5 mg/kg/day po); and Group 7 served as controls. The nerve tumor induction rate (number of tumors/number of animals alive at tumor appearance) was 9/21 in Group 1, 2/29 in Group 2, 1/15 in Group 3, 6/12 in Group 4, 1/12 in Group 5, 7/14 in Group 6, and 3/16 in Group 7. The mean latent time was 485, 531, 786, 456, 751, 442, and 720 days, respectively. The tumors were identified as neurinomas (16), gliosarcomas (4), neurogenic sarcomas (14), and sympathetic ganglioblastomas (5). They were localized in the sciatic nerve, sympathetic plexus, lumbosacral plexus, and trigeminal nerve. Neoplasms of the brain, breast, ovaries, and thyroids were found in 14 rabbits only. The findings indicate that castration and thyroidin treatment significantly reduced the tumor induction rate and prolonged the latent period. (14 refs)

78-5644 Effect of Dietary Undegraded Carrageenan on Colon Carcinogenesis in F344 Rats Treated with Azoxymethane or Methylnitrosourea. (Eng) Watanabe, K. (Maylor Dana Inst. Disease Prevention, American Health Foundation, Valhalla, NY, 10595); Reddy, B. S.; Wong, C. C.; Weisburger, J. H. *Cancer Res* 38(12): 4427-4430; 1978.

The modifying effect of undegraded carrageenan (CG) on colon carcinogenesis induced by azoxymethane (AOM) and methylnitrosourea (MNU) was studied in weanling (5 wk) female F344 rats fed a high-fat diet containing 0% or 0.5% CG. At age 8 wk, 30 rats in each dietary group (excluding controls) received 1 weekly sc injection of AOM (8 mg/kg) for 10 wk. In a separate series, 30 rats on each diet were given intrarectal injections of MNU (2 mg/rat 2 x/wk for 3 wk). The AOM groups were autopsied 40 wk and the MNU 30 wk after the first injection. No tumors developed in untreated rats fed the control diet; however, one untreated rat fed the CG diet had a single colon adenoma. The animals fed the CG diet and treated with AOM or MNU had a significantly higher incidence of colorectal carcinoma (both number of tumor-bearing rats and number of tumors per rat) than those subjected to similar carcinogen treatments but fed the control diet. It is concluded that irrespective of the type of carcinogen (AOM and MNU differ in mechanism of activation), undegraded carrageenan has an enhancing effect on colorectal carcinogenesis. (20 refs)

78-5645 Some Effects of N-Methylnitrosourea on the Oral and Odontogenic Tissues of the Syrian Golden Hamster. (Eng) Edwards, M. B. (Dept. Dental Science, Royal Coll. Surgeons of England, Lincoln's Inn Fields, London WC2A 3PN, England). *Arch Oral Biol* 23(7): 515-524; 1978.

The oral and dental changes induced by the administration of N-methyl-nitrosourea (NMU, 1 mg 2x/wk for 4 wk by stomach tube) to 10 1-mo-old male and female Syrian golden hamsters were studied. The eight animals that survived 3-15 mo after treatment showed wt loss, ascites, severe epistaxis, premalignant papillomatosis of the forestomach (7/8 animals), and premalignant lesions of the oral epithelium (7/8). The latter lesions became malignant in two animals. Between 4 and 18 wk after treatment, the lower incisors became stunted, often permanently. Three hamsters developed odontogenic cysts in incisor enamel organs that had undergone squamous metaplasia. Other changes included dental dysplasia and the growth of islands of odontogenic epithelium in the periodontal ligament. Permanent dental malformation appeared to result from repetitive cytotoxic injury and the resultant hamartoplasia, complicated by a secondary metabolic deficiency of systemic origin and by permanent disturbance of induction in odontogenesis. The proliferation of epithelium in the periodontium resembled early neoplastic changes suggestive of ameloblastoma. Further consideration of the use of NMU in models of odontogenic tumor formation is warranted. (66 refs)

78-5646 Studies on Carcinogen Chromatin-DNA Interaction: Inhibition of N-Methyl-N-nitrosourea-induced Methylation of Chromatin-DNA by Spermine and Distamycin A. (Eng) Rajalakshmi, S. (Dept. Pathology, Univ. Toronto, Medical Sciences Building, Toronto, Ontario, Canada M5S 1A8); Rao, P. M.; Sarma, D. S. *Biochemistry* 17(21): 4515-4518; 1978.

To determine whether carcinogens interact with specific regions of DNA, the effects of spermine and distamycin on the N-methyl-N-nitrosourea (MNU)-induced in vitro methylation of rat liver chromatin were studied. Both spermine and distamycin A inhibited the MNU-induced methylation of chromatin DNA at the N⁷ and O⁶ positions of guanine and the N³ position of adenine. The inhibition increased with increasing concentrations of inhibitor. Under conditions producing a 90% inhibition of chromatin DNA, the methylation of 2-deoxy-5-guanylic acid was unaffected. The results indicate that spermine and distamycin A exert their inhibitory effect by altering the structure and/or net charge of the DNA. It is suggested that MNU alkylates regions of DNA that are readily denaturable, probably A-T-rich regions located in the minor groove at or near the DNA binding sites of these two compounds. (23 refs)

78-5647 Activities of Enzymes of the Hexose Monophosphate Pathway in Nervous System Tumors In-

duced by Ethylnitrosourea. (Eng) Coleman, M. T. (Div. Neurology, Coll. Medicine, Ohio State Univ., 466 W. 10th Ave., Columbus, OH, 43210); Allen, N. *Acta Neuropathol (Berl)* 41(3): 223-227; 1978.

The activities of three enzymes of the hexose monophosphate pathway (HMP) were measured in samples of normal rat cerebral cortex and ethylnitrosourea-induced tumors of the rat nervous system (neurinomas, astrocytic gliomas, mixed gliomas, ependymomas, and undetermined types). In most tumors, adequate amounts of glucose-6-phosphate dehydrogenase (G6PDH), 6-phosphogluconate dehydrogenase (6PGDH), and transketolase (TK) were present so as not to be rate-limiting for utilization of the HMP. When enzyme activities were expressed on the basis of fresh wt, TK was increased in most tumors as compared with the cerebral cortex. G6PDH and 6PGDH were more variable, showing either higher or lower activities in tumors compared with controls. In general, the protein content of the tumors was lower than that of the cerebral cortices. When the activities were expressed in terms of this protein content, the mean values for the dehydrogenases in all tumor types were higher than the av value in the cortex. This indicated that G6PDH and 6PGDH were spared in tumors relative to other proteins. TK activities in tumors, expressed in terms of cytosol protein, were lower than or in the same range as those in the cortex. Increased activities of HMP enzymes in some tumors indicated that the potential activity of HMP in some (but not all) nervous system tumors is greater than that of the cerebral cortex. (21 refs)

78-5648 **Oxygens in DNA Are Main Targets for Ethylnitrosourea in Normal and Xeroderma Pigmentosum Fibroblasts and Fetal Rat Brain Cells.** (Eng) Singer, B. (Dept. Molecular Biology, Univ. California, Berkeley, CA, 94720); Bodell, W. J.; Cleaver, J. E.; Thomas, G. H.; Rajewsky, M. F.; Thon, W. *Nature* 276(5683): 85-88; 1978.

Experimental evidence that pyrimidine oxygen atoms are the major sites of reaction in cells treated with ethylnitrosourea (ENU) is presented. Three types of mammalian cells were treated with ¹⁴C-ENU in cell culture in conditions of low toxicity: SV40-transformed human fibroblasts, SV40-transformed Xeroderma Pigmentosum fibroblasts, and fetal (18th day) BD IX rat brain cells. The nature and proportion of alkyl products formed in the DNA of ENU-treated cells in culture were remarkably similar to those found when DNA is alkylated with ENU to a high extent in vitro. The ethylation pattern was similar for all three cell types. The products quantified in this study were ethylphosphotriesters, 7-ethylguanine, 3-ethyladenine, O⁶-ethyldeoxyguanosine, O²-ethylthymidine, O⁴-ethylthymidine, and O²-ethylcytosine. The similarity found between the in vitro and in vivo products (with over a 1,000-fold difference in the extent of reaction) suggests the possibility of predicting in vivo reaction products of an ultimate carcinogen reacting with nucleic acids from the more easily performed in vitro studies. The formation of

O-alkyl bases is likely to cause mispairing during transcription or DNA replication and may be the initiating event in cellular mutagenesis or neoplastic transformation. (18 refs)

78-5649 **Experimental Brain Tumors Induced Transplacentally by Ethylnitrosourea. V. Ultrastructural Studies Using Transmission and Scanning Electron Microscopy.** (Jpn) Taomoto, K. (Dept. Neurological Surgery, K Univ. Sch. Medicine, Ikuta-ku, Kobe 650, Japan); Fujiwara, K.; Tamaki, N.; Matsumoto, S. *No To Shinkei* 30(8): 911-1978.

The ultrastructure of gliomas induced transplacentally in SD-JCL rats with a single ip dose of 50 mg/kg ethylnitrosourea (ENU) was studied by transmission and scanning electron microscopy. The ENU-exposed subependymal layers of the fetus showed irregular and rough arrangements with expanded extracellular spaces. These changes were not seen in control rats. The subependymal layer seemed to form the "microtumor." One of the microtumors found in an 8-wk-old rat, was composed of small round cells. The fine structure of these cells was characteristic of a primitive oligodendroglioma. The fine structure of the subependymal glioma cells was often pleomorphic. These gliomas contained a mixture of primitive oligodendrocytes, ependymal cells plus anaplastic glial cells. With increasing size, the gliomas became more pleomorphic, containing a mixture of neoplastic oligodendrocytes, astrocytes, ependymal cells and ependymomalike cells with neither junctional complexes. The abnormal vascular structure of the tumor was confirmed by injection replica scanning electron microscopy. The differences between the surface of the astrocytoma and oligodendroglioma cells were clearly demonstrated. (55 refs)

78-5650 **Studies of Organ Cultures of Embryonic Brain Tissue (Hippocampus Region) of Rats Treated Transplacentally with N-Ethylnitrosourea.** (Rus) Golub, I. (Dept. Carcinogenic Agents, Oncological Res. Center, Acad. Medical Sciences USSR, Moscow, USSR); Shabat, M. *Biull Eksp Biol Med* 86(10): 496-498; 1978.

The transplacental effect of N-ethylnitrosourea (ENU: single dose of 30 or 60 mg/kg iv), administered to mongrel rats during the last week of pregnancy, was studied in organ cultures prepared from the hippocampus of embryos extracted on the 20th day of pregnancy. The brain tissue cultures proved to be highly sensitive to ENU as shown by the decreased survival rate of the experimental cultures compared with the controls (82% vs 51.9%, $p < 0.001$) and the appearance of epithelial proliferations of the vascular plexus resembling adenomas. The frequency of the morphological changes was dose-dependent: changes were found in 11/13 explants from embryos treated with 30 mg/kg ENU and in 37/132 explants from embryos treated with 60 mg/kg ENU. (10 refs)

- 651 **The Effect of Simultaneous Treatment of *E. Coli* by Chemical Mutagens and Radiation on Mutation Rate (Meeting Abstract).** (Eng) Hotz, G. (Forschungszentrum Karlsruhe, Institut für Strahlenbiologie, D 7500 Karlsruhe, W. Germany). *Mutat Res* 53(2): 1978. (no refs)
- 652 **The Effect of Expression Time on the Frequency of Carcinogen-induced Mutants at the TK HGPRT Loci in L5178Y Mouse-Lymphoma Cells (Meeting Abstract).** (Eng) Brown, M. M. (Univ. North Carolina, Chapel Hill, NC, 27514); Clive, D. *Mutat Res* 53(2): 159; 1978. (2 refs)
- 653 **Mutagenicity of Pesticides Evaluated by Means of Gene Conversion in *S. cerevisiae* and *A. nidulans* (Meeting Abstract).** (Eng) De Bertoldi, M. (Istituto di Microbiologia agraria, Università di Pisa, Pisa, Italy); Barale, R.; Giovannetti, M. *Mutat Res* 53(2): 174-175; 1978. (no refs)
- 654 **Variation in the Induction of Mitotic Gene Conversion by Chemical Mutagens (Meeting Abstract).** (Eng) Davies, P. J. (Dept. Genetics, Univ. Coll. Swansea, Singleton Park, Swansea SA2 8PP, England); Tipper, R. S.; Parry, J. M. *Mutat Res* 53(2): 173; 1978. (no refs)
- 655 **Detection of Chemically Induced Somatic Mutation in Tissues of Chinese Hamsters (Meeting Abstract).** (Eng) Dean, B. J. (Sittingbourne Res. Center, Sittingbourne, Kent ME9 8AG, England); Senner, K. R. *Mutat Res* 53(2): 173-174; 1978. (1 ref)
- 656 **Effects of Mixtures of Ethyl Methanesulfonate and Caffeine or Maleic Hydrazide on Lethal Frequencies in *Drosophila melanogaster* (Meeting Abstract).** (Eng) Nawar, M. M. (Dept. Botany, Faculty Agriculture, Suez Canal Univ., Shebeen Elkom, Egypt); Hamza, H. A. *Mutat Res* 53(2): 240; 1978. (no refs)
- 657 **Mutagenicity Studies on Chemical Carcinogens (Meeting Abstract).** (Eng) Clive, D. (Genetic Toxicology Lab., Burroughs Wellcome Co., Research Triangle Park, NC). *Mutat Res* 53(2): 169-170; 1978. (no refs)
- 658 **The Use of Marked Y Chromosome for Mutagenicity Testing in *Drosophila* (Meeting Abstract).** (Eng) Lim, J. K. (Univ. Wisconsin-Eau Claire, Eau Claire, WI, 54701). *Mutat Res* 53(2): 220-221; 1978. (no refs)
- 78-5659 **Quantitative Studies of Alkyl Methanesulfonate Induced Mutation in *Salmonella typhimurium* and Human Lymphoblasts (Meeting Abstract).** (Eng) Hoppe, H. (Dept. Genetics and Toxicology, Massachusetts Inst. Technology 18664, Cambridge, MA, 02139); Skopek, T. R.; Liber, H. L.; Thilly, W. G. *Mutat Res* 53(2): 200; 1978. (no refs)
- 78-5660 **Comparative Studies on the Relation Between Reaction Kinetics and Mutagenic Action of Monofunctional Alkylating Agents in *Drosophila* (In Vivo) and Chinese Hamster Cells (In Vitro) (Meeting Abstract).** (Eng) Vogel, E. (Dept. Radiation Genetics and Chemical Mutagenesis, Univ. Leiden, Leiden, Netherlands); Natarajan, A. T. *Mutat Res* 53(2): 279-280; 1978. (no refs)
- 78-5661 **Mutagen-sensitive Mutations of *Drosophila melanogaster* (Meeting Abstract).** (Eng) Smith, P. D. (Dept. Biology, Emory Univ., Atlanta, GA). *Mutat Res* 53(2): 265-266; 1978. (6 refs)
- 78-5662 **Sensitivity of Dominant Lethal Assay in Male Mice (Meeting Abstract).** (Eng) Ehling, U. H. (Abteilung für Genetik, Gesellschaft für Strahlen- und Umweltforschung, D-8042 Neuherberg, W. Germany). *Mutat Res* 53(2): 183; 1978. (no refs)
- 78-5663 **Purification and Characterisation of an *Escherichia coli* Enzyme that Releases Free 3-Methyladenine from Alkylated DNA (Meeting Abstract).** (Eng) Riazuddin, S. (Dept. Chemistry, Karolinska Inst., 104 01 Stockholm, Sweden); Lindahl, T. *Mutat Res* 53(2): 253; 1978. (no refs)
- 78-5664 **Somatic Mosaicism Induced by Some Carcinogens in Soybean (*Glycine max*) (Meeting Abstract).** (Eng) Arenaz, P. (Univ. Nevada, Reno, NV, 89557); Vig, B. K. *Mutat Res* 53(2): 147; 1978. (1 ref)
- 78-5665 **Interactions of Ionizing Radiation, Nitrosamines, Sulfonylalkanes and Antioxidants as**

They Affect Carcinogenesis and Survival in Mice. (Eng) Clapp, N. K. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830). *Am Ind Hyg Assoc J* 39(6): 448-453; 1978.

The effects of butylated hydroxytoluene (BHT) treatment on leukemogenesis and tumorigenesis were studied in BALB/c mice. BHT (0.75%) was added to the diet during weeks 8-12 of life; at 12 wk of age, dimethylnitrosamine (DMN), diethylnitrosamine (DEN), methyl methanesulfonate (MMS), or ethyl methanesulfonate (EMS) was given ip or the animals were x-irradiated (525 or 600 R). In another experiment, BHT was given in the feed continuously starting at 8 wk of age. In some mice, DEN (4 mg/100 ml in drinking water from weeks 11 to 18) and/or x-rays (250 R at week 11) were also administered. BHT protected the mice against acute (30-day) lethality resulting from x-irradiation or treatment with DNM, DEN, or EMS; it did not protect against the lethal effects of MMS. Survival was decreased in irradiated mice compared with nonirradiated controls, and the incidence of leukemia was increased in all irradiated groups. The incidence of leukemia was reduced in BHT-treated, irradiated animals relative to non-BHT-treated, irradiated animals. In mice treated with BHT throughout life, the compound had no effect on survival or lung tumor incidence. Survival among female mice given DEN alone was 43%; in females given DEN + BHT, it was 68%; the difference was due to a significant reduction in squamous cell carcinomas of the forestomach. BHT had no such effects in males. Radiation treatment was associated with decreased survival and an increased incidence of pulmonary adenomas; concomitant treatment with BHT nullified these effects. Treatment with DEN increased the lethality of radiation treatment, but radiation had no effect on DEN-induced pulmonary adenomas. The data suggest that the interactions among BHT and chemical and physical agents regarding late effects may occur by different mechanisms. (18 refs)

78-5666 A Linear Relationship Between Tumorigenic Potency In Vivo and Mutagenic Potency at the Heterozygous Thymidine Kinase (TK +/−) Locus of L5178Y Mouse Lymphoma Cells Coupled with Mammalian Metabolism (Meeting Abstract). (Eng) Clive, D. (Genetic Toxicology Lab., Burroughs Wellcome Co., Research Triangle Park, NC). *Mutat Res* 53(2): 170-171; 1978. (no refs)

78-5667 Application of Alkaline Sucrose Gradient Sedimentation to the Study of DNA Damage and its Repair in Mammalian Cells Treated with Methylmethanesulfonate and 4-Nitroquinoline-1-oxide. (Eng) Pavlis, R. (Dept. Biochemistry, Univ. Western Ontario London, Ontario, N6A 5C1, Canada); Yatscoff, R. W.; Sridhar, R.; Walker, I. G. *Chem Biol Interact* 23(1): 31-44; 1978.

KB human epidermoid carcinoma cells and strain L mouse cells were treated with methylmethanesulfonate (MMS,

0.1-2.0 mM) or 4-nitroquinoline-1-oxide (4 NQO, 0.05-1 μ M), and the resulting DNA damage and subsequent repair were studied using sedimentation in an alkaline sucrose gradient. The sedimentation profiles obtained resulted from a complex interaction between drug dosage, duration of the lysis period, and the repair capacity of the cells. If the number of single-strand breaks generated during the lysis period did not exceed a minimal value (on the order of 0.1 per 165S unit), the size of the DNA produced remained at 165S. At a greater frequency of breaks, DNA less than 165S was produced due to the effect of less alkali-labile lesions. It was concluded that 4NQO and MMS produce two kinds of modifications of DNA that show up as single-strand breaks during sedimentation. One is readily alkali-labile and one is less alkali-labile. Both are repaired by KB cells, but only the readily alkali-labile modification can be repaired by L cells. (14 refs)

78-5668 A Structure-Carcinogenicity Study of 4-Nitroquinoline 1-Oxides Using the SIMCA Method of Pattern Recognition. (Eng) Dunn, W. J. (Res. Group Chemometrics, Inst. Chemistry, Umea Univ., S-901 87 Umea, Sweden); Wold, S. *J Med Chem* 21(10): 1001-1007; 1978.

Structure-carcinogenicity data for a series of 33 4-nitro-4-hydroxyaminoquinoline 1-oxides were analyzed by the SIMCA method of pattern recognition. Eighteen of the compounds had consistently given positive results in animal carcinogenicity tests and 10 had been consistently negative. Using physicochemically based substituent constants to describe each compound, a principal components model was derived for the carcinogens. Using this model, 16/18 of the active and 7/10 of the inactive compounds were correctly predicted. For the seven 6-substituted compounds analyzed, a significant relationship was observed between structural parameters associated with carcinogenic potential and ability to stimulate unscheduled DNA synthesis. These results indicate that pattern recognition methods can be used to assess the carcinogenic potential of chemical compounds on the basis of a few measured variables. The differentiation between carcinogenic and noncarcinogens can be made on the basis of structural variables obtained from tabulated physicochemically based substituent constants. (29 refs)

78-5669 Mechanism of the Promotion of 4-Nitroquinoline 1-Oxide-induced Lung Tumors in Mice by Cigarette Smoke Condensate. (Jpn) Yamane, Y. (Faculty Pharmaceutical Sciences, Chiba Univ., Yayoi-cho, Chiba 280, Japan); Sakai, K.; Katsuki, H.; Okamoto, T. *Hai* 18(2): 141-147; 1978.

The tumor-promoting effect of the neutral fraction of cigarette smoke condensate (CSC) on 4-nitroquinoline 1-oxide (4-NQO)-induced mouse lung tumors was examined. Investigation of the effect of this fraction on 4-NQO metabolism

mouse lung and liver showed that 4-NQO reductase activity was strikingly enhanced in the mouse lung. This suggests that stimulation of 4-NQO reductase activity by the al fraction of CSC might be the mechanism whereby the tumors are promoted. (26 refs)

670 Carcinogenicity of 4-Nitrosoquinoline 1-Oxide and Its Possible Role in Carcinogenesis by Quinoline 1-Oxide. (Eng) Takahashi, K. (Faculty Pharmaceutical Sciences, Nagoya City Univ., Tanabedori, Nishiku, Nagoya 467, Japan); Kawazoe, Y.; Tada, M.; Ito, N.; Okada, M. *Gann* 69(4): 499-505; 1978.

carcinogenicity of 4-nitrosoquinoline 1-oxide (4-NOQO) in male ddY mice and its mutagenicity and toxicity in *Escherichia coli* were assayed. The mice received 10 weekly injections of 0.15 mg 4-NOQO sc. Fibrosarcomas were induced in 11 mice, and they were palpable between 150 and 300 days after exposure. A UV-type lesion was induced in *E. coli* after treatment with 4-NOQO, and *E. coli* mutagenicity and toxicity increased with increasing dosage, similar to the results obtained with 4-nitroquinoline 1-oxide (4-NQO). The effects of 4-NOQO were stronger than those of 4-NQO. DNA adducts in sarcoma-180 ascites cells grown in female ddY mice in vitro and incubated with 2×10^{-4} M 4-NOQO and calf thymus DNA incubated with 5×10^{-4} M in vitro were analyzed chromatographically following acid hydrolysis. The DNA adducts of 4-nitroquinoline adducts proved to be similar with those obtained by the action of 4-NQO in vivo. Because of the similarities of 4-NQO and 4-NOQO, it is assumed that these two compounds share a common ultimate metabolite, hypothesized to be O-aminoacyl-4-hydroxyaminoquinoline 1-oxide. (26 refs)

671 Misrepair Model for Carcinogenesis by 4-Nitroquinoline 1-oxide Based on Its Parallelism with Mutagenesis in *Escherichia coli* (Meeting Abstract). (Eng) Kondo, S. (Dept. Fundamental Radiology, Faculty of Medicine, Osaka Univ., Osaka 530, Japan). *Mutat Res* 53(2): 212; 1978. (no refs)

672 Carcinogenicity of 8-Nitroquinoline in Sprague-Dawley Rats. (Eng) Takahashi, M. (First Department of Pathology, Nagoya City Univ. Medical Sch., 1-1 Gokiso-cho, Mizuho-ku, Nagoya 467, Japan); Takahashi, T.; Fukushima, S.; Hosoda, K.; Yoshida, S.; Ito, N. *Cancer Lett* 4(5): 265-270; 1978.

carcinogenicity of 8-nitroquinoline given po was tested in inbred Sprague-Dawley rats. The rats were fed a semisynthetic diet supplemented with 250, 500, 1,000 or 2,500 ppm 8-nitroquinoline for 48 wk. Twenty-eight of 180 rats died during treatment due to toxicity. In the animals that survived

for 48 wk, squamous cell carcinomas of the upper digestive tract (tongue and esophagus) and, especially, of the forestomach occurred among the higher concentration groups. Squamous cell carcinomas of the forestomach occurred in 3/11 rats of the 2,500-ppm group and 7/35 rats of the 1,000-ppm group. Three cases of lung adenoma were also observed in the 2,500-ppm group. These results indicate that 8-nitroquinoline is carcinogenic in rats when administered po in a semisynthetic diet. (15 refs)

78-5673 Inhibition of DNA Replicon Initiation by 4-Nitroquinoline 1-Oxide, Adriamycin, and Ethyleneimine. (Eng) Painter, R. B. (Lab. Radiobiology, Univ. California, San Francisco, CA, 94143). *Cancer Res* 38(12): 4445-4449; 1978.

The effects of three different chemical carcinogens, 4-nitroquinoline (4-NQ), adriamycin (AM), and ethyleneimine (EI), on DNA synthesis in HeLa S3 cells were investigated by pulse labeling the nucleic acid with tritiated thymidine and sedimentation analysis. When administered in doses that reduced the rate of DNA replication to 30%-60% of control values, only EI produced damage that was manifest in lower mol wts of parental DNA. All three compounds inhibited replicon initiation but to differing extents. The primary causes of synthesis inhibition were as follows: 4-NQ, inhibition of replicon initiation; AM, slowing of chain elongation rate; and EI, blocked chain elongation. It is concluded that inhibition of replicon initiation is an important mechanism in the action of DNA-damaging agents in mammalian cells and furthermore, that control of DNA replication depends on the structural integrity of a chromosomal subunit that consists of several replicons. (32 refs)

78-5674 Effect of Aluminium Chloride on Binding of 4-Hydroxyaminoquinoline 1-Oxide to Nucleic Acids. (Eng) Yamane, Y. (Faculty Pharmaceutical Sciences, Chiba Univ., 1-33, Yayoi-cho, Chiba 280, Japan); Ohtawa, M. *Gann* 69(4): 477-486; 1978.

Aluminum chloride depressed the binding of 4-hydroxyaminoquinoline 1-oxide (4-HAQO) to DNA and RNA in the lung and liver of male dd mice and male Wistar rats. When 7 micromoles (μ mol) $AlCl_3$ were preincubated with the DNA or RNA in an ice bath for 1 hr and then incubated with 0.1 ml of an HAQO reaction mixture in the presence of 3-4 μ mol seryl-AMP, binding of the mouse and rat nucleic acids was depressed to approx 50% and 56%-68% of control values, respectively. Increasing concentrations of $AlCl_3$ decreased binding further. Sc injection of 1 mg $AlCl_3$ into mice, extraction of RNA and DNA from lungs and liver, and incubation with 4-HAQO resulted in a suppression of binding of lung DNA and RNA to 71.5% and 74.1% of control values. Corresponding figures for liver were 62.3% and 63.6%. Rats injected similarly with 50 mg/kg $AlCl_3$ displayed lung DNA

and RNA binding rates that were 60.2% and 44.7% of control values; liver DNA binding was 39.7% of control value. Binding rates with rat liver RNA were not significantly affected. When AlCl_3 (1 mg/mouse or 50 mg/kg rat) and ^{14}C -N-nitroquinoline 1-oxide (NQO) (1 mg/1 μCi /mouse or 50 mg/25 μCi /kg rat) were simultaneously injected sc on either side of the back, radioactivity measured 1 hr later was markedly depressed in lung DNA and RNA, but not in liver nucleic acids. Column chromatography demonstrated that 4-NQO severed rat lung DNA to a low-mol-wt fraction, ip or sc injection of 50 mg/kg AlCl_3 suppressed the DNA damage induced by ip or sc injection of 50 mg/kg 4-NQO. (23 refs)

78-5675 Use of the Tester Strains N23 and N24 for the Detection of the Mutagenicity of Environmental Agents and Chemical Carcinogens in *Neurospora crassa* (Meeting Abstract). (Eng) Ong, T. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC, 27709); Whong, W.; Guthrie, J. *Mutat Res* 53(2): 246; 1978. (no refs)

78-5676 Reduction of N-Methyl-N'-nitro-N-nitrosoguanidine-induced Mutations by Cobalt Chloride in *Escherichia coli*. (Eng) Kada, T. (Dept. Induced Mutation, Natl. Inst. Genetics, 1111 Yata, Misima, Sizuoka 411, Japan); Kanematsu, N. *Proc Jpn Acad Ser B Phys Biol Sci* 54(5): 234-237; 1978.

The effect of cobalt chloride on N-methyl-N'-nitro-N-nitrosoguanidine (MNNG)-induced mutations in *Escherichia coli* was investigated. *E. coli* B/r WP2, a strain that requires tryptophan (Try) for growth and reversion to prototrophy, was exposed to 60 $\mu\text{g}/\text{ml}$ MNNG for 30 min and then plated on agar that was pretreated with 5-20 $\mu\text{g}/\text{ml}$ CoCl_2 . CoCl_2 did not affect the viability of bacteria cells not treated with MNNG and it decreased the number of Try⁺ colonies per plate considerably, indicating that the compound itself is not mutagenic. With MNNG-treated bacteria, survival decreased slightly with increasing Co concentration, but mutation induction was reduced markedly. The number of Try⁺ revertants per 10^8 cells decreased from 686.8 without CoCl_2 to 640.8 with 5 $\mu\text{g}/\text{ml}$ CoCl_2 to 10.0 with 20 $\mu\text{g}/\text{ml}$ of the compound. Additional experiments showed that the effect of Co was not due to the direct inactivation of MNNG, to inhibition of the expression of Try⁺ revertants, or to the effects of crowding. It is suggested that CoCl_2 blocks an error-prone function induced by the mutagen in a way similar to that occurring in mutagenesis with UV light. (6 refs)

78-5677 Granuloma Pouch Assay. I. Induction of Ouabain Resistance by MNNG In Vivo. (Eng) Manser, P. (Inst. Toxicology, Federal Inst. Technology and Univ.

Zurich, Schorenstrasse 16, 8603 Schwerzenbach, Switzerland); Manser, P.; Zbinden, G. *Mutat Res* 54(2): 159-178.

The granuloma pouch assay was used to test the induction of ouabain resistance to N-methyl-N'-nitro-N-nitrosoguanidine (MNNG) in vivo in male albino ZUR:SIV-Z rats. MNNG (0.6 mg) was injected into the sc air pouch 24, 48, and 72 hr after induction of cell growth by croton oil (5 μl of a 0.25% soln injected sc into the air pouch); and the granulation tissue was excised 4, 24, and 48 hr later. Ouabain-resistant cells were demonstrated in all cases examined when MNNG was administered after 48 hr and the tissue was excised 4 hr later. The highest mutation frequencies were recorded when MNNG was administered 24 hr after the induction of cell growth and the tissue was excised 48 hr later. No ouabain-resistant cells were found when MNNG was administered po or ip (50 mg/kg) 36 hr after induction of growth and the granulation tissue was excised 36 hr later. These data demonstrate that MNNG has mutagenic effects in vivo in mammals if it can interact directly with the target population. (11 refs)

78-5678 Rapid Detection of DNA Damage by Carcinogenic Agents Using Immunofluorescence and Ultracentrifugation. (Eng) Neubort, S. (David and Lucile Packard Foundation, Dept. Radiology, Albert Einstein Coll. Medicine, Yeshiva Univ., Bronx, NY, 10461); Liebeskind, D.; Leifer, A.; Bases, R. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer (New York, NY): Vol. 2 (Part 1), 2404 pp.; 2029-2040; 1976.

The capability of immunofluorescence and ultracentrifugation to detect DNA damage induced by chemical carcinogens was investigated. In untreated HeLa cell cultures, only a small fraction of the cells ($\leq 15\%$) were immunoreactive with antinucleoside antibodies. G1 HeLa cells isolated by selective detachment were treated with x-rays (1,000 rads) or chemical carcinogens such as N-methyl-N'-nitro-N-nitrosoguanidine (MNNG: 4 $\mu\text{g}/\text{ml}$ for 60 min). Fixed cells were then stained with antinucleoside fluorescent antibodies and observed by UV microscopy. The fraction of cells exhibiting nucleoside fluorescence was greatly enhanced by carcinogen treatment. When HeLa cells were labeled with ^3H -thymidine, grown, lysed, and sedimented in alkaline sucrose gradients, the DNA sedimented rapidly as a large macromolecular complex. In contrast, treatment with radiation, MNNG, or one of several chemical carcinogens caused the release of slowly sedimenting single-stranded DNA. Both methods were sensitive to MNNG dose-dependent. The two approaches yielded similar results for all agents tested, and they were simple, rapid, and sensitive. Good correlation was generally obtained between the results of the two methods. (21 refs)

679 **The Role of Pre-replication and Post-replication Processes in Mutation Induction in *Haemophilus influenzae* by N-Methyl-N'-nitro-N-nitrosoguanidine.** (Eng) Kimball, R. F. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Perdue, S. W.; Boling, M. E. *Mutat Res* 52(1): 57-72; 1978.

Repair and fixation of premutational damage induced in *Haemophilus influenzae* by N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), were studied with the use of a temperature-sensitive DNA elongation mutant (*dna9*) and its combinations with mutants defective in pyrimidine dimer excision (*uvr1*, *uvr2*) and in recombination (*rec1*). The *dna9* mutant was leaky, allowing about 1% of the normal rate of DNA synthesis at the restrictive temperature. Repair of mutational lesions was detected by a decline in mutation frequency with increasing delay in DNA replication in *dna9* at the restrictive temperature. This repair was unaffected by the pyrimidine dimer excision system. Mutation fixation was detected by the ability of DNA from treated and then lysed cells to transfer mutants to recipient cells by transformation. The fixation occurred at the restrictive temperature, but the rate was much less than that occurring at the nonrestrictive temperature, suggesting that an appreciable minority of mutations resulted from lesions introduced near the replication fork but that most mutations arose from lesions introduced at some distance from the fork, perhaps randomly. DNA synthesized immediately after MNNG treatment had a lower mol wt than normal DNA, but the mol wt returned to normal with time. This return was blocked in the *dna9* mutant, suggesting that recombination is involved. (38 refs)

5680 **Mutagenesis of S49 Mouse Lymphoma Cells (Meeting Abstract).** (Eng) Coffino, P. (Univ. of California, San Francisco, CA); Friedrich, U.; Steinberg, R. *Mutat Res* 53(2): 171-172; 1978. (no refs)

5681 **NG-Induced Mutability Within Constitutively Derepressed Histidine Operon of *Salmonella typhimurium* (Meeting Abstract).** (Eng) Topisirovic, L. (Lab. of Molecular Biology, Inst. 'Boris Kidrich', PO Box 522, Belgrade, Yugoslavia); Krsmanovic-Simic, D. *Mutat Res* 53(2): 173-174; 1978. (1 ref)

5682 **A Study of the Mutagenic Potential of N-Methyl-N'-Nitro-N-Nitrosoguanidine (MNNG) in the Translocation Test (Meeting Abstract).** (Eng) Benson, T. A. (Stanford Res. Inst., Menlo Park, CA); Hershbrock, C. J.; Newell, G. W.; Green, S. *Mutat Res* 53(2): 175-176; 1978. (no refs)

78-5683 **Comparison of Carcinogenicity of N-Alkyl-N'-nitro-N-nitrosoguanidines.** (Eng) Sano, T. (Div. Biochemistry, Natl. Cancer Center Res. Inst., Tsukiji 5-1-1, Chuo-ku, Tokyo 104, Japan); Kawachi, T.; Matsukura, N.; Nakadate, M.; Sugimura, T.; Moriyama, M. *Gann* 69(2): 277-279; 1978.

The carcinogenicity of seven homologs of N-alkyl-N'-nitro-N-nitrosoguanidine, namely, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), N-propyl-N'-nitro-N-nitrosoguanidine (PrNNG), N-butyl-N'-nitro-N-nitrosoguanidine (BNNG), N-isobutyl-N'-nitro-N-nitrosoguanidine (INNG), N-pentyl-N'-nitro-N-nitrosoguanidine (PeNNG), and N-hexyl-N'-nitro-N-nitrosoguanidine (HNNG), was examined by injecting the compounds sc into male Wistar rats 2x/wk for 5 wk, for a total dose of 85 micromoles. MNNG and BNNG were strongly carcinogenic; PrNNG, INNG, and PeNNG also induced tumors at the site of injection, but at a lower rate. HNNG was not carcinogenic. Differences in membrane permeability (due to differences in the length of the alkyl chain), differences in the interaction of the various alkyl groups with nucleic acids, and differences in the repair of the various alkylated DNA's formed in the cell may be related to the differences in carcinogenicity of these compounds. (11 refs)

78-5684 **Effect of Bile Acids and Dietary Fat on Large Bowel Carcinogenesis in Animal Models.** (Eng) Narisawa, T. (Dept. Surgery, Akita Univ. Sch. Medicine, 1-1 Hondo 1-chome, Akita, 010 Japan); Reddy, B. S.; Weisburger, J. H. *Gastroenterol Jpn* 13(3): 206-212; 1978.

The effects of dietary fat and bile acids on chemical carcinogenesis in the large bowel were studied in rats. Female Donryu rats were given an intrarectal dose of N-methyl-N'-nitro-N-nitrosoguanidine (MNNG: 125 mg daily for 4 wk) followed by colostomy (COS) in some rats at 14 or 24 wk. By 60 wk, none of the rats with COS at 14 wk had tumors in the segment distal to the COS; tumors developed in the segment proximal to the COS at the same incidence as that in controls. Animals with COS at 24 wk had tumors in the distal segment, but they were fewer and smaller than those in the controls. Control animals without COS had multiple tumors distributed diffusely in the distal large bowel. The results suggest that luminal contents play a significant role, as cocarcinogens or promoters, in the induction of large bowel cancer. Male and female F344 rats were given a single intrarectal dose of MNNG (4 mg) followed by lithocholic acid or taurodeoxycholic acid (both 1 mg 5x/wk for 11 mo). After 78 weeks, the large bowel tumor incidence of rats treated with MNNG plus bile acid was twice that of rats treated with MNNG alone. Bile acids alone did not produce any tumors. Weanling female F344 rats were fed a diet high or normal in protein and fat. Starting at 7 wk of age, the animals received 1,2-dimethylhydrazine (DMH: 10 mg/kg sc for 20 wk). Ten wk after the last injection, animals fed high pro-

tein-high fat diets had a higher incidence of large bowel tumors than those on the normal diet. The results suggest that a diet high in fat and protein plays a significant role in large bowel carcinogenesis, and they support the concept that colon cancer does not develop, even in the presence of moderate amounts of carcinogen, without the promoting effect of bile acids. (25 refs)

- 78-5685 Inhibition of Morphological Transformation Induced with N-Methyl-N'-nitro-N-nitrosoguanidine in Cultures of Hamster Embryo Cells by 5'-Bromo-2'-deoxyuridine-Photolysis.** (Eng) Mironescu, S. G. (Dept. Pathology, Univ. South Carolina, Sch. Medicine, Columbia, SC, 29208). *Int J Cancer* 22(3): 304-314; 1978.

Syrian hamster embryo cells (HEC) were exposed to 5'-bromo-2'-deoxyuridine (BUdR: 3 µg/ml), N-methyl-N'-nitro-N-nitrosoguanidine (MNNG: 0.1-0.5 µg/ml), and fluorescent light (FL: 10-60 min) according to four variants of a protocol termed BUdR-MNNG-photolysis (BMP) to determine whether type III transformed foci induced by MNNG originate from the small subpopulation of cells stimulated by the carcinogen to enter DNA synthesis. Density-inhibited monolayers of HEC were exposed to FL during the last 30 min of the variable treatment periods. Quantitative transformation assays indicated that in nonilluminated cultures, addition of BUdR to MNNG-containing medium affected the transformation frequency of HEC. That is, incidence of type III foci did not subside at later intervals during the post-MNNG administration period as it did in the absence of BUdR. However FL irradiation of HEC treated with MNNG in the presence of BUdR inhibited the development of type III foci markedly. It is possible that BUdR is incorporated into potentially transformed cells during carcinogen-stimulated DNA synthesis and sensitizes them to FL irradiation. The events on which BUdR-induced sensitization of the cells to FL depended took place 5-15 hr after MNNG administration; ie, during the period of max carcinogen-stimulated DNA synthesis. FL and/or BUdR were not able to transform HEC. These results provide strong indirect arguments for the concept that carcinogen-induced DNA synthesis and the initiation of transformed clones are causally related. (51 refs)

- 78-5686 Factors Influencing the Frequency of Mitomycin C-induced Sister-Chromatid Exchanges in 5-Bromodeoxyuridine-substituted Human Lymphocytes in Culture.** (Eng) Ishii, Y. (Dept. Fundamental Radiology, Faculty Medicine, Osaka Univ., Kita-ku, Osaka 530, Japan); Bender, M. A. *Mutat Res* 51(3): 411-418; 1978.

Possible interactions between 5-bromodeoxyuridine (BUdR, 25 µM) and mitomycin C (MMC, in various concentrations) for the induction of chromosome aberrations and sister chromatid exchanges (SCE's) in cultured human peripheral

lymphocytes were studied. MMC induced chromatid aberrations, but there was no indication of synergism between MMC and BUdR. There was also no evidence of synergism between MMC and BUdR for the induction of SCE's. Treatment with MMC starting from the initiation of the lymphocyte cultures had more effect than treatment only during the last 12 hr of culture, and the SCE yield increased with the length of time MMC was present up to 30 hr. MMC-induced DNA lesions persisted long enough so that most of those present prior to S phase of the first cell cycle caused additional SCE's in the third cycle. (23 refs)

- 78-5687 Induction of Long-Lived Chromosome Aberrations as Manifested by Sister-Chromatid Exchanges in Lymphocytes of Animals Exposed to Mitomycin-C.** Stetka, D. G. (Biomedical Div., Lawrence Livermore National Lab., Livermore, CA, 94550); Minkler, J.; Carrano, A. V. *Mutat Res* 51(3): 383-396; 1978.

The cytogenetic effects of repeated vs acute exposure to mitomycin C (MMC, 2 or 4 mg/kg/wk for 1 or 8 wk) were determined using male New Zealand rabbits. The induction of sister chromatid exchanges (SCE's) was determined in bromodeoxyuridine-treated cultures of the animals' peripheral lymphocytes. MMC increased the SCE frequency within 1 day of treatment, the increases being accompanied by increases in the blood concentrations of lymphocytes and decreases in the concentrations of neutrophils. The SCE frequencies returned to intermediate or control levels within 3 wk; they remained at these levels in two animals, but abruptly increased in a third after 42 days. Repeated exposure to MMC was at least as effective as acute exposure to the same total dose in inducing increased SCE's. The mean SCE frequency remained at approx 2 to 2.5 times control level until at least day 175 (126 days after the last injection). Lymphocyte counts and elevated neutrophil counts returned to normal within 3 to 4 wk after the last injection. Analysis of lymphocyte SCE's may be an excellent assay for determining the results of repeated low-level exposure to mutagenic/carcinogenic chemicals. (40 refs)

- 78-5688 Morphogenesis and Micromorphology of Renal Tubular Tumors Induced in the Rat Kidney by Nitrosomorpholine. II. Tubular Glycogenosis and the Induction of Clear or Acidophilic Cell Tumors.** (Ger) Bannasch, H. (Abteilung für Cytopathologie, Institut für experimentelle Pathologie, Deutschen Krebsforschungszentrum Heidelberg, Im Neuenheimer Feld 280, D-6900 Heidelberg 1, W. Germany); Krech, R.; Zerban, H. *Z Krebsforsch* 92(1): 1-10; 1978.

Light and electron microscopy was used to study the development of clear and acidophilic (granular) cell kidney tumors in Sprague-Dawley rats given 12 or 50 mg/100 ml nitrosomorpholine (NNM) in their drinking water for 3-

At the end of treatment, the kidneys were morphologically similar to those of control rats. Some weeks later, however, excessive amounts of glycogen (glycogenesis: GGOS) were stored in single tubules. After a lag period of 22-97 wk, clear GGOS affected > 50% of the rats, and clear and/or acidophilic cell kidney tumors developed in 25%-30%. Variations in intermediate stages between GGOS (which occurred as early as 3 wk after withdrawal of NNM) and tumors were observed, which indicates that the former is a preneoplastic lesion. In addition to clear cells, the tumors contained predominantly cells that stored lipids or were rich in mitochondria. The tumor ultrastructure pointed to the differentiated tubules as the target tissue. The glycogen of the clear cells was located preferentially in the cytoplasm, but it was also found enclosed in autophagic vacuoles. The lipids formed membranous cytoplasmic bodies with a regular pattern of dark and light lines. The MC of the tumor cells, especially the acidophilic cells, were often characterized by a marked reduction in cristae. Large MC lacking in cristae but with an abundant matrix were sometimes found. Interstitial cells rich in acid mucopolysaccharides were distributed loose throughout the clear cell tumors. It is suggested that the GGOS, lipidosis, and mucopolysaccharidosis signify a disturbance of the cellular metabolism that might be closely correlated with carcinogenesis. (48 refs)

5689 Morphogenesis and Micromorphology of Epithelial Tumors Induced in the Rat Kidney by Nitrosomorpholine. III. Oncocytic Tubules and Oncocytomas. (Ger) Bannasch, P. (Abteilung für Cytopathologie, Institut für experimentelle Pathologie, Deutsches Krebsforschungszentrum Heidelberg, Im Neuenheimer Feld 280, D-6900 Heidelberg 1, W. Germany); Krech, H.; Zerbán, H. *Z. Krebsforsch* 92(1): 87-104; 1978.

Long-Evans rats were given 12 or 50 mg/100 ml nitrosomorpholine in their drinking water for 3-14 wk, and the development of renal oncocytomas was followed by light and electron microscopy. At the end of treatment, the kidneys were morphologically similar to those of control rats; however, some weeks later, oncocytic (OC) transformation of tubular epithelia and other tubular lesions were common. Normal and OC tubular segments were often directly connected. OC tubules were present in 66% of the rats after a lag period of 22-97 wk. OC microadenomas originated from

these tubules and developed in 36% of the rats during this period. Approx 20% of the control rats had single OC tubules and 6% had microadenomas. In some tubules, the oncocytes were combined with cells that stored excessive amounts of glycogen or lipids. Mitochondria rich in cristae and with diverse structural anomalies accumulated in the cytoplasm. In some mitochondria, the cristae were aligned in parallel and were connected by cross bridges 15-20 nanometers wide. Intramitochondrial aggregates of glycogen surrounded by a single membrane were noted, along with vesicles of the endoplasmic reticulum, Golgi complexes, and lipid bodies with a regular pattern of dark and light lines. Numerous glycogen particles were found in the cytoplasm of some oncocytes. A few of these cells showed a homogeneously condensed nucleus and cytoplasm. Interstitial cells rich in mucopolysaccharides were often found between the oncocytes. The oncocyte is suggested to be a special type of neoplastically transformed cell. (35 refs)

See also:

- *(Rev.): 78-5401, 78-5402, 78-5403, 78-5404, 78-5405, 78-5406, 78-5407, 78-5408, 78-5409, 78-5410, 78-5411, 78-5412, 78-5413, 78-5414, 78-5415, 78-5416, 78-5417, 78-5419, 78-5420, 78-5421, 78-5422, 78-5423, 78-5424, 78-5425, 78-5426, 78-5427, 78-5428, 78-5429, 78-5430, 78-5431, 78-5432, 78-5433, 78-5436, 78-5437, 78-5438, 78-5445, 78-5448, 78-5449, 78-5450, 78-5451.
- *(Phys.): 78-5694, 78-5695, 78-5697, 78-5699, 78-5700, 78-5701, 78-5702, 78-5708, 78-5721, 78-5730.
- *(Viral): 78-5758, 78-5808, 78-5810, 78-5847.
- *(Immun.): 78-5857, 78-5861, 78-5866, 78-5867.
- *(Path.): 78-5873, 78-5874, 78-5876, 78-5878, 78-5888, 78-5890, 78-5894, 78-5895, 78-5896, 78-5898.
- *(Epid.-Biom.): 78-5922, 78-5926, 78-5929, 79-5931, 78-5932, 78-5937, 79-5938, 78-5939, 78-5940, 78-5952, 78-5954, 78-5959, 78-5960, 78-5964, 78-5967, 78-5968, 78-5969, 78-5971.

PHYSICAL CARCINOGENESIS

- 78-5690 **The Proliferative Status and Clonogenic Capacity of Tumor Cells in a Transplantable Rhabdomyosarcoma of the Rat Before and After Irradiation with 800 Rad of X-rays.** (Eng) Hermens, A. F. (Radiobiological Inst. TNO, 151 Lange Kleiweg, Rijswijk, The Netherlands); Barendsen, G. W. *Cell Tissue Kinet* 11(1): 83-100; 1978.

R-1 rhabdomyosarcomas transplantable in the inbred strain of WAG/Rij rats were labeled semicontinuously with (³H)-thymidine (³H-TdR) and subsequently irradiated with x-rays to estimate the rate of recruitment of nonproliferative cells (Q) into the proliferative cells (P). Animals received up to seven ip injections of ³H-TdR (0.5 μ Ci/g of body wt) within 30 hr and half of the tumors received an acute dose of 800 rad of 300 kV x-rays 45 min after the completion of the last labeling in order to reduce the fraction of well-oxygenated cells to <1% of the surviving clonogenic cells. Analysis of the variation in the fraction of labeled tumor cells as a function of time and of the number of injections showed that the rate of recruitment of Q cells into P cells was <1%/hr. Comparison of data on the fractions of labeled cell foci derived for irradiated and nonirradiated tumors indicated that clonogenic Q cells were more frequently present among cells which survived the irradiation. These results suggest that in R-1 tumors of approx 800 mm³ the clonogenic Q cell population contains a larger fraction of hypoxic cells than the clonogenic P cell population. It was also demonstrated that in vitro cultured cells isolated from in vivo labeled tumors did not reutilize breakdown products of ³H-labeled DNA to an extent which would significantly influence values derived for fractions of foci originating from labeled P cells. (25 refs)

- 78-5691 **Sensitivity of Human Lymphocytes to Radiation During Various Phases of the Cell Cycle.** (Rus) Kukushkina, L. M. (Inst. Genetics and Cytology, Minsk, USSR); Mikhalevich, L. S. *Dokl Akad Nauk BSSR* 22(4): 371-374; 1978.

To evaluate the radiosensitivity of cells in various phases of the cycle, human lymphocyte cultures were treated with phytohemagglutinin (PHA) and, 1-50 hr later, with x-rays (100 R). The yield of chromosome aberrations in cells irradiated 46 and 50 hr after treatment with PHA was max, 77.8% and 83.0%, respectively, vs 66.6% in cells irradiated 20 hr and 40.0% in cells irradiated 1 hr after treatment. (10 refs)

- 78-5692 **Mitotic Cycle and Production of Chromosome Structural Changes in Human Cells (Meeting**

Abstract). (Eng) Luchnik, N. (Inst. Medical Radio Acad. Medical Sciences USSR, Obninsk 249020, Dni Kaluga, USSR). *Mutat Res* 53(2): 224; 1978. (no refs)

- 78-5693 **Gene Mutation Systems in Mammals (Meeting Abstract).** (Eng) Malling, H. V. (Lab. Environmental Mutagenesis, Natl. Inst. Environmental H Sciences, P.O. Box 12233, Research Triangle Park, 27709); Valcovic, L. R. *Mutat Res* 53(2): 227; 1978. (no refs)

- 78-5694 **Studies on the Relationship Between SCE and Chromatid Aberrations in Endoreduplicated Cells (Meeting Abstract).** (Eng) Palitti, F. (Centro di Studi di Genetica Evoluzionistica, CNR, Universita di Roma, Rome, Italy); Becchetti, A.; Perticone, P.; Diana, G. *Mutat Res* 53(2): 247; 1978. (2 refs)

- 78-5695 **Non-disjunction Induction by Mutagenic Agents in Male Cells of the Northern Volga Region of the USSR (Meeting Abstract).** (Eng) Tate, A. (Dept. Radiation Genetics and Chemical Mutagenesis, Leiden, Leiden, Netherlands). *Mutat Res* 53(2): 273; 1978. (1 ref)

- 78-5696 **Does the Capacity to Rejoin Radiation-Induced DNA Breaks Decline in Senescent Mice?** (Eng) Ono, T. (Dept. Radiation Biophysics, Faculty Medicine, Univ. Tokyo, Tokyo, Japan); Okada, S. *Int J Radiat Biol* 33(4): 403-407; 1978.

The rejoining rates of radiation-induced DNA scissions in the livers of 2-, 14-, and 22-mo-old C57BL/6 mice and in the cerebellums of 2- and 22-mo-old mice were compared. Single-strand breaks induced in the DNA of the cerebellum were rejoined at a similar rate in both 2- and 22-mo-old mice. There was a half-time of 2 hr; after this time, the percentage of breaks remained at 15%-20% for up to 24 hr. Radiation-induced single-strand scissions of DNA in the livers were also repaired quickly, with a half time of 1.2 hr in 2-mo-old mice and 1.3 hr in 14- and 22-mo-old mice. However, the size of the slow or nonrejoinable fraction in the livers of old mice was very difficult to estimate because of the frequent and experimental variation of the inherent age-associated DNA breaks in the livers of old mice (31%-47% of the total radiation-induced breaks). The results indicate that the hepatic cells of old mice were able to rejoin radiation-induced DNA breaks.

scissions as quickly as those of young mice, but that age-associated DNA scissions could not be rejoined. (22 refs)

697 **Unrepaired Single and Double DNA Strand Breaks Shown Cytogenetically Following X-irradiation of Lymphocytes from Patients with Ataxia Telangiectasia (Meeting Abstract).** (Eng) Taylor, A. M. (Dept. Cancer Studies, Medical Sch., Birmingham B15 2TJ, England). *Mutat Res* 53(2): 274; 1978. (1 ref)

698 **Ozone, UV Light, Sunbathing, and Skin Cancer (Letter to Editor).** (Dan) Thorling, E. B. (Kraenkningssinstituttet, Aarhus, Denmark). *Ugeskr Laeg* 138: 2316-2317; 1978.

Measurements of atmospheric ozone levels have failed to give consistent results, which may be due to the use of different measuring techniques. Measurements performed in Denmark have shown no noteworthy change during the last 12 years, while other measurements indicated that current ozone levels are 20% lower than 1951 levels. In view of the skin cancer-inducing effect of UV radiation, any reduction of the ozone level, which increases the intensity of UV radiation, may have serious consequences. (6 refs)

699 **Protection Against Nitrosoguanidine or UV-Radiation Lethality by Streptomycin in Cyanophytes (Meeting Abstract).** (Eng) Sinha, B. D. (Dept. Botany, Univ. Bihar, Muzaffarpur 842001, India); Sinha, H. D. *Mutat Res* 53(2): 263-264; 1978. (no refs)

700 **The Isolation and Characterisation of a Damage-dependent Endonuclease from *Micrococcus luteus* (Meeting Abstract)** (Eng) Riazuddin, S. (Dept. Biophysics, Johns Hopkins Univ., 615 N. Wolfe St., Baltimore, MD, 21205); Grossman, L. *Mutat Res* 53(2): 252-253; 1978. (no refs)

701 **Mutation Studies with *Micrococcus radiodurans* Auxotrophs: Effects of Exposure to Radiation, Chemicals, and Combined Radiation/Chemical Treatment (Meeting Abstract).** (Eng) MacPhee, D. G. (Dept. Genetics, Monash Univ., Bundoora, Victoria 3083, Australia); Kaminskas, M. *Mutat Res* 53(2): 238-239; 1978. (no refs)

702 **Mutagenicity of 8-Methoxy-psoralen and Long-Wave Ultraviolet Irradiation (Meeting**

Abstract). (Eng) Simmons, J. W. (Dept. Radiation Genetics and Chemical Mutagenesis, State Univ. Leiden, Leiden, Netherlands); Burger, P. M. *Mutat Res* 53(2): 263; 1978. (no refs)

78-5703 **Chromosomes after Psoralen/UVA Treatment In Vitro and In Vivo (Psoriasis Patients) (Meeting Abstract).** (Eng) Brogger, A. (Genetics Lab., Norsk Hydro's Inst. Cancer Res., Montebello, Oslo, Norway); Waksman, H.; Thune, P. *Mutat Res* 53(2): 157-158; 1978. (no refs)

78-5704 **Lack of Cytotoxicity and Mutagenicity of Ultraviolet Radiation in Normal Diploid Human Fibroblasts as Contrasted with Cells Deficient in DNA Repair (Meeting Abstract).** (Eng) Maher, V. M. (Michigan State Univ., East Lansing, MI, 48824); McCormick, J. J. *Mutat Res* 53(2): 225-226; 1978. (no refs)

78-5705 **Photoreactivation of Pyrimidine Dimers in Human Cells (Meeting Abstract).** (Eng) Mortelmans, K. (Stanford Res. Inst., Menlo Park, CA); Friedberg, E. C.; Brorsson, M. *Mutat Res* 53(2): 234; 1978. (2 refs)

78-5706 **Co-operative Denaturing Effect of Heat and UV on DNA in Human Squamous Cells (Meeting Abstract).** (Eng) Roth, D. (Inst. Environmental Medicine, New York Univ., New York, NY). *Mutat Res* 53(2): 254-255; 1978. (no refs)

78-5707 **Technical and Theoretical Considerations in Evaluating Individual Variations in Human Unscheduled DNA Synthesis (UDS) (Meeting Abstract).** (Eng) Delozier, D. (Dept. Medical Genetics, Indiana Univ. and Methodist Hosp., Indianapolis, IN); Antley, R. *Mutat Res* 53(2): 176-177; 1978. (no refs)

78-5708 **Synergistic Interaction of Hydrogen Peroxide and Near-Ultraviolet Radiation (Meeting Abstract).** (Eng) Eisenstark, A. (Div. Biological Sciences, Univ. Missouri, Columbia, MO); Ananthaswamy, H. N.; Hartman, P. S. *Mutat Res* 53(2): 184; 1978. (no refs)

78-5709 **The Kinetics of Thymine Dimer Excision in Ultraviolet-irradiated Human Cells.** (Eng) Ehmann, U. K. (Lab. Experimental Oncology, Dept. Pathology,

Stanford Univ. Sch. Medicine, Stanford, CA, 94305); Cook, K. H.; Friedberg, E. C. *Biophys J* 22(2): 249-264; 1978.

The kinetics of thymine dimer loss from the acid-insoluble fraction of UV-irradiated human fibroblast cells was investigated, and the findings were compared with data on repair synthesis. UV fluences of 10-40 joules/m² produced an av of 21-85 x 10³ thymine dimers per cell and an eventual maximal loss per cell of 12-20 x 10³ thymine dimers. The time for half-maximal loss of dimers ranged from 12 to 22 hr after UV irradiation. In contrast, the time for half-maximal repair synthesis of DNA measured by autoradiography was 4.5 hr. This figure is in good agreement with reported half-maximal repair synthesis times, which range from 0.5 to 3.6 hr. The discrepancy in the kinetics of the loss of thymine dimers from DNA and repair synthesis may be explained by one of the following: (1) the loss of thymine dimers from the DNA was not equated with thymine dimer excision, because it is possible that pyrimidine dimers are excised as acid-precipitable oligonucleotides that are then reduced to acid-soluble products by enzymatic degradation; (2) the different treatment of the cells inherent in the techniques used in measurement of repair synthesis and dimer excision; (3) repair synthesis may not reflect pyrimidine dimer excision in mammalian cells, and (4) the rapid kinetics of repair synthesis compared with the slower loss of thymine dimers may reflect a patch and cut-type mechanism in which repair synthesis occurs before the enzymatic removal of the dimers. (58 refs)

78-5710 Cancer of the Colon as a Late Sequel of Pelvic Irradiation. (Eng) Greenwald, R. (Dept. Medicine, Div. Gastroenterology, Univ. Miami Sch. Medicine, P.O.B. 520875 Biscayne Annex, Miami, FL, 33152); Barkin, J. S.; Hensley, G. T.; Kalser, M. H. *Am J Gastroenterol* 69(2): 196-198; 1978.

A 36-yr old woman developed a carcinoma of the rectum 9 yr after therapeutic radiation for a pelvic malignancy. Prolonged follow-up is urged for 5-yr survivors of pelvic irradiation, since the symptoms of radiation-induced proctocolitis may mimic those of carcinoma of the colon and rectum. (7 refs)

78-5711 Fibrosarcoma in a Previously Irradiated Larynx. (Eng) Donaldson, I. (Manchester Royal Infirmary, Oxford Road, Manchester M13 9WL, England). *J Laryngol Otol* 92(5): 425-428; 1978.

A 68 yr-old man developed a laryngeal tumor at a site that had been treated by radiotherapy for a squamous cell carcinoma. The patient had received a tumor dose of 5,600 R delivered in 12 treatments 11 yr previously. It is suggested that the neoplasm was induced by the radiation. (18 refs)

78-5712 Parathyroid Adenoma and Pre Head-and-Neck Irradiation. (Eng) Schachar, S. H. (Fairfax Hosp., Falls Church, VA); Hall, A. *Ann Intern Med* 88(6): 804; 1978.

The occurrence of a parathyroid adenoma in a 48-yr-old man and a 35-yr-old woman who had previously been treated by head and neck radiation is reported. The man had been irradiated extensively for facial acne at age 23, and the woman had received radiation therapy for thymic enlargement in infancy and later for acne. (5 refs)

78-5713 Sensitivity of the Thyroid and Parathyroid Glands to Radiation. (Eng) Clark, O. H. (Surgical Service (112), Veterans Admin. Hosp., 4150 Clement St., San Francisco, CA, 94121); Rollo, D.; Stroop, J.; Casper, B.; Rehfeld, S. J.; Loken, H. F.; Defetos, L. J. *J Surg Res* 24(4): 374-379; 1978.

The effect of radiation on male Sprague-Dawley rat thyroid and parathyroid glands, the sensitivity of thyroid follicular and parafollicular cells to this radiation, and the effect of thyroid-stimulating hormone (TSH) on the sensitivity of the gland to radiation were determined. All animals were treated by hemiparathyroidectomy and hemithyroidectomy; half the animals had a ¹²⁵I seed implanted next to the remaining thyroid gland. All animals were then placed on one of three diets: (1) low-iodine diet (LID) and deionized water, (2) lab chow and tap water, (3) lab chow and L-sodium thyroxine (6 µg/100 g/day) in the drinking water. Blood samples were obtained at 4 and 10 wk; at 10 wk, all animals were removed and the animals placed on lab chow and tap water. Blood samples were taken again 2, 4, and 9 wk after removal and the animals were sacrificed. ¹²⁵I seeds selectively inhibited thyroid function compared with parathyroid function. Stimulation of thyroid tissue by TSH and suppression of this tissue by T₄ did not alter its sensitivity to radiation, since serum TSH concentrations were not different in the T₄-suppressed, control, and TSH-stimulated animals 9 wk after the radiation was discontinued. TSH stimulation by the LID and TSH suppression by exogenous T₄ did alter the function of the parafollicular C cells, since serum calcitonin concentrations were increased with the former treatment and decreased with the latter. Serum Ca concentrations were also increased in the LID group but the reason for this increase is not known. (23 refs)

78-5714 Irradiation-induced Polyglandular Neoplasia of the Head and Neck. (Eng) Swelstad, J. A. (Surgical Oncology, Northwestern Univ. Medical Center, Evanston, IL, 60201); Scanlon, E. F.; Oviedo, M. A.; Hersh, N. E. *Am J Surg* 135(6): 820-824; 1978.

The occurrence of 13 salivary gland tumors and 8 parathyroid adenomas in 18 patients with a history of radiotherapy to the head and neck is reported. Nine of these patients

had thyroid neoplasms, 37% of which were malignant. 37% of which showed some degree of radiation thyroiditis. The salivary gland tumors and thyroid nodules were often discovered simultaneously in the same patient. Nine of the salivary gland tumors were of parotid origin. All three parotid gland tumors and six of the parotid tumors were malignant. The latent period for the salivary tumors ranged 30 yr between irradiation and diagnosis. Of the 10 patients with parathyroid adenomas, one had three hyperplastic parathyroid glands and seven had single adenomas. The latent period for the adenomas averaged 27 yr. All 10 patients remained normocalcemic after surgical management, although the presurgical hypercalcemia in four was asymptomatic. Examination of the salivary glands and thyroid recommended for patients with a history of radiation to head and neck. (10 refs)

5715 Radiation-induced Cancer of the Thyroid. (Eng) Sonenberg, M. (Endocrinology Service, Dept. of Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY); Leeper, R. D. *Clin Bull* 8(1): 29-33; 1978.

Among 64 persons with a history of head and neck radiation, 10 had thyroid abnormalities were found in 5/11 irradiated patients, 2/12 for hemangioma, 1/9 for adenoids, and 4/11 for miscellaneous reasons. None of 21 patients treated for an enlarged thymus had abnormalities. The thyroid in the acroparathyroidism ranged from 0.5 to 2 cm. (12 refs)

5716 Radioiodine Treated Hyperthyroidism and Thyroid Carcinoma. (Eng) Appell, R. (Dept. Surgery, George Washington Univ. Medical Center, 2150 Pennsylvania Ave., N.W., Washington, DC, 20037); Tsangaris, A.; Spiegel, J. *Am Surg* 44(8): 537-540; 1978.

Papillary thyroid carcinoma developed 4 yr after radioiodine therapy for hyperthyroidism in a 48-yr-old woman. Statistics on 14 similar literature cases are tabulated. (22 refs)

5717 A Nine-Year Cytogenetic Follow-up of a Patient Injected with Thorotrast. (Eng) Ishihara, T. (Div. Radiation Health, Natl. Inst. Radiological Sciences, Chiba 280, Japan); Kohno, S.; Hayata, I.; Kumatori, T. *Hum Genet* 42(1): 99-108; 1978.

A 62-yr-old man who had been given > 15 ml Thorotrast in 1939 was subjected to cytogenetic follow-up for 9 yr before his death in 1974. No malignancy resulted, but there was extensive Thorotrast deposition in the liver and several lymph nodes. The estimated doses of radiation absorbed by liver, spleen, and bone marrow were 32, 653, and 4 rad/yr, respectively. Both Cu (dicentric, rings, and acentric frag-

ments) and Cs (reciprocal translocations, inversions, deletions, duplications) types of chromosomal abnormalities were observed in the peripheral T lymphocytes at frequencies of 4.5%-9.0% and 6.0-11%, respectively, throughout the follow-up period. All of the chromosomal abnormalities in the bone marrow cells were of the Cs type, and they ranged in frequency from 2.17% to 5.4% during the first three samplings. An increase in frequency (up to 26%) at the last sampling was mainly due to cells lacking a Y chromosome (19% of all cells). The 50 aberrant bone marrow cells analyzed consisted primarily of cells lacking a Y chromosome (19 cells) and cells of clone A (25). Fifteen of the cells lacking a Y chromosome possessed an additional reciprocal translocation involving chromosomes 12 and 17. Clone A cells were observed at an average frequency of about 4%. The karyotypes of clone A cells from the bone marrow and peripheral lymphocytes were identical, involving translocation of part of the short arm of chromosome 2 onto the short arm of 22. Clone B occurred in the peripheral lymphocytes at an average frequency of 0.43%; it was also identified in one bone marrow cell. Cells lacking a Y chromosome were not identified in the peripheral lymphocytes. The finding of Clone A suggests the existence of a genetic factor related to the development of chronic myelocytic leukemia. (27 refs)

78-5718 Thyroid Cancer Risk from Iodine-131 Treatment? (Ger) Creutzig, H. (Department für Radiologie, Medizinische Hochschule, Postfach 610 180, D-3000 Hannover 61, W. Germany); Hundeshagen, H. *Med Klin* 72(19): 855-857; 1977.

The case histories of 240 thyroid carcinoma patients revealed that 5 had ¹³¹I treatment for benign thyroid diseases 0.5-5 yr before the diagnosis of carcinoma and 1 had x-irradiation of the thyroid gland 30 yr before the appearance of carcinoma. For comparison, the prevalence of thyroid carcinoma in a region without endemic goiter was found to be 2% to 5%. The findings indicate that there is no relationship between the incidence of clinically manifest thyroid carcinoma and ¹³¹I therapy. (17 refs)

78-5719 Occupational Exposure to Ionising Radiation: The Risk in Perspective. (Eng) Bonnell, J. A. (Nuclear Health and Safety Dept., Central Electricity Governing Board, London, England); Harte, G. *Lancet* 1(8072): 1032-1034; 1978.

The risk of somatic and genetic disease incurred by exposure to ionizing radiation was calculated using the International Commission on Radiological Protection risk factors and a simple risk-time relationship. For an individual exposed at 10 millisieverts (mSv)/yr from age 20, the risk of death from radiation-induced leukemia is between 1 in 100,000 and 1 in 50,000. (One sievert equals 100 rem.) If the worker has children at age 40, the risk of a serious genetic disease in his

descendants is 1 in 500 for the first two generations and 1 in 250 for subsequent generations. If a person is exposed from birth to 1 mSv/yr from nuclear plant operations and he receives a total of 25 mSv by age 25, the risk of genetic disease in his descendants will be 1 in 2,000. (8 refs)

78-5720 Occupational Exposure to Ionising Radiation (3 Letters to Editor). (Eng) Lindop, P. J. (Dept. Radiobiology, Medical Coll. St. Bartholomew's Hosp., London EC1M 6BQ, England); Mole, R. H.; Reissland, J. A.; Dolphin, G. W. *Lancet* 1(8074): 1155-1157; 1978.

A previous paper indicated that an individual exposed to 10 millisieverts (1 Sievert = 100 rem) from age 20 incurs a risk of death from radiation-induced leukemia of 1 in 100,000 to 1 in 50,000. This is actually the annual risk; the total risk is 1 in 4,000. Expert testimony on the effects of occupational exposure to radiation on cancer incidence is challenged. Finally, the results of a study of cancer incidence in Portsmouth Naval Shipyard workers should not be cited as fact because of the incompleteness of the data and the uncertainty in the categorization of radiation and nonradiation workers. (5 refs)

78-5721 Low Dose-Rate Experiment with Tritiated Thymidine as a Simulator of Chemical Mutagens Using Silkworm Oocyte System (Meeting Abstract). (Eng) Tazima, Y. (Natl. Inst. Genetics, Misima, Japan). *Mutat Res* 53(2): 274-275; 1978. (no refs)

78-5722 Effects of Whole-Body Irradiation on Neonatally Thymectomized Mice: Incidence of Benign and Malignant Tumors. (Eng) Anderson, R. E. (Dept. Pathology, Univ. New Mexico Sch. Medicine, Albuquerque, NM, 87131); Howarth, J. L.; Troup, G. M. *Am J Pathol* 91(2): 217-228; 1978.

The individual and cumulative effects of neonatal thymectomy (nTx) and irradiation on benign and malignant tumor incidence were determined in germ-free Charles River mice. The mice underwent either thymectomy or sham thymectomy (STx) within 24 hr of birth; at 6 wk of age, mice from both groups were exposed to 700 rads' whole-body irradiation. nTx increased the incidence of benign and malignant tumors and the prevalence of multiple primary neoplasms in individual mice, but it had no effect on the latent period. Irradiation further increased the incidence of benign and malignant tumors in both groups, but the increment was much more pronounced the STx mice than the nTx mice. In addition, irradiation accelerated the appearance of these tumors in both groups, but the effect was more pronounced in STx mice. Thus, the cumulative effects of thymectomy plus irradiation were less pronounced than the sum of the individual effects. Some of the carcinogenic effects of

whole-body irradiation may be mediated by suppression of the thymic-dependent component of the immune system (11 refs)

78-5723 Rat Skin Tumor Incidence Following Single Fractionated Exposures to Proton Radiation. (Eng) Burns, F. J. (Inst. Environmental Medicine, New York Univ. Medical Center, New York, NY, 10016); Strickland, P.; Vanderlaan, M.; Albert, R. E. *Radiat Res* 74(1): 152-157; 1978.

The dose-response curve and degree of split-dose recovery of tumor induction and hair follicle survival were determined in male albino CD rat skin irradiated with protons. A proton beam from a cyclotron was modified to provide an approximately linear decrease in dose with depth in tissue to a maximum of 1.3 mm. Six doses were given in the range 75-750 rads. Tumor yield at 100 wk postirradiation increased as 2.8 times the dose (D). Split doses consisted of an initial dose of 150 rads followed 24 hr later by graded doses in the range 150-600 rads. Split doses produced a sparing effect on follicle kill and on yield of tumors. The magnitude of the reduction in tumor yield was consistent with complete recovery. Comparisons with previous results indicated that protons were 2.0 times as effective as electrons for tumor induction and that the shapes of the dose-response curves were generally similar. (11 refs)

78-5724 Reactions of Radicals with Lecithin Bilayers. (Eng) Barber, D. J. (Chemistry Dept., Univ. Notre Dame, Notre Dame, IN, 46556); Thomas, J. K. *Radiat Res* 74(1): 51-65; 1978.

The reaction rates of hydroxyl radicals (OH) and the hydrated electron $e(aq)^-$ with lecithin bilayers were determined. The rate for OH + lecithin was $5.1 \pm 0.9 \times 10^8$ /M, while that for $e(aq)^-$ + lecithin was very slow. With pyrene in lecithin, the reaction rates for OH and $e(aq)^-$ were $1.65 \pm 0.12 \times 10^9$ and 7×10^7 /m/sec, respectively. These rates were lower than those observed for similar reactions in homogeneous solution. The difference is attributable to the protective effect of the bilayer and to the restricted diffusion imposed on the reactive species by the bilayer. Long-term alteration in the model membrane following OH attack is manifested by damage to the head group (which increases the water penetration of the bilayer) and by cross-linking within the membrane (which restricts motion in the bilayer interior). These results are relevant to studies of the effects of low linear-energy-transfer radiation on biological membranes. (27 refs)

78-5725 Chromosome Aberrations Induced in Human Lymphocytes by Radiation from ^{252}Cf . (Eng) Lloyd, D. C. (Natl. Radiological Protection Board, Harwell, UK).

ot, Oxfordshire OX11 ORQ, England); Purrott, R. J.;
er, E. J.; Edwards, A. A.; Dolphin, G. W. *Int J Radiat*
34(2): 177-186; 1978.

induction of chromosome aberrations in lymphocytes
a healthy adult male by ^{252}Cf neutron
tion was studied and compared with aberration induc-
by other types of radiation. An analysis of the dose-effect
in which the dose (D) of Cf neutrons was related to
ield (Y) of neutron-induced dicentric aberrations re-
d that for the two highest energy neutron curves (7.6
4.7 million electron volts), the best fit was obtained with
ear-quadratic expression in which Y was a function of
us D^2 . For lower energy neutron curves, the best fit was
ned with a simple linear expression in which Y was a
ion of D. The relative biological effectiveness (RBE) for
with respect to ^{60}Co was about 27 at very low doses,
easing to 6 at an aberration Y equivalent to 400 rads of
ads/hr gamma radiation. A profile of chromosome aber-
on induction with depth was obtained by placing blood
ples at distances of 0.65-2.05 cm from the Cf source. The
E of Cf radiation relative to ^{226}Ra gamma radiation
eased with distance from the source and more than dou-
over the distance range. This increasing RBE indicates
Cf is more effective at a greater distance in producing
romosome aberrations and, presumably, in destroying the
ty of cells to divide. This effect may be an advantage in
treatment of large tumors. (16 refs)

5726 **Metabolism and Microdistribution of Californi-
um-252 and Its Effects on Pulmonary Paren-
ma During Intratracheal Administration to Rats.** (Rus)
rova, E. S. (No affiliation given.); Stepanov, S. V. *Radi-
logiia* 18(5): 724-729; 1978.

andom-bred albino rats were given a single intratracheal
e of californium-252 (0.016, 0.004, or 0.0005 $\mu\text{Ci/g}$).
hin 1-600 days of exposure, the rats were sacrificed and
g specimens were studied histologically and radiometri-
y. Structural changes in the bronchial epithelium and in
ndothelium of the vessels (karyopyknosis, vacuolization,
desquamation) were seen on days 7-14 after Cf252 ad-
istration (rats treated with 0.016 $\mu\text{Ci/g}$ also showed de-
ection of their alveolar epithelium). Starting from days 30,
and 210, rats treated with 0.016, 0.004, and 0.0005 $\mu\text{Ci/g}$,
ectively, had sclerotic changes in the lung tissue, hyper-
ia of the bronchial and alveolar epithelium, and meta-
ia of the columnar epithelium. Of six rats who were ex-
ed to 0.0005 $\mu\text{Ci/g}$ and survived 600 days, four developed
g tumors (1 adenocarcinoma, 2 squamous cell carcinomas,
acrofollicular lymphoblastoma). (3 refs)

5727 **Effect of Combined Alginate Treatments on the
Distribution and Excretion of an Old Radios-
tium Contamination.** (Eng) Vanderborght, O. L. (Miner-

al Metabolism Lab., Radiobiology Dept., Belgian Nuclear
Centre, S.C.K./C.E.N., B 2400 Mol, Belgium); Van Puym-
broeck, S.; Babakova, I. *Health Phys* 35(2): 255-258; 1978.

Mobilization of ^{85}Sr from the skeleton was studied in mice
through combined treatment with sodium alginate incor-
porated in a dough plus alginate injected ip as an aqueous
solution. Two-month-old C57 Black male mice were injected
ip with 40 μCi $^{85}\text{SrCl}_2$. After 9 wk, mice were divided into
five experimental groups: 1 was fed alginate dough containing
100 g sodium alginate and injected with NaCl; 2 was fed
alginate dough and injected ip with 2 mg sodium alginate
2x/day; 3 was fed agar dough and injected with NaCl; 4 was
fed agar dough and injected with sodium alginate; 5 was fed
standard pellets and injected with NaCl. Treatment lasted 3
days during which urine and excreta were collected and
analyzed for ^{85}Sr content. After the 3 days, mice were killed
and ^{85}Sr content of blood, liver, spleen, kidney, and femur was
determined. Levels in blood, liver and kidney increased when
either alginate dough or injection was given; excretion in the
feces was increased in mice fed alginate; and excretion in the
urine increased in mice injected with alginate. Mice in group
2 had a significantly high increase in ^{85}Sr in blood without
an equivalent increase in fecal ^{85}Sr , indicating that the ^{85}Sr
mobilized by injected alginate was not fully available for in-
testinal excretion. However, about half the ^{85}Sr mobilized in
the blood by alginate injection was available for renal excre-
tion. Combined treatment overcame the decrease in urinary
 ^{85}Sr excretion obtained with alginate in diet only. The ra-
dionuclide content of liver, kidney, and spleen was increased
four- to six-fold by the combined treatment. (9 refs)

78-5728 **Antigenicity of ^{90}Sr Induced Tumors (2 Letters
to Editor).** (Eng) Nilsson, A. (Dept. Pathology,
Coll. Veterinary Medicine, Swedish Univ. Agricultural
Sciences, S-750 07 Uppsala, Sweden); Revesz, L.; Larsson,
S. E.; Lorentzon, R.; Boquist, L. *Acta Pathol Microbiol Scand*
(A) 86(4): 331-332; 1978.

Facts underlining the difficulties of interpreting and compar-
ing two studies of the immunostimulating effect of BCG on
 ^{90}Sr -induced osteosarcomas are given. Factors such as time
of BCG treatment, ^{90}Sr dose, and duration of animal survival
differed between the two and affected tumor incidence. (4
refs)

78-5729 **Radium-induced Malignant Tumors of the Mas-
toid and Paranasal Sinuses.** (Eng) Littman, M.
S. (Center Human Radiobiology, Argonne Natl. Lab., Ar-
gonne, IL, 60439); Kirsh, I. E.; Keane, A. T. *Am J Roent-*
genol 131(5): 773-785; 1978.

A survey of records kept at the Center for Human Radiobi-
ology (Argonne National Laboratory, Argonne, IL) revealed
21 patients with mastoid carcinoma (MC) and 11 with malig-

nant paranasal sinus tumors (PST) among 5,058 persons occupationally or therapeutically exposed to radium. People were exposed via the luminous watch-dial industry, the industrial and research processing of Ra, and medical treatment. Most of the patients (28) were women employed in the dial-painting industry. The four men received Ra therapeutically. Tumor induction times were 21-50 yr for the MC (median, 33) and 19-52 yr for the PST (median, 34). Of the 27 patients whose body burdens of Ra were measured, the calculated skeletal dose ranged from 605 to 19,519 rads; 25 of these patients had skeletal doses > 1,000 rads. Of the total 2,036 persons (including the 27 patients) whose Ra body burdens were measured, 200 had a skeletal dose > 1,000 rads; the frequency of the two cancers among the latter was relatively constant with dose and averaged 13%. There was a high proportion of mucoepidermoid carcinoma, comprising 38% of the MC and 36% of the PST. Three patients had antecedent bone sarcoma at 20, 11 and 5 yr, respectively, and a bone sarcoma was discovered at autopsy in a fourth patient. Radiographic changes in the mastoid and paranasal sinuses were similar to those in non-Ra malignant tumors. More than 800 persons exposed to Ra before 1930 and another group of unknown size who received Ra therapeutically are still at risk of developing MC and PST. (46 refs)

78-5730 Carcinogenesis in Tissue Culture. 29: Neoplastic Transformation of a Normal Human Diploid Cell Strain, WI-38, with Co-60 Gamma Rays. (Eng) Namba, M. (Dept. Pathology, Kawasaki Medical Sch., Kurashiki 701-01, Japan); Nishitani, K.; Kimoto, T. *Jpn J Exp Med* 48(4): 303-311; 1978.

Neoplastic transformation of a normal human diploid cell strain, WI-38, by irradiation with ^{60}Co γ -rays is described. In preliminary studies, a series of potentially carcinogenic agents was tested for transforming ability with human lymphocytes. 4-Nitroquinoline 1-oxide (3 μM for 1 hr) was the most effective chemical examined for the production of chromosome abnormalities (approx 10% aberrations), but its effectiveness was limited by its cytotoxicity. ^{60}Co γ -rays, however, induced very prominent chromosome aberrations with a dose-effect relationship. WI-38 cells irradiated four times with 300-400 rads for a total dose of 1,400 rads showed neoplastic transformation starting about 150 days after the last irradiation. The transformed cells exhibited an epitheliallike morphology, abnormal karyology, and the B-type isoenzyme pattern of glucose-6-phosphate dehydrogenase, and they produced sarcomas when transplanted into the cheek pouches of hamsters treated with anti-hamster thymocyte serum. They have been cultured for 13 mo without showing aging phenomena. (25 refs)

78-5731 Removal of Plutonium from Beagles Using Ca-DTPA and Zn-DTPA: Effects of Initial DTPA Injection. (Eng) Taylor, G. N. (Radiobiology Lab.,

Univ. Utah, Salt Lake City, UT, 84112); Lloyd, R. D.; Farman, J. J.; Atherton, D. R.; Mays, C. W. *Health Phys* 35(2): 201-210; 1978.

The use of trisodium calcium diethylenetriaminepentaacetate (Ca-DTPA) and zinc diethylenetriaminepentaacetate (Zn-DTPA) for prompt treatment of radioactive plutonium (^{239}Pu) exposure was compared in the dog. Sixteen beagles about 23 mo of age were each injected with 1 μCi ^{239}Pu as Pu(IV) citrate and then divided into four groups of two males and two females each. About 1/2 hr later, Group A dogs were injected iv with 30 μmol Ca-DTPA/kg, Group B and C dogs with 300 μmol Zn-DTPA/kg, and dogs in Group D dogs with 34 μmol Zn-DTPA/kg. Subsequent treatments for Groups A, B, C, and D were single daily sc injections of 34 μmol /kg Zn-DTPA. Group B received 6.8 μmol Zn-DTPA/kg 5 x/day (total dose, 34 μmol /kg/day). Pu retention in liver and nonliver tissue was determined 3, 7, and 14 days after radionuclide injection. Retention in livers and kidneys was also measured after the dogs were killed at 14 days. There were no significant differences in total-body retention at 2 wk for Groups A, B, and C (190, 230, and 26%, respectively); in each of these groups retention was significantly lower than in Group D (48%). Liver ^{239}Pu content was significantly lower in Groups A, B, and C (5.50, 4.30, and 4.4%, respectively) than in Group D (14.4%) or controls (35%) at 14 days. Kidney content did not differ among the groups. Prompt treatment with 300 μmol Zn-DTPA/kg or 30 μmol Ca-DTPA/kg provided superior clearance of radioactive Pu. It did not matter whether the subsequent daily treatment with 34 μmol Zn-DTPA/kg was given in a single injection or in five injections. (40 refs)

78-5732 Early Retention of ^{239}Pu and ^{241}Pu in Male Beagles. (Eng) Lloyd, R. D. (Radiobiology Lab., Univ. Utah, Salt Lake City, UT, 84112); McFarland, S.; Atherton, D. R.; Bruenger, F. W.; Taylor, G. N.; Mays, C. W. *Health Phys* 35(2): 211-215; 1978.

Early retention, excretion, and tissue distribution of $^{239}\text{Pu(IV)}$ citrate (0.05-0.1 μCi /kg iv), to which tracer amount of the photon-emitter ^{237}Pu had been added, were studied in five beagle dogs, age 57-84 mo. Pu retention in liver and nonliver tissue, mainly skeletal, was measured periodically in living dogs for nearly 4 mo. The mature beagles retained about 30% Pu in the liver and 50% in the skeleton about 10-120 days after injection (3 mo-old beagles have been found to retain 12% in the liver and 68% in the skeleton). The relative distribution of skeletal Pu among various bones in the mature animals was similar to that seen previously in 23 mo-old dogs, but different from that of 3 mo-old beagles. An exception was the humerus for which there was a significant difference ($p > 0.2$) in the percentage of retained skeletal Pu for juvenile, young adult, and mature dogs. About 14%-20% of the injected Pu was excreted in urine and feces in the first 3 wk after injection. Less uniformity in excretion

retention was observed in mature animals than has been reported for younger dogs. (15 refs)

7733 Decorporation from Beagles of a Mixture of Monomeric and Particulate Plutonium using DTPA and Zn-DTPA: Dependence upon Frequency of Administration. (Eng) Lloyd, R. D. (Radiobiology Lab., Univ. Utah, Salt Lake City, UT, 84112); Boseman, J. J.; Taylor, G. N.; Bruenger, F. W.; Atherton, D. R.; Stevens, W.; Smith, C. W. *Health Phys* 35(2): 217-227; 1978.

Use of Zn-diethylenetriaminepentaacetate (DTPA) and Ca-DTPA in the removal of radioactive plutonium was studied in 14 young adult beagles (av age, 19.8 mo). The dogs were injected with 0.9 $\mu\text{Ci/kg}$ polymeric ^{239}Pu to which tracer amounts of gamma-ray emitting ^{237}Pu had been added. Two weeks later, four dogs were injected with 30 $\mu\text{moles/kg}$ Ca-DTPA/day and four dogs, with 30 $\mu\text{moles/kg}$ Zn-DTPA per wk. Total ^{237}Pu excretion in the first 24 hr averaged 10.4% after Ca-DTPA, 10.4% after Zn-DTPA, and 7.1% in controls. Corresponding levels after 28 days, corrected for radioactive decay, were 38.2%, 49.4%, and 12.1%, respectively. In the untreated dogs, there was a gradual translocation of ^{237}Pu from the liver to nonliver tissue (mainly skeletal) over a 6-23 wk period. Biological retention of ^{237}Pu in nonliver tissue remained constant for 6 wk in Ca-DTPA treated beagles, but the radionuclide in the liver decreased from 60% to 44%. In Zn-DTPA treated animals, activity from liver and nonliver tissue declined from 60% to 25% and 28% to 17%, respectively, over 6-wk period. It is concluded that daily injections of Zn-DTPA are more effective in the removal of incorporated plutonium than are weekly injections of Ca-DTPA. (38 refs)

7734 Plutonium Retention, Excretion, and Distribution in Juvenile Beagles Soon after Injection. (Eng) Lloyd, R. D. (Radiobiology Lab., Univ. Utah, Salt Lake City, UT, 84112); McFarland, S. S.; Atherton, D. R.; Smith, C. W. *Radiat Res* 75(3): 633-641; 1978.

In order to determine the possible effects of environmental plutonium on humans, ^{239}Pu was injected iv into six 3-mo-old beagles. Dogs were injected with 0.4 μCi ^{239}Pu as the citrate complex and 0.4 μCi ^{237}Pu to facilitate measurement of the excretion. All excreta were collected for the first 3 wk after injection, and Pu content determined by NaI spectrometry. About 10% of the injected Pu was excreted in the first 21 days, 5% in urine and 6% in feces. Retention in liver and in nonliver tissue (mainly skeleton) was measured serially in living dogs by a combination of total- and partial- body counting, utilizing the 60- and 100 kiloelectron volt photons from ^{237}Pu . The dogs were sacrificed at 7, 14, 28, 56, 89, and 119 days after injection, respectively. Total excretion for living dogs for the first 21-days collection period was roughly 11%, with about 5% in urine and 6% in feces. Early retention of Pu in liver

was lower in these dogs (11%-15%) than that observed previously in adults aged 17 mo (31%-37%), but early skeletal retention was greater for juveniles (62%-75%) than for adults (50%-60%). Pu content of most other tissues was similar in adults and juveniles. Compared with adults, juveniles had a significantly greater fraction (2- to 5-fold) of skeletal Pu in the paws and lower leg bones and a significantly lower fraction (2-fold) in the rib cage and thoracic and lumbar vertebrae. (12 refs)

78-5735 Trophic Transfer by Chironomids and Distribution of Plutonium-239 in Simple Aquatic Microcosms. (Eng) Trabalka, J. R. (Environmental Sciences Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Frank, M. L. *Health Phys* 35(3): 492-494; 1978.

The transfer of plutonium (Pu) from labeled organic sediments to chironomids, the most important component in the diet of juveniles and adults of many freshwater fish, was determined. Adult *Chironomus riparius* from a laboratory culture were stocked in aerated aquatic microcosms that had been spiked with plutonium-239 nitrate. Three microcosms were spiked twice, at the start and on day 69, with 0.01, 0.10, and 1.0 $\mu\text{Ci/cm}^3$, respectively. Patterns of Pu concentration with time were similar at all concentrations following both additions. The distribution coefficient (ratio of Pu in sediments to that in water) at equilibrium was 7.88, 1.22, and 1.78×10^4 in the microcosms that received spikes of 0.01, 0.1 and 1.0 $\mu\text{Ci/cm}^3$, respectively. Eleven *Chironomus* larvae were collected on day 73, and trophic transfer factors (TTF) for Pu to *Chironomus* from labeled sediments were calculated. The TTF values were similar to those for other invertebrates and fish taken from a more complex system contaminated with ^{237}Pu . The actual amount of Pu absorbed from food by *Chironomus* apparently was 0.1%-1% of that in the diet. The results indicate that organic sediments are not a sink for Pu but rather a pool with a significant, though small, available fraction for transfer to the next trophic level. (11 refs)

See also:

- *(Rev.): 78-5434, 78-5444, 78-5445, 78-5446, 78-5447, 78-5451.
- *(Chem.): 78-5477, 78-5513, 78-5521, 78-5633, 78-5651, 78-5661, 78-5665, 78-5671, 78-5678, 78-5685.
- *(Viral): 78-5769, 78-5810, 78-5853.
- *(Immun.): 78-5858, 78-5860.
- *(Path.): 78-5872, 78-5873, 78-5897, 78-5898.
- *(Epid.-Biom.): 78-5922, 78-5926, 78-5959, 78-5970, 78-5972, 78-5973, 78-5975, 78-5976.

VIRAL CARCINOGENESIS

- 78-5736 Identification of an Avian Oncovirus Polypeptide in Uninfected Chick Cells.** (Eng) Eisenman, R. (Fred Hutchinson Cancer Res. Center, 1124 Columbia St., Seattle, WA, 98104); Shaikh, R.; Mason, W. S. *Cell* 14(1): 89-104; 1978.

Studies of the nature of the *gag* gene product synthesized in uninfected chicken embryo fibroblasts (CEF) positive for the expression of avian oncovirus group-specific antigen (gs+) are reported. This product was shown to be a polypeptide with a mol wt of 120,000 daltons (P120). Its presence could not be correlated with the phenotypic expression of endogenous oncovirus envelope glycoprotein. Immunoprecipitation experiments indicated that it contains antigenic determinants of the virion core proteins (products of the *gag* gene) and reverse transcriptase, but not of the envelope glycoprotein gp85. Chromatography of the tryptic digests indicated that P120 contains amino acid sequences related to the virion core proteins p27, p19, and p12 of Rous-associated virus type 0 (RAV-0), but that it lacks the tryptic peptides of p15. Further studies were consistent with the idea that P120 is equivalent to a *gag-pol* polypeptide that is missing both p15 and significant portions of reverse transcriptase. In pulse-chase experiments, P120 did not appear to be specifically cleaved to generate virion structural proteins. Furthermore, P120 was not detected in RAV-0-producing line 7 and line 15 CEF. Instead, only Pr76-*gag* (the direct precursor to virion core proteins in productively infected cells) was observed. Thus, P120 is unlikely to act as a functional intermediate in virus maturation. Infection of gs+ CEF with exogenous avian sarcoma and leukemia viruses resulted in a significant but variable decrease in the level of P120. (71 refs)

- 78-5737 Protein Kinase Activity Associated with the Avian Sarcoma Virus *src* Gene Product.** (Eng) Collett, M. S. (Dept. Pathology, Univ. Colorado Medical Center, Denver, CO, 80262); Erikson, R. L. *Proc Natl Acad Sci USA* 75(4): 2021-2024; 1978.

Evidence is presented for the association of protein kinase activity with the avian sarcoma virus (ASV) *src* gene product (p60-*src*). Incorporation of phosphorus from (γ - ^{32} P)ATP into protein was catalyzed by specific immunoprecipitates from ASV-transformed avian and mammalian cells. This incorporation was observed only when antiserum from tumor-bearing rabbits able to precipitate p60-*src* specifically was used to immunoprecipitate antigens from transformed cell lysates. Immunoprecipitates of extracts from normal cells or cells infected with a transformation-defective ASV mutant showed no activity in this assay, nor did any immune complexes formed with normal rabbit serum or any of the cell extracts tested. The expression of protein kinase activity (ATP:protein

phosphotransferase) was growth temperature-dependent in cells infected with an ASV mutant temperature-sensitive for transformation. It is speculated that protein phosphorylation may be involved in viral transformation by impeding normal assembly of cellular components. (21 refs)

- 78-5738 Purification of Virus-specific RNA from Chicken Cells Infected with Avian Sarcoma Virus. Identification of Genome-Length and Subgenome-Length Viral RNAs.** (Eng) Parsons, J. T. (Dept. Microbiology, Univ. Virginia Sch. Medicine, Charlottesville, VA, 22901); LePesch, P.; Dierks, P. *J Virol* 27(1): 227-238; 1978.

Avian sarcoma virus (ASV)-specific RNA was purified from ASV-infected chick embryo (CE) cells by hybridizing polydeoxycytidylic acid-elongated complementary DNA to ASV RNA and then chromatographing the annealing mixture on polyinosinic acid-Sephadex columns. The purity and nucleotide sequence of the purified virus-specific RNA were established by rehybridization experiments and analysis of labile RNase T₁-resistant oligonucleotides by two-dimensional polyacrylamide gel electrophoresis. Cultured CE cells were infected with Prague ASV, subgroup B or C (PrC); with Bratislava ASV, subgroup C; or with transformation-defective (td) PrC (a virus with a deletion of the sarcoma gene). Polyadenylic acid-containing RNA purified from ASV-infected cells contained approx 1%-4% virus-specific RNA, compared with 0.06%-0.15% in uninfected cells. Sucrose gradient analysis of the virus-specific RNA isolated from infected cells revealed two major classes of polyadenylated viral RNA with sedimentation values of 36S and 26S-28S. Cells infected with td PrC contained 34S and 20S-22S viral RNA species. Double-label experiments employing infected cells labeled initially for 48 hr with ^{3} H-uridine and then for 30, 60, or 120 min with ^{32} P-phosphate showed that the intracellular accumulation of genome-length RNA (36S) was significantly faster than that of the subgenomic 26S-28S viral RNA species. The possible functions of the different species are discussed. (33 refs)

- 78-5739 Peptide Analysis of the Transformation-specific Antigen from Avian Sarcoma Virus-transformed Cells.** (Eng) Brugge, J. (Dept. Pathology, Univ. Colorado Medical Center, Denver, CO, 80262); Erikson, R. L.; Collett, M. S.; Erikson, R. L. *J Virol* 26(3): 773-782; 1978.

Virus-specific proteins from chicken, hamster (2 lines), and field vole cells transformed by the Schmidt-Ruppin strain of avian sarcoma virus (SR-ASV) were isolated by immunoprecipitation and subjected to peptide analysis. Sera from

r-bearing rabbits (TBR) precipitated two proteins with
wts of 76,000 daltons (76K) and 60K from all
V-transformed cell lines. One-dimensional peptide anal-
of the 60K proteins indicated that the proteins from all
r transformed cell lines were identical. Two-dimensional
peptide analysis of the Pr76 and 60K proteins indicated that
60K protein was not related to Pr76 and that the observed
migration of the four tryptic peptides of Pr76 and 60K was
tuitous. When cell extracts from ASV-transformed chick
is radiolabeled with ^{32}P were immunoprecipitated with
R serum, a phosphorylated protein believed to be identical
60K was immunoprecipitated. Precipitation of this protein
s not blocked when TBR serum was first adsorbed with
rupted virions, although precipitation of the ^{32}P -labeled
precursor to group-specific antigens, Pr76, was blocked un-
these conditions. The tryptic phosphopeptides of 60K
protein from transformed chicken and field vole cell lines
grated identically during two-dimensional peptide anal-
is, and their migration differed from that of the principal
osphopeptide of Pr76. The acid hydrolysates of Pr76 and
K contained phosphoserine, and 60K contained a small
mount of phosphothreonine. The data show that the 60K
protein is transformation-specific, virus-coded, and unrelated
the structural proteins of ASV. (25 refs)

8-5740 **Identification of Tumor-Specific Cell Surface
Antigens from Quail Fibroblasts Transformed
by Avian Retro-Viruses and Chemical Carcinogens.** (Eng)
ubsamen, H. (Institut für Virologie der Justus-Liebig-Uni-
versität Giessen, Bereich Humanmedizin Frankfurter Strasse
71, 63 Giessen, W. Germany); Ignjatovic, J.; Hayami, M.;
auer, H. In: *Advances in Comparative Leukemia Research*
1977. *Proceedings of the VIIIth International Symposium on*
Comparative Research on Leukemia and Related Diseases,
held in Amsterdam, The Netherlands on 22-26 August, 1977.
entvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York:
Elsevier/North Holland Biomedical Press): 503 pp.; 351-352;
1977.

four tumor-specific antigens were detected on the surface of
quail fibroblasts transformed by avian sarcoma virus or by
methylcholanthrene (MC) in an in vitro cellular mi-
crocytotoxicity (CMC) assay: s-gp85, the viral envelope
glycoprotein; g-gp85, an additional antigenic site of gp85 ex-
posed only on tumor cells; a tumor-specific antigen of
virus-transformed cells; and an embryonic antigen expressed
in both MC- and virus-transformed cells. Soluble extracts of
these cells blocked the CMC reaction. (3 refs)

8-5741 **Biochemical Properties of p15-associated Pro-
tease in an Avian RNA Tumor Virus.** (Eng)
Dittmar, K. J. (Max-Planck-Institut für Molekulare Genetik,
1000 Berlin 33, W. Germany); Moelling, K. *J Virol* 28(1):
106-118; 1978.

the biochemical properties of a proteolytic activity associat-

ed with avian myeloblastosis virus structural protein p15
were studied. Ion-exchange column chromatography and size
fractionation analysis confirmed that all protease activity re-
sided in the p15 fragment. The purity of p15 was assured by
an isoelectric focusing procedure in polyacrylamide gels.
Upon serological analysis with monospecific antibodies, p15
reacted with anti-p15 serum only and not with anti-p12 or
anti-p27 sera. The proteolytic activity of p15 was demonst-
rated with bovine serum albumin (BSA), concanavalin A, ca-
sein, and ovalbumin. These substrates were denatured before
p15 treatment by heat and sodium dodecyl sulfate. The op-
timum pH for this activity was 5.7. The effects of p15 were
similar to those of papain, but papain had a much stronger
proteolytic activity and it could not be applied over the same
concentration range (0.1-2 μg) as p15. The parallel behavior
of p15 papain was further demonstrated by their responses
to proteolytic inhibitors. When p15 was incubated with poly-
protein precursors precipitated from lysed Rous-associat-
ed-virus (RAV-6)-infected chicken cells, precursor Pr76 was
cleaved. If anti-p15 serum was added, cleavage of Pr76 was
inhibited. This indicates that p15 has a direct action on Pr76.
(45 refs)

78-5742 **Virus-coded Origin of a 32,000-Dalton Protein
from Avian Retrovirus Cores: Structural Relat-
edness of p32 and the β Polypeptide of the Avian Retrovirus
DNA Polymerase.** (Eng) Schiff, R. D. (St. Louis Univ. Medi-
cal Center, Inst. Molecular Virology, St. Louis, MO, 63110);
Grandgenett, D. P. *J Virol* 28(1): 279-291; 1978.

A 32,000-dalton polypeptide (p32) located in avian retrovirus
cores was characterized by radioimmunoprecipitation (RIP)
and biochemical techniques and was hypothesized to be relat-
ed to the β polypeptide of the $\alpha\beta$ form of avian myeloblasto-
sis virus (AMV) DNA polymerase. ^{35}S -methionine-labeled
AMV virions were radioimmunoprecipitated (RIPD) with
sarc III antiserum derived from Syrian hamsters bearing avi-
an sarcoma virus-induced tumors. Precipitation of the DNA
polymerase and p32 indicated that they were encoded by the
viral genome. ^{125}I -labeled AMV p32 purified directly from
plasma AMV was also RIPD by sarc III antiserum. AMV
p32 was RIPD by goat and rabbit anti-AMV $\alpha\beta$ DNA
polymerase; polypeptides α and β were RIPD by rabbit an-
ti-AMV p32 protein, but polypeptide β was precipitated to
a much greater extent. Therefore, AMV p32 and the β poly-
peptide of AMV DNA polymerase shared common antigenic
determinants. Peptide mapping with sodium dodecyl sul-
fate-polyacrylamide gel electrophoresis after partial chymo-
tryptic or *Staphylococcus aureus* V-8 protease digestion
showed that all peptides demonstrable from AMV p32 were
also present in the digests of the β polypeptide. Conversely,
no common sequences were found between p32 and the α
polypeptide. These results indicate that p32 is derived from
the β polypeptide of AMV $\alpha\beta$ DNA polymerase by proteo-
lytic cleavage, but that it is structurally unrelated to the α
polypeptide. (46 refs)

- 78-5743 Discrimination of DNA Polymerase and RNase H Activities in Reverse Transcriptase of Avian Myeloblastosis Virus.** (Eng) Gorecki, M. (Dept. Organic Chemistry, Weizmann Inst. Science, Rehovot, Israel); Panet, A. *Biochemistry* 17(12): 2438-2442; 1978.

The functional amino acid residues in the active sites of avian myeloblastosis virus (AMV) reverse transcriptase were examined. Viral DNA polymerase activity was 3.6-fold more sensitive to the sulfhydryl- and amino-group-specific reagent N-ethylmaleimide (NEM) than was RNase H activity. Incubation of the enzyme in the presence of polynucleotides resulted in partial protection of both activities against NEM, whereas nucleotide triphosphate did not affect the inactivation rate. The interaction of NEM with DNA polymerase was temperature-dependent. NEM inactivated DNA polymerase with pK_a 's of 5.5 and 6.5 and RNase H with a pK_a of 7.5. The hydrophobic agents 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), p-hydroxymercuribenzoic acid, and 2-nitro-5-thiocyanobenzoate (NTCB) were potent inhibitors of DNA polymerase activity, but they did not affect RNase H. Hydrophilic agents such as iodoacetic acid, iodoacetamide, and ethylenimine had no effect on AMV reverse transcriptase. Since pretreatment of DNA polymerase with DTNB did not prevent NEM from reacting with a residue necessary for the activity of this enzyme, two different reactive groups are probably involved with this activity. RNase H was completely inactivated by short preincubation at pH < 4.5, but it was only 50% inactivated at pH 5.0. DNA polymerase was unaffected at pH 4.3-8.3, and sharp inactivation was observed only below pH 4.3. The inactivation caused by preincubation at low pH was irreversible and was not time-dependent. The data indicate that the DNA polymerase and RNase H activities reside on independent active sites that may partially overlap but can be distinguished by suitable chemical modifications. (22 refs)

- 78-5744 A Virus-induced Non-virion Antigen Specific for Transformation at the Surface of RSV-transformed Fibroblasts.** (Eng) Comoglio, P. M. (Dept. Human Anatomy, Univ. Torino Medical Sch., 10126 Torino, Italy); Prat, M.; Bertini, M. *Nature* 273(5661): 381-383; 1978.

Studies were made of the antigenic composition of BHK-C13/21 hamster fibroblasts transformed in vitro by Rous sarcoma virus (RSV) mutants either defective for the synthesis of envelope glycoproteins or temperature-sensitive for the ability to transform the host cell. These cells were found to express a tumor-specific antigen with the following properties: it is expressed at the surface of hamster cells transformed by RSV but not by unrelated RNA or DNA oncogenic viruses, it is expressed by RSV-transformed cells of different animal species, and it is not one of the known structural virion proteins. It was named the virus cell-surface antigen (VCSA). Experiments with the temperature-sensitive mutant indicated a strict correlation between the appearance of VCSA at the cell surface and expression of the transforming

sarc gene. VCSA may be the product of the *sarc* gene or may be coded by a cellular gene under control of this gene. (26 refs)

- 78-5745 *Ts pol* Mutants of Avian Sarcoma Viruses: Mapping and Demonstration of Single Cycle Recombination.** (Eng) Alevy, M. C. (Dept. Microbiology, U. Southern California Sch. Medicine, 2025 Zonal Ave., Los Angeles, CA, 90033); Vogt, P. K. *Virology* 87(1): 21-33; 1978.

Twenty-seven early coordinate temperature-sensitive (*ts*) mutants of the Prague strain of Rous sarcoma virus were isolated, and the position of the mutants in the *pol* gene was determined by three-factor crosses using the known polymerase mutant *LA337* and the *env* marker. The effect of temperature shifts on the transforming abilities of the mutants indicated that each mutant contains an early coordinate lesion which could interfere with the initiation but not maintenance of transformation. All mutants behaved in a qualitatively similar manner except *LA354*, which appeared to be a multiple mutant containing both late transformation and early coordinate lesions. Temperature shifts had similar effects on the replicative abilities of all mutants. The data indicated that the early coordinate lesion acted at 42°C to abort the infection cycle before a stable association between virus and host cell could be formed. The polymerase activity of each mutant was more thermolabile than that of wild-type virus in vitro. Three-factor crosses between *LA337* B and the new mutants *LA337* to *LA377* suggested that wild-type progeny clones obtained from the crosses were true recombinants and that recombination between the *ts* markers occurred at low frequency. The relative proportions of each class of progeny remained fairly constant between 48 and 96 hr after the infected cells were mixed, the wild-type recombinants having been formed by 48 hr after mixing. All of the new mutants mapped between *LA337* and the *env* gene. The data suggest that the *ts* isolates were all polymerase mutants and that heterozygotes are not obligatory precursors of recombinants. (39 refs)

- 78-5746 Disk Gel Electrophoresis and Infrared Spectroscopy of the Proteins in Rous Sarcoma Tissue.** (Rus) Klenina, N. V. (Ukrainian Res. Inst. Experimental Veterinary Medicine, Kharkov, USSR); Antonov, V. I.; Chorna, L. D. *Ukr Biokhim Zh* 50(4): 520-525; 1978.

Disk gel electrophoresis and infrared spectroscopy were used to analyze the proteins in Rous sarcoma tissue. Thirty-day-old Leghorn chickens were inoculated intramuscularly with Rous sarcoma virus homogenate (0.2 ml); tumor specimens were analyzed on days 3, 6, 10, and 15 postinoculation. The electrophoretogram of normal thoracic muscle consisted of 18 protein fractions. Tumor development resulted in major changes in all these fractions, especially in fast-protein fractions. The albumin level increased from 1.32% to 15% on day 15 and the prealbumin level increased from 1.68% to 3.5%.

compared with control thoracic muscle specimens, Rous sarcoma tissue lacked two absorption bands at 1,100 and 800 cm^{-1} . (12 refs)

78-5747 Characterization of the *env* Gene in Avian Oncoviruses by Heteroduplex Mapping. (Eng) Hu, S. (Dept. Microbiology, Univ. Southern California Sch. of Medicine, Los Angeles, CA, 90033); Lai, M. M.; Vogt, P. K. *J. Virol* 27(3): 667-676; 1978.

The *env*-specific nonhomology between ring-necked pheasant virus (RPV) and chicken leukosis and sarcoma viruses was used as the basis for a study of the *env* gene by heteroduplex mapping using electron microscopy. The *env*-specific region of nonhomology was 1.5-1.7 kilobases (kb) long. Its 3' boundary was at 0.6-0.7 kb from the 3' end of the genome transformation-defective (*td*) viruses and at about 2.5 kb from nondefective avian sarcoma viruses. This estimate is similar to that obtained by oligonucleotide fingerprinting. Comparison of several strains of avian oncoviruses showed that the 3' half of this *env* region is conserved and the 5' half is more divergent. A small area at the very 3' end of the *env* region showed divergence between the different avian oncovirus strains tested. The data argue against the possibility of an unrecognized gene between *env* and *src*. In heteroduplex molecules between *td* and nondefective sarcoma virus genomes, the *src* deletions and the *env* nonhomology region are contiguous, and their size is equal to the sum of the *src* and *env* genes. However, the possible presence of a very short region of spacer sequences between *env* and *src* could not be ruled out. Comparison of the glycoproteins from various avian oncoviruses by gel electrophoresis showed that those of RPV (gp60, gp33) and Chinese quail virus (gp55, gp29) differed in mol wt from the glycoproteins of the chicken oncoviruses (gp85, gp37). (32 refs)

78-5748 Lack of Serologic Identity of Marek's Disease Associated Tumor Surface Antigen (MATSA) in Various Marek's Disease Cell Lines. (Eng) Sharma, J. M. (Regional Poultry Res. Lab., ARS, U.S.D.A., 3606 E. Mt. Hope Road, East Lansing, MI, 48823); Nazerian, K.; Stephens, E. A.; Witter, R. L. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 191-192; 1978.

Serologic relationships between Marek's disease (MD) tumor-associated surface antigen (MATSA) on MD cell lines (MSB-1, RPL-1, and HPRS-1) and an MD-derived transplant (MDT) were studied by complement-independent antibody cytotoxicity assay. There was a one-way reaction between MSB-1 and RPL-1 cells; RPL-1 antiserum reacted

with MSB-1 cells, but MSB-1 antiserum did not cross-react with RPL-1 cells. Because of possible contamination of MSB-1 and RPL-1 sera with antibody to histocompatibility antigens, the reactivity of these sera could not be related directly to MATSA. However, the results with MDT antiserum, which lacked histocompatibility antibody, clearly indicated that MDT shared a common surface antigen with MSB-1 and HPRS-1 but not with RPL-1. The RPL-1 antiserum cross-reacted with MDT and HPRS-1 as well as with MSB-1. (5 refs)

78-5749 Demonstration of Cells with Marek's Disease Tumor-associated Surface Antigen (MATSA) in Peripheral Blood of Chickens with Marek's Disease Lymphoma. (Eng) Sato, T. (Hokuwa Livestock Hygiene Service Center, Nara, Japan); Wataki, K.; Matsuda, H.; Kato, S. *Biken J* 21(1): 33-35; 1978.

The existence of Marek's disease (MD) tumor-associated surface antigen (MATSA)-positive cells in the peripheral blood of chickens with MD lymphoma is reported. Eight of 10 chickens given 10⁷ MD tumor cells (line MSB-1) developed typical MD lymphoma between 46 and 140 days of age. MATSA-positive cells were found in the peripheral blood of four of these eight chickens and in one of the two chickens without lymphoma, but not in the peripheral blood of controls. In the chickens with lymphomas, the percentages of MATSA-positive cells among peripheral WBC ranged from 0.2% to 1.5%. Autopsy studies of commercial chickens suspected of having MD or lymphoid leukosis (LL) revealed 26 cases of MD and 5 of LL. MATSA-positive cells were found in the peripheral blood of 22/26 MD chickens but in none of the LL chickens. The percentage of positive cells in the peripheral blood of the 22 chickens ranged from 0.3% to 5%. No MATSA-positive cells were found in the peripheral blood of healthy chickens. The correlation between MD lymphoma incidence and MATSA-positive cells in the peripheral blood suggests that the latter may be useful in diagnosing MD. (9 refs)

78-5750 Interaction of Marek's Disease and Rous Sarcoma Viruses During Mixed Infection in Chickens. (Rus) Sedunov, Z. A. (All-Union Res. Veterinary Inst. Poultry Farming, Moscow, USSR); Korovin, R. N.; Neidbailik, I. E. *Veterinariia* (7): 43-46; 1978.

Characteristic features of the interaction of Marek's disease virus (MDV) and Rous sarcoma viruses (RSV) during mixed infection were analyzed by inoculating white Leghorn chickens with RSV (0.1 ml, im) and/or MDV (0.2 ml, ip). All chickens inoculated with RSV alone died within 19 days; 36.6% of the chickens inoculated with MDV died within 60 days (necropsy showed lymphoid infiltrates in 46.6% of these birds). Simultaneous injection of RSV + MDV induced both tumors (60%) and lymphomas (43.3%). The duration of sur-

vival of chickens inoculated with RSV + MDV was 11-26 days longer than that of chickens inoculated with RSV alone. (no refs)

- 78-5751 Identification of the *Mtv-2* Gene Responsible for the Early Appearance of Mammary Tumors in the GR Mouse by Nucleic Acid Hybridization.** (Eng) Michalides, R. (Div. Genetics, Antoni van Leeuwenhoekhuis, Netherlands Cancer Inst., Amsterdam, Netherlands); van Deemter, L.; Nusse, R.; van Nie, R. *Proc Natl Acad Sci USA* 75(5): 2368-2372; 1978.

The number of mammary tumor virus (MTV) proviral sequences associated with the *Mtv-2* gene was investigated by nucleic acid hybridization between MTV ³H-complementary DNA (cDNA) and DNA from GR, C57BL/10JA (B10), and GR-*Mtv-2* mice. Both B10 and GR-*Mtv-2* mice lack *Mtv-2* gene expression. The DNA of GR mice contained 12 copies of MTV proviral sequences, whereas only four copies were present in the DNA of B10 and GR-*Mtv-2* mice, leading to the conclusion that the *Mtv-2* gene in the GR mouse strain is associated with eight additional MTV proviral sequences. These results indicate that the GR genome contains two sets of MTV sequences. One is associated with the *Mtv-2* gene and controls the early appearance of mammary tumors in the GR mouse, and the other is responsible for MTV RNA expression in GR-*Mtv-2* mice. Different amounts of MTV RNA were detected at first lactation in the mammary gland of B10 and GR-*Mtv-2* mice, even though both contain four copies of MTV proviral sequences. This indicates a difference between the two strains either in the regulation of expression of these MTV proviral sequences or in their location in the murine genome. (19 refs)

- 78-5752 Cultivation of Mouse Mammary Tumor Cells Derived from DD/Tr.** (Eng) Iwai, Y. (No affiliation given); Iwai, M.; Takamori, Y.; Okumoto, M. *Annual Report Radiation Center Osaka Prefecture* 18: 113-117; 1978.

The relationship between changes in cell-surface structure and the production of murine mammary tumor virus (MuMTV) by a cell line derived from a spontaneous mammary tumor in a female DD/Tbr mouse was studied in the presence and absence of insulin (INS) or dexamethasone (DXM). Most of the cells cultured with INS or DXM were epithelioid, but fibroblastic cells always appeared in the domes. RNA-dependent DNA polymerase (RDDP) activity was detected in the culture fluid of cells grown in the presence of the hormones, but not in the culture fluid of cells grown in their absence. This activity began to increase about 11 days after seeding; it reached a max level after 14-17 days and then declined. Murine leukemia virus was not detected in the cells by XC-plaque tests or by a focus-forming assay using mink S + L- cells. Scanning electron microscopy of the mammary tumor cell surfaces in the early cultures revealed many short

microvilli and viruslike particles that increased with the increase in RDDP activity. Thus, there is a close relation between the appearance of short microvilli and viruslike particles on the cell surface and the release of virus into culture medium. (5 refs)

- 78-5753 Changes in MuMTV DNA and RNA Levels in Balb/c Mammary Epithelial Cells During Malignant Transformation by Exogenous MuMTV and Hormones.** (Eng) McGrath, C. M. (Dept. Biology, Michigan Cancer Foundation, Detroit, MI, 48201); Marineau, E.; Voyles, B. A. *Virology* 87(2): 339-353; 1978.

Concentrations of murine mammary tumor virus (MuMTV) sequences in the DNA and RNA of normal BALB/c mammary epithelial cells were compared with concentrations of virus- and hormone-induced premalignant and malignant derivatives to determine if changes in steady-state MuMTV RNA levels occur during malignant transformation. Normal mammary cells contained multiple (5 or 6) copies of MuMTV sequences in DNA. Malignant mammary cells, transformed after MuMTV infection, contained two to three times the number of MuMTV sequence copies. These extra MuMTV sequences were not detected in the DNA of normal mammary epithelial cells of MuMTV-infected females or MuMTV-induced premalignant cells. Neither malignant nor premalignant mammary cells transformed after prolonged exposure to mammotropic hormones contained in the additional MuMTV sequences. MuMTV sequences detected in malignant BALB/c mammary epithelial cells, regardless of transforming agent, were indistinguishable by DNA-complementary DNA (DNA-cDNA) hybrid thermal denaturation from the MuMTV sequences in the DNA of normal cells. Mammary adenocarcinomas induced by MuMTV or mammotropic hormones contained 30- to 50-fold higher concentrations of MuMTV RNA than normal mammary cells of young animals. Similarly high concentrations were detected in premalignant cells. It is concluded that changes in steady-state MuMTV RNA concentrations take place during premalignant and malignant stages of transformation. (refs)

- 78-5754 Mammary Tumor Virus Oncogenesis and Immunogenicity in Three Sublines of C3H Mouse.** (Eng) Vaage, J. (Dept. Cancer Therapy Development, Pondville Hosp., Walpole, MA, 02081); Medina, J. *Cancer Res* 38(8): 2443-2447; 1978.

Mammary tumorigenesis and mammary tumor transplantation immunogenicity were compared in three C3H mouse sublines: standard mammary tumor virus (MTV-S)-infected C3H/He mice; MTV-S-infected C3H/Ki mice; and MTV-S-free C3H/He mice. The age at appearance of the first tumor, the growth rate of the tumors in their first transplant generation, and the immunogenicity of each tumor in s

c female recipients were determined for the first tumor appear in each of 25 breeding females from each of the sublines. The early appearance of the tumors was related to the presence of the MTV-S. The genetically dissimilar lines C3H/He and C3H/Ki, both infected with MTV-S, developed mammary tumors at an average age of 10 mo, 11 mo in MTV-S-free C3H/He mice. The tumor characteristics immunogenicity and growth stimulation were related to genetic factors. The genetically similar sublines C3H/He C3Hf/He developed similar proportions of immunogenic growth stimulating mammary tumors; the genetically divergent C3H/Ki subline developed tumors that were immunogenic and that tended to be strongly growth stimulating. (20 refs)

78-5755 Observations on the Question of Horizontal Transmission of Mouse Mammary Tumor (Eng) Moore, D. H. (Dept. Microbiology and Immunology, Hahnemann Medical Coll. and Hosp., Philadelphia, PA, 19102); Holben, J. A. *Cancer Res* 38(8): 5-2457; 1978.

Antigen and tumor incidences in weanling BALB/c and C57BL mice living for 5 wk in cages with murine mammary virus (MuMTV)-infected females were compared with control BALB/c and C57BL mice living in the same laboratory. All mice were bred continuously, and third-lactation mice were tested for MuMTV antigen by the Ouchterlony immunodiffusion test. Although the antigen incidence was significantly greater, mammary tumor incidence in the females was not significantly different from that in controls. Mice inoculated with phosphate-buffered salt solution (0.02 M phosphate, pH 7.4; 0.15 M NaCl; and 0.1% bovine serum albumin) and sham-inoculated mice also had an elevated antigen incidence. Repeat tests of milks at the fourth or fifth lactations indicated that > 50% of those positive at the first became negative at later lactations. Factors that may have stimulated MuMTV antigen secretion are discussed. (20 refs)

78-5756 In Vivo Activities of B-tropic Isolates from Mice of Strain C57BL/Ka. (Eng) Decleve, A. (Cancer Biology Res. Lab., Dept. Radiology, Stanford Univ. Medical Center, Stanford, CA, 94305); Lieberman, M.; Travis, M.; Kaplan, H. S. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 1978.

The detection of virus-specific cytoplasmic antigens by indirect immunofluorescence was used to follow virus replication in the lymphoid and hematopoietic tissues of C57BL/Ka

mice inoculated with three B-tropic viruses isolated from that strain: radiation leukemia virus (RadLV), BL/Ka(6), and BL/Ka(B). Virus doses were 6×10^4 particles/inoculum in newborns and 5×10^5 particles/inoculum in animals aged 1 wk or older. The BL/Ka(B) and BL/Ka(6) isolates were essentially nonthymotropic after iv or ip inoculation. After direct intrathymic inoculation, BL/Ka(6) was able to infect and transform the thymic lymphocytes, but BL/Ka(B) lacked this ability. However, BL/Ka(B) could productively infect hematopoietic or lymphoid tissues when injected in neonates, which suggests that nonleukemogenic viruses may contribute to leukemogenesis in C57BL/Ka mice. The thymotropic and leukemogenic activity of BL/Ka(6) was 10^5 times lower than that of RadLV. (7 refs)

78-5757 Presence of Oncornavirus-like Particles in the P388 Murine Leukemic Cell Line. (Eng) Schmidt, J. M. (Dept. Botany and Microbiology, Arizona State Univ., Tempe, AZ, 85281); Pettit, G. R. *Experientia* 34(5): 659-660; 1978.

P388 murine lymphoblastoid cells were examined for the presence of virus like particles. Almost every sectioned cell observed had C-type oncornavirus like particles budding at the plasma membrane or, occasionally, into an intracytoplasmic vesicle. Intracytoplasmic A-type and immature B-type particles were also observed. The discovery of oncornavirus like particles in the P388 cell line increases the utility of this neoplastic system for detecting potential antineoplastic agents. (3 refs)

78-5758 Endogenous RNA Tumor Viruses Are Activated During Chemical Induction of Murine Plasmacytomas. (Eng) Armstrong, M. Y. (Dept. Epidemiology and Public Health, Yale Univ. Sch. Medicine, 333 Cedar St., New Haven, CT, 06510); Ebenstein, P.; Konigsberg, W. H.; Richards, F. F. *Proc Natl Acad Sci USA* 75(9): 4549-4552; 1978.

A study was made of the endogenous expression of murine leukemia virus (MuLV) in BALB/cAn female mice inoculated with the potent plasmacytoma-inducing oil pristane (2,6,10,14-tetramethylpentadecane: 0.5 ml ip) at age 6, 14, and 23 wk. Spleen, mesenteric lymph node, thoracic lymph node, and peritoneal exudate cells were prepared from treated and control mice over a 10-mo period, and the cell suspensions were assayed for infectious MuLV's. Tumors appeared in the treated mice after age 32 wk. From 28 to 50 wk, the percentage of ecotropic virus-positive mice and the number of virus-positive cell samples rose sharply in the treated mice (24/36 mice, 54/105 samples) compared with controls (4/22 mice, 11/58 samples). The percentage of xenotropic virus-positive samples was also consistently higher in treated mice, and virus was isolated from these mice throughout the observation period. Xenotropic MuLV could not be detected

in any control samples obtained at 28, 32, and 39 wk. Both ecotropic and xenotropic endogenous MuLV were expressed in those sites where foci of transformed plasma cells subsequently appear; ie, peritoneal exudate cells and thoracic lymph node cells. The significance of the preferential expression of ecotropic and xenotropic MuLV in target cell populations following administration of a carcinogen is discussed in terms of the possible formation of an oncogenic variant virus. (37 refs)

- 78-5759 Serological Characterization of B-tropic Viruses of C57BL Mice. Possible Origin by Recombination of Endogenous N-tropic and Xenotropic Viruses.** (Eng) Benade, L. E. (Cancer Biology Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501); Ihle, J. N.; Declève, A. *Proc Natl Acad Sci USA* 75(9): 4553-4557; 1978.

The serological characteristics of two independent sets of N- and B-tropic viruses isolated from C57BL/6 mice and C57BL/Ka mice were examined. Radioimmune competition assays for various virion components revealed that the N-tropic and B-tropic viruses were readily distinguishable by assays for the viral proteins p15 and p12, whereas these viruses were serologically identical by competition assays for proteins gp71 and p30. In addition, the N-tropic viruses had p12's serologically identical to that of AKR murine leukemia virus (MuLV), but the B-tropic viruses had p12's serologically identical to that of B-tropic radiation leukemia virus VL-3. The C57BL/6 N-tropic virus had a p15 identical to AKR MuLV p15, whereas the C57BL/Ka N- and B-tropic viruses and the C57BL/6 B-tropic virus had a serologically distinct p15. The results suggest that the endogenous N-tropic virus of C57BL mice undergoes recombination with the endogenous xenotropic virus and that this mechanism is involved in the generation of B-tropic viruses in these mice. (30 refs)

- 78-5760 Abrogation of Fv-1b Restriction with Murine Leukemia Viruses Inactivated by Heat or by Gamma Irradiation.** (Eng) Bassin, R. H. (Lab. Tumor Virus Genetics, NCI, Bethesda, MD, 20014); Duran-Troise, G.; Gerwin, B. I.; Rein, A. *J Virol* 26(2): 306-315; 1978.

The mechanism by which Fv-1b restriction in BALB/3T3 cells is abrogated following infection with N-tropic murine leukemia virus was investigated by comparing the inactivation rates for viral infectivity and for the ability of the same virus to abrogate Fv-1 restriction. Inactivation of the abrogating ability of N-tropic murine leukemia virus following graduated doses of gamma radiation ($0-2.5 \times 10^6$ rads) proceeded at half the rate of that for viral infectivity. This indicated that viral RNA must function in abrogating Fv-1b restriction but that only a portion of the viral genome is required. The inactivation kinetics of N-tropic murine leukemia virus were determined following incubation of virus at 43 C. Abrogation of Fv-1b restriction by N-tropic murine leukemia virus was found to be about six times as stable under these conditions as was viral infectivity. Virion-associated reverse transcriptase activity was inactivated at the same rate as was viral infectivity, indicating that this enzyme may not be needed during abrogation. When virus was heated to 43 C, abrogation was transient, requiring 6-9 hr after virus infection to become maximally effective and then beginning to disappear after approx 18 hr. These findings confirm the idea that abrogation of Fv-1 restriction can be separated experimentally from viral replication. (33 refs)

78-5761 **Differences in the Properties of Cytoplasmic Particles and in the Expression of Virus-specific Information in the Thymus of Normal and Lymphomatous AKR Mice.** (Eng) Gurgo, C. (Inst. Molecular Virology, Louis Univ. Medical Center, St. Louis, MO, 63110); Bondurant, M. C.; Bridges, S. H. *Cancer Res* 38(8): 2473-2479; 1978.

The properties of viruslike cytoplasmic particles in thymic tissue from normal and lymphomatous AKR mice and expression of leukemia-related sequences in these tissues were investigated. The retrovirus nature of the particles was indicated by the presence of RNA-dependent DNA polymerase (reverse-transcriptase) and by the chromatographic properties of the purified enzyme. Equilibrium centrifugation studies revealed a difference in the buoyant densities of the particles extracted from the two types of tissue. A density of 1.18 g/ml was estimated for particles of lymphomatous thymic tissue and for the Gross passage A virus, a leukemogenic AKR murine leukemia virus MuLV, compared with 1.14 g/ml for thymic particles from normal (1- and 2-month-old) mice. Molecular hybridization studies aimed at detecting differences in the expression of viral-specific information in RNA from the normal and lymphomatous thymic tissues were carried out with a Moloney MuLV complementation DNA (cDNA) probe. From 40% to 50% of the probe hybridized with the RNA from normal thymic tissue with an equivalent value of the product of the time (sec) of hybridization and the RNA concentration (moles of nucleotides/liter) at which 50% hybridization was reached being indicative of approx 100 genome equivalents of endogenous viral RNA per cell. About 80% of the probe hybridized with the RNA from lymphomatous tissue. Viral-related RNA in the lymphomatous tissue consisted of two abundance classes: one class (homologous with 50% of the probe) was hybridized at a rate suggestive of about 2,400 viral genome equivalents/cell. Thus, in the lymphomatous thymus, virus-specific RNA expression was not only increased quantitatively (relative to normal thymus) it was characterized by the expression of qualitatively 'new' sequences. The melting temperature of hybrids of MuLV cDNA with RNA from normal and lymphomatous tissue was 83 C. A melting temperature of 91 C was observed for hybrids composed of the probe and homologous Moloney MuLV 70S RNA. (32 refs)

5762 **Structural Analysis of the Surface Glycoproteins (GP70's) of Recombinant Murine C-type Viruses: Evidence for Envelope Gene Recombination** (Eng) Elder, J. H. (Scripps Clinic and Res. Foundation, 36 N. Torrey Pines Road, La Jolla, CA); Jensen, F. C.; Tsch, J. W.; Lerner, R. A.; Vogt, M. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the 5th International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J. H.; John, D. S., eds. (New York: Elsevier/North Holland Medical Press): 503 pp.; 156-159; 1978.

surface glycoprotein gp70 from parental and recombinant murine C-type viruses was analyzed by tryptic peptide fingerprinting. The recombinant viruses tested included R-247, HIX, MoMCF22, and the radiation-induced leukemia virus 136.5. The results showed that the recombinants are the products of recombination between ecotropic and xenotropic C-type viruses and that recombination took place within the envelope gene encoding gp70. (12 refs)

5763 **Scanning Electron Microscopy of Lymphoid Cells from Leukemia Virus-infected Mice.** (Eng) Farber, P. A. (Dept. Pathology, Temple Univ. Sch. of Medicine, Philadelphia, PA, 19122); Specter, S.; Friedman, R. In: *Infect Immun* 20(3): 816-826; 1978.

spleen, lymph node, bone marrow, and thymus cells from Friend leukemia virus (FLV)-infected inbred male BALB/c mice were examined at various times after infection by scanning electron microscopy. The presence of cells with surface immunoglobulin (SIg) molecules, theta antigen, and leukemia virus-associated or -induced surface antigens was determined by immunofluorescence. Within 3-5 days after infection, the number and percentage of spleen cells with villous projections decreased and, concomitantly, the number of cells that were larger and smoother increased. Within 10-20 days after infection, the majority of splenocytes were smooth, large cells showing many distinct morphological changes, including surface holes and a spongy appearance. By 25 to 35 days after infection, most splenocytes were abnormal in appearance. Similar changes occurred in the lymph nodes, but at a slower rate. Thymus and bone marrow cells showed little change in surface morphology until late in infection, and even then only a few cells appeared abnormal. Changes in the spleen cells but not the lymph node cells paralleled a rapid decrease in the percentage of cells that were positive for SIg and theta antigen. FLV antigen appeared rapidly on spleen cells after infection; fewer lymph node cells were positive and only low numbers of marrow and thymus cells. The morphological changes detected among the lymphoid cell populations appear to correlate directly with the immunosuppression induced by the virus. These results, therefore, suggest that there is a direct relationship between infection and alteration of lymphoid cell type and morphology. (26 refs)

78-5764 **Requirement of Protein Synthesis for the Degradation of Host mRNA in Friend Erythroleukemia Cells Infected with Herpes Simplex Virus Type 1.** (Eng) Nishioka, Y. (Dept. Microbiology, Coll. Physicians and Surgeons, Columbia Univ., New York, NY, 10032); Silverstein, S. *J Virol* 27(3): 619-627; 1978.

Friend erythroleukemia (FL) cells were used to elucidate the viral mechanisms responsible for inhibition of the synthesis of host-specific proteins in cells infected with herpes simplex virus. Shortly after infection, cellular polyribosomes (PR) dissociated and cellular messenger RNA (mRNA) degraded. Analysis of infected cell extracts on sucrose density gradients demonstrated that most of the PR had dissociated to monoribosomes at 2 hr postinfection. Physical measurements of infected cells RNA's supported this conclusion and also showed that the polyadenylated RNA's had decreased in size. The degradation of mRNA was apparently stochastic, as the relative abundance of adenylated globin mRNA remained constant at 2 and 4 hr postinfection. However, it is possible that different cellular mRNA's may be degraded at different rates. In experiments on the effects of pactamycin and sodium fluoride on PR and globin mRNA, it was found that dissociation of PR was not sufficient to account for the rapid degradation of globin mRNA, as the amount of globin mRNA in uninfected cells did not change when cells were treated with these substances at concentrations sufficient to dissociate all PR. In cells infected with UV-irradiated virus, PR dissociated but globin mRNA did not degrade, which suggests that it is possible to separate the two events. It is concluded that the dissociation of PR appears to be mediated by a heat-labile, virion-associated function and that degradation of the mRNA requires expression of the virus genome. This conclusion is substantiated by additional experiments with superinfected FL cells, irradiated virus, and cycloheximide. (37 refs)

78-5765 **A Neutralization Bioassay for Friend Leukemia Virus.** (Spa) Basombrio, M. A. (Seccion Leucemia Experimental, Instituto de Investigaciones Hematologicas, Academia Nacional de Medicina, Buenos Aires, Argentina); Genovese, J. *Medicina (B Aires)* 38: 245-249; 1978.

A method for the titration of Friend Leukemia Virus (FLV) in mice was standardized. The method is based on the direct relationship, at high virus doses, between the virus dose injected and spleen wt 14 days later. If, before inoculation, the virus is incubated with specific antiserum for 1 hr at 37 C, neutralization occurs, and the prevention of splenomegaly is proportional to the concentration of virus-specific antibodies in the serum. Regressing sarcomas induced in mice by a sarcoma virus pseudotype of FLV induced resistance to challenge with FLV. Sera from these animals strongly neutralized the virus. Mice in which leukemia was induced by FLV also had virus-neutralizing antibodies, but to a lesser extent than the immunized mice. A preliminary search was carried out for antibodies to FLV in 20 normal human sera and in 29

sera from patients with leukemia. Although the serum from some subjects neutralized the virus, no significant differences were detected between the normal and leukemic samples. (11 refs)

- 78-5766 Intracellular Restriction of Ecotropic Murine Leukemia Virus in Rat NRK Cells and Its Abolishment by Adaptation.** (Eng) Yoshikura, H. (Dept. Genetics, Inst. Medical Science, Univ. Tokyo, Tokyo, Japan); Yoshida, M. *J Virol* 27(3): 612-618; 1978.

Ecotropic murine leukemia viruses (MuLV-E) are partially restricted in normal rat kidney (NRK) cells. This is true for both the N-tropic FN-2 (purified helper component of Friend leukemia virus) and the B-tropic WNB-2 (purified WN1802B BALB/c-derived endogenous virus) viruses. Successive passages of the viruses in NRK cells resulted in a stable adaptation that was manifested in increased plaquing efficiency and the formation of clear and large plaques in NRK cells. The NRK cells were more susceptible to FN-2 virus than to WNB-2 virus, and the adaptation was more complete in the former. The adaptation seemed to occur mainly at the level of intracellular growth of MuLV-E in NRK cells, and no changes in the mobilities of the viral proteins gp69/71 and p30 in sodium dodecyl sulfate-polyacrylamide gel electrophoresis were observed after the adaptation of FN-2 in NRK cells. It appeared that the restriction of MuLV-E in NRK cells was not due to penetration but mainly to intracellular growth, although the former may not be excluded completely. Since both FN-2 and WNB-2 were subjected to two successive limiting dilutions in mouse SC-1 cells and were obtained from cells infected with the virus diluted 2 logarithmic units higher than the limit of plaque formation in NRK cells, the adaptation was not due to a simple selection of NRK-nonrestricted virus that had existed before NRK passage. The adaptation seemed to be initially induced by a host modification, followed by selection. Possible commitment of xenotropic virus in the adaptation was excluded. The host-induced changes in FN-2 and WNB-2 were stable even if the adapted viruses were propagated back into SC-1 cells. The adaptation was host-specific; ie, MuLV-E adapted in NRK cells failed to replicate efficiently in other rat cell lines of different origins. This suggests that host cell-specific RNA or DNA sequences are involved in the adaptation. (22 refs)

- 78-5767 Generation of Oncogenic Type C Viruses: Rapidly Leukemogenic Viruses Derived from C3H Mouse Cells In Vivo and In Vitro.** (Eng) Rapp, U. R. (Lab. Viral Carcinogenesis, NCI, NIH, Bethesda, MD, 20014); Todaro, G. J. *Proc Natl Acad Sci USA* 75(5): 2468-2472; 1978.

The generation of oncogenic C-type variants from a poorly infectious virus isolated from C3H/10T1/2 mouse cells (low-leukemia strain) is reported. The parent virus did not

cause tumors upon inoculation into newborn NIH mice, but variants with increased infectivity for mouse were derived both in vivo and in vitro by selecting for variants able to grow to high titers. The highly infectious variants were able to induce mouse fibroblasts to grow in soft agar. Inoculation of the viruses into newborn NIH Swiss mice resulted in 100% mortality from leukemia within 4 months; tumors developed at the injection site. Both mouse-tropic dual-tropic viruses were isolated from the leukemic mice and plaque-purified. The first group of viruses produced syncytial plaques on rat XC cells and did not grow in mouse cells. The viruses of the other group replicated well in mouse and mink cells, producing morphologic changes similar to transformation but not XC syncytia. This indicates that they are members of the MCF class of murine C viruses. Isolates from either group were highly leukemogenic on retesting: the mean latent period was 67 days for a mouse-tropic virus and 105 days for a dual-tropic virus. It was concluded that there is a positive correlation between the ability of a mouse C-type virus to grow in cell culture and its leukemogenic potential. Furthermore, it is possible to convert a weakly infectious, nonleukemogenic virus and to convert it to a rapidly replicating, highly leukemogenic virus by passage either in cell culture or in the animal. (16 refs)

- 78-5768 The Friend Spleen Focus-forming Virus (SFFV) Genome: Fractionation and Analysis of SFFV and Helper Virus-related Sequences.** (Eng) Mak, T. W. (Ontario Cancer Inst., Univ. Toronto, Toronto, Ontario, Canada); Penrose, D.; Gamble, C.; Bernstein, A. *Virology* 81: 73-80; 1978.

High titers of Friend spleen focus-forming virus (SFFV) and its helper lymphoid leukemia virus (LLV) propagated in NIH/3T3 fibroblast cell cultures were analyzed by density fractionation and molecular hybridization to determine the composition of the SFFV genome. When DNA complementary to helper LLV RNA (cDNA-LLV) was hybridized to viral RNA from stocks containing SFFV in apparent excess and to helper LLV RNA, the results indicated that there were two populations of helper LLV sequences present in the RNA extracted from the Friend virus complex. cDNA complementary to sequences unique to the SFFV genome was fractionated by hydroxyapatite chromatography and was shown to be homologous to sequences found only on the SFFV, and not the LLV, genome. Three cDNA probes were used to determine virus-specific sequences in SFFV nonproducer mouse and cell clones: cDNA-LLV, SFFV, cDNA-LLV, and cDNA-SFFV. The results with the first probe suggested that the RNA of cells producing SFFV contained sequences found in cells producing LLV. The results with the second probe demonstrated that the two SFFV nonproducer clones expressed helper LLV sequences, but they suggested that the SFFV genome contained only a portion of the sequences found on the LLV genome. cDNA-SFFV hybridized only to RNA from cells productively or nonproductively infected with SFFV. The data suggest that the SFFV genome contains

a SFFV-specific sequences and a portion, but not all, of helper LLV sequences. (24 refs)

5769 **Antigenic Modulation of Leukemic Cells Induced by X-irradiation of the Host.** (Eng) Ioannides, H. L. (Dept. Pathology, Lenox Hill Hosp., New York, NY); Pearce, A.; Sabbath, M. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., Eds. (New York: Elsevier/North Holland Biomedical Press): 349-350; 1977.

Antigenic modulation was induced in vitro by adding the sera of X-irradiated rats to an established tissue culture line of leukemic cells that replicates Gross leukemia virus and is antigenically positive. Indirect immunofluorescence and immunoelectron microscope techniques showed that murine leukemia virus antigens formed patches and caps and gradually decreased in staining intensity. Virus particles were localized to only a few areas of the cell membrane. These changes were reversible. (9 refs)

5770 **Viral Gene Expression in Murine Sarcoma Virus (Murine Leukemia Virus)-infected Cells.** (Eng) Dina, D. (Dept. Genetics, Albert Einstein Coll. Medicine, New York, NY, 10461); Penhoet, E. E. *J Virol* 27(3): 873-875; 1978.

Levels of viral gene expression were analyzed in Moloney murine sarcoma virus (murine leukemia virus) 124 (MSV(MLV) 124)-infected NIH 3T3 cells, and the results were compared with those in MSV- and MLV-infected and uninfected cells. The MSV(MLV) complex was prepared from the supernatant of infected clone 124 TB cells. The MSV(MLV)-infected cells produced virions that contained about 99% MSV RNA and 1% MLV RNA. Differences between these concentrations were also present in the nucleus, cytoplasm, and polysomes. However, MLV RNA was expressed at much higher levels in cells infected by MLV alone. The most likely explanation for this situation is that, as a consequence of MSV(MLV) infection, MSV proviral DNA is present in excess, forcing MLV into secondary integration sites that are transcribed at a much lower level. In MSV(MLV)-infected cells, three to four copies of proviral DNA from each virus were integrated into the cellular DNA. Measurements of RNA specific for each of the genomes by hybridization to specific complementary DNA reagents revealed a 10- to 15-fold difference in concentration in both nuclear and polysomal RNA fractions, with MSV RNA predominating in both cases. The results support the idea that the production of MSV and MLV RNA's is regulated at the level of transcription, and they show that the basic differences in concentration found in the viral particles originate at least partially in the cell nucleus. (38 refs)

78-5771 **Characterization of Viral-Type Nucleotide Sequences Detected in the RNA of Human Leukemia Cells by Probes of Murine and Simian Origin.** (Fre) Tavitian, A. (Laboratoire d'Hematologie experimentale, Hopital Saint-Louis, 2, place du Dr. Fournier, F75475 Paris Cedex, France); Hamelin, R.; Larsen, C. J.; Marty, M.; Peries, J.; Boiron, M. *Bull Cancer (Paris)* 65(1): 25-30; 1978.

Molecular hybridization techniques were used to detect viral genomes in the RNA of cells from patients with acute myelocytic leukemia (AML), acute lymphocytic leukemia (ALL), and chronic myelocytic leukemia (CML) and in the RNA of normal WBC. Complementary DNA (cDNA) prepared from Moloney murine sarcoma-leukemia virus [M-MSV(MLV)] and from simian sarcoma virus and its associated virus [SSV-(SSAV)] was used as the molecular probe. Hybridization was slight or absent in cells from the blood and spleen of the four normal subjects. In cells from the 15 leukemia patients, hybridization ranged from 0% to 13.5% for the SSV-(SSAV) probe and from 0% to 43% for the M-MSV(MLV) probe. The probes did not react exclusively with any one type of leukemia cell. Nucleotide sequences common to both probes, isolated and used as a single probe, failed to detect viral RNA in leukemia cells that had hybridized with the complete cDNA from either M-MSV(MLV) or SSV-(SSAV). A recycled probe used in cells from three AML patients succeeded in hybridizing; therefore, the complementary nucleotide sequences are probably distributed along the viral genome. (10 refs)

78-5772 **5'-Terminal Nucleotide Sequences of Mammalian Type C Helper Viruses Are Conserved in the Genomes of Replication-defective Mammalian Transforming Viruses.** (Eng) Tronick, S. R. (Lab. RNA Tumor Viruses, NCI, Bethesda, MD, 20014); Cabradilla, C. D.; Aaronson, S. A.; Haseltine, W. A. *J Virol* 26(3): 570-576; 1978.

The RNA's of replication-defective murine and primate C-type transforming viruses were analyzed for nucleotide sequences homologous to the genomes of their helper viruses using complementary DNA (cDNA) probes generated from the total or 5'-terminal regions of Moloney murine leukemia virus and woolly monkey virus RNA. The defective viruses had been shown previously to vary in their ability to express helper *gag* gene proteins. With total RNA cDNA probes, the transforming viruses varied greatly in their representation of helper virus sequences (15%-60% hybridization). In contrast, the 5'-terminal-specific sequences of the helper virus were conserved in the RNA's of every transforming virus tested (> 80% hybridization). These findings suggest a critical role of these sequences in the life cycle of the defective transforming viruses. (41 refs)

78-5773 **A Comparative Study of Two Rat Viral Complexes: MSV(RaVL) and KiMSV(RHHV).**

(Eng) Yang, S. S. (NCI, Bethesda, MD, 20014); Ting, R. C. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977*. Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 264-266; 1978.

A Moloney sarcoma MSB-1 virus rat leukemia virus pseudotype and a Kirsten sarcoma virus rat hepatoma helper virus pseudotype were found to have a common sedimentation value of 1.145-1.155 g/cm³. They also demonstrated common antigenic determinants, using antisera specific against rat leukemia virus; no reactivity was demonstrated against mouse C-type virus antisera. Proteins of mol wt 110,000 (110K)-120K, 70K, 30K, and 7K-10K daltons were present in variant amounts; the nature of these proteins is outlined. (4 refs)

78-5774 Moloney Leukemia Virus-determined Cell Membrane Antigens. (Eng) Fenyo, E. M. (Dept. Tumor Biology, Karolinska Inst., 104 01 Stockholm 60, Sweden); Troy, F. A.; Siegert, W.; Klein, E. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977*. Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 331-336; 1978.

A cell surface antigen (MCSA) that is present in Moloney leukemia virus-induced lymphomas was separated from viral proteins and histocompatibility antigens (H-2) by a sequence of purification procedures. Sodium dodecyl sulfate-gel electrophoresis revealed two subunits with a mol wt of 92,000 and 50,000 daltons, respectively. A Moloney lymphoma cell line characterized by low MCSA expression produced a variant virus deficient in MCSA-inducing activity. This suggests that MCSA is virally determined. (14 refs)

78-5775 Production of 'Rapid-Harvest' Moloney Murine Leukemia Virus by Continuous Cell Culture on Synthetic Capillaries. (Eng) Ratner, P. L. (Dept. Biochemistry, Univ. Kentucky Medical Center, Lexington, KY, 40506); Cleary, M. L.; James, E. *J Virol* 26(2): 536-539; 1978.

The use of synthetic capillaries for the continuous large-scale recovery of Moloney murine leukemia virus (M-MuLV) from chronically infected NIH/3T3 cells is reported. With this method, the M-MuLV could be automatically harvested within 60 min of release from the cells in quantities of approx 100 ml/24 hr/capillary unit. Virus production and infectivity were measured by determination of reverse transcriptase activity and by the XC syncytia assay. The optimum period for harvesting the virus was from days 4 to 11 inclusive. A max

infectious titer of virus ($> 10^9$ plaque-forming units/ml) noted by day 8. These results indicate that the use of synthetic capillary units for the production of rapid-harvest RNA tumor virus is an alternative to the roller bottle and flask methods. (24 refs)

78-5776 Relationship of Retrovirus Polypeptide Cages to Virion Maturation Studied with Temperature-sensitive Murine Leukemia Virus Mutants. (Eng) Witte, O. N. (Dept. Biology, Massachusetts Inst. Technology, Cambridge, MA, 02139); Baltimore, D. *J Virol* 27: 750-761; 1978.

The processing of murine leukemia virus (MuLV) polypeptides in cells infected by the temperature-sensitive (*ts*) mutants Moloney *ts3* and Rauscher *ts4* was studied. Both *ts3* and *ts4*-infected cells grown at the restrictive temperature (32.5°C) accumulated large numbers of late-budding structures and small numbers of early-budding forms. Cells infected with wild-type Moloney MuLV or *ts3* or *ts4* at the permissive temperature (32.5°C) contained budding structures at all stages of development. At the restrictive temperature, *ts* mutants accumulated the core precursor polypeptides Pr65(gag) and Pr180(gag-pol); little Pr80(env) or Pr90(gag-pol) was cleaved, but a large fraction was degraded. After a shift to the permissive temperature, Pr65(gag) and Pr180(gag-pol) were cleaved and released into virions. The final products p30 and p85, built up with time in the virions, and p30 and p85, was maintained at a steady-state level in the infected cells. Uncleaved Pr65, Pr180, and Pr130 were all reduced by incubation at the permissive temperature in serum-free medium, and the appearance of finished reverse transcriptase was evident. Reduction of the gag intermediates and an increase in p30 also occurred. Reverse transcriptase as an active enzyme appeared only when Pr180(gag-pol) had been cleaved to form the active p85 protein. This cleavage occurred not only, if not entirely, after the release of virions from the cells. (39 refs)

78-5777 Identification of Virus Found in Mouse Lymphomas Induced by HIX Murine Oncornavirus. (Eng) Fischinger, P. J. (Lab. Viral Carcinogenesis, NCI, Bethesda, MD, 20014); Dunlop, N. M.; Blevins, C. S. *J Virol* 26(2): 532-535; 1978.

Four lymphomas induced by pure HIX murine leukemia virus, two each in BALB-2 and NIH Swiss mice, were found to contain large amounts of virus with amphotropic properties. Analysis of tumor-derived virus purified by limiting dilution techniques indicated that its interference and neutralization spectra were essentially identical to parental HIX virus and different from ecotropic and xenotropic viruses. Examination of virus progeny from many individual foci induced by virus derived directly from lymphomas indicated that each infectious unit contained only HIX virus. No evidence of ecotropic virus was observed. (13 refs)

778 **Structural Studies on Rauscher Murine Leukemia Virus: Isolation and Characterization of Envelopes.** (Eng) Van de Ven, W. J. (Dept. Biochemistry, Univ. Nijmegen, Geert Grooteplein Noord 21, Nijmegen, Netherlands); Vermorken, A. J.; Onnekink, C.; Bloemers, H. Bloemendal, H. *J Virol* 27(3): 595-603; 1978.

Isolation and characterization of viral envelope structures from disrupted virions of Rauscher murine leukemia virus are reported. The virions were isolated from tissue culture medium of chronically infected NIH Swiss 3T3 mouse fibroblast cells. Disruption of the virions was achieved with 1% dodecyl sulfate or Nonidet P-40 in combination with 1% Triton X-100. Fractionation into a viral core fraction and a viral envelope fraction was performed through flotation in a discontinuous sucrose gradient. The amount of the *gag* and *env* gene-encoded structural polypeptides in the virions and the isolated envelope fraction was compared using sensitive competition radioimmunoassays or sodium dodecyl sulfate-polyacrylamide gel electrophoresis after immunoprecipitation with polyvalent and monospecific antisera directed against Rauscher murine leukemia virus proteins. The dominant viral structural polypeptides in the purified envelope fraction were the *env* gene-encoded polypeptides gp70, gp12(E), and p12(E). Only relatively small amounts of the *gag* gene-encoded structural polypeptides p30, p12, and p10 were present in the envelope fraction; however, a relatively high amount of the *gag* gene production p15 was detected in this fraction. (30 refs)

5779 **Immune Lysis of Type C Viruses by Antisera to Purified Viral Subunits.** (Eng) Oroszlan, S. (National Cancer Institute, Frederick Cancer Res. Center, P.O. Box B, Frederick, MD, 21501); Gilden, R. V. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 90-93; 1978.

Immune virolysis was used to study the localization of specific viral polypeptides, evaluate the specificity of antisera, and determine the presence of host cell antigens in the viral envelope. Complement-dependent specific lysis of labeled Rauscher murine leukemia virus was obtained with anti-gp70 and anti-p15(E) sera, confirming the surface localization of these subunits. Lysis by antisera to normal cells and by specific antisera to serologically defined alloantigens demonstrated that host antigens are incorporated into the viral envelope. (12 refs)

5780 **Impaired Gene Expression of Rauscher Murine Leukemia Virus.** (Eng) van de Ven, W. J. (Dept. Biochemistry, Univ. Nijmegen, Geert Grooteplein Noord 21,

Nijmegen, Netherlands); van Zaane, D.; Vermorken, A. J.; Bloemers, H. P. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 163-167; 1978.

The impaired expression of the genes coding for the structural polypeptides (*gag* and *env* gene products) of Rauscher murine leukemia virus (R-MuLV) was studied. In NIH/3T3 cells infected with temperature-sensitive R-MuLV mutants, impaired expression of the *gag* gene was observed at the restrictive temperature. *Gag*-pr65 accumulated, but *gag*-pr75 disappeared and p30 was not produced. Production of the *env*-related polypeptide p12(E) was also inhibited at the restrictive temperature. In R-MuLV-infected JLS-V9 cells, cytochalasin B (25 µg/ml) reversibly inhibited the synthesis of the glycosylated precursor peptide *env*-pr82 and the production of virus-specific gp69/71, p15(E), and p12(E). A similar but irreversible inhibition occurred in the presence of 2-deoxy-D-glucose (15 mM). (11 refs)

78-5781 **Origin of the Minor Glycoproteins of Murine Leukemia Viruses.** (Eng) Kemp, M. C. (Dept. Microbiology, Univ. Alabama in Birmingham, University Station, Birmingham, AL, 35294); Wise, K. S.; Edlund, L. E.; Acton, R. T.; Compans, R. W. *J Virol* 28(1): 84-94; 1978.

An attempt was made to prove that the minor virus-associated glycoproteins from Rauscher murine leukemia virus (R-MuLV) and from AKR thymic lymphoblastoid BW5147 cell membranes are proteolytic cleavage products of the major glycoprotein gp70. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis of R-MuLV polypeptides grown in JLS-V9 cells and labeled with ³H-glucosamine or ³H-galactose + ¹⁴C amino acids indicated the presence of two glycosylated polypeptides. The larger one weighed 72,000 (gp70), the smaller one 52,000 (gp52). Antiserum to gp70 precipitated both, indicating that they were antigenically related. When labeled R-MuLV virions were digested with increasing concentrations of trypsin and analyzed with SDS-PAGE, the amount of gp70 was reduced, with a corresponding direct increase in a smaller peptide weighing 52,000. The R-MuLV infectivity titer decreased by 69% upon digestion with 40 µg/ml trypsin. When gp70 and gp52 were digested with Pronase and treated with neuraminidase, gp70 was split into two polypeptides of mol wt 5,100 and 2,900. The 5,100-dalton fragment was the only significant entity resulting from gp52 digestion. Glycoproteins from detergent-extracted AKR thymic lymphoblastoid cell membranes were labeled with ¹²⁵I and subjected to radioimmuno-precipitation with various antisera. Goat anti-R-MuLV gp70 precipitated components of 40,000-50,000 mol wt and, to a lesser extent, a component of 30,000 mol wt. Tryptic digestion of iodinated glycoproteins from BW5147 cells produced

similar results. The appearance of similar glycoprotein components in AKR membrane preparations and R-MuLV preparations supports the contention that these minor glycoproteins arise from a cleavage of gp70. (39 refs)

78-5782 Immune Response to Rauscher Virus-induced Leukemia in DBA Mice. II. Correlation Between Antigenic Expression and Pathogenesis. (Eng) Toth, F. D. (Inst. Microbiology, Univ. Medical Sch., 4012 Debrecen, Hungary); Vaczi, L. *Neoplasma* 25(3): 265-272; 1978.

Immune responses to Rauscher virus-induced leukemia in DBA mice were studied to determine the mechanism leading to the lethal exacerbation of the disease. On the basis of changes in spleen wt, three stages of leukemia were distinguished: early progression, regression, and exacerbation. A significant tumor-specific (TS) antibody titer was first detected on the 12th after virus infection. This was followed by a fast rise to day 40, when the serum antibody level began to decline; no significant titer could be detected after day 60. During early progression (days 10-18) and regression (days 35-40), the sera of leukemic mice contained circulating immune complexes, with TS antibody in excess. This soluble antigen was identical to one of the two soluble TS antigens in the serum of Rauscher leukemic BALB/c mice. During exacerbation (days 50-65), soluble antigen was not found in the serum; thus, its accumulation in the circulation was not responsible for the lethal progression of the disease. Two types of disappearance of TS antigen were observed. Immediately before the beginning of regression, antigen could not be detected even after neuraminidase treatment. This and a rapid decrease in the number of tumor cells indicated that the antigenic receptor sites were lost as a consequence of the cytotoxic reaction mediated by TS antibodies. The presence of antigen-antibody complexes in the serum provided further evidence for this assumption. On the other hand, a new population of leukemic cells emerged during regression, and it appeared that these cells lost their specific surface antigenicity, probably because of masking of the receptor sites by sialic acid. The results illustrate that the simultaneous loss of immunosensitivity and immunogenicity of a virus-induced leukemia can occur spontaneously. The correlation between antigenic expression and pathogenesis raises questions about in vivo demasking of TS antigen receptors. (24 refs)

78-5783 Retention of Transformant Specific Type III Collagen in Dibutyl cAMP Treated Kirsten Sarcoma Virus Transformed BALB 3T3 Cells and in a Flat Revertant. (Eng) Hata, R. (Dept. Tissue Physiology, Medical Res. Inst., Tokyo Medical and Dental Univ., Tokyo, Japan); Peterkofsky, B. *J Cell Physiol* 95(3): 343-352; 1978.

Kirsten sarcoma virus-transformed BALB 3T3 cells (Ki-3T3 cells) were treated with dibutyl cyclic AMP (dbcAMP) to determine if the collagen phenotype could be restored to that

of nontransformed cells. Ki-3T3 cells contained mainly I collagen [$\alpha_1(I)$] $_{\alpha_2}$, and approx 30% of a collagen designated type III [$\alpha_1(III)$] $_{\alpha_2}$. BALB 3T3 cells contained mainly I collagen and about 25% of another type (X) composed of three α_1 chains. Procollagen labeled with radioactive proline was isolated from the medium or cells of cultures and converted to collagen with pepsin. The collagen was analyzed by carboxymethylcellulose (CMC) chromatography or gel electrophoresis under denaturing conditions. Ki-3T3 cells treated with 0.5 mM dbcAMP continued to accumulate type III collagen, but there was an increase in the number of α_1 chains eluting from CMC columns in the same position as $\alpha_1(I)$, suggesting increased accumulation of type X collagen. Although the revertant was similar to dbcAMP-treated cells, that it exhibited a flattened morphology and a high relative rate of collagen synthesis, the collagen profile was similar to that of the transformant, consisting mainly of types I and X. These findings indicate that accumulation of type III collagen is unaffected by dbcAMP, but they suggest that cAMP may be involved in the regulation of type X collagen. The failure of dbcAMP or reversion to affect the occurrence of type X collagen supports the mechanism of cell selection as a means of explaining the specific occurrence of type III collagen in Ki-3T3 cells. (18 refs)

78-5784 Helper-independent Transformation by Unintegrated Harvey Sarcoma Virus DNA. (Eng) Lowy, D. R. (Dermatology Branch, NCI, Bethesda, MD 20014); Rands, E.; Scolnick, E. M. *J Virol* 26(2): 291-297; 1978.

A study was made of the unintegrated infectious DNA of Harvey sarcoma virus (Ha-SV) and Moloney leukemia virus (M-MuLV) obtained from the Hirt supernatant fraction of cells acutely infected with these viruses. To obtain a quantitative assay for infectious viral DNA, recipient mouse cells were first exposed to calcium phosphate-precipitated viral DNA and then treated with dimethyl sulfoxide. Infectivity was monitored by focus formation for Ha-SV and plaque formation for M-MuLV. The viral DNA titration curve followed single-hit kinetics for both foci and plaques, indicating that a single molecule carried information for both functions. Focus-forming and plaque-forming activities were present in different molecules, since these two biological activities could be separated from each other by agarose gel electrophoresis. The focus-forming molecule was linear DNA with a mol wt of approx 4×10^6 daltons. The focus-forming activity of the viral DNA was sensitive to *EcoRI* and resistant to *XhoI* restriction endonucleases; plaque-forming activity was resistant to the former but sensitive to the latter. The generation of helper-independent foci indicates that Ha-SV DNA can transform mouse cells in the absence of helper virus or its proteins. (36 refs)

78-5785 The Occurrence of Feline Leukaemia Virus Neutralizing Antibodies in Cats. (Eng) Russell, J. W.

I. (ICRF Tumour Immunology Unit, Dept. Zoology, Coll. London, London W.C.1E 6BT, England); Jarrett, *Int J Cancer* 22(3): 351-357; 1978.

occurrence of feline leukemia virus-neutralizing antibodies (ab's) to subgroups A (FeLV-A), B (FeLV-B), and C (FeLV-C) were determined in four cat populations: specific-pathogen-free (SPF) cats, urban and suburban free-range cats, cats from FeLV-infected multiple cat households (MCH's), and leukemic cats. The prevalence of anti-FeLV-A was closely related to the frequency of isolation of FeLV-A in each population, except in leukemic cats. Thus, it was found in 0% of SPF cats, 4% of FR cats, and 42% of MCH cats. In individual MCH cats, there was a strong correlation between the presence of ab's to FeLV-A and the presence of active FeLV viremia. Anti-FeLV-B ab's were encountered less frequently, although FeLV-B was present in approximately half of the FeLV-B isolates. This was likely due to restricted growth of FeLV-B in cats. Surprisingly, in view of the rare isolation of FeLV-C, the prevalence of anti-FeLV-C ab's was high: 11% in FR, 51% in MCH, and 100% in leukemic cats (none were found in SPF cats). Their occurrence was not related to whether or not a cat was viremic. Anti-FeLV-C ab's are likely evoked by an antigen that is expressed in cats during infection with FeLV-A: a possible candidate is a determinant coded for by endogenous FeLV-like genes present in cat cells. An interesting finding was that anti-FeLV-C ab's in cats with lymphosarcoma, especially FeLV-negative cases, which implies that the cats had been exposed at some time to FeLV-A. To determine how these infections might have arisen, cats from all the populations were experimentally infected by contact (C) or inoculation and examined for the presence of virus and neutralizing antibodies. The outcome of the infection was determined within 1 month in I cats and 10 wk in C cats. In I cats, there was a marked positive correlation between the development of viremia and the appearance of anti-FeLV-A ab's. However, the three leukemic cats, anti-FeLV-C ab's were found in the absence of FeLV-A. (21 refs)

78-5786 **Feline Oncornavirus Associated Cell Membrane Antigen: A Tumor Specific Cell Surface Marker** (Eng) Essex, M. (Dept. Microbiology, Harvard Sch. Public Health, Boston, MA, 02115); Sliski, A. H.; Hardy, W. D.; Noronha, F.; Cotter, S. M. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 26 August, 1977*. Bentvelzen, P.; Hilgers, J.; Yohn, D. S., Eds. (New York: Elsevier/North Holland Biomedical Press): 337-340; 1978.

Cultured cells infected with feline leukemia virus (FeLV) or transformed with feline sarcoma virus (FeSV) were examined for the FeLV structural proteins p30 and gp70 and for feline oncornavirus-associated cell membrane antigen (FOCMA). FOCMA expression was found to be transformation-specific

and independent of gp70 or p30 expression. Feline lymphoma and leukemia cells also expressed FOCMA, even in the absence of FeLV p30 and gp70. Lymphoid cells from normal cats, even those infected with FeLV, were negative for FOCMA. (9 refs)

78-5787 **Presence of 5'-Terminal Cap Structures in Virus-specific RNA from Feline Leukemia Virus-infected Cells.** (Eng) Thomason, A. R. (Dept. Biology, Univ. California at Los Angeles, Los Angeles, CA, 90024); Friderici, K. H.; Velicer, L. F.; Rottman, F. *J Virol* 26(2): 226-235; 1978.

Experiments were conducted to determine whether intracellular Rickard strain Friend leukemia virus (R-FeLV)-specific RNA (vRNA) acting as messenger RNA (mRNA) is capped even though virion subunit RNA is not capped. F-422 feline thymus tumor cells, chronically infected with R-FeLV, were labeled with ^{32}P and the total cytoplasmic RNA was isolated. The RNA was centrifuged through sucrose gradients, and R-FeLV vRNA was located by hybridization of portions of the gradient fractions to R-FeLV complementary DNA. vRNA classes with sedimentation coefficients of approximately 36S, 28S, 23S, and 15S were identified. Each class of RNA was recovered and hybridized with mercurated R-FeLV complementary DNA, and the hybrids were chromatographed on columns of sulfhydryl-Sepharose to separate them from unhybridized cellular RNA. Although insufficient amounts of 36S and 28S vRNA were obtained for further analysis, the 23S and 15S vRNA classes were analyzed to determine the nature of their 5' termini. Each of these vRNA classes contained stoichiometric amounts of cap structures per unit length of RNA, consistent with the presence of one cap per molecule. The structure of the 23S vRNA cap was $\text{m}^7\text{G}^{\text{ppp}}\text{5}'\text{GmpAp}$, and that of the 15S vRNA cap was $\text{m}^7\text{G}^{\text{ppp}}\text{5}'\text{GmpGp}$. (34 refs)

78-5788 **Distribution of Bovine Leukemia Virus Proviral DNA Sequences in Tissues of Animals with Enzootic Bovine Leukosis.** (Eng) Kettmann, R. (Departement de Biologie Moléculaire, Université Libre de Bruxelles, Brussels, Belgium); Burny, A.; Cleuter, Y.; Ghysdael, J.; Mammerickx, M. *Arch Int Physiol Biochim* 85(5): 989-991; 1977.

Bovine leukemia virus (BLV) single-stranded complementary DNA (cDNA; ^3H -thymidine-labeled DNA complementary to BLV 70S RNA) was used to study the distribution of proviral DNA sequences in tissues (normal or malignant) of bovine, ovine, porcine, and human origin. After recycling against normal bovine DNA, BLV ^3H -cDNA hybridized to bovine enzootic tumor DNA but failed to hybridize to normal bovine DNA, indicating that BLV is an exogenous RNA oncogenic virus and that enzootic bovine leukosis (EBL) is an infectious disease. Proviral BLV sequences were also detected in buffy-coat cells of animals with persistent lymphocytosis

and carrying antibodies to BLV but no tumors. In animals at the tumor stage of EBL, the proviral sequences were found in buffy-coat cells, in solid tumors (lymphosarcomas), and in organs infiltrated with tumoral lymphoid cells but not in apparently normal organs. No hybridization above background was observed between BLV 'H-cDNA and DNA's extracted from buffy-coat cells and tumors corresponding to sporadic forms of bovine leukosis, porcine leukemia, and some leukemias and sarcomas. (3 refs)

- 78-5789 Localization of Papillomavirus and Virus-specific Antigens in the Skin of Tumor-bearing *Mastomys natalensis* (GRA Giessen).** (Eng) Reinacher, M. (Veterinar-Pathologisches Institut, Justus-Liebig-Universität Giessen, Frankfurter Strasse 96, D-6300 Giessen, W. Germany); Muller, H.; Thiel, W.; Rudolph, R. L. *Med Microbiol Immunol (Berl)* 165(2): 93-99; 1978.

Examination of the skin tumors of *Mastomys natalensis* (yellow, red-eyed Giessen) by electron microscope and immunofluorescence studies demonstrated *M. natalensis* papillomavirus particles and viral antigens in distinct epithelial layers in all tumors. Twenty-six spontaneous tumors (13 keratoacanthomas, 3 squamous cell carcinomas, 10 papillomas) and 24 experimentally induced tumors (20 keratoacanthomas, 4 squamous cell carcinomas) were collected from 33 male and 17 female mice. Upon EM investigation, virus particles were seen in the upper layers of the thickened stratum granulosum and in the stratum lucidum and the stratum corneum. In intact cells, virus particles were present exclusively in the nucleus; with increased keratinization, however, particles were distributed throughout the organelle-damaged cells. Particles were never seen in the stratum basale, stratum spinosum, or in the lower layers of the stratum granulosum. Antisera from tumor-bearing animals and laboratory rabbits produced identical fluorescence in each tumor type. The fluorescent areas corresponded to the location of the virus particles identified by the EM. All virus-specific material was restricted to the same distinct cell layers in all tumors, sites identical to those where papillomavirus-specific material has been demonstrated in papillomas of other species, including humans. (7 refs)

- 78-5790 Virus-induced Papillomas of the Alimentary Tract of Cattle.** (Eng) Jarrett, W. F. (Dept. Veterinary Pathology, Univ. Glasgow Veterinary Sch., Bearsden, Glasgow, G61 1QH, Scotland); Murphy, J.; O'Neil, B. W.; Laird, H. M. *Int J Cancer* 22(3): 323-328; 1978.

The incidence of alimentary papillomas (AP) in 7,746 cattle of a wide age range from northern England and Scotland was determined in an abattoir survey. The overall incidence of AP was 19%, and the tumors were found at four sites, the tongue, palate, oropharynx, and esophagus. Most (75%) of the tumor-bearing animals (TBA) had one AP and only 8% had

more than three; in addition, 95% of the TBA had tumors at one site and only 5% had them at two sites. Tracing farms of origin indicated that TBA could be as old as 10 and as young as 7 mo, that most of the animals did not come from upland farms with bracken infestation, and that AP could be found in both beef and dairy cattle from lowland farms. The four AP sites were identical to those at which carcinoma have been noted in animals from an upland farm area with a high alimentary carcinoma incidence. However, the incidence of AP, the number of sites involved per animal, and the number of AP per animal were much higher in the high-cancer-area animals, indicating the presence of an amplifier of AP, possibly bracken fern. Inclusion bodies, identified by electron microscopy as viruses, were found in cell nuclei, and a typical papilloma virus was purified from the tumors. The histology of the tumors is described, and the possible plurality of bovine papilloma viruses is discussed in the light of recent findings in human viruses. It appears that virus-induced AP are widespread in the general cattle population but at a low tumor multiplicity. Under certain environmental circumstances, there is an amplification of AP. Tumors and a proportion of animals develop alimentary carcinoma at the AP sites. The relationship between ingestion of bracken fern (and other mutagenic and immunosuppressive agents) and a variety of isolates of bovine papilloma viruses in inducing tumors is being studied. (15 refs)

- 78-5791 The Occurrence of Human Wart-Virus Antibodies in Dogs, Pigs and Cattle.** (Eng) Pyhälä, S. (Lab. Viral Immunopathology, Dept. Virology, University of Helsinki, Helsinki, Finland); Neuvonen, E. *Arch Virol* 297-305; 1978.

Dog, pig, cattle, horse, and reindeer sera were screened for antibodies against human papilloma virus (wart virus) by immunodiffusion. Human wart virus antigen was prepared from surgically removed skin warts. Antibodies against human wart virus were detected in 28/114 dog sera. Investigation of the history of nine dogs showed that two had typical papillomas. The prevalence of antibodies increased with the age of the dog. The prevalence was similar to, but generally less than, that in human children of the same age. A faint precipitation line corresponding to that for the antigen was found by 10/97 cattle sera and 5/40 pig sera. No precipitation was seen in 59 horses or 100 reindeer sera. The specificity of antibody reaction could also be demonstrated by electron microscopy. For comparison, 50 human sera were studied for the presence of bovine papilloma virus antibodies. Nine formed antibodies against a virus proved to be bovine papilloma virus by electron microscopy. These results question the concept of strict species specificity of the various papilloma viruses. (23 refs)

- 78-5792 Nucleotide Sequence Studies of Polyoma Virus. The *Hpa* II 3/5 Junction to the *Hpa* II**

18 Junction, Encoding the Origin of DNA Replication at the 5' End of the Early Region. (Eng) Friedmann, T. (pt. Pediatrics, Univ. California, San Diego, CA); Late, P.; Esty, A. *J Biol Chem* 253(18): 6561-6567; 1978.

The nucleotide sequence of polyoma DNA from near the *Hpa* I/5 junction to the *Hpa* II 4/*Hae* III 18 junction was determined by a modified chemical method. The sequence contains 878 base pairs, including the origin of DNA replication and the region known to encode the hr-t function. The region corresponding to the origin of DNA replication contains several short repeated sequences and palindromes. There is a 30-base pair region with striking similarity to the corresponding region of simian virus 40 (SV40); as in SV40, a portion of that sequence is capable of forming a stable hair-loop. In the region encoding the hr-t function, there is apparently a single open reading frame extending from position 188 to the *Hpa* II 4/*Hae* III 18 junction. The potential translation product of this open frame begins with an initiation codon starting at position 188, and the first five amino acids of the product are methionine-asparagine-arginine-valine-leucine. This sequence is similar to the NH₂-terminal five amino acids of SV40 small t antigen known from nucleotide sequence. Amino acid sequencing to be methionine-asparagine-lysine-valine-leucine. Comparison of the nucleotide and amino acid homologies between the early regions of polyoma virus and SV40 will shed light on the mechanisms of cell transformation by these viruses and on the RNA processing events involved in the biosynthesis of small and large T antigens. (27 refs)

5793 Radioimmunoassay of Type D Oncovirus from Continuous J-96 Cells. (Eng) Vlasenkova, N. K. (I. Ivanovsky Inst. Virology, USSR Acad. Medical Sciences, 123098 Moscow, USSR); Kitsak, V. Y.; Altstein, D.; Zhdanov, V. M. *Acta Virol (Praha)* 22(4): 257-262; 1978.

The preparation of extracellular J-96 virus (produced by continuous cultures of human J-96 cells) and an extract of J-96 cells were examined by radioimmunoassay (RIA) for the presence of antigenic determinants of the major internal protein (p25) of Mason-Pfizer monkey oncovirus (M-PMV) and interspecies antigenic determinants of the major internal protein (p30) of murine leukemia virus (MuLV). The J-96 virus preparation replaced ¹²⁵I-p25 M-PMV almost completely (83%), and the competition curves obtained with M-PMV, J-96 virus, and the J-96 cell extract had the same slope. These data indicate that J-96 and M-PMV have identical antigenic determinants of their major internal proteins, as detected by RIA. No group-specific antigenic determinants of p30 MuLV and none of three groups of interspecies determinants characteristic of mammalian C-type viruses (¹²⁵I-p30 MuLV-anti-p27 feline leukemia virus; ¹²⁵I-p30 MuLV-anti-p28 baboon endogenous virus) were detected in J-96 virus or the J-96 cell extract. Failure to detect determinants characteristic of mammalian C-type viruses may be explained by a lack of perma-

nent production of C-type virions by J-96 cells, by the presence of C-type virions in a concentration below the threshold of sensitivity of the RIA, or by the presence of a C-type virus that has no interspecies determinants characteristic of most mammalian C-type viruses. (15 refs)

78-5794 Morphological Studies of Hamster Brain Cells Transformed In Vitro by a Human Papovavirus BK. (Eng) Tanaka, R. (Brain Res. Inst., Niigata Univ., 1 Asahimachi, Niigata 951, Japan); Sekiguchi, K.; Ueki, K.; Gilden, D. H.; Koprowski, H. *Acta Neuropathol (Berl)* 41(3): 235-239; 1978.

The morphological characteristics of in vitro BK virus-transformed hamster brain cells (HBBK) were investigated. These cells had a rapid rate of growth and an increased capacity to persist in serial subcultures. HBBK cells were round or polygonal, with a high nucleo-cytoplasmic ratio, marked pleomorphism, and a tendency to epithelial arrangement and multilayered growth. The cells had a round to oval nucleus with numerous nucleoli; mitotic figures and multinucleated cells were frequent. When cell growth was suppressed, the HBBK cells developed long, delicate processes. Specific immunofluorescence for glial fibrillary acidic protein was diffusely distributed in the cytoplasm of almost all HBBK cells. Sc injection of 10⁶ HBBK cells into newborn hamsters resulted in poorly differentiated gliomas in 100% of the animals within 4-5 wk. These findings suggest that the virus selectively transformed hamster brain glial cells. (18 refs)

78-5795 Cleavage Map of BK Virus DNA with Restriction Endonucleases *Mbo* I and *Hae* III. (Eng) Yang, R. C. (Dept. Biochemistry, Cornell Univ., Ithaca, NY, 14853); Wu, R. *J Virol* 27(3): 700-712; 1978.

The specific cleavage and physical mapping of the DNA of a variant of BK virus BKV(MM), with the use of the restriction enzymes *Mbo*I and *Hae*III are described. *Mbo*I, which recognizes a tetranucleotide sequence, was first used to cleave BKV(MM) DNA to completion. The resulting 10 *Mbo*I fragments were ordered and mapped on the genome by two techniques: *Hind*III-*Mbo*I reciprocal digestions of the uniformly ³²P-labeled DNA fragments and by *Mbo*I partial digestions of the single-end-labeled *Hind*III subfragments. Cleavage of BKV(MM) DNA with *Hae*III produced 21 fragments. Determination of the *Mbo*I sites and the sites from eight other enzymes of BKV(MM) DNA allowed the 21 sites of *Hae*III to be mapped by similar techniques. Even though the BKV(MM) stock contained three species of related viruses, the two techniques allowed the location of the variant DNA to be identified. The region of variation was positioned between map positions 0.64 to 0.71, corresponding to *Hae*III-E. Thus, the specific cleavage map of the 41 sites with their relative positions marked on the genome was constructed. The size of the resultant fragments ranges from 6 to 614

nucleotides in length. At least 80% of the fragments obtainable from the present map are suitable for complete sequence analysis. (24 refs)

- 78-5796 Rescue of Infectious Virus from Permissive Monkey Cells Containing Simian Virus 40 DNA Fragments.** (Eng) Moyer, R. C. (Thorman Cancer Res. Lab., Trinity Univ., San Antonio, TX, 78284); Moyer, M. P.; Gero-detti, M. H. *J Virol* 26(2): 272-280; 1978.

Studies were conducted to determine whether defined simian virus 40 (SV40) DNA subgenomes had any detectable biological activity in permissive TC7 monkey cells and whether subviral information could be maintained by the cells in a potentially biologically active form. Permissive cells were separately transfected with SV40 *EcoRI/Hap II A* (74% genome) DNA fragments and *EcoRI/Hap II B* (26% genome) DNA fragments in the presence of diethylaminoethyl-dextran. Fusion of the cells receiving the A fragment [TC7 (SV40/74) cells], with cells receiving the B fragment [TC7 (SV40/26) cells] resulted in SV40 rescue. TC7 (SV40/74 + 26) cells, which had simultaneously received both complementary subgenomes, either spontaneously produced SV40 upon subculture or yielded virus upon treatment with iododeoxyuridine. Furthermore, fusion of rat cells containing the *EcoRI/Hap II A* fragment with TC7 (SV40/26) cells resulted in SV40 rescue. The nature of the released virus was verified by cytopathology, V-antigen production, neutralization, and electron microscopy. No infectious virus was produced when the combinations of cells fused did not total a complete SV40 genome equivalent. (42 refs)

- 78-5797 The Relationship of SV40 Replicating Chromosomes to Two Forms of the Non-replicating SV40 Chromosomes.** (Eng) Seidman, M. M. (Dept. Biochemical Science, Princeton Univ., Princeton, NJ, 08540); Garon, C. F.; Salzman, N. P. *Nucleic Acids Res* 5(8): 2877-2893; 1978.

The relationship of the structure of replicating (R) simian virus 40 (SV40) chromosomes (Chr) to the two conformations of the Form I Chr [nonreplicating (NR) SV40 minichromosomes] was investigated by extracting R-Chr and studying them before and after incubation in an in vitro replication system in which replicative DNA synthesis occurs and progeny Form I Chr are generated. R-Chr and NR-Chr were extracted from the nuclei of infected BSC-1 cells using a detergent-free hypotonic buffer soln consisting of 0.5 mM NaH_2PO_4 (pH 7.5), 0.05 M sucrose, and 1 mM dithiothreitol. This procedure produced 20%-30% of the R-Chr with no loss of endogenous Form I DNA during the in vitro replication assays. The R-Chr and NR-Chr were analyzed many times on sucrose gradients. Two forms of the NR-Chr were extracted, one sedimenting at 50S-55S and the other, which had a more compact structure, sedimenting at 80S-85S. The

R-Chr sedimented between the two conformations of the Form I Chr. Electron microscopy of the R-Chr suggested they replicated in a conformation more closely related to the 100-A, 55S nucleofilament structure than to the 80S-85S one. Nucleosomes were present on both sides of the replication forks. In the in vitro assays, all regions of the SV40 genome were synthesized, and a significant fraction of the R-Chr completed replication. The progeny Form I Chr cosedimented with the 50S-55S Chr, which were present before or to incubation. The sedimentation coefficients and relative amounts of the two forms of the mature Chr were unaffected by the incubation. (18 refs)

- 78-5798 Human Diploid Lung Fibroblast Cell Lines WI 26 and WI 38 Exhibit Isozyme Shift of Alkaline Phosphatase after Viral Transformation.** (Eng) Knaack, J. (Institut für Organische Chemie, Biochemie und Isotopentechnik, Universität Stuttgart, Pfaffenwaldring 55, Stuttgart 80, W. Germany); Pfeleiderer, G.; Bayreuther, H. *Clin Chim Acta* 88(2): 375-383; 1978.

Immunotitration curves of membrane-associated alkaline phosphatase (AP) revealed that there is a shift from a fetal type of AP cultures of the normal human diploid fibroblast strains WI 26 and WI 38 to placental-like variant after transformation of these cells by simian virus 40. This was confirmed by comparing the apparent Michaelis constant and heat stability of the AP from the normal and transformed cells. Hepatic AP was heat labile and had a K_m for p-nitrophenylphosphate of about 4.0×10^{-4} M. The placental-like AP was heat stable and had a K_m of about 2.2×10^{-4} M. Transformed cells contained the placental-like form of AP exclusively, which is indicative of an all-or-nothing change in the occurrence of the placental-like enzyme is similar to the appearance of the placental-like Regan enzyme of AP in human tumors in vivo. The change detected in the transformed cells is most likely caused by genetic alteration of the cells induced by the transforming virus. (32 refs)

- 78-5799 Colony Formation by Simian Virus 40 Transformed Human Parapharyngeal Cells on Semisolid Agar.** (Eng) Ruben, R. L. (Dept. Anatomical Basic Medical Sciences, Univ. Illinois Medical Center, 600 W. Polk St., Chicago, IL, 60680); Rafferty, K. A. *Cancer Inst* 61(4): 993-1000; 1978.

A human epithelioid cell line (parapharyngeal cells) transformed by simian virus 40 (SV40) was tested for its ability to grow in the absence of an anchoring substrate and for phenotypic changes occurring in sublines isolated by anchorage independence. Line SV-TGo was plated on agar medium, and resulting colonies were tested for efficiency in liquid and for virus productivity. After population doublings, the average cloning efficiency of the cells was 12.3% in liquid and 2.2% in agar, vs 0.002% in

negative controls (F2408 rat cells). Agar colonies appeared as spherical aggregates of tightly packed cells (<0.3 mm), and they were well-dispersed. Twenty-seven sublines isolated for transfer to Leighton tubes were identical to the parental line, including epithelioid morphology, organoid growth pattern, and extreme sensitivity to density-dependent inhibition of growth. Seven colonies tested were positive for production of infectious SV40 in 14 days. Most sublines grew vigorously and required subculturing in the 19 days. Senescence was noted after 39-44 total population doublings. The ability of this cell line to grow unaided in agar without phenotype change was therefore demonstrated. (36 refs)

78-5800 Cell-Free Synthesis of Simian Virus 40 T-Antigens. (Eng) Paucha, E. (Translation Lab., Imperial Cancer Res. Fund, Lincoln's Inn Fields, London WC2A 3PX, England); Harvey, R.; Smith, A. E. *J Virol* 28(1): 154-170; 1978.

In vitro synthesis of two forms of simian virus 40 (SV40) T-antigen (large T and small t) in response to messenger RNA (mRNA) isolated from infected CV1 cells was demonstrated by polyacrylamide gel electrophoresis and tryptic peptide fingerprint analysis of the proteins produced in a cell-free system derived from L cells and immunoprecipitated with SV40 anti-T serum. The mRNA's for both antigens were sedimented at about 17.5S, with the mRNA for small t sedimenting marginally, but reproducibly, ahead of that for large T. Hybridization experiments using *Hae*III fragment and *Hind*II/II fragment B showed that all fractions active in the cell-free synthesis of the two antigens hybridized equally to both fragments. This suggests that the mRNA's for both T antigens are at least partly virus-coded and that most of the early SV40 mRNA contains sequence information for both ends of the early region. The data suggest that the early T mRNA is spliced. SV40 complementary RNA (cRNA) was also translated in the L-cell system, and it gave rise to families of polypeptides that specifically immunoprecipitated with anti-T serum. The small-t family included polypeptides indistinguishable by gel electrophoresis and fingerprinting from small t isolated from cells. The 60K family was a major component of 60,000-mol wt and included other polypeptides with mol wts of 14,000-70,000. The 60K family polypeptides in common with large T but not small t. The peptides of both families accounted for all the methionine peptides of SV40 large T. It is concluded that small t is probably entirely and large T at least predominantly virus-coded, and the small t and 60K families represent the translation products of two different portions of the early region of SV40 cRNA, and that although most, if not all, of the large-T and small-t peptides are present in the cell-free product, some rearrangement of sequence arrangement of SV40 cRNA prevents the isolation of full-length large T and results instead in the synthesis of fragments. The absence of a splice in cRNA is probably responsible for this result. (57 refs)

78-5801 Extraction and Fingerprint Analysis of Simian Virus 40 Large and Small T-Antigens. (Eng) Smith, A. E. (Translation Lab., Imperial Cancer Res. Fund, London WC2A 3PX, England); Smith, R.; Paucha, E. *J Virol* 28(1): 140-153; 1978.

The different forms of simian virus 40 (SV40) tumor antigen (T-ag) isolated from infected CV1 cells by a variety of extraction procedures were analyzed by peptide mapping. In addition, parameters involved in the generation of these T-ag forms were examined. Upon sodium dodecyl sulfate-polyacrylamide gel electrophoresis, the highest mol wt form of T-ag (large T) isolated from CV1 cells comigrated with the large T isolated from SV40-transformed H65-90B hamster cells. Other faster-migrating forms of large T were generated during the extraction by a protease that is active at low pH, but these forms are probably experimental artifacts. After extraction under conditions that minimize proteolytic degradation of large T, a further form of T-ag was isolated. It had a mol wt of 15,000-20,000 and was termed small t. Fingerprint analysis of ³⁵S-methionine-labeled SV40 proteins showed that small t has 10-12 methionine peptides, but large T has 15-18. All but two of the methionine tryptic peptides in small t were also present in large T. Fingerprint data also showed that T-ag's have no peptides in common with SV40 VP1. Reagents that inhibit posttranslational cleavage of encephalomyocarditis virus polypeptides did not affect the synthesis of small t, which suggests that it is not made by proteolytic cleavage of large T in vivo. An alternative model, which proposes that large T and small t are synthesized independently, is discussed in terms of the fingerprint data and the number of methionine tryptic peptides predicted from the primary sequence of SV40 DNA. (43 refs)

78-5802 Simian Virus 40 DNA Extracted from Infected Cells with Sodium Deoxycholate No Longer Reflects Its In Vivo Superhelix Density. (Eng) Germond, J. E. (Dept. Biochemistry, Stanford Univ. Medical Center, Stanford, CA, 94305). *J Virol* 26(3): 681-685; 1978.

The superhelix density of simian virus 40 (SV40) DNA was determined using viral DNA extracted from infected African green monkey kidney CV-1 cells with low concentrations (0.25%-0.45%) of sodium deoxycholate (DOC). The superhelix density was reduced compared with that of SV40 DNA extracted from infected cells in the presence of sodium dodecyl sulfate or from purified viral particles; the max reduction was at 0.35% DOC. No reduction was seen after extraction in the presence of 1% DOC. Even at low concentrations of DOC, no minor population of DNA was selected. Up to 0.1% DOC did not prevent the relaxation of SV40 DNA form I by the untwisting extract (UE). The number of nucleosomes of SV40 minichromosomes was reduced in the presence of low concentrations of DOC; histones were extracted from DNA under these conditions. This deproteinization released the nucleosomal constraint imposed on the DNA by the histones, which appeared as superhelical turns susceptible

to UE. When SV40 minichromosomes were incubated with UE in the presence or absence of DOC, the DNA remained fully supercoiled after deproteinization. It is concluded that in the presence of low concentrations of DOC, the nucleosomal structure of SV40 minichromosomes is first damaged by a combined effect of high ionic strength and anionic detergent, allowing the resulting complex to acquire superhelical turns that are then removed by an intranuclear untwisting activity. The DOC-induced changes in the superhelical state of the DNA can be mistaken for its true physiological configuration in the cells. (13 refs)

- 78-5803 Discontinuous DNA Replication: Accumulation of Simian Virus 40 DNA at Specific Stages in Its Replication.** (Eng) Tapper, D. P. (Dept. Biological Chemistry, Harvard Medical Sch., Boston, MA, 02115); DePamphilis, M. L. *J Mol Biol* 120(3): 401-422; 1978.

Agarose gel electrophoresis was used to show that replication of simian virus 40 (SV40) DNA molecules is a linear function of the extent of replication from molecules that replicated 15% of the genome to molecules that replicated 95%. This method allowed quantitation of the relative number of DNA molecules throughout replication. SV40 replicating DNA containing uniformly labeled nascent DNA chains was isolated from intact CV-1 cells at the peak of viral DNA synthesis. The DNA pool contained 2.0 ± 0.5 times more 60%-90% replicated molecules than 20%-50% replicated molecules. This asymmetrical distribution did not occur continuously throughout DNA replication, but began with 60%-70% replicated molecules. The late-replicating molecules were found predominantly in a major peak at 89% replication and two smaller peaks at 81% and 79% replication. These results were confirmed by experiments in which viral replicating DNA, briefly labeled with α - ^{32} P-deoxycytidine triphosphate (dCTP) in isolated nuclei, was allowed to continue replication in an excess of unlabeled dCTP. The initial distribution of SV40 replicating DNA was similar to that observed in intact cells. Upon continued incubation, 40%-60% of the molecules matured into circular viral DNA by the time DNA synthesis had ceased. Of the replicating DNA molecules that remained, < 10% had not progressed to 50% replication; > 70% of the molecules were at least 80% replicated. The latter molecules had accumulated in two prominent peaks at 89% and 80% replication; a third peak was sometimes seen at 77% replication. DNA was located by measuring the radioactivity in gel slices and by autoradiography. These data demonstrate discontinuities in the rate that DNA replication forks advance as they approach each other to terminate replication. Consequently, SV40 replicating DNA molecules accumulate in at least two or more discrete stages during replication. (50 refs)

- 78-5804 Sister-Chromatid Exchange in Virus Transformation (Meeting Abstract).** (Eng) Nichols, W. W. (Inst. Medical Res., Camden, NJ); Bradt, C.; Toji, L. H.; Segawa, M. *Mutat Res* 53(2): 241; 1978. (no refs)

- 78-5805 Selective DNA Conservation and Chromatin Assembly after Injection of SV40 DNA into *Xenopus* Oocytes.** (Eng) Wyllie, A. H. (MRC Lab. Molecular Biology, Hills Road, Cambridge, England); Laskey, J. A.; Finch, J.; Gurdon, J. B. *Dev Biol* 64(1): 178-188; 1978.

The fate of simian virus 40 (SV40) DNA injected into the nucleus or cytoplasm of *Xenopus laevis* oocytes is described. The cytoplasm degraded all DNA forms; the nuclei degraded single-stranded and linear double-stranded DNA, but conserved double-stranded circular DNA in a 55S nucleoprotein complex. The correlation between the efficient transcription of SV40 DNA within oocytes and the presence of the stable nucleoprotein complex is discussed. (20 refs)

- 78-5806 Nuclear Viral Transcripts in a Clone of Rat Cells Transformed by SV40.** (Rus) Chumakov, P. M. (Inst. Molecular Biology, Moscow, USSR); Lukatskiy, E. M.; Georgiev, G. P. *Dokl Akad Nauk SSSR* 239(4): 984-987; 1978.

The sequence of viral RNA transcription was studied in a culture of simian virus 40 (SV40)-transformed rat cells (SVRE-17, which contains a single intact copy of SV40 DNA). Incubation of highly labeled uncoiled chains of SV40 RNA fragments with unlabeled nuclear RNA fragments derived from transformed rat cells showed maximal hybridization with early DNA fragment (fragment A). (12 refs)

- 78-5807 Tumorigenicity and Its Suppression in Cybrids of Mouse and Chinese Hamster Cell Lines.** (Eng) Howell, A. N. (Dept. Molecular Genetics, Sloan-Kettering Cancer Inst., 44 Binney St., Boston, MA, 02115); Farber, R. *Proc Natl Acad Sci USA* 75(5): 2358-2362; 1978.

The effect of cytoplasm on the expression of tumorigenicity was examined with a pair of mouse and a pair of Chinese hamster cell lines in intraspecies cybrids formed by reciprocal fusions between either tumorigenic or nontumorigenic cells and cytoplasts derived from them. With the mouse cell line 3T3 and the simian virus 40-transformed line SVT2, cybrid clones were tumorigenic when SVT2 cells were fused with 3T3 cytoplasts, but not in the reciprocal fusion. With the hamster cells, CHEF/18 and the spontaneous transformant CHEF/16, tumorigenicity was partially suppressed in cybrid clones formed by fusion of tumorigenic CHEF/16 cells with CHEF/18 cytoplasts; cybrids were nontumorigenic in the reciprocal fusion. These results indicate that tumorigenicity is not cytoplasmically transmitted in these two cell pairs, but suppression of tumor-forming ability can be cytoplasmically transmitted in the hamster cybrids. The fact that the cybrids have maintained their suppressed phenotype for 20-30 doublings probably rules out the role of well-understood gene products, such as messenger RNA and proteins, as the basis of cytoplasmic suppression. (28 refs)

08 **Siderophore-like Growth Factor Synthesized by SV40-transformed Cells Adapted to Picolinic Stimulates DNA Synthesis in Cultured Cells.** (Eng) Andez-Pol, J. A. (Nuclear Medicine Lab., VA Hosp., St. MO, 63125). *FEBS Lett* 88(2): 345-348; 1978.

mechanism of action of picolinic acid, which induces arrest in cultured cells, was examined using mutant lines of simian virus 40-transformed BALB/3T3 cells adapted to grow in picolinic acid (SVT2Pi3). Medium conditioned by contact with SVT2Pi3 cells contained a highly specific iron-binding ligand. The material, siderophorelike growth factor (SGF), was also produced by SVT2Pi3-A cells, a line of SVT2Pi3 cells that multiplies in the absence of iron. SGF was characterized as an ionophorepeptide of approximately 1,600 mol wt. SGF stimulated both Fe^{3+} uptake and DNA synthesis in the mutant cells. Therefore, SGF may be a mammalian siderophore growth factor that plays an important role in the control of DNA synthesis in the mutant cell line. The results suggest that cellular or extracellular chelate-like SGF, may have a relevant function as components of the transformed cell micronutrient transport system for Fe and other trace metal ions. They also support the hypothesis that neoplastic cells produce chelating agents that may be important determinants of the uncontrolled growth and invasiveness of transformed cells. (16 refs)

809 **Detection of Mason-Pfizer Monkey Virus Infection by Syncytia Formation of Human Cells Doubly Transformed by Rous Sarcoma Virus and Simian Virus 40.** Brief Report. (Eng) Ogura, H. (Dept. Virology, Cancer Inst., Okayama Univ. Medical Sch., 2-5-1 Shikata-cho, Okayama 700, Japan); Tanaka, T.; Ocho, M.; Kuwata, T. *Arch Virol* 57(2): 195-198; 1978.

Human embryonic cells doubly transformed by Rous sarcoma virus (RSV) and simian virus 40 (SV40) can be used as indicator cells for the detection of Mason-Pfizer monkey virus (MPMV), as they form syncytia upon cocultivation with MPMV-producing cells. The cells, named RSB cells, contain both the group-specific antigen of RSV and the tumor antigen of SV40. The cell fusion was blocked by anti-MPMV serum, indicating that this phenomenon is MPMV-specific. RSB cells were successfully used for MPMV infectivity assays in the same manner as KC (human glioma) cells. Although RSB and KC cells carry the RSV genome, it is not known if this genome is indispensable for indicator activity. (10 refs)

810 **Temperature-sensitive Mutants of Simian Virus 40. I. Isolation and Preliminary Characterization of B/C Gene Mutants.** (Eng) Noonan, C. A. (Dept. Virology and Epidemiology, Baylor Coll. Medicine, Houston, TX, 77030); Butel, J. S. *Intervirology* 10(3): 181-195; 1978.

Sixteen temperature-sensitive mutants of simian virus 40 (SV40) were isolated after treatment of wild-type virus with UV light, nitrous acid, or N-methyl-N'-nitro-N-nitrosoguanidine. All of the mutants were assigned to the B complementation group on the basis of quantitative complementation analyses. Characterization of several representative mutants revealed that under restrictive conditions, they produced immunologically identifiable tumor and viral capsid antigens, induced the synthesis of cellular DNA, and replicated infectious viral DNA. In addition, mutant particles produced at 33°C were more heat-labile than wild-type virus. This collection of mutants will be useful in an analysis of SV40 particle morphogenesis. (41 refs)

78-5811 **Chromosome of the Mature Virion of Simian Virus 40 Contains H1 Histone.** (Eng) Nedospasov, S. A. (Inst. Molecular Biology, USSR Acad. Sciences, Vavilov Str. 32, Moscow 117312, USSR); Bakayev, V. V.; Georgiev, G. P. *Nucleic Acids Res* 5(8): 2847-2860; 1978.

Minichromosomes (small free chromosomes) and mature virions were isolated from nuclear extracts of green monkey kidney CV1 cells infected with simian virus 40 (SV40, strain 776), and their protein composition was analyzed by gel electrophoresis (EP). A conventional isolation procedure and a modified one that makes available large amounts of mature virion particles directly from nuclear extracts were used. In the latter, the isolated nuclei of labeled CV1 cells taken at different intervals after infection were extracted with 0.12 M NaCl, 10 mM Na-EDTA, and 10 mM triethanolamine-HCl at pH 8.0 and ultracentrifuged in a sucrose density gradient. The centrifugation time was reduced to 4 hr, which resulted in the appearance of a fast-sedimenting peak containing labeled DNA in addition to the approx 70S peak corresponding to the minichromosomes. Analysis of the protein composition of the virions by sodium dodecyl sulfate-polyacrylamide gel EP in 15% or 5% gels, by acetic acid-urea EP, and by two-dimensional EP revealed the presence of a complete set of histones, including H1. H1 histone was present in the same stoichiometric ratio to the other histones as that in host chromatin. Capsid proteins VP1, VP2, and VP3 were also detected. Virions isolated from nuclear extracts were indistinguishable from those isolated by the conventional method with respect to electron microscopic appearance, buoyant density, and infectivity. The minichromosomes contained all the histones but practically no viral proteins. It is concluded that the SV40 minichromosome is organized in a compact particle of approx 300 Å in diameter and that it is incorporated into the virion as a whole, without the loss of any histone fraction. (20 refs)

78-5812 **Differences in the Capacity of Simian Virus 40 (SV40) Tumor Antigens on Cells, Membranes and in Soluble Form to Induce Transplantation Immunity in Hamsters and Mice.** (Eng) Law, L. W. (Lab. Cell Biology,

NCI, Bethesda, MD, 20014); Takemoto, K. K.; Rogers, M. J.; Henriksen, O.; Ting, R. C. *Int J Cancer* 22(3): 315-322; 1978.

The induction of simian virus 40 (SV40) transplantation immunity was compared in BALB/c mice and Syrian hamsters with the use of intact cells, cell membranes and nuclei, and solubilized membranes from several species (mouse, hamster, human, and sheep). All transformed lines were tumor-antigen-positive. The immunosensitive mKSA line of BALB/c mice and the immunosensitive SV34 line of a Syrian hamster were used for tumor challenge. All materials, regardless of source and type of preparation, induced a strong and specific transplantation immunity in the mice, but only SV40 and viable, intact SV34 cells provided protection against tumor cell challenge in the hamsters. In another limited study, BK papovavirus-transformed hamster cells and purified cell membranes and JC papovavirus-transformed hamster cells were immunogenic by the tumor rejection assay in mice but not in hamsters. SV40 immunization did not protect the hamsters against hamster cells transformed by these two viruses. The findings are best explained on the basis that SV40-transformed hamster cells (at least the SV34 cells used here) express a unique tumor-specific transplantation antigen (designated TSTA₁), identified by its immunogenicity in the hamster, that is absent from the cells of all the other transformed lines assayed. However, these other transformed lines (ie, mKSA and the human and sheep lines) express another antigen (TSTA₂), identified by its immunogenicity in the mouse. Another distinguishing feature is the fact that TSTA₂ is relatively stable after cell disruption but TSTA₁ is relatively unstable. Therefore, the TSTA induced by SV40 is heterogeneous and comprises at least two antigenic specificities. Explanations of the postulated heterogeneity of TSTA at the molecular level are given. (26 refs)

78-5813 Epidermal Growth Factor: Relationship Between Receptor Regulation and Mitogenesis in 3T3 Cells. (Eng) Aharonov, A. (Dept. Biological Chemistry, UCLA Sch. Medicine, Los Angeles, CA, 90024); Pruss, R. M.; Herschman, H. R. *J Biol Chem* 253(11): 3970-3977; 1978.

The time course and concentration dependence of epidermal growth factor (EGF)-induced ³H-thymidine (³H-TdR) uptake and cell division were compared with those required for receptor binding and down regulation (DR) in Swiss 3T3 mouse cells and in simian virus 40 (SV40)-transformed 3T3 cells. Exposure of confluent nondividing 3T3 cells to 10 nM EGF at 37 C resulted in a 70%-85% reduction in the binding capacity for ¹²⁵I-EGF. This reduction was due to a decrease in the number of available EGF receptors per cell without any change in the affinity of the receptors for EGF (DR). DR was dependent on EGF concentration, temperature, time, and the physiological state of the cell. Receptor loss occurred at physiological EGF concentrations in the range mitogenic for 3T3 cells (0.1-10 nM). Max stimulation of ³H-TdR uptake or cell division occurred at 1 nM EGF, a concentration at

which only 20% of the EGF-receptor sites are occupied. DR is only 55% complete. Low EGF concentrations (1 nM) resulted in DR of unoccupied EGF receptors. DR of EGF receptors also occurred in SV40-transformed 3T3 cells. In contrast to confluent cells, dividing 3T3 cells rapidly replaced EGF receptors on the cell surface in the presence of EGF. When EGF was removed from the medium, the receptor was quickly replenished. The initial massive internalization and concomitant receptor DR do not appear to be sufficient for EGF mitogenesis. The major role of reduction of EGF receptors by DR may be to adjust the subsequent sensitivity to EGF. (28 refs)

78-5814 Molecular Diversity among Five Different Endogenous Primate Retroviruses. (Eng) Berman, M. L. (Lab. Viral Carcinogenesis, NCI, Bethesda, MD, 20014); Sherr, C. J.; Sen, A.; Todaro, G. J. *J Virol* 26: 300-313; 1978.

The endogenous primate viral structural proteins of C-type retroviruses isolated from baboons (M7), macaques (MAC-1), and owl monkeys (OMC-1) and of the D-type viruses isolated from langurs (PO-1-Lu) and squirrel monkeys (SMRV, M534) were compared biochemically and immunologically. Also compared were the polypeptides of the prototype primate D-type virus of Old World monkeys, simian-Pfizer monkey virus (MPMV). The RNA genomes of C- and D-type retroviruses were unrelated to one another by nucleic acid hybridization criteria, and they coded for unique viral structural proteins that could be distinguished by rose gel filtration and polyacrylamide gel electrophoresis under denaturing conditions. The major structural protein of each virus had a distinct primary structure, as determined by two-dimensional tryptic peptide analysis, and it was antigenically different from the others. The major virion proteins of the C-type viruses (pp15) also differed from those of D-type viruses (pp13-pp14). Of the three D-type viruses studied, PO-1-Lu and M534 were endogenous to their respective species, but MPMV, which is transmitted horizontally in rhesus monkeys, may have been derived from a virus endogenous to langurs or to a related species. MPMV and PO-1-Lu contained related RNA genomes and coded for antigenically related structural proteins. Although certain radioimmunoassays detected interspecies antigenic determinants common to the p30 or gp70 proteins of some of these viruses, no one assay detected all groups of endogenous primate retroviridae. Data are given that show that, unlike Old World C-type viruses, the three D-type viruses may have been derived from one set of primordial gene sequences. The genomes contain at least three distinct sets of retroviral genes: two coding for C-type and one coding for D-type virus proteins. (28 refs)

78-5815 Effect of Splenectomy on Herpesvirus Infection and L₁C Leukemia in Guinea Pigs. (Eng) Berman, M. L. (Lab. Viral Carcinogenesis, NCI, Bethesda, MD, 20014); Sherr, C. J.; Sen, A.; Todaro, G. J. *J Virol* 26: 300-313; 1978.

G. D. (Virology Lab., Veterans Admin. Hosp., West Haven, CT, 06516); Gonzalez-Serva, A.; Hastings, K.; Cheng, C. K. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 1977; 1978.

ensive hyperplasia, accompanied by high virus titers, was present in the spleen of guinea pig herpeslike virus (HLV)-infected Hartley guinea pigs 3 wk postinoculation thereafter. Removal of the spleen after GPHLV inoculation reduced the virus titer in the peripheral blood. In addition, 3/9 animals that received a suspension of L₂C leukemia showed evidence of leukemia in their lymph nodes. The mechanism that permitted development of the leukemia is unclear. (2 refs)

5816 *Herpesvirus ateles* Malignant Lymphoma in Owl Monkeys. (Eng) Hunt, R. D. (Harvard Medical School, New England Regional Primate Res. Center, 1 Pine Hill Road, Southborough, MA, 01572); Barahona, H.; Daniel, M. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 198-200; 1978.

Malignant lymphomas developed in four owl monkeys inoculated with *Herpesvirus ateles* (titers of $10^{4.5}$ or 10^5 /ml of infectious TCD₅₀). In addition, the virus recovered from these monkeys induced lymphomas when reinjected into three other monkeys. All the animals died or were killed in moribund condition 27-37 days postinoculation. Microscopically, the neoplastic infiltrates were confined to the kidney, spleen, lymph nodes, and bone marrow. The liver and other tissues were spared. Leukemia was not present. (8 refs)

5817 *Herpesvirus Papio* A Lymphotropic Herpesvirus of Baboons that is Related to Epstein-Barr Virus. (Eng) Falk, L. (Rush-Presbyterian-St. Luke's Medical Center, Chicago, IL); Deinhardt, F.; Nonoyama, M.; Schuchman, A.; Wolfe, L.; Bergholz, C.; Lapin, B.; Yakovleva, L.; Vukobratovic, V.; Henle, G.; Henle, W. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 201-204; 1978.

Primate lymphoblastoid cell lines (LCL) latently harboring

a herpesvirus (*Herpesvirus papio*) related to Epstein-Barr virus (EBV) were studied for viral antigen expression, degree of genome homology with EBV DNA (approx 40%), production of transforming virus, and immunologic characteristics. The LCL contained antigens cross-reacting with EBV but lacked detectable Epstein Barr nuclear antigen in anticomplement immunofluorescence tests. Marmoset lymphocytes were transformed in vitro by x-irradiated LCL cells, but not by cell-free supernatants, suggesting little if any production of transforming virus. (8 refs)

78-5818 **Analysis of dCMP Deaminase and CDP Reductase Levels in Hamster Cells Infected by Herpes Simplex Virus.** (Eng) Langelier, Y. (Institut du Cancer de Montreal, Montreal, Quebec, Canada H2L 4M1); Dechamps, M.; Buttin, G. *J Virol* 26(3): 547-553; 1978.

Deoxycytidine monophosphate (dCMP) deaminase, cytidine diphosphate (CDP) reductase, and thymidine kinase (TK) activities were measured in dCMP deaminase-deficient Chinese hamster cells (IA1) or Syrian hamster cells (BHK-21/C13) infected with herpes simplex virus type 1 (HSV-1) or type 2 (HSV-2). Cell cultures were infected at an input multiplicity of 20-50 plaque-forming units/cell. The two viruses enhanced TK and CDP reductase activities in both cell lines, but they did not induce dCMP deaminase activity in IA1 cells or increase the low basal enzyme activity in quiescent BHK-21/C13 cells. In the IA1 cells, TK levels increased 25 times after HSV-2 infection and 10 times after HSV-1 infection; CDP reductase levels, measured only in HSV-1-infected cells, increased 8 times. In the BHK-21/C13 cells, the two viruses resulted in a 25- to 45-fold increase in TK and a 25-fold increase in CDP reductase levels. In contrast to the host enzyme, the virus-induced CDP reductase activity of IA1 and BHK-21/C13 cells was insensitive to inhibition by deoxyATP and deoxythymidine triphosphate. This property should be useful in developing a selection system for HSV-induced CDP reductase activity. (38 refs)

78-5819 **A Host Range Difference Between Herpes Simplex Virus Types 1 and 2.** (Eng) Goldfarb, P. (Inst. Genetics, Univ. Glasgow, Church St., Glasgow G11 5JR, Scotland); Timbury, M. *J Gen Virol* 39(2): 351-355; 1978.

The infectivities of several strains of herpes simplex virus types 1 and 2 (HSV-1, -2) were measured by plating on Chinese hamster lung (Don line) and human fetal lung (HFL) fibroblasts. HSV-1 strain 17 syn had a 10^4 -fold lower infectivity on Don cells than on human cells; this low infectivity on Don cells was also seen with other HSV-1 strains. Although adsorption and penetration of the virus appeared normal, there was little or no virus eclipse, and viral thymidine kinase was not induced. For all HSV-2 strains tested, the difference in titer between HFL and Don cells was only 10- to 100-fold.

It thus appears that there is a significant difference between the infectivity of HSV-1 and HSV-2 on Don cells. It is not known if this difference is related to antigenic differences between the two serotypes. Four HSV-1 and HSV-2 intertypic recombinants were also tested for infectivity on HFL and Don cells. Three of the four produced differences in titer similar to those observed with HSV-1, but the fourth behaved like HSV-2. (26 refs)

78-5820 Anatomy of Herpes Simplex Virus (HSV) DNA. X. Mapping of Viral Genes by Analysis of Polypeptides and Functions Specified by HSV-1 x HSV-2 Recombinants. (Eng) Morse, L. S. (Marjorie B. Kovler Viral Oncology Labs., Univ. Chicago, Chicago, IL, 60637); Pereira, L.; Roizman, B.; Schaffer, P. A. *J Virol* 26(2): 389-410; 1978.

The segregation patterns of polypeptides specified by herpes simplex virus types 1 and 2 recombinants (HSV-1 x HSV-2) were determined and compared with the DNA sequences of the recombinants. HSV-1 and HSV-2 polypeptides with mol wts of 20,000 to 200,000 were enumerated and matched in infected HEp-2 cells. Infected-cell polypeptides (ICP) that underwent rapid posttranslational processing and ICP that varied intratypically in mol wt were identified. Autoradiographic studies of electrophoretically separated ICP revealed that they fell into three groups according to the kinetics of their synthesis. The synthesis of α ICP peaked 2-4 hr postinfection, that of β ICP between 5-7 hr, and that of γ ICP between 12-15 hr. The templates specifying 26 ICP and several viral functions were mapped by correlating their segregation patterns with their DNA sequences. The α ICP mapped at the termini of the L and S components of HSV DNA. Although α ICP mapped entirely within the reiterated sequence of the L component, the template for α ICP 4 appeared to lie only partly within the reiterated sequence of the S component. Cells infected with a recombinant that contains HSV-1 and HSV-2 sequences in the S component produced α ICP 4 of both HSV-1 and HSV-2. Templates specifying β and γ ICP's mapped in the L component and were randomly distributed. Thymidine kinase and resistance to phosphonoacetic acid mapped in the L component. The rapid inhibition of host protein synthesis characteristic of HSV-2 infections and syncytial plaque morphology were used to map these functions in the L component. Implications of the template arrangements in HSV DNA are discussed. (41 refs)

78-5821 A Nonlytic Transforming Mutant of Herpes Simplex Virus Type 2. (Eng) Howett, M. K. (Dept. Microbiology, Milton S. Hershey Medical Center, Pennsylvania State Univ. Coll. Medicine, Hershey, PA, 17033); Rapp, F. *J Med Virol* 2(1): 29-38; 1978.

A previously isolated small-plaque mutant (No. 69) of herpes simplex virus type 2 (HSV-2) strain 333 produces a high ratio

of noninfectious to infectious particles when grown at nonpermissive temperature (39 C) in hamster embryo fibroblasts (HEF). An attempt was made to take advantage of the defective nature of this mutant in order to develop a system in which cells are transformed by a herpesvirus that has been inactivated. The noninfectious particles were tested in a transformation assay that detects biochemical transformation by the HSV-specific thymidine kinase (TK) gene. After inactivation, virus particles grown at 39 C in HEF cells were able to transform TK- mouse cells (N clA cl10) infected with an estimated particle input of 2 of the noninfectious mutant had a transformation frequency of 5 x 10⁻⁶. In comparison, cells infected at an original multiplicity of infection of 2 with HSV-2 strain 333, irradiated to yield only noninfectious particles, had a transformation frequency of 8 x 10⁻⁶. Clones obtained after incubating mutant-infected mouse cells for 3 wk contained herpes-specific antigens, as measured by immunofluorescence. The mutant contained a DNA genome that had a density identical to the DNA of wild-type HSV-2. Virus and cell DNA synthesis after infection with the mutant was similar at both the permissive (33 C) and nonpermissive temperatures. Parental virus. Preliminary experiments in which virus grown at 39 C in HEF cells were injected into newborn mice indicate that the virus is not infectious; in addition, no tumors were produced. (30 refs)

78-5822 Antibodies to Herpes Simplex Virus Type 2 in the Husbands of Patients with Cervical Carcinoma. (Letter to Editor). (Eng) Kunkel, M. (Frauenklinik I, Universitätsklinik für Medizinische Mikrobiologie und Epidemiologie, Albert-Ludwigs-Universität, 25 Rostock, E. Germany); Ceder, H.; Gartner, L.; Hofmann, R.; Ratz, K. H.; Heide, H. *Lancet* 2(8089): 585-586; 1978.

Serum antibodies against the early antigen of herpes simplex virus type 2, as measured by immunofluorescence, were found in 43/100 cervical carcinoma patients, 5/50 healthy controls, 10/36 husbands of the patients, and 2/36 male controls. (4 refs)

78-5823 Identification of a Herpesvirus Isolate from Cytomegalovirus-transformed Human Cells. (Eng) Geder, L. (Dept. Microbiology, Milton S. Hershey Medical Sch., Pennsylvania State Univ. Coll. Medicine, Hershey, PA, 17033); Hyman, R. W.; Figueroa, M.; O'Connell, E.; Iltis, J. P.; Dawson, M. S.; Rapp, F. *J Virol* 27(3): 700-704; 1978.

Human embryo lung cells transformed by cytomegalovirus (CMV) and transplanted to athymic nude mice yielded a cytopathic virus, Hershey Medical Center virus (HMCV). Following prolonged in vitro passage of the tumor cell virus is a double-enveloped herpesvirus, is sensitive to UV light and is inhibited by iododeoxyuridine. No significant antitumor activity was observed. (10 refs)

relationship to herpes simplex virus (HSV) was detected using HSV-immune sera in neutralization and immunofluorescence tests, but indirect immunofluorescence tests revealed V-related antigenicity. Further immunological tests revealed that HMCV is antigenically indistinguishable from infectious bovine rhinotracheitis virus. Thus, HMCV appears to be an infectious bovine rhinotracheitis virus, which presumably appeared in the cell culture as a contaminant from calf serum. (27 refs)

5824 The Interaction of Hydrocortisone, Lymphoid Cells and Epstein-Barr Virus. (Eng) Leyritz, M. (Dept. Microbiology and Immunology, Ste-Justine Hosp., Montreal, P. Q., Canada); Joncas, J. H. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 209-211; 1978.

Twenty-one human lymphoid cell lines of various origin were treated with hydrocortisone (HC: 20 µg/ml) for 3 days and then examined for cell viability, percentage of Epstein-Barr virus (EBV) antigens, and interferon synthesis. Unlike cells derived from the peripheral blood of normal individuals and infectious mononucleosis patients, Burkitt's lymphoma cell lines were characterized by a decrease in cell viability, an increase in EBV early antigen-positive cells, and poor interferon response. The response of the lymphoma cells to HC was consistent with the reported in vivo sensitivity of lymphoid cells to the lethal effect of glucocorticoids. (7 refs)

5825 Studies on Epstein-Barr (EBV)-like Viruses of Old World Nonhuman Primates. (Eng) Rabin, H. (Viral Oncology Program, NCI Frederick Cancer Research Center, P.O. Box B, Frederick, MD, 21501); Neubauer, R. A.; Hopkins, R. F. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 205-208; 1978.

Human, orangutan, chimpanzee, gorilla, and baboon sera were tested for antibodies to virus-associated nuclear antigens in human, orangutan, and chimpanzee cells. Of the human sera, 18/26 stained the nuclei of human Raji (B) lymphoid cells and 2/18 also stained the nuclei of orangutan CP-81 and chimpanzee CH888 cells. Of the orangutan sera, 5/9 stained Raji cell nuclei, 3/5 stained CH888 nuclei, and 2/5 stained Raji cell nuclei. Thus, certain human and orangutan sera contain antibodies that cross-react with the nuclear antigens of these three species. Chimpanzee sera cross-reacted with oran-

gutan antigen but were negative on Raji cell nuclei. Gorilla and baboon sera were negative on all three lines. Baboon virus transformed lymphocytes from several Old World nonhuman primates, but orangutan virus and Epstein-Barr virus from a cotton-topped marmoset line transformed only gibbon lymphocytes. The cells transformed in vitro had the properties of B lymphocytes, and they are being evaluated for possible malignant potential. (10 refs)

78-5826 Suppression of In Vitro Epstein-Barr Virus Infection: A New Role for Adult Human T Lymphocytes. (Eng) Thorley-Lawson, D. (Sidney Farber Cancer Inst., Boston, MA, 02115); Chess, L.; Strominger, J. L. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 219-222; 1978.

The in vitro transformation of human lymphocytes from adult peripheral and fetal cord blood by Epstein-Barr virus (EBV) was studied. B cells from these sources showed similar levels of ³H thymidine incorporation at all times post-EBV infection and showed the same efficiency (100%) and time to outgrowth (13-14 days). However, whole adult peripheral blood lymphocytes required an average of 25 days to grow out. EBV had no effect on the immunoglobulin (Ig)-negative cell population. Reconstitution of lymphocytes to the in vivo ratio of 20:80 B:Ig-negative cells resulted in a marked suppression of B-cell proliferation that occurred early in infection. The suppressor cells were shown to be the TH₁-subpopulation of T cells. The data indicate that the TH₁-T-lymphocyte subpopulation from adults but not fetal cord can suppress the outgrowth of in vitro EBV-infected B lymphocytes. This could reflect in vivo mechanisms involved in recovery from infectious mononucleosis and in the in vivo suppression of EBV-infected lymphocytes harbored by seropositive individuals. (9 refs)

78-5827 Interaction of Epstein-Barr Virus (EBV) with EBV-Genome-negative, Receptor-positive Human Lymphoid Cell Lines (LCL) Derived from Acute Lymphoblastic Leukemia (ALL) and Lymphoma Patients. (Eng) Menezes, J. (Dept. Microbiology and Immunology, Pediatric Res. Center, Univ. Montreal, Ste-Justine Hosp., Montreal H3T 1C5, Canada); Patel, P.; Bourkas, A. E. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 216-218; 1978.

The interaction of Epstein-Barr virus (EBV) with

EBV-genome-negative, receptor-positive human lymphoid cell lines (LCL) derived from patients with acute lymphoblastic leukemia (ALL) or lymphoma was studied. The results indicated that an LCL may bear receptors for only one type of EBV strain and not for the other, suggesting the possible existence of two types of EBV receptors on at least some LCL. An LCL may also express receptors for EBV but not for complement (C'), and an LCL bearing receptors for EBV may prove nonpermissive to EBV infection. These observations relate to a unique and previously unknown type of interaction between EBV and EBV-receptor-positive, genome-negative human LCL derived from ALL and lymphoma patients. It is highly probable that such an interaction also occurs in vivo in man, and it appears to be independent of C' receptors. (10 refs)

- 78-5828 Early Events in Epstein-Barr Virus-induced Transformation of Human Lymphocytes, Studied by Simultaneous Observations of Immunofluorescence, Cellular Morphology and Autoradiography.** (Eng) Takada, K. (Dept. Virology, Cancer Inst., Hokkaido Univ. Sch. Medicine, Sapporo, Japan); Aya, T.; Osato, T. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 212-215; 1978.

A method for simultaneously observing Epstein-Barr virus (EBV) marker expression, cellular morphology, and autoradiography findings was used to study early events in the transformation of human B lymphocytes by EBV. In human umbilical cord blood B-cell-rich fractions exposed to EBV and then incubated with growth medium containing ³H-thymidine, synthesis of EBV-determined nuclear antigen (EBNA) occurred prior to the blastogenic response and to DNA synthesis. EBNA-positive cells first appeared 12 hr after infection, reaching a max of 17% at 24 hr, whereas blastogenesis became progressively evident after 18 hr and DNA synthesis was noted at 36 hr. In the presence of 10 to 20 µg/ml cytosine arabinoside, EBNA synthesis was observed in 6% of cells in culture 24 hr after EBV infection, at which time EBNA-positive cells were seen in 10% of control infected cultures. Exposure of cells to cycloheximide (10 to 20 µg/ml) for the first 6 hr after infection inhibited EBNA synthesis to one-third of the rate seen in control cultures. Cycloheximide also inhibited morphological transformation. The results indicate that the early events in EBV-induced transformation of human lymphocytes occur sequentially from EBNA synthesis, to blastogenesis, to DNA synthesis. The crucial step in EBNA synthesis and EBV-induced morphological transformation is probably related to certain protein syntheses occurring in the first few hours after EBV infection. (5 refs)

- 78-5829 The Epstein-Barr Virus (EBV) in Human Pathology. II. Serologic Profiles of EBV Infections.** (Eng) Didier, J. (Laboratoire de Bacteriologie-Virologie, Universite Paul Sabatier, Faculte de Medecine Toulouse Rangueil, Chemin du Vallon, 31054 Toulouse Cedex, France); Dalens, M.; Chabanon, G.; Icart, J.; Enjalbert, L. *Immunomedicine* 28(1): 54-62; 1978.

Sera from 245 patients, including 139 with infectious mononucleosis (IM), were tested for antibodies to Epstein-Barr virus (EBV) capsid antigen (VCA), early antigen (EA), nuclear antigen, and VCA-IgM antibodies. In IM, efficiency of EBV serodiagnosis was lower than that of Paul Bunnell Davidsohn (PBD) reaction. Of the PBD-positive IM patients, only 80% had a serologic profile indicative of primary EBV infection. However, EBV serodiagnosis was useful in PBD-negative IM and in benign and malignant lymphoproliferative diseases. Of the 13 patients with lymphoproliferative malignancies, almost all were positive for EBV antibodies. In successive sera, these antibodies were frequently associated with VCA-IgM antibodies. (41 refs)

- 78-5830 Presence of EBNA in Nasopharyngeal Carcinoma and Control Patient Tissues Related to EBV Serology.** (Eng) Huang, D. P. (Medical and Health Dept., Inst. Radiology and Oncology, Queen Elizabeth Hosp., Kowloon, Hong Kong); Ho, H. C.; Henle, W.; Hsu, G.; Saw, D.; Lui, M. *Int J Cancer* 22(3): 266-274; 1978.

Biopsy specimens from 26 nasopharyngeal carcinoma (NPC) patients, 39 patients with other tumors of the head and neck, and the normal nasopharyngeal mucosa of 32 patients in whom NPC had been excluded, and the mucosa from the enlarged but nonneoplastic tonsils of 12 patients were analyzed for Epstein-Barr virus (EBV)-associated nuclear antigen (EBNA) by anticomplement immunofluorescence. Blood samples were also taken from the first three patient groups at the time of biopsy, which was prior to radiation therapy. EBNA-positive cells were detected in preparations from undifferentiated and poorly differentiated squamous cell carcinoma provided they contained tumor cell aggregates. These were not detected in two undifferentiated carcinoma preparations without aggregates and in the only case of well-differentiated papillary adenocarcinoma. In the head and neck tumor group, positive cells were found in only three undifferentiated carcinomas of the nasal fossa. None of the nasopharyngeal or tonsillar mucosa preparations contained positive EBNA. These results confirm the unique association of EBNA with undifferentiated and poorly differentiated squamous cell NPC. In EBV-related serological studies, there was a correlation between titers of IgG and IgA antibodies to capsid antigen and to the diffuse component of early antigen in patients with undifferentiated and poorly differentiated squamous cell NPC. Therefore, this simple blood test may be useful in screening for NPC in people with a high risk of this neoplasm. (33 refs)

831 Isolation of Epstein-Barr Virus from Cultures of Lymphoblastoid Cells. (Rus) Zhdanov, V. M. (Central Inst. Hematology and Blood Transfusion, Moscow, USSR); Skorikova, A. S.; Skurkovich, S. V.; Guschin, B. V.; Novikova, U. A.; Bykovskii, A. F.; Al'shtein, A. D.; Mazurenko, N. P. *Probl Gematol Pereliv Krovi* 23(6): 49-55; 1978.

An attempt was made to isolate both Epstein-Barr virus (EBV) and oncornavirus from the continuous lymphoblastoid cell line L-101 (derived from a patient with acute leukemia). The presence of EBV in the culture was confirmed by electron microscopy and by fluorescence assay: treatment of L-101 cells with serum containing antibodies against EBV resulted in specific fluorescence that was absent when the cells were treated with normal human sera. Density gradient centrifugation in sucrose gradient revealed a peak of labeled EBV at 1.24-1.26 g/ml (characteristic buoyant density of EBV). Although sedimentation in the sucrose gradient showed the presence of ³H-uridine-labeled RNA with a buoyant density of 1.15 g/ml, data on the persistence of oncornavirus in L-101 cells were nonconclusive. (8 refs)

832 Athymic Nude Mice: Induction of Tumors Containing Epstein-Barr Virus Using Burkitt's-related Cell Lines. (Eng) Ablashi, D. V. (NCI, Bethesda, Landow Building, Room 9A06, Bethesda, MD, 20014); Serlin, R.; Easton, J. M.; Nonoyama, M.; Armstrong, G. R. *J Hematol* 6(4): 365-374; 1978.

Cell development was studied in athymic nude mice inoculated with D98/HR-1 cells (1.0×10^6 cells), an epithelial cell hybrid containing the Epstein-Barr virus (EBV) genome, or with the parental D98 and HR-1 cells that were derived, respectively, from a human cervical carcinoma and Burkitt's lymphoma. The inoculum consisted of cells grown in primary culture, cells from tumors induced by the cell lines, or cells from tumor-derived cell lines. The D98/HR-1 cells induced differentiated carcinomas; the D98 cells, carcinomas; and HR-1 cells, poorly differentiated lymphomas. The number of EBV genome equivalents was sharply reduced in D98/HR-1 and HR-1 tumor cells, but subsequent in vitro passage of the cells resulted in partial recovery of the EBV genome in HR-1 cells and complete recovery in D98/HR-1 cells. Factors influencing EBV genome expression in tumors are discussed. It is concluded that the nude mouse is a useful model for studying EBV-associated tumors. (30 refs)

833 Demonstration of a Surface Antigen on Epstein-Barr Virus Genome-carrying Lymphoid Cells: Distinction from the Virus-determined Membrane Antigen. (Eng) Sakamoto, K. (Dept. Microbiology, Kumamoto Univ. Medical Sch., Kumamoto, 860, Japan); Hinuma, Y. *J Immunol* 120(5): 1702-1708; 1978.

Surface antigen (SA) was detected on Epstein-Barr virus

(EBV)-carrying lymphoid cell lines by indirect membrane immunofluorescence with an antiserum from a rabbit immunized with Raji cells. The antiserum had been extensively absorbed with normal human blood and tonsil cells. The SA was not detected on normal human umbilical cord and adult peripheral blood lymphocytes or EBV-negative cell lines. SA is distinct from EBV-determined membrane antigen (MA) as shown by the fact that five cell lines [Raji, NC-37 (C-6), OAT, P3HR-1 (NP), and MAT] that were consistently positive for SA by staining with anti-Raji serum were not stained by anti-MA human serum. Furthermore, HeHR-1 (P) and NHAd-60 lines, known to be positive for MA, were negative for SA. SA was also distinguished from MA in differential absorption tests. The results of cross-absorption tests of antiserum against Raji cells or P3HR-1 cells suggested that SA may contain more than one antigenic determinant. It is suggested that SA is a virally induced but not virally coded antigen. (35 refs)

78-5834 Interaction of the B95-8 and P3HR-1 Substrains of Epstein-Barr Virus (EBV) with Peripheral Human Lymphocytes. (Eng) Steinitz, M. (Dept. Tumor Biology, Karolinska Institutet, S-104 01 Stockholm 60, Sweden); Bakacs, T.; Klein, G. *Int J Cancer* 22(3): 251-257; 1978.

The interaction of the B95-8 and P3HR-1 substrains of Epstein-Barr virus (EBV) with normal peripheral blood lymphocytes were explored. B95-8 virus has transforming properties, but P3HR-1 does not. P3HR-1 fails to induce Epstein-Barr nuclear antigen (EBNA) or to stimulate DNA synthesis in normal lymphocytes, although it uses the same receptor for attachment as B95-8. To distinguish between the simple absence of a transforming and DNA-stimulating function and an active inhibition of its host cell, the effect of P3HR-1 pretreatment on subsequent DNA stimulation by B95-8 and by a number of mitogens was studied. P3HR-1 pretreatment completely inhibited the DNA stimulating effect and EBNA induction of subsequently added B95-8. When P3HR-1 was added after B95-8, it diminished DNA stimulation by the latter in a time-dependent fashion. Stimulation of DNA synthesis by the B-cell mitogen *Staphylococcus aureus* was also inhibited by P3HR-1. In contrast, the largely T-cell-stimulating effects of phytohemagglutinin and of an allogeneic mixed lymphocyte culture reaction were uninhibited or only slightly inhibited, which excludes the possibility that P3HR-1 had a nonspecific cytotoxic effect. These results suggest that P3HR-1 inhibits the same lymphocyte fraction that is stimulated by B95-8 and the B-cell mitogen. The effect of P3HR-1 is probably akin to its previously demonstrated growth-inhibitory effect on EBV-receptor-positive lymphoid lines. Early antigen induction is another unique property of P3HR-1, not shown by any transforming virus strain. It is possible that this property is related to the equally unique inhibitory effect of P3HR-1 on host cell DNA synthesis. (33 refs)

- 78-5835 Primary Epstein-Barr Virus Infections in African Infants. I. Decline of Maternal Antibodies and Time of Infection.** (Eng) Biggar, R. J. (Dept. Child Health, Univ. Ghana Medical Sch., Accra, Ghana); Henle, W.; Fleisher, G.; Bocker, J.; Lennette, E. T.; Henle, G. *Int J Cancer* 22(3): 239-243; 1978.

Thirty-one newborns in Accra, Ghana, an area endemic for Burkitt's lymphoma, were examined at monthly intervals for the first 15 mo and again at 21 mo to provide basic epidemiologic, clinical, and serologic data on primary Epstein-Barr virus (EBV) infections at an early age. Sera were tested for antibodies to EBV capsid antigen (VCA), EBV-induced early antigen (EA), an EBV-associated nuclear antigen (EBNA) by immunofluorescence techniques and for EBV-neutralizing antibodies and antibody-dependent cell-mediated cytotoxicity. All infants had maternal IgG antibodies to VCA at 1 mo of age and most also had antibodies detected by the other tests, except anti-EA. The anti-VCA titers subsequently declined with a half-life of nearly 5 wk. Depending on initial titers, all infants became seronegative for all antibodies by 2-8 mo, with anti-VCA persisting longest. No seroconversions were noted as long as antibodies to VCA remained measurable or for at least 2 mo after the infants became seronegative. Apparently, maternal antibodies protect infants against primary EBV infections during the first several mo of life. By 12 mo, 44% of the infants had seroconverted; these values increased to 62% and 81% at 15 and 21 mo, respectively. There was no seasonal variation in the incidence of infections. Infants from families of the lowest socio-economic level tended to become infected somewhat earlier than those from higher levels, but no other factors, such as household size or crowding, seemed to be associated with enhanced risk of early infection. Although the route of transmission remains obscure, salivary contact is the most likely source of infection. (24 refs)

- 78-5836 Primary Epstein-Barr Virus Infections in African Infants. II. Clinical and Serological Observations During Seroconversion.** (Eng) Biggar, R. J. (Dept. Child Health, Univ. Ghana Medical Sch., Accra, Ghana); Henle, G.; Bocker, J.; Lennette, E. T.; Fleisher, G.; Henle, W. *Int J Cancer* 22(3): 244-250; 1978.

Twenty-six newborns in Accra, Ghana, an area endemic for Burkitt's lymphoma, were examined monthly for 15 mo and again at 21 mo to record clinical, hematologic, and serologic responses during seroconversion (SC). Twelve infants acquired primary Epstein-Barr virus infections (EBV) by 1 yr of age and 9 others seroconverted between 13 and 21 mo of age. The infections were not accompanied by significant signs, symptoms, or hematologic changes that would have suggested a diagnosis of infectious mononucleosis (IM). Only 6/12 seroconverters showed late, minor, or insignificant heterophil antibody (ab) responses. Thus, the primary EBV infections were essentially silent. The 12 early seroconverters showed transient IgM ab responses to EB viral capsid antigen

(VCA) that, in some cases, initially exceeded the corresponding IgG ab's in titer. Peak VCA-specific IgG titers were noted 1 mo after SC, and they were comparable to those seen in IM. Most infants developed ab's to the EBV-induced early antigen complex. However, these ab's were directed against the restricted component, as observed in Burkitt's lymphoma, and not against the diffuse component, as noted in IM. Low titers of VCA-specific IgA ab's emerged in five infants. Usually, EBV-neutralizing ab's were already present in the first anti-VCA-positive serum, but ab's to EBV-associated nuclear antigen and ab-dependent cell-mediated cytotoxicity became detectable only months after SC. Attempts to demonstrate EBV by exposure of cord blood lymphocytes to filtered throat swab specimens collected from 1 mo before to 3-5 mo after SC met with limited success, which suggests that there is a low degree of viral excretion during silent SC. Possible explanations for the differences among clinical, hematologic, and serologic responses to primary EBV infections in infancy and later in life are provided. (29 refs)

- 78-5837 Hepatitis B Virus DNA in Primary Hepatocellular Carcinoma Tissue.** (Eng) Summers, J. (F. Chase Center, Inst. Cancer Res., Philadelphia, PA, 19111); O'Connell, A.; Maupas, P.; Goudeau, A.; Coursaget, P.; Drucker, J. *J Med Virol* 2(3): 207-214; 1978.

Primary hepatocellular carcinomas (PHC) from patients in which evidence of past or current infection with hepatitis virus (HBV) existed and from patients in a geographical area in which epidemiological evidence indicates a correlation between HBV infection and PHC were examined for the presence of HBV DNA by nucleic acid hybridization. Specimens of tumor, cirrhotic, and metastatic tissue were taken at autopsy from four male Senegalese patients, three of whom were positive for hepatitis B surface antigen (HBsAg). Tumors from two of these three patients contained HBV DNA sequences at a level of 1-2 genomes per cell; specimens from the third patient contained small but detectable amounts of viral DNA (< 0.1 genomes/10 cells). All three specimens from the HBsAg-negative patients contained no detectable HBV DNA. This patient showed evidence of prior HBV infection by virtue of the presence of anti-HBs and anti-core antibody in the serum. The absence of HBV DNA sequences in his tumor tissue indicates that the persistence of viral genes in every cell is not required to maintain the oncogenic state of the tumor cells. However, there is no epidemiologic evidence that correlates the presence of anti-HBs with PHC in Senegal. (15 refs)

- 78-5838 Viral DNA Sequences and Gene Products in Hamster Cells Transformed by Adenovirus Type 2.** (Eng) Johansson, K. (Dept. Microbiology, Biomedical Center, Univ. Uppsala, Uppsala, Sweden); Persson, L.; Lewis, A. M.; Pettersson, U.; Tibbetts, C.; Philipson, T. *J Virol* 27(3): 628-639; 1978.

plementary strand-specific adenovirus DNA of full length or from endonuclease *Bam*HI fragments was used as probe to analyze five LSH hamster cell lines (Ad2HE1-5) transformed by UV-inactivated adenovirus type 2 (Ad2) for fraction of viral DNA sequences represented and the copy number of these sequences per diploid equivalent of cellular DNA. No cell line contained a complete copy of the viral genome, and the fraction present varied from 44% in line Ad2HE5 to 84% in line Ad2HE3. The number of viral DNA copies per diploid cell equivalent ranged from 1.8 in Ad2HE1 to 7.1 in Ad2HE4. Serological analyses indicated variation in the expression of early viral antigen in all five lines. In vivo labeling of cell extracts early after infection with ^{35} S-methionine followed by immunoprecipitation with specific antisera revealed the presence of virus-specific polypeptides with mol wt of 42,000-58,000 in all five lines. Several other early viral polypeptides were detected in some of the Ad2-transformed cells. Thus, in spite of the presence of large fractions of the viral genome in these lines, they demonstrated considerable variation in the number of polypeptides expressed. A heterogeneous class of polypeptides appears to be present in all the lines, and it may constitute at least part of the Ad2 tumor antigens. (57 refs)

78-5839 Transcription of Viral Genes by RNA Polymerase II in Nuclei Isolated from Adenovirus 2 Transformed Cells. (Eng) Bitter, G. A. (Dept. Biological Chemistry, Div. Biology and Biomedical Sciences, Washington Univ. Sch. Medicine, St. Louis, MO, 63110); Roeder, R. *Biochemistry* 17(11): 2198-2205; 1978.

An in vitro system was developed that is capable of detecting the synthesis of viral-specific RNA in isolated nuclei from adenovirus 2 (Ad2)-transformed cells containing several (3-6) integrated copies of only the left 14% of the adenovirus genome. The α -amanitin sensitivities of the endogenous RNA polymerases demonstrated that a class II enzyme transcribes the viral genes. The synthesis of viral RNA appeared to consist entirely of elongation of RNA chains that were initiated in vivo. Under optimal conditions, the transcription of the integrated viral genes was detected by hybridization with adenovirus DNA. In nuclei isolated from B1 cells, viral-specific RNA represented about 0.01% of the total RNA synthesized in vitro. In isolated nuclei, the synthesis was completely (>95%) inhibited by concentrations of α -amanitin (6 μ g/ml) that completely and selectively inhibit RNA polymerase II. These results demonstrate the function of an RNA polymerase II in the transcription of viral genes in cells transformed by DNA tumor viruses. The transcription of the viral genes was further analyzed by hybridization of RNA synthesized in isolated nuclei to the purified separated strands of Ad2 DNA. While the stable cytoplasmic viral RNA is complementary to only the *r* strand of the viral genome, both strands of the integrated viral genes are transcribed. (43 refs)

78-5840 Evidence for an Adenovirus Type 2-coded Early Glycoprotein. (Eng) Jeng, Y. H. (Inst. Molecular Virology, St. Louis Univ. Sch. Medicine, St. Louis, MO, 63110); Wold, W. S.; Green, M. *J Virol* 28(1): 314-323; 1978.

Strong evidence that the 20,000- to 21,000-mol wt (20/21 K) polypeptide (pp) induced by adenovirus type 2 (Ad2) in human KB cells is an AD2-coded early glycopolypeptide is presented. The pp could be labeled with ^{3} H-glucosamine, and it bound to concanavalin A (Con A)-Sephadex columns and was eluted with methyl- α -D-mannoside. When labeled with ^{35} S-methionine for short periods (30 min-3 hr), it appeared as a single, sharp band with an apparent mol wt of approx 21K. However, if labeled for long periods or if pulse-labeled and then chased, the pp appeared as multiple bands with apparent mol wts of 20-21K. Partial proteolysis data indicated that the pulse-labeled 21K and the pulse-chase-labeled 20K pp's were highly related, although not identical. Both pulse (30 min) and pulse (30 min)-chase (12 hr in complete minimal essential medium) forms of ^{35} S-labeled 20/21K pp bound to Con A-Sephadex and, therefore, were glycosylated. The 20/21K pp and a 44K pp, labeled with ^{35} S-methionine or ^{3} H-glucosamine in Ad2-infected human cells, were precipitated by rat antiserum against the Ad2-transformed rat cell line T2C4, but not by antisera against three other Ad2-transformed rat cell lines or by serum from nonimmune rats. The partial proteolysis patterns of the 20/21K and 44K pp's were indistinguishable, indicating that the two are highly related and suggesting that the latter might be a dimer of the former. The 20/21K pp was also induced in Ad2-early infected monkey and hamster cells, which implies that the 20/21 K pp is synthesized in Ad2-infected human, monkey, and hamster cells and in one, but not all, Ad2-transformed rat cells. (34 refs)

78-5841 Purification and Characterization of an Early Protein (E14K) from Adenovirus Type 2-infected Cells. (Eng) Persson, H. (Dept. Microbiology, Biomedical Center, S-751 23 Uppsala, Sweden); Oberg, B.; Philipson, L. *J Virol* 28(1): 119-139; 1978.

The purification and characterization of an adenovirus type 2 (Ad2) early protein with an apparent mol wt of 14,000 (E14K), as estimated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis, are described. The protein was obtained from infected HeLa cells. Purification involved subcellular fractionation of cytoplasmic extracts, precipitation by dialysis at low pH, and diethylaminoethylcellulose, phosphocellulose, and hydroxylapatite chromatography. The yield was approx 12 μ g/10⁹ cells. Two Ad2 DNA binding proteins with mol wts of 75,000 and 45,000 were also purified, but tryptic peptide analyses indicated that E14K is unrelated to these proteins. Purified E14K has a high content of basic amino acids and a sedimentation coefficient, in native form, of 5.5S, corresponding to a mol wt of around 95,000.

Pulse-chase experiments suggested that E14K is a primary translation product. Immunoprecipitation with a monospecific antiserum against E14K revealed that it is located exclusively in the cytoplasm of infected cells. Synthesis of E14K began at 2 hr postinfection, and it was max at 4-6 hr postinfection. Immunoprecipitation of cell extracts from four AD2-transformed hamster embryo cell lines revealed that only one expressed this protein. E24K synthesis was not evident in an adenovirus-simian virus 40 hybrid virus, which suggests that the E14K gene is deleted in the genome of this hybrid. (46 refs)

78-5842 Adenovirus Type 2 Assembly Analyzed by Reversible Cross-linking of Labile Intermediates.

(Eng) D'Halluin, J. C. (Unite de Recherches de Virologie no. 102, Institut National de la Sante et de la Recherche Medicale, Lille, France); Martin, G. R.; Torpier, G.; Boulanger, P. A. *J Virol* 26(2): 357-363; 1978.

Dimethyl-4,4'-dithiobisbutyrimidate dihydrochloride was used as a cleavable cross-linking reagent to maintain the structure of labile intermediates in adenovirus type 2 assembly so that these intermediates could be characterized. Sucrose gradient analysis of nuclear adenovirus particles revealed two size classes, with sedimentation rates of 750S and 600S. After reversible fixation with the diimido ester, the different classes were further separated on CsCl gradients and characterized with regard to their buoyant density, DNA content, and polypeptide composition. The 750S particles banded at 1.345 g/cm³ in CsCl, contained a DNA with a sedimentation coefficient of 34S in alkaline sucrose gradients, and had a polypeptide composition similar to that of young virions. The 600S population consisted of two types of particles with buoyant densities of 1.315 and 1.37 g/cm³. The former contained a DNA fragment of 7S-11S and lacked the core proteins V and VII; precursors P VI and P VIII and two nonvirion proteins with a mol wt of 50,000 (50K) and 39K daltons were found in their place. 34S DNA was present in the 1.37-g/cm³ particles, which also lacked core proteins V and VII as well as the 50K and 39K proteins. Pulse-chase labeling kinetics showed that radioactivity was first detected in the 1.315-g/cm³ particles, followed by the 1.37- and then the 1.345-g/cm³ particles. The release of both 50K and 39K proteins was required for entry of viral DNA. (15 refs)

78-5843 Temperature-sensitive Mutant of Adenovirus Type 2 Blocked in Virion Assembly: Accumulation of Light Intermediate Particles. (Eng) D'Halluin, J. C. (Unite de Recherches de Virologie no. 102, Institut National de la Sante et de la Recherche Medicale, Lille, France); Milleville, M.; Boulanger, P. A.; Martin, G. R. *J Virol* 26(2): 344-356; 1978.

The assembly of human type 2 adenovirus (Ad2) was studied by biochemical and electron microscopic analysis of different

classes of particles obtained with *tsl12*, a temperature-sensitive mutant defective in a late stage required for virus maturation. At restrictive temperatures (39.5 °C) this mutant cumulated light precursor particles that were able to mature into infectious virions upon shift of the temperature to 33 °C. A mild extraction procedure and a reversible fixation by cleavable diimido ester were used to isolate the labile intermediates in adenovirus assembly. These particles had a sedimentation coefficient of about 600S and a buoyant density of 1.315 g/cm³ in CsCl. The addition of cycloheximide (10 µg/ml) to a KB cell culture infected with this virus drastically reduced the assembly of the precursor particles. The particles contained a DNA fragment of 7S-11S and two nonvirion proteins with a mol wt of 50,000 (50K) and 39K, respectively. These particles resembled the light intermediate particles found in wild-type Ad2 in both composition and morphology. The wild-type particles were identified as precursors of heavy intermediates, preceding the young virions. Apparently, the 112 lesion is located at the exit of either the 50K and/or 39K proteins and at the encapsidation of viral DNA. (32 refs)

78-5844 Tumorigenicity of Adenovirus-5-transformed BHK Cell Lines with Various Transformation-associated Properties. (Eng) Meager, A. (Dept. Biological Sciences, Univ. Warwick, Coventry, England); Nairn, Hughes, R. C.; Garrett, A. J.; Reeson, D. E. *Br J Cancer* 37(4): 566-570; 1978.

The ability of variant BHK21-C13 cell lines that survive lytic infection with adenovirus 5 to induce tumors in hamsters and immunosuppressed mice was determined. Transformed cell lines differed in their properties: some had a fibroblastic morphology (A1/3 x D, A1/3 x D1), and (F1/3A4) was composed of small round cells early in passage and fibroblasts upon continued culture. Injection of up to 10⁶ normal BHK21-C13 cells into Syrian hamsters produced no tumors, but injection of each of the variants produced tumors, although at different rates. The tumors grew progressively to a palpable size usually within 1 mo. Early-pass F1/3A4 cells were less tumorigenic than late-pass F1/3A4 cells and less tumorigenic than the other variants. In immunosuppressed mice, normal cells produced tumors when ≥10⁵ cells were injected sc. The variant A1/3 x D was more tumorigenic, producing tumors with inocula of 10⁴ cells. Cells derived from tumors produced by lines A1/3 x D and A1/3 x D1 had the same morphology as the parent cells. Cells derived from tumors produced by F1/3A4, however, were predominantly fibroblastic. A1/3 x D- and A1/3 x D1-derived tumor cells resembled the parent cells in saturation density and ability to grow in agar suspension or 0.1% serum. A tumor line obtained by injection of A1/3 x D cells into immunosuppressed mice also had the morphology and growth characteristics of the parent cells. There was no correlation between expression of a 250,000-dalton protein or other growth-related properties and tumorigenicity in the variant lines. (8 refs)

5845 **Adenovirus-induced Inhibition of Cellular DNase.** (Eng) Nass, K. (Dept. Microbiology and Immunology, Neil Hellman Medical Res. Building, Albany Medical Coll., Union Univ., Albany, NY, 12208); Frenkel, D. *J Virol* 26(2): 540-543; 1978.

Progressive decrease in the level of cellular DNase activity and single-stranded DNA occurred during the productive infection of KB cells by adenovirus type 5 (Ad5). In contrast, level of DNA polymerase remained relatively constant throughout the infection. The decrease could be prevented by inhibiting protein synthesis with cycloheximide ($\mu\text{g/ml}$). The inhibition of DNase activity did not occur after infection by Ad5 *ts125*, a DNA-negative mutant that fails to induce the adenovirus-specific DNA-binding protein. In contrast, infection by AD5 *ts125*, a DNA-negative mutant that complements *ts125*, did not result in decreased levels of DNase. It is suggested that an adenovirus-coded DNA binding protein acts as an inhibitor of DNase by binding to the single-stranded DNA substrate and thus protects the viral replicative intermediates from degradation. (33 refs)

5846 **Adenovirus Type 12 Gene 401 Function in Transforming Infection.** (Eng) Ledinko, N. (Dept. Biology, Univ. Akron, Akron, OH, 44325). *J Virol* 26(3): 828-831; 1978.

The transforming ability of wild-type adenovirus 12 (Ad12) was compared with that of the temperature-sensitive *ts401* and *ts406* mutants using secondary cultures of hamster embryo brain (HEB) cells and a cloned rat 3Y1 cell line. The *ts401* mutant was defective in its ability to transform hamster and rat cells at the respective nonpermissive temperatures (41.5 and 40°C). The transformation efficiency of WT Ad12 at the restrictive temperature was 70% and 90%, respectively, of that at the permissive temperature. The reduced efficiency of transformation by *ts401* at the nonpermissive temperature was not due to cell death. At the permissive temperature (38.5°C), *ts401* transformed HEB cells with six times the efficiency of WT virus, whereas at 41.5°C, the mutant was defective for transformation compared with WT virus. The *ts406* mutant transformed cells with approximately the same efficiency as WT at both temperatures. The *ts401* mutant also had a higher transformation efficiency than WT at 38.5°C. With 3Y1 cells, there was essentially no difference in the clear fluorescent staining characteristics of WT- and mutant-infected cells at the permissive and restrictive temperatures. Both mutants also induced cellular DNA synthesis after infection of 3Y1 cells at 40°C; *ts406* was somewhat more efficient in this respect than WT or *ts401*. It is suggested that defects of temperature-sensitive mutants may contain a higher proportion of transformation-defective genomes than WT virus. (15 refs)

5847 **Isozymic Forms of Some Energy Metabolism Enzymes During Oncovirus-induced Cell Trans-**

formation. (Eng) Ageenko, A. I. (Lab. Virology, Herten Res. Inst. Oncology, Moscow, USSR); Vitorgan, E.; Gorozhanskaya, G. *Folia Biol (Praha)* 24(2): 128-135; 1978.

Changes in the isozyme spectrum and total activity of hexokinase (HK), lactate dehydrogenase (LDH), and malate dehydrogenase (MDH) that occur during viral carcinogenesis were studied using rat embryo fibroblasts (REF) infected with an oncogenic variant of human adenovirus type 12 (Ad12) or an infectious variant of adenovirus type 6 (Ad6), uninfected REF cultures in different stages of growth, and Ad12-induced hamster sarcoma A12 and rat reticulosarcoma 321-RRS cell cultures. When the REF cultures were infected with Ad6, the activity of HK in the nuclear fraction and cytoplasm was not affected, but those of LDH and MDH were, especially that of LDH-5, which was lost in the cytoplasm. The change appeared as early as the first few days of culture-virus interaction. When the REF cultures were infected with Ad12, however, there were significant changes in all three enzymes in both the nuclear fraction and the cytoplasm. These changes were much more pronounced than those induced by Ad6. It is concluded that both infectious and oncogenic viruses rearrange the isozyme spectrum of the energy metabolism enzymes in the nucleus and cytoplasm, but the latter viruses to a greater degree. Cross analysis revealed changes in HK, LDH, and MDH in the soluble nuclear fraction and cytoplasm of Ad12-infected REF cultures and sarcoma A12 and reticulosarcoma 321-RRS cultures. These changes may serve as objective signs of cell transformation. (16 refs)

78-5848 **Initiation and Termination Sites of Adenovirus 12 DNA Replication II. Analysis of Pulse-labeled Oligonucleotides Derived from 5' Termini in the DNA Strand.** (Eng) Ariga, H. (Dept. Tumor Virus Res., Inst. Medical Science, Univ. Tokyo, 4-6-1, Shirokane-dai, Minato-ku, Tokyo 108, Japan); Shimojo, H. *Virology* 85(1): 98-108; 1978.

Pulse-labeled oligonucleotides (ON) from the 5' termini of the adenovirus type 12 (Ad12) DNA strand were studied to determine whether the initiation site of Ad12 DNA replication is located close to or exactly at the 5' end of the strand. Ad12 DNA pulse-labeled at the initiation site with ^3H -thymidine was mixed with U- ^{32}P -labeled Ad12 DNA and successively digested with the appropriate endonuclease and exonuclease to leave only the 5'-end ON intact or to digest the 5'-end ON selectively. The ON were separated from the mononucleotides by gel filtration or diethylaminoethylcellulose-Sephadex column chromatography. A temperature-sensitive mutant of Ad12, *tsA275*, which is defective in the initiation of viral DNA replication at the restrictive temperature, was used in these experiments. The ^3H label was not associated with the 5'-end ON of Ad12 DNA when the pulse-labeled DNA was digested completely with spleen DNase II, whereas a significant amount of ^{32}P -labeled Ad12 DNA was mixed with the ^3H -DNA as a control. The results

suggest that DNA replication of Ad12 is initiated at a short distance from the 5' end of the strand. The exclusion patterns of the ON in gel filtration suggest the location of the initiation site to be a short distance (about 20 to 50 nucleotides) from the 5' end. However, the replication of Ad12 DNA probably terminates right at the 3' end of the strand. (18 refs)

- 78-5849 Analysis of Mammalian Leukemia and Sarcoma Virus gag Gene-coded Translational Products: Implications Regarding Sarcoma Virus Structure.** (Eng) Stephenson, J. R. (NCI, NIH, Bethesda, MD, 20014); Reynolds, R. K. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 168-170; 1978.

The gag gene of mammalian C-type RNA tumor viruses codes for a 65,000-mol wt protein that, upon posttranslational cleavage, gives rise to four virion structural proteins with a mol wt of 30,000 (p30), 15,000 (p15), 12,000 (p12), and 10,000 (p10). Analysis of the intermediate cleavage products isolated from NIH/3T3 cells infected with Rauscher murine leukemia virus (R-MuLV) mutants defective in gag gene precursor processing yielded the sequence p15-p12-p30-p10. The gag gene expression occurred between the 5' and the 3' ends of the viral genome. Independent conformation of the R-MuLV gag sequence was obtained by analysis of gag gene-coded proteins of cells nonproductively transformed by replication-defective mammalian sarcoma virus isolates. The possibility that sarcoma viruses arise by recombination between leukemia viruses and host genetic sequences is discussed. (9 refs)

- 78-5850 Radioimmunologic Characterization of RD-114 Reverse Transcriptase: Evolutionary Relatedness of Mammalian Type C Viral pol Gene Products.** (Eng) Krakower, J. M. (Lab. RNA Tumor Viruses, NCI, Bethesda, MD, 20014); Aaronson, S. A. *Virology* 86(1): 127-137; 1978.

The reverse transcriptase derived from the endogenous mammalian C-type virus RD-114 was purified and characterized by biochemical and immunologic techniques. Using radioimmunoprecipitation assays, specific antibodies to reverse transcriptase were detected in the sera of animals naturally or experimentally infected with mammalian C-type viruses. Competition radioimmunoassays using the purified enzyme as a probe detected type-specific antigenic determinants of RD-114 reverse transcriptase, as well as group-specific determinants shared with baboon endogenous virus, the virus to which it was most closely related immunologically. The pol

gene products of all mammalian C-type viruses were shown to share interspecies antigenic determinants not detected among functionally analogous reverse transcriptases of retroviruses. These findings, together with previous immunologic evidence, strongly favor a common progenitor in the evolution of mammalian C-type viruses. (46 refs)

- 78-5851 Mutagenic Action of Viruses in Drosophila (Meeting Abstract).** (Eng) Gershenson, S. (Molecular Biology and Genetics, Academy Sciences Ukrainian SSR, Kiev, USSR). *Mutat Res* 53(2): 191-192; 1978 (refs)

- 78-5852 Fine Structure of Viruses of Murine Myeloma MOPC-21 and MSPC-1.** (Jpn) Kimura, Y. (Inst. Applied Medicine, Wakayama Medical Coll., Wakayama 640, Japan). *Saibo* 10(4): 543-546; 1978

Both A- and C-type viral particles were detected in murine myelomas MOPC-21 and MSPC-1 by electron microscopy. The A-type particles were located in the Golgi apparatus of the MOPC-21 myelomas, there were > 20 A-type particles per tumor and budding was apparent. In the MSPC-1 myeloma approx 2-5 A-type particles/tumor were observed under electron microscope, and multiple budding was not found. There were only a few C-type particles in both tumor strains, and they occurred in the extracellular space. Budding was not seen. (7 refs)

- 78-5853 Cell-associated Infectivity in the Blood of C57Bl Mice Bearing a Radiation-induced Leukemia (RadVL/D).** (Eng) Baugnet-Mahieu, L. (Dept. Radiobiology, CEN/SCK, B 2400, Mol, Belgium); Baes, C.; Coudrier, R.; Maisin, J. R. *Biomedicine* 28(1): 28-35; 1978.

The leukemogenic activity of blood constituents was studied in C57Bl mice bearing a reticulosarcoma of the spleen and lymph nodes induced by a radiation leukemia virus (RadVL/D). Viral C particles could hardly be detected in plasma, and the infectious titer (ED₅₀ at 45 days) remained very low up to 63 days after inoculation. Attempts to purify and concentrate a viral fraction from plasma were unsuccessful. From 14 to 63 days post-inoculation, infectious titers of the whole blood, lymphocyte and RBC fractions were of the same order of magnitude as that of an extract of lymphoid tissues: 10²-10³. Because RBC were not involved in the evolution of the disease, care was taken to eliminate possible contaminants (1

cytes or adsorbed C particles) from the RBC suspensions. activity of the RBC remained unaltered after differential centrifugation on Ficoll/Urografine, sulfoethylcellulose-*Se-*dex chromatography, and trypsinization. Lysis by osmotic shock or immune hemolysis always resulted in partial or total loss of biological activity, indicating that to be infectious structure of the RBC must be preserved. As has been described for other types of oncornavirus-induced tumors, it appears that blood cells, including RBC and/or their precur-

sors, play a role in the transmission of RadLV/D-induced leukemia. (20 refs)

See also:

*(Rev.): 78-5415, 78-5440, 78-5441, 78-5442, 78-5443.

*(Chem.): 78-5535.

*(Immun.): 78-5864, 78-5865, 78-5866.

*(Path.): 78-5885, 78-5886, 78-5888.

*(Epid.-Biom.): 78-5963.

- 78-5854 Enhancement and Metastasis after Immunotherapy of Ovine Squamous-Cell Carcinoma.** (Eng) Jun, M. H. (Dept. Tropical Veterinary Science, James Cook Univ. of North Queensland, Townsville, Queensland 4811, Australia); Johnson, R. H.; Maguire, D. J.; Hopkins, P. S. *Br J Cancer* 38(3): 382-391; 1978.

Ovine squamous cell carcinoma (OSCC) extracts, prepared by different procedures, were inoculated im into 184 OSCC-bearing sheep. At a concentration of 0.5 mg, OSCC protein significantly enhanced tumor growth, and the enhancement increased directly with increasing concentrations of protein. Tumor growth enhancement was associated with increases in metastases in lymphatic vessels draining primary lesions, in local lymph nodes, and in lungs. In control sheep receiving 0.15 M saline-phenol or 3 M potassium chloride extracts containing 3 mg total protein from normal sheep skin, DNA-enriched extracts from human squamous cell carcinoma, or complete Freund's adjuvant alone, tumor growth did not differ significantly from that in inoculated OSCC-bearing sheep. Sheep inoculated with 3 M potassium chloride extract of ovine fetal skin containing 3 mg total protein showed significant tumor growth enhancement and metastasis. All groups inoculated with 0.5 mg or more of tumor protein showed significantly more enhancement of Stage II than of Stage III tumors; no differences in the growth rates of Stage II and III tumors were noted in any of the control groups. The mechanisms involved in the enhancement of OSCC by tumor extracts are not known. (18 refs)

- 78-5855 Kaposi's Sarcoma and Immunosuppressive Therapy: An Appraisal.** (Eng) Gange, R. W. (Dept. Histopathology, Inst. Dermatology, St. John's Hosp. Diseases Skin, Lisle St., Leicester Square, London WC2H 7BJ, England); Jones, E. W. *Clin Exp Dermatol* 3(2): 135-146; 1978.

Three male patients (aged, 26, 60, and 78 yr) who developed Kaposi's sarcoma during steroid therapy are compared to 16 others who developed the disease in association with other forms of powerful immunosuppressive treatment. The ages of the 19 patients at the onset of Kaposi's sarcoma were similar to the normal age for development of this tumor, but there was a greater proportion of women (7/19) than is generally observed. Treatments in most cases included steroids, the average duration of treatment being 35.2 mo. Seven of the 19 patients were renal transplant patients. Seven patients, 4 clinically tumor-free, are still alive after an average period of 25.6 mo since diagnosis. Postmortem examination revealed evidence of residual Kaposi's sarcoma in only 5/11 patients. Reducing or stopping immunosuppressive therapy was followed by tumor regression in 3/4 patients. Since spontaneous remission

of Kaposi's sarcoma is rare, it would appear to be more common in tumors arising as a result of immunosuppression, more likely, to be due to a reduction in therapy, thus eroding host resistance. The data support a lymphoid origin of the spindle cells of this tumor, which appears to be especially prone to arise as a result of immunosuppressive therapy (refs)

- 78-5856 Role of Intracisternal A-Particles and of RNA Molecules in Plasmacytoma-associated Immunodeficiency.** (Eng) Katzmann, J. (Section Hematology, Mayo Clinic/Foundation, Rochester, MN, 55905); Bhoopalani, N.; Heller, P.; Hwang, L. T.; Ostro, M.; Lippman, D.; Giacomoni, D. *Cancer Res* 38(8): 2555-2561; 1978.

The origin of the RNA fractions from murine plasmacytoma (PC's) known to have immunosuppressive activity, the relationship of these molecules to fractions that induce clonal surface immunoglobulins (SIg) were determined. Subcellular fractions of the PC's MOPC-300, MOPC-104E, and J558 were transferred sc into normal BALB/c mice. Mice significantly diminished immune response to sheep red blood cells in mice given injections of precipitate B. This fraction copurified with intracisternal A particles. The RNA extracted from precipitate B was also immunosuppressive. The activity was due to the RNA molecules that contained poly(A) and poly(U) acid. Sucrose gradient fractionation showed that the RNA populations had immunosuppressive activity: 18S and 40S-50S RNA. The latter population was a thermally stable aggregate of molecules that contained the 12S-18S RNA molecules. In a previous study with normal mouse peritoneal macrophages, PC-derived material with similar physicochemical characteristics had induced SIg with the PC idiotype, supporting the possibility that one of the mechanisms responsible for the development of immunological deficiency is conversion of the SIg of nonmalignant B cells. Apparent immunosuppression and SIg conversion are due to the RNA fractions and molecules. (27 refs)

- 78-5857 Correlation of Differences in Antigenic Specificity of Four 3-Methylcholanthrene-induced Tumors in Syngeneic Mice with the Susceptibility of Tumors to Immunosuppression by PS-K.** (Eng) Abe, S. (Faculty of Pharmaceutical Sciences, Univ. Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113, Japan); Yamazaki, M.; Mizuno, D. *Gann* 69: 223-228; 1978.

The growth inhibitory effects of the immunopotentiator PS-K on syngeneic 3-methylcholanthrene-induced tumors

DDD mice were compared with differences in antigenicity among the tumors. Mice were immunized by implantation of tumor fragments in the tail, after which the tail was amputated. Two weeks later, immunized and control mice were challenged with 1.0×10^6 cells from the same tumor. The ratios of the tumor diameters of the immunized to control mice indicated that the original MSC-1 and MCA-4 tumors had high antigenicity, MCA-DDD and MCS-5 had intermediate antigenicity, and MCS-3, MCS-7, and MCS-9 had low antigenicity. MCS-8, MCS-6, and MCS-10 may have high antigenicity. Immunization with MCS-1 or MCS-8 completely inhibited tumor growth following challenge, whereas immunization with MCS-9 did not inhibit tumor growth after challenge at any dose. MCS-1 and MCS-8 thus appeared to be highly antigenic and MCS-9 appeared to be weakly antigenic. Mice immunized with MCS-1 or MCS-8 were not resistant to challenge with MCA-DDD, indicating that the tumor resistance was tumor-specific. The effect of PS-K on the growth of these tumors was significantly correlated with their antigenicity, but there was no significant correlation between antigenicity and the effect of the mitotic poison, mercaptopurine, on tumor growth. The data suggest that antitumor activity of PS-K may be used as an indicator of antigenicity in an experimental system. (12 refs)

78-5858 Further Studies on the Tumor-specific Suppressor Cells Induced by Ultraviolet Radiation. (Eng) Fisher, M. S. (Cancer Biology Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501); Kripke, M. *J Immunol* 121(3): 1139-1144; 1978.

Evidence that suppressor lymphocytes are present in UV-irradiated mice is given, along with the characteristics and specificity of these cells. Specific pathogen-free mice of inbred strains C3H/HeN(MTV-)(C3H-) and C57BL/6N were UV-irradiated with sun lamps that delivered an average rate of 2.8 joules/m²/sec over the wavelength range 280-340 nm. Dorsal hair was removed 1 x/wk, and mice were irradiated for 1 hr, 3 x/wk, for a minimum of 3 mo. Lymph nodes and spleens from normal and UV-irradiated mice were removed, cell suspensions were prepared, and lethally irradiated mice were inoculated with 5 to 10 x 10⁷ lymphoid cells (LC's). Reconstitution of lethally x-irradiated mice with LC's from syngeneic UV-irradiated donors failed to confer protection against challenge with a syngeneic UV-induced tumor, whereas reconstitution with LC's from normal donors was protective. When LC's from UV-irradiated mice were mixed with normal LC's before transfer, the recipients were again unable to resist tumor challenge, and it was shown that LC's from UV-irradiated donors actively suppressed the response of normal LC's. Furthermore, the suppressor LC's were enriched after incubation on nylon wool columns, they were inactive after treatment with anti- θ serum and complement, and they were effective for at least 7 wk after transfer to lethally x-irradiated mice. Splenectomy of UV-irradiated mice before tumor challenge did not restore their antitumor activity. The UV-induced suppressor cells appeared to be

specific for syngeneic UV-induced tumors, since they did not suppress the rejection of an allogeneic UV-induced tumor or two chemically induced syngeneic tumors. (21 refs)

78-5859 Tumor Specific Fluorescent and Complement-Dependent Cytotoxic Antibodies in the Serum of Rats with Chemically Induced Brain Gliomas. (Eng) Stavrou, D. (Lehrstuhl für Allgemeine Pathologie und Neuropathologie, Institut für Tierpathologie der Ludwig-Maximilians-Universität München, Veterinarstrasse 13, D-8000 München 22, E. Germany); Anzil, A. P.; Elling, H. *Acta Neuropathol (Berl)* 43(1-2): 111-117; 1978.

The biologic activity of antibodies in the sera of Sprague-Dawley and Long-Evans rats with N-methyl-N-nitrosourea-induced gliomas was investigated using a fluorescent antibody staining method and a complement-dependent microcytotoxicity assay. Of the chemically induced tumors, a mixed glioma, an astrocytoma, and an oligodendroglioma were established in vitro, and the first two were successfully cloned. Specific antibodies in the sera of rats with primary brain tumors were determined using the cultured cells as targets. Fluorescent and cytotoxic antibodies were demonstrated in the sera of the mixed glioma- and astrocytoma-bearing animals. However, a variable proportion of the cloned cell lines showed no reaction with the corresponding sera, and cytotoxic antibodies had a lytic effect on the autologous glioma cells only in the presence of rabbit complement. The data suggest that some cells in chemically induced tumors have antigenic properties and others do not. (15 refs)

78-5860 Lymphocytes in Ataxia Telangiectasia: Implications of the DNA Repair Defect. (Eng) Lavin, M. F. (Dept. Biochemistry, Univ. Queensland, Brisbane 4067, Australia); Kidson, C. *Clin Immunol Immunopathol* 10(3): 325-329; 1978.

The DNA repair capacity of phytohemagglutinin (PHA)-stimulated lymphocytes from four patients with ataxia telangiectasia (AT) and 11 healthy individuals was investigated after the cells were exposed to ionizing radiation. At radiation doses up to 50 krads, there was a 6- to 10-fold reduction of repair synthesis in AT lymphocytes compared with control lymphocytes. The marked impairment of repair observed in PHA-stimulated lymphocytes from AT patients provides a suitable and rapid clinical laboratory assay for AT homozygotes. The high incidence of tumors in AT homozygotes could be related to both a generalized DNA repair defect per se and to the associated lymphocyte defect if the latter results in decreased immune surveillance. (16 refs)

78-5861 *Corynebacterium parvum* Enhances Colonic Cancer in Dimethylhydrazine-treated Rats.

(Eng) Cruse, J. P. (Surgical Unit, Univ. Coll. Hosp. Medical Sch., University St., London WC1E 6JJ, England); Lewin, M. R.; Clark, C. G. *Br J Cancer* 37(4): 639-643; 1978.

The effect of *Corynebacterium parvum*, a bacterial vaccine used as immunotherapy following surgery for colorectal cancer in humans, on dimethylhydrazine (DMH)-induced colon tumors in female Wistar rats was determined. The rats were divided into four groups: Group A received DMH 20 mg/kg/wk for 20 wk; Group B received DMH + *C. parvum* (4 doses, 3.5 mg ip) at weeks 13, 16, 19, and 22; Group C received only *C. parvum*; and Group D received saline (1 mg/kg/wk, sc). *C. parvum* was first administered 13 wk after the first DMH injection to provide a population of activated macrophages at a time when the colonic epithelial cells are known to show abnormal morphological changes. All Groups C and D animals were alive at 36 wk. Group B animals had a significantly increased mortality compared with Group A animals. All Groups A and B animals examined postmortem had malignant colonic tumors, but Group B showed earlier signs of tumor presentation, greater numbers of tumors per animal, and a greater incidence of metastases than Group A animals. The extent and incidence of metastases were striking, since metastases are generally rare in this model system. Furthermore, 4/9 Group B animals developed primary cancers at sites other than the intestine (ie, squamous cell carcinomas of ear canal). The mechanism by which *C. parvum* enhances cancer in this model system is not known. (33 refs)

78-5862 Lung Tumor-associated Derepressed Alloantigen Coded for by the K Region of the H-2 Major Histocompatibility Complex. (Eng) Gipson, T. G. (Div. Virology, Bureau Biologics, Dept. Health, Education, and Welfare, Food and Drug Admin., Bethesda, MD, 20014); Imamura, M.; Conliffe, M. A.; Martin, W. J. *J Exp Med* 147(5): 1363-1373; 1978.

The strain and tissue distribution of the lung tumor-associated alloantigen that induces radioresistant immunity against lung tumor 85 in C3Hf mice was studied. Lung, liver, and kidney tissues from strain A and C3H mice and spleen cells of (C3HF x A)_{F1} mice evoked radioresistant anti-lung tumor immunity in C3Hf mice; no tissue from C3Hf mice provided antitumor protection. C3Hf mice were also protected by preimmunization with 15-day C3H embryo tissue but not by C3Hf embryo tissue. Mice that shared the *H-2a* haplotype with strain A mice or the *H-2k* haplotype with C3H mice induced highly significant immunity in C3Hf mice; tissue from mice with other *H-2* haplotypes was not effective. The gene coding for the expression of the alloantigen was therefore linked to the major histocompatibility complex (MHC). The haplotypes common to all strains conferring antitumor immunity were *H-2Kk* and *H-2IAk*. Anti-lung tumor immunity was not induced by tissue from A.TL (*H-2Ks*) or B10.AQR (*H-2Kq*) mice, and immunity was not conferred in C3H mice by tissues from the strains that induced immuni-

ty in C3Hf mice. The finding of a tumor-associated antigen in the expression of an *H-2K* region-coded antigen is consistent with the concept that MHC-coded antigens may be targets for the immunological surveillance of tumors. (10 refs)

78-5863 An Association Between Actin and the Histocompatibility Antigen H-2. (Eng) G. L. (Medical Res. Council, Lab. Molecular Biology, Rd., Cambridge, England); Smith, M. J. *Nature* 273(5667): 274-278; 1978.

P815 cells, a murine mastocytoma, shed large amounts of plasma membrane-derived material (exfoliate), in which is the major protein. The exfoliate also contains the histocompatibility antigen H-2. The association between actin and H-2 is stable even in the presence of detergent, indicating that the association depends solely on protein-protein interactions. The close parallel in the rates and extent of shedding of H-2 and actin and the absence of actin in the exfoliate argue against the possibility that the two come together after the shedding. It is therefore assumed that the H-2-actin association exists in the cell before the exfoliation occurs and that studies on the isolated H-2-actin complex are likely to be relevant to the arrangement of these components in the intact cell. (35 refs)

78-5864 Common Antigens Found on Fetal Cells and on a Rat Transformed Adult Prostatic Tissue. Javadpour, N. (Surgery Branch, Building 10, Room 101, NCI, Bethesda, MD, 20014); Hyatt, C. L.; Barone, J. *Surg Oncol* 10(3): 245-250; 1978.

A study was conducted to examine the immunologic similarity of hamster fetal antigens to antigens expressed on SV40 (SV40)-transformed hamster prostatic tissue and to assess the protection conferred on the adult host by fetal antigen immunization against growth of the virally induced transplantable prostatic tumor. Immunization with fetal antigens significantly decreased the incidence of tumor growth from 60% to 0%. Complement-dependent humoral cytotoxicity against SV40 tumor cells in microtest plates was used to demonstrate the presence of significant antibody titer in anti-fetal-antigen immunized hamsters. When the immune sera were adsorbed with SV40-transformed prostatic tissue, cytotoxicity indices were markedly lower than those of the unadsorbed immune sera. The transformed prostatic tissue had an elevated tartrate-inhibited acid phosphatase activity indicating the presence of epithelial components in the transformed tissue. These results demonstrate in vivo and in vitro cross-reactivity between fetal antigens and tumor associated antigens of transformed prostatic tissue, and they suggest that there is a similarity between fetal antigens and antigens of a prostatic organ tissue (prostate) transformed by a DNA onco-virus. (11 refs)

5865 Human Leukemia Associated Antigens (HLAA) on Cells from an AML Patient (HL23) and on Tissue Culture Cells Infected with HL23 Virus (HL23V) and Murine Oncornaviruses. (Eng) Metzgar, R. S. (Dept. Microbiology and Immunology, Duke Univ. Medical Center, Durham, NC, 27710); Mohanakumar, T.; Gallagher, R. E. *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 371-374; 1977.

Cultured and uncultured cells from a patient (HL23) with acute myelogenous leukemia (AML) were tested for their reactivity with cytotoxic primate antisera to human leukemia-associated antigens (HLAA). The cells did not react with antisera to chronic lymphocytic leukemia (CLL) or acute lymphoblastic leukemia, but they did react with antisera to AML. Two tissue culture cell lines (CCL 88, derived from human lung, and A204, derived from a human rhabdomyosarcoma) were nonreactive with the antisera until they were infected with secondary virus isolates (HL23V) from patient HL23. The antigenic reactivity of the secondarily infected cells differed from that of the cultured and uncultured cells from patient HL23. A204 and CCL 88 cells infected with bovine endogenous virus (BaEV) did not react with simian antisera to CLL but did react with antisera to myeloid cells. The reaction pattern of the BaEV-infected cells was similar to that seen after HL23V infection; however, it differed markedly from that of A204 and CCL 88 cells infected with human sarcoma virus or gibbon ape lymphoma virus. The use of secondarily infected cell lines offers a new approach to studying the nature of the cross-reactivity between simian antisera and HLAA. (8 refs)

78-5866 Expression of Oncofetal Antigens on Murine Sarcomas Characterized for Expression of Endogenous MuLV. (Eng) Pollack, S. B. (Fred Hutchinson Cancer Res. Center, 1124 Columbia St., Seattle, WA, 98104). *Int J Cancer* 22(3): 344-350; 1978.

Methylcholanthrene (3-MC)-induced BALB/c mouse fibrosarcomas previously characterized for the expression of endogenous murine leukemia virus (MuLV) were tested for the presence of oncofetal antigens (OFA) using rabbit antiserum to day 10-14 BALB/c fetuses. Iodinated protein A from *Staphylococcus aureus* was used to quantitate binding of the antiserum to cultured tumor or fetal cells or to saline extracts of tumors and fetuses. After absorption in vitro with normal adult mouse tissues and in vivo in adult mice, the antiserum bound to cells or extracts of fetuses and sarcomas to varying degrees but not to endogenous MuLV (BALB virus 1). This indicates that OFA on 3-MC-induced sarcomas exist as serologically recognizable antigens distinct from MuLV antigens. The anti-OFA activity of the serum could be absorbed with fetal cells and could be competitively inhibited by fetal extract but not by an extract of adult BALB/c lungs or by bovine serum albumin. The serum bound to some but not all lines known to express MuLV. The antiserum bound equally well to MuLV-negative and MuLV-positive sublines of the 3-MC-induced sarcomas 1420 and 1414 but not to Moloney sarcoma cells and the 3-MC-induced sarcoma 1386. Thus, the absorbed antiserum detects a class of common cross-reacting antigens that are serologically distinct from MuLV-associated antigens. (19 refs)

78-5867 Effect of Embryonic Tissue Immunization on Chemically Induced Gastrointestinal Tumors in Rats. I. Can Embryonic Antigens Act as Rejection Antigens? (Eng) Bansal, B. R. (Dept. Surgery, Alma Dea Morani Lab. Surgical Immunobiology, Medical Coll. Pennsylvania, 3300 Henry Ave., Philadelphia, PA, 19129); Mark, R.; Rhoads, J. E.; Bansal, S. C. *J Natl Cancer Inst* 61(1): 189-201; 1978.

The relationship of antigens shared by embryonic tissue and 1,2-dimethylhydrazine dihydrochloride (DMH)-induced primary gastrointestinal (GI) tumors was studied in inbred WF rats. The rats were inoculated with crude suspensions prepared from liver and gut tissue of 12- to 15-day fetuses of the same strain. Rats previously unsensitized to syngeneic embryonic tissue were inoculated with fetal material sc three times during DMH administration (15 mg base/kg/wk sc, in 2 divided doses). The optimal time period for immunization was when < 30 mg of the total DMH had been administered; ie, prior to the development of microtumors in the GI tract. Properly timed immunization inhibited the development, growth, and metastasis of primary GI tumors. The effect was observed in both sexes, but it was more pronounced in male rats. Seven of nine WF rats with DMH-induced carcinomas of the GI tract rejected syngeneic embryonal tissue implants, demonstrating the presence of a concomitant immune response to embryonic antigen(s). The two rats in which fetal tissue grew out to palpable nodules had multiple GI tumors with metastasis and extra-GI tumors, ie, a massive tumor load. Ten other rats with DMH-induced GI tumors displayed a delayed hypersensitivity reaction to embryonic antigens after being treated with unblocking serum and inoculated id with syngeneic fetal tissue. These results indicate that embryonic antigens common to fetal and tumor cells can induce immunity in an autochthonous host and can act as rejection antigens. (41 refs)

78-5868 The Isozyme Pattern of Cyclic AMP-dependent Protein Kinase and the Distribution of a Cervicovaginal Antigen in Experimental Carcinoma of the Cervix Uteri of Mice. (Eng) Doskeland, S. O. (Inst. Anatomy, Univ. Bergen, Arstadvei 19, 5000 Bergen, Norway); Kalland, T.; Breistein, L. S.; Fossberg, T. M.; Ueland, P. M. *Acta Pathol Microbiol Scand (A)* 86(2): 121-130; 1978.

Tissue sections from 12 3-methylcholanthrene-induced carcinomas of the cervix uteri of mice were tested for the presence of an antigen normally confined to the cervicovaginal epithelium. The antigen was detected in 10/12 tumors examined by indirect immunofluorescence and in all 7 tumors examined by the more sensitive method of mixed hemagglutination. The concentration of the antigen was generally higher in the well-differentiated areas of the tumors, but it was also found to be associated with solitary tumor cells that apparently were invading the stroma. The presence of cervicovaginal antigen in the tumors suggests that tumor cells originated from the cervicovaginal epithelium. The cyclic AMP-dependent protein kinase in the tumor cytosols was studied by chromatography on agarose and diethylaminoethylcellulose. The enzyme had the same properties as those of normal vaginal epithelium. Thus, the tumor cells contained an apparently normal complement of this enzyme, which is believed to be

responsible for most of the intracellular action of cyclic AMP. (36 refs)

See also:

- *(Rev.): 78-5438, 78-5439, 78-5440, 78-5443, 78-5462.
- *(Chem.): 78-5499, 78-5520, 78-5524, 78-5527, 78-5531, 78-5538, 78-564.
- *(Phys.): 78-5722, 78-5728.
- *(Viral): 78-5740, 78-5748, 78-5749, 78-5754, 78-5763, 78-5769, 78-5774, 78-5779, 78-5785, 78-5786, 78-5789, 78-5791, 78-5815, 78-5825, 78-5828, 78-5830, 78-5834, 78-5835, 78-5836.
- *(Path.): 78-5888, 78-5895, 78-5896.
- *(Epid.-Biom.): 78-5920, 78-5940.

PATHOGENESIS

5869 **Familial Intraocular Malignant Melanoma. A Case Report.** (Eng) Green, G. J. (Oncology Sec., Medical Service, Boston Veterans Admin. Hosp., 150 Huntington Ave., Boston, MA, 02130); Hong, W. K.; Evans, J. R.; Bhutani, R.; Amick, R. M. *Cancer* 41(6): 1-2483; 1978.

Family with multiple malignancies, including intraocular malignant melanoma (IMM), is described. Both the 50-yr-old proband and his father had IMM of their left eyes. A third generation was probably involved, since the proband's paternal grandmother had an eye enucleated because of a tumor in her 50's. The relationship of IMM to multiple malignancies is discussed. (20 refs)

5870 **Segregation of Chromosome 13 in Retinoblastoma (Letter to Editor).** (Eng) Knight, L. A. (Div. Cytogenetics, Dept. Pathology, Toronto General Hosp., Toronto M5G 1L7, Canada); Gardner, H. A.; Gallie, L. *Lancet* 1(8071): 989; 1978.

Family in which the mother and one of two living children had retinoblastoma was examined karyotypically to determine if chromosome 13 was associated with the retinoblastoma. The two children had inherited different maternal chromosomes 13, suggesting that the child with retinoblastoma had inherited it from her mother via this chromosome. The other karyotypic examination was not performed on a third child who had died of malignant retinoblastoma at the age of 3. (3 refs)

5871 **Duplication of the Long Arm of Chromosome 1 in a Malignant Vaginal Tumour.** (Eng) Atkin, N. B. (Dept. Cancer Res., Mount Vernon Hosp., Northwood, Middlesex, England); Baker, M. C. *Br J Cancer* 38(3): 468-471; 1978.

Chromosome studies were performed on metaphase specimens of an anaplastic, malignant, small round cell tumor removed from the vagina of a 68-yr-old nulliparous woman. Ten years previously, she had undergone surgery for a well-differentiated endometrial adenocarcinoma; no chemotherapy or radiotherapy had been given. A total of 26/34 metaphases had 47 chromosomes; 1 had 49, 1 had 48, 4 had 46, and 2 had 45 chromosomes. G-banded preparations and karyotypic analysis of 5 metaphases with 47 chromosomes suggested an additional chromosome resembling chromosome 1 in length and centromeric position, but with an abnormal pattern on its short arm. The metaphase with

48 chromosomes differed only in having an additional chromosome 10; that with 49 had an additional C-group and chromosome 3. The abnormal chromosome was present in at least one of 46-chromosome metaphases, which had only one normal chromosome 1. C banding showed that, of the two normal and one abnormal chromosomes 1, two had a pericentric inversion of the heterochromatic region. These findings, combined with previous ones, suggest that a chromosomal imbalance resulting from excess critical gene loci on the long arm of chromosome 1 plays an important role in determining the neoplastic properties of malignant tumor cells. (9 refs.)

78-5872 **Mixed Intracranial Sarcomas: Rare Forms and a New Association with Previous Radiation Therapy.** (Eng) Averback, P. (Dept. Pathology, Univ. Cambridge, Tennis Court Road, Cambridge CB2 1QP, England). *Ann Neurol* 4(3): 229-233; 1978.

The first two cases of mixed sarcoma-glioblastoma with a history of irradiation to the head are reported. Both occurred within 1 yr of radiation therapy for pituitary adenoma and residual meningioma, respectively. In addition, two unique variants of brain mixed sarcomas with marked tumor metaplasia are described, and problems concerning the origin of mixed-population brain tumors are listed. (43 refs)

78-5873 **Melanoma in a Patient Treated for Psoriasis.** (Eng) Durkin, W. (Univ. Alabama, Univ. Station, Birmingham, AL, 35294); Sun, N.; Link, J.; Paroly, W.; Schweitzer, R. *South Med J* 71(6): 732-733; 1978.

The development of melanoma in a 32-yr-old white man who had been treated for psoriasis for 20 yr is discussed. Generalized psoriasis was diagnosed at age 12, and the patient was treated with a variety of regimens, including coal tar and UV light. At age 32 a polypoid, invasive melanoma was diagnosed. During the next 2 yr, metastasis to the right axillary node and left lung was observed. This is the only case of melanoma occurring in a patient treated for psoriasis reported in the world literature. (19 refs)

78-5874 **Liver Tumors and Oral Contraceptives.** (Eng) Ansari, A. H. (Atlanta Fertility Inst., Baptist Professional Building, Suite 641, 340 Boulevard, N.E., Atlanta, GA, 30312); Ravry, M. J.; Stanton, P. E.; Hanafi, M. *Fertil Steril* 29(6): 643-650; 1978.

Three cases of hepatocellular adenoma are reported that support the existence of an association between these tumors and oral contraceptive usage. The three women, aged 25, 48, and 37 yr, had taken oral contraceptives continuously for 7, 10, and 13 yr, respectively. Clinical features and procedures for a correct diagnosis are described. (15 refs)

78-5875 Pleural Mesothelioma: Clinical and Epidemiological Aspects. (Fre) Caillard, J. F. (Service de pneumologie, C.H.R. de Lille, Lille, France); Lahoutte, C.; Frimat, P.; Romon, M. *Arch Mal Prof* 38(12): 1040-1047; 1977.

The role of exposure to asbestos fibers in four cases of pleural mesothelioma is discussed. A brief case report is presented for each of the patients (3 men, ages 51, 54, and 39 yr; and 1 woman, age 70 yr), and the diagnostic methods are described. Two of the men, a ship inspector and a ceramics worker, had been occupationally exposed to asbestos. In a retrospective study of 47 patients with mesotheliomas (23 histologically proved, 13 highly probable, and 7 pleurisy cases of probable mesothelioma origin) observed since 1965, exposure to asbestos was definitely established in 5 and considered probable in 13. (no refs)

78-5876 The Respiratory Epithelium. IV. Histogenesis of Epidermoid Metaplasia and Carcinoma In Situ in the Hamster. (Eng) Becci, P. J. (Life Sciences Res., IIT Res. Inst., 10 W. 35th St., Chicago, IL, 60616); McDowell, E. M.; Trump, B. F. *J Natl Cancer Inst* 61(2): 577-586; 1978.

The histogenesis of epidermoid metaplasia (EMP) in 108 female Syrian golden hamsters was studied in an attempt to demonstrate direct conversion of mucous cells to epidermoid cells. Seventy-two animals received 15 weekly intratracheal instillations of 3 mg benzo(a)pyrene (BP) + 3 mg ferric oxide in 0.2 ml saline. Trachea and bronchi from four airway levels were studied by light microscopy, histochemistry, and electron microscopy. Within 24 hr of the first instillation, foci of goblet cell hyperplasia and several dividing small mucous granule cells (SMGC) were observed, and some ciliated cells were extruded. Extrusion became widespread at 72 hr, with further increases in goblet cells and SMGC. By 1 wk, the latter were replaced by a layer of nonciliated cuboidal SMGC above a single basal cell layer. One week after the third and fifth instillations, the basal cell layer was covered by two-three layers of metaplastic cuboidal cells that, in turn, were covered by two-three layers of cuboidal and/or flattened surface mucous cells (EMP). These upper two-three layers proved to be SMGC. Extensive keratinization occurred at 10 wk (cornifying EMP). The luminal cornifying epidermoid cells contained PAS-positive, diastase-resistant mucosubstances. Localized areas of marked atypical metaplasia and carcinoma in situ were observed 20-50 wk after the first in-

stallation. They were characterized by 1-2 layers of basal cells and 5-10 layers of pleomorphic cuboidal cells. The surface cells had the characteristics of mucous cells, and cells near the basal lamina showed features of epidermoid cells. Mucous-filled intraepithelial cysts were observed. The results show that stratification, EMP, and carcinoma in situ can arise from differentiated mucous cells in hamster tracheal bronchial epithelium, a valid model for studies of human lung carcinoma. (48 refs)

78-5877 The Respiratory Epithelium. V. Histogenesis of Lung Carcinomas in the Human. (Eng) McDowell, E. M. (Dept. Pathology, Univ. Maryland Sch. Medicine, 31 S. Greene St., Baltimore, MD, 21201); McLaughlin, J.; Merenyl, D. K.; Kieffer, R. F.; Harris, C. C.; Trump, B. F. *J Natl Cancer Inst* 61(2): 587-606; 1978.

Following an analysis of 100 human primary lung carcinomas, a histogenetic classification of lung carcinomas based on cell of origin was proposed, and this classification was compared with that devised by the World Health Organization (WHO). Tumors were grouped under the major headings of: tumors from basal and/or mucous cells, tumors from neurosecretory cells, and tumors from Clara cells. Basal/mucous cell tumors, comprising 88% of this sample, were further classified as epidermoid carcinomas (21%), combined epidermoid and adenocarcinomas (AC, 46%), and adenocarcinomas (21%). The criteria for epidermoid differentiation included the presence of tonofilament bundles, poorly developed endoplasmic reticulum (ER) and Golgi apparatus, and well-developed desmosomes. The criteria for adenoid differentiation included well-developed ER and Golgi apparatus, poorly developed desmosomes, extracellular and/or intracellular alveoli, and/or other evidence of cellular secretion such as secretory granules. In AC other extracellular alveoli, typical junctional complexes were also present at the luminal aspect where the cell apexes bordered the alveolus. A histogenetic classification of lung tumors may be of value in current and future studies of epidemiologic risk factors, prognosis, prevention, and treatment of lung cancer. (78 refs)

78-5878 The Respiratory Epithelium. VI. Histogenesis of Lung Tumors Induced by Benzo(a)pyrene-Ferric Oxide in the Hamster. (Eng) Becci, P. J. (Life Sciences Res., IIT Res. Inst., 10 W. 35th St., Chicago, IL, 60616); McDowell, E. M.; Trump, B. F. *J Natl Cancer Inst* 61(2): 607-618; 1978.

Lung tumors induced in female golden Syrian hamsters by 15 weekly intratracheal instillations of benzo(a)pyrene (BP)-ferric oxide (each, 3 mg) were examined by light and electron microscopy and classified according to a histogenetic scheme based on cell of origin. Most of the tumors that resulted were derived from mucous cells and/or basal cells. There were 14 well- or moderately well-differentiated epidermoid

nomas, 6 epidermoid papillomas, and 20 combined epidermoid and adenocarcinomas. The latter had a wide spectrum of epidermoid and adeno components. The epidermoid component was characterized in well-differentiated tumors by the presence of intercellular bridges and/or keratinization. Well-developed desmosomes and numerous tonofilament bundles were observed ultrastructurally. The adeno component was defined by intracellular and/or extracellular alveoli, a well-developed Golgi complex and endoplasmic reticulum, and/or evidence of mucous secretion. A tumor composed of neurosecretory cells that resembled a bronchial carcinoid in humans was also found. Nests of uniform, small, polygonal cells with round-to-oval nuclei were seen light microscopically, and dense-core secretory granules 1,100-2,200 nm in diameter were present in the cytoplasm. There were also fibrosarcomas that had a very cellular structure, being composed of densely packed ovoid or spindle-shaped cells that resembled fibroblasts. The results strongly support the histogenetic classification scheme; ie, a classification based on the cell of origin. (39 refs)

5879 Mechanisms of Late Recurrence after Radical Surgery for Gastric Carcinoma. (Eng) Iwanaga, T. (Dept. Surgery, Center for Adult Diseases, Osaka, 1-3-3 Kamichi, Higashinari-ku, Osaka 537, Japan); Koyama, H.; Fukawa, H.; Taniguchi, H.; Wada, A.; Tateishi, R. *Am J Surg* 135(5): 637-640; 1978.

Factors involved in the recurrence of gastric carcinoma were investigated in 25 patients with late recurrence who died 5 or more after surgery and in 129 with early recurrence who died within 2 yr of surgery. The late recurrence rate was high in the residual stomach or gastric stump, but the early recurrence rate was high in the peritoneum and liver. In the late recurrence group, the routes of cancer spread were contiguous invasion and lymphatic spread. In the early recurrence group, dissemination and hematogenous spread were also important routes. There was a high correlation between both rates of recurrence and the following conditions: the amount of residual cancer remaining after surgery; the spread of the residual cancer; the rate of cancer proliferation; and host resistance, as represented by cellular stromal reactions, sinus stasis, and lymph node metastasis. The prevention of late recurrence is contingent on the complete removal of all cancerous tissue combined with adjuvant chemo- or immunotherapy. (12 refs)

5880 Polypoid Gastritis. Report of a Case Associated with Gastric Adenocarcinoma and Review of the Literature. (Eng) Sanner, C. J. (Dept. Internal Medicine, Section Gastroenterology, Univ. Arizona Medical Center, Tucson, AZ, 85724); Saltzman, D. A.; Mueller, J. C. *Am J Dig Dis* 23(5, Suppl): 19s-24s; 1978.

Examination of biopsy specimens from a 46-yr-old man who

presented with gastric polyposis revealed chronic superficial gastritis. Directed brushing for cytology and gastric wash cytology revealed no malignant cells. However, follow-up examination 6 mo later revealed not only areas of chronic superficial gastritis, but also adenocarcinoma in an area of erosion. Total gastrectomy was performed; adenocarcinoma was found in 3/5 omental lymph nodes and in one celiac lymph node. Because of the high incidence of gastric carcinoma in patients with atrophic hyperplastic gastritis, a correct diagnosis and long-term follow-up of polypoid gastritis patients are necessary. A review of the literature is presented with emphasis on the association between polypoid gastritis and gastric carcinoma. (43 refs)

78-5881 Increased In Vitro Tetraploidy: Tissue Specific within the Heritable Colorectal Cancer Syndromes with Polyposis Coli. (Eng) Danes, B. S. (Lab. Cell Genetics, Dept. Medicine, Cornell Univ. Medical Coll., New York, NY, 10021). *Cancer* 41(6): 2330-2334; 1978.

The occurrence of tetraploidy in cultures established from surgical tissue specimens from patients with familial polyposis coli (FPC) and Gardner's syndrome was studied. The cultures established from all tissue (skin, mesentery, ascending colon, normal mucosa, and colonic wall) obtained from four normal subjects not at risk for either syndrome showed 0%-4% tetraploidy. For the two patients with FPC, 0%-2% tetraploidy was observed in specimens of epidermis, dermis, mesentery, and colonic wall; 12%-20% tetraploidy was observed in samples of normal colonic mucosa; and 18%-22% tetraploidy was observed in samples of colonic polyps. The occurrence of tetraploidy in the eight patients with Gardner's syndrome was 11%-29% in the epidermis, 0%-3% in the dermis, 0%-1% in the mesentery, 0%-2% in the colonic wall, 14%-26% in the normal colonic mucosa, and 12%-32% in the colonic polyps. The occurrence of tetraploidy in the two patients with a variant form of Gardner's syndrome was 0%-1% in the epidermis; 0%-2% in the dermis, mesentery, and colonic wall; 10%-16% in the normal colonic mucosa; and 15%-20% in the polyps. The results suggest that in Gardner's syndrome and FPC, there is a constant population of tetraploid cells, at least in culture, that may be relevant to the multistep process of carcinogenesis. (12 refs)

78-5882 Multiple Lymphoid Polyposis and Familial Polyposis of the Colon: A Genetic Relationship. (Eng) Venkitachalam, P. S. (Dept. Surgery, Jewish Hosp. Brooklyn, 555 Prospect Place, Brooklyn, NY, 11238); Hirsch, E.; Elguezabal, A.; Littman, L. *Dis Colon Rectum* 21(5): 336-341; 1978.

The second literature case of multiple lymphoid polyposis (MLP) of the colon and terminal ileum occurring in a member of a family with a history of congenital familial polyposis (FP) is reported. The grandfather and aunt of the patient, a

12-yr-old boy, had died of carcinoma and polyposis of the colon; his mother had undergone total colectomy and ileostomy for FP. The family history and sigmoidoscopic and x-ray findings strongly suggested a case of congenital FP, and subtotal colectomy was performed without prior biopsy. Microscopically, all lesions examined postoperatively were mucosal lymphoid polyps and not adenomatous polyps. MLP is a disease that occurs in younger patients, who have diffuse involvement of the entire mucosa by small polyps that have a benign course. There is a genetic factor in the etiology of the condition, and a definite link with FP of the colon is found to exist. MLP and congenital FP are accurately distinguishable only by multiple biopsy. Literature cases in which an association between the conditions is mentioned are reviewed. (31 refs)

- 78-5883 **Pathogenesis of Polyposis of the Large Intestine.** (Rus) Malyshev, Iu. I. (Dept. Oncology, D. I. Ul'ianov Medical Inst., Kuibyshev, USSR); Katorkin, E. N. *Vopr Onkol* 24(4): 35-38; 1978.

In an attempt to evaluate the etiology and pathogenesis of polyposis of the large intestine, various manifestations of the disease were reviewed in 400 patients. The duration of polyposis ranged from 1 mo to 20 yr. Of 62 patients who underwent aspiration biopsy of the small intestine, 28 had clinical symptoms of small intestine involvement and 34 had symptoms of colitis. Villous atrophy of the small intestinal mucosa was detected in 20/62 patients (17 had mild and 3 had marked atrophy). All 180 patients who underwent x-ray examination had symptoms of chronic enteritis. Clinical manifestations of polyposis of the large intestine were correlated with disturbances of absorption in the small intestine. (7 refs)

- 78-5884 **Colonic Cancer in Inflammatory Bowel Disease (Letter to Editor).** (Eng) Sauer, W. G. (No affiliation given). *Mayo Clin Proc* 53(9): 618; 1978.

The recommendation of prophylactic proctocolectomy for patients with ulcerative colitis, in view of their increased risk of colon cancer with time, is meant only for patients whose ulcerative colitis began in childhood and is of a certain duration and extent, and not for all patients with this disease. (9 refs)

- 78-5885 **Cytomegalovirus D.N.A. and Adenocarcinoma of the Colon: Evidence for Latent Viral Infection.** (Eng) Huang, E. S. (Dept. Medicine, Univ. North Carolina at Chapel Hill, Chapel Hill, NC); Roche, J. K. *Lancet* 1(8071): 957-960; 1978.

Normal and carcinomatous human colon specimens were ex-

amined for the presence of human cytomegalovirus (CMV) DNA. A total of 28 specimens from 14 patients were analyzed by complementary RNA-DNA hybridization. Nine patients had adenocarcinoma of the large intestine, 2 had familial polyposis, and 3 had Crohn's disease. Of the 14 patients with carcinoma of the colon, three were definitely positive for CMV DNA (≥ 2 genome equivalents/cell) and one showed a consistently low level of CMV DNA hybridization in the range of 1-2 genome equivalents/cell. None of the macroscopically normal tissue (5 cm from tumor) from the 14 patients demonstrate CMV DNA, with the exception of one patient who had a recurrence within 2 yr of surgery. One specimen from a patient with familial polyposis had 1 genome equivalent/cell. Histologically normal and abnormal colon specimens from the two rectal carcinoma patients or from the three patients with Crohn's disease were uniformly negative for CMV DNA. These findings indicate an association between CMV and carcinoma of the colon: human CMV by itself or with other factors may induce carcinoma of the colon or human CMV may be prone to multiply and become reactivated in carcinomatous and/or associated inflammatory cells. (18 refs)

- 78-5886 **Rat Ileocecal Immunocytoma. An Ultrastructural Study with Special Attention to the Presence of Viral Particles.** (Eng) Burtonboy, G. (Laboratoire Virologie, Faculte de Medecine, Universite Catholique de Louvain, 30, Clos Chapelle aux Champs, boîte 3055, B-1200 Brussels, Belgium); Beckers, A.; Rodhain, J.; Bazin, H.; Lamy, M. E. *J Natl Cancer Inst* 61(2): 477-484; 1978.

Electron microscopic (EM) examination was made of spontaneous sarcomas in Lou/C/Wsl rats. These tumors were named ileocecal immunocytomas because of their unusual site of origin and their capacity to synthesize and secrete monoclonal rat immunoglobulins (Ig's). Although necrosis was prevalent, well-preserved zones were found that consisted of large, rounded, isolated cells with slightly eccentric ovoid nuclei. One or two large nucleoli were often evident, and chromatin was condensed in a thin layer along the perinuclear cisterna and around the nucleolus. Large Golgi complexes and polyribosomes were obvious. The development of the rough endoplasmic reticulum (RER) varied from cell to cell and was well-developed in only a few cells. In these cells, the RER contained abundant mitochondria. In all tumors examined, small numbers of intracisternal A-type virus-like particles were found in the cavities of RER, in smooth vacuoles, or in the perinuclear cisterna. Furthermore, anomalous ribbonlike prolongations consisting of folds of nuclear membrane surrounding a leaflet of dense material were frequently noted. Many cells displayed nuclear degeneration. The sera of 10 tumor-bearing animals had up to 40 mg/ml monoclonal Ig, and 2 other rats had Bence Jones protein in their urine. It was not possible, however, to distinguish secretory from nonsecretory tumors by EM. The tumors resembled plasmacytomas induced in mice and certain human tumors such as Burkitt's lymphoma. (32 refs)

5887 **Origin of Intestinal Lymphomas (Letter to Editor).** (Eng) Isaacson, P. (Dept. Pathology, Southampton Univ. Medical Sch., Southampton SO9 4XY, and); Wright, D. H. *Lancet* 1(8071): 984-985; 1978.

Study of 66 primary intestinal lymphomas indicated that were of histiocytic origin; no plasma cell tumors were. This is in contrast to a previous study that claimed 39% of the intestinal lymphomas were plasmacytic in. Immunoperoxidase studies were performed on 7/23 of Middle East lymphoma; 6 had the morphological and immunohistochemical characteristics of histiocytic lymphoma. It is suggested that the sequence of progressive hyperplasia and plasma cell proliferation in the gut induces malignancy in the histiocytes of the lamina propria. (10 refs)

5888 **Carcinoma-In-Situ of the Cervix and Vulval Warts Complicating Malignant Lymphoma.** (Eng) El-Maraghy, M. A. (Dept. Midwifery and Gynaecology, Queen's Univ. Belfast, Belfast, Ireland); Lamki, H.; Pinkerton, J. H.; Mawhinney, H. *Ir Med J* 71(5): 159-161; 1978.

33-yr-old woman with malignant lymphoma was treated with radiotherapy to the supraclavicular mass of nodes, excision of the nodes, and chemotherapy with vinblastine, mustard, Natulan, and prednisone. Approx 3 yr later, she was found to have vulval warts and carcinoma in situ of the cervix. Laboratory studies were suggestive of a partial T-cell or humoral-mediated immunity defect in this patient. The immunosuppressive treatment may have made her more susceptible to viral infection. (8 refs)

5889 **The Morphology of the Earliest Invasive Cell in Low Genital Tract Epidermoid Neoplasia.** (Eng) Genadry, R. (Dept. Gynecology and Obstetrics, Johns Hopkins Hosp., Baltimore, MD, 21205); Olson, J.; Parmley, J. Woodruff, J. D. *Obstet Gynecol* 51(6): 718-722; 1978.

The light and electron microscopic characteristics of the distinct eosinophilic microinvasive cell in lower genital tract epidermoid neoplasia are described. Ten cases each of grossly invasive vulvar, vaginal, and cervical tumors and 12 cases of microinvasive disease (4 vulvar and 8 cervical neoplasms) were examined, and cells at the periphery of 3 vulvar and 3 cervical invasive cancers were subjected to electron microscopy. The cells forming the initial protrusion of invasive cancer were characterized by large amounts of eosinophilic cytoplasm, regardless of the degree of differentiation of the rest of the tumor. The eosinophilic quality of the invasive cell was associated at the ultrastructural level with an accumulation of contractile protein microfilaments, which apparently presents the basis of the cell's invasive capacity. It is speculated that intraepithelial neoplasms that show large, apparently mature cells near the basement membrane should be viewed with more concern than those that demonstrate a

full-thickness change of small, dark, basal and parabasal cells without apparent cytoplasmic differentiation. (14 refs)

78-5890 **Electron Microscopic Observations of Vaginal Development in Untreated and Neonatally Estrogenized Balb/c Crgl Mice.** (Eng) Cunha, G. R. (Anatomy Dept., Sch. Medicine, Univ. Colorado, Denver, CO); Lee, A. K.; Lung, B. *Am J Anat* 152(3): 343-381; 1978.

The developing Mullerian vaginae of untreated and neonatally estrogenized [20 µg estradiol (E) given daily for 5 days] 1- to 12-day-old BALB/c Crgl mice were examined by light and electron microscopy. The vaginae of untreated 1-day-old mice had an undifferentiated, pseudostratified columnar epithelium that slowly differentiated into an apparently mucus-secreting epithelium. By day 12, the rough endoplasmic reticulum (RER), Golgi apparatus, and secretion granules were poorly developed features of the apical epithelial cells, but the stromal cells had developed prominent RER and Golgi zones that were involved in the synthesis and secretion of collagen. The collagen accumulated in the intercellular spaces. By day 2, the E-treated mice showed a slight increase in the amount of RER and Golgi structures over untreated controls, and by day 4, there was a significant increase in the number of desmosomes. E accelerated the differentiation of mucus-secreting apical epithelial cells, whose apices became filled with secretory granules, and it promoted the formation of a cornifying epithelium from the underlying basal cells. The latter process was associated with an increase in cellular proliferation and in the number of desmosomes and associated tonofilaments, and with extensive interdigitation of the plasma membranes. In both untreated and E-treated mice, the vaginal epithelium was separated from the stroma by a continuous basal lamina, which suggests that vaginal morphogenesis may be mediated by extracellular materials at the surface between these tissues. (67 refs)

78-5891 **Recurrent Extreme Breast Involution Following Pregnancy and Susceptibility to Breast Cancer: A Hypothesis.** (Eng) Petrakis, N. L. (G. W. Hooper Foundation, Univ. California, San Francisco, CA, 94143). *Med Hypotheses* 4(3): 268-272; 1978.

Nine multiparous Oriental women whose breasts had regressed to a state of flat-chestedness following pregnancy and lactation were found during a breast cancer screening project in the San Francisco Bay Area. During their pregnancies, the breasts of all nine women had developed and enlarged, and five had successfully nursed for ≥2 mo (4 did not attempt to breastfeed). Following cessation of lactation in the women who had breastfed and after delivery in those who had not, the breasts completely regressed to their prepregnant flat-chested state. All of the women had normal menstrual cycles and none suffered from major organic disease. It is

hypothesized that this extreme regression of the breast tissue may represent a marked reduction in genetically determined sensitivity of alveolar and ductal tissue to estrogen and other hormonal factors implicated in the maintenance of breast function during the nonpregnant state. The lower incidence of breast cancer reported in Oriental women may be associated with a relative insensitivity of the breast epithelium to estrogen during the nonpregnant state, which results in a decreased exposure and responsiveness of the breast to secreted exogenous and endogenous carcinogens. (13 refs)

- 78-5892 Familial Predisposition to Endometrial Cancer.** (Rus) Shvedova, L.A. (Moldavian Res. Inst. Oncology, Kishinev, USSR). *Akush Ginekol (Mosk)* (6): 58; 1978.

Papillary endometrial carcinoma developed in a 56-yr-old woman whose mother died of cancer and elder sister died of acute leukemia. Of three other sisters, one had papillary endometrial carcinoma; one had uterine fibromyoma, endometrial polyposis, and endocervicosis; and one had uterine fibromyoma with endometrial polyposis. Thus, first-degree relatives of patients with cancer of the corpus uteri constitute a group of increased risk. (1 ref)

- 78-5893 Abnormal Chromosomes and Number 1 Heterochromatin Variants Revealed in C-banded Preparations from 13 Bladder Carcinomas.** (Eng) Atkin, N. B. (Dept. Cancer Res., Mount Vernon Hosp., Northwood, Middlesex HA6 2RN, England); Baker, M. C. *Cytobios* 18(70): 101-109; 1978.

Chromosome preparations from 13 patients (11 men, 2 women aged 40-84 yr) with carcinomas of the bladder were subjected to C and G banding and then studied. Six of the patients had received previous surgical treatment for their tumors, but none had received radiotherapy. Variations in the areas of the C bands of the no. 1 chromosomes suggested a constitutional heteromorphism in eight patients; peripheral blood studies in three of the patients confirmed this heteromorphism. Possible pericentric inversions of the heterochromatic regions of the no. 1 chromosomes were present in four patients. Grossly abnormal chromosomes containing major parts of the no. 1 chromosomes were present in seven patients. They included possible isochromosomes for the long arm of no. 1 in two cases and additional C bands on the long arm of no. 1 in two cases. Minute chromosomes that appeared to consist mainly of heterochromatin were present in four patients. (6 refs)

- 78-5894 Bladder Carcinoma Associated with Long-Term Treatment with Cyclophosphamide for Multiple Myeloma.** (Ger) Lenzin, A. (Urologische Universitätsklinik,

Inselspital, Anna-Seiler-Haus, CH-3008 Bern, Switzerland). Cavalli, F.; Sonntag, R.; Zimmermann, A. *Urologe [A]* 17(10): 105-108; 1978.

A 70-yr-old woman with multiple myeloma developed bleeding extended squamous epithelial carcinoma of the urinary bladder immediately after treatment with cyclophosphamide (total dose 135 g po) for several years. (29 refs)

- 78-5895 Chronic Granulocytic Leukemia Following Successful Renal Transplantation.** (Eng) Adler, R. (Div. Hematology, Depts. Medicine and Surgery, Albany Medical Coll., Albany, NY); Lempert, N.; Scharfman, W. *Cancer* 41(6): 2206-2208; 1978.

Philadelphia chromosome-positive chronic granulocytic leukemia (CGL) occurred in a renal transplant recipient (22-yr-old man) on immunosuppressive therapy. At age 22 the patient received and subsequently rejected a sibling kidney despite irradiation to the transplant site and treatment with immunosuppressive drugs. Two years later, the kidney was removed and he received a second renal transplant from a cadaver donor. The patient achieved normal renal function and was discharged on azathioprine (150 mg/day) and prednisolone (12 mg/day). Nine months later, a progressive thrombocytosis, leukocytosis, and hyperuricemia were noted, although the patient remained asymptomatic. CGL was diagnosed, and chromosome studies showed the presence of the Philadelphia chromosome (G22) with translocation of chromosome 7 of the C group. The patient's impaired renal function was readily reversed by lowering his dosage of immunosuppressive drugs. He was continued on these drugs, however, and alkylating agents and plateletpheresis were used to regulate his peripheral blood count. His CGL has been in remission for 10 mo, and normal kidney function has been maintained. A review of transplantation centers revealed a total of 5 cases of CGL among a population of 25 renal transplant patients, a fivefold increased incidence over the general population. Immunosuppressive drugs should be used cautiously in nonmalignant disease. (15 refs)

- 78-5896 Acute Leukemia after Prolonged Immunosuppressive Therapy for Glomerulonephritis.** (Fr) Carcassonne, Y. (Clinique des Maladies du Sang, Univ. Marseille, Marseille, France); Gastaut, J. A.; Blanc, A. P. *Cancer Treat Rep* 62(7): 1110-1111; 1978.

Two male patients (15 and 56 yr old) developed acute leukemia 6 and 4 yr, respectively, after the onset of chlorambucil therapy for glomerulonephritis. The duration of chemotherapy was 3 and 2 yr, respectively, for total doses of 8.9 and 6 g; both patients also received small doses of indomethacin (1.5 and 3 g). (6 refs)

97 **Busulphan Haemorrhagic Cystitis.** (Eng) Millard, R. J. (Central Middlesex Hosp., London, Eng.). *Br J Urol* 50(3): 210; 1978.

Haemorrhagic cystitis developed in a 59-yr-old man who had been taking busulfan (total dose, 1.98 kg) since 1973 for chronic myeloid leukemia. The man had presented with hematuria of several days' duration and no other symptoms. Side effects of busulfan and cyclophosphamide, a drug known to cause hemorrhagic cystitis and bladder tumor, are pointed out. (3 refs)

98 **Stem Cell Leukemia Following Intensive Radiotherapy and Chemotherapy for Hodgkin's Disease.** (Ger) Trux, F. (Medizinische Klinik, Medizinische Hochschule Erfurt, Nordhauser Strasse 74, DDR-50 Erfurt, Germany); Fink, R.; Wutke, K. *Folia Haematol (Leipz)* 35: 354-366; 1978.

Stem cell leukemia developed in 2/106 patients treated for Hodgkin's disease (HD) during an 8-yr period. A 59-yr-old man with Stage IIIA HD (nodular sclerosis type) responded to split-course telecobalt therapy (total dose 4,000 rads to supraclavicular field, 4,400 rads to the mediastinum, 4,000 rads to a paraaortic abdominal field, and repeated supraclavicular irradiation with 4,000 rads) and chemotherapy with vincristine, cyclophosphamide, and prednisolone. However, the patient developed stem cell leukemia approximately 1 year after the last irradiation and, despite combination chemotherapy, died from a cerebral hemorrhage in 20 days. A 59-yr-old man with Stage IIA HD (mixed cell type) received radiotherapy (4,500 rads initially, later 4,300 rads) and chemotherapy according to a modified version of the De Vita protocol, in which Mustargen was replaced by cyclophosphamide and vincristine by vinblastine. He developed monocytic leukemia and died 5 days after diagnosis. A poorly differentiated mesenchymal neoplasm was found at the site of an earlier irradiation. Lymphogranulomatosis was not found at autopsy in either patient. (34 refs)

99 **The Blast Crisis of Chronic Granulocytic Leukemia: Megakaryoblastic Nature of Cells Revealed by the Presence of Platelet-Peroxidase--A Cytological Ultrastructural Study.** (Eng) Breton-Gorius, J. (S.E.R.M., U. 91 Hopital Henri Mondor, 94010 Creteil, France); Reyes, F.; Vernant, J. P.; Tulliez, M.; Dreyfus, B. *Haematol* 39(3): 295-303; 1978.

A cytochemical ultrastructural study was made of the megakaryoblastic nature of the cells in blast crisis of chronic granulocytic leukemia in a 34-yr-old woman. She had presented, 4 yr earlier, with postpartum hemorrhage, an enlarged spleen, and hepatomegaly. Circulating blast cells were characterized by a high nucleocytoplasmic ratio, dense chromatin, lack of granules, and a negative peroxidase (PO)

reaction on smears. Accordingly, they were termed lymphoid or undifferentiated cells. Most (90%) of these cells could be identified as megakaryoblasts, since they contained PO-like activity in the endoplasmic reticulum (ER) that was distinct from myeloperoxidase. The PO activity had the following similarities to platelet PO: (1) the activity was very sensitive to inhibition by an aldehyde fixative under conditions in which PO activity is preserved in granulocytic precursors; (2) the ER localization was similar to that found in the normal megakaryocytic series; and (3) in the blasts and megathrombocytes, the Golgi apparatus was devoid of activity except for one single cisterna. In addition, this Golgi-associated positive cisterna often exhibited a much higher PO content than the ER cisternae. These data emphasize the possible megakaryoblastic nature of cells in chronic granulocytic leukemia blast crisis. (56 refs)

78-5900 **Diffusion Chamber Culture in Irradiated Rats of Myeloblasts and Promyelocytes from 84 Untreated Patients with Myeloid Leukemias or Refractory Dysmyelopoietic Anemia: Comparison to Continuous Tissue Culture Lines.** (Eng) Greenberger, J. S. (Div. Hematology, Peter Bent Brigham Hosp., Boston, MA); Gans, P. J.; King, V.; Muse, M. B.; Karpas, A.; Moloney, W. C. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 299-302; 1978.

Bone marrow or buffy coat specimens from 55 untreated patients with myeloid leukemias, 24 with refractory dysmyelopoietic anemia (RDA), and 5 normal subjects were grown in diffusion chambers (DC) implanted into irradiated (750 rads) Sprague-Dawley rats. Fifty-four specimens grew for > 21 days: 22/31 acute granulocytic leukemias (AGL), 8/10 chronic myelogenous leukemias in blast crisis, 8/14 chronic myelocytic leukemias, 14/24 RDA's, and 2/5 normal specimens. Immature cells (myeloblasts, promyelocytes, and myelocytes) increased over input numbers in 24 specimens; macrophages increased in 78. Two continuous lines were established from AGL cells explanted from the DC to tissue culture. Twenty other cell lines established directly in vitro showed myeloid cell-associated enzyme increases after butyric acid or DC culture. The DC results indicate that alteration of host environment, as well as intrinsic cellular defects, may regulate the differentiation, proliferation, and malignant transformation of human myeloid cells. (3 refs)

78-5901 **Acute Megakaryocytic Leukemia: Description of a Case Initially Seen as Preleukemia Syndrome.** (Eng) Balducci, L. (Div. Hematology and Oncology, Dept. Medicine, Univ. Mississippi, Jackson, MS); Weitzner,

S.; Beghe, C.; Morrison, F. S. *Arch Intern Med* 138(5): 794-795; 1978.

A 52-yr-old man with sideroblastic anemia that was unresponsive to pyridoxine hydrochloride developed acute megakaryocytic leukemia 3 mo after diagnosis. The leukemia treatment regimen included doxorubicin hydrochloride, cytarabine, vincristine sulfate, and prednisone, but none were effective; he died 1 mo after diagnosis of the leukemia. Autopsy revealed acute megakaryocytic leukemia with involvement of the bone marrow, liver, spleen, adrenals, kidneys, and thyroid. Chromosomal analysis revealed the absence of both diploid and Philadelphia chromosomes. A mode of 45 chromosomes and aneuploidy were present. This is the first known case in which a patient with acute megakaryocytic leukemia had a recognized preleukemia syndrome. (15 refs)

78-5902 Three Cases of Acute Leukemia in One Pedigree. (Jpn) Dan, K. (First Dept. Internal Medicine, Tokyo Medical and Dental Univ., Tokyo, Japan); Kuriya, S.; Matsui, Y.; Kudo, H.; Kawada, K.; Nomura, T.; Sano, T. *Jpn J Clin Hematol* 19(6): 844-849; 1978.

Acute leukemia developed in three members of one pedigree over a period of 7 yr, an unusual phenomenon. Case 1, a 59-yr-old man, was diagnosed with acute myeloblastic leukemia (AML) in June 1959. His daughter (Case 2) was hospitalized with complaints of fever, malaise, and petechiae at age 29, and acute lymphoblastic leukemia was diagnosed in January 1976. Investigation of the family history revealed a case of AML (diagnosed in December 1970) in a son of the niece of case 1. There was no evidence of consanguinity in this family. Cases 1 and 2 lived in the same house until the occurrence of leukemia in the former; case 3 lived in a different location. Common environmental factors such as ionizing radiation and toxic chemicals could not be detected among the three cases. The occurrence and significance of familial leukemia in relation to the pathogenesis of leukemia are discussed. (13 refs)

78-5903 Discordant Chronic Myelocytic Leukemia in Identical Twins (Letter to Editor). (Eng) Goh, K. (Rochester, NY); Bauman, A. W. *Arch Intern Med* 138(8): 1303; 1978.

The Philadelphia (Ph¹) chromosome could not be demonstrated in two normal subjects whose identical chronic myelocytic leukemia (CML)-diseased twins had typical Ph¹ chromosome. This chromosomal abnormality may be acquired postzygotically, and it may be extended to show no 9q+ or other translocated chromosomes in a normal, identical twin of a CML patient. (5 refs)

78-5904 Production of Mesenchymal Tumors in Nude Mice by Ph¹ Negative Fibroblasts Obtained

from a Ph¹ Positive CML Patient: A Preliminary Report. (Eng) Wilson, F. D. (Radiobiology Lab., Univ. California, Davis, CA, 95616); Greenberg, B.; Spangler, W.; Shifrin, M.; Gershwin, M. E.; Dyck, J. *Exp Hematol* 6(6): 541-544; 1978.

The role of hematopoietic stromal elements in the regulation of hematopoiesis and in the pathogenesis of myelofibrosis and myeloproliferative disorders was investigated in an experimental model system permitting the independent study of stromal vs hematopoietic elements in vivo. The model was based on the simultaneous application of three techniques: (1) growth of bone marrow (BM)-derived fibroblastic colonies in vitro; (2) cytogenetic demonstration of marker chromosomes associated with hematopoietic malignancies; and (3) transplantation of isolated stromal elements into athymic (nude) mice. Injection of $5-9 \times 10^6$ Philadelphia chromosome (Ph¹)-negative fibroblasts from the BM of a patient with Ph¹-positive chronic myelogenous leukemia (CML) produced mesenchymal tumors in 2/5 mice. Mesenchymal tumors were also induced by BM-derived fibroblasts from a patient with aplastic anemia who was transplanted successfully with BM transplantation (2/4 mice) and by a normal human (1/2 mice). Morphologic, cytogenetic, and electron microscopic studies of BM mesenchymal elements in culture and of tumors induced in nude mice by the fibroblasts indicated that the tumor cells were of human origin and negative for the Ph¹ chromosome. These results, obtained with the use of human materials, provide the first in vivo morphological and cytogenetic support of the hypothesized relationship of precursors of in vitro fibroblast colonies to BM stromal elements. (17 refs)

78-5905 Philadelphia Chromosome-positive Chronic Myeloid Leukemia and Polycythemia (Letter to Editor). (Fre) Pris, J. (Service d'hématologie, Hôpital Dieulafoy, CHU Purpan, Place du Dr. Baylac, F 31062 Toulouse Cedex, France); Launais, B.; Capdeville, J.; Merle, J.; Bourrouilloux, G.; Duchayne, E.; Colombies, P. *Presse Med* 7(13): 1114-1115; 1978.

A 70-yr-old woman developed Philadelphia chromosome-positive chronic myeloid leukemia (CML) 2 yr after an initial diagnosis of polycythemia vera. The CML was not attributed to treatment of the polycythemia with ³²P because of the dose (7 mCi) involved and the short latent period of the leukemia (2 yr). (6 refs)

78-5906 Trisomy 8 in the Chronic Phase of Philadelphia Chromosome-Negative Chronic Myelocytic Leukemia. (Eng) Lindquist, R. (Section Oncology and Haematology, Huddinge Univ. Hosp., S-14186 Huddinge, Sweden); Gahrton, G.; Friberg, K.; Zech, L. *Scand J Haematol* 21(2): 109-114; 1978.

case is reported of a 71-year-old woman with Philadelphia chromosome (Ph¹)-negative chronic myelocytic leukemia (CML) who possessed trisomy 8 at diagnosis and throughout the chronic stage of her disease. The patient had history of thyroid gland enlargement and thyroiditis, and a 5 cm benign thyroid adenoma was diagnosed on admission (8/30/74). CML was confirmed at this time but left untreated. In 1975, the patient's WBC count increased to 115,000/liter, and busulfan treatment was initiated. In December 1975, leukemic skin infiltrations appeared; and in July 1976, therapy-resistant blast formation was noted. Chromosomes were examined on four occasions, beginning with first admission, and 100% of bone marrow metastases always showed trisomy of chromosome 8. Although trisomy 8 is common in Ph¹-positive CML in the blast phase, this anomaly in Ph¹-negative patients is unusual, especially in the early course of the disease. If trisomy 8 can appear at any stage in Ph¹-positive and Ph¹-negative patients, the anomaly would have little prognostic value. (38 refs)

78-5907 Down's Syndrome and Leukemia: Mechanism of Additional Chromosomal Abnormalities. (Eng) Goh, K. (Monroe Community Hosp., Univ. Rochester School of Medicine and Dentistry, 435 E. Henrietta Road, Rochester, NY, 14603); Lee, H.; Miller, G. *Am J Ment Defic* 82(6): 548; 1978.

Cytogenetic findings are reported for five Down's syndrome patients, four with acute leukemia and one with abnormal maturation of granulopoiesis. All patients exhibited marked polyploidy and three had morphologically abnormal chromosomes. Several metaphases from each patient had chromosome abnormalities that suggested a clonal evolution. The frequency of the affected chromosome groups, excluding those characteristic of Down's syndrome, showed that chromosome A₁ was involved 9 times; A₂, 6; A₃, 6; C, 105; D, 14; E, 48; F, 56; G, 56; and abnormal chromosomes, 17. Eight G-banding chromosomes had trisomy 21 and/or 22. The abnormal chromosome patterns in these patients could be due to nondisjunction in mitosis, an abnormality that resulted in the phenotypical expression of acute leukemia. (23 refs)

78-5908 A Near Haploid Clone: 24,XY,t(9;22)(q34;q11) from a Patient in Blast Crisis of Chronic Myeloid Leukaemia. (Eng) Daniel, A. (Cytogenetics and Cell Biology Unit, Prince of Wales Hosp., Sydney, Australia); Francis, S. E.; Stewart, L. A.; Barber, S. *Scand J Haematol* 21(2): 103; 1978.

The case of a 58-year-old man with chronic myeloid leukemia who displayed a markedly hypodiploid chromosome complement is reported. Eight days after admission, marrow chromosome studies revealed a single clone of 24,XY,t(9;22)(q34;q11), indicative of Philadelphia chromo-

soma. Busulfan treatment (2 mg/day) was initiated but discontinued when the patient developed leukopenia. Bone marrow aspirate revealed 98% blasts. The blast cells were very small compared with promyelocytes. The patient died within 2 mo of the removal of a ruptured spleen and the onset of cervical adenopathy. Postmortem marrow chromosome studies revealed a major clone of 24,XY,t(9;22), a minor clone of 46,XY,t(9;22), and possible clones of 47,XY,t(9;22)+del 22(q11), and 48,XY,t(9;22)+C,+de 122(q11). There were no clones with a karyotype intermediate between those with 24 and 46 chromosomes. A haploidy event rather than progressive hypodiploid loss may have accounted for the clone with 24 chromosomes. (9 refs)

78-5909 A Note on Banded Karyotypes in Spontaneous Leukemias of the AKR Mouse. (Eng) Kodama, Y. (Chromosome Res. Unit, Faculty Science, Hokkaido Univ., Sapporo, Japan); Yoshida, M. C.; Sasaki, M. *Proc Jpn Acad Ser B Phys Biol Sci* 54(5): 222-227; 1978.

Banded karyotypes were analyzed in 20 cases of spontaneous leukemia in AKR mice. The 784 cells examined originated from a thymic tumor in 8 mice, from leukemic spleen in 8, and from both in 4. Cells from metastatic lymph nodes (4 mice) and bone marrow (3 mice) were also examined. Analyses showed that 545 of the cells had 40 chromosomes that were karyotypically normal, 234 had 41 chromosomes, and 5 had 42 chromosomes. The cells with 42 chromosomes were individually trisomic for chromosome 8, 10, 12, 17, and the Y chromosomes, in addition to being trisomic for chromosome 15. Of the total 239 hyperdiploid cells, 220 were trisomic for chromosome 15. Structural abnormalities were rare. Therefore, among 20 cases studied, 12 showed a trisomic 15 cell line coexisting with a karyotypically normal line, although the proportion of the former varied from case to case. These findings agreed with those of earlier studies, except for the rarity of structural abnormalities. The significance of trisomy 15 in the development of leukemia is unknown. (9 refs)

78-5910 Secondary Malignant Neoplasms in Patients with Hodgkin's Disease. (Eng) Neufeld, H. (Univ. Manitoba, Winnipeg, Manitoba, Canada); Weinerman, B. H.; Kemel, S. *JAMA* 239(23): 2470-2471; 1978.

A retrospective analysis of 232 cases of Hodgkin's disease (HD) treated at the University of Manitoba between January 1, 1965 and December 31, 1973, revealed secondary malignant neoplasms in five patients. The average follow-up was 44.3 mo. Forty-nine patients had total nodal radiotherapy; 75, upper mantle; 12, inverted Y; 60, both upper mantle and inverted Y at different times in their course; and 34, local radiotherapy alone. A total of 174 patients who received radiotherapy subsequently received chemotherapy with mechlorethamine hydrochloride, vincristine sulfate, procarb-

zine hydrochloride, and prednisone. The five secondary cancers included two cases of acute leukemia (expected incidence was 0.05); both of these patients had received radiotherapy and chemotherapy. The other tumors included one glioblastoma multiforme, one squamous cell carcinoma of the skin, and one prostatic cancer. All neoplasms developed in patients with the mixed cellularity type of HD; this group represented only 28% of all HD cases. Although the therapy may have increased the leukemia incidence in these patients, the frequency was not high enough to abandon combined modality treatment. (11 refs)

78-5911 Systemic Lupus Erythematosus and Lymphoma. (Eng) Green, J. A. (Dept. Medicine and Therapeutics, Univ. Aberdeen, Aberdeen, Scotland); Dawson, A. A.; Walker, W. *Lancet* 2(8093): 753-756; 1978.

Four patients with established systemic lupus erythematosus (SLE) in whom lymphoma subsequently developed are described. All four patients were women ranging in age from 21 to 56 yr at the onset of SLE. Hodgkin's (2 cases) or non-Hodgkin's (2 cases) lymphoma was diagnosed 2 mo to 13 yr after the onset of SLE. An association between the two diseases was previously reported in 14 patients, 6 of whom developed lymphoma either before or at the same time as the autoimmune disease. The 12 patients in whom SLE preceded lymphoma were women with a mean age of 36 yr at the first SLE symptom and a mean age of 46 yr at diagnosis of lymphoma. The common factor underlying the association between the two diseases could be immunodeficiency. The primary lesion in SLE is probably suppressor T-cell dysfunction, which would predispose to abnormal B-cell proliferation in response to an extrinsic antigen or autoantigen. Early biopsy of suspect lymph nodes is recommended for SLE patients. (26 refs)

78-5912 Isolated Hepatic Metastasis from Renal Vein Leiomyosarcoma (Letter to Editor). (Eng) Radhakrishnan, J. (Chicago, IL); Alrenga, D. P.; Ghosh, B. C. *Arch Pathol Lab Med* 102(11): 606; 1978.

An unusual case of liver metastasis from a renal vein leiomyosarcoma occurred in a 93-yr-old woman who had undergone nephrectomy for the primary tumor 5 yr previously. A left hepatic lobectomy was performed; the right lobe was tumor-free. Pathological examination of the resected tissue confirmed the diagnosis of metastatic leiomyosarcoma. (1 ref)

78-5913 Ultrastructural Findings in a Pituitary Tumor Associated with Forbes-Albright Syndrome. (Jpn) Miyagami, M. (Dept. Neurological Surgery, Nihon

Univ., Sch. Medicine, Itabashi-ku, Tokyo 173, Japan); Nakamura, S.; Moriyasu, N. *Nippon Rinsho Denshi Kenbun* 10(3/4): 235-243; 1977.

Several ultrastructural characteristics were studied in a pituitary tumor associated with the Forbes-Albright syndrome. In the tumor cells, secretory granules with a single limiting membrane were markedly irregular in shape and averaged 500-800 m μ in diameter. Secretory granules appeared in the Golgi zone or near the periphery of the cell. The Golgi complex was prominent and enlarged, as were the mitochondria. The rough endoplasmic reticulum was oriented to be located near the periphery of the cell. There were many processes with microvilli on the cell membrane, and myelinlike bodies were found in the cytoplasm or intercellular space. (15 refs)

78-5914 Hyperparathyroidism, Chemodectoma, Thymoma, and Myasthenia Gravis. (Eng) Palmer, J. (Southern California Permanente Medical Group, 900 Sierra Ave., Fontana, CA, 92335); Sawyers, T. M. *Arch Intern Med* 138(9): 1402-1403; 1978.

Two patients with hyperparathyroidism, one (a 52-yr-old man) with an associated chemodectoma and the other (a 57-yr-old man) with an associated thymoma and myasthenia gravis, are described. These cases may represent an incomplete form of multiple endocrine neoplasia (MEN) syndrome type IIa and MEN type I, respectively. (11 refs)

See also:

*(Rev.): 78-5405, 78-5420, 78-5421, 78-5422, 78-5435, 78-5436, 78-5453, 78-5454, 78-5456, 78-5457, 78-5458.

*(Chem.): 78-5471, 78-5475, 78-5484, 78-5486, 78-5488, 78-5489, 78-5496, 78-5498, 78-5506, 78-5537, 78-5542, 78-5569, 78-5581, 78-5583, 78-5584, 78-5591, 78-5640, 78-5642, 78-5658, 78-5660, 78-5686, 78-5688, 78-5689.

*(Phys.): 78-5694, 78-5695, 78-5697, 78-5711, 78-5714, 78-5717, 78-5730.

*(Viral): 78-5751, 78-5794, 78-5804, 78-5807, 78-5851.

*(Immun.): 78-5854, 78-5855, 78-5861.

*(Epid.-Biom.): 78-5918, 78-5925, 78-5927, 78-5935, 78-5936, 78-5937, 78-5961, 78-5963, 78-5965, 78-5966.

EPIDEMIOLOGY AND BIOMETRY

5915 **Cancer Epidemiology.** (Spa) Carda Aparici, P. (Instituto Nacional de Oncologia, Ciudad Universitaria, Madrid, Spain). *Rev Mex Cir Ginecol Cancer* 5(5): 140-155; 1977.

Cancer epidemiology data are presented for Spain. The cancer mortality rate was 118/100,000 in 1960 and 143/100,000 in 1973. Men accounted for 52% of the total in 1960 and for 51% in 1973. In 1960, cancer mortality was highest in northern Spain, lowest in the south. In 1972, mortality was highest in the northeast. In men, stomach cancer mortality accounted for 19.2%, lung cancer 18%, prostate cancer 8.7%, liver cancer 7.2%, laryngeal cancer 5.2%, lymphoma and leukemia 4.9%, bladder cancer 4.9%, colon cancer 4.8%, and rectal cancer for 2.8% of the total cancer mortality. In women, stomach cancer accounted for 17.4%, breast cancer 12%, lung cancer 11%, uterine cancer 9.8%, colon cancer 7.9%, lymphoma and leukemia 5%, lung cancer 4.6%, and rectal cancer for 3.2% of the total. (140 refs)

5916 **Cancer Mortality in the State of Sao Paulo, Brazil.** (Por) Pastorelo, E. F. (Departamento de Epidemiologia, Faculdade de Saude Publica da USP, Avenida Dr. Arnaldo, 715, Sao Paulo, Brazil); Gotlieb, S. L. *Rev Saude Publica* 12(1): 1-15; 1978.

The distribution, during 1970-1972, of cancer mortality according to the main anatomical sites was determined for the administrative regions of Sao Paulo, Brazil. Cancer mortality averaged 8.42% in men and 8.80% in women, and regional differences were a function of industrialization. (9 refs)

5917 **Morbidity due to Head and Neck Malignant Tumors in Vinnitsa Oblast, USSR, During 1960-1974.** (Rus) Verigina, A. D. (Dept. Hosp. Surgery, Medical Inst., Vinnitsa, USSR). *Zh Ushn Nos Gorl Bolezn* 1: 57; 1978.

Analysis was made of the incidence of head and neck malignant tumors in Vinnitsa Oblast, USSR, between 1960 and 1974. Morbidity due to head and neck tumors comprised 13% of the total cancer morbidity. Standardized morbidity rates (per 100,000 individuals) were 22.43 for skin tumors, 1.5 for cancer of the lip, 1.7 for cancer of the larynx, 0.54 for cancer for the tongue and oral mucosa, 0.48 for cancer of the jaws, 0.33 for cancer of the pharynx, and 0.27 for cancer of the salivary glands. Morbidity was max in the 50-69 yr age group. Men had a significantly higher incidence of cancer of the lip (20.0 vs 3.27) and laryngeal (4.69 vs 0.13) cancer than women. (7 refs)

78-5918 **Relative High Frequency of Childhood Hodgkin's Disease in the North of Portugal.** (Eng) Sobrinho-Simoes, M. A. (Laboratorio de Anatomia Patologica, Hospital de S. Joao, Porto, Portugal); Areias, M. A. *Cancer* 42(4): 1952-1956; 1978.

Sex and age factors and the histopathological characteristics of Hodgkin's disease were studied in 322 patients in northern Portugal, particularly the 35 patients < 10 yr of age. The male:female ratio in these 35 patients was 2:1. Mixed cellularity was present in 18 patients, nodular sclerosis in 10. The incidence of childhood Hodgkin's disease was 18.2% during 1930-1944, 7.5% during 1945-1959, and 10.8% during 1960-1974. From 1970-1974, however, the frequency decreased to 4.5%. Most children belonged to large families of low socioeconomic status. All but 1/14 patients whose clinical records were reviewed presented with cervical lymphadenopathy as the initial symptom. Nine of these patients had constitutional symptoms at the time of diagnosis, and seven were classified in Stages III and IV. These findings are similar to those previously described in some developing countries of tropical and subtropical regions. (18 refs)

78-5919 **Epidemiology of Lymphogranulomatosis (Hodgkin's Disease). A Review.** (Ger) Berndt, H. (Klinik für innere Medizin, Bezirkskrankenhaus, Dr.-Wilhelm-Kulz-Strasse, DDR-20 Neubrandenburg, E. Germany). *Arch Geschwulstforsch* 48(5): 435-444; 1978.

Epidemiological data on the occurrence of Hodgkin's disease (HD) in East Germany are presented. The incidence standardized for the European population was 3.2/100,000 in men and 1.8/100,000 in women during 1964-1966, and 3.5/100,000 in men and 2.1/100,000 in women during 1968-1972. The population aged < 20 yr accounts for about 10% of the cases. The overall incidence has a first peak of 3.5/100,000 in the age bracket 25-30 yr and a second in the age bracket 65-80 yr. The mortality runs a fairly parallel course. Epidemiological studies have demonstrated familial and space/time clustering, an association with the histocompatibility antigen, and certain occupational factors (increased incidence among woodworkers and persons exposed to benzene), but no influence of ionizing radiation. Primary multiple malignant neoplasms are strikingly frequent in patients with HD, which may also be due to immunosuppressive therapy. HD may not be a nosological entity but rather a clinical and morphological manifestation of two diseases; namely, infectious juvenile granuloma and the true neoplasm in the elderly. (116 refs)

78-5920 **Ethnic and Clinical Aspects of Chronic Lymphocytic Leukemia in Israel. A Survey on 288**

Patients. (Eng) Bartal, A. (Dept. Hematology, Hebrew Univ.-Hadassah Medical Sch., Jerusalem, Israel); Bentwich, Z.; Manny, N.; Izak, G. *Acta Haematol (Basel)* 60(3): 161-171; 1978.

The prevalence of chronic lymphocytic leukemia (CLL) among the large number of ethnic groups in Israel was studied, and clinical findings for each patient were reviewed. The study population consisted of 288 patients (159 men, 129 women) from five medical centers who were diagnosed during 1960-1970. Almost two-thirds of the patients were in their fifth and sixth decades of life at diagnosis. Most (80.5%) of the patients were of East European descent, 14.5% originated from Asia and Africa, and 5% were born in Israel. Furthermore, 63% of all the CLL patients were immigrants from Poland and Rumania. Based on the ethnic distribution of the adult population of Israel during the decade, it is suggested that CLL occurs more frequently in European-born Jews than in Asian and African Jews. Coombs-positive hemolytic anemia was found in only 15/204 patients tested; all but 2 were of East European origin. Estimation of serum immunoglobulin (Ig) levels in 68 patients revealed low IgG levels in 29.4%, low IgA in 61.2%, and low IgM in 51.5%; at least one of these Ig's was below normal range in 74%. The values tended to decrease during the course of the disease. An additional primary malignant tumor was found in 26/206 patients, all of East European origin. Most of these tumors were cancer of the skin and breast. A third malignant neoplasia was diagnosed in 6/26 patients. Hypogammaglobulinemia was more frequent in patients with additional primary malignant tumors and in patients with advanced disease. Possible reasons for the ethnic distribution of CLL in Israel are discussed. (32 refs.)

78-5921 Leukemia in the Guinea Savanna of Northern Nigeria. (Eng) Fleming, A. F. (Dept. Hematology, Ahmadu Bello, Zaria, Nigeria). In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 53-54; 1978.

From 1970 to 1976, the following types of leukemia were diagnosed in 136 patients in Zaria and Kaduna, Nigeria: acute (A) lymphatic (21 patients), A myeloid (28), adult chronic (C) myeloid (36), juvenile C myeloid (4), and C lymphatic (47). Men predominated except among rural C lymphatic leukemia patients <45 yr old: depression of cell-mediated immunity during pregnancy may render women in rural Africa susceptible to agents causing this type of leukemia. (6 refs)

78-5922 Environmental Factors in the Epidemiology of Lymphosarcoma. (Eng) Goldsmith, J. R. (Epidemiological Studies Lab., California State Dept. Health,

Berkeley, CA); Guidotti, T. L. *Pathology Annual* 12(part 4): 411-425; 1977.

Data collected from ongoing studies of the role of environmental factors in the epidemiology of lymphosarcoma are summarized. Geographic mortality analyses suggest that environmental exposure, including occupational exposure, contributes to the development of lymphosarcoma. Some types of farming and food processing and distribution jobs, painting and paper work, and electrical work may carry higher than normal risks. Proportional mortality studies suggest that some building trades and some types of clerical work may have an excess risk. Workers in the rubber industry may be at a high risk, but lymphatic and myeloid leukemia had higher mortality ratios. Solvent exposure in rubber workers appears to be related to excess mortality for a variety of lymphatic tumors. With electrical workers, exposure to solder fumes, actinic radiation and its products, and polychlorinated biphenyls (PCBs) used in transformers may play a role in the high risk of lymphosarcoma. Similarly, exposure of farm workers to arsenic-containing pesticides may be associated with higher risk. Based on rates for men in large occupational groups that seem unlikely to have specific exposure, it is possible to estimate that approx 33% of cases may have an environmental exposure. Examination of body burden indices for lead, arsenic, and other metals, and PCBs may facilitate the definition of environmental hazards. Compilations of occupational and exposure histories for cases of lymphosarcoma in cancer registries may produce further data on occupational exposures in lymphosarcoma. Social security disability data can also be used to observe any trends in incidence. (11 refs)

78-5923 Dilemma of Toxic Risks (Letter to Editor). (Eng) Hartford, W. H. (Dept. Chemistry, Belmont Abbey Coll., Belmont, NC). *Chem Eng News* 56(15): 5; 1978.

A more logical mathematical modeling of the cancer risk from low concentrations of carcinogens is advocated. Calculations of the approx hypothetical probability of cancer at various doses above background are tabulated according to four models. (no refs)

78-5924 Occupational Health and Safety in Metallurgical Plants. (Eng) Taylor, J. C. (Jan H. Reid and Associates Ltd., Oakville, Ontario, Canada). *Health Affairs* 71(794): 141-157; 1978.

The general mechanisms by which ore is processed in metallurgical plants are discussed, and practical solutions to occupational health hazards associated with these processes are outlined. Careful design of the plant can eliminate many of the hazards. Technology is available to meet realistic regulations to control emissions and provide a safe, healthy environment in these plants. (29 refs)

5925 **Incidence and Long Term Survival of Children with Intracranial Tumours Treated in Denmark 1959.** (Eng) Gjerris, F. (Univ. Clinic Neurosurgery, Hospitalet, Copenhagen, Denmark); Harmsen, A.; Jensen, L.; Reske-Nielsen, E. *Br J Cancer* 38(3): 442-451;

on tumor incidence and long-term survival of 533 Danish children <15 yr of age with intracranial tumors diagnosed during 1935-1959 are presented. The av annual tumor incidence was 21/10⁶ children/yr during the 25-yr period. Treatment consisted of complete macroscopic surgical removal in 35% of the cases, partial removal or, more infrequently, biopsy in 50%, and radiotherapy (usually postoperatively) in 36%. In 75% of the children receiving radiotherapy, the skin doses were between 2,500 and 6,000 rads. Operative mortality was 28% during the 25-yr period. Of the 345 children who survived >1 mo after surgery, 119 (34%) were alive in 1974. Thirty-five percent of children surviving >1 mo with supratentorial midline astrocytomas and 25% of those with hemispherical astrocytomas had a 25-yr survival. Only 5/19 children with a supratentorial ependymoma survived >20 yr. The long-term prognosis was poor for children with craniopharyngiomas, with many recurrences developing after 10 yr. Among children with infratentorial tumors, only those with cerebellar astrocytomas had a high long-term survival rate. Children with brain stem tumors, ependymomas, or medulloblastomas had a poor prognosis. There were no significant differences in long-term survival between patients with supratentorial vs infratentorial tumors or between children with midline vs lateral supratentorial tumors. The histological diagnoses in survivors who received radiotherapy were so heterogeneous that it was impossible to evaluate the influence of treatment on the long-term prognosis. (40 refs)

5926 **The Epidemiology of Intracranial Meningiomas in Women in Los Angeles County.** (Eng) Presburg, R.; Martin, S. (Univ. California, Los Angeles, CA). *Dissemination Int B* 39(5): 2238-2239; 1978. (no refs)

5927 **Aetiological Factors in Brain Tumours (Meeting Abstract).** (Eng) Stortebecker, P. (Stockholm, Sweden). *Acta Neurochir (Wien)* 42(3/4): 252; 1978. (no refs)

5928 **Cancer Epidemiology. Part IV: Lung Cancer.** (Eng) Harberg, J. (Wisconsin Clinical Cancer Center, 1900 University Ave., Madison, WI, 53705); Buchan, D.; Davidson, D. J. *Wis Med J* 77(9): 18; 1978.

risk factors for lung cancer (including socioeconomic status; cigarette smoking; certain occupations; and ex-

posure to asbestos, uranium, chromates, or radiation) are surveyed briefly. (12 refs)

78-5929 **Occurrence of Lung Cancer at a Special Steel Manufacturing Plant.** (Swe) Rottner, B. (Fagersta AB, Fagersta, Sweden). *Lakartidningen* 75(20): 2022-2023; 1978.

Lung cancer mortality among employees of Fagersta AB, a manufacturer of special high-chromium steel, during 1956-1976 was determined in a case-control study. Metallic and trivalent chromium occurs in various technological processes. The overall mortality in the town of Fagersta over the 10-yr period was 3,112, but there were only 23 deaths from lung cancer. Compared with the national av, lung cancer mortality is underrepresented in Fagersta, which may be due to the small number of inhabitants. Twelve of the 23 lung cancer patients had been employed by Fagersta AB; 6 were office workers. Twenty-two of the 42 controls had been employed by the steel plant. The findings show no influence of employment at the chromium steel plant on lung cancer mortality. (7 refs)

78-5930 **Incidence of Cancer in Patients with Active Pulmonary Tuberculosis.** (Jpn) Fukutome, T. (Fukuoka-Higashi Natl. Hosp., Koga-Cho, Fukuoka-ken, Japan). *Fukuoka Igaku Zasshi* 69(3): 98-102; 1978.

Cancer incidence in 1,358 pulmonary tuberculosis patients treated at a Fukuoka, Japan, hospital from January 1971 to April 1977 was evaluated by the person-year method. The incidence was 6.0 vs an expected incidence of 6.74. This difference was not statistically significant. In 13/14 patients, the interval between the diagnosis of pulmonary tuberculosis and cancer appearance was >5 yr. Apparently, tuberculosis lasting >5 yr has no suppressive effect on cancer development. (16 refs)

78-5931 **Lung Cancer in Workers in a Nickel Refinery.** (Eng) Kreyberg, L. (Inst. Pathology, Univ. Oslo, Oslo, Norway). *Br J Ind Med* 35(2): 109-116; 1978.

The occurrence of lung cancer among nickel workers was studied based on 44 cases of lung cancer in persons employed in a nickel refinery during 1927-1974. Among those who began their employment between 1927 and 1939 (Series I), 5 developed cancer after 17-22 yr and 13 developed cancer after 31-40 yr. The cases with the longest development time had the lowest exposure to nickel. Among those who began their employment after 1945 (Series II), 17 developed tumors after 4-19 yr and 7 developed cancer after 20-25 yr. At diagnosis, the mean age of the series I and II workers was 57.6 and 56.0 yr, respectively. Among the 32 subjects with Group I tumors

(epidermoid and small cell anaplastic carcinoma), 29 were smokers; among the 7 subjects with Group II tumors (adenocarcinomas and other histological types), only 3 were smokers. These data suggest that tobacco smoking is an important additional factor in lung cancer in nickel workers. Among the employees not exposed to nickel, there was one with a Group I tumor and one with a Group II tumor. This study suggests that the risk of lung cancer may be higher in nickel workers than in the general population. (13 refs)

- 78-5932 Air Pollution and Cancer in Houston, Texas: A Causal Relationship?** (Eng) Marmor, M. (Inst. Environmental Medicine, New York Univ. Medical Center, 550 First Ave., New York, NY, 10016). *J Am Med Wom Assoc* 33(6): 275-277; 1978.

Respiratory cancer mortality rates from a previous study of cancer mortality and air pollution in Houston, Texas, were recast to allow investigation of possible confounding due to socioeconomic status and occupation. Cross tabulation of white male respiratory cancer mortality rates by air pollution categories and income levels for 1965-1969 yielded data that were consistent with a very strong health effect of air pollution. However, a similar recasting of the female mortality rates revealed a lack of association between air pollution levels and mortality rates. A synergistic effect between occupational and residential air pollution exposures could explain the association in men and the lack of an association in women. The possibility that geographic smoking patterns may have created a spurious relationship between air pollution and respiratory cancer cannot be eliminated. Selective migration and current air pollution patterns (compared with those of 20 yr ago) may also be confounding variables. In conclusion, it is difficult to argue that air pollution is responsible for the patterns of respiratory cancer mortality in Houston or any other city. (5 refs)

- 78-5933 Identification of Lung Cancer Risk Groups by Directed Mass Screening.** (Hun) Abraham, E. (Tudogondozo Intezet, Fovaros Bajcsy-Zsilinszky Korhaz-Rendelointezet, Budapest, Hungary). *Orv Hetil* 119(17): 1033-1036; 1978.

The results of a 1975 mass screening of 10,339 men and 10,089 women for lung tumors are presented. The mass screening was complemented by a computer-evaluated questionnaire on risk factors (age > 40 yr, smoking habit, occupational and other exposure, radiological change on previous chest x-rays, 2 or more respiratory complaints, and major surgery within the last 3 yr). During the first 1.5 yr following the mass screening, 78 cases of lung cancer were detected in the screened population (64 in men, 14 in women; incidence 619/100,000 in men, 107/100,000 in women). Among men, the incidence was 405/100,000 for those with negative previous chest x-rays, 1,069/100,000 for those with minor radio-

logical changes, and 915/100,000 for those with major x-ray changes. The corresponding incidences in women were 54/100,000, 157/100,000, and 403/100,000. In the risk group (changes on previous x-rays, smoked > 100,000 cigarettes), incidence was 1,342/100,000 in men and 985/100,000 in women. Additional harmful occupational exposures caused the incidences to be 1,105/100,000 in men and 1,093/100,000 in women. In the no-risk group (negative previous x-ray, work in unpolluted air), the lung cancer incidence was 0/1,124 in men and 1/5,129 in women. The findings indicate that the difference in lung cancer incidence between men and women is due to differences in smoking habits and occupational exposure; 10% of the men and 40% of the women screened were in the no-risk group. (13 refs)

- 78-5934 Occupational Distribution of Patients with Lung Cancer and Those with Chronic Bronchial Obstruction, Bronchial Asthma, and Sarcoidosis in Bern, Switzerland.** (Ger) Scherrer, M. (Pneumologische Abteilung, Medizinische Universitätsklinik, Inselspital, CH-3010 Bern, Switzerland); Zeller, C.; Zweifel, J. *Schweiz Med Wochenschr* 107(46): 1656-1661; 1978.

The occupational distribution of 553 patients with lung cancer, 233 with chronic bronchial obstruction, 1,397 with bronchial asthma, and 217 with sarcoidosis was studied with respect to smoking habits, degree of physical labor, and occupational and climatic exposures. Statistical analysis showed a definite prevalence of subjects who worked indoors among patients with sarcoidosis and bronchial asthma, regardless of smoking habits. In contrast, chronic obstructive lung disease and, especially, lung cancer seemed to occur predominantly in subjects whose jobs involved hard physical labor and heavy inhalative and climatic influences, regardless of smoking habits. Indoor workers tended to be spared from lung cancer, as were workers in the metals, synthetics, and chemical industries. An increased incidence of lung cancer was seen among outdoor workers, such as construction and farm workers, who are exposed to exhaust gases and cigarette and cigar smoke while performing hard physical labor. Because of continuous physical exertion, their breathing is deep and frequent, and the effect of hazardous air pollution may thus be considerably increased. (24 refs)

- 78-5935 Pneumoconiosis and Lung Cancer in Japan--Statistical Analysis of 1,172 Autopsied Cases During 1958-1974.** (Jpn) Suzuki, T. (Dept. Pathology, Sapporo Medical Coll., Sapporo-shi, Hokkaido 060, Japan); Kikuchi, I.; Okuda, M. *Jpn J Cancer Clin* 24(3): 179-181; 1978.

Cases of pneumoconiosis recorded in the Annual of Pathological Autopsy Cases in Japan during 1958-1974 (Group 1) were reviewed and compared with those occurring in patients autopsied at Iwamizawa Rosai Hospital during 1956-1974 (Group 2). Most of the silicotic patients on the island of Ho-

in Northern Japan were treated at this hospital where not all death cases are autopsied. Lung cancer was found in 179/1,172 Group 1 patients and in 49/327 Group 2 patients. The incidences of lung cancer in both groups were significantly higher than that in all men > 35 yr old autopsied in Japan during 1917-1971 and the lung cancer mortality rate was higher than the rate for all men > 35 yr old during 1967-1971. Noteworthy that the incidence of lung cancer in Groups 1 and 2 was almost identical (15.3%, 15.0%). The results suggest that there is a significant correlation between silicosis and lung cancer. (19 refs)

5936 **Epidemiologic Study of Precancer and Cancer of the Larynx.** (Bul) Dukhlev, S. (Regional Oncological Dispensary, Stara Zagora, Bulgaria). *Onkologiya* 1978; 8-12; 1978.

Factors were evaluated in 234 patients with cancer of the larynx, 240 patients with precancer of the larynx, and 743 healthy controls. Residence in urban areas, smoking, alcohol consumption, and occupational exposure to various environmental pollutants increased the risk of cancer of the larynx. (19 refs)

5937 **Cancer of the Larynx and Environmental Factors.** (Rus) Mgaloblishvili, G. N. (Dept. Otorhinolaryngology, Res. Inst. Oncology, Tbilisi, USSR); Gogeliani, N. A.; Gogeliani, G. K.; Gobronidze, I. A.; Ivanidze, D. V.; Ivanidze, T. G. *Vopr Onkol* 24(5): 66-69; 1978.

The incidence of laryngeal carcinoma in the Georgian Republic of the USSR is discussed. Between 1971 and 1975, cancer of the larynx was diagnosed in 572 patients (2.4/10⁵). The group included 535 men and 37 women (420 men and 26 women were in the 40- to 69-yr-old age group). There were 121 patients with Stage I, 121 with Stage II, 405 with Stage III, and 23 with Stage IV cancer. Histologically, most of the patients had squamous cell carcinoma. When these patients were compared with 169 matched healthy controls, it was found that the patients consumed more hot food, onions, garlic, and alcohol and they had greater exposure to various chemicals, dust, and smoke. (12 refs)

5938 **Liver Cirrhosis, Tobacco, Alcohol, and Cancer among Blacks.** (Eng) Keller, A. Z. (Epidemiologic Res., Dept. Medicine, and Surgery, Veterans Affairs Medical Center, 151 G, 941 N. Capital St., NE, Room 6303, Washington, DC, 20431). *J Natl Med Assoc* 70(8): 575-580; 1978.

Tobacco use, alcohol consumption, and age were characterized in relation to distribution of cancers by anatomical site among male black cancer patients with (n = 32) and without (n = 73) cirrhosis. Noncirrhotic (NC) black and white male

cancer patients were also characterized with respect to these variables. All patients in the series were discharged from US veterans hospitals. Among the black patients, the mean age of the cirrhotics (C's) was significantly lower than that of the NC's. Esophageal cancer was more than three times as frequent but stomach cancer was only one-fourth as frequent in the latter. However, only the excess of liver and gallbladder cancer (42.83% vs 1.50%) and lack of prostatic (26.2% vs 22.10%) and pulmonary cancer (8.31% vs 32.28%) within the C group were statistically significant. Smoking patterns between the two groups were not significantly different, but only one-fifth as many C patients were nonsmokers, and notably more C's smoked ≥ 20 cigarettes/day (75% vs 59.1%). C's drank whiskey as well as other alcoholic beverages significantly more often than NC's and there was a significant deficit of nondrinkers and an excess of heavy drinkers in this group. In a comparison of the 73 black NC's with 374 white NC cancer patients, ratios of liver, stomach, and prostate cancer were more than twice as high for blacks. However, esophageal cancer occurred at a similar frequency in both groups. Significantly more black patients smoked < 20 cigarettes/day (23.9% vs 9.5%), and significantly more white patients smoked < 20 cigarettes/day (74.8% vs 60.6%). Although blacks and whites exhibited similar profiles of drinking behavior by quantity, whiskey drinkers were significantly more numerous among the blacks. Thus, factors associated with smoking and drinking (especially whiskey), if not these habits themselves, are probably related to the increased risks of blacks for stomach and liver cancer compared with NC whites and for esophageal and hepatic cancer compared with NC blacks. (18 refs)

78-5939 **Increases in Laryngeal Cancer in Britain and Australia in Relation to Alcohol and Tobacco Consumption Trends.** (Eng) McMichael, A. J. (Div. Human Nutrition, Commonwealth Scientific and Industrial Res. Organisation, Kintore Ave., Adelaide, S. Australia 5000). *Lancet* 1(8076): 1244-1246; 1978.

The recent increase in the incidence of esophageal and laryngeal cancer in Britain and Australia, particularly in younger women, is examined. The time trends in sex-specific mortality from cancers of the larynx and esophagus throughout this century, in both Britain and Australia, show strong correlations with alcohol consumption. Since World War II, per capita alcohol consumption has almost tripled in Australia and it has increased by 70% in Britain. The time trends in relative cancer mortality between the two countries follow the trend in relative alcohol consumption quite closely, with a lag of 15 yr. The recent greater percentage increase in laryngeal cancer risk in British women than men, especially for younger women, is consistent with a joint effect of alcohol and tobacco. In women, consumption of both alcohol and cigarettes has increased in the past 30 yr, whereas the consumption of cigarettes by men has not. These results provide evidence for the close relationship between alcohol consumption and cancer of the larynx. (25 refs)

- 78-5940 The Carcinoembryonic Antigen (CEA) in Urban Populations: Epidemiological Studies.** (Ger) Schlipkoter, H. W. (Medizinische Institut für Lufthygiene und Silikoseforschung, Gurlittstrasse 53, D-4000 Düsseldorf, W. Germany); Baginski, B.; Kramer, U. *Zentralbl Bakteriologie (Orig B)* 166(2/3): 136-143; 1978.

Serum carcinoembryonic antigen (CEA) levels were measured during 1975-1976 in 50-yr-old men living in various areas of North Rhine-Westphalia, West Germany. High CEA levels were correlated with the number of cigarettes smoked and with chronic bronchitis (CB); even nonsmokers with CB had significantly higher levels than those without the disease. There was also a positive correlation between CEA levels and length of residence (> 5 yr) in an industrialized city such as Duisburg, but not in a rural town such as Gummersbach. Repeated determinations in over the 2 yr showed that the high CEA values of smokers remained stable. A graphical representation of CEA values in Cologne residents demonstrated that levels were higher in those living in industrial areas than in those living in nonindustrial areas of the city. The feasibility of using only nonsmokers to evaluate the effect of air pollution on CEA levels is discussed. (9 refs)

- 78-5941 Preliminary Study of the Mortality due to Malignant Gastrointestinal Tumors in Different Districts of Krakow, Poland, in Relation to Dietary Habits.** (Pol) Jedrychowski, W. (Zakład Epidemiologii, Instytut Medycyny Społecznej AM, Kopernika 7, 31-501 Krakow, Poland); Popiela, T.; Elgiet, W.; Cholewka, K. *Pol Tyg Lek* 33(18): 725-729; 1978.

A preliminary study was made of the 1974 mortality due to gastrointestinal neoplasms in four different districts of Krakow, Poland, and its possible relationship to dietary habits. The standardized mortality rates (SMR) for all cancers were 131.1 (144.2 for men and 199.1 for women) in Krowodrza (1); 139.9 (159.1 for men and 122.4 for women) in Nowa Huta (2); 141.2 (154 for men and 129.6 for women) in Podgórze (3) and 202.7 (201.6 for men and 203.8 for women) in the center city (4). The corresponding SMR for gastrointestinal tumors were 60.7 (65.6 for men and 56.3 for women) in district 1, 74.8 (83 for men and 67.4 for women) in district 2, 52.5 (47.7 for men and 56.9 for women) in district 3, and 94.7 (108.1 for men and 82.4 for women) in district 4. Analysis of the types of food consumed showed that the differences in overall cancer mortality and in gastrointestinal cancer mortality cannot be explained unequivocally by dietary habits. The higher cancer mortality in the center city may be due to higher air pollution levels. (14 refs)

- 78-5942 Epidemiology of Gastric Adenocarcinomas.** (Fre) Audigier, J. C. (C.N.R.S. Laboratoire 005440, pavillon H, hôpital Edouard-Herriot, 1, avenue Rockefeller, 69374 Lyon Cedex 2, France); Lambert, R. *Rev Prat* 28(45): 3409-3417; 1978.

The risk of gastric cancer is higher in the elderly, in men, subjects with blood group A, and in subjects of the low socioeconomic classes. The incidence is distributed unevenly among various countries and within the same country, even in rural and urban areas. Dietary factors probably play a major role in the etiology of this cancer. (11 refs)

- 78-5943 Talc-coated Rice as a Risk Factor for Stomach Cancer.** (Eng) Stemmermann, G. N. (Kuakini Medical Center, 347 N. Kuakini St., Honolulu, HI, 96813); Kolonel, L. N. *Am J Clin Nutr* 31(11): 2017-2019; 1978.

The hypothesis that talc-coated rice (TCR) increases the risk of stomach cancer was tested by comparing the cancer incidence rates of indigenous Japanese, who do not consume TCR and Hawaiian-Japanese, who consume TCR almost exclusively. The results show that the Japanese in Japan have very high rates of stomach cancer but the Hawaiian-Japanese have intermediate rates. The additional finding that Filipinos in Hawaii have very low gastric cancer rates yet consume the greatest amounts of TCR convincingly demonstrated that there is no correlation between frequency of exposure to TCR and gastric cancer incidence. (14 refs)

- 78-5944 Precancerous Lesions of the Stomach in Costa Rica.** (Spa) Salas, J. (Servicio de Patología, Hospital San Juan de Dios, San Jose, Costa Rica). *Patología (Mex)* 15(2): 63-79; 1977.

Lesions of the gastric mucosa were studied in 500 stomachs from 500 consecutive autopsies performed on subjects born in different provinces of Costa Rica. The patients included 254 males and 246 females aged 0-45+ yr. Intestinal metaplasia of the gastric mucosa was found in 247 patients (123 males and 117 females). The frequency of intestinal metaplasia was 9/91 in patients aged 0-14 yr, 21/54 in those aged 15-29 yr, 33/64 in those aged 30-44 yr, and 184/291 in those aged ≥45 yr. The frequency was higher in subjects born in high gastric cancer risk areas (central valley: 137/249; those born in low gastric cancer risk areas (major cities, highlands and coastal areas: 91/225), and it remained unchanged for subjects who were born in high-risk areas and then moved to low-risk areas. Localities with a higher prevalence of metaplasia showed a higher incidence of gastric cancer as well. Since intestinal type gastric cancer is predominant in Costa Rica, these findings support the hypothesis that intestinal metaplasia of the gastric mucosa is a precancerous lesion. (14 refs)

- 78-5945 Breast Cancer and Diet among the Japanese in Hawaii.** (Eng) Nomura, A. (Japan-Hawaii Cancer Study, 347 N. Kuakini St., Honolulu, HI, 96817); Folsom, B. E.; Lee, J. *Am J Clin Nutr* 31(11): 2020-2025; 1978.

evaluate the hypothesis that breast cancer may be associated with the intake of dietary fat, cholesterol, and animal protein, the dietary history of 86 Hawaiian-Japanese men whose wives had breast cancer was compared with that of 10 Japanese men whose wives did not have the disease. Due to the lack of a standardized method of dietary data collection, three different techniques were employed: (1) comprehensive 24-hr dietary recall, (2) weekly food frequency survey, and (3) an estimate of the weekly quantitative consumption of various food items. The results were consistent from one method to another and indicated that, although the two groups did not differ significantly in daily nutrient intake, breast cancer spouses consumed more American foods such as beef, pork, butter, margarine, cheese, corn, and wieners. In contrast, the control group consumed more Japanese items, such as green tea, Japanese soups, nori, and other seaweeds. It is concluded that further study of the American diet may be vital in relating nutritional factors to breast cancer. (16 refs)

5946 **Epidemiology of Breast Cancer with Special Reference to the Role of Diet.** (Eng) Hirayama, T. (Epidemiology Div., Natl. Cancer Center Res. Inst., 5-1-1 Chome, Chuo-ku, Tokyo, Japan). *Prev Med* 7(2): 151-155; 1978.

The low incidence of breast cancer in Japan, the recent increase of this disease (especially in women aged 45-59 yr), and the role of dietary fat intake and body size in these developments are discussed. Epidemiological studies of 142,857 women aged >40 yr indicated that the risk was 8.5 times higher in women of high socioeconomic strata who eat meat daily than in women of low socioeconomic strata who do not eat meat daily. A high positive correlation was noted between per capita fat intake and adjusted breast cancer death rates in different districts of Japan. It was estimated that the breast cancer death rate will rise to the US level when Japanese dietary fat intake approaches current US levels. The close relation with fat intake resulted mainly from the consumption of pork and animal fat. The recent increase in breast cancer death rates was also found to be under the combined influence of the daily intake of animal fat and 2-furyl-3-(5-nitro-2-furyl)acrylamide, a highly mutagenic food additive widely used in Japan from 1965 to 1975 and known to produce mammary carcinoma in rats. A series of case-control studies revealed an increased risk of breast cancer with increasing body size, especially in postmenopausal women. The recent breast cancer increase could therefore be a reflection of the fact that Japanese women are becoming heavier, especially after age 30. (13 refs)

5947 **Barrier Contraceptive Practice and Male Infertility as Related Factors to Breast Cancer in Married Women. Preliminary Results.** (Eng) Gjorgov, A. N. (Dept. Obstetrics and Gynecology, Hosp. Univ. Pennsylvania, 3400 Spruce St., Philadelphia, PA, 19104). *Oncology* 33(3): 97-100; 1978.

A case control study was conducted to test the hypothesis that a reduced exposure to human seminal factors in the early reproductive life of women is a risk factor in breast cancer. A total of 153 consecutive breast cancer patients, aged 35-60 yr, were matched by age, social status, and method of contraceptive practice with 168 controls. The relative risk of exposure to the hypothetical semen-factor deficiency was 4.7 times greater for the breast cancer patients than for the controls. When male infertility factors were excluded from the analysis, the risk of breast cancer in the same population in the US was estimated as five times greater for women who use barrier contraceptive methods (condom and other) than for women who use nonbarrier methods (diaphragm and other). The reduction in breast cancer incidence by eliminating the barrier contraceptive techniques would be not < 50% in married white women. (36 refs)

78-5948 **Significance of Hysterectomy in the Genesis of Breast Tumors.** (Rus) Ismailov, A. Kh. (Medical Inst., Blagoveshchensk, USSR); Kositsyna, M. V.; Palchik, E. A. *Akush Ginekol (Mosk)* (6): 56; 1978.

Of 232 women who underwent hysterectomy for uterine fibromyoma, 18.4% developed mastopathy within 5 yr of surgery and 37.9% developed mastopathy within 10 yr of surgery. (2 refs)

78-5949 **Menstrual Cycle Patterns and Breast Cancer Risk Factors.** (Eng) Wallace, R. B. (Dept. Preventive Medicine, Univ. Iowa, Iowa City, IA, 52242); Sherman, B. M.; Bean, J. A.; Leeper, J. P.; Treloar, A. E. *Cancer Res* 38(11, part 2): 4021-4024; 1978.

The relationship between abnormal menstrual cycle patterns and age at menarche, menopause, and first parity, three variables related to breast cancer incidence, was examined using a data set of women who, beginning in 1934, recorded menstrual, contraceptive, reproductive, and related medical events through life. Many daughters of the original subjects were also enrolled and, to date, nearly 5,000 women have participated. The results include data collected through 1975. Subjects taking oral contraceptives or with any gynecological surgery that might interfere with cycle patterns were excluded. Women with late onset of menarche had longer and more variable cycles in the 10 yr after menarche than those with early onset. Cumulative fertility in women after marriage did not differ according to cycle length and variance. Late menopause may be a breast cancer risk factor because of relative estrogen excess and progesterone lack, as reflected in longer, more varied cycle patterns. Late menopausal age is also a risk factor for endometrial cancer, which may also be related to persistent estrogen stimulation. In this analysis, only simple measures of cycle patterns, means, and variance were examined. A single or small number of episodes of prolonged amenorrhea may possibly be of more pathophysiological importance than deviations from average cycle length of, for example, 1 wk. (12 refs)

- 78-5950 Dormancy and Spontaneous Recurrence of Human Breast Cancer In Vitro.** (Eng) Yuhas, J. M. (Cancer Res. and Treatment Center, Univ. New Mexico, 900 Camino de Salud, N.E., Albuquerque, NM, 87131); Tarleton, A. E. *Cancer Res* 38(11, part 1): 3584-3589; 1978.

The hypothesis that hormone-dependent tumor cells survive endocrine therapy in a viable but nongrowing state, adapt to the absence of hormones genetically or physiologically, and then resume growth was demonstrated in vitro using a cell line derived from a brain metastasis of a human mammary adenocarcinoma (MDA-361). The responses of monolayers and multiple tumor spheroids (MTS's) of the MDA-361 line were compared in two types of media: one in which exogenous insulin (I: 10 µg/ml) had been added (I+) and one in which the only I present was that remaining after dialysis (I-). The monolayer cultures required I for growth and for maintenance of viability, as evidenced by the rapid and complete degeneration of the cells after they were transferred to the I- medium. Detachment occurred within 24-48 hr, and the few cells that remained attached (<0.1%) doubled every 3-4 wk. Upon transfer to I- medium, the MTS's became less dependent on and more responsive to this hormone and, after a dormant period of approx 6 wk, eventually resumed growth. During culture in I- medium, the MTS's appeared normal but reduced in size (max of 12%-40% loss of volume). The spontaneous regrowth in the absence of I continued at least through 3 mo. In terms of overall kinetics, the MTS's paralleled the behavior of residual tumors in vivo. In both cases, progressive tumor growth occurs in the presence of a specific growth factor (ie, I), shrinkage occurs upon its removal, stability persists for a prolonged period, and then growth resumes. (22 refs)

- 78-5951 Reproductive Factors and Risk for Breast Cancer in Iceland.** (Eng) Tulinius, H. (Icelandic Cancer Registry, P.O. Box 523, 101, Reykjavik, Iceland); Day, N. E.; Johannesson, G.; Bjarnason, O.; Gonzales, M. *Int J Cancer* 21(6): 724-730; 1978.

The risk of breast cancer was compared with various aspects of a woman's reproductive history in a prospective study of 31,453 Icelandic women born between 1906 and 1945. Among these subjects, 216 had developed breast cancer by 1975. Age at menarche had a slight influence on breast cancer risk (increased risk associated with earlier menarche), and this influence remained after further adjustment for parity and age at first pregnancy. Risk also increased with increasing age at first pregnancy and with nulliparity. The risk decreased with larger number of children, the effect being independent of age at first pregnancy. It is suggested that, given the linearity of increase in breast cancer risk with interval to first pregnancy, before a woman becomes pregnant her breast tissue is in a condition particularly susceptible to the action of some carcinogenic influence. Further, the mechanism of action of this putative carcinogenic influence appears to be

different from the mechanism mediating the increased susceptibility among certain populations. (24 refs)

- 78-5952 The Epidemiology of Epithelial Ovarian Cancer in Women Aged 25-49.** (Eng) Casagrande, C. (Univ. California, Los Angeles, CA). *Diss Abstr Int B* 3: 2235-2236; 1978. (no refs)

- 78-5953 Epidemiological Study of Endometrial Cancer.** (Jpn) Masubuchi, K. (Dept. Gynecology, Cancer Inst. Hosp., Toshima-ku, Tokyo 170, Japan); Nemoto, T. *Gan No Rinsho* 24(8): 766-771; 1978.

Examination of 197 endometrial cancer patients at the Cancer Institute Hospital, Tokyo, Japan, during 1970-1974 revealed that women at high risk for endometrial cancer were those >50 yr of age (3.4% were <39 yr old), nulliparous, having a max blood pressure >160 mm Hg, weighing >70 kg (7.3% weighed <39 kg), and complaining of genital bleeding. The diagnostic accuracy was 81.2% using endometrial aspiration cytology. Among 13 cases diagnosed as noncancerous at first biopsy, 8 were suspected of being malignant on aspiration cytology. (23 refs)

- 78-5954 Mortality of Female Workers in a Rubber Manufacturing Plant.** (Eng) Andjelkovich, J. (Occupational Health Studies Group, Sch. Public Health, Dept. Epidemiology, Univ. North Carolina, Chapel Hill, NC 27514); Taulbee, J.; Blum, S. *J Occup Med* 20(6): 409-413; 1978.

The 10-yr mortality experience of a cohort of white female production workers in a large rubber manufacturing plant was examined. The cohort consisted of 1,649 white female production workers who were 40-84 yr old on January 1, 1964, and either active or retired as of the same date; 99% of them had at least 10 yr employment service within the plant under study. An analysis of the observed and expected numbers of death from numerous causes revealed a slightly elevated overall standardized mortality ratio, with an excess of deaths from both lung cancer and myocardial infarction and a deficit of deaths from breast cancer. There were no deaths due to cancer of the buccal cavity and pharynx, cancer of the liver and intrahepatic bile ducts, Hodgkin's disease, or leukemia. About two-thirds of all the women had worked for the longest period of their employment, in the manufacture of industrial products; the remainder were engaged mostly in the manufacture of tires. A more detailed analysis investigation is required before nonoccupational risk factors such as smoking, can be ruled out. (20 refs)

- 78-5955 Cervical Cancer in Younger Women.** (Eng) Andrews, F. J. (Dept. Clinical Cytology, Brigham Young Univ., Salt Lake City, UT 84143).

eral Hosp., Brighton, Sussex BN2 3EW, England); Line-
J. J.; Melcher, D. H. *Lancet* 2(8093): 776-778; 1978.

incidence of cervical cancer was studied among 21,216
hton, England, women from whom cervical smears were
ined between May 1974 and May 1975. Of 134 patients
hom cervical cancer was detected and histologically con-
ed, 78 were < 35 yr old and 4 were < 20 yr old. Most
of the tumors were in situ carcinomas. Abnormal
ars and histologically malignant cervical cancers peaked
ges 26 to 30; however, abnormal smears and cancer were
id in 24 patients aged 16-19 yr. The mean age of Brighton
ents with cervical carcinoma was 50 yr in 1967 and 40
n 1974. Until the etiology of this cancer is established, all
ng women should be encouraged to have cervical smears.
(19 refs)

5956 **Mortality from Carcinoma of Cervix Uteri in
Britain.** (Eng) MacGregor, J. E. (Cytology Unit,
ot. Pathology, Univ. Aberdeen, Aberdeen, Scotland); Tep-
S. *Lancet* 2(8093): 774-776; 1978.

e-related trends in the incidence of mortality from cervical
cer in England, Wales, and Scotland were studied. For
men aged 35 yr or more, mortality from this cancer de-
ed in Britain and Wales between 1968 and 1976. A similar
nd was observed in Scotland, and the death rates were
er for two areas with established screening programs
rmpian and Tayside) than for the rest of the country.
wever, the mortality rates among English and Welsh
men < 35 yr old increased between 1968 and 1976. A simi-
pattern was observed for Scotland when the data for
mpian and Tayside were disregarded. Until cervical can-
is better understood, broad-based screening, as practiced
these two areas of Scotland, is probably justified. (10 refs)

5957 **Hysterectomy-Rates and Cervical Cancer (2
Letters to Editor).** (Eng) Rhodes, P. (35 Elm-
ld Road, Newcastle upon Tyne NE3 4BA, England);
acgregor, J. E. *Lancet* 2(8097): 1000; 1978.

he fall in cervical cancer mortality might be partly due to
increasing hysterectomy rate and an attendant decrease
the population at risk. However, in the sample previously
ported, there was no change in the number or age-distribu-
on of hysterectomies. Most were done on older women of
pper social class; this group is at relatively low risk for cervi-
l cancer. (no refs)

8-5958 **Adenocarcinoma of the Uterine Cervix in Jew-
ish Women. A Distinct Epidemiological Entity.**
(Eng) Menczer, J. (Dept. Obstetrics-Gynecology, Chaim
neba Medical Center, Tel Hashomer, Israel); Modan, B.;

Oelsner, G.; Sharon, Z.; Steintiz, R.; Sampson, S. *Cancer*
41(6): 2464-2467; 1978.

To determine whether the etiology of adenocarcinoma of the
uterine cervix (AUC) differs from that of squamous cell car-
cinoma (SUC), the records of 532 Israeli Jewish women with
cervical cancer diagnosed between 1961 and 1971 were exam-
ined. The mean annual incidence of invasive cervical cancer
among Israeli Jews during the 11-yr period was 4.2/100,000.
AUC was diagnosed in 52 women, SUC in 441, and cancers
of other histological subcategories in the remainder. The rela-
tive frequency of AUC increased steadily with age, from 6%
among those aged 20-39 yr to 15% in those aged 70 yr or
more. The incidence of SUC was significantly higher in wom-
en born in North Africa than in those of other ethnic groups.
However, there were only minimal ethnic differences in the
incidence of AUC. The incidence of AUC decreased by 50%
during 1966-1971, compared with the incidence during
1961-1965, and the incidence of SUC decreased by 24%. The
findings support the contention that the etiologies of AUC
and SUC are different. (19 refs)

78-5959 **Role of Cu-IUD in Cervical Carcinogenesis: A
Follow Up of 36 Months.** (Eng) Luthra, U. K.
(Cytology Section, Indian Registry Pathology, ICMR,
Maulana Azad Medical Coll., New Delhi-110002, India); Mi-
tra, A. B.; Prabhakar, A. K.; Bhatnagar, P.; Agarwal, S. '
Indian J Med Res 68: 78-83; 1978.

Cytological monitoring of the female genital tract was per-
formed in 2,603 Indian women who were registered as having
had copper-containing intrauterine devices (IUD) inserted.
Detailed cytologic analyses were made of 2,244 women who
had adequate preinsertional exocervical smears. The rate of
dysplasia in these women at the onset of use of the IUD was
2.5%; at 12, 24, and 36 mo, the rates were 2.5%, 2.3%, and
6.3%, respectively. No difference in the rate of dysplasia after
the use of the IUD was observed in women who had either
negative or inflammatory initial smears. Among 59 dysplasia
cases followed, 49 regressed to a normal condition and 10
persisted as dysplasia. No dysplastic lesion progressed to
higher grades of dysplasia or to carcinoma during the study
period. The above data indicate that continued use of cop-
per-containing IUD for up to 36 mo produces no adverse
effect on female genital tract epithelium. (13 refs.)

78-5960 **Risk Factors in Scrotal Epithelioma.** (Eng) Fay,
H. T. (Works Medical Office, Renold Ltd.,
Didsbury, Manchester 20, England). *J R Soc Med* 71(10):
741-747; 1978.

The occurrence of 6 cases of squamous cell carcinoma of the
scrotum, 5 fatal, plus 10 cases of forearm keratoses in em-
ployees of a light engineering factory between 1955 and 1975

prompted an investigation of possible occupational risk factors responsible for this unusual concentration. The length of employment ranged from 17-38 yr. The keratoses, which are considered a premalignant condition, developed due to exposure of the forearms to mineral oil. These men should be regarded as having a higher risk of developing scrotal cancer. The use of neat cutting oils and the use of oil reclamation plants at the factory are implicated in the scrotal cancer cases. These plants cannot be expected to remove any polycyclic aromatic hydrocarbons (PAH) formed during prolonged use of neat oils. Results of analyses of new, used, and reclaimed oils for PAH in three different laboratories are given. They show that the prolonged use of neat cutting oils and of conventional oil reclamation plants leads to a dangerous accumulation of carcinogens such as benzo(a)pyrene. Other possible risk factors considered are genetic factors, personal hygiene, protective clothing, and the high ambient temperatures in the machine shops. (26 refs)

- 78-5961 Epidemiologic Characteristics of Urinary System Tumors and Balkan Nephropathy in an Endemic Region of Bulgaria.** (Eng) Nicolov, I. G. (Inst. Oncology, Medical Acad., Sofia 1156, Bulgaria); Chernozemsky, I. N.; Petkova-Bocharova, T.; Stoyanov, I. S.; Stoichev, I. I. *Eur J Cancer* 14(11): 1237-1242; 1978.

The epidemiologic similarity between endemic nephropathy (EN) and urinary tract tumors (UTT) (especially neoplasms of the renal pelvis and ureter) in an endemic region of Bulgaria (Vratza district) was investigated. These two disorders correlated geographically, had high incidence rates, showed a familial aggregation, and predominated in women and middle-aged people. In hyperendemic villages, the annual age-adjusted incidence rates per 100,000 were 43.5 in men and 74.2 in women for tumors of the renal pelvis and ureter and 38.7 and 24.6, respectively, for tumors of the urinary bladder. UTT were often multiple, especially in EN patients, and generally originated in the uroepithelium. Among UTT patients, 27.3% also had EN, and UTT were found in 6.5% of EN patients. Among patients with both EN and UTT, the former occurred first in 51% of the cases vs the latter in 18% of the cases. It is concluded that the epidemiologic similarity between EN and UTT, their location in the urinary system, and the fact that they frequently occur in combination is proof of a close etiologic and pathological relationship. (23 refs)

- 78-5962 Etiological Factors Involved in Primary Hepatocellular Carcinoma in Abidjan (Ivory Coast).** (Fre) Le Bras, M. (Service de Clinique Medicale, C.H.U. de Treichville, Abidjan, Ivory Coast); Ticolat, R.; Clerc, M.; Loubiere, R.; Beda, B.; Bertrand, E. *Gastroenterol Clin Biol* 2(8/9): 679-688; 1978.

Etiological factors related to the development of hepatocellular carcinoma (HC) were analyzed in 252 Ivory Coast pa-

tients diagnosed in the period 1972-1974. HC represented 13.6% of liver diseases observed in patients hospitalized during this period. There was no significant correlation with etiologic factors (blood group or Hb type), season when admitted, alcoholism, or history of hepatitis (only 8%). The ratio of men to women was 5.4/1, compared with the overall hospital admission ratio of 1.5/1. Hepatitis B surface (HBs) antigen was found in the serum of 42% of the 50 HC patients, of whom it was measured, a higher percentage than that in patients with cirrhosis without HC or hepatitis. Cirrhosis was associated with HC in 77.3% of the patients. Geography appeared to be an etiologic factor since two-thirds of the patients lived in the savannah rather than forest areas. Patients who were workers or farmers from lower economic level outnumbered patients from other social and professional categories. It is concluded that environment may play a role in development of HC in the Ivory Coast. (41 refs)

- 78-5963 Occurrence of Cirrhosis and Primary Liver Cancer in an Eskimo Population Hyperendemic for Infected with Hepatitis B Virus (Meeting Abstract).** (Dan) Skinhoj, P. (Dept. Infectious Diseases, Rigshospitalet, Copenhagen, Denmark); Hansen, J. P.; Nielsen, N. H.; Kelsen, F. *Dan Med Bull* 25(5): 223; 1978. (no refs)

- 78-5964 Epidemic of Toxic Hepatitis in India of Possible Mycotoxic Origin.** (Eng) Tandon, H. D. (Dept. Pathology, All India Inst. Medical Sciences, New Delhi-110016, India); Tandon, B. N.; Ramalingaswami, V. *Pathol Lab Med* 102(7): 372-376; 1978.

An epidemic of liver disease characterized by onset with fever, rapidly progressive jaundice, and ascites occurred in a rural area of India. Several hundred people were affected and mortality was high. The epidemic was heralded by appearance of similar features in the village dogs. Liver biopsy specimens from eight patients and autopsy material from one patient and two dogs were studied histologically. Characteristic features were centrilobular scarring, hepatocellular necrosis, sinusoidal occlusion, ductular proliferation and cholestasis, syncytial giant cell transformation of hepatocytes, and cellular fibrosis. Toxic quantities of aflatoxin B₁ were found in corn samples from the local food stores. The etiology of the disease could not be established unequivocally, but aflatoxins, perhaps in combination with other factors, may have been the cause. (36 refs)

- 78-5965 Hepatoma Risk Among HBsAg Carriers (Meeting Abstract).** (Eng) Beasley, R. P. (Univ. of Washington, St. Louis, MO); Lin, C. C. *Am J Epidemiol* 108(3): 247; 1978. (no refs)

5966 **A Statistical Analysis of Mortality Due to Liver Cancer and Liver Cirrhosis in Yamanashi Prefecture, Japan, with Special Emphasis on Relationship to the Prevalence of Schistosomiasis.** (Jpn) Inaba, Y. (Dept. Hygiene and Preventive Medicine, Faculty Medicine, Univ. of Tokyo, Bunkyo-ku, Tokyo 113, Japan); Takahashi, E. Y.; Kouchi, N. *Jpn J Public Health* 24(12): 811-815; 1977.

1970-1975 mortality data for liver cancer and liver cirrhosis in Yamanashi Prefecture, Japan, were analyzed, and, in view of the fact that this prefecture contains an area endemic for schistosomiasis (SSM), an attempt was made to determine if the data were related to the occurrence of SSM. There were 704 deaths due to liver cancer during this period; however, the rate of primary liver cancer (4.7%) was significantly lower than the national rate (8.2%). There were 990 deaths due to cirrhosis, including 17 cases (1.7%) of alcoholic liver cirrhosis, a figure significantly lower than that for the nation (3.7%). The age-specific mortality rate for both liver cancer and liver cirrhosis was relatively high in the elderly, compared with the national figure, but mortality due to liver cirrhosis was relatively low for men in their fifties. During the study period, the geographic distribution of age-adjusted mortality due to both conditions correlated significantly with the prevalence of SSM in 1948-1952. The standardized mortality ratio (SMR) of liver cirrhosis in the area endemic for SSM was significantly high, but the ratio for the nonendemic area was almost the same as the national ratio. However, the SMR of liver cancer in the endemic area was approx 50% higher than that of the national figure. These findings indicate that the higher mortality due to liver cirrhosis in Yamanashi Prefecture is caused by SSM but that due to liver cancer is caused by SSM and other unknown etiological factors. (6 refs)

5967 **Angiosarcoma of the Liver in Vinyl Chloride/Polyvinyl Chloride Workers: 1977 Update of the NIOSH Register.** (Eng) Spirtas, R. (Illness Effects Section, SB/DSHEFS/NIOSH, Robert A. Taft Labs., 4676 Columbia Parkway, Cincinnati, OH, 45226); Kaminski, R. *J Occup Med* 20(6): 427-429; 1978.

Data on 64 cases of angiosarcoma among vinyl chloride (VC) polymerization workers reported as of October 1977 are summarized. The material comprises 23 US and 41 foreign cases from 11 different countries. The age at diagnosis ranged from 40 to 71 yr, with a median of 49 yr. The latency period ranged from 9 to 38 yr (median of 21 yr), duration of exposure from 1 to 31 yr (median of 18 yr). The data suggest that the report of VC-induced angiosarcoma cases gradually increased over time. The 20- to 30-yr lag between the growth of the industry and the World War II and the increase in the number of diagnosed angiosarcomas among VC/polyvinyl chloride (PVC) workers during the 1970's roughly coincides with the median latency period of 21 yr. Recent studies have also found VC to be associated with excesses in brain cancer and neoplasms of the respiratory and lymphatic systems. (6 refs)

78-5968 **Epidemiologic Investigation of Employees Chronically Exposed to Methylene Chloride. Mortality Analysis.** (Eng) Friedlander, B. R. (Health, Safety and Human Factors Lab., Eastman Kodak Co., Rochester, NY, 14650); Hearne, T.; Hall, S. *J Occup Med* 20(10): 657-666; 1978.

The 1964-1976 mortality experience of a relatively large male employee population that had been working with methylene chloride on a continuous basis for up to 30 yr was studied. The mortality of the study group was consistent with industrial controls and less than that in the New York State general population. The number of deaths due to malignant neoplasms was significantly less (7) than the number expected for industrial controls (12.5) and New York State controls (16.2). (47 refs)

78-5969 **Cohort Study of the Causes of Death in Ethylene Oxide Production Workers (Letter to Editor).** (Swe) Bengtsson, G. (Orebro, Sweden); Hogstedt, C.; Rohlen, O.; Berndtsson, B. S.; Axelson, O.; Ehrenberg, L. *Lakartidningen* 75(38): 3285-3287; 1978.

Causes of death were determined in 66 workers (Group A) with no exposure to ethylene oxide (EO), 86 workers (Group B) with occasional exposure, and 86 workers (Group C + D) who had been exposed to EO for at least 1 yr 10 or more yr earlier. Other substances to which the workers in Groups C + D and B were exposed were ethylenechlorhydrine (5 mg/m³), ethylene chloride (100 mg/m³), bis(2-chloroethyl)ether (0.05 mg/m³), and ethylene (600 mg/m³). The overall mortality was 23/89 in Group C + D, 14/86 in Group B, and 10/66 in Group A. Tumors were the cause of death in 9/89 in Group C + D subjects; they included gastric cancer (3 cases), cancer of the large intestine (1 case), malignant melanoma (2), kidney cancer (1), chronic lymphocytic leukemia (1) and acute myeloid leukemia (1). The expected tumor mortality was 3.6 for all tumors, 0.4 for gastric cancer, and 0.14 for leukemia. The differences between the expected and the observed incidence of gastric cancer and leukemia were statistically significant ($p < 0.01$). Tumors were the cause of death in 3/86 Group B subjects and 1/66 Group A subjects. The observed cancer mortality in these groups did not differ significantly from the expected mortality. The findings indicate a significant risk of gastric cancer and leukemia in the workers exposed to EO. (21 refs)

78-5970 **Cancer from Burn Scars.** (Por) Sucena, R. C. (Servico de Queimados do Hospital das Clinicas, FMUSP, Sao Paulo, Brazil). *An Paul Med Cir* 104(3): 203-226; 1978.

The malignant transformation of burn scars was studied in the 103 patients (68 men, 35 women) among 20,194 burn victims in whom this process occurred. The period between

the burn and tumor formation ranged from < 1 yr (1 patient) to > 61 yr (3 patients). Epidermoid carcinoma was diagnosed in 94 patients (Stage I in 20, Stage II in 49, Stage III in 11, and Stage IV in 14), epidermoid carcinoma + basal cell carcinoma in 1, epidermoid carcinoma + fibrosarcoma in 1, and fibrosarcoma in 7. The tumor was localized to the head in 6 patients, to the trunk in 25, to the upper limbs in 37, and to the lower limbs in 35. The time required for cicatrization of the burn wound was 1-3 mo in 28 patients, 4-6 mo in 23, 7-12 mo in 24, 13-60 mo in 19, and > 60 mo in 9. The initial thermal shock, slow cicatrization, and secondary trauma are believed to play etiological roles in the malignant transformation of burn scars. (20 refs)

- 78-5971 Concurrence of Bronchial Carcinoma and Anthracosilicosis in the Lung. Silicotic Carcinoma in Scar Tissue of the Lung.** (Ger) Bohm, E. (Pathologisches Inst., Hunscheidtstrasse 1, D-4630 Bochum, W. Germany). *Med Klin* 73(18): 659-663; 1978.

The existence of a causal relationship between anthracosilicosis and bronchial carcinoma was explored by evaluating the autopsy records of 3,452 miners with silicosis, 551 miners without silicosis, and 605 nonminers. Of the total of 4,608, there were 501 cases of bronchial carcinoma, 425 of which were associated with silicosis. The incidence of both afflictions was not related to the length of employment, and there was no correlation between the severity of anthracosilicosis and frequency of cancer incidence. Of the concurrent cases, 2.8% were diagnosed as silicotic carcinoma of the scar tissue, with another 6.8% as possible; only in these cases did a causal relationship seem probable. Histologically, 52.9% of the 425 carcinomas were diagnosed as the squamous epithelial type, 28.7% as the small cell type, and 2.1% as polymorphocellular carcinoma; 37.6% of the tumors metastasized to the adrenal gland, 35.3% to the liver, and 24.2% to the spine. (24 refs)

- 78-5972 Radiation-related Thyroid Carcinoma.** (Eng) Cerletty, J. M. (Milwaukee County General Hosp., 8700 W. Wisconsin Ave., Milwaukee, WI, 53226); Guansing, A. R.; Engbring, N. H.; Hagen, T. C.; Kim, H. J.; Shetty, K. R.; Rosenfeld, P. S.; Wilson, S. *Arch Surg* 113(9): 1072-1076; 1978.

Evaluatory studies were conducted on 1,825 subjects who had a history of head or neck irradiation. Thyroid abnormalities were found in 358 of these subjects: single or multiple thyroid nodules in 165, diffuse thyromegaly in 153, and previous thyroid surgery in 40. These 40 previous surgery patients were excluded from further study. Surgery was performed on 113 patients with thyroid nodules, and thyroid carcinoma was found in 34 (papillary adenocarcinoma in 17, mixed papillary-follicular type in 10, follicular type in 6, and a mixed papillary-follicular carcinoma in 1). Clinical examination

of the neck was the most valuable method of detecting abnormalities. Detection of nodules was not enhanced significantly by thyroid imaging studies. Thyroid function studies were helpful in determining thyroid abnormalities, and they were discontinued after 500 patients. A defect in the delayed hypersensitivity response was not demonstrated. (17 refs)

- 78-5973 Malignancies in Portuguese Thorotrast Patients.** (Eng) Horta, J. (Faculty Medicine, University of Lisbon, Lisbon, Portugal); Horta, M. E.; Motta, L.; Tavares, M. H. *Health Phys* 35(1): 137-151; 1978.

Partial results of a follow-up study of 1,237/2,434 individuals who received Thorotrast in Portugal, mostly from 1930 to 1950, are reported. As of March 1974, 931 of these patients had died: 16 from local Thorotrast granulomas, 25 from liver cirrhosis or fibrosis, 121 from malignant solid tumors (including 6 on the edge of large Thorotrast deposits, 75 tumors of the liver and other reticuloendothelial tissues, 5 bronchial carcinomas, and 4 bone tumors), and 23 from fatal blood diseases (including 12 leukemias). Hemangioendotheliomas were the most frequently diagnosed tumors in the Thorotrast patients; cholangiocarcinomas of the liver were also numerous. The Thorotrast patients, compared with a control group of 829 individuals injected with a nonradioactive contrast drug and with the general Portuguese population, had an increased incidence of liver, bone, and lung tumors and leukemias. (23 refs)

- 78-5974 Melanoma of the Skin in New Jersey.** (Eng) Stein, D. (Dept. Preventive Medicine and Community Health, New Jersey Medical Sch., Coll. Medicine and Dentistry New Jersey, Newark, NJ, 07103); Thind, I.; Louria, D. B. *J Med Soc NJ* 75(5): 391-393; 1978.

The incidence of malignant melanoma of the skin in residents of Newark and Morris and Hunterdon Counties, New Jersey, between 1970 and 1974 was determined and compared with the incidence rates from the Connecticut tumor registry and the Third National Cancer Survey. Because only five cases of cutaneous melanoma were noted in New Jersey blacks, they were excluded from the rest of the survey. Data from New Jersey, Connecticut, and the National Cancer Survey showed a progressive increase in age-specific incidence with increase in the age of the patients in both sexes. Relative few cases occurred before the age of 40. For males in Morris County, the age-specific incidence increased from 0 for those aged 0-9 yr to 20.90 for those > 80 yr of age; similar increases were found for both sexes in the other areas studied. The age-adjusted incidence rates for men were 2.78 in Newark, 3.59 in Morris County, 4.59 in Hunterdon County, 5.1 in Connecticut, and 4.6 in the National Cancer Survey. The age-adjusted incidence rates for women were 2.14 in Newark, 6.51 in Morris County, 2.35 in Hunterdon County, 4.65 in Connecticut, and 4.4 in the National Cancer Survey. (21 refs)

5975 **Skin Cancer and Ultraviolet Radiation.** (Eng) Rundel, R. D. (Space and Life Sciences Directorate, Natl. Aeronautics and Space Admin., Lyndon B. Johnson Space Center, Houston, TX, 77058); Nachtwey, D. *Photochem Photobiol* 28(3): 345-356; 1978.

model for calculating the increased incidence of non-melanoma skin cancer in Caucasians resulting from ozone reduction is presented. Third National Cancer Survey data on skin cancer incidence as a function of age were used to derive a dose-response model that was then used to predict the effects of increased biologically harmful UV radiation (BHUV) resulting from ozone reduction. A biological amplification factor was calculated separately for each geographic region for which skin cancer incidence data were available. Use of the latitudinal variation of incidence was avoided. The model postulates that the probability of skin cancer incidence is distributed log-normally as a function of total accumulated lifetime dose of BHUV and that for any given location, this dose to an individual is proportional to his age. The effect on skin cancer incidence of an increase in BHUV can then be calculated directly from the extent to which each individual's lifetime accumulated dose is thereby increased. The average result of such a perturbation would be to cause skin cancer to appear to an earlier age. Since skin cancer is predominantly a disease of the elderly, a shift to younger ages has the effect, when integrated over the entire population, of increasing the overall total number of cases of skin cancer. For the US as a whole, it is estimated that a 1% permanent decrease in the average stratospheric ozone thickness, which produces about a 2% increase in BHUV, would ultimately lead to an 8% increase in skin cancer incidence. (30 refs)

5976 **Background Dose Rate and Leukemia Mortality in Japan.** (Jpn) Sakka, M. (Tohoku Univ. School of Medicine, Sendai, Japan). *Nippon Acta Radiol* 38(3): 272-276; 1978.

The relationship between the natural background radiation dose and leukemia mortality in 27 Japanese prefectures was studied by linear regression analysis. No linear dose-effect relationship could be demonstrated over the range studied. (9 refs)

78-5977 **Characterization of Normal Human Embryo Cells Grown to Over 100 Population Doublings.** (Eng) Poiley, J. A. (Chemical Carcinogenesis Program, NCI, Frederick Cancer Res. Center, Frederick, MD, 21501); Schuman, R. F.; Pienta, R. J. *In Vitro* 14(5): 405-412; 1978.

Normal human embryo fibroblasts were subcultured for > 100 population doublings in unmodified basic medium. Their growth rate, confluent density, chromosome stability, growth in soft agar, ability to hydrolyze casein, and tumorigenicity indicated that they possessed the characteristics of normal cells. The batch of fetal bovine serum used to supplement the medium was of primary importance in the long-term growth of this culture. (32 refs)

See also:

- *(Rev.): 78-5401, 78-5402, 78-5403, 78-5404, 78-5406, 78-5411, 78-5414, 78-5423, 78-5425, 78-5426, 78-5429, 78-5431, 78-5432, 78-5435, 78-5447, 78-5459, 78-5460, 78-5461, 78-5462, 78-5463.
- *(Chem.): 78-5486, 78-5489, 78-5501, 78-5537, 78-5549, 78-5555, 78-5615, 78-5629.
- *(Phys.): 78-5715, 78-5718, 78-5719, 78-5720, 78-5729.
- *(Viral): 78-5785, 78-5790, 78-5822, 78-5835, 78-5836, 78-5837.
- *(Path.): 78-5875.

MISCELLANEOUS

- 78-5978 **Repopulation of Denuded Tracheal Grafts with Normal, Preneoplastic, and Neoplastic Epithelial Cell Populations.** (Eng) Terzaghi, M. (Biology Div., Oak Ridge Natl. Lab., P.O. Box Y, Oak Ridge, TN, 37830); Nettesheim, P.; Williams, M. L. *Cancer Res* 38(12): 4546-4553; 1978.

In vivo growth characteristics of nonmalignant and malignant epithelial (ept) cell cultures maintained in vitro for various lengths of time were studied. Three types of ept cell cultures from Fischer 344 rats were used: primary cell cultures and two established tracheal cell lines, one neoplastic (BP₁) and the other (1000 M) preneoplastic at passage 7 and neoplastic at passage 20. The cultures were inoculated into isolated tracheas that were then grafted to syngeneic recipients. At least 10⁴ ept cells were required to reestablish a complete ept lining in denuded tracheal grafts, whether normal or tumorigenic cell cultures were used. Inoculation of denuded tracheas with 1-wk-old primary cultures of normal tracheal epithelium resulted in reestablishment of a nearly normal mucociliary tracheal lining within 1-2 wk. Injection of primary ept cultures older than 1 wk failed to reestablish an ept lining; instead, the tracheal lumen became obliterated with connective tissue. In immunosuppressed recipients, inoculation with the preneoplastic (passage by) cell line resulted in the establishment of a well-organized, keratinizing squamous epithelium that remained stable for at least 6 wk if the recipient was immunosuppressed but was rejected at 4 wk in immunocompetent hosts. Inoculation with line 1000 M at passage 20 resulted in the establishment of a well-organized squamous epithelium for 3-4 wk and the development of disorganization, exophytic growth, and invasion at 4-6 wk. This occurred only in immunosuppressed recipients. In immunocompetent recipients, the ept lining was completely rejected within 2 wk. When cells from the highly malignant squamous cell carcinoma line (BP₁) were inoculated into trachea with an intact ept lining, only a few isolated nests of malignant cells were observed at 6 wk. Repopulation of denuded tracheas with cells from the same tumor line resulted in the establishment of an atypical squamous lining at 1 wk. At 2 wk, invasion was widespread, resulting in destruction of the trachea grafts shortly thereafter. The results show that this new "in vivo culture system" is suited to study of the growth and differentiation of carcinogen-altered or preneoplastic ept cells. It will complement existing in vitro culture systems. (27 refs)

- 78-5979 **Selection and Biologic Properties of Malignant Variants of a Murine Lymphosarcoma.** (Eng) Brunson, K. W. (Dept. Developmental and Cell Biology, Univ. California, Irvine, CA, 92717); Nicholson, G. L. *J Natl Cancer Inst* 61(6): 1499-1503; 1978.

The murine lymphosarcoma cell line RAW117 was used as a source of parental cells for the in vivo sequential selection of stable malignant variants in BALB/c mice. Sequential in vivo selections were performed by iv tumor cell inoculation, removal of resulting solid liver tumor nodules for growth in vitro, and reinjection of the cultured cells iv. After 10 sequential selections for implantation, invasion, survival, growth in liver, a lymphosarcoma cell line was obtained (RAW117-H10) that formed 200-fold more gross liver tumor nodules than did the parental tumor cell line in comparative biologic assays and displayed enhanced malignant properties when monitored by time of host death. Very few lung nodules or tumors were present at other sites with either the parental or any of the selected RAW117 lines. No differences in liver colonization were found between lymphosarcoma-selected lines cultured for a short term (2 wk) and cell lines cultured for a long term (3 mo). The enhanced malignancy of the tumor variant probably does not occur by temporal adaptation of the entire cell population to a different growth environment. Conditions in vivo may have resulted in the emergence of more aggressive and/or less antigenic variants from the parental population; this would be consistent with a recently proposed model for clonal evolution of neoplasia. (16 refs)

- 78-5980 **Hymenolepis nana in Retroperitoneal Sarcoma of an Albino Rat.** (Spa) Perez-Arroyo, J. (Laboratorio de Investigacion, Hospital Militar Regional, Guadalajara, Jal, Mexico); Lagarriga-Attias, J.; Corrales-Soltero, A. *Patologia (Mex)* 15(2): 111-117; 1977.

An adult form of *Hymenolepis nana* was found in the course of an undifferentiated retroperitoneal sarcoma of a laboratory albino rat. The parasitic origin of some tumors is reviewed. The relationship between parasites and the development of neoplasia is not fully understood. (24 refs)

- 78-5981 **Magnesium in Leukocyte Cells in Relation to Leukemogenesis.** (Eng) Hazdra, J. J. (Dept. Biochemistry, Illinois Benedictine Coll., Lisle, IL, 60532); Brunson, K. W.; Neenan, J. P. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands, 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D., eds. (New York: Elsevier/North Holland Biomedical Press) 503 pp.; 320-321; 1978.

Magnesium levels (micromoles/10⁹ WBC) were 7.0 in normal subjects, 5.3 in granulocytic leukemia patients, and 12.37

phocytic leukemia patients. The Mg/ATP ratio was 3.5 normal WBC, 2.5 in granulocytic WBC, and 10.74 in lymphocytic WBC. There appears to be an inverse correlation between Mg and ATP in the WBC of the leukemia patients. (6 refs)

78-5982 Genetics of Bovine Leukemia. II. Effect of Parents on the Frequency of Leukemia in the Progeny. (Rus) Ernst, L. K. (Dept. Genetics and Selection of Agricultural Animals, Agricultural Inst., Novosibirsk, USSR); Petukhov, V. L. *Genetika* 14(7): 1247-1256; 1978.

An attempt was made to estimate the heritability of a predisposition or resistance to leukemia in cattle. A pedigree analysis of 14,000 animals (progeny of 554 sires), of whom 2,060 had leukemia, showed that the contribution of fathers to the leukemia morbidity of their daughters ranged from 7% to 15%. The incidence of leukemia among the offspring of leukemic sires was 26.6%, compared with the average leukemia incidence of 20.8%. (22 refs)

78-5983 Terminal Differentiation of Human Promyelocytic Leukemia Cells Induced by Dimethyl Sulfoxide and Other Polar Compounds. (Eng) Collins, S. J. (Laboratory of Cell Biology, NCI, NIH, Bethesda, MD, 20014); Rusti, F. W.; Gallagher, R. E.; Gallo, R. C. *Proc Natl Acad Sci USA* 75(5): 2458-2462; 1978.

The effect of dimethyl sulfoxide (Me_2SO) on a human leukemic cell line (HL-60) established from the peripheral WBC of a patient with acute promyelocytic leukemia is reported. This cell line displays a distinct morphological and histochemical commitment toward myeloid differentiation; the cultured cells are predominantly promyelocytes. The addition of Me_2SO (1.25%) to the culture induced these cells to differentiate into myelocytes, metamyelocytes, and banded and segmented neutrophils. All 150 clones developed from the HL-60 culture showed similar morphological differentiation in the presence of Me_2SO . Unlike the morphologically immature promyelocytes, the Me_2SO -induced mature cells exhibited functional maturity, as exemplified by their phagocytic activity. A number of other compounds previously known to induce erythroid differentiation of mouse erythroleukemia (Friend) cells (acetamide, N-methylacetamide, N,N-dimethylacetamide, N-methylformamide, N,N-dimethylformamide, piperidone, 1-methyl-2-piperidone, and ethylene glycol) induced analogous maturation of the HL-60 cells. The marked similarity in behavior of HL-60 cells and Friend cells in the presence of these inducing agents suggests that similar molecular mechanisms are involved in the induction of differentiation of these human myeloid and murine erythroid leukemic cells. (28 refs)

78-5984 Clonal Analysis of Chronic Myeloid Leukemic Cells Using Agar Cultures. (Eng) Metcalf, D. (Cancer Res. Unit, Walter and Eliza Hall Inst., Royal Melbourne Hosp., P. O. 3050, Victoria, Australia); Johnson, G. R.; Kolber, S.; Dresch, C. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 307-310; 1978.

Myeloblasts in the peripheral blood of chronic myeloid leukemia (CML) patients were capable of forming neutrophilic and eosinophilic colonies. Analysis of feeder layer activity showed that unfractionated CML cells have relatively low feeder layer activity and that both CML and acute ML (AML) cells depend on stimulation for proliferation. AML and CML cells exhibited lower autologous stimulating activity than nonleukemic populations. The results support the concept that in CML, there is extreme hyperplasia of the granulocytic progenitor cells that remain responsive to regulator control. (6 refs)

78-5985 Growth and Differentiation of Human Myelogenous Leukemia Cells in Conditioned Medium from Human Embryo Cultured Cells. (Eng) Gallagher, R. (NCI, Bethesda, MD, 20014); Ruscetti, F.; Collins, S.; Gallo, R. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 303-306; 1978.

Conditioned medium (CM) from early passage fibroblast cultures of human embryonic origin stimulated extended growth and differentiation in suspension cultures of myeloid cells from patients with myelogenous leukemia. The stimulatory activity probably derives from a subpopulation of embryonic cells since (1) it was not recovered from adult cell cultures, (2) it was infrequent even in embryonic cultures; and (3) it was labile with respect to culture passaging and freezing. The possibility that the active factor(s) interacted with specific leukemic cell receptors (eg, fetal receptors) was suggested by its selective stimulation of myelogenous leukemia WBC and, in some cases, by cytogenetic identification of cultured leukemia cells. The ability of embryonic CM to provide replicating WBC suspension cultures should be of value in studies of myeloid cell differentiation. (7 refs)

78-5986 Heterotransplantation of the K562 Cell Line Derived from a Patient with Chronic Myeloid Leukemia. (Eng) Machado, E. A. (Center Health Sciences, Univ. Tennessee Memorial Res. Center, 1924 Alcoa High-

way, Knoxville, TN, 37920); Lozzio, B. B.; Lozzio, C. B.; Aggio, M. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 418-421; 1977.

Chronic myelogenous leukemia (CML) cells from the K562 cell line, which is positive for the Philadelphia chromosome, were transplanted sc in nude mice. The recipient mice developed solid vascularized tumors termed myelosarcomas. The myelosarcoma cells retained the original chromosome and antigen markers, and they actively incorporated ^3H -thymidine. The growth of the myelosarcomas was inhibited by reconstituting the nude mice with isogenic thymuses. The tumor, which can be maintained by serial passage, provides a useful bioassay system for therapeutic trials and studies of cell kinetics. (7 refs)

78-5987 Expression of Cholera Toxin Binding Sites on Normal and Malignant Haemopoietic Cells.

(Eng) De Cicco, D. (Imperial Cancer Res. Fund, Lincoln's Inn Fields, London WC2A 3PX, England); Greaves, M. F.; Van Den Engh, G.; Warner, N. L. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 357-362; 1977.

The expression of G-M1 gangliosides on the surface of normal and malignant hemopoietic stem cells was studied by determining the number of ^{125}I -cholera toxin binding sites per cell. There were no alterations in G-M1 expression that could be correlated with hemopoietic malignancy per se. Preliminary experiments with the Fluorescence Activated Cell Sorter (FACS) indicated that normal immature myeloid cells had a lower density of toxin binding sites than granulocytes. The correlation of G-M1 expression with maturation was also apparent within a single clone of Philadelphia chromosome-positive chronic myeloid leukemia cells entering acute blast crisis. Separation of the cells by FACS according to fluorescence intensity revealed a clear distinction between the relatively mature myeloid elements and the blast cells. Mouse cells had a receptor density two orders of magnitude higher than human lymphoid cells. This difference may explain the previously reported differential effects of cholera toxin on lymphocyte proliferation. (16 refs)

78-5988 Effect on Androgen Depletion on Growth and Androgen Dependency of Shionogi Carcinoma

115. (Eng) Kitamura, Y. (Inst. Cancer Res., Osaka U. Medical Sch., Kita-ku, Osaka, 530, Japan); Okamoto, Uchida, N.; Yamaguchi, K.; Matsumoto, K. *Cancer* 38(12): 4711-4716; 1978.

To determine whether the androgen (ADG) dependence of the Shionogi carcinoma 115 (SC115) varies in different phases of growth, the fate of SC115 tumors in an ADG-depleted state was studied in DS mice. The SC115 tumors were implanted in castrated male mice, and testosterone propionate (100 μg , sc) was injected daily for 60 days starting various times after grafting. Inoculated SC115 cells showed evidence of necrosis within 5 days after implantation in absence of ADG. After initiation of growth by ADG, subsequent fate of the tumors following ADG removal depended on the size of the tumor: small, medium, and large tumors showed complete, temporary, and no regression, respectively. In contrast to the partial and complete lack of ADG dependence of medium and large tumors in situ, tumor seeds taken from the medium and large tumors before ADG removal grew only in males when transplanted into male and female mice. On the other hand, tumor seeds taken from about one-half of the regrown tumors after ADG removal grew in both male and female hosts. One type of the ADG-independent tumors contained cytosol ADG receptor (AR) and consisted of the original SC115 cells (medullary cancer) and spindle-shaped cells. Another type of independent tumor contained no cytosol AR and consisted only of spindle-shaped cells. These findings suggest that the effect of ADG depletion on growth of the tumor, which consists of AR-positive and ADG-dependent cells, varies in different growth phases and that the ADG dependence of tumor cells can be changed after ADG removal. The SC115 tumor seems to be a good model for elucidating hormone-dependent cancer growth and for establishing endocrine treatment in hormone-dependent cancers. (26 refs)

78-5989 Thyroid Disease and Function in Breast Cancer Patients and Non-hospitalized Controls Evaluated by Determination of TSH, T3, rT3, and T4 Levels in Serum.

(Eng) Adami, H. O. (Dept. Surgery, Uppsala Univ. Uppsala, Sweden); Rimsten, A.; Thoren, L.; Vegelius, Wide, L. *Acta Chir Scand* 144(2): 89-97; 1978.

Earlier thyroid disease and thyroid function were compared in 179 consecutively diagnosed breast cancer patients and 179 age-matched controls. Thyroid function was assessed by measuring the serum levels of thyrotropin (TSH), triiodothyronine (T3), reverse triiodothyronine (rT3), and thyroxine (T4). The overall occurrence of thyroid disease, 15/179 in the patient group and 17/179 in the control group, was similar, and two women from each group had a history of T4 treatment. Slight but significant differences were found in the patient group for three serum hormone levels, with a higher mean value for TSH (2.5 milliunits/liter in the patient group vs 1.5

controls) and rT3 [0.38 vs 0.33 nanomole(nmol)/liter] a lower mean value for T3 (2.16 vs 2.25 nmol/liter). T4 concentration in the serum did not differ significantly. In a random sample of 50 patients and controls, T3-resin uptake was significantly higher in the patient group than in the control group. The changes in thyroid hormone levels in the breast cancer patients were similar to those noted in a number of other nonthyroidal diseases; eg, hepatic cirrhosis, myocardial infarction, acute febrile illness, and starvation. The pattern of changes in the patient group may represent secondary, probably extrathyroidal changes in thyroid hormone metabolism. There is no indication that thyroid disease is an etiologic factor in breast cancer. (53 refs)

5990 Thyroid-stimulating Hormone and Prolactin Levels in Breast Cancer. (Eng) Aldinger, K. A. Department of Endocrinology and Medical Breast Service, Dept. of Medicine, Univ. Texas System Cancer Center, M.D. Anderson Hosp. and Tumor Inst., 6723 Bertner Ave., Houston, TX, 77030; Schultz, P. N.; Blumenschein, G. R.; Samaan, N. A. *Arch Intern Med* 138(11): 1638-1641; 1978.

Thyroid-stimulating hormone (TSH) and prolactin (PRL) levels in 148 breast carcinoma (BC) patients were measured by radioimmunoassay before and after iv administration of protirelin (PTR: 500 µg). The patients, of whom 42 were premenopausal and 106 postmenopausal, were divided into metastatic or primary cancer groups and matched with normal controls. Basal TSH levels were elevated in 36% of the BC patients and basal PRL levels in 22%. Although differences in mean TSH and PRL levels between the patients and controls were significant, there was no correlation between serum levels of the two hormones. Of 53 patients with elevated basal TSH, 30 did not demonstrate an elevated peak after PTR stimulation. In the metastatic BC group, 13% of the patients with normal basal PRL responded to PTR stimulation. All but two of the patients with elevated basal TSH levels had normal serum thyroxine levels and normal T₃ resin uptake. Previous irradiation to the upper chest area did not affect TSH levels; and although the mean survival and disease-free interval were shorter for patients with elevated TSH and PRL, the results were not significant. (39 refs)

5991 Relationship Between Urine and Plasma Estrogen Ratios. (Eng) Longcope, C. (Worcester Foundation Experimental Biology, Shrewsbury, MA, 01545); Pratt, J. H. *Cancer Res* 38(11, part 2): 4025-4028; 1978.

An attempt was made to determine whether women with differing urinary estrogen ratios had different blood production rates (BPR) of the estrogens. Normal young women col-

lected 24-hr urine specimens during the follicular and luteal phases of their cycle, and urinary estrone, estradiol, and estriol levels were measured. The women were placed into two groups according to whether their urinary estrogen ratios were > 1.3 or < 0.7. Normal postmenopausal women collected 24-hr specimens and were grouped according to whether their ratios were > 3.2 or < 2.1. An additional group studied without regard to urinary ratios consisted of women who had had breast cancer but had no clinical evidence of the disease. Between the respective high- and low-ratio groups, there were no significant differences in circulating levels of estriol, metabolic clearance rates of estriol, or BPR of estriol, estrone, or estradiol. The breast cancer women and normal controls also had similar BPR of the three estrogens. Ratios of the BPR of estriol to estrone and estradiol were similar in the high- and low-ratio groups and also in the breast cancer and control groups. Therefore, the differences in urinary ratios reflect differences in the pathways of metabolism of the free circulating estrogens and not variations in BPR. Estriol was produced at only 10% the rate of estrone and estradiol and thus would probably play only a minor role in the overall biological activity of the estrogens at the level of the target tissues. (24 refs)

78-5992 Prolactin and the Development and Progression of Early Neoplastic Mammary Gland Lesions. (Eng) Welsch, C. W. (Dept. Anatomy, Michigan State Univ., E. Lansing, MI, 48824). *Cancer Res* 38(11, part 2): 4054-4058; 1978.

The effect of chronic suppression of prolactin (PL) on the initiation and progression of hyperplastic alveolar nodules (HAN) was determined in nulliparous 2-mo-old and multiparous 8-mo-old C3H/HeJ mice. Also determined was the growth stimulatory effect of human placental lactogen (HPL), a peptide chemically and physiologically similar to PL, on hyperplastic (benign) human breast tissues. Chronic suppression of PL by 2-bromo-α-ergocryptine (CB-154: 0.1 mg/day for up to 1 yr, sc) sharply suppressed or totally prevented the formation of HAN and neoplastic mammary gland lesions in young mice and markedly suppressed the progression of HAN to carcinomas in mature mice. Most of these mice lived an entire life-span free of mammary tumors, during which they had normal ovarian activity, normal body wt gains, and no signs of morbidity. Treatment of mice with 17β-estradiol or the oral contraceptive Enovid also suppressed HAN development and subsequent mammary tumorigenesis. CB-154-induced PL suppression was more effective than ovariectomy in suppressing HAN development and comparable to this procedure in suppressing mammary carcinoma development. HPL promoted growth both in vitro (organ culture) and in vivo (athymic nude mouse) of the epithelium in benign human breast tumors. Whether or not human pituitary PL is capable of mimicking the mammotrophic action of HPL and whether a dysplastic, PL-sensitive lesion

comparable to HAN exists in the human breast remain to be determined. (16 refs)

- 78-5993 A Specific Progesterone-binding Component of Human Breast Cyst Fluid: Its Isolation and Characterization (Meeting Abstract).** (Eng) Pearlman, W. H. (Dept Pharmacology, Univ. North Carolina Sch. Medicine, Chapel Hill, NC, 27514); Peng, L. H.; Mazoujian, G.; Haagenzen, D. E.; Wells, S. A.; Kister, S. J. *J Endocrinol* 75(3): 19-20; 1977. (3 refs)

- 78-5994 Effect of Serum on Cells Cultured from Human Mammary Tumors.** (Eng) Gaffney, E. V. (Pennsylvania State Univ., 5101 Freer Building, University Park, PA, 16802); Pigott, D. *In Vitro* 14(5): 451-457; 1978.

Cultures established from human mammary ductal carcinomas were studied to determine cell yield from tumors, survival time in primary culture, and the response of the cells to different sera. Clusters of the cells formed two morphologically distinct epithelial colonies in culture, E and E'. From 0.3% to 13.0% of E' cell clusters attached and formed colonies. Attachment was independent of tumor grade. The survival of E' cell colonies was limited to 10 days with rapid cell degeneration commencing at 7 days. A comparison of sera (fetal bovine, agamma bovine, whole bovine, human, swine, horse, and chicken) showed that colony formation by the cells during the first week of culture was max in the presence of fetal bovine serum (FBS). Human serum alone was 70%-100% less effective in promoting E' colonies. In the presence of FBS, human serum from normal donors inhibited colony development with E' but not with E cells. This inhibitory activity could not be attributed to either antagonistic hormones or the source of human serum. The results indicate that normal human serum contains a factor that exhibits an inhibitory activity specific for E' human epithelial cells derived from malignant tumors. (16 refs)

- 78-5995 Inhibition of Pregnancy-dependent Mammary Tumorigenesis by a Single Treatment of Neonatal GR/A Mice with Monosodium Glutamate.** (Eng) Nagasawa, H. (Pharmacology Div., Natl. Cancer Center Res. Inst., Tsukiji 5-1-1, Chuo-ku, Tokyo, 104, Japan); Yanai, R. *Proc Soc Exp Biol Med* 158(1): 128-129; 1978.

The effects of neonatal administration of monosodium glutamate (MSG) on the development of pregnancy-dependent mammary tumors and on reproduction were examined in GR/A mice. A single sc injection of 4 mg MSG was given on the day of birth. Mice were mated on day 60, and size and number of palpable mammary tumors were counted at parturition. MSG treatment markedly inhibited the appearance of pregnancy-dependent mammary tumors and decreased the number of tumors per mouse. Reproduction was not affected. These results suggest that the development of pregnancy-dependent mammary tumors is affected to a greater extent by subtle changes in the hormonal milieu, especially in progesterone level, than is the maintenance of pregnancy. (13 refs)

- 78-5996 Multicellular Tumor Spheroid Formation by Breast Cancer Cells Isolated from Different Sites.** (Eng) Yuhas, J. M. (Cancer Res. and Treatment Center, Univ. Mexico, 900 Camino de Salud N.E., Albuquerque, NM, 87131); Tarleton, A. E.; Molzen, K. B. *Cancer Res* 38(8): 2486-2491; 1978.

The possibility that strong natural or artificial selection for growth in suspension might produce lines that are unable to form multicellular tumor spheroids (MTS) was investigated using breast cancer lines derived from malignant pleural effusions or from solid growths. Cells from 14 breast cancer lines (8 human, 5 rat, and 1 mouse) were plated in liquid medium over an agar medium base. Within 1-3 wk, 8/14 lines formed MTS, and all of these had been isolated from solid tumors or metastases. The remaining six lines, which were isolated from malignant pleural effusions, failed to form MTS, with one exception, failed to form aggregates > 10-50 cells. The exception formed aggregates of up to 100 μ m, but failed to grow and rapidly degenerated. Similarly artificial selection for an ascites variant of the MTS-forming rat 13762-A adenocarcinoma line produced the 13762-A line, which could no longer form MTS. These results suggest that breast cancer cells from pleural effusions are genetically different from most tumor cells in solid breast cancer specimens, that they are unable to grow in true solid form, and that these differences persist in spite of prolonged propagation in tissue culture. (19 refs)

- 78-5997 Population Density as a Factor in the Evolution of Neoplastic Cell Lines.** (Eng) Taylor, W. (Lab. Cellular and Molecular Biology, NCI, Building 36, Room 4D-18, Bethesda, MD, 20014); Camalier, R. F.; Tu

R. W.; Sanford, K. K. *Cancer Res* 38(11, part 1): 0-3846; 1978.

role of population density in the progression from the neoplastic to the neoplastic state was studied in two nonneoplastic mouse cell lines, NCTC 7914 and 7915. Sublines from each clone were transferred every 3-4 days. Inoculum sizes selected to minimize or maximize cell-cell contact, 1×10^5 or 4×10^5 cells/flask. In vivo assays of tumorigenicity were made at 3-mo intervals. Tumor production was observed in all lines, regardless of inoculum size. However, sublines carried at the high inoculum size produced tumors with significantly shorter latent periods than those carried at the low size. By passage 55, 85%-100% of the colonies were diagnosed morphologically as neoplastic. By 10 days in vitro, all sublines grew in agarose as colonies ≥ 2 mm in diameter, but the number of colonies was greater in sublines grown at the higher inoculum size than those grown at the lower size. Cells derived from the neoplastic sublines were susceptible to BCG-activated macrophages; the degree of susceptibility increased as the tumor latent period increased. The parent clones did not form colonies in agarose, nor were they killed by activated macrophages. Saturation density increased and population doubling time decreased with period in culture. There was no increase in aerobic glycolysis with time in culture or neoplastic transformation. The results indicate that carrying mouse cells at low density does not prevent neoplastic transformation and that certain morphological changes, growth in agarose, and susceptibility to activated macrophages are associated with neoplastic transformation. (14 refs)

78-5998 **An Electronic Theory of Cancer.** (Eng) Szent-Gyorgyi, A. (Workshop Natl. Foundation for Cancer Res., Marine Biological Lab., Woods Hole, MA). *Int J Quantum Chem* 12(Suppl 1): 407-414; 1978.

The history of life is divided into two parts by the appearance of light: an anaerobic, dark proliferative period (α) and an aerobic period of light-regulated growth (β). The transition between the two ($\alpha \leftrightarrow \beta$) is still an everyday occurrence. However, a cell may lose its ability to build up and stabilize its β state and may thus return permanently to the proliferative α state; this state is cancer, characterized by a low cohesion and proliferation. Thus, the cancer problem may be boiled down to the chemical mechanism of the $\alpha \leftrightarrow \beta$ transformation. The central event of this transformation is the electronic desaturation of protein, the first step of which is production of an electron acceptor that can take over the electrons from the protein. The proposed acceptor is methylglyoxal. The formation of a charged radical in the processes connected with the electronic desaturation of proteins is avoided by incorporating the methylglyoxal into the donor protein molecule before transferring charges. The incorpora-


tion of methylglyoxal is essentially a p -type doping. This view suggests that the cancer problem is indeed finite. (15 refs)

78-5999 **Thermographic Tumor Detection Enhancement Using Microwave Heating.** (Eng) Thompson, J. E. (Coll. Engineering, Univ. South Carolina, Columbia, SC, 29208); Simpson, T. L.; Caulfield, J. B. *Trans Microwave Theory Tech* 26(8): 573-580; 1978.

The sensitivity of thermography for tumor detection was shown to be improved by concurrent irradiation of the examined area with 2,450-gigahertz microwaves. The microwave radiation increases the temperature differential between healthy and tumor tissue substantially. Strain-2 guinea pigs bearing an sc line-10 hepatoma in the left shoulder were irradiated for 75 sec, and thermographic measurements were made every 15 sec during both irradiation and cooling. Contour maps depicting areas that experienced the same change in temperature were constructed. These maps clearly indicated the tumor location, particularly as the duration of irradiation increased. The max temperature increase in the tumor tissue was 5.5 C, that in healthy tissue was 2.5 C. Although microwave irradiation increased the spatial temperature nonuniformity of all tissue, the max change in the healthy shoulder was 1.5 C, compared with 3.5 C in the tumor. Therefore, increased nonuniformity of the spatial temperature distribution is indicative of an existing tumor. The rates of heating and cooling of the tumor tissue were also greater than those of healthy tissue. The thermal response of breast tumors, which are likely to be less cellular and vascular than hepatomas, is presently unknown. (24 refs)

78-6000 **Use of Computerized Methods to Predict Metabolic Pathways and Metabolites.** (Eng) Spann, M. L. (Natl. Library Medicine, Bethesda, MD, 20014); Chu, K. C.; Wipke, W. T.; Ouchi, G. *J Environ Pathol Toxicol* 2(1): 123-131; 1978.

Computer programs that can predict biologic activities from metabolism of a compound are being developed. The project has two stages: (1) to incorporate metabolic reactions into a computer program to predict probable metabolites and (2) to recognize structural units that may be related to specific biologic activities. Computer analysis may aid in (1) screening compounds to determine if their metabolites are probable carcinogens; (2) identifying unknowns isolated in metabolic studies, and (3) designing appropriate radioactive tracer experiments for elucidating a specific metabolic pathway. (7 refs)



Author Index

- aronson, S. A., 78-5772, 78-5850
 Abdul-Baki, A., 78-5609
 Abe, S., 78-5857
 Abenova, U. A., 78-5831
 Ablashi, D. V., 78-5832
 Abraham, E., 78-5933
 Acton, R. T., 78-5781
 Adami, H. O., 78-5989
 Adeloze, A., 78-5456
 Adler, K. R., 78-5895
 Agarwal, S. S., 78-5959
 Ageenko, A. I., 78-5847
 Aggio, M., 78-5986
 Agrba, V., 78-5817
 Aharonov, A., 78-5813
 Ahmed, S. I., 78-5525
 Akasaka, K., 78-5565
 Al'shtein, A. D., 78-5831
 Albert, R. E., 78-5723
 Aldinger, K. A., 78-5990
 Aleksandrov, V. A., 78-5541
 Alevy, M. C., 78-5745
 Alexandrov, V. A., 78-5638
 Allen, N., 78-5647
 Alrenga, D. P., 78-5912
 Althoff, J., 78-5523
 Altstein, A. D., 78-5793
 Amick, R. M., 78-5869
 Ananthaswamy, H. N., 78-5708
 Anderson, R. E., 78-5722
 Andjelkovich, D., 78-5954
 Andrews, F. J., 78-5955
 Angeles, R. M., 78-5626
 Angsubhakorn, S., 78-5602
 Ansari, A. H., 78-5874
 Antley, R., 78-5707
 Antonov, V. S., 78-5746
 Anukarahanonta, T., 78-5470
 Anzil, A. P., 78-5859
 Appell, R., 78-5716
 Arai, H., 78-5583
 Archer, M. C., 78-5612
 Areias, M. A., 78-5918
 Arenaz, P., 78-5664
 Ariga, H., 78-5848
 Armstrong, G. R., 78-5832
 Armstrong, M. Y., 78-5758
 Arnold, S. D., 78-5636
 Aronson, I., 78-5490
 Asano, T., 78-5494
 Ashwood-Smith, M. J., 78-5563
 Atherton, D. R., 78-5731, 78-5732
 78-5733, 78-5734
 Atkin, N. B., 78-5871, 78-5893
 Audigier, J. C., 78-5942
 Autrup, H., 78-5508, 78-5557, 78-5562
 78-5590
 Averbach, P., 78-5872
 Axelsson, O., 78-5969
 Aya, T., 78-5828
 Azuma, I., 78-5532
 Bababunmi, E. A., 78-5469
 Babakova, I., 78-5727
 Baes, C., 78-5853
 Baginski, B., 78-5940
 Bailone, A., 78-5448
 Baird, W. M., 78-5513
 Bakacs, T., 78-5834
 Bakayev, V. V., 78-5811
 Baker, J. K., 78-5628
 Baker, M. C., 78-5871, 78-5893
 Balducci, L., 78-5901
 Baldwin, J. E., 78-5577
 Baldy, W. J., 78-5478
 Baltimore, D., 78-5776
 Bannasch, P., 78-5688, 78-5689
 Bansal, B. R., 78-5867
 Bansal, S. C., 78-5867
 Bapat, C. V., 78-5540
 Barahona, H., 78-5816
 Barale, R., 78-5653
 Baranova, L. N., 78-5541
 Barber, D. J., 78-5724
 Barber, S., 78-5908
 Barbin, A., 78-5492
 Barendsen, G. W., 78-5690
 Barkin, J. S., 78-5710
 Barkla, D. H., 78-5496
 Barone, J. J., 78-5864
 Barrett, L. A., 78-5508, 78-5557
 Bartal, A., 78-5920
 Bartsch, H., 78-5492
 Bases, R., 78-5678
 Basombrio, M. A., 78-5765
 Bassin, R. H., 78-5760
 Bassir, O., 78-5469
 Bauer, H., 78-5740
 Baugnet-Mahieu, L., 78-5853
 Bauman, A. W., 78-5903
 Bayreuther, K., 78-5798
 Bazin, H., 78-5886
 Bazlova, L. S., 78-5495
 Bean, J. A., 78-5949
 Beasley, R. P., 78-5965
 Becchetti, A., 78-5694
 Becci, P. J., 78-5582, 78-5876, 78-5878
 Beckers, A., 78-5886
 Beda, B., 78-5962
 Beghe, C., 78-5901
 Beland, F. A., 78-5502, 78-5512
 Belser, W. L., 78-5568
 Benade, L. E., 78-5759
 Bender, M. A., 78-5686
 Benderson, J., 78-5561
 Benfield, J. R., 78-5569
 Bengtsson, G., 78-5969
 Beniashvili, D. Sh., 78-5643
 Bentwich, Z., 78-5920
 Bergholz, C., 78-5817
 Berndt, H., 78-5461, 78-5919
 Berndtsson, B. S., 78-5969
 Bernstein, A., 78-5768
 Berry, D. L., 78-5567
 Bertini, M., 78-5744
 Bertrand, E., 78-5962
 Bhamarapravati, N., 78-5470, 78-5602
 Bhat, A. V., 78-5540
 Bhatnager, P., 78-5959
 Bhoopalani, N., 78-5856
 Bhutani, R., 78-5869
 Biggar, R. J., 78-5835, 78-5836
 Bishop, W. W., 78-5613
 Bitter, G. A., 78-5839
 Bjarnason, O., 78-5951
 Blackstock, R., 78-5438
 Blanc, A. P., 78-5896
 Blaufuss, E. M., 78-5637
 Blevins, C. S., 78-5777
 Bloemendal, H., 78-5778
 Bloemers, H. P., 78-5778, 78-5780
 Blum, S., 78-5954
 Blumenschein, G. R., 78-5990
 Bocker, J., 78-5835, 78-5836
 Bodell, W. J., 78-5648
 Bohm, E., 78-5971
 Boiron, M., 78-5771
 Boling, M. E., 78-5679
 Bondurant, M. C., 78-5761
 Bonnell, J. A., 78-5719
 Bonner, J., 78-5551
 Boquist, L., 78-5728
 Boseman, J. J., 78-5731, 78-5733
 Boulanger, P. A., 78-5842, 78-5843
 Bourkas, A. E., 78-5827
 Bourrouilloux, G., 78-5905
 Bracken, W. M., 78-5567
 Bradt, C., 78-5804
 Branz, S. E., 78-5577
 Bratslavsky, V., 78-5424
 Breistein, L. S., 78-5868
 Breton-Gorius, J., 78-5899
 Bridges, B. A., 78-5480
 Bridges, J. W., 78-5480
 Bridges, S. H., 78-5761
 Brogger, A., 78-5703
 Bronstein, R. E., 78-5544
 Brorsson, M., 78-5705
 Brown, C. C., 78-5408, 78-5582
 Brown, M. M., 78-5652
 Bruce, W. R., 78-5629
 Bruenger, F. W., 78-5732, 78-5733
 Brugge, J., 78-5739
 Brunson, K. W., 78-5979
 Bryant, M. L., 78-5814
 Buchanan-Davidson, D. J., 78-5928
 Bucheler, J., 78-5536
 Bugno, T. J., 78-5981
 Burger, P. M., 78-5702
 Burns, F. J., 78-5723
 Burny, A., 78-5788
 Burtonboy, G., 78-5886
 Bushnell, D. E., 78-5482
 Butel, J. S., 78-5810
 Buttin, G., 78-5818
 Bykovskii, A. F., 78-5831
 Cabradilla, C. D., 78-5772
 Caillard, J. F., 78-5875
 Camalier, R. F., 78-5997
 Campbell, T. C., 78-5473

- Capdeville, J., 78-5905
 Capel, I. D., 78-5560
 Carcassonne, Y., 78-5896
 Carda Aparici, P., 78-5915
 Cario, A. A., 78-5431
 Carrano, A. V., 78-5687
 Casagrande, J. T., 78-5952
 Castegnaro, M., 78-5597, 78-5616
 Castner, B., 78-5713
 Castro, J. A., 78-5603
 Caulfield, J. B., 78-5999
 Cavalieri, E., 78-5523, 78-5561
 Cavalli, F., 78-5894
 Cerletty, J. M., 78-5972
 Chabanon, G., 78-5829
 Challis, B. C., 78-5617
 Chapleo, M. R., 78-5599
 Cheng, H. F., 78-5534
 Chernozemsky, I. N., 78-5961
 Chess, L., 78-5826
 Cholewka, K., 78-5941
 Chorna, L. D., 78-5746
 Chouroulinkov, I., 78-5418
 Christiansen, R., 78-5579
 Chu, K. C., 78-6000
 Chumakov, P. M., 78-5806
 Clapp, N. K., 78-5665
 Clark, C. G., 78-5861
 Clark, O. H., 78-5713
 Clausen, O. P., 78-5526
 Cleary, M. L., 78-5775
 Cleaver, J. E., 78-5451, 78-5648
 Clerc, M., 78-5962
 Cleuter, Y., 78-5788
 Clive, D., 78-5652, 78-5657, 78-5666
 Cocuzzi, E., 78-5539
 Coffino, P., 78-5680
 Cohen, A. H., 78-5569
 Cohen, M. M., 78-5422
 Cohen, S. N., 78-5434
 Coleman, M. T., 78-5647
 Collett, M. S., 78-5737, 78-5739
 Collins, S., 78-5985
 Collins, S. J., 78-5983
 Colombies, P., 78-5905
 Comoglio, P. M., 78-5744
 Compans, R. W., 78-5781
 Conliffe, M. A., 78-5862
 Conney, A. H., 78-5548
 Connor, R., 78-5557
 Conomy, J. P., 78-5457
 Cook, K. H., 78-5709
 Corrales-Salcedo, A., 78-5980
 Cotter, S. M., 78-5786
 Coursaget, P., 78-5837
 Craighead, J. E., 78-5521
 Creutzig, H., 78-5718
 Croisy, A., 78-5492
 Cruse, J. P., 78-5861
 Cunha, G. R., 78-5890
 Curnen, M. G., 78-5463
 D'Halluin, J. C., 78-5842, 78-5843
 Dahms, R. P., 78-5510
 Dalens, M., 78-5829
 Dan, K., 78-5902
 Danes, B. S., 78-5881
 Daniel, A., 78-5908
 Daniel, M. D., 78-5816
 Davies, P. J., 78-5654
 Davies, P. W., 78-5633
 Davies, R., 78-5630
 Dawson, A. A., 78-5911
 Dawson, M. S., 78-5823
 Day, N. E., 78-5951
 De Bertoldi, M., 78-5653
 De Cicco, D., 78-5987
 Dean, B. J., 78-5655
 Debry, G., 78-5634
 Decat, G., 78-5489
 Dechamps, M., 78-5818
 Decleve, A., 78-5756, 78-5759
 Deftos, L. J., 78-5713
 Degen, G. H., 78-5472
 Deinhardt, F., 78-5817
 Delaforge, M., 78-5519
 Delorme, F., 78-5485, 78-5486
 Delozier, D., 78-5707
 Denda, A., 78-5580
 Dennis, M. J., 78-5630
 deNoronha, F., 78-5786
 DePamphilis, M. L., 78-5803
 Deschner, E. E., 78-5498
 Devoret, R., 78-5448
 Diana, G., 78-5694
 Diaz Gomez, M. I., 78-5603
 Didier, J., 78-5829
 Dierks, P., 78-5738
 Diessner, H., 78-5536
 Dina, D., 78-5770
 Dioguardi, D., 78-5527
 Dittmar, K. J., 78-5741
 Doerjer, G., 78-5536
 Doerr, R. C., 78-5622
 Dolphin, G. W., 78-5720, 78-5725
 Donaldson, I., 78-5711
 Dorange, J. L., 78-5519
 Doskeland, S. O., 78-5868
 Drake, J. W., 78-5450
 Dresch, C., 78-5984
 Drevon, C., 78-5593
 Dreyfus, B., 78-5899
 Drucker, J., 78-5837
 Duchayne, E., 78-5905
 Dukhlev, S., 78-5936
 Dunlop, N. M., 78-5777
 Dunn, W. J., 78-5668
 Dunne, F. B., 78-5505
 Duran-Troise, G., 78-5760
 Durkin, W., 78-5873
 Dyck, J., 78-5904
 Dymarskii, L. Iu., 78-5460
 Dzordzhadze, N. A., 78-5937
 Easton, J. M., 78-5832
 Ebenstein, P., 78-5758
 Ebralidze, D. V., 78-5937
 Edlund, L. E., 78-5781
 Edwards, A., 78-5617
 Edwards, A. A., 78-5725
 Edwards, M. B., 78-5645
 Ehling, U. H., 78-5662
 Ehmann, U. K., 78-5709
 Ehrenberg, L., 78-5969
 Eisenman, R., 78-5736
 Eisenstark, A., 78-5708
 El-Maraghy, M. A., 78-5888
 Elder, J. H., 78-5762
 Elespuru, R. K., 78-5515
 Elgiet, W., 78-5941
 Elguezabal, A., 78-5882
 Elling, H., 78-5859
 Endo, H., 78-5588
 Engbring, N. H., 78-5972
 Englund, F., 78-5533
 Enjalbert, L., 78-5829
 Epler, J. L., 78-5514, 78-5589
 Epstein, M. A., 78-5441
 Erickson, J., 78-5493
 Erikson, E., 78-5739
 Erikson, R. L., 78-5737, 78-5739
 Ernst, L. K., 78-5982
 Essex, M., 78-5786
 Esty, A., 78-5792
 Everett, J. R., 78-5869
 Falk, L., 78-5817
 Fan, T. Y., 78-5627
 Farber, P. A., 78-5763
 Farquhar, M. N., 78-5474
 Farsund, T., 78-5584
 Fay, H. T., 78-5960
 Fazio, T., 78-5623
 Feig, S. A., 78-0690
 Feinendegen, L. E., 78-5444
 Feingold, N., 78-5462
 Fenyo, E. M., 78-5774
 Fernandez-Pol, J. A., 78-5808
 Feron, V. J., 78-5571
 Fevolden, S. E., 78-5487
 Fiddler, W., 78-5622
 Figueroa, M., 78-5823
 Fikes, L. E., 78-5547
 Finch, J., 78-5805
 Fine, D. H., 78-5414, 78-5627
 Fink, R., 78-5898
 Fischer, S. M., 78-5567
 Fischinger, P. J., 78-5777
 Fisher, M. S., 78-5858
 Fitz, D. R., 78-5568
 Fleig, I., 78-5488, 78-5491
 Fleisher, G., 78-5835, 78-5836
 Fleming, A. F., 78-5921
 Fong-Cheng, C. K., 78-5815
 Fong, L. Y., 78-5587
 Fossberg, T. M., 78-5868
 Francis, S. E., 78-5908
 Frank, A. L., 78-5557
 Frank, M. L., 78-5735
 Freeman, J., 78-5434
 Frei, J. V., 78-5639
 Frenkel, G. D., 78-5845
 Friberg, K., 78-5906
 Fribush, H. M., 78-5625
 Friderici, K. H., 78-5787
 Fridlianskaia, I. I., 78-5503
 Friedberg, E. C., 78-5705, 78-5709
 Friedlander, B. R., 78-5968
 Friedman, H., 78-5763
 Friedmann, T., 78-5792
 Friedrich, U., 78-5680

Primat, P., 78-5875
 Prith, C. H., 78-5476
 Fritz, W., 78-5555
 Pu, P. P., 78-5512
 Fujiwara, K., 78-5649
 Fujiwara, M., 78-5483, 78-5484
 Fukushima, S., 78-5672
 Fukutome, T., 78-5930
 Furrer, R., 78-5629
 Furukawa, H., 78-5879
 Gaconnet, N., 78-5634
 Gaffield, W., 78-5619
 Gaffney, E. V., 78-5994
 Gahrton, G., 78-5906
 Galbraith, A. I., 78-5599
 Gallagher, R., 78-5985
 Gallagher, R. E., 78-5865, 78-5983
 Gallie, B. L., 78-5870
 Gallo, R., 78-5985
 Gallo, R. C., 78-5983
 Gamble, C., 78-5768
 Gange, R. W., 78-5855
 Gans, P. J., 78-5900
 Gardner, H. A., 78-5870
 Gardner, R. L., 78-5453
 Garon, C. F., 78-5797
 Garren, L., 78-5597
 Garrett, A. J., 78-5844
 Garry, V. F., 78-5401
 Gartner, L., 78-5822
 Gastaut, J. A., 78-5896
 Gautsch, J. W., 78-5762
 Geder, L., 78-5823
 Gelboin, H. V., 78-5566
 Genadry, R., 78-5889
 Genovese, J., 78-5765
 Georgiev, G. P., 78-5806, 78-5811
 Germond, J. E., 78-5802
 Gerodetti, M. H., 78-5796
 Gershenson, S., 78-5851
 Gershwin, M. E., 78-5538, 78-5904
 Gerwin, B. I., 78-5760
 Ghosh, B. C., 78-5912
 Ghysdael, J., 78-5788
 Giacomoni, D., 78-5856
 Gichner, T., 78-5641
 Gilden, D. H., 78-5794
 Gilden, R. V., 78-5779
 Gindhart, T., 78-5455
 Giovannetti, M., 78-5653
 Gipson, T. G., 78-5862
 Gjerris, F., 78-5925
 Gjorgov, A. N., 78-5947
 Glaser, R., 78-5832
 Gobronidze, I. A., 78-5937
 Godoy, H. M., 78-5603
 Gogeliani, G. K., 78-5937
 Goggelmann, W., 78-5417
 Goh, K., 78-5903, 78-5907
 Gokhale, J. A., 78-5540
 Goldfarb, P., 78-5819
 Goldsmith, J. R., 78-5922
 Golub, N. I., 78-5650
 Gonzales, M., 78-5951
 Gonzalez-Serva, A., 78-5815
 Goodall, C. M., 78-5605

Goodman, D. G., 78-5535
 Gopalan, H. N., 78-5635
 Gorecki, M., 78-5743
 Gorozhanskaya, G., 78-5847
 Gotlieb, S. L., 78-5916
 Goudeau, A., 78-5837
 Gough, T. A., 78-5416, 78-5591
 78-5595, 78-5621
 Goutier, R., 78-5853
 Grandgenett, D. P., 78-5742
 Grandjean, C., 78-5523
 Graves, J. M., 78-5534
 Greaves, M. F., 78-5987
 Green, G. J., 78-5869
 Green, J. A., 78-5911
 Green, L., 78-5577
 Green, M., 78-5840
 Green, M. H., 78-5480
 Green, S., 78-5682
 Green, U., 78-5575
 Greenberg, B., 78-5904
 Greenberger, J. S., 78-5900
 Greenwald, R., 78-5710
 Greim, H., 78-5417
 Grigg, G. W., 78-5433
 Grilli, S., 78-5415
 Grosjean, D., 78-5568
 Grossman, L., 78-5700
 Grover, P. L., 78-5504, 78-5511
 78-5530
 Grubbs, C. J., 78-5582
 Grundmann, E., 78-5499
 Guansing, A. R., 78-5972
 Guidotti, T. L., 78-5922
 Gurdon, J. B., 78-5805
 Gurgo, C., 78-5761
 Gurtoo, H. L., 78-5510
 Guschin, B. V., 78-5831
 Guthrie, J., 78-5675
 Gyorkey, F., 78-5454
 Haagenzen, D. E., 78-5993
 Habs, M., 78-5573
 Hadjiolov, D. C., 78-5542
 Hagen, T. C., 78-5972
 Hall, A., 78-5712
 Hall, S., 78-5968
 Hamburger, A. W., 78-5439
 Hamelin, R., 78-5771
 Hamza, H. A., 78-5656
 Hanafi, M., 78-5874
 Hananouchi, M., 78-5583
 Hansen, J. P., 78-5963
 Hara, E., 78-5479
 Harberg, J., 78-5928
 Hard, G. C., 78-5604
 Hardy, W. D., 78-5786
 Harmsen, A., 78-5925
 Harris, C. C., 78-5508, 78-5557
 78-5562, 78-5590, 78-5877
 Harte, G., 78-5719
 Hartford, W. H., 78-5923
 Hartman, P. S., 78-5708
 Harvey, R., 78-5800
 Harvey, R. G., 78-5505, 78-5512
 Haseltine, W. A., 78-5772
 Hashem, M. M., 78-5547

Hastings, K., 78-5815
 Hata, R., 78-5783
 Hatano, H., 78-5565
 Hauser, E., 78-5618
 Havery, D. C., 78-5623
 Hayami, M., 78-5740
 Hayashi, N., 78-5631
 Hayata, I., 78-5717
 Hayes, J. R., 78-5473
 Hazdra, J. J., 78-5981
 Hearne, T., 78-5968
 Heintz, N. H., 78-5471
 Heise, H., 78-5822
 Heiz, H. J., 78-5618
 Heller, P., 78-5856
 Hellstrom, I., 78-5524
 Hellstrom, K. E., 78-5524
 Henderson, B. E., 78-5945
 Henle, G., 78-5443, 78-5817, 78-5830
 78-5835, 78-5836
 Henle, W., 78-5443, 78-5817, 78-5830
 78-5835, 78-5836
 Henriksen, O., 78-5812
 Hensley, G. T., 78-5710
 Hermens, A. F., 78-5690
 Herschman, H. R., 78-5813
 Hewer, A., 78-5530
 Heyns, K., 78-5610
 Hicks, R. M., 78-5591
 Hidaka, K., 78-5588
 Hidmark, A., 78-5533
 Hilf, R., 78-5545
 Hill, M. J., 78-5632
 Hill, R. M., 78-5434
 Hillesund, t., 78-5550
 Hinuma, Y., 78-5833
 Hirakawa, T., 78-5552
 Hirao, F., 78-5532
 Hirayama, T., 78-5946
 Hirono, I., 78-5465
 Hirose, M., 78-5481, 78-5583
 Hirsch, E., 78-5882
 Ho, H. C., 78-5830
 Ho, T., 78-5514
 Hoffmann, D., 78-5409
 Hofmann, R., 78-5822
 Hogstedt, C., 78-5969
 Holben, J. A., 78-5755
 Hong, W. K., 78-5869
 Hong, Y. S., 78-5478
 Hopkins, P. S., 78-5854
 Hopkins, R. F., 78-5825
 Hoppe, H., 78-5659
 Horikoshi, T., 78-5506
 Horta, J., 78-5973
 Horta, M. E., 78-5973
 Hosoda, K., 78-5672
 Hotz, G., 78-5651
 Hovis, J. S., 78-5613
 Howard, J. W., 78-5623
 Howarth, J. L., 78-5722
 Howell, A. N., 78-5807
 Howett, M. K., 78-5821
 Hsiung, G. D., 78-5815
 Hu, S. S., 78-5747
 Huang, D. P., 78-5830

- Huang, E. S., 78-5885
 Hubbard, S. A., 78-5480
 Huberman, E., 78-5518
 Huff, J. E., 78-5421
 Hug, O., 78-5444
 Hughes, R. C., 78-5844
 Hugo, N. E., 78-5714
 Hui, G., 78-5600, 78-5607
 Hundeshagen, H., 78-5718
 Hunma, R. R., 78-5617
 Hunt, R. D., 78-5816
 Hurtubise, R. J., 78-5549
 Hwang, L. T., 78-5856
 Hyatt, C. L., 78-5864
 Hylden, J. L., 78-5556
 Hyman, R. W., 78-5823
 Hynds, P. M., 78-5568
 Icart, J., 78-5829
 Ichimura, S., 78-5586
 Ignjatovic, J., 78-5740
 Ihle, J. N., 78-5759
 Iizuka, T., 78-5586
 Ikeda, R., 78-5538
 Ikeda, T., 78-5580
 Iltis, J. P., 78-5823
 Ilus, T., 78-5466
 Imamura, M., 78-5862
 Inaba, Y., 78-5966
 Joachim, H. L., 78-5769
 Isaacson, P., 78-5887
 Ishihara, T., 78-5717
 Ishii, Y., 78-5686
 Ishiwata, H., 78-5631
 Ishizawa, M., 78-5588
 Ismailov, A. Kh., 78-5948
 Ito, N., 78-5481, 78-5583, 78-5670
 78-5672
 Itzhaki, R. F., 78-5599
 Ivanetich, K. M., 78-5490
 Ivanidze, T. G., 78-5937
 Ivankovic, S., 78-5573
 Iversen, O. H., 78-5584
 Iwai, M., 78-5752
 Iwai, Y., 78-5752
 Iwanaga, T., 78-5879
 Iwasaki, I., 78-5506
 Izak, G., 78-5920
 Jackson, F., 78-5508
 Jacobi, W., 78-5444
 James, E., 78-5775
 Janiaud, P., 78-5519
 Jarrett, O., 78-5785
 Jarrett, W. F., 78-5790
 Javadpour, N., 78-5864
 Jedrychowski, W., 78-5941
 Jeffrey, A. M., 78-5562
 Jeng, Y. H., 78-5840
 Jensen, F. C., 78-5762
 Jerina, D. M., 78-5548
 Jeter, J. R., 78-5482
 Johannesson, G., 78-5951
 Johansson, K., 78-5838
 Johnson, G. R., 78-5984
 Johnson, R. H., 78-5854
 Joncas, J. H., 78-5824
 Jones, E. W., 78-5855
 Jones, P. W., 78-5501
 Jones, S. M., 78-5633
 Jorgenson, T. A., 78-5682
 Jun, M. H., 78-5854
 Kada, T., 78-5676
 Kagan, B. M., 78-5434
 Kaiser, R., 78-5432
 Kalland, T., 78-5868
 Kalser, M. H., 78-5710
 Kamataki, T., 78-5564
 Kamien-Sarbinowsky, M., 78-5701
 Kaminski, R., 78-5967
 Kanematsu, N., 78-5676
 Kanisawa, M., 78-5468, 78-5522
 Kannan, R., 78-5547
 Kanter, P., 78-5510
 Kaplan, H. S., 78-5442, 78-5756
 Karle, J. M., 78-5548
 Karlowski, K., 78-5636
 Karpas, A., 78-5900
 Kato, K., 78-5465
 Kato, R., 78-5564
 Kato, S., 78-5749
 Kato, T., 78-5464
 Katowski, P., 78-5561
 Katorkin, E. N., 78-5883
 Katsuki, H., 78-5522, 78-5669
 Katz, I. D., 78-5490
 Katzmann, J., 78-5856
 Kauffman, R. E., 78-5434
 Kaufman, D. G., 78-5554
 Kawabata, A., 78-5580
 Kawabata, T., 78-5611
 Kawachi, T., 78-5586, 78-5683
 Kawada, K., 78-5902
 Kawajiri, K., 78-5479
 Kawazoe, Y., 78-5670
 Keane, A. T., 78-5729
 Keefer, L. K., 78-5626
 Kellen, J. A., 78-5539
 Keller, A. Z., 78-5938
 Kemel, S., 78-5910
 Kemp, M. C., 78-5781
 Kennedy, A., 78-5563
 Ketkar, M., 78-5575
 Kettmann, R., 78-5788
 Khodosova, I. A., 78-5503
 Kidson, C., 78-5860
 Kidwell, W. R., 78-5543
 Kieffer, R. F., 78-5877
 Kikuchi, I., 78-5935
 Kim, H. J., 78-5972
 Kimball, R. F., 78-5679
 Kimoto, T., 78-5730
 Kimura, M., 78-5852
 Kinchin, G. H., 78-5446
 King, V., 78-5900
 Kirsh, I. E., 78-5729
 Kishimoto, S., 78-5532
 Kister, S. J., 78-5993
 Kitamura, M., 78-5546
 Kitamura, Y., 78-5988
 Kitsak, V. Y., 78-5793
 Kleihues, P., 78-5536
 Klein, D., 78-5634
 Klein, E., 78-5774
 Klein, G., 78-5834
 Klenina, N. V., 78-5746
 Klinken, L., 78-5925
 Knaup, G., 78-5798
 Knazek, R., 78-5543
 Knight, L. A., 78-5870
 Knudson, G. B., 78-5568
 Kobayashi, M., 78-5608
 Kobayashi, N., 78-5522
 Koch, G. L., 78-5863
 Kodama, Y., 78-5909
 Kohno, S., 78-5717
 Kolber, S., 78-5984
 Kolonel, L. N., 78-5943
 Kondo, H., 78-5580
 Kondo, S., 78-5671
 Konigsberg, W. H., 78-5758
 Konishi, Y., 78-5580
 Korprowski, H., 78-5794
 Korovin, R. N., 78-5750
 Kositsyna, M. V., 78-5948
 Kostiukovskii, I. L., 78-5413
 Kostiukovskii, Ia. L., 78-5624
 Koyama, H., 78-5879
 Krakower, J. M., 78-5850
 Kramer, U., 78-5940
 Krech, R., 78-5688, 78-5689
 Kreyberg, L., 78-5931
 Kripke, M. L., 78-5858
 Krsmanovic-Simic, D., 78-5681
 Krull, I. S., 78-5627
 Kruysse, A., 78-5571
 Kudo, H., 78-5902
 Kukushkina, L. M., 78-5691
 Kumar, H. D., 78-5699
 Kumatori, T., 78-5717
 Kunkel, M., 78-5822
 Kurata, H., 78-5631
 Kuriya, S., 78-5902
 Kuroki, T., 78-5593
 Kusyk, C., 78-5454
 Kuwata, T., 78-5809
 Kwiram, A. L., 78-5474
 Kyrtopoulos, S. A., 78-5617
 Lagarriga-Attias, J., 78-5980
 Lahoutte, C., 78-5875
 Lai, M. M., 78-5747
 Laird, H. M., 78-5790
 Lakowicz, J. R., 78-5533, 78-5556
 Lambert, R., 78-5942
 Lamki, H., 78-5888
 Lamy, M. E., 78-5886
 Land, P. C., 78-5629
 Lange, C. E., 78-5435
 Langelier, Y., 78-5818
 Lapin, B., 78-5817
 LaPorte, P., 78-5792
 Larimer, F. W., 78-5514
 Larsen, C. J., 78-5771
 Larsson, S. E., 78-5728
 Laskey, R. A., 78-5805
 Launais, B., 78-5905
 Laurentaci, G., 78-5527
 Lauen-Wetje, A. K., 78-5609
 Lavelle, D., 78-5856
 Lavin, M. F., 78-5860

aw, L. W., 78-5812
 awley, P. D., 78-5639
 e Bras, M., 78-5962
 edinko, N., 78-5846
 ee, A. K., 78-5890
 ee, H., 78-5907
 ee, J., 78-5945
 eeper, J. P., 78-5949
 eeper, R. D., 78-5715
 egator, M., 78-5427
 eifer, A., 78-5678
 empert, N., 78-5895
 ennette, E. T., 78-5835, 78-5836
 enzin, A., 78-5894
 eonard, A., 78-5489
 eonard, E. D., 78-5489
 erner, R. A., 78-5762
 evey, G. S., 78-5500
 evi, P., 78-5519
 evin, W., 78-5548
 evine, L., 78-5509
 ewin, M. R., 78-5861
 ewis, A. M., 78-5838
 ewis, P., 78-5738
 eyritz, M., 78-5824
 iber, H. L., 78-5659
 iebberman, M., 78-5756
 iebeskind, D., 78-5678
 ijinsky, W., 78-5572, 78-5589, 78-5598
 im, J. K., 78-5658
 in, C. C., 78-5965
 Lindahl, T., 78-5663
 Lindop, P. J., 78-5720
 Lindquist, R., 78-5906
 Linehan, J. J., 78-5955
 Link, J., 78-5873
 Linnainmaa, K., 78-5466
 Lippman, M. E., 78-5543
 Littman, L., 78-5882
 Littman, M. S., 78-5729
 Liu, Y. C., 78-5474
 Lloyd, D. C., 78-5725
 LLoyd, R. D., 78-5731, 78-5732
 78-5733, 78-5734
 Loepky, R. N., 78-5579
 Loken, H. F., 78-5713
 Long, F. C., 78-5498
 Longcope, C., 78-5991
 Loprieno, N., 78-5421
 Lorentzon, R., 78-5728
 Lotlikar, P. D., 78-5478
 Loubiere, R., 78-5962
 Louria, D. B., 78-5974
 Loven, D. P., 78-5534
 Lowy, D. R., 78-5784
 Lozzio, B. B., 78-5986
 Lozzio, C. B., 78-5986
 Luchnik, N., 78-5692
 Lui, M., 78-5830
 Lukanidin, E. M., 78-5806
 Lundin, R. E., 78-5619
 Lung, B., 78-5890
 Luthra, U. K., 78-5959
 Lutz, W. K., 78-5553
 Lyle, G. G., 78-5625
 Lyle, R. E., 78-5625

Lynch, H. T., 78-5459
 Lynch, P. M., 78-5459
 Ma, C. Y., 78-5628
 MacGregor, J. E., 78-5956, 78-5957
 Machado, E. A., 78-5986
 MacPhee, D. G., 78-5701
 Magee, P. N., 78-5411
 Maguire, D. J., 78-5854
 Maher, V. M., 78-5704
 Maier, P., 78-5677
 Maisin, J. R., 78-5853
 Mak, T. W., 78-5768
 Malaveille, C., 78-5492
 Malling, H. V., 78-5693
 Malyshev, Iu. I., 78-5883
 Mammerickx, M., 78-5788
 Manny, N., 78-5920
 Manser, P., 78-5677
 Marineau, E. J., 78-5753
 Mark, R., 78-5867
 Markov, D. V., 78-5542
 Marmor, M., 78-5932
 Marsh, S., 78-5523
 Martin, G. R., 78-5842, 78-5843
 Martin, W. J., 78-5862
 Marty, M., 78-5771
 Maruchi, N., 78-5966
 Mason, W. S., 78-5736
 Mass, M. J., 78-5554
 Massey, R. C., 78-5630
 Masubuchi, K., 78-5953
 Masuda, A., 78-5479
 Matsuda, H., 78-5749
 Matsui, Y., 78-5902
 Matsukura, N., 78-5683
 Matsumoto, K., 78-5988
 Matsumoto, S., 78-5649
 Matsumura, K., 78-5569
 Matter, B. E., 78-5428
 Mattern, I. E., 78-5517
 Maupas, P., 78-5837
 Mawhinney, H., 78-5888
 Mays, C. W., 78-5731, 78-5732
 78-5733, 78-5734
 Mazoujian, G., 78-5993
 Mazurenko, N. P., 78-5831
 McCormick, J. J., 78-5704
 McDowell, E. M., 78-5557, 78-5876
 78-5877, 78-5878
 McFarland, S., 78-5732
 McFarland, S. S., 78-5734
 McGrath, C. M., 78-5753
 McGuire, L. J., 78-5592
 McLaughlin, D., 78-5523
 McLaughlin, J. S., 78-5877
 McMichael, A. J., 78-5939
 McPhail, M. F., 78-5595
 McWeeny, D. J., 78-5630
 Meager, A., 78-5844
 Medina, D., 78-5754
 Medvedev, F. A., 78-5624
 Meek, E. S., 78-5534
 Melamed, D. B., 78-5413, 78-5624
 Melcher, D. H., 78-5955
 Menczer, J., 78-5958
 Mendel, J., 78-5637

Menezes, J., 78-5827
 Merabishvili, V. M., 78-5460
 Merenyl, D. K., 78-5877
 Metcalf, D., 78-5984
 Metzgar, R. S., 78-5865
 Mgaloblishvili, G. N., 78-5937
 Michalides, R., 78-5751
 Mikhalevich, L. S., 78-5691
 Mikkelsen, F., 78-5963
 Millard, R. J., 78-5897
 Miller, A. B., 78-5445
 Miller, D. R., 78-5567
 Miller, G., 78-5907
 Miller, J. A., 78-5477
 Miller, R. W., 78-5405
 Milleville, M., 78-5843
 Minkler, J., 78-5687
 Mironescu, S. G., 78-5685
 Mirvish, S. S., 78-5636
 Mishra, K. P., 78-5565
 Mitchell, R. H., 78-5563
 Mitra, A. B., 78-5959
 Miyagami, M., 78-5913
 Miyoshi, T., 78-5497
 Mizuno, D., 78-5857
 Mizushiro, H., 78-5631
 Modan, B., 78-5958
 Moelling, K., 78-5741
 Mohanakumar, T., 78-5865
 Mohr, U., 78-5570
 Mole, R. H., 78-5720
 Moller, M. E., 78-5487
 Moloney, W. C., 78-5900
 Molzen, K. B., 78-5996
 Monaco, M. E., 78-5543
 Monnier, J., 78-5905
 Montesano, R., 78-5593
 Moon, R. C., 78-5582
 Moore, D. H., 78-5755
 Moreau, P., 78-5448
 Mori, H., 78-5465
 Morioka, M., 78-5497
 Moriyama, M., 78-5683
 Moriyasu, N., 78-5913
 Morreal, C. E., 78-5544
 Morrison, F. S., 78-5901
 Morse, L. S., 78-5820
 Mortelmans, K., 78-5705
 Mossman, B. T., 78-5521
 Motta, L. C., 78-5973
 Moyer, M. P., 78-5796
 Moyer, R. C., 78-5796
 Mueller, J. C., 78-5880
 Mukubo, K., 78-5528
 Muller, H., 78-5789
 Murasaki, G., 78-5481, 78-5583
 Murphy, J., 78-5790
 Muse, M. B., 78-5900
 Muth, H., 78-5444
 Nachtwey, D. S., 78-5975
 Nafstad, I., 78-5606
 Nagao, M., 78-5559
 Nagasawa, H., 78-5995
 Nagata, C., 78-5565
 Nairn, R., 78-5844
 Nakadate, M., 78-5683

- Nakamura, S., 78-5913
 Nakanishi, K., 78-5481
 Namba, M., 78-5730
 Narisawa, T., 78-5684
 Nass, K., 78-5845
 Natarajan, A. T., 78-5660
 Nawar, M. M., 78-5656
 Nazerian, K., 78-5748
 Nedospasov, S. A., 78-5811
 Neenan, J. P., 78-5981
 Neidbailik, I. E., 78-5750
 Nelson, N., 78-5410
 Nemoto, H., 78-5953
 Nemoto, N., 78-5552, 78-5559, 78-5566
 Nettesheim, P., 78-5640, 78-5642
 78-5978
 Neubauer, R. H., 78-5825
 Neubort, S., 78-5678
 Neufeld, H., 78-5910
 Neumann, H. G., 78-5472
 Neuser, D., 78-5461
 Neuvonen, E., 78-5791
 Newberne, P. M., 78-5473, 78-5587
 Newell, G. W., 78-5682
 Newman, M. S., 78-5547
 Nichols, W. W., 78-5804
 Nicholson, G. L., 78-5979
 Nicolov, I. G., 78-5961
 Nielsen, N. H., 78-5963
 Nilsson, A., 78-5728
 Nishi, M., 78-5581
 Nishie, K., 78-5594
 Nishikawa, H., 78-5532
 Nishioka, Y., 78-5764
 Nishitani, K., 78-5730
 Nix, C. E., 78-5514
 Njagi, G. D., 78-5635
 Nomura, A., 78-5945
 Nomura, K., 78-5499
 Nomura, T., 78-5902
 Nonoyama, M., 78-5817, 78-5832
 Noonan, C. A., 78-5810
 Noronha, R. F., 78-5605
 Nosaka, Y., 78-5565
 Nukaga, H., 78-5506
 Nusse, R., 78-5751
 O'Connell, A., 78-5837
 O'Neil, B. W., 78-5790
 Oakes, J. E., 78-5823
 Oberender, H., 78-5822
 Oberg, B., 78-5841
 Oberhausen, E., 78-5444
 Obidoa, O., 78-5469
 Ocho, M., 78-5809
 Oda, T., 78-5809
 Oelsner, G., 78-5958
 Ogura, H., 78-5809
 Ogura, T., 78-5532
 Ohhara, T., 78-5497
 Ohshima, H., 78-5611
 Ohtawa, M., 78-5674
 Ohuchi, K., 78-5509
 Okada, M., 78-5670
 Okada, S., 78-5696
 Okamoto, S., 78-5988
 Okamoto, T., 78-5522, 78-5669
 Okigaki, T., 78-5551
 Okita, M., 78-5522
 Okuda, M., 78-5935
 Okumoto, M., 78-5752
 Okun, J. D., 78-5612
 Olson, J., 78-5889
 Ong, T., 78-5675
 Onnekink, C., 78-5778
 Ono, T., 78-5696
 Oroszlan, S., 78-5779
 Osato, T., 78-5828
 Osborne, J. W., 78-5534
 Ostro, M., 78-5856
 Otsuji, N., 78-5588
 Ouchi, G., 78-6000
 Outram, J. R., 78-5617
 Oviedo, M. A., 78-5714
 Padieu, P., 78-5519
 Painter, R. B., 78-5673
 Pal'chik, E. A., 78-5948
 Pal, K., 78-5511
 Palitti, F., 78-5694
 Palmer, F. J., 78-5914
 Panda, B. B., 78-5467
 Panet, A., 78-5743
 Paquin, L., 78-5455
 Parmley, T., 78-5889
 Paroly, W., 78-5873
 Parry, J. M., 78-5654
 Parsons, J. T., 78-5738
 Pascone, M., 78-5527
 Pastorelo, E. F., 78-5916
 Patel, P., 78-5827
 Patil, K., 78-5523
 Paucha, E., 78-5800, 78-5801
 Pavlis, R., 78-5667
 Pearlman, W. H., 78-5993
 Pearse, A., 78-5769
 Pegg, A. E., 78-5600, 78-5601, 78-5607
 Peng, L. H., 78-5993
 Penhoet, E. E., 78-5770
 Penrose, D., 78-5768
 Perdue, S. W., 78-5679
 Pereira, L., 78-5820
 Perez-Arroyo, R., 78-5980
 Peries, J., 78-5771
 Persson, H., 78-5838, 78-5841
 Perticone, P., 78-5694
 Peterkofsky, B., 78-5783
 Petkova-Bocharova, T., 78-5961
 Petrakis, N. L., 78-5891
 Pettersson, U., 78-5838
 Pettit, G. R., 78-5757
 Petukhov, V. L., 78-5982
 Pfeiffer, E. H., 78-5403
 Pfleiderer, G., 78-5798
 Philipson, L., 78-5838, 78-5841
 Phillip, J. D., 78-5549
 Phillips, D. H., 78-5504
 Pienta, R. J., 78-5550, 78-5977
 Pignatelli, B., 78-5597
 Pigott, D., 78-5994
 Pinkerton, J. H., 78-5888
 Piotrowski, E. G., 78-5622
 Pitts, J. N., 78-5568
 Planche, G., 78-5492
 Podolnaya, M., 78-5424
 Poiley, J. A., 78-5550, 78-5977
 Pokorný, V., 78-5641
 Pokrovskii, A. A., 78-5624
 Pollack, S. B., 78-5866
 Pollard, M., 78-5494
 Popiela, T., 78-5941
 Poullain, B., 78-5634
 Pound, A. W., 78-5592
 Prabhakar, A. K., 78-5959
 Prat, M., 78-5744
 Pratt, J. H., 78-5991
 Preston-Martin, S., 78-5926
 Preussmann, R., 78-5406
 Pris, J., 78-5905
 Prodi, G., 78-5415
 Pruitt, A. W., 78-5434
 Pruss, R. M., 78-5813
 Purrott, R. J., 78-5725
 Purtilo, D. T., 78-5455
 Pyrhonen, S., 78-5791
 Rabin, H., 78-5825
 Radchenko, L., 78-5424
 Radhakrishnan, J., 78-5912
 Rafferty, K. A., 78-5799
 Raineri, R., 78-5550
 Rajalakshmi, S., 78-5646
 Rajewsky, M. F., 78-5648
 Ramalingaswami, V., 78-5964
 Ramos de Uribe, M. H., 78-5431
 Rands, E., 78-5784
 Rao, K. V., 78-5540
 Rao, M. S., 78-5574
 Rao, P. M., 78-5646
 Rao, T. K., 78-5514, 78-5589
 Rapp, F., 78-5821, 78-5823
 Rapp, U. R., 78-5767
 Ratner, P. L., 78-5775
 Rattle, H., 78-5530
 Ratz, K. H., 78-5822
 Ravry, M. J., 78-5874
 Reddy, B. S., 78-5644, 78-5684
 Reddy, J. K., 78-5574
 Reeder, E. J., 78-5725
 Reeson, D. E., 78-5844
 Rehfeld, S. J., 78-5713
 Reiffers, J., 78-5430
 Rein, A., 78-5760
 Reinacher, M., 78-5789
 Reissland, J. A., 78-5720
 Reske-Nielsen, E., 78-5925
 Reusch, K., 78-5419
 Revazova, J., 78-5424
 Revesz, L., 78-5728
 Reyes, F., 78-5899
 Reynolds, R. K., 78-5849
 Reznik-Schuller, H., 78-5570, 78-5585
 Rhim, J. S., 78-5507
 Rho, J. H., 78-5551
 Rhoads, J. E., 78-5867
 Rhodes, P., 78-5957
 Riazuddin, S., 78-5663, 78-5700
 Richards, F. F., 78-5758
 Rimsten, A., 78-5989
 Rinkus, S., 78-5427
 Roche, J. K., 78-5885

- Rodhain, J., 78-5886
 Roeder, R. G., 78-5839
 Rogan, E., 78-5561
 Rogers, K. J., 78-5607
 Rogers, M. J., 78-5812
 Rohlen, O., 78-5969
 Roizman, B., 78-5820
 Roller, P. P., 78-5626
 Rollo, D., 78-5713
 Romon, M., 78-5875
 Roper, H., 78-5610
 Rosenfeld, P. S., 78-5972
 Ross, R. D., 78-5627
 Roth, D., 78-5706
 Roth, H. J., 78-5609
 Roth, R., 78-5523, 78-5561
 Rottman, F., 78-5787
 Rottner, B., 78-5929
 Rovere, L. E., 78-5500
 Ruben, R. L., 78-5799
 Rubsamen, H., 78-5740
 Ruddell, W. S., 78-5632
 Rudolph, R. L., 78-5789
 Rule, J. E., 78-5476
 Rundel, R. D., 78-5975
 Ruscetti, F., 78-5985
 Ruscetti, F. W., 78-5983
 Rushbrook, C. J., 78-5682
 Russell, P. H., 78-5785
 Russo, I. H., 78-5537
 Russo, J., 78-5537
 Ryan, D., 78-5548
 Saavedra, J. E., 78-5578, 78-5625
 Sabbath, M., 78-5769
 Sager, R., 78-5807
 Sahaphong, S., 78-5602
 Sahu, R. K., 78-5467
 Sakai, K., 78-5669
 Sakamoto, K., 78-5833
 Sakka, M., 78-5976
 Salas, J., 78-5944
 Salmon, J. M., 78-5558
 Salmon, S. E., 78-5439
 Saltzman, D. A., 78-5880
 Salzman, N. P., 78-5797
 Samaan, N. A., 78-5990
 Sampson, S., 78-5958
 Sams, J. P., 78-5636
 Sanford, K. K., 78-5997
 Sankaran, V., 78-5547
 Sanner, C. J., 78-5880
 Sano, T., 78-5683, 78-5902
 Sansone, E. B., 78-5596
 Sarma, D. S., 78-5646
 Sasaki, M., 78-5909
 Sasaoka, I., 78-5465
 Sato, K., 78-5506
 Sato, T., 78-5749
 Sauer, W. G., 78-5884
 Savage, A., 78-5633
 Saw, D., 78-5830
 Sawyers, T. M., 78-5914
 Scanlon, E. F., 78-5714
 Schachner, S. H., 78-5712
 Schaeffer, W. I., 78-5471
 Schaffer, P. A., 78-5820
 Scharfman, W. B., 78-5895
 Scherrer, M., 78-5934
 Schiff, R. D., 78-5742
 Schlake, W., 78-5499
 Schlipkoter, H. W., 78-5940
 Schmahl, D., 78-5573, 78-5598
 Schmid, J. P., 78-5568
 Schmidt, C. G., 78-5420
 Schmidt, J. M., 78-5757
 Schneiderman, M. A., 78-5408
 Schudel, A., 78-5817
 Schuller, P. L., 78-5615
 Schultz, P. N., 78-5990
 Schuman, R. F., 78-5977
 Schwarz, H., 78-5637
 Schweitzer, R., 78-5873
 Scolnick, E. M., 78-5535, 78-5784
 Scott, A., 78-5577
 Sedunov, Z. A., 78-5750
 Segal, S., 78-5434
 Segawa, M., 78-5804
 Seidman, M. M., 78-5797
 Sekiguchi, K., 78-5794
 Sen, A., 78-5814
 Senior, A. E., 78-5545
 Senner, K. R., 78-5655
 Serrou, B., 78-5558
 Shabad, L. M., 78-5650
 Shaikh, R., 78-5736
 Shamsuddin, A. K., 78-5508
 Sharma, C. B., 78-5467
 Sharma, J. M., 78-5748
 Sharon, Z., 78-5958
 Sherman, B. M., 78-5949
 Sherr, C. J., 78-5814
 Shetty, K. R., 78-5972
 Shields, R., 78-5452
 Shifrine, M., 78-5904
 Shiloh, Y., 78-5422
 Shimojo, H., 78-5848
 Shirai, T., 78-5583, 78-5672
 Shoji, Y., 78-5580
 Shors, E. C., 78-5569
 Shraiber, D., 78-5638
 Shvedova, L. A., 78-5892
 Siciliano, M. J., 78-5454
 Siegert, W., 78-5774
 Silverstein, S., 78-5764
 Simmons, J. W., 78-5702
 Simpson, T. L., 78-5999
 Sims, P., 78-5504, 78-5511, 78-5530
 Singer, B., 78-5648
 Sinha, B. D., 78-5699
 Sinkovics, J. G., 78-5454
 Skaare, J. U., 78-5606
 Skar, G. T., 78-5549
 Skinhoj, P., 78-5963
 Skopek, T. R., 78-5659
 Skorikova, A. S., 78-5831
 Skurkovich, S. V., 78-5831
 Slaga, T. J., 78-5567
 Sliski, A. H., 78-5786
 Smith, A. E., 78-5800, 78-5801
 Smith, M. J., 78-5863
 Smith, P. D., 78-5661
 Smith, R., 78-5801
 Smith, R. D., 78-5545
 Smuckler, E. A., 78-5474
 Snart, R. S., 78-5576
 Sobrinho-Simoes, M. A., 78-5918
 Solt, D. B., 78-5516
 Sonenberg, M., 78-5715
 Sonntag, R., 78-5894
 Soos, K., 78-5555
 Sorsa, M., 78-5466
 Soyka, L. F., 78-5434
 Spangler, W., 78-5904
 Spann, M. L., 78-6000
 Specter, S., 78-5763
 Speedy, G., 78-5520
 Spiegel, J., 78-5716
 Spirtas, R., 78-5967
 Sporn, M. B., 78-5582
 Squire, R. A., 78-5582
 Sridhar, R., 78-5667
 Stanton, P. E., 78-5874
 Stavrou, D., 78-5859
 Stein, D., 78-5974
 Stein, G., 78-5435
 Steinberg, R., 78-5680
 Steinitz, M., 78-5834
 Steintiz, R., 78-5958
 Stemmermann, G. N., 78-5943
 Stepanov, S. V., 78-5726
 Stephany, R. W., 78-5614, 78-5615
 Stephens, E. A., 78-5748
 Stephenson, J. R., 78-5849
 Stetka, D. G., 78-5687
 Stevens, R. H., 78-5534
 Stevens, W., 78-5733
 Stewart, A., 78-5447
 Stewart, L. A., 78-5908
 Stoichev, I. I., 78-5961
 Stoner, G. D., 78-5508
 Stortebecker, P., 78-5927
 Stoyanov, I. S., 78-5961
 Streilein, J. W., 78-5440
 Strickland, P., 78-5723
 Strominger, J. L., 78-5826
 Strong, L. C., 78-5458
 Stroop, J., 78-5713
 Sucena, R. C., 78-5970
 Sugimura, T., 78-5426, 78-5683
 Sugiura, M., 78-5564
 Summers, J., 78-5837
 Sun, N., 78-5873
 Sunderman, F. W., 78-5429
 Suzuki, S., 78-5468
 Suzuki, T., 78-5935
 Swelstad, J. A., 78-5714
 Swenson, D. H., 78-5639
 Syrotuck, J. A., 78-5525
 Szent-Gyorgyi, A., 78-5998
 Tada, M., 78-5670, 78-5670
 Tagashira, Y., 78-5479
 Takada, K., 78-5828
 Takahashi, E. Y., 78-5966
 Takahashi, K., 78-5670
 Takahashi, M., 78-5672
 Takami, T., 78-5531
 Takamori, Y., 78-5752

- Takayama, S., 78-5483, 78-5484
 78-5552, 78-5559, 78-5566
 Takehisa, S., 78-5475
 Takemoto, K. K., 78-5812
 Tamaki, N., 78-5649
 Tanaka, R., 78-5794
 Tanaka, T., 78-5809
 Tandon, B. N., 78-5964
 Tandon, H. D., 78-5964
 Taniguchi, H., 78-5879
 Tanimura, A., 78-5631
 Taniyama, T., 78-5532
 Tannenbaum, S. R., 78-5577, 78-5613
 Taomoto, K., 78-5649
 Tapper, D. P., 78-5803
 Tarleton, A. E., 78-5950, 78-5996
 Tateishi, R., 78-5879
 Tatematsu, M., 78-5481
 Tate, A. D., 78-5695
 Taulbee, J., 78-5954
 Tavares, M. H., 78-5973
 Tavitian, A., 78-5771
 Taylor, A. M., 78-5697
 Taylor, G. N., 78-5731, 78-5732
 78-5733
 Taylor, H. W., 78-5572
 Taylor, J. C., 78-5924
 Taylor, W. G., 78-5997
 Tazima, Y., 78-5721
 Tchan, Y. T., 78-5608
 Tei, N., 78-5546
 Temcharoen, P., 78-5470
 Teper, S., 78-5956
 Terzaghi, M., 78-5978
 Tewari, Y. B., 78-5596
 Thamavit, W., 78-5602
 Thiel, W., 78-5789
 Thierry, C., 78-5558
 Thiess, A. M., 78-5488, 78-5491
 Thilly, W. G., 78-5659
 Thind, I. S., 78-5974
 Thomas, G. H., 78-5648
 Thomas, J. K., 78-5724
 Thomas, P. E., 78-5548
 Thomason, A. R., 78-5787
 Thompson, H. J., 78-5582
 Thompson, J. E., 78-5999
 Thon, W., 78-5648
 Thoren, L., 78-5989
 Thorley-Lawson, D., 78-5826
 Thorling, E. B., 78-5698
 Thorne, J., 78-5576
 Thorpe, J. J., 78-5437
 Thune, P., 78-5703
 Thust, R., 78-5637
 Tibbetts, C., 78-5838
 Ticolat, R., 78-5962
 Tierney, B., 78-5511, 78-5530
 Timbury, M., 78-5819
 Ting, R. C., 78-5773, 78-5812
 Tippins, R. S., 78-5654
 Todaro, G. J., 78-5767, 78-5814
 Toji, L. H., 78-5804
 Tominaga, T., 78-5546
 Topisirovic, L., 78-5681
 Torpier, G., 78-5842
 Toth, B., 78-5493
 Toth, F. D., 78-5782
 Trabalka, J. R., 78-5735
 Travis, M., 78-5756
 Treloar, A. E., 78-5949
 Tronick, S. R., 78-5772
 Troup, G. M., 78-5722
 Troy, F. A., 78-5774
 Truhaut, R., 78-5412
 Trump, B. F., 78-5508, 78-5557
 78-5562, 78-5590, 78-5876, 78-5877
 78-5878
 Trux, F., 78-5898
 Tsangaris, N., 78-5716
 Tsuchimoto, T., 78-5428
 Tucker, R. W., 78-5997
 Tulinus, H., 78-5951
 Tulliez, M., 78-5899
 Tutton, P. J., 78-5496
 Uchida, N., 78-5988
 Ueki, K., 78-5794
 Ueland, P. M., 78-5868
 Uhm, S. J., 78-5626
 Umezawa, K., 78-5559
 Uribe, M., 78-5431
 Ushimaru, Y., 78-5465
 Uzunov, P., 78-5529
 Vaage, J., 78-5754
 Vaczi, L., 78-5782
 Vainio, H., 78-5436
 Vajrasthira, S., 78-5602
 Valcovic, L. R., 78-5693
 van Broekhoven, L. W., 78-5614
 Van Cauwenberghe, K. A., 78-5568
 Van de Ven, W. J., 78-5778, 78-5780
 van Deemter, L., 78-5751
 Van Den Engh, G., 78-5987
 van Haaften, C., 78-5557
 van Nie, R., 78-5751
 Van Puymbroeck, S., 78-5727
 van Zaane, D., 78-5780
 Vanderborcht, O. L., 78-5727
 Vanderlaan, M., 78-5723
 Varghese, A. J., 78-5629
 Vaught, J. B., 78-5510
 Vegelius, J., 78-5989
 Veleminsky, J., 78-5641
 Velicer, L. F., 78-5787
 Veltman, G., 78-5435
 Venitt, S., 78-5425
 Venkitachalam, P. S., 78-5882
 Verigina, A. D., 78-5917
 Vermorken, A. J., 78-5778, 78-5780
 Vernant, J. P., 78-5899
 Vesely, D. L., 78-5500
 Vessey, M. P., 78-5423
 Viaje, A., 78-5567
 Viallet, P., 78-5558
 Vickers, S. M., 78-5434
 Vig, B. K., 78-5664
 Vitorgan, E., 78-5847
 Viviani, A., 78-5553
 Vlasenkova, N. K., 78-5793
 Vlasov, V., 78-5529
 Vogel, E., 78-5660
 Vogt, M., 78-5762
 Vogt, P. K., 78-5745, 78-5747
 Voyles, B. A., 78-5753
 Wada, A., 78-5879
 Waksvik, H., 78-5703
 Walker, E. A., 78-5597, 78-5616
 Walker, I. G., 78-5667
 Walker, W., 78-5911
 Wallace, R. B., 78-5949
 Walters, C. L., 78-5591, 78-5632
 Warner, N. L., 78-5987
 Warren, W., 78-5639
 Warzok, R., 78-5637
 Wassom, J. S., 78-5421
 Wataki, K., 78-5749
 Watanabe, K., 78-5631, 78-5644
 Watkins, P. A., 78-5449
 Webb, K. S., 78-5595
 Weber, W. W., 78-5407
 Weinerman, B. H., 78-5910
 Weisburger, J. H., 78-5644, 78-5684
 Weitzner, S., 78-5901
 Wells, S. A., 78-5993
 Welsch, C. W., 78-5992
 Wenk, M. L., 78-5535
 Whong, W., 78-5675
 Wide, L., 78-5989
 Williams, D. C., 78-5560
 Williams, M. L., 78-5978
 Wilson, F. D., 78-5904
 Wilson, S., 78-5972
 Wipke, W. T., 78-6000
 Wise, K. S., 78-5781
 Wishnok, J. S., 78-5613
 Witte, O. N., 78-5776
 Witter, R. L., 78-5748
 Wold, S., 78-5668
 Wold, W. S., 78-5840
 Wolf, M. H., 78-5627
 Wolfe, L., 78-5817
 Wolff, S., 78-5475
 Wolff, T., 78-5417
 Wong, C. Q., 78-5644
 Wood, A. W., 78-5548
 Woodruff, J. D., 78-5889
 Woodruff, M., 78-5520
 Worthington, B. S., 78-5525
 Wright, D. H., 78-5887
 Wu, R., 78-5795
 Wutke, K., 78-5898
 Wyllie, A. H., 78-5805
 Wynder, E., 78-5404
 Wynder, E. L., 78-5402, 78-5409
 Yager, J. D., 78-5477
 Yagi, H., 78-5548
 Yakovleva, L., 78-5817
 Yamaguchi, K., 78-5988
 Yamamura, Y., 78-5532
 Yamane, Y., 78-5669, 78-5674
 Yamazaki, M., 78-5857
 Yamazoe, Y., 78-5564
 Yanai, R., 78-5995
 Yang, H. S., 78-5612
 Yang, R. C., 78-5795
 Yang, S. S., 78-5773
 Yarita, T., 78-5640, 78-5642
 Yarmolinsky, M. B., 78-5515

aki, S., 78-5532
tscoff, R. W., 78-5667
ekawa, H., 78-5479
hida, M., 78-5766
hida, M. C., 78-5909
hida, S., 78-5672
shikura, H., 78-5766
shimoto, T., 78-5532


Young, H. A., 78-5535
Young, J. A., 78-5589
Young, J. C., 78-5620
Yuhas, J. M., 78-5950, 78-5996
Zbinden, G., 78-5677
Zech, L., 78-5906
Zeller, C., 78-5934
Zerban, H., 78-5688, 78-5689

Zhdanov, V. M., 78-5793, 78-5831

Zhorova, E. S., 78-5726

Zimmermann, A., 78-5894

Zweifel, J., 78-5934



Subject Index

- ormalities**
- Anesthetics**
 - Epidemiology, Review, 78-5423
- Lysergic Acid Diethylamide**
 - Teratogenicity, Review, 78-5422
- amide**
- Leukemia, Myeloblastic**
 - Cell Differentiation, 78-5983
- amide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-**
 - DNA Repair
 - Hepatocytes, Rat, 78-5477
- amide, *N*-Butyl-**
 - Nitrous Acid, Sodium Salt
 - Nitrosation, 78-5636
- amide, *N,N*-Dimethyl-**
 - Leukemia, Myeloblastic
 - Cell Differentiation, 78-5983
- amide, *N*-Fluoren-2-yl-**
 - Acetohydroxamic Acid, *N*-Fluoren-2-yl-
 - Carcinogenic Metabolite, 78-5478
 - 7,8-Benzoflavone
 - RNA, Binding, 78-5479
 - Bladder Neoplasms
 - Carcinoma, Transitional Cell, 78-5476
 - Hyperplasia, 78-5476
 - Precancerous Conditions, 78-5476
 - Cholanthrene, 3-Methyl-
 - Hydroxylation, 78-5478
 - RNA, Binding, 78-5479
 - Chromatids
 - Hamster Ovary Cells, 78-5475
 - Cytochromes
 - Metabolism, Review, 78-5407
 - Deoxyguanosine Adducts
 - Computer Model, 78-5502
 - Diethylamine, *N*-Nitroso-
 - Co-carcinogenic Effect, 78-5516
 - DNA Repair
 - Hepatocytes, Rat, 78-5477
 - Enzyme Activation
 - Carcinogenic Metabolite, Review, 78-5407
 - Mutagenic Activity, 78-5480
 - Mutagenic Activity, Review, 78-5407
 - Erythroleukemia
 - Bone Marrow Cells, 78-5483
 - Histological Study, Rat, 78-5483
 - Leukemia, Myelocytic
 - Bone Marrow Cells, 78-5483
 - Histological Study, Rat, 78-5483
 - Liver Cells, Intact
 - Mutagenic Activity, 78-5480
 - Liver Neoplasms
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5481
 - 78-5516
 - 5,6-Benzoflavone, 78-5481
 - Cholanthrene, 3-Methyl-, 78-5516
 - Co-carcinogenic Effect, 78-5516
 - Hyperplasia, 78-5481
 - 2,4-Oxazolidinedione, 3-(3,5-Dichlorophenyl)-5,5-dimethyl-, 78-5481
 - Polychlorobiphenyl Compounds, 78-5481
- Acetamide, *N*-Fluoren-2-yl- (cont'd)**
 - Microsomes, Liver
 - Chromatids, 78-5475
 - Hydroxylation, 78-5478
 - RNA, Binding, 78-5479
 - Nucleic Acids
 - Circadian Rhythm, 78-5482
 - Metabolism, Rat Liver, 78-5482
 - Oxidoreductases
 - Enzyme Activation, 78-5479
 - Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Acetamide, *N*-Fluoren-4-yl-**
 - Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Acetamide, *N,N*-Fluorene-2,7-diylbis-**
 - Leukemia, Myelocytic
 - Neoplasm Metastasis, 78-5484
- Acetamide, *N*-Methyl-**
 - Leukemia, Myeloblastic
 - Cell Differentiation, 78-5983
- Acetamide, *N*-Propyl-**
 - Methylamine, α -Acetoxybenzyl-*N*-nitroso-
 - Hydrolysis, Methanolysis, 78-5577
- Acetic Acid, ((Carboxymethylimino)bis(ethylenenitrilo))tetra-**
 - Citric Acid, Plutonium Salt
 - Body Burden, Dog, 78-5731
 - Drug Therapy, Cadmium/Zinc Chelates, 78-5731
 - Plutonium
 - Body Burden, Dog, 78-5731, 78-5733
 - Drug Therapy, Cadmium/Zinc Chelates, 78-5731
 - 78-5733
- Acetic Acid, (Ethylenedinitrilo)tetra-**
 - Cholanthrene, 3-Methyl-
 - Dihydrodiol Formation, 78-5530
- Acetic Acid, *o*-Hydroxyphenyl-**
 - Coumarin
 - Hepatocarcinogenicity, Review, 78-5433
- Acetic Acid, Lead Salt**
 - Piperidine, 1-Nitroso-
 - Acetoxy Derivatives, 78-5578
 - Pyrrolidine, 1-Nitroso-
 - Acetoxy Derivatives, 78-5578
- Acetic Acid, (2,4,5-Trichlorophenoxy)-**
 - Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
 - Toxic Contaminant, Review, 78-5421
- Acetohydroxamic Acid, *N*-Fluoren-2-yl-**
 - Acetamide, *N*-Fluoren-2-yl-
 - Carcinogenic Metabolite, 78-5478
 - Cholanthrene, 3-Methyl-
 - Hydroxylation, 78-5478
 - DNA Repair
 - Hepatocytes, Rat, 78-5477
 - Microsomes, Liver
 - RNA, Binding, 78-5479
- Acetone**
 - Nitrosamines
 - Permeability, Laboratory Gloves, 78-5596

Acetyltransferases

- Isonicotinic Acid Hydrazide
Metabolism, Review, 78-5407

Achlorhydria

- Stomach Neoplasms
Nitrous Acid, 78-5633

Acid Phosphatase

- Prostatic Neoplasms
Virus, SV40, 78-5864

Acridine Mustard

- Neurospora crassa*
Mutagenic Activity, 78-5675

Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-

- Breast Neoplasms
Epidemiology, Japan, 78-5946
- Thymidine Kinase
Mutagenic Activity, 78-5666

Actin

- Sarcoma, Mast Cell
Cells, Cultured, 78-5863
- Histocompatibility Antigens, 78-5863

Adenine, 3-Methyl-

- Glycoside Hydrolases
DNA, Alkylation, 78-5663
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
Glycoside Hydrolases, 78-5663
- Methanesulfonic Acid, Methyl Ester
Glycoside Hydrolases, 78-5663
- Sulfuric Acid, Dimethyl Ester
Glycoside Hydrolases, 78-5663
- Urea, Methyl Nitroso-
Glycoside Hydrolases, 78-5663

Adenocarcinoma

- Breast Neoplasms
Cells, Cultured, 78-5950
- Insulin, 78-5950
- Cervix Neoplasms
Epidemiology, Israel, 78-5958
- Ethnic Groups, 78-5958
- Colonic Neoplasms
Hydrazine, 1,2-Dimethyl-, 78-5496, 78-5497
78-5498
- Virus, Cytomegalo, 78-5885
- Ear Neoplasms
Diethylamine, 2,2-Dichloro-*N*-nitroso-, 78-5572
- Gastrointestinal Neoplasms
Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5867
- Intestinal Neoplasms
Bracken Fern, 78-5464, 78-5465
- Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499
- Virus, Cytomegalo, 78-5885
- Kidney Neoplasms
Diethylamine, *N*-Nitroso-, 78-5573
- Ochratoxin A, 78-5468
- Liver Neoplasms
Ochratoxin A, 78-5468
- Lung Neoplasms
Animal Model, Hamster, 78-5878
- Azocine, Octahydro-1-nitroso-, 78-5585
- Benzo(a)pyrene, 78-5878
- Californium, 78-5726
- Classification, 78-5877
- Hydrazine, 1,2-Diformyl-, 78-5493
- Nickel, 78-5931

Adenocarcinoma (cont'd)

- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
- Ultrastructural/Histochemical Study, 78-5877
- Mammary Neoplasms, Experimental
Benz(a)anthracene, 7,12-Dimethyl-, 78-5535
78-5537, 78-5539
- Prolactin, 78-5753
- RNA, Viral, 78-5535
- Testosterone, Propionate, 78-5988
- Transplantation Immunology, 78-5539
- Virus, Murine Mammary Tumor, 78-5753
- Nose Neoplasms
Diethylamine, *N*-Nitroso-, 78-5575
- Ovarian Neoplasms
Age Factors, 78-5952
- Precancerous Conditions, 78-5952
- Pancreatic Neoplasms
Benz(a)anthracene, 7,12-Dimethyl-, 78-5534
- Stomach Neoplasms
Epidemiology, 78-5942
- Gastritis, 78-5880
- Thyroid Neoplasms
Radiation, Ionizing, 78-5972
- Tracheal Neoplasms
Cholanthrene, 3-Methyl-, 78-5521
- Urea, Methyl Nitroso-, 78-5640, 78-5642

Adenocarcinoma, Papillary

- Nasopharyngeal Neoplasms
Virus, Epstein-Barr, 78-5830
- Thyroid Neoplasms
Radiation, Ionizing, 78-5972

Adenofibroma

- Mammary Neoplasms, Experimental
Benz(a)anthracene, 7,12-Dimethyl-, 78-5535
78-5537

Adenoma

- Brain Neoplasms
Urea, Ethyl Nitroso-, 78-5650
- Colonic Neoplasms
Carrageen, 78-5644
- Hydrazine, 1,2-Dimethyl-, 78-5497, 78-5498
- Intestinal Neoplasms
Bracken Fern, 78-5464, 78-5465
- Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499
- Kidney Neoplasms
Aflatoxin B1, 78-5468
- Diethylamine, *N*-Nitroso-, 78-5573, 78-5574
- Morpholine, *N*-Nitroso-, 78-5688, 78-5689
- Ochratoxin A, 78-5468
- WY-14,643, 78-5574
- Leukemia, Myelocytic
Thyroid Neoplasms, 78-5906
- Liver Neoplasms
Aflatoxin B1, 78-5468
- Ochratoxin A, 78-5468
- Lung Neoplasms
Azocine, Octahydro-1-nitroso-, 78-5585
- Benzo(a)pyrene, 78-5570
- Diethylamine, *N*-Nitroso-, 78-5665
- Dimethylamine, *N*-Nitroso-, 78-5570
- Hydrazine, 1,2-Diformyl-, 78-5493
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
- Quinoline, 8-Nitro-, 78-5672
- Radiation, Ionizing, 78-5665
- Parathyroid Neoplasms
Case Report, 78-5712

oma (cont'd)

- Radiation, Ionizing, 78-5712, 78-5714
- Thyroid Neoplasms
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580

oma, Chromophobe

- Pituitary Neoplasms
 - Epidemiology, Nigeria, 78-5456

oma, Eosinophilic

- Kidney Neoplasms
 - Morpholine, *N*-Nitroso-, 78-5688

osine Cyclic 3',5' Monophosphate

- Cervix Neoplasms
 - Cholanthrene, 3-Methyl-, 78-5868
- Pancreatic Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5534
 - Tissue Levels, 78-5534

osine Triphosphatase

- Aflatoxin B2
 - Mitochondria, Liver, 78-5469
- Aflatoxin G2
 - Mitochondria, Liver, 78-5469
- Benz(a)anthracene, 7,12-Dimethyl-
 - Calcium, 78-5503
 - Muscle, Connective Tissue, Rat, 78-5503
- Carbamic Acid, Ethyl Ester
 - Calcium, 78-5503
 - Muscle, Connective Tissue, Rat, 78-5503
- Dibenz(a,h)anthracene
 - Calcium, 78-5503
 - Muscle, Connective Tissue, Rat, 78-5503
- Dimethylamine, *N*-Nitroso-
 - Calcium, 78-5503

osine Triphosphate

- Benzo(a)pyrene
 - Ames Test, 78-5559
- Leukemia, Lymphocytic
 - Magnesium, 78-5981
- Leukemia, Myelocytic
 - Magnesium, 78-5981
- Polychlorobiphenyl Compounds
 - Aryl Hydrocarbon Hydroxylases, 78-5559

iamycin

- Chromosomes
 - DNA Replication, 78-5673
- HeLa Cells
 - DNA Replication, 78-5673

toxin B1

- Aflatoxin B1, 2,3-Dihydro-2-(*S*-glutathionyl)-3-hydroxy-
 - Detoxification Product, 78-5472
 - Liver, Bile, Rat, 78-5472
- Aflatoxin B1 2,3-Oxide
 - Carcinogenic Metabolite, 78-5510
- Aflatoxin M1
 - Carcinogenic Metabolite, 78-5510
- Aryl Hydrocarbon Hydroxylases
 - Strain Difference, Mouse, 78-5510
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Metabolism, Mouse, 78-5510
- Carcinogen, Environmental
 - Metabolism, Review, 78-5406
- Cell Transformation, Neoplastic
 - Cells, Cultured, 78-5471
 - Liver, Rat, 78-5471
- Cells, Cultured

Aflatoxin B1 (cont'd)

- Prostaglandins, 78-5509
- Tumorigenicity, 78-5471
- Cholanthrene, 3-Methyl-
 - Aryl Hydrocarbon Hydroxylases, 78-5510
 - Carcinogenic Metabolite, 78-5510
- Choline
 - Cytochrome P-450, 78-5473
 - Oxidoreductases, 78-5473
- Colon
 - DNA, Binding, 78-5508
 - Metabolism, Rat, 78-5508
- Cytochrome P-450
 - Carcinogenic Metabolite, 78-5510
- Dietary Fats
 - Glutathione, 78-5473
 - Hepatocarcinogenicity, Rat, 78-5473
 - Nucleic Acids, Binding, 78-5473
- Dietary Proteins
 - Hepatocarcinogenicity, 78-5470
- DNA, Binding
 - Magnetic Resonance Spectra, 78-5474
- Folic Acid
 - Cytochrome P-450, 78-5473
 - Oxidoreductases, 78-5473
- Glutathione
 - Metabolism, Rat, 78-5472
- Hepatitis
 - Epidemiology, India, 78-5964
 - Histological Study, 78-5964
- Hepatoma
 - Cells, Cultured, 78-5471
 - Dietary Proteins, 78-5470
 - Vitamin B12, 78-5470
- Karyotyping
 - Cells, Cultured, 78-5471
- Kidney Neoplasms
 - Adenoma, 78-5468
 - Ochratoxin A, 78-5468
- Liver
 - DNA, Binding, 78-5474
- Liver Neoplasms
 - Adenoma, 78-5468
 - Ochratoxin A, 78-5468
- Methionine
 - Cytochrome P-450, 78-5473
 - Oxidoreductases, 78-5473
- Microsomes, Liver
 - Metabolism, Mouse, 78-5510
- Ochratoxin A
 - Co-carcinogenic Effect, 78-5468
- Oxidoreductases
 - Microsomes, Liver, 78-5473
- Vitamin B12
 - Hepatocarcinogenicity, 78-5470

Aflatoxin B2

- DNA, Binding
 - Magnetic Resonance Spectra, 78-5474
- Liver
 - DNA, Binding, 78-5474
- Mitochondria, Liver
 - Adenosine Triphosphatase, 78-5469
 - Cell Membrane Permeability, 78-5469
 - Cytochrome Oxidase, 78-5469

Aflatoxin B1, 2,3-Dihydro-2-(*S*-glutathionyl)-3-hydroxy-**Aflatoxin B1**

**Aflatoxin B1, 2,3-Dihydro-2-(S-glutathionyl)-3-hydroxy-
(cont'd)**

Detoxification Product, 78-5472
Liver, Bile, Rat, 78-5472

Aflatoxin B1 2,3-Oxide
Detoxification Product, 78-5472

Aflatoxin B1 2,3-Oxide

Aflatoxin B1
Carcinogenic Metabolite, 78-5510
Aflatoxin B1, 2,3-Dihydro-2-(S-glutathionyl)-3-hydroxy-
Detoxification Product, 78-5472
Carcinogenic Metabolite
Liver, Bile, Rat, 78-5472
Cytochrome P-450
Metabolism, Mouse, 78-5510

Aflatoxin G1

Dietary Proteins
Hepatocarcinogenicity, 78-5470
Hepatoma
Dietary Proteins, 78-5470
Vitamin B12, 78-5470
Vitamin B12
Hepatocarcinogenicity, 78-5470

Aflatoxin G2

Mitochondria, Liver
Adenosine Triphosphatase, 78-5469
Cell Membrane Permeability, 78-5469
Cytochrome Oxidase, 78-5469

Aflatoxin M1

Aflatoxin B1
Carcinogenic Metabolite, 78-5510
Cytochrome P-450
Metabolism, Mouse, 78-5510

Agammaglobulinemia

Leukemia, Lymphocytic
Neoplasms, Multiple Primary, 78-5920

Aging

Breast Neoplasms
Endocrine Glands, Review, 78-5411
Carcinogen, Chemical
DNA Repair, Review, 78-5411
Enzyme Activation, Review, 78-5411
Fibroblasts
Cells, Cultured, 78-5977
Prostatic Neoplasms
Endocrine Glands, Review, 78-5411

Air Pollution

Carcinoembryonic Antigen
Serum Levels, 78-5940
Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
Mutagenic Activity, Review, 78-5421
Gastrointestinal Neoplasms
Epidemiology, Poland, 78-5941
Lung Diseases, Obstructive
Epidemiology, 78-5934
Lung Neoplasms
Epidemiology, 78-5934
Epidemiology, Review, 78-5409
Nitrosamines
Epidemiology, Review, 78-5409
Respiratory Tract Neoplasms
Epidemiology, Houston, 78-5932
Occupational Hazard, 78-5932
Smoking

Air Pollution (cont'd)

Co-carcinogenic Activity, 78-5932

Alanine, Phenyl-

Neoplasms, Experimental
Cholanthrene, 3-Methyl-, 78-5525

Albumins

Sarcoma
Virus, Rous Sarcoma, 78-5746

Alginate Acid, Sodium Salt

Strontium Radioisotopes
Body Burden, Rat, 78-5727
Metabolism, Rat, 78-5727

Alkaline Phosphatase

Fibroblasts
Cell Transformation, Neoplastic, 78-5798
Virus, SV40
Cell Membrane, 78-5798
Cell Transformation, Neoplastic, 78-5798
Enzymatic Activity, 78-5798

Alpha Particles

Plutonium Oxide
Carcinogenic Potential, Review, 78-5444

Aluminum Chloride

Quinoline, 4-(Hydroxyamino)-, 1-Oxide
Nucleic Acids, Binding, 78-5674

α -Amanitine

Virus, Adeno 2
RNA, Viral, 78-5839

Ames Test

2-Anthracenamide
Aroclor 1254, 78-5517
Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
5,6-Benzoflavone, 78-5517
Aroclor 1254
S-9 Fraction, 78-5517
Azo Compounds
Evaluation, Review, 78-5427
Benz(a)anthracene, 7,12-Dimethyl-
Aroclor 1254, 78-5517
Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
5,6-Benzoflavone, 78-5517
Benzene, (Epoxyethyl)-
Mutagenic Activity, Review, 78-5436
Benzo(a)pyrene
Adenosine Triphosphate, 78-5559
Aroclor 1254, 78-5517
Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
5,6-Benzoflavone, 78-5517
Glutathione, 78-5559
Microsomes, Liver, 78-5563
Nitric Acid, 78-5568
Nitro Derivatives, 78-5568
Nitrogen Oxide, 78-5568
Uridine Diphosphate Sugars, 78-5559
Benzo(a)pyrene, 12b,12c-Dihydro-12b,12c-dimethyl-
Microsomes, Liver, 78-5563
Benzo(b,def)chrysene, 13b,13c-Dihydro-13b,13c-dimethyl-
Microsomes, Liver, 78-5563
Benzo(e)pyrene, 12b,12c-Dihydro-12b,12c-dimethyl-
Microsomes, Liver, 78-5563
5,6-Benzoflavone
S-9 Fraction, 78-5517
1,3-Butadiene, 1-Chloro-

Test (cont'd)

- Mutagenic Activity, 78-5492
- 1,3-Butadiene, 2-Chloro-
 - Mutagenic Activity, 78-5492
- 1-Butene, 3,4-Dichloro-
 - Carcinogenic Metabolite, 78-5492
 - Mutagenic Activity, 78-5492
- 2-Butene, 1,4-Dichloro-
 - Mutagenic Activity, 78-5492
- Carcinogen, Chemical
 - Evaluation, Review, 78-5427
- Colchicine
 - Evaluation, Review, 78-5427
- Cyclophosphamide
 - Aroclor 1254, 78-5517
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
 - 5,6-Benzoflavone, 78-5517
- 1*H*-1,4-Diazepine, Hexahydro-1,4-dinitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Ethidium Bromide
 - Aroclor 1254, 78-5517
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
 - 5,6-Benzoflavone, 78-5517
- Ethylene, Bromo-
 - Mutagenic Activity, 78-5492
- Ethylene, 1,1-Dichloro-
 - Mutagenic Activity, 78-5492
- Ethylene Oxide, Chloro-
 - Mutagenic Activity, 78-5492
- Fluoren-2-amine
 - Aroclor 1254, 78-5517
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
 - 5,6-Benzoflavone, 78-5517
- Methanesulfonic Acid, Butyl Ester
 - Mutagenic Activity, 78-5659
- Methanesulfonic Acid, Ethyl Ester
 - Mutagenic Activity, 78-5659
- Methanesulfonic Acid, Methyl Ester
 - Mutagenic Activity, 78-5659
- Methanesulfonic Acid, Propyl Ester
 - Mutagenic Activity, 78-5659
- Morpholine, *N*-Nitroso-
 - Aroclor 1254, 78-5517
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5517
 - 5,6-Benzoflavone, 78-5517
- Mutagens
 - Feces, 78-5629
 - In Vitro Test Systems, Review, 78-5417
- Perylene
 - Nitric Acid, 78-5568
 - Nitro Derivatives, 78-5568
 - Nitrogen Oxide, 78-5568
- Pesticides
 - Evaluation, Review, 78-5427
- Piperazine, 4-Benzoyl-3,5-dimethyl-1-nitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Piperazine, 2,5-Dimethyl-1,4-dinitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Piperazine, 2,6-Dimethyl-1,4-dinitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Piperazine, 1,4-Dinitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589

Ames Test (cont'd)

- Piperazine, 1,4-Dinitroso-2-methyl-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Piperazine, 1-Methyl-4-nitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Piperazine, 1-Nitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Piperazine, 2,3,5,6-Tetramethyl-1,4-dinitroso-
 - Aroclor 1254, 78-5589
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5589
- Polycyclic Hydrocarbons
 - Synthetic Fuels, 78-5514
- Polyethylene
 - Mutagenic Activity, 78-5487
- 5*H*-Pyrido(4,3-*b*)indole, 3-Amino-1-methyl-
 - Mutagens, 78-5426
- Radiation, Ionizing
 - Evaluation, Review, 78-5427

Amino Acids

- Cholanthrene, 3-Methyl-
 - Transplantation Immunology, 78-5525
- Neoplasms, Experimental
 - Transplantation Immunology, 78-5525
- Virus, Avian Myeloblastosis
 - Reverse Transcriptase, 78-5743
- Virus, Avian Sarcoma
 - Viral Proteins, 78-5739
- Virus, Polyoma
 - Nucleotide Sequence, 78-5792

Aminophylline

see Theophylline

Aminotransferases

- Dimethylamine, *N*-Nitroso-
 - Selenium, 78-5606
 - Vitamin E, 78-5606

Amyloidosis

- Cholanthrene, 3-Methyl-
 - BCG*, 78-5532
 - Thymectomy, 78-5532
- Quinoline, 4-Nitro-, 1-Oxide
 - BCG*, 78-5532
 - Thymectomy, 78-5532

Anemia, Aplastic

- Mesenchymoma
 - Transplantation, Heterologous, 78-5904

Anemia, Hemolytic

- Leukemia, Lymphocytic
 - Epidemiology, Israel, 78-5920

Anemia, Sideroblastic

- Leukemia
 - Case Report, 78-5901
 - Megakaryocytes, 78-5901

Anesthetics

- Abnormalities
 - Epidemiology, Review, 78-5423
- Neoplasms
 - Epidemiology, Review, 78-5423
- Occupational Hazard
 - Epidemiology, Review, 78-5423

Angiosarcoma

- Ethylene, Chloro-
 - Chromosome Aberrations, 78-5491
 - Epidemiology, 78-5486, 78-5967
 - Occupational Hazard, 78-5486, 78-5491
 - Occupational Hazard, Review, 78-5435
 - Precancerous Conditions, Review, 78-5435
- Ethylene, Chloro- Polymer
 - Epidemiology, 78-5967
- Liver Neoplasms
 - Ethylene, Chloro-, 78-5485, 78-5486, 78-5967
 - Ethylene, Chloro- Polymer, 78-5967

Aniline, 4-Fluoro-

- Saccharomyces cerevisiae*
 - Mutagenic Activity, 78-5654

Aniline, N-Methyl-4-nitroso-

- Cysteine, N-Nitroso-, Hydrochloride
 - Catalysis, 78-5630

Aniline, N-Methyl-N-nitroso-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Mutagenic Activity, Hamster V79 Cells, 78-5593

Aniline, p-(Phenylazo)-

- Soy Beans
 - Somatic Mosaicism, 78-5664

2-Anthracenamide

- Aroclor 1254
 - Ames Test, 78-5517
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Ames Test, 78-5517
- 5,6-Benzoflavone
 - Ames Test, 78-5517

Anthracosilicosis

- Lung Neoplasms
 - Carcinoma, Epidermoid, 78-5971
 - Epidemiology, 78-5971
 - Occupational Hazard, 78-5971

Anti-Antibodies

- Intestinal Neoplasms
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499
 - Neoplasm Metastasis, 78-5499

Antibodies, Viral

- Burkitt's Lymphoma
 - Immune Response, Infant, 78-5836
- Cervix Neoplasms
 - Virus, Herpes Simplex 2, 78-5822
- Infectious Mononucleosis
 - Virus, Epstein-Barr, 78-5829
- Leukemia
 - Virus, Epstein-Barr, 78-5829
- Lymphoma
 - Virus, Epstein-Barr, 78-5829
- Lymphosarcoma
 - Virus, Feline Leukemia, 78-5785
- Nasopharyngeal Neoplasms
 - Virus, Epstein-Barr, 78-5830
- Virus, Epstein-Barr
 - IgM, 78-5829
 - Immune Response, Infant, 78-5835, 78-5836
 - Seroepidemiology, Ghana, 78-5835
- Virus, Feline Leukemia
 - Epidemiology, 78-5785
 - Immune Response, 78-5785
 - Virus Subgroup, 78-5785

Antibodies, Viral (cont'd)

- Virus, Friend Murine Leukemia
 - Titration Method, 78-5765
- Virus, Papilloma
 - Animals, Domestic, 78-5791
 - Immunodiffusion, 78-5791

Antibody Formation

- Brain Neoplasms
 - Urea, Methyl Nitroso-, 78-5859
- Prostatic Neoplasms
 - Carcinoembryonic Antigen, 78-5864

Antibody Specificity

- Astrocytoma
 - Urea, Methyl Nitroso-, 78-5859
- Brain Neoplasms
 - Urea, Methyl Nitroso-, 78-5859
- Fibrosarcoma
 - Antigens, Viral, 78-5866
 - Carcinoembryonic Antigen, 78-5866
- Glioma
 - Urea, Methyl Nitroso-, 78-5859
- Virus, Epstein-Barr
 - Infant, 78-5836
- Virus, RD-114
 - Reverse Transcriptase, 78-5850

Antigen-Antibody Complex

- Erythroleukemia
 - Virus, Rauscher Murine Leukemia, 78-5782
- Virus, Papilloma
 - Ultrastructural Study, Dog, 78-5791
- Virus, Rauscher Murine Leukemia
 - Cell Transformation, Neoplastic, 78-5782
 - Immune Response, 78-5782
 - Immune Serums, 78-5782
 - Neoplasm Regression, Spontaneous, 78-5782

Antigen-Antibody Reactions

- Antigens, Viral
 - Human/Simian Sera, 78-5825
- Leukemia, Myeloblastic
 - Cells, Cultured, 78-5865
 - Primate Antisera, 78-5865
- Rhabdomyosarcoma
 - Primate Antisera, 78-5865
 - Virus, Baboon, 78-5865
 - Virus, Gibbon Ape Lymphoma, 78-5865
 - Virus, Simian Sarcoma, 78-5865
- Sarcoma
 - Fetal Globulins, 78-5531
- Virus, Avian Myeloblastosis
 - Antigenic Determinants, 78-5741
 - Antigens, Viral, 78-5741
- Virus, Marek's Disease Herpes
 - Histocompatibility Antigens, 78-5748
- Virus, Murine Leukemia
 - Viral Proteins, 78-5759

Antigenic Determinants

- Fibrosarcoma
 - Cholanthrene, 3-Methyl-, 78-5866
- Virus, Avian Leukosis-Sarcoma
 - Chick Embryo, 78-5736
 - Viral Proteins, 78-5736
- Virus, Avian Myeloblastosis
 - Antigen-Antibody Reactions, 78-5741
 - Viral Proteins, 78-5742
 - Virus, Rous Sarcoma, 78-5741

Antigen Determinants (cont'd)

- Virus, Avian Sarcoma
 - Cell Transformation, Neoplastic, 78-5739
- Virus, Cytomegalo
 - Virus, Herpes, 78-5823
 - Virus, Rhino, 78-5823
- Virus, Herpes Papio
 - Virus, Epstein-Barr, 78-5817
- Virus, Polyoma, JC
 - Radioimmunoassay, 78-5793
- Virus, Baboon, 78-5793
- Virus, Feline Leukemia, 78-5793
- Virus, Mason-Pfizer Monkey, 78-5793
- Virus, Murine Leukemia, 78-5793
- Virus, Rauscher Murine Leukemia
 - Glycoproteins, 78-5781
 - Viral Proteins, 78-5778
 - Virus, Murine Leukemia, 78-5781
- Virus, RD-114
 - Reverse Transcriptase, 78-5850
- Virus, Baboon, 78-5850
- Virus, Rhino
 - Virus, Herpes, 78-5823
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-5744
- Virus, SV40
 - Histocompatibility Antigens, 78-5812

Antigens

- Cervix Neoplasms
 - Cholanthrene, 3-Methyl-, 78-5868
 - Isolation and Characterization, 78-5868
- Virus, Epstein-Barr
 - Cell Membrane, 78-5833
 - Cell Transformation, Neoplastic, 78-5833
 - Isolation and Characterization, 78-5833
- Virus, Rauscher Murine Leukemia
 - Cell Transformation, Neoplastic, 78-5782
 - Sialic Acid, 78-5782

Antigens, Neoplasm

- Carcinoma, Epidermoid
 - Immunotherapy, 78-5854

Antigens, Viral

- Antigen-Antibody Reactions
 - Human/Simian Sera, 78-5825
- Avian Leukosis
 - Virus, Marek's Disease Herpes, 78-5749
- Burkitt's Lymphoma
 - Virus, Epstein-Barr, 78-5824, 78-5827
- Fibrosarcoma
 - Antibody Specificity, 78-5866
 - Virus, Murine Leukemia, 78-5866
- Head and Neck Neoplasms
 - Virus, Epstein-Barr, 78-5830
- Infectious Mononucleosis
 - Virus, Epstein-Barr, 78-5824
- Leukemia, Lymphoblastic
 - Virus, Epstein-Barr, 78-5827
- Lymphoma
 - Virus, Marek's Disease Herpes, 78-5749
 - Virus, Moloney Murine Leukemia, 78-5774
- Sarcoma
 - Carcinogen, Chemical, 78-5438
- Virus, Avian Myeloblastosis
 - Antigen-Antibody Reactions, 78-5741
 - Casein, 78-5741
 - Concanavalin A, 78-5741

Antigens, Viral (cont'd)

- Ovalbumin, 78-5741
- Papain, 78-5741
- Peptide Hydrolases, 78-5741
- Serum Albumin, Bovine, 78-5741
- Virus, Avian Sarcoma
 - Cell Transformation, Neoplastic, 78-5740
- Virus, Epstein-Barr
 - Cortisone, 78-5824
 - Cycloheximide, 78-5828
 - DNA Replication, 78-5828
 - Lymphocyte Transformation, 78-5828
- Virus, Feline Leukemia
 - Cell Membrane, 78-5786
 - Cell Transformation, Neoplastic, 78-5786
- Virus, Friend Murine Leukemia
 - Lymphoid Tissue, 78-5763
- Virus, Gross Murine Leukemia
 - Cells, Cultured, 78-5769
 - Immune Serums, 78-5769
 - Radiation, Ionizing, 78-5769
- Virus, Marek's Disease Herpes
 - Blood Cells, 78-5749
- Virus, Moloney Murine Leukemia
 - Isolation and Characterization, 78-5774
- Virus, Murine Mammary Tumor
 - Horizontal Transmission, 78-5755
- Virus, Rous Sarcoma
 - Cell Membrane, 78-5744
 - Temperature Sensitive Mutants, 78-5744
- Virus, SV40
 - Cell-Free Synthesis, 78-5800
 - Cell Transformation, Neoplastic, 78-5801
 - DNA-RNA Hybridization, 78-5800
 - DNA, Viral, 78-5801
 - Immunization, 78-5812
 - Immunoprecipitation, 78-5801
 - Isolation and Characterization, 78-5800
 - Prostate, Mouse, 78-5864
 - RNA, Messenger, 78-5800
 - Tryptic Peptide Analysis, 78-5800, 78-5801

Antilymphocyte Serum

- Sarcoma
 - T-Lymphocytes, 78-5730

Antineoplastic Agents

- Escherichia coli*
 - Coliphages, 78-5515

Antipain

- Bladder Neoplasms
 - Papilloma, 78-5583
- Hepatoma
 - Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer, 78-5583

Antipyrine

- Stress
 - Enzyme Activation, 78-5560
 - Plasma Levels, 78-5560

Antipyrine, 4-(Dimethylamino)-

- Nitrous Acid, Sodium Salt
- Nitrosation, 78-5636

Aprotinin

- see Kallikrein-Trypsin Inhibitor

Aroclor 1254

- Ames Test

Aroclor 1254 (cont'd)

- S-9 Fraction, 78-5517
- 2-Anthracenamide
 - Ames Test, 78-5517
- Benz(a)anthracene, 7,12-Dimethyl-
 - Ames Test, 78-5517
- Benzene, 4-Allyl-1,2-(methylenedioxy)-
 - Carcinogenic Metabolite, 78-5519
- Benzo(a)pyrene
 - Ames Test, 78-5517
- Carcinogen, Chemical
 - Enzyme Activation, 78-5480
- Cyclophosphamide
 - Ames Test, 78-5517
- 1*H*-1,4-Diazepine, Hexahydro-1,4-dinitroso-
 - Ames Test, 78-5589
- Dibenz(a,h)anthracene
 - Mutagenic Activity, 78-5548
- Ethidium Bromide
 - Ames Test, 78-5517
- Fluoren-2-amine
 - Ames Test, 78-5517
- Morpholine, *N*-Nitroso-
 - Ames Test, 78-5517
- Piperazine, 4-Benzoyl-3,5-dimethyl-1-nitroso-
 - Ames Test, 78-5589
- Piperazine, 2,5-Dimethyl-1,4-dinitroso-
 - Ames Test, 78-5589
- Piperazine, 2,6-Dimethyl-1,4-dinitroso-
 - Ames Test, 78-5589
- Piperazine, 1,4-Dinitroso-
 - Ames Test, 78-5589
- Piperazine, 1,4-Dinitroso-2-methyl-
 - Ames Test, 78-5589
- Piperazine, 1-Methyl-4-nitroso-
 - Ames Test, 78-5589
- Piperazine, 1-Nitroso-
 - Ames Test, 78-5589
- Piperazine, 2,3,5,6-Tetramethyl-1,4-dinitroso-
 - Ames Test, 78-5589

Arsenic

- Lymphosarcoma
 - Occupational Hazard, 78-5922

Arthritis, Rheumatoid

- Sarcoma, Kaposi's
 - Prednisone, 78-5855

Aryl Hydrocarbon Hydroxylases

- Aflatoxin B₁
 - Cholanthrene, 3-Methyl-, 78-5510
 - Strain Difference, Mouse, 78-5510
- Benzo(a)pyrene
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5553
 - Cholanthrene, 3-Methyl-, 78-5553
 - Dieldrin, 78-5553
 - DNA, Binding, 78-5553
- Cortisol
 - Stress, 78-5560
- Polychlorobiphenyl Compounds
 - Adenosine Triphosphate, 78-5559
 - S-9 Liver Fraction, 78-5559

Asbestos

- Benzo(a)pyrene
 - Adsorption, 78-5556
- Gastrointestinal Neoplasms
 - Epidemiology, Review, 78-5445

Asbestos (cont'd)

- Occupational Hazard, Review, 78-5445
- Lung Neoplasms
 - Air Pollution, Review, 78-5409
 - Epidemiology, 78-5928
- Mesothelioma
 - Case Report, 78-5875
- Pleural Neoplasms
 - Mesothelioma, 78-5875
- Smoking
 - Co-carcinogenic Effect, 78-5928

Ascorbic Acid

- Cholanthrene, 3-Methyl-
 - Dihydrodiol Formation, 78-5530
- Dimethylamine, *N*-Nitroso-
 - Minocycline, 78-5610
 - Vibramycin, 78-5610
- Mutagens
 - Feces, 78-5629
- Nitrosamines
 - Metabolism, Rat, 78-5594
- Nitroso Compounds
 - Mutagens, 78-5629

Ascorbic Acid, Monosodium Salt

- Urea, Methyl-
 - Nitrosation, 78-5636

Aspergillus nidulans

- 4-Cyclohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-
 - Saccharomyces cerevisiae*, 78-5653
- Methanesulfonic Acid, Methyl Ester
 - Saccharomyces cerevisiae*, 78-5653
- Pesticides
 - Saccharomyces cerevisiae*, 78-5653

Astrocytoma

- Brain Neoplasms
 - Urea, Ethyl Nitroso-, 78-5649
 - Urea, Methyl Nitroso-, 78-5859
- Child
 - Epidemiology, Denmark, 78-5925
- Urea, Methyl Nitroso-
 - Antibody-Dependent Cytotoxicity, 78-5859
 - Antibody Specificity, 78-5859
 - Urea, Ethyl Nitroso-, 78-5638

Ataxia Telangiectasia

- Lymphocytes
 - DNA Repair, 78-5697, 78-5860
 - Plant Agglutinins, 78-5860
- Radiation, Ionizing
 - Chromosome Aberrations, 78-5697
 - DNA Repair, 78-5697, 78-5860

Australia Antigen

- Hepatoma
 - DNA, Viral, 78-5837
 - Epidemiology, China, 78-5965
 - Epidemiology, Ivory Coast, 78-5962
 - Liver Cirrhosis, 78-5965
 - Virus, Hepatitis, 78-5837

Avian Leukosis

- Virus, Marek's Disease Herpes
 - Antigens, Viral, 78-5749

Azathioprine

- Leukemia, Myelocytic

hioprine (cont'd)

Immunosuppression, 78-5895

Compounds**Ames Test**

Evaluation, Review, 78-5427

ine, Octahydro-1-nitroso-**Lung Neoplasms**

Adenocarcinoma, 78-5585

Adenoma, 78-5585

Carcinoma, Epidermoid, 78-5585

Neoplasms, Multiple Primary, 78-5585

Ultrastructural Study, Hamster, 78-5585

Nose Neoplasms

Carcinogenic Activity, Hamster, 78-5585

illus brevis**1-Hexanamine, N-Hexyl-N-nitroso-**

Nitrosation, 78-5612

illus thuringiensis**Endotoxins**

Isolation and Characterization, 78-5467

Mitosis, 78-5467

Exotoxin

Isolation and Characterization, 78-5467

Mitosis

Exotoxin, 78-5467

teria**Carcinogen, Chemical**

Mutagenic Activity, Review, 78-5418

Carcinogen, Environmental

Mutagenic Activity, 78-5426

Nitric Acid, Sodium Salt

Mouth, Monkey, 78-5631

teriophages**Carcinogen, Chemical**

Derepression, Review, 78-5448

Mutagenic Activity, Review, 78-5448

bituric Acid, 5-Ethyl-5-phenyl-**Aflatoxin B1**

Metabolism, Mouse, 78-5510

Aniline, N-Methyl-N-nitroso-

Mutagenic Activity, Hamster V79 Cells, 78-5593

2-Anthracenamide

Ames Test, 78-5517

Benz(a)anthracene, 7,12-Dimethyl-

Ames Test, 78-5517

Benzene, 4-Allyl-1,2-(methylenedioxy)-

Carcinogenic Metabolite, 78-5519

Benzo(a)pyrene

Ames Test, 78-5517

Aryl Hydrocarbon Hydroxylases, 78-5553

DNA, Binding, 78-5553

Benzo(a)pyrene 4,5-Oxide

NADH, NADPH Oxidoreductases, 78-5564

Cyclophosphamide

Ames Test, 78-5517

1H-1,4-Diazepine, Hexahydro-1,4-dinitroso-

Ames Test, 78-5589

Diethylamine, N-Nitroso-

Mutagenic Activity, Hamster V79 Cells, 78-5593

Dimethylamine, N-Nitroso-

Mutagenic Activity, Hamster V79 Cells, 78-5593

Nucleic Acids, Binding, 78-5603

Dipropylamine, N-Nitroso-

Mutagenic Activity, Hamster V79 Cells, 78-5593

Barbituric Acid, 5-Ethyl-5-phenyl- (cont'd)**Ethidium Bromide**

Ames Test, 78-5517

Ethylene, Chloro-

Cytochrome P-450, 78-5490

Mutagenic Activity, Hamster V79 Cells, 78-5593

Fluoren-2-amine

Ames Test, 78-5517

Liver Neoplasms

Acetamide, N-Fluoren-2-yl-, 78-5481, 78-5516

Morpholine, N-Nitroso-

Ames Test, 78-5517

Mutagenic Activity, Hamster V79 Cells, 78-5593

Piperazine, 4-Benzoyl-3,5-dimethyl-1-nitroso-

Ames Test, 78-5589

Piperazine, 2,5-Dimethyl-1,4-dinitroso-

Ames Test, 78-5589

Piperazine, 2,6-Dimethyl-1,4-dinitroso-

Ames Test, 78-5589

Piperazine, 1,4-Dinitroso-

Ames Test, 78-5589

Piperazine, 1,4-Dinitroso-2-methyl-

Ames Test, 78-5589

Piperazine, 1-Methyl-4-nitroso-

Ames Test, 78-5589

Mutagenic Activity, Hamster V79 Cells, 78-5593

Piperazine, 1-Nitroso-

Ames Test, 78-5589

Piperazine, 2,3,5,6-Tetramethyl-1,4-dinitroso-

Ames Test, 78-5589

Pyrrolidine, 1-Nitroso-

Mutagenic Activity, Hamster V79 Cells, 78-5593

BCG**Amyloidosis**

Cholanthrene, 3-Methyl-, 78-5532

Quinoline, 4-Nitro-, 1-Oxide, 78-5532

Cell Transformation, Neoplastic

Macrophages, 78-5997

Lung Neoplasms

Cell-Wall Skeleton, 78-5532

Cholanthrene, 3-Methyl-, 78-5532

Quinoline, 4-Nitro-, 1-Oxide, 78-5532

Sarcoma, Osteogenic

Immune Response, Mouse, 78-5728

Strontium Radioisotopes, 78-5728

Bence Jones Protein**Intestinal Neoplasms**

Lymphosarcoma, 78-5886

Benz(a)anthracene**Lecithins**

Membrane Uptake, 78-5533

Silica

Adsorption, 78-5533

Co-carcinogenic Effect, 78-5533

Membrane Uptake, 78-5533

Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-7-methyl-**Chromatids**

Mutagenic Activity, Hamster Ovary Cells, 78-5511

Benz(a)anthracene, 3,9-Dihydroxy-7,12-dimethyl-**Benz(a)anthracene, 7,12-Dimethyl-**

Diphenolic Derivative, 78-5544

Naphthalene, 2-Bromo-6-methoxy-

Synthesis, 78-5544

Phthalic Anhydride, 4-Methoxy-

Synthesis, 78-5544

Benz(a)anthracene, 4,7-Dimethyl-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benz(a)anthracene, 7,12-Dimethyl-
Adenosine Triphosphatase
Muscle, Connective Tissue, Rat, 78-5503
Aroclor 1254
Ames Test, 78-5517
Barbituric Acid, 5-Ethyl-5-phenyl-
Ames Test, 78-5517
Benz(a)anthracene, 3,9-Dihydroxy-7,12-dimethyl-
Diphenolic Derivative, 78-5544
5,6-Benzoflavone
Ames Test, 78-5517
Brain Neoplasms
Cells, Cultured, 78-5507
Sarcoma, Osteogenic, 78-5507

Calcium
Adenosine Triphosphatase, 78-5503

Cell Division
Nucleic Acids, Binding, 78-5550

Cell Transformation, Neoplastic
Embryo Cells, 78-5550
Nucleic Acids, Binding, 78-5550

Cells, Cultured
Mutagenic Activity, 78-5518
Nucleic Acids, Binding, 78-5550
Prostaglandins, 78-5509
Proteins, Binding, 78-5550

Colon
DNA, Binding, 78-5508
Metabolism, Rat, 78-5508

DNA, Binding
Brain, Fetus, 78-5536
Skin, Mouse, 78-5504

Fluorescence
DNA, Binding, 78-5513
Hydrocarbon-DNA Adducts, 78-5513

Mammary Neoplasms, Experimental
Acetate Incorporation, 78-5543
Adenocarcinoma, 78-5535, 78-5537, 78-5539
Adenofibroma, 78-5535, 78-5537
Carcinoembryonic Antigen, 78-5539
DNA Replication, 78-5537
Estrogens, 78-5546
Parity, 78-5537
Precancerous Conditions, 78-5537
Prolactin, 78-5545
RNA, Viral, 78-5535
Thyroxine, 78-5545
Transplantation Immunology, 78-5539
Uracil, 6-Propyl-2-thio-, 78-5545
Virus, C-Type RNA Tumor, 78-5535

Maternal-Fetal Exchange
Carcinogenic Metabolite, 78-5536
DNA, Binding, 78-5536
Tissue Distribution, Rat, 78-5541

Mucopolysaccharidosis
Cell Transformation, Neoplastic, 78-5540
Co-carcinogenic Effect, 78-5540
Fibroblasts, 78-5540

Nucleic Acids, Binding
Embryo Cells, 78-5550

Pancreatic Neoplasms
Adenocarcinoma, 78-5534
Adenosine Cyclic 3',5' Monophosphate, 78-5534
Guanosine Cyclic 3',5' Monophosphate, 78-5534

Benz(a)anthracene, 7,12-Dimethyl- (cont'd)
Papilloma

Immunosuppression, 78-5538
Nude/Asplenic Mouse, 78-5538

Radiation
Enzyme Activation, 78-5518
Mutagenic Activity, 78-5518

Rhabdomyosarcoma
Cell Differentiation, 78-5542
Precancerous Conditions, 78-5542
Ultrastructural Study, Hamster, 78-5542

Sarcoma
Carcinogenic Potential, Rat, 78-5547

Sarcoma, Osteogenic
Cell Transformation, Neoplastic, 78-5507
Histological Study, Rabbit, 78-5506
Tumorigenicity, Clone Cells, 78-5507

Skin Neoplasms
Papilloma, 78-5538

Theophylline
Mutagenic Activity, 78-5518

Ultraviolet Rays
Co-carcinogenic Effect, 78-5540

Benz(a)anthracene, 7,12-Dimethyl-1-fluoro-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benz(a)anthracene, 7,12-Dimethyl-2-fluoro-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benz(a)anthracene, 7,12-Dimethyl-4-fluoro-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benz(a)anthracene-7,12-dione, 5-Fluoro-
Sarcoma
Carcinogenic Potential, Rat, 78-5547

Benz(a)anthracene, 2-Fluoro-7-methyl-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benz(a)anthracene-7-methanol, 5-Fluoro-12-methyl-
Sarcoma
Carcinogenic Potential, Rat, 78-5547

Benz(a)anthracene-7-methanol, 5-Fluoro-12-methyl-, Acetate-
Sarcoma
Carcinogenic Potential, Rat, 78-5547

Benz(a)anthracene, 7-Methyl-
Chromatids
Mutagenic Activity, Hamster Ovary Cells, 78-5518

Benz(a)anthracene, 2,7,12-Trimethyl-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benz(a)anthracene, 4,7,12-Trimethyl-
Structure-Activity Relationship
Carcinogenic Activity, Rat, 78-5505

Benzene
Hodgkin's Disease
Occupational Hazard, 78-5919
Leukemia, Myeloblastic
Epidemiology, Review, 78-5437
Lung Neoplasms
Air Pollution, Review, 78-5409
Occupational Hazard
Carcinogenic Potential, Review, 78-5437

azene (cont'd)

Epidemiology, Review, 78-5410

azene, 4-Allyl-1,2-(methylenedioxy)-

Aroclor 1254

Carcinogenic Metabolite, 78-5519

Barbituric Acid, 5-Ethyl-5-phenyl-

Carcinogenic Metabolite, 78-5519

Cholanthrene, 3-Methyl-

Carcinogenic Metabolite, 78-5519

Microsomes, Liver

Carcinogenic Metabolite, 78-5519

azene, (Epoxyethyl)-

Ames Test

Mutagenic Activity, Review, 78-5436

Nucleic Acids, Binding

Carcinogenic Activity, 78-5436

azo(a)pyren-3-ol

Benzo(a)pyrene

Glucuronidase, 78-5552

Sulfuric Acid, Disodium Salt

Conjugation Product, 78-5566

azo(a)pyren-6-ol

Benzo(a)pyrene, 6-Oxy-

Free Radicals, 78-5565

azo(a)pyren-9-ol

Benzo(a)pyrene

Glucuronidase, 78-5552

azo(a)pyrene

Adenosine Triphosphate

Ames Test, 78-5559

Ames Test

Nitro Derivatives, 78-5568

Aroclor 1254

Ames Test, 78-5517

Aryl Hydrocarbon Hydroxylases

DNA, Binding, 78-5553

Asbestos

Adsorption, 78-5556

Barbituric Acid, 5-Ethyl-5-phenyl-

Ames Test, 78-5517

Aryl Hydrocarbon Hydroxylases, 78-5553

DNA, Binding, 78-5553

Benzo(a)pyren-3-ol

Glucuronidase, 78-5552

Benzo(a)pyren-9-ol

Glucuronidase, 78-5552

Benzo(a)pyrene, 9,10-Dihydro-9,10-dihydroxy-

Carcinogenic Metabolite, 78-5554

Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-

tetrahydro-

Carcinogenic Metabolite, 78-5551

5,6-Benzoflavone

Ames Test, 78-5517

7,8-Benzoflavone

DNA, Binding, 78-5557

Bile Acids and Salts

DNA, Binding, 78-5562

Bronchial Neoplasms

Carcinoma, Epidermoid, 78-5522

Transplantation, Autologous, 78-5522

Carcinogen, Environmental

Metabolism, Review, 78-5406

Cells, Cultured

Prostaglandins, 78-5509

Benzo(a)pyrene (cont'd)5 β -Cholan-24-oic Acid, 3 α -Hydroxy-

DNA, Binding, 78-5562

Cholanthrene, 3-Methyl-

Aryl Hydrocarbon Hydroxylases, 78-5553

DNA, Binding, 78-5553

Chromatids

Mutagenic Activity, Hamster Ovary Cells, 78-5511

Chromatin, Binding

Lung, Hamster Embryo, 78-5551

Colon

Carcinogenic Metabolite, 78-5562

DNA, Binding, 78-5508

Metabolism, Rat, 78-5508

Colonic Neoplasms

Carcinogenic Metabolite, 78-5562

Deoxyguanosine Adducts

Computer Model, 78-5502

Dieldrin

Aryl Hydrocarbon Hydroxylases, 78-5553

DNA, Binding, 78-5553

Disulfide, Bis(diethylthiocarbamoyl)-

DNA, Binding, 78-5557

DNA, Binding

Bronchi, 78-5557

Liver, Rat, 78-5553

Lung, Hamster Embryo, 78-5551

Skin, Mouse, 78-5504

Fluorescence

DNA, Binding, 78-5513

Food Contamination

Smoked Products, 78-5555

2-Furaldehyde

Co-carcinogenic Effect, 78-5571

Glucuronidase

Carcinogenic Metabolite, 78-5552

Metabolism, Dose-Response Study, 78-5552

Glutathione

Ames Test, 78-5559

Lung Neoplasms

Adenocarcinoma, 78-5878

Adenoma, 78-5570

Carcinoid Tumor, 78-5878

Carcinoma, Epidermoid, 78-5878

Fibrosarcoma, 78-5878

Iron Oxide, 78-5878

Papilloma, 78-5878

Precancerous Conditions, 78-5570

Ultrastructural Study, Hamster, 78-5570, 78-5878

Lymphocytes

Metabolism, Fluorescence Study, 78-5558

Macrophages

Metabolism, Fluorescence Study, 78-5558

Microsomes, Liver

Ames Test, 78-5563

DNA, Binding, 78-5561

Polyribonucleotides, Binding, 78-5561

Nitric Acid

Ames Test, 78-5568

Nitrogen Oxide

Ames Test, 78-5568

Oxygenases

Enzymatic Activity, 78-5554

Prostatic Neoplasms

Occupational Hazard, 78-5960

Respiratory Tract Neoplasms

2-Furaldehyde, 78-5571

Benzo(a)pyrene (cont'd)

- Histological Study, Dog, 78-5569
- Iron Oxide, 78-5569, 78-5876
- Precancerous Conditions, 78-5569, 78-5876
- Ultrastructural/Histochemical Study, Hamster 78-5876
- Urea, Methyl Nitroso-, 78-5569
- Retinol
 - DNA, Binding, 78-5557
- Silica
 - Adsorption, 78-5556
- Skin Neoplasms
 - 7,8-Benzoflavone, 78-5567
 - Flavone, 3,3',4',5,7-Pentahydroxy-, 78-5567
 - 2-Imidazoline, 2-Benzyl-, 78-5567
 - Papilloma, 78-5567
 - 12-O-Tetradecanoylphorbol-13-acetate, 78-5567
- Stress
 - DNA, Binding, 78-5560
 - Enzyme Activation, 78-5560
- Sulfuric Acid, Disodium Salt
 - Conjugation, Phenol/Quinone Metabolites, 78-5566
- Taurodeoxycholic Acid
 - DNA, Binding, 78-5562
- Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Trachea
 - Carcinogenic Metabolite, 78-5554
 - Metabolism, Microsomes, 78-5554
 - Metabolism, Organ Culture, 78-5554
- Tritium Loss
 - Binding Sites, 78-5561
- Uridine Diphosphate Sugars
 - Ames Test, 78-5559
- Vitamin E
 - DNA, Binding, 78-5557
- Water Pollution
 - Quantitation Method, 78-5549
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-**
 - Cells, Cultured
 - Carcinogenic Metabolite, 78-5552
 - Chromatids
 - Mutagenic Activity, Hamster Ovary Cells, 78-5511
 - Colon
 - Carcinogenic Metabolite, 78-5562
- Benzo(a)pyrene, 9,10-Dihydro-9,10-dihydroxy-**
 - Benzo(a)pyrene
 - Carcinogenic Metabolite, 78-5554
 - Cells, Cultured
 - Carcinogenic Metabolite, 78-5552
 - Colon
 - Carcinogenic Metabolite, 78-5562
 - Trachea
 - Carcinogenic Metabolite, 78-5554
- Benzo(a)pyrene, 12b,12c-Dihydro-12b,12c-dimethyl-**
 - Microsomes, Liver
 - Ames Test, 78-5563
- Benzo(a)pyrene, 7,8-Dihydro-7,8,9-trihydroxy-**
 - Colon
 - Carcinogenic Metabolite, 78-5562
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-**
 - Benzo(a)pyrene
 - Carcinogenic Metabolite, 78-5551
 - Carcinogenic Metabolite
 - Lung, Hamster Embryo, 78-5551
- Benzo(a)pyrene, 7,10-Dimethyl-**
 - Structure-Activity Relationship
 - Carcinogenic Activity, Rat, 78-5505
- Benzo(a)pyrene-6-methanol**
 - Skin Neoplasms
 - 7,8-Benzoflavone, 78-5567
 - 12-O-Tetradecanoylphorbol-13-acetate, 78-5567
- Benzo(a)pyrene, 6-Methyl-**
 - Skin Neoplasms
 - 7,8-Benzoflavone, 78-5567
 - 12-O-Tetradecanoylphorbol-13-acetate, 78-5567
- Benzo(a)pyrene, 7-Methyl-**
 - Structure-Activity Relationship
 - Carcinogenic Activity, Rat, 78-5505
- Benzo(a)pyrene, 10-Methyl-**
 - Structure-Activity Relationship
 - Carcinogenic Activity, Rat, 78-5505
- Benzo(a)pyrene 4,5-Oxide**
 - Cytochrome P-450
 - NADH, NADPH Oxidoreductases, 78-5564
 - Microsomes, Liver
 - NADH, NADPH Oxidoreductases, 78-5564
 - NADH, NADPH Oxidoreductases
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5564
 - Cholanthrene, 3-Methyl-, 78-5564
 - Riboflavine, 78-5564
- Benzo(a)pyrene, 6-Oxy-**
 - Benzo(a)pyren-6-ol
 - Free Radicals, 78-5565
 - Caffeine
 - DNA, Binding, 78-5565
 - Nuclear Magnetic Resonance, 78-5565
 - DNA
 - Nuclear Magnetic Resonance, 78-5565
- Benzo(a)pyrene, 7,8,9,10-Tetrahydro-7,8,9,10-tetrahydroxy-**
 - Colon
 - Carcinogenic Metabolite, 78-5562
- Benzo(b,def)chrysene, 13b,13c-Dihydro-13b,13c-dimethyl-**
 - Microsomes, Liver
 - Ames Test, 78-5563
- Benzo(b)triphenylene**
 - Cells, Cultured
 - Nucleic Acids, Binding, 78-5550
 - Proteins, Binding, 78-5550
- Benzo(e)pyrene**
 - Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Benzo(e)pyrene, 12b,12c-Dihydro-12b,12c-dimethyl-**
 - Microsomes, Liver
 - Ames Test, 78-5563
- 5,6-Benzoflavone**
 - Ames Test
 - S-9 Fraction, 78-5517
 - 2-Anthracenamide
 - Ames Test, 78-5517
 - Benz(a)anthracene, 7,12-Dimethyl-
 - Ames Test, 78-5517
 - Benzo(a)pyrene
 - Ames Test, 78-5517
 - Cyclophosphamide
 - Ames Test, 78-5517

Benzo(a)fluorene (cont'd)

- Ethidium Bromide
 - Ames Test, 78-5517
- Fluorene-2-amine
 - Ames Test, 78-5517
- Liver Neoplasms
 - Acetamide, *N*-Fluorene-2-yl-, 78-5481
- Morpholine, *N*-Nitroso-
 - Ames Test, 78-5517

Benzo(a)fluorene

- Acetamide, *N*-Fluorene-2-yl-
 - RNA, Binding, 78-5479
- Benzo(a)pyrene
 - DNA, Binding, 78-5557
- Skin Neoplasms
 - Benzo(a)pyrene, 78-5567
 - Benzo(a)pyrene-6-methanol, 78-5567
 - Benzo(a)pyrene, 6-Methyl-, 78-5567
 - 2-Imidazoline, 2-Benzyl-, 78-5567

Benzo(a)fluorene-3,3'-Dithiobis(6-nitro)-

- Virus, Avian Myeloblastosis
 - DNA Polymerase, 78-5743
 - Ribonuclease, 78-5743

Benzo(a)fluorene-3,3'-Dithiobis(6-nitro)-

- Virus, Avian Myeloblastosis
 - DNA Polymerase, 78-5743
 - Ribonuclease, 78-5743

Benzo(a)fluorene-2-Nitro-5-thiocyano-

- Virus, Avian Myeloblastosis
 - DNA Polymerase, 78-5743
 - Ribonuclease, 78-5743

Benzo(a)fluorene-2-Nitro-5-thiocyano-

- Sodium Salt
 - Chromosomes
 - Mutagenic Activity, 78-5635
- Vicia faba*
 - Mutagenic Activity, 78-5635

Benzo(a)fluorene-2,3,5,6-Tetrakis(1-aziridinyl)-

- Neurospora crassa*
 - Mutagenic Activity, 78-5675

Benzo(a)fluorene-2-Nitro-5-thiocyano-

- Esophageal Neoplasms
 - Nitrous Acid, Sodium Salt, 78-5586
 - Precancerous Conditions, 78-5586

Benzo(a)fluorene-2-Nitro-5-thiocyano-

- Esophageal Neoplasms
 - Precancerous Conditions, 78-5586
 - Ulcer, 78-5586
 - Zinc, 78-5587
- Esophagus
 - Organ Distribution, Rat, 78-5586

Benzo(a)fluorene-2-Nitro-5-thiocyano-

- Benzo(a)pyrene
 - DNA, Binding, 78-5562
- Intestinal Neoplasms
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5684

Bladder Diseases

- Cystitis
 - Nitrosamines, 78-5591

Bladder Neoplasms

- Acetamide, *N*-Fluorene-2-yl-
 - Precancerous Conditions, 78-5476
- Bracken Fern

Bladder Neoplasms (cont'd)

- Carcinogenic Activity, Rat, 78-5464
- 1-Butanol, 4-(Butylnitrosamino)-
 - Chromosomes, 78-5581
 - Precancerous Conditions, 78-5583
 - Retinoic Acid, 78-5582
 - Ultrastructural Study, Rat, 78-5581
- Carcinoma
 - Chromosome Abnormalities, 78-5893
 - Cyclophosphamide, 78-5894
- Carcinoma, Epidermoid
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5581, 78-5582
- Carcinoma, Transitional Cell
 - Acetamide, *N*-Fluorene-2-yl-, 78-5476
 - Bracken Fern, 78-5465
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5581, 78-5582
- Chromosome Abnormalities
 - Chromosomes, Human, 1-3, 78-5893
- Clone Cells
 - Ultrastructural Study, Rat, 78-5581
- Cyclophosphamide
 - Carcinogenic Effect, Review, 78-5420
 - Case Report, 78-5894
- Dimethylamine, *N*-Nitroso-
 - Urine, 78-5591
- Hemangioma
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5583
- Heterochromatin
 - Chromosome Abnormalities, 78-5893
- Hyperplasia
 - Acetamide, *N*-Fluorene-2-yl-, 78-5476
- Multiple Myeloma
 - Drug Therapy, 78-5894
- Neoplasms, Multiple Primary
 - Bracken Fern, 78-5465
- Nitrosamines
 - Urine, 78-5591
- Occupational Hazard
 - Epidemiology, Review, 78-5401
- Papilloma
 - Antipain, 78-5583
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5582, 78-5583
 - Leupeptins, 78-5583
 - Pepstatins, 78-5583
- Retinoic Acid
 - Neoplasm Invasiveness, 78-5582
- Schistosoma haematobium*
 - Dimethylamine, *N*-Nitroso-, 78-5591
 - Nitrosamines, 78-5591

Blood Cells

- Virus, Marek's Disease Herpes
 - Antigens, Viral, 78-5749

Blood Groups

- Stomach Neoplasms
 - Epidemiology, 78-5942

Blood Platelets

- Leukemia, Myelocytic
 - Peroxidases, 78-5899

Bone and Bones

- Plutonium
 - Body Burden, Dog, 78-5732, 78-5734

Bone Marrow Cells

- Erythroleukemia
 - Acetamide, *N*-Fluorene-2-yl-, 78-5483
- Ethylene, Chloro- Polymer

Bone Marrow Cells (cont'd)

- Chromosome Aberrations, 78-5488
- Leukemia, Myeloblastic
 - Cell Differentiation, 78-5900
- Leukemia, Myelocytic
 - Acetamide, *N*-Fluoren-2-yl-, 78-5483
 - Cell Differentiation, 78-5900
 - Neoplasm Metastasis, 78-5484
- Thorium Dioxide
 - Case Report, 78-5717

Bone Neoplasms

- Carcinoma
 - Radium, 78-5729
- Radium
 - Epidemiology, 78-5729
 - Mastoid, 78-5729
 - Occupational Hazard, 78-5729
- Thorium Dioxide
 - Epidemiology, Portugal, 78-5973

Bracken Fern

- Bladder Neoplasms
 - Carcinogenic Activity, Rat, 78-5464
 - Carcinoma, Transitional Cell, 78-5465
 - Neoplasms, Multiple Primary, 78-5465
- Boiling Water Extract
 - Carcinogenic Activity, 78-5465
- Intestinal Neoplasms
 - Adenocarcinoma, 78-5464, 78-5465
 - Adenoma, 78-5464, 78-5465
 - Carcinogenic Activity, Rat, 78-5464
 - Neoplasms, Multiple Primary, 78-5465
 - Precancerous Conditions, 78-5464
 - Sarcoma, 78-5464
- Virus, Papilloma
 - Co-carcinogenic Activity, 78-5790
 - Epidemiology, Cattle, 78-5790

Brain Neoplasms

- Adenoma
 - Urea, Ethyl Nitroso-, 78-5650
- Astrocytoma
 - Urea, Ethyl Nitroso-, 78-5649
 - Urea, Methyl Nitroso-, 78-5859
- Benz(a)anthracene, 7,12-Dimethyl-
 - Cells, Cultured, 78-5507
- Burkitt's Lymphoma
 - Neoplasm Metastasis, 78-5456
- Carcinogen, Chemical
 - Occupational Hazard, 78-5926
- Child
 - Epidemiology, Denmark, 78-5925
- Choriocarcinoma
 - Neoplasm Metastasis, 78-5456
- Glioblastoma Multiforme
 - Epidemiology, Sweden, 78-5927
- Glioma
 - Epidemiology, Sweden, 78-5927
 - Radiation, Ionizing, 78-5872
 - Urea, Ethyl Nitroso-, 78-5649
 - Urea, Methyl Nitroso-, 78-5859
 - Virus, Polyoma, BK, 78-5794
- Meningioma
 - Epidemiology, Nigeria, 78-5456
 - Epidemiology, Sweden, 78-5927
 - Epidemiology, Women, 78-5926
 - Nitrous Acid, Sodium Salt, 78-5926
 - Radiation, Ionizing, 78-5926

Brain Neoplasms (cont'd)

- Mouth Diseases
 - Microcirculation, 78-5927
- Neoplasm Metastasis
 - Epidemiology, Nigeria, 78-5456
- Oligodendroglioma
 - Urea, Ethyl Nitroso-, 78-5649
 - Urea, Methyl Nitroso-, 78-5859
- Radiotherapy
 - Prognosis, Child, 78-5925
- Sarcoma, Osteogenic
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5507
- Surgery, Operative
 - Prognosis, Child, 78-5925
- Urea, Ethyl Nitroso-
 - Organ Culture, 78-5650
 - Transplacental Carcinogenesis, 78-5649, 78-5650
 - Ultrastructural Study, 78-5649
- Urea, Methyl Nitroso-
 - Antibody-Dependent Cytotoxicity, 78-5859
 - Antibody Formation, 78-5859
 - Antibody Specificity, 78-5859
- Virus, Polyoma, BK
 - Ultrastructural Study, Hamster, 78-5794

Breast

- Pregnancy
 - Tissue Regression, 78-5891

Breast Diseases

- Cysts
 - Proteins, 78-5993
- Progesterone
 - Isolation and Characterization, Binding Component 78-5993

Breast Neoplasms

- Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-
 - Epidemiology, Japan, 78-5946
- Adenocarcinoma
 - Cells, Cultured, 78-5950
 - Insulin, 78-5950
- Aging
 - Endocrine Glands, Review, 78-5411
- Carcinogen, Environmental
 - Epidemiology, Review, 78-5463
- Carcinoma, Ductal
 - Culture Media, 78-5994
 - Growth Substances, 78-5994
- Cells, Cultured
 - Growth, 78-5950
 - Multicellular Spheroid Formation, 78-5996
- Contraceptive Devices
 - Epidemiology, 78-5947
- Diet
 - Epidemiology, Hawaii, 78-5945
- Dietary Fats
 - Epidemiology, Japan, 78-5946
 - Epidemiology, Review, 78-5404
- Epidemiology
 - Spain, 78-5915
 - USSR, 78-5460
- Estradiol
 - Plasma/Urine Ratios, 78-5991
- Estriol
 - Plasma/Urine Ratios, 78-5991
- Estrogens
 - Epidemiology, Chinese, 78-5891
 - Epidemiology, Review, 78-5432

east Neoplasms (cont'd)

- Genetics, 78-5891
 - Menopause, 78-5949
 - Estrone
 - Plasma/Urine Ratios, 78-5991
 - Ethnic Groups
 - Epidemiology, Hawaii, 78-5945
 - Genetics
 - Classification, Review, 78-5455
 - Growth Substances
 - Epithelial Cells, 78-5994
 - Hysterectomy
 - Epidemiology, 78-5948
 - Insulin
 - Growth, 78-5950
 - Leukemia, Lymphocytic
 - Neoplasms, Multiple Primary, 78-5920
 - Menarche
 - Epidemiology, Iceland, 78-5951
 - Menopause
 - Epidemiology, 78-5949
 - Menstruation
 - Epidemiology, 78-5949
 - Parity
 - Epidemiology, Iceland, 78-5951
 - Pleural Effusion
 - Multicellular Spheroid Formation, 78-5996
 - Progesterone
 - Menopause, 78-5949
 - Prolactin
 - Neoplasm Metastasis, 78-5990
 - Serum Levels, 78-5990
 - Protirelin
 - Prolactin, 78-5990
 - Thyrotropin, 78-5990
 - Rubber
 - Epidemiology, 78-5954
 - Occupational Hazard, Women, 78-5954
 - Semen
 - Prostaglandins, 78-5947
 - Sterility, Male
 - Epidemiology, 78-5947
 - Thyroid Diseases
 - Epidemiology, 78-5989
 - Thyroid Hormones
 - Serum Levels, 78-5989
 - Thyronine, 3,3',5-Triiodo-
 - Metabolism, 78-5990
 - Thyrotropin
 - Neoplasm Metastasis, 78-5990
 - Radiotherapy, 78-5990
 - Serum Levels, 78-5989, 78-5990
 - Thyroxine
 - Serum Levels, 78-5990
- ## Bronchial Neoplasms
- Benzo(a)pyrene
 - Transplantation, Autologous, 78-5522
 - Carcinoma, Epidermoid
 - Benzo(a)pyrene, 78-5522
 - Cholanthrene, 3-Methyl-, 78-5522
 - Precancerous Conditions, 78-5522
 - Cholanthrene, 3-Methyl-
 - Transplantation, Autologous, 78-5522
 - Precancerous Conditions
 - Histological/Ultrastructural Study, Dog, 78-5522

Bronchitis

- Carcinoembryonic Antigen
 - Epidemiology, 78-5940
 - Serum Levels, 78-5940
- Smoking
 - Carcinoembryonic Antigen, 78-5940

Burkitt's Lymphoma

- Antibodies, Viral
 - Immune Response, Infant, 78-5836
- Brain Neoplasms
 - Neoplasm Metastasis, 78-5456
- Cells, Cultured
 - Cortisone, 78-5824
- Cortisone
 - Cell Survival, 78-5824
 - Interferon, 78-5824
- Hybrid Cells
 - Transplantation, Heterologous, 78-5832
- Transplantation, Heterologous
 - Mouse, Nude, 78-5832
- Virus, Epstein-Barr
 - Antigens, Viral, 78-5824, 78-5827
 - Cortisone, 78-5824
 - Epidemiology, Review, 78-5441
 - Receptors, Complement, 78-5827
 - Receptors, Viral, 78-5827
 - Seroepidemiology, Review, 78-5443

1,3-Butadiene, 1-Chloro-

- Ames Test
- Mutagenic Activity, 78-5492

1,3-Butadiene, 2-Chloro-

- Ames Test
- Mutagenic Activity, 78-5492
- Occupational Hazard
 - Epidemiology, Review, 78-5410
- Pyridine, 4-(p-Nitrobenzyl)-
 - Carcinogenic Metabolite, 78-5492

1-Butanamine, N-Butyl-N-nitroso-

- Cell Transformation, Neoplastic
 - Bladder, Mouse, 78-5584
- Dimethylamine, N-Nitroso-
 - Purine, 2-Amino-6-methoxy-, 78-5601
- DNA Replication
 - Bladder, Mouse, 78-5584
 - Cell Cycle Kinetics, 78-5584
- Food Contamination
 - Quantitation Method, 78-5616
- Occupational Hazard
 - Permeability, Laboratory Gloves, 78-5595
 - Permeability, Rubber Gloves, 78-5597

1,4-Butanediol, Dimethylsulfonate

- Cystitis
 - Case Report, 78-5897
- Leukemia, Myelocytic
 - Cystitis, 78-5897
 - Hemorrhage, 78-5897

1-Butanol, 4-(Butylnitrosamino)-

- Bladder Neoplasms
 - Carcinoma, Epidermoid, 78-5581, 78-5582
 - Carcinoma, Transitional Cell, 78-5581, 78-5582
 - Chromosomes, 78-5581
 - Hemangioma, 78-5583
 - Papilloma, 78-5582, 78-5583
 - Precancerous Conditions, 78-5583

1-Butanol, 4-(Butylnitrosamino)- (cont'd)
Retinoic Acid, 78-5582
Ultrastructural Study, Rat, 78-5581

1-Butene, 3,4-Dichloro-
Ames Test
Carcinogenic Metabolite, 78-5492
Mutagenic Activity, 78-5492

2-Butene, 1,4-Dichloro-
Ames Test
Mutagenic Activity, 78-5492

Butylamine, N-Methyl-N-nitroso-
Occupational Hazard
Permeability, Laboratory Gloves, 78-5596

Butyric Acid, 4-(p-Bis(2-chloroethyl)aminophenyl)-
Leukemia, Myeloblastic
Case Report, 78-5896
Glomerulonephritis, 78-5896

Butyrophene, γ-Chloro-p-fluoro-
Escherichia coli
DNA Repair, 78-5449
Salmonella typhimurium
DNA Repair, 78-5449

Bux-Ten, N-Nitroso-
Stomach Neoplasms
Carcinoma, 78-5598
Papilloma, 78-5598

Caffeine
Benzo(a)pyrene, 6-Oxy-
DNA, Binding, 78-5565
Nuclear Magnetic Resonance, 78-5565
Drosophila melanogaster
Mutagenic Activity, 78-5656
Methanesulfonic Acid, Ethyl Ester
Mutagenic Activity, 78-5654
Nitrous Acid
Mutagenic Activity, 78-5654
Psoralen, 8-Methoxy-
Chromosome Aberrations, 78-5703
Quinoline, 4-Nitro-, 1-Oxide
Mutagenic Activity, 78-5671
Ultraviolet Rays
Chromosome Aberrations, 78-5703

Calcium
Benz(a)anthracene, 7,12-Dimethyl-
Adenosine Triphosphatase, 78-5503
Carbamic Acid, Ethyl Ester
Adenosine Triphosphatase, 78-5503
Dibenz(a,h)anthracene
Adenosine Triphosphatase, 78-5503
Dimethylamine, N-Nitroso-
Adenosine Triphosphatase, 78-5503
Iodine Radioisotopes
Serum Levels, Rat, 78-5713

Californium
Chromosome Aberrations
Dose-Response Study, 78-5725
Lymphocytes, 78-5725
Lung Neoplasms
Adenocarcinoma, 78-5726
Carcinogenic Activity, Rat, 78-5726
Carcinoma, Epidermoid, 78-5726
Lymphosarcoma, 78-5726
Radioisotopes, 78-5726

Californium (cont'd)
Radioisotopes
Chromosome Aberrations, 78-5725

Captan
see 4-Cyclohexene-1,2-dicarboximide, N-(Trichloromethyl)thio-

Carbamic Acid, Ethyl Ester
Adenosine Triphosphatase
Muscle, Connective Tissue, Rat, 78-5503
Calcium
Adenosine Triphosphatase, 78-5503

Carbamic Acid, Ethyl-, Ethyl Ester
Nitrous Acid, Sodium Salt
Nitrosation, 78-5636
Urea, Methyl Nitroso-
Nitrous Acid, Sodium Salt, 78-5636

Carbamic Acid, N-Methyl-N-nitroso-, o-Isopropoxyphenyl Ester
Stomach Neoplasms
Carcinoma, 78-5598
Papilloma, 78-5598

Carbamic Acid, N-Methyl-N-nitroso-, 1-Naphthyl Ester
Stomach Neoplasms
Carcinoma, 78-5598
Papilloma, 78-5598

Carbamic Acid, N-Methyl-N-nitroso-, Phenyl Ester
Stomach Neoplasms
Carcinoma, 78-5598
Papilloma, 78-5598

Carbamic Acid, N-Methyl-N-nitroso-, Trimethylphenyl Ester
Stomach Neoplasms
Carcinoma, 78-5598
Papilloma, 78-5598

Carbofuran, N-Nitroso-
Stomach Neoplasms
Carcinoma, 78-5598
Papilloma, 78-5598

Carbohydrates
Virus, Murine Leukemia
Glycoproteins, 78-5781
Virus, Rauscher Murine Leukemia
Glycoproteins, 78-5781

Carbon Dioxide
Dimethylamine, N-Nitroso-
Metabolism, Rat, 78-5603
Nucleic Acids, Binding, 78-5603

Carbon Tetrachloride
Dimethylamine, N-Nitroso-
Metabolism, Rat, 78-5603
Nucleic Acids, Binding, 78-5603
Liver Neoplasms
Diethylamine, N-Nitroso-, 78-5592
Hepatectomy, 78-5592

Carcinoembryonic Antigen
Air Pollution
Serum Levels, 78-5940
Bronchitis
Epidemiology, 78-5940
Serum Levels, 78-5940
Smoking, 78-5940
Carcinoma, Epidermoid

Carcinoembryonic Antigen (cont'd)

Immunotherapy, 78-5854

Cholanthrene, 3-Methyl-

Cell Transformation, Neoplastic, 78-5740

Fibrosarcoma

Antibody Specificity, 78-5866

Cholanthrene, 3-Methyl-, 78-5866

Gastrointestinal Neoplasms

Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5867

Hypersensitivity, Delayed, 78-5867

Immune Response, 78-5867

Immunization, 78-5867

Neoplasm Metastasis, 78-5867

Mammary Neoplasms, Experimental

Benz(a)anthracene, 7,12-Dimethyl-, 78-5539

Immunization, 78-5539

Transplantation Immunology, 78-5539

Prostatic Neoplasms

Antibody Formation, 78-5864

Immunization, 78-5864

Virus, SV40, 78-5864

Sarcoma

Carcinogen, Chemical, 78-5438

Smoking

Serum Levels, 78-5940

Virus, Avian Sarcoma

Cell Transformation, Neoplastic, 78-5740

Carcinogen, Chemical**Aging**

DNA Repair, Review, 78-5411

Enzyme Activation, Review, 78-5411

Ames Test

Evaluation, Review, 78-5427

Aroclor 1254

Enzyme Activation, 78-5480

Bacteria

Mutagenic Activity, Review, 78-5418

Bacteriophages

Derepression, Review, 78-5448

Mutagenic Activity, Review, 78-5448

Brain Neoplasms

Occupational Hazard, 78-5926

Chromosomes

Mutagenic Activity, Review, 78-5418

DNA, Binding

Carcinogenic Activity, Review, 78-5418

Enzyme Activation

Carcinogenic Metabolite, Review, 78-5419

Liver Cells, Intact, 78-5480

Hypoxanthine Phosphoribosyltransferase

Mutation, 78-5652

Lymphoma

Mutation, 78-5652

Metabolism

Computer Analysis, 78-6000

Metallurgical Plants

Epidemiology, 78-5924

Nucleic Acids

Carcinogenic Activity, Review, 78-5415

Carcinogenic Metabolite, Review, 78-5419

Mutagenic Activity, Review, 78-5415

Sarcoma

Antigens, Review, 78-5438

Antigens, Viral, 78-5438

Carcinoembryonic Antigen, 78-5438

Histocompatibility Antigens, 78-5438

Structure-Activity Relationship**Carcinogen, Chemical (cont'd)**

Computer Analysis, 78-6000

Thymidine Kinase

Mutation, 78-5652

Carcinogen, Environmental**Aflatoxin B1**

Metabolism, Review, 78-5406

Bacteria

Mutagenic Activity, 78-5426

Benzo(a)pyrene

Metabolism, Review, 78-5406

Breast Neoplasms

Epidemiology, Review, 78-5463

Chromosome Abnormalities

Genetics, Review, 78-5458

Chromosomes

Mutagenic Activity, Review, 78-5455

Colonic Neoplasms

Epidemiology, Review, 78-5463

Diet

Epidemiology, Review, 78-5402

Dimethylamine, N-Nitroso-

Epidemiology, Review, 78-5414

Dose-Response Study

Animal Model, Review, 78-5408

Data Extrapolation, Review, 78-5408

Mathematical Model, 78-5923

Threshold Values, Review, 78-5412

Ethyl Alcohol

Epidemiology, Review, 78-5402

Ethylene, Chloro-

Metabolism, Review, 78-5406

Genetics

Cell Transformation, Neoplastic, Review, 78-5455

Epidemiology, Review, 78-5405

Immunologic Deficiency Syndromes

Risk Factors, Review, 78-5458

Laryngeal Neoplasms

Epidemiology, 78-5936

Epidemiology, Georgian SSR, 78-5937

Precancerous Conditions, 78-5936

Lung Neoplasms

Epidemiology, Review, 78-5463

Neoplasms

Epidemiology, Review, 78-5463

Nitrosamines

Isolation and Characterization, Review, 78-5413

Quantitation Method, Review, 78-5416

Nitroso Compounds

Metabolism, Review, 78-5406

Prostatic Neoplasms

Epidemiology, Review, 78-5463

Rectal Neoplasms

Epidemiology, Review, 78-5463

Risk Factors

Data Extrapolation, Review, 78-5408

Mathematical Model, 78-5923

Screening Tests, Review, 78-5401

Smoking

Epidemiology, Review, 78-5402

Teratogens

Epidemiology, Review, 78-5405

Carcinoid Tumor**Lung Neoplasms**

Benzo(a)pyrene, 78-5878

Classification, 78-5877

Carcinoma

- Bladder Neoplasms
 - Chromosome Abnormalities, 78-5893
 - Cyclophosphamide, 78-5894
- Bone Neoplasms
 - Radium, 78-5729
- Cell Line
 - Ultrastructural Study, Review, 78-5454
- Cholanthrene, 3-Methyl-
 - Transplantation, Heterologous, 78-5529
- Chromosomes
 - Tissue Culture, Review, 78-5454
- Colonic Neoplasms
 - Radiation, Ionizing, 78-5710
- Growth
 - Tissue Culture, Review, 78-5454
- Head and Neck Neoplasms
 - Virus, Epstein-Barr, 78-5830
- HeLa Cells
 - Mouse, Nude, 78-5832
- Lung Neoplasms
 - Nickel, 78-5931
- Nasopharyngeal Neoplasms
 - Virus, Epstein-Barr, 78-5830
- Paranasal Sinus Neoplasms
 - Radium, 78-5729
- Salivary Gland Neoplasms
 - Radiation, Ionizing, 78-5714
- Stomach Neoplasms
 - Bux-Ten, *N*-Nitroso-, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, *o*-Isopropoxyphenyl Ester, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, Phenyl Ester, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, Trime-thylphenyl Ester, 78-5598
 - Carbofuran, *N*-Nitroso-, 78-5598
 - Gastrectomy, 78-5879
 - Methomyl, *N*-Nitroso-, 78-5598
 - Neoplasm Recurrence, 78-5879
- Tracheal Neoplasms
 - Cholanthrene, 3-Methyl-, 78-5521
 - Urea, Methyl Nitroso-, 78-5640, 78-5642
- Vaginal Neoplasms
 - Chromosome Aberrations, 78-5871
- Virus, Epstein-Barr
 - Mouse, Nude, 78-5832
- Virus, Murine Leukemia
 - Isolation and Characterization, 78-5767

Carcinoma, Basal Cell

- Cicatrix
 - Burns, 78-5970

Carcinoma, Ductal

- Breast Neoplasms
 - Culture Media, 78-5994
 - Growth Substances, 78-5994

Carcinoma, Epidermoid

- Antigens, Neoplasm
 - Immunotherapy, 78-5854
- Bladder Neoplasms
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5581, 78-5582
- Bronchial Neoplasms
 - Benzo(a)pyrene, 78-5522
 - Cholanthrene, 3-Methyl-, 78-5522

Carcinoma, Epidermoid (cont'd)

- Precancerous Conditions, 78-5522
- Carcinoembryonic Antigen
 - Immunotherapy, 78-5854
- Cervix Neoplasms
 - Epidemiology, Israel, 78-5958
 - Ethnic Groups, 78-5958
- Cholanthrene, 1,2-Dehydro-3-Methyl-
 - Neoplasm Metastasis, 78-5523
- Cicatrix
 - Burns, 78-5970
- Colonic Neoplasms
 - Hydrazine, 1,2-Dimethyl-, 78-5498
- Digestive System Neoplasms
 - Quinoline, 8-Nitro-, 78-5672
- Ear Neoplasms
 - Hydrazine, 1,1-Dimethyl-, 78-5861
- Hodgkin's Disease
 - Neoplasms, Multiple Primary, 78-5910
- Laryngeal Neoplasms
 - Epidemiology, Georgian SSR, 78-5937
- Lung Neoplasms
 - Animal Model, Hamster, 78-5878
 - Anthracosilicosis, 78-5971
 - Azocine, Octahydro-1-nitroso-, 78-5585
 - Benzo(a)pyrene, 78-5878
 - Californium, 78-5726
 - Cholanthrene, 3-Methyl-, 78-5528
 - Classification, 78-5877
 - Nickel, 78-5931
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
 - Ultrastructural/Histochemical Study, 78-5877
- Methanesulfonic Acid, Methyl Ester
 - DNA Repair, 78-5667
- Nasopharyngeal Neoplasms
 - Virus, Epstein-Barr, 78-5830
- Neoplasm Metastasis
 - Immunotherapy, 78-5854
 - Sheep, 78-5854
- Neoplasms, Multiple Primary
 - Cicatrix, 78-5970
- Prostatic Neoplasms
 - Epidemiology, 78-5960
 - Occupational Hazard, 78-5960
- Quinoline, 4-Nitro-, 1-Oxide
 - DNA Repair, 78-5667
- Skin Neoplasms
 - Cicatrix, 78-5970
 - Virus-Like Particles, 78-5789
- Stomach Neoplasms
 - Urea, 1-Methyl-1-nitroso-3-phenyl-, 78-5637
- Tracheal Neoplasms
 - Cholanthrene, 3-Methyl-, 78-5521
 - Precancerous Conditions, 78-5978
 - Transplantation, Homologous, 78-5978
 - Urea, Methyl Nitroso-, 78-5640, 78-5642

Carcinoma In Situ

- Cervix Neoplasms
 - Drug Therapy, 78-5888
 - Epidemiology, England, 78-5955
 - Epidemiology, Review, 78-5461
 - Radiotherapy, 78-5888

Carcinoma, Mucinous

- Colonic Neoplasms
 - Hydrazine, 1,2-Dimethyl-, 78-5497
- Intestinal Neoplasms

noma, Mucinous (cont'd)
Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499

noma, Oat Cell
Lung Neoplasms
Classification, 78-5877

noma, Papillary
Thyroid Neoplasms
Iodine Radioisotopes, 78-5716
Radiation, Ionizing, 78-5715
Uterine Neoplasms
Genetics, 78-5892

noma, Transitional Cell
Bladder Neoplasms
Acetamide, *N*-Fluoren-2-yl-, 78-5476
Bracken Fern, 78-5465
1-Butanol, 4-(Butylnitrosamino)-, 78-5581, 78-5582

inosarcoma
Cholanthrene, 3-Methyl-
Transplantation, Heterologous, 78-5529

ageen
Colonic Neoplasms
Adenoma, 78-5644
Methane, Azoxy-, 78-5644
Urea, Methyl Nitroso-, 78-5644

rier Proteins
Virus, Adeno 2
DNA, Viral, 78-5841

ein
Virus, Avian Myeloblastosis
Antigens, Viral, 78-5741

Aggregation
Guanidine, 1-Methyl-3-nitro-1-nitroso-
Cell Transformation, Neoplastic, 78-5507
Virus, Kirsten Murine Sarcoma
Cell Transformation, Neoplastic, 78-5507

Differentiation
Cell Transformation, Neoplastic
Cells, Cultured, 78-5900
Estradiol
Vagina, Neonatal Mouse, 78-5890
Leukemia, Myeloblastic
Acetamide, 78-5983
Acetamide, *N,N*-Dimethyl-, 78-5983
Acetamide, *N*-Methyl-, 78-5983
Bone Marrow Cells, 78-5900
Cell Line, 78-5900
Cells, Cultured, 78-5900
Formamide, *N,N*-Diethyl-, 78-5983
Formamide, *N*-Methyl-, 78-5983
Methane, Sulfinylbis-, 78-5983
2-Piperidone, 78-5983
Triethylene Glycol, 78-5983
Leukemia, Myelocytic
Bone Marrow Cells, 78-5900
Cells, Cultured, 78-5900, 78-5985
Hematopoietic Stem Cells, 78-5904
Mesenchymoma
Hematopoietic Stem Cells, 78-5904
Phorbol 12,13-Didecanoate
5,8,11,14-Eicosatetranoic Acid, 78-5509
Rhabdomyosarcoma
Benz(a)anthracene, 7,12-Dimethyl-, 78-5542
Radiation, Ionizing, 78-5690

Cell Differentiation (cont'd)
Sex Chromosomes
Mouse Embryo, Review, 78-5453
Teratoid Tumor
Mouse Embryo, Review, 78-5453
Neoplasm Transplantation, Review, 78-5453
12-*O*-Tetradecanoylphorbol-13-acetate
5,8,11,14-Eicosatetranoic Acid, 78-5509
Tracheal Neoplasms
Cells, Cultured, 78-5978

Cell Division
Benz(a)anthracene, 7,12-Dimethyl-
Nucleic Acids, Binding, 78-5550
Fibroblasts
Epidermal Growth Factors, 78-5813

Cell Fusion
Virus, Mason-Pfizer Monkey
Virus, Rous Sarcoma, 78-5809
Virus, SV40, 78-5809
Virus, SV40
Transformation, Genetic, 78-5796
Virus Replication, 78-5796

Cell Membrane
Fibroblasts
Epidermal Growth Factors, 78-5813
Lecithins
Hydroxy Radicals, 78-5724
Leukemia
Gangliosides, 78-5987
Lymphocytes
Gangliosides, 78-5987
Lymphoma
Gangliosides, 78-5987
Plasmacytoma
Gangliosides, 78-5987
Radiation Effects
Model System, 78-5724
Virus, Epstein-Barr
Antigens, 78-5833
Virus, Feline Leukemia
Antigens, Viral, 78-5786
Virus, Moloney Murine Leukemia
Viral Proteins, 78-5776
Virus, Rauscher Murine Leukemia
Viral Proteins, 78-5776
Virus, Rous Sarcoma
Antigens, Viral, 78-5744
Virus, SV40
Alkaline Phosphatase, 78-5798
Epidermal Growth Factors, 78-5813

Cell Membrane Permeability
Aflatoxin B2
Mitochondria, Liver, 78-5469
Aflatoxin G2
Mitochondria, Liver, 78-5469

Cell Transformation, Neoplastic
Aflatoxin B1
Cells, Cultured, 78-5471
Liver, Rat, 78-5471
Alkaline Phosphatase
Fibroblasts, 78-5798
BCG
Macrophages, 78-5997
Benz(a)anthracene, 7,12-Dimethyl-
Embryo Cells, 78-5550

Cell Transformation, Neoplastic (cont'd)

- Mucopolysaccharidosis, 78-5540
- Nucleic Acids, Binding, 78-5550
- 1-Butanamine, *N*-Butyl-*N*-nitroso-
 - Bladder, Mouse, 78-5584
- Cell Differentiation
 - Cells, Cultured, 78-5900
- Cells, Cultured
 - Cell Density, 78-5997
 - Macrophages, 78-5997
- Cholanthrene, 3-Methyl-
 - Carcinoembryonic Antigen, 78-5740
 - Lung Explant, Mouse, 78-5529
 - Transplantation, Heterologous, 78-5529
- Collagen
 - Isolation and Characterization, 78-5783
- Cytoplasm
 - Hybrid Cells, 78-5807
- Dibutyl Cyclic AMP
 - Collagen, 78-5783
- Growth Factors
 - DNA Replication, 78-5808
 - Iron, Metabolism, 78-5808
 - Siderophore, 78-5808
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Cell Aggregation, 78-5507
 - DNA Replication, 78-5685
 - Fluorescence, 78-5685
 - Uridine, 5-Bromo-2'-deoxy-, 78-5685
- Hereditary Diseases
 - DNA Repair, Review, 78-5451
- Propanal, 2-Oxo-
 - Electron Acceptors, 78-5998
 - Proteins, 78-5998
- Quinoline, 4-Nitro-, 1-Oxide
 - Cells, Cultured, 78-5671
- Radiation, Ionizing
 - Chromosome Abnormalities, 78-5730
 - Fibroblasts, 78-5730
 - Glucosephosphate Dehydrogenase, 78-5730
- Sarcoma
 - Virus, Adeno 12, 78-5847
- Sarcoma, Osteogenic
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5507
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5507
- Urea, Ethyl Nitroso-
 - DNA, Alkylation, 78-5648
- Virus, Adeno 2
 - DNA, Viral, 78-5838
 - RNA, Viral, 78-5839
 - Ultraviolet Rays, 78-5838
 - Viral Proteins, 78-5838
- Virus, Adeno 5
 - Glycoproteins, 78-5844
 - Surface Properties, 78-5844
- Virus, Adeno 12
 - DNA, Viral, 78-5846
 - Hexokinase, 78-5847
 - Lactate Dehydrogenase Isoenzymes, 78-5847
 - Malate Dehydrogenase, 78-5847
 - Temperature Sensitive Mutants, 78-5846
- Virus, Avian Sarcoma
 - Antigenic Determinants, 78-5739
 - Antigens, Viral, 78-5740
 - Carcinoembryonic Antigen, 78-5740
 - Protein Kinase, 78-5737
- Virus, Bovine Leukemia

Cell Transformation, Neoplastic (cont'd)

- DNA, Viral, 78-5788
- Virus, Cytomegalo
 - Cell Culture Contamination, 78-5823
- Virus, Epstein-Barr
 - Antigens, 78-5833
- Virus, Feline Leukemia
 - Antigens, Viral, 78-5786
 - Viral Proteins, 78-5786
- Virus, Feline Sarcoma
 - Viral Proteins, 78-5786
- Virus, Harvey Murine Sarcoma
 - DNA, Viral, 78-5784
 - Virus, Helper, 78-5784
- Virus, Herpes
 - Cells, Cultured, 78-5823
- Virus, Herpes Simplex 2
 - Temperature Sensitive Mutants, 78-5821
 - Thymidine Kinase Phenotype, 78-5821
- Virus, Kirsten Murine Sarcoma
 - Cell Aggregation, 78-5507
 - Collagen, 78-5783
- Virus, Murine Leukemia
 - Glycoproteins, 78-5781
- Virus, Murine Mammary Tumor
 - Nucleic Acids, 78-5753
- Virus, Polyoma, BK
 - Neuroglia, 78-5794
- Virus, Rauscher Murine Leukemia
 - Antigen-Antibody Complex, 78-5782
 - Antigens, 78-5782
- Virus, Rous Sarcoma
 - Antigenic Determinants, 78-5744
 - Temperature Sensitive Mutants, 78-5744
- Virus, SV40, 78-5809
- Virus, SV40
 - Alkaline Phosphatase, 78-5798
 - Antigens, Viral, 78-5801
 - Chromatids, 78-5804
 - Colony Formation, 78-5799
 - DNA-RNA Hybridization, 78-5806
 - Epidermal Growth Factors, 78-5813
 - Fibroblasts, 78-5648
 - Growth Factors, 78-5808
 - Growth Factors, Review, 78-5452
 - Hybrid Cells, 78-5807
 - Prostate, Mouse, 78-5864

Cells, Cultured

- Aflatoxin B1
 - Cell Transformation, Neoplastic, 78-5471
 - Karyotyping, 78-5471
 - Prostaglandins, 78-5509
 - Tumorigenicity, 78-5471
- Benz(a)anthracene, 7,12-Dimethyl-
 - Mutagenic Activity, 78-5518
 - Nucleic Acids, Binding, 78-5550
 - Prostaglandins, 78-5509
 - Proteins, Binding, 78-5550
- Benzo(a)pyrene
 - Prostaglandins, 78-5509
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Carcinogenic Metabolite, 78-5552
- Benzo(a)pyrene, 9,10-Dihydro-9,10-dihydroxy-
 - Carcinogenic Metabolite, 78-5552
- Benzo(b)triphenylene
 - Nucleic Acids, Binding, 78-5550
 - Proteins, Binding, 78-5550

Cells, Cultured (cont'd)

- Brain Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5507
- Breast Neoplasms
 - Adenocarcinoma, 78-5950
 - Growth, 78-5950
 - Multicellular Spheroid Formation, 78-5996
- Burkitt's Lymphoma
 - Cortisone, 78-5824
- Cell Transformation, Neoplastic
 - Cell Density, 78-5997
 - Cell Differentiation, 78-5900
 - Macrophages, 78-5997
- Cholanthrene, 3-Methyl-
 - Prostaglandins, 78-5509
- Dibenz(a,h)anthracene
 - Nucleic Acids, Binding, 78-5550
 - Proteins, Binding, 78-5550
- Diethylamine, *N*-Nitroso-
 - Metabolism, Colonic Mucosa, 78-5590
- Dimethylamine, *N*-Nitroso-
 - Metabolism, Colonic Mucosa, 78-5590
- Fibroblasts
 - Aging, 78-5977
 - Caseinolysis, 78-5977
- Hepatoma
 - Aflatoxin B1, 78-5471
- Leukemia
 - Virus, Epstein-Barr, 78-5831
 - Virus, RNA Tumor, 78-5831
- Leukemia, Myeloblastic
 - Antigen-Antibody Reactions, 78-5865
 - Cell Differentiation, 78-5900
- Leukemia, Myelocytic
 - Cell Differentiation, 78-5900, 78-5985
 - Embryo, 78-5985
- Liver Neoplasms
 - Lymphosarcoma, 78-5979
- Lung Neoplasms
 - Lymphosarcoma, 78-5979
- Lymphosarcoma
 - Neoplasm Invasiveness, 78-5979
 - Virus, Abelson Murine Leukemia, 78-5979
- Mammary Neoplasms, Experimental
 - Multicellular Spheroid Formation, 78-5996
 - Ultrastructural Study, 78-5752
- Methanesulfonic Acid, Ethyl Ester
 - Mutagenic Activity, 78-5660
- Methanesulfonic Acid, Methyl Ester
 - Mutagenic Activity, 78-5660
- Nicotine, 1'-Nitroso-1'-demethyl-
 - Metabolism, Colonic Mucosa, 78-5590
- Oncogenic Viruses
 - Isolation and Characterization, Review, 78-5442
- Peptides
 - Prostaglandins, 78-5509
- Phorbol 12,13-Didecanoate
 - Lipids, Deacylation, 78-5509
 - Prostaglandins, 78-5509
- Piperidine, 1-Nitroso-
 - Metabolism, Colonic Mucosa, 78-5590
- Polycyclic Hydrocarbons
 - Azaguanine Resistance, 78-5518
 - Mutagenic Activity, 78-5518
 - Ouabain Resistance, 78-5518
 - Temperature Resistance, 78-5518
- Prostaglandins

Cells, Cultured (cont'd)

- Kidney, Dog, 78-5509
- Prostatic Neoplasms
 - Virus, SV40, 78-5864
- Psoralen, 8-Methoxy-
 - Mutagenic Activity, 78-5702
- Pyrrolidine, 1-Nitroso-
 - Metabolism, Colonic Mucosa, 78-5590
- Quinoline, 4-Nitro-, 1-Oxide
 - Cell Transformation, Neoplastic, 78-5671
- Sarcoma
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5507
 - Radiation, Ionizing, 78-5730
- Sarcoma, Mast Cell
 - Actin, 78-5863
- Sarcoma, Osteogenic
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5507
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Lipids, Deacylation, 78-5509
 - Prostaglandins, 78-5509
- Tracheal Neoplasms
 - Cell Differentiation, 78-5978
 - Transplantation, Homologous, 78-5978
- Ultraviolet Rays
 - Mutagenic Activity, 78-5702
- Urea, Ethyl Nitroso-
 - Mutagenic Activity, 78-5660
- Urea, Methyl Nitroso-
 - Mutagenic Activity, 78-5660
- Virus, Adeno 5
 - Surface Properties, 78-5844
 - Tumorigenicity, 78-5844
- Virus, C-Type RNA Tumor
 - Antigenic Determinants, Viral Complex, 78-5773
- Virus, Epstein-Barr
 - Virus Replication, 78-5831
- Virus, Gross Murine Leukemia
 - Antigens, Viral, 78-5769
 - Virus-Like Particles, 78-5769
- Virus, Herpes
 - Cell Transformation, Neoplastic, 78-5823
- Virus, Murine Mammary Tumor
 - Virus Replication, 78-5752
- Virus, Rhino
 - Isolation and Characterization, 78-5823

Cellular Inclusions

- Pituitary Neoplasms
 - Ultrastructural Study, 78-5913

Cervix Neoplasms

- Adenocarcinoma
 - Epidemiology, Israel, 78-5958
 - Ethnic Groups, 78-5958
- Age Factors
 - Epidemiology, England, 78-5955
 - Epidemiology, Great Britain, 78-5956
- Antigens
 - Isolation and Characterization, 78-5868
- Carcinoma, Epidermoid
 - Epidemiology, Israel, 78-5958
 - Ethnic Groups, 78-5958
- Carcinoma In Situ
 - Drug Therapy, 78-5888
 - Epidemiology, England, 78-5955
 - Epidemiology, Review, 78-5461
 - Radiotherapy, 78-5888
- Cholanthrene, 3-Methyl-

Cervix Neoplasms (cont'd)

- Adenosine Cyclic 3',5' Monophosphate, 78-5868
- Antigens, 78-5868
- Protein Kinase, 78-5868

Contraceptive Devices

- Copper, 78-5959
- Precancerous Conditions, 78-5959

Eosinophils

- Neoplasm Invasiveness, 78-5889

Ethnic Groups

- Epidemiology, Israel, 78-5958

Hysterectomy

- Epidemiology, 78-5957

Immunosuppression

- Case Report, 78-5888

Neoplasm Invasiveness

- Ultrastructural Study, 78-5889

Neoplasms, Multiple Primary

- Case Report, 78-5888

Precancerous Conditions

- Epidemiology, Review, 78-5461

Socioeconomic Factors

- Epidemiology, 78-5957

Virus, Herpes Simplex 2

- Antibodies, Viral, 78-5822
- Epidemiology, 78-5822

Chemodectoma

- see Paraganglioma, Nonchromaffin

Chlorambucil

- see Butyric Acid, 4-(p-Bis(2-chloroethyl)aminophenyl)-

Chlorine**Food Contamination**

- Carcinogenic Potential, Review, 78-5403

Neoplasms, Experimental

- Carcinogenic Potential, Review, 78-5403

Water Supply

- Carcinogenic Potential, Review, 78-5403

5 β -Cholan-24-oic Acid, 3 α ,12 α -Dihydroxy-, Monosodium Salt

- Virus, SV40
- DNA, Viral, 78-5802

5 β -Cholan-24-oic Acid, 3 α -Hydroxy-**Benzo(a)pyrene**

- DNA, Binding, 78-5562

Colon

- Carcinogenic Metabolite, 78-5562

Intestinal Neoplasms

- Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5684

Cholangioma**Diethylamine, N-Nitroso-**

- Carcinogenic Activity, Cat, Chicken, 78-5573

Dimethylamine, N-Nitroso-

- Opisthorchis viverrini*, 78-5602

- Trematoda, 78-5602

Liver Neoplasms

- Diethylamine, N-Nitroso-, 78-5575
- Thorium Dioxide, 78-5973

Cholanthren-1-ol, 3-Methyl-**Skin Neoplasms**

- Carcinogenic Activity, Mouse, 78-5523

Cholanthren-2-ol, 3-Methyl-**Skin Neoplasms**

- Carcinogenic Activity, Mouse, 78-5523

Cholanthren-1-one, 3-Methyl-**Skin Neoplasms**

- Carcinogenic Activity, Mouse, 78-5523

Cholanthren-2-one, 3-Methyl-**Skin Neoplasms**

- Carcinogenic Activity, Mouse, 78-5523

Cholanthrene, 1,2-Dehydro-3-Methyl-**Carcinoma, Epidermoid**

- Neoplasm Metastasis, 78-5523

Skin Neoplasms

- Carcinogenic Activity, Mouse, 78-5523

Cholanthrene, 7,8-Dihydro-7,8-dihydroxy-3-methyl-**Microsomes, Liver**

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 9,10-Dihydro-9,10-dihydroxy-3-methyl-**Microsomes, Liver**

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 11,12-Dihydro-11,12-dihydroxy-3-methyl-**Microsomes, Liver**

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 1,2-Dihydroxy-3-methyl-**Microsomes, Liver**

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 3-Methyl-**Acetamide, N-Fluoren-2-yl-**

- Hydroxylation, 78-5478

- RNA, Binding, 78-5479

Acetic Acid, (Ethylenedinitrilo)tetra-

- Dihydrodiol Formation, 78-5530

Acetohydroxamic Acid, N-Fluoren-2-yl-

- Hydroxylation, 78-5478

Aflatoxin B1

- Aryl Hydrocarbon Hydroxylase, 78-5510

- Carcinogenic Metabolite, 78-5510

Amino Acids

- Transplantation Immunology, 78-5525

Amyloidosis

- BCG, 78-5532

- Thymectomy, 78-5532

Ascorbic Acid

- Dihydrodiol Formation, 78-5530

Benzene, 4-Allyl-1,2-(methylenedioxy)-

- Carcinogenic Metabolite, 78-5519

Benzo(a)pyrene

- Aryl Hydrocarbon Hydroxylases, 78-5553

- DNA, Binding, 78-5553

Benzo(a)pyrene 4,5-Oxide

- NADH, NADPH Oxidoreductases, 78-5564

Bronchial Neoplasms

- Carcinoma, Epidermoid, 78-5522

- Transplantation, Autologous, 78-5522

Carcinoembryonic Antigen

- Cell Transformation, Neoplastic, 78-5740

Carcinoma

- Transplantation, Heterologous, 78-5529

Carcinosarcoma

- Transplantation, Heterologous, 78-5529

Cell Cycle Kinetics

- Skin, Mouse, 78-5526

Cell Transformation, Neoplastic

- Lung Explant, Mouse, 78-5529

- Transplantation, Heterologous, 78-5529

Cells, Cultured

- Prostaglandins, 78-5509

Cholanthrene, 3-Methyl- (cont'd)

- Cervix Neoplasms
 - Adenosine Cyclic 3',5' Monophosphate, 78-5868
 - Antigens, 78-5868
 - Protein Kinase, 78-5868

- Chromium
 - Co-carcinogenic Activity, 78-5528

- DNA
 - Basal Cells, 78-5526

- DNA, Binding
 - Skin, Mouse, 78-5504

- DNA Replication
 - Cell Cycle Kinetics, 78-5526
 - Skin, Mouse, 78-5526

- Ethylene, Chloro-
 - Cytochrome P-450, 78-5490

- Fibrosarcoma
 - Antigenic Determinants, 78-5866
 - Carcinoembryonic Antigen, 78-5866
 - Corynebacterium parvum*, 78-5520
 - Histological Study, Rabbit, 78-5506

- Kallikrein-Trypsin Inhibitor
 - Immune Response, Rat, 78-5527

- Liver Neoplasms
 - Acetamide, N-Fluoren-2-yl-, 78-5516

- Lung Neoplasms
 - BCG, 78-5532
 - Carcinoma, Epidermoid, 78-5528
 - Chromium, 78-5528
 - Nickel, 78-5528
 - Precancerous Conditions, 78-5528
 - Thymectomy, 78-5532

- Microsomes, Liver
 - Carcinogenic Metabolite, 78-5530

- Neoplasms, Experimental
 - Alanine, Phenyl-, 78-5525
 - Isoleucine, 78-5525
 - Leucine, 78-5525
 - Polysaccharides, 78-5857
 - Purine-6-thiol, 78-5857
 - Tyrosine, 78-5525

- Nickel
 - Co-carcinogenic Activity, 78-5528

- Sarcoma
 - Cyclophosphamide, 78-5524
 - Fetal Globulins, 78-5531
 - Kallikrein-Trypsin Inhibitor, 78-5527
 - Radiation, Ionizing, 78-5524
 - Transplantation, Heterologous, 78-5529

- Sarcoma, Osteogenic
 - Histological Study, Rabbit, 78-5506

- Skin Neoplasms
 - Carcinogenic Activity, Mouse, 78-5523

- Sulfuric Acid, Iron Salt
 - Dihydrodiol Formation, 78-5530

- Tracheal Neoplasms
 - Adenocarcinoma, 78-5521
 - Carcinoma, 78-5521
 - Carcinoma, Epidermoid, 78-5521
 - Dose-Response Study, 78-5521
 - Fibrosarcoma, 78-5521
 - Neoplasm Transplantation, 78-5521
 - Papilloma, 78-5521

Cholera Toxin

- Leukemia
 - Binding Sites, 78-5987
- Leukemia, Myelocytic

Cholera Toxin (cont'd)

- Binding Sites, 78-5987
- Lymphocytes
 - Binding Sites, 78-5987
- Lymphoma
 - Binding Sites, 78-5987
- Plasmacytoma
 - Binding Sites, 78-5987

Cholestasis

- Contraceptives, Oral
- Epidemiology, 78-5431

Choline

- Aflatoxin B1
 - Cytochrome P-450, 78-5473
 - Oxidoreductases, 78-5473

Chondrosarcoma

- Quinoline, 4-Nitro-, 1-Oxide
 - Histological Study, Rabbit, 78-5506

Choriocarcinoma

- Brain Neoplasms
 - Neoplasm Metastasis, 78-5456
- Spinal Cord Neoplasms
 - Neoplasm Metastasis, 78-5456

Chromate

- Lung Neoplasms
 - Epidemiology, 78-5928
- Smoking
 - Co-carcinogenic Effect, 78-5928

Chromatids

- Acetamide, N-Fluoren-2-yl-
 - Hamster Ovary Cells, 78-5475
 - Microsomes, Liver, 78-5475
- Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-7-methyl-
 - Mutagenic Activity, Hamster Ovary Cells, 78-5511
- Benz(a)anthracene, 7-Methyl-
 - Mutagenic Activity, Hamster Ovary Cells, 78-5511
- Benzo(a)pyrene
 - Mutagenic Activity, Hamster Ovary Cells, 78-5511
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Mutagenic Activity, Hamster Ovary Cells, 78-5511
- Mitomycin C
 - Cell Cycle Kinetics, 78-5686, 78-5694
 - Dose-Response Study, Rabbit, 78-5687
 - Lymphocytes, 78-5686
 - Uridine, 5-Bromo-2'-deoxy-, 78-5686, 78-5687
- Quinoline, 4-Nitro-, 1-Oxide
 - Cell Cycle Kinetics, 78-5694
- Radiation, Ionizing
 - Cell Cycle Kinetics, 78-5694
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-5804
 - Chromosome Aberrations, 78-5804

Chromatin

- Urea, Methyl Nitroso-
 - DNA, Methylation, 78-5646
 - Liver, Rat, 78-5646
- Virus, SV40
 - Nucleoprotein Complex, *Xenopus laevis* Oocytes, 78-5805

Chromium

- Cholanthrene, 3-Methyl-
 - Co-carcinogenic Activity, 78-5528
- Lung Neoplasms

Chromium (cont'd)

- Cholanthrene, 3-Methyl-, 78-5528
- Epidemiology, 78-5929
- Occupational Hazard, 78-5929

Chromosome Aberrations

- Angiosarcoma
 - Ethylene, Chloro-, 78-5491
- Ataxia Telangiectasia
 - Radiation, Ionizing, 78-5697
- Californium
 - Dose-Response Study, 78-5725
 - Lymphocytes, 78-5725
 - Radioisotopes, 78-5725
- Cobalt Radioisotopes
 - Lymphocytes, 78-5725
- Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
 - Toxicity, Review, 78-5421
- Drugs
 - Bone Marrow Cells, Mouse, 78-5424
 - Micronucleus Test, 78-5428
- Ethylene, Chloro-
 - Lymphocytes, 78-5489
 - Occupational Hazard, 78-5489
 - Occupational Hazard, Review, 78-5436
- Ethylene, Chloro- Polymer
 - Bone Marrow Cells, 78-5488
 - Lymphocytes, 78-5488
 - Occupational Hazard, 78-5488
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Cytogenetics, Mouse, 78-5682
- Hematologic Diseases
 - Down's Syndrome, 78-5907
- Leukemia
 - Down's Syndrome, 78-5907
- Leukemia, Myelocytic
 - Blast Crises, 78-5908
 - Case Report, 78-5906, 78-5908
 - Chromosomes, Human, 6-12, 78-5906
 - Chromosomes, Human, 21-22, 78-5903, 78-5908
 - Trisomy 8, 78-5906
- Methanesulfonic Acid, Ethyl Ester
 - Drosophila melanogaster*, 78-5660
- Methanesulfonic Acid, Methyl Ester
 - Drosophila melanogaster*, 78-5660
- Mitomycin C
 - Cell Cycle Kinetics, 78-5694
 - Uridine, 5-Bromo-2'-deoxy-, 78-5686
- Psoralen, 8-Methoxy-
 - Caffeine, 78-5703
- Psoralen, 8-Methoxy-
 - Lymphocytes, 78-5703
- Quinoline, 4-Nitro-, 1-Oxide
 - Cell Cycle Kinetics, 78-5694
- Radiation, Ionizing
 - Cell Cycle Kinetics, 78-5691, 78-5692, 78-5694
 - Lymphocytes, 78-5691, 78-5692
 - Uridine, 2'-Deoxy-5-fluoro-, 78-5692
- Radium
 - Lymphocytes, 78-5725
 - Radioisotopes, 78-5725
- Styrene
 - Occupational Hazard, Review, 78-5436
- Thorium Dioxide
 - Case Report, 78-5717
 - Chromosomes, Human, 1-3, 78-5717
 - Chromosomes, Human, 21-22, 78-5717
 - T-Lymphocytes, 78-5717

Chromosome Aberrations (cont'd)

- Ultraviolet Rays
 - Caffeine, 78-5703
 - Lymphocytes, 78-5703
 - Psoralen, 8-Methoxy-, 78-5703
 - Psoralen, 4,5',8-Trimethyl-, 78-5703
- Urea, Ethyl Nitroso-
 - Drosophila melanogaster*, 78-5660
- Urea, Methyl Nitroso-
 - Drosophila melanogaster*, 78-5660
- Vaginal Neoplasms
 - Carcinoma, 78-5871
 - Case Report, 78-5871
 - Chromosomes, Human, 1-3, 78-5871
- Virus, SV40
 - Chromatids, 78-5804

Chromosome Abnormalities

- Bladder Neoplasms
 - Carcinoma, 78-5893
 - Chromosomes, Human, 1-3, 78-5893
 - Heterochromatin, 78-5893
- Carcinogen, Environmental
 - Genetics, Review, 78-5458
- Colonic Neoplasms
 - Genetics, 78-5881
 - Polyps, 78-5881
 - Precancerous Conditions, 78-5881
- Leukemia
 - Megakaryocytes, 78-5901
 - Trisomy 15, AKR Mice, 78-5909
- Lysergic Acid Diethylamide
 - Lymphocytes, Review, 78-5422
- Quinoline, 4-Nitro-, 1-Oxide
 - Lymphocytes, 78-5730
- Radiation, Ionizing
 - Cell Transformation, Neoplastic, 78-5730
 - Dose-Response Study, 78-5730

Chromosomes

- Adriamycin
 - DNA Replication, 78-5673
- Benzoic Acid, Sodium Salt
 - Mutagenic Activity, 78-5635
- Bladder Neoplasms
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5581
- Carcinogen, Chemical
 - Mutagenic Activity, Review, 78-5418
- Carcinogen, Environmental
 - Mutagenic Activity, Review, 78-5455
- Carcinoma
 - Tissue Culture, Review, 78-5454
- Ethylenimine
 - DNA Replication, 78-5673
- Melanoma
 - Tissue Culture, Review, 78-5454
- Quinoline, 4-Nitro-, 1-Oxide
 - DNA Replication, 78-5673
- Sarcoma
 - Tissue Culture, Review, 78-5454
- Virus, SV40
 - DNA, Viral, 78-5797, 78-5811
 - Histones, 78-5811
 - Isolation and Characterization, 78-5811
 - Ultrastructural Study, 78-5797
 - Viral Proteins, 78-5811
 - Virus Replication, 78-5797

Chromosomes, Human, 1-3

- Bladder Neoplasms
- Chromosome Abnormalities, 78-5893
- Thorium Dioxide
- Chromosome Aberrations, 78-5717
- Vaginal Neoplasms
- Chromosome Aberrations, 78-5871

Chromosomes, Human, 6-12

- Leukemia, Myelocytic
- Chromosome Aberrations, 78-5906

Chromosomes, Human, 13-15

- Retinoblastoma
- Genetics, 78-5870

Chromosomes, Human, 21-22

- Leukemia, Myelocytic
- Chromosome Aberrations, 78-5903, 78-5908
- Gangliosides, 78-5987
- Genetics, 78-5903
- Kidney Transplantation, 78-5895
- Polycythemia Vera, 78-5905
- Transplantation, Heterologous, 78-5904, 78-5986
- Thorium Dioxide
- Chromosome Aberrations, 78-5717

Carcinoma

- Carcinoma, Basal Cell
- Burns, 78-5970
- Carcinoma, Epidermoid
- Burns, 78-5970
- Neoplasms, Multiple Primary, 78-5970
- Fibrosarcoma
- Burns, 78-5970
- Neoplasms, Connective Tissue
- Fibrosarcoma, 78-5970
- Skin Neoplasms
- Carcinoma, Epidermoid, 78-5970

Citric Acid, Plutonium Salt

- Acetic Acid,
- ((Carboxymethylimino)bis(ethylenenitrilo))tetra-
- Body Burden, Dog, 78-5731
- Drug Therapy, Cadmium/Zinc Chelates, 78-5731
- Plutonium
- Body Burden, Dog, 78-5732

Clofibrate

- see Propionic Acid, 2-(p-Chlorophenoxy)-2-methyl-, Ethyl Ester

Coal Tar

- Melanoma
- Psoriasis, 78-5873

Cobalt Chloride

- Escherichia coli*
- Tryptophan Reversion, 78-5676
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
- Mutagenic Activity, 78-5676

Cobalt Radioisotopes

- Chromosome Aberrations
- Lymphocytes, 78-5725

Colchicine

- Ames Test
- Evaluation, Review, 78-5427

Coliphages

- Antineoplastic Agents

Coliphages (cont'd)

- Escherichia coli*, 78-5515
- Mycotoxins
- Escherichia coli*, 78-5515
- Nitroso Compounds
- Escherichia coli*, 78-5515
- Polycyclic Hydrocarbons
- Escherichia coli*, 78-5515

Colitis, Ulcerative

- Colonic Neoplasms
- Precancerous Conditions, 78-5884
- Precancerous Conditions
- Age Factors, 78-5884
- Surgery, Operative, 78-5884

Collagen

- Cell Transformation, Neoplastic
- Isolation and Characterization, 78-5783
- Dibutyryl Cyclic AMP
- Cell Transformation, Neoplastic, 78-5783
- Estradiol
- Vagina, Neonatal Mouse, 78-5890
- Virus, Kirsten Murine Sarcoma
- Cell Transformation, Neoplastic, 78-5783
- Dibutyryl Cyclic AMP, 78-5783
- Revertant Cells, 78-5783

Colon

- Aflatoxin B1
- DNA, Binding, 78-5508
- Metabolism, Rat, 78-5508
- Benz(a)anthracene, 7,12-Dimethyl-
- DNA, Binding, 78-5508
- Metabolism, Rat, 78-5508
- Benzo(a)pyrene
- Carcinogenic Metabolite, 78-5562
- DNA, Binding, 78-5508
- Metabolism, Rat, 78-5508
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
- Carcinogenic Metabolite, 78-5562
- Benzo(a)pyrene, 9,10-Dihydro-9,10-dihydroxy-
- Carcinogenic Metabolite, 78-5562
- Benzo(a)pyrene, 7,8-Dihydro-7,8,9-trihydroxy-
- Carcinogenic Metabolite, 78-5562
- Benzo(a)pyrene, 7,8,9,10-Tetrahydro-7,8,9,10-tetrahydroxy-
- Carcinogenic Metabolite, 78-5562
- 5 β -Cholan-24-oic Acid, 3 α -Hydroxy-
- Carcinogenic Metabolite, 78-5562
- Hydrazine, 1,2-Dimethyl-
- DNA, Binding, 78-5508
- Metabolism, Rat, 78-5508

Colonic Neoplasms

- Adenocarcinoma
- Hydrazine, 1,2-Dimethyl-, 78-5496, 78-5497
- 78-5498
- Virus, Cytomegalo, 78-5885
- Adenoma
- Carrageen, 78-5644
- Hydrazine, 1,2-Dimethyl-, 78-5497, 78-5498
- Benzo(a)pyrene
- Carcinogenic Metabolite, 78-5562
- Carcinogen, Environmental
- Epidemiology, Review, 78-5463
- Carcinoma
- Radiation, Ionizing, 78-5710
- Carcinoma, Epidermoid

- Colonic Neoplasms (cont'd)**
 Hydrazine, 1,2-Dimethyl-, 78-5498
 Carcinoma, Mucinous
 Hydrazine, 1,2-Dimethyl-, 78-5497
 Chromosome Abnormalities
 Precancerous Conditions, 78-5881
 Colitis, Ulcerative
 Precancerous Conditions, 78-5884
 Corynebacterium parvum
 Neoplasm Metastasis, 78-5861
 Dietary Fats
 Epidemiology, Review, 78-5404
 Genetics
 Chromosome Abnormalities, 78-5881
 Classification, Review, 78-5455
 Hydrazine, 1,1-Dimethyl-
 Corynebacterium parvum, 78-5861
 Neoplasm Metastasis, 78-5861
 Hydrazine, 1,2-Dimethyl-
 Cell Cycle Kinetics, 78-5496
 DNA Replication, 78-5498
 Histological Study, Rat, 78-5497
 Neoplasm Metastasis, 78-5497
 Precancerous Conditions, 78-5497, 78-5498
 Ultrastructural Study, Rat, 78-5496
 Lymphoma
 Case Report, 78-5882
 Genetics, 78-5882
 Neoplasms, Multiple Primary, 78-5882
 Methane, Azoxy-
 Carrageen, 78-5644
 Nitrosamines
 DNA, Binding, 78-5590
 Polyps
 Chromosome Abnormalities, 78-5881
 Radiotherapy
 Case Report, 78-5710
 Urea, Methyl Nitroso-
 Carrageen, 78-5644
 Virus, Cytomegalo
 DNA-RNA Hybridization, 78-5885
 DNA, Viral, 78-5885
- Concanavalin A**
 Virus, Adeno 2
 Glycoproteins, 78-5840
 Virus, Avian Myeloblastosis
 Antigens, Viral, 78-5741
- Contraceptive Devices**
 Breast Neoplasms
 Epidemiology, 78-5947
 Cervix Neoplasms
 Copper, 78-5959
 Precancerous Conditions, 78-5959
 Copper
 Carcinogenic Potential, 78-5959
- Contraceptives, Oral**
 Cholestasis
 Epidemiology, 78-5431
 Hepatitis
 Epidemiology, 78-5431
 Hepatoma
 Case Report, 78-5874
 Diagnosis, 78-5874
 Epidemiology, 78-5431
 Ovarian Neoplasms
 Epidemiology, 78-5952
- Copper**
 Cervix Neoplasms
 Contraceptive Devices, 78-5959
 Contraceptive Devices
 Carcinogenic Potential, 78-5959
- Coriolus versicolor***
 Neoplasms, Experimental
 Polysaccharides, 78-5857
- Corticosterone**
 Hepatoma
 Diethylamine, *N*-Nitroso-, 78-5576
 Tyrosine Aminotransferase, 78-5576
 Tyrosine Aminotransferase
 Liver, Rat, 78-5576
- Corticotropin**
 Diethylamine, *N*-Nitroso-
 Metabolism, Rat, 78-5576
- Cortisol**
 Phorbol 12,13-Didecanoate
 Prostaglandins, 78-5509
 Stress
 Aryl Hydrocarbon Hydroxylases, 78-5560
 Plasma Levels, 78-5560
 12-*O*-Tetradecanoylphorbol-13-acetate
 Prostaglandins, 78-5509
- Cortisone**
 Burkitt's Lymphoma
 Cell Survival, 78-5824
 Cells, Cultured, 78-5824
 Interferon, 78-5824
 Virus, Epstein-Barr, 78-5824
 Infectious Mononucleosis
 Interferon, 78-5824
 Virus, Epstein-Barr, 78-5824
 Virus, Epstein-Barr
 Antigens, Viral, 78-5824
- Corynebacterium parvum***
 Colonic Neoplasms
 Hydrazine, 1,1-Dimethyl-, 78-5861
 Neoplasm Metastasis, 78-5861
 Ear Neoplasms
 Hydrazine, 1,1-Dimethyl-, 78-5861
 Fibrosarcoma
 Cholanthrene, 3-Methyl-, 78-5520
 Macrophage Activation, 78-5520
- Coumarin**
 Acetic Acid, *o*-Hydroxyphenyl-
 Hepatocarcinogenicity, Review, 78-5433
 Dose-Response Study
 Hepatocarcinogenicity, Review, 78-5433
 Mammary Neoplasms, Experimental
 Immune Response, Review, 78-5433
 Ultraviolet Rays
 DNA Repair, Review, 78-5433
- Coumarin, 3-(α -Acetonylbenzyl)-4-hydroxy-**
 Drug Therapy
 Immune Response, Review, 78-5433
 Metabolism, Liver
 Species Difference, Review, 78-5433
- Coumarin, 3,3'-Methylenebis(4-hydroxy)-**
 Drug Therapy
 Immune Response, Review, 78-5433

opharyngioma

Child
Epidemiology, Denmark, 78-5925

sol, 2,6-Di-tert-butyl-

Diethylamine, *N*-Nitroso-
Carcinogenic Activity, Mouse, 78-5665

Dimethylamine, *N*-Nitroso-

Carcinogenic Activity, Mouse, 78-5665

Leukemia

Diethylamine, *N*-Nitroso-, 78-5665
Dimethylamine, *N*-Nitroso-, 78-5665
Methanesulfonic Acid, Ethyl Ester, 78-5665
Methanesulfonic Acid, Methyl Ester, 78-5665
Radiation, Ionizing, 78-5665

Lung Neoplasms

Diethylamine, *N*-Nitroso-, 78-5665
Radiation, Ionizing, 78-5665
Methanesulfonic Acid, Ethyl Ester
Carcinogenic Activity, Mouse, 78-5665
Methanesulfonic Acid, Methyl Ester
Carcinogenic Activity, Mouse, 78-5665
Radiation, Ionizing
Carcinogenic Activity, Mouse, 78-5665

sol, *p*-Nitroso-

Pyrrolidine, 1-Nitroso-
Catalysis, 78-5630

on Oil

Granuloma
Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5677

amide, Methylnitroso-

Escherichia coli
Mutagenic Activity, 78-5588
Tryptophan Reversion, 78-5588

ohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer

Hepatoma
Antipain, 78-5583
Leupeptins, 78-5583
Pepstatins, 78-5583
Precancerous Conditions, 78-5583

clohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-

Saccharomyces cerevisiae
Aspergillus nidulans, 78-5653

lobeximide

Erythroleukemia
RNA, Messenger, 78-5764
Phorbol 12,13-Didecanoate
Prostaglandins, 78-5509
12-*O*-Tetradecanoylphorbol-13-acetate
Prostaglandins, 78-5509
Virus, Adeno 5
Deoxyribonuclease, 78-5845
Virus, Epstein-Barr
Antigens, Viral, 78-5828

lophosphamide

Aroclor 1254
Ames Test, 78-5517
Barbituric Acid, 5-Ethyl-5-phenyl-
Ames Test, 78-5517
5,6-Benzoflavone
Ames Test, 78-5517
Bladder Neoplasms
Carcinogenic Effect, Review, 78-5420
Carcinoma, 78-5894

Cyclophosphamide (cont'd)

Case Report, 78-5894
Leukemia
Hodgkin's Disease, 78-5898
Leukemia, Myelocytic
Hodgkin's Disease, 78-5898
Sarcoma
Cholanthrene, 3-Methyl-, 78-5524
Tumor Latency, 78-5524

Cystadenocarcinoma

Ovarian Neoplasms
Age Factors, 78-5952
Precancerous Conditions, 78-5952

Cysteine, *N*-Nitroso-, Hydrochloride

Aniline, *N*-Methyl-4-nitroso-
Catalysis, 78-5630

Cystitis

Bladder Diseases
Nitrosamines, 78-5591
1,4-Butanediol, Dimethylsulfonate
Case Report, 78-5897
Leukemia, Myelocytic
1,4-Butanediol, Dimethylsulfonate, 78-5897
Hemorrhage, 78-5897

Cysts

Breast Diseases
Proteins, 78-5993

Cytochalasin B

Virus, Rauscher Murine Leukemia
Viral Proteins, 78-5780

Cytochrome Oxidase

Aflatoxin B2
Mitochondria, Liver, 78-5469
Aflatoxin G2
Mitochondria, Liver, 78-5469

Cytochrome P-450

Aflatoxin B1
Carcinogenic Metabolite, 78-5510
Choline, 78-5473
Folic Acid, 78-5473
Methionine, 78-5473
Aflatoxin B1 2,3-Oxide
Metabolism, Mouse, 78-5510
Aflatoxin M1
Metabolism, Mouse, 78-5510
Benzo(a)pyrene 4,5-Oxide
NADH, NADPH Oxidoreductases, 78-5564
Ethylene, Chloro-
Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5490
Cholanthrene, 3-Methyl-, 78-5490
Metabolism, 78-5490

Cytochromes

Acetamide, *N*-Fluoren-2-yl-
Metabolism, Review, 78-5407

Cytoplasm

Cell Transformation, Neoplastic
Hybrid Cells, 78-5807

Cytosine, 1- β -*D*-Arabinofuranosyl-

Ultraviolet Rays
DNA Repair, 78-5477

Deoxyribonuclease

- Virus, Adeno 5
 - Cycloheximide, 78-5845
 - DNA Nucleotidyltransferases, 78-5845
 - DNA, Single-Stranded, 78-5845
 - DNA, Viral, 78-5845
 - Virus Replication, 78-5845

Dexamethasone

- DNA, Binding
 - Liver, Rat, 78-5576
- Hepatoma
 - Diethylamine, *N*-Nitroso-, 78-5576
 - Receptors, Hormone, 78-5576
 - Tyrosine Aminotransferase, 78-5576
- Mammary Neoplasms, Experimental
 - Surface Properties, 78-5752
- Receptors, Hormone
 - Liver, Rat, 78-5576
- Tyrosine Aminotransferase
 - Liver, Rat, 78-5576

Diazepam

- Liver Neoplasms
 - Diethylamine, *N*-Nitroso-, 78-5575
- Nose Neoplasms
 - Diethylamine, *N*-Nitroso-, 78-5575

1*H*-1,4-Diazepine, Hexahydro-1,4-dinitroso-

- Aroclor 1254
 - Ames Test, 78-5589
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Ames Test, 78-5589

Dibenz(a,c)anthracene

- see Benzo(b)triphenylene

Dibenz(a,h)anthracene

- Adenosine Triphosphatase
 - Muscle, Connective Tissue, Rat, 78-5503
- Aroclor 1254
 - Mutagenic Activity, 78-5548
- Calcium
 - Adenosine Triphosphatase, 78-5503
- Cells, Cultured
 - Nucleic Acids, Binding, 78-5550
 - Proteins, Binding, 78-5550
- Dibenz(a,h)anthracene, 3,4-Dihydro-3,4-dihydroxy-
 - Enzyme Activation, 78-5548
- Dibenz(a,h)anthracene, 5,6-Dihydro-5,6-dihydroxy-
 - Enzyme Activation, 78-5548
- Enzyme Activation
 - Carcinogenic Metabolite, 78-5548
- Microsomes, Liver
 - Mutagenic Activity, 78-5548
- Salmonella typhimurium*
 - Mutagenic Activity, 78-5548

Dibenz(a,h)anthracene, 3,4-Dihydro-3,4-dihydroxy-

- Dibenz(a,h)anthracene
 - Enzyme Activation, 78-5548

Dibenz(a,h)anthracene, 5,6-Dihydro-5,6-dihydroxy-

- Dibenz(a,h)anthracene
 - Enzyme Activation, 78-5548

Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-

- Acetic Acid, (2,4,5-Trichlorophenoxy)-

Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro- (cont'd)

- Toxic Contaminant, Review, 78-5421
- Air Pollution
 - Mutagenic Activity, Review, 78-5421
- Chromosome Aberrations
 - Toxicity, Review, 78-5421
- Occupational Hazard
 - Mutagenic Activity, Review, 78-5421
- Phenol, 2,2'-Methylenebis(3,4,6-trichloro)-
 - Toxic Contaminant, Review, 78-5421
- Phenol, Pentachloro-
 - Toxic Contaminant, Review, 78-5421

1,2:5,6-Dibenzoanthracene

- see Dibenz(a,h)anthracene

Dibutyl Cyclic AMP

- Collagen
 - Cell Transformation, Neoplastic, 78-5783
- Virus, Kirsten Murine Sarcoma
 - Collagen, 78-5783

Dieldrin

- Benzo(a)pyrene
 - Aryl Hydrocarbon Hydroxylases, 78-5553
 - DNA, Binding, 78-5553

Dienestrol

- Nervous System Neoplasms
 - Urea, Methyl Nitroso-, 78-5643

Diet

- Breast Neoplasms
 - Epidemiology, Hawaii, 78-5945
- Carcinogen, Environmental
 - Epidemiology, Review, 78-5402
- Esophageal Neoplasms
 - Zinc, 78-5587
- Gastrointestinal Neoplasms
 - Epidemiology, Poland, 78-5941
- Iodine Radioisotopes
 - Thyroid Gland Function, 78-5713
 - Thyroxine, 78-5713
- Laryngeal Neoplasms
 - Epidemiology, Georgian SSR, 78-5937
- Stomach Neoplasms
 - Epidemiology, 78-5942

Dietary Fats

- Aflatoxin B1
 - Glutathione, 78-5473
 - Hepatocarcinogenicity, Rat, 78-5473
 - Nucleic Acids, Binding, 78-5473
- Breast Neoplasms
 - Epidemiology, Japan, 78-5946
 - Epidemiology, Review, 78-5404
- Colonic Neoplasms
 - Epidemiology, Review, 78-5404
- Intestinal Neoplasms
 - Hydrazine, 1,2-Dimethyl-, 78-5684

Dietary Proteins

- Aflatoxin B1
 - Hepatocarcinogenicity, 78-5470
- Aflatoxin G1
 - Hepatocarcinogenicity, 78-5470
- Hepatoma
 - Aflatoxin B1, 78-5470
 - Aflatoxin G1, 78-5470
- Intestinal Neoplasms
 - Hydrazine, 1,2-Dimethyl-, 78-5684

thylamine, Bis(2,2'-diethoxy)-N-nitroso-
Drinking Water
Carcinogenic Activity, Rat, 78-5572

thylamine, 2,2-Dichloro-N-nitroso-
Ear Neoplasms
Adenocarcinoma, 78-5572
Stomach Neoplasms
Papilloma, 78-5572

thylamine, 2,2'-Diethoxy-N-nitroso-
Liver Neoplasms
Carcinogenic Activity, Rat, 78-5572

thylamine, 2,2'-Dimethoxy-N-nitroso-
Liver Neoplasms
Carcinogenic Activity, Rat, 78-5572

thylamine, N-Nitroso-
Acetamide, N-Fluoren-2-yl-
Co-carcinogenic Effect, 78-5516
Barbituric Acid, 5-Ethyl-5-phenyl-
Mutagenic Activity, Hamster V79 Cells, 78-5593
Cells, Cultured
Metabolism, Colonic Mucosa, 78-5590

Cholangioma
Carcinogenic Activity, Cat, Chicken, 78-5573
Corticotropin
Metabolism, Rat, 78-5576

p-Cresol, 2,6-Di-tert-butyl-
Carcinogenic Activity, Mouse, 78-5665
Derivatives

Carcinogenic Activity, Rat, 78-5572
Dimethylamine, N-Nitroso-
Purine, 2-Amino-6-methoxy-, 78-5601

Esophageal Neoplasms
Carcinogenic Activity, Rat, 78-5572

Ethylene, Chloro-
Permeability, Laboratory Gloves, 78-5595

Food Contamination
Concentration Levels, 78-5621
Degradation, 78-5618
Quantitation Method, 78-5616

2-Furaldehyde
Co-carcinogenic Effect, 78-5571

Hemangioma
Carcinogenic Activity, Cat, Chicken, 78-5573

Hepatoma
Carcinogenic Activity, Cat, Chicken, 78-5573
Corticosterone, 78-5576
Dexamethasone, 78-5576
Propionic Acid, 2-(p-Chlorophenoxy)-2-methyl-,
Ethyl Ester, 78-5574
Tyrosine Aminotransferase, 78-5576
WY-14,643, 78-5574

Kidney Neoplasms
Adenocarcinoma, 78-5573
Adenoma, 78-5573, 78-5574
Carcinogenic Activity, Cat, Chicken, 78-5573

Leukemia
p-Cresol, 2,6-Di-tert-butyl-, 78-5665

Liver Neoplasms
Carbon Tetrachloride, 78-5592
Carcinogenic Activity, Rat, 78-5572
Cholangioma, 78-5575
Diazepam, 78-5575
Hepatectomy, 78-5592
Precancerous Conditions, 78-5516
Ultrastructural Study, Gerbil, 78-5575

Diethylamine, N-Nitroso- (cont'd)

Uracil, 2-Thio-, 78-5575
WY-14,643, 78-5574

Lung Neoplasms
Adenoma, 78-5665
p-Cresol, 2,6-Di-tert-butyl-, 78-5665

Mutagenic Activity
Azaguanine Resistance, 78-5655
Lung, Hamster, 78-5655
Ouabain Resistance, 78-5655

Nitric Acid, Ammonium Salt
Kidney Calculi, 78-5615

Nitric Acid, Potassium Salt
Stomach, Cow, 78-5614

Nitric Acid, Sodium Salt
Food Contamination, 78-5618

Nose Neoplasms
Adenocarcinoma, 78-5575
Diazepam, 78-5575
Ultrastructural Study, Gerbil, 78-5575
Uracil, 2-Thio-, 78-5575

Occupational Hazard
Permeability, Laboratory Gloves, 78-5595, 78-5596
Permeability, Rubber Gloves, 78-5597

Pheochromocytoma
Carcinogenic Activity, Cat, Chicken, 78-5573

Respiratory Tract Neoplasms
2-Furaldehyde, 78-5571

Thymidine Kinase
Mutagenic Activity, 78-5666

Ultraviolet Rays
Photolysis Products, 78-5620

WY-14,643
Co-carcinogenic Effect, 78-5574

Diethylenetriaminepentaacetic Acid

see Acetic Acid, ((Carboxymethylimino)bis(ethylenetri-
trilo))tetra-

Digestive System Neoplasms

Carcinoma, Epidermoid
Quinoline, 8-Nitro-, 78-5672
Quinoline, 8-Nitro-
Dose-Response Study, Rat, 78-5672

Dimethylamine, N-Nitroso-

Barbituric Acid, 5-Ethyl-5-phenyl-
Mutagenic Activity, Hamster V79 Cells, 78-5593
Nucleic Acids, Binding, 78-5603

Bladder Neoplasms
Schistosoma haematobium, 78-5591
Urine, 78-5591

1-Butanamine, N-Butyl-N-Nitroso-
Purine, 2-Amino-6-methoxy-, 78-5601

Calcium
Adenosine Triphosphatase, 78-5503

Carbon Dioxide
Metabolism, Rat, 78-5603
Nucleic Acids, Binding, 78-5603

Carbon Tetrachloride
Metabolism, Rat, 78-5603
Nucleic Acids, Binding, 78-5603

Carcinogen, Environmental
Epidemiology, Review, 78-5414

Cells, Cultured
Metabolism, Colonic Mucosa, 78-5590

Cholangioma
Opisthorchis viverrini, 78-5602
Trematoda, 78-5602

Dimethylamine, N-Nitroso- (cont'd)

- p*-Cresol, 2,6-Di-*tert*-butyl-
 - Carcinogenic Activity, Mouse, 78-5665
- Diethylamine, *N*-Nitroso-
 - Purine, 2-Amino-6-methoxy-, 78-5601
- Dipropylamine, *N*-Nitroso-
 - Purine, 2-Amino-6-methoxy-, 78-5601
- DNA, Binding
 - Metabolism, Colonic Mucosa, 78-5590
- Ethylene, Chloro-
 - Permeability, Laboratory Gloves, 78-5595
- Food Contamination
 - Degradation, 78-5618
 - Quantitation Method, 78-5616, 78-5623
- Guanine, 7-Methyl-
 - DNA, Alkylation, 78-5599, 78-5600
 - DNA, Liver, 78-5607
 - Kidney, Liver, Rat, 78-5600
 - Liver, Rat, 78-5599
- Hypophysectomy
 - DNA, Liver, 78-5607
- Kidney Neoplasms
 - DNA Replication, 78-5604
 - Thyroidectomy, 78-5605
- Leukemia
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Liver
 - Metabolism, Rat, 78-5603
 - Nucleic Acids, Binding, 78-5603
- Liver Neoplasms
 - Hepatectomy, 78-5592
 - Rhodopseudomonas capsulata*, 78-5608
 - Thyroidectomy, 78-5605
- Lung Neoplasms
 - Adenoma, 78-5570
 - APUD Cells, 78-5570
 - Metaplasia, 78-5570
 - Precancerous Conditions, 78-5570
 - Thyroidectomy, 78-5605
 - Ultrastructural Study, Hamster, 78-5570
- Methanamine, *N,N*-Dimethyl-
 - Nitrous Acid, Sodium Salt, 78-5611
- Methanamine, *N,N*-Dimethyl-, *N*-Oxide
 - Nitrous Acid, Sodium Salt, 78-5611
- Minocycline
 - Ascorbic Acid, 78-5610
- Neurospora crassa*
 - Mutagenic Activity, 78-5675
- Nitric Acid, Potassium Salt
 - Stomach, Cow, 78-5614
- Nitric Acid, Sodium Salt
 - Food Contamination, 78-5618
- Nitrous Acid, Sodium Salt
 - Minocycline, 78-5610
 - Oxytetracycline, 78-5610
 - Sludge, 78-5608
 - Tetracycline, 78-5610
 - Vibramycin, 78-5610
 - Water Pollution, 78-5608
- Occupational Hazard
 - Permeability, Laboratory Gloves, 78-5595, 78-5596
 - Permeability, Rubber Gloves, 78-5597
- Phenothiazine, 10-(2-Dimethylamino)propyl-
 - Nitrosation Product, 78-5609
 - Nitrous Acid, Sodium Salt, 78-5609
- Purine, 2-Amino-6-methoxy-
 - DNA, Alkylation, 78-5600

Dimethylamine, N-Nitroso- (cont'd)

- DNA, Liver, 78-5601, 78-5607
- Kidney, Liver, Rat, 78-5600
- Selenium
 - Aminotransferases, 78-5606
 - Hepatotoxicity, Rat, 78-5606
- Somatotropin
 - DNA, Liver, 78-5607
- Soy Beans
 - Somatic Mosaicism, 78-5664
- Thymidine
 - DNA, Alkylation, 78-5599
- Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Ultraviolet Rays
 - Photolysis Products, 78-5620
- Valeric Acid, 2,2-Diphenyl-, 2-(Diethylamino)ethyl Ester HCl
 - Metabolism, Rat, 78-5603
 - Nucleic Acids, Binding, 78-5603
- Vibramycin
 - Ascorbic Acid, 78-5610
- Vitamin E
 - Aminotransferases, 78-5606
 - Antioxidant Levels, Liver, 78-5606
 - Hepatotoxicity, Rat, 78-5606

Diphenylamine

- Nitric Acid
 - Nitrosation, Review, 78-5414

Dipropylamine, N-Nitroso-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Mutagenic Activity, Hamster V79 Cells, 78-5593
- Dimethylamine, *N*-Nitroso-
 - Purine, 2-Amino-6-methoxy-, 78-5601
- Ethylene, Chloro-
 - Permeability, Laboratory Gloves, 78-5595
- Occupational Hazard
 - Permeability, Laboratory Gloves, 78-5595, 78-5596
 - Permeability, Rubber Gloves, 78-5597
- Thymidine Kinase
 - Mutagenic Activity, 78-5666

Disgerminoma

- Child
 - Epidemiology, Review, 78-5459

Distamycin A

- Urea, Methyl Nitroso-
 - DNA, Methylation, 78-5646

Disulfide, Bis(diethylthiocarbamoyl)-

- Benzo(a)pyrene
 - DNA, Binding, 78-5557

DNA

- Benzo(a)pyrene, 6-Oxy-
 - Nuclear Magnetic Resonance, 78-5565
- Cholanthrene, 3-Methyl-
 - Basal Cells, 78-5526
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Alkaline Sucrose Gradient Sedimentation, 78-5678
 - HeLa Cells, 78-5678
 - Immunoreactivity, Antinucleoside Antibodies, 78-5678
- Heat
 - Squamous Cells, 78-5706
- Methanesulfonic Acid, Ethyl Ester
 - Alkylation Products, 78-5639

(cont'd)

Radiation, Ionizing

- Alkaline Sucrose Gradient Sedimentation, 78-5678
- HeLa Cells, 78-5678
- Immunoreactivity, Antinucleoside Antibodies 78-5678

Ultraviolet Rays

- Nucleic Acid Denaturation, 78-5706
- Squamous Cells, 78-5706

Urea, Ethyl Nitroso-

- Alkylation Products, 78-5639

Urea, Methyl Nitroso-

- Alkylation Products, 78-5639

Bacterial

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Haemophilus influenzae*, 78-5679
- Temperature Sensitive Mutants, 78-5679

Nucleotidyltransferases

Virus, Adeno 5

- Deoxyribonuclease, 78-5845

Polymerase

Virus, Avian Myeloblastosis

- Benzoic Acid, 3,3'-Dithiobis(6-nitro)-, 78-5743
- Benzoic Acid, *p*-Hydroxymercuri-, 78-5743
- Benzoic Acid, 2-Nitro-5-thiocyano-, 78-5743
- Proteolysis, 78-5742
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-, 78-5743
- Viral Proteins, 78-5742

Repair

Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-

- Hepatocytes, Rat, 78-5477

Acetamide, *N*-Fluoren-2-yl-

- Hepatocytes, Rat, 78-5477

Acetohydroxamic Acid, *N*-Fluoren-2-yl-

- Hepatocytes, Rat, 78-5477

Ataxia Telangiectasia

- Lymphocytes, 78-5697, 78-5860
- Radiation, Ionizing, 78-5697, 78-5860

Butyrophene, γ -Chloro-*p*-fluoro-

- Escherichia coli*, 78-5449
- Salmonella typhimurium*, 78-5449

Carcinoma, Epidermoid

- Methanesulfonic Acid, Methyl Ester, 78-5667
- Quinoline, 4-Nitro-, 1-Oxide, 78-5667

Endonucleases

- Micrococcus luteus*, 78-5700

Guanidine, 1-Butyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Rat, 78-5683

Guanidine, 1-Hexyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Rat, 78-5683

Guanidine, 1-Isobutyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Rat, 78-5683

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Rat, 78-5683

Guanidine, 1-Pentyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Rat, 78-5683

Guanidine, 1-Propyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Rat, 78-5683

Methanesulfonic Acid, Ethyl Ester

- Escherichia coli*, 78-5651

Methanesulfonic Acid, Methyl Ester

- L Cells, 78-5667

Mitomycin C

- Mutagenic Activity, 78-5426

Mutagens

DNA Repair (cont'd)

Screening Test, Review, 78-5450

- 2-Propenoic Acid, 2-Methyl-, 1,2-Ethanediyl Ester
Escherichia coli, 78-5449

***Salmonella typhimurium*, 78-5449**

Quinoline, 4-(Hydroxyamino)-, 1-Oxide

- Structure-Activity Relationship, 78-5668

Quinoline, 4-Nitro-, 1-Oxide

- L Cells, 78-5667
- Structure-Activity Relationship, 78-5668

Radiation, Ionizing

- Age Factors, 78-5696
- Liver/Cerebellum, Mouse, 78-5696
- Methanesulfonic Acid, Ethyl Ester, 78-5651

Ultraviolet Rays

- Cytosine, 1- β -*D*-Arabinofuranosyl-, 78-5477
- Dose-Response Study, 78-5707
- Ethidium Bromide, 78-5477
- Fibroblasts, 78-5704, 78-5705, 78-5707, 78-5709
- Hepatocytes, Rat, 78-5477
- Photoreactivation, 78-5705
- Thymine Dimer Excision, 78-5709
- Xeroderma Pigmentosum, 78-5704

Urea, Methyl Nitroso-

- DNA, Single-Stranded, 78-5641
- Sodium Azide, 78-5641

Xeroderma Pigmentosum

- Fibroblasts, 78-5704

DNA Replication

Adriamycin

- Chromosomes, 78-5673
- HeLa Cells, 78-5673

1-Butanamine, *N*-Butyl-*N*-nitroso-

- Bladder, Mouse, 78-5584
- Cell Cycle Kinetics, 78-5584

Cell Transformation, Neoplastic

- Growth Factors, 78-5808

Cholanthrene, 3-Methyl-

- Cell Cycle Kinetics, 78-5526
- Skin, Mouse, 78-5526

Colonic Neoplasms

- Hydrazine, 1,2-Dimethyl-, 78-5498

Ethylenimine

- Chromosomes, 78-5673
- HeLa Cells, 78-5673

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Cell Transformation, Neoplastic, 78-5685
- Haemophilus influenzae*, 78-5679
- Uridine, 5-Bromo-2'-deoxy-, 78-5685

Kidney Neoplasms

- Dimethylamine, *N*-Nitroso-, 78-5604
- Precancerous Conditions, 78-5604

Leukemia, Myelocytic

- Transplantation, Heterologous, 78-5986

Mammary Neoplasms, Experimental

- Benz(a)anthracene, 7,12-Dimethyl-, 78-5537

Quinoline, 4-Nitro-, 1-Oxide

- Chromosomes, 78-5673
- HeLa Cells, 78-5673

Virus, Adeno 12

- DNA, Viral, 78-5848
- Temperature Sensitive Mutants, 78-5846

Virus, Epstein-Barr

- Antigens, Viral, 78-5828
- Interactions, B95-8/P3HR-1 Substrains, 78-5834
- Lymphocytes, 78-5834
- B-Lymphocytes, 78-5826

DNA Replication (cont'd)

- Staphylococcus aureus*, 78-5834
- Virus, Herpes Simplex 2
 - Temperature Sensitive Mutants, 78-5821
- Virus, SV40
 - Agarose Gel Electrophoresis, 78-5803
 - Deoxycytosine Nucleotides, 78-5803
 - DNA, Viral, 78-5803
 - Picolinic Acid, 78-5808
 - Temperature Sensitive Mutants, 78-5810

DNA, Single-Stranded

- Urea, Methyl Nitroso-
 - DNA Repair, 78-5641
 - Mutagenic Activity, Plants, 78-5641
- Virus, Adeno 5
 - Deoxyribonuclease, 78-5845

DNA, Viral

- Colonic Neoplasms
 - Virus, Cytomegalo, 78-5885
- Enteritis, Regional
 - Virus, Cytomegalo, 78-5885
- Hepatoma
 - Australia Antigen, 78-5837
- Intestinal Neoplasms
 - Virus, Cytomegalo, 78-5885
- Intestinal Polyps
 - Virus, Cytomegalo, 78-5885
- Lymphosarcoma
 - Virus, Bovine Leukemia, 78-5788
- Mammary Neoplasms, Experimental
 - Virus, Murine Mammary Tumor, 78-5751
- Virus, Adeno 2
 - Carrier Proteins, 78-5841
 - Cell Transformation, Neoplastic, 78-5838
 - DNA-DNA Hybridization, 78-5838
 - Viral Proteins, 78-5842
- Virus, Adeno 5
 - Deoxyribonuclease, 78-5845
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5846
 - DNA Replication, 78-5848
 - Initiation and Termination Sites, 78-5848
 - Oligonucleotides, 78-5848
- Virus, Bovine Leukemia
 - Cell Transformation, Neoplastic, 78-5788
 - DNA-DNA Hybridization, 78-5788
 - Tissue Distribution, 78-5788
- Virus, Harvey Murine Sarcoma
 - Cell Transformation, Neoplastic, 78-5784
 - Kinetics, Plaque and Focus Formation, 78-5784
 - Transformation, Genetic, 78-5784
- Virus, Herpes Simplex 1
 - Peptides, 78-5820
 - Phosphonoacetic Acid, 78-5820
 - Thymidine Kinase, 78-5820
 - Virus, Recombinant, 78-5820
- Virus, Herpes Simplex 2
 - Peptides, 78-5820
 - Phosphonoacetic Acid, 78-5820
 - Thymidine Kinase, 78-5820
 - Virus, Recombinant, 78-5820
- Virus, Murine Mammary Tumor
 - Virus Replication, 78-5751
- Virus, Polyoma
 - Nucleotide Sequence, 78-5792
- Virus, Polyoma, BK

DNA, Viral (cont'd)

- Cleavage Sites, 78-5795
- Endonucleases, 78-5795
- Virus, SV40
 - Antigens, Viral, 78-5801
 - 5 β -Cholan-24-oic Acid, 3 α ,12 α -Dihydroxy-, Monosodium Salt, 78-5802
 - Chromosomes, 78-5797, 78-5811
 - DNA Replication, 78-5803
 - Nucleoprotein Complex, *Xenopus laevis* Oocytes, 78-5805
 - Nucleotide Sequence, 78-5792
 - Superhelical Turns, 78-5802
 - Transformation, Genetic, 78-5796

Down's Syndrome

- Hematologic Diseases
 - Chromosome Aberrations, 78-5907
- Leukemia
 - Case Report, 78-5907
 - Chromosome Aberrations, 78-5907

Drosophila

- Mycotoxins
 - Mutagenic Activity, 78-5466
- Virus, DNA
 - Mutagenic Activity, 78-5851
- Virus, RNA
 - Mutagenic Activity, 78-5851

Drosophila melanogaster

- Caffeine
 - Mutagenic Activity, 78-5656
- Drugs
 - Mutagenic Activity, 78-5424
- Methanesulfonic Acid, Ethyl Ester
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5656, 78-5660
 - Sex Chromosomes, 78-5658
- Methanesulfonic Acid, Methyl Ester
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5660, 78-5661
 - Recombination-Deficient Mutants, 78-5661
- 3,6-Pyridazinedione, 1,2-Dihydro-
 - Mutagenic Activity, 78-5656
- Radiation, Ionizing
 - Mutagenic Activity, 78-5661
 - Recombination-Deficient Mutants, 78-5661
- Ultraviolet Rays
 - Mutagenic Activity, 78-5661
- Urea, Ethyl Nitroso-
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5660
- Urea, Methyl Nitroso-
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5660

Drug Therapy

- Bladder Neoplasms
 - Multiple Myeloma, 78-5894
- Cervix Neoplasms
 - Carcinoma In Situ, 78-5888
- Coumarin, 3-(α -Acetonylbenzyl)-4-hydroxy-
 - Immune Response, Review, 78-5433
- Coumarin, 3,3'-Methylenebis(4-hydroxy)-
 - Immune Response, Review, 78-5433
- Hodgkin's Disease
 - Neoplasms, Multiple Primary, 78-5888, 78-5910
 - Risk Factors, 78-5910

gs

- Chromosome Aberrations
 - Bone Marrow Cells, Mouse, 78-5424
 - Micronucleus Test, 78-5428
- Drosophila melanogaster*
 - Mutagenic Activity, 78-5424

Anal Ulcer

- Gastrectomy
 - Nitrous Acid, 78-5633

Neoplasms

- Adenocarcinoma
 - Diethylamine, 2,2-Dichloro-*N*-nitroso-, 78-5572
- Carcinoma, Epidermoid
 - Hydrazine, 1,1-Dimethyl-, 78-5861
- Hydrazine, 1,1-Dimethyl-
 - Corynebacterium parvum*, 78-5861

11,14-Eicosatetranic Acid

- Phorbol 12,13-Didecanoate
 - Cell Differentiation, 78-5509
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Cell Differentiation, 78-5509

onucleases

- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Micrococcus luteus*, 78-5700
- Micrococcus luteus*
 - DNA Repair, 78-5700
 - Isolation and Characterization, 78-5700
- Ultraviolet Rays
 - Micrococcus luteus*, 78-5700
- Virus, Polyoma, BK
 - DNA, Viral, 78-5795

otoxins

- Bacillus thuringiensis*
 - Isolation and Characterization, 78-5467
 - Mitosis, 78-5467

vid

- Mammary Neoplasms, Experimental
 - Prolactin, 78-5992

eritis

- Intestinal Polyps
 - Epidemiology, 78-5883

eritis, Regional

- Virus, Cytomegalo
 - DNA, Viral, 78-5885

inophils

- Cervix Neoplasms
 - Neoplasm Invasiveness, 78-5889
- Leukemia, Myeloblastic
 - Colony Stimulating Factor, 78-5984
- Leukemia, Myelocytic
 - Colony Stimulating Factor, 78-5984
- Vaginal Neoplasms
 - Neoplasm Invasiveness, 78-5889
- Vulvar Neoplasms
 - Neoplasm Invasiveness, 78-5889

ndymoma

- Child
 - Epidemiology, Denmark, 78-5925
- Urea, Methyl Nitroso-
 - Urea, Ethyl Nitroso-, 78-5638

ocryptine, 2-Bromo-

- Mammary Neoplasms, Experimental

Ergocryptine, 2-Bromo- (cont'd)

- Precancerous Conditions, 78-5992

Erythrocytes

- Virus, Radiation Leukemia
 - Virus Replication, 78-5853

Erythroleukemia

- Acetamide, *N*-Fluorenyl-2-yl-
 - Bone Marrow Cells, 78-5483
 - Histological Study, Rat, 78-5483
- RNA, Messenger
 - Cycloheximide, 78-5764
 - Pactamycin, 78-5764
 - Sodium Fluoride, 78-5764
- Virus, Friend Murine Leukemia
 - Polyribosomes, 78-5764
 - RNA, Messenger, 78-5764
- Virus, Herpes Simplex 1
 - Polyribosomes, 78-5764
 - RNA, Messenger, 78-5764
- Virus, Rauscher Murine Leukemia
 - Antigen-Antibody Complex, 78-5782
 - Neoplasm Regression, Spontaneous, 78-5782

Escherichia coli

- Antineoplastic Agents
 - Coliphages, 78-5515
- Butyrophene, γ -Chloro-*p*-fluoro-
 - DNA Repair, 78-5449
- Cobalt Chloride
 - Tryptophan Reversion, 78-5676
- Cyanamide, Methylnitroso-
 - Mutagenic Activity, 78-5588
 - Tryptophan Reversion, 78-5588
- Fibrosarcoma
 - Mutagenic Activity, 78-5670
- Glycoside Hydrolases
 - Isolation and Characterization, 78-5663
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Mutagenic Activity, 78-5588, 78-5676
- 1-Hexanamine, *N*-Hexyl-*N*-nitroso-
 - Nitrosation, 78-5612
- Methanesulfonic Acid, Ethyl Ester
 - DNA Repair, 78-5651
- Mycotoxins
 - Coliphages, 78-5515
- Nitroso Compounds
 - Coliphages, 78-5515
- Polycyclic Hydrocarbons
 - Coliphages, 78-5515
- 2-Propenoic Acid, 2-Methyl-, 1,2-Ethanediyl Ester
 - DNA Repair, 78-5449
- Quinoline, 4-Nitro-, 1-Oxide
 - Mutagenic Activity, 78-5671
- Quinoline, 4-Nitroso-, 1-Oxide
 - Mutagenic Activity, 78-5670
- Radiation, Ionizing
 - Methanesulfonic Acid, Ethyl Ester, 78-5651
- Urea, Methyl Nitroso-
 - Mutagenic Activity, 78-5588

Esophageal Neoplasms

- Benzylamine, *N*-Methyl-
 - Nitrous Acid, Sodium Salt, 78-5586
 - Precancerous Conditions, 78-5586
- Benzylamine, *N*-Methyl-*N*-nitroso-
 - Precancerous Conditions, 78-5586
- Zinc, 78-5587

Esophageal Neoplasms (cont'd)

- Diethylamine, *N*-Nitroso-
Carcinogenic Activity, Rat, 78-5572
- Ethnic Groups
Epidemiology, 78-5938
- Ethyl Alcohol
Epidemiology, 78-5939
Epidemiology, Review, 78-5404
- Liver Cirrhosis
Ethyl Alcohol, 78-5938
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
Carcinogenic Activity, Rat, 78-5580
- Smoking
Epidemiology, 78-5939
Epidemiology, Review, 78-5404
- Ulcer
Benzylamine, *N*-Methyl-*N*-nitroso-, 78-5586
- Virus, Papilloma
Epidemiology, Cattle, 78-5790
- Zinc
Diet, 78-5587

Esophagus

- Benzylamine, *N*-Methyl-*N*-nitroso-
Organ Distribution, Rat, 78-5586

Estradiol

- Age Factors
Metabolism, 78-5991
- Breast Neoplasms
Plasma/Urine Ratios, 78-5991
- Cell Differentiation
Vagina, Neonatal Mouse, 78-5890
- Collagen
Vagina, Neonatal Mouse, 78-5890
- Mammary Neoplasms, Experimental
Prolactin, 78-5992
- Vagina, Neonatal Mouse
Ultrastructural Study, 78-5890

Estrinol

- Age Factors
Metabolism, 78-5991
- Breast Neoplasms
Plasma/Urine Ratios, 78-5991

Estrogens

- Breast Neoplasms
Epidemiology, Chinese, 78-5891
Epidemiology, Review, 78-5432
Genetics, 78-5891
Menopause, 78-5949
- Mammary Neoplasms, Experimental
Benz(a)anthracene, 7,12-Dimethyl-, 78-5546
Castration, 78-5546
Receptors, Hormone, 78-5546
- Uterine Neoplasms
Epidemiology, Review, 78-5432

Estrone

- Age Factors
Metabolism, 78-5991
- Breast Neoplasms
Plasma/Urine Ratios, 78-5991

Ethane, 1,2-Dichloro-

- Occupational Hazard
Epidemiology, 78-5969

Ethanol, 2-Chloro-

- Occupational Hazard

Ethanol, 2-Chloro- (cont'd)

- Epidemiology, 78-5969

Ethanol, 2-Mercapto-

- B-Lymphocytes
Colony Formation, Review, 78-5439
- Multiple Myeloma
Colony Formation, Review, 78-5439

Ether, Bis(2-chloroethyl)-

- Occupational Hazard
Epidemiology, 78-5969

Ether, Bis(chloromethyl)-

- Lung Neoplasms
Air Pollution, Review, 78-5409
Epidemiology, Review, 78-5410

Ethidium Bromide

- Aroclor 1254
Ames Test, 78-5517
- Barbituric Acid, 5-Ethyl-5-phenyl-
Ames Test, 78-5517
- 5,6-Benzoflavone
Ames Test, 78-5517
- Ultraviolet Rays
DNA Repair, 78-5477

Ethyl Alcohol

- Carcinogen, Environmental
Epidemiology, Review, 78-5402
- Esophageal Neoplasms
Epidemiology, 78-5939
Epidemiology, Review, 78-5404
Liver Cirrhosis, 78-5938
- Ethnic Groups
Epidemiology, 78-5938
- Laryngeal Neoplasms
Epidemiology, 78-5936, 78-5939
- Liver Cirrhosis
Epidemiology, Japan, 78-5966
- Liver Neoplasms
Liver Cirrhosis, 78-5938
- Nitrosamines
Nitrogen Oxide, 78-5617
Permeability, Laboratory Gloves, 78-5596
- Prostatic Neoplasms
Ethnic Groups, 78-5938
- Respiratory Tract Neoplasms
Epidemiology, Review, 78-5404
- Smoking
Co-carcinogenic Effect, Review, 78-5404

Ethylamine, *N*-Methyl-*N*-nitroso-

- Food Contamination
Concentration Levels, 78-5621
- Liver Neoplasms
Hepatectomy, 78-5592
- Occupational Hazard
Permeability, Laboratory Gloves, 78-5596

Ethylene

- Occupational Hazard
Epidemiology, 78-5969

Ethylene, Bromo-

- Ames Test
Mutagenic Activity, 78-5492

Ethylene, Chloro-

- Angiosarcoma
Chromosome Aberrations, 78-5491

ethylene, Chloro- (cont'd)

- Epidemiology, 78-5486, 78-5967
- Occupational Hazard, 78-5486, 78-5491
- Occupational Hazard, Review, 78-5435
- Precancerous Conditions, Review, 78-5435
- Barbituric Acid, 5-Ethyl-5-phenyl-
Cytochrome P-450, 78-5490
- Mutagenic Activity, Hamster V79 Cells, 78-5593
- Carcinogen, Environmental
Metabolism, Review, 78-5406
- Cholanthrene, 3-Methyl-
Cytochrome P-450, 78-5490
- Chromosome Aberrations
Occupational Hazard, 78-5489
- Occupational Hazard, Review, 78-5436
- Cytochrome P-450
Metabolism, 78-5490
- Diethylamine, N-Nitroso-
Permeability, Laboratory Gloves, 78-5595
- Dimethylamine, N-Nitroso-
Permeability, Laboratory Gloves, 78-5595
- Dipropylamine, N-Nitroso-
Permeability, Laboratory Gloves, 78-5595
- Enzyme Activation
Mutagenic Activity, Review, 78-5436
- Hepatoma
Case Report, 78-5485
- Neoplasms, Multiple Primary, 78-5485
- Liver Neoplasms
Angiosarcoma, 78-5485, 78-5486, 78-5967
- Lung Neoplasms
Air Pollution, Review, 78-5409
- Lymphocytes
Chromosome Aberrations, 78-5489
- Microsomes, Liver
Metabolism, 78-5490
- Occupational Hazard
Epidemiology, Review, 78-5410
- Piperidine, 1-Nitroso-
Permeability, Laboratory Gloves, 78-5595
- Pyrrolidine, 1-Nitroso-
Permeability, Laboratory Gloves, 78-5595

ethylene, Chloro- Polymer

- Angiosarcoma
Epidemiology, 78-5967
- Bone Marrow Cells
Chromosome Aberrations, 78-5488
- Chromosome Aberrations
Occupational Hazard, 78-5488
- Liver Neoplasms
Angiosarcoma, 78-5967
- Lymphocytes
Chromosome Aberrations, 78-5488

ethylene, 1,1-Dichloro-

- Ames Test
Mutagenic Activity, 78-5492
- Pyridine, 4-(p-Nitrobenzyl)-
Carcinogenic Metabolite, 78-5492

ethylene, 1,1-Difluoro-

- Pyridine, 4-(p-Nitrobenzyl)-
Carcinogenic Metabolite, 78-5492

ethylene Glycol Dimethacrylate

- see 2-Propenoic Acid, 2-Methyl-, 1,2-Ethanediyl Ester

ethylene Oxide

- Intestinal Neoplasms

Ethylene Oxide (cont'd)

- Occupational Hazard, 78-5969
- Kidney Neoplasms
Occupational Hazard, 78-5969
- Leukemia, Lymphocytic
Epidemiology, 78-5969
- Occupational Hazard, 78-5969
- Leukemia, Myeloblastic
Epidemiology, 78-5969
- Occupational Hazard, 78-5969
- Melanoma
Occupational Hazard, 78-5969
- Occupational Hazard
Epidemiology, 78-5969
- Stomach Neoplasms
Epidemiology, 78-5969
- Occupational Hazard, 78-5969

Ethylene Oxide, Chloro-

- Ames Test
Mutagenic Activity, 78-5492
- Nucleic Acids, Binding
Carcinogenic Activity, 78-5436
- Pyridine, 4-(p-Nitrobenzyl)-
Carcinogenic Metabolite, 78-5492

Ethylene, Trichloro-

- Pyridine, 4-(p-Nitrobenzyl)-
Carcinogenic Metabolite, 78-5492

Ethylenimine

- Chromosomes
DNA Replication, 78-5673
- HeLa Cells
DNA Replication, 78-5673

Eye Neoplasms

- Melanoma
Genetics, 78-5869

Feces

- Ascorbic Acid
Mutagens, 78-5629
- Mutagens
Ames Test, 78-5629
- Nitroso Compounds
Mutagens, 78-5629

Fenuron

- see Urea, 1,1-Dimethyl-3-phenyl-

Fetal Globulins

- Sarcoma
Antigen-Antibody Reactions, 78-5531
- Cholanthrene, 3-Methyl-, 78-5531
- Transplantation Immunology, 78-5531

Fibroblasts

- Aging
Cells, Cultured, 78-5977
- Alkaline Phosphatase
Cell Transformation, Neoplastic, 78-5798
- Benz(a)anthracene, 7,12-Dimethyl-
Mucopolysaccharidosis, 78-5540
- Cell Division
Epidermal Growth Factors, 78-5813
- Cell Membrane
Epidermal Growth Factors, 78-5813
- Cells, Cultured
Caseinolysis, 78-5977
- Growth

Fibroblasts (cont'd)

- Culture Media, 78-5977
- Leukemia, Myelocytic
 - Hematopoietic Stem Cells, 78-5904
- Mesenchymoma
 - Hematopoietic Stem Cells, 78-5904
- Radiation, Ionizing
 - Cell Transformation, Neoplastic, 78-5730
- Ultraviolet Rays
 - DNA Repair, 78-5704, 78-5705, 78-5707, 78-5709
- Urea, Ethyl Nitroso-
 - DNA, Alkylation, 78-5648
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-5648
- Xeroderma Pigmentosum
 - DNA Repair, 78-5704

Fibrosarcoma

- Antigens, Viral
 - Antibody Specificity, 78-5866
- Carcinoembryonic Antigen
 - Antibody Specificity, 78-5866
- Cholanthrene, 3-Methyl-
 - Antigenic Determinants, 78-5866
 - Carcinoembryonic Antigen, 78-5866
 - Corynebacterium parvum*, 78-5520
 - Histological Study, Rabbit, 78-5506
- Cicatrix
 - Burns, 78-5970
- Corynebacterium parvum*
 - Macrophage Activation, 78-5520
- Escherichia coli*
 - Mutagenic Activity, 78-5670
- Laryngeal Neoplasms
 - Radiation, Ionizing, 78-5711
 - Radiotherapy, 78-5711
- Lung Neoplasms
 - Benzo(a)pyrene, 78-5878
- Neoplasms, Connective Tissue
 - Cicatrix, 78-5970
- Nervous System Neoplasms
 - Urea, Methyl Nitroso-, 78-5643
- Quinoline, 4-Nitro-, 1-Oxide
 - Carcinogenic Activity, Mouse, 78-5670
 - Histological Study, Rabbit, 78-5506
- Quinoline, 4-Nitroso-, 1-Oxide
 - Carcinogenic Activity, Mouse, 78-5670
- Somatomedin
 - Growth Factors, Review, 78-5452
- Tracheal Neoplasms
 - Cholanthrene, 3-Methyl-, 78-5521
- Ultraviolet Rays
 - Transplantation Immunology, 78-5858
- Virus, Murine Leukemia
 - Antigens, Viral, 78-5866

Flavone, 3,3',4',5,7-Pentahydroxy-

- Skin Neoplasms
 - Benzo(a)pyrene, 78-5567

Flukes

- see Trematoda

Fluoren-2-amine

- Aroclor 1254
 - Ames Test, 78-5517
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Ames Test, 78-5517
- 5,6-Benzoflavone

Fluoren-2-amine (cont'd)

- Ames Test, 78-5517

Fluorescence

- Benzo(a)anthracene, 7,12-Dimethyl-
 - DNA, Binding, 78-5513
 - Hydrocarbon-DNA Adducts, 78-5513
- Benzo(a)pyrene
 - DNA, Binding, 78-5513
- Glycine, *N*-(Phosphonomethyl)-*N*-nitroso-
 - Quantitation Method, 78-5620
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Cell Transformation, Neoplastic, 78-5685
- Nitrosamines
 - Quantitation Method, 78-5620

5-Fluoro-7,12-benz(a)anthraquinone

- see Benzo(a)anthracene-7,12-dione, 5-Fluoro-

Folic Acid

- Aflatoxin B1
 - Cytochrome P-450, 78-5473
 - Oxidoreductases, 78-5473

Food Contamination

- Benzo(a)pyrene
 - Smoked Products, 78-5555
- 1-Butanamine, *N*-Butyl-*N*-nitroso-
 - Quantitation Method, 78-5616
- Chlorine
 - Carcinogenic Potential, Review, 78-5403
- Diethylamine, *N*-Nitroso-
 - Concentration Levels, 78-5621
 - Degradation, 78-5618
 - Nitric Acid, Sodium Salt, 78-5618
 - Quantitation Method, 78-5616
- Dimethylamine, *N*-Nitroso-
 - Degradation, 78-5618
 - Nitric Acid, Sodium Salt, 78-5618
 - Quantitation Method, 78-5616, 78-5623
- Ethylamine, *N*-Methyl-*N*-nitroso-
 - Concentration Levels, 78-5621
- Hepatitis
 - Epidemiology, India, 78-5964
- Nitric Acid, Sodium Salt
 - Metabolism, 78-5615
- Nitrosamines
 - Concentration Levels, 78-5621
 - Fragmentation and Condensation, 78-5579
 - Nitric Acid, Sodium Salt, 78-5615
 - Nitrous Acid, Sodium Salt, 78-5615
 - Quantitation Method, 78-5621, 78-5624
- Nitrous Acid, Sodium Salt
 - Metabolism, 78-5615
- Piperidine, 1-Nitroso-
 - Degradation, 78-5618
 - Nitric Acid, Sodium Salt, 78-5618
- Polycyclic Hydrocarbons
 - Smoked Products, 78-5555
- Proline, 4-Hydroxy-1-nitroso-
 - Quantitation Method, 78-5616
- Proline, 1-Nitroso-
 - Quantitation Method, 78-5616
- Pyrrolidine, 1-Nitroso-
 - Concentration Levels, 78-5621
 - Degradation, 78-5618
 - Nitric Acid, Sodium Salt, 78-5618
 - Quantitation Method, 78-5616, 78-5623
- Sarcosine, *N*-Nitroso-

Food Contamination (cont'd)

Quantitation Method, 78-5616

Stomach Neoplasms

Talc, 78-5943

Reber-Albright Syndrome

see Lactation Disorders

Mamamide, N,N-Dimethyl-

Leukemia, Myeloblastic

Cell Differentiation, 78-5983

Mamamide, N-Methyl-

Leukemia, Myeloblastic

Cell Differentiation, 78-5983

Furfuraldehyde

Benzo(a)pyrene

Co-carcinogenic Effect, 78-5571

Diethylamine, N-Nitroso-

Co-carcinogenic Effect, 78-5571

Respiratory Tract Neoplasms

Benzo(a)pyrene, 78-5571

Diethylamine, N-Nitroso-, 78-5571

Furfural

see 2-Furfuraldehyde

Sararium

Mycotoxins

Mutagenic Activity, 78-5466

Toxicity, Mouse, 78-5466

Angliioneuroma

Nervous System Neoplasms

Urea, Methyl Nitroso-, 78-5643

Angliosides

Leukemia

Cell Membrane, 78-5987

Leukemia, Myelocytic

Chromosomes, Human, 21-22, 78-5987

Lymphocytes

Cell Membrane, 78-5987

Lymphoma

Cell Membrane, 78-5987

Plasmacytoma

Cell Membrane, 78-5987

Gastrectomy

Duodenal Ulcer

Nitrous Acid, 78-5633

Nitrous Acid

Gastric Juice, 78-5633

Stomach Neoplasms

Carcinoma, 78-5879

Neoplasm Recurrence, 78-5879

Nitrous Acid, 78-5633

Gastric Juice

Gastrectomy

Nitrous Acid, 78-5633

Nitric Acid, Sodium Salt

Nitric Acid, 78-5634

Nitrous Acid, 78-5634

Nitrous Acid

Pentagastrin, 78-5632

Peptic Ulcer

Nitrous Acid, 78-5632

Smoking

Thiocyanic Acid, 78-5632

Stomach Neoplasms

Gastric Juice (cont'd)

Hydrogen-Ion Concentration, 78-5632

Nitrous Acid, 78-5632

Thiocyanic Acid

Pentagastrin, 78-5632

Gastritis

Stomach Neoplasms

Adenocarcinoma, 78-5880

Case Report, 78-5880

Polyps, 78-5880

Gastrointestinal Neoplasms

Adenocarcinoma

Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5867

Air Pollution

Epidemiology, Poland, 78-5941

Asbestos

Epidemiology, Review, 78-5445

Occupational Hazard, Review, 78-5445

Carcinoembryonic Antigen

Hypersensitivity, Delayed, 78-5867

Immune Response, 78-5867

Immunization, 78-5867

Neoplasm Metastasis, 78-5867

Diet

Epidemiology, Poland, 78-5941

Epidemiology

Spain, 78-5915

Ethnic Groups

Epidemiology, 78-5938

Hydrazine, 1,2-Dimethyl-, Dihydrochloride

Carcinoembryonic Antigen, 78-5867

Genetics

Breast Neoplasms

Classification, Review, 78-5455

Estrogens, 78-5891

Carcinogen, Environmental

Cell Transformation, Neoplastic, Review, 78-5455

Epidemiology, Review, 78-5405

Colonic Neoplasms

Chromosome Abnormalities, 78-5881

Classification, Review, 78-5455

Lymphoma, 78-5882

Eye Neoplasms

Melanoma, 78-5869

Hodgkin's Disease

Epidemiology, 78-5919

Intestinal Polyps

Virus, Cytomegalo, 78-5885

Leukemia, Lymphoblastic

Case Report, 78-5902

Leukemia, Myeloblastic

Case Report, 78-5902

Leukemia, Myelocytic

Case Report, 78-5903

Chromosomes, Human, 21-22, 78-5903

Lung Neoplasms

Classification, Review, 78-5455

Melanoma

Case Report, 78-5869

Neoplasms, Multiple Primary, 78-5869

Neoplastic Endocrine-Like Syndromes

Classification, Review, 78-5455

Nervous System Neoplasms

Classification, Review, 78-5455

Neurofibromatosis

Classification, Review, 78-5455

Genetics (cont'd)

- Radiation, Ionizing
 - Occupational Hazard, 78-5719
- Retinoblastoma
 - Case Report, 78-5870
 - Chromosomes, Human, 13-15, 78-5870
- Urogenital Neoplasms
 - Classification, Review, 78-5455
 - Kidney Diseases, 78-5961
- Uterine Neoplasms
 - Carcinoma, Papillary, 78-5892
 - Case Report, 78-5892
 - Leiomyoma, 78-5892
- Virus, Bovine Leukemia
 - Immune Response, 78-5982

Glioblastoma Multiforme

- Brain Neoplasms
 - Epidemiology, Sweden, 78-5927
- Hodgkin's Disease
 - Neoplasms, Multiple Primary, 78-5910
- Urea, Methyl Nitroso-
 - Urea, Ethyl Nitroso-, 78-5638

Glioma

- Brain Neoplasms
 - Epidemiology, Sweden, 78-5927
 - Radiation, Ionizing, 78-5872
 - Urea, Ethyl Nitroso-, 78-5649
 - Urea, Methyl Nitroso-, 78-5859
 - Virus, Polyoma, BK, 78-5794
- Radiation, Ionizing
 - Case Report, 78-5872
- Urea, Methyl Nitroso-
 - Antibody-Dependent Cytotoxicity, 78-5859
 - Antibody Specificity, 78-5859
- Virus, Polyoma, BK
 - Proteins, Acidic, 78-5794

Gliosarcoma

see Glioma

Globin

- Polyribosomes
 - RNA, Messenger, 78-5764

Glomerulonephritis

- Leukemia, Myeloblastic
 - Butyric Acid, 4-(p-Bis(2-chloroethyl)aminophenyl)-
78-5896

Glucose, 2-Deoxy-

- Virus, Rauscher Murine Leukemia
- Viral Proteins, 78-5780

Glucosephosphate Dehydrogenase

- Nervous System Neoplasms
 - Urea, Ethyl Nitroso-, 78-5647
- Radiation, Ionizing
 - Cell Transformation, Neoplastic, 78-5730

Glucuronidase

- Benzo(a)pyrene
 - Benzo(a)pyren-3-ol, 78-5552
 - Benzo(a)pyren-9-ol, 78-5552
 - Carcinogenic Metabolite, 78-5552
 - Metabolism, Dose-Response Study, 78-5552

Glutamic Acid, Monosodium Salt

- Mammary Neoplasms, Experimental
- Pregnancy, 78-5995
- Progesterone, 78-5995

Glutamyl Transpeptidase

- Nitrosamines
 - Dose-Response Study, 78-5594
 - Serum Levels, 78-5594

Glutathione

- Aflatoxin B1
 - Dietary Fats, 78-5473
 - Metabolism, Rat, 78-5472
- Benzo(a)pyrene
 - Ames Test, 78-5559

Glycine, N-(Phosphonomethyl)-N-nitroso-

- Fluorescence
 - Quantitation Method, 78-5620

Glycogen

- Nitrosamines
 - Hepatotoxicity, 78-5594

Glycoproteins

- Virus, Adeno 2
 - Concanavalin A, 78-5840
 - Immunoprecipitation, 78-5840
 - Isolation and Characterization, 78-5840
 - Methylglycosides, 78-5840
 - Proteolysis, 78-5840
- Virus, Adeno 5
 - Cell Transformation, Neoplastic, 78-5844
- Virus, C-Type RNA Tumor
 - Tryptic Peptide Analysis, 78-5762
- Virus, Recombinant, 78-5762
- Virus, Murine Leukemia
 - Carbohydrates, 78-5781
 - Cell Transformation, Neoplastic, 78-5781
 - Mouse, AKR, 78-5781
 - Virus, Recombinant, 78-5762
- Virus, Radiation Leukemia
 - Virus, Recombinant, 78-5762
- Virus, Rauscher Murine Leukemia
 - Antigenic Determinants, 78-5781
 - Carbohydrates, 78-5781
 - Isolation and Characterization, 78-5781

Glycoside Hydrolases

- Adenine, 3-Methyl-
 - DNA, Alkylation, 78-5663
- Escherichia coli*
 - Isolation and Characterization, 78-5663
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Adenine, 3-Methyl-, 78-5663
- Methanesulfonic Acid, Methyl Ester
 - Adenine, 3-Methyl-, 78-5663
- Sulfuric Acid, Dimethyl Ester
 - Adenine, 3-Methyl-, 78-5663
- Urea, Methyl Nitroso-
 - Adenine, 3-Methyl-, 78-5663

Granuloma

- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Croton Oil, 78-5677
 - Ouabain Resistance, Rat, 78-5677
- Head and Neck Neoplasms
 - Thorium Dioxide, 78-5973
- Thorium Dioxide
 - Epidemiology, Portugal, 78-5973

Growth

- Breast Neoplasms
 - Cells, Cultured, 78-5950
 - Insulin, 78-5950

with (cont'd)

- Carcinoma
 - Tissue Culture, Review, 78-5454
- Fibroblasts
 - Culture Media, 78-5977
- Mammary Neoplasms, Experimental
 - Androgen Dependency, 78-5988
 - Testosterone, Propionate, 78-5988
- Melanoma
 - Tissue Culture, Review, 78-5454
- Sarcoma
 - Tissue Culture, Review, 78-5454
- Tracheal Neoplasms
 - Immunosuppression, 78-5978

with Factors

- Cell Transformation, Neoplastic
 - DNA Replication, 78-5808
 - Iron, Metabolism, 78-5808
 - Siderophore, 78-5808
- Neoplasms
 - Synthesis, Review, 78-5452
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-5808
 - Picolinic Acid, 78-5808

with Substances

- Breast Neoplasms
 - Carcinoma, Ductal, 78-5994
 - Epithelial Cells, 78-5994
- Guanidine, 1-Butyl-3-nitro-1-nitroso-
 - DNA Repair
 - Carcinogenic Activity, Rat, 78-5683
- Guanidine, 1-Ethyl-3-nitro-1-nitroso-
 - Skin Neoplasms
 - Carcinogenic Activity, Mouse, 78-5598

- Guanidine, 1-Hexyl-3-nitro-1-nitroso-
 - DNA Repair
 - Carcinogenic Activity, Rat, 78-5683

- Guanidine, 1-Isobutyl-3-nitro-1-nitroso-
 - DNA Repair
 - Carcinogenic Activity, Rat, 78-5683

- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Adenine, 3-Methyl-
 - Glycoside Hydrolases, 78-5663

- Cell Aggregation
 - Cell Transformation, Neoplastic, 78-5507
- Chromosome Aberrations
 - Cytogenetics, Mouse, 78-5682

- Cobalt Chloride
 - Mutagenic Activity, 78-5676

- DNA
 - Alkaline Sucrose Gradient Sedimentation, 78-5678
 - Immunoreactivity, Antinucleoside Antibodies 78-5678

- DNA, Bacterial
 - Temperature Sensitive Mutants, 78-5679

- DNA Repair
 - Carcinogenic Activity, Rat, 78-5683

- DNA Replication
 - Cell Transformation, Neoplastic, 78-5685

- Endonucleases
 - Micrococcus luteus*, 78-5700

- Escherichia coli*
 - Mutagenic Activity, 78-5588, 78-5676

Fluorescence

Guanidine, 1-Methyl-3-nitro-1-nitroso- (cont'd)

- Cell Transformation, Neoplastic, 78-5685
- Granuloma
 - Croton Oil, 78-5677
 - Ouabain Resistance, Rat, 78-5677
- Haemophilus influenzae*
 - DNA, Bacterial, 78-5679
 - DNA Replication, 78-5679
 - Mutagenic Activity, 78-5679
- HeLa Cells
 - DNA, 78-5678
- Histidine Reversion
 - Mutagenic Activity, 78-5681
- Intestinal Neoplasms
 - Bile Acids and Salts, 78-5684
 - 5 β -Cholan-24-oic Acid, 3 α -Hydroxy-, 78-5684
 - Taurodeoxycholic Acid, 78-5684
- Micrococcus radiodurans*
 - Mutagenic Activity, 78-5701
 - Tryptophan Auxotrophs, 78-5701
- Neurospora crassa*
 - Mutagenic Activity, 78-5675
- Salmonella typhimurium*
 - Histidine Reversion, 78-5681
- Sarcoma
 - Cells, Cultured, 78-5507
- Sarcoma, Osteogenic
 - Cell Transformation, Neoplastic, 78-5507
 - Cells, Cultured, 78-5507
 - Tumorigenicity, Clone Cells, 78-5507
- Skin Neoplasms
 - Carcinogenic Activity, Mouse, 78-5598
- Soy Beans
 - Somatic Mosaicism, 78-5664
- Streptomycin
 - Lethality, Algae, 78-5699
- Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Uridine, 5-Bromo-2'-deoxy-
 - Cell Transformation, Neoplastic, 78-5685
 - DNA Replication, 78-5685
 - Photosensitivity, 78-5685
- Virus, SV40
 - Temperature Sensitive Mutants, 78-5810

Guanidine, Nitroso-

- Lymphoma
 - Mutagenic Activity, S49 Cells, 78-5680
 - Ouabain Resistance, 78-5680

Guanidine, 1-Pentyl-3-nitro-1-nitroso-

- DNA Repair
 - Carcinogenic Activity, Rat, 78-5683

Guanidine, 1-Propyl-3-nitro-1-nitroso-

- DNA Repair
 - Carcinogenic Activity, Rat, 78-5683

Guanine, 7-Methyl-

- Dimethylamine, N-Nitroso-
 - DNA, Alkylation, 78-5599, 78-5600
 - DNA, Liver, 78-5607
 - Kidney, Liver, Rat, 78-5600
 - Liver, Rat, 78-5599

Guanosine Cyclic 3',5' Monophosphate

- Pancreatic Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5534
 - Tissue Levels, 78-5534

Guanyl Cyclase

- Hydrazine
 - Enzymatic Activity, 78-5500
- Hydrazine, 1,1-Dimethyl-
 - Enzymatic Activity, 78-5500
- Hydrazine, Sulfate
 - Enzymatic Activity, 78-5500
- Isonicotinic Acid Hydrazide
 - Enzymatic Activity, 78-5500

Haemophilus influenzae

- Guanidine, 1-Methyl-3-nitro-1-nitroso-DNA, Bacterial, 78-5679
- DNA Replication, 78-5679
- Mutagenic Activity, 78-5679

Hair Dyes

- o*-Phenylenediamine, 4-Nitro-
 - Carcinogenic Activity, Review, 78-5425
 - Mutagenic Activity, Review, 78-5425
- p*-Phenylenediamine, 2-Nitro-
 - Carcinogenic Activity, Review, 78-5425
 - Mutagenic Activity, Review, 78-5425
- o*-Phenylenediamine
 - Mutagenic Activity, Review, 78-5425
- Toluene-2,4-diamine
 - Carcinogenic Activity, Review, 78-5425
 - Mutagenic Activity, Review, 78-5425

Head and Neck Neoplasms

- Age Factors
 - Epidemiology, USSR, 78-5917
- Carcinoma
 - Virus, Epstein-Barr, 78-5830
- Granuloma
 - Thorium Dioxide, 78-5973
- Virus, Epstein-Barr
 - Antigens, Viral, 78-5830

HeLa Cells

- Adriamycin
 - DNA Replication, 78-5673
- Carcinoma
 - Mouse, Nude, 78-5832
- Ethylenimine
 - DNA Replication, 78-5673
- Guanidine, 1-Methyl-3-nitro-1-nitroso-DNA, 78-5678
- Quinoline, 4-Nitro-, 1-Oxide
 - DNA Replication, 78-5673
- Radiation, Ionizing
 - DNA, 78-5678
- Virus, Adeno 2
 - Viral Proteins, 78-5841

Hemangioendothelioma

- Liver Neoplasms
 - Thorium Dioxide, 78-5973

Hemangioma

- Bladder Neoplasms
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5583
- Diethylamine, *N*-Nitroso-
 - Carcinogenic Activity, Cat, Chicken, 78-5573
- Liver Neoplasms
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580

Hematologic Diseases

- Down's Syndrome
 - Chromosome Aberrations, 78-5907

Hematopoietic Stem Cells

- Leukemia, Myelocytic
 - Cell Differentiation, 78-5904
 - Fibroblasts, 78-5904
- Mesenchymoma
 - Cell Differentiation, 78-5904
 - Fibroblasts, 78-5904

Hematopoietic System

- Virus, Murine Leukemia
 - Virus Replication, 78-5756
- Virus, Radiation Leukemia
 - Virus Replication, 78-5756

Hemolysis

- Virus, Radiation Leukemia
 - Blood Cells, Leukemogenic Activity, 78-5853

Hemorrhage

- Leukemia, Myelocytic
 - 1,4-Butanediol, Dimethylsulfonate, 78-5897
 - Cystitis, 78-5897

Hepatitis

- Aflatoxin B1
 - Epidemiology, India, 78-5964
 - Histological Study, 78-5964
- Contraceptives, Oral
 - Epidemiology, 78-5431
- Food Contamination
 - Epidemiology, India, 78-5964

Hepatoma

- Aflatoxin B1
 - Cells, Cultured, 78-5471
 - Dietary Proteins, 78-5470
 - Vitamin B12, 78-5470
- Aflatoxin G1
 - Dietary Proteins, 78-5470
 - Vitamin B12, 78-5470
- Australia Antigen
 - DNA, Viral, 78-5837
 - Epidemiology, China, 78-5965
 - Epidemiology, Ivory Coast, 78-5962
- Contraceptives, Oral
 - Case Report, 78-5874
 - Diagnosis, 78-5874
 - Epidemiology, 78-5431
- Corticosterone
 - Tyrosine Aminotransferase, 78-5576
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
 - Antipain, 78-5583
 - Leupeptins, 78-5583
 - Pepstatins, 78-5583
 - Precancerous Conditions, 78-5583
- Dexamethasone
 - Receptors, Hormone, 78-5576
 - Tyrosine Aminotransferase, 78-5576
- Diethylamine, *N*-Nitroso-
 - Carcinogenic Activity, Cat, Chicken, 78-5573
 - Corticosterone, 78-5576
 - Dexamethasone, 78-5576
 - Propionic Acid, 2-(*p*-Chlorophenoxy)-2-methyl-, Ethyl Ester, 78-5574
 - Tyrosine Aminotransferase, 78-5576
 - WY-14,643, 78-5574
- Ethylene, Chloro-
 - Case Report, 78-5485
- Geographic Factors
 - Epidemiology, Ivory Coast, 78-5962

atoma (cont'd)

- Liver Cirrhosis
 - Australia Antigen, 78-5965
 - Epidemiology, Ivory Coast, 78-5962
- Microwaves
 - Thermography, 78-5999
- Neoplasms, Multiple Primary
 - Case Report, 78-5485
 - Ethylene, Chloro-, 78-5485
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
 - Carcinogenic Activity, Rat, 78-5580
- Thermography
 - Diagnosis, 78-5999
- Virus, Hepatitis
 - Australia Antigen, 78-5837
 - Epidemiology, Senegal, 78-5837

editary Diseases

- Cell Transformation, Neoplastic
 - DNA Repair, Review, 78-5451
- Radiation, Ionizing
 - DNA Repair, Review, 78-5451
- Ultraviolet Rays
 - DNA Repair, Review, 78-5451

erochromatin

- Bladder Neoplasms
 - Chromosome Abnormalities, 78-5893

achlorophene

- see Phenol, 2,2'-Methylenebis(3,4,6-trichloro)-

exanamine, *N*-Hexyl-*N*-nitroso-

- Bacillus brevis*
 - Nitrosation, 78-5612
- Escherichia coli*
 - Nitrosation, 78-5612
- Saccharomyces cerevisiae*
 - Nitrosation, 78-5612
- Saccharomycopsis lipolytica*
 - Nitrosation, 78-5612

okinase

- Sarcoma
 - Virus, Adeno 12, 78-5847
- Sarcoma, Reticulum Cell
 - Virus, Adeno 12, 78-5847
- Virus, Adeno 6
 - Enzymatic Activity, 78-5847
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5847

ocompatibility Antigens

- Hodgkin's Disease
 - Antigen Frequency, Review, 78-5462
- Leukemia, Lymphoblastic
 - Antigen Frequency, Review, 78-5462
- Lung Neoplasms
 - Isoantigens, 78-5862
- Nasopharyngeal Neoplasms
 - Antigen Frequency, Review, 78-5462
- Sarcoma
 - Carcinogen, Chemical, 78-5438
- Sarcoma, Mast Cell
 - Actin, 78-5863
- Transplantation Immunology
 - Hamster, Review, 78-5440
- Virus, Marek's Disease Herpes
 - Antigen-Antibody Reactions, 78-5748
- Virus, Rauscher Murine Leukemia

Histocompatibility Antigens (cont'd)

- Viral Proteins, 78-5779
- Virus, SV40
 - Antigenic Determinants, 78-5812
 - Transplantation Immunology, 78-5812

Histones

- Virus, SV40
 - Chromosomes, 78-5811

Hodgkin's Disease

- Age Factors
 - Epidemiology, East Germany, 78-5919
- Benzene
 - Occupational Hazard, 78-5919
- Carcinoma, Epidermoid
 - Neoplasms, Multiple Primary, 78-5910
- Child
 - Epidemiology, Portugal, 78-5918
 - Histological Study, 78-5918
- Drug Therapy
 - Risk Factors, 78-5910
- Genetics
 - Epidemiology, 78-5919
- Glioblastoma Multiforme
 - Neoplasms, Multiple Primary, 78-5910
- Histocompatibility Antigens
 - Antigen Frequency, Review, 78-5462
- Leukemia
 - Cyclophosphamide, 78-5898
 - Neoplasms, Multiple Primary, 78-5910
 - Prednisolone, 78-5898
 - Radiation, Ionizing, 78-5898
- Leukemia, Myelocytic
 - Cyclophosphamide, 78-5898
 - Radiation, Ionizing, 78-5898
 - Vinblastine, 78-5898
- Lupus Erythematosus, Systemic
 - Case Report, 78-5911
- Neoplasms, Multiple Primary
 - Drug Therapy, 78-5888, 78-5910
 - Epidemiology, 78-5919
 - Radiotherapy, 78-5888, 78-5910
- Occupational Hazard
 - Epidemiology, 78-5919
- Prostatic Neoplasms
 - Neoplasms, Multiple Primary, 78-5910
- Radiation, Ionizing
 - Risk Factors, 78-5910
- Rubber
 - Occupational Hazard, Women, 78-5954

Hybrid Cells

- Burkitt's Lymphoma
 - Transplantation, Heterologous, 78-5832
- Cell Transformation, Neoplastic
 - Cytoplasm, 78-5807
- Virus, Epstein-Barr
 - Transplantation, Heterologous, 78-5832
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-5807

Hycanthone

- Thymidine Kinase
 - Mutagenic Activity, 78-5666

Hydantoin, 5,5-Diphenyl-

- Lymphoma
 - Epidemiology, Review, 78-5430

- Hydrazine**
 Guanyl Cyclase
 Enzymatic Activity, 78-5500
- Hydrazine, 1,2-Diformyl-**
 Lung Neoplasms
 Adenocarcinoma, 78-5493
 Adenoma, 78-5493
 Carcinogenic Activity, Mouse, 78-5493
- Hydrazine, 1,1-Dimethyl-**
 Colonic Neoplasms
Corynebacterium parvum, 78-5861
 Neoplasm Metastasis, 78-5861
 Ear Neoplasms
 Carcinoma, Epidermoid, 78-5861
Corynebacterium parvum, 78-5861
 Guanyl Cyclase
 Enzymatic Activity, 78-5500
- Hydrazine, 1,2-Dimethyl-**
 Colon
 DNA, Binding, 78-5508
 Metabolism, Rat, 78-5508
 Colonic Neoplasms
 Adenocarcinoma, 78-5496, 78-5497, 78-5498
 Adenoma, 78-5497, 78-5498
 Carcinoma, Epidermoid, 78-5498
 Carcinoma, Mucinous, 78-5497
 Cell Cycle Kinetics, 78-5496
 DNA Replication, 78-5498
 Histological Study, Rat, 78-5497
 Neoplasm Metastasis, 78-5497
 Precancerous Conditions, 78-5497, 78-5498
 Ultrastructural Study, Rat, 78-5496
 Intestinal Neoplasms
 Dietary Fats, 78-5684
 Dietary Proteins, 78-5684
 Microflora, 78-5494
 Strain Difference, Rat, 78-5494
 Uterine Neoplasms
 Neoplasm Transplantation, 78-5495
 Sarcoma, 78-5495
- Hydrazine, 1,2-Dimethyl-, Dihydrochloride**
 Gastrointestinal Neoplasms
 Adenocarcinoma, 78-5867
 Carcinoembryonic Antigen, 78-5867
 Intestinal Neoplasms
 Adenocarcinoma, 78-5499
 Adenoma, 78-5499
 Anti-Antibodies, 78-5499
 Carcinoma, Mucinous, 78-5499
 Histological Study, Rat, 78-5499
 Neoplasm Metastasis, 78-5499
- Hydrazine, Sulfate**
 Guanyl Cyclase
 Enzymatic Activity, 78-5500
- Hydrogen Peroxide**
 Ultraviolet Rays
 Cell Survival, 78-5708
 Mutagenic Activity, 78-5708
 Synergistic Activity, 78-5708
- 1-Hydroxy-3-methylcholanthrene**
 see Cholanthren-1-ol, 3-Methyl-
- Hymenolepis nana**
 Retroperitoneal Neoplasms
- Hymenolepis nana (cont'd)**
 Sarcoma, 78-5980
- Hyperparathyroidism**
 Paraganglioma, Nonchromaffin
 Case Report, 78-5914
 Thymoma
 Case Report, 78-5914
- Hyperplasia**
 Bladder Neoplasms
 Acetamide, *N*-Fluoren-2-yl-, 78-5476
 Liver Neoplasms
 Acetamide, *N*-Fluoren-2-yl-, 78-5481
 Mammary Neoplasms, Experimental
 Precancerous Conditions, 78-5992
 Virus, Herpes
 Spleen, Guinea Pig, 78-5815
- Hypersensitivity, Delayed**
 Gastrointestinal Neoplasms
 Carcinoembryonic Antigen, 78-5867
- Hypertension**
 Uterine Neoplasms
 Epidemiology, Japan, 78-5953
- Hyperthyroidism**
 Mammary Neoplasms, Experimental
 Prolactin, 78-5545
 Thyroid Neoplasms
 Radiotherapy, 78-5716
- Hyponitrous Acid**
 Methylamine, α -Acetoxybenzyl-*N*-nitroso-
 Hydrolysis, Methanolysis, 78-5577
- Hypothyroidism**
 Mammary Neoplasms, Experimental
 Prolactin, 78-5545
- Hypoxanthine Phosphoribosyltransferase**
 Carcinogen, Chemical
 Mutation, 78-5652
 Lymphoma
 Methanesulfonic Acid, Ethyl Ester, 78-5652
 Methanesulfonic Acid, Ethyl Ester
 Mutagenic Activity, 78-5657
 Mutation, 78-5652
- Hysterectomy**
 Breast Neoplasms
 Epidemiology, 78-5948
 Cervix Neoplasms
 Epidemiology, 78-5957
- ICR 191**
 Lymphoma
 Mutagenic Activity, S49 Cells, 78-5680
 Thioguanine Resistance, 78-5680
- IgM**
 Virus, Epstein-Barr
 Antibodies, Viral, 78-5829
- 2-Imidazoline, 2-Benzyl-**
 Skin Neoplasms
 Benzo(a)pyrene, 78-5567
 7,8-Benzoflavone, 78-5567
- Immune Serums**
 Leukemia
 Virus, Friend Murine Leukemia, 78-5765
 Sarcoma

Murine Serums (cont'd)

- Virus, Friend Murine Leukemia, 78-5765
- Virus, Friend Murine Leukemia
 - Virus Neutralization, 78-5765
- Virus, Gross Murine Leukemia
 - Antigens, Viral, 78-5769
- Virus, Rauscher Murine Leukemia
 - Antigen-Antibody Complex, 78-5782

Munity

- Slow Virus Diseases
 - Hamster, Review, 78-5440

Neoplasms

- Gastrointestinal Neoplasms
 - Carcinoembryonic Antigen, 78-5867
- Prostatic Neoplasms
 - Carcinoembryonic Antigen, 78-5864
- Virus, SV40
 - Antigens, Viral, 78-5812
 - Virus, Polyoma, BK, 78-5812
 - Virus, Polyoma, JC, 78-5812

Neonoglobulins

- Intestinal Neoplasms
 - Lymphosarcoma, 78-5886

Neonoglobulins, Surface

- Plasmacytoma
 - Immunosuppression, 78-5856
 - B-Lymphocytes, 78-5856
- Virus, Friend Murine Leukemia
 - Lymphoid Tissue, 78-5763

Neonologic Deficiency Syndromes

- Carcinogen, Environmental
 - Risk Factors, Review, 78-5458

Neonoppression

- Cervix Neoplasms
 - Case Report, 78-5888
- Leukemia, Lymphocytic
 - Pregnancy, 78-5921
- Leukemia, Myelocytic
 - Azathioprine, 78-5895
 - Kidney Transplantation, 78-5895
 - T-Lymphocytes, 78-5895
 - Prednisolone, Methyl-, 78-5895
- Neoplasms, Experimental
 - Radiation, Ionizing, 78-5722
- Papilloma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5538
- Plasmacytoma
 - Immunoglobulins, Surface, 78-5856
 - RNA, 78-5856
 - Virus-Like Particles, 78-5856
- Radiation, Ionizing
 - T-Lymphocytes, 78-5722
- Tracheal Neoplasms
 - Growth, 78-5978
 - Transplantation, Homologous, 78-5978
- Ultraviolet Rays
 - T-Lymphocytes, 78-5858
 - Lymphoid Tissue, 78-5858
- Virus, Friend Murine Leukemia
 - Virus Replication, 78-5763

Neonoppressive Agents

- Sarcoma, Kaposi's
 - Epidemiology, 78-5855

Infectious Mononucleosis

- Cortisone
 - Interferon, 78-5824
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-5829
 - Antigens, Viral, 78-5824
 - Cortisone, 78-5824
 - Seroepidemiology, Review, 78-5443

Insulin

- Breast Neoplasms
 - Adenocarcinoma, 78-5950
 - Growth, 78-5950
- Mammary Neoplasms, Experimental
 - Surface Properties, 78-5752

Interferon

- Burkitt's Lymphoma
 - Cortisone, 78-5824
- Infectious Mononucleosis
 - Cortisone, 78-5824

Intestinal Neoplasms

- Adenocarcinoma
 - Bracken Fern, 78-5464, 78-5465
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499
 - Virus, Cytomegalo, 78-5885
- Adenoma
 - Bracken Fern, 78-5464, 78-5465
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499
- Anti-Antibodies
 - Neoplasm Metastasis, 78-5499
- Bracken Fern
 - Carcinogenic Activity, Rat, 78-5464
 - Precancerous Conditions, 78-5464
- Carcinoma, Mucinous
 - Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499
- Ethylene Oxide
 - Occupational Hazard, 78-5969
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Bile Acids and Salts, 78-5684
 - 5 β -Cholan-24-oic Acid, 3 α -Hydroxy-, 78-5684
 - Taurodeoxycholic Acid, 78-5684
- Hydrazine, 1,2-Dimethyl-
 - Dietary Fats, 78-5684
 - Dietary Proteins, 78-5684
 - Microflora, 78-5494
 - Strain Difference, Rat, 78-5494
- Hydrazine, 1,2-Dimethyl-, Dihydrochloride
 - Anti-Antibodies, 78-5499
 - Histological Study, Rat, 78-5499
 - Neoplasm Metastasis, 78-5499
- Lymphosarcoma
 - Bence Jones Protein, 78-5886
 - Immunoglobulins, 78-5886
 - Ultrastructural Study, Rat, 78-5886
 - Virus-Like Particles, 78-5886
- Neoplasms, Multiple Primary
 - Bracken Fern, 78-5465
- Sarcoma
 - Bracken Fern, 78-5464
- Sarcoma, Reticulum Cell
 - Epidemiology, 78-5887
 - Immunohistochemical Study, 78-5887
- Virus, Cytomegalo
 - DNA-RNA Hybridization, 78-5885
 - DNA, Viral, 78-5885

Intestinal Polyps

- Enteritis
 - Epidemiology, 78-5883
- Genetics
 - Virus, Cytomegalo, 78-5885
- Intestine, Large
 - Epidemiology, 78-5883
- Intestine, Small
 - Malabsorption Syndromes, 78-5883
- Malabsorption Syndromes
 - Epidemiology, 78-5883
- Virus, Cytomegalo
 - DNA, Viral, 78-5885

Intestine, Large

- Intestinal Polyps
 - Epidemiology, 78-5883

Intestine, Small

- Intestinal Polyps
 - Malabsorption Syndromes, 78-5883

Intrauterine Devices

- see Contraceptive Devices

Iodine

- Nitrosamines
 - Nitrogen Oxide, 78-5617

Iodine Radioisotopes

- Calcium
 - Serum Levels, Rat, 78-5713
- Diet
 - Thyroid Gland Function, 78-5713
- Thyroid Neoplasms
 - Carcinoma, Papillary, 78-5716
 - Case Report, 78-5716, 78-5718
 - Epidemiology, 78-5718
- Thyrotropin
 - Serum Levels, Rat, 78-5713
 - Thyroid Gland Function, 78-5713
- Thyroxine
 - Diet, 78-5713
 - Serum Levels, Rat, 78-5713

Iron Oxide

- Lung Neoplasms
 - Benzo(a)pyrene, 78-5878
- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-5569, 78-5876

Isoantigens

- Lung Neoplasms
 - Histocompatibility Antigens, 78-5862
 - Immunity, Radioresistant, 78-5862
 - Strain/Tissue Distribution, Mouse, 78-5862
- Virus, Rauscher Murine Leukemia
 - Viral Proteins, 78-5779

Isoleucine

- Neoplasms, Experimental
 - Cholanthrene, 3-Methyl-, 78-5525

Isonicotinic Acid Hydrazide

- Acetyltransferases
 - Metabolism, Review, 78-5407
- Enzyme Activation
 - Carcinogenic Metabolite, Review, 78-5407
- Guanyl Cyclase
 - Enzymatic Activity, 78-5500

Kallikrein-Trypsin Inhibitor

- Cholanthrene, 3-Methyl-
 - Immune Response, Rat, 78-5527
- Sarcoma
 - Cholanthrene, 3-Methyl-, 78-5527

Karyotyping

- Aflatoxin B1
 - Cells, Cultured, 78-5471

Keratoacanthoma

- Skin Neoplasms
 - Virus-Like Particles, 78-5789

Keratoses

- Petroleum
 - Occupational Hazard, 78-5960

Kidney Calculi

- Diethylamine, *N*-Nitroso-
 - Nitric Acid, Ammonium Salt, 78-5615
- Nitrosamines
 - Nitric Acid, Ammonium Salt, 78-5615
- Pyrrolidine, 1-Nitroso-
 - Nitric Acid, Ammonium Salt, 78-5615

Kidney Diseases

- Urogenital Neoplasms
 - Epidemiology, Bulgaria, 78-5961
 - Genetics, 78-5961

Kidney Neoplasms

- Adenocarcinoma
 - Diethylamine, *N*-Nitroso-, 78-5573
 - Ochratoxin A, 78-5468
- Adenoma
 - Aflatoxin B1, 78-5468
 - Diethylamine, *N*-Nitroso-, 78-5573, 78-5574
 - Morpholine, *N*-Nitroso-, 78-5688, 78-5689
 - Ochratoxin A, 78-5468
 - WY-14,643, 78-5574
- Adenoma, Eosinophilic
 - Morpholine, *N*-Nitroso-, 78-5688
- Diethylamine, *N*-Nitroso-
 - Carcinogenic Activity, Cat, Chicken, 78-5573
- Dimethylamine, *N*-Nitroso-
 - DNA Replication, 78-5604
 - Thyroidectomy, 78-5605
- DNA Replication
 - Precancerous Conditions, 78-5604
- Ethylene Oxide
 - Occupational Hazard, 78-5969
- Leiomyosarcoma
 - Neoplasm Metastasis, 78-5912
- Morpholine, *N*-Nitroso-
 - Precancerous Conditions, 78-5688
 - Ultrastructural Study, Rat, 78-5688, 78-5689
- Ochratoxin A
 - Aflatoxin B1, 78-5468

L Cells

- Methanesulfonic Acid, Methyl Ester
 - DNA Repair, 78-5667
- Quinoline, 4-Nitro-, 1-Oxide
 - DNA Repair, 78-5667

Lactate Dehydrogenase Isoenzymes

- Virus, Adeno 6
 - Enzymatic Activity, 78-5847
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5847

itation Disorders

- Pituitary Neoplasms
Ultrastructural Study, 78-5913

abda Phage

- see Coliphages

yngeal Neoplasms

- Age Factors
Epidemiology, USSR, 78-5917
- Carcinogen, Environmental
Epidemiology, 78-5936
Epidemiology, Georgian SSR, 78-5937
Precancerous Conditions, 78-5936
- Carcinoma, Epidermoid
Epidemiology, Georgian SSR, 78-5937
- Diet
Epidemiology, Georgian SSR, 78-5937
- Ethyl Alcohol
Epidemiology, 78-5936, 78-5939
- Fibrosarcoma
Radiation, Ionizing, 78-5711
Radiotherapy, 78-5711
- Radiotherapy
Case Report, 78-5711
- Smoking
Epidemiology, 78-5936, 78-5939

ithins

- Benz(a)anthracene
Membrane Uptake, 78-5533
- Cell Membrane
Hydroxy Radicals, 78-5724
- Pulse Radiolysis
Hydroxy Radicals, 78-5724
- Pyrene
Hydroxy Radicals, 78-5724

omyoma

- Uterine Neoplasms
Genetics, 78-5892

omyosarcoma

- Kidney Neoplasms
Neoplasm Metastasis, 78-5912
- Liver Neoplasms
Case Report, 78-5912
Neoplasm Metastasis, 78-5912

icine

- Neoplasms, Experimental
Cholanthrene, 3-Methyl-, 78-5525

leukemia

- Anemia, Sideroblastic
Case Report, 78-5901
- Cell Membrane
Gangliosides, 78-5987
- Cholera Toxin
Binding Sites, 78-5987
- Chromosome Abnormalities
Trisomy 15, AKR Mice, 78-5909
- Diethylamine, *N*-Nitroso-
p-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Dimethylamine, *N*-Nitroso-
p-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Down's Syndrome
Case Report, 78-5907
Chromosome Aberrations, 78-5907
- Drug Therapy
Case Report, 78-5898

Leukemia (cont'd)

- Epidemiology
Spain, 78-5915
- Hodgkin's Disease
Cyclophosphamide, 78-5898
Neoplasms, Multiple Primary, 78-5910
Prednisolone, 78-5898
Radiation, Ionizing, 78-5898
- Lysergic Acid Diethylamide
Carcinogenic Potential, Review, 78-5422
- Megakaryocytes
Anemia, Sideroblastic, 78-5901
Chromosome Abnormalities, 78-5901
- Methanesulfonic Acid, Ethyl Ester
p-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Methanesulfonic Acid, Methyl Ester
p-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Occupational Hazard
Dose-Response Study, 78-5719
- Radiation, Ionizing
Case Report, 78-5898
p-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
Dose-Response Study, 78-5719, 78-5976
Epidemiology, Japan, 78-5976
Occupational Hazard, 78-5719
- Rubber
Occupational Hazard, Women, 78-5954
- Thorium Dioxide
Epidemiology, Portugal, 78-5973
- Virus, Epstein-Barr
Antibodies, Viral, 78-5829
Cells, Cultured, 78-5831
Serodiagnosis, 78-5829
- Virus, Friend Murine Leukemia
Immune Serums, 78-5765
- Virus, Herpes
Immune Response, Guinea Pig, 78-5815
- Virus, Murine Leukemia
Virus-Like Particles, 78-5757
- Virus, RNA Tumor
Cells, Cultured, 78-5831

Leukemia, Lymphoblastic

- Genetics
Case Report, 78-5902
- Histocompatibility Antigens
Antigen Frequency, Review, 78-5462
- Sex Factors
Epidemiology, Nigeria, 78-5921
- Virus, Epstein-Barr
Antigens, Viral, 78-5827
Receptors, Complement, 78-5827
Receptors, Viral, 78-5827
- Virus, Moloney Murine Sarcoma
DNA-RNA Hybridization, 78-5771
- Virus, Simian Sarcoma
DNA-RNA Hybridization, 78-5771

Leukemia, Lymphocytic

- Anemia, Hemolytic
Epidemiology, Israel, 78-5920
- Breast Neoplasms
Neoplasms, Multiple Primary, 78-5920
- Ethnic Groups
Epidemiology, Israel, 78-5920
- Ethylene Oxide
Epidemiology, 78-5969
Occupational Hazard, 78-5969

Leukemia, Lymphocytic (cont'd)

- Magnesium
 - Adenosine Triphosphate, 78-5981
- Neoplasms, Multiple Primary
 - Agammaglobulinemia, 78-5920
- Occupational Hazard
 - Epidemiology, 78-5922
- Pregnancy
 - Immunosuppression, 78-5921
- Rubber
 - Epidemiology, 78-5922
- Sex Factors
 - Epidemiology, Nigeria, 78-5921
- Skin Neoplasms
 - Neoplasms, Multiple Primary, 78-5920
- Virus, Murine Leukemia
 - Isolation and Characterization, 78-5767

Leukemia, Myeloblastic

- Acetamide
 - Cell Differentiation, 78-5983
- Acetamide, *N,N*-Dimethyl-
 - Cell Differentiation, 78-5983
- Acetamide, *N*-Methyl-
 - Cell Differentiation, 78-5983
- Antigen-Antibody Reactions
 - Cells, Cultured, 78-5865
 - Primate Antisera, 78-5865
- Benzene
 - Epidemiology, Review, 78-5437
- Bone Marrow Cells
 - Cell Differentiation, 78-5900
- Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
 - Case Report, 78-5896
 - Glomerulonephritis, 78-5896
- Cell Differentiation
 - Cell Line, 78-5900
- Cells, Cultured
 - Cell Differentiation, 78-5900
- Eosinophils
 - Colony Stimulating Factor, 78-5984
- Ethylene Oxide
 - Epidemiology, 78-5969
 - Occupational Hazard, 78-5969
- Formamide, *N,N*-Dimethyl-
 - Cell Differentiation, 78-5983
- Formamide, *N*-Methyl-
 - Cell Differentiation, 78-5983
- Genetics
 - Case Report, 78-5902
- Leukemia, Myelocytic
 - Colony Stimulating Factor, 78-5984
- Methane, Sulfinylbis-
 - Cell Differentiation, 78-5983
 - Phagocytosis, 78-5983
- Neutrophils
 - Colony Stimulating Factor, 78-5984
- 2-Piperidone
 - Cell Differentiation, 78-5983
- Sex Factors
 - Epidemiology, Nigeria, 78-5921
- Triethylene Glycol
 - Cell Differentiation, 78-5983
- Virus, Moloney Murine Sarcoma
 - DNA-RNA Hybridization, 78-5771
- Virus, Simian Sarcoma
 - DNA-RNA Hybridization, 78-5771

Leukemia, Myelocytic

- Acetamide, *N*-Fluorene-2-yl-
 - Bone Marrow Cells, 78-5483
 - Histological Study, Rat, 78-5483
- Acetamide, *N,N*-Fluorene-2,7-diylbis-
 - Neoplasm Metastasis, 78-5484
- Azathioprine
 - Immunosuppression, 78-5895
- Blood Platelets
 - Peroxidases, 78-5899
- Bone Marrow Cells
 - Cell Differentiation, 78-5900
- 1,4-Butanediol, Dimethylsulfonate
 - Cystitis, 78-5897
 - Hemorrhage, 78-5897
- Cell Differentiation
 - Cells, Cultured, 78-5985
- Cells, Cultured
 - Cell Differentiation, 78-5900
 - Embryo, 78-5985
- Cholera Toxin
 - Binding Sites, 78-5987
- Chromosome Aberrations
 - Blast Crises, 78-5908
 - Case Report, 78-5906, 78-5908
 - Chromosomes, Human, 6-12, 78-5906
 - Chromosomes, Human, 21-22, 78-5903
 - Trisomy 8, 78-5906
- Chromosomes, Human, 21-22
 - Chromosome Aberrations, 78-5908
 - Gangliosides, 78-5987
 - Kidney Transplantation, 78-5895
 - Transplantation, Heterologous, 78-5904, 78-5986
- Cystitis
 - Hemorrhage, 78-5897
- Drug Therapy
 - Case Report, 78-5898
- Eosinophils
 - Colony Stimulating Factor, 78-5984
- Genetics
 - Case Report, 78-5903
 - Chromosomes, Human, 21-22, 78-5903
- Hematopoietic Stem Cells
 - Cell Differentiation, 78-5904
 - Fibroblasts, 78-5904
- Hodgkin's Disease
 - Cyclophosphamide, 78-5898
 - Radiation, Ionizing, 78-5898
 - Vinblastine, 78-5898
- Immunosuppression
 - Kidney Transplantation, 78-5895
 - T-Lymphocytes, 78-5895
- Kidney Transplantation
 - Case Report, 78-5895
- Leukemia, Myeloblastic
 - Colony Stimulating Factor, 78-5984
- Magnesium
 - Adenosine Triphosphate, 78-5981
- Megakaryocytes
 - Blast Crisis, 78-5899
 - Cytochemical/Ultrastructural Study, 78-5899
- Mesenchymoma
 - Radiation, Ionizing, 78-5898
 - Transplantation, Heterologous, 78-5904
- Neoplasm Metastasis
 - Bone Marrow Cells, 78-5484
 - Trisomy 8, 78-5906

Leukemia, Myelocytic (cont'd)

- Neoplasm Transplantation
- Neoplasm Metastasis, 78-5484
- Occupational Hazard
- Epidemiology, 78-5922
- Peroxidases
- Blast Crisis, 78-5899
- Polycythemia Vera
- Case Report, 78-5905
- Chromosomes, Human, 21-22, 78-5905
- Prednisolone, Methyl-
- Immunosuppression, 78-5895
- Radiation, Ionizing
- Case Report, 78-5898
- Rubber
- Epidemiology, 78-5922
- Sarcoma
- Transplantation, Heterologous, 78-5986
- Sex Factors
- Epidemiology, Nigeria, 78-5921
- Thyroid Neoplasms
- Adenoma, 78-5906
- Transplantation, Heterologous
- DNA Replication, 78-5986
- Mouse, Nude, 78-5904, 78-5986
- Trisomy 8
- Case Report, 78-5906
- Virus, Moloney Murine Sarcoma
- DNA-RNA Hybridization, 78-5771
- Virus, Simian Sarcoma
- DNA-RNA Hybridization, 78-5771

Leukemia, Radiation-Induced

- Radiation, Ionizing
- Occupational Hazard, Review, 78-5720

Neuropeptides

- Bladder Neoplasms
- Papilloma, 78-5583
- Hepatoma
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
- 78-5583

Lipids

- Mammary Neoplasms, Experimental
- Acetate Incorporation, 78-5543

Lithocholic Acid

- see 5 β -Cholan-24-oic Acid, 3 α -Hydroxy-

Liver

- Aflatoxin B1
- DNA, Binding, 78-5474
- Aflatoxin B2
- DNA, Binding, 78-5474
- Dimethylamine, *N*-Nitroso-
- Metabolism, Rat, 78-5603
- Nucleic Acids, Binding, 78-5603
- Plutonium
- Body Burden, Dog, 78-5732, 78-5734

Liver Cirrhosis

- Esophageal Neoplasms
- Ethyl Alcohol, 78-5938
- Ethyl Alcohol
- Epidemiology, Japan, 78-5966
- Hepatoma
- Australia Antigen, 78-5965
- Epidemiology, Ivory Coast, 78-5962
- Liver Neoplasms

Liver Cirrhosis (cont'd)

- Ethyl Alcohol, 78-5938
- Smoking, 78-5938
- Schistosomiasis
- Epidemiology, Japan, 78-5966
- Virus, Hepatitis
- Epidemiology, Greenland, 78-5963

Liver Neoplasms

- Acetamide, *N*-Fluorenyl-2-yl-
- Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5481
- 78-5516
- 5,6-Benzoflavone, 78-5481
- Cholanthrene, 3-Methyl-, 78-5516
- Co-carcinogenic Effect, 78-5516
- Hyperplasia, 78-5481
- 2,4-Oxazolidinedione, 3-(3,5-Dichlorophenyl)-5,5-dimethyl-, 78-5481
- Polychlorobiphenyl Compounds, 78-5481
- Adenocarcinoma
- Ochratoxin A, 78-5468
- Adenoma
- Aflatoxin B1, 78-5468
- Ochratoxin A, 78-5468
- Angiosarcoma
- Ethylene, Chloro-, 78-5485, 78-5486, 78-5967
- Ethylene, Chloro- Polymer, 78-5967
- Carbon Tetrachloride
- Hepatectomy, 78-5592
- Cholangioma
- Diethylamine, *N*-Nitroso-, 78-5575
- Thorium Dioxide, 78-5973
- Diethylamine, 2,2'-Diethoxy-*N*-nitroso-
- Carcinogenic Activity, Rat, 78-5572
- Diethylamine, 2,2'-Dimethoxy-*N*-nitroso-
- Carcinogenic Activity, Rat, 78-5572
- Diethylamine, *N*-Nitroso-
- Carbon Tetrachloride, 78-5592
- Carcinogenic Activity, Rat, 78-5572
- Diazepam, 78-5575
- Hepatectomy, 78-5592
- Precancerous Conditions, 78-5516
- Ultrastructural Study, Gerbil, 78-5575
- Uracil, 2-Thio-, 78-5575
- WY-14,643, 78-5574
- Dimethylamine, *N*-Nitroso-
- Hepatectomy, 78-5592
- Rhodopseudomonas capsulata*, 78-5608
- Thyroidectomy, 78-5605
- Ethnic Groups
- Epidemiology, 78-5938
- Ethylamine, *N*-Methyl-*N*-nitroso-
- Hepatectomy, 78-5592
- Hemangioendothelioma
- Thorium Dioxide, 78-5973
- Hemangioma
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
- Leiomyosarcoma
- Case Report, 78-5912
- Neoplasm Metastasis, 78-5912
- Liver Cirrhosis
- Ethyl Alcohol, 78-5938
- Smoking, 78-5938
- Lymphosarcoma
- Cells, Cultured, 78-5979
- Neoplasms, Multiple Primary
- Case Report, 78-5485
- Ochratoxin A

Liver Neoplasms (cont'd)

- Aflatoxin B1, 78-5468
- Precancerous Conditions
 - Hepatectomy, 78-5592
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
 - Carcinogenic Activity, Rat, 78-5580
- Schistosomiasis
 - Epidemiology, Japan, 78-5966
- Thorium Dioxide
 - Epidemiology, Portugal, 78-5973
- Virus, Hepatitis
 - Epidemiology, Greenland, 78-5963

LSD-25

- see Lysergic Acid Diethylamide

Lung Diseases, Obstructive

- Air Pollution
 - Epidemiology, 78-5934
- Occupational Hazard
 - Epidemiology, 78-5934
- Smoking
 - Epidemiology, 78-5934

Lung Neoplasms

- Adenocarcinoma
 - Animal Model, Hamster, 78-5878
 - Azocine, Octahydro-1-nitroso-, 78-5585
 - Benzo(a)pyrene, 78-5878
 - Californium, 78-5726
 - Classification, 78-5877
 - Hydrazine, 1,2-Diformyl-, 78-5493
 - Nickel, 78-5931
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
 - Ultrastructural/Histochemical Study, 78-5877
- Adenoma
 - Azocine, Octahydro-1-nitroso-, 78-5585
 - Benzo(a)pyrene, 78-5570
 - Diethylamine, *N*-Nitroso-, 78-5665
 - Dimethylamine, *N*-Nitroso-, 78-5570
 - Hydrazine, 1,2-Diformyl-, 78-5493
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
 - Quinoline, 8-Nitro-, 78-5672
 - Radiation, Ionizing, 78-5665
- Air Pollution
 - Epidemiology, 78-5934
 - Epidemiology, Review, 78-5409
- Anthracosilicosis
 - Epidemiology, 78-5971
 - Occupational Hazard, 78-5971
- Asbestos
 - Air Pollution, Review, 78-5409
 - Epidemiology, 78-5928
- Azocine, Octahydro-1-nitroso-
 - Ultrastructural Study, Hamster, 78-5585
- BCG
 - Cell-Wall Skeleton, 78-5532
- Benzene
 - Air Pollution, Review, 78-5409
- Benzo(a)pyrene
 - Iron Oxide, 78-5878
 - Precancerous Conditions, 78-5570
 - Ultrastructural Study, Hamster, 78-5570, 78-5878
- Californium
 - Carcinogenic Activity, Rat, 78-5726
 - Radioisotopes, 78-5726
- Carcinogen, Environmental
 - Epidemiology, Review, 78-5463
- Carcinoid Tumor

Lung Neoplasms (cont'd)

- Benzo(a)pyrene, 78-5878
- Classification, 78-5877
- Carcinoma
 - Nickel, 78-5931
- Carcinoma, Epidermoid
 - Animal Model, Hamster, 78-5878
 - Anthracosilicosis, 78-5971
 - Azocine, Octahydro-1-nitroso-, 78-5585
 - Benzo(a)pyrene, 78-5878
 - Californium, 78-5726
 - Cholanthrene, 3-Methyl-, 78-5528
 - Classification, 78-5877
 - Nickel, 78-5931
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
 - Ultrastructural/Histochemical Study, 78-5877
- Carcinoma, Oat Cell
 - Classification, 78-5877
- Cholanthrene, 3-Methyl-
 - BCG, 78-5532
 - Chromium, 78-5528
 - Nickel, 78-5528
 - Precancerous Conditions, 78-5528
 - Thymectomy, 78-5532
- Chromate
 - Epidemiology, 78-5928
- Chromium
 - Epidemiology, 78-5929
 - Occupational Hazard, 78-5929
- Diethylamine, *N*-Nitroso-
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Dimethylamine, *N*-Nitroso-
 - APUD Cells, 78-5570
 - Metaplasia, 78-5570
 - Precancerous Conditions, 78-5570
 - Thyroidectomy, 78-5605
 - Ultrastructural Study, Hamster, 78-5570
- Epidemiology
 - Spain, 78-5915
- Ether, Bis(chloromethyl)-
 - Air Pollution, Review, 78-5409
 - Epidemiology, Review, 78-5410
- Ethylene, Chloro-
 - Air Pollution, Review, 78-5409
- Fibrosarcoma
 - Benzo(a)pyrene, 78-5878
- Genetics
 - Classification, Review, 78-5455
- Hydrazine, 1,2-Diformyl-
 - Carcinogenic Activity, Mouse, 78-5493
- Isoantigens
 - Histocompatibility Antigens, 78-5862
 - Immunity, Radioresistant, 78-5862
 - Strain/Tissue Distribution, Mouse, 78-5862
- Lymphosarcoma
 - Californium, 78-5726
 - Cells, Cultured, 78-5979
- Neoplasms, Multiple Primary
 - Azocine, Octahydro-1-nitroso-, 78-5585
- Nickel
 - Epidemiology, 78-5931
 - Occupational Hazard, 78-5931
 - Smoking, 78-5931
- Occupational Hazard
 - Epidemiology, 78-5933, 78-5934
- Papilloma
 - Benzo(a)pyrene, 78-5878

Neoplasms (cont'd)

- Plutonium Oxide
 - Carcinogenic Potential, Review, 78-5444
 - Epidemiology, Review, 78-5444
- Pneumoconiosis
 - Epidemiology, Japan, 78-5935
- Precancerous Conditions
 - Radiography, 78-5933
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
 - Carcinogenic Activity, Rat, 78-5580
- Quinoline, 4-Nitro-, 1-Oxide
 - BCG, 78-5532
 - Smoke Condensate, 78-5669
 - Thymectomy, 78-5532
- Radiation, Ionizing
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Rubber
 - Epidemiology, 78-5954
 - Occupational Hazard, Women, 78-5954
- Smoke Condensate
 - Oxidoreductases, 78-5669
- Smoking
 - Epidemiology, 78-5933, 78-5934
 - Epidemiology, Review, 78-5409
- Thorium Dioxide
 - Epidemiology, Portugal, 78-5973
- Tuberculosis, Pulmonary
 - Epidemiology, 78-5930
- Uranium
 - Epidemiology, 78-5928
- Erythematous, Systemic
 - Hodgkin's Disease
 - Case Report, 78-5911
- Lymphoma
 - Case Report, 78-5911

Lymphocyte Transformation

- Virus, Baboon
 - B-Lymphocytes, 78-5825
 - Trans-Species Infection, 78-5825
- Virus, Epstein-Barr
 - Antigens, Viral, 78-5828
 - Interactions, B95-8/P3HR-1 Substrains, 78-5834
 - B-Lymphocytes, 78-5825, 78-5826, 78-5828
 - T-Lymphocytes, 78-5826
 - Trans-Species Infection, 78-5825
- Virus, Herpes Papio
 - Marmoset, 78-5817
 - Radiation, Ionizing, 78-5817
- Virus, Murine Leukemia
 - Thymus Gland, 78-5756
- Virus, Radiation Leukemia
 - Thymus Gland, 78-5756

Lymphocytes

- Ataxia Telangiectasia
 - DNA Repair, 78-5697, 78-5860
 - Plant Agglutinins, 78-5860
- Benzo(a)pyrene
 - Metabolism, Fluorescence Study, 78-5558
- Californium
 - Chromosome Aberrations, 78-5725
- Cell Membrane
 - Gangliosides, 78-5987
- Cholera Toxin
 - Binding Sites, 78-5987
- Cobalt Radioisotopes
 - Chromosome Aberrations, 78-5725

Lymphocytes (cont'd)

- Ethylene, Chloro-
 - Chromosome Aberrations, 78-5489
 - Ethylene, Chloro- Polymer
 - Chromosome Aberrations, 78-5488
 - Mitomycin C
 - Blood Cell Count, 78-5687
 - Chromatids, 78-5686
 - Psoralen, 8-Methoxy-
 - Chromosome Aberrations, 78-5703
 - Quinoline, 4-Nitro-, 1-Oxide
 - Chromosome Abnormalities, 78-5730
 - Radiation, Ionizing
 - Chromosome Aberrations, 78-5691, 78-5692
 - Plant Agglutinins, 78-5691
 - Radium
 - Chromosome Aberrations, 78-5725
 - Ultraviolet Rays
 - Chromosome Aberrations, 78-5703
 - Virus, Epstein-Barr
 - DNA Replication, 78-5834
- ## B-Lymphocytes
- Ethanol, 2-Mercapto-
 - Colony Formation, Review, 78-5439
 - Plasmacytoma
 - Immunoglobulins, Surface, 78-5856
 - Virus, Baboon
 - Lymphocyte Transformation, 78-5825
 - Virus, Epstein-Barr
 - DNA Replication, 78-5826
 - Lymphocyte Transformation, 78-5825, 78-5826, 78-5828
 - Malaria, 78-5441
 - Virus, Herpes
 - Immune Response, Guinea Pig, 78-5815

T-Lymphocytes

- Leukemia, Myelocytic
 - Immunosuppression, 78-5895
- Neoplasms, Experimental
 - Radiation, Ionizing, 78-5722
- Radiation, Ionizing
 - Immunosuppression, 78-5722
- Sarcoma
 - Antilymphocyte Serum, 78-5730
- Thorium Dioxide
 - Chromosome Aberrations, 78-5717
- Ultraviolet Rays
 - Immunosuppression, 78-5858
 - Transplantation Immunology, 78-5858
- Virus, Epstein-Barr
 - Lymphocyte Transformation, 78-5826

Lymphoid Tissue

- Ultraviolet Rays
 - Immunosuppression, 78-5858
- Virus, Friend Murine Leukemia
 - Antigens, Viral, 78-5763
 - Immunoglobulins, Surface, 78-5763
 - Surface Properties, 78-5763
 - Ultrastructural Study, 78-5763
- Virus, Murine Leukemia
 - Virus Replication, 78-5756
- Virus, Radiation Leukemia
 - Virus Replication, 78-5756

Lymphoma (General and Unspecified)

- Carcinogen, Chemical

Lymphoma (General and Unspecified) (cont'd)

- Mutation, 78-5652
- Cell Membrane
 - Gangliosides, 78-5987
- Cholera Toxin
 - Binding Sites, 78-5987
- Colonic Neoplasms
 - Case Report, 78-5882
 - Genetics, 78-5882
 - Neoplasms, Multiple Primary, 78-5882
- Epidemiology
 - Spain, 78-5915
- Guanidine, Nitroso-
 - Mutagenic Activity, S49 Cells, 78-5680
 - Ouabain Resistance, 78-5680
- Hydantoin, 5,5-Diphenyl-
 - Epidemiology, Review, 78-5430
- ICR 191
 - Mutagenic Activity, S49 Cells, 78-5680
 - Thioguanine Resistance, 78-5680
- Immune Response
 - Hamster, Review, 78-5440
- Lupus Erythematosus, Systemic
 - Case Report, 78-5911
- Methanesulfonic Acid, Ethyl Ester
 - Hypoxanthine Phosphoribosyltransferase, 78-5652
 - Mutation, 78-5652, 78-5657
 - Thymidine Kinase, 78-5652
- Thymus Neoplasms
 - Methanesulfonic Acid, Ethyl Ester, 78-5639
 - Urea, Ethyl Nitroso-, 78-5639
 - Urea, Methyl Nitroso-, 78-5639
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-5829
 - Serodiagnosis, 78-5829
- Virus, Herpes Ateles
 - Owl Monkey, 78-5816
- Virus, Marek's Disease Herpes
 - Antigens, Viral, 78-5749
- Virus, Moloney Murine Leukemia
 - Antigens, Viral, 78-5774
 - Virus, Helper, 78-5777
- Virus, Murine Leukemia
 - RNA, Viral, 78-5761
 - Virus-Like Particles, 78-5761
- Virus, Rous Sarcoma
 - Virus, Marek's Disease Herpes, 78-5750

Lymphoma, Histiocytic

see Sarcoma, Reticulum Cell

Lymphosarcoma

- Arsenic
 - Occupational Hazard, 78-5922
- Cells, Cultured
 - Neoplasm Invasiveness, 78-5979
- Intestinal Neoplasms
 - Bence Jones Protein, 78-5886
 - Immunoglobulins, 78-5886
 - Ultrastructural Study, Rat, 78-5886
 - Virus-Like Particles, 78-5886
- Liver Neoplasms
 - Cells, Cultured, 78-5979
- Lung Neoplasms
 - Californium, 78-5726
 - Cells, Cultured, 78-5979
- Occupational Hazard
 - Epidemiology, 78-5922

Lymphosarcoma (cont'd)

- Polychlorobiphenyl Compounds
 - Occupational Hazard, 78-5922
- Rubber
 - Epidemiology, 78-5922
- Ultraviolet Rays
 - Occupational Hazard, 78-5922
- Virus, Abelson Murine Leukemia
 - Cells, Cultured, 78-5979
 - Neoplasm Transplantation, 78-5979
- Virus, Bovine Leukemia
 - DNA, Viral, 78-5788
- Virus, Epstein-Barr
 - Receptors, Viral, 78-5827
- Virus, Feline Leukemia
 - Antibodies, Viral, 78-5785

Lysergic Acid Diethylamide

- Abnormalities
 - Teratogenicity, Review, 78-5422
- Chromosome Abnormalities
 - Lymphocytes, Review, 78-5422
- Dose-Response Study
 - Mutagenic Activity, Review, 78-5422
- Leukemia
 - Carcinogenic Potential, Review, 78-5422

Macrophages

- Benzo(a)pyrene
 - Metabolism, Fluorescence Study, 78-5558
- Cell Transformation, Neoplastic
 - BCG, 78-5997
 - Cells, Cultured, 78-5997
- Multiple Myeloma
 - Colony Formation, Review, 78-5439
- Petroleum
 - Colony Formation, Review, 78-5439

Magnesium

- Leukemia, Lymphocytic
 - Adenosine Triphosphate, 78-5981
- Leukemia, Myelocytic
 - Adenosine Triphosphate, 78-5981

Malabsorption Syndromes

- Intestinal Polyps
 - Epidemiology, 78-5883
 - Intestine, Small, 78-5883

Malaria

- Virus, Epstein-Barr
 - Co-carcinogenic Activity, Review, 78-5441
 - B-Lymphocytes, 78-5441

Malate Dehydrogenase

- Virus, Adeno 6
 - Enzymatic Activity, 78-5847
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5847

Maleimide, N-Ethyl-

see 1*H*-Pyrrole-2,5-dione, 1-Ethyl-

Mammary Neoplasms, Experimental

- Adenocarcinoma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5535
 - 78-5537, 78-5539
 - Prolactin, 78-5753
 - RNA, Viral, 78-5535
 - Testosterone, Propionate, 78-5988
 - Transplantation Immunology, 78-5539

Mammary Neoplasms, Experimental (cont'd)

- Virus, Murine Mammary Tumor, 78-5753
- Adenofibroma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5535
 - 78-5537
- Benz(a)anthracene, 7,12-Dimethyl-
 - Acetate Incorporation, 78-5543
 - Carcinoembryonic Antigen, 78-5539
 - DNA Replication, 78-5537
 - Estrogens, 78-5546
 - Parity, 78-5537
 - Precancerous Conditions, 78-5537
 - Prolactin, 78-5545
 - RNA, Viral, 78-5535
 - Thyroxine, 78-5545
 - Transplantation Immunology, 78-5539
 - Uracil, 6-Propyl-2-thio-, 78-5545
 - Virus, C-Type RNA Tumor, 78-5535
- Carcinoembryonic Antigen
 - Immunization, 78-5539
- Cells, Cultured
 - Multicellular Spheroid Formation, 78-5996
 - Ultrastructural Study, 78-5752
- Coumarin
 - Immune Response, Review, 78-5433
- Dexamethasone
 - Surface Properties, 78-5752
- Enovid
 - Prolactin, 78-5992
- Ergocryptine, 2-Bromo-
 - Precancerous Conditions, 78-5992
- Estradiol
 - Prolactin, 78-5992
- Estrogens
 - Castration, 78-5546
 - Receptors, Hormone, 78-5546
- Growth
 - Androgen Dependency, 78-5988
- Hyperplasia
 - Precancerous Conditions, 78-5992
- Hyperthyroidism
 - Prolactin, 78-5545
- Hypothyroidism
 - Prolactin, 78-5545
- Insulin
 - Surface Properties, 78-5752
- Lipids
 - Acetate Incorporation, 78-5543
- Oxytocin
 - Acetate Incorporation, 78-5543
- Placental Lactogen
 - Precancerous Conditions, 78-5992
- Pregnancy
 - Glutamic Acid, Monosodium Salt, 78-5995
- Progesterone
 - Glutamic Acid, Monosodium Salt, 78-5995
- Prolactin
 - Acetate Incorporation, 78-5543
 - Liver, Binding, 78-5545
 - Precancerous Conditions, 78-5992
 - RNA, Viral, 78-5753
- Somatotropin
 - Acetate Incorporation, 78-5543
- Testosterone, Propionate
 - Growth, 78-5988
 - Neoplasm Transplantation, 78-5988
- Transplantation Immunology

Mammary Neoplasms, Experimental (cont'd)

- Carcinoembryonic Antigen, 78-5539
- Vasopressin
 - Acetate Incorporation, 78-5543
- Virus, C-Type RNA Tumor
 - Isolation and Characterization, 78-5535
- Virus, Murine Mammary Tumor
 - DNA-RNA Hybridization, 78-5751
 - DNA, Viral, 78-5751
 - Horizontal Transmission, 78-5755
 - Immunogenetics, 78-5754
 - Mouse, GH, 78-5751
 - Reverse Transcriptase, 78-5752
 - RNA, Viral, 78-5753
 - Transplantation Immunology, 78-5754
- Medulloblastoma**
 - Child
 - Epidemiology, Denmark, 78-5925
- Megakaryocytes**
 - Leukemia
 - Anemia, Sideroblastic, 78-5901
 - Chromosome Abnormalities, 78-5901
 - Leukemia, Myelocytic
 - Blast Crisis, 78-5899
 - Cytochemical/Ultrastructural Study, 78-5899
- Melanoma**
 - Age Factors
 - Epidemiology, New Jersey, 78-5974
 - Cell Line
 - Ultrastructural Study, Review, 78-5454
 - Chromosomes
 - Tissue Culture, Review, 78-5454
 - Coal Tar
 - Psoriasis, 78-5873
 - Ethylene Oxide
 - Occupational Hazard, 78-5969
 - Eye Neoplasms
 - Genetics, 78-5869
 - Genetics
 - Case Report, 78-5869
 - Geographic Factors
 - Epidemiology, New Jersey, 78-5974
 - Growth
 - Tissue Culture, Review, 78-5454
 - Neoplasms, Multiple Primary
 - Genetics, 78-5869
 - Psoriasis
 - Case Report, 78-5873
 - Ultraviolet Rays
 - Radiotherapy, 78-5873
- Menarche**
 - Breast Neoplasms
 - Epidemiology, Iceland, 78-5951
- Meningioma**
 - Brain Neoplasms
 - Epidemiology, Nigeria, 78-5456
 - Epidemiology, Sweden, 78-5927
 - Epidemiology, Women, 78-5926
 - Nitrous Acid, Sodium Salt, 78-5926
 - Radiation, Ionizing, 78-5926
- Menopause**
 - Breast Neoplasms
 - Epidemiology, 78-5949
 - Estrogens, 78-5949

- Menopause (cont'd)**
Progesterone, 78-5949
- Menstruation**
Breast Neoplasms
Epidemiology, 78-5949
- Mercury, Methyl-**
Teratogens
Monitoring, Review, 78-5405
- Mescalolyl**
Nervous System Neoplasms
Urea, Methyl Nitroso-, 78-5643
- Mesenchymoma**
Anemia, Aplastic
Transplantation, Heterologous, 78-5904
Hematopoietic Stem Cells
Cell Differentiation, 78-5904
Fibroblasts, 78-5904
Leukemia, Myelocytic
Radiation, Ionizing, 78-5898
Transplantation, Heterologous, 78-5904
- Mesothelioma**
Asbestos
Case Report, 78-5875
Pleural Neoplasms
Asbestos, 78-5875
Occupational Hazard, 78-5875
- Metals**
Occupational Hazard
Epidemiology, 78-5924
- Metaplasia**
Lung Neoplasms
Dimethylamine, *N*-Nitroso-, 78-5570
Respiratory Tract Neoplasms
Precancerous Conditions, 78-5876
Stomach Neoplasms
Epidemiology, Costa Rica, 78-5944
Precancerous Conditions, 78-5944
- Methanamine, *N,N*-Dimethyl-**
Dimethylamine, *N*-Nitroso-
Nitrous Acid, Sodium Salt, 78-5611
- Methanamine, *N,N*-Dimethyl-, *N*-Oxide**
Dimethylamine, *N*-Nitroso-
Nitrous Acid, Sodium Salt, 78-5611
- Methane, Azoxy-**
Colonic Neoplasms
Carrageen, 78-5644
- Methane, Dichloro-**
Neoplasms
Occupational Hazard, 78-5968
Nitrosamines
Permeability, Laboratory Gloves, 78-5596
Occupational Hazard
Epidemiology, 78-5968
- Methane, Iodo-**
Thymidine Kinase
Mutagenic Activity, 78-5666
- Methane, Sulfinylbis-**
Leukemia, Myeloblastic
Cell Differentiation, 78-5983
Phagocytosis, 78-5983
- Methanesulfonic Acid, Butyl Ester**
Ames Test
Mutagenic Activity, 78-5659
Lymphoblasts
Mutagenic Activity, 78-5659
- Methanesulfonic Acid, Ethyl Ester**
Ames Test
Mutagenic Activity, 78-5659
Caffeine
Mutagenic Activity, 78-5654
Cells, Cultured
Mutagenic Activity, 78-5660
p-Cresol, 2,6-Di-*tert*-butyl-
Carcinogenic Activity, Mouse, 78-5665
DNA
Alkylation Products, 78-5639
Drosophila melanogaster
Chromosome Aberrations, 78-5660
Mutagenic Activity, 78-5656, 78-5660
Sex Chromosomes, 78-5658
Escherichia coli
DNA Repair, 78-5651
Hypoxanthine Phosphoribosyltransferase
Mutagenic Activity, 78-5657
Mutation, 78-5652
Leukemia
p-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
Lymphoblasts
Mutagenic Activity, 78-5659
Lymphoma
Hypoxanthine Phosphoribosyltransferase, 78-5652
Mutation, 78-5652, 78-5657
Thymidine Kinase, 78-5652
Micrococcus radiodurans
Mutagenic Activity, 78-5701
Mutagenic Activity
Azaguanine Resistance, 78-5655
Lung, Hamster, 78-5655
Ouabain Resistance, 78-5655
Neurospora crassa
Mutagenic Activity, 78-5675
Radiation, Ionizing
DNA Repair, 78-5651
Escherichia coli, 78-5651
Saccharomyces cerevisiae
Mutagenic Activity, 78-5654
Sex Chromosomes
Mutagenic Activity, 78-5658
Thymidine Kinase
Mutagenic Activity, 78-5657, 78-5666
Mutation, 78-5652
Thymus Neoplasms
Lymphoma, 78-5639
- Methanesulfonic Acid, Methyl Ester**
Adenine, 3-Methyl-
Glycoside Hydrolases, 78-5663
Ames Test
Mutagenic Activity, 78-5659
Carcinoma, Epidermoid
DNA Repair, 78-5667
Cells, Cultured
Mutagenic Activity, 78-5660
p-Cresol, 2,6-Di-*tert*-butyl-
Carcinogenic Activity, Mouse, 78-5665
Dose-Response Study
Mutagenic Activity, Mouse, 78-5662

thanesulfonic Acid, Methyl Ester (cont'd)

- Drosophila melanogaster*
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5660, 78-5661
 - Recombination-Deficient Mutants, 78-5661

L Cells

- DNA Repair, 78-5667

Leukemia

- p*-Cresol, 2,6-Di-*tert*-butyl-, 78-5665

Microtus oeconomus

- Mutagenic Activity, 78-5695

Saccharomyces cerevisiae

- Aspergillus nidulans*, 78-5653

Sex Chromosomes

- Microtus oeconomus*, 78-5695

Soy Beans

- Somatic Mosaicism, 78-5664

thanesulfonic Acid, Propyl Ester**Ames Test**

- Mutagenic Activity, 78-5659

Lymphoblasts

- Mutagenic Activity, 78-5659

thionine**Aflatoxin B1**

- Cytochrome P-450, 78-5473
- Oxidoreductases, 78-5473

thomyl, N-Nitroso-**Stomach Neoplasms**

- Carcinoma, 78-5598
- Papilloma, 78-5598

thylamine, α -Acetoxybenzyl-N-nitroso-**Acetamide, N-Propyl-**

- Hydrolysis, Methanolysis, 78-5577

Hyponitrous Acid

- Hydrolysis, Methanolysis, 78-5577

Methylcholanthrene

- see Cholanthrene, 1,2-Dehydro-3-Methyl

thylene Chloride

- see Methane, Dichloro-

thylglycosides**Virus, Adeno 2**

- Glycoproteins, 78-5840

thylglyoxal

- see Propanal, 2-Oxo-

Micrococcus luteus**Endonucleases**

- DNA Repair, 78-5700
 - Isolation and Characterization, 78-5700
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
- Endonucleases, 78-5700
- Ultraviolet Rays
- Endonucleases, 78-5700

Micrococcus radiodurans**Guanidine, 1-Methyl-3-nitro-1-nitroso-**

- Mutagenic Activity, 78-5701
- Tryptophan Auxotrophs, 78-5701

Methanesulfonic Acid, Ethyl Ester

- Mutagenic Activity, 78-5701

2-Oxetanone

- Mutagenic Activity, 78-5701

Radiation, Ionizing

- Mutagenic Activity, 78-5701

Micrococcus radiodurans (cont'd)**Sulfuric Acid, Diethyl Ester**

- Mutagenic Activity, 78-5701

Ultraviolet Rays

- Mutagenic Activity, 78-5701

Microsomes, Liver**Acetamide, N-Fluoren-2-yl-**

- Chromatids, 78-5475
- Hydroxylation, 78-5478
- RNA, Binding, 78-5479

Acetohydroxamic Acid, N-Fluoren-2-yl-

- RNA, Binding, 78-5479

Aflatoxin B1

- Metabolism, Mouse, 78-5510
- Oxidoreductases, 78-5473

Benzene, 4-Allyl-1,2-(methylenedioxy)-

- Carcinogenic Metabolite, 78-5519

Benzo(a)pyrene

- Ames Test, 78-5563
- DNA, Binding, 78-5561
- Polyribonucleotides, Binding, 78-5561

Benzo(a)pyrene, 12b,12c-Dihydro-12b,12c-dimethyl-

- Ames Test, 78-5563

Benzo(a)pyrene 4,5-Oxide

- NADH, NADPH Oxidoreductases, 78-5564

Benzo(b,def)chrysene, 13b,13c-Dihydro-13b,13c-dimethyl-

- Ames Test, 78-5563

Benzo(e)pyrene, 12b,12c-Dihydro-12b,12c-dimethyl-

- Ames Test, 78-5563

Cholanthrene, 7,8-Dihydro-7,8-dihydroxy-3-methyl-

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 9,10-Dihydro-9,10-dihydroxy-3-methyl-

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 11,12-Dihydro-11,12-dihydroxy-3-methyl-

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 1,2-Dihydroxy-3-methyl-

- Carcinogenic Metabolite, 78-5530

Cholanthrene, 3-Methyl-

- Carcinogenic Metabolite, 78-5530

Dibenz(a,h)anthracene

- Mutagenic Activity, 78-5548

Ethylene, Chloro-

- Metabolism, 78-5490

Minocycline**Dimethylamine, N-Nitroso-**

- Ascorbic Acid, 78-5610
- Nitrous Acid, Sodium Salt, 78-5610

Mitochondria, Liver**Aflatoxin B2**

- Adenosine Triphosphatase, 78-5469
- Cell Membrane Permeability, 78-5469
- Cytochrome Oxidase, 78-5469

Aflatoxin G2

- Adenosine Triphosphatase, 78-5469
- Cell Membrane Permeability, 78-5469
- Cytochrome Oxidase, 78-5469

Mitomycin C**Chromatids**

- Cell Cycle Kinetics, 78-5686, 78-5694
- Dose-Response Study, Rabbit, 78-5687
- Lymphocytes, 78-5686

Chromosome Aberrations

- Cell Cycle Kinetics, 78-5694

DNA Repair

- Mutagenic Activity, 78-5426

Mitomycin C (cont'd)

- Dose-Response Study
 - Mutagenic Activity, Mouse, 78-5662
- Lymphocytes
 - Blood Cell Count, 78-5687
- Neutrophils
 - Blood Cell Count, 78-5687
- Saccharomyces cerevisiae*
 - Mutagenic Activity, 78-5654
- Uridine, 5-Bromo-2'-deoxy-
 - Chromatids, 78-5686, 78-5687
 - Chromosome Aberrations, 78-5686

Mitosis

- Bacillus thuringiensis*
 - Endotoxins, 78-5467
 - Exotoxin, 78-5467

Morpholine

- Microorganisms
 - Nitrosation, 78-5612

Morpholine, N-Nitroso-

- Aroclor 1254
 - Ames Test, 78-5517
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Ames Test, 78-5517
 - Mutagenic Activity, Hamster V79 Cells, 78-5593
- 5,6-Benzoflavone
 - Ames Test, 78-5517
- Kidney Neoplasms
 - Adenoma, 78-5688, 78-5689
 - Adenoma, Eosinophilic, 78-5688
 - Precancerous Conditions, 78-5688
 - Ultrastructural Study, Rat, 78-5688, 78-5689

Mouth Neoplasms

- Age Factors
 - Epidemiology, USSR, 78-5917
- Papilloma
 - Urea, Methyl Nitroso-, 78-5645
- Rubber
 - Occupational Hazard, Women, 78-5954
- Virus, Papilloma
 - Epidemiology, Cattle, 78-5790

Mucopolysaccharidosis

- Benz(a)anthracene, 7,12-Dimethyl-
 - Cell Transformation, Neoplastic, 78-5540
 - Co-carcinogenic Effect, 78-5540
 - Fibroblasts, 78-5540
- Ultraviolet Rays
 - Co-carcinogenic Effect, 78-5540

Multiple Myeloma

- Bladder Neoplasms
 - Drug Therapy, 78-5894
- Ethanol, 2-Mercapto-
 - Colony Formation, Review, 78-5439
- Macrophages
 - Colony Formation, Review, 78-5439
- Neoplasms, Multiple Primary
 - Case Report, 78-5894
 - Drug Therapy, Review, 78-5420

Muscle Proteins

- Sarcoma
 - Virus, Rous Sarcoma, 78-5746

Mutagens

- Ames Test

Mutagens (cont'd)

- In Vitro Test Systems, Review, 78-5417
- Ascorbic Acid
 - Feces, 78-5629
- DNA Repair
 - Screening Test, Review, 78-5450
- Enzyme Activation
 - In Vitro Test Systems, Review, 78-5417
- Feces
 - Ames Test, 78-5629
- Nitroso Compounds
 - Ascorbic Acid, 78-5629
 - Feces, 78-5629
- 5H-Pyrido(4,3-b)indole, 3-Amino-1-methyl-
 - Ames Test, 78-5426

Mutation

- Carcinogen, Chemical
 - Carcinogen, Chemical, 78-5652
- Lymphoma
 - Carcinogen, Chemical, 78-5652
 - Methanesulfonic Acid, Ethyl Ester, 78-5652
- Methanesulfonic Acid, Ethyl Ester
 - Hypoxanthine Phosphoribosyltransferase, 78-5652
 - Thymidine Kinase, 78-5652

Myasthenia Gravis

- Thymoma
 - Case Report, 78-5914

Mycotoxins

- Drosophila*
 - Mutagenic Activity, 78-5466
- Escherichia coli*
 - Coliphages, 78-5515
- Fusarium*
 - Mutagenic Activity, 78-5466
 - Toxicity, Mouse, 78-5466

NADH, NADPH Oxidoreductases

- Benzo(a)pyrene 4,5-Oxide
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-5564
 - Cholanthrene, 3-Methyl-, 78-5564
 - Cytochrome P-450, 78-5564
 - Microsomes, Liver, 78-5564
 - Riboflavine, 78-5564

Naphthalene, 2-Bromo-6-methoxy-

- Benz(a)anthracene, 3,9-Dihydroxy-7,12-dimethyl-
 - Synthesis, 78-5544

1-Naphthylamine

- Deoxyguanosine Adducts
 - Computer Model, 78-5502

Nasopharyngeal Neoplasms

- Adenocarcinoma, Papillary
 - Virus, Epstein-Barr, 78-5830
- Carcinoma
 - Virus, Epstein-Barr, 78-5830
- Carcinoma, Epidermoid
 - Virus, Epstein-Barr, 78-5830
- Histocompatibility Antigens
 - Antigen Frequency, Review, 78-5462
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-5830
 - Seroepidemiology, Review, 78-5443

son Syndrome

Pituitary Neoplasms

Adrenalectomy, Review, 78-5457

Neoplasm Metastasis

Brain Neoplasms

Burkitt's Lymphoma, 78-5456

Choriocarcinoma, 78-5456

Epidemiology, Nigeria, 78-5456

Breast Neoplasms

Prolactin, 78-5990

Thyrotropin, 78-5990

Carcinoma, Epidermoid

Cholanthrene, 1,2-Dehydro-3-Methyl-, 78-5523

Immunotherapy, 78-5854

Sheep, 78-5854

Colonic Neoplasms

Corynebacterium parvum, 78-5861

Hydrazine, 1,1-Dimethyl-, 78-5861

Hydrazine, 1,2-Dimethyl-, 78-5497

Gastrointestinal Neoplasms

Carcinoembryonic Antigen, 78-5867

Intestinal Neoplasms

Anti-Antibodies, 78-5499

Hydrazine, 1,2-Dimethyl-, Dihydrochloride, 78-5499

Kidney Neoplasms

Leiomyosarcoma, 78-5912

Leukemia, Myelocytic

Acetamide, *N,N*-Fluorene-2,7-diylbis-, 78-5484

Bone Marrow Cells, 78-5484

Neoplasm Transplantation, 78-5484

Trisomy 8, 78-5906

Liver Neoplasms

Leiomyosarcoma, 78-5912

Spinal Cord Neoplasms

Choriocarcinoma, 78-5456

Neoplasm Regression, Spontaneous

Erythroleukemia

Virus, Rauscher Murine Leukemia, 78-5782

Virus, Rauscher Murine Leukemia

Antigen-Antibody Complex, 78-5782

Neoplasm Transplantation

Leukemia, Myelocytic

Neoplasm Metastasis, 78-5484

Lymphosarcoma

Virus, Abelson Murine Leukemia, 78-5979

Mammary Neoplasms, Experimental

Testosterone, Propionate, 78-5988

Tracheal Neoplasms

Cholanthrene, 3-Methyl-, 78-5521

Uterine Neoplasms

Hydrazine, 1,2-Dimethyl-, 78-5495

Neoplasms (General and Unspecified)

Anesthetics

Epidemiology, Review, 78-5423

Carcinogen, Environmental

Epidemiology, Review, 78-5463

Child

Epidemiology, Review, 78-5459

Genetics, Review, 78-5459

Epidemiology

Sao Paulo, Brazil, 78-5916

Growth Factors

Synthesis, Review, 78-5452

Methane, Dichloro-

Occupational Hazard, 78-5968

Neoplasms (General and Unspecified) (cont'd)

Microwaves

Thermography, 78-5999

Oncogenic Viruses

Isolation and Characterization, Review, 78-5442

Tuberculosis, Pulmonary

Epidemiology, 78-5930

Neoplasms, Connective Tissue

Fibrosarcoma

Cicatrix, 78-5970

Neoplasms, Experimental

Amino Acids

Transplantation Immunology, 78-5525

Chlorine

Carcinogenic Potential, Review, 78-5403

Cholanthrene, 3-Methyl-

Alanine, Phenyl-, 78-5525

Isoleucine, 78-5525

Leucine, 78-5525

Polysaccharides, 78-5857

Purine-6-thiol, 78-5857

Tyrosine, 78-5525

Polysaccharides

Coriolus versicolor, 78-5857

Transplantation Immunology, 78-5857

Purine-6-thiol

Transplantation Immunology, 78-5857

Radiation, Ionizing

Immunosuppression, 78-5722

T-Lymphocytes, 78-5722

Thymectomy, 78-5722

Virus, Adeno 5

Surface Properties, 78-5844

Virus, Epstein-Barr

Animal Model, Mouse, 78-5832

Neoplasms, Multiple Primary

Bladder Neoplasms

Bracken Fern, 78-5465

Carcinoma, Epidermoid

Cicatrix, 78-5970

Cervix Neoplasms

Case Report, 78-5888

Colonic Neoplasms

Lymphoma, 78-5882

Hepatoma

Case Report, 78-5485

Ethylene, Chloro-, 78-5485

Hodgkin's Disease

Carcinoma, Epidermoid, 78-5910

Drug Therapy, 78-5888, 78-5910

Epidemiology, 78-5919

Glioblastoma Multiforme, 78-5910

Leukemia, 78-5910

Prostatic Neoplasms, 78-5910

Radiotherapy, 78-5888, 78-5910

Intestinal Neoplasms

Bracken Fern, 78-5465

Leukemia, Lymphocytic

Agammaglobulinemia, 78-5920

Breast Neoplasms, 78-5920

Skin Neoplasms, 78-5920

Liver Neoplasms

Case Report, 78-5485

Lung Neoplasms

Azocine, Octahydro-1-nitroso-, 78-5585

Melanoma

- Neoplasms, Multiple Primary (cont'd)**
 Genetics, 78-5869
 Multiple Myeloma
 Case Report, 78-5894
 Drug Therapy, Review, 78-5420
 Parathyroid Neoplasms
 Radiation, Ionizing, 78-5714
 Salivary Gland Neoplasms
 Radiation, Ionizing, 78-5714
 Thyroid Neoplasms
 Radiation, Ionizing, 78-5714
 Vulvar Neoplasms
 Warts, 78-5888
- Neoplasms, Radiation-Induced**
 Nuclear Reactors
 Risk Factors, 78-5446
 Plutonium
 Epidemiology, Review, 78-5447
- Neoplastic Endocrine-Like Syndromes**
 Genetics
 Classification, Review, 78-5455
 Paraganglioma, Nonchromaffin
 Case Report, 78-5914
 Thymoma
 Case Report, 78-5914
- Nervous System Neoplasms**
 Fibrosarcoma
 Urea, Methyl Nitroso-, 78-5643
 Ganglioneuroma
 Urea, Methyl Nitroso-, 78-5643
 Genetics
 Classification, Review, 78-5455
 Neurilemmoma
 Urea, Methyl Nitroso-, 78-5643
 Neuroblastoma
 Urea, Methyl Nitroso-, 78-5643
 Neurofibroma
 Urea, Methyl Nitroso-, 78-5643
 Urea, Ethyl Nitroso-
 Enzymatic Activity, 78-5647
 Glucosephosphate Dehydrogenase, 78-5647
 Phosphogluconate Dehydrogenase, 78-5647
 Transketolase, 78-5647
 Transplacental Carcinogenesis, 78-5638
 Urea, Methyl Nitroso-
 Castration, 78-5643
 Dieneol, 78-5643
 Immunization, 78-5643
 Mescasolyl, 78-5643
 Thyroidin, 78-5643
 Transplacental Carcinogenesis, 78-5638
- Neurilemmoma**
 Nervous System Neoplasms
 Urea, Methyl Nitroso-, 78-5643
 Urea, Methyl Nitroso-
 Urea, Ethyl Nitroso-, 78-5638
- Neuroblastoma**
 Nervous System Neoplasms
 Urea, Methyl Nitroso-, 78-5643
- Neurofibroma**
 Nervous System Neoplasms
 Urea, Methyl Nitroso-, 78-5643
- Neurofibromatosis**
 Genetics
- Neurofibromatosis (cont'd)**
 Classification, Review, 78-5455
- Neuroglia**
 Virus, Polyoma, BK
 Cell Transformation, Neoplastic, 78-5794
- Neurospora crassa**
 Acridine Mustard
 Mutagenic Activity, 78-5675
p-Benzoquinone, 2,3,5,6-Tetrakis(1-aziridinyl)-
 Mutagenic Activity, 78-5675
 Dimethylamine, *N*-Nitroso-
 Mutagenic Activity, 78-5675
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Mutagenic Activity, 78-5675
 Methanesulfonic Acid, Ethyl Ester
 Mutagenic Activity, 78-5675
 Quinoline, 4-Nitro-, 1-Oxide
 Mutagenic Activity, 78-5675
 Urea, Methyl Nitroso-
 Mutagenic Activity, 78-5675
- Neutrophils**
 Leukemia, Myeloblastic
 Colony Stimulating Factor, 78-5984
 Mitomycin C
 Blood Cell Count, 78-5687
- Nickel**
 Cholanthrene, 3-Methyl-
 Co-carcinogenic Activity, 78-5528
 Lung Neoplasms
 Adenocarcinoma, 78-5931
 Carcinoma, 78-5931
 Carcinoma, Epidermoid, 78-5931
 Cholanthrene, 3-Methyl-, 78-5528
 Epidemiology, 78-5931
 Occupational Hazard, 78-5931
 Smoking, 78-5931
- Nickel Carbonyl**
 Occupational Hazard
 Epidemiology, Review, 78-5410
- Nickel Sulfide**
 Carcinogenic Activity
 Review, 78-5429
- Nicotine, 1'-Nitroso-1'-demethyl-**
 Cells, Cultured
 Metabolism, Colonic Mucosa, 78-5590
- Nitric Acid**
 Benzo(a)pyrene
 Ames Test, 78-5568
 Diphenylamine
 Nitrosation, Review, 78-5414
 Nitric Acid, Sodium Salt
 Blood/Salivary Concentrations, Monkey, 78-5631
 Gastric Juice, 78-5634
 Saliva, 78-5634
 Nitrosamines
 Food Contamination, Review, 78-5414
 Perylene
 Ames Test, 78-5568
 Vicia faba
 Mutagenic Activity, 78-5635
- Nitric Acid, Ammonium Salt**
 Diethylamine, *N*-Nitroso-
 Kidney Calculi, 78-5615

ric Acid, Ammonium Salt (cont'd)

- Nitrosamines
 - Kidney Calculi, 78-5615
- Pyrrolidine, 1-Nitroso-
 - Kidney Calculi, 78-5615

ric Acid, Potassium Salt

- Diethylamine, *N*-Nitroso-
 - Stomach, Cow, 78-5614
- Dimethylamine, *N*-Nitroso-
 - Stomach, Cow, 78-5614
- Pyrrolidine, 1-Nitroso-
 - Stomach, Cow, 78-5614

ric Acid, Sodium Salt

- Bacteria
 - Mouth, Monkey, 78-5631
- Diethylamine, *N*-Nitroso-
 - Food Contamination, 78-5618
- Dimethylamine, *N*-Nitroso-
 - Food Contamination, 78-5618
- Food Contamination
 - Metabolism, 78-5615
- Nitric Acid
 - Blood/Salivary Concentrations, Monkey, 78-5631
 - Gastric Juice, 78-5634
 - Saliva, 78-5634
- Nitrosamines
 - Food Contamination, 78-5615
- Nitrous Acid
 - Blood/Salivary Concentrations, Monkey, 78-5631
 - Gastric Juice, 78-5634
 - Saliva, 78-5634
- Piperidine, 1-Nitroso-
 - Food Contamination, 78-5618
- Polyethylene Glycol
 - Metabolism, 78-5634
- Pyrrolidine, 1-Nitroso-
 - Food Contamination, 78-5618

trogen Oxide

- Benzo(a)pyrene
 - Ames Test, 78-5568
- Nitrosamines
 - Ethyl Alcohol, 78-5617
 - Hydrogen-Ion Concentration, 78-5617
 - Iodine, 78-5617
- Perylene
 - Ames Test, 78-5568

trosamines

- Acetone
 - Permeability, Laboratory Gloves, 78-5596
- Air Pollution
 - Epidemiology, Review, 78-5409
- Ascorbic Acid
 - Metabolism, Rat, 78-5594
- Bladder Diseases
 - Cystitis, 78-5591
- Bladder Neoplasms
 - Schistosoma haematobium*, 78-5591
 - Urine, 78-5591
- Carcinogen, Environmental
 - Isolation and Characterization, Review, 78-5413
 - Quantitation Method, Review, 78-5416
- Chiral Alcohols
 - Hydrogen Bonding, 78-5625
- Colonic Neoplasms
 - DNA, Binding, 78-5590

Nitrosamines (cont'd)

- Enzyme Activation
 - Structure-Activity Relationship, 78-5625
 - Ethyl Alcohol
 - Permeability, Laboratory Gloves, 78-5596
 - Fluorescence
 - Quantitation Method, 78-5620
 - Food Contamination
 - Concentration Levels, 78-5621
 - Fragmentation and Condensation, 78-5579
 - Quantitation Method, 78-5621, 78-5624
 - Glutamyl Transpeptidase
 - Dose-Response Study, 78-5594
 - Serum Levels, 78-5594
 - Glycogen
 - Hepatotoxicity, 78-5594
 - Hepatotoxicity
 - Dose-Response Study, 78-5594
 - Methane, Dichloro-
 - Permeability, Laboratory Gloves, 78-5596
 - Nitric Acid
 - Food Contamination, Review, 78-5414
 - Nitric Acid, Ammonium Salt
 - Kidney Calculi, 78-5615
 - Nitric Acid, Sodium Salt
 - Food Contamination, 78-5615
 - Nitrogen Oxide
 - Ethyl Alcohol, 78-5617
 - Hydrogen-Ion Concentration, 78-5617
 - Iodine, 78-5617
 - Nitrous Acid, Sodium Salt
 - Food Contamination, 78-5615
 - Solvents, 78-5626
 - Nucleic Acids
 - Carcinogenic Activity, Review, 78-5415
 - Occupational Hazard
 - Epidemiology, Review, 78-5414
 - Phenobarbital-Induced Sleeping Time
 - Dose-Response Study, 78-5594
 - Piperazine, 1-Methyl-
 - Hydrogen-Ion Concentration, 78-5617
 - Potassium Hydroxide
 - Fragmentation and Condensation, 78-5579
 - 2-Propanone, 1,1,1,3,3,3-Hexafluoro-
 - Spectrum Analysis, 78-5619
 - Thermal Energy Analyzer
 - Quantitation Method, 78-5622
 - Thiocyanic Acid
 - Nitrous Acid, 78-5613
 - Ultraviolet Rays
 - Photolysis Products, 78-5620
 - Water
 - Permeability, Laboratory Gloves, 78-5596
 - Water Pollution
 - Water, Deionized, 78-5626
-
- Nitroso Compounds**
 - Carcinogen, Environmental
 - Metabolism, Review, 78-5406
 - Escherichia coli*
 - Coliphages, 78-5515
 - Mutagens
 - Ascorbic Acid, 78-5629
 - Feces, 78-5629
 - Thermal Energy Analyzer
 - Quantitation Method, 78-5627, 78-5628

Nitrosoheptamethyleneimine
see Azocine, Octahydro-1-nitroso-

Nitrosomethylphenylcarbamate
see Carbamic Acid, *N*-Methyl-*N*-nitroso-, Phenyl Ester

Nitrous Acid
Caffeine
 Mutagenic Activity, 78-5654
Duodenal Ulcer
 Gastrectomy, 78-5633
Gastrectomy
 Gastric Juice, 78-5633
Nitric Acid, Sodium Salt
 Blood/Salivary Concentrations, Monkey, 78-5631
 Gastric Juice, 78-5634
 Saliva, 78-5634
Nitrosamines
 Thiocyanic Acid, 78-5613
Pentagastrin
 Gastric Juice, 78-5632
Peptic Ulcer
 Gastric Juice, 78-5632
Saccharomyces cerevisiae
 Mutagenic Activity, 78-5654
Stomach Neoplasms
 Achlorhydria, 78-5633
 Gastrectomy, 78-5633
 Gastric Juice, 78-5632
 Hydrogen-Ion Concentration, 78-5632
 Precancerous Conditions, 78-5633
Vicia faba
 Mutagenic Activity, 78-5635
Virus, SV40
 Temperature Sensitive Mutants, 78-5810

Nitrous Acid, Sodium Salt
Acetamide, *N*-Butyl-
 Nitrosation, 78-5636
Antipyrine, 4-(Dimethylamino)-
 Nitrosation, 78-5636
Brain Neoplasms
 Meningioma, 78-5926
Carbamic Acid, Ethyl-, Ethyl Ester
 Nitrosation, 78-5636
Dimethylamine, *N*-Nitroso-
 Methanamine, *N,N*-Dimethyl-, 78-5611
 Methanamine, *N,N*-Dimethyl-, *N*-Oxide, 78-5611
 Minocycline, 78-5610
 Oxytetracycline, 78-5610
 Phenothiazine, 10-(2-Dimethylamino)propyl)-
 78-5609
 Sludge, 78-5608
 Tetracycline, 78-5610
 Vibramycin, 78-5610
 Water Pollution, 78-5608
Esophageal Neoplasms
 Benzylamine, *N*-Methyl-, 78-5586
Food Contamination
 Metabolism, 78-5615
Nitrosamines
 Food Contamination, 78-5615
 Solvents, 78-5626
Pyrrolidine, 1-Nitroso-
 Catalysis, 78-5630
Stomach Neoplasms
 Urea, 1-Methyl-3-phenyl-, 78-5637
Urea, Ethyl-
 Nitrosation, 78-5636

Nitrous Acid, Sodium Salt (cont'd)
Urea, Hexyl-
 Nitrosation, 78-5636
Urea, Methyl Nitroso-
 Carbamic Acid, Ethyl-, Ethyl Ester, 78-5636
 Stomach, Rat, 78-5636
Urea, 1-Methyl-3-phenyl-
 Carcinogenic Metabolite, 78-5637

Nose Neoplasms
Adenocarcinoma
 Diethylamine, *N*-Nitroso-, 78-5575
Azocine, Octahydro-1-nitroso-
 Carcinogenic Activity, Hamster, 78-5585
Diethylamine, *N*-Nitroso-
 Diazepam, 78-5575
 Ultrastructural Study, Gerbil, 78-5575
Uracil, 2-Thio-, 78-5575

Nuclear Reactors
Neoplasms, Radiation-Induced
 Risk Factors, 78-5446
Risk Factors
 Design Criteria, Review, 78-5446

Nucleic Acids
Acetamide, *N*-Fluoren-2-yl-
 Circadian Rhythm, 78-5482
 Metabolism, Rat Liver, 78-5482
Carcinogen, Chemical
 Carcinogenic Activity, Review, 78-5415
 Carcinogenic Metabolite, Review, 78-5419
 Mutagenic Activity, Review, 78-5415
Nitrosamines
 Carcinogenic Activity, Review, 78-5415
Oncogenic Viruses
 Carcinogenic Activity, Review, 78-5415
Polycyclic Hydrocarbons
 Carcinogenic Activity, Review, 78-5415
Virus, Murine Mammary Tumor
 Cell Transformation, Neoplastic, 78-5753

Nucleoside Deaminases
Virus, Herpes Simplex 1
 Enzymatic Activity, 78-5818
Virus, Herpes Simplex 2
 Enzymatic Activity, 78-5818

Nucleosides
Virus, Herpes Simplex 1
 Oxidoreductases, 78-5818
Virus, Herpes Simplex 2
 Oxidoreductases, 78-5818

Occupational Hazard
Anesthetics
 Epidemiology, Review, 78-5423
Angiosarcoma
 Ethylene, Chloro-, 78-5486, 78-5491
Benzene
 Carcinogenic Potential, Review, 78-5437
 Epidemiology, Review, 78-5410
Bladder Neoplasms
 Epidemiology, Review, 78-5401
Bone Neoplasms
 Radium, 78-5729
Brain Neoplasms
 Carcinogen, Chemical, 78-5926
1,3-Butadiene, 2-Chloro-
 Epidemiology, Review, 78-5410

Occupational Hazard (cont'd)

- 1-Butanamine, *N*-Butyl-*N*-nitroso-
 - Permeability, Laboratory Gloves, 78-5595
 - Permeability, Rubber Gloves, 78-5597
- Butylamine, *N*-Methyl-*N*-nitroso-
 - Permeability, Laboratory Gloves, 78-5596
- Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
 - Mutagenic Activity, Review, 78-5421
- Diethylamine, *N*-Nitroso-
 - Permeability, Laboratory Gloves, 78-5595, 78-5596
 - Permeability, Rubber Gloves, 78-5597
- Dimethylamine, *N*-Nitroso-
 - Permeability, Laboratory Gloves, 78-5595, 78-5596
 - Permeability, Rubber Gloves, 78-5597
- Dipropylamine, *N*-Nitroso-
 - Permeability, Laboratory Gloves, 78-5595, 78-5596
 - Permeability, Rubber Gloves, 78-5597
- Ethane, 1,2-Dichloro-
 - Epidemiology, 78-5969
- Ethanol, 2-Chloro-
 - Epidemiology, 78-5969
- Ether, Bis(2-chloroethyl)-
 - Epidemiology, 78-5969
- Ethylamine, *N*-Methyl-*N*-nitroso-
 - Permeability, Laboratory Gloves, 78-5596
- Ethylene
 - Epidemiology, 78-5969
- Ethylene, Chloro-
 - Chromosome Aberrations, 78-5489
 - Epidemiology, Review, 78-5410
- Ethylene, Chloro- Polymer
 - Chromosome Aberrations, 78-5488
- Ethylene Oxide
 - Epidemiology, 78-5969
- Hodgkin's Disease
 - Benzene, 78-5919
 - Epidemiology, 78-5919
- Intestinal Neoplasms
 - Ethylene Oxide, 78-5969
- Keratosi s
 - Petroleum, 78-5960
- Kidney Neoplasms
 - Ethylene Oxide, 78-5969
- Leukemia
 - Dose-Response Study, 78-5719
 - Radiation, Ionizing, 78-5719
- Leukemia, Lymphocytic
 - Epidemiology, 78-5922
 - Ethylene Oxide, 78-5969
- Leukemia, Myeloblastic
 - Ethylene Oxide, 78-5969
- Leukemia, Myelocytic
 - Epidemiology, 78-5922
- Lung Diseases, Obstructive
 - Epidemiology, 78-5934
- Lung Neoplasms
 - Anthraco silicosis, 78-5971
 - Chromium, 78-5929
 - Epidemiology, 78-5933, 78-5934
 - Nickel, 78-5931
- Lymphosarcoma
 - Arsenic, 78-5922
 - Epidemiology, 78-5922
 - Polychlorobiphenyl Compounds, 78-5922
 - Ultraviolet Rays, 78-5922
- Melanoma
 - Ethylene Oxide, 78-5969

Occupational Hazard (cont'd)

- Metals
 - Epidemiology, 78-5924
 - Methane, Dichloro-
 - Epidemiology, 78-5968
 - Neoplasms
 - Methane, Dichloro-, 78-5968
 - Nickel Carbonyl
 - Epidemiology, Review, 78-5410
 - Nitrosamines
 - Epidemiology, Review, 78-5414
 - Paranasal Sinus Neoplasms
 - Radium, 78-5729
 - Pentylamine, *N*-Methyl-*N*-nitroso-
 - Permeability, Rubber Gloves, 78-5597
 - Piperidine, 1-Nitroso-
 - Permeability, Laboratory Gloves, 78-5595
 - Permeability, Rubber Gloves, 78-5597
 - Pleural Neoplasms
 - Mesothelioma, 78-5875
 - Plutonium
 - Epidemiology, Review, 78-5447
 - Plutonium Oxide
 - Epidemiology, Review, 78-5444
 - Prostatic Neoplasms
 - Benzo(a)pyrene, 78-5960
 - Carcinoma, Epidermoid, 78-5960
 - Pyrrolidine, 1-Nitroso-
 - Permeability, Laboratory Gloves, 78-5595
 - Permeability, Rubber Gloves, 78-5597
 - Radiation, Ionizing
 - Genetics, 78-5719
 - Respiratory Tract Neoplasms
 - Air Pollution, 78-5932
 - Sarcoidosis
 - Epidemiology, 78-5934
 - Skin Neoplasms
 - Epidemiology, 78-5960
 - Stomach Neoplasms
 - Ethylene Oxide, 78-5969
 - Sulfide, Bis(2-chloroethyl)-
 - Epidemiology, Review, 78-5410
- Ochratoxin A**
- Aflatoxin B1
 - Co-carcinogenic Effect, 78-5468
 - Kidney Neoplasms
 - Adenocarcinoma, 78-5468
 - Adenoma, 78-5468
 - Aflatoxin B1, 78-5468
 - Liver Neoplasms
 - Adenocarcinoma, 78-5468
 - Adenoma, 78-5468
 - Aflatoxin B1, 78-5468
- Odontogenic Tumor**
- Urea, Methyl Nitroso-
 - Histological Study, Mouse, 78-5645
- Oligodendroglioma**
- Brain Neoplasms
 - Urea, Ethyl Nitroso-, 78-5649
 - Urea, Methyl Nitroso-, 78-5859
 - Urea, Methyl Nitroso-
 - Urea, Ethyl Nitroso-, 78-5638
- Oncocytoma**
- see Adenoma

Oncogenic Viruses

- Cells, Cultured
 - Isolation and Characterization, Review, 78-5442
- Neoplasms
 - Isolation and Characterization, Review, 78-5442
- Nucleic Acids
 - Carcinogenic Activity, Review, 78-5415

Opisthorchis viverrini

- Cholangioma
 - Dimethylamine, *N*-Nitroso-, 78-5602

Ovalbumin

- Virus, Avian Myeloblastosis
 - Antigens, Viral, 78-5741

Ovarian Neoplasms

- Adenocarcinoma
 - Age Factors, 78-5952
 - Precancerous Conditions, 78-5952
- Contraceptives, Oral
 - Epidemiology, 78-5952
- Cystadenocarcinoma
 - Age Factors, 78-5952
 - Precancerous Conditions, 78-5952
- Ethnic Groups
 - Epidemiology, 78-5952
- Parity
 - Epidemiology, 78-5952

2,4-Oxazolidinedione, 3-(3,5-Dichlorophenyl)-5,5-dimethyl-

- Liver Neoplasms
 - Acetamide, *N*-Fluoren-2-yl-, 78-5481

2-Oxetanone

- Micrococcus radiodurans*
 - Mutagenic Activity, 78-5701
- Thymidine Kinase
 - Mutagenic Activity, 78-5666

Oxidoreductases

- Acetamide, *N*-Fluoren-2-yl-
 - Enzyme Activation, 78-5479
- Aflatoxin B1
 - Choline, 78-5473
 - Folic Acid, 78-5473
 - Methionine, 78-5473
 - Microsomes, Liver, 78-5473
- Lung Neoplasms
 - Smoke Condensate, 78-5669
- Virus, Herpes Simplex 1
 - Enzymatic Activity, 78-5818
 - Nucleosides, 78-5818
- Virus, Herpes Simplex 2
 - Enzymatic Activity, 78-5818
 - Nucleosides, 78-5818

Oxygen

- Urea, Ethyl Nitroso-
 - DNA, Alkylation, 78-5648

Oxygenases

- Benzo(a)pyrene
 - Enzymatic Activity, 78-5554

Oxytetracycline

- Dimethylamine, *N*-Nitroso-
 - Nitrous Acid, Sodium Salt, 78-5610

Oxytocin

- Mammary Neoplasms, Experimental
 - Acetate Incorporation, 78-5543

Ozone

- Atmospheric Levels
 - Denmark, 78-5698

Pactamycin

- Erythroleukemia
 - RNA, Messenger, 78-5764

Pancreatic Neoplasms

- Adenocarcinoma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5534
- Adenosine Cyclic 3',5' Monophosphate
 - Tissue Levels, 78-5534
- Benz(a)anthracene, 7,12-Dimethyl-
 - Adenosine Cyclic 3',5' Monophosphate, 78-5534
 - Guanosine Cyclic 3',5' Monophosphate, 78-5534
- Guanosine Cyclic 3',5' Monophosphate
 - Tissue Levels, 78-5534

Papain

- Virus, Avian Myeloblastosis
 - Antigens, Viral, 78-5741

Papilloma

- Benz(a)anthracene, 7,12-Dimethyl-
 - Immunosuppression, 78-5538
 - Nude/Asplenic Mouse, 78-5538
- Bladder Neoplasms
 - Antipain, 78-5583
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5582, 78-5583
 - Leupeptins, 78-5583
 - Pepstatins, 78-5583
- Lung Neoplasms
 - Benzo(a)pyrene, 78-5878
- Mouth Neoplasms
 - Urea, Methyl Nitroso-, 78-5645
- Skin Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5538
 - Benzo(a)pyrene, 78-5567
 - Virus-Like Particles, 78-5789
- Stomach Neoplasms
 - Bux-Ten, *N*-Nitroso-, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, *o*-Isopropoxyphenyl Ester, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, Phenyl Ester, 78-5598
 - Carbamic Acid, *N*-Methyl-*N*-nitroso-, Trime-thylphenyl Ester, 78-5598
 - Carbofuran, *N*-Nitroso-, 78-5598
 - Diethylamine, 2,2-Dichloro-*N*-nitroso-, 78-5572
 - Methomyl, *N*-Nitroso-, 78-5598
 - Urea, Methyl Nitroso-, 78-5645
 - Urea, 1-Methyl-1-nitroso-3-phenyl-, 78-5637
- Tracheal Neoplasms
 - Cholanthrene, 3-Methyl-, 78-5521
 - Urea, Methyl Nitroso-, 78-5642

Paraganglioma, Nonchromaffin

- Hyperparathyroidism
 - Case Report, 78-5914
- Neoplastic Endocrine-Like Syndromes
 - Case Report, 78-5914

Paranasal Sinus Neoplasms

- Carcinoma
 - Radium, 78-5729
- Radium
 - Epidemiology, 78-5729

nasal Sinus Neoplasms (cont'd)
Occupational Hazard, 78-5729

sites
Retroperitoneal Neoplasms
Sarcoma, 78-5980
thyroid Neoplasms
Adenoma
Case Report, 78-5712
Radiation, Ionizing, 78-5712, 78-5714
Neoplasms, Multiple Primary
Radiation, Ionizing, 78-5714
Radiation, Ionizing
Case Report, 78-5714

ity
Breast Neoplasms
Epidemiology, Iceland, 78-5951
Ovarian Neoplasms
Epidemiology, 78-5952
Uterine Neoplasms
Epidemiology, Japan, 78-5953

mphigus
Sarcoma, Kaposi's
Prednisone, 78-5855

atadecane, 2,6,10,14-Tetramethyl-
Virus, Murine Leukemia
Virus Activation, 78-5758

ntagastrin
Nitrous Acid
Gastric Juice, 78-5632
Thiocyanic Acid
Gastric Juice, 78-5632

ntylamine, N-Methyl-N-nitroso-
Occupational Hazard
Permeability, Rubber Gloves, 78-5597

psstatins
Bladder Neoplasms
Papilloma, 78-5583
Hepatoma
Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
78-5583

ptic Ulcer
Nitrous Acid
Gastric Juice, 78-5632

ptide Hydrolases
Virus, Avian Myeloblastosis
Antigens, Viral, 78-5741

ptides
Cells, Cultured
Prostaglandins, 78-5509
Virus, Herpes Simplex 1
DNA, Viral, 78-5820
Isolation and Characterization, 78-5820
Virus, Recombinant, 78-5820
Virus, Herpes Simplex 2
DNA, Viral, 78-5820
Isolation and Characterization, 78-5820
Virus, Recombinant, 78-5820

eroxidases
Leukemia, Myelocytic
Blast Crisis, 78-5899
Blood Platelets, 78-5899

Perylene
Ames Test
Nitro Derivatives, 78-5568
Nitric Acid
Ames Test, 78-5568
Nitrogen Oxide
Ames Test, 78-5568

Pesticides
Ames Test
Evaluation, Review, 78-5427
Saccharomyces cerevisiae
Aspergillus nidulans, 78-5653

Petroleum
Keratosis
Occupational Hazard, 78-5960
Macrophages
Colony Formation, Review, 78-5439
Skin Neoplasms
Precancerous Conditions, 78-5960

Phagocytosis
Leukemia, Myeloblastic
Methane, Sulfinylbis-, 78-5983

Pharyngeal Neoplasms
Rubber
Occupational Hazard, Women, 78-5954
Virus, Papilloma
Epidemiology, Cattle, 78-5790

Phenol, 2,2'-Methylenebis(3,4,6-trichloro)-
Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
Toxic Contaminant, Review, 78-5421

Phenol, Pentachloro-
Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-
Toxic Contaminant, Review, 78-5421

Phenothiazine, 10-(2-Dimethylamino)propyl-
Dimethylamine, *N*-Nitroso-
Nitrosation Product, 78-5609
Nitrous Acid, Sodium Salt, 78-5609

***o*-Phenylenediamine, 4-Nitro-**
Hair Dyes
Carcinogenic Activity, Review, 78-5425
Mutagenic Activity, Review, 78-5425

***p*-Phenylenediamine, 2-Nitro-**
Hair Dyes
Carcinogenic Activity, Review, 78-5425
Mutagenic Activity, Review, 78-5425

***o*-Phenylenediamine**
Hair Dyes
Mutagenic Activity, Review, 78-5425

Phenytoin
see Hydantoin, 5,5-Diphenyl-

Pheochromocytoma
Diethylamine, *N*-Nitroso-
Carcinogenic Activity, Cat, Chicken, 78-5573

Phorbol 12,13-Didecanoate
Cells, Cultured
Lipids, Deacylation, 78-5509
Prostaglandins, 78-5509
Cortisol
Prostaglandins, 78-5509
Cycloheximide
Prostaglandins, 78-5509

- Phorbol 12,13-Didecanoate (cont'd)**
 5,8,11,14-Eicosatetranoic Acid
 Cell Differentiation, 78-5509
 Prostaglandins
 Kidney, Dog, 78-5509
- Phosphogluconate Dehydrogenase**
 Nervous System Neoplasms
 Urea, Ethyl Nitroso-, 78-5647
- Phosphonoacetic Acid**
 Virus, Herpes Simplex 1
 DNA, Viral, 78-5820
 Virus, Herpes Simplex 2
 DNA, Viral, 78-5820
- Phthalic Anhydride, 4-Methoxy-**
 Benz(a)anthracene, 3,9-Dihydroxy-7,12-dimethyl-
 Synthesis, 78-5544
- Picolinic Acid**
 Virus, SV40
 DNA Replication, 78-5808
 Growth Factors, 78-5808
- Pinealoma**
 Urea, Methyl Nitroso-
 Urea, Ethyl Nitroso-, 78-5638
- Piperazine, 4-Benzoyl-3,5-dimethyl-1-nitroso-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
- Piperazine, 2,5-Dimethyl-1,4-dinitroso-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
- Piperazine, 2,6-Dimethyl-1,4-dinitroso-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
- Piperazine, 1,4-Dinitroso-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
 Soy Beans
 Somatic Mosaicism, 78-5664
- Piperazine, 1,4-Dinitroso-2-methyl-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
- Piperazine, 1-Methyl-**
 Nitrosamines
 Hydrogen-Ion Concentration, 78-5617
- Piperazine, 1-Methyl-4-nitroso-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
 Mutagenic Activity, Hamster V79 Cells, 78-5593
- Piperazine, 1-Nitroso-**
 Aroclor 1254
- Piperazine, 1-Nitroso- (cont'd)**
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
- Piperazine, 2,3,5,6-Tetramethyl-1,4-dinitroso-**
 Aroclor 1254
 Ames Test, 78-5589
 Barbituric Acid, 5-Ethyl-5-phenyl-
 Ames Test, 78-5589
- Piperidine**
 Microorganisms
 Nitrosation, 78-5612
- Piperidine, 1-Nitroso-**
 Acetic Acid, Lead Salt
 Acetoxy Derivatives, 78-5578
 Cells, Cultured
 Metabolism, Colonic Mucosa, 78-5590
 Ethylene, Chloro-
 Permeability, Laboratory Gloves, 78-5595
 Food Contamination
 Degradation, 78-5618
 Nitric Acid, Sodium Salt
 Food Contamination, 78-5618
 Occupational Hazard
 Permeability, Laboratory Gloves, 78-5595
 Permeability, Rubber Gloves, 78-5597
- 2-Piperidone**
 Leukemia, Myeloblastic
 Cell Differentiation, 78-5983
- Pituitary Neoplasms**
 Adenoma, Chromophobe
 Epidemiology, Nigeria, 78-5456
 Cellular Inclusions
 Ultrastructural Study, 78-5913
 Lactation Disorders
 Ultrastructural Study, 78-5913
 Nelson Syndrome
 Adrenalectomy, Review, 78-5457
- Placental Lactogen**
 Mammary Neoplasms, Experimental
 Precancerous Conditions, 78-5992
- Plant Agglutinins**
 Ataxia Telangiectasia
 Lymphocytes, 78-5860
 Radiation, Ionizing
 Lymphocytes, 78-5691
- Plasmacytoma**
 Cell Membrane
 Gangliosides, 78-5987
 Cholera Toxin
 Binding Sites, 78-5987
 Immunoglobulins, Surface
 Immunosuppression, 78-5856
 B-Lymphocytes
 Immunoglobulins, Surface, 78-5856
 RNA
 Immunosuppression, 78-5856
 Poly A, 78-5856
 Virus-Like Particles, 78-5856
 Virus-Like Particles
 A- and C-Type Particles, 78-5852
 Immunosuppression, 78-5856
 Ultrastructural Study, Mouse, 78-5852

Plasmacytoma (cont'd)

- Virus, Murine Leukemia
- Virus Activation, 78-5758

Al Neoplasms**Mesothelioma**

- Asbestos, 78-5875
- Occupational Hazard, 78-5875

Cadmium

- Acetic Acid,
((Carboxymethylimino)bis(ethylenenitrilo))tetra-
ra-
Body Burden, Dog, 78-5731, 78-5733
- Drug Therapy, Cadmium/Zinc Chelates, 78-5731
78-5733

Bone and Bones

- Body Burden, Dog, 78-5732, 78-5734

Chironomus riparius

- Trophic Transfer, 78-5735

Citric Acid, Plutonium Salt

- Body Burden, Dog, 78-5732

Liver

- Body Burden, Dog, 78-5732, 78-5734

Neoplasms, Radiation-Induced

- Epidemiology, Review, 78-5447

Occupational Hazard

- Epidemiology, Review, 78-5447

Water Pollution

- Body Burden, Fish, 78-5735

Plutonium Oxide**Alpha Particles**

- Carcinogenic Potential, Review, 78-5444

Lung Neoplasms

- Carcinogenic Potential, Review, 78-5444
- Epidemiology, Review, 78-5444

Occupational Hazard

- Epidemiology, Review, 78-5444

Mononucleosis**Lung Neoplasms**

- Epidemiology, Japan, 78-5935

A**Plasmacytoma**

- RNA, 78-5856

Virus, Avian Sarcoma

- RNA, Viral, 78-5738

Chlorobiphenyl Compounds**Adenosine Triphosphate**

- Aryl Hydrocarbon Hydroxylases, 78-5559

Aryl Hydrocarbon Hydroxylases

- S-9 Liver Fraction, 78-5559

Liver Neoplasms

- Acetamide, *N*-Fluoren-2-yl-, 78-5481

Lymphosarcoma

- Occupational Hazard, 78-5922

Teratogens

- Monitoring, Review, 78-5405

Cyclic Hydrocarbons**Ames Test**

- Synthetic Fuels, 78-5514

Arene Oxides

- Isomerization, 78-5512

Cells, Cultured

- Azaguanine Resistance, 78-5518
- Mutagenic Activity, 78-5518
- Ouabain Resistance, 78-5518

Polycyclic Hydrocarbons (cont'd)

- Temperature Resistance, 78-5518

Escherichia coli

- Coliphages, 78-5515

Food Contamination

- Smoked Products, 78-5555

Industrial Effluents

- Quantitation Method, 78-5501

Nucleic Acids

- Carcinogenic Activity, Review, 78-5415

Radiation

- Enzyme Activation, 78-5518

Structure-Activity Relationship

- Arene Oxides, 78-5512

Theophylline

- Mutagenic Activity, 78-5518

Polycythemia Vera

- Leukemia, Myelocytic

- Case Report, 78-5905

- Chromosomes, Human, 21-22, 78-5905

Polyethylene

- Ames Test

- Mutagenic Activity, 78-5487

Polyethylene Glycol

- Nitric Acid, Sodium Salt

- Metabolism, 78-5634

Polyp, Lymphoid

- see Lymphoma

Polyps

- Colonic Neoplasms

- Chromosome Abnormalities, 78-5881

- Stomach Neoplasms

- Gastritis, 78-5880

- Tracheal Neoplasms

- Urea, Methyl Nitroso-, 78-5642

Polyribosomes

- Erythroleukemia

- Virus, Friend Murine Leukemia, 78-5764

- Virus, Herpes Simplex 1, 78-5764

- Globin

- RNA, Messenger, 78-5764

- Virus, Herpes Simplex 1

- RNA, Messenger, 78-5764

Polysaccharides

- Neoplasms, Experimental

- Cholanthrene, 3-Methyl-, 78-5857

- Coriulus versicolor*, 78-5857

- Transplantation Immunology, 78-5857

Potassium Hydroxide

- Nitrosamines

- Fragmentation and Condensation, 78-5579

Precancerous Conditions

- Bladder Neoplasms

- Acetamide, *N*-Fluoren-2-yl-, 78-5476

- 1-Butanol, 4-(Butylnitrosamino)-, 78-5583

- Bronchial Neoplasms

- Carcinoma, Epidermoid, 78-5522

- Histological/Ultrastructural Study, Dog, 78-5522

- Cervix Neoplasms

- Contraceptive Devices, 78-5959

- Epidemiology, Review, 78-5461

- Colitis, Ulcerative

- Age Factors, 78-5884

Precancerous Conditions (cont'd)

- Surgery, Operative, 78-5884
- Colonic Neoplasms
 - Chromosome Abnormalities, 78-5881
 - Colitis, Ulcerative, 78-5884
 - Hydrazine, 1,2-Dimethyl-, 78-5497, 78-5498
- Esophageal Neoplasms
 - Benzylamine, *N*-Methyl-, 78-5586
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-5586
- Hepatoma
 - Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer 78-5583
- Intestinal Neoplasms
 - Bracken Fern, 78-5464
- Kidney Neoplasms
 - DNA Replication, 78-5604
 - Morpholine, *N*-Nitroso-, 78-5688
- Laryngeal Neoplasms
 - Carcinogen, Environmental, 78-5936
- Liver Neoplasms
 - Diethylamine, *N*-Nitroso-, 78-5516
 - Hepatectomy, 78-5592
- Lung Neoplasms
 - Benzo(a)pyrene, 78-5570
 - Cholanthrene, 3-Methyl-, 78-5528
 - Dimethylamine, *N*-Nitroso-, 78-5570
 - Radiography, 78-5933
- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5537
 - Ergocryptine, 2-Bromo-, 78-5992
 - Hyperplasia, 78-5992
 - Placental Lactogen, 78-5992
 - Prolactin, 78-5992
- Ovarian Neoplasms
 - Adenocarcinoma, 78-5952
 - Cystadenocarcinoma, 78-5952
- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-5569, 78-5876
 - Metaplasia, 78-5876
 - Ultrastructural/Histochemical Study, Hamster 78-5876
 - Urea, Methyl Nitroso-, 78-5569
- Rhabdomyosarcoma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5542
- Skin Neoplasms
 - Petroleum, 78-5960
- Stomach Neoplasms
 - Metaplasia, 78-5944
 - Nitrous Acid, 78-5633
- Thyroid Neoplasms
 - Radiation, Ionizing, 78-5715, 78-5972
- Tracheal Neoplasms
 - Carcinoma, Epidermoid, 78-5978
 - Urea, Methyl Nitroso-, 78-5640
- Virus, Murine Mammary Tumor
 - RNA, Viral, 78-5753

Prednisolone

- Leukemia
 - Hodgkin's Disease, 78-5898

Prednisolone, Methyl-

- Leukemia, Myelocytic
 - Immunosuppression, 78-5895

Prednisone

- Sarcoma, Kaposi's
 - Arthritis, Rheumatoid, 78-5855
 - Case Report, 78-5855

Prednisone (cont'd)

- Pemphigus, 78-5855

Pregnancy

- Breast
 - Tissue Regression, 78-5891
- Leukemia, Lymphocytic
 - Immunosuppression, 78-5921
- Mammary Neoplasms, Experimental
 - Glutamic Acid, Monosodium Salt, 78-5995

Pristane

- see* Pentadecane, 2,6,10,14-Tetramethyl-

Progesterone

- Breast Diseases
 - Isolation and Characterization, Binding Compone 78-5993
- Breast Neoplasms
 - Menopause, 78-5949
- Mammary Neoplasms, Experimental
 - Glutamic Acid, Monosodium Salt, 78-5995

Prolactin

- Breast Neoplasms
 - Neoplasm Metastasis, 78-5990
 - Protirelin, 78-5990
 - Serum Levels, 78-5990
- Mammary Neoplasms, Experimental
 - Acetate Incorporation, 78-5543
 - Adenocarcinoma, 78-5753
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5545
 - Enovid, 78-5992
 - Estradiol, 78-5992
 - Hyperthyroidism, 78-5545
 - Hypothyroidism, 78-5545
 - Liver, Binding, 78-5545
 - Precancerous Conditions, 78-5992
 - RNA, Viral, 78-5753

Proline, 4-Hydroxy-1-nitroso-

- Food Contamination
 - Quantitation Method, 78-5616

Proline, 1-Nitroso-

- Food Contamination
 - Quantitation Method, 78-5616

Propanal, 2-Oxo-

- Cell Transformation, Neoplastic
 - Electron Acceptors, 78-5998
- Proteins
 - Cell Transformation, Neoplastic, 78-5998

2-Propanol, 1,1'-Iminodi-*N*-nitroso-

- Esophageal Neoplasms
 - Carcinogenic Activity, Rat, 78-5580
- Hepatoma
 - Carcinogenic Activity, Rat, 78-5580
- Liver Neoplasms
 - Carcinogenic Activity, Rat, 78-5580
 - Hemangioma, 78-5580
- Lung Neoplasms
 - Adenocarcinoma, 78-5580
 - Adenoma, 78-5580
 - Carcinogenic Activity, Rat, 78-5580
 - Carcinoma, Epidermoid, 78-5580
- Thyroid Neoplasms
 - Adenoma, 78-5580
 - Carcinogenic Activity, Rat, 78-5580
- Urogenital Neoplasms

panol, 1,1'-Iminodi-N-nitroso- (cont'd)
Carcinogenic Activity, Rat, 78-5580

panone, 1,1,1,3,3,3-Hexafluoro-
Nitrosamines
Spectrum Analysis, 78-5619

penoic Acid, 2-Methyl-, 1,2-Ethanediyl Ester
Escherichia coli
DNA Repair, 78-5449
Salmonella typhimurium
DNA Repair, 78-5449

ionic Acid, 2-(p-Chlorophenoxy)-2-methyl-, Ethyl Ester
Hepatoma
Diethylamine, N-Nitroso-, 78-5574

ionitrile, N-Nitrosoiminodi-
Drinking Water
Carcinogenic Activity, Rat, 78-5572

taglandins
Aflatoxin B1
Cells, Cultured, 78-5509
Benz(a)anthracene, 7,12-Dimethyl-
Cells, Cultured, 78-5509
Benzo(a)pyrene
Cells, Cultured, 78-5509
Breast Neoplasms
Semen, 78-5947
Cells, Cultured
Kidney, Dog, 78-5509
Cholanthrene, 3-Methyl-
Cells, Cultured, 78-5509
Peptides
Cells, Cultured, 78-5509
Phorbol 12,13-Didecanoate
Cells, Cultured, 78-5509
Cortisol, 78-5509
Cycloheximide, 78-5509
Kidney, Dog, 78-5509
12-O-Tetradecanoylphorbol-13-acetate
Cells, Cultured, 78-5509
Cortisol, 78-5509
Cycloheximide, 78-5509
Kidney, Dog, 78-5509

tatic Neoplasms
Aging
Endocrine Glands, Review, 78-5411
Benzo(a)pyrene
Occupational Hazard, 78-5960
Carcinoembryonic Antigen
Antibody Formation, 78-5864
Immunization, 78-5864
Carcinogen, Environmental
Epidemiology, Review, 78-5463
Carcinoma, Epidermoid
Epidemiology, 78-5960
Occupational Hazard, 78-5960
Epidemiology
Spain, 78-5915
Ethyl Alcohol
Ethnic Groups, 78-5938
Hodgkin's Disease
Neoplasms, Multiple Primary, 78-5910
Virus, SV40
Acid Phosphatase, 78-5864
Carcinoembryonic Antigen, 78-5864
Cells, Cultured, 78-5864

Protein Kinase
Cervix Neoplasms
Cholanthrene, 3-Methyl-, 78-5868
Virus, Avian Sarcoma
Cell Transformation, Neoplastic, 78-5737
Enzymatic Activity, 78-5737
Genes, Viral, 78-5737
Temperature Sensitive Mutants, 78-5737

Proteins
Breast Diseases
Cysts, 78-5993
Propanal, 2-Oxo-
Cell Transformation, Neoplastic, 78-5998

Protirelin
Breast Neoplasms
Prolactin, 78-5990
Thyrotropin, 78-5990

PS-K
see Polysaccharides

Psoralen
Ultraviolet Rays
DNA, Review, 78-5434
Mutagenic/Carcinogenic Activity, Review, 78-5434

psoralen methoxy 8
Ultraviolet Rays
Chromosome Aberrations, 78-5703

Psoralen, 8-Methoxy-
Caffeine
Chromosome Aberrations, 78-5703
Cells, Cultured
Mutagenic Activity, 78-5702
Lymphocytes
Chromosome Aberrations, 78-5703
Mutagenic Activity
Dose-Response Study, 78-5702
Ultraviolet Rays
Mutagenic Activity, 78-5702

Psoralen, 4,5',8-Trimethyl-
Ultraviolet Rays
Chromosome Aberrations, 78-5703

Psoriasis
Melanoma
Case Report, 78-5873
Coal Tar, 78-5873

Purine, 2-Amino-6-methoxy-
Dimethylamine, N-Nitroso-
1-Butanamine, N-Butyl-N-Nitroso-, 78-5601
Diethylamine, N-Nitroso-, 78-5601
Dipropylamine, N-Nitroso-, 78-5601
DNA, Alkylation, 78-5600
DNA, Liver, 78-5601, 78-5607
Kidney, Liver, Rat, 78-5600

Purine-6-thiol
Neoplasms, Experimental
Cholanthrene, 3-Methyl-, 78-5857
Transplantation Immunology, 78-5857

Pyrene
Lecithins
Hydroxy Radicals, 78-5724

3,6-Pyridazinedione, 1,2-Dihydro-
Drosophila melanogaster

3,6-Pyridazinedione, 1,2-Dihydro- (cont'd)
Mutagenic Activity, 78-5656

Pyridine, 4-(p-Nitrobenzyl)-

- 1,3-Butadiene, 2-Chloro-
Carcinogenic Metabolite, 78-5492
- Ethylene, 1,1-Dichloro-
Carcinogenic Metabolite, 78-5492
- Ethylene, 1,1-Difluoro-
Carcinogenic Metabolite, 78-5492
- Ethylene Oxide, Chloro-
Carcinogenic Metabolite, 78-5492
- Ethylene, Trichloro-
Carcinogenic Metabolite, 78-5492

5H-Pyrido(4,3-b)indole, 3-Amino-1-methyl-
Mutagens

Ames Test, 78-5426

1H-Pyrrole-2,5-dione, 1-Ethyl-

- Virus, Avian Myeloblastosis
DNA Polymerase, 78-5743
- Ribonuclease, 78-5743

Pyrrolidine, 1-Nitroso-

- Acetic Acid, Lead Salt
Acetoxy Derivatives, 78-5578
- Barbituric Acid, 5-Ethyl-5-phenyl-
Mutagenic Activity, Hamster V79 Cells, 78-5593
- Cells, Cultured
Metabolism, Colonic Mucosa, 78-5590
- o*-Cresol, *p*-Nitroso-
Catalysis, 78-5630
- Ethylene, Chloro-
Permeability, Laboratory Gloves, 78-5595
- Food Contamination
Concentration Levels, 78-5621
- Degradation, 78-5618
- Quantitation Method, 78-5616, 78-5623
- Nitric Acid, Ammonium Salt
Kidney Calculi, 78-5615
- Nitric Acid, Potassium Salt
Stomach, Cow, 78-5614
- Nitric Acid, Sodium Salt
Food Contamination, 78-5618
- Nitrous Acid, Sodium Salt
Catalysis, 78-5630
- Occupational Hazard
Permeability, Laboratory Gloves, 78-5595
- Permeability, Rubber Gloves, 78-5597
- Ultraviolet Rays
Photolysis Products, 78-5620

Quercetin

see Flavone, 3,3',4',5,7-Pentahydroxy-

Quinoline, O-Aminoacyl-4-(hydroxyamino)-, 1-Oxide

- Quinoline, 4-Nitro-, 1-Oxide
Carcinogenic Metabolite, 78-5670
- Quinoline, 4-Nitroso-, 1-Oxide
Carcinogenic Metabolite, 78-5670

Quinoline, 4-(Hydroxyamino)-, 1-Oxide

- Aluminum Chloride
Nucleic Acids, Binding, 78-5674
- Nucleic Acids, Binding
Liver, Lung, Mouse, 78-5674
- Structure-Activity Relationship
Carcinogenic Activity, 78-5668
- DNA Repair, 78-5668

Quinoline, 8-Nitro-

- Digestive System Neoplasms
Carcinoma, Epidermoid, 78-5672
- Dose-Response Study, Rat, 78-5672
- Lung Neoplasms
Adenoma, 78-5672

Quinoline, 4-Nitro-, 1-Oxide

- Amyloidosis
BCG, 78-5532
- Thymectomy, 78-5532
- Caffeine
Mutagenic Activity, 78-5671
- Carcinoma, Epidermoid
DNA Repair, 78-5667
- Cells, Cultured
Cell Transformation, Neoplastic, 78-5671
- Chondrosarcoma
Histological Study, Rabbit, 78-5506
- Chromatids
Cell Cycle Kinetics, 78-5694
- Chromosome Aberrations
Cell Cycle Kinetics, 78-5694
- Chromosome Abnormalities
Lymphocytes, 78-5730
- Chromosomes
DNA Replication, 78-5673
- Escherichia coli*
Mutagenic Activity, 78-5671
- Fibrosarcoma
Carcinogenic Activity, Mouse, 78-5670
- Histological Study, Rabbit, 78-5506
- HeLa Cells
DNA Replication, 78-5673
- L Cells
DNA Repair, 78-5667
- Lung Neoplasms
BCG, 78-5532
- Smoke Condensate, 78-5669
- Thymectomy, 78-5532
- Neurospora crassa*
Mutagenic Activity, 78-5675
- Quinoline, *O*-Aminoacyl-4-(hydroxyamino)-, 1-Oxide
Carcinogenic Metabolite, 78-5670
- Sarcoma, Osteogenic
Histological Study, Rabbit, 78-5506
- Structure-Activity Relationship
Carcinogenic Activity, 78-5668
- DNA Repair, 78-5668

Quinoline, 4-Nitroso-, 1-Oxide

- Escherichia coli*
Mutagenic Activity, 78-5670
- Fibrosarcoma
Carcinogenic Activity, Mouse, 78-5670
- Quinoline, *O*-Aminoacyl-4-(hydroxyamino)-, 1-Oxide
Carcinogenic Metabolite, 78-5670
- Sarcoma 180, Crocker
DNA Adduct, 78-5670

Radiation

- Benz(a)anthracene, 7,12-Dimethyl-
Enzyme Activation, 78-5518
- Mutagenic Activity, 78-5518
- Polycyclic Hydrocarbons
Enzyme Activation, 78-5518
- Thymidine
Mutagenic Activity, 78-5721

ation, Ionizing

- Ames Test
 - Evaluation, Review, 78-5427
- Ataxia Telangiectasia
 - Chromosome Aberrations, 78-5697
 - DNA Repair, 78-5697, 78-5860
- Brain Neoplasms
 - Glioma, 78-5872
 - Meningioma, 78-5926
- Chromatids
 - Cell Cycle Kinetics, 78-5694
- Chromosome Aberrations
 - Cell Cycle Kinetics, 78-5691, 78-5692, 78-5694
- Chromosome Abnormalities
 - Cell Transformation, Neoplastic, 78-5730
 - Dose-Response Study, 78-5730
- Colonic Neoplasms
 - Carcinoma, 78-5710
- p*-Cresol, 2,6-Di-*tert*-butyl-
 - Carcinogenic Activity, Mouse, 78-5665
- DNA
 - Alkaline Sucrose Gradient Sedimentation, 78-5678
 - HeLa Cells, 78-5678
 - Immunoreactivity, Antinucleoside Antibodies 78-5678
- DNA Repair
 - Age Factors, 78-5696
 - Liver/Cerebellum, Mouse, 78-5696
- Drosophila melanogaster*
 - Mutagenic Activity, 78-5661
 - Recombination-Deficient Mutants, 78-5661
- Fibroblasts
 - Cell Transformation, Neoplastic, 78-5730
- Genes
 - Mutagenic Activity, Mouse, 78-5693
- Glioma
 - Case Report, 78-5872
- Glucosephosphate Dehydrogenase
 - Cell Transformation, Neoplastic, 78-5730
- Hereditary Diseases
 - DNA Repair, Review, 78-5451
- Hodgkin's Disease
 - Risk Factors, 78-5910
- Laryngeal Neoplasms
 - Fibrosarcoma, 78-5711
- Leukemia
 - Case Report, 78-5898
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
 - Dose-Response Study, 78-5719, 78-5976
 - Epidemiology, Japan, 78-5976
 - Hodgkin's Disease, 78-5898
 - Occupational Hazard, 78-5719
- Leukemia, Myelocytic
 - Case Report, 78-5898
 - Hodgkin's Disease, 78-5898
 - Mesenchymoma, 78-5898
- Leukemia, Radiation-Induced
 - Occupational Hazard, Review, 78-5720
- Lung Neoplasms
 - Adenoma, 78-5665
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-5665
- Lymphocytes
 - Chromosome Aberrations, 78-5691, 78-5692
 - Plant Agglutinins, 78-5691
- T-Lymphocytes
 - Immunosuppression, 78-5722
- Methanesulfonic Acid, Ethyl Ester

Radiation, Ionizing (cont'd)

- DNA Repair, 78-5651
- Escherichia coli*, 78-5651
- Micrococcus radiodurans*
 - Mutagenic Activity, 78-5701
- Microtus oeconomus*
 - Mutagenic Activity, 78-5695
- Neoplasms, Experimental
 - Immunosuppression, 78-5722
 - T-Lymphocytes, 78-5722
 - Thymectomy, 78-5722
- Occupational Hazard
 - Genetics, 78-5719
- Parathyroid Neoplasms
 - Adenoma, 78-5712, 78-5714
 - Case Report, 78-5714
 - Neoplasms, Multiple Primary, 78-5714
- Rhabdomyosarcoma
 - Cell Differentiation, 78-5690
 - DNA, Metabolism, 78-5690
- Salivary Gland Neoplasms
 - Carcinoma, 78-5714
 - Case Report, 78-5714
 - Neoplasms, Multiple Primary, 78-5714
- Sarcoma
 - Cells, Cultured, 78-5730
 - Cholanthrene, 3-Methyl-, 78-5524
- Sex Chromosomes
 - Microtus oeconomus*, 78-5695
- Skin Neoplasms
 - Dose-Response Study, Rat, 78-5723
 - Protons, 78-5723
- Thyroid Neoplasms
 - Adenocarcinoma, 78-5972
 - Adenocarcinoma, Papillary, 78-5972
 - Carcinoma, Papillary, 78-5715
 - Case Report, 78-5714, 78-5718
 - Epidemiology, 78-5715, 78-5972
 - Neoplasms, Multiple Primary, 78-5714
 - Precancerous Conditions, 78-5715, 78-5972
- Ultraviolet Rays
 - Transplantation Immunology, 78-5858
- Uridine, 2'-Deoxy-5-fluoro-
 - Chromosome Aberrations, 78-5692
- Virus, Gross Murine Leukemia
 - Antigens, Viral, 78-5769
 - Virus Replication, 78-5769
- Virus, Herpes Papio
 - Lymphocyte Transformation, 78-5817
- Virus, Murine Leukemia
 - Fv-1 Restriction, 78-5760
 - Virus Replication, 78-5760

Radioisotopes

- Californium
 - Chromosome Aberrations, 78-5725
- Lung Neoplasms
 - Californium, 78-5726
- Radium
 - Chromosome Aberrations, 78-5725

Radium

- Bone Neoplasms
 - Carcinoma, 78-5729
 - Epidemiology, 78-5729
 - Mastoid, 78-5729
 - Occupational Hazard, 78-5729
- Chromosome Aberrations

Radium (cont'd)

- Lymphocytes, 78-5725
- Paranasal Sinus Neoplasms
 - Carcinoma, 78-5729
 - Epidemiology, 78-5729
 - Occupational Hazard, 78-5729
- Radioisotopes
 - Chromosome Aberrations, 78-5725

Receptors, Hormone

- Dexamethasone
 - Liver, Rat, 78-5576
- Hepatoma
 - Dexamethasone, 78-5576
- Mammary Neoplasms, Experimental
 - Estrogens, 78-5546

Rectal Neoplasms

- Carcinogen, Environmental
 - Epidemiology, Review, 78-5463

Respiratory Tract Neoplasms

- Air Pollution
 - Epidemiology, Houston, 78-5932
 - Occupational Hazard, 78-5932
- Benzo(a)pyrene
 - 2-Furaldehyde, 78-5571
 - Histological Study, Dog, 78-5569
 - Iron Oxide, 78-5569, 78-5876
 - Precancerous Conditions, 78-5569, 78-5876
 - Ultrastructural/Histochemical Study, Hamster 78-5876
 - Urea, Methyl Nitroso-, 78-5569
- Diethylamine, *N*-Nitroso-
 - 2-Furaldehyde, 78-5571
- Ethyl Alcohol
 - Epidemiology, Review, 78-5404
- Metaplasia
 - Precancerous Conditions, 78-5876
- Precancerous Conditions
 - Ultrastructural/Histochemical Study, Hamster 78-5876
- Smoking
 - Epidemiology, Review, 78-5404
- Socioeconomic Factors
 - Epidemiology, Houston, 78-5932
- Urea, Methyl Nitroso-
 - Histological Study, Dog, 78-5569
 - Precancerous Conditions, 78-5569

Retinoblastoma

- Genetics
 - Case Report, 78-5870
 - Chromosomes, Human, 13-15, 78-5870

Retinoic Acid

- Bladder Neoplasms
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-5582
 - Neoplasm Invasiveness, 78-5582

Retinol

- Benzo(a)pyrene
 - DNA, Binding, 78-5557

Retroperitoneal Neoplasms

- Sarcoma
 - Hymenolepis nana*, 78-5980
 - Parasites, 78-5980
 - Rat, 78-5980

Reverse Transcriptase

- Mammary Neoplasms, Experimental
 - Virus, Murine Mammary Tumor, 78-5752
- Virus, Avian Myeloblastosis
 - Amino Acids, 78-5743
- Virus, Moloney Murine Leukemia
 - Temperature Sensitive Mutants, 78-5776
 - Virus Cultivation, 78-5775
- Virus, Murine Leukemia
 - Virus-Like Particles, 78-5761
 - Virus, Moloney Murine Leukemia, 78-5761
- Virus, Rauscher Murine Leukemia
 - Temperature Sensitive Mutants, 78-5776
- Virus, RD-114
 - Antibody Specificity, 78-5850
 - Antigenic Determinants, 78-5850
 - Isolation and Characterization, 78-5850
- Virus, Rous Sarcoma
 - Temperature Sensitive Mutants, 78-5745

Rhabdomyosarcoma

- Antigen-Antibody Reactions
 - Primate Antisera, 78-5865
- Antigens, Leukemia
 - Primate Antisera, 78-5865
- Benzo(a)anthracene, 7,12-Dimethyl-
 - Cell Differentiation, 78-5542
 - Precancerous Conditions, 78-5542
 - Ultrastructural Study, Hamster, 78-5542
- Radiation, Ionizing
 - Cell Differentiation, 78-5690
 - DNA, Metabolism, 78-5690
- Virus, Baboon
 - Antigen-Antibody Reactions, 78-5865
- Virus, Gibbon Ape Lymphoma
 - Antigen-Antibody Reactions, 78-5865
- Virus, Simian Sarcoma
 - Antigen-Antibody Reactions, 78-5865

Rhodopseudomonas capsulata

- Liver Neoplasms
 - Dimethylamine, *N*-Nitroso-, 78-5608

Riboflavine

- Benzo(a)pyrene 4,5-Oxide
 - NADH, NADPH Oxidoreductases, 78-5564

Ribonuclease

- Virus, Avian Myeloblastosis
 - Benzoic Acid, 3,3'-Dithiobis(6-nitro)-, 78-5743
 - Benzoic Acid, *p*-Hydroxymercuri-, 78-5743
 - Benzoic Acid, 2-Nitro-5-thiocyano-, 78-5743
 - 1*H*-Pyrrole-2,5-dione, 1-Ethyl-, 78-5743

RNA

- Plasmacytoma
 - Immunosuppression, 78-5856
 - Poly A, 78-5856
 - Virus-Like Particles, 78-5856

RNA, Messenger

- Erythroleukemia
 - Cycloheximide, 78-5764
 - Pactamycin, 78-5764
 - Sodium Fluoride, 78-5764
 - Virus, Friend Murine Leukemia, 78-5764
 - Virus, Herpes Simplex 1, 78-5764
- Polyribosomes
 - Globin, 78-5764
- Virus, Herpes Simplex 1

A, Messenger (cont'd)

- Polyribosomes, 78-5764
- Ultraviolet Rays, 78-5764
- Virus, SV40
- Antigens, Viral, 78-5800

A, Polymerase

- Virus, Adeno 2
- RNA, Viral, 78-5839

A, Viral

- Lymphoma
 - Virus, Murine Leukemia, 78-5761
- Mammary Neoplasms, Experimental
 - Adenocarcinoma, 78-5535
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5535
 - Prolactin, 78-5753
 - Virus, Murine Mammary Tumor, 78-5753
- Thymus Neoplasms
 - Virus, Feline Leukemia, 78-5787
- Virus, Adeno 2
 - α -Amanitine, 78-5839
 - Cell Transformation, Neoplastic, 78-5839
 - DNA-RNA Hybridization, 78-5839
 - RNA Polymerase, 78-5839
- Virus, Avian Sarcoma
 - DNA-RNA Hybridization, 78-5738
 - Isolation and Characterization, 78-5738
 - Nucleotide Sequence, 78-5738
 - Poly A, 78-5738
 - Transformation-Defective Mutants, 78-5738
- Virus, C-Type RNA Tumor
 - DNA-RNA Hybridization, 78-5814
 - Replication-Defective Viruses, 78-5772
 - Virus, Helper, 78-5772
- Virus, D-Type RNA Tumor
 - DNA-RNA Hybridization, 78-5814
- Virus, Feline Leukemia
 - Cap Structures, 78-5787
 - DNA-RNA Hybridization, 78-5787
 - Isolation and Characterization, 78-5787
- Virus, Friend Spleen Focus-Forming
 - Virus, Helper, 78-5768
- Virus, Moloney Murine Leukemia
 - DNA-RNA Hybridization, 78-5770
- Virus, Moloney Murine Sarcoma
 - DNA-RNA Hybridization, 78-5770
- Virus, Moloney Murine Sarcoma-Leukemia
 - DNA-RNA Hybridization, 78-5770
 - Virus Replication, 78-5770
- Virus, Murine Leukemia
 - DNA-RNA Hybridization, 78-5761
 - Thymus Gland, 78-5761
- Virus, Murine Mammary Tumor
 - DNA-RNA Hybridization, 78-5753
 - Precancerous Conditions, 78-5753
- Virus, Ring-Necked Pheasant
 - DNA-RNA Hybridization, 78-5747
- Virus, SV40
 - DNA-RNA Hybridization, 78-5806

Rubber

- Breast Neoplasms
 - Epidemiology, 78-5954
 - Occupational Hazard, Women, 78-5954
- Hodgkin's Disease
 - Occupational Hazard, Women, 78-5954
- Leukemia
 - Occupational Hazard, Women, 78-5954

Rubber (cont'd)

- Leukemia, Lymphocytic
 - Epidemiology, 78-5922
- Leukemia, Myelocytic
 - Epidemiology, 78-5922
- Lung Neoplasms
 - Epidemiology, 78-5954
 - Occupational Hazard, Women, 78-5954
- Lymphosarcoma
 - Epidemiology, 78-5922
- Mouth Neoplasms
 - Occupational Hazard, Women, 78-5954
- Pharyngeal Neoplasms
 - Occupational Hazard, Women, 78-5954

Saccharomyces cerevisiae

- Aniline, 4-Fluoro-
 - Mutagenic Activity, 78-5654
- 4-Cyclohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-
 - Aspergillus nidulans*, 78-5653
- 1-Hexanamine, *N*-Hexyl-*N*-nitroso-
 - Nitrosation, 78-5612
- Methanesulfonic Acid, Ethyl Ester
 - Mutagenic Activity, 78-5654
- Methanesulfonic Acid, Methyl Ester
 - Aspergillus nidulans*, 78-5653
- Mitomycin C
 - Mutagenic Activity, 78-5654
- Nitrous Acid
 - Mutagenic Activity, 78-5654
- Pesticides
 - Aspergillus nidulans*, 78-5653

Saccharomycopsis lipolytica

- 1-Hexanamine, *N*-Hexyl-*N*-nitroso-
 - Nitrosation, 78-5612

Safrole

- see* Benzene, 4-Allyl-1,2-(methylenedioxy)-

Saliva

- Nitric Acid, Sodium Salt
 - Nitric Acid, 78-5634
 - Nitrous Acid, 78-5634
- Virus, Epstein-Barr
 - Horizontal Transmission, Infant, 78-5835

Salivary Gland Neoplasms

- Age Factors
 - Epidemiology, USSR, 78-5917
- Carcinoma
 - Radiation, Ionizing, 78-5714
- Neoplasms, Multiple Primary
 - Radiation, Ionizing, 78-5714
- Radiation, Ionizing
 - Case Report, 78-5714

Salmonella typhimurium

- Butyrophene, γ -Chloro-*p*-fluoro-
 - DNA Repair, 78-5449
- Dibenz(a,h)anthracene
 - Mutagenic Activity, 78-5548
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Histidine Reversion, 78-5681
- 2-Propenoic Acid, 2-Methyl-, 1,2-Ethanediy Ester
 - DNA Repair, 78-5449

Sarcoidosis

- Occupational Hazard
 - Epidemiology, 78-5934

Sarcoma

- Benz(a)anthracene, 7,12-Dimethyl-
 - Carcinogenic Potential, Rat, 78-5547
- Benz(a)anthracene-7,12-dione, 5-Fluoro-
 - Carcinogenic Potential, Rat, 78-5547
- Benz(a)anthracene-7-methanol, 5-Fluoro-12-methyl-
 - Carcinogenic Potential, Rat, 78-5547
- Benz(a)anthracene-7-methanol, 5-Fluoro-12-methyl-, Acetate
 - Carcinogenic Potential, Rat, 78-5547
- Carcinogen, Chemical
 - Antigens, Review, 78-5438
 - Antigens, Viral, 78-5438
 - Carcinoembryonic Antigen, 78-5438
 - Histocompatibility Antigens, 78-5438
- Cell Line
 - Ultrastructural Study, Review, 78-5454
- Cholanthrene, 3-Methyl-
 - Cyclophosphamide, 78-5524
 - Fetal Globulins, 78-5531
 - Kallikrein-Trypsin Inhibitor, 78-5527
 - Radiation, Ionizing, 78-5524
 - Transplantation, Heterologous, 78-5529
- Chromosomes
 - Tissue Culture, Review, 78-5454
- Cyclophosphamide
 - Tumor Latency, 78-5524
- Fetal Globulins
 - Antigen-Antibody Reactions, 78-5531
 - Transplantation Immunology, 78-5531
- Growth
 - Tissue Culture, Review, 78-5454
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Cells, Cultured, 78-5507
- Intestinal Neoplasms
 - Bracken Fern, 78-5464
- Leukemia, Myelocytic
 - Transplantation, Heterologous, 78-5986
- T-Lymphocytes
 - Antilymphocyte Serum, 78-5730
- Radiation, Ionizing
 - Cells, Cultured, 78-5730
- Retroperitoneal Neoplasms
 - Hymenolepis nana*, 78-5980
 - Parasites, 78-5980
 - Rat, 78-5980
- Uterine Neoplasms
 - Hydrazine, 1,2-Dimethyl-, 78-5495
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-5847
 - Hexokinase, 78-5847
- Virus, Friend Murine Leukemia
 - Immune Serums, 78-5765
- Virus, Murine Leukemia
 - Isolation and Characterization, 78-5767
- Virus, Rous Sarcoma
 - Albumins, 78-5746
 - Muscle Proteins, 78-5746
 - Virus, Marek's Disease Herpes, 78-5750
- Sarcoma 180, Crocker**
 - Quinoline, 4-Nitroso-, 1-Oxide
 - DNA Adduct, 78-5670
- Sarcoma, Kaposi's**
 - Arthritis, Rheumatoid
 - Prednisone, 78-5855
 - Immunosuppressive Agents

Sarcoma, Kaposi's (cont'd)

- Epidemiology, 78-5855
- Pemphigus
 - Prednisone, 78-5855
- Prednisone
 - Case Report, 78-5855
- Sarcoma, Mast Cell**
 - Actin
 - Cells, Cultured, 78-5863
 - Histocompatibility Antigens, 78-5863
- Sarcoma, Osteogenic**
 - BCG
 - Immune Response, Mouse, 78-5728
 - Benz(a)anthracene, 7,12-Dimethyl-
 - Cell Transformation, Neoplastic, 78-5507
 - Histological Study, Rabbit, 78-5506
 - Tumorigenicity, Clone Cells, 78-5507
 - Brain Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5507
 - Cholanthrene, 3-Methyl-
 - Histological Study, Rabbit, 78-5506
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Cell Transformation, Neoplastic, 78-5507
 - Cells, Cultured, 78-5507
 - Tumorigenicity, Clone Cells, 78-5507
 - Quinoline, 4-Nitro-, 1-Oxide
 - Histological Study, Rabbit, 78-5506
 - Strontium Radioisotopes
 - BCG, 78-5728
- Sarcoma, Reticulum Cell**
 - Intestinal Neoplasms
 - Epidemiology, 78-5887
 - Immunohistochemical Study, 78-5887
 - Virus, Adeno 12
 - Enzymatic Activity, 78-5847
 - Hexokinase, 78-5847
 - Virus, Radiation Leukemia
 - Blood Cells, Leukemogenic Activity, 78-5853
- Sarcosine, N-Nitroso-**
 - Food Contamination
 - Quantitation Method, 78-5616
- Schistosoma haematobium**
 - Bladder Neoplasms
 - Dimethylamine, N-Nitroso-, 78-5591
 - Nitrosamines, 78-5591
- Schistosomiasis**
 - Liver Cirrhosis
 - Epidemiology, Japan, 78-5966
 - Liver Neoplasms
 - Epidemiology, Japan, 78-5966
- Selenium**
 - Dimethylamine, N-Nitroso-
 - Aminotransferases, 78-5606
 - Hepatotoxicity, Rat, 78-5606
- Semen**
 - Breast Neoplasms
 - Prostaglandins, 78-5947
- Serum Albumin, Bovine**
 - Virus, Avian Myeloblastosis
 - Antigens, Viral, 78-5741
- Sex Chromosomes**
 - Cell Differentiation

Chromosomes (cont'd)

- Mouse Embryo, Review, 78-5453
- Methanesulfonic Acid, Ethyl Ester
 - Drosophila melanogaster*, 78-5658
 - Mutagenic Activity, 78-5658
- Methanesulfonic Acid, Methyl Ester
 - Microtus oeconomus*, 78-5695
- Radiation, Ionizing
 - Microtus oeconomus*, 78-5695

c Acid

- Virus, Rauscher Murine Leukemia
- Antigens, 78-5782

a

- Benz(a)anthracene
 - Adsorption, 78-5533
 - Co-carcinogenic Effect, 78-5533
 - Membrane Uptake, 78-5533
- Benzo(a)pyrene
 - Adsorption, 78-5556

- 525A
 - see Valeric Acid, 2,2-Diphenyl-, 2-(Diethylamino)ethyl Ester, HCl

Neoplasms

- Benzo(a)pyrene
 - 7,8-Benzoflavone, 78-5567
 - Flavone, 3,3',4',5,7-Pentahydroxy-, 78-5567
 - 2-Imidazoline, 2-Benzyl-, 78-5567
 - 12-O-Tetradecanoylphorbol-13-acetate, 78-5567
- Benzo(a)pyrene-6-methanol
 - 7,8-Benzoflavone, 78-5567
 - 12-O-Tetradecanoylphorbol-13-acetate, 78-5567
- Benzo(a)pyrene, 6-Methyl-
 - 7,8-Benzoflavone, 78-5567
 - 12-O-Tetradecanoylphorbol-13-acetate, 78-5567
- 7,8-Benzoflavone
 - 2-Imidazoline, 2-Benzyl-, 78-5567
- Carcinoma, Epidermoid
 - Cicatrix, 78-5970
 - Virus-Like Particles, 78-5789
- Cholanthren-1-ol, 3-Methyl-
 - Carcinogenic Activity, Mouse, 78-5523
- Cholanthren-2-ol, 3-Methyl-
 - Carcinogenic Activity, Mouse, 78-5523
- Cholanthren-1-one, 3-Methyl-
 - Carcinogenic Activity, Mouse, 78-5523
- Cholanthren-2-one, 3-Methyl-
 - Carcinogenic Activity, Mouse, 78-5523
- Cholanthrene, 1,2-Dehydro-3-Methyl-
 - Carcinogenic Activity, Mouse, 78-5523
- Cholanthrene, 3-Methyl-
 - Carcinogenic Activity, Mouse, 78-5523
- Guanidine, 1-Ethyl-3-nitro-1-nitroso-
 - Carcinogenic Activity, Mouse, 78-5598
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Carcinogenic Activity, Mouse, 78-5598
- Keratoacanthoma
 - Virus-Like Particles, 78-5789
- Leukemia, Lymphocytic
 - Neoplasms, Multiple Primary, 78-5920
- Occupational Hazard
 - Epidemiology, 78-5960
- Papilloma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5538
 - Benzo(a)pyrene, 78-5567
 - Virus-Like Particles, 78-5789

Skin Neoplasms (cont'd)

- Petroleum
 - Precancerous Conditions, 78-5960
- Radiation, Ionizing
 - Dose-Response Study, Rat, 78-5723
 - Protons, 78-5723
- Ultraviolet Rays
 - Dose-Response, Mathematical Model, 78-5975
 - Epidemiology, 78-5975
 - Ozone Depletion, 78-5698, 78-5975
- Urea, Methyl Nitroso-
 - Carcinogenic Activity, Mouse, 78-5598
- Virus-Like Particles
 - Ultrastructural/Immunofluorescent Study, Mouse 78-5789
- Virus, Papilloma
 - Virus-Like Particles, 78-5789

Smoke Condensate

- Lung Neoplasms
 - Oxidoreductases, 78-5669
 - Quinoline, 4-Nitro-, 1-Oxide, 78-5669

Smoking

- Air Pollution
 - Co-carcinogenic Activity, 78-5932
- Asbestos
 - Co-carcinogenic Effect, 78-5928
- Bronchitis
 - Carcinoembryonic Antigen, 78-5940
- Carcinoembryonic Antigen
 - Serum Levels, 78-5940
- Carcinogen, Environmental
 - Epidemiology, Review, 78-5402
- Chromate
 - Co-carcinogenic Effect, 78-5928
- Esophageal Neoplasms
 - Epidemiology, 78-5939
 - Epidemiology, Review, 78-5404
- Ethnic Groups
 - Epidemiology, 78-5938
- Ethyl Alcohol
 - Co-carcinogenic Effect, Review, 78-5404
- Laryngeal Neoplasms
 - Epidemiology, 78-5936, 78-5939
- Liver Neoplasms
 - Liver Cirrhosis, 78-5938
- Lung Diseases, Obstructive
 - Epidemiology, 78-5934
- Lung Neoplasms
 - Epidemiology, 78-5933, 78-5934
 - Epidemiology, Review, 78-5409
 - Nickel, 78-5931
- Respiratory Tract Neoplasms
 - Epidemiology, Review, 78-5404
- Thiocyanic Acid
 - Gastric Juice, 78-5632
 - Saliva, 78-5632
- Uranium
 - Co-carcinogenic Effect, 78-5928

Sodium Alginate

- see Alginic Acid, Sodium Salt

Sodium Azide

- Urea, Methyl Nitroso-
 - DNA Repair, 78-5641

Sodium Fluoride

- Erythroleukemia
- RNA, Messenger, 78-5764

Somatomedin

- Fibrosarcoma
- Growth Factors, Review, 78-5452

Somatotropin

- Dimethylamine, *N*-Nitroso-
- DNA, Liver, 78-5607
- Mammary Neoplasms, Experimental
- Acetate Incorporation, 78-5543

Spermine

- Urea, Methyl Nitroso-
- DNA, Methylation, 78-5646

Spinal Cord Neoplasms

- Choriocarcinoma
- Neoplasm Metastasis, 78-5456

Staphylococcus aureus

- Virus, Epstein-Barr
- DNA Replication, 78-5834

Stomach Neoplasms

- Achlorhydria
- Nitrous Acid, 78-5633
- Adenocarcinoma
- Epidemiology, 78-5942
- Gastritis, 78-5880
- Blood Groups
- Epidemiology, 78-5942
- Carcinoma
- Bux-Ten, *N*-Nitroso-, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, *o*-Isopropoxyphenyl Ester, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, Phenyl Ester, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, Trime-thylphenyl Ester, 78-5598
- Carbofuran, *N*-Nitroso-, 78-5598
- Gastrectomy, 78-5879
- Methomyl, *N*-Nitroso-, 78-5598
- Neoplasm Recurrence, 78-5879
- Carcinoma, Epidermoid
- Urea, 1-Methyl-1-nitroso-3-phenyl-, 78-5637
- Diet
- Epidemiology, 78-5942
- Ethylene Oxide
- Epidemiology, 78-5969
- Occupational Hazard, 78-5969
- Gastrectomy
- Neoplasm Recurrence, 78-5879
- Nitrous Acid, 78-5633
- Gastric Juice
- Hydrogen-Ion Concentration, 78-5632
- Gastritis
- Case Report, 78-5880
- Polyps, 78-5880
- Metaplasia
- Epidemiology, Costa Rica, 78-5944
- Precancerous Conditions, 78-5944
- Nitrous Acid
- Gastric Juice, 78-5632
- Hydrogen-Ion Concentration, 78-5632
- Precancerous Conditions, 78-5633

Stomach Neoplasms (cont'd)**Papilloma**

- Bux-Ten, *N*-Nitroso-, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, *o*-Isopropoxyphenyl Ester, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, Phenyl Ester, 78-5598
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, Trime-thylphenyl Ester, 78-5598
- Carbofuran, *N*-Nitroso-, 78-5598
- Diethylamine, 2,2-Dichloro-*N*-nitroso-, 78-5572
- Methomyl, *N*-Nitroso-, 78-5598
- Urea, Methyl Nitroso-, 78-5645
- Urea, 1-Methyl-1-nitroso-3-phenyl-, 78-5637

Talc

- Epidemiology, Hawaii, Japan, 78-5943
- Food Contamination, 78-5943
- Urea, 1-Methyl-1-nitroso-3-phenyl-
- Carcinogenic Activity, Rat, 78-5637
- Urea, 1-Methyl-3-phenyl-
- Nitrous Acid, Sodium Salt, 78-5637

Streptomycin

- Guanidine, 1-Methyl-3-nitro-1-nitroso-
- Lethality, Algae, 78-5699
- Ultraviolet Rays
- Lethality, Algae, 78-5699

Stress

- Antipyrine
- Enzyme Activation, 78-5560
- Plasma Levels, 78-5560
- Benzo(a)pyrene
- DNA, Binding, 78-5560
- Enzyme Activation, 78-5560
- Cortisol
- Aryl Hydrocarbon Hydroxylases, 78-5560
- Plasma Levels, 78-5560

Strontium Radioisotopes

- Alginic Acid, Sodium Salt
- Body Burden, Rat, 78-5727
- Metabolism, Rat, 78-5727
- Sarcoma, Osteogenic
- BCG, 78-5728

Styrene

- Chromosome Aberrations
- Occupational Hazard, Review, 78-5436
- Enzyme Activation
- Mutagenic Activity, Review, 78-5436

Styrene Oxide

- see Benzene, (Epoxyethyl)-

Sulfide, Bis(2-chloroethyl)-

- Occupational Hazard
- Epidemiology, Review, 78-5410

Sulfite

- Vicia faba*
- Mutagenic Activity, 78-5635

Sulfuric Acid, Diethyl Ester

- Micrococcus radiodurans*
- Mutagenic Activity, 78-5701

Sulfuric Acid, Dimethyl Ester

- Adenine, 3-Methyl-
- Glycoside Hydrolases, 78-5663

uric Acid, Disodium Salt

- Benzo(a)pyren-3-ol
Conjugation Product, 78-5566
- Benzo(a)pyrene
Conjugation, Phenol/Quinone Metabolites, 78-5566

uric Acid, Iron Salt

- Cholanthrene, 3-Methyl-
Dihydrodiol Formation, 78-5530

Stomach Neoplasms

- Epidemiology, Hawaii, Japan, 78-5943
- Food Contamination, 78-5943

rodeoxycholic Acid

- Benzo(a)pyrene
DNA, Binding, 78-5562
- Intestinal Neoplasms
Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-5684

atogens

- Carcinogen, Environmental
Epidemiology, Review, 78-5405
- Mercury, Methyl-
Monitoring, Review, 78-5405
- Polychlorobiphenyl Compounds
Monitoring, Review, 78-5405

atoid Tumor

- Cell Differentiation
Mouse Embryo, Review, 78-5453
- Neoplasm Transplantation, Review, 78-5453

osterone, Propionate

- Mammary Neoplasms, Experimental
Adenocarcinoma, 78-5988
- Growth, 78-5988
- Neoplasm Transplantation, 78-5988

acycline

- Dimethylamine, *N*-Nitroso-
Nitrous Acid, Sodium Salt, 78-5610

0-Tetradecanoylphorbol-13-acetate

- Cells, Cultured
Lipids, Deacylation, 78-5509

Cortisol

- Prostaglandins, 78-5509

Cycloheximide

- Prostaglandins, 78-5509
- 5,8,11,14-Eicosatetranoic Acid
Cell Differentiation, 78-5509

Prostaglandins

- Cells, Cultured, 78-5509
- Kidney, Dog, 78-5509

Skin Neoplasms

- Benzo(a)pyrene, 78-5567
- Benzo(a)pyrene-6-methanol, 78-5567
- Benzo(a)pyrene, 6-Methyl-, 78-5567

ophylline

- Benz(a)anthracene, 7,12-Dimethyl-
Mutagenic Activity, 78-5518
- Polycyclic Hydrocarbons
Mutagenic Activity, 78-5518

ocyanic Acid

- Nitrosamines
Nitrous Acid, 78-5613
- Pentagastrin
Gastric Juice, 78-5632

Thiocyanic Acid (cont'd)

- Smoking
Gastric Juice, 78-5632
- Saliva, 78-5632

Thorium Dioxide

- Bone Marrow Cells
Case Report, 78-5717
- Bone Neoplasms
Epidemiology, Portugal, 78-5973
- Chromosome Aberrations
Case Report, 78-5717
- Chromosomes, Human, 1-3, 78-5717
- Chromosomes, Human, 21-22, 78-5717
- Granuloma
Epidemiology, Portugal, 78-5973
- Head and Neck Neoplasms
Granuloma, 78-5973
- Leukemia
Epidemiology, Portugal, 78-5973
- Liver Neoplasms
Cholangioma, 78-5973
- Epidemiology, Portugal, 78-5973
- Hemangioendothelioma, 78-5973
- Lung Neoplasms
Epidemiology, Portugal, 78-5973
- T-Lymphocytes
Chromosome Aberrations, 78-5717

Thymidine

- Dimethylamine, *N*-Nitroso-
DNA, Alkylation, 78-5599
- Oocytes
Mutagenic Activity, 78-5721
- Radiation
Mutagenic Activity, 78-5721

Thymidine Kinase

- Acetamide, *N*-Fluoren-2-yl-
Mutagenic Activity, 78-5666
- Acetamide, *N*-Fluoren-4-yl-
Mutagenic Activity, 78-5666
- Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-
Mutagenic Activity, 78-5666
- Benzo(a)pyrene
Mutagenic Activity, 78-5666
- Benzo(e)pyrene
Mutagenic Activity, 78-5666
- Carcinogen, Chemical
Mutation, 78-5652
- Diethylamine, *N*-Nitroso-
Mutagenic Activity, 78-5666
- Dimethylamine, *N*-Nitroso-
Mutagenic Activity, 78-5666
- Dipropylamine, *N*-Nitroso-
Mutagenic Activity, 78-5666
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
Mutagenic Activity, 78-5666
- Hycanthone
Mutagenic Activity, 78-5666
- Lymphoma
Methanesulfonic Acid, Ethyl Ester, 78-5652
- Methane, Iodo-
Mutagenic Activity, 78-5666
- Methanesulfonic Acid, Ethyl Ester
Mutagenic Activity, 78-5657, 78-5666
- Mutation, 78-5652
- 2-Oxetanone
Mutagenic Activity, 78-5666

Thymidine Kinase (cont'd)

- Uracil, 5-(Bis(2-chloroethyl)amino)-
Mutagenic Activity, 78-5666
- Virus, Herpes Simplex 1
DNA, Viral, 78-5820
Enzymatic Activity, 78-5818
- Virus, Herpes Simplex 2
DNA, Viral, 78-5820
Enzymatic Activity, 78-5818

Thymoma

- Hyperparathyroidism
Case Report, 78-5914
- Myasthenia Gravis
Case Report, 78-5914
- Neoplastic Endocrine-Like Syndromes
Case Report, 78-5914

Thymus Gland

- Virus, Murine Leukemia
Lymphocyte Transformation, 78-5756
RNA, Viral, 78-5761
Virus-Like Particles, 78-5761
- Virus, Radiation Leukemia
Lymphocyte Transformation, 78-5756

Thymus Neoplasms

- Lymphoma
Methanesulfonic Acid, Ethyl Ester, 78-5639
Urea, Ethyl Nitroso-, 78-5639
Urea, Methyl Nitroso-, 78-5639
- Virus, Feline Leukemia
RNA, Viral, 78-5787

Thyroid Diseases

- Breast Neoplasms
Epidemiology, 78-5989

Thyroid Hormones

- Breast Neoplasms
Serum Levels, 78-5989

Thyroid Neoplasms

- Acne
Radiotherapy, 78-5715
- Adenocarcinoma
Radiation, Ionizing, 78-5972
- Adenocarcinoma, Papillary
Radiation, Ionizing, 78-5972
- Adenoma
2-Propanol, 1,1'-Iminodi-*N*-nitroso-, 78-5580
- Carcinoma, Papillary
Iodine Radioisotopes, 78-5716
Radiation, Ionizing, 78-5715
- Hyperthyroidism
Radiotherapy, 78-5716
- Iodine Radioisotopes
Case Report, 78-5716, 78-5718
Epidemiology, 78-5718
- Leukemia, Myelocytic
Adenoma, 78-5906
- Neoplasms, Multiple Primary
Radiation, Ionizing, 78-5714
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
Carcinogenic Activity, Rat, 78-5580
- Radiation, Ionizing
Case Report, 78-5714, 78-5718
Epidemiology, 78-5715, 78-5972
Precancerous Conditions, 78-5715, 78-5972

Thyroidin

- Nervous System Neoplasms
Urea, Methyl Nitroso-, 78-5643

Thyronine, 3,3',5-Triiodo-

- Breast Neoplasms
Metabolism, 78-5990

Thyrotropin

- Breast Neoplasms
Neoplasm Metastasis, 78-5990
Protirelin, 78-5990
Radiotherapy, 78-5990
Serum Levels, 78-5989, 78-5990
- Iodine Radioisotopes
Serum Levels, Rat, 78-5713
Thyroid Gland Function, 78-5713

Thyroxine

- Breast Neoplasms
Serum Levels, 78-5990
- Iodine Radioisotopes
Diet, 78-5713
Serum Levels, Rat, 78-5713
- Mammary Neoplasms, Experimental
Benz(a)anthracene, 7,12-Dimethyl-, 78-5545

***p*-Toluanide, *N*-Isopropyl- α -(2-methylhydrazino)-, HCl**

- Dose-Response Study
Mutagenic Activity, Mouse, 78-5662

Toluene-2,4-diamine

- Hair Dyes
Carcinogenic Activity, Review, 78-5425
Mutagenic Activity, Review, 78-5425

Trachea

- Benzo(a)pyrene
Carcinogenic Metabolite, 78-5554
Metabolism, Microsomes, 78-5554
Metabolism, Organ Culture, 78-5554
- Benzo(a)pyrene, 9,10-Dihydro-9,10-dihydroxy-
Carcinogenic Metabolite, 78-5554

Tracheal Neoplasms

- Adenocarcinoma
Cholanthrene, 3-Methyl-, 78-5521
Urea, Methyl Nitroso-, 78-5640, 78-5642
- Carcinoma
Cholanthrene, 3-Methyl-, 78-5521
Urea, Methyl Nitroso-, 78-5640, 78-5642
- Carcinoma, Epidermoid
Cholanthrene, 3-Methyl-, 78-5521
Precancerous Conditions, 78-5978
Transplantation, Homologous, 78-5978
Urea, Methyl Nitroso-, 78-5640, 78-5642
- Cell Differentiation
Cells, Cultured, 78-5978
- Cells, Cultured
Transplantation, Homologous, 78-5978
- Cholanthrene, 3-Methyl-
Dose-Response Study, 78-5521
Neoplasm Transplantation, 78-5521
- Fibrosarcoma
Cholanthrene, 3-Methyl-, 78-5521
- Immunosuppression
Growth, 78-5978
Transplantation, Homologous, 78-5978
- Papilloma
Cholanthrene, 3-Methyl-, 78-5521
Urea, Methyl Nitroso-, 78-5642

heal Neoplasms (cont'd)

- Polyps
 - Urea, Methyl Nitroso-, 78-5642
- Urea, Methyl Nitroso-
 - Dose-Response Study, Hamster, 78-5640, 78-5642
- Precancerous Conditions, 78-5640

infection

see Transformation, Genetic

Information, Genetic

- Virus, Harvey Murine Sarcoma
 - DNA, Viral, 78-5784
- Virus, Moloney Murine Leukemia, 78-5784
- Virus, SV40
 - Cell Fusion, 78-5796
 - DNA, Viral, 78-5796
 - Uridine, 2'-Deoxy-5-iodo-, 78-5796

ketolase

- Nervous System Neoplasms
 - Urea, Ethyl Nitroso-, 78-5647

plantation, Autologous

- Bronchial Neoplasms
 - Benzo(a)pyrene, 78-5522
 - Cholanthrene, 3-Methyl-, 78-5522

plantation, Heterologous

- Burkitt's Lymphoma
 - Hybrid Cells, 78-5832
 - Mouse, Nude, 78-5832
- Carcinoma
 - Cholanthrene, 3-Methyl-, 78-5529
- Carcinosarcoma
 - Cholanthrene, 3-Methyl-, 78-5529
- Cholanthrene, 3-Methyl-
 - Cell Transformation, Neoplastic, 78-5529
- Leukemia, Myelocytic
 - Chromosomes, Human, 21-22, 78-5904, 78-5986
 - DNA Replication, 78-5986
 - Mouse, Nude, 78-5904, 78-5986
- Mesenchymoma
 - Anemia, Aplastic, 78-5904
 - Leukemia, Myelocytic, 78-5904
- Sarcoma
 - Cholanthrene, 3-Methyl-, 78-5529
 - Leukemia, Myelocytic, 78-5986
- Virus, Epstein-Barr
 - Hybrid Cells, 78-5832

plantation, Homologous

- Tracheal Neoplasms
 - Carcinoma, Epidermoid, 78-5978
 - Cells, Cultured, 78-5978
 - Immunosuppression, 78-5978

plantation Immunology

- Cholanthrene, 3-Methyl-
 - Amino Acids, 78-5525
- Fibrosarcoma
 - Ultraviolet Rays, 78-5858
- Histocompatibility Antigens
 - Hamster, Review, 78-5440
- Mammary Neoplasms, Experimental
 - Adenocarcinoma, 78-5539
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5539
 - Carcinoembryonic Antigen, 78-5539
 - Virus, Murine Mammary Tumor, 78-5754
- Neoplasms, Experimental
 - Amino Acids, 78-5525

Transplantation Immunology (cont'd)

- Polysaccharides, 78-5857
- Purine-6-thiol, 78-5857
- Sarcoma
 - Fetal Globulins, 78-5531
- Ultraviolet Rays
 - T-Lymphocytes, 78-5858
 - Radiation, Ionizing, 78-5858
- Virus, SV40
 - Histocompatibility Antigens, 78-5812
 - Mouse, Hamster, 78-5812

Trematoda

- Cholangioma
 - Dimethylamine, *N*-Nitroso-, 78-5602

s-Triazine, 2,4,6-Tris(1-aziridinyl)-

- Genes
 - Mutagenic Activity, Mouse, 78-5693

Triethylene Glycol

- Leukemia, Myeloblastic
 - Cell Differentiation, 78-5983

Trimethylamine

- see Methanamine, *N,N*-Dimethyl-

Trimethylaminoxide

- see Methanamine, *N,N*-Dimethyl-, *N*-Oxide

Tuberculosis, Pulmonary

- Lung Neoplasms
 - Epidemiology, 78-5930
- Neoplasms
 - Epidemiology, 78-5930

Tumor, Grawitz

- see Adenocarcinoma

Tyrosine

- Neoplasms, Experimental
 - Cholanthrene, 3-Methyl-, 78-5525

Tyrosine Aminotransferase

- Corticosterone
 - Liver, Rat, 78-5576
- Dexamethasone
 - Liver, Rat, 78-5576
- Hepatoma
 - Corticosterone, 78-5576
 - Dexamethasone, 78-5576
 - Diethylamine, *N*-Nitroso-, 78-5576

Ulcer

- Esophageal Neoplasms
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-5586

Ultraviolet Rays

- Benz(a)anthracene, 7,12-Dimethyl-
 - Co-carcinogenic Effect, 78-5540
- Caffeine
 - Chromosome Aberrations, 78-5703
- Cells, Cultured
 - Mutagenic Activity, 78-5702
- Coumarin
 - DNA Repair, Review, 78-5433
- Cytosine, 1- β -*D*-Arabinofuranosyl-
 - DNA Repair, 78-5477
- Diethylamine, *N*-Nitroso-
 - Photolysis Products, 78-5620
- Dimethylamine, *N*-Nitroso-
 - Photolysis Products, 78-5620
- DNA

Ultraviolet Rays (cont'd)

- Nucleic Acid Denaturation, 78-5706
- Squamous Cells, 78-5706
- DNA Repair
 - Dose-Response Study, 78-5707
 - Hepatocytes, Rat, 78-5477
 - Photoreactivation, 78-5705
 - Thymine Dimer Excision, 78-5709
- Drosophila melanogaster*
 - Mutagenic Activity, 78-5661
- Endonucleases
 - Micrococcus luteus*, 78-5700
- Ethidium Bromide
 - DNA Repair, 78-5477
- Fibroblasts
 - DNA Repair, 78-5704, 78-5705, 78-5707, 78-5709
- Fibrosarcoma
 - Transplantation Immunology, 78-5858
- Heat
 - Nucleic Acid Denaturation, 78-5706
- Hereditary Diseases
 - DNA Repair, Review, 78-5451
- Hydrogen Peroxide
 - Cell Survival, 78-5708
 - Mutagenic Activity, 78-5708
 - Synergistic Activity, 78-5708
- Immunosuppression
 - Lymphoid Tissue, 78-5858
- Lymphocytes
 - Chromosome Aberrations, 78-5703
- T-Lymphocytes
 - Immunosuppression, 78-5858
 - Transplantation Immunology, 78-5858
- Lymphosarcoma
 - Occupational Hazard, 78-5922
- Melanoma
 - Radiotherapy, 78-5873
- Micrococcus radiodurans*
 - Mutagenic Activity, 78-5701
- Mucopolysaccharidosis
 - Co-carcinogenic Effect, 78-5540
- Mutagenic Activity
 - Dose-Response Study, 78-5702
- Nitrosamines
 - Photolysis Products, 78-5620
- Psoralen
 - DNA, Review, 78-5434
 - Mutagenic/Carcinogenic Activity, Review, 78-5434
- Psoralen, 8-Methoxy-
 - Chromosome Aberrations, 78-5703
 - Mutagenic Activity, 78-5702
- Psoralen, 4,5',8-Trimethyl-
 - Chromosome Aberrations, 78-5703
- Pyrrolidine, 1-Nitroso-
 - Photolysis Products, 78-5620
- Radiation, Ionizing
 - Transplantation Immunology, 78-5858
- Skin Neoplasms
 - Dose-Response, Mathematical Model, 78-5975
 - Epidemiology, 78-5975
 - Ozone Depletion, 78-5698, 78-5975
- Streptomycin
 - Lethality, Algae, 78-5699
- Thymine Dimer Excision
 - Kinetics, 78-5709
- Virus, Adeno 2
 - Cell Transformation, Neoplastic, 78-5838

Ultraviolet Rays (cont'd)

- Virus, Herpes Simplex 1
 - RNA, Messenger, 78-5764
- Virus, SV40
 - Temperature Sensitive Mutants, 78-5810
- Xeroderma Pigmentosum
 - DNA Repair, 78-5704
- Uracil, 5-(Bis(2-chloroethyl)amino)-
 - Thymidine Kinase
 - Mutagenic Activity, 78-5666
- Uracil, 6-Propyl-2-thio-
 - Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-5545
- Uracil, 2-Thio-
 - Liver Neoplasms
 - Diethylamine, *N*-Nitroso-, 78-5575
 - Nose Neoplasms
 - Diethylamine, *N*-Nitroso-, 78-5575
- Uranium
 - Lung Neoplasms
 - Epidemiology, 78-5928
 - Smoking
 - Co-carcinogenic Effect, 78-5928
- Urea, 1,1-Dimethyl-3-phenyl-
 - Urea, 1-Methyl-1-nitroso-3-phenyl-
 - Carcinogenic Metabolite, 78-5637
- Urea, Ethyl-
 - Nitrous Acid, Sodium Salt
 - Nitrosation, 78-5636
- Urea, Ethyl Nitroso-
 - Astrocytoma
 - Urea, Methyl Nitroso-, 78-5638
 - Brain Neoplasms
 - Adenoma, 78-5650
 - Astrocytoma, 78-5649
 - Glioma, 78-5649
 - Oligodendroglioma, 78-5649
 - Organ Culture, 78-5650
 - Transplacental Carcinogenesis, 78-5649, 78-5650
 - Ultrastructural Study, 78-5649
 - Cell Transformation, Neoplastic
 - DNA, Alkylation, 78-5648
 - Cells, Cultured
 - Mutagenic Activity, 78-5660
 - DNA
 - Alkylation Products, 78-5639
 - Drosophila melanogaster*
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5660
 - Ependymoma
 - Urea, Methyl Nitroso-, 78-5638
 - Fibroblasts
 - DNA, Alkylation, 78-5648
 - Glioblastoma Multiforme
 - Urea, Methyl Nitroso-, 78-5638
 - Nervous System Neoplasms
 - Enzymatic Activity, 78-5647
 - Glucosephosphate Dehydrogenase, 78-5647
 - Phosphogluconate Dehydrogenase, 78-5647
 - Transketolase, 78-5647
 - Transplacental Carcinogenesis, 78-5638
 - Neurilemmoma
 - Urea, Methyl Nitroso-, 78-5638
 - Oligodendroglioma

Urea, Methyl Nitroso- (cont'd)

- Urea, Methyl Nitroso-, 78-5638
- Oxygen
 - DNA, Alkylation, 78-5648
- Pinealoma
 - Urea, Methyl Nitroso-, 78-5638
- Thymus Neoplasms
 - Lymphoma, 78-5639
- Urea, Methyl Nitroso-
 - Transplacental Carcinogenesis, 78-5638
- Virus, SV40
 - DNA, Alkylation, 78-5648
- Xeroderma Pigmentosum
 - DNA, Alkylation, 78-5648
- Urea, Ethyl Nitroso-**
 - Nitrous Acid, Sodium Salt
 - Nitrosation, 78-5636
- Urea, Methyl Nitroso-**
 - Ascorbic Acid, Monosodium Salt
 - Nitrosation, 78-5636
- Urea, Methyl Nitroso-**
 - Adenine, 3-Methyl-
 - Glycoside Hydrolases, 78-5663
- Astrocytoma
 - Antibody-Dependent Cytotoxicity, 78-5859
 - Antibody Specificity, 78-5859
 - Urea, Ethyl Nitroso-, 78-5638
- Brain Neoplasms
 - Antibody-Dependent Cytotoxicity, 78-5859
 - Antibody Formation, 78-5859
 - Antibody Specificity, 78-5859
 - Astrocytoma, 78-5859
 - Glioma, 78-5859
 - Oligodendroglioma, 78-5859
- Cells, Cultured
 - Mutagenic Activity, 78-5660
- Chromatin
 - DNA, Methylation, 78-5646
 - Liver, Rat, 78-5646
- Colonic Neoplasms
 - Carrageen, 78-5644
- Distamycin A
 - DNA, Methylation, 78-5646
- DNA
 - Alkylation Products, 78-5639
- DNA Repair
 - DNA, Single-Stranded, 78-5641
- DNA, Single-Stranded
 - Mutagenic Activity, Plants, 78-5641
- Drosophila melanogaster*
 - Chromosome Aberrations, 78-5660
 - Mutagenic Activity, 78-5660
- Ependymoma
 - Urea, Ethyl Nitroso-, 78-5638
- Escherichia coli*
 - Mutagenic Activity, 78-5588
- Glioblastoma Multiforme
 - Urea, Ethyl Nitroso-, 78-5638
- Glioma
 - Antibody-Dependent Cytotoxicity, 78-5859
 - Antibody Specificity, 78-5859
- Mouth Neoplasms
 - Papilloma, 78-5645
- Nervous System Neoplasms
 - Castration, 78-5643
 - Dienestrol, 78-5643

Urea, Methyl Nitroso- (cont'd)

- Fibrosarcoma, 78-5643
- Ganglioneuroma, 78-5643
- Immunization, 78-5643
- Mescasolyl, 78-5643
- Neurilemmoma, 78-5643
- Neuroblastoma, 78-5643
- Neurofibroma, 78-5643
- Thyroidin, 78-5643
- Transplacental Carcinogenesis, 78-5638
- Neurilemmoma
 - Urea, Ethyl Nitroso-, 78-5638
- Neurospora crassa*
 - Mutagenic Activity, 78-5675
- Nitrous Acid, Sodium Salt
 - Carbamic Acid, Ethyl-, Ethyl Ester, 78-5636
 - Stomach, Rat, 78-5636
- Odontogenic Tumor
 - Histological Study, Mouse, 78-5645
- Oligodendroglioma
 - Urea, Ethyl Nitroso-, 78-5638
- Pinealoma
 - Urea, Ethyl Nitroso-, 78-5638
- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-5569
 - Histological Study, Dog, 78-5569
 - Precancerous Conditions, 78-5569
- Skin Neoplasms
 - Carcinogenic Activity, Mouse, 78-5598
- Sodium Azide
 - DNA Repair, 78-5641
- Soy Beans
 - Somatic Mosaicism, 78-5664
- Spermine
 - DNA, Methylation, 78-5646
- Stomach Neoplasms
 - Papilloma, 78-5645
- Thymus Neoplasms
 - Lymphoma, 78-5639
- Tracheal Neoplasms
 - Adenocarcinoma, 78-5640, 78-5642
 - Carcinoma, 78-5640, 78-5642
 - Carcinoma, Epidermoid, 78-5640, 78-5642
 - Dose-Response Study, Hamster, 78-5640, 78-5642
 - Papilloma, 78-5642
 - Polyps, 78-5642
 - Precancerous Conditions, 78-5640
- Urea, Ethyl Nitroso-
 - Transplacental Carcinogenesis, 78-5638
- Urea, 1-Methyl-1-nitroso-3-phenyl-**
 - Stomach Neoplasms
 - Carcinogenic Activity, Rat, 78-5637
 - Carcinoma, Epidermoid, 78-5637
 - Papilloma, 78-5637
 - Urea, 1,1-Dimethyl-3-phenyl-
 - Carcinogenic Metabolite, 78-5637
- Urea, 1-Methyl-3-phenyl-**
 - Nitrous Acid, Sodium Salt
 - Carcinogenic Metabolite, 78-5637
 - Stomach Neoplasms
 - Nitrous Acid, Sodium Salt, 78-5637
- Uridine, 5-Bromo-2'-deoxy-**
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Cell Transformation, Neoplastic, 78-5685
 - DNA Replication, 78-5685
 - Photosensitivity, 78-5685

- Uridine, 5-Bromo-2'-deoxy- (cont'd)**
 Mitomycin C
 Chromatids, 78-5686, 78-5687
 Chromosome Aberrations, 78-5686
- Uridine, 2'-Deoxy-5-fluoro-**
 Radiation, Ionizing
 Chromosome Aberrations, 78-5692
- Uridine, 2'-Deoxy-5-iodo-**
 Virus, Murine Leukemia
 Virus Cultivation, 78-5767
 Virus, SV40
 Transformation, Genetic, 78-5796
- Uridine Diphosphate Glucuronic Acid**
 see Uridine Diphosphate Sugars
- Uridine Diphosphate Sugars**
 Benzo(a)pyrene
 Ames Test, 78-5559
- Urogenital Neoplasms**
 Genetics
 Classification, Review, 78-5455
 Kidney Diseases
 Epidemiology, Bulgaria, 78-5961
 Genetics, 78-5961
 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
 Carcinogenic Activity, Rat, 78-5580
- Uterine Neoplasms**
 Age Factors
 Epidemiology, Japan, 78-5953
 Carcinoma, Papillary
 Genetics, 78-5892
 Epidemiology
 Spain, 78-5915
 Estrogens
 Epidemiology, Review, 78-5432
 Genetics
 Case Report, 78-5892
 Hydrazine, 1,2-Dimethyl-
 Neoplasm Transplantation, 78-5495
 Hypertension
 Epidemiology, Japan, 78-5953
 Leiomyoma
 Genetics, 78-5892
 Parity
 Epidemiology, Japan, 78-5953
 Sarcoma
 Hydrazine, 1,2-Dimethyl-, 78-5495
- Vaginal Neoplasms**
 Carcinoma
 Chromosome Aberrations, 78-5871
 Chromosome Aberrations
 Case Report, 78-5871
 Chromosomes, Human, 1-3, 78-5871
 Eosinophils
 Neoplasm Invasiveness, 78-5889
 Neoplasm Invasiveness
 Ultrastructural Study, 78-5889
- Valeric Acid, 2,2-Diphenyl-, 2-(Diethylamino)ethyl Ester, HCl**
 Dimethylamine, *N*-Nitroso-
 Metabolism, Rat, 78-5603
 Nucleic Acids, Binding, 78-5603
- Vasopressin**
 Mammary Neoplasms, Experimental
 Acetate Incorporation, 78-5543
- Vibramycin**
 Dimethylamine, *N*-Nitroso-
 Ascorbic Acid, 78-5610
 Nitrous Acid, Sodium Salt, 78-5610
- Vinblastine**
 Leukemia, Myelocytic
 Hodgkin's Disease, 78-5898
- Viral Proteins**
 Virus, Adeno 2
 Cell Transformation, Neoplastic, 78-5838
 Diimido Ester Cleavage, 78-5842
 DNA, Viral, 78-5842
 HeLa Cells, 78-5841
 Isolation and Characterization, 78-5838, 78-5841
 78-5842
 Temperature Sensitive Mutants, 78-5843
 Virus Assembly, Ultrastructural Study, 78-5842
 Virus, Adeno 2 - SV40 Hybrid
 Virus, Adeno 2, 78-5841
 Virus, Avian Leukosis
 Electrophoretic Mobility, 78-5747
 Virus, Avian Leukosis-Sarcoma
 Antigenic Determinants, 78-5736
 Isolation and Characterization, 78-5736
 Virus, Avian Myeloblastosis
 Antigenic Determinants, 78-5742
 DNA Polymerase, 78-5742
 Immunoprecipitation, 78-5742
 Isolation and Characterization, 78-5742
 Proteolysis, 78-5742
 Virus, Avian Sarcoma
 Amino Acids, 78-5739
 Isolation and Characterization, 78-5739
 Virus, Baboon
 Antigenic Determinants, 78-5814
 Virus, C-Type RNA Tumor
 Antigenic Determinants, 78-5814
 Genes, Structural, 78-5849
 Virus, D-Type RNA Tumor
 Antigenic Determinants, 78-5814
 Virus, Feline Leukemia
 Cell Transformation, Neoplastic, 78-5786
 Virus, Feline Sarcoma
 Cell Transformation, Neoplastic, 78-5786
 Virus, Kirsten Murine Sarcoma
 Virus, Helper, 78-5773
 Virus, Mason-Pfizer Monkey
 Antigenic Determinants, 78-5814
 Virus, Moloney Murine Leukemia
 Cell Membrane, 78-5776
 Virus, Moloney Murine Sarcoma
 Virus, Helper, 78-5773
 Virus, Murine Leukemia
 Antigen-Antibody Reactions, 78-5759
 B-Tropism, 78-5759
 Radioimmunoassay, 78-5759
 Virus, Recombinant, 78-5759
 Virus, Rauscher Murine Leukemia
 Antibody Specificity, 78-5779
 Antigenic Determinants, 78-5778
 Cell Membrane, 78-5776
 Cytochalasin B, 78-5780
 Genes, Structural, 78-5849
 Glucose, 2-Deoxy-, 78-5780
 Histocompatibility Antigens, 78-5779
 Immune Virolysis Competition Assay, 78-5779

al Proteins (cont'd)

- Isoantigens, 78-5779
- Isolation and Characterization, 78-5778
- Temperature Sensitive Mutants, 78-5780
- Virus, Ring-Necked Pheasant**
 - Electrophoretic Mobility, 78-5747
- Virus, Rous Sarcoma**
 - Electrophoretic Mobility, 78-5747
- Virus, Squirrel Monkey**
 - Antigenic Determinants, 78-5814
- Virus, SV40**
 - Chromosomes, 78-5811

rus, Abelson Murine Leukemia

- Lymphosarcoma
 - Cells, Cultured, 78-5979
 - Neoplasm Transplantation, 78-5979

rus, Adeno 2

- Concanavalin A
 - Glycoproteins, 78-5840
- DNA, Viral**
 - Carrier Proteins, 78-5841
 - Cell Transformation, Neoplastic, 78-5838
 - DNA-DNA Hybridization, 78-5838
 - Viral Proteins, 78-5842
- Glycoproteins**
 - Immunoprecipitation, 78-5840
 - Isolation and Characterization, 78-5840
 - Proteolysis, 78-5840
- HeLa Cells**
 - Viral Proteins, 78-5841
- Methylglycosides**
 - Glycoproteins, 78-5840
- RNA, Viral**
 - α -Amanitine, 78-5839
 - Cell Transformation, Neoplastic, 78-5839
 - DNA-RNA Hybridization, 78-5839
 - RNA Polymerase, 78-5839
- Ultraviolet Rays**
 - Cell Transformation, Neoplastic, 78-5838
- Viral Proteins**
 - Cell Transformation, Neoplastic, 78-5838
 - Diimido Ester Cleavage, 78-5842
 - Isolation and Characterization, 78-5838, 78-5841, 78-5842
 - Temperature Sensitive Mutants, 78-5843
 - Virus Assembly, Ultrastructural Study, 78-5842
- Virus, Adeno 2 - SV40 Hybrid**
 - Viral Proteins, 78-5841
- Virus Replication**
 - Temperature Sensitive Mutants, 78-5843

irus, Adeno 5

- Cell Transformation, Neoplastic
 - Surface Properties, 78-5844
- Cells, Cultured
 - Surface Properties, 78-5844
 - Tumorigenicity, 78-5844
- Cycloheximide**
 - Deoxyribonuclease, 78-5845
- Deoxyribonuclease**
 - DNA Nucleotidyltransferases, 78-5845
 - DNA, Single-Stranded, 78-5845
 - Virus Replication, 78-5845
- DNA, Viral**
 - Deoxyribonuclease, 78-5845
- Glycoproteins**
 - Cell Transformation, Neoplastic, 78-5844

Virus, Adeno 5 (cont'd)

- Neoplasms, Experimental
 - Surface Properties, 78-5844

Virus, Adeno 6

- Hexokinase**
 - Enzymatic Activity, 78-5847
- Lactate Dehydrogenase Isoenzymes**
 - Enzymatic Activity, 78-5847
- Malate Dehydrogenase**
 - Enzymatic Activity, 78-5847

Virus, Adeno 12

- Cell Transformation, Neoplastic
 - Temperature Sensitive Mutants, 78-5846
- DNA Replication**
 - Temperature Sensitive Mutants, 78-5846
- DNA, Viral**
 - Cell Transformation, Neoplastic, 78-5846
 - DNA Replication, 78-5848
 - Initiation and Termination Sites, 78-5848
 - Oligonucleotides, 78-5848
- Hexokinase**
 - Cell Transformation, Neoplastic, 78-5847
- Lactate Dehydrogenase Isoenzymes**
 - Cell Transformation, Neoplastic, 78-5847
- Malate Dehydrogenase**
 - Cell Transformation, Neoplastic, 78-5847
- Sarcoma**
 - Cell Transformation, Neoplastic, 78-5847
 - Hexokinase, 78-5847
- Sarcoma, Reticulum Cell**
 - Enzymatic Activity, 78-5847
 - Hexokinase, 78-5847

Virus, Adeno 2 - SV40 Hybrid

- Virus, Adeno 2**
 - Viral Proteins, 78-5841

Virus, Avian Leukosis

- Viral Proteins**
 - Electrophoretic Mobility, 78-5747

Virus, Avian Leukosis-Sarcoma

- Antigenic Determinants**
 - Chick Embryo, 78-5736
- Viral Proteins**
 - Antigenic Determinants, 78-5736
 - Isolation and Characterization, 78-5736

Virus, Avian Myeloblastosis

- Antigen-Antibody Reactions**
 - Antigenic Determinants, 78-5741
- Antigens, Viral**
 - Antigen-Antibody Reactions, 78-5741
 - Casein, 78-5741
 - Concanavalin A, 78-5741
 - Ovalbumin, 78-5741
 - Papain, 78-5741
 - Peptide Hydrolases, 78-5741
 - Serum Albumin, Bovine, 78-5741
- DNA Polymerase**
 - Benzoic Acid, 3,3'-Dithiobis(6-nitro)-, 78-5743
 - Benzoic Acid, p-Hydroxymercuri-, 78-5743
 - Benzoic Acid, 2-Nitro-5-thiocyano-, 78-5743
 - Proteolysis, 78-5742
 - 1H-Pyrrole-2,5-dione, 1-Ethyl-, 78-5743
- Reverse Transcriptase**
 - Amino Acids, 78-5743
- Ribonuclease**

Virus, Avian Myeloblastosis (cont'd)

- Benzoic Acid, 3,3'-Dithiobis(6-nitro)-, 78-5743
- Benzoic Acid, *p*-Hydroxymercuri-, 78-5743
- Benzoic Acid, 2-Nitro-5-thiocyano-, 78-5743
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-, 78-5743

Viral Proteins

- Antigenic Determinants, 78-5742
- DNA Polymerase, 78-5742
- Immunoprecipitation, 78-5742
- Isolation and Characterization, 78-5742
- Proteolysis, 78-5742

Virus, Rous Sarcoma

- Antigenic Determinants, 78-5741

Virus, Avian Sarcoma**Antigenic Determinants**

- Cell Transformation, Neoplastic, 78-5739

Antigens, Viral

- Cell Transformation, Neoplastic, 78-5740

Carcinoembryonic Antigen

- Cell Transformation, Neoplastic, 78-5740

Protein Kinase

- Cell Transformation, Neoplastic, 78-5737
- Enzymatic Activity, 78-5737
- Genes, Viral, 78-5737
- Temperature Sensitive Mutants, 78-5737

RNA, Viral

- DNA-RNA Hybridization, 78-5738
- Isolation and Characterization, 78-5738
- Nucleotide Sequence, 78-5738
- Poly A, 78-5738
- Transformation-Defective Mutants, 78-5738

Viral Proteins

- Amino Acids, 78-5739
- Isolation and Characterization, 78-5739

Virus, Baboon**Lymphocyte Transformation**

- Trans-Species Infection, 78-5825

B-Lymphocytes

- Lymphocyte Transformation, 78-5825

Rhabdomyosarcoma

- Antigen-Antibody Reactions, 78-5865

Viral Proteins

- Antigenic Determinants, 78-5814

Virus, Polyoma, JC

- Antigenic Determinants, 78-5793

Virus, RD-114

- Antigenic Determinants, 78-5850

Virus, Squirrel Monkey

- DNA-RNA Hybridization, 78-5814

Virus, Bovine Leukemia**DNA, Viral**

- Cell Transformation, Neoplastic, 78-5788
- DNA-DNA Hybridization, 78-5788
- Tissue Distribution, 78-5788

Genetics

- Immune Response, 78-5982

Horizontal Transmission

- Epidemiology, 78-5982

Lymphosarcoma

- DNA, Viral, 78-5788

Virus, C-Type RNA Tumor**Cells, Cultured**

- Antigenic Determinants, Viral Complex, 78-5773

Glycoproteins

- Tryptic Peptide Analysis, 78-5762

Virus, C-Type RNA Tumor (cont'd)

- Virus, Recombinant, 78-5762

Mammary Neoplasms, Experimental

- Benz(a)anthracene, 7,12-Dimethyl-, 78-5535
- Isolation and Characterization, 78-5535

RNA, Viral

- DNA-RNA Hybridization, 78-5814
- Replication-Defective Viruses, 78-5772

Viral Proteins

- Antigenic Determinants, 78-5814
- Genes, Structural, 78-5849

Virus, D-Type RNA Tumor

- Antigenic Determinants, 78-5814
- DNA-RNA Hybridization, 78-5814

Virus, Helper

- DNA-RNA Hybridization, 78-5772
- RNA, Viral, 78-5772

Virus, Sarcoma

- Recombination, Genetic, 78-5849

Virus, Cytomegalo**Cell Transformation, Neoplastic**

- Cell Culture Contamination, 78-5823

Colonic Neoplasms

- Adenocarcinoma, 78-5885
- DNA-RNA Hybridization, 78-5885
- DNA, Viral, 78-5885

Enteritis, Regional

- DNA, Viral, 78-5885

Intestinal Neoplasms

- Adenocarcinoma, 78-5885
- DNA-RNA Hybridization, 78-5885
- DNA, Viral, 78-5885

Intestinal Polyps

- DNA, Viral, 78-5885
- Genetics, 78-5885

Virus, Herpes

- Antigenic Determinants, 78-5823
- Cell Culture Contamination, 78-5823

Virus, Rhino

- Antigenic Determinants, 78-5823
- Isolation and Characterization, 78-5823

Virus, D-Type RNA Tumor**RNA, Viral**

- DNA-RNA Hybridization, 78-5814

Viral Proteins

- Antigenic Determinants, 78-5814

Virus, C-Type RNA Tumor

- Antigenic Determinants, 78-5814
- DNA-RNA Hybridization, 78-5814

Virus, Mason-Pfizer Monkey

- Antigenic Determinants, 78-5814

Virus, DNA***Drosophila***

- Mutagenic Activity, 78-5851

Virus, Epstein-Barr**Antibodies, Viral**

- IgM, 78-5829
- Immune Response, Infant, 78-5835, 78-5836
- Seroepidemiology, Ghana, 78-5835

Antibody Specificity

- Infant, 78-5836

Antigens

- Cell Membrane, 78-5833
- Cell Transformation, Neoplastic, 78-5833
- Isolation and Characterization, 78-5833

Epstein-Barr (cont'd)

- Antigens, Viral
 - Cortisone, 78-5824
 - Cycloheximide, 78-5828
 - DNA Replication, 78-5828
 - Lymphocyte Transformation, 78-5828
- Burkitt's Lymphoma
 - Antigens, Viral, 78-5824, 78-5827
 - Cortisone, 78-5824
 - Epidemiology, Review, 78-5441
 - Receptors, Complement, 78-5827
 - Receptors, Viral, 78-5827
 - Seroepidemiology, Review, 78-5443
- Carcinoma
 - Mouse, Nude, 78-5832
- DNA Replication
 - Interactions, B95-8/P3HR-1 Substrains, 78-5834
- Head and Neck Neoplasms
 - Antigens, Viral, 78-5830
 - Carcinoma, 78-5830
- Hybrid Cells
 - Transplantation, Heterologous, 78-5832
- Infant
 - Seroepidemiology, Ghana, 78-5836
- Infectious Mononucleosis
 - Antibodies, Viral, 78-5829
 - Antigens, Viral, 78-5824
 - Cortisone, 78-5824
 - Seroepidemiology, Review, 78-5443
- Leukemia
 - Antibodies, Viral, 78-5829
 - Cells, Cultured, 78-5831
 - Serodiagnosis, 78-5829
- Leukemia, Lymphoblastic
 - Antigens, Viral, 78-5827
 - Receptors, Complement, 78-5827
 - Receptors, Viral, 78-5827
- Lymphocyte Transformation
 - Interactions, B95-8/P3HR-1 Substrains, 78-5834
 - Trans-Species Infection, 78-5825
- Lymphocytes
 - DNA Replication, 78-5834
- B-Lymphocytes
 - DNA Replication, 78-5826
 - Lymphocyte Transformation, 78-5825, 78-5826
 - 78-5828
- T-Lymphocytes
 - Lymphocyte Transformation, 78-5826
- Lymphoma
 - Antibodies, Viral, 78-5829
 - Serodiagnosis, 78-5829
- Lymphosarcoma
 - Receptors, Viral, 78-5827
- Malaria
 - Co-carcinogenic Activity, Review, 78-5441
 - B-Lymphocytes, 78-5441
- Nasopharyngeal Neoplasms
 - Adenocarcinoma, Papillary, 78-5830
 - Antibodies, Viral, 78-5830
 - Carcinoma, 78-5830
 - Carcinoma, Epidermoid, 78-5830
 - Seroepidemiology, Review, 78-5443
- Neoplasms, Experimental
 - Animal Model, Mouse, 78-5832
- Saliva
 - Horizontal Transmission, Infant, 78-5835
- Socioeconomic Factors

Virus, Epstein-Barr (cont'd)

- Seroepidemiology, Ghana, 78-5835
 - Staphylococcus aureus*
 - DNA Replication, 78-5834
 - Virus, Herpes Papio
 - Antigenic Determinants, 78-5817
 - DNA-RNA Hybridization, 78-5817
 - Virus Replication
 - Cells, Cultured, 78-5831
- Virus, Feline Leukemia**
- Antibodies, Viral
 - Epidemiology, 78-5785
 - Immune Response, 78-5785
 - Virus Subgroup, 78-5785
 - Antigens, Viral
 - Cell Membrane, 78-5786
 - Cell Transformation, Neoplastic, 78-5786
 - Lymphosarcoma
 - Antibodies, Viral, 78-5785
 - RNA, Viral
 - Cap Structures, 78-5787
 - DNA-RNA Hybridization, 78-5787
 - Isolation and Characterization, 78-5787
 - Thymus Neoplasms
 - RNA, Viral, 78-5787
 - Viral Proteins
 - Cell Transformation, Neoplastic, 78-5786
 - Virus, Polyoma, JC
 - Antigenic Determinants, 78-5793

Virus, Feline Sarcoma

- Viral Proteins
 - Cell Transformation, Neoplastic, 78-5786

Virus, Friend Murine Leukemia

- Antibodies, Viral
 - Titration Method, 78-5765
 - Erythroleukemia
 - Polyribosomes, 78-5764
 - RNA, Messenger, 78-5764
 - Immune Serums
 - Virus Neutralization, 78-5765
 - Immunosuppression
 - Virus Replication, 78-5763
 - Leukemia
 - Immune Serums, 78-5765
 - Lymphoid Tissue
 - Antigens, Viral, 78-5763
 - Immunoglobulins, Surface, 78-5763
 - Surface Properties, 78-5763
 - Ultrastructural Study, 78-5763
 - Sarcoma
 - Immune Serums, 78-5765
 - Virus, Friend Spleen Focus-Forming
 - DNA-RNA Hybridization, 78-5768
 - Virus, Helper
 - Rat Kidney Cells, 78-5766
 - Virus Replication
 - Rat Kidney Cells, 78-5766
- Virus, Friend Spleen Focus-Forming**
- RNA, Viral
 - Virus, Helper, 78-5768
 - Virus, Friend Murine Leukemia
 - DNA-RNA Hybridization, 78-5768
 - Virus, Helper
 - DNA-RNA Hybridization, 78-5768

- Virus, Gibbon Ape Lymphoma**
 - Rhabdomyosarcoma
 - Antigen-Antibody Reactions, 78-5865
- Virus, Gross Murine Leukemia**
 - Antigens, Viral
 - Cells, Cultured, 78-5769
 - Immune Serums
 - Antigens, Viral, 78-5769
 - Radiation, Ionizing
 - Antigens, Viral, 78-5769
 - Virus Replication, 78-5769
 - Virus-Like Particles
 - Cells, Cultured, 78-5769
- Virus, Harvey Murine Sarcoma**
 - DNA, Viral
 - Cell Transformation, Neoplastic, 78-5784
 - Kinetics, Plaque and Focus Formation, 78-5784
 - Transformation, Genetic, 78-5784
 - Virus, Helper
 - Cell Transformation, Neoplastic, 78-5784
 - Virus, Moloney Murine Leukemia
 - Transformation, Genetic, 78-5784
- Virus, Helper**
 - Lymphoma
 - Virus, Moloney Murine Leukemia, 78-5777
 - Virus, C-Type RNA Tumor
 - DNA-RNA Hybridization, 78-5772
 - RNA, Viral, 78-5772
 - Virus, Friend Murine Leukemia
 - Rat Kidney Cells, 78-5766
 - Virus, Friend Spleen Focus-Forming
 - DNA-RNA Hybridization, 78-5768
 - RNA, Viral, 78-5768
 - Virus, Harvey Murine Sarcoma
 - Cell Transformation, Neoplastic, 78-5784
 - Virus, Kirsten Murine Sarcoma
 - Antigenic Determinants, Viral Complex, 78-5773
 - Viral Proteins, 78-5773
 - Virus, Moloney Murine Leukemia
 - DNA-RNA Hybridization, 78-5772
 - Virus, Moloney Murine Sarcoma
 - Antigenic Determinants, Viral Complex, 78-5773
 - Viral Proteins, 78-5773
 - Virus, Murine Leukemia, 78-5773
 - Virus, Murine Leukemia
 - Virus Replication, 78-5756
 - Virus, Woolly Monkey
 - DNA-RNA Hybridization, 78-5772
- Virus, Hepatitis**
 - Hepatoma
 - Australia Antigen, 78-5837
 - Epidemiology, Senegal, 78-5837
 - Liver Cirrhosis
 - Epidemiology, Greenland, 78-5963
 - Liver Neoplasms
 - Epidemiology, Greenland, 78-5963
- Virus, Herpes**
 - Cell Transformation, Neoplastic
 - Cells, Cultured, 78-5823
 - Hyperplasia
 - Spleen, Guinea Pig, 78-5815
 - Leukemia
 - Immune Response, Guinea Pig, 78-5815
 - B-Lymphocytes
 - Immune Response, Guinea Pig, 78-5815
- Virus, Herpes (cont'd)**
 - Virus, Cytomegalo
 - Antigenic Determinants, 78-5823
 - Cell Culture Contamination, 78-5823
 - Virus, Rhino
 - Antigenic Determinants, 78-5823
- Virus, Herpes Ateles**
 - Lymphoma
 - Owl Monkey, 78-5816
- Virus, Herpes Papio**
 - Lymphocyte Transformation
 - Marmoset, 78-5817
 - Radiation, Ionizing
 - Lymphocyte Transformation, 78-5817
 - Virus, Epstein-Barr
 - Antigenic Determinants, 78-5817
 - DNA-RNA Hybridization, 78-5817
- Virus, Herpes Simplex 1**
 - DNA, Viral
 - Peptides, 78-5820
 - Phosphonoacetic Acid, 78-5820
 - Thymidine Kinase, 78-5820
 - Virus, Recombinant, 78-5820
 - Erythroleukemia
 - Polyribosomes, 78-5764
 - RNA, Messenger, 78-5764
 - Nucleoside Deaminases
 - Enzymatic Activity, 78-5818
 - Nucleosides
 - Oxidoreductases, 78-5818
 - Oxidoreductases
 - Enzymatic Activity, 78-5818
 - Peptides
 - Isolation and Characterization, 78-5820
 - Virus, Recombinant, 78-5820
 - Polyribosomes
 - RNA, Messenger, 78-5764
 - Thymidine Kinase
 - Enzymatic Activity, 78-5818
 - Ultraviolet Rays
 - RNA, Messenger, 78-5764
 - Virus, Herpes Simplex 2
 - Virus, Recombinant, 78-5819, 78-5820
 - Virus Replication
 - Host Range, 78-5819
 - Virus, Recombinant, 78-5819
- Virus, Herpes Simplex 2**
 - Cell Transformation, Neoplastic
 - Temperature Sensitive Mutants, 78-5821
 - Thymidine Kinase Phenotype, 78-5821
 - Cervix Neoplasms
 - Antibodies, Viral, 78-5822
 - Epidemiology, 78-5822
 - DNA Replication
 - Temperature Sensitive Mutants, 78-5821
 - DNA, Viral
 - Peptides, 78-5820
 - Phosphonoacetic Acid, 78-5820
 - Thymidine Kinase, 78-5820
 - Virus, Recombinant, 78-5820
 - Nucleoside Deaminases
 - Enzymatic Activity, 78-5818
 - Nucleosides
 - Oxidoreductases, 78-5818
 - Oxidoreductases

Herpes Simplex 2 (cont'd)
 Enzymatic Activity, 78-5818

Peptides
 Isolation and Characterization, 78-5820
 Virus, Recombinant, 78-5820

Thymidine Kinase
 Enzymatic Activity, 78-5818

Virus, Herpes Simplex 1
 Virus, Recombinant, 78-5819, 78-5820

Virus Replication
 Host Range, 78-5819
 Virus, Recombinant, 78-5819

Kirsten Murine Sarcoma
 Cell Aggregation
 Cell Transformation, Neoplastic, 78-5507

Collagen
 Cell Transformation, Neoplastic, 78-5783
 Revertant Cells, 78-5783

Dibutyryl Cyclic AMP
 Collagen, 78-5783

Virus, Helper
 Antigenic Determinants, Viral Complex, 78-5773
 Viral Proteins, 78-5773

Virus, Moloney Murine Sarcoma
 Antigenic Determinants, Viral Complex, 78-5773

is-Like Particles
 Intestinal Neoplasms
 Lymphosarcoma, 78-5886

Leukemia
 Virus, Murine Leukemia, 78-5757

Lymphoma
 Virus, Murine Leukemia, 78-5761

Plasmacytoma
 A- and C-Type Particles, 78-5852
 Immunosuppression, 78-5856
 RNA, 78-5856
 Ultrastructural Study, Mouse, 78-5852

Skin Neoplasms
 Carcinoma, Epidermoid, 78-5789
 Keratoacanthoma, 78-5789
 Papilloma, 78-5789
 Ultrastructural/Immunofluorescent Study, Mouse
 78-5789
 Virus, Papilloma, 78-5789

Virus, Gross Murine Leukemia
 Cells, Cultured, 78-5769

Virus, Murine Leukemia
 Isolation and Characterization, AKR Mouse
 78-5761
 Reverse Transcriptase, 78-5761
 Thymus Gland, 78-5761
 Ultrastructural Study, P388 Cells, 78-5757

rus, Marek's Disease Herpes
 Antigens, Viral
 Blood Cells, 78-5749

Avian Leukosis
 Antigens, Viral, 78-5749

Histocompatibility Antigens
 Antigen-Antibody Reactions, 78-5748

Lymphoma
 Antigens, Viral, 78-5749
 Virus, Rous Sarcoma, 78-5750

Sarcoma
 Virus, Rous Sarcoma, 78-5750

Virus, Rous Sarcoma
 Co-carcinogenic Activity, 78-5750

Virus, Mason-Pfizer Monkey
 Viral Proteins
 Antigenic Determinants, 78-5814

Virus, D-Type RNA Tumor
 Antigenic Determinants, 78-5814

Virus, Polyoma, JC
 Antigenic Determinants, 78-5793

Virus Replication
 Embryonic Cells, Human, 78-5809

Virus, Rous Sarcoma
 Cell Fusion, 78-5809
 Virus Replication, 78-5809

Virus, Squirrel Monkey
 DNA-RNA Hybridization, 78-5814

Virus, SV40
 Cell Fusion, 78-5809
 Virus Replication, 78-5809

Virus, Moloney Murine Leukemia
 Antigens, Viral
 Isolation and Characterization, 78-5774

Cell Membrane
 Viral Proteins, 78-5776

Lymphoma
 Antigens, Viral, 78-5774
 Virus, Helper, 78-5777

Reverse Transcriptase
 Temperature Sensitive Mutants, 78-5776

RNA, Viral
 DNA-RNA Hybridization, 78-5770

Virus Cultivation
 Plaque Assay, 78-5775
 Reverse Transcriptase, 78-5775
 Synthetic Capillary Method, 78-5775

Virus, Harvey Murine Sarcoma
 Transformation, Genetic, 78-5784

Virus, Helper
 DNA-RNA Hybridization, 78-5772

Virus, Murine Leukemia
 Reverse Transcriptase, 78-5761

Virus, Recombinant
 Isolation and Characterization, 78-5777

Virus Replication
 Temperature Sensitive Mutants, 78-5776

Virus, Moloney Murine Sarcoma
 Leukemia, Lymphoblastic
 DNA-RNA Hybridization, 78-5771

Leukemia, Myeloblastic
 DNA-RNA Hybridization, 78-5771

Leukemia, Myelocytic
 DNA-RNA Hybridization, 78-5771

RNA, Viral
 DNA-RNA Hybridization, 78-5770

Virus, Helper
 Antigenic Determinants, Viral Complex, 78-5773
 Viral Proteins, 78-5773

Virus, Kirsten Murine Sarcoma
 Antigenic Determinants, Viral Complex, 78-5773

Virus, Murine Leukemia
 Virus, Helper, 78-5773

Virus, Moloney Murine Sarcoma-Leukemia
 RNA, Viral
 DNA-RNA Hybridization, 78-5770
 Virus Replication, 78-5770

Virus, Murine Leukemia
 Carcinoma

Virus, Murine Leukemia (cont'd)

- Isolation and Characterization, 78-5767
- DNA-RNA Hybridization
 - Rat Kidney Cells, 78-5766
- Fibrosarcoma
 - Antigens, Viral, 78-5866
- Glycoproteins
 - Carbohydrates, 78-5781
 - Cell Transformation, Neoplastic, 78-5781
 - Mouse, AKR, 78-5781
 - Virus, Recombinant, 78-5762
- Leukemia
 - Virus-Like Particles, 78-5757
- Leukemia, Lymphocytic
 - Isolation and Characterization, 78-5767
- Lymphoma
 - RNA, Viral, 78-5761
 - Virus-Like Particles, 78-5761
- Pentadecane, 2,6,10,14-Tetramethyl-
 - Virus Activation, 78-5758
- Plasmacytoma
 - Virus Activation, 78-5758
- Radiation, Ionizing
 - Fv-1 Restriction, 78-5760
 - Virus Replication, 78-5760
- RNA, Viral
 - DNA-RNA Hybridization, 78-5761
 - Thymus Gland, 78-5761
- Sarcoma
 - Isolation and Characterization, 78-5767
- Thymus Gland
 - Lymphocyte Transformation, 78-5756
- Uridine, 2'-Deoxy-5-iodo-
 - Virus Cultivation, 78-5767
- Viral Proteins
 - Antigen-Antibody Reactions, 78-5759
 - B-Tropism, 78-5759
 - Radioimmunoassay, 78-5759
 - Virus, Recombinant, 78-5759
- Virus Cultivation
 - Nonleukemogenic Virus, 78-5767
 - Tumorigenicity, 78-5767
- Virus, Helper
 - Virus Replication, 78-5756
- Virus-Like Particles
 - Isolation and Characterization, AKR Mouse 78-5761
 - Reverse Transcriptase, 78-5761
 - Thymus Gland, 78-5761
 - Ultrastructural Study, P388 Cells, 78-5757
- Virus, Moloney Murine Leukemia
 - Reverse Transcriptase, 78-5761
- Virus, Moloney Murine Sarcoma
 - Virus, Helper, 78-5773
- Virus, Polyoma, JC
 - Antigenic Determinants, 78-5793
- Virus, Rauscher Murine Leukemia
 - Antigenic Determinants, 78-5781
- Virus Replication
 - Hematopoietic System, 78-5756
 - Host Range, 78-5767
 - Lymphoid Tissue, 78-5756
 - Rat Kidney Cells, 78-5766

Virus, Murine Mammary Tumor

- Antigens, Viral
 - Horizontal Transmission, 78-5755
- DNA, Viral

Virus, Murine Mammary Tumor (cont'd)

- Virus Replication, 78-5751
- Immunogenetics
 - Milk-Transmitted Virus, 78-5754
 - Vertical Transmission, 78-5754
- Mammary Neoplasms, Experimental
 - Adenocarcinoma, 78-5753
 - DNA-RNA Hybridization, 78-5751
 - DNA, Viral, 78-5751
 - Horizontal Transmission, 78-5755
 - Immunogenetics, 78-5754
 - Mouse, GH, 78-5751
 - Reverse Transcriptase, 78-5752
 - RNA, Viral, 78-5753
 - Transplantation Immunology, 78-5754
- Nucleic Acids
 - Cell Transformation, Neoplastic, 78-5753
- RNA, Viral
 - DNA-RNA Hybridization, 78-5753
 - Precancerous Conditions, 78-5753
- Virus Replication
 - Cells, Cultured, 78-5752
- Virus, Papilloma**
 - Antibodies, Viral
 - Animals, Domestic, 78-5791
 - Immunodiffusion, 78-5791
 - Antigen-Antibody Complex
 - Ultrastructural Study, Dog, 78-5791
 - Bracken Fern
 - Co-carcinogenic Activity, 78-5790
 - Epidemiology, Cattle, 78-5790
 - Esophageal Neoplasms
 - Epidemiology, Cattle, 78-5790
 - Mouth Neoplasms
 - Epidemiology, Cattle, 78-5790
 - Pharyngeal Neoplasms
 - Epidemiology, Cattle, 78-5790
 - Skin Neoplasms
 - Virus-Like Particles, 78-5789
- Virus, Polyoma**
 - Amino Acids
 - Nucleotide Sequence, 78-5792
 - DNA, Viral
 - Nucleotide Sequence, 78-5792
- Virus, Polyoma, BK**
 - Brain Neoplasms
 - Glioma, 78-5794
 - Ultrastructural Study, Hamster, 78-5794
 - DNA, Viral
 - Cleavage Sites, 78-5795
 - Endonucleases, 78-5795
 - Glioma
 - Proteins, Acidic, 78-5794
 - Neuroglia
 - Cell Transformation, Neoplastic, 78-5794
 - Virus, SV40
 - Immunization, 78-5812
- Virus, Polyoma, JC**
 - Antigenic Determinants
 - Radioimmunoassay, 78-5793
 - Virus, Baboon
 - Antigenic Determinants, 78-5793
 - Virus, Feline Leukemia
 - Antigenic Determinants, 78-5793
 - Virus, Mason-Pfizer Monkey

Polyoma, JC (cont'd)
 Antigenic Determinants, 78-5793

Virus, Murine Leukemia
 Antigenic Determinants, 78-5793

Virus, SV40
 Immunization, 78-5812

s, Radiation Leukemia
 Erythrocytes
 Virus Replication, 78-5853

Glycoproteins
 Virus, Recombinant, 78-5762

Hemolysis
 Blood Cells, Leukemogenic Activity, 78-5853

Sarcoma, Reticulum Cell
 Blood Cells, Leukemogenic Activity, 78-5853

Thymus Gland
 Lymphocyte Transformation, 78-5756

Virus Replication
 Hematopoietic System, 78-5756
 Lymphoid Tissue, 78-5756

s, Rauscher Murine Leukemia
 Antigen-Antibody Complex
 Cell Transformation, Neoplastic, 78-5782
 Immune Response, 78-5782
 Immune Serums, 78-5782
 Neoplasm Regression, Spontaneous, 78-5782

Antigens
 Cell Transformation, Neoplastic, 78-5782
 Sialic Acid, 78-5782

Cell Membrane
 Viral Proteins, 78-5776

Cytochalasin B
 Viral Proteins, 78-5780

Erythroleukemia
 Antigen-Antibody Complex, 78-5782
 Neoplasm Regression, Spontaneous, 78-5782

Genes, Viral
 Temperature Sensitive Mutants, 78-5780

Glucose, 2-Deoxy-
 Viral Proteins, 78-5780

Glycoproteins
 Antigenic Determinants, 78-5781
 Carbohydrates, 78-5781
 Isolation and Characterization, 78-5781

Reverse Transcriptase
 Temperature Sensitive Mutants, 78-5776

Viral Proteins
 Antibody Specificity, 78-5779
 Antigenic Determinants, 78-5778
 Genes, Structural, 78-5849
 Histocompatibility Antigens, 78-5779
 Immune Virolysis Competition Assay, 78-5779
 Isoantigens, 78-5779
 Isolation and Characterization, 78-5778
 Temperature Sensitive Mutants, 78-5780

Virus, Murine Leukemia
 Antigenic Determinants, 78-5781

Virus Replication
 Temperature Sensitive Mutants, 78-5776

Virus, Sarcoma
 Recombination, Genetic, 78-5849

irus, RD-114
 Reverse Transcriptase
 Antibody Specificity, 78-5850
 Antigenic Determinants, 78-5850
 Isolation and Characterization, 78-5850

Virus, RD-114 (cont'd)
 Virus, Baboon
 Antigenic Determinants, 78-5850

Virus Replication
 Virus, Adeno 2
 Temperature Sensitive Mutants, 78-5843

Virus, Adeno 5
 Deoxyribonuclease, 78-5845

Virus, Epstein-Barr
 Cells, Cultured, 78-5831

Virus, Friend Murine Leukemia
 Immunosuppression, 78-5763
 Rat Kidney Cells, 78-5766

Virus, Gross Murine Leukemia
 Radiation, Ionizing, 78-5769

Virus, Herpes Simplex 1
 Host Range, 78-5819
 Virus, Recombinant, 78-5819

Virus, Herpes Simplex 2
 Host Range, 78-5819
 Virus, Recombinant, 78-5819

Virus, Mason-Pfizer Monkey
 Embryonic Cells, Human, 78-5809
 Virus, Rous Sarcoma, 78-5809
 Virus, SV40, 78-5809

Virus, Moloney Murine Leukemia
 Temperature Sensitive Mutants, 78-5776

Virus, Moloney Murine Sarcoma-Leukemia
 RNA, Viral, 78-5770

Virus, Murine Leukemia
 Hematopoietic System, 78-5756
 Host Range, 78-5767
 Lymphoid Tissue, 78-5756
 Radiation, Ionizing, 78-5760
 Rat Kidney Cells, 78-5766
 Virus, Helper, 78-5756

Virus, Murine Mammary Tumor
 Cells, Cultured, 78-5752
 DNA, Viral, 78-5751

Virus, Radiation Leukemia
 Erythrocytes, 78-5853
 Hematopoietic System, 78-5756
 Lymphoid Tissue, 78-5756

Virus, Rauscher Murine Leukemia
 Temperature Sensitive Mutants, 78-5776

Virus, SV40
 Cell Fusion, 78-5796
 Chromosomes, 78-5797
 Parapharyngeal Cells, 78-5799

Virus, Rhino
 Cells, Cultured
 Isolation and Characterization, 78-5823

Virus, Cytomegalo
 Antigenic Determinants, 78-5823
 Isolation and Characterization, 78-5823

Virus, Herpes
 Antigenic Determinants, 78-5823

Virus, Ring-Necked Pheasant
 Heteroduplex Formation
 Ultrastructural Study, 78-5747

RNA, Viral
 DNA-RNA Hybridization, 78-5747

Viral Proteins
 Electrophoretic Mobility, 78-5747

Virus, Rous Sarcoma
 Heteroduplex Formation, 78-5747

- Virus, RNA**
Drosophila
 Mutagenic Activity, 78-5851
- Virus, RNA Tumor**
 Leukemia
 Cells, Cultured, 78-5831
- Virus, Rous Sarcoma**
 Antigenic Determinants
 Cell Transformation, Neoplastic, 78-5744
 Antigens, Viral
 Cell Membrane, 78-5744
 Temperature Sensitive Mutants, 78-5744
 Cell Transformation, Neoplastic
 Temperature Sensitive Mutants, 78-5744
 Lymphoma
 Virus, Marek's Disease Herpes, 78-5750
 Reverse Transcriptase
 Temperature Sensitive Mutants, 78-5745
 Sarcoma
 Albumins, 78-5746
 Muscle Proteins, 78-5746
 Virus, Marek's Disease Herpes, 78-5750
 Temperature Sensitive Mutants
 Genes, Viral, 78-5745
 Viral Proteins
 Electrophoretic Mobility, 78-5747
 Virus, Avian Myeloblastosis
 Antigenic Determinants, 78-5741
 Virus, Marek's Disease Herpes
 Co-carcinogenic Activity, 78-5750
 Virus, Mason-Pfizer Monkey
 Cell Fusion, 78-5809
 Virus Replication, 78-5809
 Virus, Recombinant
 Temperature Sensitive Mutants, 78-5745
 Virus, Ring-Necked Pheasant
 Heteroduplex Formation, 78-5747
 Virus, SV40
 Cell Transformation, Neoplastic, 78-5809
- Virus, Sarcoma**
 Virus, C-Type RNA Tumor
 Recombination, Genetic, 78-5849
 Virus, Rauscher Murine Leukemia
 Recombination, Genetic, 78-5849
- Virus, Simian Sarcoma**
 Leukemia, Lymphoblastic
 DNA-RNA Hybridization, 78-5771
 Leukemia, Myeloblastic
 DNA-RNA Hybridization, 78-5771
 Leukemia, Myelocytic
 DNA-RNA Hybridization, 78-5771
 Rhabdomyosarcoma
 Antigen-Antibody Reactions, 78-5865
- Virus, Squirrel Monkey**
 Viral Proteins
 Antigenic Determinants, 78-5814
 Virus, Baboon
 DNA-RNA Hybridization, 78-5814
 Virus, Mason-Pfizer Monkey
 DNA-RNA Hybridization, 78-5814
- Virus, SV40**
 Alkaline Phosphatase
 Cell Transformation, Neoplastic, 78-5798
 Enzymatic Activity, 78-5798
- Virus, SV40 (cont'd)**
 Antigens, Viral
 Cell-Free Synthesis, 78-5800
 Cell Transformation, Neoplastic, 78-5801
 DNA-RNA Hybridization, 78-5800
 Immunization, 78-5812
 Immunoprecipitation, 78-5801
 Isolation and Characterization, 78-5800
 Prostate, Mouse, 78-5864
 Tryptic Peptide Analysis, 78-5800, 78-5801
 Cell Fusion
 Virus Replication, 78-5796
 Cell Membrane
 Alkaline Phosphatase, 78-5798
 Epidermal Growth Factors, 78-5813
 Cell Transformation, Neoplastic
 Colony Formation, 78-5799
 DNA-RNA Hybridization, 78-5806
 Epidermal Growth Factors, 78-5813
 Growth Factors, Review, 78-5452
 Prostate, Mouse, 78-5864
 Chromatids
 Cell Transformation, Neoplastic, 78-5804
 Chromosome Aberrations, 78-5804
 Chromatin
 Nucleoprotein Complex, *Xenopus laevis* Oocytes
 78-5805
 Chromosomes
 Histones, 78-5811
 Isolation and Characterization, 78-5811
 Ultrastructural Study, 78-5797
 Viral Proteins, 78-5811
 Virus Replication, 78-5797
 Colony Formation
 Parapharyngeal Cells, 78-5799
 DNA Replication
 Agarose Gel Electrophoresis, 78-5803
 Deoxycytosine Nucleotides, 78-5803
 Temperature Sensitive Mutants, 78-5810
 DNA, Viral
 Antigens, Viral, 78-5801
 5 β -Cholan-24-oic Acid, 3 α ,12 α -Dihydroxy-,
 Monosodium Salt, 78-5802
 Chromosomes, 78-5797, 78-5811
 DNA Replication, 78-5803
 Nucleoprotein Complex, *Xenopus laevis* Oocytes
 78-5805
 Nucleotide Sequence, 78-5792
 Superhelical Turns, 78-5802
 Transformation, Genetic, 78-5796
 Fibroblasts
 Cell Transformation, Neoplastic, 78-5648
 Growth Factors
 Cell Transformation, Neoplastic, 78-5808
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Temperature Sensitive Mutants, 78-5810
 Histocompatibility Antigens
 Antigenic Determinants, 78-5812
 Transplantation Immunology, 78-5812
 Hybrid Cells
 Cell Transformation, Neoplastic, 78-5807
 Nitrous Acid
 Temperature Sensitive Mutants, 78-5810
 Picolinic Acid
 DNA Replication, 78-5808
 Growth Factors, 78-5808
 Prostatic Neoplasms

s, SV40 (cont'd)

- Acid Phosphatase, 78-5864
- Carcinoembryonic Antigen, 78-5864
- Cells, Cultured, 78-5864

RNA, Messenger

- Antigens, Viral, 78-5800

RNA, Viral

- DNA-RNA Hybridization, 78-5806

Temperature Sensitive Mutants

- B/C Gene, 78-5810
- Isolation and Characterization, 78-5810

Transformation, Genetic

- Cell Fusion, 78-5796
- Uridine, 2'-Deoxy-5-iodo-, 78-5796

Transplantation Immunology

- Mouse, Hamster, 78-5812

Ultraviolet Rays

- Temperature Sensitive Mutants, 78-5810

Urea, Ethyl Nitroso-

- DNA, Alkylation, 78-5648

Virus, Mason-Pfizer Monkey

- Cell Fusion, 78-5809
- Virus Replication, 78-5809

Virus, Polyoma, BK

- Immunization, 78-5812

Virus, Polyoma, JC

- Immunization, 78-5812

Virus Replication

- Parapharyngeal Cells, 78-5799

Virus, Rous Sarcoma

- Cell Transformation, Neoplastic, 78-5809

us, Woolly Monkey

Virus, Helper

- DNA-RNA Hybridization, 78-5772

itamin B12

Aflatoxin B1

- Hepatocarcinogenicity, 78-5470

Aflatoxin G1

- Hepatocarcinogenicity, 78-5470

Hepatoma

- Aflatoxin B1, 78-5470
- Aflatoxin G1, 78-5470

itamin E

Benzo(a)pyrene

- DNA, Binding, 78-5557

Dimethylamine, N-Nitroso-

- Aminotransferases, 78-5606
- Antioxidant Levels, Liver, 78-5606
- Hepatotoxicity, Rat, 78-5606

Vulvar Neoplasms

Eosinophils

- Neoplasm Invasiveness, 78-5889

Neoplasm Invasiveness

- Ultrastructural Study, 78-5889

Neoplasms, Multiple Primary

- Warts, 78-5888

Warfarin

- see Coumarin, 3-(α -Acetonylbenzyl)-4-hydroxy-

Warts

Vulvar Neoplasms

- Neoplasms, Multiple Primary, 78-5888

Water Pollution

Benzo(a)pyrene

- Quantitation Method, 78-5549

Dimethylamine, N-Nitroso-

- Nitrous Acid, Sodium Salt, 78-5608

Nitrosamines

- Water, Deionized, 78-5626

Plutonium

- Body Burden, Fish, 78-5735

WY-14,643

Diethylamine, N-Nitroso-

- Co-carcinogenic Effect, 78-5574

Hepatoma

- Diethylamine, N-Nitroso-, 78-5574

Kidney Neoplasms

- Adenoma, 78-5574

Liver Neoplasms

- Diethylamine, N-Nitroso-, 78-5574

Xeroderma Pigmentosum

Fibroblasts

- DNA Repair, 78-5704

Ultraviolet Rays

- DNA Repair, 78-5704

Urea, Ethyl Nitroso-

- DNA, Alkylation, 78-5648

Zinc

Esophageal Neoplasms

- Benzylamine, N-Methyl-N-nitroso-, 78-5587
- Diet, 78-5587

1

Chemical Abstracts Service Registry Number Index

0-02-2, 78-5576, 78-5752

0-06-6, 78-5481, 78-5510, 78-5517
78-5519, 78-5553, 78-5564
78-5589, 78-5593

0-07-7, 78-5426, 78-5654, 78-5662
78-5686, 78-5687, 78-5694

0-18-0, 78-5420, 78-5517, 78-5524
78-5894, 78-5898

50-22-6, 78-5576

50-23-7, 78-5509, 78-5560

50-24-8, 78-5898

50-27-1, 78-5991

50-28-2, 78-5890, 78-5991

50-32-8, 78-5406

50-37-3, 78-5422

50-44-2, 78-5857

50-56-6, 78-5543

50-81-7, 78-5530

50-89-5, 78-5599, 78-5721

50-91-9, 78-5692

51-18-3, 78-5693

51-48-9, 78-5545, 78-5713, 78-5990

51-52-5, 78-5545

51-79-6, 78-5503

53-03-2, 78-5855

53-06-5, 78-5824

53-16-7, 78-5991

53-70-3, 78-5503, 78-5548, 78-5550

53-95-2, 78-5477

53-96-3, 78-5407

54-42-2, 78-5767, 78-5796

54-85-3, 78-5407, 78-5500

55-18-5, 78-5516, 78-5571, 78-5572
78-5573, 78-5574, 78-5575
78-5576, 78-5590, 78-5592
78-5593, 78-5595, 78-5596
78-5597, 78-5601, 78-5614
78-5615, 78-5616, 78-5618
78-5620, 78-5621, 78-5655
78-5665, 78-5666

55-98-1, 78-5897

56-23-5, 78-5592, 78-5603

56-49-5, 78-5478, 78-5479, 78-5490
78-5504, 78-5506, 78-5509
78-5510, 78-5516, 78-5519
78-5520, 78-5522, 78-5523
78-5524, 78-5525, 78-5526
78-5527, 78-5528, 78-5529
78-5530, 78-5531, 78-5532
78-5553, 78-5564, 78-5740
78-5857, 78-5866, 78-5868

56-55-3, 78-5533

56-57-5, 78-5506, 78-5532, 78-5667
78-5668, 78-5669, 78-5670
78-5671, 78-5673, 78-5675
78-5694, 78-5730

56-65-5, 78-5559

57-14-7, 78-5500, 78-5861

57-41-0, 78-5430

57-57-8, 78-5666, 78-5701

57-83-0, 78-5949, 78-5993, 78-5995

57-85-2, 78-5988

57-92-1, 78-5699

57-97-6, 78-5503, 78-5504, 78-5506
78-5507, 78-5508, 78-5509
78-5513, 78-5517, 78-5518
78-5534, 78-5535, 78-5536
78-5537, 78-5538, 78-5539
78-5540, 78-5541, 78-5542
78-5543, 78-5544, 78-5545
78-5546, 78-5547, 78-5550

58-08-2, 78-5565, 78-5654, 78-5656
78-5671, 78-5703

58-15-1, 78-5636

58-55-9, 78-5518

59-02-9, 78-5557, 78-5606

59-14-3, 78-5685, 78-5686, 78-5687

59-30-3, 78-5473

59-51-8, 78-5473

59-89-2, 78-5517, 78-5593, 78-5688
78-5689

59-98-3, 78-5567

60-00-4, 78-5530

60-09-3, 78-5664

60-18-4, 78-5525

60-24-2, 78-5439

60-35-5, 78-5983

60-54-8, 78-5610

60-57-1, 78-5553

60-80-0, 78-5560

60-87-7, 78-5609

60-92-4, 78-5534

61-90-5, 78-5525

62-49-7, 78-5473

62-50-0, 78-5639, 78-5651, 78-5652
78-5654, 78-5655, 78-5656
78-5657, 78-5658, 78-5659
78-5660, 78-5665, 78-5666
78-5675, 78-5701

62-68-0, 78-5603

62-75-9, 78-5414, 78-5503, 78-5570
78-5590, 78-5591, 78-5592
78-5593, 78-5595, 78-5596
78-5597, 78-5599, 78-5600
78-5601, 78-5602, 78-5603
78-5604, 78-5605, 78-5606
78-5607, 78-5608, 78-5609
78-5610, 78-5611, 78-5614
78-5616, 78-5618, 78-5620
78-5664, 78-5665, 78-5666
78-5675

63-68-3, 78-5473

63-91-2, 78-5525

64-17-5, 78-5402, 78-5404, 78-5592
78-5596, 78-5617, 78-5621
78-5936, 78-5938, 78-5939
78-5966

64-67-5, 78-5701

64-86-8, 78-5427

66-27-3, 78-5653, 78-5659, 78-5660
78-5661, 78-5662, 78-5663
78-5664, 78-5665, 78-5667
78-5695

66-75-1, 78-5666

66-76-2, 78-5433

66-81-9, 78-5509, 78-5764, 78-5828
78-5845

66-97-7, 78-5434

67-43-6, 78-5731, 78-5733

67-64-1, 78-5596

67-68-5, 78-5983

68-12-2, 78-5983

68-19-9, 78-5470

68-26-8, 78-5557

69-78-3, 78-5743

70-18-8, 78-5472, 78-5473, 78-5559

70-25-7, 78-5507, 78-5588, 78-5598
 78-5663, 78-5664, 78-5666
 78-5675, 78-5676, 78-5677
 78-5678, 78-5679, 78-5681
 78-5682, 78-5683, 78-5684
 78-5685, 78-5699, 78-5700
 78-5701, 78-5810
 70-30-4, 78-5421
 71-43-2, 78-5409, 78-5410, 78-5437
 78-5919
 71-44-3, 78-5646
 73-32-5, 78-5525
 74-85-1, 78-5969
 74-88-4, 78-5666
 75-01-4, 78-5406, 78-5409, 78-5410
 78-5435, 78-5436, 78-5485
 78-5486, 78-5489, 78-5490
 78-5491, 78-5593, 78-5595
 78-5967
 75-09-2, 78-5596, 78-5968
 75-21-8, 78-5969
 75-35-4, 78-5492
 75-38-7, 78-5492
 75-50-3, 78-5611
 77-78-1, 78-5663
 78-98-8, 78-5998
 79-01-6, 78-5492
 79-16-3, 78-5983
 79-57-2, 78-5610
 81-81-2, 78-5433
 83-43-2, 78-5895
 83-88-5, 78-5564
 84-17-3, 78-5643
 87-86-5, 78-5421
 91-64-5, 78-5433
 93-76-5, 78-5421
 94-59-7, 78-5519
 95-54-5, 78-5425
 95-80-7, 78-5425
 96-09-3, 78-5436
 97-77-8, 78-5557
 97-90-5, 78-5449
 98-01-1, 78-5571
 98-98-6, 78-5808
 99-56-9, 78-5425
 100-42-5, 78-5436
 100-75-4, 78-5578, 78-5590, 78-5595
 78-5597, 78-5618

101-42-8, 78-5637
 103-67-3, 78-5586
 107-06-2, 78-5969
 107-07-3, 78-5969
 109-01-3, 78-5617
 110-89-4, 78-5612
 110-91-8, 78-5612
 111-44-4, 78-5969
 112-27-6, 78-5983
 117-39-5, 78-5567
 122-39-4, 78-5414
 123-33-1, 78-5656
 123-39-7, 78-5983
 124-38-9, 78-5603
 126-99-8, 78-5410, 78-5492
 127-19-5, 78-5983
 128-37-0, 78-5665
 128-53-0, 78-5743
 129-00-0, 78-5724
 131-48-6, 78-5782
 133-06-2, 78-5653
 134-03-2, 78-5636
 134-32-7, 78-5502
 140-79-4, 78-5589, 78-5664
 141-90-2, 78-5575
 142-47-2, 78-5995
 146-59-8, 78-5675
 147-94-4, 78-5477
 151-56-4, 78-5673
 153-78-6, 78-5517
 154-17-6, 78-5780
 198-55-0, 78-5568
 215-58-7, 78-5550
 298-81-7, 78-5702, 78-5703
 302-01-2, 78-5500
 302-79-4, 78-5582
 302-95-4, 78-5802
 303-47-9, 78-5468
 304-28-9, 78-5484
 305-03-3, 78-5896
 306-37-6, 78-5499, 78-5867
 319-84-6, 78-5583
 362-74-3, 78-5783
 363-24-6, 78-5509

366-70-1, 78-5662
 371-40-4, 78-5654
 434-13-9, 78-5562, 78-5684
 443-79-8, 78-5525
 446-86-6, 78-5895
 463-56-9, 78-5613, 78-5632
 505-60-2, 78-5410
 506-32-1, 78-5509
 532-32-1, 78-5635
 540-73-8, 78-5494, 78-5495, 78-5496
 78-5497, 78-5498, 78-5508
 78-5684
 542-88-1, 78-5409, 78-5410
 546-67-8, 78-5578
 551-11-1, 78-5509
 578-76-7, 78-5599, 78-5600, 78-5607
 593-60-2, 78-5492
 598-50-5, 78-5636
 604-59-1, 78-5479
 607-35-2, 78-5672
 613-13-8, 78-5517
 614-00-6, 78-5593
 614-75-5, 78-5433
 621-64-7, 78-5593, 78-5595, 78-5596
 78-5597, 78-5601, 78-5666
 623-78-9, 78-5636
 625-52-5, 78-5636
 628-36-4, 78-5493
 636-47-5, 78-5646
 637-07-0, 78-5574
 671-16-9, 78-5662
 674-81-7, 78-5680
 675-20-7, 78-5983
 684-16-2, 78-5619
 684-93-5, 78-5569, 78-5588, 78-5598
 78-5636, 78-5638, 78-5639
 78-5640, 78-5641, 78-5642
 78-5643, 78-5644, 78-5645
 78-5646, 78-5660, 78-5663
 78-5664, 78-5675, 78-5859
 745-65-3, 78-5509
 759-73-9, 78-5638, 78-5639, 78-5647
 78-5648, 78-5649, 78-5650
 78-5660
 760-23-6, 78-5492
 764-41-0, 78-5492
 865-21-4, 78-5898

924-16-3, 78-5584, 78-5595, 78-5597
 78-5601, 78-5616
 930-55-2, 78-5578, 78-5590, 78-5593
 78-5595, 78-5597, 78-5614
 78-5615, 78-5616, 78-5618
 78-5620, 78-5621, 78-5623
 78-5630
 937-40-6, 78-5586, 78-5587
 1007-36-9, 78-5637
 1083-48-3, 78-5492
 1119-49-9, 78-5636
 1126-48-3, 78-5743
 1130-69-4, 78-5670
 1162-65-8, 78-5406
 1184-78-7, 78-5611
 1239-45-8, 78-5477, 78-5517
 1309-37-1, 78-5569, 78-5876, 78-5878
 1310-58-3, 78-5579
 1314-20-1, 78-5717, 78-5973
 1332-21-4, 78-5409
 1385-95-1, 78-5470
 1405-46-5, 78-5764
 1746-01-6, 78-5421
 1912-31-8, 78-5659
 1912-32-9, 78-5659
 1921-70-6, 78-5758
 2158-11-4, 78-5636
 2381-39-7, 78-5567
 2541-69-7, 78-5511
 3105-97-3, 78-5666
 3308-64-3, 78-5523
 3342-98-1, 78-5523
 3343-07-5, 78-5523
 3343-08-6, 78-5523
 3343-10-0, 78-5523
 3343-12-2, 78-5530
 3688-53-7, 78-5946
 3719-39-9, 78-5548
 3719-40-2, 78-5548
 3817-11-6, 78-5581, 78-5582, 78-5583
 3902-71-4, 78-5703
 4239-06-9, 78-5675
 4245-77-6, 78-5598
 4408-78-0, 78-5820
 4637-56-3, 78-5668, 78-5674
 5142-23-4, 78-5663

5307-14-2, 78-5425
 5461-85-8, 78-5683
 5534-95-2, 78-5632
 5632-47-3, 78-5589
 6051-87-2, 78-5481
 6098-44-8, 78-5477
 6484-52-2, 78-5615
 6613-85-0, 78-5575
 6795-23-9, 78-5510
 6893-02-3, 78-5989, 78-5990
 6949-28-6, 78-5612
 7004-09-3, 78-5525
 7068-83-9, 78-5596
 7090-25-7, 78-5598
 7220-81-7, 78-5469
 7241-98-7, 78-5469
 7390-95-6, 78-5530
 7439-95-4, 78-5981
 7440-02-0, 78-5528, 78-5931
 7440-07-5, 78-5447, 78-5732, 78-5733
 78-5734
 7440-14-4, 78-5729
 7440-38-2, 78-5922
 7440-47-3, 78-5528, 78-5929
 7440-50-8, 78-5959
 7440-61-1, 78-5928
 7440-65-5, 78-5587
 7440-66-6, 78-5587
 7440-70-2, 78-5503, 78-5713
 7446-70-0, 78-5674
 7553-56-2, 78-5617
 7631-86-9, 78-5533, 78-5556
 7631-99-4, 78-5615, 78-5618, 78-5631
 78-5634
 7632-00-0, 78-5586, 78-5608, 78-5609
 78-5610, 78-5611, 78-5615
 78-5626, 78-5630, 78-5636
 78-5637, 78-5926
 7646-79-9, 78-5676
 7665-99-8, 78-5534
 7681-49-4, 78-5764
 7697-37-2, 78-5414, 78-5568, 78-5631
 78-5634, 78-5635
 7720-78-7, 78-5530
 7722-84-1, 78-5708
 7757-79-1, 78-5614

7757-82-6, 78-5566
 7763-77-1, 78-5436, 78-5492
 7782-44-7, 78-5648
 7782-49-2, 78-5606
 7782-50-5, 78-5403
 7782-77-6, 78-5613, 78-5631, 78-5632
 78-5633, 78-5634, 78-5635
 78-5654, 78-5810
 8001-24-9, 78-5643
 8001-28-3, 78-5677
 8002-05-9, 78-5439, 78-5960
 8002-43-5, 78-5533, 78-5724
 8007-45-2, 78-5873
 8015-30-3, 78-5992
 8015-95-0, 78-5644
 8063-94-3, 78-5488, 78-5967
 9000-07-1, 78-5644
 9000-71-9, 78-5741
 9001-45-0, 78-5552
 9001-51-8, 78-5847
 9001-73-4, 78-5741
 9001-77-8, 78-5864
 9001-78-9, 78-5798
 9001-99-4, 78-5743
 9002-60-2, 78-5576
 9002-62-4, 78-5543, 78-5545, 78-5753
 78-5990, 78-5992
 9002-71-5, 78-5713, 78-5989, 78-5990
 9002-72-6, 78-5543, 78-5607
 9002-86-2, 78-5488, 78-5967
 9002-88-4, 78-5487
 9003-98-9, 78-5845
 9004-10-8, 78-5752, 78-5950
 9005-38-3, 78-5727
 9005-79-2, 78-5594
 9006-04-6, 78-5922, 78-5954
 9008-11-1, 78-5824
 9035-50-1, 78-5473, 78-5490, 78-5510
 78-5564
 9038-70-4, 78-5452
 9054-75-5, 78-5500
 10024-97-2, 78-5568, 78-5617
 10028-15-6, 78-5698
 10034-93-2, 78-5500
 10043-66-0, 78-5716, 78-5718
 10098-97-2, 78-5728

10118-90-8, 78-5610
 10198-40-0, 78-5725
 10595-51-4, 78-5630
 10595-95-6, 78-5592, 78-5596, 78-5621
 11000-17-2, 78-5543
 11028-71-0, 78-5741
 11085-36-2, 78-5992
 11097-69-1, 78-5480
 12001-28-4, 78-5409
 12001-29-5, 78-5409
 12035-72-2, 78-5429
 12059-95-9, 78-5444
 12172-73-5, 78-5409
 12268-14-3, 78-5928
 13010-07-6, 78-5683
 13010-08-7, 78-5683
 13010-10-1, 78-5683
 13256-07-0, 78-5597
 13256-22-9, 78-5616
 13345-21-6, 78-5552
 13463-39-3, 78-5410
 13907-45-4, 78-5928
 13907-47-6, 78-5928
 13967-73-2, 78-5727
 13981-17-4, 78-5725, 78-5726
 13982-63-3, 78-5725
 14265-45-3, 78-5635
 14448-38-5, 78-5577

14807-96-6, 78-5943
 14930-96-2, 78-5780
 15117-48-3, 78-5731, 78-5732, 78-5733
 78-5734, 78-5735
 16056-34-1, 78-5405
 16339-07-4, 78-5589, 78-5593
 16543-55-8, 78-5590
 16561-29-8, 78-5509, 78-5567
 16812-54-7, 78-5429
 17068-78-9, 78-5409
 17070-44-9, 78-5680
 17144-19-3, 78-5713, 78-5716, 78-5718
 17573-29-4, 78-5552
 20535-83-5, 78-5600, 78-5601, 78-5607
 20917-49-1, 78-5585
 21150-20-9, 78-5839
 21247-98-3, 78-5567
 21373-20-6, 78-5839
 21561-99-9, 78-5637
 22775-54-8, 78-5610
 23109-05-9, 78-5839
 23214-92-8, 78-5673
 24201-58-9, 78-5481
 24305-27-9, 78-5990
 24909-09-9, 78-5552
 24928-17-4, 78-5509
 25322-68-3, 78-5634
 25614-03-3, 78-5992
 25843-45-2, 78-5644

26628-22-8, 78-5641
 28281-76-7, 78-5544
 28322-02-3, 78-5666
 30310-80-6, 78-5616
 33868-17-6, 78-5588
 33953-73-0, 78-5565
 35089-70-4, 78-5683
 35187-18-9, 78-5505
 35187-23-6, 78-5505
 35215-31-7, 78-5505
 37114-16-2, 78-5616
 37574-47-3, 78-5564
 37691-11-5, 78-5583
 38777-13-8, 78-5598
 50892-23-4, 78-5574
 55380-34-2, 78-5589
 55556-88-2, 78-5589
 55557-00-1, 78-5589
 56484-47-0, 78-5552
 57117-24-5, 78-5598
 57303-99-8, 78-5552
 57679-08-0, 78-5598
 59963-01-8, 78-5551
 62593-23-1, 78-5598
 62641-70-7, 78-5511

Wiswesser Line Notation Index

S, 78-5922
A, 78-5503, 78-5713
F, 78-5725, 78-5726
O, 78-5725
O..G2, 78-5676
R, 78-5528, 78-5929
U, 78-5959
E..S-O4, 78-5530
E2.O3, 78-5569, 78-5876, 78-5878
78-5716, 78-5718
I, 78-5617
2, 78-5713, 78-5716, 78-5718
A..N-O3, 78-5614
MG, 78-5981
NA..F, 78-5764
NA..N-O-Q, 78-5586, 78-5608, 78-5609, 78-5610, 78-5611
78-5615, 78-5626, 78-5630, 78-5636, 78-5637
78-5926
NA..N-O3, 78-5615, 78-5618, 78-5631, 78-5634
NA2.S-O2-Q2, 78-5566
NI, 78-5528, 78-5931
NI3.S2, 78-5429
PU, 78-5447, 78-5731, 78-5732, 78-5733, 78-5734, 78-5735
PU..O2, 78-5444
RA, 78-5725, 78-5729
SE, 78-5606
SL..O2, 78-5533, 78-5556
SR, 78-5727, 78-5728
TH..O2, 78-5717, 78-5973
UR, 78-5928
ZN, 78-5587
HG-1, 78-5405
/*O2*/ , 78-5634
/*2*/ , 78-5487
AL G3, 78-5674
EIUI, 78-5492
FXFFVXFFF, 78-5619
FYFU1, 78-5492
GXGGG, 78-5592, 78-5603
GYGU1, 78-5492
GYGU1G, 78-5492
G1G, 78-5596, 78-5968
G1O1G, 78-5409, 78-5410

G1U1, 78-5406, 78-5409, 78-5410, 78-5435, 78-5436, 78-5485
78-5486, 78-5489, 78-5490, 78-5491, 78-5593
78-5595, 78-5967
G2, 78-5403
G2G, 78-5969
G2O2G, 78-5969
G2S2G, 78-5410
G2U2G, 78-5492
I1, 78-5666
KA Q, 78-5579
L B656 HHJ CMV1, 78-5666
L B656 HHJ EMV1, 78-5407
L B656 HHJ EMV1 KMV1, 78-5484
L B656 HHJ ENQV1, 78-5477
L B666J HHJ EZ, 78-5517
L B677 MV&T&J CO1 DO1 EO1 JMV1 NO1, 78-5427
L C666J EZ, 78-5517
L D6 B666J, 78-5533
L D6 B666J C J, 78-5503, 78-5504, 78-5506, 78-5507
78-5508, 78-5509, 78-5513, 78-5517, 78-5518
78-5534, 78-5535, 78-5536, 78-5537, 78-5538
78-5539, 78-5540, 78-5541, 78-5542, 78-5543
78-5544, 78-5545, 78-5546, 78-5547, 78-5550
L D6 B666J C J O, 78-5505
L D6 B666J C J Q, 78-5505
L D6 B666J J, 78-5511
L D6 B666J J QF, 78-5505
L D6 B6666 SAB TJ OQ, 78-5552
L D6 B6666 2AB TJ, 78-5406
L D6 B6666 2AB TJ FQ, 78-5552
L D6 B6666 2AB TJ GQ HQ, 78-5552
L D6 B6666 2AB TJ H, 78-5505, 78-5567
L D6 J6 C666J, 78-5550
L E5 B666 FVTTT&J E OQ, 78-5991
L E5 B666 GV OV MUTJ A CQ E FV1Q -B&ACEF
78-5824
L E5 B666 OV MU PUTJ A1 CQ E1 FV1Q FQ L1 -
B&ACEF
78-5895
L E5 B666 OV MUTJ A CQ E FV1Q FQ -B&ACEF
78-5509, 78-5560
L E5 B666 OV MUTJ A1 E1 FV1 -B&AEF, 78-5949
78-5993, 78-5995
L E5 B666TJ A1 E1 FY1&2VQ OQ -B&AEFMO, 78-5562
78-5684
L E5 B666TTT&J E FQ GQ OQ, 78-5991

L E5 B666TTT&J E FQ OQ, 78-5890, 78-5991
 L E6 C666 BV FV CU GUTTT&J DQ EQ GVZ HQ IN1&1
 MQ M RQ
 78-5610
 L E6 D6656 1A T OV PHJ R, 78-5523
 L E6 D6656 1A T PV OHJ R, 78-5523
 L E6 D6656 1A T&&&T&J OQ R, 78-5523
 L E6 D6656 1A T&&&T&J PQ R, 78-5523
 L E6 D6656 1A T&&&T&J R, 78-5478, 78-5479, 78-5490
 78-5504, 78-5506, 78-5509, 78-5510, 78-5516
 78-5519, 78-5520, 78-5522, 78-5523, 78-5524
 78-5525, 78-5526, 78-5527, 78-5528, 78-5529
 78-5530, 78-5531, 78-5532, 78-5553, 78-5564
 78-5740, 78-5857, 78-5866, 78-5868
 L G6 D6 B666J, 78-5503, 78-5548, 78-5550
 L6TJ AG BG CG DG EG FG *ALPHA, 78-5583
 L6UTJ A A BIUIY&UZUIYUIVQ C -T, 78-5582
 L6UTJ A BL/UIY&U2/ 2Q C C -T, 78-5557
 L66J BZ, 78-5502
 L666 B6 2AB PJ, 78-5724
 NCN1&NO, 78-5588
 N2 O, 78-5568, 78-5617
 OC 4-NI-, 78-5410
 OCO, 78-5603
 ONN1&R, 78-5593
 ONN1&1, 78-5414, 78-5503, 78-5570, 78-5590, 78-5591
 78-5592, 78-5593, 78-5595, 78-5596, 78-5597
 78-5599, 78-5600, 78-5601, 78-5602, 78-5603
 78-5604, 78-5605, 78-5606, 78-5607, 78-5608
 78-5609, 78-5610, 78-5611, 78-5614, 78-5616
 78-5618, 78-5620, 78-5664, 78-5665, 78-5666
 78-5675
 ONN1&1R, 78-5586, 78-5587
 ONN2&2, 78-5516, 78-5571, 78-5572, 78-5573, 78-5574
 78-5575, 78-5576, 78-5590, 78-5592, 78-5593
 78-5595, 78-5596, 78-5597, 78-5601, 78-5614
 78-5615, 78-5616, 78-5618, 78-5620, 78-5621
 78-5655, 78-5665, 78-5666
 ONN3&3, 78-5593, 78-5595, 78-5596, 78-5597, 78-5601
 78-5666
 ONN4&1, 78-5596
 ONN4&4, 78-5584, 78-5595, 78-5597, 78-5601, 78-5616
 ONN5&1, 78-5597
 ONO, 78-5613, 78-5631, 78-5632, 78-5633, 78-5634, 78-5635
 78-5654, 78-5810
 ON1&UN1, 78-5644
 OO, 78-5648
 OOO, 78-5698
 OS1&1, 78-5983
 OVR &-NA-, 78-5635


OV1 & 4-PB-, 78-5578
 OV1XQVO&1VO &-PU-, 78-5731, 78-5732, 78-5734
 QR B DNO, 78-5630
 QR BG DG EG F- 21, 78-5421
 QR-/G 5, 78-5421
 QVYZ1R DQ, 78-5525
 QVYZ1Y -L, 78-5525
 QVYZ2S1, 78-5473
 QVYZ2VQ &-NA- -DL, 78-5995
 QVIN1&NO, 78-5616
 QVIN1VQ 22, 78-5530
 QVIN1VQ2 2N1VQ, 78-5731, 78-5733
 QV1OR BG DG EG, 78-5421
 QV3R DN2G2G, 78-5896
 QV4U3U3U3U6, 78-5509
 Q2, 78-5402, 78-5404, 78-5592, 78-5596, 78-5617, 78-5621
 78-5936, 78-5938, 78-5939, 78-5966
 Q2G, 78-5969
 Q2K &Q, 78-5473
 Q2O2OQ, 78-5983
 Q4N4&NO, 78-5581, 78-5582, 78-5583
 R, 78-5409, 78-5410, 78-5437, 78-5919
 RMR, 78-5414
 SHCN, 78-5613, 78-5632
 SH2Q, 78-5439
 T B666 HKJ EZ H2 IR& LZ &E &9/26, 78-5477, 78-5517
 T C566 DO LVOJ BO1, 78-5702, 78-5703
 T C6 B5665 2AB S BX IN QN NU JH&&TTTTJ FO1 I
 KVO1 KQ LOV1 M2 E- NT F6 E596 A BN
 LM&&TTJ NVO1 Q Q2-, 78-5898
 T C6656 1A P GN LM CUTT&&J EVN2&2 G -B, 78-5422
 T C666 BN DNVMV INJ B1YQYQYQ1Q L M -BBB
 78-5564
 T C666 BN ISJ B1Y&N1&1, 78-5609
 T D3 B556 BN EM JV MVTTT&J GO1 H1OVZ KZ L
 78-5426, 78-5654, 78-5662, 78-5686, 78-5687
 78-5694
 T D36 I666 B6 2AB U EOT&&&&J, 78-5564
 T E3 D5 C555 A D- FO KUTJ AG AG BG JG KG LG
 78-5553
 T E3 D6 B6666 2AB U FOTT&&&&J HQ IQ, 78-5551
 T F5 C6 B655 DOV GV OO QO RUT&&TTJ LO1, 78-5400
 T F5 C6 B655 DOV GV OO QOT&&TTJ LO1, 78-5469
 T F6 C6 B655 DOV GVO PO ROT&&TTJ MO1, 78-5469
 T H3 F5 C6 B655 DOV GV IO PO RO SUTT&&TTJ MO1
 78-5472
 T3MTJ, 78-5673

OTJ, 78-5969
 OTJ BG, 78-5436, 78-5492
 OTJ BR, 78-5436
 OVTJ, 78-5666, 78-5701
 M CN BUTJ B1R, 78-5567
 MVMV EHJ ER& ER, 78-5430
 NNVJ A BR& DN1&1 E, 78-5636
 NNVJ A BR& E, 78-5560
 NTJ ANO, 78-5578, 78-5590, 78-5593, 78-5595, 78-5597
 78-5614, 78-5615, 78-5616, 78-5618, 78-5620
 78-5621, 78-5623, 78-5630
 SOJ BVH, 78-5571
 SOJ BYVZU1- BT5OJ ENW, 78-5946
 SOV EHJ CQ DQ EYQ1Q, 78-5530
 SVNJV B2, 78-5743
 56 BM DN FN HNJ GZ IO1, 78-5600, 78-5601, 78-5607
 56 BM DN FN HNJ ISH, 78-5857
 56 BM DN FNVNVJ F H, 78-5518
 56 BN DN FNVNVJ B F H, 78-5565, 78-5654, 78-5656
 78-5671, 78-5703
 56 BO DO CHJ G2U1, 78-5519
 56 BVNV GUTJ CSXGGG, 78-5653
 6M DNTJ D, 78-5617
 6M DNTJ DNO, 78-5589
 6M DOTJ, 78-5612
 6MPOTJ BO BN2G2G, 78-5420, 78-5517, 78-5524, 78-5894
 78-5898
 6MTJ, 78-5612
 6MVMVJ EN2G2G, 78-5666
 6MYMVJ BUS, 78-5575
 6MYMVJ BUS F3, 78-5545
 6N CN ENJ B- D- F-/- AT3NTJ 3, 78-5693
 6N DNTJ ANO B1 DNO F1, 78-5589
 6N DNTJ ANO DNO, 78-5589, 78-5664
 6N DNTJ A1 DNO, 78-5589, 78-5593
 6N DOTJ ANO, 78-5517, 78-5593, 78-5688, 78-5689
 6NJ C- BT5NTJ ANO, 78-5590
 6NJ DVMZ, 78-5407, 78-5500
 6NTJ ANO, 78-5578, 78-5590, 78-5595, 78-5597, 78-5618
 6NVMVJ EE A- ET5OTJ B1Q CQ -A&C, 78-5685
 78-5686, 78-5687
 6NVMVJ EF A- ET5OTJ B1Q CQ, 78-5692
 6NVMVJ E1 A- ET5OTJ B1Q CQ -A&C, 78-5767, 78-5796
 6NVNJ DZ A- BT5OTJ CQ DQ E1Q, 78-5477
 6OTJ CM1 DQ EQ F1Q BO- CT5OTJ DVH DQ E BO-
 AL6TJ BQ CQ CMYZUM EQ FMYZUM
 78-5699


T6VMMVJ, 78-5656
 T6VMMVJ FHJ F2 FR, 78-5481, 78-5510, 78-5517, 78-5519
 78-5553, 78-5564, 78-5589, 78-5593
 T6VMTJ E1YQ- BL6VTJ D F, 78-5509, 78-5764, 78-5828
 78-5845
 T66 BM DN FN HNJ IS- ET5N ONJ DNW, 78-5895
 T66 BN DN GN JNJ CZ EQ H1MR DVMYVQ2VQ
 78-5473
 T66 BNJ BO EMQ, 78-5668, 78-5674
 T66 BNJ BO ENW, 78-5506, 78-5532, 78-5667, 78-5668
 78-5669, 78-5670, 78-5671, 78-5673, 78-5675
 78-5694, 78-5730
 T66 BNJ JNW, 78-5672
 T66 BO EVJ CR CQ DQ& DQ GQ IQ, 78-5567
 T66 BOVJ, 78-5433
 T66 BOVJ DYR&1V1 EQ, 78-5433
 T66 BOVJ EQ D- 21, 78-5433
 T66 BVOT&J D GG IVMYVQ1R& JQ, 78-5468
 T67 GNV JN IHJ CG G KR, 78-5575
 T8NTJ ANO, 78-5585
 VHN1&1, 78-5983
 VHIYQYQYQ1Q -BAA -D, 78-5780
 WNMVUM&N1&NO, 78-5507, 78-5588, 78-5598, 78-5663
 78-5664, 78-5666, 78-5675, 78-5676, 78-5677
 78-5678, 78-5679, 78-5681, 78-5682, 78-5683
 78-5684, 78-5685, 78-5699, 78-5700, 78-5701
 78-5810
 WNMVUM&N2&NO, 78-5598
 WNMVUM&N3&NO, 78-5683
 WSQQ &ZZ, 78-5500
 WS1&O1, 78-5653, 78-5659, 78-5660, 78-5661, 78-5662
 78-5663, 78-5664, 78-5665, 78-5667, 78-5695
 WS1&O2, 78-5639, 78-5651, 78-5652, 78-5654, 78-5655
 78-5656, 78-5657, 78-5658, 78-5659, 78-5660
 78-5665, 78-5666, 78-5675, 78-5701
 WS1&O2 2U -C, 78-5897
 ZN1&1, 78-5500, 78-5861
 ZQ, 78-5500
 ZR BNW DZ, 78-5425
 ZR BZ, 78-5425
 ZR BZ DNW, 78-5425
 ZR CZ D, 78-5425
 ZR DF, 78-5654
 ZR DNUNR, 78-5664
 ZVM1, 78-5636
 ZVM6, 78-5636

ZVN1&NO, 78-5569, 78-5588, 78-5598, 78-5636, 78-5638
78-5639, 78-5640, 78-5641, 78-5642, 78-5643
78-5644, 78-5645, 78-5646, 78-5660, 78-5663
78-5664, 78-5675, 78-5859
ZVN2&NO, 78-5638, 78-5639, 78-5647, 78-5648, 78-5649
78-5650, 78-5660
ZVO2, 78-5503
ZV1R, 78-5983
Z3M4M3Z, 78-5646
1MM1, 78-5494, 78-5495, 78-5496, 78-5497, 78-5498, 78-5508
78-5684
1MM1 &GH &GH, 78-5499, 78-5867
1M1R, 78-5586
1N1&1, 78-5611

1OSWO1, 78-5663
1UYG1U1, 78-5410, 78-5492
1U1, 78-5969
1U1R, 78-5436
1VN1&1, 78-5983
1V1, 78-5596
1X&&R BQ, 78-5665
1Y&MVR D1MM1 &GH, 78-5662
2N2&YUS&S 2, 78-5557
2OSWO2, 78-5701
2OVM2, 78-5636
3XR&R&VO2N2&2 &GH, 78-5603



ALBANY, N. Y. 1884.



1871

185

185

THE UNIVERSITY OF CHICAGO

405
12-10-11

Bro 1.

CARCINOGENESIS ABSTRACTS

BIOLOGY ABSTRACTS
J01 600-12-11

JUN 26 1979

VOLUME 16,
ISSUE 11



THE FRANKLIN INSTITUTE PRESS™

THE UNIVERSITY OF CHICAGO

CARCINOGENESIS ABSTRACTS

A monthly publication sponsored by the
NATIONAL CANCER INSTITUTE

EDITOR

GEORGE P. STUDZINSKI, M.D., Ph.D.
COLLEGE OF MEDICINE AND DENTISTRY
OF NEW JERSEY, NEWARK

ASSOCIATE EDITOR

JUSSI J. SAUKKONEN, M.D.
JEFFERSON MEDICAL COLLEGE
PHILADELPHIA

NCI STAFF CONSULTANTS

ELIZABETH WEISBURGER, Ph.D.
JOAN W. CHASE, M.S.

Literature Selected, Abstracted, and Indexed
by

Information Services Department, Science Information Services Organization

BRUCE H. KLEINSTEIN, Ph.D., J.D., Assistant Director, Biomedical Projects

RUTHANN E. AUCHINLECK, Managing Editor

The Franklin Research Center
The Benjamin Franklin Parkway, Philadelphia, Pennsylvania 19103

All rights reserved. No part of this journal may be reproduced in any form, for any purpose or by any means, abstracted, or entered into any data base, electronic or otherwise, without specific permission in writing from the publisher.

Copyright © 1979 The Franklin Institute Press

CARCINOGENESIS ABSTRACTS

VOLUME 16, ISSUE 11

CONTENTS

	Cross Reference Abbreviations	Article Numbers	Page
REVIEW	(Rev)	78-6001-78-6098	2273
CHEMICAL CARCINOGENESIS	(Chem)	78-6099-78-6355	2295
PHYSICAL CARCINOGENESIS	(Phys)	78-6356-78-6386	2349
VIROLOGICAL CARCINOGENESIS	(Viral)	78-6387-78-6486	2357
IMMUNOLOGY	(Immun)	78-6487-78-6505	2384
PATHOGENESIS	(Path)	78-6506-78-6542	2390
EPIDEMIOLOGY AND BIOMETRY	(Epid-Biom)	78-6543-78-6578	2399
MISCELLANEOUS	(Misc)	78-6579-78-6600	2409
AUTHOR INDEX			2415
SUBJECT INDEX			2423
CHEMICAL ABSTRACTS SERVICE REGISTRY NUMBER INDEX			2519
CAS REGISTRY LINE NOTATION INDEX			2525

GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations or citations of significant carcinogenesis articles collected from the current major biomedical sources of world literature. This service is provided by the National Cancer Institute through a contract with the Franklin Research Center for preparation of the publication, under Contract No. NOI-CP-75885 with the National Cancer Institute, U.S. Department of Health, Education and Welfare. Published and distributed by the Franklin Institute PressSM.

SUBSCRIPTIONS

For sale by The Franklin Institute Press, Box 2266, Philadelphia, PA 19103. Subscription rate for Volume 16, consisting of 12 monthly issues and a cumulative index is \$75.00, Domestic and \$90.00, Foreign. Check should be made payable to The Franklin Institute Press.

Maruzen Co., Ltd., 3-10, Nihonbashi 2-Chome, Tokyo 103, Japan, is the exclusive distributor of CARCINOGENESIS ABSTRACTS in Japan.

Handwritten text in the left margin, possibly a page number or date.

GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations, or citations of significant cancer articles collected from the current major biomedical sources of the world literature. This service is provided by the National Cancer Institute through a contract with Franklin Research Center for preparation of the publication.

FREQUENCY OF PUBLICATION: *CARCINOGENESIS ABSTRACTS*, Volume 16, is published in 12 monthly issues with indexes, plus a cumulative index.

ABBREVIATIONS

JOURNAL names are abbreviated according to the *List of Journals Indexed in Index Medicus, Abbreviation Listing*. If the journal is not listed in this, abbreviations are derived from the *International List of Periodical Title and Abbreviations*.

LANGUAGE of the article is indicated in parentheses after the title and is represented by a three-letter code. The source for these codes is *MARC Manuals Used by the Library of Congress*, pages 183-187.

ABBREVIATIONS used in abstracts:

angstrom(s)	mOsm	milliosmolar
adrenocorticotrophic hormone	max	maximum
adenosine diphosphate	mEq	milliequivalent(s)
adenosine monophosphate	min	minute(s)
adenosine triphosphate	ml	milliliter(s)
approximately	μl	microliter(s)
average	mm	millimeter(s)
bacillus Calmette-Guerin	mo	month(s)
twice daily	mol wt	molecular weight
degree(s) centigrade	N	normal concentration
calorie(s)	NAD	nicotinamide adenine dinucleotide
kilocalorie(s)	NADH	reduced nicotinamide adenine dinucleotide
cubic centimeter(s)	NADP	nicotinamide adenine dinucleotidephosphate
curie(s)	NADPH	reduced nicotinamide adenine dinucleotidephosphate
millicurie(s)	NCI	National Cancer Institute
microcurie(s)	NIH	National Institutes of Health
centimeter(s)	PAS	periodic acid-Schiff
central nervous system	po	orally
counts per minute	ppb	parts per billion
deoxyribonucleic acid	ppm	parts per million
median effective dose	qid	four times daily
ethylenediamine tetraacetic acid	qod	every other day
gram(s)	QO ₂	oxygen quotient
kilogram(s)	R	roentgen
milligram(s)	RBC	red blood cells (erythrocytes)
microgram(s)	RNA	ribonucleic acid
hemoglobin	rpm	revolutions per minute
hour(s)	sc	subcutaneous
intra-arterial	sec	second(s)
intra-dermal	SGOT	serum glutamic-oxaloacetic transaminase
Immunoglobulin A	SGPT	serum glutamic-pyruvic transaminase
Immunoglobulin B	soln	solution
Immunoglobulin G	TCD	tissue culture dose
Immunoglobulin M	TCD ₅₀	median tissue culture dose
increased life span	tid	three times daily
intramuscular	UV	ultraviolet
intraperitoneal	WBC	white blood cells (leukocytes)
International Unit(s)	wk	week(s)
intravenous	wt	weight
Michaelis constant	X	times
lethal dose	yr	year(s)
median lethal dose		
molar		
micromolar		

Handwritten text in the left margin, possibly a page number or reference.

REVIEW

6001 **Report of the Conference on Modification of Mutagenic and Carcinogenic Activity, Williamsburg, Virginia, October 24-26, 1977.** (Eng) De Serres, J. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC). *Mutat Res* 54(2): 197-202; 1978.

Discussion of the results of a joint US/Japanese conference discuss factors that can modify the mutagenic and/or carcinogenic activity of environmental chemicals indicated that these factors are wide-ranging and that they include genes, enzymes, disease, several compounds, metal ions, sex hormones, and vegetable extracts. (no refs.)

6002 **Occupational Carcinogens and Their Interaction with Other Aetiological Factors.** (Eng) Wainhouse, J. A. (Regional Cancer Registry, Queen Elizabeth Medical Centre, Birmingham, England). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer (New York, NY): Vol. 2(Part 1), 2404 pp.; 1601-1608; 1978.

Studies of occupational carcinogenesis must consider, in addition to duration of exposure to the carcinogen, individual factors (age, sex, ethnic origin, personal hygiene, smoking, alcohol intake, and dietary habits), and environmental factors (pollutants in the atmosphere, water or food), climatic and geographic factors, and the urban/rural subdivision. These studies are reviewed that correlate malignancies with various working situations. A study of bladder cancer incidence in the dyestuffs industry showed that the age at onset is related to the age of first exposure. A study of lung cancer in asbestos workers indicates a synergistic effect between asbestos and cigarette smoking. It is also possible for a single substance to act as a multiple carcinogen, as evidenced by a study relating scrotal epithelioma to industrial mineral oil exposure, which also induced primary carcinomas of the testis, skin and stomach. (8 refs.)

6003 **Current Views on the Mechanisms of Chemical Carcinogenesis.** (Rus) Pliss, G. B. (N. N. Petrov Ecological Res. Inst., USSR Ministry Health, Leningrad, USSR). *Vopr Onkol* 24(9): 95-107; 1978.

Current concepts on the mechanism of chemical carcinogenesis are reviewed. In the first step of chemical carcinogenesis, the carcinogen is circulated in the body and metabolized. Metabolic activation to an ultimate carcinogen is a characteristic of practically all chemical carcinogens. Enzymes are involved in the metabolism of most carcinogens. In the second step of carcinogenesis, the ultimate carcinogen is bound to the biological substrate (nucleic acids, especially DNA) of the target cells, thereby causing cell damage. Under the effect of carcinogens, the enzyme spectrum often becomes similar to the embryonal enzyme spectrum. The contact of a cell with a carcinogen usually does not lead to immediate formation of a cancer cell, but rather to the appearance of a population of precancer cells whose malignant transformation requires further exposure to carcinogenic or cocarcinogenic factors. (92 refs.)

78-6004 **Chemical Carcinogens: How Dangerous Are Low Doses?** (Eng) Maugh, T. H. (No affiliation given). *Science* 202(4363): 37-41; 1978.

The controversy over the effects of low carcinogen doses on subsequent tumor initiation is reviewed. One view is that the dose-response curve is a straight line; that is, that a tumor can be induced at any dosage. Restated, this is the "one-hit" hypothesis: cancer is a replicable change in cellular genetics resulting from the interaction of one carcinogen molecule with a critical receptor in one cell. The opposing view is that there is a threshold, or no-effect, dose of carcinogen below which tumor initiation occurs only at a chance rate. This view is the one presently held legislatively. The "one-hit" hypothesis is supported by data on radiation-induced cancer, which follows a linear dose-response to an incidence of 0.1%. A review of 151 chemicals studied at several doses in animals showed that none were consistent with a threshold model. Proponents of the threshold hypothesis state that humans are surrounded by carcinogens, and the fact that everyone does not develop cancer indicates there must be protective mechanisms. In fact, compounds such as glutathione and vitamins A and E are able to detoxify small amounts of carcinogens in the body, as long as these systems are not overwhelmed. The body itself needs small amounts of carcinogens, such as heavy metal ions or estrogens, for proper functioning. It is generally accepted that the latent period increases as the dose is reduced. Threshold hypothesis supporters claim that at small doses, the latent period may be longer than the life of the organism. Preliminary results of a large study (the megamouse study) on the effects of low doses of carcinogen indicate that more than one mechanism of carcinogenesis may be operating. (no refs.)

- 78-6005 The Salmonella/Microsome Test for Detection of Carcinogens as Mutagens.** (Eng) Ames, B. N. (Dept. Biochemistry, Univ. California at Berkeley, Berkeley, CA, 94720); McCann, J. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 2013-2017; 1978.

Advantages of the *Salmonella typhimurium*/microsome test for detecting chemical carcinogens as mutagens are reviewed. Several improvements have made the bacteria orders of magnitude more sensitive than a simple system that just detects reversion of any mutation back to wild-type or a simple forward-mutation system. It is highly probable that chemicals found to be mutagenic in this test will turn out to be carcinogens. (19 refs.)

- 78-6006 Philadelphia Scientist Suggests an Alternate Way to Detect Nitrosamines.** (Eng) Carson, H. C. (No affiliation given). *Household & Personal Products Industry* 15(10): 59-61; 1978.

Differential pulse polarography is proposed as a suitable additional means of detecting nitrosamines and nitrite ion. (3 refs.)

- 78-6007 Practical and Administrative Aspects of the Use of Carcinogenicity Screening Tests (Letter to Editor).** (Dan) Knudsen, I. (Inst. Toxicology, State Food Res. Inst., Denmark). *Ugeskr Laeger* 140(41): 2515-2517; 1978.

The reliability of short-term in vitro carcinogenicity screening tests of chemical substances is reduced by false-positive and false-negative findings. Substances suspect in the in vitro tests cannot be justifiably banned. (15 refs.)

- 78-6008 Relationship Between Mutagenesis and Carcinogenesis.** (Eng) Trosko, J. E. (Dept. Human Development, Coll. Human Medicine, Michigan State Univ., East Lansing, MI); Chang, C. C. *Photochem Photobiol* 28(2): 157-168; 1978.

Genetic influences on the regulation of cell proliferation are discussed, and an integrative theory of carcinogenesis is offered to reconcile the mutation vs epigenetic theories of carcinogenesis, the two-stage theory of carcinogenesis, and Comings' general theory of carcinogenesis. Carcinogenic initiation seems to be the result of the induction of DNA damage, its subsequent error-prone repair and/or replication, and muta-

tion fixation. Carcinogenic promotion appears to involve depression of previously mutated but repressed genes. Therefore, a few neoplasms could arise from a mutagenic event alone if the mutated gene is in the derepressed state at a time of mutation fixation. Many tumors would arise only after there had been a mutagenic event that is followed by an epigenetic alteration of these mutated genes (2-stage theory). Some tumors can arise simply by the abnormal derepression of genes in tissues at a particular developmental stage that prevents the normal regulation of genes that influence cell division. (102 refs.)

- 78-6009 Volatile Carcinogens: Occurrence, Formation, and Analysis.** (Eng) Hoffmann, D. (Natl. Cancer Inst. Disease Prevention, American Health Foundation, Valhalla, NY, 10595); Schmeltz, I.; Hecht, S. S.; Brunemann, K. D.; Wynder, E. L. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1943-1949; 1978.

Carcinogenicity data (humans and animals); levels in occupational environments, polluted air, and cigarette smoke; analytical methods for several volatile chemical carcinogens including vinyl chloride, chlorinated hydrocarbons, bis(chloromethyl)ether, N-nitrosamines, and hydrazines, are reviewed. (33 refs.)

- 78-6010 Analytical Methods for Polynuclear Aromatic Hydrocarbons.** (Eng) Schmeltz, I. (Natl. Cancer Inst. Disease Prevention, American Health Foundation, Valhalla, NY, 10595); Brunemann, K. D.; Hoffmann, D. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1973-1992; 1978.

Techniques for analyzing polynuclear aromatic hydrocarbons in tobacco smoke, air, and soil are reviewed, and the methods include various chromatographic and spectroscopic procedures, gas chromatography/mass spectrometry, and isotopic dilution. Detection data are tabulated, and some biological data are also presented. (68 refs.)

- 78-6011 Carcinogenicity Testing of Chemicals with Particular Reference to Organochlorine Pesticides.** (Eng) Reuber, M. D. (NCI Frederick Cancer Res. Center

Frederick, MD, 21501). *Sci Total Environ* 10(2): 105-115; 1978.

The testing of chemicals, particularly organochlorine pesticides, for carcinogenicity is reviewed. Classification of hepatic lesions in mice and rats, diagnostic criteria for malignant hepatic tumors in these animals, toxicity vs carcinogenicity, comparison of carcinomas and cirrhosis of the liver in animals and humans, and the significance of laboratory findings to human health are covered. (66 refs.)

8-6012 **Scared of Cancer? Don't Blame Modern Agriculture.** (Eng) Grierson, W. (Univ. Florida, Agricultural Res. and Education Center, Lake Alfred, FL). *J Farm Chemicals* 141(9): 69-76; 1978.

Whatever the causes of cancer, they long antecede the use of modern agricultural chemicals or food additives. This statement is supported by the facts that all forms of cancer except lung cancer are static or declining in the US, life expectancy has increased over the past 50 yr, there is no documented case of anyone in the US ever having been made sick by a pesticide residue legally used, and there is evidence that DDT is an anticarcinogen. (39 refs.)

8-6013 **Induced Plasma Cell Neoplasms.** (Pol) Salwa, J. (Zaklad Immunologii Nowotworow, Instytut Immunologii i Terapii Doswiadczalnej PAN, Czerna 12, 53-114 Wroclaw, Poland). *Postepy Hig Med Dosw* 31(6): 803-807; 1977.

The induction and antigenic properties of plasmacytomas are reviewed. Plasmacytomas can be induced in mice by 2,6,10,14-tetramethylpentadecane, 2,6,10,14-tetramethylhexadecane, and Abelson's virus. (51 refs.)

8-6014 **Health Problems Associated with Selected Petrochemicals: Benzene, Chlorinated Hydrocarbons, and Styrene-Butadiene Rubber.** (Eng) Lemen, R. A. (Div. Surveillance, Hazard Evaluations and Field Studies, Natl. Inst. Occupational Safety and Health, U.S. Post Office and Courthouse Building, Cincinnati, OH, 45202); Brown, P. P.; Rinsky, R. A.; Young, R. J.; Meinhardt, T. J. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1979-1891; 1978.

Hazards associated with occupational exposure to benzene, a recently recognized leukemogen; chlorinated hydrocarbons, known to be hepatotoxic and potentially carcinogenic and mutagenic to humans; and styrene-butadiene rubber, which has been connected with an increase in lymphatic and hematopoietic cancer, are reviewed. (66 refs.)

78-6015 **Structure and Carcinogenic Activity Relationships in the Benz(a)anthracene Series.** (Eng) Newman, M. S. (Chemistry Dept., Ohio State Univ., Columbus, OH). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 2093-2096; 1978.

The relationship between the structure and carcinogenicity of the 12 monomethylbenz(a)anthracenes and 9 of the possible nuclear monofluorinated-7-methylbenz(a)anthracenes is considered. It is hypothesized that two types of metabolism, a detoxification process and a carcinogenic process, are involved. Because 7-methylbenz(a)anthracene (7-MBA) is the most potent of the monomethyl derivatives, the 7-position is the spot at which detoxification is initiated. Because only 5-fluoro-7-methylbenz(a)anthracene is completely inactive, the 5-position is the spot at which the carcinogenic process is initiated. The fact that 1-, 2-, 3-, and 4-fluoro-7-methylbenz(a)anthracene are only slightly active indicates that the angular ring may also be involved in carcinogenesis. Steric effects are invoked to account for the moderate activity of 6- and 8-methylbenz(a)anthracene, for the fact that 7,12-dimethylbenz(a)anthracene is a more potent carcinogen than 7-MBA (the added methyl group at the 12-position produces strain due to overcrowding, which renders the molecule more reactive), and for the carcinogenicity of 12-methylbenz(a)anthracene, which does not contain a methyl group at the 6-, 7-, and 8-positions. The carcinogenicity of 7-MBA is caused by oxidative metabolism on the 7-methyl group, which results in the formation of an alkylating agent. The hypothesis is supported by metabolic studies using *Escherichia coli* and by carcinogenicity tests. (12 refs.)

78-6016 **Effects in Animals of Chronic Administration of Spironolactone--A Review.** (Eng) Lumb, G. (G. D. Searle & Co., 4901 Searle Pkwy., Skokie, IL, 60077); Newberne, P.; Rust, J. H.; Wagner, B. *J Environ Pathol Toxicol* 1(5): 641-660; 1978.

Studies of the effect of chronic administration of spironolactone to rats, dogs, and monkeys are reviewed. The animals were treated for up to 104 wk with daily dose levels (0-500 mg/kg) that were often higher than 100 times the recommended human dose. The pituitary, adrenals, and kidneys of

all animals showed no significant changes. Histological changes were noted in the rat liver, thyroid, and male internal genitalia. There were also alterations in monkey testes and male mammary glands. There was no evidence that spironolactone is tumorigenic or carcinogenic. (33 refs.)

- 78-6017 The Crudeness of Coal Tar.** (Eng) Rasmussen, J. E. (Dept. Dermatology, State Univ. New York at Buffalo, Children's Hosp. Buffalo, Buffalo, NY). *Prog Dermatol* 12(5): 23-29; 1978.

The toxicology of coal tars is reviewed with emphasis on their carcinogenicity. Coal tars are known to contain four types of carcinogens: complete carcinogens, organ-specific carcinogens, tumor-promoting agents and cocarcinogens, and lymphoma-inducing naphthylenes. Industrial exposure to coal tar has been associated with an increased incidence of cutaneous malignancies, primarily on the sun-exposed areas of the skin but also on the scrotum. Larger studies are needed, however, to determine the risk factor for both external and internal cancers following coal tar exposure. (69 refs.)

- 78-6018 Health Effects of Petroleum Extender Oils.** (Eng) Low, H. (Shell Oil Co., One Shell Plaza, Houston, TX, 77001). *J Elastomers Plast* 10(3): 229-240; 1978.

Experimental and epidemiological studies of the health effects of extender oils, high-boiling (315-538 C) fractions of petroleum that have been put through several refining steps, are reviewed. The skin and the pulmonary system are most likely to be exposed to these oils. Both skin painting and inhalation studies with various animal species have confirmed their carcinogenic potential. Epidemiological studies of workers exposed to oil, oil vapor, and oil mists in various occupational settings (ie, oil refineries and printing and metal working industries) have given ambiguous results. The health effects of exposure range from no apparent effect to excessive mortality. Significantly elevated mortality from cancer of the buccal cavity and pharynx was observed in one study of newspaper pressmen in the early and middle working years (20-54 yr old). Commercial pressmen showed little excess mortality from cancer of the buccal cavity, but they had a significant excess of cancer of the pancreas between ages 20 and 54 and rectal cancer after age 65. However, in another study involving 460 pressmen and 700 controls, no statistically significant increase in mortality rates was found. In a study of refinery workers, mortality from digestive cancer and all malignant neoplasms was higher in men with the lowest estimated exposure to atmospheric hydrocarbons, but respiratory cancer increased with increased exposure. (26 refs.)

- 78-6019 Lung Cancer, Smoking, and Atmospheric Pollution.** (Eng) Hansford-Miller, F. (No affiliation given). *Clean Air* 8(30): 27-31; 1978.

The premise is put forth that the increase in lung cancer during the last hundred years is not due primarily to cigarette smoking but rather to the increase in CO₂ in the atmosphere as a result of the industrial revolution and other associated factors related to the earth's atmosphere. Combustion of fossil fuels increases the amount of CO₂ discharged into the atmosphere by 2 x 10⁹ tons/yr. Up to half the amount of carbon trapped in virgin grassland is lost upon cultivation. The burning of lime in cement production is another major contributor of atmospheric CO₂. The current CO₂ level of 330 ppm in the atmosphere is estimated to be increasing at a rate of 1 ppm/yr. Human lungs evolved at a constant CO₂ level, since 1850 this level has changed radically, as have the ratios of CO₂/O₂ and 14C/12C. The latter is due to the fact that fossil fuels are essentially devoid of C14. Graphs showing there is a very strong correlation between the incidence of lung cancer mortality and the combustion of fossil fuels is suggested that the inability of humans to adapt to the rapid changing atmospheric levels has resulted in current lung cancer increases. (27 refs.)

- 78-6020 The Role of Cytochrome P-450 in the Toxicity of Xenobiotics.** (Eng) Lampe, J. (Central Institute of Molecular Biology, Dept. Biocatalysis, Acad. Sciences GDR, Lindenberger Weg 70, DDR-115 Berlin, E. Germany); Kerschak, G. *Pharmazie* 33(7): 407-411; 1978.

Initiation of the formation of active metabolites from halogenated aliphatic and aromatic hydrocarbons, polycyclic aromatic hydrocarbons, aflatoxins, nitrosamines, and aromatic amines by the cytochrome P-450-dependent enzyme system is reviewed. (101 refs.)

- 78-6021 Chapter XVII--Occupational Safety and Health Administration, Department of Labor. Part 1910 - Occupational Safety and Health Standards: Occupational Exposure to Acrylonitrile (Vinyl Cyanide).** (Eng) Brinkerhoff, G. (Office Compliance Programs, Rm. N-3112, Occupational Safety and Health Admin., Third St. and Constitution Ave., NW, Washington, DC, 20210). *Federal Register* 43(192): 45762-45819; 1978.

The final standard established by the Occupational Safety and Health Administration (OSHA) for exposure to acrylonitrile (AN; vinyl cyanide), which becomes effective November 2, 1978, is delineated in detail. Based on the results of experimental studies that have shown excess tumor incidences in AN-exposed animals and on indications of a cancer excess

Among workers exposed to AN, OSHA concluded that the compound poses a carcinogenic risk to workers. In addition, AN is mutagenic, an effect observed for other carcinogenic chemicals. The permissible limit of occupational exposure to AN was set at 2 ppm, as an 8-hr time-weighted av concentration with a ceiling level of 10 ppm for any 15-min period during an 8-hr day. In addition, an action level of 1 ppm was established as an 8-hr time-weighted av. Provision is made for specific exemptions from the standard for the processing, handling, and use of acrylic and modacrylic fibers, N-butadiene-styrene/styrene-AN resins, nitrile barrier resins, and nitrile rubbers when these materials are in the form of finished polymers. The standard requires that employee exposures be controlled by engineering and work practices, supplemented by respiratory and other protective equipment in limited situations. It also requires the measurement of employee exposure, employee training, medical surveillance, and recordkeeping. (no refs.)

78-6022 The Health of Women in the Pharmaceutical and Chemical Industries. (Eng) Cuthbert, J. W. May & Baker, Ltd., Dagenham Essex RM10 7XS, England). *Occup Med* 20(9): 601-604; 1978.

Changes in the pattern of women's employment, expansion of the chemical industry, and changes in legislation have increased the exposure of women to potential chemical mutagens, teratogens, and carcinogens, and advanced knowledge in medicine and toxicology has heightened the awareness of this exposure. (19 refs.)

78-6023 Pleural Reactions to Environmental Agents. (Eng) Anderson, H. A. (Mount Sinai Sch. Medicine, City Univ. New York, New York, NY, 10029); Selikoff, J. *Fed Proc* 37(11): 2496-2500; 1978.

Radiographically evident pleural reactions to asbestos are described. Both pleural calcification (PC) and pleural thickening (PT) are seen in chest x-rays of asbestosis patients. Several studies have indicated that there is a long latent period before the appearance of pleural changes following occupational exposure to asbestos. Both PC and PT are uncommon with < 10 yr of exposure. PT and parenchymal fibrosis begin to be seen after 10-11 yr, and PC becomes common after 30 yr. In a study of 626 household contacts of workers employed in an asbestos factory, PT was the most prevalent radiographic abnormality (19%). PC was present in 8%, minimal parenchymal changes in 16%. These changes were significantly more prevalent among the household contacts than among an age- and sex-matched control group from the same area without known asbestos exposure. The prevalence of abnormalities among household contacts increased with duration of the workers' employment. It should become evident in the next 10 yr whether the levels of asbestos exposure in people living in the vicinity of asbestos plants and in household con-

tacts of workers were sufficient to cause serious disease. (30 refs.)

78-6024 Role of Pineal Gland in Etiology and Treatment of Breast Cancer. (Eng) Cohen, M. (Clinical Pharmacology Branch, NCI, NIH, Bethesda, MD, 20014); Lippman, M.; Chabner, B. *Lancet* 2(8094): 814-816; 1978.

The hypothesis is put forth that diminished pineal gland function and the subsequent inhibited secretion of melatonin may promote the development of human breast cancer (BC). BC has a known relationship with increased levels of secreted estrogen: 65% of BC cases have estrogen and progesterone receptors, and a relationship has been shown between BC and factors indicating a long exposure to estrogens, such as early menarche, late menopause, and few pregnancies. The pineal gland regulates gonadal function by the feedback inhibitory effect of melatonin on gonadal development and maturation; in fact, a specific melatonin receptor in peripheral gonadal tissue has recently been demonstrated. Therefore, it can be assumed that altered pineal function would release estrogen from the suppressive action of melatonin, and this excess estrogen could induce BC. Several lines of evidence have been developed to support this hypothesis. The rate of pineal calcification in populations and the incidence of BC are strongly correlated. Precocious sexual maturation is seen in children with pineal tumors. Female psychiatric populations, treated with serum-melatonin-raising phenothiazines, have a lower incidence of breast cancer. Finally, studies show that melatonin may influence the development of tumors in animals. If melatonin levels are found to be related to BC, this relationship would be of great value in screening for individuals at high risk. (35 refs.)

78-6025 Radiation Leukemia in Humans. (Fre) Tubiana, M. (Institut Gustave-Roussy, F-94800 Villejuif, France). *Schweiz Med Wochenschr* 108(41): 1563-1568; 1978.

The occurrence of leukemia in humans exposed to various doses and types of radiation is reviewed. The risk of leukemia from low doses of radiation such as those used in radiodiagnosis and from total body irradiation with doses exceeding 100 rads of x-rays or lower doses of neutrons was compared. Quantitative data on the incidence of leukemia in survivors of the atomic bomb explosions in Japan, in patients irradiated for ankylosing spondylitis or gynecological pathology, or in radiologists are reviewed. The dose-effect relationship between radiation and leukemia development appears to be curvilinear for large doses. The frequency of leukemia in subjects exposed to low doses of radiation, such as persons exposed occupationally or those living in the vicinity of a nuclear power station, is tabulated from literature studies. Linear extrapolation should not be used to evaluate the risk of small doses of radiation (doses in the range of a millirad). Studies of the effect of radiation in utero give conflicting results, although it appears that the small doses used for radiodiagnosis

do not present a risk. Literature reports on the increased rate of leukemia among patients treated with combination radiation and chemotherapy are reviewed. (17 refs.)

- 78-6026 Hormonal Steroid Contraceptives: A Further Review of Adverse Reactions.** (Eng) McQueen, E. G. (Dept. Pharmacology, Univ. Otago Medical Sch., P.O. Box 913, Dunedin, New Zealand). *Drugs* 16(4): 322-357; 1978.

The adverse effects of oral contraceptives (OC) are reviewed extensively. Reports of benign hepatomas in women taking these drugs are numerous. A small number of these tumors have shown malignant change. A possible etiological association with steroid-induced hyperplasia of intracellular organelles is suggested. A positive association of sequential OC and endometrial adenocarcinoma and a negative association of OC and breast cancer are also suggested. (144 refs.)

- 78-6027 Role of Exogenous Female Hormones in Altering the Risk of Benign and Malignant Neoplasms in Humans.** (Eng) Thomas, D. B. (Fred Hutchinson Cancer Res. Center, Seattle, WA, 98104). *Cancer Res* 38(11, part 2): 3991-4000; 1978.

Clinical and epidemiological evidence for the association of exogenous estrogens and the development of neoplasms of the breast, female genital system, or liver is reviewed. In utero exposure to diethylstilbestrol can cause clear cell adenomas of the vagina and cervix. Oral contraceptives very probably cause hepatic cell adenomas but reduce the risk of cystic disease and fibroadenomas of the breast. There is also strong evidence that various forms of estrogens given around the time of menopause increase the risk of endometrial carcinoma and that sequential oral contraceptives may also be a cause of this neoplasm. There is no consistent or conclusive evidence that any form of exogenous estrogens alters the risk of neoplasms of the breast or ovary or that oral contraceptives cause focal nodular hyperplasia of the liver or cervical neoplasia. However, recent studies suggest that the risk of these neoplasms may be increased, at least in some subgroups, by exposure to one or more forms of exogenous estrogens. (98 refs.)

- 78-6028 Steroid Hormones and Endometrial Cancer.** (Eng) Siiteri, P. K. (Dept. Obstetrics, Gynecology and Reproductive Sciences, Univ. California at San Francisco Sch. Medicine, San Francisco, CA, 94143). *Cancer Res* 38(11, part 2): 4360-4366; 1978.

Factors that increase the risk of endometrial cancer, including obesity, nulliparity, late menopause, diabetes mellitus, hypertension, polycystic ovarian disease, hepatic disease, and ovarian tumors, are reviewed. Exogenous estrogens may play a causal role in the development of endometrial carcinoma

in some individuals. The increased risk among estrogen users ranges from 3 to 8 and appears to be dose- and duration-dependent. Estrogen users who develop cancer do not exhibit any of the increased risk factors. Many of the conditions that place women at increased risk of endometrial cancer (eg, obesity, hepatic disease) are associated with an elevated production of estrone in peripheral (nonendocrine) tissues from circulating androstenedione. Neither the exact tissue site nor mechanism of conversion has been established. The fact that the constitutional features that have long been associated with an increased risk of endometrial cancer are also associated with increased peripheral estrogen production is a strong argument for the participation of endogenous estrogens in the development of endometrial cancer. (74 refs.)

- 78-6029 Cervical Cancer and Its Prophylaxis (Letter to the Editor).** (Dan) Clemmesen, J. (Cancerregisteret, Copenhagen). *Ugeskr Laeg* 140(38): 2341-2343; 1978.

Epidemiological data on cervical cancer morbidity are reviewed, and the etiology of this cancer is discussed. The morbidity rates per 100,000 persons, found in the 1950's, were 38.3 in Copenhagen, 17.2 in rural areas of Denmark, 4.4 among Jewish women in New York, 17.1 among other whites in New York, and 32.8 among whites and 70.4 among blacks in other parts of the US. The low incidence of cervical cancer among Jewish women cannot be explained by circumcision. Epidemiological studies and various observations clearly show that early sexual activity and promiscuity are predisposing factors for cervical cancer. (15 refs.)

- 78-6030 Identification of Tumor Susceptibility of Individuals at High Risk for the Development of Cancer.** (Eng) Rawson, R. W. (Univ. Texas System Cancer Center, Houston, TX). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer, New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1493-1512; 1978.

Factors that can identify high-cancer-risk individuals and/or populations are reviewed, and they include host or intrinsic factors (genetic, congenital, and immune deficiency disorders), acquired nonneoplastic diseases, previous cancers), environmental factors (demographic, geographic, occupational, radiation, air pollution, viruses), and individual living patterns (tobacco, alcohol, diet, drugs, sexual practices). (7 refs.)

- 78-6031 An Epidemiological Perspective of the Biology of Cancer.** (Eng) Doll, R. (Univ. Oxford, Oxford, England). *Cancer Res* 38(11, part 1): 3573-3583; 1978.

cancer epidemiology data are reviewed to determine if they contribute to defining a mechanism of carcinogenesis. There is no evidence for genetic susceptibility to cancer as a whole, but there are differences in susceptibility to individual types of cancer. Epidemiological evidence suggests a possible role for an infectious viral agent in African Burkitt's lymphoma, Hodgkin's disease, cervical cancer, and chondrosarcomas. The incidence of several types of cancer increases after 2 yr of treatment with immunosuppressive drugs. The incidence of leukemia and lymphoreticular tumors is much greater in patients with certain immunodeficiency syndromes than in normal persons, but it is not clear whether the cancers are caused by the immunodeficiency or associated with it because of a common cause. Susceptibility to radiation-induced cancer increases with age, perhaps as a result of a progressive breakdown in the efficiency of immunosurveillance with age. Studies on tobacco smoking and lung cancer incidence indicate that the incidence increases with duration of exposure and dose, independently of age. The trend in smokers who stop smoking shows an initial fall in incidence followed by an increase, which keeps the rate ahead of that in lifelong nonsmokers. The simplest explanation for these data is that cancer production is a multistage process and that the various stages represent a series of discrete cellular changes that cumulatively result in the production of a malignant clone. (45 refs.)

6032 **Cancer and the Environment.** (Eng) Thomas, H. F. (MRC Epidemiology Unit, Cardiff, Wales). *Environ Health* 86(10): 232-233; 1978.

The relationships between the environment and human cancers, 75% to 90% of which are estimated to be environmentally caused, are considered. The cultural environment is the most important factor in cancer etiology, and it comprises smoking, drinking, and eating habits. Agents in the general and occupational environments do not appear to be important causes of cancer. In one report, it is argued that many of the most potent carcinogens have not been in use for the probable latent interval (25-30 yr) before chemical cancers become apparent. Likewise, cancers due to congenital, hereditary, iatrogenic, or viral causes appear to account for a very small proportion of the overall cancer incidence. The interaction of different agents, eg, asbestos and cigarettes, may increase the risk synergistically. (14 refs.)

6033 **Some Current Concepts in Intestinal Bacteriology.** (Eng) Moore, W. E. (Anaerobe Lab., Virginia Polytechnic Inst. and State Univ., Blacksburg, VA, 24061); Cato, E. P.; Holdeman, L. V. *Am J Clin Nutr* 31(10): 833-842; 1978.

Intestinal bacteriology and its possible relevance to colon cancer etiology are discussed. Recent studies indicate the presence in all human populations of intestinal bacteria that are

capable of metabolizing sterols to carcinogens. However, comparisons of fecal flora in high- and low-colon cancer risk populations suggest that the relative proportions of certain bacterial species within a genera may be correlated with colon cancer incidence. In low-risk African and Japanese populations and high-risk North Americans, a positive correlation was noted between colon cancer and *Bacteroides vulgatus*, *Bacteroides distasonis*, and *Peptostreptococcus productus*. There was an inverse relationship between colon cancer and *Eubacterium aerofaciens* II, *Bacteriodes fragilis*, and *Escherichia coli*. Because flora of the transverse and descending colon and rectum are in the stationary growth phase, fecal bacteria adequately represent colonic flora. Bacterial response can be affected by epinephrine release during fear or anger reactions. Diet manipulation does not appreciably change intestinal flora, but high-fat diets may be involved in colon carcinogenesis by providing different substrates for bacterial action. Although technical problems make this type of study extremely difficult, future work must concern itself with the identification of carcinogens in the intestine, the bacteria responsible for their presence, and the factors that control their production. (59 refs.)

78-6034 **Common Intestinal Bacteria in Etiology of Colon Cancer (Meeting Abstract).** (Eng) Storstebecker, T. P. (Akerbyvagen 282, Taby, Stockholm, Sweden). *Acta Med Scand Suppl* (621): 46; 1978. (2 refs.)

78-6035 **Fecal Flora in Different Populations, with Special Reference to Diet.** (Eng) Finegold, S. M. (Infectious Disease Section, Wadsworth Veterans Admin. Hosp., Los Angeles, CA, 90073); Sutter, V. L. *Am J Clin Nutr* 31(10): S116-S122; 1978.

Studies of the fecal microbial flora of different populations are summarized. Two groups of Japanese-Americans, one on a Japanese diet and the other a western diet, exhibited significant differences in the occurrence of seven intestinal species or genera. Similarly, patients with multiple adenomatous polyposis differed significantly from controls in nine types of organisms. In both studies, however, there was little overlap in the species present in high- vs low-incidence colon cancer groups, so the differences could have been the result of chance variation. In neither study was there a significant difference in presence of *Clostridium paraputrificum* organisms. In a study of Seventh-Day Adventists (SDA's), *Clostridium septicum* and *Clostridium tertium* were significantly more common in non-SDA controls than in either vegetarian or non-vegetarian SDA's. Nonvegetarian SDA's, compared with other previously studied nonvegetarian groups, showed a much higher incidence of lactobacilli and a lower incidence of *Clostridium perfringens*. These and other studies demonstrated that fecal flora are remarkably constant, with only 21 organisms or groups of organisms differing in frequency between high- and low-incidence colon cancer populations.

These organisms should be studied for their ability to produce carcinogens or cocarcinogens from substrates such as bile acids. (21 refs)

- 78-6036 Role of Diet and Nutrition in Cancer Cause, Prevention and Treatment.** (Eng) Gori, G. B. (Diet, Nutrition and Cancer Program, NCI, Building 31, Room 11A-03, Bethesda, MD, 20014). *Bull Cancer (Paris)* 65(2): 115-126; 1978.

The roles of diet and nutrition as etiologic factors in cancer incidence are reviewed. Epidemiologic evidence, including that obtained from studies of several migrant populations and ethnic groups, supports the hypothesis that different dietary constituents rather than other environmental contaminants are important in the cause of certain types of cancer. Fat consumption has been associated with some cancers, particularly bowel cancer. According to one theory, an excessive fat intake modifies the metabolism of cholesterol, bile acids, and neutral steroids in the intestine, as well as the metabolism and secretion of steroid hormones in the circulation. Bile acids secreted in the intestine could be degraded by intestinal bacteria to form carcinogens, or the altered metabolism of steroid hormones could impose unnatural burdens on cellular receptors of specific target tissues (uterus or breast). Animal studies strongly confirm the results of epidemiologic studies in humans. (74 refs)

- 78-6037 Diet and Breast Cancer: A Review.** (Eng) Hankin, J. H. (Sch. Public Health, Univ. Hawaii, 1960 East-West Road, Honolulu, HI, 96822); Rawlings, V. *Am J Clin Nutr* 31(11): 2005-2016; 1978.

Human breast cancer (BC) is a multifactorial disease that is probably influenced by dietary intake. Studies of geographical variations in diet have shown that North America and northern Europe are high risk areas for BC, but countries in Africa and Asia are relatively low risk areas. The association of early menarche with higher risk and early first pregnancy with lower risk suggest that tumor initiation or inhibition may be determined in adolescence. Obesity may contribute to BC by influencing hormone synthesis and metabolism. Although dietary fat has been implicated in BC, dietary proteins, cholesterol, carbohydrates, and calories may also be involved. Studies of mammary tumor development in rodents indicate that tumorigenesis is associated with polyunsaturated rather than saturated fats; however, nutritional factors, including fat type, are more likely to act as promoters rather than initiators of tumorigenesis. Endocrinologic data indicate that women with a low ratio of estradiol to estrone and estrone are at high risk for BC. In rats, a high-fat diet increased serum prolactin levels and promoted mammary tumor development by 7,12-dimethylbenz(a)anthracene, but studies of prolactin levels in BC patients have yielded inconsistent results. The

need for further research in all of these areas is emphasized. (91 refs)

- 78-6038 Workshop V--Fiber and Cancer: Summary and Recommendations.** (Eng) Burkitt, D. P. (Filiation given). *Am J Clin Nutr* 31(10): S213-S215; 1978.

Mechanisms whereby lack of fiber may promote colorectal cancer or whereby fiber may protect against this cancer are reviewed briefly. Applicable fecal parameters include neutral steroids, bile salts, and lipids; ammonia levels; pH; enzyme activities; mutagenicity/carcinogenicity; transit time; and water content; and microbial flora. (no refs)

- 78-6039 Recent Progress in Dietary Fiber (Plantix) and Human Nutrition.** (Eng) Spiller, G. A. (Syntex Res., Palo Alto, CA); Shipley, E. A.; Blake, J. A. *Crit Rev Food Sci Nutr* 10(1): 31-90; 1978.

Research performed since 1975 on the role of dietary fiber (plantix) in human nutrition is reviewed. A high low-plantix diet has been correlated with a high incidence of colorectal cancer. The significance of plantix may not be an affecter of transit time/constipation but rather as a diluent of colonic contents. However, other factors such as naturally occurring antioxidants (vitamins E and C, selenium, etc.) and vitamin A may also play a key role in the etiology of colorectal cancer. Many plantix polymers have a great capacity to interact with other dietary components. A study of 198 I colon cancer patients showed that they had a significantly lower fiber consumption. Study of the etiology of colorectal cancer in Chile, where the incidence is very low, revealed that the daily diet is a high-carbohydrate, low-protein, low-fat, one rich in plant fibers. One hypothesis advanced is that plantix-associated factors, ie, lectin, may be involved in protecting against colon cancer. Several possible mechanisms by which the body's response to undigested lectin in the colon would protect against intestinal cancer are given. Although many factors probably contribute to colon cancer, a higher intake of plantix and antioxidants may help to prevent the disease. (161 refs)

- 78-6040 Interaction of Dietary Fiber with Other Dietary Components: a Possible Factor in Certain Colorectal Cancer Etiologies.** (Eng) Spiller, G. A. (Syntex Res., 3401 View Ave., Palo Alto, CA, 94304). *Am J Clin Nutr* 31(10): S231-S232; 1978.

The concept that dietary fiber, or "plantix" (ie, all polymers of plant materials that are indigestible by human digestive enzymes) is an integral part of gastrointestinal physiology which cannot be studied purely on the basis of its presence or absence, is discussed. Plantix must be considered in the

its interaction with other nutrients, and this pattern must be precisely identified. Plantix may affect the feces by altering pH, general composition, or transit time. It may affect colonic microflora directly, serving as a nutrient for certain organisms, or it may carry other constituents to the colon that would not arrive there otherwise. It may alter pH, the elimination rate of microfloral metabolites or toxins, or dilute carcinogens. Finally, it can alter the location and absorption rate of nutrients and the reabsorption of metabolites along the intestine. All these factors suggest that a more complex means of studying plantix is necessary, one that considers the interactive facets of this dietary component. (3 refs)

78-6041 The Relationship Between Bowel Cancer and Fiber Content in the Diet. (Eng) Walker, A. R. (Human Biochemistry Res. Unit, South African Inst. Medical Res., P.O. Box 1038, Johannesburg 2000, South Africa). *Am J Clin Nutr* 31(10): S248-S251; 1978.

A review of data on colon cancer in South African blacks (SAB) indicates that the disease is very uncommon in this population. This rarity is not attributed to lack of adequate case reporting since high and/or increased incidence rates have been reported for cancers at certain other sites. There has been an increase in standard of living and urbanization of SAB; but in spite of the westernization in diet and way of life, the risk of colon cancer remains the same. Since the dietary changes which accompany urbanization (decreased crude fiber in diet, longer fecal transit time, lower frequency of defecation and less feces voided per day) have been correlated with increases in colon cancer elsewhere, intrinsic factors seem to be protecting SAB. The existence of this protective factor is suggested by the unchanged incidence of appendicitis, another disease which is usually seen to increase after urbanization affects diet. Further research is needed on bowel physiology fecal chemical composition, and enzymatic markers of intestinal microfloral activity in SAB. (19 refs)

78-6042 Diet and Cell Growth Modulation by Ammonia. (Eng) Visek, W. J. (Sch. Basic Science, Dept. Food Science, Univ. Illinois, Urbana, IL, 61801). *Am J Clin Nutr* 31(10): S216-S220; 1978.

The role of dietary fiber in the production and metabolism of ammonia by host cells, digestive enzymes, and gastrointestinal microflora is discussed. Fiber passes through the small intestine to the cecum and colon, where bacteria use it as an energy source for protein synthesis, in which the necessary nitrogen is supplied by ammonia. By facilitating the incorporation of ammonia into protein or its dilution, fiber reduces the harmful effects of ammonia on the gastrointestinal mucosa. Furthermore, fiber fermentation yields fatty acids that decrease colon pH and provide anions to trap free ammonia. Ammonia can induce cell proliferation, which enhances neoplastic transformation by increasing errors in DNA copying,

increasing code breaking actions of mutagens present in sub-threshold doses, or unmasking latent DNA changes caused by mutagens. Ammonia also alters cell life-span and metabolism, and enhances virus production. Although it does not transform cells per se, the incidence of polyps and cancers is highest in colonic areas with the highest ammonia concentrations. The role of fiber in removing free ammonia from the colon may explain the low incidence of colon cancer in populations with high-dietary fiber. (48 refs)

78-6043 Colonic-Rectal Cancer: Fiber and Other Dietary Factors. (Eng) Burkitt, D. P. (Unit Geographical Pathology, St. Thomas' Hosp., London SE1 7EH, England). *Am J Clin Nutr* 31(10): S58-S64; 1978.

The epidemiological evidence supporting the low fiber-high fat hypothesis for colon cancer etiology is comprehensively reviewed. Large bowel cancer bears a direct and constant relationship to economic development and westernization. Furthermore, within high incidence locales, there is an urban-rural correlation, with colon cancer being more prevalent in the urban areas. Chronological changes have also been documented; eg, the incidence rate in American blacks has increased significantly in the past generation. Groups that have immigrated from low to high incidence areas, notably the Japanese in Hawaii and California, show an increase in cancer incidence. Occurrence of other diseases, such as adenomatous polyps, diverticular disease, hiatus hernia and appendicitis, correlates with appearance of colon cancer, suggesting that all may result, at least to some degree, from common factors. Investigation has frequently centered on differences in fecal composition as a result of westernization and the accompanying decrease in dietary fiber and increase in fat intake. Hypotheses that have been put forth to explain how these dietary changes could result in cancer include decreased transit time, decreased dietary dilution, accumulation of toxins and carcinogens, or interactions between bacterial flora and dietary constituents. (52 refs)

78-6044 Models Relating Ultraviolet Light and Non-melanoma Skin Cancer Incidence. (Eng) Green, A. E. (ICAAS, Univ. Florida, Gainesville, FL, 32611); Hedinger, R. A. *Photochem Photobiol* 28(2): 283-291; 1978.

Various approaches used to relate UV light doses to non-melanoma skin cancer (SC) incidence are reviewed. Three spectral response functions (DNA, Coblenz-Stair erythema response, and Robertson-Berger meter response) that have been used in epidemiological UV-SC studies are illustrated. If the DNA response function is the relevant one in SC, as seems most reasonable from a biophysical viewpoint, then a true dose-response relationship could depart significantly from an apparent relationship obtained by comparing meter readings with incidence data. Based on standard statistical

or epidemiological approaches, a double power law model seems to be as good a dose-response relationship as can be determined from Third National Cancer Survey data, which span a narrow range of annual doses. Various inductive approaches to dose-response relations are also reviewed. Despite the inconsistency of epidemiological data collected in different parts of the world, the data tend to favor multiple cause or age-exposure models over reciprocity, dose potency, and separable models. Both favored models are biophysically reasonable. Thus, the notion that there is more than one cause of SC is also suggested by evidence pointing to chemical carcinogens. The idea that age specific incidence rates of SC might depend on age and exposure is compatible with modern ideas on the degradation of immunological defenses. The application of the mathematical machinery of Reliability Theory to models relating UV light and nonmelanoma SC incidence is also illustrated. (54 refs)

- 78-6045 Ultraviolet Light Mutagenesis in *Escherichia coli* (Meeting Abstract).** (Eng) Volkert, M. R. (Dept. Molecular Biology, Univ. California, Berkeley, CA, 94720). *Can J Genet Cytol* 20(3): 438; 1978. (no refs)

- 78-6046 Solar Radiation and Skin Cancer.** (Nor) Fyrand, O. (Rikshospitalet, Hudavdelingen, Norway). *Tidsskr Nor Laegeforen* 98(24): 1139-1140; 1978.

In this literature review, the relationship between exposure to UV rays, natural or artificial, and precancerous changes in the skin and skin cancer is confirmed. The effect of the exposure is cumulative. A causal relationship between exposure to light and malignant skin melanoma is probable. (13 refs)

- 78-6047 Comparison of Relative Risk from Radiation Exposure and Other Common Hazards.** (Eng) Kelsey, C. A. (Radiology Dept., Univ. New Mexico, Albuquerque, NM, 87131). *Health Phys* 35(2): 428-430; 1978.

Three comparisons are presented that have proven useful in putting effects from accidental or occupational exposure to radiation in perspective and in assisting radiation workers in evaluating the relative risks due to radiation exposure: (1) comparison of hazards from radiation exposure to hazards from various accidents in the US, (2) relative accident rates for various occupations in the US, and (3) comparison of the health hazards associated with cigarette smoking and radiation. Results are reported in terms of hazard/rem (roentgen equivalent man) exposure, assuming that there is no recovery from radiation injury and that the effect of 1 rem is the same whether delivered over periods of 1 mo, 1 yr, or 5 yr. Deaths per yr per million persons are 9 for each rem of radiation compared to 4,780 for cardiovascular disease. Among 15 oc-

cupations, radiation workers have the lowest risk of job-related accidental death (0.02 relative death rate) and miners highest (6.2). The life shortening effect of smoking one cigarette is equivalent to exposure to 2.5 mm radiation. (10 refs)

- 78-6048 A Stochastic Model of Carcinogenesis Incorporating Certain Observations from Chemical and Radiation Dose-Response Data.** (Eng) Brodsky, A. (Office of Standards Development, U.S. Nuclear Regulatory Commission, Washington, DC, 20555). *Health Phys* 35(3): 421-428; 1978.

A stochastic model of carcinogenesis that incorporates phenomena regarding the sequential interactions of carcinogens is presented. There is preliminary evidence from animal experiments that this model is consistent with observed quantitative dose-response relationships. An example is dose-response relationships for radiation carcinogenesis (tumors in mice implanted with Ru^{106} pellets or administered Ce^{144} intratracheally). The two-step sequential stochastic model may provide an alternative model to the log-normal model for interpreting a broad range of radiobiological toxicological effects. (9 refs)

- 78-6049 The Combined Effects of Ionizing Radiation and Smoking in the Development of Bronchogenic Carcinoma.** (Ger) Schuttmann, W. (Hauptabteilung Strahlenschutzmedizin, Staatlichen Amtes für Atomsicherheit Strahlenschutz der DDR, Waldowallee 117, DDR-1157 Berlin, E. Germany). *Z Erkr Atmungsorgane* 150(3): 243-247; 1978.

Evidence that smoking increases the probability of respiratory tract cancer developing in individuals previously exposed to ionizing radiation is reviewed. Uranium miners who smoked had a respiratory tract cancer rate of 17/10,000 person-yr, but nonsmoking miners had a rate of only 1.7/10,000 person-yr. Miners who smoked more than one pack of cigarettes/day had a significantly shorter average tumor-latency period than nonsmokers. This type of effect is typical of that of a tumor promoter. The finding that the same proportion of small cell and epidermoid lung cancers occurs in smoking and nonsmoking uranium miners also suggests that smoking has a promoting or cocarcinogenic effect. These findings have significance for the nonmining population. The average annual radiation dose to the lungs from natural sources (ie, burning of fossil fuels, presence of ^{210}Pb and ^{210}Po in tobacco smoke) is 250.5 millirems (rem = roentgen-equivalent-man) for the general population. Thus, every smoker suffers the combined effects of radiation and smoking. (15 refs)

- 78-6050 Can Animal and In Vitro Studies Give New Relevant Answers to Questions Concerning the Mechanisms of Carcinogenesis?** (Eng) Brodsky, A. (Office of Standards Development, U.S. Nuclear Regulatory Commission, Washington, DC, 20555). *Health Phys* 35(3): 429-430; 1978.

Mammographic Screening for Human Breast Cancer? (Eng) Thlefsen, L. A. (Section Radiation and Tumor Biology, Dept. Radiology, Univ. Utah Medical Center, Salt Lake City, UT, 84132); Brown, J. M.; Carrano, A. V.; Nandi, S. *J Natl Cancer Inst* 61(6): 1537-1545; 1978.

This report of an ad hoc committee appointed by the National Cancer Task Force reviews current arguments surrounding the role of mammography in breast cancer development and recommends directions for future research using animal or in vitro models. Human carcinogenesis studies have shown that the risk of breast cancer from ionizing radiation is not diminished by dose fractionation and is highly age dependent, resulting in an excess risk of 7.5 cases/10⁶ women/rad for women <35. In rats, fractionation of a 500 R dose did not change the overall incidence of mammary tumors, but it increased adenocarcinoma incidence and decreased fibroadenoma incidence. Hormonal factors modify radiation-induced mammary cancer in rats, and postpubertal animals are at higher risk than prepubertal animals. On the basis of current knowledge of cell transformation mechanisms and the results of present study protocols, two specific areas of future research are suggested: the specific effects of radiation dose fractionation on the efficiency of cell transformation, and the relationship between DNA repair processes and transformation. The mammary fat pad transplantation techniques and established in vitro cellular transformation systems is suggested. No further animal studies aimed at quantifying the tumor risk per rad should be initiated at this time. (58 refs)

8-6051 Iatrogenic Carcinogenesis. (Hun) Schuler, D. (II. Gyermekklinika, Semmelweis Orvostudományi Egyetem, Budapest, Hungary). *Orv Hetil* 119(37): 2239-2243; 1978.

Ionizing radiation and drugs with carcinogenic, cocarcinogenic or immunosuppressive effects lead to iatrogenic carcinogenesis. These drugs include cyclophosphamide, busulfan, chlorambucil, melphalan, 6-azauridine, chlornaphazin, arsenicals, hydantoin derivatives, phenacetin, diethylstilbestrol, and 18-methylnorethisterone. (92 refs)

8-6052 Multiple Pathways of DNA Repair in Bacteria and Their Roles in Mutagenesis. (Eng) Smith, K. C. (Dept. Radiology, Stanford Univ. Sch. Medicine, Stanford, CA, 94305). *Photochem Photobiol* 28(2): 121-129; 1978.

Photoreactivation, excision repair, and postreplication repair of DNA in bacteria are reviewed in relation to mutagenesis, which appears to be the result of error-prone DNA repair. There is growing evidence that carcinogenesis is also the result of error-prone DNA repair. One pathway of postreplication repair has been shown to be blocked by postirradiation treatment with the protein synthesis inhibitor chloram-

phenicol (CAP), suggesting that protein synthesis is required after UV irradiation before this pathway can function. This pathway requires functional *uvrD*, *recB*, and *lexA* genes. UV-induced mutagenesis is blocked by postirradiation treatment with CAP, suggesting that the *uvrD*, *lexA*, and *recB* gene products are involved in mutagenic repair. The fact that UV-induced mutations are not observed in *lexA* strains seems to support the idea that the CAP-inhibitable pathway is the mutagenic pathway. An area of mutagenesis research that is currently receiving considerable attention is based on the concept that error-prone (mutagenic) repair is due to a new repair function (the so-called SOS repair system) that is induced by damage produced in DNA. Since UV-induced mutagenesis is prevented by a short postirradiation treatment with CAP, postirradiation protein synthesis is apparently required for the expression of UV-induced mutations. The *recA* and *lexA* strains are not UV mutable; therefore, *recA* + *lexA* + functions are probably essential for expressing this inducible error-prone repair function. (75 refs)

78-6053 Evaluation of DNA Damage and Repair in Mammalian Cells Exposed to Chemical Carcinogens. Methods Potentially Useful as Short Term Pre-screening Tests. (Eng) Brambilla, G. (Dept. Pharmacology, Univ. Genoa, Genoa, Italy); Cavanna, M.; Parodi, S. *Pharmacol Res Commun* 10(8): 693-717; 1978.

Autoradiographic techniques for detecting unscheduled incorporation of labeled nucleotides into mammalian DNA and techniques for detecting single-strand DNA breaks in alkali are discussed in relation to their ability to predict the oncogenic potential of chemical agents. Although potentially good indicators of carcinogenic risks in humans, carcinogenicity tests in rodents present problems related to possible species differences in the metabolic activation of procarcinogens. This method is also time-consuming and expensive. Mutagenicity tests such as the Ames assay provide rapid information, and a short-term test based on DNA damage-repair in mammalian cells would be a useful complementary test. Of the short-term tests, the evaluation of DNA damage in vivo by the alkaline elution technique may prove to be the method of choice. To confirm its predictive value, the test should be extended to the evaluation of several hundred carcinogenic and noncarcinogenic compounds of different chemical structures. (122 refs)

78-6054 A Re-evaluation of Evidence for Carcinogen-induced DNA "Repair" Synthesis. (Eng) Melzer, M. S. (2550 Kemper Road, Shaker Heights, OH, 44120). *Chem Biol Interact* 22(2/3): 359-363; 1978.

The argument that carcinogenesis involves primary attack on nuclear DNA in the crucial reaction of the transforming process is based mainly on studies concerning 'repair' or unscheduled DNA synthesis. If it can be demonstrated that

upon treatment with a carcinogen, there is additional DNA synthesis (increased incorporation of labeled thymidine into DNA) of a nonreplicative type, this would indicate that a lesion has been introduced into the DNA that was being repaired, suggesting direct carcinogen-DNA interaction as the crucial step in carcinogenesis. This study evaluates the evidence for such carcinogen-induced repair synthesis and suggests that some of it is distinctly flawed. In fact, some of the evidence could just as well be interpreted as evidence for carcinogen-induced replicative DNA synthesis, which would point up the fact that carcinogens, in their initial effect on target cells, induce increases in (or even initiation of) new replicative DNA synthesis and/or new mitosis. It is hoped that the points raised will stimulate further thinking among carcinogenesis researchers about possible alternative mechanisms of initial cellular attack by chemical carcinogens. (27 refs)

- 78-6055 DNA Repair and Mutagenesis in Mammalian Cells.** (Eng) Hart, R. W. (Dept. Radiology, Ohio State Univ. Sch. Medicine, Columbus, OH, 43210); Hall, K. Y.; Daniel, F. B. *Photochem Photobiol* 28(2): 131-155; 1978.

Mammalian cell DNA repair is reviewed in relation to mutagenesis and carcinogenesis. Although DNA damage leads to measurable mutations in mammalian cells, the means by which this damage results in an inheritable change in base sequencing is not known. Depending on the type of insult and its location within the DNA, inheritable errors could arise during excision repair, DNA replication, or postreplication repair. Among the clinical syndromes associated with mutagenesis that have been studied most extensively are xeroderma pigmentosum (XP), dyskeratosis congenita, Fanconi's anemia, ataxia telangiectasia, and Werners, Bloom's, Chidiak-Higoshi, Hutchinson-Gilford, and Down's syndromes, all of which have a genetic predisposition to cancer and/or other specific syndrome anomalies. XP is characterized by hypersensitivity of the skin to solar radiation, a high incidence of multiple carcinomas, and deficiencies in DNA repair. The predisposition to cancer among XP patients is inversely proportional to their ability to repair UV radiation- and acetoxyacetylaminofluorene-induced damage. An explanation for the high frequency of mutation in XP variants, which are slow in postreplication repair but have normal excision repair, involves the possibility that the DNA polymerase III-like complex is altered, leading to the observed mutation frequencies. In bacterial systems, mutations in the gene for DNA polymerase III can lead to enhancement of spontaneous mutations in vivo. (326 refs)

- 78-6056 Glycopeptide Changes and Malignant Transformation: A Possible Role for Carbohydrate in Malignant Behavior.** (Eng) Warren, L. (Wistar Inst. Anatomy and Biology, 36th St. at Spruce, Philadelphia, PA,

19104); Buck, C. A.; Tuszyński, G. P. *Biochim Biophys Acta* 516(1): 97-127; 1978.

Studies comparing glycopeptides (GP) isolated from control and transformed animal cells are reviewed. Sephadex chromatography of Pronase-treated GP yields, among other fractions, groups A and B GP, which elute at mol wt approx 4,600 and 3,700, respectively; upon removal of the sugar, these peptides are reduced to a common size but retain a common substructure. After Pronase digestion, glycoproteins producing groups A and B GP have been found associated with nuclei, mitochondria, lysosomes and endoplasmic reticulum. Group A material is increased in malignant cells regardless of transforming agent or species of origin; it is strongly growth-dependent. Its presence has been related with tumorigenesis and noted both in vivo and in vitro. The total number of GP may be constant in normal and transformed cells, and as the number of group A GP increases, the number of group B GP decreases. Studies made of GP derived from individual glycoproteins of identical mol wt and isoelectric point from control and transformed cells. Among 24 different pairs of homologous glycoproteins, 20 differences of varying degrees of severity were noted. Apparently, the gross increase of group A GP seen upon Sephadex chromatography is the cumulative result of several changes in the carbohydrate components of many glycoproteins in crude fractions. It is hypothesized that these changes, particularly in the cell membrane, where most glycoprotein activity is found, do not endanger premalignant cells but alter their relationships with external factors and influences. This alteration could then lead to malignant change, perhaps by removing the cells' sensitivity to these external controlling factors. (188 refs)

- 78-6057 High Molecular Weight, Cell Surface-associated Glycoprotein (Fibronectin) Lost in Malignant Transformation.** (Eng) Vaheri, A. (Dept. Virology, Univ. Helsinki, 00290 Helsinki 29, Finland); Mosher, D. *Biochim Biophys Acta* 516(1): 1-25; 1978.

The literature on fibronectin (FBN), a polymorphic glycoprotein found in blood and tissues of vertebrates and in structures of adherent vertebrate cells, is reviewed, and the significance of its loss from surfaces of transformed cells in vivo and the relationship of FBN to the biology of cancer cells in vivo are discussed. Two forms of FBN have been identified: a high-mol wt, cell-surface-associated, transformation-sensitive form (I) and a high-mol-wt circulating form (II). Form I is decreased consistently in cells transformed by virus, including virus mutants temperature-sensitive for transformation. Cells from human glioblastomas, fibrosarcomas, rhabdomyosarcomas, and osteosarcomas produce FBN but have little form I FBN. Compared with normal parental cells, form I is also decreased in spontaneously and chemically transformed rodent cells. Among adenovirus-transformed embryo cells, there is a close correlation between a decrease in cells positive for FBN immunofluorescence at cell-cell con-

ect and an increase in tumorigenicity in syngeneic rats and mice. Although, there are exceptions, the conclusion that malignant cells may have a normal complement of surface-associated FBN seems premature when these exceptions are compared with well-characterized virally transformed or tumor-derived cells of genetically stable species. Transformation appears to be associated with alterations of FBN, sulfatoglycosaminoglycans, and collagen, but it is not known how these components interact in vivo and in vitro. A decreased association to matrix components by malignant cells may be crucial to unregulated growth. It is suggested that transformed cells have defective interactions and associations with matrix elements and thus are free to divide and move. Little is known of the role of form II FBN in neoplasia. Changes of this form are not common in patients with malignancies. (92 refs)

78-6058 Stress and Cancer: Fresh Perspectives. (Eng) Riley, V. (Pacific Northwest Res. Foundation, Seattle, WA, 98104). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1769-1776; 1978.

Studies of the relationship between stress and cancer are reviewed. According to epidemiological studies, religious groups such as Mormons and Seventh-Day Adventists have a reduced incidence of cancer even at anatomical sites not generally associated with the effects of smoking or alcohol consumption. This suggests the beneficial influences of a less stressful life-style upon pathological processes. Psychosocial screening questionnaires have been employed in an effort to correlate personality traits and cancer risk. A multihormonal approach for deciphering the effects of stress has led to the quantification and integration of the behavior of a broad spectrum of interrelated hormones associated with stressful events. The known interference of stress with immunocompetence appears to be the most promising rationale for the induction of disease by acute, chronic, or intermittent stress. Certain varieties of stress, however, suppress specific neoplastic processes rather than enhancing them. There is no explanation for this paradox. The enhancement of tumor growth or of tumor incidence by stress probably can only be observed in experimental circumstances in which the host exerts a total or partial controlling influence on the establishment or growth rate of the tumor. It is the disruption of such host control that permits stress to enhance pathological processes. (17 refs)

78-6059 Etiology of Cancer as Studied in the Platyfish-Swordtail System. (Eng) Anders, A. (Genetisches Institut der Justus-Liebig-Universität Giessen, Hein-

rich-Buff-Ring 58-62, D-6300 Giessen, W. Germany); Anders, F. *Biochim Biophys Acta* 516(1): 61-95; 1978.

Research in cancer etiology utilizing the platyfish-swordtail (Xiphophorus fish) tumor system is reviewed. Although inbreeding of wild-type populations produces only normal offspring, interspecific and interpopulational crossings produce progeny that spontaneously develop various neoplasms, including melanomas, pterinophoromas, neuroblastomas, thyroid carcinomas, kidney carcinomas, and reticulosarcomas. Genetic information coding for neoplastic transformation is represented by a "tumor gene" (Tu), which may be present in several copies on different chromosomes, including one copy on the terminal portion of the sex chromosome. Tu is normally repressed by cell-type-specific systems of regulating genes. Four types of neoplasm-triggering mechanisms have been recognized in this model, all mediated by Tu. These include altered cell differentiation, somatic mutation, germ line mutation, and hybrid crossings. However, neoplasms are usually initiated by a combination of tumor-inducing methods, and the ultimate type of cancer is always determined by the last step in the multistep process. The different cell stages of transformation in the melanophore system to melanoma have been well-characterized. Certain melanomas are preferentially located in certain compartments of the fish body, and 13 different genes governing 13 different compartments of melanoma expression are linked to the Tu. Likewise, the malignancy and benignancy of melanoma are apparently regulated by presence of the single gene R-Diff. Experiments on neoplasia etiology in this tumor system support generally accepted concepts for the origin of cancer and disclose relationships among the different concepts. (177 refs)

78-6060 Specificity of Arrest, Survival, and Growth of Selected Metastatic Variant Cell Lines. (Eng) Nicolson, G. L. (Dept. Developmental and Cell Biology, Univ. California, Irvine, CA, 92717); Brunson, K. W.; Fidler, I. J. *Cancer Res* 38(11, part 2): 4105-4111; 1978.

Animal tumor models for blood-borne metastasis that were developed by in vitro cloning or in vivo selection of malignant tumor cell populations to obtain organ-preferring variant tumor cell lines with altered arrest, survival, invasion, and growth properties are reviewed. Selection and some tumor cell characteristics of lung-, brain-, and ovary-colonizing metastatic B16 melanoma variants, liver-colonizing RAW117 lymphosarcoma variants, and lung-colonizing sarcoma variants are discussed. Cloning and fluctuation tests with the B16-F1 melanoma cell line and with a vasoformative sarcoma, MSV3T3-LN, demonstrated the presence of populations of high and low metastatic potential before in vivo selection, suggesting that each in vivo selection results in enrichment of more metastatic cells in each subsequent cell line established in culture. Organ preference in arrest and colonization may be explained by adhesive interactions of circulating tumor cells with noncirculating host cells in the target organ or by adhesive interactions leading to arrest in the mi-

circulation at specific locations. Tumor cell invasion appears to occur by a combination of mechanical and enzymatic mechanisms. In addition to unique tumor cell characteristics, host properties such as immune response can influence metastasis. Therefore, both tumor cell and host properties probably determine whether metastasis occurs in each tumor-host system. (48 refs)

- 78-6061 Cell Membrane Ionic Permeability, Calcium Ion, Mitochondria, and Carcinogenesis.** (Eng) Anghileri, L. J. (Department de Protection, Boite Postale No. 6, 92260 Fontenay-Aux-Roses, France). *Arch Geschwulstforsch* 48(6): 497-503; 1978.

Cumulative evidence in support of a hypothesis of carcinogenesis at the cellular level is reviewed. The hypothesis states that a carcinogen, chemical or viral, alters cell membrane permeability, which upsets the normal intracellular ionic composition of the cytoplasm and allows excess calcium ions to flow into the cell. Calcium is sequestered in the mitochondria, resulting in damage to its oxidative phosphorylation mechanism and also to its genetic component. Therefore, the cell will shift toward glycolytic means of energy production. Moreover, if released from damaged mitochondria into the cytoplasm, mitochondrial DNA may become incorporated into the nuclear genome and may induce malignant transformation. The role of mitochondrial DNA in the specification of membrane characteristics is also important. Damage to this function may lead to the loss of growth inhibition and communications with neighboring cells and effectors that is noted in transformed cells. (55 refs)

- 78-6062 Pathogenesis of Attic Cholesteatomas.** (Eng) Beales, P. (Royal Infirmary, Doncaster, England). *J R Soc Med* 71(10): 707-708; 1978.

The congenital, migration, and metaplasia theories of the pathogenesis of attic cholesteatomas (AC) are reviewed. Although the migration theory is the one most commonly accepted, the evidence of contact inhibition of stratified squamous epithelium and the absence of documentation of the evolution of AC from a preexisting perforation or from marginal perforations argue against this theory. The exact mechanism of AC formation is not known. (11 refs)

- 78-6063 Small Cell Carcinoma of the Lung: Cellular Origin and Relationship to Other Neoplasms.** (Eng) Tischler, A. S. (Dept. Pathology, Tufts Univ. Sch. Medicine, 126 Harrison Ave., Boston, MA, 02111). *Semin Oncol* 5(3): 244-252; 1978.

The origin and behavior of small cell carcinoma of the lung are considered in relation to its similarity to APUD (amine

precursor uptake and decarboxylation) cells. Cells from small cell carcinomas of the lung and from the more benign pulmonary carcinoid tumors contain secretory granules that are morphologically and, to some extent, cytochemically similar to those in normal pulmonary APUD cells. Both tumors produce amine and peptide hormones. Carcinoids have been observed in association with increased numbers of APUD cells that might represent hyperplasia or neoplasia in situ; such observations have not been documented for small cell carcinomas. Although small cell carcinomas have many similarities to carcinoid tumors and to normal pulmonary APUD cells, it cannot be stated that they arise from a specific cell type. Because the embryologic lineages of normal pulmonary APUD cells have not yet been determined with certainty, small cell carcinomas would have an unclear origin even if they had been observed histologically to arise from such cells. Some tumors that currently are classified as small cell carcinomas probably do not have APUD cell characteristics, but others appear to have both APUD and non-APUD features. A subclassification of small cell carcinomas based on a combination of physiological and morphological features might prove of prognostic and therapeutic value. (48 refs)

- 78-6064 Nasopharyngeal Carcinoma. Aetiological Aspects and Electronmicroscopic Observations.** (Eng) Prasad, U. (Dept. Otolaryngology, Univ. Malaya, Kuala Lumpur, Malaya). *J R Coll Surg Edinb* 23(2): 199-207; 1978.

Current knowledge of the etiology of nasopharyngeal carcinoma (NPC) is reviewed, and the results of an electron microscope study of 60 cases are presented. The study supports the view that NPC originates from squamous epithelial cells, and it indicates that, in addition to genetic factors, environmental factors play a significant role in the etiology of NPC. (48 refs)

- 78-6065 Metastatic Potential of Malignant Tumors.** (Eng) Carter, R. L. (Inst. Cancer Res., Sutton, England). *Invest Cell Pathol* 1(3): 275-286; 1978.

The metastatic potential (MP) of malignant tumors and contributing factors are reviewed based on clinical and experimental observations. Morphological factors include tumor site, size, and histology and the pattern of local tumor spread. This information can be supplemented by techniques such as immunoperoxidase staining to determine MP. However, more functional, nonmorphological features of the primary tumor are likely to be of greater importance. These features are the functional heterogeneity of tumor cells with respect to MP; the elaboration of certain tumor-associated products that may potentiate tumor spread; and immunological factors in the form of local host cell infiltrates in and around primary tumors. Tumor-associated products include substances that

originate largely or wholly from tumor cells and host cells. Tumor-specific antigens, ectopically produced hormones, and tumor angiogenesis factor may thus be contrasted with prostaglandins and collagenases. Irrespective of their cellular origin, however, both types of products contribute to MP. Thus, some host factors may act as determinants of metastasis, a view compatible with evidence that shows that host cells sometimes make up a far larger proportion of tumor populations than would be expected from conventional morphology. It is concluded that the MP of tumors cannot be assessed in terms of conventional tumor pathology alone. (122 refs)

6066 Constitutional Factors and Endometrial Carcinoma. (Eng) Lynch, H. T. (Dept. Preventive Medicine, Creighton Univ., Omaha, NB); Lynch, P. M. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 05-2117; 1978.

Several etiological factors in endometrial carcinoma (EMC) are identified by a review of the medical records of 154 EMC patients treated at one hospital from 1946 to 1965. The most common finding was obesity (123/154), which was massive in some patients. Other constitutional factors included hypertension (100/154) and diabetes mellitus (66/154). The combination of obesity, hypertension, and diabetes was seen in 46 women. A review of the literature indicated that nulliparity and late natural menopause are also risk factors for EMC. In an autopsy series of 30 patients who died from EMC revealed significant aberrations in pituitary, ovarian, pancreatic, and renal histology. Initial studies have not revealed any synergistic interaction between constitutional factors and exogenous estrogen. Multiple genotypes have been postulated to explain the occurrence of EMC in several familial cancer syndromes. It is suggested that the inherited gene(s) defect in certain instances of familial EMC may initially express itself in the endocrine system, possibly involving the pituitary-adrenal axes. Speculation that the adrenals might be the site of the inherited endocrine lesion is supported by statistical indications that adrenal estrogen production may have a heritable basis. (32 refs)

78-6067 An Overview of Hormone-associated Cancers. (Eng) Miller, A. B. (NCIC-Epidemiology Unit, Faculty Medicine, Univ. Toronto, Toronto, Ontario M5S 1A8, Canada). *Cancer Res* 38(11, part 2): 3985-3990; 1978.

Data on the etiology of hormone-associated cancers, particularly breast cancer, are reviewed. Although many breast cancer risk factors point to the relevance of hormonal factors, the findings are not uniform. If age at first pregnancy is the

prime epidemiological indicator of some largely internally influenced hormonal mechanism and if the risk ratio of approx 1.7 from the literature is appropriate, then the attributable risk (AR) for this factor is approx 25%. Family history of breast disease, with a risk ratio of 2.4, results in an AR of 20%. Finally, total fat intake, by a weighted average over pre- and postmenopausal women, results in an AR of 27%. The AR's for age at first pregnancy, which may be overestimated, and total fat intake, which may be underestimated are the most doubtful. There also may be an interaction between the two, and neither may adequately reflect the true risk factor. However, evidence points to the importance of dietary factors, and one study appears to confirm an association with high total fat intake. Dietary factors, although possibly mediated by a hormonal mechanism, may be found to play a major role in the etiology of breast cancer. Studies of endometrial and ovarian cancer also show the importance of hormonally associated factors, although not always the same way as for breast cancer. For both these cancers and for cancer of the prostate, dietary factors may also be relevant in their etiology. (47 refs)

78-6068 Genetic Factors in the Etiology of Breast Cancer. (Fre) Sancho-Garnier, H. (Departement de Statistique medicale, Institut Gustave-Roussy, 16 bis, avenue P.-V.-Couturier, F 94800 Villejuif, France); Henry-Amar, M.; Le, M. *Bull Cancer (Paris)* 65(2): 127-130; 1978.

Literature studies on the relationship of genetic factors to the development of breast cancer are reviewed. A higher risk is present for women with mothers or sisters with premenopausal breast cancer. Only a few studies have evaluated the incidence of breast cancer in twins but a concordance of 0.28 has been found for monozygotic twins vs 0.12 for dizygotic twins. The incidence of breast cancer in various ethnic groups changes upon migration, which indicates that environmental factors are involved. Attempts have been made to identify specific markers such as blood group, human leukocyte antigens (HLA), or fingerprints that may be associated with high-risk populations. An increased incidence has been found for young patients with type O blood and older patients with type A. The HLA haplotypes 2-12 and 10 have been found in families with tendency to develop breast cancer. One unconfirmed study relates cancer incidence to fingerprint characteristics. Apparently, families with breast cancer are more likely to be carriers of mammary tumor virus. As for many types of cancer, the development of breast cancer is probably due to a multigenetic transmission of susceptibility and to the action of environmental factors. (27 refs)

78-6069 Tumors of Dysgenetic Gonads in Females with Y Chromosome. (Pol) Boczkowski, K. (Samodzielna Pracownia Genetyki, Instytut Ginekologii i Poloznict-

wa AM, pl. Starynkiewicz 1, 02-015 Warsaw, Poland). *Postepy Hig Med Dosw* 32(3): 381-390; 1978.

Studies of the frequency, genetic basis, and origin of tumors of dysgenetic gonads in women with Y chromosomes are reviewed. Dysgerminoma of the left ovary was diagnosed in a 21-yr-old eunuchoid woman with a 46,XY karyotype. (31 refs)

78-6070 Chromosomes and Cancer. (Eng) Brown, J. K. (Radiation Biology Section, Australian Atomic Energy Commission, Private Mail Bag, Sutherland, New South Wales 2232, Australia). *Cancer Forum* 14: 97-100; 1978.

The origin of cancer may be due to a genetically unbalanced chromosome complement that predisposes a cell to malignant transformation. On the other hand, chromosome anomalies may be of a secondary nature, merely accompanying the more significant changes. Many, but not all, cancer cells have chromosome abnormalities. Clonal evolution is common in neoplasms, particularly leukemias. At present, only a few specific chromosome abnormalities characteristic of particular neoplasms have been found. (no refs)

78-6071 Hereditary Retinoblastoma: Delayed Mutation or Host Resistance? (Eng) Matsunaga, E. (Dept. Human Genetics, Natl. Inst. Genetics, 411 Mishima, Shizuoka-Ken, Japan). *Am J Hum Genet* 30(4): 406-424; 1978.

A review of 261 pedigrees with 2 or more cases of retinoblastoma taken from 11 different sources indicates that host resistance explains the variation in the expression of the retinoblastoma locus and that there is little need to postulate delayed mutation. Both penetrance and expressivity in the gene carrier can be defined as a variable determined by genetic and environmental factors, not by Poisson distribution of tumors formed. Of individuals who receive a new mutation from a healthy parent, approx 13% do not manifest retinoblastoma; the heritability of the host resistance is estimated at about 90%. Significant associations of human WBC antigens with retinoblastoma have been reported, and the cell-mediated immune response is thought to play a role in host resistance. The host resistance model suggests that nonhereditary retinoblastoma, due to somatic mutations, occurs in the most susceptible group of the general population, whereas unaffected carriers, including spontaneously regressed cases, are the most resistant group. Therefore, a genetic marker correlated with host resistance to retinoblastoma would be found with decreasing frequency in unaffected carriers, hereditary unilateral cases, the general population, and bilateral and nonhereditary cases. The detection of such a marker would be maximized by comparing unaffected carriers with nonhereditary cases. (36 refs)

78-6072 The Relation of Gene Control Mechanisms to Cancer. (Eng) Busch, H. (Dept. Pharmacology, Baylor Coll. Medicine, Houston, TX, 77030). *Z Krebsforsch* 92(2): 123-135; 1978.

Recent experiments on the involvement of genetic mechanisms in the cancer cell phenotype (growth and cell division, invasiveness, and metastasis) are reviewed. Support for the idea that gene control of cancer cells is aberrant emerges from studies that suggest that aberrations in gene control can be rectifiable. Combined teratoma cells and normal embryonic cells produced normal heterozygous guinea pig embryos. Normal controls produced in normal sequence in the biological clock were apparently capable of converting teratoma cells to normal cells. The direct controls of gene function are considered to be the nonhistone nuclear proteins. Studies showed that the nonhistone nuclear proteins of experimental hepatomas and nontumor tissues differ. In other studies, different antigens were found in the nuclear proteins of tumors and nontumor tissues. One was nuclear antigen NAg-1, the first fetal chromosomal protein. NAg-1 or No-Ag-2, a nucleolar protein, may be special stimuli that result in the transcription and translation of messenger RNA species for the cancer cell phenotype. There is uniform agreement that cancer cells exhibit characteristics that are uncontrolled by systems available to the host. However, it is possible that substances responsible for normal maturation at specific times of embryogenesis would function in tumor cells to reorder or eliminate the specific readouts related to the cancer process. Fetal readouts that may be responsible for the cancer phenotype are likely to be timing errors of dislocated genomic control which proper timing may be restored by the substances normally involved in this reordering process. (24 refs)

78-6073 Some Aspects of Human Tumor Genetics. (Rus) Pogosiants, E. E. (Oncological Res. Center, Moscow, USSR). *Vopr Onkol* 24(9): 15-24; 1978.

The role of genetic factors in cancer etiology is reviewed. In most human tumors, genetic effects are of little importance but environmental factors are of paramount importance. However, there are some rare tumors that are dependent on specific inherited genes. In some cases, genetic markers are responsible for the development of the tumor proper; in others, they are responsible for the complex syndromes in which a malignant tumor may be only one of the obligatory features. In other cases, a syndrome that predisposes to the development of tumors and/or leukemia is inherited. The recent discovery of the genetic heterogeneity of tumors at the same site (including common tumors such as breast and renal cancer) is of particular theoretical and clinical interest. (22 refs)

78-6074 Mendelian Genetics and Cancer. (Fre) Auer, J. L. (Service de Medecine A, Institut G.

ve-Roussy, 16 bis, avenue Paul-Vaillant-Couturier, F 94800
illejuif, France). *Bull Cancer (Paris)* 65(2): 131-136; 1978.

Approx 1,142 simple genetic patterns known to be associat-
ed with 2,336 syndromes, about 200 are associated with high
cancer risk. However, other factors such as the environment
may interact with these genes to affect cancer risk. It is some-
times difficult to distinguish between these factors. In
anconi's anemia and xeroderma pigmentosum, there is
predisposition to cancer development because of
chromosome instability. The role of the histocompati-
bility system in evolution of certain cancers is discussed.
The most common hereditary syndromes associated with
higher cancer risk are tabulated. (22 refs)

78-6075 **Malignant Neoplasms of Genetic Origin in
Drosophila melanogaster.** (Eng) Gateff, E. (Bi-
ologisches Institut I-Zoologie, Albert-Ludwigs-Universitat,
8 Freiburg im Breisgau, W. Germany). *Science* 200(4349):
1445-1451; 1978.

Malignant neoplasms that develop in recessive-lethal larval
mutants of *Drosophila melanogaster* are described. There are
x nonallelic mutants that develop malignant neuroblas-
tomas in the adult optic centers. These mutants, on chromo-
somes 1, 2, and 3, represent developmental genes that prevent
differentiation of the adult optic neuroblasts into adult optic
neurons by causing an uncontrolled, autonomous, invasive,
lethal, and transplantable growth of the adult optic neuro-
blasts and ganglion mother cells. Some neoplastic cells, even
though they are incapable of differentiation, retain parts of
their behavioral repertoire, whereas other neoplastic cells
become unresponsive to hormonal signals. Eight recess-
ive-lethal mutants develop imaginal disk neoplasms: five
form imaginal discs that grow in a compact, noninvasive fash-
ion, and three form neoplasms that grow in an invasive pat-
tern. Morphological variations exist among imaginal
disks from different mutant stocks, and neoplastic disks
in situ as well as after transplantation possess qualities
of both malignant and benign neoplasms. The presence
of *y f* clones on the integument indicates nonautonomy
and a capacity of the homozygous tumor cell for reversal
to the differentiating state; the absence of *y f* clones
shows autonomy and an incapacity for reversal. A third
type of tumor mutation induces malignant neoplastic
transformation of the presumptive blood cells in the
larval blood-forming organs (lymph glands). The mutant
genes are located on chromosomes 1, 2, and 3.
Plasmatocyte-type cells are the malignant component,
and they grow in an autonomous way; crystal cells do
not participate in the growth after transplantation. The
neoplastic state of the mutant blood cells in situ is not
autonomous, but may result from a normal factor else-
where in the animal. (54 refs)

78-6076 **Tumors Arising in Organ Transplant Reci-
pients.** (Eng) Penn, I. (Dept. Surgery, Univ.

Colorado Sch. Medicine, Denver, CO). *Adv Cancer Res* 28:
31-61; 1978.

The development of cancer in organ transplant patients is
reviewed. These patients can be classified into three groups:
(1) those with inadvertently transplanted neoplasms; (2)
those with cancer that arose de novo at some time after trans-
plantation; and (3) those with tumors present before graft-
ing. Data were collected on 65 patients who received organs
(1 heart, the rest kidneys) from donors who had cancer. Six
of the 13 organs from living donors contained neoplasms; one
of these tumors spread locally and one caused distant metas-
tases. Of the 52 cadaver organs, 20 contained cancer cells.
Five remained localized to the kidney, 1 spread locally, and
14 caused distant metastases that were histologically identical
to the original donor tumors. After transplantation, 5%-7%
of patients develop tumors arising de novo. The incidence is
100 times greater than that in the general population of the
same age range. Data are given for 548 recipients who devel-
oped 572 de novo tumors (24 patients had > 1 type of can-
cer). These tumors were not related to the 12 preexisting
recipient tumors treated from several weeks to 17 yr before
transplantation or to any of the 6 tumors found among the
650 donors. Of 164 patients who had tumors removed at the
time of or 5 yr prior to transplantation, 92 had no recurrence,
63 had a recurrence or metastases, 10 developed de novo
cancers, and 1 case remained unchanged. Cancer in organ
transplant recipients probably develops as a result of a com-
plex interplay of multiple factors, including alterations in im-
munity, immunosuppressive agents or other drugs, genetic
factors, and oncogenic viruses. (143 refs)

78-6077 **Tumor Immunity. An Overview.** (Eng) Siegel, B.
V. (Dept. Pathology, Univ. Oregon Health
Sciences Center, Portland, OR). *Am J Pathol* 93(2): 515-523;
1978.

From attempts to formulate specific modes of cancer therapy
based on the ability of the immune system to recognize and
destroy foreign antigens, it has become clear that tumorigene-
sis is affected by immunologic and nonimmunologic factors.
Tumors may evade immunosurveillance by a number of
mechanisms, including sneaking through, immunosuppres-
sion, antigenic modulation, serum blocking factor, antigen
shielding, genetic resistance, and immunostimulation. An
integral part of the immune response is the activation of mac-
rophages, possibly through interferon, to a specifically cyto-
toxic state, which may be associated with the prevention of
metastatic spread. Since much of the knowledge of im-
munologic mechanisms in tumor-cell killing has been gained
from in vitro experiments, the relevancy of such observations
to events in vivo remains unclear. However, tumor immunity
may be enhanced by (1) amplification of cytotoxic T cell
pathways; (2) modification of genetic restriction on immune
response to tumor antigens; and (3) use of immunopotential-
tators to stimulate macrophage cytotoxicity and phagocytic ac-
tivity. (37 refs)

- 78-6078 Acute Lymphocytic Leukemia Involving Pre-B Cells.** (Eng) Vogler, L. B. (Cellular Immunobiology Unit Tumor Inst., Dept. Pediatrics & Microbiology, Univ. Alabama, Birmingham, AL); Crist, W. M.; Cooper, M. D.; Lawton, A. R. In: *Human Lymphocyte Differentiation: Its Application to Cancer. Proceedings of the International Symposium on Human Lymphocyte Differentiation: Its Application to Cancer, held in Montpellier (France) by Institut National de la Sante et de la Recherche Medicale and Delegation Generale a la Recherche Scientifique et Technique, 2-4 March, 1978.* Serrou, B., Rosenfeld, C., eds. (Amsterdam: North Holland Publishing Company): 423 pp.; 407-413; 1978.

Some of the characteristics of precursor B cells (pre-B cells) are described, and their participation in the proportion of childhood acute lymphoblastic leukemias is discussed. Pre-B cells contain cytoplasmic IgM (cIgM+) but they appear to lack functional antigen receptors and detectable surface IgM (sIgM). Large pre-B cells in humans divide rapidly, giving rise to smaller pre-B cells that are the direct precursors of sIg+ B lymphocytes; they resemble the sIg- lymphocytes that are the precursors of B lymphocytes in mice. Generation of antibody diversity appears to begin at the pre-B cell stage of differentiation. Lymphoblasts from 4/22 children with acute lymphocytic leukemia had the pre-B phenotype. Cells from all four patients bore the D locus related p23,30 antigen; a proportion of cIgM+ lymphoblasts from patients 2 and 4 expressed barely detectable sIgM, and Fc receptors were present on virtually all cells from patients 1 and 2. In patient 2, cIgM+ blasts from a lymph node were devoid of sIgM, whereas 75% of cIgM+ blasts from bone marrow stained immunoglobulins were not found in any patient. Chemotherapy resulted in remissions associated with the disappearance of cIgM+sIg- lymphoblasts from the blood in all four patients. (27 refs)

- 78-6079 Macrophage Turnover and Traffic: A Review.** (Eng) Spector, W. G. (Dept. Pathology, St. Bartholomew's Hosp. Medical Coll., London, England). In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 15-18; 1977.

Current knowledge of the turnover rate of macrophages (MP) in different body states is reviewed. In steady-state conditions, the MP pool in sites such as the liver and peritoneum is maintained by a slow turnover of fresh MP from circulation and also by a process of mitotic division in situ. In inflammatory states, monocytes migrate to the site at a constant rate through interendothelial gaps in venules, and, once inside a lesion, they are converted to MP with enhanced phagocytic activity. Both processes probably depend on serum and humoral factors. Factors that accelerate monocyte release from marrow or initiate peritoneal MP mitosis in cell cultures have been discovered. Although there is a close correlation be-

tween hypersensitivity and chronic inflammation, it is not known whether MP mobilization is linked to T-lymphocyte activity. Complement, however, may be involved. MP occurs primarily by three mechanisms: endocytosis of foreign material, mitosis with lethal chromosome aberrations, or formation of multinucleate giant cells. Both resident steady-state populations and inflammatory MP are probably involved in carcinogenesis. (13 refs)

- 78-6080 Mononuclear Phagocytes and Antitumor Resistance: A Discussion.** (Eng) Keller, R. (Immunobiology Res. Group, Univ. Zurich, Schonleinstrasse 7, CH-8032 Zurich, Switzerland). In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 31-49; 1977.

The contribution of macrophages (MP) and MP-like cells to host tumor resistance is reviewed. The numerous effects of MP on tumor growth in vivo and in vitro are discussed. Mechanisms by which MP alter the growth of other cells are reviewed. MP activation by specific or nonspecific means. This activation, expressed as cytotoxic function, usually requires a latent period of 1-2 hr. An activated MP is capable of killing a variety of target cells (in vitro). This effect transcends intra- and interspecific compatibility barriers, and it is independent of the target cell type or growth characteristics. However, MP-mediated cytotoxicity shows some selectivity for transformed cells. The mechanism of MP-mediated cytotoxicity is not understood, but the process is resistant to x-rays and several metabolic inhibitors, requires energy, and possibly involves complement activation and proteases. In vivo studies have shown that passive transfer of activated MP can inhibit tumor growth, and research is underway regarding the in vivo modification of MP functions in the tumor-bearing host. Not only MP but also nonphagocytic peritoneal cells, phagocytic lymphoid cells, cytotoxic lymphoreticular cells, and natural killer cells have been implicated in antitumor resistance. (106 refs)

- 78-6081 The Influence of Tumour Cell Products on Macrophage Function In Vitro and In Vivo.** (Eng) James, K. (Dept. Surgery, Univ. Edinburgh Medical School, Teviot Place, Edinburgh EH8 9AG, Scotland). In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 225-246; 1977.

The effects of viable tumors, tumor culture supernatants, and tumor culture extracts on macrophage development and function are reviewed. Animal studies in vivo and in vitro have shown that accumulation of macrophages and their chemotactic responses can be impaired by tumors or their soluble products. Some findings indicate that tumors may

pair immunosurveillance against them or other tumors. Conversely, macrophage number, activity, or chemotaxis may be enhanced by tumors. In vitro studies have usually included that chemotactic response of human blood monocytes is depressed by tumors. However, the factors influencing macrophage function could also be of host cell or viral origin. The possible modes of tumor-macrophage interaction are manifold: tumor products could interfere with the production of factors that modulate macrophage function, prevent the necessary interaction of such factors, or tumor products themselves could have direct effects on macrophages. Normal macrophages may be sequestered at the tumor site and, therefore, rendered inactive, or tumor products may inhibit macrophage activity within a tumor. In any event, macrophage development and motility can be adversely affected, placing the host at a distinct disadvantage. (61 refs)

6082 Anti-tumour Effects of Macrophages Injected into Animals: A Review. (Eng) Den Otter, W. (Dept. Pathology, State Univ., Pasteurstraat 2, Utrecht, Netherlands); Dullens, H. F.; Van Loveren, H.; Pels, E. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 119-141; 1977.

Concerning cytotoxic macrophages, the local antitumor effect of macrophages injected into animals along with tumor cells, and the antitumor effect of macrophages injected at a distance from that of the tumor cells are reviewed. (67 refs)

6083 Cancer of the Large Bowel with Special Reference to Incidence and Etiology. (Eng) Baughn, B. B. (401 W. Main St., Frankfort, KY, 40601). *Am J Surg* 44(9): 548-553; 1978.

In addition to lung cancer, large bowel cancer (LBC) is the second most common cancer in the US, and it is much more common in people in the higher socioeconomic strata. This finding seems to be related to the fact that persons with diets high in fats and proteins also show an increased incidence of LBC. Three premalignant conditions are associated with LBC: adenomatous polyps, ulcerative colitis, and congenital multiple polyposis. Operative levels of carcinoembryonic antigen have been shown to correlate with the extent of the disease and prognosis for recurrence. (34 refs)

6084 Etiology and Pathogenesis of Tumors. (Rus) Diad'kova, A. M. (N. N. Petrov Oncological

Res. Inst., USSR Ministry Public Health, Leningrad, USSR). *Vopr Onkol* 24(9): 3-7; 1978.

Recent studies on the etiology and pathogenesis of tumors are reviewed with special regard to viral carcinogenesis. Penetration of the carcinogenic agent (virus, chemical substance, or radiation) into the body, primary reactions of the body to the agent, cellular changes, and the appearance or existence of conditions necessary for or facilitating tumor development are the basic steps of carcinogenesis. These steps can be interrelated and simultaneous, but none of them alone determines the fate of the cellular changes. Herpes virus type 2 and Epstein-Barr virus are suspected etiological agents in cervical carcinoma and Burkitt's lymphoma, respectively, in humans. (no refs)

78-6085 Pathogenesis of Malignant Neoplasms Induced in Hamsters by the Oncogenic DNA Simian Virus 40 (SV40). (Eng) Diamandopoulos, G. T. (Dept. Pathology, Harvard Medical Sch., Boston, MA, 02115). *Fed Proc* 37(7): 2082-2084; 1978.

Factors responsible for the morphogenesis of neoplasms induced in hamsters by simian virus 40 (SV40) were determined by inoculating random-bred Syrian golden hamsters with SV40 iv [$10^{6.5}$ median tissue culture infective dose ($TCID_{50}$)]. Six to 12 mo later, approx 90% of the hamsters developed lymphocytic leukemia, lymphoblastic lymphosarcoma, reticulum cell sarcoma, osteogenic sarcoma, and/or anaplastic sarcoma. Controls were tumor-free. Tumor antigen, tumor-specific transplantation antigen, and antibody to tumor antigen were found in the tumor-bearers. Similar results were obtained with three inbred hamster lines. Reticulum cell sarcoma cells lost their original morphology soon after dissociation into suspensions, but preserved their phagocytic activity throughout many in vitro passages. Osteogenic sarcoma cells lost both their differentiated phenotype and their capacity to form osteoid during serial passage. Both the lymphocytic and the lymphoblastic neoplastic cells preserved their lymphoid morphology and their ability to produce immunoglobulin after an indefinite number of in vitro passages. The effect of hamster age and virus dose on SV40-induced tumor development was studied. Random-bred 3-wk- to 12-mo-old hamsters received $10^{8.5}$ $TCID_{50}$ of SV40 iv, and groups of 3-wk-old hamsters were inoculated iv with SV40 aliquots of progressively decreasing titers. Tumor incidence and, to a lesser extent, tumor latency were directly dependent on the age of the animals at the time of virus exposure and on virus dose. The results suggested that cell replication at the time of oncogenic exposure enhances its susceptibility to transformation. Three strains of 3-wk-old inbred hamsters received $10^{7.5}$ $TCID_{50}$ iv, and tumor incidence and latency were approx the same in all three; however, random-bred hamsters were almost completely resistant under identical conditions. These results could be due to genetic differences, control at the cell-response level, or differences in immunological capacity. (15 refs)

- 78-6086 Virus Oncogenes and Carcinogenesis.** (Rus) Zhdanov, V. M. (D. I. Ivanovskii Inst. Virology, Acad. Medical Sciences USSR, Moscow, USSR); Al'tshtein, A. D. *Vopr Onkol* 24(9): 8-15; 1978.

Recent studies of viral oncogenes are reviewed with special regard to those of polyoma viruses, adenoviruses, and oncornaviruses. Each nondefective oncornavirus has three genes that are necessary for its reproduction: (1) *env*, coding for the precursor of the principal glycoprotein and low-mol-wt membrane proteins; (2) *pol*, coding for the synthesis of reverse transcriptase; and (3) *gag*, coding for the precursor of the internal structural proteins of the virion. Defective viruses are replicated by complementation with a helper virus. The simian virus 40 genome has only three genes: *A*, which codes for the nonstructural proteins synthesized before replication of virus DNA; *BC*, which codes for the synthesis of the main structural protein VP-1; and *D*, which codes for the synthesis of structural proteins VP-2 and VP-3. Most adenoviruses, including those that are not oncogenic in vivo, are able to transform cells in vitro. Virus-specific T and S antigens are always present in the adenovirus-transformed cells. These antigens appear to be coded by the adenovirus oncogene, and they are responsible for the cell transformation. The characteristics of the viral oncogenes suggest that they are cellular in origin. As a result of these studies, it is suggested that nonviral oncogenesis is due to a disturbance in the functioning of certain cellular genes, as a result of which they are transformed into endogenous oncogenes. (53 refs)

- 78-6087 The Gene Order of Avian RNA Tumor Viruses Derived from Biochemical Analyses of Deletion Mutants and Viral Recombinants.** (Eng) Wang, L. H. (Rockefeller Univ., New York, NY, 10021). *Annu Rev Microbiol* 32: 561-592; 1978.

The various experimental approaches that led to determination of the gene order of avian RNA tumor viruses are reviewed. Through biochemical analyses of deletion mutants and viral recombinants, the gene order 5'-gag-pol-env-src-c-3' can be drawn for sarcoma viruses and 5'-gag-pol-env-c-3' for leukosis and transformation-defective viruses. (180 refs)

- 78-6088 Bovine Leukemia Virus Involvement in Enzootic Bovine Leukosis.** (Eng) Burny, A. (Dept. Molecular Biology, Univ. Brussels, 1640-Rhode St-Genese, Brussels, Belgium); Bex, F.; Chantrenne, H.; Cleuter, Y.; Dekegel, D.; Ghysdael, J.; Kettmann, R.; Leclercq, M.; Leunen, J.; Mammerickx, M.; Portetelle, D. *Adv Cancer Res* 28: 251-311; 1978.

The major achievements of research over the last 20 yr on

the role of bovine leukemia virus (BLV) in enzootic bovine leukosis are reviewed. According to its high-mol-wt RNA-reverse transcriptase complex, structural proteins, their mode of maturation, and general morphology, BLV is considered a retrovirus belonging to the subgroup C mammalian oncornaviruses. However, this virus is unique in its way of budding, the strict Mg^{2+} requirement of its reverse transcriptase, and its property of inducing syncytia in monolayer cultures of nontransformed cells. In addition, neither molecular hybridization nor antigenic cross-reactivity occurs with any of the known RNA oncogenic viruses. BLV is an enzootic infectious agent. Persistent lymphocytosis (PL) is a clinical form of enzootic bovine leukosis—a percentage of lymphocytes are tumor cells containing BLV proviral DNA sequences. Infected animals actively synthesize antibodies to BLV, and there are several serologic methods of detecting BLV antibodies. BLV produced in culture can infect calves, sheep, goats, and chimpanzees, and it has induced tumors in PL in calves, sheep, and goats. Only a fraction of infected animals develop tumors. Virus dose, age at infection, and genetic make-up of the recipient appear important in determining the final outcome of the interaction between host and virus. (334 refs)

- 78-6089 Epstein-Barr Virus: The Ravager.** (Eng) Gross, R. (J. C. (Div. Otolaryngology, Albany Medical Center, Albany, NY, 12208); Parulekar, S. S.; Paonesu, D. F.; Rapoport, A.; Kouyoumgian, J. *Ann Otol Rhinol Laryngol* 87(5, part 1): 729-735; 1978.

The role of Epstein-Barr virus in infectious mononucleosis, Burkitt's lymphoma, and nasopharyngeal carcinoma is reviewed. (26 refs)

- 78-6090 Malignant Tumors Induced by Human Papilloma Virus.** (Pol) Jablonska, S. (Klinika Dermatologiczna AM, ul. Koszykowa 82a, 02-008 Warsaw, Poland); Orth, G.; Jarzabek-Chorzelska, M.; Rzeska, G.; Obal, S.; Glinski, W.; Favre, M.; Croissant, O. *Przegl Dermatol* 65(5): 497-507; 1978.

Studies of human papilloma viruses (HPV) are reviewed. HPV 3 induces both flat warts and a form of epidermodysplasia verruciformis (EV) characterized by lesions similar to flat warts. HPV 3 has slight, if any, oncogenic potential, but HPV 4, which also induces EV, is oncogenic. EV due to HPV 4 is characterized by very flat, reddish wartlike lesions, red plaques flush with the skin, hyperpigmentation reminiscent of pityriasis versicolor, and depigmentation. HPV 1, which induces plantar warts, and HPV 2, which induces common warts, do not seem to be oncogenic. Depressed cell-mediated immunity is of great significance in the dissemination and

istence of warts, especially flat warts, and EV. However, malignant transformation is due mainly to HPV 4. The possibility that specific vaccines can prevent transformation in patients with HPV-4 induced EV is discussed. (60 refs)

78-6091 The Human Papillomaviruses. (Eng) Orth, G. (Unite de Recherches sur l'etiologie virale des cancers humains, Laboratoire no. 147 associe au CNRS, 140 de l'INSERM, Institut Gustave-Roussy, 16 bis, rue P. Vaillant-Couturier, F 94800 Villejuif, France); Jansz, S.; Breitburd, F.; Favre, M.; Croissant, O. *Bull Cancer (Paris)* 65(2): 151-164; 1978.

Data on the properties and pathogenicity of human papillomaviruses (HPV's), the lesions induced by HPV's, in vivo virus-cell interactions, and factors involved in the production of carcinomas by papillomaviruses are reviewed. Recent biochemical and serological studies have shown the existence of at least four distinct types of HPV's that cause benign skin lesions. These viruses show hardly any antigenic relationships; their DNA's differ by their sensitivity to restriction endonucleases and show little, if any, sequence homology in molecular hybridization experiments using complementary DNA's transcribed in vitro. Data on the pathogenicity of HPV's are incomplete, but they indicate that some types of HPV's are preferentially associated with certain types of HPV. Epidermodysplasia verruciformis (EV) is associated with two types of HPV, HPV-3 and HPV-4. The lesions produced by these two viruses differ. Of a series of 13 EV patients, malignant transformation occurred in only 1/8 with HPV-3, which induces flat, wart-type lesions and hyperpigmented plaques. Intraepidermal or invasive carcinomas were observed in 5 patients infected with HPV-4, which produces reddish and pityriasis versicolor-like plaques. This suggests that at least one HPV may have oncogenic potential, as do rabbit (Shope) papillomavirus and bovine alimentary tract papillomavirus. (82 refs)

78-6092 Polyoma Virus, as a Model for Viral Skin Carcinogenesis. (Eng) Meyer, G. (Unite 119 de l'INSERM, 27 boulevard Lei-Roure, F 13009 Marseille, France). *Bull Cancer (Paris)* 65(2): 145-150; 1978.

Polyoma (Py) virus, the simplest model of transformation by DNA virus, is characterized along with Py virus-sensitive cells, virus/cell interactions, perpetuation of viral information, and the mechanism of Py virus-induced malignant transformation. Py virus is a small (45-m μ) DNA virus with icosahedral symmetry and without an envelope. Its capsid is composed of 72 capsomers forming an icosahedral structure. The virus contains a nucleic acid having a mol wt of 3.6×10^6

daltons, and the DNA is supercoiled. Three types of cells are sensitive to Py virus: cells that allow a complete virus cycle to take place, cells that do not allow maturation of infectious virions, and cells with intermediate properties. Three possible virus/cell interactions are reviewed: lytic infection, abortive infection, and malignant transformation. Perpetuation of viral information begins after penetration of viral DNA into the nucleus, with the production of early proteins (tumor and surface antigens), the induction of cellular DNA synthesis followed by viral DNA synthesis, the production of late proteins (virus antigen), and cell lysis. Nuclear and surface antigenic modifications are correlated with the mechanism of transformation in vitro and in vivo. (9 refs)

78-6093 Structural Organization of Papovavirus and Adenovirus DNA. (Rus) Roninson, I. B. (D. I. Ivanovskii Inst. Virology, Acad. Medical Sciences USSR, Moscow, USSR); Naroditskii, B. S. *Vopr Med Khim* 24(5): 579-591; 1978.

Studies of the structural organization and specific fragmentation of DNA in papova- and adenoviruses are reviewed with special regard to simian virus 40 and adenovirus 2. (142 refs.)

78-6094 Structure and Morphogenesis of Type-C Retroviruses. (Eng) Montelaro, R. C. (Dept. Surgery, Duke Univ. Medical Center, Durham, NC, 27710); Bolognesi, D. P. *Adv Cancer Res* 28: 63-89; 1978.

Models for the structure and morphogenesis of avian and murine C-type retroviruses were derived from an extensive review of literature concerning retrovirus morphogenesis. The models are based on the molecular arrangement of virion structural components in the particle as well as on available data relating to the biosynthesis of these products. (128 refs)

78-6095 Herpes Simplex Virus Type 2 and Cervical Cancer. (Dut) te Velde, E. R. (Afdelingen Gynaecologie, Academisch Ziekenhuis, Utrecht, Netherlands); The, T. H. *Ned Tijdschr Geneesk* 122(33): 1226-1230; 1978.

Epidemiological and virological studies show a correlation between herpes simplex virus 2 (HSV-2) infection and carcinoma of the uterine cervix, which suggests that HSV-2 may be involved in the etiology of this carcinoma. (12 refs)

- 78-6096 Chemical-Viral Interactions in Carcinogenesis.** (Eng) Weinstein, I. B. (Inst. Cancer Research, Columbia Univ. Coll. Physicians and Surgeons, New York, NY). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1609-1620; 1978.

Studies on the possible role of RNA viruses in the transformation of epithelial cells by chemical carcinogens are reviewed. In 1972, it was discovered that cell cultures established from rat hepatomas induced by aromatic amine carcinogens synthesize RNA viruslike particles. Electron micrographs revealed that these included intracisternal A particles, enveloped A particles, and, more rarely, mature C particles. Almost all rodent cell cultures established from tumors induced by aromatic amine carcinogens, polycyclic aromatic hydrocarbons, or asbestos synthesized RNA viruslike particles, as did rat epithelial cell cultures transformed in vitro. All attempts to detect RNA viruses in human cancers were negative. Further studies indicated that viruslike particles associated with chemically induced rodent tumors lacked biologic activity. This result is consistent with studies indicating that although RNA viruslike particles or their biochemical markers can be demonstrated frequently in spontaneous and radiation- or chemically-induced rodent tumors, these particles only rarely induce neoplasia when introduced into normal animals. There is no direct evidence that virus activation plays a causal role in the induction of carcinoma in animals or humans by radiation or chemical carcinogens. The viruslike particles and virus-related sequences in spontaneous and chemical carcinogen-induced neoplasms may play the role of a cofactor in carcinogenesis. (30 refs)

- 78-6097 Recombinant Origins of Leukemogenic Murine Viruses (Parke-Davis Award Lecture, 1978).** (Eng) Lerner, R. A. (Dept. Cellular and Developmental Immunology, Res. Inst. Scripps Clinic, 10666 N. Torrey Pines Rd., La Jolla, CA, 92037). *Am J Pathol* 93(1): 9-26; 1978.

Four points concerning retroviral genes and their hosts are derived using data on murine systems, although much of the information can be generalized to other species, including primates. (1) Retroviral genes, as proviruses, comprise an extensive multigene family. (2) As with host genes, proviral expression is regulated during development and differentiation. (3) Members of the retroviral gene family are constructed out of functionally distinct modules. (4) Viruses with new or altered pathogenicity may be constructed from nonpathogenic members of the multigene family. The last point is the

most important. Within the infinite reassortment and recombination processes that occur by recombination, biologically distinct viruses with unique pathogenicity can be constructed. The generation of virion diversity is said to be by a somatic recombination of proviruses. What makes this concept more interesting is that the provirus encodes no differentiation molecules of the mouse. Thus, recombination not only generates virus but also alters differentiation molecules within the adult lifetime of the mouse. Neoplastic version is envisioned to take place within this framework; a somatic mechanism is responsible for changing the cell face in mice. (20 refs)

- 78-6098 Recent Studies on the Role of the Immune Response in Resistance to Virus-induced Leukemias and Lymphomas.** (Eng) Meruelo, D. (Dept. Pathology, New York Univ. Medical Center, 550 First Ave., New York, NY, 10016); McDevitt, H. O. *Semin Hematol* 15: 399-419; 1978.

The contributions of immunogenetics to the understanding of the influence of immune surveillance on neoplasia, particularly leukemias and lymphomas, are reviewed. The murine histocompatibility complex, *H-2*, plays a large role in the susceptibility and resistance of mice to virus-induced leukemia. The *H-2k/k* haplotype results in >90% incidence of leukemia in one crossed strain. *H-2*-linked immune response genes (involved in the immune response to virus, tumor-specific transplantation antigens) seem to be directly involved in conferring resistance to the development of malignancy. The *H-2* complex plays a major role in determining the specificity of T cells sensitized against a variety of antigens, including viruses, tumor-associated antigens, antigens induced by chemical modification of the cell surface, autoantigens, and non-*H-2* histocompatibility antigens. Two models have been proposed to explain the requirement for *H-2* compatibility in these cytotoxic T-cell responses. A product of the *H-2K* or *H-2D* region may interact with a virus or hapten on a cell membrane, resulting in a new antigen recognized by effector T cells, or the interaction of T cells with target cells, may involve a dual recognition system, with one needing an *H-2* gene product. A second immune response gene locus, separate from *Rgv-1*, may be responsible for conferring resistance specifically to Friend and radiation leukemia virus-induced neoplasia. Following intrathymic radiation inoculation, there is an increase in the cell-surface expression of *H-2K* and *H-2D* molecules on thymocytes of both susceptible and resistant mice, but *H-2D* molecules are increased a longer period of time in resistant mice. Immunogenetic studies have begun to make an impact on human disease, evidenced by the finding that children with acute lymphoblastic leukemia (ALL) expressing the *HL-A9* antigen had increased survival over other ALL patients. (127 refs)

CHEMICAL CARCINOGENESIS

6099 **The Need for a Mammalian Test System for Chemical Mutagens and Carcinogens** (Meeting abstract). (Eng) Stich, H. F. (Environmental Carcinogenesis Unit, British Columbia Cancer Res. Centre, Vancouver, B. C., Canada); Wei, L.; Whiting, R. *Can J Genet Cytol* 20(3): 1-7; 1978. (no refs)

6100 **International Programme for the Evaluation of Short-Term Tests for Carcinogenicity.** (Eng) Ashby, J. (Central Toxicology Lab., Imperial Chemical Industries Ltd., Alderley Park, Nr. Macclesfield, Cheshire SK10 4TJ, England); de Serres, F. J. *Mutat Res* 54(2): 13-206; 1978.

An international collaborative study was initiated to evaluate 5 mutagenicity assay systems (prokaryotic, eukaryotic, andongenetic) for their ability to predict chemical carcinogenicity, using 42 reference carcinogens and noncarcinogens. The response of each assay will be compared with carcinogenicity findings for each chemical, and divergences between individual assay results and carcinogenicity data will be viewed within the context of the response of other assays. (4 refs)

6101 **Organic Trace Analysis Using Direct Probe Sample Introduction and High Resolution Mass Spectrometry.** (Eng) Haddon, W. F. (Western Regional Res. Center, US Dept. Agriculture, Albany, CA, 94710). In: *Symposium on High Performance Mass Spectrometry: Chemical Applications held in Lincoln, November 3-5, 1976*. University of Nebraska, National Science Foundation, AEI Scientific, and INCOS Corp. 358 pp.; 97-119; 1978.

High-resolution selected ion monitoring measurements on samples introduced on a direct probe can provide accurate and highly specific qualitative and quantitative information about compounds present in trace amounts, even compounds of low volatility or chemical instability. Highest sensitivities are obtained for substances with large mass defects, such as halogenated pesticides and mycotoxins. (40 refs)

6102 **Analytical and Biological Analyses of Test Materials from the Synthetic Fuel Technologies. II. Extended Genetic and Biochemical Studies with Mutagenic Fractions.** (Eng) Rao, T. K. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Young, J. A.; Har-

degree, A. A.; Winton, W.; Epler, J. L. *Mutat Res* 54(2): 185-191; 1978.

Certain highly mutagenic fractions of the crude synthetic fuel Synfuel A-3 were subjected to extensive genetic and biochemical analyses. Slight (not more than two-fold) mutagenic activity was noted with all fractions when the *Salmonella typhimurium* strain that detects base-pair substitution (TA1535) was used; metabolic activation was required. Strains that detect frameshift mutations (TA1537, TA1538, and TA98) were more sensitive to the fractionated Synfuel. TA1538 and TA98 exhibited a 20- to 75-fold increase over the spontaneous mutation rate when metabolic activation with various induced rat liver microsomal fractions (S-9) was included. Qualitative differences were seen with different induced S-9 preparations, and rat liver enzymes were the rate-limiting factor in the activation of the highly active basic fraction. The data indicate that the mutagenicity of Synfuel A-3 occurs by a frameshift mechanism and that the fuel contains chemical mutagens requiring metabolic transformation for conversion into an active form. (12 refs)

78-6103 **High-Pressure Liquid Chromatographic Analysis of Arylhydroxylamines after Derivatization with Methyl Isocyanate.** (Eng) Sternson, L. A. (Dept. Pharmaceutical Chemistry, McCollum Labs., Univ. Kansas, Lawrence, KS, 66044); DeWitte, W. J.; Stevens, J. G. *J Chromatogr* 153(2): 481-488; 1978.

A method for the analysis of micromolar levels of arylhydroxylamines in liver homogenates is described. The arylhydroxylamine is extracted with dichloromethane and then converted to the corresponding N-hydroxyurea by reaction with methyl isocyanate prior to high-pressure liquid chromatographic separation. This procedure renders the reaction product stable in soln, and it converts the hydroxylamine to a species that can be distinguished from other amine metabolites and metabonates. (18 refs)

78-6104 **Further Evaluation of a Hamster Embryo Cell Carcinogenesis Bioassay.** (Eng) Pienta, R. J. (NCI Frederick Cancer Res. Center, Frederick, MD, 21501); Pooley, J. A.; Leberherz, W. B. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group

for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1993-2011; 1978.

A standardized Syrian hamster embryo cell carcinogenesis bioassay was established, and results of its evaluation with 77 known carcinogens showed a high correlation (89%) with the reported carcinogenicities of these chemicals. No false-positive results were obtained with noncarcinogens, and spontaneous transformation was never observed in untreated cultures. However, until further evaluations are made, chemicals with negative responses cannot necessarily be considered noncarcinogenic. (11 refs)

- 78-6105 Mutagenicity of Ozonated, Recycled Water.** (Eng) Gruener, N. (Gulf South Res. Inst., New Orleans, LA, 70186). *Bull Environ Contam Toxicol* 20(4): 522-526; 1978.

The mutagenicity of ozonated and nonozonated recycled water was tested in the *Salmonella*/microsome system. Recycled water was concentrated by low-pressure, low-temperature (37-38 C) distillation, and the recovery of organics [measured as total organic carbon (TOC)] in the concentrates was 95%. The mutagenicity of this concentrated water, containing 20-250 µg TOC/plate, was tested in *Salmonella typhimurium* strains TA100 and TA98 with and without microsomes. Results were negative at these levels; higher levels of TOC could not be tested because of toxicity. Water ozonated for 15 min contained 25% less organics and was mutagenic to both TA100 and TA98. Microsomal activation was not required in either strain but with strain TA98, the presence of rat liver microsomes resulted in an additional increase in the number of mutants. The specific mutants formed during ozonation were not identified. (8 refs)

- 78-6106 Rat Liver Microsome-mediated N-Demethylation and Mutagenicity of Azoxymethane.** (Eng) Campbell, R. L. (Procter and Gamble Co., Miami Valley Labs., P.O. Box 39175, Cincinnati, OH, 45247); Supnick, J. D.; Hettrick, J. M.; Nigro, N. D. *Cancer Res* 38(12): 4585-4590; 1978.

The rat liver microsome-mediated metabolism of azoxymethane (AOM) to formaldehyde was studied in vitro. A mutagen reverting *Salmonella typhimurium* strains HisG46, TA1530, TA1535, and TA100 was detected in incubations of AOM (13.6 mM) with liver microsomes and NADPH. Mutagen production was dependent on the length of incubation and on microsomal protein and AOM concentration. Detection of the mutagen was most sensitive with strain TA100. The N-demethylation (N-DM) of AOM to formaldehyde by liver microsomes required NADPH and O₂, thus exhibiting mixed-function oxidase characteristics. Phenobarbital (80 mg/kg/day, 3 days) or 3-methylcholanthrene (40 mg/kg/day, 2 days) pretreatment of the rats did not significantly

alter the specific activity of microsomal AOM N-DM but it did increase microsomal cytochrome P-450 levels 164% and 40%, respectively. Cobaltous chloride (60 mg/sc) reduced AOM N-DM by 49% and cytochrome P-450 69%. SKF-525A in vitro inhibited AOM N-DM (43% in activity at 3 mM), but 7,8-benzoflavone did not. Disulfiram reduced microsomal AOM N-DM by 70% when given 0.25% (wt/wt) in the diet, but it did not alter cytochrome P-450 levels. The specific activity of microsomal AOM N-DM was 86% greater when animals were fed a rat chow diet than when they were fed a semisynthetic diet containing 56% dextrose, 25% casein, 7% salt mix, 5% corn oil, 5% bran, and 2% vitamin supplement. Compared with a semisynthetic diet control, AOM N-DM was increased 55% and 175%, respectively, in animals fed a semisynthetic diet or rat chow diet and then starved 18 hr. The addition of 30% beef fat, however, did not alter the level of AOM N-DM. Since beef fat enhances AOM intestinal carcinogenesis, the mechanism by which dietary fat enhances tumorigenesis may not be related to metabolism of the carcinogen but rather promotional factors. (28 refs)

- 78-6107 Inhibition of the Metabolism of the Colon Carcinogen, Azoxymethane, by Pyrazole.** (Eng) Fiala, E. S. (Naylor Dana Inst., American Health Foundation, Valhalla, NY, 10595); Kulakis, C.; Christiansen, C.; Weisburger, J. H. *Cancer Res* 38(12): 4515-4521; 1978.

To provide further evidence for the function of alcohol dehydrogenase in the metabolism of methylazoxymethane (MAM) and its precursor, azoxymethane (AOM), the effect of pyrazole (PZ), an inhibitor of this enzyme, on AOM metabolism were studied in vivo and in vitro. Male F344 rats treated with ¹⁴C-AOM (10.5 mg/kg, sc) alone (controls) exhaled approx 36% of the dose in 6 hr as ¹⁴CO₂, whereas rats pretreated with PZ (40 mg/kg, ip) exhaled approx 3.5%. Pretreatment of rats with 360 mg/kg PZ inhibited exhalation of ¹⁴CO₂ for at least 6 hr. After 48 hr, the amount of ¹⁴CO₂ exhaled comprised approx 50% of the dose in controls and 26% and 20% in the 40- and 360-mg/kg PZ-pretreated rats, respectively. Pretreatment with 40 mg/kg PZ increased urinary MAM levels 2.5-fold compared with controls; pretreatment with 360 mg/kg PZ increased the urinary AOM levels 2.5- to 3-fold. In contrast to controls and 40-mg/kg PZ-pretreated rats, 360-mg/kg PZ-pretreated animals showed increased urinary excretion of ¹⁴C-methylamine and continued ¹⁴C-MAM excretion during the 24- to 48-hr period after labeled AOM administration. Washed liver microsomes from PZ-treated rats converted AOM to MAM at the same rate as controls but the hydroxylation was inhibited by adding PZ to the incubation medium. These results show that PZ inhibits the metabolism of AOM by at least two steps. At 40 mg/kg, PZ inhibits MAM oxidation, presumably by blocking alcohol dehydrogenase. At 360 mg/kg, it blocks this reaction and inhibits the hydroxylation of AOM. (27 refs)

108 Failure of Ethyl Alcohol to Induce Dominant Lethal Mutations in Wistar Rats or to Synergize the Effect of X-rays in Swiss Mice (Meeting Abstract).

J. Chauhan, P. S. (Bio-Medical Group, Bhabha Atomic Centre, Trombay, Bombay 400085, India); Aravindakumar, M.; Sundaram, K. *Mutat Res* 53(2): 165-166; 1978. (no

6109 Deuterium Isotope Effect in Bioactivation and Hepatotoxicity of Chloroform. (Eng) Pohl,

R. (Lab. Chemical Pharmacology, Natl. Heart, Lung, Blood Inst., Bethesda, MD, 20014); Krishna, G. *Sci* 23(10): 1067-1072; 1978.

details of the formation of phosgene (COCl_2) from chloroform (CHCl_3), including the electron transfer pathway and rate-determining step, were monitored by the use of the deuterium isotope. When unlabeled CHCl_3 was incubated with liver microsomes from male Sprague-Dawley rats given intraperitoneal injections of phenobarbital (80 mg/kg), COCl_2 was readily produced. This reaction was NADPH-dependent and was inhibited by carbon monoxide or SKF 525-A, which suggests that cytochrome P-450 catalyzed formation of COCl_2 . When deuterium-labeled chloroform (CDCl_3) was substituted for CHCl_3 , COCl_2 production was significantly curtailed; therefore, the cleavage of the C-H bond was the rate-determining step of the reaction. The hepatotoxicity of CDCl_3 and CHCl_3 (2.49 and 4.98 nanomoles/kg ip) was determined by measuring SGPT levels and by histologic examination. CDCl_3 was less hepatotoxic than CHCl_3 , producing only minor histological changes (as opposed to highly elevated levels). These findings implicate the oxidative dechlorination mechanism in the bioactivation and subsequent carcinogenicity of chloroform. (29 refs)

6110 Carcinogenic Effects of Chronic Exposure to Very Low Levels of Toxic Substances. (Eng)

W. O. R. (Dept. Regius Professor Medicine, Radcliffe Infirmary, Oxford OX2 6HE, England). *Environ Health Perspect* 155-159; 1978.

The effects of chronic exposure to very low levels of carcinogens are examined from a statistical point of view. The results indicate that at low dose rates, the age distribution of extra tumors is independent of the dose rate and that the expected number of extra tumors is simply proportional to the dose rate. Because of statistical uncertainties in the differences between the numbers of tumors found in a control group and a treated group of animals, cancer risks $< 10\%$ cannot be accurately characterized. Because of this, much work has gone into the fitting of mathematical models. Although the results of the models in any particular case will depend on the details of the models they happen to fit, the qualitative conclusions reached would have been reached whatever class of

models had been fitted. Therefore, these conclusions are valid as a basis for legislation. (3 refs)

78-6111 Best-Estimate Low-Dose Extrapolation of Carcinogenicity Data. (Eng) Guess, H. A. (Environmental Biometry Branch, Natl. Inst. Environmental Health

Sciences, Research Triangle Park, NC, 27709); Crump, K. S. *Environ Health Perspect* 22: 149-152; 1978.

A low-dose extrapolation technique was developed for dichotomous data and used to analyze carcinogenicity data for vinyl chloride, dieldrin, 1,1,1-trichloro-2,2-bis-(p-chlorophenyl)ethane (DDT), dimethylnitrosamine, and ionizing radiation. The results suggest that when the very flat and the gradually sloping (linear or nearly linear) dose-response curves are considered together, it is extremely difficult on mathematical grounds to reject the hypothesis that the dose-response curve is nearly linear in the dose range corresponding to increased risks over a background of $\leq 10^{-6}$. These results have implications both for test design and for risk assessment. They suggest that if the hypothesis of a nearly linear dose-response curve in the low-dose range cannot be ruled out by assumption, then it seems questionable that it can be rejected by the data, even in cases in which it may be false. (no refs)

78-6112 Interactions of Vinyl Chloride with Rat-Liver DNA In Vivo. (Eng) Green, T. (Imperial

Chemical Industries Ltd., Central Toxicology Lab., Alderley Park, Cheshire SK10 4TJ, England); Hathway, D. E. *Chem Biol Interact* 22(2/3): 211-224; 1978.

The reaction of vinyl chloride (VC) with liver DNA in Wistar rats and the reaction of chloroacetaldehyde (CAA) and calf thymus DNA were investigated. 9β -D-2'-Deoxyribofuranosylimidazo(2,1-i)purine (ethenodeoxyadenosine) and 1β -D-2'-deoxyribofuranosyl-1,2-dihydro-2-oxoimidazo(1,2-c)pyrimidine (ethenodeoxycytidine) were identified in the enzyme hydrolysates obtained (1) from calf thymus DNA that had been modified by chemical reaction with CAA and (2) from liver DNA prepared from rats that had been exposed *po* to VC in their drinking water (250 ppm) for approx 2 yr. Thus, VC-derived chloroethylene oxide and/or CAA behaved as a bifunctional alkylating agent toward deoxyadenosine and deoxycytidine residues of DNA. It is concluded that modification of the DNA structure by VC, through imidazo-cyclization of deoxyadenosine and deoxycytidine residues and through depurination of the resulting ethenodeoxyadenosine residues, is related to the mutagenicity of VC. The carcinogenicity of VC in animals and in humans and its mutagenic properties after biotransformation (by microsomal enzymes) into the active metabolites chloroethylene oxide and CAA, which show electrophilic reactivity toward DNA, would appear to conform with the

idea of a relationship between carcinogenesis and mutagenesis. (41 refs)

- 78-6113 **The Mutagenicity and Metabolism of 1,2-Dichloroethane (Meeting Abstract).** (Eng) Ran-nug, U. (Environmental Toxicology Unit, Wallenberg Lab., Univ. Stockholm, Stockholm, Sweden); Ramel, C. *Mutat Res* 53(2): 251-252; 1978. (1 ref)

- 78-6114 **Report on Carcinogenesis Bioassay of 1,2-Dichloroethane (EDC).** (Eng) National Cancer Institute (U.S. Dept. Health, Education, and Welfare, NIH, NCI, Bethesda, MD, 20014). *Am Ind Hyg Assoc J* 39(11): A26-A30; 1978.

The halogenated solvent 1,2-dichloroethane (ethylene dichloride, or EDC) was administered to rats and mice for 78 weeks at time-weighted av dosages of 95 mg/kg/day (high dose) and 47 mg/kg/day (low dose) in a carcinogenesis bioassay. In male rats, EDC caused forestomach cancers, multiple hemangiosarcomas, and subcutaneous fibromas. In female rats, EDC induced mammary cancers, appearing in some high-dose animals as early as the 20th week. EDC also caused breast and uterine cancers in female mice and respiratory tract cancers in both male and female mice. (no refs)

- 78-6115 **Macromolecular Binding and Metabolism of the Carcinogen 1,2-Dibromoethane.** (Eng) Hill, D. L. (Kettering-Mayer Lab., Southern Res. Inst., 2000 Ninth Ave., S. Birmingham, AL, 35205); Shih, T. W.; Johnston, T. P.; Struck, R. F. *Cancer Res* 38(8): 2438-2442; 1978.

Biochemical mechanisms related to the activation and inactivation of the carcinogen 1,2-dibromoethane (1,2-DBE) were investigated. In rats inoculated ip with 1,2-¹⁴C-1,2-DBE [30.4 μ Ci; 4.2 micromoles (μ mol)]: 0.2 ml ethanol, measurable amounts of radioactivity became bound to protein, RNA, and DNA of all major tissues. For each macromolecules class, the largest amounts of bound radioactivity were in the liver and kidneys. Optimum conditions for the rat liver glutathione S-transferase that uses 1,2-DBE as a substrate were established. The pH optimum was 8.2, the Km for 1,2-DBE was 25 mM, and Vmax was 2.1 μ mol/min/g liver. Considerable enzyme activity was found in rat kidney; detectable activity was present in lung, testis, spleen, and heart. Enzymatic activity leading to the irreversible binding of radioactivity from ¹⁴C-1,2-DBE to proteins in the reaction system was present in rat liver microsomes. The activity was inducible by phenobarbital but not by benz[a]anthracene. NADPH was required for activity of both the noninduced and the induced reactions; MgCl₂ stimulated both reactions. Bromoacetaldehyde, a reactive compound probably involved in the irreversible binding, was identified as a metabolite formed in the reaction system.

Radioactivity from ¹⁴C-1,2-DBE also became bound to chromosomal proteins by a nonenzymatic chemical reaction. Enzymatic reaction leading to binding to macromolecules and/or the chemical binding reaction may be involved in biological activity of 1,2-DBE. (22 refs)

- 78-6116 **Mutagenicity of Industrial Compounds Evaluated by Means of Yeast Genetic Systems (Meeting Abstract).** (Eng) Fumero, S. (Istituto di Ricerche mediche 'A. Marxer', Ivrea, Italy); Mondino, A.; Barale, Loprieno, N. *Mutat Res* 53(2): 189; 1978. (no refs)

- 78-6117 **Preliminary Report on the Possible Use of Neuronal Tissue Culture in Carcinogenesis In Vitro.** (Eng) Boulos, B. M. (Sch. Public Health, Univ. Illinois Medical Center, Chicago, IL, 60680). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 2019-2028; 1978.

The in vitro effect of the food additive thiodipropionic acid (TDPA) on frog (*Rana pipiens*) spinal cord explants was investigated. The explants were taken from tadpoles at stages V and X, and incubated in a solution containing water and minimum essential medium (9:1). Six days later, addition of 10⁻³ μ M TDPA to the culture for 24 hr caused retraction of neuronal fibers. Addition of TDPA for 5 days caused growth and thickening of the fibers, a granular appearance, and a large number of fibroblasts. Lowering the TDPA concentration to 10⁻⁴ μ M caused increased thickening and lengthening of the fibers plus increased branching and granularity. At 10⁻¹ μ M, TDPA caused degeneration and neurite retraction. The neuronal cultures with TDPA required no addition of nutrients and showed growth up to 40-42 days. Control cultures were unable to grow without serum, and they grew for only 30-32 days with serum. The neuronal tissue culture system may prove useful in the in vitro identification of chemical carcinogens. (43 refs)

- 78-6118 **Mutagenic Assays with Bis(2-Chloroisopropyl)ether (Meeting Abstract).** (Eng) Simon, V. F. (Stanford Res. Inst., Menlo Park, CA); Hedderley, L.; Poole, D.; Tardiff, R. *Mutat Res* 53(2): 261-263; 1978. (no refs)

- 78-6119 **Trace Analysis: A Requirement for Toxicological Research with Carcinogens and Hazardous**

stances. (Eng) Bowman, M. C. (Food and Drug Admin., Natl. Center Toxicological Res., Jefferson, AR, 72079). *J Assoc Off Anal Chem* 61(5): 1253-1262; 1978.

scheme for analyzing an admixture of 13 carcinogens and related compounds in human urine, wastewater, and environmental surveillance samples (swabs, air filters, etc) is based on electron capture gas chromatography of their pentafluorophenyl derivatives. (6 refs)

-6120 Effects of Caffeine on the Frequencies of Chromosomal Aberrations and Sister-Chromatid Exchanges Induced by Mutagenic Agents in Root Tips of *Vicia faba* (Meeting Abstract). (Eng) Sturelid, S. (Dept. Genetics and Plant Breeding, Royal Agricultural Coll. Sweden, S-750 07 Uppsala 7, Sweden); Kihlman, B. A. *Mutat Res* 270: 270; 1978. (no refs)

-6121 Caffeine Inhibition of Liquid-holding Recovery from Ultraviolet-induced Budding Delay in Yeast Cells. (Eng) Dawson, P. C. (Dept. Physics, Univ. Victoria, Victoria, British Columbia, Canada); Johnson, R. G.; Rodin, L. J. *Radiat Res* 74(1): 191-198; 1978.

Examination of liquid-holding recovery from UV-induced budding delay in yeast cells (*Saccharomyces cerevisiae*) indicated that addition of 20 mg/ml caffeine to the liquid-holding medium completely inhibited the reversal of budding delay. When liquid-holding in caffeine was followed by photoreactivation, the budding delay was greatly reduced. These findings suggest that caffeine interferes with the excision of pyrimidine dimers. (20 refs)

-6122 Evaluation of Possible Mutagenicity of Ginger, Turmeric, Asafoetida, Clove and Cinnamon Administered Alone and in Combination with Caffeine or Theophylline in *Drosophila melanogaster* (Meeting Abstract). (Eng) Abraham, S. K. (Sch. Life Sciences, Jawaharlal Nehru Univ., New Delhi-110057, India); Kesavan, P. C. *Mutat Res* 53(2): 142; 1978. (no refs)

-6123 Testicular Atrophy and Impaired Spermatogenesis in Rats Fed High Levels of the Methylxanthines Caffeine, Theobromine, or Theophylline. (Eng) Reinberger, M. A. (Div. Pathology, Bureau Foods, Food and Drug Admin., Washington, DC, 20204); Friedman, L.; Farber, T. M.; Moreland, F. M.; Peters, E. L.; Gilmore, C. E.; Huan, M. A. *J Environ Pathol Toxicol* 1(5): 669-688; 1978.

Studies were conducted to determine whether simultaneous administration of sodium nitrite and the methylxanthines caffeine

(CF), theobromine (TB), or theophylline (TP) would produce lesions similar to those seen in animals treated with other amines and sodium nitrite. Osborne-Mendel male rats (4-6 wk old) were given the methylxanthines at a dietary level of 0.5% for 14-75 wk with and without sodium nitrite (also at 0.5%). The compounds did not produce neoplastic or preneoplastic lesions, but severe bilateral testicular atrophy with aspermatogenesis or oligospermatogenesis occurred in 85%-100% of the rats fed CF and TB. The methylxanthines were fed to male Holtzman rats (4 wk old) at a dietary level of 0.5% for 19 wk to determine whether they would induce testicular atrophy in a second rat strain. The testicular effects were even more pronounced in these rats: CF and TB induced testicular injury in nearly all animals; TP induced severe testicular atrophy in 14% and mild to moderate atrophy in 71%, and it had no effect in only 15% of the rats. There was a significantly reduced number of mitotic cells in testes from the CF-treated group vs those from the TP-treated group. Plasma testosterone concentrations were significantly elevated in the TB group and somewhat elevated in the CF group; plasma cholesterol levels were significantly increased in the CF and TB groups. The toxic effect of CF and TB appeared to be specific for the germinal epithelium of the seminiferous tubules; the Leydig cells in many severely atrophied testes appeared to be hyperplastic. (29 refs)

78-6124 Modification of DNA by Aflatoxin B₁ Creates Alkali-labile Lesions in DNA at Positions of Guanine and Adenine. (Eng) D'Andrea, A. D. (Sidney Farber Cancer Inst., Harvard Medical Sch., 44 Binney St., Boston, MA, 02115); Haseltine, W. A. *Proc Natl Acad Sci USA* 75(9): 4120-4124; 1978.

Terminally labeled (³²P) DNA fragments of known sequence were used to study the base specificity of modification of DNA by aflatoxin B₁ (AFB₁). The DNA used was a 117- or 168-nucleotide-long DNA fragment that contained the control region, the operator and promoter sequences, of the *lac*-*ose* operon. The 168-nucleotide DNA was incubated with 300 μM AFB₁ in the presence of activating enzymes supplied by a crude rat liver extract. This incubation modified the DNA in such a way that subsequent treatment with 1 M piperidine for 30 min or 0.1 M NaOH for 15 min at 90 C resulted in cleavage of the DNA at sites corresponding to the position of guanine residues. The lengths of the cleavage products were the same as those produced by dimethyl sulfate in an alkali-induced reaction. The 117-nucleotide fragment was incubated with 300 μM AFB₁ in the presence of a normal mouse liver extract, and again all possible guanine residues were accessible to modification by AFB₁, but the rate for each individual guanine varied. When the 168-nucleotide fragment was incubated with a high concentration of AFB₁ (1 mM) in the presence of purified rat liver microsomes, deproteinized, and incubated with 1 M piperidine at 90 C, extensive cleavage of DNA was noted, not only at guanine but also adenine residues within the sequence. The DNA products following AFB₁ treatment were identical in length to those

pretreated with Me_2SO_4 , a result that suggests that similar mechanisms are involved in the piperidine cleavage of both. (26 refs)

- 78-6125 Urinary Excretion of Aflatoxin M_1 after Administration of Aflatoxin B_1 in Sucrose- or Starch-rich Diets.** (Eng) Wise, A. (Regional Mycotoxin Center, Univ. Isfahan, Isfahan, Iran); Suzangar, M.; Messripour, M.; Mohammadi, J. *Br J Nutr* 40(2): 397-401; 1978.

The urinary excretion of aflatoxin M_1 (AFM₁) was compared in two groups of male Sprague-Dawley rats fed a sucrose- or starch-rich diet for 200 days. The diets contained (g/kg): casein (250), corn oil (50), mineral mix (60), vitamin mix (10), and starch or sucrose (630). In addition, 2 mg/kg aflatoxin B_1 was added to the diet. Animals were sacrificed at 75, 145, or 200 days, and livers were examined microscopically. In the last week of life, urine was collected from each rat for 24 hr and analyzed for AFM₁. Although the food intake of both groups was similar, the excretion of AFM₁ ($\mu\text{g/kg/day}$) was significantly less in sucrose-fed rats (mean value: 1.7 vs 2.3 in starch-fed rats). The livers of sucrose-fed rats were a larger proportion of body weight (49.5 vs 39.1 g/kg), providing more available tissue for metabolism. Therefore, the excretion per gram of liver showed an even larger deficit in sucrose-fed rats, the mean value being 270 vs 560 $\mu\text{g/kg/g}$ liver/day. There was no histological difference between the two groups in the liver slices analyzed. It is suggested that sucrose decreases the activity of the drug-metabolizing enzyme. (14 refs)

- 78-6126 Antacids, Bile-Salts, and Colon Cancer (Letter to Editor).** (Eng) Nomura, A. (Japan-Hawaii Cancer Study, Kuakini Medical Center, Honolulu, HI 96817); Stemmermann, G. N.; Guber, G. A. *Lancet* 2(8093): 785; 1978.

Of 8,006 men examined from 1965 to 1968 in the Japan-Hawaii Cancer Study, 345 had had a gastrectomy for gastric or duodenal ulcer. Many of these patients were probably former users of antacids based on aluminum hydroxide. During follow-up, 4 gastrectomy and 94 nongastrectomy men were diagnosed with large bowel cancer. The data do not support the view that aluminum hydroxide increases the risk for colorectal cancer. (1 ref)

- 78-6127 Lung Carcinogenesis Induced by Inhaled High-fired Oxides of Beryllium and Plutonium.** (Eng) Sanders, C. L. (Pacific Northwest Lab., Richland, WA, 99352); Cannon, W. C.; Powers, G. J. *Health Phys* 35(2): 193-199; 1978.

The cocarcinogenicity of inhaled, submicron, high-fired

$^{239}\text{PuO}_2$ and beryllium oxide particles was investigated in 600 Wistar rats. The BeO was calcined at 1000 C and $^{239}\text{PuO}_2$ at 750 C. The ultrafilterability of $^{239}\text{PuO}_2$ was <0 , and it did not increase with time in water suspension. Female rats were exposed via nose only to BeO and $^{239}\text{PuO}_2$ aerosols for a single period of 30-180 min. There were 12 exposure groups: two unexposed control groups (128 rats); three groups exposed to $^{239}\text{PuO}_2$ only (184); three groups exposed to BeO only (184); and four groups exposed to both oxides (119). The amount of oxide in the lung was determined in 3-5 rats in each group 24 hr after the single exposure, it ranged from 1.0 to 91 μg for BeO and from 4 to 100 μg for $^{239}\text{PuO}_2$. No unexposed rat developed a lung tumor. Only one of the 184 rats exposed to BeO alone developed a tumor. Lung tumor incidences were 10% at an alveolar deposition of 5 nCi $^{239}\text{PuO}_2$, 37% at 45 nCi, and 58% at 180 nCi. Tumor incidences after exposure to both oxides were 15% at 7.6 nCi $^{239}\text{PuO}_2$ and 14 μg BeO, 37% at 76 nCi $^{239}\text{PuO}_2$ and 14 μg BeO, 12% at 3.8 nCi $^{239}\text{PuO}_2$ and 91 μg BeO, and 58% at 59 nCi $^{239}\text{PuO}_2$ and 91 μg BeO. The alveolar clearance of ^{239}Pu decreased and the translocation of the radionuclide to thoracic lymph nodes increased in rats previously exposed to BeO. It is concluded that previous exposure to the BeO did not significantly change the incidence of lung tumors induced by $^{239}\text{PuO}_2$, despite the effect of BeO on alveolar clearance. (12 refs)

- 78-6128 Mutagenicity of Metal Compounds (Meeting Abstract).** (Eng) Kanematsu, K. (Second Department of Oral Surgery, Gifu Dental Coll., Hosumi, Gifu-ken, Japan); Kada, T. *Mutat Res* 53(2): 207-208; 1978. (no refs)

- 78-6129 The Mutagenicity of Hexavalent Chromium Decreased by Microsomal Metabolism.** (Eng) Lofroth, G. (Wallenberg Lab., Univ. Stockholm, S-106 21 Stockholm, Sweden). *Naturwissenschaften* 65(4): 207-208; 1978.

The mutagenicity of hexavalent chromium for *Salmonella typhimurium* strains TA1978 and Ta92, as measured with the Ames plate-incorporation assay, was decreased by the action of a microsomal activation system containing NADPH. No increase occurred in the absence of the dinucleotide, indicating that it was the main factor contributing to Cr(VI) metabolism in the microsomal system. Oxidation and reduction states did not change when Cr(VI) and NADPH were mixed in aqueous soln, which implied that at least one additional factor in the microsomal system was necessary for the reaction. (8 refs)

- 78-6130 Metabolic Deactivation of Hexavalent Chromium Mutagenicity.** (Eng) Petrilli, F. L. (Inst. I

ne, Univ. Genoa, Via Pastore 1, 16132 Genoa, Italy); De
ora, S. *Mutat Res* 54(2): 139-147; 1978.

the mutagenicities of various hexavalent chromium compounds (sodium dichromate, potassium chromate, chromic acid, basic zinc chromate, and basic lead chromate) for *Salmonella typhimurium* were tested in the presence of rat tissue preparations and human blood components. All hexavalent chromium compounds were weakly mutagenic for *S. typhimurium* his₅ strains and were more mutagenic for strain A100. Addition of the microsomal fraction from rat liver, human RBC lysates, or, to a lesser extent, the S-9 fraction from rat lung reduced the mutagenicity of all divalent chromium compounds. Human serum or plasma and the microsomal fraction from rat muscle had no effect on the mutagenicities of these compounds. In the presence of the microsomal fraction from rat liver or human RBC lysates, the addition of ascorbic acid, Na₂SO₃, reduced glutathione, ADH, or NADPH prevented hexavalent chromium mutagenicity, whereas KMnO₄ slightly increased it. The data could be relevant to the elective localization of chromium-induced human lung tumors and could account for the results of animal carcinogenicity tests, which generally show the development of tumors only at implant sites. (24 refs)

-6131 **A Study of Cancer in Chromate Production Workers.** (Eng) Hayes, R. B. (Johns Hopkins University, Baltimore, MD). *Diss Abstr Int B* 39(5): 37-2238; 1978. (no refs)

-6132 **Studies on the Mutagenic Properties of Bleaching Effluents.** (Eng) Ander, P. (Dept. Chemistry, Box 5604, Swedish Forest Products Res. Lab., S-114 86 Stockholm, Sweden); Eriksson, K. E.; Kolar, M. C.; Kringstad, K.; Rannug, U.; Ramel, C. *Svensk Papperstidning* (14): 454-458; 1977.

samples of industrial effluents from the bleaching of softwood kraft pulps were taken from six mills and tested for mutagenicity using a streptomycin-dependent strain (Sd-4) of *Escherichia coli* and different histidine-requiring mutant strains (TA1535, TA1537, TA1538, TA100) of *Salmonella typhimurium*. None of the effluents exhibited any significant mutagenic effects or toxicity toward *E. coli*. However, effluents from the chlorination (C) stages exhibited definite mutagenic effects on TA1535 and TA100; concentrated C-stage effluents also had a mutagenic effect on TA1537. A weak mutagenic effect was also produced by the hypochloride stage effluent, but the other effluents had little or no mutagenic activity. The addition of liver microsomes decreased the mutagenic activity of the C-stage effluent, indicating a metabolic detoxification of the mutagen(s). The mutagenic effects of the C-stage effluents were almost completely restricted to substances with low relative molecular mass. Part of the mutagenicity remained after ether extraction; the part that

was ether-soluble could be destroyed by alkali treatments. The C-stage effluents appear to contain compounds with alkylating and intercalating properties that lead to both base-substitution and frame-shift mutations. (6 refs)

78-6133 **Sarcoma at the Injection Site of Gold Sodium Thiomalate.** (Eng) Payne, B. J. (Dept. Pathology and Toxicology, Smith Kline & French Labs., Philadelphia, PA). *Vet Pathol* 15(Suppl 5): 39-50; 1978.

The local effects on rats following injections of gold sodium thiomalate were studied. Groups of 10 male and 10 female rats were injected in the rear legs with 3, 6, 9, 12, 24, 48, 96, or 192 mg/kg/day, with the right and left legs being injected alternately, every other day. The three highest doses were injected undiluted, the others were diluted with water to make a 2.5% soln. Doses of 48 mg/kg or more produced a considerable local reaction and rats became increasingly belligerent; therefore, rats in dose groups 192, 96, and 48 mg/kg were sacrificed on days 57, 112, and 212, respectively. No tumors were seen in these animals. Doses of 24 mg/kg were generally tolerated without local swelling, although two rats developed hematomas at 7 and 9 mo. Upon sacrifice of the remaining rats of this group (at 357 days) and of the other lower-dose groups (at 514-748 days), a total of nine fibrosarcomas and one rhabdomyosarcoma were found. Unlike the fibrosarcomas, the rhabdomyosarcoma did not invade surrounding muscle. Morphologically, the tumors resembled those found in other studies of the injection of heavy metals into rats. It is suggested that gold be added to the list of heavy metals that produce sarcomas in rats. (10 refs)

78-6134 **Mutagenic, Potentiating and Antimutagenic Activity of Certain Metallic Ions in the Rice Genetic System (Letter to Editor).** (Eng) Reddy, T. P. (Dept. Genetics, Osmania Univ., Hyderabad 500 007, India); Vaidyanath, K. *Curr Sci (Bangalore)* 47(14): 513-515; 1978.

Sets of rice seeds were treated with one of nine metallic salts and/or γ-rays. Seven metals produced chlorophyll mutations in the following order of potency: Ba, Cd (both equal to γ-rays), Cu, Hg, Sr, Fe, and Pb. Mn and Ca were not mutagenic. Sequential treatments suggested that radiation- and metal-induced genetic lesions interacted synergistically in mutation induction. (18 refs)

78-6135 **Cytogenetic and Biochemical Study of Lead-exposed Population (Meeting Abstract).** (Eng) Horvat, D. J. (Inst. Medical Res. and Occupational Health, Zagreb, Yugoslavia); Prpic-Majic, D. *Mutat Res* 53(2): 200-201; 1978. (no refs)

78-6136 Chromosome Analysis of Children Exposed to the Emission of Lead-Smelting Plant (Meeting Abstract). (Eng) Dresch, J. (Institut für Biologie, Gesellschaft für Strahlen- und Umweltforschung mbH München, D-8000 München, W. Germany); Bauchinger, M.; Schmid, E.; Englert, N. *Mutat Res* 53(2): 181; 1978. (no refs)

78-6137 Manganese Toxicity and Mutagenesis in Two Blue-Green Algae. (Eng) Singh, S. P. (Dept. Botany, Banaras Hindu Univ., Varanasi 221005, India); Kashyap, A. K. *Environ Exp Bot* 18(1): 47-53; 1978.

The toxicity and mutagenicity of manganese were examined in the coccoid blue-green alga *Chroococcus limneticus* and the filamentous blue-green alga *Plectonema boryanum*. *C. limneticus* responded slowly to $MnCl_2$ concentrations < 6 mM; at higher concentrations, the survival pattern declined exponentially. Only 1.6% of the cells could form colonies on nutrient agar in the presence of 0.01 M Mn. However, doses causing $< 50\%$ lethality (6 mM) were the most efficient at inducing the formation of abnormal cells. Higher doses were more lethal than teratogenic, although all survivors of 9 and 10 mM had abnormalities. The induction of phage- and streptomycin-resistant mutants of *P. boryanum* was investigated at $MnCl_2$ concentrations of 1.0×10^{-6} to 5.0×10^{-4} M. The resistant mutants arose 10-12 days after plating at frequencies ranging from 2.6 to 28.0×10^{-8} for phage resistance and 0.9 to 1.8×10^{-8} for streptomycin resistance. When the algae were exposed to graded concentrations of $MnCl_2$, a sigmoidal pattern of inactivation was observed. The streptomycin-resistant mutants showed increased sensitivity to $MnCl_2$ compared with those of phage-resistant and parent strains. In contrast to *C. limneticus*, microscopic observations revealed no abnormal cell formation or an increase in cell size. When the parent strain of *P. boryanum* was grown in the presence of low concentrations of $MnCl_2$, the phage-resistant mutants did vary in their sensitivity to Mn. The induced frequencies were concluded to be due to the induction of phage-resistant mutants. The max mutant frequencies were approx 10 times that of the background level. Streptomycin-resistant mutants were not induced to a significant level. (22 refs)

78-6138 In Vivo Mutagenesis Investigations of Four GRAS Chemicals (Meeting Abstract). (Eng) Jorgenson, T. A. (Stanford Res. Inst., Menlo Park, CA); Rushbrook, C. J.; Newell, G. W.; Green, S. *Mutat Res* 53(2): 205; 1978. (no refs)

78-6139 Mutagenic Activity of Ethylmercurichloride and Its Disintegration Product Mercury (Meeting Abstract). (Eng) Lekevichus, R. (Dept. Botany and Genetics, State Univ. Vilnius, Lithuanian SSR, USSR); Vijan-

skaja, T.; Stukiene, R. *Mutat Res* 53(2): 218-219; 1978. (no refs)

78-6140 Chromosome Distribution Studies in Phenyl Mercury Acetate Exposed Subjects and Age-related Controls. (Eng) Verschaeve, L. (Laboratorium voor Antropogenetica, Vrije Universiteit Brussel, Brussels, Belgium); Kirsch-Volders, M.; Hens, L.; Susanne, C. *Mutat Res* 57(3): 335-347; 1978.

The distribution of metaphase chromosomes in the peripheral lymphocytes of 16 men exposed to phenyl mercury acetate was compared with that in 12 unexposed men. One hundred metaphases from both populations were trypsin-banded and karyotyped. The relative position of the metaphase chromosomes was studied by means of centromere-centromere distances (Δ^2) and centromere-metaphase center distances obtained by computer-aided mathematical transformation of the individual metaphase coordinates. Blood mercury levels were within normal limits in both groups: 0 to 3.5 $\mu g/l$ in the controls and 0 to 5.6 $\mu g/l$ in the exposed subjects. Cytogenetic analysis showed only a significant increase in hypodiploidy in the exposed subjects. There were no translocations. However, there were 18 significantly different Δ^2 values between the two groups. For 12 chromosome groups, the values were greater among the exposed subjects, and for 6 groups the values were smaller for the exposed subjects. Pair 1-1 was the only one for which the value in the exposed subjects was obviously smaller. The alterations in chromosome distribution due to phenyl mercury acetate were particularly evident among the D-group chromosomes. (18 refs)

78-6141 Nickel-induced Erythrocytosis: Efficacies of Nickel Compounds and Susceptibilities of Strains. (Eng) Hopfer, S. M. (Dept. Lab. Medicine, University of Connecticut Sch. Medicine, Farmington, CT, 06032); Siderman, F. W.; Fredrickson, T. N.; Morse, E. E. *Ann N Y Acad Sci* 306: 396-402; 1978.

Dusts of nickel subsulfide (αNi_3S_2), nickel sulfide (αNi_3S_2), nickel monosulfide (βNiS), amorphous NiS , and metallic nickel were administered to male Fischer-344 rats intrarenally (ir) to assess the relative efficacies of the compounds for erythrocytosis (EC) induction. The dusts, which were suspended in 0.2 ml glycerol diluted 1:1 with NaCl soln (0.9% mM), were injected into the right kidney. From 4 to 10 wk postinjection, significant differences were observed among the mean blood packed cell volume (PCV) values with the order being $\alpha Ni_3S_2 > \beta NiS > \alpha Ni_3S_2 > Ni > amorphous NiS$. Crystalline βNiS induced an EC that was almost equal to that for αNi_3S_2 , but amorphous NiS did not induce any EC. To study in vitro rates of dissolution of these compounds, blood samples from Fischer-344 rats were spun and equivalent amounts of Ni in the above compounds and solutions were incubated. Analyses were performed at 2, 3, 8, and 10 days after

ubation. After 10 days, the concentrations of dissolved Ni serum ($\mu\text{g-atom/ml}$) were 5.86 amorphous NiS, 3.50 αNi 1.53 $\alpha\text{Ni}_2\text{Se}_3$, 0.62 βNiS , and 0.38 Ni dust. Six male and female mice of the Fischer-344 (Group H), NIH Black (Group I), Wistar-Lewis (Group J), and Long-Evans (Group K) strains were injected with 5 mg $\alpha\text{Ni}_2\text{S}_3$ ir to study strain differences in EC induction. At 2 to 8 wk, significant differences in mean PCV were seen in the following hierarchy: Group H > I > J > K. At 4 wk postinjection, significant differences were seen in the mean reticulocyte count: Group H > I > K > J. It is suggested that induction of EC following injection could provide a screening technique to test the carcinogenicities of Ni compounds. (16 refs)

78-6142 Nickel and Strontium Distribution in Some Mouse Tissues; Passage Through Placenta and Mammary Glands. (Eng) Jacobsen, N. (Dept. Microbiology, Dental Faculty, Univ. Oslo, Blindern, Oslo 3, Norway); Alfheim, I.; Jonsen, J. *Res Commun Chem Pathol Pharmacol* 3(3): 571-584; 1978.

The distribution and retention patterns of nickel and strontium in the soft and hard tissues of female mice were studied. Isotopes of Sr (0.12 $\mu\text{g/g}$) and Ni (0.14 $\mu\text{g/g}$) were injected as the chlorides. The Sr concentration was comparable in the liver, kidney, and heart tissue, but brain tissue showed a concentration of one-eighth of those of the other soft tissues. The radioactivity in the calvaria and long bones was eight times greater than that in the liver, and the radioactivity in the lower incisors was two to five times greater than that in the bones. The distribution in fetal tissues was similar to that of the maternal tissue, but the concentrations were reduced. The concentrations in sucklings nursing from a nonradioactive female decreased more rapidly than those in sucklings nursing from the biological, radioactive mother. Ni was recovered in all soft tissues, the highest concentrations occurring in the kidney and brain, respectively. Ni had no apparent affinity for the mineralized tissues, and it was transferred through the placenta to the progeny; in many tissues, the progeny had higher concentrations of radioactivity than the mother. Ni was also transferred to sucklings through the mother's milk. The results indicate that Sr is a bone-seeking mineral that forms a hard tissue reservoir, whereas there is a prolonged retention of Ni in mineralized tissue. (16 refs)

78-6143 The Induction of Malignant Tumors in Rats by Nickel-containing Aerosols. (Rus) Saknyn, A. (Res. Inst. Vocational Hygiene and Occupational Diseases, Sverdlovsk, USSR); Blokhin, V. A. *Vopr Onkol* 24(4): 48; 1978.

The results of a toxicologic experiment on the long-term effects of nickel-containing industrial dusts (NCID) are presented. Random-bred albino rats were (1) subjected to daily inhalations of NCID (av content of dust in the chamber, 70

mg/m^3), 5 hr/day, 5x/wk, for 6 mo; (2) inoculated ip with NCID soln (90-150 $\text{mg}/1.5\text{ ml}$); or (3) administered NCID soln intratracheally (20-40 $\text{mg}/0.6\text{ ml}$). Within 6-15 mo of ip NCID administration, 6/39 rats developed sarcomas (3 reticulosarcomas of the abdominal lymph nodes, 3 fibrosarcomas of the abdominal wall). After ip and intratracheal administration of NCID, 1/5 and 1/26 rats, respectively, developed lung cancer. The latent period of lung tumor development in both cases was 17 mo. (12 refs)

78-6144 Evaluation of Tris (2,3-dibromopropyl)phosphate for Mutagenic and Transforming Activity in a Battery of Test Systems (Meeting Abstract). (Eng) Brunsick, D. (Dept. Genetics, Litton Bionetics, Inc., Kensington, MD, 20795); Matheson, D.; Bakshi, K. *Mutat Res* 53(2): 160; 1978. (no refs)

78-6145 Test of O,O-Dimethyl(1-hydroxy-2,2,2-trichloroethyl)phosphonate (Trichlorofon) for Carcinogenic Activity in Rats Following Oral (Esophageal-Gastric Intubation) and Intraperitoneal Application. (Rus) Teichmann, B. (Zentralinstitut für Krebsforschung, Akademie der Wissenschaften der DDR, Lindenberger Weg 80, DDR-1115 Berlin-Buch, E. Germany); Hauschild, F.; Eckelmann, A. *Arch Geschwulstforsch* 48(2): 112-119; 1978.

The carcinogenicity of O,O-dimethyl(1-hydroxy-2,2,2-trichloroethyl)phosphonate (Trichlorofon) was examined in albino rats following po (esophageal-gastric intubation: 22 mg/kg 2x/wk) and ip (12 mg/kg 2x/wk) administration. Controls were given po or ip NaCl soln. All animals were treated for 90 wk; the experiment was terminated at 118 wk. There was no statistically significant difference in tumor incidence between treated and control animals by either route of application. (41 refs)

78-6146 Selenium and Cancer: Effects of Selenium and of the Diet on the Genesis of Spontaneous Mammary Tumors in Virgin Inbred Female C₃H/St Mice. (Eng) Schrauzer, G. N. (Dept. Chemistry, Univ. California at San Diego, La Jolla, CA, 92093); White, D. A.; Schneider, C. J. *Bioinorg Chem* 8(5): 387-396; 1978.

The efficacy of dietary selenium in reducing the incidence of spontaneous mammary tumors was studied in inbred female C₃/St mice, which show an 80% to 100% incidence of spontaneous tumor development. The animals were maintained on a basal diet (main source of protein, fishmeal) containing 0.45 ppm Se. Se supplementation was started at 2 mo of age and continued throughout life. Mice receiving 0.1 and 0.5 ppm Se in the drinking water gained wt faster than those receiving no Se in the drinking water (controls) or those receiving 1.0 ppm; the difference was significant at 1 yr. After

24 mo, tumor incidence in the control group was 42%, which was significantly lower than that in mice fed a diet containing 0.15 ppm Se, with meat and dried skimmed milk as major source of protein (80%-100%, previous study). The incidence of tumors declined progressively and the age at first tumor appearance increased progressively with increasing Se supplementation. Tumor growth rates did not differ significantly among the groups. A higher frequency of mesentery tumors was observed among controls (5/7) than among Se-supplemented animals (3/14). There was no histological evidence of liver damage in any group. The data suggest that a human diet rich in Se-containing seafoods and cereals could lower the cancer risk compared with a predominantly meat-based diet. (9 refs)

78-6147 Mutagenicity of Heavy Metals (Meeting Abstract). (Eng) Deknudt, G. (Dept. Radiobiology, Lab. Genetics, S.C.K.-C.E.N., B-2400 Mol, Belgium). *Mutat Res* 53(2): 176; 1978. (no refs)

78-6148 Relation Between Gastric Cancer Mortality Rate and Amounts of Zn in Rice in Nara Prefecture, Japan. (Jpn) Ikeda, K. (Dept. Hygiene, Nara Medical Univ., Kashiwara-shi, Nara Prefecture 634, Japan). *J Nara Med Assoc* 28(6): 909-916; 1977.

The relationship between gastric cancer mortality and zinc levels in the rice was determined for Nara Prefecture, Japan. This region has the highest gastric cancer mortality rate in Japan, although the rate has been declining in recent years; in one of its districts (Uda), however, the rate has been rising in the elderly. Unhulled rice from 129 areas in Nara Prefecture was analyzed for trace metals. Zn, Fe, and Cu levels were higher than the av values for the nation, and that of Ca was lower. The coefficient of variation was higher for Fe, Cu, and Cd than for Mg, Zn, and Ca. The distribution curves for Mg, Ca, and Fe skewed to the left, that of Zn had two peaks. The ratio of the metal concentration in unhulled and hulled rice was 0.08 for Mg, 0.54 for Zn, and 0.90 for Ca. Classification according to district showed that Uda had higher levels of Mg, Zn, and Ca in its rice than the other districts. Apparently, the high level of Zn in rice is related to the high gastric mortality rate. (9 refs)

78-6149 Short-term Toxicity Study of Chocolate Brown HT in Pigs. (Eng) Hendy, R. J. (British Industrial Biological Res. Assoc., Woodmansterne Road, Carshalton, Surrey SM5 4DS, England); Butterworth, K. R.; Gaunt, I. F.; Hooson, J.; Grasso, P. *Toxicology* 11(2): 189-192; 1978.

The toxicity of Chocolate Brown HT a food coloring agent, was studied in 10-wk-old male and female Large White pigs. The coloring was incorporated in the diet for 13 wk at a dose

level of 0, 5, 20, or 100 mg/kg/day. There were no abnormalities in the condition or appearance of the pigs, body wt gain, absolute organ wts, or urine. Feces of pigs given 100 mg/kg/day were dark brown and occasionally maroon in color. Significantly reduced Hb concentrations in all groups of treated male and elevated WBC counts in a single female of a single male were not considered to be related to treatment with Chocolate Brown HT. Scattered significant differences in relative organ wts between control and treated pigs were also considered to be unrelated to Chocolate Brown HT. On the basis of these results, an acceptable daily intake of Chocolate Brown HT for man would be 1 mg/kg or about 70 mg for an adult. (6 refs)

78-6150 The Mutagenicity of a Sauna Pine Scent Product in *Salmonella typhimurium* and *Saccharomyces cerevisiae* (Meeting Abstract). (Eng) De Marini, D. M. (Dept. Biological Sciences, Illinois State Univ., Normal, IL). *Mutat Res* 53(2): 178; 1978. (no refs)

78-6151 Chromosomal Damage and Hair Dyes (2 Letters to Editor). (Eng) Feinstein, A. R. (Dept. Medicine, Yale Univ. Sch. Medicine, New Haven, CT, 06510); Kirkland, D. J.; Lawler, S. D.; Venitt, S. *Lancet* 2(8059): 627-629; 1978.

Criticisms concerning an investigation of the possible carcinogenicity of hair dyes are listed, as are, in a subsequent letter, the replies to these criticisms. In the reply, it is repeated that an excess of chromosome damage (over that in controls) was found in a group of 60 professional hair-dyers. However, the excess was not definitive. (7 refs)

78-6152 On the Mutagenicity of Some Azo-Dyes (Meeting Abstract). (Eng) Busk, L. (Toxicology Lab., Swedish Natl. Food Admin., Box 622, S-751 26 Uppsala, Sweden); Albanus, L. *Mutat Res* 53(2): 161; 1978. (no refs)

78-6153 Investigations into Genetic Activity of Food Color Dyes (Meeting Abstract). (Eng) Smith, R. (Dept. Biological Sciences, Portsmouth Polytechnic, Portsmouth, England); Combes, R. D. *Mutat Res* 53(2): 201; 1978. (no refs)

78-6154 Correlation Between the Carcinogenicity of Organic Substances and Their Spectral Characteristics. (Eng) Veljkovic, V. (Boris Kidric Inst. Nuclear Sciences, Lab. Theoretical Physics, P.O. Box 522, 11001 Belgrade, Yugoslavia); Lalovic, D. I. *Experientia* 34(10): 1342-1343; 1978.

ce the UV radiation that is most effective in inducing can-
lies in the region 233-278 nanometers (nm), the UV ab-
sorption spectra of chemical carcinogens were analyzed to
determine whether they have peaks in this spectral region and
whether they differ in this respect from noncarcinogens. A
strong correlation was found between the UV spectral char-
acteristics and carcinogenicity of a variety of organic sub-
stances. The organic carcinogens had an absorption peak(s)
in the 206- to 248-nm region. These findings and those from
preceding work resulted in the formulation of the following
principle: if an organic chemical has an av quasi-valence
number < 3.20 and if its spectrum shows an absorption
peak(s) in the wavelength region 206-248 nm, it should be
considered carcinogenic. This rule was fulfilled for 91% of
chemicals tested. A mechanism of carcinogenesis is
proposed that assumes that cancer associated-DNA is a
blocked part of normal cell DNA. The most important step
in carcinogenesis is deblocking of cancer-associated DNA.
The deblocking mechanism is independent of the nature of
the carcinogenic agent and must be due to some characteristic
energy features. (6 refs)

78-6155 **Effects of Harman and Norharman on Spon-
taneous and Ultraviolet Light-induced Muta-
genesis in Cultured Chinese Hamster Cells.** (Eng) Chang, C.
(Dept. Human Development, Coll. Human Medicine, Mi-
chigan State Univ., East Lansing, MI, 48824); Castellazzi,
G.; Glover, T. W.; Trosko, J. E. *Cancer Res* 38(12):
27-4533; 1978.

The effects of nontoxic concentrations of two nonmutagenic
carboline derivatives, harman (HM) and norharman
(NH), on DNA repair and mutagenesis were studied in
cultured Chinese hamster (V79) cells. Cytotoxicity studies
showed that the colony-forming ability of the cells was unaf-
fected by 5 $\mu\text{g/ml}$ HM and decreased to only 85% by 10 $\mu\text{g/ml}$.
20 $\mu\text{g/ml}$ completely eliminated colony-forming ability,
whereas this dosage of NH inhibited plating efficiency to
only 65%. The survival of UV- or x-ray-damaged V79 cells
exposed to 30 or 50 μM HM was significantly reduced com-
pared with cells not treated with HM, and the effect was
concentration-dependent. The effect was not detectable at 10
 μM . The unscheduled DNA synthesis induced by UV light
(200-900 R) was slightly, but consistently, inhibited by HM,
but normal DNA synthesis was unaffected. The frequency
of spontaneous or UV light-induced ouabain-resistant or
thioguanine-resistant mutations was reduced by HM. The
effect of HM on survival and mutagenesis was greater than
that of NH and was detected primarily when cells were
exposed to HM immediately after UV irradiation. The muta-
tion frequency was reduced significantly by 28 μM NH at
3.5 days after UV irradiation and by 30 μM at 0-1.5 days
after irradiation. The data indicate that HM decreases the
capacity of V79 cells to repair DNA damage and fix muta-
tions, possibly because of its intercalation properties. (28 refs)

78-6156 **Perspective in Chronobiology of Air Pollution.**
(Eng) Stupfel, M. (French Natl. Inst. Health

(INSERM), 101 rue de Tolbiac, Paris 75013, France); Hal-
berg, F.; Mordelet-Dambrine, M.; Magnier, M. *Chronobi-
ologia* 4(4): 333-351; 1978.

Chronobiological experiments with rodents that determine
how cycles in air pollutant emissions synchronize, reinforce,
or interfere with the internal or endogenous rhythms charac-
teristic of human physiology or physiopathology are re-
viewed. (56 refs)

78-6157 **The Mutagenicity of a Dinitro-o-cresol-contain-
ing Pesticide.** (Hun) Nehez, M. (Kozegesz-
segugyi es Jarvanytani Intezet, SZOTE, Szeged, Hungary);
Selyes, A.; Paldy, A. *Egeszsegtudomány* 21(3): 237-243;
1978.

The mutagenicity of Krezonit E, which contains 50% dini-
tro-o-cresol (DNOC) as the active agent, was studied in vitro
in normal human WBC cultures and in vivo in bone marrow
cell cultures of treated male albino mice. The numbers of
human WBC with chromosome aberrations increased with
DNOC concentration, from 113/200 at 0.02 $\mu\text{g/ml}$ to
125/196 at 0.2 $\mu\text{g/ml}$ and to 129/200 at 2 $\mu\text{g/ml}$. Chroma-
tid-type aberrations accounted for 35.7% of all aberrations,
and their frequency was DNOC concentration-dependent;
chromosome-type aberrations accounted for 25%, and their
frequency was DNOC concentration-independent. Numerical
aberrations occurred at a frequency of 0.3%. The spon-
taneous aberration rate was 2.5%. In mice, the number of
cells with chromosome aberrations was (a) 75/100 in bone
marrow cultures from those treated with a single ip injection
of DNOC (20 mg/kg) and (b) 76-95/100 in those treated with
five ip injections of 1, 5, or 10 mg/kg on days 1, 3, 5, 7, and
9 before sacrifice on day 10, which indicates that the
mutagenicity of DNOC is higher after subacute treatment.
Most of the aberrations were of the chromosome type; labile
aberrations accounted for 4%-7%, stable aberrations for
68%-85%. Chromatid-type aberrations accounted for
1%-10%. The spontaneous aberration frequency was 1/100.
The findings show that DNOC is mutagenic. (24 refs)

78-6158 **Evaluation of Organic Extracts of Drinking
Water by Bacterial Mutagenesis (Meeting Ab-
stract).** (Eng) Loper, J. C. (Dept. Microbiology, Univ. Cin-
cinnati Medical Center, Cincinnati, OH); Schoeny, R. S.;
Tardiff, R. G. *Mutat Res* 53(2): 223-224; 1978. (2 refs)

78-6159 **Galactose-Resistant Forward Mutation Induc-
tion in *E. coli* K-12 and *S. typhimurium* galE or
galU Strains by Environmental Mutagens (Meeting Ab-
stract).** (Eng) Clarke, C. H. (Sch. Biological Sciences, Univ.
East Anglia, Norwich, NR4 4TJ, England); Shankel, D. M.
Mutat Res 53(2): 168; 1978. (no refs)

- 78-6160 Co-operative Mutagenic Actions of Bisulfite and Nitrogen Nucleophiles (Meeting Abstract).** (Eng) Hayatsu, H. (Faculty Pharmaceutical Sciences, Univ. Tokyo, Bunkyo-ku, Tokyo 113, Japan). *Mutat Res* 53(2): 197; 1978. (1 ref)

- 78-6161 The Detection of Mutation in Human Diploid Fibroblasts after Mutagen Treatment Using Non-selective Cloning and Enzyme Electrophoresis.** (Eng) Nelson, R. L. (Dept. Surgery, Univ. Illinois Hosp., P.O. Box 6998, Chicago, IL, 60680); Harris, H. *Mutat Res* 50(2): 277-283; 1978.

Human diploid fibroblasts from a 21-yr-old man (RM3), an 11-yr-old boy (FS1), and a male infant (PM1) were exposed to ethyl methanesulfonate (EMS): 400-1,100 $\mu\text{g}/\text{ml}$ and examined for mutation using nonselective cloning and enzyme electrophoresis. RM3 cultures had three doses of EMS, and between each dose the cells were cultured in fresh medium. The RM3 cells maintained a normal morphology and a reasonable incidence of survivors with doses of EMS up to 1,100 $\mu\text{g}/\text{ml}$. PM1 cells, however, died at levels of EMS > 400 $\mu\text{g}/\text{ml}$. The FS1 culture was intermediate in its sensitivity to EMS. Clones that were isolated following mutagen treatment were analyzed by starch-gel electrophoresis. Two mutants were identified: an electrophoretic variant of peptidase D (PEPD) was observed in an FS1 clone and a phenotypic alteration involving phosphoglucosyltransferase (PGM₁) was found in an RM3 clone. (8 refs)

- 78-6162 The Evaluation of Sixteen Carcinogens in the Rat Using the Micronucleus Test.** (Eng) Trzos, R. J. (Upjohn Co., 301 Henrietta St., Kalamazoo, MI, 49001); Petzold, G. L.; Brunden, M. N.; Swenberg, J. A. *Mutat Res* 58(1): 79-86; 1978.

The ability of the micronucleus test to detect clastogenic promutagens/procarcinogens was evaluated using 16 known carcinogens, 15 of which require tissue-specific metabolic activation. Groups of three male TUC(SD) spf (Sprague-Dawley) rats were administered a single ip injection of one of three doses/compound. The direct-acting mutagen ethyl methanesulfonate and the promutagens/procarcinogens cyclophosphamide and 4-nitroquinoline 1-oxide produced statistically significant dose-response increases in the number of micronucleated polychromatic RBC (MP-RBC). A marginally significant dose response was exhibited by dimethylnitrosamine. Aflatoxin B₁ significantly increased the number of MP-RBC at two doses, but a dose-response relationship was not found. The other agents, 2-acetylaminofluorene, 4-aminobiphenyl, benzidine, diethylnitrosamine, 9,10-dimethylbenzanthracene, 1,2-dimethylhydrazine, ethyl carbamate, β -naphthylamine, ethionine, hexametaphol, and metronidazole, did not increase the number of micronuclei significantly. Thus, the micronucleus test appears to lack the

sensitivity for detecting procarcinogen/promutagens that require tissue-specific metabolic activation or are rapidly inactivated. (22 refs)

- 78-6163 Mutagenicity of Some Environmental Chemicals in Salmonella Test Systems without Chromosomal Activation (Meeting Abstract).** (Eng) Adams, J. (Regional Science Teaching Center, Regional Staff Development Center, Graduate Sch., Notre Dame of Marbel C. Koronadal, South Cotabato 9708, Philippines). *Mutat Res* 53(2): 142-143; 1978. (no refs)

- 78-6164 Comparison of Toxicity and Mutagenicity of Butyl Methanesulfonate among Human Lymphoblast Lines.** (Eng) Slapikoff, S. A. (Dept. Biology, Tufts Univ., Medford, MA, 02155); Andon, B. M.; Thilly, W. *Mutat Res* 54(2): 193-196; 1978.

A comparison was made of the toxicity and mutagenicity of butyl methanesulfonate (BMS) among four diploid human lymphoblast lines: MIT-2, from a female mononucleosis patient; GM 130, from a normal man; MGL8B-2, a clone derivative of line MGL8 isolated from a normal man; and WI-L2, from a man with hereditary spherocytosis. Each line was exposed to 0-3 mM BMS for 24 hr. A period of ≥ 12 d was allowed for expression of the mutant phenotype, lack of hypoxanthine-guanine phosphoribosyltransferase activity. The mutant fraction was determined by soft-agar plating of the cultures in the presence and absence of 10 $\mu\text{g}/\text{ml}$ 6-thioguanine. BMS was toxic and mutagenic to lines MGL8B-2, MIT-2, and WI-L2, particularly to the latter two lines. In contrast, GM 130 cells demonstrated almost complete resistance to these effects of BMS, even at the highest concentration tested. Line GM 130 was also highly resistant to the toxic and mutagenic effects of methylnitrosourea and methyltrinitrosoguanidine, at concentrations highly toxic and mutagenic to line MIT-2. The reason(s) for the resistance of line GM 130 is not known. (9 refs)

- 78-6165 The Potential Carcinogenicity of Methyl Fluorosulphonate ($\text{CH}_3\text{OSO}_2\text{F}$; Magic Methyl).** (Eng) Ashby, J. (Central Toxicology Lab., Imperial Chemical Industries Ltd., Alderley Park, Macclesfield, Cheshire, England); Anderson, D.; Styles, J. A. *Mutat Res* 51(2): 285-287; 1978.

The potential carcinogenicity of methyl fluorosulfonate (MFS) was evaluated in two short-term in vitro tests, the Salmonella reverse mutation assay and in a cell transformation assay. In the former, MFS was mutagenic to *Salmonella* strain TA1538 in a dose-dependent manner at concentrations up to 500 $\mu\text{g}/\text{plate}$ in the absence of a metabolizing system. It produced a max of 140 revertants/plate (vs 20/plate

control cultures). MFS resulted in 100% lethality of baby hamster kidney 21/C13 cells at a concentration of 2.5 $\mu\text{g/ml}$, dose that also induced a max of 900 transformants/ 10^6 survivors (vs 60/ 10^6 in control cultures). Both cell survival and transformation induction were dose-dependent. The chemical activity of MFS and the results of these assays are consistent with this compound being a possible carcinogen. However, if not all, chemical alkylating agents are probably capable of producing positive results in *in vitro* assays, and they may not all prove to be carcinogenic *in vivo*. (9 refs)

6166 **Carcinogenicity and Mutagenicity of 47 Heteropolycyclic Compounds (Meeting Abstract).** (Eng) Glatt, H. R. (Pharmakologisches Institut der Universität, D6500 Mainz, W. Germany); Schwind, H.; Zajack, F.; Jacquignon, P. C.; Oesch, F. *Mutat Res* 53(2): 1-193; 1978. (no refs)

6167 **Mutagenicity of Chlorinated Cyclopentadienes Due to Metabolic Activation (Meeting Abstract).** (Eng) Goggelmann, W. (Gesellschaft für Strahlenforschung, Ingolstädter Landstrasse 1, 8042 Neuherberg, W. Germany); Greim, H.; Bonse, G.; Henschler, D. *Mutat Res* 53(2): 193-194; 1978. (no refs)

6168 **Asbestos Exposure and Related Neoplasia. The 28 Year Experience of a Major Urban Hospital.** (Eng) Hasan, F. M. (Div. Respiratory Sciences, Arizona Health Sciences Center, Univ. Arizona, Tucson, AZ, 85724); Ash, G.; Kazemi, H. *Am J Med* 65(4): 649-654; 1978.

Forty-nine asbestosis patients were seen at Massachusetts General Hospital between 1945 and 1972, 30 of whom also had cancer: 17 mesothelioma, 12 lung cancer (6 adenocarcinoma), and 1 prostate cancer. Only four asbestosis cases were diagnosed before 1960, and the diagnosis has been made with increasing frequency since then. Forty-eight patients had been exposed occupationally to asbestos, and 38 of these patients had a history of significant exposure in naval shipyards. One asbestosis patient with adenocarcinoma had no known history of exposure. The mean age of the patients was 67.7 yr, 45/49 were men, and only 4 were nonsmokers. In 37 patients, the exposure to asbestos was > 15 yr. Most patients with brief exposure had an associated neoplasm, and they presented with symptoms related to the tumor and not asbestosis. In 9/12 cancer patients who gave a history of exposure of < 5 yr, the neoplasm was mesothelioma. Unlike patients with uncomplicated asbestosis who often had worked with asbestos until the time of their illness, in patients with asbestos-related neoplasm there was a long interval between last exposure and emergence of symptoms. The mean latent period between onset of exposure and diagnosis of asbestos-related neoplasia was 26.3 yr. When pathologic data on 20 pa-

tients with autopsy studies were correlated with premortem roentgenologic changes, the latter seemed to underestimate the underlying disease. (26 refs)

78-6169 **Aryl Hydrocarbon Hydroxylase in Asbestos Workers.** (Eng) Naseem, S. M. (Channing Lab., Dept. Medicine, Harvard Medical Sch., 180 Longwood Ave., Boston, MA, 02115); Tishler, P. V.; Anderson, H. A.; Selikoff, I. J. *Am Rev Respir Dis* 118(4): 693-700; 1978.

The hypothesis that significant asbestos exposure may lead not only to long-term pulmonary accumulation of aromatic hydrocarbons, but also to increased inducibility (Ind) of aryl hydrocarbon hydroxylase (AHH) was tested by investigating lymphoblast AHH Ind in 47 asbestos workers (all men) and 31 control subjects (11 men, 20 women). Controls included 20 relatives of the exposed subjects, 12 of whom were spouses. Mean Ind [3-methylcholanthrene (3-mc)-dibenz(a,h)anthracene (DBA)-induced divided by basal enzyme activity] was significantly greater in the asbestos workers than in controls. The estimated change in the mean value of MC Ind was 0.88, that of DBA Ind 1.59. The increased Ind appeared to result almost exclusively from an increase in induced enzyme activity; there was little change in basal enzyme activity. Factors such as age, smoking status, year of first exposure, yield of lymphoblasts in culture, and blastogenesis rate (measured by incorporation of ^3H -thymidine into nucleic acid) had no effect on differences in AHH Ind ratios. The fact that spouses with potential household exposure did not have increased AHH Ind suggests that some minimal dose of asbestos is essential for this effect. Increased AHH Ind may bear some relation to the previously determined greatly increased risk (≥ 90 -fold) of bronchogenic carcinoma in asbestos workers who smoked cigarettes vs workers who did not smoke. (47 refs)

78-6170 **Cytogenetic Effects of the Gaseous Phase of Cigarette Smoke on Root-Tip Cells of *Allium sativum* L.** (Eng) Pandey, K. N. (T. H. Morgan Sch. Biological Sciences, Univ. Kentucky, Lexington, KY, 40506); Benner, J. F.; Sabharwal, P. S. *Environ Exp Bot* 18(1): 67-75; 1978.

The cytogenetic effects of the gaseous phase of cigarette smoke from five types of experimental cigarettes on the root tip cells of garlic, *Allium sativum* L., were determined. The root tips were exposed to 1, 2, 3, 4, 5, 10, 15, or 20 35-ml puffs of smoke at 7-min intervals. Chromosome abnormalities in the form of breakages (mostly fragments), bridges, lags, stickiness, and differential condensation were observed. Multinucleate cells, polyploid cells, and multipolar mitotic divisions were also noted. In general, the percentage of abnormalities increased when root tips were exposed to higher numbers of puffs. The mitotic index increased at low levels of smoke exposure, but it declined sharply at higher levels

of exposure. An analysis of variance indicated that the results were highly significant for cigarette types and exposure times with respect to the mitotic index as well as the total number of abnormally dividing cells. (24 refs)

- 78-6171 Analysis of a Tumor-inhibiting Fraction of Tobacco Smoke Condensate.** (Eng) Chamberlain, W. J. (Tobacco Lab., Science and Education Admin., Federal Reserve, U.S. Dept. Agriculture, P.O. Box 5677, Athens, GA, 30604); Snook, M. E.; Haebeler, A. F. *J Anal Toxicol* 2(4): 138-140; 1978.

A tumor-inhibiting fraction of cigarette smoke condensate containing ether-soluble neutrals was partially characterized by gel filtration and gas chromatography. In addition to solanesol, sterols, and other neutral oxygenated compounds such as acetophenones and fluorenones, this fraction contained 1,3-indandione, a compound new to cigarette smoke. (10 refs)

- 78-6172 A Comparison of Short Term Bioassay Results with Carcinogenicity of Experimental Cigarettes.** (Eng) Basrur, P. K. (Dept. Biomedical Sciences, Univ. Guelph, Guelph, Ontario, Canada); McClure, S.; Zilkey, B. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 2041-2048; 1978.

Four experimental cigarettes with known tumor induction potential in long-term tests were examined in two short-term bioassays, the *Salmonella typhimurium* mutagenicity assay and the mouse skin response test, and the results were compared. In the mutagenicity assay, rankings using condensates and smoke on a concentration basis were similar for the extremes, but the intermediate rankings were reversed. The mutagenicity ranks of condensate calculated on a per cigarette basis paralleled the total tumor potency trend established by long-term tests. The mutagenicity values obtained for fresh smoke on a per cigarette basis did not follow the same pattern. All of the tests, including the sebaceous gland suppression response in mouse skin with 10% condensates, detected the same cigarette as the most active. The epidermal hyperplasia response in mouse skin did not distinguish the extremes with either of the two concentrations of condensate (10% and 20%) used. It is concluded that the epidermal hyperplasia response, assumed to be a measure of the promoters in cigarette smoke, was not helpful in rating cigarettes whereas the sebaceous gland suppression response, which reflects the tumor initiator levels, and the mutagenicity response may be used as indices of biological insult correlated with the tumor-inducing potency of cigarettes. (8 refs)

- 78-6173 In Vitro Transformation of Hamster Embryo Cells by Betel Tobacco Extracts.** (Eng) Umehara, K. (Dept. Experimental Pathology, Cancer Inst., 1-3-3 Kami-Ikebukuro, Toshima-ku, Tokyo 170, Japan); Takama, S.; Fujie, S.; Matsushima, T.; Sugimura, T. *Toxicol* 2(4): 243-246; 1978.

The transforming effects of the ethanol and ethyl acetate extracts of betel tobacco on cryopreserved hamster embryo cells were studied. The ethanol extract induced one transformed colony at 100 $\mu\text{g/ml}$, and the ethyl acetate extract induced two transformed colonies at 5 $\mu\text{g/ml}$ and two colonies at 10 $\mu\text{g/ml}$. The positive results in this assay were significant, that no transformation was observed in control cultures. The ethyl acetate extract of betel tobacco may contain carcinogenic principle(s) of betel. (5 refs)

- 78-6174 Aryl Hydrocarbon Hydroxylase Induction in Human Lymphocytes and Sensitivity of Several Hematopoietic Parameters to Cigarette Smoking.** (Eng) Shultz, T. D. (Res. Section., Mountain States Tumor Inst., Boise, ID, 83702); Smith, C. E.; Steuart, C. D. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1677-1687; 1978.

Aryl hydrocarbon hydroxylase (AHH) activity was assayed in cultured lymphocytes obtained from 30 healthy smokers and 30 healthy nonsmokers. Total WBC, Hb, and hematocrit values were also compared for the smokers and nonsmokers. Smokers had significantly higher total WBC and Hb levels than nonsmokers; the hematocrit differences were not significant. Mean AHH activity was twice as high in smokers (0.058 unit) as in nonsmokers (0.029 unit). There was a negative correlation between the number of cigarettes consumed per day and the level of AHH induction. Arbitrary low, intermediate, and high subgroups within the smoker and nonsmoker populations suggested a trimodal distribution of absolute AHH levels, but the significance of this relationship could not be determined because of the small number of individuals in each subgroup. The lack of a correlation between the number of cigarettes consumed per day and AHH induction suggests that the latter is under genetic control rather than being a quantitative response to polycyclic hydrocarbons. (50 refs)

- 78-6175 Metabolic α -Hydroxylation of the Tobacco-Specific Carcinogen, N'-Nitrosornicotine.** (Eng) Chen, C. B. (Div. Environmental Carcinogenesis, National Cancer Inst. Disease Prevention, American Health Found., Bethesda, MD, 20892).

n, Valhalla, NY, 10595); Hecht, S. S.; Hoffmann, D. *Cancer Res* 38(11, part 1): 3639-3645; 1978.

The metabolism of N'-nitrosornicotine (NNN), a carcinogen that occurs in cigarette and cigar tobacco, mainstream and sidestream cigarette and cigar smoke, chewing tobacco, and snuff, was investigated in male F-344 rats, with emphasis on the α -hydroxylation pathway. Since α -hydroxynitrosamines are unstable, the model compounds 2'- and 5'-acetoxy-NNN were hydrolyzed to determine the decomposition products of 2'- and 5'-hydroxy-NNN. The major products of 2'-acetoxy-NNN were myosmine (MM: 50%-60%) and hydroxy-1-(3-pyridyl)-1-butanone (HPB: 5%-10%), that of 5'-acetoxy-NNN was 2-hydroxy-5-(3-pyridyl)-tetrahydrofuran (HPTHF: 60%-70%). Hydrolysis of a third model compound, 4-(N-carbethoxy-N-nitrosamino)-1-(3-pyridyl)-1-butanone, also gave HPB (50%-60%). These results support the intermediate role of electrophilic diazohydroxides and/or carbonium ions in the decomposition of 2'- and 5'-hydroxy-NNN. These electrophiles may be active forms of NNN, since all three model compounds were mutagenic toward *Salmonella typhimurium* TA100 without enzymatic activation. Upon incubation of ¹⁴C-NNN with rat liver microsomes, O₂, and NADPH, HPB and HPTHF were produced as the 2,4-dinitrophenylhydrazone derivatives. MM was formed in incubation mixtures analyzed without derivative formation. When rats were inoculated with 87.3 μ g ¹⁴C-NNN sc, 73%-85% of the dose appeared in the 48-hr urine. The major metabolites were 4-(3-pyridyl)-4-oxobutyric acid, 4-hydroxy-4-(3-pyridyl)butyric acid, and 5-(3-pyridyl)-tetrahydrofuran-2-one, the oxidation products of HPB and HPTHF. The results are consistent with the hypothesis that NNN is activated by α -hydroxylation. (36 refs)

78-6176 **Chemical Studies on Tobacco Smoke. II: Studies on Non-volatile Nitrosamines in Tobacco.** (Eng) Hecht, S. S. (Naylor Dana Inst. Disease Prevention, American Health Foundation, Dana Road, Valhalla, NY, 10595); Chen, C. B.; Dong, M.; Orna, R. M.; Hoffmann, D.; Iso, T. C. *Beitr Tabakforsch* 9(1): 1-6; 1977.

The isolation of the Z (syn) and E (anti) isomers of N'-nitrosornicotine (NNN) from tobacco, the influence of curing and stalk position on NNN levels in tobacco, and the synthesis of two new nitrosamines that may also be present in tobacco are described. Commercial and noncommercial (Burley, Bright) tobacco samples were used. The E/Z ratio in fine-cut chewing tobacco was determined by high-pressure liquid chromatography to be similar to that observed in solution at similar pH (7.6) and temperature (25 C). The interconversion rate was first-order and reversible, and it increased at lower pH. NNN was not detected (detection limit, 40 nanograms/g dry tobacco) in uncured tobacco. In tobacco grown in high N fertilization, the presence of NNN was confirmed beginning at the half-cured stage. NNN levels were higher in the high-N than in the low-N-fertilized samples, in which NNN was detected only after complete curing. No significant

correlation was observed between stalk position and NNN levels in one variety of Bright tobacco. 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-butanal and 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone, two additional nitrosamines that, theoretically, could also arise from nicotine were synthesized, and a detailed description of the process is given. (21 refs)

78-6177 **Long-Term Fate of [¹⁴C]Nicotine in the Mouse: Retention in the Bronchi, Melanin-Containing Tissues and Urinary Bladder Wall.** (Eng) Szuts, T. (Dept. Toxicology, Univ. Uppsala, Biomedical Center, Box 573, S-751 23 Uppsala, Sweden); Olsson, S.; Lindquist, N. G.; Ullberg, S.; Pilotti, A.; Enzell, C. *Toxicology* 10(3): 207-220; 1978.

The long-term fate of N-methyl-¹⁴C- and 2'-¹⁴C-labeled nicotine in pigmented C57BL and albino NMRI mice was studied using whole-body autoradiography, liquid scintillation counting, and thin-layer chromatography (TLC). Thirty days after injection, N-methyl-¹⁴C-nicotine persisted in the pigmented uveal tract, respiratory mucosal lining, and urinary bladder wall of the pigmented mice. Late-gestation fetuses showed a high accumulation in the eye, inner ear, and respiratory tract. The distribution was similar in the albino mice, except that no accumulation was found in structures corresponding to the melanin-containing tissues of the pigmented mice. The activity levels in the bronchi decreased faster when 2'-¹⁴C-nicotine was used. Quantitative measurements of the retention of the two ¹⁴C-labeled nicotine preparations confirmed the autoradiographic findings. TLC studies showed that no nicotine was present in the lungs after 24 hr and that only unmetabolized nicotine was present in melanin after 4 hr. Some reactive nicotine metabolites may be responsible for the retention in the urinary bladder wall, which persisted even after rinsing. The accumulation and long-term retention of nicotine in melanin-containing structures might accelerate the development of drug-induced or senile changes in these tissues. An accumulatory mechanism might also participate in the pathogenesis of urinary bladder cancer. (23 refs)

78-6178 **Dietary Fibre and Experimental Colon Cancer (Letter to Editor).** (Eng) Heaton, K. W. (Dept. Medicine, Bristol Royal Infirmary, Bristol BS2 8HW, England); Williamson, R. C. *Lancet* 2(8093): 784-785; 1978.

The protective effect of bran against colon cancer induced by cycasin derivatives in rats, an effect not observed with a diet containing three times the ordinary amount of dietary fiber, may be due at least in part to its lower energy value. Bran-fed animals grew more slowly and, for most of the experiment, their body wt (and thus their carcinogen dose) was only half that of rats receiving the fiber-enriched chow. (11 refs)

78-6179 Protective Effect of Dietary Fibre Against Chemically Induced Bowel Tumours in Rats. (Eng) Fleiszer, D. (McGill Univ. Surgical Clinic and Dept. Pathology, Montreal General Hosp., Quebec, Canada); MacFarlane, J.; Murray, D.; Brown, R. A. *Lancet* 2(8089): 552-553; 1978.

The effects of different levels of dietary fiber (0-28 g/100 g diet) on stool production and on the incidence of dimethylhydrazine-induced (20 mg/kg/wk for 20 wk, sc) large intestinal carcinomas were studied using inbred male Chester Beatty rats. Stool mass and frequency decreased with decreasing dietary fiber, the differences in stool frequency and mass between the highest fiber level group and the other groups being statistically significant ($p < 0.05$). The incidence of colonic carcinoma increased with decreasing fiber content except that the rats given 5 g fiber/100 g diet had more carcinomas (70.6%) than the group given 0 g fiber/100 g diet (42.1%). The difference in carcinoma incidence between the group given the greatest amount of fiber (5.9%) and the other groups was significant. The results support the hypothesis that a diet with high fiber content leads to frequent, bulky stools and a reduced incidence of colonic cancer. (14 refs)

78-6180 Dietary Fibre and Experimental Colon Cancer (Letter to Editor). (Eng) Cruse, J. P. (Surgical Unit, Univ. Coll. Hosp. Medical Sch., London WC1E 6JJ, England); Lewin, M. R.; Clark, C. G. *Lancet* 2(8094): 843-844; 1978.

Several errors in a previous report of a study of the protective effects of dietary fiber against chemically induced rat colon cancer are pointed out. In similar experiments of the authors, dietary fiber did not affect time to tumor or survival of dimethylhydrazine-treated rats significantly. Furthermore, the incidence of colon carcinoma was 100% at 1 yr in all dietary groups, regardless of fiber content. (4 refs)

78-6181 Mutagenic Properties of Ethylidene Gyromitrin and Its Metabolites in Microsomal Activation Tests and in the Host-Mediated Assay. (Eng) von Wright, A. (Food Res. Lab., Technical Res. Center Finland, Biologinkuja 1, SF-02150 Espoo 15, Finland); Niskanen, A.; Pyysalo, H. *Mutat Res* 54(2): 167-173; 1978.

The mutagenicity of the main poisonous hydrazine derivative of the edible mushroom false morel (ethylidene gyromitrin: EG) and its metabolites (N-methyl-N-formylhydrazine and N-methylhydrazine: MFH and MH, respectively) was tested by the microsomal and host-mediated assay. In the microsomal activation tests, the microsomal fraction of mouse liver did not increase the mutagenic or bacteriocidal properties of the EG and its metabolites to *Salmonella typhimurium* strain TA100. None of the compounds caused deaths due to poisoning when given by stomach tube to NMRI mice, although

two mice showed typical neurological symptoms after treatment with EG. EG and MH, but not MFH, inhibited growth of bacteria injected ip into the experimental mice. There were no increases in mutation frequency in the bacteria grown in treated mice compared with control bacteria. The data confirm the mutagenicity of MH and suggest that a bacteriocidal metabolite is formed from EG. (15 refs)

78-6182 Relationships Between the Mutagenic and Carcinogenic Effects of Hydrazine Derivatives. (Jpn) Shimizu, H. (Dept. Public Health, Jikei Univ. School of Medicine, Tokyo, Japan); Hayashi, K.; Takemura, N. *J. Hyg* 33(3): 474-485; 1978.

The mutagenic and carcinogenic effects of hydrazine derivatives were investigated. The microbial mutagenicity of 26 hydrazine compounds was tested with *Salmonella typhimurium* TA98, TA100, TA1537, in the presence or absence of S-9 mix. Eighteen of the 26 compounds were known carcinogens but the carcinogenicity of the remaining compounds was unknown. Five carcinogens such as 2-hydroxyethylhydrazine, butylhydrazine oxalate, phenylhydrazine HCl, beta-phenylthylhydrazine sulfate, and p-tolylhydrazine HCl, were mutagenic to TA100, with and without the S-9 mix. Two carcinogens, hydrazine hydrate and hydrazine sulfate, were mutagenic to TA100 only in the presence of the S-9 mix. 2,4-Dinitrophenylhydrazine and p,p'-oxybisbenzidine were mutagenic to TA100, with and without the S-9 mix. Phenylhydrazine HCl, p-tolylhydrazine HCl, and 2,4-dinitrophenylhydrazine were mutagenic to TA1537, with and without the S-9 mix. Phenylhydrazine HCl and p-tolylhydrazine HCl were mutagenic to TA98 with the S-9 mix, but 2,4-dinitrophenylhydrazine was mutagenic to TA98 without the S-9 mix. (78 refs)

78-6183 Transplacental Effect of Urethane on DNA Synthesis in Organ Cultures of Murine Embryonic Lung Tissues. (Rus) Kolesnichenko, T. S. (Dept. Carcinogenic Agents, Oncological Res. Center, Acad. Medical Sciences USSR, Moscow, USSR); Gusina, N. B. *Biull Eksp Biol Med* 86(10): 426-428; 1978.

The effect of transplacental exposure to urethane on DNA synthesis in embryonal lung tissue cultures was studied in line A mice. The pregnant mice embryonal lung tissue culture was studied in line A mice. The pregnant mice received single sc injection of 1 mg/kg urethane on day 15, 17, or 19 of embryogenesis. The embryos were extracted 1 day after injection, cultured, and examined 1, 7, 15, and 21 days later. DNA synthesis was determined from the ³H-thymidine labeling index. Embryonal lung tissue cultures from untreated pregnant mice served as controls. Urethane inhibited DNA synthesis during the first 24 hr; the labeling index was one-third of that found in the controls. After a normalization

the labeling index peaked on day 7 of culture; ie, 8 days after transplacental exposure to urethane. The labeling index decreased by day 15 of culture, but it was still higher than that of controls. On day 21 of culture, DNA synthesis was seen in the experimental explants only, but not in controls. (11 refs)

78-6184 Effect of Butylated Hydroxytoluene and Paraquat on Urethan Tumorigenesis in Mouse Lung. (Eng) Bojan, F. (Inst. Hygiene and Epidemiology, Univ. Medical Sch., 4012 Debrecen, Hungary); Nagy, A.; Herman, J. *Bull Environ Contam Toxicol* 20(4): 573-576; 1978.

The effect of administration of butylated hydroxytoluene (BHT) or Paraquat on urethan carcinogenesis was studied in the mouse lung. Female CFLP mice (30-35 days old) were treated with BHT (100-800 mg/kg ip) and injected with urethan (1 mg/g ip) 2 or 26 days before, 2-14 days after, or concomitantly with BHT. Other mice received a urethan injection (1 mg/g ip) concomitantly with or 1-4 days after Paraquat (5, 10, or 20 mg/kg ip). When given alone, BHT and Paraquat stimulated mitotic activity in type II alveolar cells and enhanced the carcinogenicity of urethan in the mouse lung. The potentiating effect of BHT was significant when the urethan was given 6-7 days after BHT; the yield of induced lung tumors increased in proportion to the BHT dose. Paraquat also significantly enhanced the number of lung tumors induced by urethan given 2 days after Paraquat; this effect was also dose-dependent. Previous reports have shown that lung tumors induced by urethan develop from type II alveolar cells. These results suggest that BHT and Paraquat enhance the carcinogenicity of urethan by increasing the number of proliferating target cells in the lung. (13 refs)

78-6185 Vinyl Carbamate as a Promutagen and a More Carcinogenic Analog of Ethyl Carbamate. (Eng) Ahl, G. A. (McArdle Lab. Cancer Res., Univ. Wisconsin Center Health Sciences, Madison, WI, 53706); Miller, J. A.; Miller, E. C. *Cancer Res* 38(11, part 1): 3793-3804; 1978.

This study was undertaken to test the hypothesis that vinyl carbamate (VC) is a proximate carcinogenic metabolite of ethyl carbamate (EC). When administered topically or ip at the same doses, VC was 10-50 times more active than EC in the initiation of skin tumors and lung adenomas in CD-1 and A/Jax mice. VC was mutagenic to *Salmonella typhimurium* A1535 and TA100, but only in the presence of ADPH-fortified rat or mouse liver mitochondrial supernatant fractions. This mutagenicity was strongly inhibited by cytochrome P-450 inhibitors. EC was not mutagenic for *S. typhimurium* in the presence or absence of liver fractions. Tests to detect VC as an in vivo metabolite of EC in 5-day-old CD-1 and adult A/Jax mice, performed under a variety of conditions, were unsuccessful. However, desaturation of the ethyl group during metabolism is suggested by the lower $^3\text{H}/$

^{14}C ratios in the hepatic DNA, ribosomal RNA, and protein of adult A/Jax mice administered [ethyl-1,2- ^3H ; 1- ^{14}C]-EC vs the $^3\text{H}/^{14}\text{C}$ ratios of the EC. The qualitatively similar, but much higher, carcinogenic activity of VC compared to that of EC suggests that the metabolic pathways of these two carbamates may converge in the formation of similar or identical electrophilic reactants that bind covalently to macromolecules in vivo and initiate carcinogenesis. (40 refs)

78-6186 Tolerances and Exemptions from Tolerances for Pesticide Chemicals in or on Raw Agricultural Commodities. (Eng) Johnson, E. L. (Office of Pesticide Programs, EPA, 401 M St. SW, Washington, DC, 20460). *Fed Regist* 43(219): 52486-52487; 1978.

An 0.2 ppm tolerance for residues of the herbicide isopropyl *m*-chlorocarbamate (CIPC) on soybeans has been established. The toxicological data supporting the proposed tolerance included a rat po LD study with an LD of 5-7.5 g/kg, a 2-yr rat feeding study with a no-observable-effect level (NOEL) of 2,000 ppm, a 1-yr dog feeding study with NOEL of 2,000 ppm, two 18-mo mouse carcinogenicity studies at a feeding level of 160 mg/kg/day (negative), a 116-wk mouse carcinogenicity study at a feeding level of 1,000 ppm (negative), a 33-mo hamster carcinogenicity study at a feeding level of 2,000 ppm, and a three-generation rat reproduction study (negative at 1,000 ppm). Desirable data that are lacking are a teratology study, a delayed neurotoxicity study, and an adequate study for effects on cholinesterase. The proposed 0.2 ppm tolerance of soybeans will not contribute residues to the human diet. The 2,000 ppm NOEL in the 2-yr rat feeding study supports an intake of 20 ppm in the human diet using a 100-fold safety factor. A 50 ppm tolerance was previously established for CIPC residues on potatoes, and interim 50 ppm tolerances were established residues on the hay of alfalfa, clover, and grass. (no refs)

78-6187 Chromosome Investigations of Persons Exposed to Dimethylcarbamoyl Chloride and Diethylcarbamoyl Chloride. (Eng) Fleig, I. (Occupational Medicine and Health Protection, BASF Aktiengesellschaft, 6700 Ludwigshafen/Rhein, W. Germany); Thiess, A. M. *J Occup Med* 20(11): 745-746; 1978.

Chromosome analyses were made of 10 chemical plant employees who had been exposed to dimethylcarbamoyl chloride (DMCC) and diethylcarbamoyl chloride (DECC) for 4-17 yr and 10 age-matched controls. The frequency of aberrant metaphases in the exposed group was 1.9%, while that in the controls was 1.3%. Chromosome-type aberrations comprised 1.1% of the aberrant metaphases in the exposed subjects and 0.8% in the controls. The differences were not statistically significant, indicating that DMCC and DECC exposure does not cause chromosome damage. (3 refs)

- 78-6188 Acetaldehyde Induces Cross-Links in DNA and Causes Sister-Chromatid Exchanges in Human Cells.** (Eng) Ristow, H. (Institut für Genetik, Freie Universität Berlin, Arnimalle 5-7, D-1000 Berlin 33, W. Germany); Obe, G. *Mutat Res* 58(1): 115-119; 1978.

Evidence is presented that acetaldehyde (AA), the first metabolite of ethanol, induces cross-links between calf thymus complementary DNA strands and induces sister chromatid exchanges (SCE's) in human lymphocytes. The DNA was incubated for 30 min at 37 C with 1 M AA and dialyzed overnight at 4 C. The DNA was then heated to 90 C, rapidly cooled, and reheated. The thermal transition midpoint (T_m) of AA-treated DNA was 66.2 C during the first heating and 66.6 C during the second. The T_m of control, untreated DNA was 70 C during the first heating, and no characteristic thermal transition was noted during the second. Thus, after heating and cooling, the hyperchromicity of AA-treated DNA was 60% of the original. In contrast, there was only an approx 20% loss of hyperchromicity with control DNA. It is suggested that cross-links were induced between the DNA strands by AA, leading to spontaneous reannealing of part of the DNA after the first heating and resulting in a partial reversal of the hyperchromic effect. Control DNA showed less spontaneous renaturing, possibly because of a lack of cross-links. Hyperchromicity after thermal denaturation was lowered as a function of AA treatment time; spontaneous reannealing after cooling was also related to treatment time. Both processes were dependent on AA concentration. Addition of 0.0005%-0.002% AA 24 or 48 hr before fixation of human lymphocyte cultures induced SCE's in a dose-dependent manner. The number of SCE's produced by the 24-hr treatment was almost doubled when cells were treated for 48 hr. These results indicate that ethanol, via AA, may be mutagenic in humans. (22 refs)

- 78-6189 Effect of Colostomy on Intestinal Carcinogenesis of Methylazoxymethanol Acetate in Rats.** (Jpn) Matsubara, N. (First Dept. Surgery, Sch. Medicine, Gifu Univ., Tsukasa-cho, Gifu 500, Japan). *Gifu Daigaku Igakubu Kiyo* 25(6): 501-520; 1977.

The effect of colostomy (COS) on the intestinal carcinogenicity of methylazoxymethanol acetate (MAM acetate) was studied in 5-wk-old Sprague-Dawley rats. Groups 1-3 rats underwent double COS, single COS, and no surgery, respectively, plus treatment with MAM acetate (25 mg/kg/wk x 10, iv). Group 4 rats served as untreated controls. Rats in Groups 1-3 developed multiple intestinal tumors, and the tumor incidence in these groups was similar. Tumor multiplicity was higher in Group 3 than in Group 2. The incidence of colon tumors in Group 1 was 64.5% in the oral one-third of the colon, 83.8% at the COS site, and 41.9% in the anal two-thirds; the mean number of tumors per rat in these locations was 1.4, 1.5, and 0.6, respectively. In Group 2 rats the tumor incidence was 28% in the oral section, 84.0% at the COS site, and 80.0% in the anal section; the mean number

of tumors per rat was 0.2, 1.4, and 1.0, respectively. MAM tumors in the anal section were at the oral stump of the colon and they were larger than those in other areas. In Group 1 tumor incidence was 72.2% in the oral section and 100% in the anal section, and the mean number of tumors per rat was 1.3 and 4.5, respectively. It was concluded that the carcinogenic metabolite of MAM acetate can reach the intestinal mucosa via the vascular system as well as through the biliary tract. (24 refs)

- 78-6190 Separation and Analysis of Various Carcinogenic Aromatic Amines and Their Metabolites.** (Eng) Weisburger, E. K. (Carcinogen Metabolism and Toxicology Branch, NCI, Bethesda, MD, 20014); Grantham, H.; Benjamin, T. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, NY) Vol. 2(Part 1), 2404 pp.; 1961-1972; 1978.

The continued application of various chromatographic techniques, including paper, thin-layer, gas, and high-pressure liquid chromatography, has led to improvements in the separation and identification of metabolites of three carcinogenic aromatic amines, N-2-fluorenylacetamide, 6-aminochrysene, and 2,4-toluenediamine. This in turn has increased knowledge of their metabolic pathways. (24 refs)

- 78-6191 The Distribution of DNA Repair Synthesis in Chromatin and its Rearrangement Following DNA Damage with N-Acetoxy-2-acetylaminofluorene.** (Eng) Tlsty, T. D. (Dept. Pathology, Washington Univ. Sch. Medicine, St. Louis, MO, 63110); Lieberman, M. W. *Nucleic Acids Res* 5(9): 3261-3273; 1978.

The distribution of DNA repair synthesis in the chromatin of confluent human diploid fibroblasts damaged with N-acetoxy-2-acetylaminofluorene (NA-AAF, 5-100 μ M) was studied. The amount of repair synthesis was dependent on NA-AAF concentration up to 10 μ M and much less at concentrations > 25 μ M. Kinetic analysis of staphylococcal nuclease digestion data showed that more repair synthesis occurred in nuclease-sensitive regions of chromatin than in nuclease-resistant regions. Data obtained by gel electrophoresis were in good agreement with the kinetic data, and they indicated that more repair synthesis occurred in linker DNA than in core DNA. Continuous-labeling and pulse-chase experiments showed that with time, an increasing proportion of nucleotides inserted during repair synthesis was present in the nuclease-resistant regions of chromatin. Rearrangement involved movement of label incorporated by repair synthesis into core DNA. Repair synthesis was a biphasic process. The data suggest that there is a relationship between rates of

synthesis and chromatin rearrangement. (28 refs)

6192 **An Investigation into the Effects of the Carcinogens Benzidine and 2-Acetylaminofluorine (AF) on the Control of Cholesterol Metabolism in the Rat and the Mouse.** (Eng) Depass, L. R. (Univ. Arkansas Medical Cent., Little Rock, AR, 72201). *Diss Abstr Int B* 39(5): 6192; 1978. (no refs)

6193 **Enhancement of Rat Hepatocellular-altered Foci by the Liver Tumor Promoter Phenobarbital: Evidence that Foci Are Precursors of Neoplasms and that Phenobarbital Acts on Carcinogen-induced Lesions.** (Eng) Tanabe, K. (Kagoshima Univ. Sch. Medicine, 1208-1, Shimo-Ogino, Kagoshima 890, Japan); Williams, G. M. *J Natl Cancer Inst* 61(5): 1311-1314; 1978.

The mechanism of liver tumor promotion by phenobarbital (PB) was investigated in inbred F344 rats fed 2-fluorenylacetylamide (FAA) and an iron-loading diet. Rats fed a diet containing 0.02% FAA, 0.8% 8-hydroxyquinoline, and 2.19% ferrous gluconate (FG) for 13 wk developed numerous iron-resistant, altered foci in every sub-lobule of the liver. Following cessation of carcinogen feeding, most foci reverted to iron accumulation and could no longer be detected by 24 wk, although some persisted for this interval. At the end of 24 wk after FAA termination, 33% and 78% of two groups of rats had tumors. When 0.05% PB was added to the diet for 24 wk after termination of FAA, the iron-resistant foci persisted, and the incidence of liver lesions was 78%-89%. The same levels of PB in the iron-loading diet of rats not given FAA previously did not induce iron-resistant foci. It was concluded that PB did not induce iron-resistant foci, but acted to retard the reversion of FAA-induced foci. (33 refs)

6194 **Diffusion Chamber Cultures for Mutagenic and Carcinogenic Assays (Meeting Abstract).** (Eng) Wang, C. C. (Dept. Experimental Biology, Roswell Park Memorial Inst., Buffalo, NY, 14263); Furukawa, M. *Mutat Res* 53(2): 201-202; 1978. (no refs)

6195 **Cyclophosphamide-induced Oncogenic Transformation, Chromosomal Breakage, and Sister Chromatid Exchange Following Microsomal Activation.** (Eng) Benedict, W. F. (Div. Hematology-Oncology, Dept. Medicine, Childrens Hosp. Los Angeles, Los Angeles, CA, 90027); Banerjee, A.; Venkatesan, N. *Cancer Res* 38(9): 2229-2234; 1978.

Cyclophosphamide (CP), an extensively used cancer chemo-

therapeutic agent, requires metabolic activation through a mixed-function oxygenase system. The capacity of this agent to produce oncogenic transformation and chromosomal damage, including increases in sister chromatid exchanges, was investigated in cultures of C3H/10T1/2CL8 mouse cells and A(T) Cl-3 Syrian hamster pseudodiploid cells with or without an exogenous liver metabolic activation system. This system was prepared from the livers of Sprague-Dawley rats inoculated with Aroclor 1254 (500 mg/kg). No oncogenic transformation or chromosomal aberrations were produced by CP in the absence of metabolic activation, whereas significant transformation, chromosomal breaks, and increases in sister chromatid exchanges were observed when the activation system was incorporated into the assays. The oncogenic transformation and chromosomal changes were completely eliminated by removing glucose 6-phosphate and NADP from the system. These studies emphasize the need for incorporating an activation procedure into short-term assays used to evaluate the mutagenic and/or oncogenic potential of various chemicals. (14 refs)

78-6196 **Acute Myelocytic Leukemia 11 Years Following Successful Therapy of Hodgkin's Disease. Report on the Problems of a Second Tumor.** (Ger) Ritter, J. (Kinderklinik der Stadt Braunschweig, Holwedestrasse 16, D-3300 Braunschweig, W. Germany); Oehme, J. *Med Klin* 73(39): 1360-1362; 1978.

Therapy-resistant acute myelogenous leukemia (AML) was diagnosed in a 21-yr-old man who underwent therapy for Hodgkin's disease at the age of 10 and for recurrences at ages 11 and 14-16 yr. The first therapy comprised surgical removal of the tumor, radiation (2,000 rads' focal dose), and several cycles of cytostatics (total doses: 23.6 g cyclophosphamide, 80 mg triethylenemelamine, 40 g procarbazine, 2.3 g chlorambucil, 130 mg vinblastine). Cyclic chemotherapy was repeated with the addition of Ultralan for the recurrences and discontinued 5 yr before the diagnosis of AML. For the first recurrence, radiation (1,600 rads' focal dose) was given. After the diagnosis of AML based on a bone-marrow biopsy, vigorous chemotherapy was given but the patient did not respond. An autopsy showed all the usual signs of AML with no microscopically evident infiltration of Hodgkin's disease. This case suggests that chemotherapy for Hodgkin's disease Stage I or II should not be used without consideration of the possibility of causing a second malignancy. This does not apply to the more advanced stages. A 5% incidence of AML in patients surviving 10 yr after radiotherapy + chemotherapy for Hodgkin's disease has been reported. (13 refs)

78-6197 **Acute Leukemia Following Cyclophosphamide Therapy for Sjogren's Syndrome.** (Eng) Hochberg, M. C. (Rheumatology Div., Good Samaritan Hosp., 5601 Loch Raven Blvd., Baltimore, MD, 21239); Shulman, L. E. *Johns Hopkins Med J* 142(6): 211-214; 1978.

A 56-yr-old woman with Sjogren's syndrome (SS) developed acute myelomonocytic leukemia after 31 mo of cyclophosphamide (CP) therapy (50 mg/day initially, with the dose eventually increased to 150 mg/day). Since the leukemia occurred 12 yr after the onset of SS, CP may have been a significant etiologic factor. (22 refs)

78-6198 Isolation and Structure Determination of Mutagenic Substances in L-Glutamic Acid Pyrolysate. (Eng) Yamamoto, T. (Shizuoka Coll. Pharmacy, 2-2-1 Oshika, Shizuoka-shi, Shizuoka 422, Japan); Tsuji, K.; Kosuge, T.; Okamoto, T.; Shudo, K.; Takeda, K.; Iitaka, Y.; Yamaguchi, K.; Seino, Y.; Yahagi, T.; Nagao, M.; Sugimura, T. *Proc Jpn Acad Ser B Phys Biol Sci* 54(5): 248-250; 1978.

Two strongly mutagenic substances produced by pyrolysis of L-glutamic acid were isolated and their structures determined. Pyrolysis was performed by heating 10 kg L-glutamic acid, collecting the resulting tar, distilling the basic fraction, and then purifying this fraction by silica gel, Sephadex column, alumina column, and thin-layer chromatography. Acidification of the two resulting bands with 0.5% HBr produced two hydrobromides with differing Rf values. These compounds were designated Glu-P-1 and Glu-P-20 mutagenicity, determined by the Ames assay, increased with each purification step. The activity of the final purified compounds was 13,700 and 1,600 revertants/ μ g. Mass spectroscopy and elemental analysis determined that the formulas of Glu-P-1 and Glu-P-2 were $C_{11}H_{10}N_4$ and $C_{10}H_8N_4$, respectively. Nuclear magnetic resonance and UV spectra indicated that the two compounds differed by an aromatic methyl group. The structure of Glu-P-1 was determined to be 2-amino-6-methyldipyrido[1,2-a:3',2'-d]imidazole, that of Glu-P-2 2-aminodipyrido[1,2-a:3',2'-d]imidazole. (7 refs)

78-6199 Azaserine Carcinogenesis: Organ Susceptibility Change in Rats Fed a Diet Devoid of Choline. (Eng) Shinozuka, H. (Dept. Pathology, Univ. Pittsburgh, Sch. Medicine, Pittsburgh, PA, 15261); Katyal, S. L.; Lombardi, B. *Int J Cancer* 22(1): 36-39; 1978.

The effects of feeding a choline-deficient (CD) diet on the carcinogenicity of azaserine (AS) in the rat pancreas were investigated in groups of male Wistar rats. Groups I and II were fed a CD or choline-supplemented diet, respectively, and inoculated ip with AS (30 mg/kg in 2 ml saline) 2x/wk for the first 4 wk and 1x/wk thereafter, for a total of 14 injections. Control groups were fed either of the diets and inoculated with saline. Four to seven animals from each group were killed 1, 4, and 6 mo after the first injection, and the liver and pancreas were examined histologically. None of the controls developed hepatic or pancreatic tumors, and no hepatomas were observed in the 12 Group II rats. In contrast, 3/5 and 5/7 Group I rats killed at 4 and 6 mo, respectively, had multiple hepatomas. Both Groups I and II rats

developed multiple atypical acinar cell nodules (AACN) in the pancreas after 4 mo. However, the number of AACN was significantly smaller in Group I rats (mean, 7.6) than in Group II (mean, 16.0). It is concluded that the CD diet changed the organ susceptibility to the carcinogenic action of AS, enhancing hepatocarcinogenesis and reducing the effect of the carcinogen on the pancreas. (16 refs)

78-6200 Persistence of DNA Damage in Rat Pancreas Following Administration of Three Carcinogens and/or Mutagens. (Eng) Lilja, H. S. (Dept. Pathology, University of New Hampshire Medical Sch., Hanover, NH, 03755); Curphey, J. D.; Yager, J. D.; Longnecker, D. S. *Chem Biol Interact* 22(2): 287-295; 1978.

Alkaline sucrose gradients were used to detect and follow the persistence of DNA damage in the pancreas and liver of Fischer 344 and Wistar/Lewis rats after they were given single or multiple weekly injections of azaserine (AZ: 10 mg/kg), MNCO (N-methyl-N-nitrosocarbonyl-L-ornithine) (MNCO: 10 or 218 mg/kg/wk) or neutral red (96 or 288 mg/kg/wk). Results were compared with the carcinogenicity and mutagenicity of each compound. Damaged DNA was present in pancreas 1 hr after AZ administration, and damage persisted in both pancreas and liver for > 1 wk. Damage to pancreatic and hepatic DNA was more extensive 1 wk following 6 wk or 6 mo of weekly AZ injections than at 1 wk after a single injection. Damage to pancreatic DNA was detectable in rats 1 hr after either dose of MNCO. Pancreatic DNA damage was not apparent 1 wk after 72 mg/kg MNCO, but it was detectable 1 wk after 218 mg/kg. Significant pancreatic DNA damage was detected 1 wk after 6 wk or 6 mo of weekly injections of 72 mg/kg. Damage to hepatic DNA was detectable at either 72 or 218 mg/kg whether administered once or weekly in the above protocols. DNA damage was present in pancreas and liver 1 hr after injection of 96 mg/kg neutral red but not at 1 wk. Limited DNA damage was detectable in liver but not in pancreas 1 wk after a dose of 288 mg/kg. Persistent DNA damage was also detectable in liver but not in pancreas after six weekly injections of 96 mg/kg. The results indicate that the relative persistence of DNA damage may be an accurate indication of the organotropic carcinogenic potential of AZ, MNCO and neutral red administered at nontoxic dose levels. (17 refs)

78-6201 On Mutagens in Foods, Especially on Compounds from Pyrolysis of Protein (Meeting Abstract). (Eng) Nagao, M. (Natl. Cancer Center Res. Inst., Tokyo, Japan); Yahagi, T.; Kawachi, T.; Seino, Y.; Honda, M.; Matsukura, N.; Sugimura, T.; Wakabayashi, K.; Tsutsui, K.; Kosuge, T. *Mutat Res* 53(2): 239; 1978. (no refs)

78-6202 Characteristic Features of Tryptophan Metabolite-induced Hemoblastoses in Various Animals.

ciency. (Rus) Khrustalev, S. A. (Cancer Res. Center, Moscow, USSR). *Probl Gematol Pereliv Krovi* 23(9): 33-36; 1978.

The leukemogenic activity of 3-indolylacrylic acid (3-IAA) and 2-aminoacetophenone (2-AAP) was studied in Syrian hamsters, guinea pigs, and rats. Animals received sc injections of small doses of these tryptophan metabolites for 2-3 months. 3-IAA induced hemoblastoses in 10/60 hamsters, 4/71 guinea pigs, and 1/30 rats; 2-AAP induced hemoblastoses in 5/5 hamsters, 3/70 rats, and 0/20 guinea pigs (incidence of spontaneous hemoblastoses in hamsters, rats, and guinea pigs was 3/33, 2/60 and 1/40 respectively). (4 refs)

6203 **Significance of the Reaction of Nitrite with Tryptophan.** (Eng) Nakai, H. (Natl. Inst. Animal Industry, Chiba-Shi, 280, Japan); Cassens, R. G.; Greaser, M. L.; Woolford, G. *J Food Sci* 43(6): 1857-1860; 1978.

The reaction of sodium nitrite with tryptophan was studied spectrophotometrically to determine the significance of the reaction in cured meats. Although the tryptophan content of meat is relatively low, it is an essential amino acid and the indole nitrogen presents a secondary amine for reaction. Absorption at 330 nm indicated the formation of nitrosotryptophan. The reaction proceeded rapidly at pH 4.0, slowly at pH 5.0 and very sluggishly at pH 5.5 or higher. Experiments with dipeptides and acetyltryptophan indicated that nitrosation of the indole nitrogen was promoted if the primary amino group of tryptophan was blocked. It is concluded that, in view of the concentration of reactants and pH in cured meat, the environment for the nitrosation of tryptophan is favorable. (18 refs)

6204 **Mutagenic Properties of Nitrosated Spermidine.** (Eng) Kokatnur, M. G. (Dept. Pathology, Louisiana State Univ. Medical Center, New Orleans, LA, 70112); Murray, M. L.; Correa, P. *Proc Soc Exp Biol Med* 131(1): 85-88; 1978.

The nitrosation of spermidine resulted in direct-acting mutagens in the Ames assay using *Salmonella typhimurium* 1535. The mutagenic effect was catalyzed by thiocyanate and inhibited by ascorbic acid. The extent of formation of mutagenic compounds was dependent upon the pH of the medium, with an optimum at 4.0 and a range of 3.0-6.0. These results suggest that the nitrosated spermidine model fulfills many of the requirements of the hypothesis that gastric cancer and its precursor lesions may be induced by the intragastric formation of mutagenic/carcinogenic N-nitroso compounds. It yields a direct-acting mutagen; it has an optimum pH range very similar to that found in the gastric juice of human populations at high risk for gastric cancer, as a result of chronic atrophic gastritis; its mutagenicity may be enhanced by thiocyanate ions, which are present in human gastric juice, and spermidine is found in foods prominent in the diet of populations at high gastric cancer risk. (18 refs)

78-6205 **Effect of L-Ethionine on Macromolecular Synthesis in Mitogen-Stimulated Lymphocytes.**

(Eng) Zabos, P. (Dept. Pediatrics and Biochemistry, Mount Sinai Sch. Medicine City Univ. New York, New York, NY); Kyner, D. A.; Seide-Kehoe, R.; Acs, G.; Christman, J. K. *Biochim Biophys Acta* 520(1): 139-152; 1978.

The effects of L-ethionine (Eth) on macromolecular synthesis and modification in mitogen-stimulated human and mouse lymphocytes was studied. DNA synthesis in phytohemagglutinin- or concanavalin A-stimulated lymphocytes was completely inhibited by 2-4 mM Eth. Eth also inhibited the methylation of transfer RNA and prevented the mitogen-induced increase in the activity of histone-modifying enzymes, but it had little or no effect on commitment to DNA synthesis, rates of RNA and protein synthesis, size of the ATP and guanosine triphosphate pools, or the occurrence of morphological changes characteristic of blast formation. Thus, Eth permitted a quiescent lymphocyte to enter the G₁ stage of the cell cycle and to progress toward S, although it blocked DNA synthesis. These effects of Eth were completely reversible: removal of Eth after ≥ 24 hr of exposure resulted in the rate of RNA methylation, and an increase in the activity of histone-modifying enzymes. The results suggest that Eth may inhibit DNA synthesis by inhibiting a methylation reaction critical to initiation of that synthesis. (37 refs)

78-6206 **Some Improvements in the Mutagenicity Testing of Chemicals with *Bacillus subtilis*, *Salmonella typhimurium* and *Escherichia coli*; Results with 500 Chemicals (Meeting Abstract).** (Eng) Hirano, K. (Natl. Inst. Genetics, Misima, Sizuoka-ken, Japan); Narui, K.; Sadaie, Y.; Kada, T. *Mutat Res* 53(2): 200; 1978. (no refs)

78-6207 **Biochemical Studies on Rats with Insulin-secreting Islet Cell tumors Induced by Streptozotocin: With Special Reference to Physiological Response to Oral Glucose Load in the Course of and after Tumor Induction.** (Eng) Kazumi, T. (Second Dept. Internal Medicine, Kobe Univ. Sch. Medicine, Kobe, Japan); Yoshino, G.; Yoshida, Y.; Doi, K.; Yoshida, M.; Kaneko, S.; Baba, S. *Endocrinology* 103(5): 1541-1545; 1978.

Pancreatic islet cell tumors were induced in 32/49 male Wistar rats surviving 9 mo or longer after treatment with streptozotocin (S: 65 mg/kg iv) alone (Group 1), nicotinamide (500 mg/kg ip) + S (Group 2), or picolinamide (250 mg/kg ip) + S (Group 3). Of a total of 73 tumors, there were 14 in Group 1 rats, 47 in Group 2 rats, and 12 in Group 3 rats. In glucose (G) tolerance tests, elevation of blood G following intragastric G was depressed throughout the study in Group 2 rats, and it was significantly depressed at 7 mo. Conversely, mean plasma insulin (I) levels, which had remained unchanged up to 7 mo after treatment, rose abruptly 30 min

after G loading (44 to 167 μ U/ml) at 9 mo treatment. In tumor-bearing rats, G levels after loading were always significantly lower than those of tumor-free rats, but mean plasma I levels rose significantly from 47 to 132 μ U/ml within 30 min of G administration. Tumor-bearing rats tended to develop hypoglycemia when appearing moribund 13-26 mo after treatment. The mean I concentration in 11 islet cell tumors was 401 U/g wet wt, as opposed to the control mean of 14.0 U/g wet wt. These findings suggest that pancreatic islet cell tumors induced by streptozotocin alone or by combined treatment are insulin-secreting. (8 refs)

78-6208 Tumorigenic Action of Streptozotocin on the Pancreas and Kidney in Male Wistar Rats. (Eng) Kazumi, T. (Second Dept. Internal Medicine, Kobe Univ. Sch. Medicine, Kusunoki-cho 7-chome, Ikuta-ku, Kobe 650, Japan); Yoshino, G.; Fujii, S.; Baba, S. *Cancer Res* 38(7): 2144-2147; 1978.

The carcinogenic action of streptozotocin (STZ) was investigated in male Wistar rats. Pancreatic islet cell tumors were induced in 38 of 44 rats which survived 9-14 mo following various treatment schedules with STZ. Single iv injections of STZ alone at doses of 30, 40, 50, and 65 mg/kg produced adenomas of pancreatic islet cells in 8 of 9 (89%), 6 of 7 (86%), 2 of 4 (50%), and 1 of 2 (50%) rats, respectively. The neoplasms were seen in all of 8 rats given a single ip injection of picolinamide (250 mg/kg) 15 min before a single iv injection of STZ (65 mg/kg). Pancreatic islet cell tumors developed in 13 of 14 (95%) rats given a single ip injection of nicotinamide (500 mg/kg) 15 min before a single iv injection of STZ (65 mg/kg). Renal tumors were seen in only 3 rats treated with STZ and nicotinamide. None of the rats used in this study developed hepatic tumors. The results demonstrate that STZ, even in a dose of 30 mg/kg, has an exceedingly marked tumorigenic action on the rat pancreas, while it has little effect on the kidney and no effect on the liver. (19 refs)

78-6209 Oncogenic Effects of Antimitotic Chemotherapy. Observation of Four Cases. (Fre) Carcassonne, Y. (Clinique des maladies du sang, Hopital de la Conception, 136, rue Saint-Pierre, F 13385 Marseille Cedex 4, France); Blanc, A. P.; Gastaut, J. A.; Sebahoun, G. *Ann Med Interne (Paris)* 129(3): 181-186; 1978.

Four men developed cancer after undergoing treatment with chemotherapeutic agents. Three patients, ages 24, 53, and 64 yr, had a blood profile and symptoms indicative of acute myeloblastic leukemia after the use of chlorambucil, azathioprine, or 6-mercaptopurine for glomerulonephritis in two cases and immunosuppression prior to kidney transplant in one case. These three patients died of infection or hemorrhage. The fourth patient, age 52 yr, developed a lymphoma after 3 yr of treatment with chlorambucil for glomerulone-

phritis (total dose, 5,598 g). Combination chemotherapy and radiation have succeeded in controlling the malignancy. Possible mechanisms of action of the drugs are discussed. (no refs)

78-6210 Gastric Cancer in Rats after Chronic Intraperitoneal Application of Sap of Green Potatoes (*Solanum tuberosum* L.). (Eng) Ivankovic, M. (Inst. Toxicology and Chemotherapy, German Cancer Research Center, Im Neuenheimer Feld 280, D-69 Heidelberg, W. Germany). *Experientia* 34(5): 645; 1978.

Sap from the green tops of potatoes (*Solanum tuberosum* L.) was injected ip (0.25 ml) once weekly into 12 BD IX rats to determine its carcinogenicity. Two rats developed adenocarcinomas of the glandular stomach, and two developed papilloma and a squamous cell carcinoma of the forestomach. Control rats developed tumors. Chemical analysis (unpublished) revealed the presence of several volatile N-nitrosamines in the sap. (no refs)

78-6211 Acute Myelogenous Leukemia Following Chlorambucil Treatment. Report of Two Cases. (F) Fiere, D. (Service d'Hematologie, Hopital Edouard Herriot, 5 place d'Arsonval, F 69374 Lyon Cedex 2, France); Fellenz, P.; Vu Van, H.; Coiffier, B. *Nouv Presse Med* 7(9): 756; 1978.

Two 55-yr-old patients treated with chlorambucil for 2 and 4 yr for nephrotic syndrome and rheumatoid polyarthritis respectively, developed acute myelogenous leukemia after treatment periods of 3 and 3.5 yr. (1 ref)

78-6212 Inhibition of Deoxyribonucleic Acid Synthesis by Diisopropanolnitrosamine, N-Methylnitrosourea, Azaserine and 4-Hydroxyaminoquinoline 1-oxide after Partial Pancreatectomy in Rats. (Eng) Konishi, K. (Dept. Oncological Pathology, Cancer Center, Nara Medical Univ., Kashiwara, 634, Nara, Japan); Denda, A.; Kawabuchi, H.; Takahashi, S.; Sunagawa, M.; Kondo, H. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404-2405; 1978.

The effect of a single injection of dimethylnitrosamine (DMH: 100 mg/kg ip), ethionine (700 mg/kg ip), streptozotocin (65 mg/kg iv), diisopropanolnitrosamine (DIPN: 1,000 mg/kg ip), N-methylnitrosourea (MNUT: 1,000 mg/kg ip), azaserine (5 mg/kg ip), and 4-hydroxyaminoquinoline 1-oxide (4-HAQO: 7 mg/kg iv) on DNA

thesis of pancreatic acinar cells after partial pancreatectomy (64%) was studied in male Wistar rats. The carcinogens administered 3 days after the operation, when DNA synthesis was at its max level, and the animals were sacrificed later. DNA synthesis was not inhibited by DMN or aflatoxine, both nonpancreatic carcinogens. It was inhibited by DIPN (57.9%), MNUT (76.4%), azaserine (71.7%), and AAO (82.1%), all pancreatic nonendocrine carcinogens. The effect of streptozotocin, a pancreatic endocrine carcinogen on DNA synthesis was not clear. Dose-dependent inhibition of DNA synthesis was seen with azaserine and 4-HAQO but not with NMUT or DIPN. The inhibition of DNA synthesis by 4-HAQO lasted for at least 24 hr; it recovered to control values by 72 hr. Mechanisms involved in the inhibition of pancreatic DNA synthesis by these carcinogens require further study. (36 refs)

78-6213 Enzymatic Removal of O⁶-Methylguanine from DNA by Mammalian Cell Extracts. (Eng) Pegg, A. E. (Dept. Physiology, Milton S. Hershey Medical Center, Pennsylvania State Univ., Hershey, PA, 17033). *Biochem Biophys Res Commun* 84(1): 166-173; 1978.

Extracts from various rat tissues were incubated with methylated DNA or chromatin, and their abilities to catalyze the removal of labeled O⁶-methylguanine (OMG) from acid-precipitable DNA were compared. The four substrates were dimethylnitrosamine (DMN)-treated DNA and chromatin, isolated from the livers of rats inoculated with 50 mg/kg ³H-DMN (3.5 Ci/millimole) 20 min before death, and methyl-N-nitrosourea (NMU)-treated calf thymus DNA and NMU-treated rat liver chromatin, obtained by incubating the two substrates with 0.05 mM ³H-NMU (1.055 Ci/millimole) in vitro. In the first experiment, the substrates were incubated for 60 min at 37 C in the presence or absence of liver protein. The rat liver extracts were able to catalyze the loss of OMG from acid-precipitable DNA when incubated with all four substrates. The most convenient substrate was NMU-treated calf thymus DNA, because it was the easiest to prepare and the extracts showed max activity on this substrate. Prolonged incubation of the substrate with high concentrations of liver protein led to a loss of up to 50% OMG. In the second experiment, using extracts from different rat tissues and NMU-treated calf thymus DNA, liver extracts had the greatest activity. Kidney extracts had about 35% of the liver activity, and extracts from lung, colon, small intestine, and brain were much less active. The enzyme responsible did not appear to be an N-glycosidase, because labeled OMG could be detected in the supernatant, even though >50% of this base was lost from the DNA. The released radioactivity was methanol, which is consistent with the possibility that the reaction involves a demethylase action on the DNA substrate or on an oligonucleotide derived from it. (16 refs)

78-6214 Further Assessment of a Test for Determining the Mutagenicity of Compounds Metabolised In

Vivo in the Rat (Meeting Abstract). (Eng) Newton, M. F. (Dept. Biology as Applied to Medicine, Middlesex Hosp. Medical Sch., Cleveland St., London W1P 6DB, England); Lilly, L. J. *Mutat Res* 53(2): 241; 1978. (1 ref)

78-6215 Carcinogen Induced Alterations in DNAase I Release of N³-Methyladenine from Liver Chromatin. (Eng) Rao, P. M. (Dept. Pathology, Univ. Toronto, Medical Sciences Building, Toronto, Ontario M5S 1A8, Canada); Rajalakshmi, S.; Sarma, D. S. *Chem Biol Interact* 22(2/3): 347-351; 1978.

To determine whether carcinogens alter gene organization, Fischer rat liver chromatin DNA was labeled in vivo with ¹⁴C-dimethylnitrosamine to generate methylated bases such as N⁷-methylguanine, O⁶-methylguanine, and N³-methyladenine. The isolated chromatin was treated with pancreatic deoxyribonuclease I (DNase I) to digest 50% of the chromatin DNA, and the distribution of the methylated purines in the DNase I-digestible and nondigestible fractions was determined. Resistance to DNase I was attributed in part to the association of proteins with these regions of DNA. Accordingly, any alteration in the distribution of these dimethylnitrosamine-induced methylated bases in the chromatin fractions as a result of the subsequent administration of carcinogen (N-methyl-N-nitrosourea) was used as a parameter to indicate a change in DNA-protein association in the chromatin brought about by the carcinogen. The observed release of carcinogen-modified bases in the DNase I-digestible fractions of chromatin may represent an interaction of the carcinogen with regions of chromatin that are probably transcribable. The results also indicate that carcinogens can alter the DNA-protein association (gene organization) in such a way that DNA regions in chromatin accessible to nuclease digestion before carcinogen administration are rendered inaccessible subsequent to its administration. (18 refs)

78-6216 De Novo and Repair Replication of DNA in Liver of Carcinogen-treated Animals. (Eng) Craddock, V. M. (Toxicology Unit, Medical Res. Council Lab., Woodmansterne Road, Carshalton, Surrey, SM5 4EF, England); Henderson, A. R. *Cancer Res* 38(7): 2135-2143; 1978.

The effect of various carcinogens on de novo and repair replication of DNA in the liver of Wistar rats was investigated. Single ip injections of diethylnitrosamine (100, 200, or 250 mg/kg), ethyl methanesulfonate (300 or 500 mg/kg), aflatoxin (0.5 or 2.0 mg/kg), and retrorsine (160 mg/kg) induced DNA repair replication in vivo but did not alter de novo synthesis 2 hr after injection. Single ip injections of carbon tetrachloride (4,000 mg/kg) and ethionine (500 mg/kg) also induced repair replication but only after a delay period of 17 hr, suggesting that the repair was that of damage caused by an indirect mechanism, such as by deoxyribonuclease activity resulting from lysosomal damage (carbon tetrachloride) or

from nonenzymic reaction of DNA with a metabolite of the carcinogen that is slow to accumulate in the liver (S-adenosylmethionine after ethionine). A single ip injection of thioacetamide (150 mg/kg) did not cause detectable repair replication, a result that correlates with the lack of evidence for a reaction between thioacetamide and DNA. (54 refs)

78-6217 Hepatic Microsomal Metabolism of N,N-Dimethyl-nitrosamine To Its Methylating Intermediate. (Eng) Chin, A. E. (Univ. Rochester, Rochester, NY, 14627). *Diss Abstr Int B* 39(5): 2243; 1978. (no refs)

78-6218 Some Peculiarities in the Kinetics and Mechanism of Transformation of Paramagnetic Heme and Non-heme Iron Complexes in Carcinogenesis. (Eng) Saprin, A. N. (Dept. Kinetics Chemical and Biological Processes, Inst. Chemical Physics, Acad. Sciences USSR, Moscow, USSR). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1207-1218; 1978.

The kinetics of changes in cytochrome P-450 (CCP-450) levels in the Wistar rat liver during dimethylaminoazobenzene (DAB)- and diethylnitrosamine (DEN)-induced carcinogenesis was studied by electron spin resonance. During the precancerous stage, CCP-450 levels in the hyperplastic tissue increased to 2.5 times that of controls and then dropped at the time of tumor appearance and subsequent tumor growth. During the terminal stages of tumor growth, CCP-450 levels decreased fourfold compared with control levels. The free-radical concentration in microsomes increased more than four times during the precancerous stage, compared with normal liver. This increase was observed prior to the decrease in CCP-450 levels, which suggests that the free radicals may be responsible for the decrease. Liver tissue from rats fed both DAB and butylated hydroxytoluene (BHT), a nontoxic inhibitor of free-radical processes, showed no significant change in free-radical concentrations from those in normal tissue, and no tumors developed after 1 yr. However, DAB + BHT increased CCP-450 levels at times earlier than the increase in free-radical concentration. This seems to indicate that the anticarcinogenic properties of BHT are due to its induction of high levels of CCP-450 in rat liver, which promotes the detoxification of DAB, rather than to its inhibition of free-radical formation. (35 refs)

78-6219 Hematologic Disorder after MOPP Treatment of Hodgkin's Disease. (Eng) Moloney, W. C. (Massachusetts General Hosp., Boston, MA); Long, J. C.; Ferrucci, J. T.; Groopman, J. E.; Aisenberg, A. C.;

Galbadini, J. J.; Dermirjian, Z. N.; Hirsch, M. S.; Man, R. M.; Stossel, T. P. *N Engl J Med* 299(17): 946; 1978.

The case report of a man with Hodgkin's disease (HD: cellularity type) who, 6 yr later at age 35, developed a myelogenous or erythromyelocytic leukemia with terminal pneumocystis or fungal pneumonia is described. In addition, the patient had viral hepatitis, repeated episodes of bacterial pneumonia, osteomyelitis, cytomegalovirus infection, and hypersensitivity syndrome. After the diagnosis of HD, 4 courses of MOPP [nitrogen mustard, vincristine (Oncovin), procarbazine, and prednisone] were administered over 18 mo. The patient improved and was well and active during the next 4 yr. Throughout the final illness there was no evidence of recurrent HD. The repeated bacterial and viral infections indicated a lack of immunologic defenses. Skin tests for purified protein derivative of tuberculin, streptokinase-streptolysin, and monilia were negative. Whether the splenectomy after the first diagnosis and the MOPP therapy contributed to the lowering of host defenses is a matter of speculation. The leukemia may be related to an immunologic defect caused by the therapy or to one of many other factors, such as injury to lymphoid tissue by the immunosuppressants, or perhaps with activation of a leukemogenic virus. (10 refs)

78-6220 Dominant Lethal Effects of Triethylenemelamine in the Guppy *Poecilia reticulata*. (Eng) Mathews, J. G. (Sch. Biology, Georgia Inst. Technology, Atlanta, GA, 30332); Favor, J. B.; Crenshaw, J. W. *Mutat Res* 54(2): 149-157; 1978.

Triethylenemelamine [2,4,6-tris(aziridinyl)-s-triazine, TEM] was selected to test the potential of an experimental dye for detecting dominant lethal events in the common guppy, *Poecilia reticulata*. Dominant lethal effects were induced by ip injection of males with 0, 0.1, 0.2, or 0.4 mg/kg TEM dissolved in Fish Ringer's soln or by immersion of males in the same solns used for the injections for 24 hr. All treated males were subsequently mated to virgin females 1 day after treatment, and the females were removed 24 hr later. 7 days following the establishment of matings, the females were sacrificed and dissected, and live and dead embryos were counted. The percentage of dead embryos increased and the total number of embryos decreased with TEM dose. The injection method had a more deleterious effect than the immersion method, but the reason(s) for this finding are not clear. The results are compared with those of *Drosophila* and mammalian studies. (17 refs)

78-6221 DNA Alkylation and Neuro-Oncogenesis by 3,3-Dimethyl-1-phenyltriazene. (Eng) Coccaro, H. K. (Max-Planck-Institut für Hirnforschung, D-5000 L, 91, W. Germany); Hauenstein, E.; Kolar, G. F.; Kleihues, P. *Acta Neuropathol (Berl)* 43(1-2): 105-109; 1978.

The role of DNA alkylation in the initiation of malignant transformation by 3,3-dimethyl-1-phenyltriazene (DMPT) was studied using BD-IX rats. Following a single sc injection

¹⁴C-DMPT (100 mg/kg) into pregnant females on gestation day 21, the concentrations of 7-methylguanine and O⁶-methylguanine in the brains and livers of the fetuses were similar. Within 2 days after birth, however, the liver DNA values were significantly higher than in the brain values, and this trend continued during postnatal growth. In all organs investigated, the concentrations of 7-methylguanine in the DNA of adult mice 12 hr after the last of five weekly injections of ¹⁴C-DMPT (50 mg/kg each) were higher than those observed 12 hr after a single injection (50 mg/kg). This effect was most pronounced in the ileum, brain, and lung. The O⁶-methylguanine concentrations following multiple doses of ¹⁴C-DMPT were increased only in brain and lung DNA relative to the concentrations observed after a single dose. The data suggest that the efficient excision repair capacity of the CNS is a significant factor in the organ-specific carcinogenicity of DMPT and related carcinogens. (24 refs)

78-6222 **Expression of TEM-induced Damage to Postmeiotic Stages of Spermatogenesis of the Mouse during Early Embryogenesis. II. Cytological Investigations.** (Eng) Burki, K. (Swiss Inst. Experimental Cancer Res., CH-1066 Epalinges, Switzerland); Sheridan, W. *Mutat Res* (1): 107-115; 1978.

Analyses were made of triethylenemelamine (TEM)-induced chromosome damage to postmeiotic male germ cells during early mouse embryogenesis. Male DBA/2J mice treated with 2 or 0.4 mg/kg TEM were mated with CD-1/CR females. For the analysis of first-cleavage metaphases, pregnant females were treated with 2.5 mg/kg colchicine on day 1 of pregnancy, and the eggs flushed from their oviducts approximately 1 hr later. In addition, embryos were flushed from the oviducts (day 2), junction (day 3), or uteri (day 4) of pregnant females without colchicine pretreatment. The frequencies of first-cleavage metaphases exhibiting structural aberrations (all of the chromosome type) and the frequencies of two-cell embryos and of four- to eight-cell embryos containing nuclei accompanied by micronuclei or nuclei connected by bridges were closely correlated to frequencies of preimplantation loss of embryos recorded in a dominant lethal test. The frequencies of morulae/blastulae containing blastomeres with micronuclei showed a close correlation to the frequencies of total (pre- and postimplantation) loss of embryos. This indicates that the expression of TEM-induced chromosomal damage is delayed and, thus, the damage could persist undetected in very early stages of embryogenesis. (12 refs)

78-6223 **Mutagenicity of Arylalkyl- and Arylacetyl-triazenes in Respect to Their Reaction Mechanisms (Meeting Abstract).** (Eng) Pool, B. L. (Inst. Experimental Toxicology and Chemotherapy, German Cancer Res. Center, Heidelberg, W. Germany). *Mutat Res* 53(2): 249; 1978. (no refs)

78-6224 **Carcinogenicity of Triethanolamine in Mice and Its Mutagenicity after Reaction with Sodium Nitrite in Bacteria.** (Eng) Hoshino, H. (Radiobiology Div., Natl. Cancer Center Res. Inst., Tsukiji, Chuo-ku, Tokyo 104, Japan); Tanooka, H. *Cancer Res* 38(11, part 1): 3918-3921; 1978.

The carcinogenicity of triethanolamine (TEA) was examined in male and female ICR-JCL mice fed a diet containing 0.3% or 0.03% (wt/wt) TEA from the age of 6 wk to death or sacrifice. The incidence of various tumors was significantly higher in TEA-fed mice than in controls. Female mice showed a high incidence of lymphoid tissue tumors (22%), but these tumors were absent in males. Other types of malignant tumors found were independent of sex. Mice fed the TEA-containing diet survived as long as the controls; the survival rate was 50% at 85 wk in females and at 65 wk in males. TEA was not mutagenic to *Bacillus subtilis*, with or without liver microsomes. When TEA was incubated with sodium nitrite under acidic conditions or when the mixture was heated, a direct-acting mutagen was produced, but its mutagenic activity was destroyed by rat liver microsomes. N-Nitrosodiethanolamine (NDEA) was detected in the reaction mixture by thin-layer chromatography, but it may not be the main mutagenic product, because this product was a stable and direct mutagen and its mutagenic activity was destroyed by liver enzymes, whereas NDEA is unstable and does not induce mutation in *B. subtilis* by itself. The chemical nature of the mutagenic product in the TEA:sodium nitrite mixture has not been identified. (11 refs)

78-6225 **Analytical Investigations of Hexamethylenetetramine Samples for the Presence of Volatile Organic Compounds and of In Vitro Nitrosation Metabolites (Nitrosamines).** (Ger) Roper, H. (Institut für Organische Chemie und Biochemie, Universität Hamburg, Martin-Luther-King-Platz 6, D-2000 Hamburg 13, W. Germany); Heyns, K. *Z Lebensm Unters Forsch* 167(3): 145-149; 1978.

Analysis of four hexamethylenetetramine (HMT) samples by gas chromatography/mass spectrometry revealed the presence of formaldehyde, methylamine, chloroform, and trimethylamine N-oxide, but no secondary or tertiary aliphatic amines. No dimethylnitrosamine was formed in in vitro nitrosation reactions with HMT + sodium nitrate in acidic solution. In view of these findings, it is proposed that HMT be reconsidered for use as a food additive. (36 refs)

78-6226 **Combined Action of Alkylating Compounds on Human Chromosomes (Meeting Abstract).** (Eng) Bochkov, N. P. (Inst. Medical Genetics, USSR Acad. Medical Science, Moscow, USSR); Yakovenko, K. N.; Nazarenko, S. A. *Mutat Res* 53(2): 152-153; 1978. (no refs)

- 78-6227 **Metabolic Activation of Nitrofurans by Testis Homogenate from Mammals (Meeting Abstract).** (Eng) Maier, P. (Environmental Mutagenesis Branch, Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC); Chhabra, R. S. *Mutat Res* 53(2): 226; 1978. (1 ref)

- 78-6228 **Direct Study of Alkylating Agent-RNA Interaction by ^{13}C Nuclear Magnetic Resonance Spectroscopy.** (Eng) Chang, C. (Dept. Medicinal Chemistry and Pharmacognosy, Sch. Pharmacy and Pharmacal Sciences, Purdue Univ., W. Lafayette, IN, 47907); Lee, C. G. *Cancer Res* 38(11, part 1): 3734-3736; 1978.

The interactions of four methylating agents and *Torula* yeast RNA were studied by ^{13}C nuclear magnetic resonance (NMR) spectroscopy, a direct method that eliminates all processes that may cause decomposition and structural alteration. In addition, spectral interference is reduced, product distribution can be determined at an early alkylation stage, and the amount of alkylating agent can be reduced. Methylation reactions were carried out at room temperature and pH 7 by treating an 8.5-ml aqueous soln of 500 mg lyophilized RNA with 1.5 millimoles of methyl methanesulfonate (MMS: 130 μl), dimethyl sulfate (DMS: 142 μl), or 1-methyl-1-nitrosourea (MNNG: 221 mg) for 12 hr or with 1-methyl-3-nitro-1-nitrosoguanidine (MNNG: 221 mg) for 259 hr. Each reaction mixture was then dialyzed and lyophilized repeatedly, and NMR samples were prepared. Based on integration curves, the agents could be classified into two groups according to biological potency: (1) weakly mutagenic/carcinogenic, MMS and DMS; (2) strongly mutagenic/carcinogenic, MNU and MNNG. In conclusion, from the chemical shift analyses and the spectral integration curves, the reactive sites, the product distributions and specificity of the different methylating agents, and the degree of N-7 methylation of guanosine could be determined. The results show the potential application of ^{13}C NMR spectroscopy in establishing the mechanisms of chemical carcinogenesis and mutagenesis. (32 refs)

- 78-6229 **N-Methyl-N-nitrosourea Mediated Conversion of O⁶-Methylguanine to Xanthine.** (Eng) Rao, P. M. (Dept. Pathology, Univ. Toronto, Medical Sciences Building, Toronto, Ontario M5S 1A8, Canada); Rajalakshmi, S.; Serma, D. S.; Krepinsky, J. *Chem Biol Interact* 22(2/3): 341-346; 1978.

The in vitro alkylation by N-methyl-N-nitrosourea (MNU) of rat liver chromatin DNA prelabeled in vivo with ^{14}C -dimethylnitrosamine (^{14}C -DMN) was studied. The standard reac-

tion condition was to incubate 4.24 micromoles (μmol) of (^{14}C) methylguanine [O^6 -(^{14}C)meG] in 0.1 N HCl at 70 C 45 min or at 37 C for 16 hr. In all cases, demethylation of O^6 -meG yielded guanine. However, when treated with MNU (29.1 μmol) and 0.1 N HCl, O^6 -meG was converted mostly to xanthine. Observations that guanine is not deaminated by MNU and 0.1 N HCl as effectively as O^6 -meG and that guanine is detected in the formation of xanthine from meG suggest that the deamination of O^6 -meG precedes the demethylation step in its conversion to xanthine. It is suggested that if the conversion of O^6 -meG to xanthine occurs in vivo at the macromolecular level, the mutagenicity of N-nitroso-N-alkylamides can be due to not only the formation of a miscoding lesion of O^6 -meG, but also to its conversion to xanthine. (9 refs)

- 78-6230 **C-Type Virus Particles in Cell Cultures of Chemically Induced Glioma in Sprague-Dawley Rats.** (Rus) Anzil, A. P. (Electron Microscopy Lab., Max Planck Inst. Psychiatry, Munich, W. Germany); Stavrou, A.; Blinzinger, K. *Vopr Onkol* 24(9): 30-32; 1978.

Long-term cultures were prepared from mixed anaplastic gliomas induced by N-methyl-N-nitrosourea (6 mg/kg/wk for at least 10-12 mo) in Sprague-Dawley rats. Electron microscopy revealed budding and large numbers of cytoplasmic and extracellular C-type virus particles in many cultures. This is the first report of a chemically induced glioma associated with C-type virus particles in rats. The characteristics of this rat glioma are being studied. (10 refs)

- 78-6231 **Surface Structure of Fetal Rat Brain Cells During Neoplastic Transformation in Cell Culture.** (Eng) Haugen, A. (Dept. Pathology, Gade Inst., Univ. Bergen, N-5016 Haukeland Hosp., Bergen, Norway); Laerum, O. D. *J Natl Cancer Inst* 61(6): 1415-1422; 1978.

The surface microstructure of fetal BD IX rat brain cells undergoing neoplastic transformation in long-term cell culture after they were exposed to a single transplacental pulse of N-ethyl-N-nitrosourea (ENU; 75 $\mu\text{g/g}$) on gestation day 18 was investigated by scanning electron microscopy. The fetal brain cells (FBC) were transferred into monolayer culture 24 hr after exposure and maintained in culture for up to 1.5 yr. After 3 wk, ENU-pretreated FBC showed focal proliferation of neural cells on an underlayer of flat, epithelioid cells. The neural cells exhibited varying forms of numerous dorsal ruffles and an increased number of other surface microprojections. Between the 40th and 100th day, nodules of bipolar and multipolar neural cells with a complex surface microstructure, including many blebs and ruffles and an increased number of microvilli, were observed. After 100 days, more rapidly proliferating, morphologically altered cells

ned piled-up foci, which resulted in a homogeneous population of cells with numerous long microvilli, large ruffles, blebs over the whole surface. Attempts to establish permanent cell lines were successful in 4/13 trials. Three of these lines were tumorigenic upon retransplantation into baby at around 273 days. The surface structure of cells from solid tumors was the same as that seen in vitro, exhibiting numerous microvilli and ruffling membranes over the whole surface. Thus, surface alterations characteristic of neoplastic cells were observable >100 days before the cells became tumorigenic. (29 refs)

6232 **Mutagenicity of Several Classes of Antitumor Agents to *Salmonella typhimurium* TA98, TA100, and TA92.** (Eng) Seino, Y. (Biochemistry Div., Natl. Cancer Center Res. Inst., Tsukiji, Chuo-ku, Tokyo 104, Japan); Nagao, M.; Yahagi, T.; Hoshi, A.; Kawachi, T.; Shimura, T. *Cancer Res* 38(7): 2148-2156; 1978.

The mutagenic activities of various antitumor agents which included 19 antimetabolites, 5 antibiotics, 5 alkylating agents, 1 alkaloid, 1 enzyme, and 1 adrenal steroid hormone were tested on *Salmonella typhimurium* strains TA100, TA98, and TA92. Four of the anticancer agents, busulfan, carbazilquinone, 1-(4-amino-2-methylpyrimidine-5-yl)methyl-3-(2-chloroethyl)-3-nitrosourea hydrochloride, and pipobroman were found to be mutagenic for the first time. The known mutagenicities of five others, daunomycin hydrochloride, mitomycin C, 6-mercaptopurine, and cyclophosphamide were confirmed. Since there is a demonstrated correlation between mutagenicity in this bacterial test system and the carcinogenic potential in laboratory animals, the development of new chemotherapeutic agents that have no mutagenicity in such bacterial test systems is desirable from the standpoint of possibly decreasing the risk of latent drug-induced cancer in patients undergoing chemotherapy. (33 refs)

6233 **Alloxan Induced Leukemoid Condition in a Fresh Water Teleost *Clarias batrachus*** (Letter to Editor). (Eng) Agrawal, P. (Dept. Zoology, M.M./P.G. College, Modinagar 201 204, District Ghaziabad, Uttar Pradesh, India); Goel, K. A. *Curr Sci (Bangalore)* 47(15): 555-560; 1978.

The possible leukemogenic action of alloxan was demonstrated in 42 fish (*Clarias batrachus*) inoculated im with 50 mg/kg of the compound every 3 days. The number of WBC increased after 10-40 days, and after 40 days, 50% of the total WBC were immature. Mitosis was also observed in some WBC at this time. (2 refs)

6234 **Mutagenicity of Nitro Compounds in *Drosophila melanogaster*** (Meeting Abstract). (Eng)

Kramers, P. G. (Dept. Radiation Genetics and Chemical Mutagenesis, State Univ. Leiden, Wassenaarseweg 72, Leiden, Netherlands). *Mutat Res* 53(2): 213; 1978. (no refs)

78-6235 **Flagyl and Radiation-induced Chromosome Breakage (Letter to Editor).** (Eng) Prosser, J. S. (Natl. Radiological Protection Board, Harwell, Didcot, Oxon OOX11 0RQ, England); White, C. M. *Br J Radiol* 51(608): 654-655; 1978.

Preliminary observations of the effect of Flagyl (metronidazole) on the yield of x-ray-induced dicentric aberrations in human lymphocytes from anoxic and normal venous blood are presented. Flagyl increased the dicentric yield from irradiated anoxic blood significantly, but the dicentric levels were not as high as those observed in normal venous blood. The dicentric yield from blood at venous oxygen tension was not changed. (7 refs)

78-6236 **The Mutagenic Screening of Fourteen Imidazo Compounds Using a Modified Ames' Test.** (Eng) Turner, J. V. (Dept. Tissue Pathology, Inst. Medical and Veterinary Science, P.O. Box 14, Rundle St., Adelaide, S. Australia, 5000, Australia); Ward, A. D.; Freeman, C. G. *Mutat Res* 57(2): 135-139; 1978.

Eight imidazo[1,2-a]pyrimidine derivatives and six cyclic guanidine derivatives were synthesized and tested for mutagenic activity by a modified Ames *Salmonella typhimurium* assay using strains TA100, TA98, and TA1537, with or without microsomal activation. No mutagenicity or inhibition of growth was observed with any test compound. (6 refs)

78-6237 **High Pressure Liquid Chromatographic Assay for α Hydroxylation of N-Nitrosopyrrolidine by Isolated Rat Liver Microsomes.** (Eng) Chen, C. B. (Div. Environmental Carcinogenesis, Naylor Dana Inst. Disease Prevention, American Health Foundation, Valhalla, NY, 10595); McCoy, G. D.; Hecht, S. S.; Hoffmann, D.; Wynder, E. L. *Cancer Res* 38(11, part 1): 3812-3816; 1978.

A high-pressure liquid chromatographic assay for the α hydroxylation of N-nitrosopyrrolidine (NPY) by isolated hepatic microsomes from Aroclor-induced or noninduced male F-344 rats is described. This sensitive assay may allow determination of the role of hydroxylation in NPY carcinogenesis. Mixtures consisting of NPY, rat liver microsomes, and an NADPH-generating system were incubated at 37 C. The major α hydroxylation product, 2-hydroxytetrahydrofuran, was trapped by the addition of 2,4-dinitrophenylhydrazine to form 4-hydroxybutyraldehyde-2,4-dinitrophenylhydrazone, which was then quantitated by reverse-phase high-pressure liquid chromatography. Under optimal conditions, the α hy-

droxylation of NPY was linear for at least 90 min and had characteristics typical of the microsomal mixed-function oxidase system, such as inhibition by carbon monoxide and induction by pretreatment of the rats with Aroclor (500 mg/kg). Aroclor induction resulted in a greater than twofold increase in the α hydroxylation rate compared with noninduction. There was a close correspondence between the binding affinity and the α hydroxylation of NPY by the microsomes. This is the first available method for in vitro studies of the α hydroxylation of a cyclic nitrosamine. (28 refs)

78-6238 The Effect of the Microsomal Enzyme Inhibitor, Piperonyl Butoxide, on Mutagenesis in *Drosophila melanogaster* (Meeting Abstract). (Eng) Lindsay, R. J. (Sch. Biological Sciences, Flinders Univ. South Australia, Bedford Park, S.A. 5042 Australia); Clark, A. M. *Mutat Res* 53(2): 221; 1978. (no refs)

78-6239 Structural Requirements for Mutagenic Activities of N-Heterocyclic Bases in the Salmonella Test System. (Eng) Matsumoto, T. (Central Res. Inst., Japan Tobacco and Salt Public Corp., 6-2, Umeokaoka, Midoriku, Yokohama, Kanagawa 227, Japan); Yoshida, D.; Mizusaki, S.; Tomita, H.; Koshimizu, K. *Agric Biol Chem* 42(4): 861-864; 1978.

The effects of structural changes on the mutagenic activities of quinoline (Q), isoquinoline (IQ), phenanthridine (PA), benzo(f)quinoline [B(f)Q], benzo(h)quinoline [B(h)Q], and their α -amino derivatives were studied in the *Salmonella typhimurium* test system. All mutagenic compounds tested required the liver microsomal fraction (S-9 mix) in order to express their mutagenic activity. PA, B(f)Q, and Q were mutagenic to strains TA98 and TA100, B(h)Q was mutagenic only to TA100, and IQ was nonmutagenic to both strains. The mutagenic activities of Q and the BQ's increased greatly after amination in the α -position of the pyridine moiety, but amino-IQ was nonmutagenic to TA98 and TA100. The addition of a benzene ring to 2-amino-Q markedly increased its mutagenic activity, whereas the addition of a benzene ring to the pyridine moiety of 2-amino-Q only slightly increased mutagenic activity. The mutagenic activity of 2-amino-B(f)Q, but not those of 2-amino-B(h)Q and 9-amino-PA, was markedly stimulated by B(f)Q, B(h)Q, and PA in the presence of S-9 mix. Norharman enhanced the effect of 2-amino-B(f)Q in the presence of S-9 mix. The results indicate that among the α -amino heterocyclic bases, a three-ring system is more mutagenic than a two-ring system and the fused position of the three-ring system affects mutagenic activity and strain specificity. (15 refs)

78-6240 Carcinogenicity of 4-Chloronitrosopiperidine in Sprague-Dawley Rats. (Eng) Lijinsky, W.

(Chemical Carcinogenesis Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501); Taylor, H. W. *Z Krebsforsch* 92(2): 217-220; 1978.

The carcinogenicity of 4-chloronitrosopiperidine was tested in 15 Sprague-Dawley rats who were given the compound in their drinking water (0.005% soln) for 27 wk (total dose, 1.5 mg). Seven animals died by 41 wk, all 15 by 58 wk. Two rats had tumors: 7 had liver tumors, 5 had esophageal tumors, 2 tumors of the nonglandular stomach, and 2 tumors of the nasal turbinates. Several rats had tumors of more than one organ. The mortality pattern of these rats was almost identical with that of rats treated with a considerably higher dose (440 mg) of nitrosopiperidine. It is concluded that the addition of a single chlorine atom to the nitrosopiperidine molecule changes the chemistry so that the compound is more active in producing liver neoplasms. (5 refs)

78-6241 Formation and Biological Effect of N-Nitrosopiperazine Compound from Piperazine Pesticide Triflorine. (Eng) Borzsonyi, M. (Dept. Morphology, Natl. Inst. Public Health, Budapest, Hungary); Pinter, A.; Torok, G.; Surjan, A.; Nadasdi, L. *IARC Sci Publ* (19): 477-484; 1978.

In vitro experiments were conducted to determine the formation of the carcinogen dinitrosopiperazine (DNPZ) from the fungicide Triflorine (TF: N,N'-piperazinidylbis-2,2,2-trichloroethylidenebisformamide), and in vivo experiments were conducted in Swiss/Leiden mice to determine tumor incidence and morphology after intragastric administration of TF (300 mg/kg 2 x/wk). One hundred of the mice all received 0.05% sodium nitrite in their drinking water (Group 1), and 50 were treated with TF alone (Group 2). Tumor incidence was highest in Group 1 (104 tumors), and most of the tumors were lymphomas. The proliferating tumor cells corresponded to lymphoblasts. Ten Group 1 mice had stomach tumors, four gastric papillomas and six invasive squamous cell carcinomas. Prior to day 180, however, there were no morphological alterations in the glandular mucosa of the stomach. Most of the intestinal tumors that developed were in the duodenum and jejunum, and tumors appeared to develop from surface foci of atypical epithelial cells. Tumor incidence was similar in Group 2 animals, animals treated with sodium nitrite only, and untreated controls, and it was much less than that of Group 1 (25 tumors/401 mice). The results indicate that various tumors can be induced by DNPZ formed in vivo from TF. In the in vitro nitrosation of TF, 100 g of TF was dissolved in 270 ml 6 N HCl and 47.6 g of sodium nitrite dissolved in 270 ml distilled water was added slowly. After 24 hr at 4 C, 15.8 g of 1,4-dinitropiperazine was formed. (8 refs)

78-6242 Chemical Carcinogen Alteration of SV40 Virus Induced Transformation of Normal Human Cell Populations In Vitro. (Eng) Milo, G. E. (Dept. Physiologic

Chemistry, Coll. Medicine, Ohio State Univ., 1900 Coffey Road, Columbus, OH, 43210); Blakeslee, J. R.; Hart, R.; and others. *Chem Biol Interact* 22(2/3): 185-197; 1978.

The in vitro interactions of chemical carcinogens with a transforming DNA virus in human cells were determined to an index of the cocarcinogenicity of these agents. The frequency of simian papovavirus 40 (SV40)-induced transformation of human foreskin fibroblasts or neonatal foreskin cells was enhanced after pretreatment of the cells with naphthylacetamide (NA-2), N-methyl-N'-nitrosoguanidine (MNNG), acetyl-2-fluorenylacetylamide (N-Ac-AAF), benzo(a)pyrene (BP), aflatoxin B₁ (AFB₁), propane sultone, β -propiolactone (β PL), 4-nitroquinoline oxide (4-NQO), methylmethanesulfonate (MMS), or diethylnitrosamine (DEN). Pretreatment with 4-NQO, MMS, MNNG, or DEN inhibited transformation, but posttreatment with AFB₁, β -PL, or NA-2 did not alter transformation, similar to the action of N-Ac-AAF and BP. All carcinogens that altered transformation after pretreatment damaged cellular DNA. Pretreatment with posttreatment with 3-methylcholanthrene (3-MC) or 1,2-dimethylbenz(a)anthracene (7,12-DMBA) did not damage cellular DNA and did not enhance transformation. Moreover, pre- or posttreatment with other weak or noncarcinogens that did not damage cellular DNA also did not alter transformation-induced transformation. All foci formed in the cocarcinogen-treated cultures, whether the carcinogen inhibited or enhanced transformation, were virus-directed. Although a similar pattern of response existed for carcinogens that either enhanced or inhibit transformation, each of the carcinogens that enhanced or inhibited foci formation damaged cellular DNA. Moreover, those carcinogens that enhanced focus formation, compared with the carcinogens that inhibited focus formation, exhibited similar DNA damage profiles. (35 refs)

78-6243 Breast Cancer at a Psychiatric Hospital Before and after the Introduction of Neuroleptic Agents. (Eng) Wagner, S. (Office Medical Director, Norristown State Hosp., Norristown, PA, 19401); Mantel, N. *Cancer Res* 38(9): 2703-2708; 1978.

The possible association between breast cancer and treatment with neuroleptic agents was investigated in 5,463 women who had or underwent breast biopsy at the Norristown State Hospital (in Pennsylvania) between 1940 and 1974. The method of proportionality was used to evaluate the breast cancer incidence. Cases of breast cancer were related to deaths from other cancers and to deaths from all causes other than breast cancer. Deaths and cancers that occurred within 1 yr of hospital admission were excluded from the analysis. To allow for a 10-yr transition period and for latency in the development of breast cancer, comparison was made between the 15 yr preceding the introduction of neuroleptic agents into medical practice (1940-1954) and the second decade after their introduction (1965-1974). Although neuroleptic agents were used extensively to treat the patients at the hospital, there was no statistically or medically significant change in the indicated

incidence of breast cancer at the hospital during the neuroleptic period. The study did not rule out the possibility that neuroleptic agents might have some effect on breast cancer after a very long latency period. (no refs)

78-6244 Investigation on the Tester Strain of *Saccharomyces cerevisiae* for Primary Screening of Environmental Mutagens (Meeting Abstract). (Eng) (Div. Genetics, Natl. Inst. Radiological Sciences, Chiba, Japan); Machida, I.; Saeki, T.; Yoshikawa, K. *Mutat Res* 53(2): 240; 1978. (no refs)

78-6245 Neoplastic Transformation of Human Diploid Cells In Vitro after Chemical Carcinogen Treatment. (Eng) Milo, G. E. (Dept. Physiological Chemistry and Veterinary Pathobiology, Ohio State Univ., 1900 Coffey Road, Columbus, OH, 43210); DiPaolo, J. A. *Nature* 275(5676): 130-132; 1978.

The in vitro neoplastic transformation of primary or low-passage cells from 10 human foreskin cell populations by the carcinogens propane sultone (PS: 5 μ g/ml), β -propiolactone (PL: 13 μ g/ml), aflatoxin B₁ (AFB₁: 10 μ g/ml), N-methyl-N'-nitro-N-nitrosoguanidine (MNNG: 0.1 μ g/ml), 4-nitroquinoline 1-oxide (4-NQO: 0.002 μ g/ml), and ethyl methane sulfonate (EMS: 10.0 μ g/ml) is reported. To obtain transformation, the normal cells were blocked in the G₁/S interphase, released, and then treated with carcinogen during the S phase. Removal of the carcinogen 12 hr later coincided with the end of the S phase. After treatment, the proliferation of the unaffected cell population was purposely inhibited. Posttreatment population doubling No. 20 was found to be suitable for determining the ability of the transformed cells to grow in 0.5% soft agar. Cells transformed by PS, PL, and AFB₁ had 100% colony-forming ability in soft agar, whereas only 25%-70% of those transformed by 4-NQO, EMS, and MNNG formed colonies. A close correlation between growth in soft agar and tumor formation in nude mice was obtained. Injection of cells into the subcapsular area of the mice produced sc tumors 18-25 days later. The tumors were composed of multiple vacuolated cell types having predominantly a syncytial network with an eosinophilic cytoplasm and ovoid basophilic nuclei. The tumors were classified as undifferentiated mesenchymal tumors. To date, tumors have been serially subpassaged three times. (15 refs)

78-6246 Rat Liver Microsome-Mediated Binding to DNA of 3-Amino-1-methyl-5H-pyrido[4,3-b]indole, a Potent Mutagen Isolated from Tryptophan Pyrolysate. (Eng) Hashimoto, Y. (Faculty Pharmaceutical Sciences, Univ. Tokyo, Bunkyo-ku, Tokyo, Japan); Takeda, K.; Shudo, K.; Okamoto, T.; Sugimura, T.; Kosuge, T. *Chem Biol Interact* 23(1): 137-140; 1978.

The binding of 3-amino-1-methyl-5H-pyrido(4,3-b)indole (Trp-P-2) to calf thymus DNA was measured after incubation with liver microsomes from male Wistar rats treated with polychlorinated biphenyl (500 mg/kg, ip). The radioactivity from ^3H -Trp-P-2 bound tightly to DNA in the presence of rat liver microsomes. The binding was not affected by reprecipitation, dialysis, or gel chromatography. The amount of Trp-P-2 bound to DNA was about 60 $\mu\text{mol/mol}$ P, which increased to 230 $\mu\text{moles/mole}$ P after three successive incubations of the DNA with Trp-P-2 and microsomes. Trp-P-2 bound significantly to poly G in the presence of microsomes, but it did not bind to poly A, poly C, or poly U in the presence or absence of microsomes. After acid hydrolysis, DNA with bound Trp-P-2 gave more than two fluorescent spots on thin layer chromatography. (8 refs)

78-6247 Guanyl O⁶-Arylation and O⁶-Arylation of DNA by the Carcinogen N-Hydroxy-1-naphthylamine. (Eng) Kadlubar, F. F. (Natl. Center Toxicological Res., Jefferson, AR, 72079); Miller, J. A.; Miller, E. C. *Cancer Res* 38(11, part 1): 3628-3638; 1978.

Two O⁶-substituted deoxyguanosine adducts formed by the reaction of N-hydroxy-1-naphthylamine (N-HO-1-NA), a carcinogen for liver, urinary bladder, and other tissues of humans and animals, with DNA at pH 5 were isolated and characterized. N-HO-1-NA reacted with nucleic acids and protein at pH 5 to form covalently bound derivatives with 3-20 naphthyl residues/1,000 monomer units. The level of binding was in the following order: DNA > polyguanylic acid > denatured DNA and ribosomal RNA > serum albumin > transfer RNA > polyadenylic acid. The carcinogen did not react with nucleosides and nucleotides. Binding of N-HO-1-NA to DNA was not inhibited by the addition of nucleosides, nucleotides, methionine, or glutathione. Enzymatic hydrolysis of the DNA-containing naphthyl residues yielded three nucleoside arylamine adducts, N-(deoxyguanosin-O⁶-yl)-1-naphthylamine, 2-(deoxyguanosin-O⁶-yl)-1-naphthylamine, and the decomposition product of the latter. Direct evidence for acid-dependent arylnitrenium ion formation was obtained by isotope exchange upon solvolysis of N-HO-1-NA in acidic H₂¹⁸O, and carbocation formation was indicated by formation of the solvolysis products 1-amino-2-naphthol and 1-amino-4-naphthol. The potential role of the guan-O⁶-yl naphthylamine adducts in the initiation of urinary bladder carcinogenesis through the formation of promutagenic lesions in DNA is discussed. (64 refs)

78-6248 Follicular and Parafollicular Adenomas in the Thyroids of Rats Treated with Thiamazole. (Fre) Stoll, R. (Laboratoire de la Fondation Bergonie, 180 rue de St. Genes, F 33076 Bordeaux Cedex, France); Faucouneau, N.; Maraud, R. *Ann Endocrinol (Paris)* 39(3): 179-189; 1978.

The development of follicular cell (FC) and parafollicular (PFC) adenomas in the thyroids of rats treated with the thyroid drug thiamazole (TMZ: 1-methyl-2-mercaptimidazole) was followed. Three groups of male Wistar rats, ages 3, 6, and 8 mo respectively, were administered 0.1% 0.2% TMZ in their drinking water for 1, 3, 8, 9, 12, or 17 mo. The rats were killed and their thyroids examined immediately after the end of treatment or at various times to 17 mo later. Control rats of same age were also sacrificed. Immediately after treatment, animals on TMZ for 1-3 mo had thyroid glands that were larger and heavier than controls. Hypertrophy and hyperplasia were evident, but no adenomas were observed. Rats sacrificed immediately after 8 mo treatment had one or more FC adenomas, some of which occupied an entire lobe of the thyroid. PFC adenomas developed in rats starting 6 mo after the cessation of TMZ treatment. In rats treated for 3 mo, the PFC adenomas were only type observed after cessation of treatment; in rats treated for 8 mo or more, PFC adenomas coexisted with FC adenomas. Among the 72 rats treated for 9-14 mo, 64 had adenomas, 46 FC, 7 PFC, and 11 FC + PFC types. The percentage of C cells (or parafollicular cells, of which the PFC adenomas are composed) increased with age of rat. C-cell hyperplasia occurred in rats during TMZ treatment, but no PFC adenomas were formed until treatment had stopped. It is concluded that both types of adenomas are related to the hypothyroid condition and its duration, but the hormonal mechanism involved differs. (19 refs)

78-6249 3-Hydroxyxanthine: Transplacental Effect on Ontogeny of Related Sulfate Metabolism in Rats and Mice. (Eng) Anderson, L. M. (Memorial Sloan-Kettering Cancer Center, Walker Lab., Rye, NY 10580); McDonald, J. J.; Budinger, J. M.; Mountain, I. B.; Brown, G. B. *J Natl Cancer Inst* 61(6): 1405-1410; 1978.

Sulfotransferase activity toward the carcinogenic parent N-oxide 3-hydroxyxanthine (3-OH-X) was detected in fetal Sprague-Dawley (SD) rats and (C57BL/6 x BALB/c)F₁ mice delivered near term at a rate of approx 25% of adult values. The activities of 3-OH-X sulfotransferase, 3'-phosphoadenosine-5'-phosphosulfate (PAPS) sulfotransferase and ^3S -p-nitrophenol (PNP) sulfotransferase were compared in an assay using homogenized fetal livers. At day 21 of gestation, 3-OH-X sulfotransferase activity in fetal livers was significantly lower than adult values, and treatment of pregnant females with 3-OH-X (25 mg/day 4x) did not alter the enzyme levels. After birth, the activity rose linearly to adult values at 6 wk. PNP sulfotransferase was low or absent in fetal livers, but after birth, it eventually reached adult levels at 6 wk. PAPS sulfotransferase was 2-3 times higher in fetal livers than in livers of pregnant females or 6-wk-old rats, and it fell steadily to adult values after birth. 3-OH-X was injected ip into pregnant rats four times at doses of 1, 5, 10, or 25 mg on days 19-21 of gestation; pregnant mice were injected ip with 50 mg/kg on days 17 and 18. Of 10 male rat offspring receiving 25 mg/kg doses prenatally, 3 had liver tumors (including 2 hepatoma

compared with 0/29 controls. Kidney disease at all doses was common in male rats, but no liver abnormalities were seen in female offspring. Prenatal administration of 3-OH-X in rats resulted in significantly increased prenatal deaths, but the survivors had fewer lung adenomas at 20 mo of age (6/28 vs 21/50 controls). The association of low levels of activating enzyme in the male fetal rat liver with the weak but significant tumorigenic effect of 3-OH-X supports the hypothesis that enzymatic activation may be a limiting factor in transplacental carcinogenesis. (25 refs)

78-6250 Increased Incidence of Mammary Tumors in Rats After Direct or Transplacental Exposure to 3-Hydroxyxanthine. (Eng) Anderson, L. M. (Memorial Sloan-Kettering Cancer Center, Walker Lab., Rye, NY, 10580); Teller, M. N.; Budinger, J. M.; Brown, G. B. *J Natl Cancer Inst* 61(6): 1411-1414; 1978.

Rats exposed directly or transplacentally to N-oxide 3-hydroxyxanthine (3-OH-X) developed mammary tumors at significantly increased rates over controls. Among female Wistar rats that received 3-OH-X sc 1x/wk in varying doses, beginning at age 9 days, 4 wk, or 10 wk, there was a significantly increased proportion of animals bearing mammary tumors and an increased number of tumors/rat in all dose groups except the highest (7 mg x 22). In that group, the rats died before mammary tumors could develop. Histologically, the tumors included fibroadenomas, adenomas, and carcinomas. Administration of 3-OH-X to males sc or to females did not increase mammary tumor incidence. Female offspring of Sprague-Dawley mothers given four ip injections of 25 mg 3-OH-X during pregnancy developed mammary tumors significantly more often than controls (93% vs. 57%). The number of tumors/rat was also significantly increased, but the number of females with at least one malignant mammary tumor (3/14) was similar to that in control group (1/16). Lower prenatal doses of 3-OH-X did not increase male mammary tumor incidence, but male tumor incidence was greater than the control incidence in all dose groups. Because the tumors initiated by transplacental exposure to 3-OH-X remained dormant for many months, prolactin may have stimulated their growth as the rats aged. (19 refs)

78-6251 Excretion and Distribution of Morpholine Salts in Rats. (Eng) Tanaka, A. (Dept. Medical Chemistry, Natl. Inst. Hygienic Sciences, 18-1, Kamiyoga 1-chome, Setagaya-ku, Tokyo, Japan); Tokieda, T.; Namikawa, S.; Osawa, M.; Yamahara, T. *J Food Hyg Soc Jpn* 19(3): 229-234; 1978.

The elimination, organ distribution, and metabolism of morpholine HCl (MPL HCl) and MPL palmitate were studied in male Wistar rats by chemical assays using gas-liquid chromatography and by radioassays using ¹⁴C-MPL. For the chemical assay, the MPL HCl doses were 500 mg/kg po and

250 mg/kg iv; for the radioassays, the doses were 200 mg/kg po and 150 mg/kg iv. ¹⁴C-MCL palmitate (400 mg/kg) was given po through a stomach tube. MPL HCl was eliminated very rapidly after po and iv administration. In the chemical assays, > 85% appeared in urine after 24 hr, and 90% was excreted over 3 days. Very small amounts were excreted in feces. Similar results were obtained for both salts by radioassay. Over 5 days, 80%-90% was excreted in urine and 3%-10% in feces, with fecal excretion of the palmitate being slightly higher than that of the HCl. Thin-layer chromatography confirmed that MPL was excreted unchanged. MPL accumulated primarily in muscle and intestine, and significant quantities remained in the intestine for 12 hr. Elimination from other organs was very rapid. Adipose tissue showed the lowest affinity for MPL, regardless of route of administration. From these findings, it would appear that MPL is an inert substance in the rat body. (15 refs)

78-6252 Chromosome Investigations on Workers Exposed to Styrene/Polystyrene. (Eng) Thiess, A. M. (Occupational Medicine and Health Protection, BASF Aktiengesellschaft, 6700 Ludwigshafen/Rhein, W. Germany); Fleig, I. *J Occup Med* 20(11): 747-749; 1978.

Cytogenetic studies were carried out in 12 chemical plant employees with 3-39 yr possible exposure to styrene, and urinary mandelic acid levels were studied in 58 additional employees. The frequency of aberrant metaphases in the chromosomes of the 12 workers studied was 1.9%; this frequency was the same as that in 12 controls. Forty-eight workers excreted < 30 mg/liter mandelic acid in the urine; 8 excreted 30-50 mg/liter, and only 2 excreted > 50 mg/liter. The concentrations of styrene at the plant were usually < 1 ppm; in areas where the concentrations were higher, but < 10 ppm, exposure lasted only 3-5 min at a time. (9 refs)

78-6253 Mutagenicity of Styrene and Styrene Oxide in Drosophila Tests (Meeting Abstract). (Eng) Sorsa, M. (Dept. Genetics, Univ. Helsinki, Helsinki, Finland); Laamanen, I.; Vainio, H. *Mutat Res* 53(2): 266-267; 1978. (no refs)

78-6254 Carcinogenicity and Mutagenicity of 5-Nitroacenaphthene and Tetrachlorobenzidine. (Eng) Takemura, N. (Dept. Public Health, Jikei Univ. Sch. Medicine, Tokyo, Japan); Hashida, C.; Shimizu, H. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1329-1336; 1978.

The carcinogenicity of 5-nitroacenaphthene (NAN) and tetrachlorobenzidine (TCB) was tested in male and female Wistar rats and Syrian golden hamsters and in female dd strain mice; their mutagenicity was tested on *Salmonella typhimurium* strains TA100 and TA98. Twelve of 30 female rats given NAN (1% of diet for 4 mo) lived > 200 days after the start of feeding and developed malignant tumors: 1 rhabdomyosarcoma, 2 sebaceous squamous cell carcinomas in the ear duct, 5 intraductal breast carcinomas, and 10 adenocarcinomas of the small intestine. Ten of 20 female rats given NAN (1% of diet for life) lived > 200 days and developed tumors in the ear duct (4), breast (3), kidney (1), and liver (1). Thirteen of 24 female hamsters given NAN (1% of diet for 6 mo) survived > 270 days, and 7/13 developed cholangiomas. None of the males developed tumors. Seventeen of the 20 mice given TCB (0.1% of diet for life) survived > 300 days and developed 14 malignant lymphomas, 2 mammary carcinomas, and 1 mammary fibrosarcoma. Both chemicals were mutagenic on *S. typhimurium* TA100 and TA98 with metabolic activation. Precautions should be taken when these chemicals are handled occupationally. (4 refs)

78-6255 Cytological Effects of the Antihelminthic Agent Oxytocanide on Mitotic Chromosomes of *Allium cepa* L. (Eng) Reddy, P. V. (Dept. Zoology, Kakatiya Univ., Vidyanayapuri Warangal 506 009, India); Murthy, D. K.; Subramanyam, S. *Indian J Exp Biol* 15(1): 1073-1074; 1977.

The effect of the antihelminth oxytocanide (Zanil, or 3,3,5,5,6-pentachloro-2,2-dihydroxybenzylidene) on *Allium cepa* roots soaked in a 0.25% soln of the agent for 2-24 hr was determined. There was a gradual decline in the mitotic index with time, with the most intense effect being observed around 24 hr. A pronounced C-mitotic effect and chromosome abnormalities were also induced. The mitotic index rose slowly and the percentage of abnormal cells decreased when the roots were allowed to recover for 24, 48, and 72 hr after an 8-hr treatment, indicating the reversibility of the genetic damage. (14 refs)

78-6256 Potential Carcinogenicity of Some Food Preservatives in the Presence of Traces of Nitrite (Meeting Abstract). (Eng) Khoudokormoff, B. (Lab. Microbial Genetics, Res. and Development, P.O. Box 1, Delft, Netherlands); Gist-Brocades, N. V. *Mutat Res* 53(2): 208-209; 1978. (no refs)

78-6257 Enhancement of Liver Microsome Epoxide Hydratase Activity in Rodents by Treatment with 2(3)-tert-Butyl-4-hydroxyanisole. (Eng) Cha, Y. N. (Dept. Pathobiology, Johns Hopkins Medical Inst., Baltimore, MD, 21205); Martz, F.; Bueding, E. *Cancer Res* 38(12): 4496-4498; 1978.

Epoxide hydratase (EH) activity was assayed in liver microsomes from female CD-1 mice fed a diet containing 0.75% of the antioxidant 2(3)-tert-butyl-4-hydroxyanisole (BHA) for 10 days. The liver microsomes were incubated with styrene oxide [200 nanomoles (nmol)], and EH activities were expressed in terms of nmol of styrene diol metabolite per mg protein per min. Three days after initiation of BHA feeding the specific activity of EH was 9.40 nmol diol/mg/protein/min, or 9 x the control value. The activity then rose more slowly, reaching 11.11 nmol/mg/protein/min at 10 days. Only two- to threefold increases in enzyme activity were induced by 3-methylcholanthrene, phenobarbital, and Aroclor 1254. When BHA (750 mg/kg/day) was administered by gastric intubation for 4 days, the increase in EH activity was 75% of that attained by dietary administration of the antioxidant. The possibility that BHA may enhance or protect against the carcinogenic effects of diol-epoxides is discussed (26 refs)

78-6258 N-Acetyl- β -glucosaminidase Activity in the Peripheral Blood WBC of Rats Subacutely Exposed to Benzene Vapor. (Pol) Starek, A. (Katedra Medycyny Pracy i Chorob Zawodowych, AM w Krakowie, 30-900 Krakow 28, Poland); Moszczynski, P. *Patol Pol* 29(1): 71-72; 1978.

The effect of a 10-day exposure to benzene vapor (6 hr/day, 27,000 mg/m³) on N-acetyl- β -glucosaminidase (AGA) activity in the peripheral blood WBC was studied in 12 male Wistar rats aged 8 mo. The exposure reduced the number of neutrophil granulocytes from 3,826/mm³ to 2,271/mm³ (p < 0.005), the AGA activity in the granulocytes from 53.8 to 16.8 units (p < 0.001), the lymphocyte count from 10,963/mm³ to 1,330/mm³ (p < 0.001), and the percentage of AGA-positive lymphocytes from 54.9% to 50.4% (p < 0.05). Within the subpopulation of AGA-positive lymphocytes, the percentage of granular-type cells decreased and that of granular-diffusion and diffusion-type cells increased. The effects are due to the damaging action of benzene on the granulocyte lysosomes and on peripheral blood lymphocytes (24 refs)

78-6259 Modified Method for Electron Capture Gas-Liquid Chromatographic Determination of Hexestrol Residues in Bovine Tissues. (Eng) Tobioka, H. (Dept. Animal Science, Coll. Agriculture, Kyoto Univ., Kyoto 606, Japan); Kawashima, R. *J Assoc Off Anal Chem* 61(5): 1054-1057; 1978.

Hexestrol residues in the tissues of cows injected with hexestrol dicaprylate were determined by electron capture gas-liquid chromatography. The method was suitable for determining residues of 0.5 ppb; the limit of sensitivity ranged from 0.1 to 0.2 ppb. (16 refs)

6260 Thiophene Analogues of the Carcinogens Benzidine and 4-Aminobiphenyl: Evaluation In Vitro.

(Eng) Ashby, J. (Central Toxicology Lab., Imperial Chemical Industries Ltd., Alderley Park, Macclesfield, Cheshire, England); Styles, J. A.; Anderson, D.; Paton, D. *Br J Cancer* 38: 521-529; 1978.

Thiophene analogs of benzidine and 4-aminobiphenyl (N-acetamidophenyl-2-thiophenamine hydrochloride (XV), 5-phenyl-2-thiophenamine hydrochloride (XIX), N-p-acetamidophenylthiophen-2-yl)acetamide (XV), and N-phenylthiophen-2-yl)acetamide (XVI), were tested for mutagenicity in vitro by the Ames reverse-mutation assay and the Styles cell-transformation assay. In Ames tests with *Salmonella typhimurium* strains TA1535, TA1538, TA98, and TA100, each of the thiophenes produced at least one positive response in TA1538 or TA98, sometimes in the absence of a rat liver S-9 mix. In two runs of the Styles assay, the acetyl derivatives XV and XVI gave a positive response, while the amine hydrochlorides XIX and XVI were negative. The acetyl derivatives possessed greater cytotoxicity. It was suggested that the relative transforming abilities of these compounds is determined by the ability of XV and XVI to form DNA-reactive hydroxylamine-ester compounds and the ability of XIX and XIV to form N-hydroxy-ester compounds. (29 refs)

6261 Di-(4-aminophenyl)-methane (MDA): 4-7 Year Dog Feeding Study. (Eng) Deichmann, W. B. (National Center for Toxicology, Dept. Pharmacology, Univ. of Miami Sch. of Medicine, P.O. Box 248216, Coral Gables, FL 33124); MacDonald, W. E.; Coplan, M.; Woods, F.; and E. *Toxicology* 11(2): 185-188; 1978.

The possible carcinogenic action of purified and crude di-(4-aminophenyl)-methane (MDA) on the urinary bladder was studied in nine female beagle dogs. The compounds were given po at a dose of 70 mg/kg/day 3 x/wk for life (up to 4 yr and 2 mo). There were no significant effects on body weight except for an occasional weight loss that was followed by quick recovery after withdrawal from treatment. Results of all laboratory tests were negative except for elevated alkaline phosphatase activity in two dogs. Moderate to severe gross morphologic changes occurred in the livers of all dogs; severe changes were seen in the kidneys (4 dogs), spleens, urinary bladders (2), lungs (3), ovaries and fallopian tubes, and uterus (1). There were no tumors of the urinary bladder or liver. (3 refs)

6262 Chlorination of Benzidine and Other Aromatic Amines in Aqueous Environments. (Eng) Jensen, R. L. (County Sanitation Districts Los Angeles County, San Jose Creek Water Quality Lab., 1965 S. Workman Mill Road, Whittier, CA, 90601); Haskins, J. E.; Carmona, L. G.; and Bird, R. B. *Arch Environ Contam Toxicol* 7(3): 301-315; 1978.

Exposure of aniline, but not N,N-dimethylaniline, to chlorinated waters yielded a solid product. For these monophenyl amines, ring chlorination was a significant depletion pathway. Extended chlorination of benzidine resulted immediately in a solid product that, upon infrared analysis, was shown to be a polymeric structure with no ring chlorination. Analysis of the supernatant revealed no ring-substituted isomers of benzidine. (18 refs)

78-6263 An Improved Method for Detecting Mutagens (Meeting Abstract). (Eng) Yahagi, T. (Natl. Cancer Res. Inst., Tokyo, Japan); Nagao, T.; Matsushima, T.; Seino, Y.; Sawamura, N.; Shirai, A.; Kawachi, T.; Sugimura, T. *Mutat Res* 53(2): 285; 1978. (no refs)**78-6264 Genetic Variation in N-Acetylation of Carcinogenic Arylamines by Human and Rabbit Liver.**

(Eng) Glowinski, I. B. (Dept. Pharmacology, Univ. Michigan, Ann Arbor, MI, 48109); Radtke, H. E.; Weber, W. W. *Mol Pharmacol* 14(5): 940-949; 1978.

The arylamine carcinogens aminofluorene, α -naphthylamine, β -naphthylamine, benzidine, and methylene bis-2-chloroaniline were found to be acetylated by the same polymorphic N-acetyltransferase (NAT) as sulfamethazine (SMZ) and isoniazid (INH). Liver (0.10 ml) NATs purified from slow and fast INH-acetylating rabbits and humans were incubated with 0.05 ml aqueous radioactive acetyl coenzyme A (2×10^{-3} M) and 0.05 ml amine substrate solution. After the radioassay of acetylating activity, K_m values were determined for enzyme-substrate interactions. Rabbit liver NAT had a smaller apparent K_m than the human liver enzyme for all substrates tested except α and β naphthylamine, and reactions with carcinogenic substrates in both enzyme systems yielded K_m 's at least an order of magnitude smaller than when SMZ was the substrate. Although K_m values within a species did not change, the carcinogenic compounds showed greater differences in acetylation rates between rapid and slow acetylators than did SMZ; therefore, they appeared to be polymorphically acetylated in both rabbit and man. Octanol-water partition coefficients measured for each substrate correlated closely with the apparent K_m values for both human and rabbit NAT, supporting the hypothesis that differences in apparent K_m 's were related to the greater hydrophobicity of the carcinogens than of SMZ. Because all the carcinogens served as substrates for polymorphic NAT, hydrophobicity and differences between slow and fast acetylators could be important in determining an individual's susceptibility to arylamine carcinogenesis. (51 refs)

78-6265 Hepatocarcinogen-induced tRNA Methylase Activity in Rat Liver. (Eng) Forrester, P. I. (Div. Medical Biochemistry, Faculty Medicine, Univ. Cal-

gary, Calgary, Canada); Hancock, R. L. *Physiol Chem Phys* 10(2): 115-123; 1978.

Dimethylaminoazobenzene and carbon tetrachloride extend the list of hepatocarcinogens capable of increasing rat liver transfer RNA (tRNA) methylase activity. Ethionine increased this activity in rats and guinea pigs but not in chicks. Aminoazobenzene, acetylaminofluorene, α - and β -naphthylisothiocyanate, dimethyl- and diethylnitrosamine, and methylcholanthrene produced no increase or slight increases. (13 refs)

- 78-6266 **Toxicity of p-Substituted Derivatives of Aniline in Experimental Rats.** (Cze) Matrká, M. (Toxikol. Odd., Vyzkumny ustav organických syntéz, 533 51 Pardubice-Rosice, Czechoslovakia); Rambousek, V.; Zverina, Z. *Cesk Hyg* 23(4): 168-172; 1978.

The acute po toxicity of aniline and six of its substituted derivatives was determined in male Wistar rats. The substances were administered in soln or in an olive oil suspension. The LD₅₀ values were 0.440 mg/kg for aniline (A), 0.417 mg/kg for 4-fluoroaniline (FA), 0.370 mg/kg for 4-chloroaniline (CA), 0.456 mg/kg for 4-bromoaniline (BA), 0.523 mg/kg for 4-iodoaniline (IA), 0.794 mg/kg for 4-methylaniline (MA), and 0.750 mg/kg for 4-nitroaniline (NA). The mole LD₅₀ values were 0.00265 (BA), 0.00239 (IA), 0.00755 (MA), and 0.00543 (NA). Hammett's σ constants were 0 (A), +0.06 (FA), +0.23 (CA and BA), +0.27 (IA), -0.17 (MA), +0.78 (NA). A good correlation was seen between the mole LD₅₀ values and Hammett's constants. The six aniline derivatives represent the last biotransformation step of the corresponding carcinogens derived from 1-phenyl-3,3-dimethyltriazene. (16 refs)

- 78-6267 **Metabolism-mediated Cytotoxicity of Chemical Carcinogens and Non-carcinogens.** (Eng) Wiebkin, P. (Inst. Industrial, Environmental Health and Safety, Univ. Surrey, Guildford, Surrey GU2 5XH, England); Fry, J. R.; Bridges, J. W. *Biochem Pharmacol* 27(14): 1849-1851; 1978.

The general applicability of a mixed cell culture system in assessing the importance of metabolism in chemically mediated cytotoxicity was tested using carcinogenic and noncarcinogenic chemicals. Rat fibroblasts were incubated with at least five concentrations of each test compound for 1 hr at 37 C in the presence or absence of viable rat hepatocytes. The samples were then diluted and cultured at 37 C for 6-8 days, after which time counts of nuclei were performed and the results calculated as ID₅₀ values; i.e., the concentration of the chemical that inhibited fibroblast growth by 50%. In the presence of hepatocytes, the most toxic pair of compounds tested was 3'-methyl-4-dimethylaminoazobenzene (3'-methyl-DAB; ID₅₀, 0.1 μ g/ml) and 2-methyl-4-dimethyl-

aminoazobenzene (2-methyl-DAB; ID₅₀, 2.0 μ g/ml). In the absence of hepatocytes, the ID₅₀ values for 3'-methyl-DAB and 2-methyl-DAB were > 125 and > 250 μ g/ml, respectively. Mitomycin C also required hepatocyte metabolism for its cytotoxicity to be fully expressed, but hepatocyte metabolism decreased the cytotoxicity of 4-nitroquinoline-1-oxide. Other compounds that showed increased cytotoxicity in the presence of hepatocytes were cyclophosphamide, dimethylnitrosamine, 2-acetylaminofluorene, 4-acetylaminofluorene, 6-aminochrysene, and 2-aminochrysene. (14 refs)

- 78-6268 **Mutagenicity of the Herbicide Trifluralin in *Drosophila melanogaster*** (Meeting Abstracts) (Eng) Murnik, M. R. (Dept. Biological Sciences, Western Illinois Univ., Macomb, IL, 61455). *Mutat Res* 53(2): 235-236; 1978. (1 ref)

- 78-6269 **Mutagenicity of Six Indian Hair Dyes Tested in *Salmonella typhimurium* Strains** (Meeting Abstracts). (Eng) Bajaj, M. H. (Biology and Agriculture Department, Bhabha Atomic Res. Centre, Bombay-400 085, India); Nigam, N. K. *Mutat Res* 53(2): 149-150; 1978. (no refs)

- 78-6270 **Liver and Urinary Bladder Tumors in Dogs from 3,3'-Dichlorobenzidine.** (Eng) Stula, E. (Haskell Lab., Elkton Road, Newark, DE, 19711); Barnard, J. R.; Sherman, H.; Reinhardt, C. F.; Zapp, J. A. *J Environ Pathol Toxicol* 1(4): 475-490; 1978.

The carcinogenic effects of po 3,3'-dichlorobenzidine (DCB) were studied in six female beagle dogs. DCB was given at a rate of 100 mg 3x/wk for 6 wk, then 5x/wk for up to 7.1 yr. Untreated controls were sacrificed after 8.3-9.0 yr. All DCB-treated dogs showed elevated plasma glutamic-pyruvate transaminase activity during the first 3 yr, and two dogs showed elevated activity throughout the test. One DCB dog sacrificed in extremis after 3.5 yr, had no tumors. Another dog sacrificed in extremis after 6.6 yr, had an undifferentiated carcinoma of the liver with metastases to many organs and a papillary transitional cell carcinoma of the urinary bladder. Three of the remaining four dogs sacrificed after 7.1 yr had hepatocellular carcinomas, and all four had transitional cell carcinomas of the urinary bladder. There were no such tumors in the control dogs. The data indicate that DCB is carcinogenic for the liver and urinary bladder in the dog and that in this species, it is more carcinogenic than benzidine. (1 refs)

- 78-6271 **Mutagenicity of Pesticides in Blue-Green Algae.** (Eng) Das, B. (Lab. Blue-Green Algae)

tral Rice Res. Inst., Cuttack 753006, Orissa, India);
gh, P. K. *Microbios Lett* 5(19/20): 103-107; 1977.

e mutagenicity of hexachlorocyclohexane (BHC) and
dichlorophenoxyacetic acid (2,4-D) to four blue-green al-
(3 N-fixing, 1 non-N-fixing) was investigated. A slight
crease in mutation frequency was noticed after treatment
h BHC, but 2,4-D treated cultures did not show any in-
ase. The results suggest that BHC is a weak mutagen for
e-green algae, whereas 2,4-D is nonmutagenic. (19 refs)

6272 **The Mutagenicity of Substituted and Unsub-**
stituted Styrene Oxides in *E. coli*: Relationship
between Mutagenic Potencies and Physico-chemical Proper-
ties. (Eng) Sugiura, K. (Inst. Ecotoxicology, Gakushuin
iv., Toshima-ku, Tokyo 171, Japan); Yamanaka, S.;
kasawa, S.; Goto, M. *Chemosphere* 7(9): 737-742; 1978.

e relationship between the mutagenic potencies and
physicochemical properties of substituted and unsubstituted
styrene oxides was studied. 3,4-Dimethylstyrene oxide was
most lethal to *Escherichia coli* strain WP100 and more muta-
genic in strains WP100 and CM571 than in strains WP2uvrA
and WP2. Styrene oxide had similar lethal and mutagenic
effects. These results indicate that the uvrA and recA genes
affect both the lethal and mutagenic responses. The
mutagenicity and lethality of these and other compounds in-
creased in the order: m-methoxystyrene oxide < styrene ox-
ide < p-methylstyrene oxide < m-methylstyrene oxide <
3,4-dimethylstyrene oxide < m-chlorostyrene oxide <
bromostyrene oxide. In reactions of substituted styrene ox-
ides with nucleophiles, the normal reaction rate was higher
with electron-withdrawing substituents and lower with elec-
tron-releasing substituents. In contrast, the rate of abnormal
reactions (attack at CH) was higher with electron-releasing
substituents. The relative LD₅₀ values of the styrene oxide
derivatives agreed with the relative partition coefficients, sug-
gesting that the rate of transport of styrene oxides to their
molecular sites of action is the rate-limiting condition. The data
demonstrate that the mutagenic effect of a chemical may vary
with the structure of its products. (11 refs)

6273 **Pathology of Esophageal Neoplasms and As-**
sociated Proliferative Lesions Induced in Rats
by N-Methyl-N-benzyl nitrosamine. (Eng) Stinson, S. F. (Tu-
mor Pathology Branch, Div. Cancer Cause and Prevention,
NCI, NIH, Public Health Service, U.S. Dept. Health, Educa-
tion and Welfare, Bethesda, MD, 21205); Squire, R. A.;
Barn, M. B. *J Natl Cancer Inst* 61(6): 1471-1475; 1978.

After receiving 20 weekly sc injections of N-methyl-N-ben-
zyl nitrosamine (MBZN: 2.5 mg/kg), 24 inbred male F344
rats developed 194 lesions of the upper gastrointestinal tract.

Grossly, most lesions were exophytic; 74% were pedun-
culated papillary growths and the remainder were sessile pro-
liferations. Localized elevated thickening of the esophageal
wall was observed in 20% of the rats. All rats developed
papillomas, 63% developed carcinomas that invaded the eso-
phageal wall, and 67% developed pedunculated papillary car-
cinomas. Of the esophageal tumors, 66% were papillomas,
17% were pedunculated papillary carcinomas, and 17% were
invasive carcinomas. The papillomas were pedunculated
(87%) or sessile (13%). The carcinomas that invaded the
esophageal wall were sessile (78%) or infiltrating (22%). All
neoplasms were well-differentiated squamous cell tumors,
with varying degrees of keratinization. Although penetration
through the esophageal wall was common, no metastases
were seen. No primary lesions were seen in sites other than
the esophagus and pharynx. These tumors were similar mor-
phologically to human specimens. Therefore, the usefulness
of this model in the study of human esophageal carcinoma
is indicated. (11 refs)

78-6274 **The Role of Glutathione in Experimental**
Chemical Hepatocarcinogenesis. (Eng) Fiala, S.
(Cell Physiology Lab., VA Center, Martinsburg, WV); Trout,
E. C.; Kettering, W. G. In: *Prevention and Detection of Can-*
cer. Proceedings of the Third International Symposium on De-
tection and Prevention of Cancer Held by the International
Study Group for the Detection and Prevention of Cancer in
New York, April 26 - May 1, 1976. International Study Group
for the Detection and Prevention of Cancer. (New York,
NY): Vol. 2(Part 1), 2404 pp.; 1291-1304; 1978.

The behavior of glutathione (GSH) in rat liver during azo
dye-induced carcinogenesis was investigated to determine
whether hepatocarcinogenesis is due to inhibited and disrupt-
ed differentiation of stem cells. The concentration of GSH
in fetal and neonatal liver was 110 and 140 mg/100 g, respec-
tively, vs 234 mg/100 g in adult liver. Following administra-
tion of 3'-methyl-4-dimethylaminoazobenzene (3'-Me-DAB;
33 mg), there was a small initial decrease in liver GSH due
to its conjugation with some of the azo dye metabolites and
its elimination in bile. This was followed by a rapid increase
in GSH synthesis as early as 8 hr after 3'-Me-DAB adminis-
tration. In later stages of hepatocarcinogenesis, γ -glutamyl-
transpeptidase (GT-ase) activity was very high, resulting in
rapid degradation of GSH. The liver of hypophysectomized
rats, which do not develop hepatomas as a result of azo dye
administration, did not form GSH conjugates and did not
have increased GSH or GT-ase activity. GSH, being nucleo-
philic, may combine with and protect against various electro-
philic metabolites. Even with an excess of carcinogen, the
hepatocyte seems to be largely immune to electrophilic attack
because the initial GSH depletion is overcompensated by its
increased rate of synthesis. The mature hepatocyte, therefore,
does not appear to be the target of proximate carcinogens.
A fetal-like hepatic stem cell with high GT-ase activity may
be the actual carcinogen-sensitive cell. (27 refs)

- 78-6275 Inhibition of 3'-Methyl-4-dimethylaminoazobenzene-induced Hepatocarcinogenesis by Portacaval Shunt.** (Eng) Fiala, S. (Cell Physiology Lab., Veterans Admin. Center, Martinsburg, WV, 25401); Pragani, B.; Reuber, M. D. *Tumori* 64(2): 131-142; 1978.

Adult male Sprague-Dawley rats who underwent an end-to-side portacaval shunt (PCS) did not develop hyperplastic nodules and hepatomas when they were fed 3'-methyl-4-dimethylaminoazobenzene (3'-Me-DAB) in a semisynthetic basal diet for up to 169 days. In contrast, all intact rats fed the same diet for only 75 days developed hyperplastic nodules. Following transfer to a normal diet for another 25 days, hepatomas developed in 100% of these animals. The amount of protein-bound 3'-Me-DAB and the glutathione (GSH) level was lower in operated rats than in intact controls. A single large dose of 3'-Me-DAB (10 mg/100 g) increased the GSH concentration by only 30% after 24-48 hr, compared with an increase of 50%-100% in intact rats. There was no gradual increase in the activity of γ -glutamyltranspeptidase (GTase) in operated rats during 5.5 mo of 3'-Me-DAB feeding. The increase in GTase activity was never > 30% higher than the level in livers of operated rats fed a basal diet without the carcinogen. This striking inhibition of GTase activity induced by 3'-Me-DAB in PCS-operated rats contrasted with a 5,000% increase in enzyme activity found in the livers of intact rats with hyperplastic nodules after 75 days of 3'-Me-DAB feeding and with the up to 10,000% increase in the hepatomas. These effects and the inhibition of 3'-Me-DAB-induced hepatocarcinogenesis, manifested by a lack of preneoplastic morphologic changes and the absence of hepatomas in rats after PCS, can best be explained by a functional deficiency of the liver to metabolize the procarcinogen 3'-Me-DAB to an activated carcinogen. (25 refs)

- 78-6276 A Study of Cyclic Nucleotide Metabolism and the Histology of Rat Liver During 3'-Methyl-4-dimethylaminoazobenzene Carcinogenesis. I. Histology.** (Eng) Boyd, H. (Dept. Pharmacology, Univ. Miami Sch. Medicine, Miami, FL, 33152); Hensley, G. T. *Tissue Cell* 10(3): 463-476; 1978.

Male Sprague-Dawley albino rats were fed 3'-methyl-4-dimethylaminoazobenzene (3'-MeDAB) ad libitum in pellets (0.06% in 0.75% corn oil) for 12 wk, and their livers were examined histologically at weekly intervals up to 13 wk. The first tumor mass was noted at 4 wk and there was a 100% incidence of cholangiocarcinomas by 10 wk. Acidophilic hepatocytes that appeared in periportal regions of liver lobules at 3 wk were smaller than normal and glycogen-deficient. After 4 wk, even smaller glycogen-deficient cells with a large nucleus/cytoplasm ratio appeared. These small cells and the acidophilic cells underwent mitosis, and there was no indication of acute or chronic inflammation or fibrosis. By 7 wk, the small cells became increasingly organized into ductular structures that were composed of columnar epithelial cells and mucous-secreting goblet cells. Proliferating epi-

thelial cells were frequently associated with necrosis and destruction of the parenchyma. Tissue that appeared to be uninvolved to the naked eye often contained 50% unorganized small cells. In this study, no hepatocellular carcinoma developed by 13 wk. (42 refs)

- 78-6277 Pre-tumor Phase Serum Alpha-Fetoprotein and Its Molecular Variants During Hepatocarcinogenesis.** (Eng) Nadworny, H. A. (Dept. Medicine, Univ. Vermont Coll. Medicine, Burlington, VT, 05401); Smith, C. Kelleher, P. C. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY) Vol. 2(Part 1), 2404 pp.; 1283-1290; 1978.

Radial immunodiffusion was used to monitor the weekly serum alpha-fetoprotein (AFP) concentrations in 39 Fischer 344 rats that were fed 0.063% 3'-methyl-4-dimethylaminoazobenzene (3'-Me-DAB) in the diet beginning at 6 wk of age. Serum AFP levels were undetectable (< $\mu\text{g/ml}$) at 6 wk of age. AFP was detectable in the serum of 30/39 rats after 3 wk of 3'-Me-DAB intake and in all surviving rats after 6 wk of intake. This first elevation of pre-tumor phase serum AFP lasted until 7 wk after the start of the diet and was followed by a second elevation starting at 10-11 wk in 35/36 rats and lasting for 4-6 wk. Max serum AFP concentrations during each period of increase varied widely among the individual rats (12-700 $\mu\text{g/ml}$ in the first period, 20-220 $\mu\text{g/ml}$ in the second). The percentages of total pre-tumor phase serum AFP not reactive with concanavalin A showed no consistent pattern either within an individual animal (14%-51%) or among individuals (11%-51%). The mechanisms responsible for the cyclic elevations of pre-tumor phase serum AFP are not known. (13 refs)

- 78-6278 Mechanisms of Immunosuppression Induced in Mice by o-Aminoazotoluene and Its Noncarcinogenic Analog.** (Rus) Kaledin, V. I. (Lab. Cancer Genetics, Inst. Cytology and Genetics, Siberian Branch, Acad. Sciences USSR, Novosibirsk, USSR); Glazko, T. T.; Krasovskiy, P. M.; Chesnokova, V. M. *Vopr. Onkol* 24(9): 46-52; 1978.

The immunosuppressive effects of single ip doses of o-aminoazotoluene (OAT: 4.5 mg) or its noncarcinogenic analog aminoazobenzene (AAB: 4 mg) were studied in male DD mice inoculated im with 10^6 Krebs-2 carcinoma cells. When OAT and AAB were administered 24 hr before tumor transplantation, tumor wts at the end of the experiment were 0.68 and 0.9 g, respectively, vs 0.55 g in controls. When OAT and AAB were injected 72 hr before tumor graft, tumor wts were 1.36 and 1.26, respectively, vs 0.91 g in controls. Immunization of the mice with irradiated (10,000 rads) tumor cells

ngly inhibited the growth of the implanted tumor, but the or wt was 3-3.5 times higher in the immunized and OAT-AAB-treated animals than it was in immunized animals out OAT or AAB treatment. The findings indicate the ly pronounced immunosuppressive effect of OAT and B. The immunosuppressive effect of AAB was due to its phocytotoxicity, whereas the effect of OAT was mediated the adrenal gland. After OAT injection, the serum oxycorticosteroid level was elevated, although production his hormone by the adrenal glands was not increased. The umulation of glucocorticoids in serum after OAT injection elieved due to reduced hormone utilization because of a ck of receptor proteins or metabolizing hepatic enzymes the carcinogen. (12 refs)

6279 **The Generation of Aerosols of Carcinogenic Aromatic Amines.** (Eng) Rappaport, S. M. (Sch. blic Health, Univ. California, Berkeley, CA, 94720); ttemy, D. J. *Am Ind Hyg Assoc J* 39(4): 287-294; 1978.

he development of an aerosol generator for the production test atmospheres of carcinogenic aromatic amines is de-ribed. The generator, a compressed-gas nebulizer, operates a temperature above the melting point of the test com-unds and uses nitrogen as a driving force. The nebulizer tput and size distributions of aerosols of 4,4'-me-ylenebis(2-chloroaniline), benzidine, and 3,3'-dichloroben-dine are given. The apparatus is capable of continuous oper- on over a minimum of 8 hr. (13 refs)

8-6280 **Mutagenicity Studies of Commercially Avail- ble Saccharins (Meeting Abstract).** (Eng) ajumdar, S. K. (Dept. Biology, Lafayette Coll., Easton, A, 18042); Viglianti, G. A.; Thatcher, J. D.; Dennis, E. H.; aharam, L. C.; Natoli, J. *Can J Genet Cytol* 20(3): 449; 1978. (no refs)

8-6281 **Absorption of Saccharin from Rat Urinary Bladder (Letter to Editor).** (Eng) Colburn, W. . (Dept. Biochemistry and Drug Metabolism, Hoffmann-La Roche, Inc., Nutley, NY, 07110). *J Pharm Sci* 67(10): 493-1494; 1978.

Measurements of plasma levels of saccharin following in vivo nstillation of 10 µg or 1 or 100 mg/kg of the compound into he urinary bladder of male Sprague-Dawley rats showed that his substance was absorbed extensively from the bladder. nificant reabsorption may decrease the elimination rate for accharin in much the same way as enterohepatic cycling and probably accounts for the retention of saccharin in urinary ladder tissue. (3 refs)

78-6282 **Saccharin Inhibits Guanylate Cyclase Activity: Possible Relationship to Carcinogenesis.** (Eng) Vesely, D. L. (Dept. Medicine, Univ. Miami Sch. Medicine, Miami, FL, 33152); Levey, G. S. *Biochem Biophys Res Com- mun* 81(4): 1384-1389; 1978.

The effect of saccharin (2,3-dihydro-3-oxobenzisul- fonazole) on guanylate cyclase activity was tested in a variety of tissues obtained from male Sprague-Dawley rats. Saccha- rin significantly inhibited guanylate cyclase in urinary blad- der, liver, stomach, colon, and kidney tissues over the concen- tration range 16-160 mM. Maximum inhibition was 60%-75%. The concentrations of saccharin decreasing gua- nylate cyclase activity in the in vitro rat urinary bladder assay (approx 3 to 9 mg/ml) were well within the range of concen- tration which can be obtained in urine of persons who use low calorie soft drinks and sugar substitutes and therefore interact in vivo with bladder tissue. The saccharin used in this study was synthesized by the Maumee process and contained no detectable amounts of o-toluenesulfonamide, the impurity most often associated with cancer. It is concluded that the question of saccharin carcinogenicity in man must be serious- ly studied regardless of its mode of synthesis. (29 refs)

78-6283 **Enhancement of Oncogenesis in C3H/10T1/2 Mouse Embryo Cell Cultures by Saccharin.** (Eng) Mondal, S. (Comprehensive Cancer Center, Univ. Southern California, Los Angeles, CA, 90033); Brankow, D. W.; Heidelberger, C. *Science* 201(4361): 1141-1142; 1978.

The transforming and promoting activities of saccharin were studied in vitro using C3H/10T1/2, clone 8, mouse embryo fibroblasts. At 2 mg/ml, saccharin alone was nontoxic and nontransforming, and it had no initiating effect when fol- lowed 4 days later by continuous treatment with 12-O-tet- radecanoylphorbol-13-acetate (TPA). However, both pure and impure samples of saccharin (100 µg/ml) showed signifi- cant cocarcinogenic activity during continuous treatment of cultures initiated with 3-methylcholanthrene (0.1 µg/ml). TPA, but not saccharin, promoted UV-initiated cells. The reason for this difference between these otherwise equivalent initiators is not clear. Nevertheless, the results suggest that saccharin acts as a promoter and that caution should be ap- plied to the indiscriminate or long-term use of saccharin in human nutrition. (22 refs)

78-6284 **Cytogenetic Hazards from Fensulfothion, Dex- on (DAPA) and its Degradation Product, Dime- thylphenylene Diamine (DMPDA) (Meeting Abstract).** (Eng) Panda, B. B. (Cytogenetics Lab., Berhampur Univ., Berhampur 760 007, India); Sahu, R. K.; Sharma, C. B. *Mutat Res* 53(2): 247-248; 1978. (no refs)

- 78-6285 **On the Mutagenic Action of 2-Methyl-3,5-dinitrobenzamide and 3,5-Dinitrobenzamide (Meeting Abstract).** (Eng) Voogd, C. E. (Natl. Inst. Public Health, PO Box 1, Bilthoven, Netherlands). *Mutat Res* 53(2): 280-281; 1978. (no refs)

- 78-6286 **Carcinogenicity of o-Aminoazotoluene in Mouse Intestine.** (Rus) Kaledin, V. I. (Lab. Cancer Genetics, Inst. Cytology and Genetics, Siberian Branch, Acad. Sciences USSR, Novosibirsk, USSR); Alekseeva, G. V.; Volkova, A. I. *Biull Eksp Biol Med* 86(10): 476-477; 1978.

The carcinogenicity of o-aminoazotoluene (OAT) was studied in 60 2-mo-old male A/He mice. OAT was implanted under the skin in 8 doses of 10 mg each at monthly intervals (total dose, 80 mg). All animals died 435 ± 13 days after the first implantation. Liver tumors were the principal cause of death, but 19/60 animals also developed intestinal tumors localized exclusively to the cecum. These tumors were identified as adenomas or muciparous adenocarcinomas. According to the literature, A/He mice do not develop spontaneous intestinal tumors, and in previous experiments with CC57BR, C57BL/6, C3H/He, and DD mice treated with OAT in the same manner, intestinal tumors failed to develop. The genetic basis for the sensitivity of the cecal tissue of A/He mice to OAT is unknown. (2 refs)

- 78-6287 **Mutagenicity of Chlorinated Phenoxyacetic Acids in *Saccharomyces cerevisiae* (Meeting Abstract).** (Eng) Zetterberg, G. (Dept. Genetics and Plant Breeding, Univ. Uppsala, S-750 07, Uppsala, Sweden). *Mutat Res* 53(2): 285-286; 1978. (2 refs)

- 78-6288 **Case Control Study: Malignant Mesenchymal Soft-Tissue Tumors and Exposure to Phenoxy Acids or Chlorophenols (Letter to Editor).** (Swe) Sandstrom, A. (Umea, Sweden) *Lakartidningen* 75(40):

Fifty-two men, aged 26-80 yr, with malignant mesenchymal soft-tissue tumors and 206 matched controls were surveyed by questionnaire for past occupational exposure to 2,4-dichlorophenoxyacetic acid (2,4-D) and/or 2,4,5-trichlorophenoxyacetic acid (2,4,5-T). The incidence of past exposure was 19/52 among the tumor patients vs 19/206 among the tumor-free controls. This difference was statistically significant ($p < 0.001$). It was not possible to evaluate the risks of 2,4-D and 2,4,5-T separately, because most patients may also have been exposed to other chemicals, such as tetrachlorodibenzodioxin. These findings suggest that exposure to phenoxy acids, chlorophenols, and/or their impurities increases the risk of soft tissue tumors. (5 refs)

- 78-6289 **The Chemistry of Phenalenium Systems. XX. Novel Ring Contraction During m-Chlorobenzoic Acid Oxidation of Mutagenic Azuleno[1,2,3-cd]phenalene. A Model for Metabolic Activation and Binding of Mutagenic Nonalternant Hydrocarbons.** (Eng) Nakasuji, K. (Dept. Chemistry, Faculty Science, Osaka Univ., Toyonaka, Osaka 560, Japan); Nakamura, Murata, I. *Tetrahedron Lett* (18): 1539-1542; 1978.

The reaction of m-chloroperbenzoic acid with azuleno[1,2,3-cd]phenalene (AP), a highly mutagenic, nonalternant isomer of benzo(a)pyrene, was investigated. The products formed were 11-oxo-11H-benzo[cd]fluoranthene and 1-oxo-1H-benzo[cd]fluoranthene. This reaction may provide a model for the mutagenic and carcinogenic activity of AP. (13 refs)

- 78-6290 **Screening of Safrole, Eugenol, Their Ninhydrin Positive Metabolites and Selected Secondary Amines for Potential Mutagenicity.** (Eng) Green, N. (Dept. Food and Nutrition, Coll. Home Economics, Florida State Univ., Tallahassee, FL, 32306); Savage, J. R. *Mutat Res* 57(2): 115-121; 1978.

The mutagenicity of safrole (4-allyl-1,2-methylenedioxybenzene), eugenol, the secondary amines with which they combine during metabolism, and the ninhydrin positive urinary metabolites of safrole and eugenol was tested in the direct bacterial assay, a microsomal mutagenesis assay (MMA), and a host-mediated assay. In the direct bacterial assay, in which four mutant strains of *Salmonella typhimurium* (TA1538, TA1531, TA1532, and TA1964) were used, all the compounds gave negative results. With the same four mutant strains, safrole and its metabolite 3-piperidyl-1-(3',4'-methylenedioxyphenyl)-1-propanone were mutagenic with strains TA1530 and TA1532 in the MMA and with strains TA1950 and TA1952 in the host-mediated assay. Dimethylamine was a weak mutagen in the MMA with strain TA1538. Negative results were obtained for the safrole metabolites 3-N,N-dimethylamino-1-(3',4'-methylenedioxyphenyl)-1-propanone and 3-pyrrolidinyl-1-(3',4'-methylenedioxyphenyl)-1-propanone, eugenol, the eugenol metabolites 3-piperidyl-1-(3'-methoxy-4'-hydroxyphenyl)-1-propanone and 3-pyrrolidinyl-1-(3'-methoxy-4'-hydroxyphenyl)-1-propanone, piperidine, piperidine acid, proline, and pyrrolidine in all three assay systems. These results suggest that the MMA is more sensitive than the host-mediated assay. (16 refs)

- 78-6291 **Conversion of the N-O-Glucuronide and N-O-Sulfate Conjugates of N-Hydroxyphenacetin to Reactive Intermediates.** (Eng) Mulder, G. J. (Lab. Chemical Pharmacology, Natl. Heart, Lung and Blood Institute, NIH, Bethesda, MD, 20014); Hinson, J. A.; Gillette, J. *Biochem Pharmacol* 27(12): 1641-1649; 1978.

stability and reactions of the reactive intermediates formed from phenacetin (PA) N-O conjugates were studied. N-O-glucuronide of ^{14}C -acetyl-N-hydroxy-PA is sufficiently stable for it to be purified but slowly degrades in aqueous solution to a reactive intermediate that can covalently bind to protein. Several lines of evidence are given that indicate the major chemically reactive intermediate formed from glucuronidation of N-hydroxy-PA is the deethylated reactive, N-acetylimidoquinone. However, the data suggest the formation of PA and 2-hydroxy-PA glucuronide occurs by different mechanisms. The N-O-sulfate of ^{14}C -acetyl-N-hydroxy-PA also breaks down to a reactive intermediate that has properties similar to those of the reactive intermediate formed from the N-O-glucuronide and thus may also be N-acetylimidoquinone. In contrast, the relative ability of various nucleophiles to prevent the covalent binding of the reactive intermediate formed from the N-O-sulfate of N-acetylaminofluorene to protein differs from the relative ability of the nucleophiles to prevent the covalent binding of the reactive intermediate of the N-O-sulfate or the N-O-glucuronide of PA, suggesting that the relative rates at which these intermediates combine with the different macromolecules may differ markedly. These studies may be relevant to other pathways of activation of PA and acetaminophen. Although the mechanism of PA-induced nephrotoxicity is not known, it seems plausible that PA N-O-glucuronide may play a role, since it would be produced primarily in the liver and concentrated in the kidney. Its conversion to a reactive intermediate, possibly accelerated by the low pH of urine, could lead to covalent binding and the renal tubular damage associated with PA abusers. (22 refs)

78-6292 **Benzo(a)pyrene Metabolism and Plasma Elimination Rates of Phenacetin; Acetanilide and Theophylline in Man.** (Eng) Kellermann, G. (Dept. Human Toxicology, Univ. Wisconsin Medical Sch., Madison, WI); Layten-Kellermann, M. *Pharmacology* 17(4): 191-200; 1978.

The plasma elimination rates of phenacetin (PA), acetanilide (AA), and theophylline (TP) were determined in 34 healthy subjects in a search for drugs whose metabolism and regulation might resemble those of benzo(a)pyrene (BP). The plasma half-lives of PA ranged from 32.0 to 88.2 min, those of AA from 70.5 to 186.5 min, and those of TP from 5.1 to 10.4 min. The metabolic clearance rate of PA varied from 11.6 to 144.9 liters/hr, of AA from 7.5 to 18.2 liters/hr, and of TP from 55.8 to 117.0 ml/hr. Statistically significant correlations were found between (a) the half-lives of PA, AA, and TP and the metabolic clearance rates of AA and TP, and (b) the inducibility of aryl hydrocarbon hydroxylase in mitogen-stimulated lymphocytes; the metabolic clearance rates of AA did not correlate. Correlations of the three drugs with the half-lives and metabolic clearance rates of antipyrine (determined previously) were equally good. In previous studies, antipyrine metabolism correlated best with BP metabolism. In this study, antipyrine half-lives and metabolic clearance rates were most closely correlated with TP half-lives and

clearance rates. The high correlations between antipyrine and PA half-lives contrast with the findings of other studies. (54 refs)

78-6293 **Analgesic Nephropathy and Transitional Cell Carcinoma.** (Eng) Moshakis, V. (Queen Mary's Hosp. for East End, West Ham Lane, Stratford, London E15 4SD, England); Hooper, A. A. *Postgrad Med J* 54(630): 285-286; 1978.

A transitional cell carcinoma of the kidney was found in a 42-yr-old woman who had taken an average of six Codeine compound tablets daily since age 13. This represented a total of 14 kg phenacetin over 26 yr. (4 refs)

78-6294 **Long-Term Safety Evaluation and Metabolism of o-Benzoin Sulfimide (Saccharin) in Rhesus Monkeys.** (Eng) Coulston, F. (Inst. Comparative and Human Toxicology, Albany Medical Coll., Albany, NY, 12208); McChesney, E.; Benitz, K. F. *Ecotoxicol Environ Saf* 2(1): 33-48; 1978.

The results of a 79-mo safety evaluation of saccharin in 7 male and 7 female monkeys are reported. Saccharin was administered by stomach tube 6 days/wk at daily doses of 20 mg/kg (4 animals), 100 mg/kg (4 animals), and 500 mg/kg (6 animals). Six monkeys of comparable age and wt served as controls. Although there were five deaths, one in each treatment group and two in the control group, none of the deaths was related to saccharin intake. The dose regimens had no effect on animal growth or on serum biochemical or whole blood hematological parameters. At necropsy, the only lesions detected grossly or microscopically occurred as frequently in controls as in any of the treated groups. Moreover, the suspected target organ, the urinary bladder, was free of lesions in all groups. Metabolic experiments demonstrated that saccharin was eliminated rapidly (estimated half-life of 9.5 hr), mostly in the urine, and in at least 98% unchanged form. It is concluded that the no-effect level of saccharin for the rhesus monkey exceeds 500 mg/kg/day, a dose that, on a wt basis, is 50 times the recommended max daily human intake. (31 refs)

78-6295 **Effect of DNA Repair Modifying-Factors and Tumor-Promoting Agents on Mutagenesis in a Mammalian Cell Culture System.** (Eng) Lankas, G. R. (Univ. Cincinnati, Cincinnati, OH, 45221). *Diss Abstr Int B* 39(5): 2203-2204; 1978. (no refs)

78-6296 **Aryl Hydrocarbon Hydroxylase Induction in an Insect, *Spodoptera eridania* (Cramer), by Poly-**

chlorinated Biphenyls (PCBs). (Eng) Anderson, R. S. (Donald S. Walker Lab., Sloan-Kettering Inst. Cancer Res., 145 Boston Post Road, Rye, NY, 10580). *Comp Biochem Physiol [C]* 60(1): 51-55; 1978.

Commercial polychlorinated biphenyl (PCB) preparations were tested for their ability to induce aryl hydrocarbon hydroxylase (AHH) activity in terminal instar southern armyworm (*Spodoptera eridania*) midgut homogenates or microsomes. The preparations included Capacitor 21, Aroclor 1016, Aroclor 1242, Aroclor 1254, and several PCB-containing immersion oils. Topical application of PCB's in doses of 2-3 mg induced significant AHH activity, as measured by increased ¹⁴C-benzo(a)pyrene substrate metabolism. The induction levels were not proportional to the chlorine content of a given PCB compound. Max induction occurred 24 hr after exposure, and the effect persisted for at least 4 days after a single application of Aroclor 1254. *Spodoptera* homogenate and protein concentrations did not increase within 48 hr of treatment, which suggested that AHH induction by PCB's precedes proliferation of the smooth endoplasmic reticulum. (20 refs)

78-6297 Effect of Intestinal Microflora on 3,2'-Dimethyl-4-aminobiphenyl-induced Carcinogenesis in F344 Rats. (Eng) Reddy, B. S. (Naylor Dana Inst. Disease Prevention, American Health Foundation, 1 Dana Rd., Valhalla, NY, 10595); Watanabe, K. *J Natl Cancer Inst* 61(5): 1269-1271; 1978.

The effect of intestinal microflora on colon and breast cancer induced by 3,2'-dimethyl-4-aminobiphenyl (DMAB) was studied in germfree (22 male, 22 female) and conventional (30 male, 30 female) F344 rats. At 7 wk of age, all animals were started on a series of DMAB injections (100 mg/kg/wk x 20, sc). Females were autopsied at 10 wk, males at 15 wk. Tumors were induced in the colons (24/60), duodena (20/60), breasts (24/30 females), ear ducts (9/60), salivary glands (6/60), and skin (36/60) of conventional rats and in the colons (9/44), breasts (9/22 females), ear ducts (6/44), salivary glands (10/44), and skin (4/44) of germfree rats. There was no consistent difference in the incidence of ear duct, salivary gland, and skin tumors between the germfree and conventional rats. However, none of the germfree rats had duodenal tumors, and the incidence of colon and breast cancer was lower in germfree than in conventional rats. In conventional rats, both bacterial and mucosal β -glucuronidase is involved in hydrolyzing the glucuronide conjugate of DMAB, whereas in germfree rats, only mucosal β -glucuronidase is involved. This might explain the difference in colon cancer susceptibility. The possibility that the lower incidence of DMAB-induced colon and breast cancer in germfree rats might be due to their lower hepatic levels of DMAB-metabolizing enzymes cannot be ruled out. (19 refs)

78-6298 A Sequential Screening of the Cytogenetic Damage Induced by Triaziquone. (Eng) Oud, J.

L. (Inst. Genetics, Univ. Amsterdam, Kruislaan 318, Amsterdam, Netherlands); Peters, P. W. *Mutat Res* 54(1): 175-184; 1978.

The suitability of six in vivo cytogenetic tests for determining chromosome damage and the mutagenicity of the alkylating agent triaziquone (TAQ) in young adult random-bred Swiss/RIV mice was determined. The mice were inoculated either once or twice (24-hr interval) with 0.125 mg/TAQ. Two injections increased the number of aberrations considerably in both the micronucleus test and the bone marrow metaphase test, but not in the spermatocyte translocation test or the spermatogonial metaphase test. In the latter, a strong cell-killing effect of TAQ was noted. Analysis of whole mounts of seminiferous tubules showed that 0.125 mg/kg was a lethal dose for all B- and intermediate-type spermatogonia and partly killed A-type spermatogonia. A single dose caused stable chromosome rearrangements in spermatids that could be demonstrated with the F₁ translocation test, and it gave rise to dominant lethality of fetuses originating from postmeiotic sperm. The micronucleus test, the results of which are similar to those of the bone marrow test, is concluded to be a reliable and easy screening method for chromosome aberrations in vivo. The spermatogonial metaphase test can be used to determine the capability of a compound to reach germ cells in seminiferous tubules. The spermatocyte translocation test is informative about stable aberrations in a permanent cell type, and the F₁ translocation and dominant lethal tests reveal the induction of inheritable chromosome rearrangements in spermatids and spermatozoa. (20 refs)

78-6299 Carcinogenic Compounds: Synthesis of Polycyclic Carbazole Derivatives by Diels-Alder Reaction. (Eng) Hiremath, S. P. (Dept. Chemistry, Karnataka Univ. Post Graduate Centre, Gulbarga-585 105, India); Kaddargi, S. S.; Purohit, M. G. *J Indian Chem Soc* 55(2): 156-157; 1978.

Polycyclic benzocarbazoles were synthesized by Diels-Alder reaction of 3-(β -nitrovinyl)benz(e)indole with various dienophiles such as 1,4-benzoquinone, 1,4-naphthoquinone, N-phenylmaleimide, dimethyl acetylenedicarboxylate, and tetracyanoethylene. These compounds are structurally similar to polycyclic hydrocarbons, and they have shown carcinogenic activity. (13 refs)

78-6300 The Tumor-producing Effect of Automobile Exhaust Condensate and Fractions Thereof. Part 1: Chemical Studies. (Eng) Grimmer, G. (Biochemisches Institut für Umweltcarcinogene, Sieker Landstrasse 19, 2070 Ahrensburg, W. Germany); Bohnke, H. *J Environ Pathol Toxicol* 1(5): 661-667; 1978.

A detection system designed to discriminate between the carcinogenic and noncarcinogenic fractions of a mixture must

onstrate the following: (1) a reproducible dose/effect relation of the carcinogen over a large range; (2) additive effects of two or more carcinogens; and (3) no or negligible influence of large amounts of ineffective substances on the carcinogenicity of the carcinogen in a complex mixture. The key for the detection of all carcinogens from a complex mixture and the estimation of their efficiencies is a suitable animal test system. Automobile exhaust gas condensate was collected and separated into polycyclic aromatic hydrocarbon (PAH)-containing and PAH-free fractions by liquid-liquid distribution between methanol + water + cyclohexane and cyclohexane + nitromethane, respectively. Carbazol and indole derivatives as well as aromatic amines remained predominantly in the PAH-free methanolic phase. Tests involving topical application to the skin of CFLP mice showed that the N-containing and other water-soluble carcinogens do not contribute significantly to carcinogenicity. The results therefore support the assumption that the carcinogenic effect of automobile exhaust condensate is due to the PAH content. (13 refs)

78-6301 Simultaneous Gas-Chromatographic Profile Analysis of Carcinogenic Polycyclic Aromatic Compounds: Polycyclic Aromatic Hydrocarbons, Carbazoles, Acridines/Aromatic Amines. (Eng) Grimmer, G. (Bismisches Institut für Umweltcarcinogene, Sieker Landstrasse 19, 2070 Ahrensburg, W. Germany); Bohnke, H.; Ujjack, K. W. *Fresenius Z Anal Chem* 290(2): 147; 1978.

termination of the carcinogenic risk of polycyclic aromatic compounds to humans by a simultaneous gas-chromatographic procedure is outlined. (6 refs)

78-6302 Importance of Inactivating Mechanisms in Chemical Mutagenesis (Meeting Abstract). (Eng) Oesch, F. (Section Molecular Pharmacology, Inst. Pharmacology, Univ., Obere Zahlbacher Strasse 67, D-6500 Mainz, W. Germany). *Mutat Res* 53(2): 244-245; 1978. (no refs)

78-6303 Apparent and Real Thresholds: A Study on Two Mutagens (Meeting Abstract). (Eng) Seiler, J. (Swiss Federal Res. Station, CH-8820 Wädenswil, Switzerland). *Mutat Res* 53(2): 260-261; 1978. (no refs)

78-6304 Genetic Activity of Lead Chromate in Bacteria and of Actinomycin D in Yeast (Meeting Abstract). (Eng) Nestmann, E. R. (Environmental Mutagenesis Section, Dept. Natl. Health and Welfare, Ottawa, Canada). *Can J Genet Cytol* 20(3): 451; 1978. (1 ref)

78-6305 Persistent Pesticides and Environmental Chemicals Are Taken Up to a Greater Extent than Non-Persistent Compounds by Cultured Human Cells. (Eng) Murakami, M. (Inst. Physical Chemical Res., Wako-shi, Saitama 351, Japan); Fukami, J. *Bull Environ Contam Toxicol* 19(4): 423-427; 1978.

Data on the cellular uptake of 11 pesticides by cultured human embryonic lung diploid cells are presented. The pesticides dichlorodiphenyltrichloroethane (DDT), dieldrin, aldrin 2,4,5,2',5'-hexachlorobiphenyl (PCB), hexachlorobenzene (HCB), carbaryl, malathion, parathion, chlorodimeform, 2,4,5-trichlorophenoxy-1-acetic acid (2,4,5-T), and 2,4-dichlorophenoxy-1-acetic acid (2,4-D) showed the following uptake levels (expressed as picomoles/mg cell protein) at 48 hr of incubation: 4,830, 4,380, 3,640, 1,740, 778, 945, 180, 22.6, 32.4, 9.9, and 3.7, respectively. Thus, persistent pesticides such as DDT, dieldrin, aldrin, PCB, and HCB were taken up by cells to a greater extent than nonpersistent pesticides. The considerable amount of carbaryl taken up by the cells suggests that the naphthalene ring in the carbaryl molecule may be involved in the incorporation potential of this compound. The effect of the above pesticides on cell growth was also investigated; most compounds were not toxic at the same concentration (4×10^{-6} M) and over the same experimental period as in the uptake experiments. The herbicides 2,4-D and 2,4,5-T, although less incorporated than the insecticidal and industrial chemicals, appeared to be more toxic. It is evident that differences in uptake are not the cause of the different degrees of cytotoxicity of the chemicals. (13 refs)

78-6306 Tumor-promoting Phorbol Esters Inhibit Binding of Epidermal Growth Factor to Cellular Receptors. (Eng) Lee, L. S. (Inst. Cancer Res., Coll. Physicians and Surgeons, Columbia Univ., New York, NY, 10032); Weinstein, I. B. *Science* 202(4365): 313-315; 1978.

Tumor-promoting phorbol esters and related plant diterpenes were tested for their ability to inhibit the binding of ^{125} I-labeled epidermal growth factor (EGF) to HeLa cells. Extremely low concentrations [0.25-50 nanograms (ng)/ml] of 12-O-tetradecanoylphorbol-13-acetate (TPA) and related compounds produced a dose-dependent decrease in ^{125} I-EGF binding. At 25 ng/ml (4.2×10^{-8} M), TPA inhibited the binding of ^{125} I-EGF by 90%. In decreasing order of effectiveness, the test compounds that inhibited EGF binding were TPA, merzerein, gnilatimacrin, gnidilatin, phorbol-12,13-didecanoate (PDD), and phorbol dibenzoate. Inhibition occurred when the test agent was added 90 min before or simultaneously with EGF. Phorbol, 4 α -phorbol-12,13-didecanoate, 4-O-methyl-12-O-tetradecanoylphorbol-13-acetate, gnidipalmin, and anthralin did not have an appreciable effect on ^{125} I-EGF binding when tested at concentrations up to 100 ng/ml. The inhibitory effect, therefore, occurs at concentrations as low as 10^{-8} - 10^{-9} M and possesses marked structural specificity and stereospecificity. These re-

sults also parallel the structural requirements for promoting activity in the two-stage mouse skin carcinogenesis assay. Inhibition of binding appears to be due to a decrease in the number of available EGF receptors rather than to a change in receptor affinity. These results suggest that certain biologic effects of tumor promoters may result from alterations in the function of cell-surface receptors involved in growth regulation. (48 refs)

- 78-6307 Blocking Tumor Promotion by Protease Inhibitors.** (Eng) Troll, W. (Dept. Environmental Medicine, New York Univ. Medical Center, New York, NY, 10016); Levitz, M. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1245-1250; 1978.

The influence of protease inhibitors on tumor promotion in mouse skin and the association between estrogens and plasminogen activation in mouse uterus were investigated. Treatment of mouse skin with 1 μ g phorbol myristate acetate (PMA) resulted in the appearance of appreciable protease activity (35 fluorescence units/mg protein/hr); this activity was inhibited by tosyllysine chloromethyl ketone, a protease inhibitor. In skin tumor studies, tumor appearance was delayed 50-100 days by the application of protease inhibitors 3x/wk. Tumors initiated on the skin of female mice with 7,12-dimethylbenz(a)anthracene and promoted 2 wk later with croton oil were suppressed by the application of glucocorticoids (dexamethasone, Schering No. 11572, prednisolone, hydrocortisone, and cortisone) at 6 and 30 μ g, 5x/wk. Tissue culture studies indicated that dexamethasone elaborates a protease inhibitor. The association between estrogens and plasminogen activation was investigated by treating ovariectomized young mice with 0.1 μ g estradiol bid for 3 days. This resulted in the appearance of plasminogen activator specifically localized in the 12,000 g pellet, perhaps bound to the plasma membrane, and the nucleus. Inoculation of normal cycling female mice with a mixture of the protease inhibitors Antipain (3.0 mg) and Leupeptin (20 mg) for 15 days caused a significant diminution in uterine wt, total DNA, and total protein. The role of plasminogen activator may be at an early stage of tumor promotion; hormones may act in a similar manner. (12 refs)

- 78-6308 The Nucleotide-Permeable *Escherichia coli* Cell, a Sensitive DNA Repair Indicator for Carcinogens, Mutagens, and Antitumor Agents Binding Covalently to DNA.** (Eng) Thielmann, H. W. (Deutsches Krebsforschungszentrum, Institut für Biochemie, Im Neuenheimer Feld 280, D-6900 Heidelberg, West Germany); Gersbach, H. *Z Krebsforsch* 92(2): 177-214; 1978.

The use of ether-permeabilized (nucleotide-permeable) *Escherichia coli* cells in assay of capability for DNA-binding and stimulation of excision repair of 11 ultimate carcinogens, 10 proximate carcinogens, 2 tumor promoters, 6 mutagens, and 12 antitumor agents was investigated. The strains *uvrA* and *uvrB* used in the study were the wild types, H 540, KMBL 1789, and the mutants *uvrA* (H 512) and *uvrB* (H 551). All the ultimate carcinogens (alkylating, alkylating, arylalkylating agents) and the mutagens such as hydrogen peroxide and acridine derivatives caused DNA excision repair in wild type cells and simultaneously inhibited replication DNA synthesis to various degrees. Proximate carcinogens such as dimethyl nitrosamine and the aflatoxins require activation for DNA-binding did not induce excision repair. Intercalating mutagens such as quinacrine and ethidium bromide inhibited replicative DNA synthesis, however they were not repair inducers. The tumor promoters 12-O-tetradecanoyl-phorbol-13- and phorbol-12,13-didecanoate did not cause excision repair, even when applied in high concentrations, nor did they inhibit excision repair synthesis stimulated by N-methyl-N-nitrosourea or N-acetoxy-2-acetylaminofluorene. The antitumor agents were classified in two groups on the basis of their effect on DNA synthesis: members of the first group known to bind covalently to DNA [bleomycin, N-bis(2-chloroethyl)-N-nitrosourea, etc] stimulated excision repair polymerization and inhibited DNA replication; the other group of antitumor agents, known to react with DNA by noncovalent binding [*cis*-Pt(II) diamine complexes, adriamycin, etc] inhibited DNA replication without stimulating repair synthesis. It is concluded that the ether-permeabilized *E. coli* cell can be successfully used as assay for DNA repair-inducing and replication-inhibiting activity. (110 refs)

- 78-6309 Effect of Tumor Promoters on the Response of Cultured Embryonic Chick Ganglia to Nerve Growth Factor.** (Eng) Ishii, D. N. (Dept. Pharmacology, Coll. Physicians and Surgeons, Columbia Univ., New York, NY, 10032). *Cancer Res* 38(11, part 1): 3886-3893; 1978.

The effects of the tumor-promoting agent 12-O-tetradecanoylphorbol-13-acetate (TPA) on nerve growth factor (NGF)-induced neurite outgrowth (NOG) were studied in cultured chick embryonic ganglia. There was no detectable NOG in the absence of NGF, but in its presence [1 nanogram(ng)/ml], a max response was attained in 24 hr and sustained for at least 48 hr. The addition of TPA (100 ng/ml, 1.7×10^{-7} M) to the culture medium inhibited further NOG for 24 hr, after which there was a gradual increase in outgrowth, even in the presence of fresh TPA. Neurites formed in the presence of NGF (10 ng/ml) + TPA (200 ng/ml) were shorter, thicker, and sparser than those of NGF-treated cultures. In cytotoxicity tests, TPA alone substantially enhanced the survival of ganglion cells. TPA + NGF appeared to increase the number of surviving cells over that produced with either agent alone. Thus, TPA is not cytotoxic to ganglion cells at concentrations that antagonize NGF. Other plant di-

enes that are tumor promoters in the mouse skin carcinogenesis system (phorbol-12,13-didecanoate, ingenol dibenzoate, and mezerein) also inhibited the ganglia response to NGF, but structurally related nonpromoting compounds were inactive. The concentration of NGF that induced a 50% max NOG response in sensory ganglia was 0.6×10^{-11} M. The antagonism of NGF by TPA was dose-dependent and noncompetitive. TPA concentrations that extensively inhibited NOG did not affect the amount of ^{125} I-labeled NGF bound to specific sites on dissociated dorsal root ganglion neurons, supporting the theory that TPA acts at a stage beyond the initial interaction of the factor with its receptive site. (51 refs)

6310 Specificity of Response in Hamster Cells Induced to Produce Plasminogen Activator by the Tumor Promoter, 12-O-Tetradecanoylphorbol-13-acetate. (Eng) Christman, J. K. (Dept. Biochemistry, Mount Sinai Medical Center, New York, NY, 10029); Copp, R. P.; Pedriotti, L.; Whalen, C. E. *Cancer Res* 38(11, part 1): 3854-3860; 1978.

The effect of the tumor-promoting agent 12-O-tetradecanoylphorbol-13-acetate (TPA) on the form of plasminogen activator (PA) produced by cells in culture was studied using several hamster cell lines. TPA enhanced PA production only in hamster cells that synthesize a PA that is immunologically identical to that of normal hamster lung cells. No enhancement of PA production was detected in hamster cells (kidneys, ovary, melanoma, or glioma) that synthesize a PA of different antigenic form. Neither the tissue of origin nor the transforming agent served as an absolute determinant of the form of PA produced. Two of three simian virus 40 (SV40)-transformed early-passage Chinese hamster lung cells, a subclone of the DON hamster lung line selected for azaguanine resistance, and one of two SV40-transformed hamster kidney lines produced the lung-type PA. Cells of the third SV40-transformed lung culture produced a PA distinct from that of normal hamster lung cells. In the presence of TPA, however, this line synthesized PA of the lung form. Thus, TPA was able to induce the reexpression of at least one parameter of the original phenotype, synthesis of the lung form of PA. These results suggest that the two antigenically distinct PA's are separate gene products. The relationship, if any, between the ability of TPA to increase PA production and its ability to promote tumor formation in mouse skin is not known. (46 refs)

78-6311 Distribution of 3-Hydroxyanthranilic Acid in Mice. (Eng) Watanabe, M. (Dept. Analytical Chemistry, Faculty Pharmaceutical Sciences, Teikyo Univ., Yagami, Kanagawa-ken 199-01, Japan). *Gann* 69(4): 585-587; 1978.

Male (BALB/c x DBA/2)F₁ mice were inoculated sc with

^{3} H-hydroxyanthranilic acid (^{3} H-HAA: 2 mg, 748 μ Ci) and, 30 and 60 min and 6 hr later, the distribution of ^{3} H was observed by whole-body autoradiography and radioactivity measurements, and metabolites in organs and urine were measured by paper and thin-layer chromatography. At 30 and 60 min after injection, the radioactivity was found mainly in the liver and kidneys; it was distributed almost uniformly in other organs. After 6 hr, the radioactivity remained in the kidneys and lymphoid organs. However, at 30 min and 6 hr after injection, neither HAA nor any carcinogenic metabolites could be detected in the liver, kidneys, spleen, or lymph nodes. Considerable amounts of HAA and its sulfate were found only in urine. These results suggest that HAA is metabolized rapidly and does not accumulate in a specific organ. However, urinary organs may be at risk of tumor development because of the rapid excretion of most of the compound in urine. (8 refs)

78-6312 Studies on the Hepatic Microsomal Metabolism of [^{14} C]Phenanthrene. (Eng) Chaturapit, S. (Dept. Pharmacy, Univ. Sydney, New South Wales 2006, Australia); Holder, G. M. *Biochem Pharmacol* 27(14): 1865-1871; 1978.

The metabolism of ^{14} C-phenanthrene by liver microsomes from Wistar rats, SW and DBA/2J mice, and guinea pigs was examined using untreated, sodium phenobarbital (PB)-pretreated, and 3-methylcholanthrene (3-MC)-pretreated animals. Metabolite profiles were obtained by comparing the thin-layer chromatographs of the reaction products with chromatographs of authentic reference compounds. Trans-Dihydrodihydroxyphenanthrenes were the major metabolites in all preparations. With all reactions except for the one in which microsomes from 3-MC-pretreated guinea pigs were used, trans-9,10-dihydro-9,10-dihydroxyphenanthrene predominated. With microsomes from the 3-MC-pretreated guinea pigs, phenanthrene 1,2-dihydrodiol was the major metabolite. Addition of cyclohexene oxide (5.0 mM) to the incubation mixture inhibited dihydrodiol formation in rat and mouse preparations by 90% and in guinea pig preparations by 70%. Phenols and phenanthrene 9,10-oxide were produced instead. Compared with control microsomes, increases in phenanthrene metabolism occurred with microsomes from all PB-pretreated animals and with microsomes from 3-MC-pretreated rats and SW mice, a pattern similar to that seen for the induction of aryl hydrocarbon hydroxylase. (43 refs)

78-6313 Excision of Hydrocarbon-DNA Adducts and Consequent Cell Survival in Normal and Repair Defective Human Cells. (Eng) McCaw, B. A. (Chemical Carcinogenesis Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501); Dipple, A.; Young, S.; Roberts, J. *J. Chem Biol Interact* 22(2/3): 139-151; 1978.

An attempt was made to demonstrate selectivity of excision in normal human fibroblasts, xeroderma pigmentosum (XP) variants that are not deficient in excision repair, and in partially excision-repair-deficient XP fibroblasts. Survival curves for UV-irradiated normal human cells and XP variant cells (XPB4BE) were indistinguishable. In comparison, cells from XP complementation group A (XP12BE) were very sensitive to UV radiation. Complementation group C (XP2BE) cells were almost as sensitive as group A cells. These survival phenomena paralleled the known unscheduled DNA synthesis responses of these cells to UV radiation, which, compared with normal cells, are: XP4BE, 100%; XP2BE, 20%; XP12BE, 2%. The relative capacities of these cells to excise 7-bromomethylbenz(a)anthracene (BMBA)-DNA adducts and to survive treatment with the carcinogen were similar to the responses to UV radiation, except that the XP2BE cell line excised and survived this damage far better than anticipated from its response to UV radiation. Moreover, although the ratio of hydrocarbon-adenine adduct to hydrocarbon-guanine adduct remaining in DNA decreased notably with excision in normal and variant cells, this ratio did not change significantly with excision in XP2BE cells. The relationship between greater excision capacity and increased cell survival in the experiments with BMBA indicates that the unexcised damage is responsible for the cell-killing action of this agent. The different relative repair and survival responses of these cell lines to UV radiation and to BMBA indicate that in at least one of these cell lines (XP2BE), and possibly all the lines, different cellular mechanisms are involved in the repair of DNA damage resulting from UV radiation and that resulting from chemical carcinogen treatment. (22 refs)

78-6314 Cytogenetic Mapping of the Trisomic Segment of Chromosome 15 in Murine T-Cell Leukemia.

(Eng) Wiener, F. (Dept. Tumor Biology, Weizmann Inst. Science, Rehovot, Israel); Ohno, S.; Spira, J.; Haran-Ghera, N.; Klein, G. *Nature* 275(5681): 658-660; 1978.

A chromosome segment located distal to the T6 breakpoint of chromosome 15 was found to be duplicated in 7,12-dimethylbenz(a)anthracene-induced T-cell leukemia in CBAT6T6 mice. These leukemias are associated with trisomy 15, a non-random cytogenetic change also associated with murine T-cell leukemia induced by viruses and x-rays. The induced leukemias were transplanted sc into recipient mice of the opposite sex, and G-banded metaphase plates were prepared from sc tumors, spleens, and bone marrow. Five of six leukemias were aneuploid, with 41 or 42 chromosomes. All the aneuploid leukemias had a trisomy T(14;15)6. The sixth tumor, however, had a banding pattern indistinguishable from that of a normal CBAT6T6 cell. The consistency of the trisomic change emphasizes the specific role of genes associated with chromosome 15 in T-cell leukemia. However, the occurrence of a diploid leukemia in one tumor shows that the change is not exclusive. (9 refs)

78-6315 Chemical Carcinogenesis in C3H/StWi Mice: a Worthwhile Experimental Model for Breast Cancer. (Eng) Smith, G. H. (Lab. Molecular Biology, NIH, Bethesda, MD, 20014); Pauley, R. J.; Socher, S. H.; Meeker, D. *Cancer Res* 38(12): 4504-4509; 1978.

To determine whether the low incidence of mammary tumors in C3H/StWi mice is due to the fact that low-oncogenic mouse mammary tumor virus MMTV(L) is not expressed in these mice, the effects of estrogenic stimulation and 7,12-dimethylbenz(a)anthracene (DMBA) on mammary tumor incidence were examined in C3H/StWi and (BALB/c x C3H/StWi)F₁ mice. None of 18 virgin C3H/StWi females in which 20% diethylstilbestrol (DES) pellets were implanted sc at 5-6 wk of age developed mammary cancer throughout their life-span, which averaged 24.5 mo. Only 2/24 hybrid mice similarly treated developed mammary cancer. In contrast, 17/37 virgin C3H/StWi mice treated with DMBA (50 mg/wk, intragastrically) over a 6-wk period developed mammary tumors at an average of 7.8 mo (range, 4.5-10 mo). Tumors were mainly papillary adenocarcinomas. Mammary dysplasias were found in 73% of the whole-gland mounts examined at 10 mo (3.6 mice). All but three dysplasias were ductal in origin. No hyperplastic alveolar nodules were found. Electron microscopic examination of 14 DMBA-induced mammary tumors failed to reveal the presence of A- or B-type particles, but 11 tumors contained budding or extracellular C-type virions. Evaluation of five tumors for MMTV-specific RNA sequences indicated levels as low or lower than those found in the mammary glands of retired multiparous C3H/StWi females (0.0035%). Much higher levels (0.2%-0.4%) of MMTV-specific RNA to total RNA occurred in MMTV-induced C3H/StWi mammary tumors. Thus, the C3H/StWi subline is different from other C3H sublines in view of its low mammary cancer incidence, low response to estrogenic stimulation, and the fact that it does not produce MMTV(L). It may represent a unique model for the study of viral, chemical, and viral-chemical interactions in the malignant transformation of mammary epithelial cells. (21 refs)

78-6316 A Technique for Localization of Tumours in Hamster Cheek-Pouch Carcinogenesis. (Eng)

MacDonald, D. G. (Dept. Oral Medicine and Pathology, Glasgow Dental Hosp. and Sch., 211 Renfrew St., Glasgow G2 3JZ, Scotland). *Arch Oral Biol* 23(7): 573-577; 1978.

A technique was developed to enable the area of exposure to a carcinogen and of subsequent tumor growth to be predetermined and restricted in the hamster cheek pouch. Thirty male Syrian hamsters received either 0.5% or 0.25% 9,10-dimethyl-1,2-benzanthracene (DMBA) topically, 3x/wk for 6 wk in a restricted (1-cm²) anterior portion of both cheek pouches just posterior to the prominent blood vessels at the pouch mouth. At 15 wk posttreatment, all animals were sacrificed and examined surgically. Tumors were first evident at 5 wk posttreatment, and by time of biopsy at 8 wk, 11 animals had

tors in 13 pouches, but there was no significant difference between the two treatment groups with respect to tumor yield distribution. At sacrifice, 19 animals had developed tumors in 30 pouches, many at multiple sites. In 13 pouches, tumors were within the experimental area; in 9, they were outside this area; and in 8, tumors were within and without defined area. Again, there was no significant difference between the two treatment groups with respect to tumor yield distribution, but the 0.25% group showed significantly higher tumor localization in terms of pouches with tumors inside the experimental area only (8/9 pouches) compared to pouches with tumors outside this area only. This means localization allows sequential studies of carcinogenesis in defined area to be undertaken. (7 refs)

6317 **Azo Dyes Prevent Hydrocarbon-induced Leukemia in the Rat.** (Eng) Huggins, C. B. (Ben May Lab. Cancer Res., Univ. Chicago, Chicago, IL, 60637); La, N.; Russo, A. *Proc Natl Acad Sci USA* 75(9): 44527; 1978.

The azo dyes 1-(p-phenylazophenylazo)-2-naphthol (Sudan III) and 1-(o-tolylazo-o-tolylazo)-2-naphthol (Sudan IV) were investigated as possible protectors against leukemia induced by multiple injections of 7,8,12-trimethylbenz(a)anthracene (7,8,12-TMBA) in Long-Evans rats. Three groups of female rats were given four 35-mg/kg injections of 7,8,12-TMBA, iv, 2-wk intervals. Two groups were fed 1 mg of either dye 24 hr before each iv injection; controls were fed same oil. In rats fed Sudan III, leukemia incidence was 60%, mammary cancer incidence 60%. Analogous figures for rats fed Sudan IV were 21% and 57%. Of control rats, 21% developed leukemia and 43%, mammary cancer. Three groups of male rats received 7,8,12-TMBA as above. Two groups received Sudan III at 48 and 24 hr prior to each injection, one (Group I) at a dose of 1 mg, and the other (Group II) at a dose of 10 mg. The incidence of leukemia in Group I was 6%, that in Group II, 0%. Controls developed leukemia at a rate of 92%. Daily feeding of 1 mg Sudan III to eight female rats for 4 days produced a progressive increase in the hepatic menadione reductase concentration, but the protein concentration was unchanged. A single iv injection of 35 mg/kg 7,8,12-TMBA produced a transient leukopenia, but no changes in WBC counts were seen in similarly treated rats protected with 1 mg Sudan III. All 16 female rats given a single injection of 75 mg/kg 7,12-DMBA expired in 1-3 days. In an identical group protected with 1 mg Sudan III 24 hr before DMBA injection, no rats died, although 50% eventually developed leukemia. Sudan III itself was not found to be carcinogenic. (18 refs)

6318 **Development of Experimental Brain Tumors under Amine Metabolism Modulation.** (Rus) Zumanis, E. A. (Dept. Pharmaceutical Chemistry, Riga

Medical Inst., Riga, USSR); Plaude, O. O.; Buikis, I. M. *Vopr Onkol* 24(9): 71-77; 1978.

9,10-Dimethyl-1,2-benzanthracene (DMBA)-induced brain tumor mortality and its relationship to brain serotonin levels were studied over a 10-mo period in Wistar rats with DMBA implanted into their cerebellum. The development of cerebellar tumors was accompanied by a depletion of the serotonin level in the cerebellum, but the level in the hypothalamus remained unchanged. An inverse correlation was found between DMBA-induced brain tumor mortality and serotonin levels. Topical administration of monoamine oxidase inhibitors before the precancerous stage (before day 120 after DMBA implantation) reduced tumor incidence and tumor mortality by reducing the number of astrocytomas and histogenetically related glioblastomas. The monoamine oxidase inhibitors had no effect on the malignant transformation of other neuroectodermal cells. The findings indicate that serotonin (the levels of which are increased by monoamine oxidase inhibitors) increases the resistance of astrocytes to malignant transformation. (21 refs)

78-6319 **Hemopexin and Aryl Hydrocarbon Hydroxylase Induction in the Mouse and in Cell Culture.**

(Eng) Owens, I. S. (Developmental Pharmacology Branch, Natl. Inst. Child Health and Human Development, NIH, Bethesda, MD, 20014); Nebert, D. W.; Liem, H. H.; Muller-Eberhard, U. *Biochem Pharmacol* 27(12): 1665-1669; 1978.

Hemopexin (HP) and aryl hydrocarbon hydroxylase (AHH) induction among inbred mouse strains and among offspring derived from appropriate genetic crosses was examined along with HP and AHH levels in several rat hepatoma-derived cell culture lines to determine whether an association exists between HP induction and the Ah allele. Treatment of three responsive mouse strains, C57BL/6N (B6), Balb/cAnN, and C3H/HeN, with a single dose (80 mg/kg, ip) of 3-methylcholanthrene (3-MC) or β -naphthoflavone (NF) increased plasma HP levels 71%-160%. The nonresponsive strains DBA/2N, NZB/BLN, and NZW/BLN had increases of 14% to 60%. In B6 mice, the greatest rise in HP levels occurred between 24 and 48 hr. AHH activity induced by 3-MC occurred during the first 24 hr. The increase in HP therefore appears to be subsequent to AHH induction by 3-MC. Administration to the mice of 0.05% phenobarbital in drinking water for up to 14 days induced similar levels of AHH activity in both responsive and nonresponsive mice. Again, the increase in HP followed AHH induction. AHH was not inducible in 6/18 3-MC treated (B6 x NZW)_{F2} offspring. In hepatoma cell lines grown in culture and treated with 13 μ M benz(a)anthracene (BA), HP production occurred in H-4-II-E but not in HTC or Hepa-1 cells, but a relationship between inducible AHH activity and plasma HP levels could not be inferred. HP appearance was equivalent in BA-treated and control H-4-II-E cultures, whereas AHH induction occurred primarily in the BA-treated cultures. Frequent remov-

al of tissue culture medium resulted in an increased secretion of intracellular HP to the surrounding medium. The reason for this effect is not known. (33 refs)

- 78-6320 The Topology of Epoxide Hydratase and Benzopyrene Monooxygenase in the Endoplasmic Reticulum of Rat Liver.** (Eng) Seidegard, J. (Arrhenius Lab., Dept. Biochemistry, Univ. Stockholm, Fack, S-106 91 Stockholm, Sweden); Moron, M. S.; Eriksson, L. C.; DePierre, J. W. *Biochim Biophys Acta* 543(1): 29-40; 1978.

The distribution of benzo(a)pyrene monooxygenase and epoxide hydratase, in the liver microsome subfractions of control and of phenobarbital (PB)- and methylcholanthrene(MC)-treated rats was investigated. Male Sprague-Dawley rats were injected ip with 20 mg/kg MC on days 5, 3, and 1 before sacrifice or with 80 mg/kg PB on days 5-1 before sacrifice. The specific activities of these enzymes in the rough and smooth microsomes from control and PB-treated rats were approx equal. After treatment with MC, however, the activity of benzo(a)pyrene monooxygenase was four times higher and that of epoxide hydratase two times higher in the rough vesicles. Differential centrifugation revealed that both enzymes were dispersed uniformly throughout the microsomal vesicles. It is concluded that these two enzyme systems may form a complex of unique stoichiometry in the microsomal membranes from control rats, not in those from MC-treated rats. Studies with proteases and diazobenzene sulfonate suggest that epoxide hydratase is located deep within the lipid phase of the membrane of the hepatic endoplasmic reticulum. (31 refs)

- 78-6321 An Experimental Approach to the Breast Cancer, Reserpine Problem.** (Eng) Homburger, F. (Bio-Res. Inst., 9 Commercial Ave., Cambridge, MA, 02141); Van Dongen, C. G.; Adams, R. A.; Soto, E. A.; Pai, K. J. *Cancer Lett* 4(3): 177-183; 1978.

The effect of reserpine on methylcholanthrene (MC)-induced breast cancer was studied in BIO 15.16 Syrian hamsters, a strain previously shown to be sensitive to the induction of mammary cancer by MC. There were seven experimental groups. Group 0 (25 hamsters) received 10 µg reserpine ip three times weekly for life. (An initial dose of 100 µg was reduced after 1 wk because of intolerance.) Group I (25) received 50 gastric gavages of 0.05 mg MC. Group II received (25) reserpine + MC. Group III received reserpine + 50 gavages of 0.1 mg MC. Group IV received 50 gastric gavages of 4 mg MC. Group V received the vehicle for reserpine and MC. Group VI received no treatment. All MC doses except the 4-mg dose were non-carcinogenic. The survival curves of all groups except Group IV were not significantly different from each other. Mammary tumors were noted only in 11 Group IV animals; other Group IV tumors included 3 adenocarcinomas of the colon and cecum, 2 lymphomas, and 2 sar-

comas. Multiple mammary tumors were a frequent finding. Metastasis of mammary gland carcinoma to the cervical, axillary, and/or inguinal lymph nodes was observed in 10 animals, metastases to the lungs in 2. No tumors were noted in Groups 0 and I. Group II had 1 sarcoma, Group III 2 lymphomas, Group V 1 sarcoma, and Group VI 1 lymphoma. The results do not conclusively rule out or demonstrate cocarcinogenicity of reserpine. (9 refs)

- 78-6322 Metabolic and Enzymatic Factors in Tumor Susceptibility.** (Eng) Kellermann, G. H. (Div. Human Oncology, Univ. Wisconsin Medical Sch., Madison, WI). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for Detection and Prevention of Cancer in New York, April 1-5, May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY) : Vol. 2 (Part I), 2404 pp.; 1517-1524; 1978.

The relationship between aryl hydrocarbon hydroxylase (AHH) activity and susceptibility to carcinogenic polycyclic aromatic hydrocarbons (PAH) was studied. Human lymphocytes were cultured for 2 days in the presence of 2 mg/ml phytohemagglutinin and pokeweed prior to the administration of 3-methylcholanthrene or benzo(a)anthracene to induce AHH activity. Individual inducibility was expressed as the ratio of AHH activity in the PAH-treated cultures to AHH activity in the cultures without treatment. The lymphocytes were obtained from 289 normal subjects, patients with bronchogenic carcinoma (50) or laryngeal carcinoma (37), and 46 patients (cancer control group) with Hodgkin's disease or cancers of the skin, gastrointestinal tract, head and neck, breast, uterus, prostate or bladder. Previous family and population studies have revealed considerable interindividual variation of this enzyme complex and demonstrated its genetic control. Ninety percent of the healthy subjects had low or intermediate AHH inducibilities in their lymphocytes. However, 96% of the lung cancer patients, all heavy smokers, had intermediate (66.0%) or high (30.0%) AHH inducibility. This distribution pattern was not seen in the laryngeal cancer patients (19.0% with high inducibility) or in the cancer control group (10.9% with higher inducibility). It is suggested that cigarette smokers with high AHH inducibility have a higher risk of developing lung cancer than persons with low inducibility. Ninety percent of the 289 healthy subjects had low (114) or intermediate (118) AHH inducibilities in their lymphocytes. (18 refs)

- 78-6323 1,2-Dihydro-1,2-dihydroxy-5-methylchrysene, A Major Activated Metabolite of the Environmental Carcinogen 5-Methylchrysene.** (Eng) Hecht, S. (Div. Environmental Carcinogenesis, Naylor Dana Inst. Disease Prevention, American Health Foundation, Valhalla, NY)

, 10595); LaVoie, E.; Mazzaresse, R.; Amin, S.; Bedenko, Hoffmann, D. *Cancer Res* 38(7): 2191-2194; 1978.

the metabolic activation of the environmental carcinogen methylchrysene was studied by combining high-pressure liquid chromatographic analysis of metabolites formed in vivo with assays of these metabolites for mutagenic activity toward *Salmonella typhimurium*. Metabolites were formed during the incubation of 5-methylchrysene with the 9000 x g supernatant from Aroclor-treated rat livers. With the use of reverse-phase columns, the metabolites were resolved into several peaks. Significant mutagenic activity was observed primarily in one peak and to a lesser extent in a second peak. The major mutagenic metabolite was identified as 1,2-dihydro-1,2-dihydroxy-5-methylchrysene (7.0% from 5-methylchrysene), and the secondary mutagenic metabolite was identified as 7,8-dihydro-7,8-dihydroxy-5-methylchrysene (2.6% from 5-methylchrysene). The results indicate that 1,2-dihydro-1,2-dihydroxy-5-methylchrysene is a proximate mutagenic metabolite of methylchrysene and support the hypothesis that carcinogenic polynuclear aromatic hydrocarbons can be activated through the formation of specific angular ring diol-epoxides. (no refs)

6324 Relationship Between DNA Damage, Cell Survival and DNA Repair in CHO Cells after In Vivo Metabolic Activation of Promutagens (Meeting Abstract). (Eng) Douglas, G. R. (Environmental Mutagenesis Section, Dept. Natl. Health and Welfare, Tunney's Pasture, Ottawa, Canada); Grant, C. E. *Can J Genet Cytol* 20(3): 442; 1978. (no refs)

6325 Carcinogen-induced DNA Repair in Nucleotide-Permeable *Escherichia coli* Cells. Analysis of DNA Repair Induced by Carcinogenic K-Region Epoxides 1,2,3,4-Diepoxybutane. (Eng) Thielmann, H. W. (Deutsches Krebsforschungszentrum, Institut für Biochemie, Im Neuenheimer Feld 280, D-6900 Heidelberg, West Germany); Thielmann, H. *Z Krebsforsch* 92(2): 157-176; 1978.

Enzymic activities catalyzing DNA repair in ether-permeabilized (nucleotide-permeable) *Escherichia coli* cells exposed to carcinogenic K-region epoxides or the aliphatic epoxide 1,2,3,4-diepoxybutane (DPB) were identified by determination of repair polymerization and size reduction of damaged DNA. The requirement for the *uvr* gene function in excision repair was tested using wild type and *uvr* A and *uvr* B mutant *E. coli* cells. When exposed to the carcinogenic K-region epoxides 7-methyl- and 7,12-dimethylbenz(a)anthracene-5,6-oxide, chrysene-5,6-oxide, and benzo(a)pyrene-4,5-oxide, the wild type cells (KMBL 1788; H 540) exhibited DNA excision repair, but the *uvr* A and *uvr* B cells did not. The K-region epoxide phenanthrene-9,10-oxide was effective in all *E. coli* strains tested. DPB and the 6,7-epoxides of the tumor promotor, 12-O-tetradecanoyl-phor-

bol-13-acetate, elicited DNA repair in the *uvr* A and *uvr* B mutant cells. None of the epoxides induced repair polymerization in a mutant that lacked the 5'-3' exonucleolytic activity of DNA polymerase I (exonuclease VI). Greatly reduced levels of repair polymerization were observed in mutants that lacked DNA polymerase I. It is concluded that excision of K-region epoxide damaged nucleotides is controlled, at least in part, by three repair enzyme activities: a UV-endonuclease (missing in the *uvr* mutants), the 5'-3' exonucleolytic activity of DNA polymerase I, and an intact DNA polymerase. Excision of DPB damaged nucleotides requires in addition a *uvr*-independent repair endonuclease, which is probably endonuclease II, DNA glycosidase(s), and/or apurinic endonuclease(s). (58 refs)

78-6326 Studies on the Metabolic Activation of Benzo(a)pyrene and Subsequent Binding to Nuclear Macromolecules. (Eng) Pezzuto, J. M. (Coll. Medicine and Dentistry, New Jersey Graduate Sch. Biomedical Sciences, Newark, NJ, 07103). *Diss Abstr Int B* 39(5): 2281-2282; 1978. (no refs)

78-6327 Benzo(a)pyrene Absorption from the Intestinal Tract of the Rat with Emphasis on Lymphatic Uptake and Tissue Distribution. (Eng) Reid, J. B. (Univ. Cincinnati, Cincinnati, OH). *Diss Abstr Int B* 39(5): 2205; 1978. (no refs)

78-6328 Antimutagenic Effects of 2(3)-tert-Butyl-4-hydroxyanisole and of Antimicrobial Agents. (Eng) Batzinger, R. P. (Dept. Pathobiology, John Hopkins Medical Inst., Baltimore, MD, 21205); Ou, S. Y.; Bueding, E. *Cancer Res* 38(12): 4478-4485; 1978.

To determine antimutagenic effects of the antioxidants 2(3)-tert-butyl-4-hydroxyanisole (BHA) and ethoxyquin (EQ), diets containing 0.75% BHA or 0.5% EQ were fed to female CD-1 mice that were subsequently treated with benzo(a)pyrene (BP: 100 mg/kg im in sesame oil), hycanthone (60 mg/kg im as the aqueous soln of methane sulfonate salt), and metronidazole, diazepam, or mebendazole (each 200 mg/kg by gastric intubation). BHA and EQ markedly decreased the level of mutagens present in BP-treated mice, as determined by a host-mediated assay and tests with *Salmonella typhimurium* strains TA100 and TA98. BHA also reduced the mutagenic activities in vivo of the four antischistosomal compounds. These effects were accompanied by increases in the thiol levels of some tissues. The production of mutagenic metabolites of two other antischistosomal drugs, 4-isothiocyano-4'-nitrodiphenylamine and oxamniquine, was not reduced by BHA treatment. However, reductions in mutagenicity were achieved by the administration of enteric antibacterial agents, which implicates intestinal microorgan-

isms in the mutagenic activation of certain chemical agents. Combined treatment of mice with BHA and enteric antimicrobial agents reduced the levels of mutagens derived from metronidazole by more than 90%, and the combined treatments were more effective than was either treatment alone. (37 refs)

- 78-6329 Selective Fluorescence Quenching of Benzo(a)-pyrene and a Mutagenic Diol Epoxide Derivative in Mouse Cells.** (Eng) Wade, C. G. (Dept. Chemistry, Univ. Texas at Austin, Austin, TX, 78712); Baker, D. E.; Bartholomew, J. C. *Biochemistry* 17(20): 4332-4337; 1978.

Fluorescence and flow cytometric studies were made of the effects of various quenchers on suspensions of cultured mouse cells containing benzo(a)pyrene (BP) and its diol epoxide (DE) 7 β , 8 α -dihydroxy-9 α , 10 α -epoxy-7,8,9,10-tetrahydrobenzo(a)pyrene. The overlap of the fluorescence spectra of BP and its metabolites, particularly the DE, had previously prevented the use of fluorescence in investigations of the impact of these carcinogens on cell populations. The cells used were NMuLi (mouse liver epithelial cells), BALB A31 3T3 HYF (mouse fibroblast cells), and MSV/MLV BALB 3T3 A31 HYF (Moloney sarcoma virus-transformed mouse fibroblast cells). The most effective quencher was the lipid-soluble N-octyl iodide, which quenched the BP emission by a factor of 6 with little change in the DE emission. At comparable molarity, methylene iodide was a relatively effective quencher for both aromatic hydrocarbons, but it was not selective. The ionic quencher KI was ineffective. The results were, in general, independent of the type of mouse cell strain. Information on the location of the hydrocarbons in the cells is given. The selectivity of n-octyl iodide may be due to the fact that collisions with BP are much more likely than collisions with the DE. Thus, the possible locations of the DE are: bound in proteins or attached near the polar head groups of lipids in membranes or attached in a nonlipid portion of the cell. The selective quenching technique should have broad application to fluorometric studies of various kinds in biological systems, including flow cytometry and cell sorting. (44 refs)

- 78-6330 Health Aspects of Water Chlorination with Special Consideration of the Cocarcinogenicity of Chlorine. II. The Cocarcinogenicity of Chlorine.** (Ger) Pfeiffer, E. H. (Hygiene-Institut, Hochhaus am Augustusplatz, D-6500 Mainz, W. Germany). *Zentralbl Bakteriol [Orig B]* 166(2/3): 185-211; 1978.

The possible cocarcinogenicity of chlorine and polycyclic aromatic hydrocarbons was investigated. A 0.22% or 0.44% acetic benzo(a)pyrene (BP) soln was applied twice weekly to the depilated skin of the backs of female NMRI mice for 10 wk to a total dose of 750 or 1,500 μ g/animal. In addition, various groups were treated with a 1% NaOCl soln either

before, during, or after the BP treatment. After 128 wk tumor development was considerably slowed or terminated prematurely in the groups given a chlorine pretreatment of 750 or 1,500 μ g BP and in the groups that had received chlorine after applications of the lower BP dose. Tumor growth in the remaining groups was not affected. Administration of chlorine before BP noticeably reduced the number of tumors considerably; only a very slight reduction was found in the remaining test groups. Chlorine treatment decreased the number of carcinomas among the tumors by about 40%, regardless of the treatment method or the dosage. The mean tumor induction times were not significantly affected by chlorine treatment, and there were no major differences in mortality rates among the groups. Possible mechanisms of action of chlorine are discussed. (162 refs)

- 78-6331 Pulmonary Metabolism of Epoxides.** (C) Smith, B. R. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC, 27709); Maguiness, H.; Ball, L. M.; Bend, J. R. *Fed Proc* 37(11): 2480-2484; 1978.

Epoxide hydrase (EH) and glutathione S-transferase (GST) activities were measured in pulmonary tissue of guinea pigs, rabbits, and rats. On the basis of total organ activity per milligram of protein, pulmonary tissue had a greater capacity to metabolize epoxides than liver tissue. Pulmonary EH and GST were refractory toward inducers such as dieldrin, zanthracene, pregnenolone-16 α -carbonitrile, 2,3,6,7-tetrachlorodibenzo-p-dioxin, and phenobarbital. In the isolated perfused rabbit lung, benzo(a)pyrene 4,5-oxide (BPO) was metabolized by EH and GST at similar initial rates, but EH activity was lost after 5 min at a dose of 1 micromole (μ M) BPO/lung and after 15 min at 5 μ M BPO/lung. Administration of a second dose of BPO (2.5 μ M) 19 min after the first temporarily restored the initial rate of dihydrodiol formation, indicating that the EH system itself was still functional. EH activity is probably lost because of inadequate substrate levels. GST from rabbit lung cytosol separated chromatographically into six peaks containing 1-chloro-2,4-dinitrobenzene (DNCB)-conjugating activity. All six peaks also displayed GST activity toward BPO, but two toward styrene oxide. GST isolated from rabbit lung supernatant fraction was also separated into six peaks, four of which conjugated glutathione with DNCB and BPO, only one of which conjugated glutathione with styrene oxide. The liver protein and total transferase activity levels were higher than those in the lung. Each of the cell types in the lungs is postulated to possess its own specific enzyme level. For the detoxification defense mechanism to be effective, each cell would have to be capable of metabolizing its internally generated epoxides. (25 refs)

- 78-6332 Nephrogenic Hypertension and Carcinogenesis.** (Rus) Litvitskii, P. F. (Dept. Pathologic Physiology,

gy, I. M. Sechenov First Moscow Medical Inst., Moscow, (SR); Kogan, A. Kh. *Vopr Onkol* 24(9): 56-60; 1978.

study was made of the effect of nephrogenic hypertension (NHT), induced by left nephrectomy and stenosis of the artery of the contralateral kidney, on the growth of benzo(a)pyrene (BP)-induced tumors and of transplanted carcinoma RS-1, Walker's carcinoma 45, Walker's carcinosarcoma, Zajdela ascites hepatoma, and Walker's metastasizing carcinosarcoma in outbred male rats. BP was injected sc, the solid tumors were transplanted sc, and Zajdela hepatoma ascites cells were injected into the abdominal cavity. Another group of rats without NHT was treated in the same manner. NHT reduced the latent period of BP-induced tumors by 21.1 days ($p < 0.001$), but it had no effect on the latent period of the transplanted tumors. In addition, NHT increased the absolute wt of all tumors from 33 to 41.4 g, the tumor/body wt ratio from 10.14 to 14.1 ($p < 0.001$), and the frequency of metastases of Zajdela ascites hepatoma and Walker's metastasizing carcinosarcoma. The findings indicate that NHT is a predisposing factor for carcinogenesis, probably as a result of the changes in immunological reactivity and the deficient blood supply and tissue oxygenation that occur during certain phases of hypertension. (14 refs)

78-6333 Elevation of Hepatic Glutathione S-Transferase Activities and Protection Against Mutagenic Metabolites of Benzo(a)pyrene by Dietary Antioxidants. (Eng) Benson, A. M. (Dept. Pharmacology, Johns Hopkins Univ. Sch. Medicine, Baltimore, MD, 21205); Batzinger, R. C.; Ou, S. Y.; Bueding, E.; Cha, Y. N.; Talalay, P. *Cancer Res* 38(12): 4486-4495; 1978.

The effects of the antioxidants 2(3)-tert-butyl-4-hydroxyanisole (BHA) and ethoxyquin (EQ) on the mutagenic metabolites of benzo(a)pyrene (BP) were studied in female C57BL/6J mice and male Sprague-Dawley rats injected im with 10 mg/kg BP. The mice were maintained for 12 days on diets containing 0.75% BHA or 0.5% EQ; alternatively, some mice received BHA (100 to 1,000 mg/kg/day for 4 days) by gavage. The rats were fed a diet containing 0.75% BHA for 8 days. The mutagenic activity of BP urinary metabolites for *Salmonella typhimurium* strains TA98 and TA100 was markedly reduced in the presence of glutathione together with the liver cytosols of animals fed BHA. The liver cytosols of mice and rats maintained on control diets were much less effective in this respect. Dietary BHA increased mouse and rat hepatic glutathione S-transferase (GST) specific activities with 1,2-dichloro-4-nitrobenzene, 1-chloro-4-nitrobenzene, p-nitrobenzylchloride, and δ^5 -androsterone-3,17-dione. The increases were larger (5- to 10-fold) in mice, and they depended on the dose and duration of BHA administration. Increases in GST specific activities were also observed in mouse hepatic cytosols after feeding of EQ. Direct addition of reduced glutathione and purified GST's A and B obtained from rat liver to the mutagenicity assay system mimicked the effect of the rodent cytosols. The

protective effects of the two antioxidants against potent carcinogens may be due, at least in part, to their ability to elevate GST's. These enzymes inactivate arene oxides and other hydrophobic electrophiles by catalyzing their conjugation with glutathione. (45 refs)

78-6334 Carcinogens and Dicumarol: Opposite Effects on Rat Liver NAD(P)H Dehydrogenation. (Eng) Schor, N. A. (Dept. Pathology, Tulane Univ. Sch. Medicine, New Orleans, LA, 70112); Boh, E.; Burke, V. T. *Enzyme* 23(4): 217-224; 1978.

The effects of four carcinogens and dicumarol (DC) on hepatic D-T-diaphorase activity were studied in several groups of Sprague-Dawley rats. Groups 1-4 received benzo(a)pyrene (BP), 7,12-dimethylbenz(a)anthracene, N-acetylaminofluorene, or DC (4.0, 3.9, 4.5, and 3.0×10^{-4} moles/kg, respectively, intragastrically) 24 hr or (DC) 24 and 48 hr before sacrifice. In Group 5, BP was administered for 6 days, DC was given on days 5 and 6, and rats were killed on day 7; in Group 6, 3-methylcholanthrene (3-MC: 7.5×10^{-5} moles/kg, ip) was given for 3 days, DC was given on day 3, and rats were killed on day 5. All carcinogens increased D-T-diaphorase activity 24 hr after administration. DC inhibited this activity, but not the aryl hydrocarbon hydroxylase activity induced by BP and 3-MC. DC potentiated the toxic effects of 3-MC and BP. None of the chemicals affected NADPH cytochrome c reductase activity. It is suggested that carcinogens stimulate two enzyme systems: the microsomal oxidases, which transform some chemicals into highly reactive and toxic compounds, and the cytosolic reductases, which may further detoxify the compounds formed in the microsomal fraction by transforming them into dihydrodiols and phenols. Apparently, there is a competition between both systems. The increase in toxicity with DC (which inhibits the cytosolic system) + BP or 3-MC is presumed due to a lack of detoxification of the two carcinogens. (19 refs)

78-6335 Effect of Anesthetic Agent on Lung Tumor Induction in Hamsters Given Benzo(a)pyrene-Ferric Oxide. (Eng) Henry, M. C. (Environmental Protection Res. Div., US Army Medical Bioengineering Res. and Development Lab., Ft. Detrick, MD, 21501); Port, C. D. *J Natl Cancer Inst* 61(5): 1221-1227; 1978.

Comparison was made of the effects of Brevital, ether, and methoxyflurane (MF) on lung tumor induction in noninbred Syrian golden hamsters given 10 weekly intratracheal instillations of a benzo(a)pyrene (BP)-ferric oxide mixture. Inhalation anesthesia with ether or MF was accomplished by a closed recirculatory system that allowed a short induction time and good control over the concentration of anesthetic. A third group was subjected to systemic induction with 0.4-0.8 ml of 1% Brevital soln ip. Survival rates during the 10 weekly instillations were least for the Brevital-treated

group and greatest for the MF-treated animals. Body wt gain was lower in the ether- and Brevital-treated groups than in the MF-anesthetized group. Animals anesthetized with Brevital had the shortest tumor latency, but tumor incidence was similar in this group and the group treated with ether. Exposure to MF and carcinogen produced a slow onset of deaths from tumors and a lower tumor incidence. The predominant tumor type in all segments of the respiratory tract was squamous cell carcinoma. The most common tumor sites were the trachea and bronchus. Thus, experimental conditions for the induction of lung carcinomas must be standardized if consistent and reproducible results are to be obtained. Inhalation anesthesia with ether does not appear to be the method of choice because of its irritant properties and possible effect on tumor formation. (14 refs)

- 78-6336 Human Skin Aryl Hydrocarbon Hydroxylase: Induction by Coal Tar.** (Eng) Bickers, D. R. (Div. Dermatology, Case Western Reserve Univ., Sch. Medicine, Cleveland, OH, 44106); Kappas, A. *J Clin Invest* 62(5): 1061-1068; 1978.

The effect of cutaneous application of a coal tar (CT) soln (crude CT diluted to 20% with ethanol and polysorbate 80) on aryl hydrocarbon hydroxylase (AHH) activity was investigated in six patients with generalized psoriasis and three with severe atopic dermatitis. CT soln (100 μ l) was applied to the skin, and the site was biopsied 24 hr later. Cutaneous AHH activity was increased two- to fivefold in treated vs untreated skin. AHH activity was also induced in human skin samples incubated with CT for 24 hr, but not to as high a level. In human skin incubated with CT in vitro, there was an initial lag phase of 8 hr followed by a 16-hr period of linearly increasing AHH activity, after which the activity diminished. Max responses both in vivo and in vitro occurred after 24 hr, and enzyme activity in vitro was time-, tissue- and substrate concentration-dependent. The application of a CT soln to neonatal rat skin resulted in a > 10-fold induction of skin AHH activity 24 hr later. Hepatic AHH activity was also markedly increased, indicating that substantial percutaneous absorption had occurred. Assay of several defined constituents of CT for AHH induction showed that benzo(a)pyrene was the most potent inducer of AHH tested followed by pyrene and anthracene. These studies indicate that topical application of CT soln in doses used to treat dermatologic disease causes induction of AHH in human skin, and they suggest that this induced enzyme activity could relate to carcinogenic responses to this agent in skin and other tissues. (44 refs)

- 78-6337 The Identification of High Molecular Weight Polynuclear Aromatic Hydrocarbons in a Biologically Active Fraction of Cigarette Smoke Condensate.** (Eng) Snook, M. E. (Tobacco Lab., Agricultural Res. Service, U.S. Dept. Agriculture, P.O. Box 5677, Athens, GA, 30604);

Severson, R. F.; Arrendale, R. F.; Higman, H. C.; Chort O. T. *Beitr Tabakforsch* 9(2): 79-101; 1977.

As a result of several gas chromatographic procedures including gel filtration and high-pressure liquid chromatography, and of mass spectral, UV absorption, and chromatographic retention data, > 200 compounds were isolated and identified in the most biologically active fraction of cigarette smoke condensate. Newly identified were 3,4-dimethylenene, 3,4-trimethylenepyrene, cyclopenta(c,d)pyrene, 4,5-triethylenetriphenylene, benzo(b)perylene, and several dibenzofluoranthenes. (21 refs)

- 78-6338 Plasminogen Activator in Chick Fibroblasts: Induction of Synthesis by Retinoic Acid; Synergism with Viral Transformation and Phorbol Ester.** (E) Wilson, E. L. (Dept. Clinical Science and Immunology, Medical Sch., Univ. Cape Town, Cape Town, South Africa); Reich, E. *Cell* 15(2): 385-392; 1978.

The effects of vitamin A and its derivatives, the retinoids, on plasminogen activator (PA) synthesis were investigated in chick embryo fibroblast (CEF) cultures. Low concentrations of retinoic acid (RA: 10^{-6} - 10^{-10} M) and the retinoids stimulated PA synthesis by up to 9- or 10-fold, a level somewhat lower than that obtained with optimal concentrations of the potent tumor promoter phorbol myristate acetate (PMA). The effect of RA required protein and messenger RNA synthesis, in contrast to enzyme induction by PMA and/or sarcoma virus transformation, retinoid effects were not significantly inhibited by elevated concentrations of cyclic AMP. The effects of RA and sarcoma virus transformation were synergistic rather than additive in inducing and/or stimulating PA production. Analogous synergism was observed between RA and PMA, but only at suboptimal concentrations of the latter. RA had no effect on PA production in normal or transformed cultures maximally stimulated by PMA. These findings may help to elucidate the role of retinoids in promoting tumor growth, tissue remodeling, and teratogenesis. (27 refs)

- 78-6339 The Induction of Differentiation in Teratocarcinoma Stem Cells by Retinoic Acid.** (E) Strickland, S. (Rockefeller Univ., New York, NY, 10021); Mahdavi, V. *Cell* 15(2): 393-403; 1978.

Plasminogen activator (PA) was used as a marker to study the effects of a number of compounds, especially retinoids, on F9 cells, a teratocarcinoma cell line that undergoes very limited differentiation under normal culture conditions. Retinoic acid at concentrations as low as 10^{-9} M induced multiple phenotypic changes in the cultures. The molecule was increasingly effective as the concentration was raised to 10^{-6} (10^{-7} M was used in most of the experiments); above this amount, it was toxic. Retinoic acid increased PA secretion more than 10-fold in 3 days of culture. There was a concomitant

change in cellular morphology, which closely resembled presumptive endoderm derived from teratocarcinoma yolk bodies. Treatment with retinoic acid alone or in combination with dibutyryl cyclic AMP (10^{-3} M) resulted in increase in intracellular PA levels that paralleled the end secretion of this enzyme, indicating that these compounds do not simply increase secretion of preformed PA but also stimulate the production of active enzyme. The percentage of colonies secreting PA was increased from < 10% to almost 100% by retinoic acid. When cultures were treated first with retinoic acid, washed well, and then treated with dibutyryl cyclic AMP, the cells secreted much more PA than if they were incubated with retinoic acid followed by control medium. Retinoic acid also increased the synthesis of collagenlike proteins and increased the alkaline phosphatase content of the cultures by 20-fold. The nature of these changes, as well as their independence of the continued presence of retinoic acid, are consistent with the proposition that retinoic acid induces differentiation of embryonal carcinoma cells into endoderm. (42 refs)

78-6340 The Effects of Organic Compounds on Mutagen- and Carcinogen-Activating Enzymes In Vivo (Meeting Abstract). (Eng) Bradshaw, J. J. (Dept. Physiology and Medical Biochemistry, Univ. Cape Town Medical School, Cape Town, South Africa); Ivanetich, K. M. *Mutat Res* 154: 154; 1978. (no refs)

78-6341 Fecal Hydroxysteroid Dehydrogenase Activities in Vegetarian Seventh-Day Adventists, Controls, and Bowel Cancer Patients. (Eng) Macdonald, J. (Dept. Medicine, Dalhousie Univ., Halifax, Nova Scotia, Canada); Webb, G. R.; Mahony, D. E. *Am J Clin Oncol* 31(10): S233-S238; 1978.

Free extracts of anaerobic bacteria from the fecal preparations of 14 vegetarian Seventh-Day Adventists (SDA), 8 patients with cancer of the large bowel, and 16 controls were assayed for NAD- and NADP-dependent 3α -, 7α -, and 12α -hydroxysteroid dehydrogenase (HSDA) activities using androsterone and various bile salts as substrates in eight substrate-cofactor combinations. NADP-dependent 12α -HSDH activity was highest in the SDA, while 7α -HSDH content was significantly increased in cancer patients over controls and also over SDA. Fecal pH values for SDA, controls, and cancer patients were 6.46 ± 0.58 , 6.66 ± 0.62 , and 7.03 ± 0.15 , respectively. The cancer pH was significantly higher than the SDA value; and when plotted against values for 7α -HSDH, fecal pH segregated 7/8 cancer patients from the other groups. Parameters for NADP-dependent 3α -HSDH activities using either type substrate were correlated, and the relationship ($r = 0.72$) suggested that this enzyme can oxidize both bile salts and neutral sterols in fecal anaerobic bacteria. Although 7α -HSDH is elevated in bowel cancer pa-

tients, it is presently unclear whether this enzyme is merely a marker, or plays some carcinogenic role itself. (30 refs)

78-6342 Rat Differences in Mammary Tumor Induction with Estrogen and Neutron Radiation. (Eng) Shellabarger, C. J. (Medical Dept., Brookhaven Natl. Lab., Upton, NY, 11973); Stone, J. P.; Holtzman, S. *J Natl Cancer Inst* 61(6): 1505-1508; 1978.

The effects of diethylstilbestrol (DES) and/or neutron radiation on mammary tumor formation were studied in female ACI and Sprague-Dawley (SD) rats. Young adult rats were given sc implants of pellets containing 5 mg DES and 15 mg cholesterol 2 days before irradiation with 0.4, 1.3, or 4.0 rads of 0.43 million electron-volt neutrons. Other rats received DES or irradiation only or no treatment; all rats were studied until death or for a max of 48 wk. DES treatment alone significantly increased the number of ACI rats (52%) with mammary adenocarcinomas (MAC) and the number of MAC/rat (2.74), but it had no effect on the SD rats. Conversely, irradiation only at 4 rads had no effect on the ACI rats, but it significantly increased the number of SD rats (32%) with mammary fibroadenomas (FAC), the number of MAF/rat, and the number of MAC/rat. Combining DES and irradiation 4 rads increased MAC incidence in ACI rats (91%), the incidence of multiple MAC (≥ 16 tumors) (65%), and the number of MAC/rat (15.10) to a total larger than the sum of the effects of the individual agents. Combined treatment in SD rats decreased MFA incidence (3%) and the number of MFA/rat (0.03) to values lower than irradiation alone. Additionally, DES treatment produced large pituitary tumors in 78%-100% of ACI rats and pyometritis in 53% of SD rats. Thus, DES and neutron irradiation acted synergistically on MAC formation in ACI rats but not in SD rats. (15 refs)

78-6343 Perphenazine and Testosterone Metabolism by Mammary Tumours in Oophorectomized Rats. (Eng) Buchan, P. (Dept. Clinical Surgery, Medical Sch., Univ. Edinburgh, Edinburgh EH8 9AG, Scotland); Miller, W. R. *Biochem Soc Trans* 6(1): 135-136; 1978.

The effects of prolactin and ovarian hormones on tumor growth and steroid metabolism were studied in female Sprague-Dawley rats with 7,12-dimethylbenzanthracene-induced (30 mg intragastrically) mammary tumors. When the tumors reached a size of > 2 cm $\times$ 2 cm, the animals were oophorectomized and placed in one of two groups. The treatment group received daily sc injections of 5 mg/kg perphenazine, and the control group received the vehicle only; all animals were killed 14 days later. Oophorectomy alone produced an immediate regression of all tumors, but the combined treatment resulted in a variable response: certain tumors increased in size while others regressed, usually after a short period of growth. Animals from the treatment group showed a signifi-

cantly higher metabolism of testosterone than controls. However, the major metabolites of testosterone in groups were 5α -dihydrotestosterone and 5α -androstenediol. Thus, in oophorectomized animals, perphenazine does not produce the stimulatory effects on mammary tumor growth and steroid metabolism that it has been shown to do in endocrine-intact animals. (3 refs)

- 78-6344 Attempted Induction of Granulosa Cell Tumor in Balb-C Mice by Gonadotropin Administration.** (Eng) Menczer, J. (Dept. Obstetrics and Gynecology, Sheba Medical Center, Tel-Hashomer, Israel); Komarov, H.; Shenboun, M.; Insler, V.; Czernobilsky, B. *Gynecol Invest* 8(5/6): 314-322; 1977.

An attempt of granulosa cell tumor (GCT) induction by prolonged administration of exogenous pregnant mare's serum gonadotropin (PMSG) to young (21-day-old), mature (6-mo-old), and middle-aged (10-mo-old) BALB/c mice resulted mainly in stromal luteal cell proliferation. This proliferation occurred in 48.9% of young mice, 64.5% of mature mice, and 65% of middle-aged mice. The effect seemed to depend on duration of treatment in young mice. Young mice inoculated for < 6 mo developed significantly less luteal cell proliferation than those inoculated for > 6 mo. Only one GCT was found. This study supports the contention that besides gonadotropin stimulation, other factors may be involved in GCT development. A large number of PMSG-inoculated mice developed generalized lymphosarcoma, the mechanism of which remains unknown. (15 refs)

- 78-6345 Covalent Binding of Ethinylestradiol and Estrone to Rat Liver DNA In Vivo.** (Eng) Jaggi, W. (Inst. Toxicology, Federal Inst. Technology and Univ. Zurich, CH-8603 Schwerzenbach, Switzerland); Lutz, W. K.; Schlatter, C. *Chem Biol Interact* 23(1): 13-18; 1978.

The covalent binding of $6,7\text{-}^3\text{H}$ -ethinylestradiol (EE) and $6,7\text{-}^3\text{H}$ -estrone (E) to the liver DNA of fasted female Sprague-Dawley rats (190 to 240 g) was measured 8 hr after the administration of 80 μg (9.2 mCi) of the estrogens by gavage. Only 1% of the total radioactivity from EE was excreted in the urine within 8 hr; 4% of the radioactivity lyophilized as tritiated water. Three percent of the radioactivity from E was excreted in the urine within 8 hr, 3% of it as tritiated water. The binding of EE and E to the liver DNA was 1.5 and 1.1, respectively, expressed as binding to DNA/dose, in units of $\mu\text{moles hormone/mol DNA phosphate/millimoles hormone/kg}$. It was on the same order of magnitude as that for benzene and about 10,000-fold lower than that for typical liver carcinogens such as aflatoxin B₁ and N,N-dimethylnitrosamine. E and its ethinyl derivative should not differ with regard to their suspected carcinogenicity. (20 refs)

- 78-6346 Histometric Study of the Pituitary in Treated Neonatally with Steroids and the Relationship Between Prolactin Cells and Mammary Tumorigenesis.** (Eng) Kawashima, S. (Zoological Inst., Faculty of Science, Univ. Tokyo, Tokyo 113, Japan); Bern, H. A.; Jones, L. A.; Mills, K. T. *Endocrinol Jpn* 25(4): 341-348; 1978.

In female BALB/cfC3H/Crgl mice, neonatal hormone treatments had a significant effect on pituitary histology and mammary tumorigenesis. Within 36 hr after birth, mice were given five daily sc doses of 5 or 20 μg 17β -estradiol (E), 100 μg progesterone (PRG), or 5 or 20 μg ovine prolactin (PRL), singly or in various combinations. Pituitary glands were studied at autopsy upon tumor development or at 1 yr. Treatment with ESD (with or without PRL), PRG + ESD, or ESD + PRG resulted in significantly greater overall mammary tumor incidence, at a significantly earlier age, than in untreated controls. Pituitary PRL cells were more numerous in tumor-bearing mice vs non-tumor-bearing mice. Ovariectomy at day 40 markedly decreased the PRL cell count and abolished the effects of neonatal steroid treatment. Neonatal ESD (with or without PRL) and PRG treatment also resulted in significant decreases in gonadotropes. PRL alone had no such effect. There was correlation between gonadotropin levels and thyrotrope levels and mammary tumor incidence. These results suggest that neonatal steroid treatment stimulates mammary tumorigenesis through increased prolactin secretion. (34 refs)

- 78-6347 Induction of Ovalbumin Synthesis in Immature Chicks by Actinomycin D and Thioacetamide.** (Eng) Sharma, O. K. (Dept. Microbiology, Univ. Colorado Medical Center, Denver, CO, 80262). *Science* 202(4368-69): 1978.

The induction of ovalbumin synthesis and of an increase in serum progesterone (PG) levels in estrogen-treated immature chicks by thioacetamide (TAM) and actinomycin D (AMD) is reported. Four-day-old Calhorm chicks were inoculated with 1 mg estradiol benzoate in sesame oil for 10 days (primary estrogen stimulation). After 3-4 wk of estrogen withdrawal, with no ovalbumin synthesis being detectable, chicks were inoculated ip with TAM (0.25 mg/g/day x 3) or AMD (0.4 μg /g/day x 1). TAM-treated chicks were sacrificed after 3 days, AMD-treated chicks after 24 hr. Oviducts were removed, and ovalbumin synthesis in the oviductal lumen was determined. Labeled ovalbumin was measured by specific immunoprecipitation with monospecific antiserum against purified ovalbumin and confirmed by gel electrophoresis. Pooled serum was analyzed for PG by radioimmunoassay. TAM and AMD induced ovalbumin synthesis from undetectable percentages of total protein synthesis to 9.1% and 8.7%, respectively. Likewise, serum PG levels (nanograms/100 mg) were increased from 89 to 296 and from 40 to 317 in two TAM treatment trials and from 100 to 400 and from 81 to 496 in two AMD trials. PG induced by TAM and AMD may account for the induction of ovalbumin synthesis.

s. How carcinogens of different structures such as TAM and AMD increase serum PG remains to be elucidated. The hormone imbalance may have implications in the mechanisms of carcinogenesis. (27 refs)

348 Metabolism of Androgens and Estrogens by Human Fecal Microorganisms. (Eng) Lombardi, P. Infectious Disease Service, Dept. Medicine, Tufts-New England Medical Center, Boston, MA, 02111; Goldin, B.; Bou-E., Gorbach, S. L. *J Steroid Biochem* 9(8): 795-801; 1978.

metabolism of androgens and estrogens by human fecal microorganisms was studied in vitro at high and low fecal concentrations and under aerobic and anaerobic conditions. The reactions were run for 0-7 days at a fecal extract concentration of 50-60 or 6-10 mg/ml; the metabolites were extracted and purified, and steroids were identified by gas-liquid chromatography. In reactions at the high fecal concentration 5 β -3-ketosteroids were reduced to 5 β -3-hydroxysteroids and 17-ketoandrogens and estrogens were reduced to hydroxy products. A low fecal concentration resulted in products formed by oxidation and by cleavage of conjugated steroids: 5 β -androsterone-3,17-dione was converted to 4-androstene-3,17-dione, and 17 β -hydroxysteroids were converted to 17-oxosteroids. Estrogen-3-sulfate and estradiol-3-glucuronide were hydrolyzed to form estrone and estradiol, respectively. These results indicate that fecal concentration determines the types of metabolites formed. The reactions occurred under aerobic and anaerobic conditions. Their occurrence under aerobic conditions may reflect the ability of fecal extract to lower the oxidation-reduction potential of the reaction mixture. This study demonstrates that human intestinal flora can carry out a variety of reductive, oxidative, and hydrolytic reactions on androgens and estrogens. (21

349 Plasma Hormone Profiles of Young Women at Risk for Familial Breast Cancer. (Eng) Fishman, J. (Inst. Steroid Res., Montefiore Hosp. and Medical Center, Bronx, NY, 10467); Fukushima, D.; O'Connor, J.; Benfeld, R. S.; Lynch, H. T.; Lynch, J. F.; Guirgis, H.; Honey, K. *Cancer Res* 38(11, part 2): 4006-4011; 1978.

Plasma hormone profiles from 30 women at high risk for familial breast cancer and from their matched low-risk controls were analyzed. The case subjects were premenopausal, pubertal women without evidence of neoplastic disease with many close relatives with breast cancer or tumors associated with breast cancer and who, by genetic analysis, were at a clearly defined high risk for the disease. The controls were matched closely for height, wt, age, and reproductive history. Plasma hormone levels were measured every second day throughout the menstrual cycle, and the results were analyzed in terms of follicular, luteal, and full-cycle mean concentrations. There were no statistically significant differ-

ences in prolactin, follicle-stimulating hormone, luteinizing hormone, estrone, estradiol, or estriol levels between the two groups, although the high-risk group had consistently lower values of all these hormones except estriol. The mean plasma estriol levels in the follicular and luteal stages were 19.7 and 24.8 picograms/ml in the high-risk women and 23.5 and 23.1 picograms/ml in the controls. The results suggest that the etiology of female and male familial breast cancer is fundamentally different, since male relatives of male breast cancer patients had substantially elevated plasma estrogen levels. (26 refs)

78-6350 Phytoestrogen Interaction with Estrogen Receptors in Human Breast Cancer Cells. (Eng) Martin, P. M. (Institut J Paoli-I. Calmettes, 13273 Marseille, France); Horwitz, K. B.; Ryan, D. S.; McGuire, W. L. *Endocrinology* 103(5): 1860-1867; 1978.

The interactions of phytoestrogens with estrogen receptors (ER) were studied in the human breast cancer cell line MCF-7. When cytosol from MCF-7 cells was incubated with 4 nanomoles (nmol) ³H-estradiol and 400 nmol unlabeled phytoestrogens, all but one compound competed with estradiol for binding to nuclear and cytoplasmic ER. The relative binding affinities (followed by approx dissociation constants) were zearalenol high-melting point isomer (HP) (1 nmol) > zearalenol low-LMP (15 nmol) > zearalenone (6 nmol) = coumestrol (2 nmol) > genistein (10 nmol) > formononetin (2 μ M), which had little effect. The phytoestrogens bound only weakly to sex steroid-binding globulin and none bound to corticosteroid-binding globulin. When whole MCF-7 cells were incubated with 1 μ M genistein, coumestrol, and zearalenol HMP for 5 min to 15 hr, the cytoplasmic ER was depleted within 5 min, and the hormone-receptor complex was translocated to the nucleus. Zearalenol was as effective as estradiol in the ER translocation. The phytoestrogen-bound receptors were processed in the nucleus at about the same rate as estradiol-bound receptor. The plant estrogens, particularly coumestrol, were markedly enhanced during the log-phase growth of MCF-7 cells. These nonsteroidal compounds apparently function in the same manner as estrogen, and they may have profound effects on estrogen target cells. (41 refs)

78-6351 Impeding Estrogens and the Etiology of Breast Cancer. (Eng) Wotiz, H. H. (Dept. Biochemistry, Boston Univ. Sch. Medicine, Boston, MA, 02118); Chatteraj, S. C.; Kudisch, M.; Muller, R. E. *Cancer Res* 38(11, part 2): 4012-4020; 1978.

New data relevant to the action of estriol (E) and related impeding estrogens, obtained from several experiments, are presented. Synthesis of progesterone receptor, induced by 2-day treatment of Sprague-Dawley rats with estradiol (ED: 0.1 μ g/day/sc), was inhibited by simultaneous injection of 10 μ g/day E or 16-epiestriol. Silastic tubing implants con-

taining E (1 mg) markedly reduced breast tumor takes and delayed the onset of first tumors in rats treated 4 days later with 7,12-dimethylbenz(a)anthracene (3 mg every other day x 3, iv). In contrast, implants containing 1 mg ED appeared to increase the number of tumor-bearers and to accelerate the onset of first tumors. Administration to pregnant rats of antibody specific to ED or E during gestation days 11-14 resulted in a slight reduction in the number of offspring for anti-ED antibody but a nearly total loss of fetuses with anti-E antibody. The contraceptive function of E cyclopentyl ether (500 μ g/kg po, gestation day 1) was reversible when 1.2 μ g/kg ED, was injected sc on gestation day 4. Injection of freshly mated rats with up to 500 μ g/kg 2-hydroxyestrone sc did not result in any antiimplantational or estrogenic activity. However, topical insertion of Silastic tubing implants containing this steroid into rabbit uteri produced complete contraceptive effects. Multiple injections of 50 μ g 2-hydroxyestrone produced max uterine growth in immature and adult ovariectomized rats. The data suggest that E, although containing some estrogenic properties, must have a significantly different physiological role in cancer development and reproduction when present in appropriate admixture with ED. (43 refs)

- 78-6352 Benign Liver Tumors and Oral Contraceptives.** (Eng) Aungst, C. W. (Roswell Park Memorial Inst., 666 Elm St., Buffalo, NY, 14263). *NY State J Med* 78(12): 1933-1934; 1978.

A hepatic adenoma developed in two women (both aged 24) who had been taking oral contraceptives for up to 4 yr and in one woman (aged 52) who had been taking conjugated estrogens for menopausal symptoms for 3 yr. One of the young women had symptoms suggesting an amebic abscess. She also had a carcinoembryonic antigen level of 18.7 nanograms/ml, a finding that may have been fortuitous. (7 refs)

- 78-6353 Hepatocellular Carcinoma Associated with Oral Contraceptives.** (Eng) Menzies-Gow, N. (Central Middlesex Hosp., Acton Lane, London, England). *Br J Surg* 65(5): 316-317; 1978.

A primary hepatic tumor occurred in a 36-yr-old woman who had been taking oral contraceptives (norethisterone acetate 3 mg, ethinylestradiol 50 μ g) for 6 yr. The patient suffered also from anemia and lactic acidosis. A partial hepatectomy was performed, and a secondary mass of tumor-containing lymphatic tissue in the region of the celiac axis was also excised. The tumor was a well-differentiated hepatocellular carcinoma with highly vascular areas and evidence of vascular invasion. The secondary tumor was histologically similar to the hepatocellular carcinoma. This case is the first with histological evidence of secondary spread of a malignant tumor associated with oral contraceptives. (13 refs)

- 78-6354 Liver Adenoma Associated with Oral Contraceptive Pill Administration.** (Eng) Baker L. (Univ. Chicago, Box 400, 950 E. 59th St., Chicago, 60637); Paloyan, D.; Simonowitz, D.; Lipson, R. M. *A Dig Dis* 23(5, Suppl): 53s-58s; 1978.

The case of a 33-yr-old woman with liver adenoma in whom an elevated serum alkaline phosphatase level led to the diagnosis is reported. The patient had been taking an oral contraceptive, Ovulen 21 (0.1 mg mestranol and 1.0 mg ethynodiol diacetate) continuously for 11 yr. (43 refs)

- 78-6355 Hepatic Estrogen and Progesterone Receptors in an Estrogen-associated Hepatic Neoplasm.** (Eng) Macdonald, J. S. (Div. Medical Oncology, Vincent Lombardi Cancer Res. Center, Washington, DC, 20004); Lippman, M. E.; Woolley, P. V.; Petrucci, P. P.; Schein, S. *Cancer Chemother Pharmacol* 1(3): 135-138; 1978.

The presence of cytosol estrogen and progesterone binding proteins in an estrogen-associated hepatic neoplasm was determined. The patient was a 53-yr-old premenopausal Caucasian woman who had received weekly im injections of diethylstilbestrol for 1 yr to control menstrual irregularity, 23 months before diagnosis of a polyploid vascular hepatic tumor (8 cm), and daily doses of an oral contraceptive containing norethynodrel + mestranol for 3 mo, 5 yr before diagnosis. Histologically, the tumor was classified as a focal nodular hyperplasia of the liver. It was found to contain high-affinity receptors for estradiol (binding capacity, 47.9 fmol/mg protein) and progesterone (binding capacity, 1.5 femtomoles/mg protein), with the binding capacity of the former being three times greater than that of normal liver. These results are consistent with the hypothesis that a small portion of normal hepatic cells have relatively high concentrations of cytosol estrogen receptors; these receptors render the cells particularly susceptible to stimulation to a neoplastic state by estrogen. (16 refs)

See also:

- * (Rev.): 78-6001, 78-6002, 78-6003, 78-6004, 78-6005, 78-6006, 78-6007, 78-6008, 78-6009, 78-6010, 78-6011, 78-6012, 78-6013, 78-6014, 78-6015, 78-6016, 78-6017, 78-6018, 78-6019, 78-6020, 78-6021, 78-6022, 78-6023, 78-6024, 78-6025, 78-6026, 78-6027, 78-6028, 78-6031, 78-6034, 78-6035, 78-6038, 78-6048, 78-6051, 78-6053, 78-6054, 78-6055, 78-6066, 78-6096.
* (Phys.): 78-6356, 78-6358, 78-6359, 78-6360, 78-6362, 78-6368, 78-6375, 78-6376.
* (Viral): 78-6395, 78-6411, 78-6426, 78-6440, 78-6441.
* (Immun.): 78-6490.
* (Path.): 78-6509, 78-6529.
* (Epid.-Biom.): 78-6549, 78-6550, 78-6552, 78-6553, 78-6556, 78-6557, 78-6561, 78-6562, 78-6566, 78-6567, 78-6574.

PHYSICAL CARCINOGENESIS

56 **Greater Promotion in Sister Chromatid Exchanges by Trimethylpsoralen than by 8-Meopsoralen in the Presence of UV-Light.** (Eng) Gaynor, (Dept. Dermatology, Yale Univ. Sch. Medicine, 333 St., New Haven, CT, 06510); Carter, D. M. *J Invest Biol* 71(4): 257-259; 1978.

8-methoxypsoralen (8-MOP) and 4,5',8-trimethylpsoralen (TMP) were compared with respect to their ability to promote sister chromatid exchanges (SCEs) in normal human lymphocytes exposed to UV light (365 nanomoles). With either chemical treatment, leukocytes had more SCEs than untreated controls. TMP consistently induced more SCEs per cell than did equal concentrations of 8-MOP. The increase in SCEs was proportional to drug concentration (3.23×10^{-7} or 10^{-8} M) and dose of UV light (0.3, 0.6, or 0.9 joule/cm²). Treatment with khellin plus UV light also caused a dose-dependent increase in the number of SCEs per cell; however, the number of SCEs was not increased at the higher concentrations of this drug. No chromosomal aberrations were observed with any of the treatments, but there were more mitoses in cells receiving higher doses of drug and longer exposures to UV light than in controls. The differences in the effects of 8-MOP and TMP on various cell functions may be related to qualitative and quantitative differences in the products they form. (22 refs)

57 **Nucleosome Rearrangement in Human Chromatin During UV-induced DNA Repair Synthesis.** Smerdon, M. J. (Dept. Pathology, Washington Univ. School of Medicine, St. Louis, MO, 63110); Lieberman, M. W. *Natl Acad Sci USA* 75(9): 4238-4241; 1978.

Nucleosome rearrangement was investigated in human diploid fibroblasts (IMR-90 cells) prelabeled with ¹⁴C-thymidine during the logarithmic phase, grown to confluence, treated with hydroxyurea, and irradiated with a UV dose of 100 J/m². Immediately after irradiation, the cells were chased with ³H-thymidine (10 or 90 min) and chased in non-radioactive medium for different times, and the distribution of UV-induced DNA repair synthesis in chromatin was measured. Immediately after irradiation, repair synthesis occurred primarily in staphylococcal nuclease-sensitive regions of chromatin. Subsequent incubation of cells in unlabeled medium resulted in the conversion of many of these repair sites to nuclease-resistant regions. Agarose gel data indicated that repair sites acquired nuclease resistance with the appearance in core DNA. Rearrangement also resulted in a more uniform distribution of repair sites in linker and core regions and not in an overrepresentation of repair sites in the latter. Rearrangement was both rapid and exten-

sive: at least 85% of the repair sites underwent rearrangement during the first few hours; the rest of the repair sites underwent rearrangement at a much slower rate. There was a striking correlation between the time course of rearrangement and repair synthesis, suggesting that nucleosome rearrangement may be induced by the repair process or that the rate of repair synthesis may be regulated by nucleosome rearrangement. (14 refs)

78-6358 **Molecular Mechanisms Involved in the Production of Chromosomal Aberrations. I. Utilization of Neurospora Endonuclease for the Study of Aberration Production in G2 Stage of the Cell Cycle.** (Eng) Natarajan, A. T. (Dept. Radiation Genetics and Chemical Mutagenesis, State Univ. Leiden, Wassenaarseweg 72, Leiden, Netherlands); Obe, G. *Mutat Res* 52(1): 137-149; 1978.

A study was made of the effect of Neurospora endonuclease (NE) on the induction of single-strand (ss) breaks in the G2 stage of the cycle of Chinese hamster ovary cells by x-rays, UV light, or chromosome-breaking chemicals. The enzyme was added in the presence of inactivated Sendai virus, which mediated the introduction of NE into living cells. NE is ss-specific and thus is expected to induce double-strand (ds) breaks in the ss regions of DNA. Addition of NE + virus immediately after treatment of the cells with 50-200 R x-rays more than doubled the frequencies of breaks and exchanges compared with x-irradiated controls. The effect was dose-dependent. This increase was due to the conversion of some of the x-ray-induced ss breaks into ds breaks by NE. NE + virus increased the number of aberrations produced by methyl methanesulfonate (0.01%) and bleomycin (10 µg/ml), but not those produced by UV light (10-50 ergs/mm²) or mitomycin C (mmc: 2 µg/ml). Addition of Micrococcus endonuclease and NE to the cells also did not alter the frequencies of UV-induced aberrations. The failure to convert UV- and MMC-induced lesions into aberrations in G2 indicated that these agents need an S-phase-dependent misreplication for aberration development. The study shows that the basic structure involved in chromosomal aberrations is DNA, ds breaks lead to chromosomal aberrations, and that the introduction of repair enzymes into living cells offers a way to probe into the molecular events of chromosomal aberration formation. (16 refs)

78-6359 **Effects of Chemical and Physical Mutagens on the Frequency of a Large Genetic Duplication in Salmonella typhimurium. I. Induction of Duplications.** (Eng) Hoffmann, G. R. (Natl. Inst. Environmental Health

Sciences, Research Triangle Park, NC, 27709); Morgan, R. W.; Harvey, R. C. *Mutat Res* 52(1): 73-80; 1978.

A simple selection procedure for detecting *Salmonella typhimurium* strains that are merodiploid (MD) for almost one-third of the chromosome is based on improved utilization of L-malate as the sole carbon source in MD strains. The spontaneous frequency of the duplication in haploid strains is approx 10^{-4} per cell plated. Following the exposure of a haploid strain to mutagenic agents, there is a dose-dependent increase in the duplication frequency above the spontaneous level. X-rays, UV light, ethyl methanesulfonate, nitrous acid, and the azaacridine half-mustard ICR-372, a group of mutagens that have a variety of different molecular mechanisms of action, induced genetic duplications. It is concluded that this effect in bacteria may be a consequence of the recombination of mutagenic agents analogous to that previously noted in yeasts. (32 refs)

78-6360 Effects of Chemical and Physical Mutagens on the Frequency of a Large Genetic Duplication in *Salmonella typhimurium*. II. Stimulation of Duplication-Loss from Merodiploids. (Eng) Hoffmann, G. R. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC, 27709); Morgan, R. W.; Kirven, R. *Mutat Res* 52(1): 81-86; 1978.

Strains of *Salmonella typhimurium* (ST) that contain a duplication of approx 30% of the genome may be obtained by a simple selective procedure. These strains are very unstable, losing the duplication when grown on nonselective medium. The system used to detect duplication formation and loss is based upon clone morphology and rates of growth on L-malate. Most *Salmonella* derived from wild-type strain LT2 grow very poorly on medium in which L-malate is the sole carbon source, but merodiploid (MD) derivatives of these bacteria exhibit improved growth on minimal malate medium. MD's are also distinguishable from haploids by of their characteristic rough colony morphology on minimal malate or minimal glucose medium. Treatment of MD bacteria with mutagenic agents stimulated the rate at which haploid segregants were obtained from MD strains. The mutagens tested for this effect were x-rays, UV light, ethyl methanesulfonate, and the azaacridine half-mustard ICR-372. Although other models for the induction of duplication loss in MD *Salmonella* strains cannot be ruled out, the variety of molecular mechanisms of genetic activity exhibited by these mutagens point to a recombinational effect analogous to induced mitotic recombination in yeasts. (21 refs)

78-6361 Does Ultraviolet Light Enhance Postreplication Repair in Mammalian Cells? (Eng) Painter, R. B. (Lab. Radiobiology, Univ. California, San Francisco, CA, 94143). *Nature* 275(5677): 243-245; 1978.

The results of a previous study in which Chinese hamster cells were exposed twice to UV and their DNA was analyzed by alkaline sucrose gradient velocity sedimentation were confirmed, but an alternative interpretation of the data is presented. The previous work postulated the existence of an enhanced postreplication repair system to explain the altered pulse-chase profiles observed for the DNA of mammary cells irradiated with 2.5 joules (J)/m² of UV light 2 hr before a second exposure to 5.0 J/m². If the profiles are plotted as total radioactivity per fixed number of cells, the inhibition of DNA synthesis caused by each treatment is demonstrated. Analysis of profiles of both pulse-labeled and pulse-chase DNA indicated that cells exposed to 2.5 J/m² had not recovered from the damage in their DNA by 2.5 hr after irradiation. The status of nascent DNA molecules in these cells was not the same as in controls; although the mol wt of the nascent DNA molecules was lower in treated cells than in control cells, a significant number of these molecules had a higher mol wt. Thus, the effects of 5.0 J/m² of UV light on DNA synthesis and of maturation of nascent Chinese hamster V79 DNA are different in cells that were irradiated earlier with 2.5 J/m² than in previously unirradiated cells. These differences can be explained entirely by the damage effects of the first exposure, which cause an abnormal distribution of sizes of nascent DNA molecules at the time of irradiation with 5.0 J/m². (4 refs)

78-6362 Chromosome Damage and DNA Repair Induced in Human Fibroblasts by UV and Cholesterol Oxide. (Eng) Parsons, P. G. (Queensland Inst. Medical Research, Herston, Queensland, Australia); Goss, P. *Aust J Exp Med Sci* 56(part 3): 287-296; 1978.

Attempts were made to transform normal melanoma-derived human fibroblasts using UV light, cholesterol oxide (ChO), and hyperthermia (40 C), singly or in combination. Chromosome and DNA damage was monitored along with transformed foci. No evidence of transformation was obtained after any of the treatments. Chromosome aberrations were induced by UV light (9.8%) and ChO (8.1%) alone, and the number of aberrations was increased to 13%-20% when hyperthermia was combined with ChO and/or UV light and to 15% when ChO and UV light were combined. All the induced aberrations were still present after 8 wk. Aberration frequencies were not significantly increased by multiple as opposed to single UV light treatment. UV light and ChO also induced DNA repair synthesis, and the former but not the latter, induced single-strand breaks and inhibited thymidine incorporation. The data suggest that factors synergistic to the later stages of carcinogenesis might be effective long after UV exposure. (22 refs)

78-6363 Anaplastic Thyroid Carcinoma Following Low-Dose Irradiation. (Eng) Komorowski, A. (Dept. Pathology, Medical Coll. Wisconsin, 8700 W. Wisconsin Ave., Milwaukee, WI, 53226); Hanson, G. A.; Gajdos, J. C. *Am J Clin Pathol* 70(2): 303-307; 1978.

2-yr-old man underwent surgical removal of a metastatic, well-differentiated papillary adenocarcinoma of the thyroid gland after treatment with low-dose irradiation for adenoid cystic carcinoma. Fifteen months later, an enlarged cervical node prompted a second surgical exploration and a modified, radical neck dissection was performed. Histologic examination revealed that a portion of the lesion was identical to the original tumor but that the majority of the lesion was an anaplastic thyroid cell tumor. Five of the 60 lymph nodes in the dissection contained well-differentiated papillary carcinoma. Three months postoperatively, 0.5 cm nodules were again palpated in the left neck. It appears that radiation-induced thyroid carcinomas, in at least a very small number of people, may be associated with aggressive carcinoma. (14 refs.)

6364 **Apurinic and/or Apyrimidinic Endonuclease Activity in Ataxia Telangiectasia Cell Extracts.** (Eng) Sheridan, R. B. (Dept. Biochemistry, Johns Hopkins Univ., Sch. Hygiene and Public Health, 615 N. Wolfe St., Baltimore, MD, 21205); Huang, P. C. *Mutat Res* 52(1): 131-136; 1978.

A hypothesis that a defective repair mechanism for ionizing radiation-induced DNA damage is associated with ataxia telangiectasia (AT) was investigated with the use of six APU/APY three control lines. The base excision repair mechanism involving the apurinic and/or apyrimidinic endonuclease (Apu/APY EN) responsible for incision at the site of base loss was examined in a nicking assay for Apu/APY activity using double-stranded, covalently closed, supercoiled Φ X174 RFI DNA modified to produce an average of one Apu site/molecule. There was no statistical difference in acid phosphatase levels (chosen as an internal control) between the AT and control lines, but Apu/APY EN activity was slightly reduced in the former (15 units, vs 214 units in control lines). However, even the lowest level of activity seen was far in excess of that detected by a growing cell. Next, the extent of incision by the Apu/APY EN of each of the cell lines was measured as a function of increasing amounts of damage in the substrate. As the number of Apu sites increased, both control and AT lines appeared capable of incising all those present. Thus, AT cells complete the incision step to the same extent as controls, even with 22% less enzyme activity. It is concluded that a deficiency in base excision repair at the level of the APU/APY specific for the site of base loss is probably not an explanation for AT. (24 refs)

6365 **Fatal Long-Term Sequela Following Radiation "Cure" for Ependymoma.** (Eng) Bachman, D. (Johns Hopkins Hosp., Baltimore, MD); Ostrow, P. T. *Ann Neurol* 4(4): 319-321; 1978.

A case of glioblastoma or severe radiation-induced encephalopathy occurred at the site of an ependymoma that had been treated by radiotherapy 5 yr previously. The patient, a 4-yr-old child, had been treated at age 13 mo with 3,960 rads to the whole brain over 6 wk followed by 2,300 rads to the spinal

cord over 6 wk. It seems likely that the changes were radiation-induced. (11 refs)

78-6366 **Non-enzymatic DNA Strand Breaks Induced in Mammalian Cells by Fluorescent Light.** (Eng) Bradley, M. O. (Lab. Molecular Pharmacology, Developmental Therapeutics Program, Div. Cancer Treatment, NCI, NIH, Bethesda, MD, 20014); Erickson, L. C.; Kohn, K. W. *Biochim Biophys Acta* 520(1): 11-20; 1978.

The relationship between the mutagenicity and toxicity of fluorescent light and the production of DNA damage was examined in male Chinese hamster lung (V-79-4) cells. The cells were irradiated with 5.4 joules/m²/sec of fluorescent light in complete medium at 1 C for 1, 2, or 3 hr. The light induced 0.041 single-strand break/10⁶ daltons/hr in the DNA of the cells (4.8×10^{-6} break/10⁶ daltons/joule⁻¹m²). The breaks were unrelated to pyrimidine dimers and were also apparently distinct from previously described chemically induced alkaline-sensitive lesions. They were presumably generated by nonenzymatic mechanisms, as shown by the facts that (1) the breaks were produced at 1 C, a temperature at which most enzymatic reactions are slow; (2) the number of breaks remained constant at 1 C for 2 hr after illumination, showing no delay between illumination and break production; and (3) the breaks were produced by fluorescent light in DNA isolated from cells that were lysed and deproteinized under conditions in which no enzymes should survive. The light probably produced the breaks by a nonenzymatic mechanism either directly in the DNA itself or by the action of some unknown sensitizer. When the cells were incubated at 37 C, the breaks rejoined within 2 hr. At least two lesions were responsible for the observed effects. One had the ability to break DNA subsequently treated with alkali but was neither toxic nor mutagenic. This lesion was due to > 350-nanometer (nm) light. The other lesion(s) produced mutagenicity and/or toxicity but did not necessarily produce strand breaks. This lesion(s) was due to < 350-nm light. (21 refs)

78-6367 **The Determination of Plutonium in Tissue by Aliquat-336 Extractions.** (Eng) Fisenne, I. M. (Environmental Measurements Lab., U.S. Dept. Energy, New York, NY, 10014); Perry, P. M. *Radiochem Radioanal Lett* 33(4): 259-264; 1978.

A rapid, reproducible method based on solvent extraction was developed to separate plutonium from environmentally exposed human tissue. Tetravalent Pu was extracted from 8.5 N HNO₃ into a 30% Aliquat-336 (methyl tricaprylyl ammonium chloride)/toluene mixture. Uranium and thorium were removed by nitric and hydrochloric acid washes, and the Pu was back-extracted with an HCl-HF soln. Electrodeposition of Pu from an acidic ammonium sulfate soln produced a plate suitable for alpha spectrometry. The amount of Pu recovered from samples of human lung, bronchial tree, and bone was 70%-80%, indicating that the

chemical yield was consistent regardless of tissue type or wt of tissue analyzed. (6 refs)

- 78-6368 **The Use of Transposable Genetic Elements in *Drosophila melanogaster* as a Fast and Sensitive System for Mutagenicity Testing (Meeting Abstract).** (Eng) Rasmuson, B. (Dept. Genetics, Univ. Umea, S-901 87 Umea, Sweden); Svahlin, H.; Rasmuson, A.; Montell, I.; Olofsson, H. *Mutat Res* 53(2): 252; 1978. (no refs)

- 78-6369 **$^{239}\text{PuO}_2$ Aerosol Inhalation Exposure of Miniature Swine.** (Eng) Karagianes, M. T. (Biology Dept., Battelle, Pacific Northwest Labs., Richland, WA, 99352); Beamer, J. L.; Clary, A. J.; Craig, D. K.; Decker, J. R.; Cannon, W. C. *Radiat Res* 74(1): 79-87; 1978.

Because of the anatomic and physiological similarities between swine and humans, techniques were developed for the inhalation exposure of miniature swine to radioactive aerosols to determine the biological effects of $^{239}\text{PuO}_2$. Anesthetized animals were exposed to $^{239}\text{PuO}_2$ aerosols in dog inhalation chambers that had been modified to accommodate the swine. A sling cart for transporting and restraining the pigs and special face masks to allow intranasal exposure only were designed and fabricated. Four swine were exposed to aerosols consisting of 20 mg ^{239}Pu /ml water and an air mixture of 5:1. Two swine were exposed to aerosols having an activity median aerodynamic diameter (AMAD) of $3.0\text{ }\mu\text{m}$ and a geometric standard deviation (GSD) of 2.1, and two were exposed to aerosols with an AMAD of $2.0\text{ }\mu\text{m}$ and a GSD of 1.8. The pigs were killed at 10, 13, 30, and 90 days postexposure. Pu was cleared from the body in a biphasic pattern, with a biological retention half-time of a few days for 90% of the initial alveolar burden and ≥ 50 days for the remainder. There was very little transfer of Pu to the systemic organs: 83%-96% of the Pu retained in the body at necropsy was in the lung and pulmonary lymph nodes. (20 refs)

- 78-6370 **Frequency of α -Particles from $^{239}\text{PuO}_2$ in Lung Cells.** (Eng) Kellerer, A. M. (Institut fur Medizinische Strahlenkunde der Universitat Wurzburg, Versbacher Landstrasse 5, D-8700 Wurzburg, W. Germany). *Radiat Environ Biophys* 15(11): 13-19; 1978.

Mathematical considerations of the effect of $^{239}\text{PuO}_2$ particles in the lung are presented. If a given activity is uniformly dispersed, a max number of cells will be traversed by α -particles and the probability of multiple traversals of one cell will be small. However, if the activity occurs in $^{239}\text{PuO}_2$ particles, the cells close to these particles will be subject to multiple traversals and the number of cells at risk will be reduced. Furthermore, malignant transformations produced by the α -particles are less likely to be expressed because it is unlikely that multiply traversed cells will retain their proliferative capacity. It is thus concluded that a given activity will be less effective when it occurs in $^{239}\text{PuO}_2$ particles. (4 refs)

- 78-6371 **Endonuclease Activities from a Permanent Established Mouse Cell Line that Act on DNA Damaged by Ultraviolet Light, Acid and Osmium Tetroxide.** (Eng) Nes, I. F. (Dept. Biochemistry, Univ. Bergen, Arstadveien 19, 5000 Bergen, Norway); Nissen-Meyer, M. *Acta Biochim Biophys Acta* 520(1): 111-121; 1978.

Damage-dependent endonuclease activities in cell-free extracts from mouse plasmacytoma cells (line MPC-11) were characterized using ϕ x 174 RFI DNA damaged by UV light ($1,000\text{ joules/m}^2$), acid, or osmium tetroxide. The extracts produced approx 1.1 nicks per 35 thymine-containing dimer in the UV-damaged DNA. The acid-treated DNA, which contained apurinic sites, was enzymatically broken at every kali-labile site. The extract was shown to contain an apurinic-specific endonuclease that was stimulated by magnesium, stimulation being lower for the UV-endonucleolytic activity than for the apurinic-endonucleolytic activity. Also, when the UV-endonuclease activity was little affected by reducing the KCl concentration, the apurinic-specific activity was most completely abolished. DNA exposed to increasing concentrations of OsO_4 (to $80\text{ }\mu\text{g/ml}$) became a progressively better substrate for the endonucleolytic activity. The response of the OsO_4 -specific endonuclease activity to KCl and Mg was qualitatively similar to that of the endonuclease activity that acted on UV-irradiated DNA. The results suggest that one enzymatic activity recognized both OsO_4 and UV-induced lesions in DNA (ie, pyrimidines with a saturated double bond). (24 refs)

- 78-6372 **Effect of DL-Tryptophan on Plastic-induced Tumorigenesis.** (Rus) Pliss, G. B. (Lab. Chemical Carcinogens, N. N. Petrov Res. Inst. Oncology, Leningrad, USSR); Eliseev, V. V. *Biull Eksp Biol Med* 86: 353-356; 1978.

The effect of DL-tryptophan (Try) on plastic carcinogenesis was studied in random-bred albino rats. A copolymer of methyl and ethyl ester of methacrylic acid was implanted in the form of a powder or plate, into the rats: 800 mg of powder was implanted into the right side and a 20- x 20- mm plate was implanted into the left side. On the next day the rats were divided into two groups: 34 rats in Group 1 received Try in sunflower oil (100 mg/day, 5x/wk, po), and 32 rats in Group 2 received sunflower oil only (0.5 ml/d, 5x/wk). In Group 2, the first acrylic plate-induced tumors were detected 12.3 mo after transplantation: 11/27 survivors developed fibrosarcomas after an av latent period of 11.3 mo. Implantation of the acrylic powder did not induce tumors. In Group 1, the first acrylic plate-induced tumors were detected 13.8 mo after transplantation: 6/19 survivors developed fibroblastic sarcomas (av latent period was 16.3 mo). Contrary to the Group 2 rats, two rats in Group 1 developed a fibrosarcoma at the site of acrylic powder implantation at 18.7 mo. (11 refs)

373 **Late Effects of Successive Massive Doses of Radiation on Parabiont Rat Pairs.** (Eng) War-S. (Cancer Res. Inst., New England Deaconess Hosp., Boston, MA, 02215); Chute, R. N.; Porter, M. W.; Brown, J. *Int J Radiat Biol* 34(3): 265-271; 1978.

Study was made of the life-span and leukemia and osteogenic sarcoma incidence in pairs of parabiont rats that survived 1,000-R doses of whole-body radiation, given first to one partner and then to the other after intervals of 5, 10, and 30 days. The degree of recovery and of residual damage in the first irradiated rat was measured by the extent of its ability to support its subsequently exposed partner. Life-span was shortened by 100-150 days following a 5-day interval and by 200-300 days following a 10-day interval, but it was not shortened following a 30-day interval. The incidence of leukemia (1.7%) was the same in sequentially irradiated groups, regardless of the time between exposures; the incidence of leukemia in control pairs was 4.8%. Shortening the interval between irradiations decreased the average age at death from leukemia; leukemic pairs had an average age at death of 690 days for those irradiated 30 days apart, 490 days for those with a 10-day interval, and only 325 days for those with a 5-day interval. None of the pairs in which both partners were irradiated developed solid lymphoid tumors. The incidence of osteogenic sarcoma (including lesions occurring in soft parts) in the parabiont pairs in which both partners were irradiated sharply increased to 12.9%, or about 8 times the rate in non-irradiated controls (1.6%), 2 times that in pairs in a hemi-irradiated series (7.0%), and 9 times that in shielded partners in the hemi-irradiated series (1.4%). (14 refs)

6374 **Radiation-induced Soft-tissue and Bone Sarcoma.** (Eng) Kim, J. H. (Dept. Radiation Therapy, Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Chu, F. C.; Woodard, H. Q.; Melamed, M. R.; Vos, A.; Cantin, J. *Radiology* 129(2): 501-508; 1978.

The records of a single hospital over the past 50 yr revealed 10 cases of soft-tissue sarcoma arising from irradiated tissues and 27 cases of radiation-induced bone sarcoma. The patients with soft-tissue sarcomas ranged in age at the time of initial treatment from 7 to 44 yr, the mean and median latent period from first irradiation to the discovery of sarcoma being 12 and 16 yr, respectively. Nine of these patients were irradiated for benign conditions, 10 were irradiated for malignancies, and one was occupationally exposed to radiation. The patients with bone sarcomas ranged in age from 1 to 56 yr, the mean and median latent period from first irradiation to the discovery of sarcoma being 11 and 12 yr, respectively. Two patients were irradiated for benign conditions and 25 were irradiated for malignancies. Twenty-four additional cases of radiation-induced osteogenic sarcoma found in the literature resembled the current patients in sex ratio, radiation dose, and induction time. The median nominal standard doses of radiation associated with the soft-tissue and bone sarcomas were 1900 and 2270 rets (rads-equivalent-therapeutic), re-

spectively. No radiation-induced tumors were seen at doses below 1100 rets. For the bone sarcomas, the latent period was longer after larger radiation doses; and children appeared to be more susceptible to the development of cancer than did adults. (30 refs)

78-6375 **Chronic Mechanical Trauma in the Aetiology of Oro-pharyngeal Carcinoma.** (Eng) Thumfart, W. (Universitäts-HNO-Klinik, Waldstrasse 1, D-8520 Erlangen, W. Germany); Weidenbecher, M.; Waller, G.; Pesch, H. *J. J Maxillofac Surg* 6(3): 217-221; 1978.

The etiology of oropharyngeal carcinoma was studied in 86 patients with tumors of the oral cavity. Tumors of the gum and palate accounted for 62% of these tumors. Most (83%) of the patients were smokers and 49% regularly consumed alcohol, some to excess. In 44% of these patients, there was a coincidence of the cancer site with the site of persistent mechanical irritation by sharp teeth edges, prominent parts of an ill-fitting denture, or similar mechanical factors. Most of the tumors were keratinizing, poorly differentiated, squamous cell carcinomas, and all carcinomas in contact with a mechanical irritation had a surrounding wall of chronic inflammatory infiltration. Dysplasias of different degrees were found in the areas around the carcinomas. (16 refs)

78-6376 **Natural History of Radiation-associated Thyroid Cancer.** (Eng) Roudebush, C. P. (Thyroid Study Unit, Dept. Medicine, Univ. Chicago, Box 138, 950 E. 59th St., Chicago, IL, 60637); Asteris, G. T.; De Groot, L. *J. Arch Intern Med* 138(11): 1631-1634; 1978.

The clinical courses of 91 thyroid cancer patients who had received x-ray therapy (IR) between 1950 and 1975 were compared with those of a control group of thyroid cancer patients who had not received radiation therapy (RT). Although the two populations displayed similar profiles, several differences were apparent. In RT cases, the mean age was lower (28.2 yr vs 36.5), and, perhaps due to this, cervical adenopathy occurred more frequently (38.4% vs 25.4%). Treatment of RT patients had been more aggressive; more had total or near-total thyroidectomy (72.5% vs 56.9%) or received radioactive iodide (41.8% vs 21.9%). Primary papillary tumors were common in the RT group (87.8%). No anaplastic or medullary tumors were found, but invasive or metastatic disease (9.9% vs 4.2%) and multifocal disease (46.5% vs 15.9%) were more prevalent in these subjects. Moreover, recurrences were more common in the RT group (6.6% vs 1.4%). The death rate was similar in both groups. Parathyroid adenomas were observed in three irradiated patients, but none occurred in the controls. (20 refs)

78-6377 **Thyroid and Parathyroid Lesions after Radiation (Letter to Editor).** (Eng) Knights, E. M.

(South Macomb Hosp., Warren, MI, 48093); Phillips, E. *Ann Intern Med* 89(5, part 1): 725-726; 1978.

A 53-yr-old woman was found to have a solitary parathyroid adenoma, a papillary adenocarcinoma of the thyroid, and a colloid nodule of the thyroid. She had been treated with radiotherapy for acne at age 18. (3 refs)

- 78-6378 Radiation-induced Thyroid Carcinoma Following Treatment for Fetal Medulloblastoma.** (Pol) Borejko, J. (Klinika Radioterapii, Instytut Radiologii AM, Warsaw, Poland); Jeleniewska, A. *Pol Przegl Radiol* 42(3): 231-235; 1978.

Two patients (a girl and a boy aged 8 yr), received conventional radiotherapy for medulloblastoma. The girl received total doses of 2,500 and 2,800 R on two fields and an additional 3,300 R for a recurrence 5 yr later. She developed papillary carcinoma of the thyroid gland 16 yr after the first irradiation. The boy received a total dose of 3,500 R in 35 sessions in 67 days. He developed a mixed (papillary and follicular) thyroid carcinoma with metastases 10 yr later. (20 refs)

- 78-6379 Occurrence of Previous Neck Radiotherapy among Patients with Associated Non-medullary Thyroid Carcinoma and Parathyroid Adenoma or Hyperplasia.** (Eng) Tisell, L. E. (Dept. Surgery II, Sahlgren's Hosp., S-41345 Goteborg, Sweden); Hansson, G.; Lindberg, S.; Ragnhult, I. *Acta Chir Scand* 144(1): 7-11; 1978.

The medical records of 10/16 consecutive patients (12 women, 4 men) with coexistent primary hyperparathyroidism (HPT) and nonmedullary thyroid carcinoma revealed history of prior external radiation treatment to the neck. The ages at the time of first irradiation ranged from 2 mo to 51 yr (mean, 18 yr). The time lapse between first exposure and diagnosis of HPT and thyroid carcinoma ranged from 14 to 47 yr (mean, 36 yr). The thyroid tumor was limited to the gland in eight patients (Stage I). Regional lymph node metastases without concomitant cancer growth beyond the thyroid capsule were found in six patients (Stage II). In two patients, the tumor involved adjacent structures (Stage III); one of these patients also had lymph node metastases. The thyroid carcinoma was multifocal in four patients. There were 13 patients with a single parathyroid adenoma: in 8, a rim of compressed normal parathyroid tissue was found at the periphery of the adenoma; in 3, the removed parathyroid glands showed hyperplasia. In some of the enlarged parathyroid glands, hemorrhages as well as zones of fibrosis were present. Microscopically, the enlarged parathyroids in irradiated patients were similar to those of nonirradiated patients. The results agree with the hypothesis that radiation can serve as an etiological factor in the development of associated thyroid carcinoma and HPT. (17 refs)

- 78-6380 Radiation-induced Cancers in the Head and Neck Region.** (Jpn) Miyahara, H. (Dept. Otolaryngology, Kumamoto Univ. Sch. Medicine, H. Kumamoto 860, Japan); Matsuoka, H.; Ohbuchi, M.; Tani, M.; Sato, T.; Maeda, K. *J Otolaryngol Jpn* 81(3): 233-238; 1978.

A survey was made of 54 cases of radiation-induced cancer in the head and neck region. In 37 patients, the radiation had been applied as therapy for benign diseases, including 14 cases of tuberculous cervical lymphadenitis and 3 cases of dermal disease; in 14 patients, the radiation had been applied to treat malignant tumors, including 9 in the larynx and 5 in the thyroid. Among the radiation-induced cancers, there were 23 hypopharyngeal tumors, 14 thyroid tumors, and 17 cervicoesophageal tumors. Patients irradiated for malignant tumors had shorter intervals between treatment and development of radiation-induced cancer (av, 9.4 yr) than patients treated for benign conditions (av, 33 yr). Most of the tumors were well-differentiated squamous cell carcinomas of the pharynx and larynx; there were also 14 papillary and follicular adenocarcinomas of the thyroid gland and 1 fibrosarcoma of the upper lip. Studies show that the incidence of radiation-induced cancer is increasing. (16 refs)

- 78-6381 Promotion of Metastasis of C3H Mouse Mammary Carcinoma by Local Hyperthermia.** (Eng) Walker, A. (Glasgow Inst. Radiotherapeutics and Oncology, Belvidere Hosp., London Rd., Glasgow G31 4PG, Scotland); McCallum, H. M.; Wheldon, T. E.; Nias, A. H.; Abde, A. S. *Br J Cancer* 38(4): 561-563; 1978.

The response to hyperthermia of an infrequently metastasizing transplantable mammary carcinoma that originally arose spontaneously in C3H mice was investigated. Using a water heater that transfers heat via a thin membrane, heat was applied to 5- to 7-mm dorsal tumors in a temperature range of 42-46°C and a time range of 1-2 hr, time-temperature combination sufficient to eradicate the primary tumor. Fifty mice had their tumors locally controlled at 100 days of observation; time of death or sacrifice, if earlier. Of these mice, 7 died unexpectedly prior to routine autopsy and 14 were autopsied because of unexpected early death or physical deterioration. All 14 autopsied mice were found to have pulmonary metastases and, in some cases, metastases at other sites (4 to kidney, 3 to brown fat, and 1 each to heart, diaphragm, spleen, and lymph node). Taking only the proved metastases, the minimum rate for metastasis in this group was 14%, which may be contrasted to the max rate found previously in 14/131 mice whose tumors were locally controlled by radiation. (11 refs)

- 78-6382 Tumors in Animals Irradiated During Embryogenesis.** (Rus) Strel'tsova, V. N. (No affiliation given); Pavlenko-Mikhailov, Iu. N. *Vopr Onkol* 24(9): 25-28; 1978.

effect of irradiation during embryogenesis on tumor development was studied in male and female white rats whose fathers were irradiated with a single ^{60}Co dose of 100 rads (dose rate, 168 rads/min) on gestation days 7, 14, or 19. The animals that died at age 200-600 days were evaluated. The tumor induction rate was 184/219 vs 9/53 controls. The tumors included 52 breast tumors, 42 cases of leukemia, 53 lymphoid tumors, 11 ovarian tumors, 4 adrenal tumors, 3 thyroid tumors, 3 lung tumors, 3 bone tumors, and 1 tumor of the skin, uterus and intestine. The tumor incidence that occurred after irradiation during the embryonal (day 7) and fetal (day 19) periods was 69/107 and 44/70, respectively. Eighteen of the 42 animals irradiated during organogenesis (day 14) developed tumors by day 400, but animals irradiated during the embryonal and fetal periods developed tumors by days 120-180. The incidence of new tumors was highest during the last third of the life-span. Male rats irradiated during embryogenesis developed mammary gland and bone tumors, and females developed ovarian and bone tumors; these tumors were not seen in the controls. (12 refs)

6383 **Bilateral Pleural Thickening: A Manifestation of Asbestos Dust Exposure.** (Eng) Sargent, E. (Dept. Radiology, Los Angeles County-Univ. Southern California Medical Center, 1200 N. State St., Los Angeles, CA 90033); Gordonson, J.; Jacobson, G.; Birnbaum, W.; Hub, M. *Am J Roentgenol* 131(4): 579-585; 1978.

Forty-six asymptomatic patients with radiologic evidence of pleural thickening were discovered to have been occupationally exposed to asbestos dust, although this fact was clinically unsuspected. Pleural thickening most frequently involved the lower lung fields and was max at the level of the fifth through eighth ribs and costophrenic angles. There was no associated involvement of the parenchyma of the lung or the costophrenic angles. Most (67%) of the patients showed thickening up to 5 mm wide, and 8% showed thickening > 10 mm wide. The extent and severity of the pleural thickening did not correlate with the total duration of or time elapsed since asbestos exposure. Pleural thickening or plaques with no other evidence of disease in a relatively asymptomatic patient is a sign of possible exposure to asbestos dust. When retrospectively correlated with a lifetime occupational history, bilateral pleural thickening might serve as an indicator of asbestos dust exposure in the general population. (23 refs)

6384 **Renal Cell Carcinoma Following Radiation Therapy for an Adrenal Cortical Carcinoma.** (Eng) Andler, W. (Dept. Pediatrics, Univ. Essen, Hufelandstr. 55, D-4300 Essen 1, W. Germany); Havers, W.; Ambolis, C.; Medrano, J.; Stollmann, B. *J Pediatr* 93(4): 4-636; 1978.

The occurrence of renal cell carcinoma in an 11-yr-old girl approx 10 yr after radiation therapy for an adrenal cortical carcinoma is reported. The renal cell carcinoma was in all likelihood a direct consequence of the radiation therapy. (10 refs)

78-6385 **Influence of Misonidazole on the Incidence of Radiation-Induced Intestinal Tumours in Mice.**

(Eng) Rostom, A. Y. (Regional Centre Radiotherapy and Oncology, St. Lukes Hosp., Warren Road, Guildford, Surrey GU1 3NT, England); Kauffman, S. L.; Steel, G. G. *Br J Cancer* 38(4): 530-536; 1978.

The combined effects of misonidazole (MSD: 1 mg/g ip) and abdominal radiation in the range 16-24 Gray (Gy) units (1 unit = 100 rads) were studied in male C57BL mice. Eighty-one of 105 mice given radiation alone survived 50 days or longer, as did 81/106 mice sensitized with MSD 45 min before irradiation. Although the proportion of mice developing intestinal adenocarcinomas was similar in the two groups (60% and 68%, respectively) the incidence of multiple tumors was significantly higher in the MSD group (41% vs 18%). The av number of tumors per mouse was higher in the MSD groups at every radiation level, but the difference in the MSD and irradiation-only groups was most marked at a 20 Gy dose. With radiation alone, the most common tumor site was the cecum (49% of tumors) followed by the rectum (27%). MSD treatment reduced the cecal tumor incidence (37%), but it increased the rectal tumor incidence (40%). Most of the adenocarcinomas were mucin-secreting and located in regions of radiation damage. Although MSD may sensitize normal intestinal cells to the carcinogenic effect of radiation, these results do not call into question its present clinical use. However, there is the possibility that chemical radiosensitizers could enhance the incidence of second cancers in patients responding to radiotherapy. (15 refs)

78-6386 **Scenarios of ^{14}C Releases from the World Nuclear Power Industry from 1975 to 2020 and the Estimated Radiological Impact.** (Eng) Killough, G. G. (Health and Safety Res. Div., Oak Ridge Natl. Lab., Oak Ridge, TN); Till, J. E. *Nucl Saf* 19(5): 602-617; 1978.

Radioactive ^{14}C is formed as a by-product of nuclear power generation and a part is released to the environment. Estimated releases of ^{14}C from presently operating nuclear power facilities are still quite small in comparison with natural and other man-made sources, but potential growth in world demand for power will raise the quantity produced and released. The radiation dose to world population from natural and man-made ^{14}C is calculated based on three possible scenarios: (1) A pessimistic scenario (P) in which all the ^{14}C projected to be produced in fuel cycles is released. (2) An optimistic scenario (O) that assumes a decontamination fac-

tor of 100 for fuel reprocessing. (3) An intermediate scenario (I) that simulates a phased improvement in effluent treatment technology. The P scenario projects a ^{14}C release rate by the nuclear industry that exceeds natural production after the year 2000. The O and I scenarios project that ^{14}C release from the nuclear industry would attain a stationary level at approx 37% of natural production by 2020. Potential effects on the incidence of cancers and genetic disorders are calculated for the three scenarios from risk factors for exposure to ionizing radiation and ^{14}C specifically estimated by the International Commission on Radiological Protection. Total number of cancers and genetic effects over infinite time (effectively 46,000 yr or about eight half-lives of ^{14}C) are given for each of the three scenarios. For the O and I scenarios the cumulative potential health effects are about 22% of those from natural ^{14}C formed from 1975-2020, about 6% of those from

weapons testing, and $<0.04\%$ of those from natural already in the environment. (45 refs)

See also:

*(Rev.): 78-6004, 78-6008, 78-6025, 78-6031, 78-6045, 78-6046, 78-6047, 78-6048, 78-6051, 78-6052, 78-6055.

*(Chem.): 78-6108, 78-6111, 78-6120, 78-6154, 78-6196, 78-6235, 78-6244, 78-6313, 78-6325, 78-63

*(Immun.): 78-6490.

*(Path.): 78-6509, 78-6512, 78-6516, 78-6529.

*(Epid.-Biom.): 78-6547, 78-6554, 78-6564, 78-65

VIRAL CARCINOGENESIS

6387 Identification of Nucleotide Sequences Which May Encode the Oncogenic Capacity of Avian Retrovirus MC29. (Eng) Sheiness, D. (Dept. Microbiology, Univ. California, San Francisco, CA, 94143); Fanshier, L.; Shop, J. M. *J Virol* 28(2): 600-610; 1978.

The genome of MC29 virus, a retrovirus that induces a variety of tumors in chickens (ie, myelocytomatosis, kidney and liver carcinomas), and transforms cultures of embryonic avian macrophages and fibroblasts, was characterized, and nucleotide sequences that may encode the oncogenic potential of the virus were identified. MC29 virus can replicate only with the assistance of a related helper virus (MCAV), and the replication defect is apparently due to a deletion in one or more viral genes. The haploid genome of MC29 virus has a mol wt of approx 1.7×10^6 , whereas the MCAV genome has a mol wt of 3.1×10^6 . Although MC29 virus transforms avian fibroblasts in culture, its genome does not contain a gene homologous to the *src* gene responsible for fibroblast transformation by avian sarcoma viruses (ASV's). Radioactive DNA complementary to nucleotide sequences found in the MC29 virus genome (cDNA-MC29), but not in that of MCAV, were prepared. Hybridization data showed that the MC29 virus genome contained approx 1,500 nucleotides not found in MCAV, ASV, or Rous-associated virus 0, an oncogenic virus of chickens. These sequences could encode part or all of the gene(s) responsible for MC29 virus oncogenesis and transformation. The use of cDNA-MC29 will help determine the evolution of MC29 and may assist in identifying its transforming functions and in relating these functions to those of other leukosis-leukemia viruses. (43 refs)

6388 OK 10 Virus, an Avian Retrovirus Resembling the Acute Leukaemia Viruses. (Eng) Oker-Blom, N. (Dept. Virology, Univ. Helsinki and Coll. Veterinary Medicine, Helsinki, Finland); Hortling, L.; Kallio, A.; Nurmiaho, E. L.; Westermarck, H. *J Gen Virol* 30(3): 623-633; 1978.

The isolation and characterization of the OK 10 virus complex, which was derived from a liver tumor in a chicken that had been inoculated iv as an embryo with a field isolate of an avian leukosis virus, are described. Interference assays indicated that this virus complex contained both transforming (OK 10 V) and nontransforming (OK 10 AV) virus, both of subgroup A. OK 10 V, which could not be isolated in vitro without OK 10 AV, induced foci in chick embryo cells under agar overlay, and the cells from these foci formed colonies in soft agar. Injection of virus or infected cells into chicks resulted in acute leukemia but no local tumors. OK 10 AV, which comprised about 99% of the OK 10 complex, did not

induce foci in chick embryo cells, but it caused leukosis in chickens 17 mo after injection. OK 10 stocks showed typical C-type virus particles that had a density of 1.16 g/ml and contained 70S RNA. The latter released type b RNA subunits after heat denaturation. The data indicate that OK 10 virus is related to the subgroup C MH 2 virus or to the acute leukemia viruses of chickens. (33 refs)

6389 Rapid Oncogenesis In Vivo by Chicken Retrovirus OK10. (Eng) Hortling, L. (Dept. Virology, Univ. Helsinki, Haartmaninkatu 3, SF-00290 Helsinki 29, Finland). *Acta Pathol Microbiol Scand (B)* 86(4): 185-192; 1978.

The pathogenesis of OK10 virus infection was studied in 2- to 4-day-old Brown Leghorn chickens injected intraabdominally with 0.5 ml OK10 virus stock (titer: 5×10^4 focus-forming units/ml). The associated virus OK10AV was detected after 1 wk in the bursa of Fabricius, thymus, and liver, but not in spleen and bone marrow. OK10AV was detected in the spleen 3 wk after infection. The transforming virus OK10V was detected only in the bursa of Fabricius 2 wk after infection and in all tissues tested except bone marrow 3 wk after infection. Neutralizing serum antibodies against the OK10 virus complex were detected after 3 wk, when the first tumor was seen in the gut mesentery. After 6 wk, tumors were found in the liver but not in any other organs. When chickens were injected iv with OK10 virus, tumors could also be detected in the liver, kidneys, and testes. The mesenteric tumors were composed of two kinds of cells: clear cells with a large nucleus, prominent nucleoli, and cytoplasmic vacuolation and surrounding lymphocytes. OK10AV was infectious for chickens and induced neutralizing serum antibodies. One of seven chickens infected with OK10AV died of leukosis after 1.5 yr. Continuous cell lines, established from OK10 virus-induced tumors, produced both OK10V and OK10AV viruses and had a blastlike morphology. It is concluded that the OK10 virus complex, composed of a very oncogenic virus OK10V and an associated weakly oncogenic virus OK10AV, may be a good model for studying the mechanism of oncogenesis in vivo and in vitro. (33 refs)

6390 Temperature-sensitive Mutant of Avian Erythroblastosis Virus Suggests a Block of Differentiation as Mechanism of Leukaemogenesis. (Eng) Graf, T. (Institut für Virusforschung, Deutsches Krebsforschungszentrum, im Neuenheimer Feld 280, D-6900 Heidelberg, W. Germany); Ade, N.; Beug, H. *Nature* 275(5680): 496-501; 1978.

Virus-induced leukemic transformation was investigated by isolating a temperature-sensitive (*ts*) mutant of avian erythroblastosis virus (AEV). Erythroblast (EB) colonies were produced by infecting bone marrow cultures with mutagen-treated AEV at 35 C and seeding them into semisolid medium. Cells from individual colonies were propagated at 35 C. Aliquots of these clones were then shifted to 41 C or kept at 35 C and assayed for benzidine-positive (B+) cells. Three clones were *ts* for Hb expression, containing > 70% strongly B+ cells at 41 C and < 10% at 35 C. Clone 34 produced virus (*ts*34 AEV) capable of inducing EB colonies *ts* for Hb expression. *ts*34 AEV (1,000 colony-forming units) was less leukemogenic than the parent virus (10 units) in 8-day-old chicks: there was a decreased incidence of animals that died of leukemia and a delay of onset of disease and death. Most of the leukemic EB's from the peripheral blood of the mutant-infected chicks exhibited *ts* properties. It is suggested that AEV induces a block of differentiation in the erythroid target cells, that this block plays a crucial part in virus-induced leukemogenesis, and that the continuous expression of viral gene product is necessary to maintain the leukemic state of infected cells. (18 refs)

- 78-6391 At Least 104 Nucleotides Are Transposed from the 5' Terminus of the Avian Sarcoma Virus Genome to the 5' Termini of Smaller Viral mRNAs.** (Eng) Cordell, B. (Dept. Biochemistry and Biophysics, Univ. California, San Francisco, CA, 94143); Weiss, S. R.; Varmus, H. E.; Bishop, J. M. *Cell* 15(1): 79-91; 1978.

Based on the previous finding that all three forms of avian sarcoma virus (ASV)-specific cytoplasmic RNA contain a nucleotide sequence that is homologous to a sequence of 101 contiguous nucleotides found at the 5' terminus of the ASV genome, but not within the 3' half of the genome, it was proposed that this nucleotide sequence may be transposed to the 5' termini of viral messenger RNA's (mRNA's) during the genesis of these RNA's. Two strategies were used to substantiate this proposal. First, the three forms of ASV mRNA were isolated by molecular hybridization. Chemical analysis was then used to demonstrate that all three forms shared a capped nucleotide sequence from the 5' terminus of the viral genome, although other portions of the 5' half of the genome were not present in the smaller viral mRNA's. Second, single-stranded DNA complementary to a sequence of 101 nucleotides at the 5' terminus of the ASV genome (cDNA-5') was prepared, and it was shown that all three forms of ASV mRNA could form RNAase-resistant duplexes with the full extent of cDNA-5'. The RNA sequences included in these duplexes were capped and had identical compositions. In contrast, fragments of RNA derived from the 3' half of the ASV genome did not form duplexes with cDNA5'. It is concluded that a sequence of > 100 contiguous nucleotides is transposed from the 5' terminus of the ASV genome to the 5' termini of smaller virus-specific RNA's during the genesis of these RNA's. (53 refs)

- 78-6392 Nucleotide Sequences Related to the Transforming Gene of Avian Sarcoma Virus Are Present in DNA of Uninfected Vertebrates.** (Eng) Spector, H. (Dept. Microbiology, Univ. California, San Francisco, CA, 94143); Varmus, H. E.; Bishop, J. M. *Proc Natl Acad Sci USA* 75(9): 4102-4106; 1978.

Purified radioactive DNA complementary to the avian sarcoma virus (ASV) transforming gene (cDNA-sarc) was annealed with the DNA of several uninfected species under conditions conducive to pairing of partially matched nucleotide sequences (1.5 M NaCl, 59 C). Denatured DNA's of 10-11-day-old chicken embryos, liver and spleen of a GR mouse, human placental tissue, calf thymus, salmon sperm, sea urchin sperm, *Drosophila* cells grown in culture, *Escherichia coli* cells, and XC cells from a tumor in an ASV-infected chicken were incubated for 96-120 hr with ³H-cDNA-sarc. Duplex formation was measured by hydroxyapatite fractionation. cDNA-sarc formed duplexes with mammalian and fish DNA's (18%-43%), chicken DNA (55%-62%), and XC cells DNA (65%), but only slight annealing occurred with sea urchin DNA and none occurred with *E. coli* or *Drosophila* DNA. Kinetics studies indicated that the nucleotide sequences related to cDNA-sarc in species in which annealing occurred were present as single or only a few copies. The failure of total hybridization in these species was most probably due to stoichiometric limitations imposed by mismatched base pairs and not due to insufficient amounts of cDNA. The degree of base mismatching, as determined by the thermal stability of the duplexes, was approx 3%-4% for chicken DNA and 8%-10% for mammalian or fish DNA. These results indicate that there is a highly conserved set of nucleotide sequences to at least a portion of the ASV sarc gene in the DNA of these vertebrates. (29 refs)

- 78-6393 Specific Restriction of Avian Sarcoma Virus Replication by a Line of Transformed Lymphoid Cells.** (Eng) Neiman, P. E. (Dept. Medicine, Div. Oncology, Univ. Washington, Seattle, WA, 98195); McMillin-Helsel, C.; Cooper, G. M. *Virology* 89(2): 360-371; 1978.

The properties of viral replication on MSB-1 cells, a line of transformed chicken lymphoid cells derived from tumors induced by Marek's disease viruses, were examined. These cells are susceptible to infection by avian leukosis virus (ALV) subgroups A and C, including transformation-defective (non-oncogenic) mutants of avian sarcoma virus (ASV). Various parameters of viral growth revealed that infection of MSB-1 cells by *td* ASV was indistinguishable from infection of chicken embryo fibroblasts (CEF) by *td* ASV. However, the growth of wild-type ASV was restricted in these cells. Further studies, employing Hirt supernatant fractionation, suggested that, following infection of MSB-1 cells with ASV, infectious proviral sequences were synthesized, but the viral DNA did not integrate into host cell DNA in a normal fashion and was lost from the cell. Since replication of *td* ASV does not seem to be restricted in MSB-1 cells, it is concluded that the failure

SV DNA to integrate normally in these cells is due to presence of *src* sequences (genes that confer the ability to transform CEF and other cell types) in the viral genome. (68 refs)

6394 Effect of Glucosamine on Virus Production and Antigen Expression in Avian Sarcoma Virus-transformed Cells. (Eng) Israel, E. (Departement de Microbiologie et d'Immunologie, Universite de Montreal, Montreal, Quebec, Canada); Mandeville, R.; Wainberg, M. *Can J Microbiol* 24(9): 1024-1031; 1978.

The effect of glucosamine (20 mM) on virus production and antigen expression in chicken embryo fibroblast (CEF) cells transformed by avian sarcoma virus was studied. Significant quantities of transforming virus were not synthesized by cells exposed to glucosamine, and glucosamine significantly inhibited viral RNA synthesis and the levels of RNA-dependent RNA polymerase. The inhibition was reduced when glucosamine treatment was carried out in the presence of glucose. Glucosamine treatment also affected antigenic expression in transformed CEF, as measured by an indirect immunofluorescence test, but the degree of inhibition was far less than in a lymphocyte-stimulation assay for the detection of cell-mediated immunity was used. (22 refs)

6395 Synergistic Effect of Tumor Virus Transformation and Tumor Promoter Treatment on the Induction of Plasminogen Activation by Chick Embryo Fibroblasts. (Eng) Goldfarb, R. H. (Dept. Microbiology and Immunology, State Univ. New York, Downstate Medical Center, Brooklyn, NY, 11203); Quigley, J. P. *Cancer Res* 38(12): 4601-4609; 1978.

Studies were carried out to determine whether the tumor promoter phorbol myristate acetate (PMA) could further enhance the protease plasminogen activator (PA) activity in cultures of chick embryo fibroblasts transformed by Rous sarcoma virus (RSVCEF). PMA caused a 12- to 13-fold enhancement of the extracellular PA activity in these cultures, which in the absence of PMA produced at least 50- to 100-fold more PA than did normal chick embryo fibroblasts. The increase in RSVCEF PA activity was first detectable at a PMA concentration of 1 nanogram (ng)/ml, but was maximal when the PMA concentration was between 25 and 100 ng/ml. Increased levels of PA activity in RSVCEF were observed as early as 1-2 hr after PMA treatment. Sodium dodecylsulfate-polyacrylamide gel electrophoresis demonstrated that the PA produced by PMA-treated cultures had a molecular weight identical to that of the PA produced by untreated cultures. Induction by PMA could not be accomplished in cell-free extracts, but required protein synthesis in intact cells. Under serum-free conditions, PMA-treated RSVCEF secreted high levels of PA for 4-6 days and underwent pronounced morphological alterations; large dense colonies of round, refrac-

tile cells formed, and it appeared that at least 80%-90% of the cells in the culture dish were in colonial aggregates. Modified culture conditions and the use of PMA to induce PA allowed the accumulation of large amounts of RSVCEF culture fluid and subsequent purification of the enzyme. The sensitivity of transformed chick embryo fibroblasts to PMA and the generation of enhanced proteolytic activity in the cellular microenvironment may provide a model system to examine the role of both PA in malignant transformation and PMA in tumor promotion. (68 refs)

78-6396 Reversion of Transformed Glycolysis to Normal by Inhibition of Protein Synthesis in Rat Kidney Cells Infected with Temperature-sensitive Mutant of Rous Sarcoma Virus. (Eng) Carroll, R. C. (Dept. Biology, Univ. California at San Diego, La Jolla, CA, 92093); Ash, J. F.; Vogt, P. K.; Singer, S. J. *Proc Natl Acad Sci USA* 75(10): 5015-5019; 1978.

Normal rat kidney cells infected with a temperature-sensitive mutant (LA23) of Rous sarcoma virus exhibit the transformed phenotype when grown at the permissive temperature (33 C) and the normal phenotype when grown at the nonpermissive temperature (39 C). Since the addition of protein synthesis inhibitors to infected cells at the lower temperature causes them to revert to the normal phenotype with respect to morphological and cytoskeletal characteristics, studies were carried out to determine if there is also a reversion in metabolic characteristics. Two sets of experiments, performed at widely separated times, gave qualitatively similar but quantitatively different results. In the first experiments, lactate production was 50%-60% greater in LA23-infected cells at the permissive temperature than in those at the nonpermissive temperature; in the other experiments, the corresponding difference was 80%-100%. In both experiments, all cells showed an increased rate of glycolysis when dinitrophenol was added to the assay, and the control rate/dinitrophenol rate ratio was greater for the transformed phenotype than for normal. In the first experiment, oligomycin inhibited the glycolytic rate of LA23-infected cells at the permissive temperature by more than 70%, but it had no effect in cells grown at the nonpermissive temperature. The glycolytic rate in the transformed cells was unchanged by oligomycin in the second set of experiments, but the drug reproducibly increased the glycolytic rate by about 20% in cells exhibiting the normal phenotype. When transformed cells grown at the 33 C were treated with cycloheximide (15 ng/ml) or abrin (200 ng/ml), the rate of glycolysis without any other additives decreased to normal values. In the first set of experiments, cycloheximide treatment also caused a return to normal of the control rate/dinitrophenol rate ratio and eliminated the oligomycin sensitivity of the rate. Other metabolic changes that might be correlated with transformation were investigated in the second set of experiments: glucose transport, hexokinase activity, and phosphofructokinase activities. The results support the thesis that transformation by the *src* gene of the Rous sarcoma virus is a pleiotropic and

reversible process, such as that involved in a pleiotypic enzymatic modification reaction and its reversal. (31 refs)

- 78-6397 The Expression of Plasminogen Activator in Rous Sarcoma Virus-transformed Cells Is Controlled Both by the Virus and the Cell.** (Eng) Wolf, B. A. (Rockefeller Univ., New York, NY, 10021); Goldberg, A. R. *Virology* 89(2): 570-577; 1978.

The ability of virus produced by clones of Rous sarcoma virus (RSV)-transformed chicken embryo fibroblasts to elevate the level of cellular plasminogen activator (PA) was analyzed by a casein-plaque assay. The genotype of the virus produced by these transformed clones, which exhibit variable low to high levels of PA, was found to be identical to that of the parent virus, regardless of the PA level of the cell in which the virus is replicated. Virus from low PA clones was not defective in its ability to produce transformed cells with increased levels of PA. This virus produced the same spectrum of plaque-forming foci and soft-agar clones as virus from intermediate or high PA clones. These findings lead to the conclusion that the level of PA is cellularly determined. Additional evidence, however, such as the increased expression of PA by superinfection of low activator clones with another strain of RSV, the ability of some RSV strains to elevate PA levels, and the ability of cells infected with some temperature-sensitive *src* mutants to express properties of transformed cells at nonpermissive temperatures, suggests that PA levels are also influenced by the *src* (transforming) gene of RSV. (33 refs)

- 78-6398 Lack of Correlation Between Tumorigenicity and Level of Plasminogen Activator in Fibroblasts Transformed by Rous Sarcoma Virus.** (Eng) Wolf, B. A. (Rockefeller Univ., New York, NY, 10021); Goldberg, A. R. *Proc Natl Acad Sci USA* 75(10): 4967-4971; 1978.

The oncogenicity of chick embryo fibroblast (CEF) clones derived from infection with the replication-defective Bryan strain of Rous sarcoma virus (RSV) was studied to determine whether there is a correlation between tumorigenicity and the level of plasminogen activator (PA) in transformed fibroblasts. CEF clones transformed by B77 and Prague strains of RSV, previously isolated from agar suspension culture, varied in the expression of PA activity, but the tumorigenicity of the transformed cells could not be evaluated with clone cells because B77 and Prague RSV replicate and cause sarcoma in chickens. The range of properties exhibited by the Bryan RSV-derived clones fell within the continuum observed for the B77 and Prague clones. Thus, all clones exhibited an altered (round) morphology, a two- to fivefold greater rate of sugar transport, and a 10^5 - to 10^6 -fold higher efficiency of growth in agar suspension, regardless of the PA level. Five clones, transformed by the Bryan strain and showing low, intermediate, or high PA activity, were tested for tumorigenicity in newborn chicks. The three clones with low

activity exhibited little caseinolysis relative to the clone with the highest activity, even though 3-4 times more cells were assayed. At both cell concentrations tested (500 and 5000 cells/chick), all clones induced tumors at the two sites of injection. Latency periods were similar, regardless of the activator level of the clone injected. Every injected clone induced tumors regardless of its activator activity prior to injection, indicating that tumors exhibiting a spectrum of plasminogen activator activity. No correlations could be made with regard to the level of PA, size of excised tumor, number of cells surviving transplantation, or morphology of explanted tumors. These results confirm previous conclusions that tumorigenicity is not directly related with the level of plasminogen activator. (42 refs)

- 78-6399 Heterogeneity of RNA Synthesized in Lysates of Rous Sarcoma Virus Particles.** (Rus) Petrov, N. N. (N. N. Petrov Oncologic Res. Inst., Leningrad, USSR); Ratovitskii, E. A.; Korobitsyn, L. P.; Kniashuk, P. G.; Seits, I. F. *Dokl Akad Nauk SSSR* 241(3): 714-717; 1978.

The RNA polymerase of lysates of Rous sarcoma virus (RSV) particles was reacted with ^3H - or ^{32}P -labeled ribonucleoside triphosphates, and the product was characterized. The synthesis product was found to be sensitive to RNase A and KOH, but less sensitive to DNase. Electrophoresis in 2.5% polyacrylamide gels revealed two radioactivity peaks at 18S-20S and 4S; there was a shift toward 4S with increasing reaction time, which may indicate the presence of endogenous RNase activity. Synthesis of heteropolynucleotides was observed in the lysates in the presence of all four ribonucleoside triphosphates. Hybridization of the ^3H -labeled product of RNA polymerase reaction, isolated from the RSV lysate with RSV RNA demonstrated that 16% of the ^3H -labeled product has complementary segments with the virion RNA. Thus, the lysates of RSV particles contain an RNA polymerase that catalyzes the synthesis of both homopolynucleotides and heteropolynucleotides, and the synthesis of at least part of the RNA is a matrix reaction, with the matrix being RSV RNA. This fact is of particular interest in light of the generally held view that only RNA-dependent DNA polymerase activity is present in oncornaviruses. (10 refs)

- 78-6400 Complementary Interaction Between Genetic and Immunologic Mechanisms in Determining the Induction of Tumors by Rous Sarcoma Virus on the Chorioallantoic Membrane of the Chick Embryo.** (Eng) Bower, K. (Dept. Botany, Univ. Arkansas at Fayetteville, Fayetteville, AR, 72701); Gyles, N. R. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1645-1658; 1978.

udies were conducted to determine whether the genetic mechanism for regression has any influence on the number of tumors on the chorioallantoic membranes (CAM) of chick embryos and on the regression of tumors induced in young chicks by Rous sarcoma virus (RSV). CAM from 322 chick embryos originating from matings between parents of the rebred Progression and Regression lines were examined. Chick embryos descended from unchallenged regressor (R) parents and from progressor (P) parents were equally susceptible to induction of tumors on CAM by RSV. Chick embryos from immune R parents and those from a parental mating in which only the dam was an immune R demonstrated a marked reduction in tumor counts compared with embryos from P parents. The data suggest that a maternal factor transmitted from the immune hen through the egg interacts with the genotype of the embryo to reduce the number of CAM tumors. Three hundred and forty-four chicks, with parents from the same three breeding groups as those used for the CAM study, were injected in the wing web with RSV at age 1 day, 2 wk, and 4 wk. Day-old chicks descended from immune R parents demonstrated the greatest ability to effect tumor regression compared with chicks from all other mating combinations. This result also demonstrates the importance of an interaction between the maternal factor from immune parents and the genotype of the newly hatched chick in facilitating tumor regression. The influence of the maternal factor diminished sharply at 2 wk of age and was not detectable at 4 wk. (16 refs)

78-6401 Processing of 60,000-Dalton *sarc* Gene Protein Synthesized by Cell-Free Translation. (Eng) Amine, J. (Dept. Biology, Massachusetts Inst. Technology, Cambridge, MA, 02139); Buchanan, J. M. *Proc Natl Acad Sci USA* 75(9): 4399-4403; 1978.

New Zealand White rabbit antiserum directed against the 60,000-dalton (60K) transformation-specific antigen of Schmidt-Ruppin subgroup D Rous sarcoma virus (RD-RSV) immunoprecipitated both the 60K and 25K transformation-specific proteins synthesized by cell-free translation of virion RNA. However, upon incubation of the antiserum with ³⁵S-methionine-labeled protein from RD-RSV-infected cells, a 25K protein and one that was approximately 2K smaller than 60K were immunoprecipitated, which suggests that the primary translation product is processed intracellularly. Peptide mapping of sodium dodecyl sulfate/polyacrylamide gels indicated that the 60K translation product and the 60K cellular protein are closely related structurally. Addition of canine pancreas membrane vesicles to the reticulocyte lysate system during cell-free translation of transforming (*sarc*) gene messenger RNA specifically cleaved a 2K segment from the 60K protein so that it comigrated with the cellular species. Secretory proteins and, probably, at least some integral membrane proteins were synthesized with short hydrophobic signal sequences at their NH₂ terminus. The segment lost from the 60K transformation-specific protein is suggested to be a signal sequence, as the membrane

vesicles processed the 60K protein only during translation and the processed protein was sequestered by the vesicles. It is concluded that the 60K *sarc* protein is synthesized as a precursor with a signal sequence that functions to insert or transport the processed protein into or through cellular membranes. (28 refs)

78-6402 Reversion of Temperature Sensitive Transformation Mutants of Rous Sarcoma Virus and its Effect on the Expression of Tumour Specific Surface Antigen. (Eng) Wyke, J. A. (Imperial Cancer Research Fund, P.O. Box 123, Lincoln's Inn Fields, London WC2A 2PX, England); Kurth, R. *J Gen Virol* 40(3): 701-704; 1978.

To determine whether transformation and the expression of tumor-specific cell-surface antigen (TSSA) are causally related, reversion for TSSA expression was studied in the Rous sarcoma virus mutants *tsLA29* and *tsLA33*, which show a temperature-sensitive (*ts*) expression of TSSA, and their wild-type revertants. Chick embryo fibroblasts infected with *ts* virus showed low TSSA expression at the restrictive temperature (41°C), but cells infected by the two revertants expressed TSSA levels at this temperature comparable to those expressed at the permissive temperature (35°C). The revertants were all wild-type for transformation, as judged by focus formation and sugar transport, and they all showed a wild-type expression of TSSA. The results indicate that it is extremely unlikely that TSSA expression and transformation are unrelated, and they suggest that TSSA is necessary and sufficient for transformation or that TSSA expression is concomitant with but insufficient to account for the appearance of other parameters of transformation. (15 refs)

78-6403 Mechanism of Transfection of Chicken Embryo Fibroblasts by Rous Sarcoma Virus DNA. (Eng) Cooper, G. M. (Sidney Farber Cancer Inst., Boston, MA, 02115); Okenquist, S. *J Virol* 28(1): 45-52; 1978.

The mechanism of transfection of chicken embryo fibroblasts (CEF) by Rous sarcoma virus (RSV) DNA was investigated by assaying DNA-mediated transformation under conditions that restrict secondary virus infection (SVI). CEF that were genetically resistant to virus infection as a result of the absence of receptors for virus penetration were also resistant to transformation by integrated or unintegrated RSV DNA, as assayed by focus formation. Since resistance to RSV infection appears to be expressed at the level of virus penetration, these results indicate that focus formation by RSV DNA required formation of extracellular progeny virus and SVI of sensitive cells. DNA of replication-defective Bryan high-titer RSV was noninfectious, and transformation by DNA of a temperature-sensitive (*ts*) DNA polymerase mutant was *ts*. These results indicate that SVI was necessary for transformation by RSV DNA. Since transformation was also assayed by colony formation in soft agar, the requirement for SVI was not an

artifact of possible difficulties in detecting foci formed only by cell division. Thus, exposure to RSV DNA did not result in genetically stable transformation of the primary DNA-treated recipient cells. The results are consistent with the hypothesis that transfection of CEF by RSV DNA proceeded by transcription of donor DNA, formation of extracellular progeny virus, and SVI of sensitive cells. (30 refs)

- 78-6404 Transfer of Duck Cell DNA Sequences to the Nucleus of 3T3 Cells by Rous Sarcoma Virus.** (Eng) Baxt, W. G. (Biology Dept., Univ. California at San Diego, La Jolla, CA, 92037); Meinkoth, J. L. *Proc Natl Acad Sci USA* 75(9): 4252-4256; 1978.

The ability of a retrovirus [Prague C Rous sarcoma virus (RSV)] to transduce incorporated host polynucleotide sequences into the nucleus of a second unrelated host cell was studied. Upon infection of Peking duck cells by Prague C RSV, polynucleotide sequences complementary to Peking duck cell nuclear DNA were detected in the high-mol-wt RNA from progeny Peking duck cell-passaged RSV (RSV-PD) by RNA-DNA hybridization. When mouse 3T3 cells were subsequently infected by RSV-PD, polynucleotide sequences complementary to Peking duck cell nuclear DNA were detected in the 3T3 cell nuclear DNA by RNA-DNA molecular hybridization. The data introduce the possibility that passage of intact genes or portions of genes between species might occur after evolutionary divergence has taken place. In this manner, the gene composition of organisms could undergo modulation at a rate considerably greater than the one that would result from random mutations alone. (38 refs)

- 78-6405 Isolation and Characterization of a Virus-specific Ribonucleoprotein Complex from Reticuloendotheliosis Virus-transformed Chicken Bone Marrow Cells.** (Eng) Wong, T. C. (Dept. Microbiology, Univ. Texas Health Science Center at Dallas, Southwestern Medical Sch., Dallas, TX, 75235); Kang, C. Y. *J Virol* 28(1): 34-44; 1978.

A ribonucleoprotein (RNP) complex containing endogenous RNA-directed DNA polymerase (RDDP) activity was isolated from the cytoplasm of reticuloendotheliosis virus (REV)-transformed chicken bone marrow cells (BMC) but not from normal BMC. This RNP complex has a density of 1.23 g/cm³ and sediments at 80S-100S in sucrose gradients, properties that indicate that it is distinct from the mature virion. RDDP activity was completely neutralized by anti-REV DNA polymerase antibody but not by anti-avian myeloblastosis virus DNA polymerase antibody. The DNA product from the endogenous RDDP reaction of the RNP complex hybridized to REV RNA but not to avian leukosis virus RNA. The RNA extracted from the RNP hybridized only to REV-specific complementary DNA (cDNA) synthe-

sized from an endogenous DNA polymerase reaction of purified REV. Max hybridization was only 40%; the significance of this apparently low hybridization efficiency is not clear. The size of the RNA in the RNP is 30S-35S, a size equivalent to one subunit of the RNA genome of the virus. No mature genomic RNA was found within the RNP complex. Thus, the formation of mature viral genomic RNA seems to occur after budding, probably during the condensation of nucleoid. (32 refs)

- 78-6406 Protection Against Marek's Disease-derived Tumor Transplants by the Nononcogenic Strain of Marek's Disease Virus.** (Eng) Schat, K. A. (Dept. Avian and Aquatic Animal Medicine, New York State College of Veterinary Medicine, Cornell Univ., Ithaca, NY, 14853); Calnek, B. W. *Infect Immun* 22(1): 225-232; 1978.

The ability of a nononcogenic strain of Marek's disease virus (SB-1) to protect against three Marek's disease-derived tumor transplants was studied in intact, embryonally bursectomized (Bx), or thymectomized (Tx) or cyclophosphamide (CP)-CP/Tx-treated S, N, P, and G-B1 chickens. The chickens were challenged with JMV, a non-virus-producing tumor transplant, or GA/Tr-1 or MDT-198, two virus-producing tumor transplants. Protection was significant when SB-1 (5 × 10⁵ focus-forming units) and JMB (10⁵ viable cells) were administered together, but protection was optimum when challenge was performed 7 days' postvaccination. Protection was abolished by increasing the number of JMV tumor cells to 10⁶ cells or by CP/Tx treatment. Tx, CP, or Bx alone did not ameliorate the immune response. Inactivated SB-1-infected cells (equivalent to 14,300 focus-forming units of live SB-1) were unable to protect against JMV challenge. SB-1 protected against GA/Tr-1, as evidenced by enhanced tumor rejection and the absence of metastases and virus-induced tumors, but not against MDT-198 in intact syngeneic and allogeneic chickens. Protection against MDT-198 was present in vaccinated Bx chickens. It is concluded that protection against JMV is T-cell-dependent and requires the induction of neoantigens not present in the inactivated SB-1 cellular preparation. Several possible explanations for the different responses to GA/Tr-1 and MDT-198 are given. (37 refs)

- 78-6407 70 Nm Particles Detected in Lymphoblastoid Cell Lines Derived from Marek's Disease Tumors.** (Eng) Nii, S. (Dept. Pathology, Res. Inst. Microbial Diseases, Osaka Univ., Yamada-kami, Suita, Osaka, Japan); Yasuda, I.; Ikuta, K. *Biken J* 20(3-4): 151-154; 1977.

The ultrastructure of the 70-nanometer (nm) particles found in two lymphoblastoid cell lines derived from Marek's disease lymphomas (MSB-1 and MOB-1) was determined. Intracellular viral capsids of about 100 nm diameter were found in a few cells from both lines, and about 25% of these capsid-bearing cells also contained the spherical 70-nm particles.

latter were not detected in duck fibroblasts infected with Marek's disease virus, although another aberrant form of the virus could be found in both infected duck embryo fibroblasts and lymphoblastoid cell lines. Most 70-nm particles appeared to consist of an inner spherical structure of high electron density and outer, radially arranged projections of moderate density. It is not certain whether the particles actually have a substructure with rod-shaped projections or whether these projections represent cross-sections of strandlike structures that coil around the inner spherical structure. It is suggested that the 70 nm particle is an aberrant form of Marek's disease virus. (8 refs)

78-6408 Induction of Intracranial Neoplasms by Intracerebral Inoculation of the Bratislava-77 Strain of Avian Sarcoma Virus in Perinatal and Adult F344 Rats: Influence of Age at Inoculation on Tumor Incidence and Morphology and Post-Inoculation Survival. (Eng) Copeand, D. D. (Duke Univ., Durham, NC). *Diss Abstr Int B* (5): 2229; 1978. (no refs)

78-6409 Modulation of Mouse Mammary Tumor Virus Production in the MJY-Alpha Cell Line. (Eng) Magi, M. J. (Dept. Neoplastic Diseases, Mount Sinai Sch. of Medicine, New York, NY, 10029); Blair, P. B.; Lane, M. A. *J Virol* 28(2): 611-623; 1978.

Production of mouse mammary tumor virus (MMTV) virions by the mouse mammary tumor cell line MJY-alpha was shown to be reversibly suppressed by in vivo implantation of the cells into isogenic MMTV-free BALB/c or MMTV-infected BALB/cfC3H mice. This suppression was rapid; the decrease in MMTV particles was observed as early as 7 days after implantation and it remained constant with further in vivo passage of the MJY-alpha tumors. Examination of thin tumor sections by electron microscopy revealed that intracytoplasmic A particles, budding particles, and cell-free MMTV B particles were all affected. However, immunofluorescence assays of the sections demonstrated the presence of MMTV viral antigens in the cells. Cell cultures initiated from the tumors also produced extremely low numbers of virions during the first two passages; after this, the level of MMTV virion synthesis increased and, in later subcultures, was similar to that of the parent cell line. Polypeptide profiles obtained by sodium dodecyl sulfate-polyacrylamide gel electrophoresis of virions purified from these cultures were identical to MMTV. Immunodiffusion demonstrated a cross-reactivity between these virions and MMTV particles obtained from mouse milk. Treatment of MJY-alpha cell cultures with rabbit anti-MMTV antiserum reduced the number of extracellular MMTV virions and altered their electrophoretic polypeptide patterns. Possible reactions for virion particle suppression in vivo are given. (25 refs)

78-6410 Mammary Tumour Virus and Casein Gene Transcription During Mouse Mammary Development. (Eng) Pauley, R. J. (Dept. Cell Biology, Baylor Coll. Medicine, Houston, TX, 77030); Rosen, J. M.; Socher, S. H. *Nature* 275(5679): 455-457; 1978.

Mouse mammary tumor virus (MMTV) RNA and casein messenger RNA (mRNA) were quantitated in developing mammary tissue from BALB/c and BALB/cfC3H mice (strains that have a low and high incidence of mammary tumors, respectively) by molecular hybridization with specific complementary DNA (cDNA) probes to MMTV 70S RNA and casein 15S mRNA. The level of MMTV RNA in the mammary tissues was below the limit of detection in virgin and early pregnant BALB/cfC3H mice and was low (0.0014% of the RNA) in mice at midpregnancy. MMTV RNA increased to 0.0125% in the BALB/cfC3H mammary gland in late pregnancy and increased twofold after parturition. The level of MMTV RNA in BALB/c mammary tissues at all the developmental stages assayed was $\leq 0.0005\%$. Casein mRNA was present at 0.0009% of the RNA in the virgin mammary gland, at 0.0066% in early pregnancy, and at 0.365% late in pregnancy. Casein mRNA increased and was maintained at 1% of the RNA during lactation. Following weaning, casein mRNA decreased to 10% of the lactating level, but this level was higher than that observed in the virgin gland. Changes in casein mRNA levels were essentially the same in BALB/c and BALB/cfC3H mice. These studies indicate that the hormonal milieu required for mammary development and lactogenesis enhanced both casein and MMTV expression in the mammary gland of BALB/cfC3H mice. Following weaning, MMTV RNA did not decrease to the same degree as casein mRNA. Therefore, the accumulation of MMTV RNA in BALB/cfC3H mammary glands is dependent upon mammary development and influenced by differentiated mammary function. (28 refs)

78-6411 Hormonal Induction of Mammary Tumor Viruses and Its Implications for Carcinogenesis. (Eng) McGrath, C. M. (Dept. Biology, Michigan Cancer Foundation, 110 E. Warren, Detroit, MI, 48201); Jones, R. F. *Cancer Res* 38(11, part 2): 4112-4125; 1978.

The possibility that hormones and mouse mammary tumor virus (MuMTV) are cocarcinogenic for mammary epithelium was studied in BALB/c and BALB/cfC3H mice. Rates of premalignant (PM) and malignant (M) transformation were compared in cells carrying genetically (BALB/c) or horizontally (BALB/cfC3H) transmitted MuMTV proviral sequences in animals with or without pituitary implants to stimulate hormonal production. The carcinogenesis rate in BALB3cfC3H females was substantially faster than that in BALB3c mice, but the final mammary carcinoma incidence was about the same. Hormonal stimulation promoted PM transformations in noninfected as well as infected females, but the efficiency of PM transformation was substantially greater in infected than in noninfected females. The

rate-limiting step, which infectious MuMTV overcomes, was in PM transformation from the normal state. Integration of new MuMTV sequences in mammary cell DNA was detected by molecular hybridization in cells containing horizontally transmitted MuMTV but not in those germinally infected. MuMTV expression was changed in mammary epithelial cells during hormonal carcinogenesis. Expression of envelope glycopeptides (glycoprotein with a mol wt of 52,000; gp52) was induced, which correlates with amplification of MuMTV RNA sequence content. MuMTV RNA and gp52 were induced by dexamethasone in PM and M cells, but not in normal cells. Both PM and M cells had altered surfaces; the alteration may include membrane incorporation of MuMTV-related determinants of gp52. A hormonal cocarcinogenesis hypothesis is proposed in which expression of germinally transmitted MuMTV genes is the proximal cause of neoplastic transformation. (60 refs)

- 78-6412 Occurrence of Antibody Against Intracytoplasmic A-Particles of Mouse Mammary Tumor Virus in Sera from Breast Cancer Patients.** (Eng) Ogawa, H. (Dept. Surgery, Faculty Medicine, Kyoto Univ., Shogoin-Kawahara-cho, Sakyo-ku, Kyoto 606, Japan); Tanaka, H. *Gann* 69(4): 539-544; 1978.

Antibody to intracytoplasmic A particles (pronucleocapsids of mouse mammary tumor virus) was detected in the sera of 26/43 Japanese breast cancer patients and 4/16 mammary fibroadenoma patients, but in only 4/37 controls. The antibody was detected by indirect immunofluorescence using acetone-fixed smears of A-particle-forming DBA/2 mouse leukemia cells. A-particle clusters fluoresced in discrete granules in the cytoplasm. The presence of the antibody was not correlated with a particular type of tumor histology, blood group, or clinical stage. Twenty of the breast cancer patients were followed after tumor resection: 10/13 antibody-positive patients became negative within 1 yr, and these 10 patients are apparently healthy. Three antibody-negative patients became positive after surgery, but two eventually reverted. Four patients remained negative, three of whom died or suffered a tumor recurrence. The antibody incidence in the breast cancer patients is higher than that detected previously in non-Japanese populations, and a possible ethnic difference is suggested. (28 refs)

- 78-6413 Characterization of a New Virus from *Mus cervicolor* Immunologically Related to the Mouse Mammary Tumor Virus.** (Eng) Schlom, J. (Lab. Viral Carcinogenesis, Div. Cancer Cause and Prevention, NCI, NIH, Bethesda, MD, 20014); Hand, P. H.; Teramoto, Y. A.; Callahan, R.; Todaro, G.; Schidlovsky, G. *J Natl Cancer Inst* 61(6): 1509-1519; 1978.

A virus similar to the murine mammary tumor viruses (MuMTV) of the laboratory mouse *Mus musculus* was identi-

fied in the milk of *M. cervicolor popaeus* mice and characterized. The virus was morphologically indistinguishable from B-type MuMTV and was given the designation MC-MTV. Radioimmunoassays for the 52,000-dalton major envelope glycoprotein and the 28,000-dalton major internal protein of MuMTV demonstrated that MC-MTV shared some antigenic determinants with both of these MuMTV proteins. The reactivity was different, however, from that observed with MuMTV tested from *M. musculus*. MC-MTV had a density of 1.16 g/ml in sucrose and a virion-associated DNA polymerase with a divalent cation preference for Mg²⁺ over Mn²⁺. Radioimmunoassay differentiated MC-MTV from the three other viruses previously identified from *M. cervicolor*. MC-MTV is the first virus from another species that is immunologically related to the MuMTV of *M. musculus*. B-type particles similar to MC-MTV were observed in a spontaneous *M. cervicolor popaeus* mammary tumor, and they are being characterized to determine their relationships to the milk-borne MC-MTV particles. (38 refs)

- 78-6414 Detection of Mammary Tumor Virus Antigen in DD/Tbr Strain Mice and the Hybrids.** (Eng) Imai, S. (Second Dept. Pathology, Nara Medical Univ., Nara, Japan). *J Nara Med Assoc* 29(2): 482-489; 1978.

Immunological methods were used to study the mode of transmission of murine mammary tumor virus (MuMTV) antigen expression in DD/Tbr (DD) mice. The incidence of mammary tumors declined in DD mice when they were raised by C57BL/6J mothers; the DD offspring thus raised were designated DDfB. MuMTV antigen was detected by immunodiffusion in the milk of DD mice, BfDD mice (C57BL/6J mice foster-nursed by DD/Tbr mothers), and (female BALB/c x male DD/Tbr)F₁ hybrids, but no antigen was detected in DDfB, BALB/c, C57BL, or (female C57BL/6J x male DDfB)F₁ mice. The transmission of MuMTV through the milk in the DD strain was thus confirmed. Radioimmunoassay revealed MuMTV antigen in the milk of 2/2 DDfB mice and 3/6 (C57BL/6J x DDfB)F₁ mice. The data suggest that DD mice might harbor endogenous MTV. (27 refs)

- 78-6415 Evidence for a Precursor-Product Relationship Between Intracytoplasmic A Particles and Mouse Mammary Tumour Virus Cores.** (Eng) Smith, G. H. (Lab. Molecular Biology, NCI, NIH, Bethesda, MD, 20014). *J Gen Virol* 41(1): 193-200; 1978.

The relationship between intracytoplasmic A particles (CAP) and mouse mammary tumor virus (MMTV) was investigated. CAP were purified from transplantable Leydig cell tumors or from primary mammary tumors in C3H/He mice. Electrophoretic and immunodiffusion techniques detected three MMTV-specific antigenic determinants in CAP (MMTV p27, p14, and p10) and provided evidence that showed that

... sizes and antigenic specificities of these determinants were similar to those of the CAP polypeptides p82, p27, p20-18 and p15-10. The results suggest that CAP possess incompletely processed virion proteins and are intracellular structural precursors to MMTV cores. It is concluded that CAP structural proteins are modified during maturation by protease, i.e., they are reduced to a size comparable to that of the major internal proteins of the virion and that this reaction is determined by the folding of CAP proteins into the morphologically intact A particle. (20 refs)

78-6416 Comparative Susceptibilities of Hybrid Mice to Four Mammary Tumor Viruses from Two Mouse Strains. (Eng) Moore, D. H. (Dept. Microbiology and Immunology, Hahnemann Medical Coll. and Hosp., Philadelphia, PA, 19102); Merryman, C. F.; Maurer, P. H.; Holbrook, J. A. *Cancer Res* 38(11, part 1): 3871-3878; 1978.

The effect of injection of milk containing one of four different strains of murine mammary tumor virus (MuMTV) into several mouse strains and their F₁ hybrids was measured by the recurrence of mammary tumors and the presence of viral antigen in the milk after the third and later litters. BALB/c mice were very susceptible to infection with strain A virus, but C57BL mice and (C57BL x BALB/c)F₁ hybrids were not. Strain Af virus had a low infectivity in all three groups of mice. RIII virus was very infectious in C57BL but less infectious in RIII mice. The susceptibility of (RIII x C57BL)F₁ hybrids was similar to that of C57BL mice. RIII virus was moderately infectious in RIII mice but poorly infectious in both C57BL mice and (RIII x C57BL)F₁ hybrids. However, RIII was highly infectious and produced tumors at an early age in BALB/c mice and (RIII x BALB/c) F₁ hybrids. These results demonstrate the marked influence that the father can exert on the susceptibility of hybrids to a particular virus. There was a good correlation between incidence of infection and incidence of mammary tumors. These experiments point out the complexity of the mouse assay for MuMTV and the erroneous conclusions that may be drawn from a single experiment. (38 refs)

78-6417 Suppression of Chemical (DEN) Carcinogenesis in SWR/J Mice by Goat Antibodies Against Endogenous Murine Leukemia Viruses. (Eng) Pottathil, R. Jackson Lab., Bar Harbor, ME, 04609; Huebner, R. J.; Meisner, H. *Proc Soc Exp Biol Med* 159(1): 65-68; 1978.

The possible involvement of endogenous murine leukemia virus (MuLV) in chemical carcinogenesis in vivo was tested using SWR/J mice treated with diethylnitrosamine (DEN: 0.0 mg/kg ip). The mice were treated with goat anti-radiation leukemia virus (RadLV) and AT124 MuLV IgG before and after DEN treatment, and their effects on the induction and latency of lung tumors were followed. Fifteen of 20 DEN-treated mice immunized with AT124 IgG died from

lung tumors; 50% mortality from lung tumors occurred at 54 wk after treatment. Thirteen of 20 unimmunized controls died with multiple lung tumors; the 50% mortality occurred by the 29th wk. Mice immunized with anti-RadLV IgG survived up to 64 wk after DEN treatment, with 18/20 mice developing lung tumors; 50% mortality occurred at 58 wk. Control mice survived only to 38 wk, with 12/20 developing lung tumors. Immunized mice had larger lung tumors than nonimmunized controls. These data indicate that the survival of antiviral IgG-treated mice was significantly prolonged over that of nonimmunized mice. The antiviral IgG's had apparently very little effect, if any, on tumor growth rate. Although the mechanism of the suppression of chemical carcinogenesis by antiviral IgG's is not clear, the findings indicate that viruses are involved in chemical carcinogenesis. (14 refs)

78-6418 Rapid In Vitro Assay for Thymotropic, Leukemogenic Murine C-Type RNA Viruses. (Eng) Lieberman, M. (Cancer Biology Res. Lab., Dept. Radiology, Stanford Univ. Sch. Medicine, Stanford, CA, 94305); Declève, A.; Kaplan, H. S. *Virology* 90(2): 274-278; 1978.

The in vitro quantitation of thymotropic and leukemogenic murine C-type RNA viruses is described. When serial dilutions of murine leukemia viruses (MuLV's) from C57BL/Ka mice were plated on a nonproducer lymphoma cell line derived from a radiation-induced lymphoma of this strain, end-point titers were obtained that correlated well with those obtained by titration methods in vivo. This rapid in vitro assay may be extended to the titration of a variety of MuLV's. (14 refs)

78-6419 Functional Macrophage Cell Lines Transformed by Abelson Leukemia Virus. (Eng) Raschke, W. C. (Developmental Biology Lab., Salk Inst., PO Box 1809, San Diego, CA, 92112); Baird, S.; Ralph, P.; Ninkovic, I. *Cell* 15(1): 261-267; 1978.

Three cloned cell lines that express properties of macrophages were established from murine tumors induced with Abelson leukemia virus (A-MuLV). Two of the three original tumors also yielded lymphocyte cell lines, one typical of the Abelson virus disease and the other a thymic lymphoma. Two of the macrophage lines were tumorigenic when placed in syngeneic BALB/c mice, producing typical A-MuLV lymphocyte neoplasms. All three macrophage lines pinocytosed neutral red, phagocytosed zymosan and latex beads, killed antibody-coated sheep RBC targets, and secreted lysozyme. The co-isolated lymphocyte lines showed none of these properties. Of two macrophage cell lines tested, neither was capable of replacing the adherent cell population required for the induction of in vitro immune responses. Bacterial lipopolysaccharide, an agent that activates normal macrophages, specifically inhibited the growth of the transformed macrophages in culture. The secretion of A-MuLV by two of the

macrophage lines provides conclusive evidence that the Abelson virus is capable of productively infecting the macrophage cell type. The other macrophage line became positive for Abelson virus production after rescue by Moloney leukemia virus. (21 refs)

- 78-6420 Translation of Type C Viral RNAs in *Xenopus laevis* Oocytes: Evidence that the 120,000-Molecular-Weight Polyprotein Expressed in Abelson Leukemia Virus-transformed Cells is Virus Coded.** (Eng) Reynolds, R. K. (Lab. Cellular and Molecular Biology, NCI, Bethesda, MD, 20014); van de Ven, W. J.; Stephenson, J. R. *J Virol* 28(2): 665-670; 1978.

The genomic RNA of two representative mammalian replication-defective transforming viruses, Abelson leukemia virus (AbLV) and woolly monkey sarcoma virus (WMSV), was purified and translated in *Xenopus laevis* oocytes. The major AbLV-specific protein synthesized was a polyprotein (pp) similar in mol wt and immunological properties to a previously described 100,000- to 130,000-dalton pp expressed in AbLV-transformed cells that contained the two amino terminal gag gene proteins p15 and p12. Translation of WMSV RNA resulted in the synthesis of a 55,000-dalton pp consisting of woolly helper virus p30, p15, and p12. The results establish that the nonstructural portion of the 110,000- to 130,000-dalton pp expressed in AbLV-transformed cells is virus-coded. The in vitro translation system may provide a means of testing sera from various sources for antibody directed against their nonstructural components. (20 refs)

- 78-6421 *Fv-1* Regulation of Lymphoma Development and of Thymic, Ecotropic and Xenotropic MuLV Expression in Mice of the AKR/J x RF/J Cross.** (Eng) Mayer, A. (Dept. Genetics and Pathology, Albert Einstein Coll. Medicine, Bronx, NY, 10461); Duran-Reynals, M. L.; Lilly, F. *Cell* 15(2): 429-435; 1978.

The incidence of spontaneous thymic lymphoma in crosses between AKR/J and RF/J mice was studied. AKR mice developed almost a 100% incidence of spontaneous lymphocytic thymic lymphoma. The incidence was much lower in the (AKR x RF)_F hybrids and in about 50% of the (AKR x RF) x AKR backcross mice. This resistance was associated with a reduction in endogenous ecotropic and xenotropic murine leukemia virus (MuLV) expression in the prelymphomatous thymus. The data indicated that the RF mice transmitted an allele at a single locus that resulted in a dominant suppression of lymphoma-inducing, AKR-transmitted genes. This allele appeared to map at the *Fv-1* locus closely linked to the *Gpd-1* locus. The results indicate that the particular *Fv-1* allele of AKR mice provides a permissive genetic background for endogenous ecotropic and xenotropic MuLV expression and that these viral activities may be etiologically involved in the development of spontaneous thymic lymphoma in the mouse. (18 refs)

- 78-6422 Age-dependent Relaxation of Gene Repression: Increase of Endogenous Murine Leukemia Virus-related and Globin-related RNA in Brain and Liver of Mice.** (Eng) Ono, T. (Gerontology Res. Center, Natl. Inst. Aging, Baltimore City Hosps., Baltimore, MD, 21224); Cline, R. G. *Proc Natl Acad Sci USA* 75(9): 4431-4435; 1978.

In order to determine whether a slow loss of differentiation by epigenetic and/or mutational mechanisms could be a common denominator in mammalian aging, any age-dependent expression of specific genes not normally expected to be expressed was investigated. ³H-labeled complementary DNA (cDNA) to the murine leukemia C-type RNA virus (MuLV) genome and globin genes were used as probes to determine the presence of RNA complementary to these genes in RNA extracted from brain and liver of young (6-7 mo), middle-aged (19-20 mo), and old (26-27 mo) C57BL/6J mice. Significant amounts of globin RNA were found in brain nuclei and cytoplasm. There was no age difference in the globin RNA sequences present, but the number of globin RNA molecules increased from 15 in 6-mo-old mice to 34 in 27-mo-old animals for nuclei and from 280 to 500 for cytoplasm. Similar results were found in the liver. RNA complementary to the MuLV cDNA probe was also found in brain and liver nuclei. In contrast to globin, the different RNA sequences increased in brain and liver nuclei from about 45% of the MuLV genome in 6-mo-old animals to 72% in those 27 mo old. The number of MuLV-related RNA molecules remained constant at about 22 and 38 per nuclei for brain and liver, respectively. Derepression of genes of this magnitude could result in a time-dependent increase of virus-related diseases and a general deterioration of the organism. (34 refs)

- 78-6423 Heteroduplex Analysis of the Sequence Relationships Between the RNAs of Mink Cell Focus-inducing and Murine Leukemia Viruses.** (Eng) Chien, Y. H. (Dept. Chemistry, California Inst. Technology, Pasadena, CA, 91125); Verma, I. M.; Shih, T. Y.; Scolnick, E. M.; Davidson, N. *J Virol* 28(1): 352-360; 1978.

Electron microscope heteroduplex analysis was used to study the sequence relationships between AKR ecotropic virus and an AKR-derived mink cell focus-inducing (MCF) isolate (AKR MCF 247), between Moloney murine leukemia virus (M-MLV) and an M-MLV MCF isolate (M-MLV_{MCF}), and between AKR and M-MLV. For the MCF viral RNA of AKR origin and the MCF viral RNA of M-MLV origin, the MCF-specific sequences were located in a region extending from 1.95 to 2.75 kilobases (kb) from the 3' end. Studies of the sequence relationship between AKR and M-MLV showed four major sequence nonhomology regions centered at 1.1, 2.3, 3.3, and 6.8 kb from the 3' end of viral RNA, with lengths of 0.5, 0.6, 0.5, and 0.5 kb, respectively. Further observations suggested that there is some degree of partial sequence mismatch in the entire region from 0.9 to 3.5 kb. However, the AKR and M-MLV sequences immediately adjacent to the 1.95- and 2.75-kb junctions with MCF-specific

ences were relatively similar in AKR and M-MLV. The results suggest that the *env* gene of MLV's maps from 1 to 5 kb from the 3' end of the genomic RNA and that the carboxyl end of the glycoprotein of each MCF strain is similar and identical to that of its ecotropic parent. Thus, the physical distance from heteroduplex structures is consistent with the hypothesis, derived from biological properties and tryptic peptide analysis, that the MCF viruses are recombinant with the *env* gene. (25 refs)

78-6424 Erythropoietin Responsiveness In Vivo of Friend Virus Target Cells for Transformation and Virus Replication. (Eng) Nasrallah, A. G. (Div. Experimental Biology, Baylor Coll. Medicine, Houston, TX, 77030); McGarry, M. P. *Cancer Res* 38(11, part 1): 4-3778; 1978.

In vivo cell transformation by Friend virus (FV) was investigated by a technique that allows selective stimulation of the erythropoietin-responsive cells (ERC) in mice with suppressed multipotential hemopoietic stem cells. DBA/2 mice previously made plethoric by hypertransfusion were administered busulfan (BU: 60 mg/kg intragastrically) in order to depress stem cell populations. On each of the following 3 days, mice received either 8 IU erythropoietin (Ep) or saline injections. Experimental and control mice were injected with equivalent doses of polycythemia-inducing FV. A comparison of the number of target cells transformed in the mice given Ep with those not given Ep provided evidence for the FV as the in vivo compartment in which the target cell for transformation resides. FV-transformed cells were determined by the spleen tumor colony-forming unit assay. Differences in plasma virus titers in Ep-stimulated and non-Ep-stimulated mice revealed that the Ep-responsive cell is predominantly responsible for the production of FV, as measured by focus-forming activity. Mice given saline and FV had very low levels of virus detectable in plasma ($p < 0.01$). Thus, the predominant in vivo target cells for FV replication and tumor colony formation reside within the ERC population. (29 refs)

78-6425 Genetic Resistance in Japanese Wild Mice (*Mus musculus molossinus*) to an NB-tropic Friend Murine Leukemia Virus. (Eng) Odaka, T. (Inst. Medical Science, Univ. Tokyo, P.O. Takanawa, Tokyo 108, Japan); Iwata, H.; Moriwaki, K.; Matsuzawa, A.; Mizuno, M.; Konno, K. *J Natl Cancer Inst* 61(5): 1301-1306; 1978.

Wild mice (*Mus musculus molossinus*; Sk, Hz-VI, Hz-IV, Hz-V, Mol.A, Fu, Te, and Sn) trapped in various areas of Japan were crossed with mice of inbred strains (C57BL/6, C57L, BALB/c, and C57BL/10), and their progeny were infected with NB-tropic Friend murine leukemia virus (F-MuLV: 2.5×10^5 plaque-forming units, ip) at 4 wk of age. Ten days after infection, the spleens were weighed, examined for macroscopic

lesions, and assayed for infectious virus by the XC test. The genotypes of the wild mice were deduced from the responses of the infected progeny. Genetic analysis indicated that 4/8 mice tested had a dominant gene that suppressed virus replication; this gene resembled the *Fv-4'* allele. No mice with the *Fv-2'* allele were found. The gene resembling the *Fv-4'* allele must have an important role in the natural occurrence and propagation of MuLV in wild mice, and it may provide an explanation for the low expression of MuLV in some populations of wild mice. (22 refs)

78-6426 Changes in Cyclic Adenosine 3':5'-Monophosphate Levels During Induction of Differentiation in Murine Erythroleukemia Cells. (Eng) Gazitt, Y. (Dept. Human Genetics and Development, Columbia Univ., Coll. Physicians and Surgeons, New York, NY, 10032); Reuben, R. C.; Deitch, A. D.; Marks, P. A.; Rifkind, R. A. *Cancer Res* 38(11, part 1): 3779-3783; 1978.

Evidence is presented, using synchronized (Syn) and nonsynchronized (NSyn) cultures of Friend virus-infected murine erythroleukemia cells (MELC), that several MELC differentiation-inducing agents induce both an increase in cyclic AMP (cAMP) levels and a transient prolongation of G_1 . The sequential high thymidine (2 mM)-hydroxyurea (0.5 mM) method was used to synchronize the cells. Upon release from cell cycle arrest, the cells go through S in 5 hr, G_2 -M in 1 hr, and G_1 in 3 hr. Syn cells exposed to the inducers hexamethylene bisacetamide, dimethyl sulfoxide, and sodium butyrate progressed through the first Syn cell cycle with the same kinetics as uninduced Syn cells, but remained in G_1 for 6-8 hr. In control cells, the cAMP level was high when cells were at the G_1 -S boundary, fell during S and G_2 -M, and then rose during G_1 . In inducer-treated cells, there was a transient five- to sixfold increase in cAMP during mid-S. In NSyn cultures, an increase in the proportion of cells in G_1 was first noted 6-8 hr after addition of inducer and, depending on the inducer used, could still be detected 10-20 hr later. A transient two- to threefold increase in cAMP occurred 3-6 hr after addition of inducer. Two potent cyclic nucleotide phosphodiesterase inhibitors, 1-methyl-3-isobutylxanthine and RO-20-1724, also increased cAMP, prolonged G_1 , and induced MELC differentiation, but two weaker inhibitors, caffeine (CF) and theophylline (TP), failed to induce differentiation. The fact that TP and CF elevated cAMP and transiently prolonged G_1 but failed to stimulate Hb synthesis suggests that these two effects may not be sufficient to trigger erythroid differentiation or that other effects of these agents prevent differentiation. The magnitude and duration of inducer-mediated prolongation of G_1 did not predict or correlate with the degree of induced differentiation. (35 refs)

78-6427 Mechanism of Genetic Resistance to Friend Virus Leukemia in Mice. V. Relevance of *Fv-3* Gene in the Regulation of In Vivo Immunosuppression. (Eng)

Kumar, V. (Dept. Pathology, Boston Univ. Sch. Medicine, 80 E. Concord St., Boston, MA, 02118); Resnick, P.; Eastcott, J. W.; Bennett, M. *J Natl Cancer Inst* 61(4): 1117-1123; 1978.

An attempt was made to determine if the single autosomal gene *Fv-3*, which regulates susceptibility to immunosuppression by Friend murine leukemia virus (F-MuLV) in vitro, has a similar effect in vivo. Infection of DBA/2, B10.DZ/n, and their progeny F₁ mice with F-MuLV suppressed humoral antibody synthesis in vivo and lymphocyte mitogenesis in vitro. Both these effects of F-MuLV were under host genetic control. In vitro suppression of lymphocyte mitogenesis was regulated by *Fv-3*. Genetic analyses, with the use of the susceptible DBA/2 and resistant B10.D2/n parents and their F₁, intercross, and backcross progeny, indicated that a single autosomal gene dominant for susceptibility regulated the in vivo susceptibility to immunosuppression by F-MuLV. Individual [(DBA/2 x B10.D2)F₁ x B10.D2] mice were typed for susceptibility to F-MuLV-induced suppression of lymphocyte mitogenesis in vitro (an *Fv-3* function) and for susceptibility to immunosuppression by F-MuLV in vivo. This analysis indicated that the same mice that were susceptible or resistant to immunosuppression in vivo were susceptible or resistant to suppression of lymphocyte mitogenesis in vitro. Spearman's rank analysis of the data also indicated that the in vivo and in vitro immunosuppressive effects of F-MuLV were correlated with and not independent of each other. Thus *Fv-3*, which regulates the effect of F-MuLV on lymphocytes in vitro, also appears to regulate the effect of F-MuLV on antibody-forming cells in vivo. (17 refs)

78-6428 Induction of Intracisternal A-Type Particles During Friend Cell Differentiation. (Eng) Ostertag, W. (Max-Planck-Institut für Experimentelle Medizin, 3400 Gottingen, W. Germany); Krieg, C. J.; Cole, T.; Pragnell, I. B.; Swetly, P.; Weimann, B. J.; Dube, S. K. *Exp Cell Res* 116(1): 31-37; 1978.

The regulation of intracisternal A-type particle synthesis during dimethylsulfoxide (Me₂SO)-induced erythroid differentiation was studied in a number of Friend cell lines under conditions which favored or restricted Friend spleen focus-forming virus (FV) production. At least three cell types were derived from the virus positive Friend cell clone F4: Class I lines (V+, A+) produced high levels of infectious FV and low numbers of A-type particles; Class II lines (V+, A++), to which Class I lines could spontaneously and reversibly shift, produced marginal FV and high numbers of A-type particles; and Class III lines (V-, A-) produced neither FV nor A-type particles. Induction of erythroid differentiation in (V+, A++) cells by Me₂SO was paralleled by an increase in C-type, but not A-type, particles. Inhibition of C-type virus production by interferon or azidothymidine was accompanied by a great increase in A-type particle production. A-type particle number was also increased during induction of differentiation in B8 cells, which show an endogenous

defect in Friend virus synthesis (V+, A++). Neither C- nor A-type particles were induced during induction of differentiation in Class III cells, although these cells show a marginal increase in Friend C-type viral RNA levels in the cytoplasm. (V+, A++) cells also showed an increase in Friend virus-related RNA during differentiation. The results suggest that intracisternal A-type particles in Friend cells are related to Friend virus and that the increase in expression of this component is linked to, but is not conditional for, thymopoietic differentiation in Friend cells. (16 refs)

78-6429 Increase in Intracisternal A-Type Particles in Friend Cells During Inhibition of Friend Virus (SFFV) Release by Interferon or Azidothymidine. (Eng) Krieg, C. J. (Max-Planck-Institut für Experimentelle Medizin, D-3400 Gottingen, W. Germany); Ostertag, W.; Clauss, U.; Pragnell, I. B.; Swetly, P.; Roesler, W.; Weimann, B. J. *Exp Cell Res* 116(1): 21-29; 1978.

The effects of interferon and azidothymidine (N₃T), which are known inhibitors of Friend virus synthesis, on the level of intracisternal A-type particles in Friend erythroleukemic (F4-6) cells were studied. Treatment with interferon (10⁶ IU/ml for 4 days) or N₃T (0.25 mM for 35 hr) inhibited spleen focus-forming virus (SFFV) production, reverse transcriptase activity, and the number of C-type particles while increasing the number of intracisternal A-type particles 1- to 20-fold. The A-type particles released from treated cells did not differ in ultrastructure from those released from untreated cells. Interferon treatment of cells, which show an a priori low level of C-type virus synthesis, did not result in an increase in A-type particles. Similarly, the number of A-type particles was not increased by interferon treatment of three C-type virus-releasing cell lines, although C-type virus release was inhibited. Viral RNA levels were not markedly increased during inhibition of C-type virus release in F4-6 cells. The presence and accumulation of A-type particles seems to be related to the production of Friend virus complex (SFFV). (41 refs)

78-6430 Effect of Infection of Mice with Friend Leukemia Complex Viruses on Background Plaque-Forming Cell Production In Vitro. (Eng) Bendinelli, L. (Inst. Microbiology, Univ. Pisa, I-56100 Pisa, Italy); Matteucci, D.; Toniolo, A.; Friedman, H. *Experientia* 34(11): 1161-1162; 1978.

After infection of BALB/c donor mice with Friend leukemia complex viruses and Rowson-Parr virus, the production in vitro of background plaque-forming cells against sheep red blood cells by spleen cells was depressed. (14 refs)

78-6431 Specific Inhibition of XC Syncytia Formation by Anti-MuLV gp69/71 Antibodies. (Eng) G...

brecht, S. (Laboratoire d'Immunologie et de Virologie des
meurs, Inserm U-152, Hopital Cochin, 27 rue du Fau-
arg Saint Jacques, F-75674 Paris Cedex 14, France); Blai-
u, C.; Levy, J. P. *Intervirology* 10(6): 338-344; 1978.

Results of XC syncytia-inhibition tests using ecotropic
murine leukemia virus (MuLV) producer cells and various
viral antisera are presented. The producer cells were:
3-FL cells infected by Gross leukemia virus (GLV);
7BL/6 mouse embryonic fibroblasts infected by Rauscher
leukemia virus (RLV); and (BALB/c x NIH)F₁ hybrid emb-
ryonic fibroblasts infected by Moloney leukemia virus
(MLV). Antisera used included rat antisera against
RLV-induced rat tumors; (W/Fu x B/N)F₁ rat anti-AT124
ecotropic C-type virus; mouse anti-FMR (Friend, Moloney,
Rauscher) sera; monospecific goat antisera against RLV
polypeptides p10, p12, p15, p30, p60, and gp69/71; and nor-
mal sera from uninoculated mice of various strains. When
MLV producer cells were treated with RLV polypeptide an-
tisera prior to coculture with XC cells, antisera against p10,
p12, p15, p30, and p60 were devoid of inhibitory activity,
whereas the two anti-gp69/71 sera were strongly inhibitory,
suggesting that gp69/71 mediates the fusion of mouse pro-
ducer and XC cells. Since this inhibitor is specific for an-
ti-gp69/71, it provides a simple, sensitive, and convenient
assay to type ecotropic C-type viruses and to detect an-
ti-gp69/71 antibodies. The results with normal mouse sera
suggest that older B6 mice and their F₁ hybrids with DBA/2
C3H/He spontaneously produce anti-GLV-gp69/71 an-
tibodies with type-specific activity. (17 refs)

78-6432 **Sequential Changes of T- and B-Cells, Virus An-
tigen Expression and Primary Histologic Tu-
mor Diagnosis in Virus-induced Lymphomagenesis of Mice.**
(Eng) Krueger, G. R. (Immunopathologisches Laboratori-
um, Pathologisches Institut der Universitat Koln, Jo-
seph-Stelzmann-Strasse 9, D-5000 Cologne, W. Germany);
Rauscher, K. M.; Flesch, H. G. *Z Krebsforsch* 92(1): 41-54;
1978.

and B-cell counts, estimation of immunoglobulin (Ig)
receptor fluidity, and expression of virus-coded antigens were
correlated with histological findings during the development
of a virus-induced mouse lymphoma. BALBc mice were in-
oculated at 4 days of age with the strongly oncogenic Moloney
leukemia virus (MLV), the moderately oncogenic (in
BALB/c mice) Gross passage A virus (GLV-A), or the
nononcogenic Gross 3T3 tissue culture virus (GLV-T). Rou-
tine autopsies were done at 2- to 4-wk intervals postinjection
(PI). Several changes were observed in mice with MLV and
GLV-A infection but not in those with GLV-T infection. Ig
receptor fluidity and expression of virus antigen were de-
creased at the initial examination (2 wk PI). There were sig-
nificant decreases in the percentage of T cells 9-12 wk PI,
accompanied by histologic atrophy of the thymus and of
thymus-dependent regions of lymphatic tissues. The first le-
sions histologically suspicious for lymphoma occurred in the

thymus 11-12 wk PI with MLV and 12-20 wk PI with
GLV-A. At 15-20 wk PI with MLV and 20-30 wk PI with
GLV-A, generalized lymphoma was seen. In ultrastructural
studies, there was no specific cell change indicating malignant
transformation of individual cells. There was some indication
of derangement and of a reduction of cytoplasmic microfila-
ments in proliferating lymphoblasts and lymphoma cells. (45
refs)

78-6433 **Replication-defective Ecotropic Murine Leu-
kemia Viruses: Detection and Quantitation of
Infectivity Using Helper-dependent XC Plaque Formation.**
(Eng) Rein, A. (Lab. Tumor Virus Genetics, NCI, Bethesda,
MD, 20014); Bassin, R. H. *J Virol* 28(2): 656-660; 1978.

An infectivity assay for a defective variant of Moloney mu-
rine leukemia virus M (MuLV-M) and for other replica-
tion-defective (R-D) ecotropic MuLV's is reported. The vari-
ant was present in a clone of mouse cells (clone 8A) isolated
from a murine sarcoma virus (MSV)-transformed derivative
of mouse 3T3FL cells superinfected with MuLV-M. The as-
say is based on the fact that some isolates of MuLV replicate
well in mouse cells but, unlike nondefective ecotropic
MuLV's, do not induce fusion in XC cells. Particles of R-D
ecotropic MuLV, which do not themselves initiate XC plaque
formation, give rise to XC plaques in cells that are coinfect-
ed with XC-negative, nondefective MuLV. Thus, fusion of XC
cells is a marker by which the defective virus can be specifi-
cally detected in the presence of a nondefective virus. In two
trials, clones 8A and NP-N (a non-plaque-forming variant
of N-tropic MuLV) produced particles that did not form
plaques in the XC test. However, these particles formed XC
plaques when amphotropic MuLV, which is XC-negative,
was added to the assay plates. This phenomenon can be used
as a quantitative infectivity assay for R-D MuLV's. (17 refs)

78-6434 **Ornithine Decarboxylase Activity in Cells
Acutely and Chronically Transformed by Mu-
rine Sarcoma Virus.** (Eng) Kilton, L. J. (NCI-VA Medical
Oncology Unit, NCI, Bethesda, MD, 20014); Gazdar, A. F.
Proc Soc Exp Biol Med 159(1): 142-147; 1978.

Three contact-inhibited or murine sarcoma virus
(MSV)-transformed clones of BALB/3T3 mouse cells were
transformed or supertransformed by MSV or its accompan-
ying nontransforming helper virus (murine leukemia virus,
MuLV), and the relationships between ornithine decarboxy-
lase ODC activity, morphology, virus production, and
growth rates were examined. Clones isolated from these lines
were also studied. All of the virus-infected lines released both
MSV and MuLV. ODC activities could not be correlated with
difference in growth rates or with release of transforming or
nontransforming viruses. The only consistent relationship
was between elevated ODC activity and acute morphological
transformation. With cell passage, ODC levels of acutely

transformed cells returned to baseline levels, perhaps because most acutely transformed cells fail to divide. These findings plus those of a previous study, which indicated that human cytomegalovirus infection of fibroblasts overrides end product repression of ODC by putrescine, suggest that the oncogenic potential of a virus may be related to its ability to interfere with normal regulatory functions of key cellular metabolic enzymes. (16 refs)

78-6435 Enhanced Susceptibility of Virus-Infected Cells to Starch-induced Peritoneal Exudate Cells.

(Eng) Goldman, R. (Dept. Zoology, Univ. Coll. London, Gower St., London, WC1E 6BT, England); Hogg, N. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 97-107; 1977.

Starch-induced peritoneal exudate cells from BALB/c mice were tested for their ability to inhibit 5-¹²⁵I-iodo-2'-deoxyuridine (IUDR) incorporation in normal, virus-infected, or virus-transformed mouse embryo fibroblasts (MEF). Preliminary experiments showed that the PEC abrogated IUDR incorporation in K-BALB cells, and that this cytostatic activity resided in the adherent cell fraction of the PEC, which included macrophages but not T or B lymphocytes. Recently explanted MEF were relatively resistant to PEC-mediated cytostasis. However, secondary MEF transformed by murine sarcoma virus or infected with murine leukemia virus, Sendai virus, or influenza virus were sensitive to PEC cytostasis. Virus-infected or -transformed MEF from four mouse strains (BALB/c, CBA, AKR, C57BL) were affected to a similar degree, indicating that there is no requirement of H-2 compatibility between effector and target cells. These results show that membrane alterations induced by a nontransforming virus infection are sufficient to render cells susceptible to activated, adherent PEC. (26 refs)

78-6436 Structural Markers on Core Protein p30 of Murine Leukemia Virus: Functional Correlation with Fv-1 Tropism. (Eng) Gautsch, J. W. (Res. Inst. Scripps Clinic, 10666 N. Torrey Pines Road, La Jolla, CA, 92037); Elder, J. H.; Schindler, J.; Jensen, F. C.; Lerner, R. A. *Proc Natl Acad Sci USA* 75(9): 4170-4174; 1978.

The primary structure of the major core protein, p30, of murine leukemia virus (MuLV) was studied in over 50 virus isolates by a simplified method of in situ radioiodination and two-dimensional peptide mapping. Although all MuLV p30's displayed a close structural similarity, minor differences in the peptide profiles did occur, and they were reproducible and served as excellent markers for certain MuLV strains. A loss of peptide spots 1 and/or 2 in the peptide map was noted only in Friend, Moloney, and Rauscher leukemia viruses, and not endogenous MuLV. Peptide 12, which varied

in its relative position more than any other peptide in the maps, exhibited one of four primary positions, and was to be functionally associated with Fv-1 tropism. On this basis a virus could be classified as N-tropic, B-tropic, xenotropic and/or from a wild mouse. Although phenotypically identical, two BALB/c B-tropic isolates, WN1802B and B could be genotypically distinguished, as could two different BALB/c xenotropic viruses, B-228 and Mk-137MC. The clone of Moloney murine sarcoma virus was seen to possess extra marker peptides derived from the COOH end of that could be sarcoma-associated. (38 refs)

78-6437 Further Evidence for Deletion of Envelope Glycoprotein (gp69/71) Sequences in Formation of Moloney-Murine Sarcoma Virus. (Eng) Peebles, P. (Lab. Tumor Virus Genetics, NCI, Bethesda, MD); Straube, M.; August, J. T. *J Gen Virol* 40(3): 503-510; 1978.

The S+L-strain of Moloney murine sarcoma virus (M-MuSV) was nonproductively cloned in S+L-mouse, dog, mink, and human cells, and expression of the Moloney murine leukemia virus (M-MuLV) 30,000-mol wt core protein (p30) and envelope glycoprotein (gp69/71) was studied by radioimmunoassay. In all cases, the p30 determinant, which has been demonstrated to be part of a larger p60 protein precursor, remained associated with the MSV genome. However, heterologous host S+L- cell line showed evidence for M-MuLV gp69/71. These data support previous biological data and suggest that the gp69/71 gene was deleted in the formation of this MSV genome. The S+L- dog cells released large amounts of a soluble p30 antigenic determinant in their culture supernatant fluids in the absence of detectable virus particle production and under optimal conditions for cell viability. This is the first example of the in vitro release of a major structural core virus protein in soluble form in the absence of virus particle production. (31 refs)

78-6438 Biochemical Studies on RNA Tumor Virus Replication and its Transmission in B16 Murine Melanoma. (Eng) Hehlmann, R. (Med. Poliklinik, Universität München, Munich, W. Germany); Erfle, V. *Int J Cancer* 22(5): 630-638; 1978.

The RNA tumor virus information of the B16 mouse melanoma system and its transmission were studied biochemically. Density-fractionated B16 melanomas contained particles possessing a 70S RNA associated with a reverse transcriptase. The particles had a density of 1.14-1.18 g/cm³ along 25%-55% sucrose gradient. The ³H-complementary DNA synthesized by these particles hybridized back to its template RNA. The B16 particle RNA showed considerable homology with the RNA's of the murine leukemia-sarcoma virus AKR-MuLV, Kirsten-MuSV, and Rauscher-MuLV and some degree of homology to the RNA's from simian sarcoma virus and RD-114 viruses. In addition, the B16 DNA con-

med particle-related sequences not present in the DNA of normal C57BL mice. (30 refs)

78-6439 Chemical Determination of the m1 Moloney Sarcoma Virus pP60gag Gene Order: Evidence for Unique Peptides in the Carboxy Terminus of the Polyprotein. (Eng) Oskarsson, M. K. (Lab. DNA Tumor Viruses, NCI, Bethesda, MD, 20014); Elder, J. H.; Gautsch, J. W.; Turner, R. A.; Vande Woude, G. F. *Proc Natl Acad Sci USA* 10(10): 4694-4698; 1978.

The gene order of the m1 isolate of Moloney sarcoma virus (MSV) specific polyprotein pP60gag (P60) was determined by direct chemical analysis. P60 was cleaved with cyanogen bromide (CNBr) into eight partial and complete fragments ranging from 10,000 daltons to 58,000 daltons. Peptide maps of these fragments were compared with maps of p15, p12, and three CNBr fragments of p30. The polarity of p15 and p12 in a CNBr fragment of P60 was determined by carboxypeptidase A digestion; the CNBr fragments of p30 were ordered by aminopeptidase digestion. The linear arrangement of P60 CNBr fragments gave the gene order of H_2 -p15-p12-p30-COOH. This linear arrangement of CNBr fragments should facilitate primary amino acid sequence analysis of the polyprotein. In addition, as demonstrated with MSV P70, this map should serve as a useful basis for comparison to other MSV or Moloney murine leukemia virus (MLV) gag polyproteins when the latter are available in sufficient quantities. The m3 isolate of MSV expressed a P70 gag protein. Peptide maps of 48,000-dalton CNBr fragments of m3 P70 and m1 P60 were similar, which suggested that these polyproteins were similar through the NH_2 -terminal two-thirds of p30. However, the presence of peptides unique to the 10,500-dalton COOH-terminal fragment of m1MSV P60, ie, peptides absent in the p30 of either m3MSV or MLV, suggested that the gag gene deletion in the m1 isolate begins at the p30 reading frame. (21 refs)

78-6440 Feline Sarcoma Virus In Vitro Infection of Human Cells. Influence of Chemical Carcinogens on Focus Formation. (Eng) Blakeslee, J. R. (Dept. Veterinary Pathobiology, Coll. Veterinary Medicine, Ohio State Univ., 25 Coffey Road, Columbus, OH, 43210); Milo, G. E. *Chem Biol Interact* 23(1): 1-11; 1978.

The effects of benzo(a)pyrene (BP), aflatoxin B₁ (AFB₁), and 2-acetoxy-2-fluorenylacemide (A-AAF) on Synder-Theil feline sarcoma virus-induced focus formation was studied using normal human foreskin fibroblasts (D550). Focus formation was inhibited by 27% to 60% in cells treated with 1.0-15.0 μ g/ml 24 hr before virus infection, by 65% to 87% in cells treated 6 hr before infection, by 59% to 87% in cells treated 2 hr before infection, and by 31% to 42% in cells treated after infection. A 43% inhibition occurred in cells treated with A-AAF (1.0 μ g/ml) 6 hr before infection,

70% in cells treated 2 hr after infection, and 55% in cells treated 24 hr after virus adsorption. AFB₁ inhibited focus formation by 39% in cells treated with 1.0 μ g/ml 6 hr before virus infection and by 46% to 59% in cells treated with 0.1 μ g/ml after infection. Focus formation was not significantly altered by treatment before or after infection with pyrene (10 μ g/ml) or 0.1 μ g/ml A-AAF or by treatment before infection with 0.1 μ g/ml AFB₁. Focus formation was not inhibited in cells treated with BP (15 μ g/ml) or AFB₁ (0.1 μ g/ml) 48 to 96 hr postinfection. Virus-directed RNA-dependent DNA polymerase activity was not reduced in cells showing significant inhibition of focus formation, and no infectious virus was detected in either treated or untreated virus-infected cells. The data indicate that the inhibitory effect on focus formation is chemically mediated, whereas the inhibition of virus synthesis is not. (14 refs)

78-6441 Endogenous Feline Oncoviruses: Genome Expression of RD-114 and Feline Leukemia Virus in Normal and Abnormal Differentiation of the Domestic Cat (*Felis catus*). (Eng) Niman, H. L. (Univ. Southern California, Los Angeles, CA, 90007). *Diss Abstr Int B* 39(5): 2131; 1978. (no refs)

78-6442 Feline Leukemia Virus Envelope Glycoprotein Vaccine: Preparation and Evaluation of Immunizing Potency in Guinea Pig and Cat. (Eng) Salerno, R. A. (Div. Virus and Cell Biology Res., Merck Inst. Therapeutic Res., West Point, PA, 19486); Lehman, E. D.; Larson, V. M.; Hilleman, M. R. *J Natl Cancer Inst* 61(6): 1487-1493; 1978.

Envelope glycoprotein (gp) of feline leukemia virus (FeLV), strain FeLV-1, was extracted with 2% Triton X-100 and 0.43 M sodium chloride in 10 mM Tris (pH 7.5) and purified by ion-exchange chromatography on phosphocellulose. Purified FeLV envelope gp75 (mol wt 75,000 daltons) induced antibodies of high titer in goats. These antibodies specifically neutralized FeLV, precipitated radiolabeled FeLV envelope gp75, and lysed FL74 cells. Administration of 20 μ g gp75 to guinea pigs in alum or adjuvant 65 induced neutralizing antibody [geometric mean titer (GMT), 8-25], radioimmune precipitating antibody (GMT, 1,382-6,450), and cytotoxic antibody (titer, ≥ 2) in all animals. Aqueous gp75 vaccine induced neutralizing antibody (GMT, < 4) in only 50% of the guinea pigs. Specific-pathogen-free cats immunized with 40 μ g gp75 in alum or Freund's adjuvant developed radioimmune antibody with GMT's of 127-207 (100% seroconversion, SC), but little or no neutralizing antibody (22%-38% SC; GMT, < 2) or cytotoxic antibody (0%-44% SC) developed. These findings indicate that oncovirus subunit vaccines will require highly effective immunologic adjuvants to enhance the immunogenicity of the antigen in the homologous host. (36 refs)

78-6443 Feline Oncornavirus-associated Cell Membrane Antigen. (Eng) Essex, M. (Dept. Microbiology, Harvard Univ. Sch. Public Health, Boston, MA, 02115); de Noronha, F.; Oroszlan, S.; Hardy, W. D. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1401-1414; 1978.

The object of this study was twofold: (1) to determine if feline leukemia virus (FeLV) proteins gp70, p27, and p15 were detectable on the surface of feline leukemia or lymphoma cells (including cell culture lines and freshly biopsied tumor cells) and (2) to determine if feline oncornavirus-associated membrane antigen (FOCMA) activity (found naturally in healthy virus-exposed cats) could be adsorbed or removed with either pure intact or fractionated FeLV. Monospecific goat or rabbit antisera to gp70, p27, and p15 indicated that these antigens are present on FOCMA-positive cells that produce FeLV, including two cultured lymphoblastoid lines and biopsied cells from cats with naturally occurring leukemia and lymphoma. However, the antibody activity from these monospecific antisera was readily removed by adsorption with either pure gp70, p27, or p15 or whole disrupted FeLV. In contrast, similar treatment of feline sera containing high titers of FOCMA antibody did not remove antibody activity directed to the surface of virus-producer tumor cells. It is concluded that FOCMA contain antigenic determinants that are not shared with the major virus structural proteins. (43 refs)

78-6444 Characterization of a Feline Sarcoma Virus-coded Antigen (FOCMA-S) by Radioimmunoassay. (Eng) Sherr, C. J. (Lab. Viral Carcinogenesis, NCI, NIH, Bethesda, MD, 20014); Todaro, G. J.; Sliski, A.; Essex, M. *Proc Natl Acad Sci USA* 75(9): 4489-4493; 1978.

The previously detected expression of a feline oncornavirus-associated cell membrane antigen (FOCMA) on feline sarcoma virus (FeSV)-transformed rat and mink nonproducer cells (FOCMA-S) provided evidence that this antigen is encoded by a feline C-type viral gene. The presence of FOCMA-S in FeSV pseudotype particles facilitated the development of a radioimmunoassay (RIA) for an FeSV gene product that uses rabbit antiserum and protein derived from viral particles produced by nonfeline (mink, rat) cell. Thus, it was possible to assay for FOCMA-S with reagents generated completely outside of the feline system. Although the polypeptide (p85) detected in pseudotype virions contained p15 and p12 antigenic determinants, antibodies to these could be adsorbed with purified feline leukemia virus (FeLV) proteins. Hence, the RIA for FOCMA-S did not detect the major viral structural protein antigens found in p85. The adsorbed rabbit antiserum bound to antigenic determinants also present in

p85 and reacted specifically in immunofluorescence tests of rat cells transformed by FeSV and with FOCMA-positive lymphoid tumor cells. Competition assays detected FOCMA-S in pseudotype C-type viruses rescued from FeSV-transformed mink and rat cells, but not in heterologous C-type helper viruses or in FeLV. A cross-reactive antigen was also detected in pseudotype of Kirsten sarcoma virus. This quantitative assay could also be used to purify viral antigens from transformed cells and to characterize these potential 'transforming proteins.' (26 refs)

78-6445 Demonstration of Two Distinct Classes of Bovine Papilloma Virus. (Eng) Lancaster, W. (Div. Otolaryngology, Case Western Reserve Univ., Cleveland, OH, 44106); Olson, C. *Virology* 89(2): 372-379; 1978.

The purpose of this study was to determine if different classes of bovine papilloma virus (BPV) exist. Virus DNA was isolated and purified from 10 naturally occurring cases of bovine fibropapillomatosis and hybridized to labeled BPV DNA. Results indicate that two distinct classes of BPV can be distinguished based on the extent of hybridization. Saturating hybridization showed that the two virus classes shared 52%-58% of their polynucleotide sequences. DNA-DNA reassociating kinetics revealed that the two virus classes shared a common base pairs along 45% of the genome. The viruses differed in their ability to hemagglutinate mouse RBCs, they were antigenically cross-reactive but not identical in complement fixation and hemagglutination inhibition. Although many of the structural proteins of the two virus classes seemed to have similar electrophoretic mobilities on sodium dodecyl sulfate-polyacrylamide gels, a more detailed analysis will be required to detect minor differences in molecular weight. There were no differences between the viruses with respect to location or morphology of papillomas or to a preference for a given bovine breed. (23 refs)

78-6446 Common Membrane Neoantigens on Bovine Papilloma Virus-induced Fibroma Cells from Cattle and Horses. (Eng) Barthold, S. W. (Section Comparative Medicine, Yale Univ. Sch. Medicine, New Haven, CT 06510); Olson, C. *Am J Vet Res* 39(10): 1643-1645; 1978.

Bovine papilloma virus (BPV)-induced fibroblastic tumors in cattle, horses, and hamsters were studied by indirect immunofluorescence in order to identify common antigens. Sixteen calves <2 wk old and six adult ponies were inoculated with 0.1-0.2 ml of 20%-30% BPV homogenate in multiple sites. Serum was collected at inoculation and at various specified times afterward. Nine outbred Syrian hamsters were inoculated sc with 0.2 ml of homogenate, and a tumor from one of these hamsters was transplanted sc into 13 other hamsters. Four calves developed fibromas, three fibropapillomas. Two ponies developed fibromas. Seven of nine hamsters developed fibrosarcomas, and all those with transplanted tumor de-

ed large sarcomas. Specific membrane IF occurred on boe and equine tumor explant cells, but not on normal derl cells, if reacted against homologous or heterologous sera. cept for two bovine sera, the reactions of bovine sera to ine tumor cells and equine tumor cells were similar. The ction pattern of equine sera was similar, but seemed to be dified by tumor size of the serum source. The reaction ild be blocked in both species by homologous, but not erologous, sera. Sera from tumor-bearing hamsters had no ectable reactivity to cell membranes of hamster, bovine, equine cells. Internal neoantigens were not found in any the acetone-fixed tumor cells, using sera from the three cies. Both tumor and normal cells were found free of BPV igen by direct IF. (30 refs)

6447 **Replication of Nondefective Parvoviruses: Lack of a Virion-associated DNA Polymerase.** (Eng) tchard, C. (Dept. Biology, Virginia Polytechnic Inst., icksburg, VA, 24061); Patton, J. T.; Bates, R. C.; Stout, R. *J Virol* 28(1): 20-27; 1978.

ur nondefective parvoviruses, bovine parvovirus (BPV), l, LuIII, and Kilham rat virus (KRV), were examined for ion-associated DNA polymerase activity. Virions were rified from neuraminidase-treated infected cell lysates by pycnic centrifugation in CsCl or from infected cell materi- by CaCl₂ precipitation and centrifugation through sucrose o CsCl. BPV and KRV preparations obtained by the for- procedure contained DNA polymerase activity but were e free of contaminating cellular proteins. In both cases, the latest amount of residual enzyme activity consisted of NA polymerase β . The latter method produced viral prepa- tions free of contaminating cellular proteins, and no DNA ymerase activity was detected in any of the four viruses. rruption of infectious virions obtained by the second meth- with ionic and nonionic detergents and sonic treatment d not result in the detection of a DNA polymerase. The ological activity and purity of each of the viruses were de- mined by hemagglutination and infectivity assays, poly- /amide gel electrophoresis, and electron microscopy. In ch case, the virions banding at a density of 1.39-1.41 g/cm³ CsCl were infectious and contained only the virion struc- al proteins. DNA polymerase activity was not detected in y of the purified virus preparations. It is concluded that RV, BPV, H-1, and LuIII do not contain an essential vin- ion-associated DNA polymerase required for productive in- tion. (25 refs)

-6448 **Viral DNA Synthesis in Cells Infected with Temperature-sensitive Mutants of Herpes Sim- plex Virus Type 1.** (Eng) Bone, D. R. (Tumor Virology Lab., lk Inst. Biological Studies, San Diego, CA, 92112); Brown, ; Crombie, I.; Francke, B. *J Virol* 28(1): 14-19; 1978.

temperature-sensitive (*ts*) mutants of herpes simplex virus

type 1 (HSV-1) representing eight DNA-negative (DNA-) complementation groups were characterized in tempera- ture-shift experiments. Viral DNA synthesis was measured in mutant-infected baby hamster kidney (BHK-21) cells after (a) shift-up to the nonpermissive temperature (39 C) to deter- mine whether synthesis continued or was inhibited and (b) shift-down to the permissive temperature (32 C) in the pres- ence and absence of inhibitors of protein and RNA synthesis. The mutants could be grouped into three categories based on the results of these experiments. (1) After shift-up, viral DNA synthesis was inhibited in cells infected with *tsB*, *tsH*, and *tsJ*, indicating that the B, H, and J proteins are continuously required for this process. Cells infected with *tsB* and *tsH* syn- thesized viral DNA after shift-down in the absence of de novo RNA and protein synthesis; *tsJ*-infected cells did not synthe- size viral DNA in the absence of protein synthesis. (2) Viral DNA synthesis continued after shift-up in cells infected with *tsD* and *tsK* but not after shift-down in the absence of RNA and protein synthesis, suggesting that these mutants are de- fective in early regulatory functions. (3) Cells infected with *tsL*, *tsS*, and *tsU* synthesized viral DNA after shift-up and after shift-down in the absence of RNA and protein synthesis. The functions of the L, S, and U proteins were not clarified. (11 refs)

78-6449 **Frequency of Anti-AG-4 Antibody in Patients with Uterine Cervical Cancer and Controls.** (Eng) Kawana, T. (Dept. Obstetrics and Gynecology, Facul- ty Medicine, Univ. Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113, Japan); Sakamoto, S.; Kasamatsu, T.; Aurelian, L. *Gann* 69(4): 589-591; 1978.

The association of herpesvirus type 2 (HSV-2) with cervical cancer was investigated by determining the prevalence of an- tibody to an early antigen (Ag-4, an antigen that is induced in Hep-2 cells by HSV-2 4 hr after infection) in 33 Japanese cancer patients and 30 age-matched controls. A microquan- titative complement fixation assay detected antibody Ag-4 in 17/33 patients and 5/30 controls. There was a tendency for antibody prevalence to increase with clinical stage: 43% at Stage I, 63% at Stage II, and 75% at Stage III. A neutraliza- tion kinetics assay showed that 11/15 anti-Ag-4-positive sera contained neutralizing antibody to HSV-2, as did 11/16 anti-Ag-4-negative sera. Eleven of 22 HSV-2-positive sera were also Ag-4-positive, which indicates that there is no correla- tion between the prevalence of Ag-4 antibody and that of HSV-2 neutralizing antibody. Thus, Ag-4 differs from viral antigens involved in neutralization. These results, similar to those of several US studies, suggest that HSV-2 is associated with cervical cancer and appears to reflect tumor progression. There are two possible explanations for the observation that only 17/34 patients were Ag-4-positive: antibody levels in the negative patients were too low for detection or the negative patients were not previously exposed to HSV-2. The latter explanation suggests that HSV-1 may be associated with cer- vical cancer in Ag-4-negative patients. (10 refs)

- 78-6450 Selective Stimulation and Differentiation of Early Antigens in Lymphoblastoid Cell Lines Producing Epstein-Barr-Like Viruses.** (Eng) Neubauer, R. H. (Viral Oncology Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Rabin, H.; Brown, A. M. *J Gen Virol* 41(2): 295-301; 1978.

The early antigens (EA) of Epstein-Barr virus (EBV), *herpesvirus papio*, *herpesvirus pongu*, and *herpesvirus pan* were examined using a procedure that combines antigen stimulation with inhibition of late virus functions and results in the selective stimulation of EA. Treatments combining 5-iodo-2'-deoxyuridine and phosphonoacetic acid (PAA) resulted in the selective stimulation of EA in lymphoblastoid cell lines that produce EBV-related viruses. The marked increase in the frequency of EA-positive cells in the relative absence of virus capsid antigens permitted the comparison of EA in related viruses. The EA's of the four viruses were distinguishable, with the greatest differences being between EBV and *h. papio*. The differences were indicated by the ability to select sera that had positive reactivity only with the homologous antigen and by differences in homologous and heterologous titers in certain cross-reacting sera. These differences were also demonstrable for both the restricted and diffuse components of EA, and, they suggest that EBV-related viruses can be distinguished by their early antigens as well as by comparisons of virus-specific nuclear antigens and DNA hybridization techniques. (11 refs)

- 78-6451 Chloroquine Enhances Epstein-Barr Virus Expression.** (Eng) Karmali, R. A. (Dept. Pathology, Sch. Medicine, East Carolina Univ., Greenville, NC, 27884); Horrobin, D. F.; Menezes, J.; Patel, P.; Musto, J. *Nature* 275(5679): 444-445; 1978.

The effect of chloroquine (CQ), an antimalarial agent, on the expression of Epstein-Barr virus (EBV) antigens was studied in the Raji cell line, which is EBV-induced nuclear antigen (EBNA)-positive, early antigen (EA)-negative, and virus capsid antigen (VCA)-negative. The cells were superinfected with the P3HR-1 strain of EBV for the induction of EA and VCA. CQ was added to the infected cells, and the cultures were incubated for 48 hr. Indirect immunofluorescence showed that 6.5% of the infected cells were EA-positive. Following treatment with 0.1, 1.0, or 5.0 $\mu\text{g}/\text{ml}$ CQ, however, 19%, 16.8%, and 18.3% of the cells were EA-positive, respectively. (Plasma concentrations of CQ in humans during malaria chemotherapy are 0.16-0.53 $\mu\text{g}/\text{ml}$.) There was no difference in VCA-positive cells in the presence of CQ. These results suggest that CQ may be acting either directly on the exogenous EBV genome or indirectly, by activating cellular mechanisms essential for its expression. Therefore, CQ may play an important part in the development of African Burkitt's lymphoma by enhancing EBV expression. (26 refs)

- 78-6452 Longitudinal Study of Epstein-Barr Virus Antibody Titers and Excretion in Pediatric Patients**

with Hodgkin's Disease. (Eng) Lange, B. (Div. Oncology, Children's Hosp. Philadelphia, Philadelphia, PA, 19104); Beter, A.; Hewetson, J.; Henle, W. *Int J Cancer* 22: 521-527; 1978.

An attempt was made to provide evidence of the latent acter of Epstein-Barr virus (EBV) infection by monitoring children with Hodgkin's disease (HD) and a history of infection for changes in the titers and spectra of antibody to EBV-determined antigens, excretion of the virus in oropharynx, EVB-carrying lymphoid cells in the peripheral blood, and clinical parameters signifying a reactivation of latent viral condition. Although the incidence and geometric mean titers of IgG antibodies to viral capsid antigen (VCA) in the HD patients were similar at diagnosis to those of controls, the anti-VCA titers rose significantly above controls during and after therapy and remained elevated throughout the 7 yr of observation. At no time were heterophil VCA-specific IgM antibodies detected. Antibodies to EBV-induced early antigens were much more common (10% vs 9%) and antibody levels to EBV-associated nuclear antigen were significantly lower in HD patients than in controls. No specific clinical alterations were correlated with the elevated antibody titers or the observed increase in EBV excretion. Although a reactivation of EBV infection by the immunosuppressive effects of HD or its therapy is presently the best explanation of these results, direct evidence has yet to be provided. (38 refs)

- 78-6453 Isolation of a Herpesvirus-specific DNA Polymerase from Tissues of an American Patient with Burkitt Lymphoma.** (Eng) Allaudeen, H. S. (Dept. Pharmacology, Yale Univ. Sch. Medicine, New Haven, CT 06510); Bertino, J. R. *Proc Natl Acad Sci USA* 75: 4504-4508; 1978.

A DNA polymerase (Pase) isolated from a 27-yr-old white American man with Burkitt's lymphoma (BL) showed specific properties of herpes simplex virus (HSV)-induced DNA polymerase and the Epstein-Barr virus (EBV) DNA Pase of Burkitt's lymphoma in culture. The enzyme was partially purified from a crude extract and separated from the cellular DNA Pases. The DNA Pase, which represented 27%-30% of the total cellular DNA activity, differed from the cellular DNA Pases. Its molecular wt was determined to be 90,000 by velocity sedimentation. The activity of the enzyme increased three to four times in the presence of high salt concentrations (100 mM), similar to HPV DNA Pase. This enzyme further resembled HSV DNA Pase in its primer template preference, high monovalent cation requirement for activity, and sensitivity to phosphonoacetate inhibition. N-Ethylmaleimide had no effect on its activity. HSV DNA Pase antiserum inhibited the activity of BL DNA Pase, whereas antibody to HeLa cell DNA Pase α or to DNA Pase γ from human lymphoblast cells did not. Although the patient's serum contained a high EBV titer, addition of the serum to the assay mixture did not have any effect on the activity of BL DNA Pase. Spleen and liver

did not contain HSV DNA. Detection of the HSV DNA in BL tissue provides additional evidence for the association of EBV with this malignancy. (17 refs)

78-6454 Induction of Nuclear Antigen (EBNA) Synthesis Without Transformation in Human Cord Lymphocytes by P3HR-1 Strain of Epstein-Barr Virus. (Eng) Yamamoto, N. (Dept. Microbiology, Kumamoto Univ. Medical Sch., Kumamoto, 860, Japan); Hinuma, Y. *Immunology* 89(2): 623-626; 1978.

The transforming potential of the P3HR-1 strain of Epstein-Barr virus (EBV) was analyzed by clonal transformation in soft agar, and EBV-determined nuclear antigen (EBNA) induction by this virus was investigated in primary cultures of normal umbilical cord lymphocytes. The B95-8 strain of EBV was used as a positive control. Experiments with six cord cell preparations from different neonates showed that all nine preparations of P3HR-1 virus gave negative transformation results, but all cell preparations infected with B95-8 virus were transformed. In addition, EBNA-positive cells were present in all P3HR-1-infected cultures. EBNA synthesis without transformation was confirmed through colcemid treatment of P3HR-1-infected cord cells. No labeled cells were found upon autoradiographic examination of cells infected with P3HR-1 and exposed to ³H-thymidine at several intervals after infection. It is concluded that EBNA synthesis does not correlate with transformation and that DNA synthesis is not necessary for EBNA induction. (3 refs)

78-6455 Epstein-Barr Virus and Human Chromosomes: Close Association of the Resident Viral Genome and the Expression of the Virus-determined Nuclear Antigen (EBNA) with the Presence of Chromosome 14 in Human-Mouse Hybrid Cells. (Eng) Yamamoto, K. (Dept. Microbiology, Cancer Inst., Hokkaido Univ. Sch. Medicine, 060 Sapporo, Japan); Mizuno, F.; Matsuo, T.; Tanaka, A.; Onoyama, M.; Osato, T. *Proc Natl Acad Sci USA* 75(10): 5551-5559; 1978.

Fourteen hybrid clones derived from fused cultures of human lymphoblastoid FV5 cells and 5-bromodeoxyuridine-resistant mouse fibroblastic MCB2 cells grown in hypoxanthine/aminopterin/thymidine selective medium were examined for the presence of Epstein-Barr virus (EBV) DNA, expression of the virus-determined nuclear antigen (EBNA), and the presence of human chromosomes. When tested 3 months after their isolation, 11 of the hybrid clones were positive for EBNA, but when tested 18 months after their isolation, only 3 clones were positive for EBNA. EBV DNA was detected only in the three clones that were positive for EBNA after 18 months. Chromosome analysis of the hybrid clones showed that human chromosome 14 was common to all three positive clones, whereas this chromosome was not present in any of the nega-

tive clones tested. In 10 subclones isolated from one of the positive clones, a concordant segregation of EBNA, EBV DNA, and chromosome 14 was evident. The data suggest that the resident EBV genome is closely associated with chromosome 14 and that the presence of this chromosome alone is sufficient for the maintenance and expression of EBV genetic information in human lymphoblastoid cells. (20 refs)

78-6456 Studies of Epstein-Barr Virus (EBV)-Associated Nuclear Antigen: Solubilization from Raji Cell Chromatin with 5 M-Urea-2 M-NaCl and Fractionation on Hydroxyapatite. (Eng) Brown, T. D. (Beatson Inst. Cancer Res., Wolfson Lab. Molecular Pathology, Bearsden, Glasgow G61 1BD, Scotland); Rickwood, D.; MacGillivray, A. J.; Klein, G. *J Gen Virol* 40(3): 511-517; 1978.

A ¹²⁵I-IgG absorption assay was used to detect Epstein-Barr virus-associated nuclear antigen (EBNA) solubilized from Raji cell chromatin by treatment of the chromatin with high concentrations of urea and salt. The chromatin was fractionated into histone and nonhistone fractions on hydroxyapatite (HAP), and the nonhistone proteins were subfractionated by a discontinuous elution system. Very little antigenic activity was present in the histone-containing HAP 1 fraction, but activity was readily detected in the three non-histone fractions (HAP 2, 3, and 4). The absorption patterns of HAP 3 and HAP 4 differed, suggesting a fractionation of antigenic specificities. HAP 3 and 4 contained the bulk of the antigenic activity but only 7% of the total chromatin protein. The data suggest that EBNA is a nonhistone chromatin-associated protein. (27 refs)

78-6457 Alkaline Phosphatase in Epstein-Barr Viral Nuclear Antigen-positive Cell Lines. (Eng) Karpas, A. (Dept. Haematological Medicine, Univ. Cambridge, Cambridge, England); Hayhoe, F. G.; Greenberger, J.; Neumann, H. *Science* 202(4365): 318-319; 1978.

Alkaline phosphatase (APase) present in Epstein-Barr viral nuclear antigen (EBNA)-positive, surface membrane immunoglobulin (SmIg)-negative cell lines established from two leukemia patients was characterized. One of the patients had acute myeloid leukemia (AML) and the other a combined B- and T-cell acute lymphoblastic leukemia (ALL). The AML-derived cells contained myeloid APase, as determined by its ability to catalyze the hydrolysis of two types of substrates, ortho-substituted monoesters of orthophosphoric acid (substrate type 1) and the S-substituted monoesters of thiophosphoric acid (substrate type 2) with equal efficiency. The ALL-derived cells contained lymphoid APase, which catalyzes the hydrolysis of substrate type 1 only. Biochemical testing for two types of APase may allow distinction between primitive lymphoid and myeloid cells and may also be useful for detecting APase in cells that are cytochemically negative for the enzyme. The presence of the myeloid-specific enzyme

in an SmIg-negative cell line suggests that the line is composed of myeloid precursor cells and that such cells may be susceptible to infection with Epstein-Barr virus. (14 refs)

- 78-6458 Translocation of a Hydrocarbon Fluorescent Probe Between Epstein-Barr Virus and Lymphoid Cells: An Assay for Early Events in Viral Infection.** (Eng) Rosenthal, K. S. (Harvard Medical Sch., Boston, MA, 02115); Yanovich, S.; Inbar, M.; Strominger, J. L. *Proc Natl Acad Sci USA* 75(10): 5076-5080; 1978.

An assay to determine early biochemical events during cell-virus interactions is described, as well as its application in measuring events following the interaction of Epstein-Barr virus (EBV) with different types of lymphoid cells. The method is based on translocation of the hydrocarbon fluorescent probe diphenylhexatriene (DPH) between viral envelopes and cellular membranes, as monitored by fluorescence polarization (P) analysis. Using a model system, the high P value (0.324) of DPH-labeled cholesterol/phosphatidylcholine liposomes and the low P value (0.157) of DPH-labeled phosphatidylcholine liposomes allowed detection of DPH-translocation between interacting liposomes. This was monitored by the change in P in either direction. The P value of DPH-labeled EBV was significantly higher (p.350-0.392) than the P value of DPH-labeled lymphoid cells (0.238-0.289). Hence, DPH translocation could be detected by changes in P following incubation of DPH-labeled EBV and nonlabeled cells. A marked decrease in P was observed after incubation of DPH-labeled EBV with nonlabeled lymphoblastoid Raji cells or fresh human B lymphocytes. However, only a slight decrease in P was obtained when DPH-labeled EBV was incubated with nonlabeled fresh human T lymphocytes or fresh rabbit T or B lymphocytes. Moreover, incubation of human B lymphocytes with purified complement 3 (a putative inhibitor for the EBV receptor) prior to the addition of DPH-labeled EBV abolished the decrease in the P value. Most of these experiments were carried out with both the P3HR-1 and B95-8 EBV strains. Therefore, DPH translocation, as determined by fluorescence polarization, measures some early event during interaction of this enveloped virus and mammalian cells. The potential applicability of this technique to other viruses is illustrated by an experiment with Semliki Forest virus. (32 refs)

- 78-6459 An Evaluation of Antibody Titers to the Epstein-Barr Virus as a Test for Susceptibility to Cancer.** (Eng) Levine, P. H. (Lab. Viral Carcinogenesis, NCI, NIH, Bethesda, MD, 20014); Connelly, R.; Heath, C. W.; Caldwell, G. G.; Terasaki, P. I.; Mickey, M. R.; Fabrizio, P. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and

Prevention of Cancer. (New York, NY): Vol. 2 (part 1), 2 pp.; 1525-1537; 1978.

Antibodies to Epstein-Barr virus (EBV) capsid antigen (VCA) were studied in 210 normal first-degree relatives of cancer patients from 31 families with multiple cancer cases (MCF). Sera from each relative were matched by age, sex, and sex to a control sample from a normal individual. They were tested for VCA antibodies by immunofluorescence using a procedure using P3HR-1 antigen and Hyland goat anti-human IgG as antisera. VCA titers were significantly higher in normal MCF relatives compared with controls, which suggest that EBV-VCA titers may be indicative of an abnormality predisposing to malignancy. A comparison of EBV-VCA titers and histocompatibility antigens HL-A among patients with Hodgkin's disease or Burkitt's lymphoma and their relatives showed that high titers were not associated with specific HL-A types (A-1, B-5, B-8) linked to lymphoproliferative disease. (27 refs)

- 78-6460 Papain Solubilization of the Epstein-Barr Virus-induced Membrane Antigen.** (Eng) Peterson, G. R. (Dept. Microbiology, Mayo Clinic/Foundation, Rochester, MN, 55901); Qualtiere, L. F. *J Virol* 28: 344-351; 1978.

Epstein-Barr virus (EBV)-induced membrane antigen (MA) was purified and characterized in a series of experiments. EBV-infected Raji cells were exposed to papain (5 to 10 units/ 1×10^7 to 2×10^7 cells) for 1 hr. Membrane immunofluorescence and antibody-dependent cellular cytotoxicity (ADCC) assays of the cells revealed that most of the MA was removed by this treatment. Soluble MA was demonstrated in papain digests through inhibition of ADCC and by inhibition of the binding of anti-MA antibodies to EBV-infected Raji cells, as detected with the use of ^{125}I -labeled staphylococcal protein A. Papain extracts were centrifuged at 100,000 $\times g$ for 1 hr and tested for MA activity in the ^{125}I -protein A test. Approx 75% of this activity remained in the supernatant, indicating that most EBV MA activity in the extract was associated with low-mol-wt material. Antiserum prepared from an owl monkey immunized with the papain extract had no detectable antibodies to viral capsid antigen or early antigen; however, an antibody reactive to EBV-induced MA was present at a titer of 80 and a neutralization titer of 320. (23 refs)

- 78-6461 Isolation of Epstein-Barr Virus (EBV) in Suspensions of Cells Obtained from Patients with Different Hemoblastoses.** (Rus) Agrba, V. Z. (Inst. Experimental Pathology and Therapy, Acad. Medical Sciences, USSR, Moscow, USSR); Iakovleva, L. A.; Lapin, B. A.; Lebedev, V. N.; Grigor'ev, A. S.; Sangulia, I. A.; Timonovskaya, V. V.; Chuvirov, G. N.; Bukaeva, I. A.; Markova,

Markarian, D. S.; Kokosha, L. V. *Vopr Onkol* 24(9): 10; 1978.

sension cultures prepared from peripheral blood lymphocytes of leukemia patients and from lymph node cells and cells from patients with other hemoblastoses were investigated electron microscopically for the presence of Epstein-Barr virus (EBV). Eight virus-producing cell lines were established: 5 from 7 patients with myelogenous leukemia, from 9 patients with lymphogranulomatosis, and 1 from a patient with a lung tumor and a leukemoid reaction. No virus-producing cell lines could be established from three patients with lymphocytic leukemia or from another patient with non-Hodgkin's lymphoma. All eight cell lines produced virus in 0.5%-3% of the cells. The cells of the virus-producing lines were lymphoblasts with B-cell characteristics. (34

462 **Further Studies on the Replication of Adenovirus Type 2 DNA.** (Eng) Raptis, L. (Departement Microbiologie, Faculte de Medecine, Centre Hospitalier Universitaire, Universite de Sherbrooke, Sherbrooke, Quebec Canada J1H 5N4); Delbecchi, L.; Bourgaux, P. *Can J Microbiol* 56(9): 857-865; 1978.

distribution of ³H-thymidine in pulse-labeled replicating adenovirus 2 (Ad2) DNA was determined, and an attempt was made to pinpoint the origin(s) of replication on the Ad2 genome. Monolayer cultures of Ad2-infected KB cells were subjected to short pulses of ³H-thymidine at the time of max viral DNA synthesis. Nascent viral DNA was isolated from the cells, purified, and fractionated into partly completed (C) molecules. The latter were cleaved by *Eco*HindIII, *Hpa*I, or by *Hind*III + *Hpa*I, and the fragments were separated electrophoretically. R molecules were also treated with these enzymes, but the double-stranded fragments were separated from the partly single-stranded fragments by chromatography on benzoylated-naphthoylated diethylaminoethylcellulose before they were subjected to electrophoresis. Analysis of C molecules revealed a strong preferential labeling of fragments derived from the ends of molecules, as expected if both ends contain termini for replication. Analysis of double-stranded fragments from R molecules revealed a predominant labeling of the center of the genome that could be reconciled with initiation occurring internally or at both ends. In addition, two gradients of labeling arising toward each end and encompassing the outer quarter of the molecule, were noted that slowly receded when pulse length was increased. This finding may reflect an overrepresentation of molecules in the late stages of replication in the pool of nascent DNA, and it is to be expected if replication slows down when it nears the termination site(s). This interpretation is consistent with the labeling kinetics of various portions of the genome observed in the pool of molecules. Because of this complication, however, the distribution of label in R molecules cannot be used to determine

the exact location of the origin(s) of replication on the Ad2 DNA. (20 refs)

78-6463 **Size Distribution of Polyadenylated Adenovirus 2 RNA Synthesized in Isolated Nuclei.** (Eng) Zimmer, S. G. (Dept. Pathology, Univ. Kentucky Medical Center, MDRF 3, Lexington, KY, 40506); Goldenberg, C. J.; Carlson, D. P.; Craig, E. A.; Raskas, H. J. *Biochemistry* 17(20): 4207-4213; 1978.

Nuclei isolated from cultures of KB cells 14 hr after infection with adenovirus 2 were studied for their ability to transcribe viral RNA with fidelity and to generate discrete viral RNA size classes from specific regions of the genome. RNA synthesized in vitro was approx 40% viral and was derived almost entirely (95%) from the viral *r* strand. Hybridization-inhibition experiments demonstrated that the in vitro RNA transcripts included two classes of sequences, those restricted to the nucleus in vivo and those in cytoplasmic RNA. Quantitation of viral sequences from defined regions of the genome was performed by exhaustive hybridization with seven viral DNA fragments generated by digestion with endonuclease *R-Eco*R1 or *R-Sma*I. The sizes of the viral RNA synthesized by the KB cell nuclei in vitro were compared with viral RNA labeled by in vivo experiments (whole cell cultures). The polyadenylated viral RNA synthesized in vitro ranged in size from 13S to 28S, with peaks of 28S, 25S, and 22S. The polyadenylated nuclear RNA labeled during a 30-min in vivo pulse was larger, 23S-36S; however, incubation of the prelabeled nuclei in RNA synthesis conditions resulted in a gradual decrease in the size of the prelabeled RNA. RNA prelabeled in vivo contained 36S, 28S, and 22S species that, when incubated under in vitro conditions, were converted to a size distribution identical to that of RNA synthesized by isolated KB nuclei under in vitro conditions. This evidence supports the view that the in vivo and in vitro syntheses of RNA are identical. (46 refs)

78-6464 **Biochemical Studies on Bovine Adenovirus Type 3. IV. Transformation by Viral DNA and DNA Fragments.** (Eng) Igarashi, K. (Biological Res. Labs., Central Res. Div., Takeda Chemical Industries, Ltd., Osaka, Japan); Sasada, R.; Kurokawa, T.; Niiyama, Y.; Tsukamoto, K.; Sugino, Y. *J Virol* 28(1): 219-226; 1978.

Using the calcium technique, mouse cell line A31 (a clone of line BALB/3T3) was transformed in vitro with intact DNA of bovine adenovirus type 3 (BAV3) and also with BAV3 DNA fragments produced by two specific restriction endonucleases. Approx 5 to 10 foci/ μ g of DNA were produced within 20 days of transformation by BAV3 DNA. All foci were composed of round, cuboidal, and densely packed cells. RNase A and Pronase had little effect on the transformation, but DNase was able to abolish it completely. The *Eco*RI fragment D, mapping at 3.6 to 19.7 units, could transform

A31 cells almost as efficiently as intact DNA. No transformation occurred with *Hind*III fragments J, E, and B, but enzymatic joining of J + E (0-11.9 map units) with a ligase restored transforming activity. These results indicate that the genetic information of BAV3 necessary for transformation is located between map units 3.6 and 11.9, comprising 8.3% of the viral genome. Cells transformed by BAV3 DNA, BAV3 DNA *Eco*RI fragment D, and BAV3 *Hind*III fragment J + E (lines Tr, TrD, and TrJE, respectively) all showed the positive nuclear immunofluorescence characteristic of BAV3-transformed cells. Compared with untransformed A31 cells, all transformed lines tested showed rapid growth, high saturation density, and anchorage-independent growth. They also contained BAV3-specific tumor antigen. Lines Tr and TrD produced tumors in 100% of nude mice injected in 2 wk, and all three transformed lines produced tumors in 63%-100% of BALB/c mice within 2 mo. (23 refs)

- 78-6465 **Biochemical Studies on Bovine Adenovirus Type 3. III Cleavage Maps of Viral DNA by Restriction Endonucleases *Eco*RI, *Bam*HI, and *Hind*III.** (Eng) Kurokawa, T. (Biological Res. Labs., Central Res. Div., Takeda Chemical Industries, Ltd., Osaka, Japan); Igarashi, K.; Sugino, Y. *J Virol* 28(1): 212-218; 1978.

The restriction endonucleases *Eco*RI, *Bam* and *Hind*III were used to construct cleavage maps of bovine adenovirus type 3 (BAV3) DNA by analyzing overlapping sets of specific fragments produced by the enzymes and by cleavage of DNA previously end-labeled by T4 DNA polymerase. After digestion, fragments were analyzed by gel electrophoresis. *Bam*HI yielded 5 fragments (A-E); *Eco*RI, 7 (A-G); and *Hind*III, 12 (A-L). The order of fragments on the the DNA molecule was determined by overlapping the specific fragments produced by two different restriction endonucleases. The order was GDACBFE for the *Eco*RI fragments, AEBDC for the *Bam*HI fragments, and JEBKACDHFGIL for the *Hind*III fragments. The BAV3 preparation contains a unique population of incomplete virus whose genome has a deletion of 13% of the complete virus genome. Restriction endonuclease digestion of the incomplete virus DNA revealed that *Eco*RI E and F, *Bam*HI C, and *Hind*III G, I, and L fragments were missing. Therefore, the deleted region of incomplete virus DNA is located near the right end of complete virus DNA, a result consistent with previous electron microscope observations of heteroduplex molecules formed between complete and incomplete BAV3 DNA. (22 refs)

- 78-6466 **Transcription of Viral Sequences in Cells Transformed by Adenovirus Type 5.** (Eng) Frolova, E. I. (Inst. Molecular Biology, Acad. Sciences USSR, Moscow, USSR); Zalmanzon, E. S. *Virology* 89(2): 347-359; 1978.

Viral genomes and their transcription and chromatin conformation were studied in five lines of adenovirus 5 (Ad5)-trans-

formed rat embryo cells. Renaturation experiments showed that all lines contained sequences homologous to *Hpa*I fragment E, but they varied in other Ad5 DNA sequences. Any viral functions required for maintaining the transforming phenotype are encoded by the left end of Ad5 DNA. All viral DNA sequences in the three lines studied were transcribed into nuclear RNA. Only part of the sequences of I fragment E in all lines were sensitive to DNase I. (23 refs)

- 78-6467 **The Relationship Between T Antigens of group A and B Adenoviruses.** (Eng) Shirokawa, H. (Inst. Medical Science, Univ. Tokyo, Shirokawa Minato-ku, Tokyo 108, Japan); Shimojo, H.; Sekikawa, K. *Virology* 89(2): 627-631; 1978.

The relationship between the tumor antigens (TAgs) of groups A and B adenoviruses was investigated by immunofluorescence and DNA-DNA hybridization experiments. Secondary cultures of human embryo kidney cell 3Y1 cells (a line derived from a Fischer rat embryo) infected with adenovirus 12 (Ad12), Ad31, Ad7, Ad3, and Ad5 (10 plaque-forming units/cell) in the presence of μ g/ml of cytosine arabinoside. After 24 hr, cells were harvested and stained by indirect immunofluorescence with anti-TAg serum. TAg was detected in the nuclei of rats bearing GY1 cell (rat line transformed by the dIII-G fragment of Ad12 DNA) tumors. TAg with the morphology of cytoplasmic flecks and nuclear fibrils was observed in Ad12- and Ad31-infected cells, but TAg predominantly in the form of nuclear fibrils was observed in Ad7 and Ad3-infected cells. Thus, the morphology of TAg seems to be determined by virus type. TAg was not detected in Ad5-infected cells. These observations indicate that TAg is common among members of subgroups A (Ad12, Ad31, and B (Ad7, Ad3) adenoviruses, a conclusion that agrees with the high homology between the *Hind*III-G fragment of Ad12 DNA and Ad31 DNA and with the high homology between the *Hind*III-IJ fragment of Ad7 DNA and Ad3 DNA. In addition, TAg of Ad7 was partially stained by immunofluorescence with antiserum to Ad12 TAg and vice versa. However, the homology between the transforming fragment of Ad12 DNA and Ad7 DNA was only 9%-15%. The discrepancy between the common antigenicity and DNA homology, which also occurs with simian virus 40 and Ad virus, is discussed. (9 refs)

- 78-6468 **Adenovirus-induced Changes in the Weight of the Mouse Spleen and Thymus.** (Rus) Ageeva, A. I. (Lab. Virology, P. A. Hertsen Res. Inst. Oncology, Moscow, USSR); Erkhov, V. S.; Sviridova, I. K. *Biull Eksp Med* 86(9): 356-358; 1978.

In an attempt to detect lymphoid cells with increased proliferative potential, newborn CBA mice were inoculated with simian adenovirus (SA7, C8, 10^4 units/0.1 ml). Twenty days later, the mice were sacrificed and the spleen cells

transferred into newborn syngeneic mice (9×10^6 cells/mouse). The effect of the transfer was measured by changes in the wt of the lymphoid organs in 10-day-old mice. The transfer of splenocytes from mice during the latent period of SA7-induced carcinogenesis resulted in a significant increase in the spleen wt:body wt ratio (5.64, compared with 3.71 after transfer of normal splenocytes and 3.72 in controls) and significant decrease in the thymus wt:body wt ratio (4.04, vs 4.88 and 6.21, respectively). Thus, adenoviral carcinogenesis mimics some manifestations of autoimmune disease. (10 refs)

6469 Transcription Map for Adenovirus Type 12 DNA. (Eng) Smiley, J. R. (Dept. Therapeutic Radiology, Radiology Labs., Yale Univ. Sch. Medicine, New Haven, CT, 06510); Mak, S. *J Virol* 28(1): 227-239; 1978.

The physical locations and strand derivation of the adenovirus type 12 (Ad12) DNA sequences expressed as early and late cytoplasmic RNA (cyRNA) were mapped, and the relative rates of accumulation of this RNA from different regions of the genome were monitored at different times after infection. ^3H -labeled, intact, complementary l- and r-strand DNA fragments were hybridized with saturating amounts of unlabeled late cyRNA extracted 24 hr after infection of human cells with Ad12. The segments of each DNA strand complementary to the RNA were then purified by S1 nuclease digestion of the hybrids. These ^3H -DNA preparations were used as templates for hybridization with unlabeled DNA fragments generated by the restriction endonucleases *EcoRI*, *BamHI*, and *HindIII*. The completed map was very similar to that of Ad2. The subset of the late cyRNA sequences that were expressed at early times was located on the map by hybridizing labeled, early cyRNA to both unlabeled DNA fragments and unlabeled complementary strands of specific fragments. Early cyRNA hybridized to the r strand of *EcoRI*-C, *BamHI*-B and to the l strand of *BamHI*-E. Hybridization with *BamHI*-C was also observed. The relative rates of accumulation of cyRNA complementary to individual restriction fragments was measured at both early and late times. Early after infection, most of the viral RNA appearing in the cytoplasm was derived from the molecular ends of the genome. Later (24-26 hr postinfection), most of the newly synthesized cyRNA was transcribed from DNA sequences mapped between 25 and 60 map units on the genome. (39 refs)

6470 Morphological Revertants of Adenovirus Type 12-Transformed Hamster Cells. (Eng) Grone, G. J. (Inst. Genetics, Univ. Cologne, Cologne, W. Germany); Sutter, D.; Soboll, H.; Doerfler, W. *J Gen Virol* 40(3): 635-645; 1978.

B₁ subline of baby hamster kidney (BHK21) cells, the T637 derived by transformation of this subline with adenovirus type 12, and two morphological revertants of T637 (F10 and G12) were characterized. In contrast to T637

cells, which were round, the revertant lines exhibited a fibroblastic morphology, as did B₁ cells. Unlike B₁ and T637 cells, the revertant cells reacted with anti-LETS (large, external, transformation-sensitive protein) serum. The revertants were also less easily agglutinable by concanavalin A and less likely to form colonies in soft agar than T637, and they resembled B₁ in their uptake of amino acids and 2-deoxyglucose. Judging from the time required to induce tumor growth, G12 and F10 grew less readily in 4-wk-old Syrian hamsters than B₁ and T637 cells. The karyotypes of the four cell lines appeared to be indistinguishable, but the patterns of integrated virus genomes in the revertants differed markedly from that of T637 cells; F10 appeared to have lost all viral DNA sequences. (22 refs)

78-6471 Activated Endogenous Viruses of Tumor Induced by Human Adenovirus: An Electron Microscope Study. (Eng) Ohmori, M. (Dept. Pathology, Okayama Univ. Medical Sch., Okayama 700, Japan). *Exp Mol Pathol* 29(2): 260-271; 1978.

Activated endogenous C-type virus particles (ECP) and intracisternal A particles (IAP) in C3H/BiF₁ mice tumors induced by human adenovirus 12 (Ad12) were studied by electron microscope freeze-fracture-etching techniques. IAP were found in the endoplasmic reticulum (ER) and perinuclear cisternae. The ER membrane was electron dense and protruded into the cisternae to form either a single particle or many particles that were released eventually into the cisternae. The ECP particles began as spherical elevations on the tumor cell membrane; as the particles matured, the surface projections were lost and the particles became polygonal in shape. The production of ECP appeared to be continuous at some cell sites. Because these particles are not seen in chemically induced or in spontaneous tumors of syngeneic mice, it is assumed that Ad12 infection supplies the incomplete, masked genome of an endogenous leukemia virus with the material needed for its proliferation. (31 refs)

78-6472 Nucleotide Sequence Relationship Between Intracisternal Type A Particles of *Mus musculus* and an Endogenous Retrovirus (M432) of *Mus cervicolor*. (Eng) Kuff, E. L. (Lab. Biochemistry, NCI, NIH, Bethesda, MD, 20014); Lueders, K. K.; Scolnick, E. M. *J Virol* 28(1): 66-74; 1978.

The purified high-mol-wt RNA of *Mus musculus* (MM) intracisternal A-type particles was tested for homology with the viral genome of M432, a retrovirus isolated from *M. cervicolor* (MC). Partial nucleotide sequence homology was observed between the 32S and 35S RNA components of A-type particles from neuroblastoma cell line N4 (from a tumor in A/Jax mice) and the 70S fraction from M432. M432 complementary DNA (cDNA) was hybridized to about the same extent (25%-30% of the homologous reaction) by MM cellular

DNA and A-particle RNA; therefore, the entire group of M432-related MM sequences could be represented in the A-particle RNA. The hybrids showed a lower thermal stability than those formed with homologous RNA. The reaction was commensurate with that found between M432 cDNA and divergent sequences in the MM genome. The capacity to hybridize M432 cDNA was correlated closely with the concentration of A-particle sequences in the cytoplasmic RNA of several MM cell types. The major RNA fraction of M432 virus showed a reciprocal partial reaction with the A-particle cDNA's; the virus, which was grown in NIH/3T3 (MM) cells, also contained a small proportion of apparently authentic A-particle nucleotide sequences. However, a portion of the A-particle sequences, the so-called abundant class, seemed to be lacking in M432 genomic RNA. Multiple copies of both M432- and A-particle-related sequences were also present in MC DNA. However, the latter seemed to contain a more complete representation of the intracisternal A-particle information than was found in the M432 virus, since A-particle 35S cDNA was hybridized to a greater extent by MC cellular DNA than would have been expected on the basis of its partial homology with M432 viral RNA, and sequences homologous to the A-particle abundant-class probe were present in the DNA, even though they had no counterpart in the M432 viral genome. (31 refs)

78-6473 Defective C-type Retrovirus Particles Secreted by L1210 Leukemia Cells. (Eng) Kennel, S. J. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Tsakeres, F.; Kelly, P. A.; Allison, D. P. *Cancer Res* 38(12): 4574-4579; 1978.

C-type retrovirus particles released from cultures of L1210 leukemia cells (L1210) were characterized as a first step in determining the effect virus may have on the biology of the L1210 model system. The virus particle was found to be nononcogenic and noninfectious in standard assays for xenotropic and ecotropic murine retroviruses. This L1210 virus had a normal complement of viral proteins except that it had no detectable glycoprotein with a mol wt of 70,000 (gp70). Quantitative radioimmunoassay for the major structural protein, the one with a mol wt of 30,000 (p30), indicated that the virus is at least a thousandfold more defective than ecotropic viruses such as AKR and Moloney leukemia virus. L1210 cells grown in vitro released about 20 times more particle-associated p30 than in vivo-cultured cells, but cell-associated p30 and gp70 were found at similar levels. Both in vitro- and in vivo-cultured cells displayed gp70 on their surfaces that appeared to be of higher mol wt than the gp70 from most ecotropic viruses. Cells grown in vitro released a high level of non-virion-associated gp70 into the growth medium, but the high concentration of gp70 in the serum of normal DBA/2 mice precluded such measurements on in vivo-cultured cells. These results indicate that L1210 virus does not act as a classical infectious leukemia virus during the growth of L1210 cells, but the presence of relatively large amounts of viral antigens in these cells may affect the immune response to them. (28 refs)

78-6474 The Etiology of Laryngeal Papillomatosis. (Eng) Quick, C. A. (Div. Otolaryngology, V. A. Medical Center, Morgantown, WV, 26506); Fawcett, A.; Krzysek, R. *Laryngoscope* 88(11): 1789-1795; 1978.

Virological techniques for isolating whole virus, free viral DNA, or integrated viral DNA were used to determine whether the virus involved in the etiology of laryngeal papillomatosis is human papilloma virus. Human papilloma virus banded at 1.34 g/ml in a cesium chloride gradient; examination of the supernatant from pooled laryngeal papilloma specimens showed no such band, indicating that no virus was present. The electrophoretic mobility of the supernatant from a cutaneous wart was compared with that of markers of known molecular size, and the results were positive for free human papillomatosis virus DNA. When the technique was repeated with laryngeal papillomas, no such band was identifiable in the agarose gel. When integrated viral DNA was assayed by molecular hybridization, human papilloma virus type V showed 100% hybridization, but no hybridization was seen with the laryngeal papillomas. The results suggest that the virus responsible for laryngeal papillomatosis is not the same as the virus responsible for cutaneous papillomatosis lesions and, for the most part, exists in integrated DNA form. (7 refs)

78-6475 Interactions of Shope Papilloma Virus with Some Other DNA Viruses. (Eng) Chardon, Y. (Unite de Virologie Fondamentale et Appliquee, SERM, Groupe de Recherche 33, CNRS, 1, place du Professeur Joseph-Renaut, F 69371 Lyon Cedex 2, France); Grell, T.; Lyon, M.; Sohier, R. *Bull Cancer (Paris)* 65: 195-206; 1978.

The interactions of Shope papilloma virus (SPV) with mouse and rabbit cells in tissue culture were investigated. Rabbit cell cultures were derived from normal epidermis, SPV-infected epidermis, SPV-induced papillomas, and SPV-associated carcinoma. None of these cell cultures, whether infected in vitro with SPV or derived from tissue infected in vivo, produced infectious SPV or detectable viral antigens. Some other DNA viruses behaved differently: cells that had been in previous contact with SPV either in vitro or in vivo. Adenovirus type 5 multiplied better in human cells infected 24 hr previously with SPV than in untreated controls. The production of infectious virions of herpes simplex virus or Shope fibroma virus was reduced in cells derived from SPV-induced papillomas or carcinoma, due to a delay in viral maturation. Of the rabbit cells, only those derived from in vivo infected tissue or those previously infected in vitro with SPV were transformed by simian virus 40. The rabbit cell lines derived from papillomas or carcinoma differed from their counterparts derived from normal epidermis or from tissue infected 24 hr before biopsy in their karyotype and in their ability to grow in soft agar. Similar karyotypic alterations were induced in cells derived from healthy epidermis by infection in vitro with SPV. (29 refs)

- 476 **Heterogeneity of Human Papilloma Viruses.** (Eng) Pfister, H. (Institut für Virologie, Zentrum für Hygiene, Hermann-Herder-Strasse 11, 7800 Freiburg, W. Germany); Gissmann, L. *Bull Cancer (Paris)* 65(2): 67; 1978.

types of human papillomaviruses, HPV-1 and HPV-4, characterized by purifying virions from single skin. Viral DNA was extracted by solubilizing the protein with sodium dodecyl sulfate and then cleaved with different restriction endonucleases. The resulting fragments separated by gel electrophoresis. The results show no relationship between HPV-1 and HPV-4 as far as the following criteria are concerned: the pattern of restriction enzyme fragments of their DNA, DNA-complementary RNA hybridization, and complement fixation with specific rabbit antiserum. Moreover, the mol wts of their major proteins differed significantly. Both virus types could be isolated from plantar warts and from verrucae vulgares. In addition, a third type of HPV was isolated from the skin warts, with HPV-1 or with HPV-4 rabbit antiserum. (10 refs)

- 477 **Mastomys natalensis Papilloma Virus (MnPV), the Causative Agent of Epithelial Proliferation: Characterization of the Virus Particle.** (Eng) Müller, H. (Institut für Virologie, Justus-Liebig-Universität Giessen, Justus-Straße 107, 6300 Lahn-Giessen, W. Germany); Gissmann, L. *J Gen Virol* 41(2): 315-323; 1978.

physicochemical and morphological characteristics of *Mastomys natalensis* papilloma virus (MnPV) were studied. Virus particles banded in CsCl gradients at densities of 1.34 g/ml (full particles) and 1.29 g/ml (empty particles). The DNA had a buoyant density of 1.7104 g/ml, and it existed in three different conformations (supercoiled circular, relaxed circular, and linear). The sedimentation values of the three conformations were 23S-24S, 16S-17S, and 15S, respectively. Although the mol wt of MnPV DNA ($\times 10^6$) was similar to that of human papilloma virus (HPV) DNA, the size of the fragments obtained after cleavage of MnPV DNA with the restriction endonuclease *Hae* III differed from that of HPV cleavage fragments. Polyacrylamide gel electrophoresis showed that virion contained different polypeptides; the major structural protein had a mol wt of 56,000. Skin tumors were induced in 36% of mice infected with purified MnPV (containing approx 150 ng of virus protein), but no cutaneous alterations were observed when the particles were inoculated in the presence of anti-MnPV serum; this indicated that MnPV was the etiologic agent in the skin proliferations. MnPV was reisolated from the experimental tumors. (22 refs)

- 478 **Conditions Leading to the Establishment of the N (a Gene Dependent) and A (a Gene Independent) Transformed States after Polyoma Virus Infection of**

Rat Fibroblasts. (Eng) Rassoulzadegan, M. (Centre de Biochimie, Université de Nice, 06034 Nice, France); Seif, R.; Cuzin, F. *J Virol* 28(2): 421-426; 1978.

Infection of normal rat fibroblasts (FR 3T3) with the early temperature-sensitive mutant (*tsa*) of polyoma virus was shown to lead to either the A or the N phenotype. The choice between A and N was made during the first 5 days at 33°C and primarily depended on the growth potential of the infected cells during that time period. Polyoma *tsa*-transformed A clones were temperature-independent for expression of the transformed phenotype, but N transformants were strongly temperature-dependent. The crucial parameter in the N vs A choice is probably the physiological state of the cells during the early part of transformation, and it is likely that no causal relationship exists between the method of selection and the subsequent mode of maintenance. A transformation occurred in both liquid and agar media when the infected cells were maintained under growth-restricting conditions, such as absence of anchorage and contact inhibition at confluency. N transformation occurred in cells maintained in active growth after virus infection (sparse cultures on a solid substratum). The N and A states were redefined as meaning polyoma *a* gene (or simian virus 40 *A* gene) dependent vs independent for maintenance of the transformed state. (12 refs)

- 78-6479 **Tumor Antigens Induced by Nontransforming Mutants of Polyoma Virus.** (Eng) Silver, J. (Dept. Hematology, Beth Israel Hosp., Boston, MA, 02115); Schaffhausen, B.; Benjamin, T. *Cell* 15(2): 485-496; 1978.

Wild-type polyoma virus and nine independently isolated hr-t mutants of polyoma virus (with an absolute defect in transformation) were studied to determine which tumor (T) antigens they induced during lytic infection in mouse cells. In a variety of cell lines, wild-type virus induced a major T-antigen species with an apparent mol wt of 100,000 daltons and four minor T-antigen species with apparent mol wts of 63,500, 56,000, 36,000, and 22,000 daltons. The hr-t mutants induced a normal 100,000-dalton T antigen but none of the minor wild-type virus species; it did induce antigens with mol wts of 6,000-9,000 daltons. "Point" hr-t mutants induced normal amounts of the 56,000-dalton antigen and reduced amounts of the 22,000-dalton antigen but neither of the other two minor antigens. Ts-a mutants, which have a temperature-sensitive defect in ability to induce stable transformation induced T antigens with the same mobility as wild-type virus, but the 100,000-dalton species was thermolabile compared with the wild-type species. A double mutant virus carrying both a ts-a mutation and a deletion hr-t mutation induced a thermolabile 100,000-dalton antigen but none of the minor species. Cell fractionation studies with productively infected cells were carried out to localize the T-antigen species. (55 refs)

- 78-6480 **Contact Behaviour and Pattern Formation of BHK and Polyoma Virus-transformed BHK Fi-**

broblasts in Culture. (Eng) Erickson, C. A. (Dept. Biology, Yale Univ., New Haven, CT, 06520). *J Cell Sci* 33: 53-84; 1978.

The contact behavior of normal (BHK21/13) and polyoma virus-transformed (BHK21) cell lines was studied using high-resolution, time-lapse cinemicrography combined with scanning electron microscopy. At high density, the BHK21/13 cells were generally organized in monolayered parallel arrays with little crisscrossing of whole cells, whereas the BHK21 cells showed a multilayered, highly crisscrossed arrangement with considerable nuclear overlap. In both cell lines, ruffling was inhibited when contact was made ruffle to ruffle. When BHK21/13 cells contacted each other ruffle to side, strong adhesions occurred with accompanying paralysis of ruffling. Then the contacting cell either changed direction, usually spreading along the side of the contacted cell, or it occasionally continued to protrude and underlap the other cell, although the original adhesions remained. The transformed cells never formed strong ruffle-to-side adhesions, and they usually underlapped each other totally. None of the cells moved over the surface of others (overlapping), and all crisscrossing occurred by underlapping. The BHK21 cells had many fewer side-to-substratum adhesions than the BHK21/13 cells, which could explain the ease with which the transformed cells underlap. (36 refs)

78-6481 Growth Factors Produced by Sarcoma Virus-transformed Cells. (Eng) Todaro, G. J. (Lab. Viral Carcinogenesis, NCI, NIH, Bethesda, MD, 20014); De Larco, J. E. *Cancer Res* 38(11, part 2): 4147-4154; 1978.

The epidermal growth factor (EGF) binding capacity of a variety of normal and transformed cell lines was determined. Simian virus 40 (SV40), polyoma virus, and avian sarcoma virus transformants showed normal levels of EGF binding. Only 5/47 chemically transformed cells had lost their EGF specific receptors. However, the binding levels of all murine sarcoma virus (MSV)-transformed lines tested (mouse, rat, and mink) were altered; in none of the cases was multiplication-stimulating activity (MSA) binding reduced. Thus, loss of EGF receptors after transformation appears to be specifically associated with sarcoma virus transformation. Of the >40 human fibrosarcomas tested, 2 had normal levels of EGF receptors but no MSA receptors. One of the MSA receptor-negative lines was producing a family of related growth polypeptides that interacted with MSA receptors but not with EGF receptors and stimulated cell division in cells of various species. The activities of both the sarcoma growth factor and the MSA-related human fibrosarcoma growth factor were heat-stable, protease-sensitive, and sensitive to disulfide-reducing agents. It appears that the production of the sarcoma growth factor is at least in part the mechanism by which sarcoma viruses transform fibroblasts. One of the ways in which cells become transformed may be by endogenous production of growth factors for which they have receptors

and to which they are capable of responding. This intracellular production of growth polypeptides serves as a constant stimulus for cell division, thereby releasing these cells from the constraints of the normal physiological controls. (29 refs)

78-6482 Replication of Polyoma DNA in Isolated Nuclei. Synthesis and Distribution of Initiator RNA. (Eng) Eliasson, R. (Medical Nobel Inst., Div. of Biochemistry, Karolinska Inst., Stockholm, Sweden); Reichard, P. *J Biol Chem* 253(20): 7469-7475; 1978.

The synthesis and distribution of initiator RNA (iRNA) was studied in nuclei isolated from polyoma virus-infected mouse fibroblasts. The nuclei were incubated for 5 min at 37°C under standard conditions, including 2 mM ATP, ¹⁴C-dATP, ³H-guanosine triphosphate (³H-GTP), and ³²P-GTP. Polyoma DNA was extracted selectively, and replicative intermediates (RI) and iRNA were purified and measured. The frequency with which ATP or GTP was incorporated into iRNA during incubation depended on the ATP/GTP ratio. The total number of 5'-ends of iRNA most equalled the total number of 3'-ends, indicating that most iRNA molecules are complete. Newly synthesized iRNA attached to short Okazaki fragments was first detected at 1.5 min. Between 1.5 and 5 min, iRNA accumulated on longer strands, but little change occurred after 5 min. At times (1.5, 5, 10, and 15 min), there was an unexplained accumulation of iRNA linked to very short strands (25-30 nucleotides in length). Only one-fourth to one-third of the Okazaki fragments contained iRNA. This value was independent of the length of the strands (270 or 95 nucleotides) or incubation time. Since the uniqueness of iRNA is not size, not its nucleotide sequence, it is unlikely that a specific base sequence of the template determines either the beginning or the end of iRNA synthesis. It is suggested that iRNA serves as a primer for the synthesis of Okazaki fragments, and that it is transcribed randomly on the genome during iRNA replication. (32 refs)

78-6483 Polyoma Virus Giant RNAs Contain Tandem Repeats of the Nucleotide Sequence of the Entire Viral Genome. (Eng) Acheson, N. H. (Swiss Inst. Experimental Cancer Res., 1066 Epalinges, Switzerland). *Proc Acad Sci USA* 75(10): 4754-4758; 1978.

The arrangement of viral nucleotide sequences in the late virus-specific RNA's synthesized in polyoma virus-infected mouse cells was studied by electron microscopy of hybrids between purified high-mol-wt viral RNA and the HindII-1 fragment of polyoma DNA, which contains 91% of the viral genome. Hybrid molecules containing a short single-stranded gap (corresponding to the 9% of viral sequence not present in HindII-1), flanked by double-stranded regions, were photographed and measured. Most hybrid molecules

tained no single-stranded loops or branches, demonstrating that all viral sequences are transcribed contiguously and that no nonviral sequences are present in the RNA. In hybrid molecules containing RNA up to 3.5 times the genome length, a repeating structure of single-stranded gaps 8% of genome length was observed interspersed with double-stranded regions 89% of the genome length; therefore, polyoma RNA's contain tandem repeats of the nucleotide sequence of the entire viral DNA. A small proportion of hybrid molecules contained single-stranded branches and deletion loops in characteristic positions, indicating that RNA splicing may occur on high-mol-wt nuclear polyoma RNA. (29 refs)

6484 Characterization of the mRNA's for the Polyoma Virus Capsid Proteins VP1, VP2, and VP3. (Eng) Hunter, T. (Tumor Virology Lab., Salk Inst., San Diego, CA, 92112); Gibson, W. *J Virol* 28(1): 240-253; 1978.

The polyoma virus capsid proteins VP1, VP2, and VP3 of the 126 kD-type polyoma virus were identified following translation of total cytoplasmic RNA in the wheat germ system in the presence of ³⁵S-methionine. Upon sodium dodecyl sulfate-polyacrylamide gel electrophoresis, bands corresponding to the three capsid proteins were evident. The bands were identified from analyses of mock infected cultures. The translation products of RNA selected from total cytoplasmic RNA of infected cells by hybridization to polyoma virus DNA showed a high degree of enrichment for VP1, VP2, and VP3. Fractionation of this virus-specific RNA showed that almost all VP1 messenger RNA (mRNA) activity contains poly-A tails, that a substantial proportion of VP2 and VP3 mRNA activity is poly-A deficient. Further proof of the identity of these proteins was afforded by close agreement of the in vitro and in vivo tryptic peptide maps and by the fact that the mobilities of the in vitro products on two-dimensional gels were the same as those of viral proteins labeled in vivo. Gel electrophoresis of in vitro products synthesized from polyoma virus-specific RNA in the presence of ³⁵S-formylmethionine-transfer RNA(met)(f) showed that all three proteins were labeled. Synthesis of all these proteins was inhibited by 80% with 1 mM-guanosine triphosphate, suggesting that their 5' ends are all capped. The sizes of the major mRNA's for these proteins were determined by gel fractionation to be 4.6 x 10³ daltons for VP1, 8.5 x 10³ daltons for VP2, and 7.4 x 10³ daltons for VP3. It is concluded that all three proteins are synthesized independently in vitro and are virally coded and that each has a discrete mRNA. (37 refs)

6485 Splicing of SV40 Late mRNA Is a Post-transcriptional Process. (Eng) Horowitz, M. (Dept. Genetics, Weizmann Inst. Science, Rehovot, Israel); Laub, J.; Bratosin, S.; Aloni, Y. *Nature* 275(5680): 558-559; 1978.

The splicing of simian virus 40 (SV40) late messenger RNA (mRNA) was investigated by analyzing R-loop structures

formed between nuclear (nu) and cytoplasmic (cy) poly(A)+ RNA's and linear SV40 DNA. R loops were obtained by annealing nu or cy 19S poly(A)+ RNA with double-stranded linear SV40 DNA obtained by cleavage of form I DNA with *Bgl*I or *Taq*I restriction endonucleases. The 5' ends of the nu RNA R loops were closer to the cleavage sites than the 5' ends of the cy RNA R loops. Free leaders were more common at the 5' ends of the cy RNA R loops than at the 5' ends of the nu RNA R loops. The ability of the precursor nu poly(A)+ viral RNA to anneal to the entire length of the late portion of SV40 DNA (approx 0.67-0.17 map units), including the corresponding interruptions between the leaders and the bodies of the cy mRNA's indicates that splicing is post-transcriptional. Splicing may occur in the nucleus before the RNA molecule is transported to the cytoplasm. However, it is possible that splicing occurs during transport or immediately after transport. The excision-ligation reaction in the precursor molecule is presumably carried out by an enzyme(s) that recognizes short base-paired sequences in this molecule. This reaction would represent a novel mechanism in the regulation of gene expression in animal cells. (17 refs)

78-6486 Immunocytochemical Evidence for SV40-Related T Antigen in Two Human Brain Tumours of Ependymal Origin. (Eng) Tabuchi, K. (Dept. Neurological Surgery, Okayama Univ. Medical Sch., 2-5-1 Shikata-Cho, Okayama 700, Japan); Kirsch, W. M.; Van Buskirk, J. J. *Acta Neurochir (Wien)* 43(3/4): 239-249; 1978.

Direct and indirect immunoperoxidase techniques were used to screen 39 human brain tumors for the presence or absence of simian virus 40 (SV40) T antigen. The tumors included 13 glioblastomas, 5 astrocytomas, 1 ependymoma, 1 oligodendroglioma, 1 choroid plexus papilloma, 2 medulloblastomas, 1 neurinoma, 10 meningiomas, 1 hemangiopericytoma, 1 angiofibroma, and 3 metastatic cancers. The nuclei of the ependymoma and the choroid plexus papilloma exhibited a strong positive reaction for T antigen. The remaining 37 tumors were negative for the antigen. In the ependymoma, the presence of T antigen was confirmed by a complement-fixation assay of tumor extract. It is noted that the majority of hamster intracerebral tumors induced by papovaviruses are ependymomas or choroid plexus papillomas. (42 refs)

See also:

*(Rev.): 78-6013, 78-6084, 78-6085, 78-6086, 78-6087, 78-6088, 78-6089, 78-6090, 78-6091, 78-6092, 78-6093, 78-6094, 78-6095, 78-6096, 78-6097, 78-6098.

*(Chem.): 78-6219, 78-6230, 78-6242, 78-6315, 78-6338.

*(Immun.): 78-6489, 78-6496, 78-6497, 78-6499, 78-6500, 78-6501, 78-6503.

*(Path.): 78-6521, 78-6534, 78-6552.

*(Epid.-Biom.): 78-6571.

- 78-6487 Evidence of Acquired Immune Deficiencies in Mediterranean Lymphoma. A Possible Aetiological Link.** (Eng) Al-Saleem, T. I. (Dept. Pathology, Coll. Medicine, Univ. Baghdad, Baghdad, Iraq). *Lancet* 2(8092, part 1): 709-712; 1978.

Cellular and humoral immunity was systematically studied in 57 patients with α heavy-chain-positive or -negative Mediterranean lymphoma (ML) and 425 controls (healthy volunteers and patients with minor illnesses or other types of lymphoma). Patients with ML had statistically significant reductions in serum IgG and IgM, whereas patients with Hodgkin's disease and nonintestinal non-Hodgkin's lymphoma did not. Impaired cellular immunity was observed in 50% of ML patients challenged with two recall antigens (mumps and purified protein derivative of tuberculin) and with dinitrochlorobenzene. Comparable anergy was found only in patients with advanced Hodgkin's disease. It is hypothesized that repeated exposure of the intestinal epithelium to *Vibrio cholerae* toxin and/or other biologically similar enterotoxins may produce local (and possibly delayed) general impairment of immunity and thus predispose the patient to the development of lymphoplasmacytic neoplasia. Animals that are genetically susceptible to autoimmune disease may be useful as experimental models to test this hypothesis. (35 refs)

- 78-6488 Role for Monocyte-Macrophage-derived Colony-stimulating Factor and Prostaglandin E in the Positive and Negative Feedback Control of Myeloid Stem Cell Proliferation.** (Eng) Kurland, J. I. (Dept. Medicine, UCLA Medical Center, Los Angeles, CA); Broxmeyer, H. E.; Pelus, L. M.; Bockman, R. S.; Moore, M. A. *Blood* 52(2): 388-407; 1978.

Female B6D2F₁ mouse peritoneal macrophages and human peripheral blood monocytes elaborated colony-stimulating factors (CSF) that stimulated the proliferation in vitro of a class of hematopoietic progenitor cells (colony-forming units-culture, CFU-C) restricted to granulocyte and monocyte-macrophage differentiation. High numbers of monocyte-macrophages had little stimulating activity and were markedly suppressive because of the accumulation of a dialyzable, non-species-specific inhibitor of CFU-C proliferation identified by radioimmunoassay as prostaglandin E (PGE). PGE levels correlated inversely with the incidence of the CFU-C, or committed myeloid stem cells. Indomethacin suppression of PGE synthesis increased the number and size of myeloid colonies stimulated, allowing more accurate estimates of CFU-C incidence. The dose-dependent stimulation of PGE synthesis by macrophages in response to increasing

concentrations of CSF was accompanied by progressive CFU-C inhibitory activity. When human monocytes were exposed to an extract of human granulocytes that decreased CSF production, monocyte PGE synthesis was reduced significantly. The findings indicate that the in vitro proliferation of committed myeloid stem cells is regulated by both positive and negative feedback controls involving monocyte-macrophage-derived CSF and PGE, respectively. The balance between CSF and PGE, factors with mutually opposing actions, is determined by an afferent feedback mechanism operating via the intrinsic modulation of macrophage PGE synthesis in response to local elevations in CSF concentration. (44 refs)

- 78-6489 Age-dependent Leukemia Produced in Thymectomized AKR Mice by Antiserum to Leukemic Cells.** (Eng) Kohn, R. R. (Inst. Pathology, Case Western Reserve Univ., Cleveland, OH, 44106); Sciotto, C. G.; Hensley, S. *Exp Gerontol* 13(3/4): 181-187; 1978.

The hypothesis that immunological attack can cause malignant transformation was tested by treating thymectomized (Th-x) AKR mice of different ages with rabbit antiserum against leukemic cells (ALS), antiserum to nonleukemic cells (ANS), and normal rabbit serum (NRS). Groups of young and old non-Th-x mice were also treated with antisera. There were three series of experiments, one with 366-day-old mice, one with 439-day-old mice, and one with seven survivors of Th-x controls from the first series treated at 500 days of age. Leukemic deaths and ages at death were determined. In the first series, by 500 days of age, 9/10 intact mice but only 3/10 Th-x mice had died of leukemia, and deaths in the latter were delayed. Three out of five Th-x mice receiving ANS and 1/5 receiving NRS died from leukemia. All five Th-x mice that received ALS died from leukemia soon after treatment. In the second series, among Th-x controls and mice receiving NRS, only one died from leukemia by 500 days of age. On the other hand, 3/7 mice receiving ANS died from leukemia by 500 days of age. In the third series, when ALS was given to 500-day-old Th-x mice, the results were equivocal: there were 2 early leukemia deaths, 2 early nonleukemic deaths, and 3 survivors. Antisera given to non-Th-x mice did not significantly affect the age at which leukemia developed. However, there was no clear leukemia-producing effect of antisera in Th-x mice > 500 days old. The leukemogenic effect of antiserum in Th-x mice correlated with the antiviral (Gross leukemia virus) activity of the serum; the ALS interacted most effectively with the virus and caused the highest incidence of leukemia. It is concluded that leukemic transformation is a response to immunological attack on the virus or on cells expressing viral antigens and that an additional age-dependent factor is required for leukemogenesis. (21 refs)

6490 Macrophages and Resistance to Tumors. II. Influence of Agents Affecting Macrophages and Delayed-Type Hypersensitivity on Resistance to a Tumor Inducing Specific 'Sinecomitant' Immunity: Acquired Resistance as an Expression of Delayed-Type Hypersensitivity. (Eng) Nelson, M. (Kolling Inst. Medical Res., Royal North Shore Hosp., Sydney, St. Leonards, New South Wales 2065, Australia); Nelson, D. S. *Cancer Immunol Immunother* 4(2): 107; 1978.

The influence of agents affecting macrophages and delayed-type hypersensitivity (DTH) on resistance to the 3-methylcholanthrene-induced fibrosarcoma H-7 (to which specific resistance can be induced by a low, non-tumorigenic dose of cells) was determined in CBA/H/WEHI mice. The expression of DTH to sheep RBC was inhibited by carrageenan (CG: 2 mg/mouse ip), niridazole (200 mg/kg po), reserpine (RP: 1 mg/kg iv), and by whole-body irradiation (450 and 550 rads, ^{60}Co source, 37 r/min). The specific resistance of immunized mice (2.5 x 10^5 viable cells sc, 17, 18, or 21 days prior to challenge) to pad challenge with H-7 cells was also inhibited by these agents, as shown by progressive tumor growth in the feet. Tumor growth in the feet of nonimmunized mice was not affected by radiation or niridazole, but it was inhibited or delayed by CG, depending on the dose of tumor cells injected. CG permitted the emergence of tumors from an inoculum of 10^4 cells, reduced tumor growth from 5×10^4 cells, had no effect on growth from 10^5 cells, and increased tumor growth from 5×10^5 cells. Also, CG treatment just before challenge with 5×10^5 cells led to tumor growth early after challenge, but CG treatment 16-19 days before challenge led to some restriction of growth. Paradoxically, RP inhibited tumor growth in nonimmune mice profoundly, which suggests that some inflammatory reaction may be necessary for survival of tumor cells injected into the footpad. It was suggested that specific, acquired resistance to H-7 is genetically akin to DTH and that mice offer some natural resistance to this tumor. (36 refs.)

6491 Tumor Induction in Host-Versus-Graft Disease. I. Clinical and Pathologic Features. (Eng) Cornejo, E. A. (Dept. Diagnostic Radiology, Yale Univ. Sch. of Medicine, 333 Cedar St., New Haven, CT, 06510). *Am J Pathol* 109(3): 675-688; 1978.

Inbred C57BL/1 mice given injections of (SJL/J x C57BL/1) F_1 spleen cells developed a runting syndrome designated host-vs-graft disease (HVGd). Mortality from the disease was related to the dose of F_1 cells. Acute pathologic changes resembled those occurring in parent- F_1 graft-vs-host disease (GVHD), although plasmacytosis was more pronounced. Mice who recovered from HVGd showed a slight increase in lymphoma incidence over control mice. Such mice were hyperreactive to the F_1 cells used to initiate the original HVGd syndrome. Most of the tumors developed in those animals receiving the initial injection of F_1 spleen cells within

24 hr of birth. Tumor incidence was unrelated to the clinical severity of HVGd. By contrast, GVHD in the same strain combination resulted in a much higher incidence of lymphomas in a much shorter time. Parental strain cells were detectable in the F_1 hosts up to the time of tumor development. HVGd has a low tumor induction potential; GVHD has a high tumor induction potential. (29 refs)

78-6492 Transfer of Viable Putative Preneoplastic Hepatocytes to the Livers of Syngeneic Host Rats. (Eng) Laishes, B. A. (McArdle Lab. Cancer Res., Univ. Wisconsin, Madison, WI, 53706); Farber, E. *J Natl Cancer Inst* 61(2): 507-512; 1978.

A new technique for the transfer of freshly isolated, carcinogen-altered hepatocytes containing the marker γ -glutamyl transpeptidase (GT) from male F344 albino rats to syngeneic host rat livers is described. Donor mice were inoculated on day 1 with diethylnitrosamine (200 mg/kg ip), fed 2-acetylaminofluorene (2-AAF: 0.02% of diet) on days 14-28, and given a two-thirds hepatectomy (PH) on day 21; cells were isolated on day 31. Hepatocyte suspensions were prepared by proteolytic enzyme perfusion, and transfer was effected by infusion of 0.3 ml of suspension into the hepatic portal vein of rats that had been fed a control diet or the 2-AAF diet for 7 days. All recipients underwent PH on day 7 or 10. Selective proliferation of GT-positive hepatocytes was induced in the livers of hosts that had been fed 2-AAF: large, macroscopic colonies of altered hepatocytes were generated within 10 days of cell transfer. Quantitation of the number of colonies revealed that prior treatment of hosts with a low dose of 2-AAF was essential for the appearance of large numbers of foci. In addition, the appearance of foci on day 10 depended on the presence of intact, GT-positive hepatocytes in the transferred cell suspensions. (24 refs)

78-6493 Tumor-associated Transplantation Antigen Distinct from H-2k-like Antigens on a BALB/c (H-2d) Fibrosarcoma. (Eng) Parmiani, G. (Div. Experimental Oncology A, Istituto Nazionale per lo Studio e la Cura dei Tumori, Via G. Venezian 1, 20133 Milan, Italy); Meschini, A.; Invernizzi, G.; Carbone, G. *J Natl Cancer Inst* 61(5): 1229-1234; 1978.

To determine whether the foreign H-2.1, H-2.5, H-2.11, H-2.23, and H-2.25 antigens on the 3-methylcholanthrene (3-MC)-induced (H-2d) fibrosarcoma C-1 of BALB/cLacDp (C) mice represent the tumor's individual tumor-associated transplantation antigen (TATA), in vivo immunogenicity tests were performed in C mice (H-2d) and in a variety of hybrids of various haplotypes. The C-1 tumor lost its immunogenicity completely in CC3F $_1$ mice (H-2d x H-2k) and partially in CAF $_1$ mice (H-2d x H-2a). Multiple immunizations with C-1 cells in C and hybrid mice showed that C mice had in vivo resistance to tumor challenges of 10^5 C-1 cells

(anti-TATA immunity) and production of antibodies to H-2k antigens, as revealed by a complement-dependent cytotoxicity assay of normal C3Hf/HeDp (C3) lymphocytes. No effector lymphocytes to H-2k antigens were found in mice that had detectable levels of anti-H-2k antibodies. CC3F₁ animals were unable to mount both anti-TATA and anti-H-2k immune responses, whereas CAKF₁, CBRF₁ (both *H-2d* x *H-2k*), and, to a lesser extent, CAF₁ mice displayed anti-TATA immunity in the absence of anti-H-2k antibody production. CD2F₁ (*H-2d* x *H-2d*) mice behaved like C mice, whereas CB6F₁ (*H-2d* x *H-2b*) and CNIHF₁ (*H-2d* x *H-2q*) crosses were resistant to tumor challenge but had lower titers of anti-H-2k antibodies than C mice. Anti-H-2k antibody production in C mice peaked after the sixth immunization, with a very low level being evident after one or two immunizations, when a clear antitumor immunity was present. It is concluded that H-2k antigens and TATA are probably independent antigenic entities on C-1 cells. (13 refs)

78-6494 Stimulating Capacity of Blast Cells from Patients with Chronic Myelocytic Leukaemia, in Blastic Crisis in 'One-Way' Mixed Lymphocyte Reaction: Lack of Evidence for T Lymphoblastic Conversion. (Eng) Han, T. (Depts. Medicine B and Immunology Res., Roswell Park Memorial Inst., New York State Dept. Health, Buffalo, NY, 14263); Gomez, G. A.; Minowada, J. *Immunology* 35(2): 299-305; 1978.

The stimulatory capacity of fresh leukemia blast cells from patients with Philadelphia chromosome (Ph¹)-positive chronic myelocytic leukemia-blastic crisis (CML-BC, 6 patients), acute myeloblastic leukemia (AML, 10), and T- and null-cell type acute lymphoblastic leukemia (ALL, 1 and 5, respectively) on allogeneic lymphocytes in one-way mixed lymphocyte reactions was studied. The blast cells from the CML-BC patients significantly stimulated the allogeneic T lymphocytes. The degree of stimulation increased with the number of blast cells, and there was no apparent difference in stimulatory capacity between myeloid-type and lymphoid-type blasts. The blast cells from the AML patients and from three with null-cell ALL also stimulated allogeneic lymphocytes, but cells from the patient with T-cell ALL and cultured leukemic T lymphoblastoid cells (2 lines) showed no stimulating capacity. The cells from two patients with null-cell ALL were not stimulatory. However, cultured leukemic null cells (4 lines) consistently exerted a strong stimulation in the mixed lymphocyte reaction. The data suggest that leukemic cells from patients with lymphoblastic type CML-BC are not T lymphoblasts, although the possibility that they are non-T/non-B lymphoblasts cannot be ruled out entirely. (22 refs)

78-6495 Cell-mediated Lympholysis: A Receptor-associated Lytic Mechanism. (Eng) Gensheimer, G. G. (Transplantation Biology Section, Immunology

Branch, NCI, NIH, Bethesda, MD, 20014); Neefe, J. R. *Immunol* 36(1): 54-64; 1978.

The mechanism of cell-mediated lympholysis was investigated. Two major hypotheses have been proposed to explain the relationship between the binding stage and the killing stage: (1) lysis is a nonspecific result of the intimacy created between killer and target by specific antigen-receptor binding and (2) lysis is effected by the killer cell receptor itself or a closely linked cell-surface component. Two populations of cytotoxic effector cells were coincubated, one relevantly sensitized to the second and the second irrelevantly sensitized to a third party; in some cases the third-party cells were added. Relevantly sensitized killers were not lysed when coincubated with either cytotoxic targets alone or with cytotoxic targets and their appropriate third-party targets. It is clear, therefore, that the lytic event in cell-mediated killing is not simply a result of the intimacy created by attacker-target binding. In addition, receptor accommodation does not produce an activated state in which killer cells secrete nonspecific soluble toxic factors. These findings suggest, rather, that the lytic event in cell-mediated lympholysis is accomplished by the receptor itself or by a geographically close linked cell-surface component. (17 refs)

78-6496 Characterization of T Antigens in Polyoma Virus-Infected and Transformed Cells. (Eng) Hutcheson, M. A. (Tumor Virology Lab., Salk Inst., PO Box 18380, San Diego, CA, 92112); Hunter, T.; Eckhart, W. *Cell* 15: 65-77; 1978.

The immunoreactive proteins [tumor (T) antigens] in polyoma (Py)-infected 3T6 cells were characterized. The virus codes for three species having apparent mol wts of 90K, 60K, and 22K, as determined by polyacrylamide gel electrophoresis. The 90K and 22K species produced by a large- and a small-plaque wild-type Py virus have similar mobilities, but the 60K species produced by the small-plaque wild-type virus has a higher mobility than the 60K species produced by the large-plaque form. The relationship of the virus-coded 90K, 60K, and 22K T antigens to each other is of interest in view of the existence of at least two different genetically defined early viral functions. These functions are defined by two classes of mutants, the temperature-sensitive tsA mutants and the host-range nontransforming hr-t mutants. The 90K species is affected by the tsA mutations, since this species is unstable at the nonpermissive temperature in cells infected by seven tsA mutants. The 60K and 22K species are stable under the same conditions. Only the 22K species is present in cells infected by a mutant carrying a deletion between roughly 98 and 3 map units in the early region of the viral genome. Tryptic peptide analysis of the isolated 90K, 60K, and 22K species shows that the three have few peptides in common. These peptides may be derived from a common N terminal region. The 60K and 22K species contain amino acid sequences not found in the 90K species, and the 60K species has several unique methionine-containing

ides not found in the 22K or 90K species. The observation that two BHK cell lines transformed by Py virus do not contain detectable amounts of the 90K T antigen is consistent with previous observations that permissive or semipermissive Poma-transformed cells fail to contain stable cytoplasmic T antigen corresponding to the distal portion of the early region of the Py genome. The tsA function may be involved in regulating the growth properties of some transformed cells, because some cells transformed by a Py tsA mutant show temperature-dependent growth properties. (27 refs)

6497 Specificity and Age of Appearance of Natural Immunity to MTV Antigens. (Eng) Lopez, D. (Lab. Virology, Dept. Microbiology, Univ. Miami, Sch. of Medicine, Miami, FL, 33152); Sigel, M. M.; Ortiz-Muniz, G.; and others. *Proc Soc Exp Biol Med* 158(1): 23-27; 1978.

The natural immune reactivity of BALB/c mice to mammary virus (MTV) antigens was examined by lymphocyte transformation and migration inhibition assays. The specificity of the lymphocyte response to RIII MTV was measured by virus neutralization reactions using anti-MTV and anti-mouse leukemia virus (abtu-RLV) sera. The former induced the blastogenic response to MTV, but the latter had no effect. RLV did not induce transformation in BALB/c lymphocytes. MTV-induced lymphocyte transformation was observed at low levels and in a small percentage of BALB/cRgl mice as early as 2 wk after birth. At 8 wk of age, the responses in the lymphocyte transformation and migration inhibition assays were intermediate in frequency and/or intensity, and by 16 wk, essentially maximal responses were seen. The nature of the antigen responsible for the lymphocyte sensitivity has not been determined, but radioimmunoassays have largely ruled out the MTV p14 antigen. It is hypothesized that lymphocyte sensitivity to MTV antigen originates from the occurrence of a viral antigen produced by the expression of part of the viral genome in the host cytoplasm. (23 refs)

6498 Leukaemia Associated Antigens: Expression by Human Cell Lines. (Eng) Bertoglio, J. (Laboratoire d'Immunologie et de Cancerologie Experimentale, Centre Leon Berard, 28 rue Laennec, 69373 Lyon Cedex 2, France); Dore, J. F. In: *Human Lymphocyte Differentiation: Application to Cancer. Proceedings of the International Symposium on Human Lymphocyte Differentiation: Its Application to Cancer, held in Montpellier (France) by Institut National de la Sante et de la Recherche Medicale and Delegation Generale a la Recherche Scientifique et Technique, 2-4 March, 1978.* Serrou, B., Rosenfeld, C., eds. (Amsterdam: North Holland Publishing Company); 423 pp.; 357-364; 1978.

Thirteen human lymphoid cell lines derived from normal B lymphocytes or from patients with leukemia, lymphoma, or

Burkitt's lymphoma were tested for expression of leukemia-associated antigens using sera from leukemia patients and normal donors cytotoxic to human leukemia cells and relatively specific for acute lymphoblastic leukemia (ALL). None of the lines derived from the normal B lymphocytes or from patients with Burkitt's lymphoma were reactive. Only four lines, derived from patients with ALL, acute myelomonocytic leukemia, the blastic crisis of chronic myeloid leukemia, and lymphosarcoma, respectively, were reactive with one or more of the 10 sera used. The Reh line, derived from a patient with non-T, non-B ALL, was the most reactive and the only one to react with patients' sera. Cells from a fifth line (derived from a patient with ALL) were able to absorb the reactivity of one serum (MOR: from the father of a leukemia patient) to ALL cells, as were all three cell lines that were reactive with serum MOR in the direct test. Thus, permanent cell lines of leukemic origin may be useful in the study of leukemia-associated antigens. (19 refs)

78-6499 Humoral and Cellular Immune Response of Prostatic Cancer Patients to Cytomegalovirus-related Antigens. (Eng) Sanford, E. J. (Div. Urology, Milton S. Hershey Medical Center, Pennsylvania State Univ., Hershey, PA, 17033); Geder, L.; Dagen, J. E.; Laychock, A.; Rohner, T. J.; Rapp, F. *J Surg Res* 24(5): 404-408; 1978.

The association between prostatic adenocarcinoma and cytomegalovirus (CMV) infection was investigated in patients with biopsy proved adenocarcinoma or benign prostatic hypertrophy (BPH). Humoral immune response studies were performed to detect antibodies against human embryonic lung cells (HEL), HEL transformed by CMV (CMV-Mj-HEL-2), and cells from a CMV-Mj-HEL-2-induced tumor in nude mice (CMV-Mj-HEL-2, T-1). Neither the cancer nor BPH patients had antibodies against HEL or CMV-Mj-HEL-2 cells, but 80% of the cancer patients and 68% of the BPH patients had antibodies against CMV-Mj-HEL-2, T-1. In cellular immune response studies, 82% of the cancer and 35% of the BPH patients had lymphocytes cytotoxic to CMV-Mj-HEL-2 cells, 82% and 90%, respectively, had lymphocytes cytotoxic to CMV-Mj-HEL-2, T-1 cells, and 20% of both groups had significant cytotoxicity to HEL cells. Sera from prostatic carcinoma patients significantly blocked the cytotoxicity of lymphocytes from prostatic cancer patients against CMV-Mj-HEL-2, T-1 to a greater degree than it did lymphocytes from nontumor controls. Sera from the nontumor controls exhibited little blocking activity. Hyperimmune rabbit anti-CMV sera did not block the cytotoxicity of cancer patient sera and, in fact, enhanced the cytotoxic effect in 2/3 patients. The results suggest that prostatic carcinoma patients possess a serum blocking factor that may be a CMV-related antigen-antibody complex. (18 refs)

78-6500 Specific Immune Response Against Tumor-associated Antigens of a Syngeneic Simian Virus

40-induced Sarcoma in Mice. (Eng) Glaser, M. (Lab. Cell Biology, Immunology Program, Div. Cancer Biology and Diagnosis, NCI, NIH, Bethesda, MD, 20014). *J Natl Cancer Inst* 61(5): 1351-1355; 1978.

An immune response against syngeneic simian virus 40 (SV40)-induced tumor-associated antigens (TAA) in (BALB/c x C57Bl/6)F₁ female mice was detected by in vivo tumor challenge and tumor cell neutralization tests and by the in vitro ⁵¹Cr-release cytotoxicity assay. All mice immunized with 1 x 10⁵ SV40-transformed cells were protected against the growth of SV40-induced tumor from 3 to 90 days after a single immunization. In the tumor cell neutralization test, spleen cells (T cells) from mice immunized with 1 x 10⁵ transformed cells prevented tumor growth from 10 to 90 days after immunization. Spleen cells from mice immunized with a minimum of 20 x 10⁴ transformed cells were cytotoxic in vitro against the relevant target cells in a 4-hr ⁵¹Cr-release cytotoxicity assay. Cytotoxic lymphocytes were detected 3 days after immunization, reached a peak activity at 8 days, and declined to low levels at 14 days. At all times after immunization, T cells were needed for cytotoxic reactivity to be eliminated by specific anti- θ serum and complement. (21 refs)

78-6501 Stimulatory Effect of Immunization on Tumor Induction by Moloney Murine Sarcoma Virus. (Eng) Murasko, D. M. (Dept. Microbiology, Medical Coll. Pennsylvania and Hosp., 3300 Henry Ave., Philadelphia, PA, 19129); Prehn, R. T. *J Natl Cancer Inst* 61(5): 1323-1327; 1978.

Adult BALB/c mice were immunized with varying doses of inactivated Moloney murine leukemia virus (M-MuLV) and challenged 8 wk later with 0.1 ml Moloney murine sarcoma virus (M-MuSV), at a dose that induces tumors in 50% of normal animals. Sera from all immunized mice demonstrated neutralizing activity 14 days after immunization in the XC assay. The ability of sera to neutralize M-MuLV was directly related to the amount of M-MuLV used for immunization. No neutralizing activity was detected in any serum sample taken 8 wk after immunization. Mice immunized with high doses (10¹⁰ particles) of M-MuLV had significantly decreased tumor incidences, whereas mice immunized with low doses (10² particles) had significantly increased tumor incidences, compared with those in nonimmunized controls. The stimulatory effect could be abrogated by irradiation of the mice with 450 rads 24 hr prior to M-MuSV challenge, but the inhibitory effect was resistant to this treatment. The results suggest that depending on dose, immunization with virus can either stimulate or inhibit virus-induced tumorigenesis. Two possible explanations for the mechanism of increased tumorigenesis are suggested: (1) direct stimulation of target cells by lymphoid cells or lymphoid cell products; and (2) suppression of an inhibitory immune response. (33 refs)

78-6502 Resurgence of Killing and In Vivo Protection Mediated by Lymphocytes Cultured from

Lymph Nodes Draining Moloney Sarcomas. (Eng) Gilles, G. Y. (Dept. Pathology, Univ. North Carolina Sch. Medicine, Chapel Hill, NC, 27514); Hansen, C. B.; Russell, S. *Br J Cancer* 38(3): 365-374; 1978.

The return of cytolytic activity to tumor-draining lymph node cells that were no longer cytotoxic in vivo was investigated. The popliteal and lumbar lymph nodes draining progressing or progressing Moloney sarcomas in male BALB/c AnCr mice were removed from the animals and minced, the lymphocytes were isolated. The lymph node cells were no longer cytotoxic in vivo spontaneously regenerated peak levels of killing after a 4-day in vitro culture in the absence of exogenous tumor antigen. Cytolytic activity which was antigenically specific, was mediated by T cells. Resurgence of cytolytic activity in vitro was accompanied by proliferative changes (DNA synthesis, blast transformation, cell division) that peaked on day 3 of culture. Although normal, nonimmune lymph node cells underwent quantitatively similar proliferative changes in culture, the killing that developed was weak and antigenically nonspecific. Transfer of cultured, tumor-draining lymph node cells to immunologically compromised, syngeneic mice conferred complete protection from Moloney sarcoma progression. Adoptive transfer could be delayed for 6 days after tumor induction without loss of protection. These findings suggest that some mechanism in Moloney sarcoma-bearing mice limits the differentiation of prekiller cells into cytolytically active T cells and that this inhibition is eliminated when the lymph node cells are explanted into culture. (31 refs.)

78-6503 Activation of Complement by Spontaneous Leukemic Cells of AKR Mice. (Eng) Okada, H. (Dept. Bacteriology, Kitasato Inst., Shirogane 5-9-1, Minato-ku, Tokyo 108, Japan); Okada, H. *Int J Cancer* 22(2): 282-287; 1978.

The extent of nonspecific activation of complement (NAC) by AKR mouse spontaneous leukemic and normal cells was compared by the human erythrocyte (E) rosette formation following treatment of cells with C4-deficient guinea pig serum in the presence of Mg²⁺/gelatin veronal-buffered saline/ethyleneglycolbis(β-aminoethyl ether) N,N'-tetraacetate (EGTA) and in the absence of Ca²⁺. Following treatment, 10%-30% of lymphoma cells and 10%-17% of thymoma cells formed rosettes, vs only 3%-4% of thymus and mesenteric lymph node cells from nonleukemic mice. The percentage of NAC-positive cells was essentially the same for the original spontaneous, transplanted (to nonleukemic mice), and cultured lymphoma cells, indicating that NAC ability does not result from antigen-antibody-complement complexes that are formed or fixed in vivo. The results suggest that NAC ability may be increased when normal AKR cells become transformed into leukemic cells and/or that leukemic cells arise from thymus cells that possess NAC ability. This interpretation was confirmed further by the finding that a higher proportion of leukemic cells than that of normal

neocytes were lysed by mouse complement in Mg^{2+} -A. The fact that NAC-positive AKR thymoma cells are lysed more effectively by homologous C3H/He mouse serum than by the guinea pig serum agrees with earlier findings that a temporary regression of leukemia occurs in leukemic AKR mice supplemented with C5. (30 refs)

6504 **Characterization of Fc Receptor in Human Malignant Tumors (Meeting Abstract).** (Eng) Mahan, E. C. (Coll. Health Science and Hosp., Univ. Kansas Medical Center, Kansas City, KS, 66103); Jewell, W. R. *Am Assoc Cancer Res* 19: 96; 1978. (no refs)

6505 **Cell Receptor Studies on Six Anaplastic Tumours of the Thyroid.** (Eng) Macaulay, R. A. (Dept. Forensic Medicine, Univ. Edinburgh, Edinburgh, Scotland); Dewar, A. E.; Langlands, A. O.; Stuart, A. E. *J Clin Pathol* 31(5): 461-468; 1978.

An attempt was made to distinguish malignant lymphoma from other thyroid tumors by examining suspensions of tumor cells for lymphocyte surface markers. The results were related with the ultrastructural appearance of the tumors, clinical extent of the disease, and response to radiotherapy. Cell suspensions were obtained from the anaplastic thyroid tumors of six patients (5 women, 1 man, aged 60-85). In five cases, many of the cells showed surface immuno-

globulin. Electron microscopy studies of three of these tumors were indicative of malignant lymphoma. In 2/5 cases, the patients had a history of previous thyroid disease. The single tumor that did not show surface immunoglobulin had an entirely different ultrastructure and was probably a carcinoma. This patient was the only one that did not achieve complete remission with radiotherapy, and she died from extensive local recurrence. It is concluded that receptor techniques are of value in distinguishing between malignant lymphoma and other anaplastic tumors of the thyroid, particularly when the results are correlated with histology. (23 refs)

See also:

- *(Rev.): 78-6013, 78-6071, 78-6072, 78-6076, 78-6077, 78-6078, 78-6079, 78-6080, 78-6081, 78-6082, 78-6083, 78-6088, 78-6098.
- *(Chem.): 78-6209, 78-6219, 78-6277, 78-6278, 78-6310, 78-6332.
- *(Viral): 78-6400, 78-6402, 78-6406, 78-6427, 78-6430, 78-6432, 78-6435, 78-6442, 78-6443, 78-6444, 78-6446, 78-6449, 78-6452, 78-6459, 78-6460, 78-6468, 78-6473, 78-6479.
- *(Path.): 78-6514.

PATHOGENESIS

- 78-6506 Genetically Determined Sex-Reversal in 46,XY Humans.** (Eng) German, J. (New York Blood Center, 310 E. 67 St., New York, NY, 10021); Simpson, J. L.; Chaganti, R. S.; Summitt, R. L.; Reid, L. B.; Merkatz, I. R. *Science* 202(4363): 53-56; 1978.

Observations of one family that constitute strong evidence for the existence in humans of an X-borne gene that in mutant form results in sex-reversal in 46,XY individuals are reported. They also provide further insight into the genetic control of normal human gonadal differentiation. Affected were two sisters, their maternal aunt, and one first cousin from another maternal aunt. All four developed normally throughout childhood. At puberty, axillary and pubic hair developed normally, but breast development and menstruation had to be induced by appropriate hormones. None of the four displayed Turner's syndrome characteristics. Genitalia were unambiguously female, and uterus and fallopian tubes were normal at prepubertal age. Upon pelvic surgery (between age 19 and 29), however, the sisters were found to have bilateral streak gonads reminiscent of 45X gonadal dysgenesis. The cousins's gonads were grossly streaklike, but were proved to be bilateral gonadoblastomas. Pathology of an abdominal tumor removed from the aunt is unknown, but suspect. Chromosome studies of the three living women (the aunt died postoperatively) ruled strongly against mosaicism. X and Y chromosomes appeared normal in banding studies. The inability of the Y chromosome alone to initiate male differentiation of the embryo was ruled out, because three of the Y chromosomes came from different fathers. Therefore, it was hypothesized that some X-Y interaction is necessary for the initiation of male differentiation and that in these women the necessary X locus was mutant, perhaps transmitted as an X-linked recessive trait. In the absence of male differentiation and in the presence of only one X, the sexual differentiation resulting in streak gonads was like that seen in 45X gonadal dysgenesis. (27 refs)

- 78-6507 Uneven Geographical Distribution of 15;17-Translocation in Acute Promyelocytic Leukaemia (Letter to Editor).** (Eng) Teerenhovi, L. (Third Dept. Medicine, Univ. Helsinki, SF-00290 Helsinki 29, Finland); Borgstrom, G. H.; Mitelman, F.; Brandt, L.; Vuopio, P.; Timonen, T.; Almqvist, A.; de la Chapelle, A. *Lancet* 2(8093): 797; 1978.

In contrast to findings in Belgian and, perhaps, US patients, none of the nine Finnish and three Swedish patients with acute promyelocytic leukemia (APL) examined (comprising all cytogenetically analyzed cases from Finland and southern Sweden) had a 15;17-translocation. Apparently, there is a

striking difference in the geographical distribution of the chromosome anomaly in APL patients. (10 refs)

- 78-6508 Simultaneous Occurrence of Non-Hodgkin Lymphoma and Spontaneous Acute Granulocytic Leukemia.** (Eng) Youness, E. (Dept. Lab. Medicine, M. D. Anderson Hosp. and Tumor Inst., 6723 Bertner Ave. Houston, TX, 77030); Ahearn, M. J.; Drewinko, B. *Am. Clin. Pathol.* 70(3): 415-420; 1978.

The second reported case of simultaneous non-Hodgkin lymphoma and spontaneous acute granulocytic leukemia occurred in a 67-yr-old Caucasian farmer. He had no remarkable family or medical history except for chronic exposure to some form of insecticide. The diagnoses of the two diseases were confirmed by examination of the patient's bone marrow and peripheral blood specimens at the light and electron microscopic level and by autopsy findings. Comparison of the patient with the previously reported patient (a woman) showed that both had no prior history of chemotherapy and/or radiotherapy, they were in their late 60's, they had relatively short histories of anemia and wt loss, and they had simultaneous involvement of lymphopoietic organs and peripheral blood. The woman had generalized lymphadenopathy and blast-containing Auer rods, but the man had lymphoid involvement confined to the liver, spleen, and bone marrow and his blasts did not show Auer rods. Chronic exposure to an unknown insecticide may have played a leukemogenic role in the latter. (32 refs)

- 78-6509 Probable Clonal Origin of Acute Myeloblastic Leukemia Following Radiation and Chemotherapy of Colon Cancer.** (Eng) Wiggans, R. G. (Dept. Medicine, Div. Medical Oncology, Georgetown Univ. Hosp., Vincent T. Lombardi Cancer Res. Center, Georgetown Univ. Washington, DC); Jacobson, R. J.; Fialkow, P. J.; Woolley, P. V.; Macdonald, J. S.; Schein, P. S. *Blood* 52(4): 659-663; 1978.

Acute myeloblastic leukemia (AML) developed in a 64-yr-old black woman after radiation and drug therapy for disseminated colon cancer. The occurrence and then progression of pulmonary metastases prompted two ⁶⁰Co-radiation treatments (3,500 and 4,200 rads) 3 and 1 yr before AML diagnosis respectively. She was also treated with cytotoxic drugs in the 2 yr prior to diagnosis, including intermittent courses of 5-fluorouracil throughout that time and, in the year preceding diagnosis, with 1-(2-chloroethyl)-3-(4-methylcyclohexyl)-1-nitrosourea (4 mo) and streptozotocin (3 wk). Approx

Amounts of B and A types of glucose-5-phosphate dehydrogenase were found in skin and cultured skin fibroblasts, only type B was detected in the abnormal WBC. In chromosome studies, cultured cells from normal tissues were normal, but all 90 unstimulated blood cells studied were missing approx two-thirds of the short arms of chromosome 12 (X,12p-). This abnormality was also detected in 18/24 unstimulated marrow cells. These data are consistent with a clonal origin of this leukemia. (23 refs)

78-6510 Multifocal Histogenesis of a Cystosarcoma Phyllodes. (Eng.) Salm, R. (Dept. Histopathology, Royal Postgraduate Medical Sch., Hammersmith Hosp., Duane Road, London W12 0HS, England). *J Clin Pathol* 31(9): 897-903; 1978.

An unusual multifocal cystosarcoma phyllodes (CSP) occurred in a 50-yr-old woman. The CSP presented as a small, well-circumscribed tumor mass, and it was associated with two separate neoplastic lesions that arose sequentially in two separate parts of the mammary disk. Microscopically, the tumor appeared to have been enlarging by the formation of isolated minute tumor nodules within the adjacent normal breast tissue. This represents a third mechanism by which CSP may arise, the other and more common modes being expansion in situ and extension of tumor projections from the periphery. (6 refs)

78-6511 Soft Tissue Sarcomas in the Cat. (Eng.) Brown, N. O. (Donaldson-Atwood Cancer Clinic, Animal Medical Center, 510 E. 62nd St., New York, NY, 10021); Malik, A. K.; Mooney, S.; Hayes, A.; Harvey, H. J.; Ewen, E. G. *J Am Vet Med Assoc* 173(6): 744-749; 1978.

A retrospective survey was made of the soft tissue sarcomas found in 23 cats at an animal medical center from 1971 to 1976. The nature of the tumors was variable. All tumors tended to develop in older cats and in various locations, although most were on the extremities. Diagnosis was made upon biopsy. No feline leukemia virus was detected in the 13 cats tested. (10 refs)

78-6512 Connection Between Accidental Trauma and Course of Malignant Melanoma of the Skin. (Ger.) Wozniak, K. D. (Universitäts-Hautklinik, Ernst-Reuter-Strasse 5/8, DDR-402 Halle [S], E. Germany). *Dermatosen Beruf Umwelt* 26(5): 166-168; 1978.

The use of malignant melanoma that may have been accelerated into rapid metastasis by trauma is described. A 38-yr-old man who had a lentil-sized nevus above his left knee for as long as he could remember presented with painful lymph node swelling in the left groin. He reported that the

nevus had been growing for 6 mo. At presentation it was cherry-sized, half-spherical, gray-black, tender, and eroded. After this tumor was removed it was identified as a melanocytoblastoma, partly round-cell and partly spindle cell in composition. The inguinal lymph nodes were also removed. They had the same histology as the tumor. The patient was wounded in the area of the nevus 4 mo before presentation; the wound caused erosion of the nevus. An inflammatory reaction resulted and lymphangitis followed soon after. The patient was symptom-free for 3 mo after treatment, but then neurological symptoms appeared. In spite of intensive therapy, the patient died of heart failure. At autopsy, generalized metastases of the melanoma were found in the iliac artery and aorta, liver, and spleen. Although trauma is no longer considered a cause of melanoma, trauma to a nevus that has begun to undergo malignant transformation may accelerate the course of the disease, as suggested by this case. (24 refs)

78-6513 Malignant Lymphoproliferative Disorders in Chronic Liver Disease. Report of Four Cases and Review of the Literature. (Eng.) Naparstek, J. (Dept. Medicine, Hadassah Univ. Hosp., Jerusalem, Israel); Elia-kim, M. *Am J Dig Dis* 23(10): 887-892; 1978.

The case reports of four patients (3 men, 1 woman 54-61 yr old) with chronic liver disease who later developed lymphoma are given, and the related literature is reviewed. Three of the patients had nonalcoholic cirrhosis and one had chronic cholangitis. Non-Hodgkin's lymphoma developed long after the presence of liver disease had been established in three patients; the fourth died of hepatic coma, and Hodgkin's disease was discovered incidentally at autopsy. No drugs could be implicated in the appearance of the liver disease or the lymphoma. The findings of these four cases within a relatively short period suggests that the association between the two conditions is not incidental. A review of the literature yielded 40 additional cases. In many, lymphoma complicated long chronic liver disease; in others, however, liver disease was secondary to hepatitis B infection or to treatment of lymphoma. Among the several mechanisms that may be operative in the pathogenesis of lymphoma in patients with liver cirrhosis, chronic antigenic stimulation, the presence of hepatitis B surface antigen, and immunosuppressive therapy seem to be the most important. (44 refs)

78-6514 An Unusual Case of Bence-Jones Myeloma with Unreleased γ Heavy Chains in the Cytoplasm of the Myeloma Cells: A Close Relationship Between Bence-Jones Myeloma and Nonsecretory Myeloma. (Jpn) Kanoh, T. (First Div., Dept Internal Medicine, Faculty of Medicine, Kyoto Univ., Kyoto, Japan); Dohmae, N.; Uchino, H.; Sugimoto, C.; Shimada, T. *Acta Haematol Jpn* 41(5): 938-944; 1978.

A case of Bence Jones (K) partially nonsecretory myeloma with unreleased γ heavy chains in the cytoplasm of the myeloma cells is presented, and the close relationship between Bence Jones myeloma and nonsecretory myeloma is discussed. Bence Jones myeloma arises from one of the following mechanisms, most of which are common to nonsecretory myeloma: (1) loss of the capacity to synthesize heavy chains (Bence Jones myeloma without heavy-chain synthesis); (2) loss of the capacity to secrete whole molecules of immunoglobulins, with secretion of light chains only (Bence Jones myeloma with heavy-chain synthesis); and (3) secretion of immunoglobulins with an abnormal heavy or light chain and their rapid extracellular degradation (Bence Jones myeloma without detectable immunoglobulin synthesis). The pathogenesis of the present case may be due to mechanism 2. The characteristic picture of myeloma cells filled with numerous Russell bodies in the present case might be related to the presence of unreleased γ heavy chains in the cytoplasm. (19 refs)

78-6515 Radium-Induced Cancer: Excessive Radium Dosage (Meeting Abstract). (Eng) Applebaum, E. L. (Glencoe, IL). *Otolaryngology* 86(4): 188-189; 1978. (no refs)

78-6516 Tumor and Tumor-like Lesions of Perilimbal Conjunctiva in Laboratory Dogs. (Eng) Hargis, A. M. (Collaborative Radiological Health Lab., Dept. Pathology, Colorado State Univ., Fort Collins, CO, 80523); Lee, A. C.; Thomassen, R. W. *J Am Vet Med Assoc* 173(9): 1185-1190; 1978.

Among 1,680 research colony beagles, proliferative and neoplastic lesions developed in the nasal or temporal limbal conjunctiva of 8 male and 6 female dogs. The mean age at observation was 5.2 yr. Acanthosis and squamous cell carcinoma of the conjunctival epithelium to vascular ectasia, hemangioma, and invasive hemangiosarcoma of the underlying lamina propria were seen. The dogs had been kept in a sunny, relatively high-altitude (1,500 m), smog-free area, suggesting that exposure to solar radiation may have been involved in tumor development. (19 refs)

78-6517 Cerebellar Medulloblastoma: An Analysis of Four Cases of Extraneural Metastasis. (Eng) Schnitzler, E. R. (Dept. Neurology, Univ. Wisconsin, 1300 University Ave., Madison, WI, 53706); Richards, M. J.; Chun, R. W. *Am J Dis Child* 132(10): 1004-1008; 1978.

The cases of 4/28 medulloblastoma patients who developed extraneural metastases are reviewed, and their clinical features are compared with those of the 24 patients without metastases. Suboccipital craniectomy, radiation therapy, and

patient longevity were not well-correlated with metastasis. However, subarachnoid tumor spread was a consistent cursor (4/4 patients with metastasis vs 7/24 without). A triculoatrial shunt may have been a factor in one patient. A serious infection that preceded diffuse metastases may have been a factor in another. Two of the four patients with metastases had the desmoplastic variety of medulloblastoma, only 1/24 without metastases had this variant. Identification of the medulloblastoma patient with an increased risk of extraneural metastasis is difficult but it has practical implications, since recent studies indicate that this metastasis is sensitive to chemotherapy. (21 refs)

78-6518 Linkage Study in a Kindred with Multiple Endocrine Neoplasia (MEN 41) (Meeting Abstract). (Eng) Forster, C. J. (Queen's Univ., Toronto, Ontario, Canada); Sears, E.; Ghent, W.; Falk, J.; Goodall Partington, M. W.; Simpson, N. E. *Can J Genet Cytol* 20: 443; 1978. (no refs)

78-6519 Squamous Cell Carcinoma Arising in a Meningiomyelocoele. (Eng) Saksun, J. M. (Div. I Pathology, Dept. Medicine, Wellesley Hosp., 160 Wellesley St., E., Toronto, Ontario M4Y 1J3, Canada); Fisher, B. *Can Med Assoc J* 119(7): 739-741; 1978.

A squamous cell carcinoma arose in a lumbosacral meningiomyelocoele of a 25-yr-old man. Only two similar cases were found in the literature. Neoplastic change in untreated meningiomyelocoeles is possible after years of mechanical irritation due to braces and chronic bacterial infection. (28 refs)

78-6520 Pathogenesis of Attic Cholesteatomas. (Eng) Sade, J. (Meir General Hosp., Kfar Saba, Israel). *J R Soc Med* 71(10): 716-732; 1978.

To determine the mode of genesis of attic cholesteatomas (AC), 22 of these tumors were investigated in situ and in toto. They were all epidermoid cysts of elaborate cellular organization, 18 of which were situated medial to the ossicular chain. This site makes self-cleansing difficult, producing secondary infection. Three AC had no connection with the external ear canal. However, whether or not AC communicate with the external canal, they have a lining that often has mixed epithelium: stratified squamous epithelium (EC) synthesizing keratin and respiratory epithelium producing mucus. This shows that a metaplastic process is involved, at least to some degree. Therefore, AC are epidermoid cysts that originate primarily in the attic from epithelial cells that metaplastic mature into keratinizing epithelium. They often present themselves secondarily in the external canal as a perforated epidermoid cyst. The favored etiology of an invasive retraction from the external ear is difficult to accept in view of the

tact inhibition of the SE or the invariable outward migration of stratified SE from the edges of retraction pockets as well as from cholesteatoma perforations. Also, there is no documentation of an evolution of AC from preexisting perforations or from marginal perforations. (47 refs)

78-6521 Intranasal Tumor of the Ethmoid Olfactory Mucosa in Sheep. (Eng) Yonemichi, H. Takikawa Animal Husbandry Experimental Station, Hokkaido, Takikawa 073, Japan; Ohgi, T.; Fujimoto, Y.; Okada, T.; Onuma, M.; Mikami, T. *Am J Vet Res* 39(10): 1599-1606; 1978.

Intranasal tumors (papillary adenomas or adenocarcinomas) of the ethmoid olfactory mucosa of sheep were investigated by light and electron microscopy. The fine structure of the tumor cells was characterized by the presence of numerous secretory granules. Viral particles that were morphologically similar to a visna-maedi virus were detected in all tumor tissues and in 3/4 cultures examined. The particles [93-97 nanometers (nm)] had an eccentrically located electron-dense core (47 nm), often surrounded by a single (intermediate) membrane; these particles also had numerous spikes (9 nm) on their surfaces. There were no viral particles in the lung of sheep with pneumonia or in the normal olfactory mucosa of two control sheep. RNA-dependent DNA polymerase (DDP) activities in the tumor tissues or in cultured tumor cells were greater than those in normal intranasal tissues or cultured cells from the choroid plexus. Viral particles similar to herpesvirus were also detected in one culture, and particles of a paravaccinia type were found in another. (28 refs)

78-6522 Familial Medullary Thyroid Carcinoma. (Eng) Hillyard, C. J. (Dept. Surgery, Royal Postgraduate Medical School, Hammersmith Hosp., Duane Road, London W12 0HS, England); Evans, I. M.; Hill, P. A.; Taylor, J. *Lancet* 1(8072): 1009-1011; 1978.

Twenty-one relatives of a 14-yr-old girl with medullary thyroid carcinoma (MCT) were screened by the plasma calcitonin assay before and after alcohol provocation. The patient's mother and two siblings had raised basal levels, and the diagnosis of MCT was confirmed at surgery; their calcitonin levels became undetectable after total thyroidectomy. Only 1/3 had a palpable thyroid mass. Three more distant relatives had detectable calcitonin levels after alcohol stimulation, and they are being rescreened at regular intervals. Cervical lymph node metastases were found in the index patient 1 mo after total thyroidectomy. Her current calcitonin level is now 36.8 pg/liter. (9 refs)

78-6523 Total Thyroidectomy: The Indications and Results of 630 Cases. (Eng) Katz, A. D. (8733 Beverly

Blvd., Los Angeles, CA, 90048); Bronson, D. *Am J Surg* 136(4): 450-454; 1978.

Indications for total thyroidectomy are illustrated in a series of 630 patients (499 females, 131 males aged 7-83 yr) in whom this operation was performed during 1958-1978 in Los Angeles County. The patients were divided into three groups: (I) 121 with Grave's disease; (II) 133 with a history of childhood head, neck, and thorax irradiation and a cold thyroid nodule; and (III) 396 with a dominant cold thyroid nodule suspected of being a carcinoma. The total cancer incidence was 193/630 patients: 183 had papillary follicular carcinoma, 4 medullary carcinoma, 5 anaplastic or small cell carcinoma and 1 reticulum cell sarcoma. Group I patients had a 5% malignancy rate, Group II a 42.5% rate, and Group III a 35% rate. Concerning Group II patients, available data indicate that at least 2% of individuals whose thyroids were exposed to 300 rads of x-rays will develop thyroid cancer within 25 yr and that 9% will have benign nodules. Irradiation of thymolymphatic tissue in the head, neck, and thorax might result in changes in lymphoid elements that subsequently reach the thyroid gland. Lymphoid elements are far more vulnerable to radiation-induced changes than thyroidal epithelium. In most of the 113 Group II patients, both lobes of the thyroid were grossly and microscopically abnormal (follicular and Hurthle cell adenomas, multicentric carcinomas, and thyroiditis). (25 refs)

78-6524 Synovial Sarcoma of the Neck. (Ita) Berton, F. (Servizio di anatomia ed istologia patologica, Ospedale Marcello Malpighi di Bologna, Bologna, Italy); Bacchini, P. *Pathologica* 70(1005/1006): 409-418; 1978.

The case of a 42-yr-old woman with synovial sarcoma of the neck is reported. The patient had symptoms of pain in the neck region, swelling, and hoarseness. Examination of the larynx revealed a polypous neoplasm on the posterior and lateral walls of the pharynx. A biopsy was performed, and a diagnosis of synovial sarcoma was made. The patient underwent a partial pharyngectomy and a total laryngectomy. No evidence of metastases to the cervical lymph nodes was found. Microscopically, fused cells were seen with voluminous nuclei and a granular-eosinophilic cytoplasm. The nuclei were polydimensional, with indented and jagged membranes. Histologically, the neoplasm was composed of two components: epitheliallike and mesenchymal. The epithelial-like components (PAS- and Alcian-positive) formed pseudoglandular formations. The mesenchymal component was pressed between these formations and was rarely PAS- or Alcian-positive. The histological aspects confirmed the diagnosis. (15 refs)

78-6525 Mammary Gland Hyperplastic Alveolar Nodulogenesis in the LEW/Mai Rat in Relation to Mitotic Activity, Estrous Cycle Status, and Age. (Eng)

Purnell, D. M. (Dept. Pathology, Sch. Medicine, Univ. Maryland, 660 W. Redwood St., Baltimore, MD, 21201); Stowers, D. J. *J Natl Cancer Inst* 61(5): 1339-1346; 1978.

The genesis of spontaneous mammary hyperplastic alveolar nodules (HAN) was studied in nulliparous female LEW/Mai rats in relation to age, estrous cycle status, and mammary gland mitotic activity. Mitotic activity in the contralateral mammary gland rose almost linearly from 30 to 108 days of age and then fell progressively, although it remained significantly higher than the activity in mammary glands of rats < 100 days old. Mitotic activity was nonrandomly distributed in the mammary glands of young pubertal rats in contrast to that in glands of prepubertal rats and in rats in persistent estrus. The percentage of rats with HAN as well as the incidence of HAN in the inguinal mammary gland increased progressively with age. At 150 days of age, 12.5% of the rats had 1-5 HAN/inguinal mammary gland; by 183 days, 18.5% had 1-19 HAN/gland; and by 215 days, 91.7% had 3-44 HAN/gland. HAN were more common near the nipple than in other regions of the mammary gland. They appeared only in rats in persistent vaginal estrus, and they were more frequent in rats that displayed this condition for the longest period of time. Rats with HAN also had significantly increased pituitary wts. These findings suggest that hormone imbalance may be an important etiologic factor in the genesis of spontaneous HAN in rats. (20 refs)

78-6526 Phagocytosis of Tumor Cells by Murine Kupffer Cells In Vivo and In the Perfused Mouse Liver. (Eng) Roos, E. (Div. Cell Biology, Antoni van Leeuwenhoekhuis, Netherlands Cancer Inst., Sarphatistraat 108, Amsterdam, Netherlands); Dingemans, K. P. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 216-224; 1977.

The behavior of Kupffer cells toward several tumor cell lines (mammary adenocarcinomas TA3/Ha and TA3/St, melanoma B16, lymphosarcoma MB VI A, and thymic leukemias GRSL-18 and GRSL-19) was studied after in vivo injection of the tumor cells in the portal system of mice or addition of these cells to a synthetic medium that perfused an isolated mouse liver. Within 30 min after their arrival into the liver, TA3 and B16 cells were partly or completely surrounded by thin, electron-lucent cell processes of Kupffer cells. Micrographs suggest that the tumor cells were surrounded by a thin rim of Kupffer cell cytoplasm that subsequently contracted to enclose part or all of the tumor cell. A dense material, probably containing lysosomal enzymes, was then released around the enclosed tumor cell and it was subsequently degenerated. These observations strongly suggest that the cells were phagocytized. For the TA3 and B16 cells, there was no difference in the percentage of partially or completely surrounded tumor cells after 30, 60, 90 or 120 min. The average percentage of the cell surface covered by the Kupffer cells

was similar in all experiments with one tumor cell line also did not tend to increase with time. MB VI A and GRSL cells were little affected by the Kupffer cells. The data from the perfusion and in vivo experiments were similar (refs)

78-6527 Effect of Parity on Recovery of Inapparent Nodule-transformed Mammary Gland Cells In Vivo. (Eng) DeOme, K. B. (Dept. Zoology, Univ. California Berkeley, CA, 94720); Miyamoto, M. J.; Osborn, R. C.; Gorman, R. C.; Lum, K. *Cancer Res* 38(11, part 2): 4050-4054; 1978.

Inapparent nodule-transformed cells were recovered from late-pregnant BALB/cfC3H mice (5 first-pregnancy females 4 mo old, 5 multiparous females 6-7 mo old) by a cell-coculture technique, aliquots of 100,000 viable cells in 0.1 ml media were injected into the gland-free mammary fat pad of 3-wk-old syngeneic hosts, and the emergence of typical nodule outgrowths (LA's) was determined. Recovery of LA's indicated the presence of nodule-transformed cells in the inoculum. The results were compared with those derived previously from virgin BALB/cfC3H mice. The percentage of LA-type outgrowths (LS's, nodule-ductal outgrowths, and nodule-tumor outgrowths) derived from both donor groups was high: 63% and 45% for those aged 4 and 6-7 mo, respectively. There was no correlation between the percentage of LA's and donor age or number of litters born. Virgin mice of the same ages averaged 11% LA's. LA's were recovered from all 10 of the parous donors, vs only 8/12 of the virgin donors. Hyperplastic alveolar nodules (HAN) and tumors were not encountered in virgins < 8-9 mo old, whereas 36% of the parous females 4-5 mo old had overt HAN and 3.6% had tumors, and 73% of the parous females 8-9 mo old had overt HAN and 73% had tumors. The incidence of overt lesions was only 40% in virgins aged 12-14 mo. The data suggest that parity significantly increases the number of nodule-transformed cells in donor tissue, decreases the time required for the emergence of nodules and tumors, and increases the number of overt nodules and tumors. (7 refs)

78-6528 The Respiratory Epithelium. III. Histogenesis of Epidermoid Metaplasia and Carcinoma In Situ in the Human. (Eng) Trump, B. F. (Dept. Pathology, Univ. Maryland Sch. Medicine, 31 S. Greene St., Baltimore, MD, 21201); McDowell, E. M.; Glavin, F.; Barrett, L. A.; Becci, P. J.; Schurch, W.; Kaiser, H. E.; Harris, C. C. *J Natl Cancer Inst* 61(2): 563-575; 1978.

In an effort to test the hypothesis that epidermoid metaplasia (EMP) in the bronchial epithelium can result from direct conversion of columnar mucous cells to keratinizing epidermoid cells, bronchial specimens of 100 patients with and without lung cancer were studied by light and electron microscopy. Of the 71 lung cancer patients, 15 showed focal

P. The metaplastic epithelium was characteristically composed of basal cells overlaid by small mucous granule cells (SMGC) that became progressively flattened toward the luminal surface. Ciliated cells were absent. EMP showing differentiation was seen in only two specimens, both from cancer patients. Epithelium showing mucous goblet cell hyperplasia and ciliated epithelium underwent abrupt transition to that of noncornifying metaplastic epithelium. Mucous goblet cell hyperplasia was seen in more than half of the specimens; a pattern of stratification was noted in some areas in one-third of these hyperplasia specimens. Layers of SMGC overlaid a row of basal cells, and the surface layer consisted of cuboidal mucous goblet cells and/or SMGC. Staining of metaplastic cells produced patterns similar to those of mucous cells in mucous goblet cells. Electron microscopy further confirmed the mucous nature of these cell types. This study emphasizes the differentiation potentialities of mucous cells that can divide and undergo goblet cell hyperplasia and EMP. It is suggested that EMP and carcinoma in situ can result from the conversion of mucous cells. (73 refs)

6529 **Carcinoma of the Lung in a Drywall Taping Worker. Report of a Case.** (Eng) Fischbein, A. (Environmental Sciences Lab., Dept. Community Medicine, Mount Sinai Sch. Medicine, City Univ. New York, Fifth Ave. and 100th St., New York, NY, 10029); Langer, A. M.; Suzuki, Y.; Selikoff, I. J. *Toxicol Lett* 2(4): 231-236; 1978.

Case of lung cancer in a 57-yr-old man who had been a heavy smoker and a drywall taping worker for 25 yr is reported. Histologically, the tumor was a well-differentiated squamous cell carcinoma. Congestion of the pulmonary parenchyma, edema, bronchopneumonia, and pulmonary fibrosis were also present. Numerous particles--sheet silicates, diatom fragments, chrysotile asbestos fibrils, and unidentified crystalline fibers--were observed throughout the lung tissue. The size of these particles was $< 1 \mu\text{m}$ in greatest dimension. This case illustrates the importance of tissue analysis in the evaluation of dust disease of the lung. (7 refs)

6530 **Giant Cell Transformation: Cerebrohepatorenal Syndrome.** (Eng) Carlson, B. R. (Dept. Pathology, Children's Medical Center, 1935 Amelia, Dallas, TX, 75235); Weinberg, A. G. *Arch Pathol Lab Med* 102(11): 596-599; 1978.

A male infant with cerebrohepatorenal syndrome of Zellweger was found, upon liver biopsy performed at 6.5 wk of age, to have extensive hepatic giant cell transformation. In addition to features atypical of the early phase of the syndrome there was widespread enlargement and multinucleation of hepatocytes. Upon autopsy at 16 wk, the liver showed early cirrhosis and rare giant cells. Zellweger's syndrome may represent another form of hepatic insult that may induce giant cell transformation. (19 refs)

78-6531 **Hepatoma Development During Long Term Follow-up Period of Liver Cirrhosis.** (Eng) Kobayashi, K. (First Dept. Internal Medicine, Sch. Medicine, Kanazawa Univ., Kanazawa, Japan); Kumagai, M.; Kameda, S.; Sugimoto, T.; Suzuki, K.; Nishimura, K.; Kato, Y.; Sugioka, G.; Hattori, N. *Acta Hepatogastroenterol (Stuttg)* 25(5): 344-349; 1978.

The relationship of hepatoma to liver cirrhosis was investigated, based on long-term (> 3 yr) follow-up studies of 20 Japanese patients with liver cirrhosis. Hepatoma was found at autopsy in 10/20 patients, all of whom had mixed-type cirrhosis. In the nonhepatoma group, two had micronodular cirrhosis and eight had the mixed type. The mean follow-up in the hepatoma and nonhepatoma patients was 5.5 and 5.0 yr, respectively. There were no heavy drinkers among the hepatoma patients, vs three in the nonhepatoma group. Blood samples were analyzed for hepatitis B surface antigen (HBsAg) and α -fetoprotein (FP) by radioimmunoassay, and localization and distribution patterns of HBsAg in the liver were determined by direct immunofluorescence and orcein staining. Six of the eight hepatoma patients tested were positive for HBsAg, vs 2/10 nonhepatoma patients. FP levels $> 20 \mu\text{g/ml}$ were detected in 6/8 hepatoma patients tested, and in four of these patients they were $> 300 \mu\text{g/ml}$. These results indicate that HBsAg-positive liver cirrhosis of the mixed type is closely related to hepatoma development in Japan. (12 refs)

78-6532 **Ultrastructural Features of Renal Carcinoma Cells.** (Rus) Dikshtein, E. A. (Dept. Pathological Anatomy, M. Gor'kii Donetsk Medical Inst., Donetsk, USSR); Vlodavskii, E. A.; Shevchenko, N. I.; Miima, J. G.; Merezko, V. A. *Arkiv Patol* 40(8): 10-16; 1978.

The ultrastructure of three renal cell carcinomas was studied. Two basic cell types were found. Clear cells, which predominated in these carcinomas, were characterized by a cytoplasm rich in lipid vacuoles and glycogen granules. Dark cells, which were scarce, contained small amounts of fat and glycogen but had well-developed mitochondria and other organelles. Both types of cells contained microvilli on the cell surface, desmosomes and interdigitations in the region of the cell contacts, and channels between the tumor cells. These features suggest that clear and dark cells have a common origin in the epithelium of the proximal tubules. (6 refs)

78-6533 **Mixed Endocrine Gastric Tumors Associated with Hypergastrinemia of Antral Origin.** (Eng) Larsson, L. I. (Inst. Medical Biochemistry, Univ. Aarhus, DK-8000 Aarhus, Denmark); Rehfeld, J. F.; Stockbrugger, R.; Blohme, G.; Schoon, I. M.; Lundqvist, G.; Kindblom, L. G.; Save-Soderberg, J.; Grimelius, L.; Olbe, L. *Am J Pathol* 93(1): 53-68; 1978.

Hypergastrinemia of antral origin associated with multiple mixed endocrine tumors of the gastric corpus mucosa occurred in a 57-yr-old diabetic woman. A screening test revealed serum gastrin levels of 4,000-5,000 picograms (pg)/ml (normal levels are < 100 pg/ml), and a pentagastrin test revealed achlorhydria. Upon gastroscopy, approx 20 polypous lesions were detected in the corpus mucosa. Following gastrectomy, serum gastrin levels decreased to < 10 pg/ml. Most (70%-80%) of the tumor cells were of the endocrine cell (so-called ECL cell) type, but enterochromaffin cells of various subtypes and agranular cells were also found. The tumors were locally invasive into the walls of submucosal blood vessels. The surrounding mucosa showed severe atrophic gastritis with intestinalization and contained numerous goblet cells, enterochromaffin cells, and cholecystokinin cells. The latter do not occur in normal oxyntic mucosa. Their presence in intestinalized gastric epithelium suggests that intestinalization also is associated with changes in endocrine cell populations. Gastrin exerts a trophic action on ECL cells. It is speculated that the primary lesion was achlorhydria; the subsequently raised serum gastrin levels, caused by decreased acid feedback inhibition, may have been instrumental in bringing about the hyperplasia and neoplasia of the ECL cells. (30 refs)

- 78-6534 **Adenocarcinoma of the Stomach with Common Variable Immunodeficiency Syndrome.** (Eng) Battle, W. M. (Section Gastroenterology, Dept. Medicine, Univ. Pennsylvania Hosp. and Sch. Medicine, 3600 Spruce St., Philadelphia, PA, 19104); Brooks, F. P. *Arch Intern Med* 138(11): 1682-1684; 1978.

Gastric adenocarcinomas developed as a late complication of the common variable immunodeficiency (CVI) syndrome in a man and a woman who were followed since age 27 and 30 yr, respectively. The syndrome was diagnosed on the basis of onset of disease in adolescence, no family history of increased susceptibility to infection, and little or no circulating IgG, IgA, or IgM. Atrophic gastritis and pernicious anemia were present in both patients. The anemia was characteristic of the CVI syndrome, with a defect in absorption of vitamin B₁₂ and a lack of correction of the Shilling test after administration of intrinsic factor. In both cases, serum gastrin levels remained normal. Adenocarcinoma of the stomach was the most common epithelial tumor in a previous series of malignancies associated with the CVI syndrome. (18 refs)

- 78-6535 **Relationship Between Chronic Schistosomiasis Japonica and Carcinoma of the Large Intestine: Studies on 12 Cases of Adenocarcinoma of Large Intestine Associated with Eggs of *Schistosoma japonicum*.** (Jpn) Inoguchi, T. (First Dept. Surgery, Kurume Univ. Sch. Medicine, Kurume, Japan); Adachi, T.; Yamauchi, Y.; Isomoto, H.; Takamori, K.; Shinohara, M. *Igaku Kenkyu* 48(2): 93-100; 1978.

The relationship between chronic infection by *Schistosoma japonicum* and carcinoma of the large intestine was investigated. In 12 of 381 patients resected for colonic cancer/cancer/parasite association was demonstrated histologically mainly in the rectum and colon, with the stromal reaction in the mucosal or submucosal layer. However, a causal relationship could not be proven. (42 refs)

- 78-6536 **Histogenesis of Malignant Transformation of the Colonic Epithelium. A Detailed Study of Semi-thin Sections.** (Eng) Mughal, S. (Dept. Histopathology, Westminster Medical Sch., London, England); Filipe, M. *Acta Gastroenterol Belg* 41(5/6): 215-225; 1978.

The histogenesis of colonic cancer was studied using semithin (1 micron) resin sections of normal mucosa from 6 patients without gastrointestinal disease, adenomatous polyps and/or carcinomas from 11 patients with familial polyposis coli (FPC), polyps from 8 patients without FPC, and 5 adenocarcinomas. For light microscopy, the sections were stained with toluidine blue (TB), periodic acid-silver methenamine (PASM), PAS, high iron diamine (HID), and alcian blue (AB). The semithin sections revealed the gradations of epithelial atypia from normal mucosa through adenoma to carcinoma to the same degree as did electron microscopy (EM). Goblet cell increases associated with changes in mucin secretory products were shown by PASM. Vesiculated and electron dense bodies were visible with PASM and PAS staining and undifferentiated cells were easily identified with TB and PASM. AB and HID highlighted variations in mucous secretions in apparently normal mucosa adjacent to neoplastic growths. Other features indicative of epithelial dysplasia were alterations in the basement cell membrane and microvilli and nuclear atypia. In FPC, the changes were more severe in the large polyps of the left colon and also in the polyps of the carcinoma patients. The advantages of semithin sections over EM and routine histology in detecting early cancer are discussed. (12 refs)

- 78-6537 **Development of Cancer in Ulcerative Colitis.** (Swe) Fredlund, P. (Kirurgiska kliniken, Lasarettet, S-221 85 Lund, Sweden); Hoevels, J.; Petersson, B. G. Schain, M. *Lakartidningen* 75(40): 3539-3542; 1978.

Four patients (two men, two women, aged 29-48 yr) with ulcerative colitis developed colonic tumors. At surgery, two of the tumors were diagnosed as carcinomas and two as adenocarcinomas. Three patients also had peritoneal carcinomatosis. The interval between the diagnosis of ulcerative colitis and that of the colonic tumor was 12-20 yr in three cases, and approx 1 yr in the fourth. The incidence of colorectal cancer in ulcerative colitis is highest in the distal segment of the intestine. These cases confirm earlier observations on the increased incidence of colonic cancer in ulcerative colitis patients. (18 refs)

6538 Heritable Colon Cancer and Solitary Adenomatous Polyps. (Eng) Lynch, P. (Dept. Preventive Medicine, Creighton Univ., Omaha, NB); Lynch H.; Harris, J.; Guirgis, H. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1573-1589; 1978.

Study was made of a large cancer-prone kindred characterized by a wide variation in age of onset of colon cancer and either a total lack or sporadic distribution of adenomatous polyps in family members who developed colon cancer. Of approx 230 members in this kindred, 32 members have reportedly developed colonic and other gastrointestinal cancers. A striking trait expressed by this kindred is the inordinately high number of primary carcinomas occurring at the right colon. Among the 24 patients whose tumor sites have been documented, the cecum was involved in 16 cases, the ascending colon in 2, and the transverse colon in 4; only 2/24 had cancer at only the descending or rectosigmoid colon. The incidence of patients affected at the right colon is significantly greater than the highest reported figure for the general population. Solitary adenomatous polyps were found in 13 colon cancer patients, but 7 had no polyps. The prevalence of synchronous and metachronous colon cancers was approx 50%, with additional primary cancers arising after segmental colectomy. The lack of an association between polyps and cancer in this kindred brings into question the role of adenomatous polyps in malignant transformation. (19 refs)

6539 Histogenesis of Cloacogenic Carcinoma: Fine Structure of Anal Transitional Epithelium and Cloacogenic Carcinoma. (Eng) Gillespie, J. J. (Dept. Anatomic Pathology, Indiana Univ. Sch. Medicine, 1100 W. Michigan St., Indianapolis, IN, 46202); MacKay, B. *Hum Pathol* 9(5): 579-587; 1978.

The fine structural morphology of the anal transitional zone is described, and the ultrastructure of the two principal variants of cloacogenic carcinoma is presented. The descriptions are based on histologic and electron microscopic evaluation of eight cases of cloacogenic carcinoma (6 women, 2 men, 40-70 yr old) and tissue from four normal anorectal junctions. Anal transitional epithelium is not highly specialized and incorporates features of both urothelium and squamous epithelium; slight urothelial differentiation is probably vestigial. The transitional variant of cloacogenic carcinoma contained clusters of small vesicles and Golgi complexes, indicating partial differentiation. Histologically, the transitional variants resembled urothelial carcinoma but could be distinguished from it by numerous, well-formed desmosomes, frequent vil-

lous projections, and a distinctly elongated form. The pleomorphic and basaloid variants differed from the transitional type chiefly in their relative lack of differentiation. The basaloid tumors resembled cutaneous basal cell carcinoma morphologically and probably represent undifferentiated basal cells originating within the basal layer of anal transitional epithelium. The cloacogenic basaloid tumor is highly metastatic. The observation of features of both urothelial and squamous structure in the transitional variant confirms that it is derived from the anal transitional zone, which also contains features of both structures. (21 refs)

78-6540 Ovarian Fibromata in Four Successive Generations. (Eng) Dumont-Herskowitz, R. A. (New England Medical Center Hosp., 171 Harrison Ave., Boston, MA, 02111); Safaii, H. S.; Senior, B. *J Pediatr* 93(4): 621-624; 1978.

Brief case reports are given for six ovarian fibroma patients from four generations of the same family. In two patients (21 and 42 yr old), the tumors were unsuspected until hysterectomies had been done because of uterine fibroids. In the remaining four patients (3-16 yr old), the tumors were large, multinodular, and calcified and were discovered before menarche. The tumors were bilateral in three subjects. No endocrine effects could be ascribed to any of the tumors. The distribution of ovarian fibromas in this family was very different from that usually observed. The tumors typically occur in an older age group and after menopause. They are multiple or bilateral in only 10% of the patients and calcified in only 8.5%. Many and possibly all female subjects, including adolescents, who suffer from the basal cell nevus syndrome have ovarian fibromas that are characteristically multinodular and calcified, but none of the patients in this family had any signs of this syndrome. The disorder in this family followed a dominant mode of transmission. The pattern was consistent with either autosomal sex-limited or X-linked sex-limited inheritance. (10 refs)

78-6541 Characteristic Features of Hormonal Balance in Patients with Uterine Myoma. (Rus) Arakelian, A. B. (Dept. Obstetrics and Gynecology, Central Inst. Advanced Training Physicians, Moscow, USSR); Bakuleva, L. P. *Sov Med* (8): 140-144; 1978.

Changes in luteinizing hormone (LH), follicle-stimulating hormone (FSH), and somatotrophic hormone (STH) levels were studied in 35 patients with uterine myoma (26 were 24-40 yr old, 9 were 40-45 yr old). On the basis of rectal temperature, 14 patients had a normal biphasic menstrual

cycle, 13 had a reduced second stage of the cycle (duration of the follicular phase was 19 days, duration of the luteinic phase was 8 days), and 8 patients had a monophasic basal temperature. In all the patients, there was no correlation between the peaks of LH and FSH, and there were additional peaks of LH and FSH during the first and second phases of the menstrual cycle. The patients with a deficient luteinic phase and an anovular menstrual cycle had a significantly decreased basal level of LH and FSH. All the patients had mildly decreased levels of STH. Fourteen patients had internal endometriosis, and a retrospective analysis revealed that these patients underwent (on av) 3.13 abortions, compared with 1.64 abortions in the patients without endometriosis. (18 refs)

78-6542 Estimation and Correlation of High-Risk Factors for Cervical Carcinoma. (Por) Do Carmo Fajardo Diestel, M. (Servico de Ginecologia, Faculdade Catolica de Medicina de Porto Alegre, Porto Alegre, Brazil); Dos Reis Peroni, R. *J Bras Ginecol* 85(5): 253-254; 1978.

Early sexual activity, young age at first childbirth, multiparity, and, especially, combinations of these criteria were identi-

fied as high-risk factors for cervical carcinoma in 28 patients (6 refs)

See also:

- *(Rev.): 78-6034, 78-6049, 78-6055, 78-6059, 78-6061, 78-6062, 78-6063, 78-6064, 78-6066, 78-6068, 78-6069, 78-6070, 78-6072, 78-6073, 78-6074, 78-6075, 78-6078-6085.
- *(Chem.): 78-6120, 78-6135, 78-6136, 78-6147, 78-6157, 78-6191, 78-6195, 78-6201, 78-6211, 78-6222, 78-6235, 78-6261, 78-6276, 78-6280, 78-6332, 78-6351-6354.
- *(Phys.): 78-6363, 78-6364, 78-6375, 78-6377-6378.
- *(Viral): 78-6408, 78-6421, 78-6427, 78-6452, 78-6453-6486.
- *(Immun.): 78-6487, 78-6489, 78-6505.
- *(Epid.-Biom.): 78-6543, 78-6547, 78-6549, 78-6552-6553, 78-6565, 78-6573, 78-6575, 78-6576-6577.

EPIDEMIOLOGY AND BIOMETRY

543 **Pre- and Postmenopausal Breast Cancer: Differences in Risk Factors.** (Eng) Vakil, D. V. (Dept. Preventive Medicine and Biostatistics, Div. Community Health, Univ. Toronto, Toronto, Ontario, Canada); Goss, R. W. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1539-1550; 1978.

Because of the possibility that pre- and postmenopausal breast cancer represent two disease processes with different etiologies, an attempt was made to identify risk factors that differ between early- and late-onset breast cancer patients. A total of 170 former breast cancer patients were identified in the Toronto General Hospital tumor registry for 1960-1971. Information of genetic, epidemiological, and pathological importance was elicited in an interview questionnaire administered to all 170 women. Results of the survey revealed that 61 patients were premenopausal and 109 were postmenopausal at the time of diagnosis. Analysis of risk factors revealed that high socioeconomic status and early age at menarche were important in the development of premenopausal breast cancer. Nulliparity, age at first pregnancy, and weight-height index were more important in the development of postmenopausal breast cancer. In addition, a higher proportion of premenopausal patients (38%) had a family history of breast cancer, compared with the postmenopausal patients (22%). Thus, genetic factors are more important in the etiology of premenopausal breast cancer. (24 refs)

544 **Endocrine Status in Normal British, Japanese and Hawaiian-Japanese Women.** (Eng) Hayflick, J. L. (Breast Unit, Guy's Hosp., London SE1 9RT, England); Greenwood, F. C.; Glober, G.; Stemmerman, G.; Brook, R. D.; Wang, D. Y.; Kumaokas, S. *Eur J Cancer* 11: 1221-1228; 1978.

The hypothesis that there are no major differences in endocrine function among women with Japanese ancestry in Hawaii, Japanese women, and British women that would account for the differences in breast cancer incidence among these racial groups was investigated. Women were divided into three menstrual groups based on menopausal status. There were no differences in plasma and urinary estrogens among premenopausal women of the three racial groups. The postmenopausal Hawaiians had a higher plasma and urinary estrone level and a lower plasma progesterone level than British

or Japanese women. In postmenopausal Hawaiian women, plasma estradiol levels were significantly lower than those in the other two groups. The differences in androgen status were very complex: plasma androstenedione and urinary 11-deoxy-17-ketosteroid and etiocholanolone levels were lower in Hawaiian than British women and resembled those in Japanese women, but plasma dehydroepiandrosterone levels were higher in Hawaiians than in Japanese and resembled those in British women. This study reinforces the conclusion that ovarian and pituitary hormone levels do not explain the differences in incidence of breast cancer among the three groups and suggests that changes of environment bring about some alterations of endocrine function, mainly in the androgens. (39 refs)

78-6545 **Cancer of the Brain in Hawaii.** (Eng) Korsak, R. K. (Cancer Center Hawaii, Univ. Hawaii, Honolulu, HI); Rashad, M. N. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 2225-2228; 1978.

The incidence of malignant brain tumors in Hawaii was studied for two periods, 1960-1964 and 1968-1972. Age-specific incidence rates and age-adjusted rates were computed by sex and race (Caucasian, Chinese, Filipino, Hawaiian, and Japanese). There was a total of 162 brain tumors: 68 occurred during the first period, 94 during the second period. The age/race-incidence rates were 1.9/100,000 women and 2.2/100,000 men for the early period. During the later period, the incidence was 2.2/100,000 women and 2.7/100,000 men. There appeared to be significant racial differences, with Caucasians and Hawaiians having higher incidence rates than the other racial groups. Mortality due to malignant brain tumors was highest among Caucasian men and Hawaiian women. These results confirm the gradient between Caucasians and Orientals for brain tumors. (11 refs)

78-6546 **Cancer among Hawaiians and Part-Hawaiians.** (Eng) Rashad, M. N. (Dept. Genetics, Cancer Center Hawaii, Univ. Hawaii, Honolulu, HI); Mi, M. P.; Budy, A. M. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detec-

tion and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 2209-2218; 1978.

Cancer incidence among Hawaiians and part-Hawaiians was analyzed for the period 1968-1972. Of the total 1,057 cancer patients, 465 were women and 592 men. Age-specific and age-adjusted incidence rates were computed for selected sites. The age-adjusted annual incidence rate per 100,000 was 243.6 for men and 258.1 for women. The most common cancer in men was cancer of the lung, which accounted for 25% of all cases, followed by cancer of the stomach (8.6%). In women, the most frequent cancers were cancer of the breast (21.3%) and cervical cancer (17.2%). Comparison with other racial groups (Caucasian and Japanese) showed that Hawaiians and part-Hawaiians have the highest incidence rate of cancer of the lung and of the corpus uteri. This study confirms earlier reports of an excess of cancer among Hawaiians and part-Hawaiians. (30 refs)

78-6547 Contribution to the Problem of the Association of Silicosis and Lung Carcinoma. (Ita) Cardesi, E. (Servizio di anatomia e istologia patologica, Ospedale San Luigi Gonzaga di Orbassano, Turin, Italy); Provana, A. *Patologica* 70(1005/1006): 377-401; 1978.

Data on 204 patients with silicosis and/or pulmonary carcinoma diagnosed at autopsy or by postoperative histological examination at an Italian Hospital during 1975-1976 were tabulated and analyzed. Age, sex, profession, tumor histology and site, type of silicosis, and the carcinoma-silicosis association were considered. There were 135 patients with pulmonary carcinomas (116 men aged 34-81 yr and 19 women aged 50-81 yr). These tumors were subdivided into epithelial carcinoma (49 men, 3 women), microcytoma (17 men, 3 women), adenocarcinoma (18 men, 5 women), anaplastic and large cell carcinoma (26 men, 3 women), and bronchial-alveolar carcinoma (6 men, 5 women). The most frequent sites were the superior (60/135) and inferior lobes (34/135). Of the 32 silicosis patients, 29 were men (aged 40-80 yr) and 3 were women (aged 44-61 yr). Silicosis was classified as massive (10 men, 2 women), nodular (8 men, 1 woman), and peribronchial-vascular (11 men). Thirty-seven patients had both conditions (34 men aged 41-80 yr and 3 women aged 62-80 yr). The most frequent sites for silicosis-carcinoma were superior lobe (15/37), inferior lobe (12/37), and principal bronchus (5/37). Epithelial carcinomas were most common, and they were associated with all types of silicosis: massive (3/37), nodular (5/37), and peribronchial-vascular (7/37). Because of a lack of information, no statistically valid conclusions could be drawn on the association of various professions with silicosis or carcinoma. No statistically valid correlation could be made between the type of silicosis and the histological type of carcinoma. An indirect causal relationship may exist between silicosis and pulmonary carcinoma. (36 refs)

78-6548 Epidemiologic Studies of Cancer in Man and Animals Sharing the Same Environment. (Eng) Schneider, R. (Animal Neoplasm Registry, Dept. Epidemiology and Preventive Medicine, Univ. California, Sch. Veterinary Medicine, Davis, CA, 95616). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol 2(part 1), 2404 pp.; 1377-1378.

The patterns of human and animal cancer in the same environment were examined by comparing the incidence rate and sex pattern for the major sites of cancer found in humans, dogs, and cats in two California counties: Alameda and Contra Costa. Animal ages were converted to human age-equivalents, and both the human and animal crude incidence rates were age-adjusted to the same population age distribution determined by the 1950 US census. Human cancer rates were at least threefold higher than dog or cat rates for six sites: lung, bladder, digestive tract, prostate, ovary, and uterus. Possible reasons for these excess cancers are given. Cancer rates were threefold higher in the gum and testis, and feline rates were threefold higher for leukemias and lymphomas. Sex rate ratios for cancer showed excess male cases for humans and cats and excess female cases for dogs. Sex rates at specific sites indicated better agreement between humans and cats than between humans and dogs. Comparison of cancer occurrence in three species under similar conditions may help elucidate how cancer is expressed at specific sites. (10 refs)

78-6549 Association of Squamous Cell Carcinoma of the Head and Neck with Cancer of the Esophagus. (Eng) Goldstein, H. M. (Dept. Diagnostic Radiology, University of Texas System Cancer Center, M.D. Anderson Hosp. and Tumor Inst., 6723 Bertner Ave., Houston, TX, 77030); Zelen, J. *Am J Roentgenol* 131(5): 791-794; 1978.

The association between esophageal cancer and squamous cell carcinoma of the head and neck region (oral cavity, pharynx, and larynx) was studied by reviewing hospital files for the period March 1944 to August 1976. Eighty-nine patients were found who had separate primary squamous cell tumors of the head and neck and of the esophagus, representing 10% of all patients with esophageal carcinoma evaluated at the institution. In >80% of the patients, the separate primary tumors were discovered asynchronously (2 mo-16 yr apart). The expected number of esophageal carcinomas among the 10,000 patients with head and neck cancer was 10, compared with the 89 observed. The number of second primary tumors of the stomach and colon among the head and neck cancer population was not greater than anticipated. Most patients

a longstanding history of considerable cigarette and alcohol use. These findings suggest that alcohol and tobacco etiologic factors linking these two primary cancer groups, they indicate the importance of complete radiological evaluation of the esophagus in head and neck cancer patients. (5 refs)

6550 **Fluoridation of Water Supplies and Cancer - A Possible Association?** (Eng) Oldham, P. D. RC Pneumoconiosis Unit, Llandough Hosp., Penarth, S. (morgan, Wales); Newell, D. J. *J R Stat Soc Ser C* 2: 125-135; 1977.

a from a recent study of the crude cancer mortality rates in 10 US cities with fluoridated water supplies and 10 US cities with nonfluoridated water were reexamined to determine whether the conclusion that nationwide fluoridation would lead to excess cancer deaths was justified. At the beginning of the study (1950), the cities with fluoridated water had fewer elderly white men and women, more nonwhites aged 60 yr, and a greater proportion of cancer deaths than those in nonfluoridated water. At the end of the study, the cities with fluoridated water had many more nonwhites of all ages and fewer whites aged <55 yr than those with nonfluoridated water. Taking these variables into account, the excess cancer rate increased by 1% over the study period in the cities with fluoridated water and by 4% in those with nonfluoridated water. In absolute terms, the excess cancer rate between 1950 and 1970 increased by 8.8/100,000 in those with nonfluoridated water. It is concluded that the study provided no evidence that fluoride increases cancer mortality. (7 refs)

6551 **Cancer Epidemiologic Surveillance in the Du Pont Company.** (Eng) Pell, S. (Medical Div., E. I. duPont de Nemours and Co., Wilmington, DE, 19898); Berg, M. T.; Karrh, B. W. *J Occup Med* 20(11): 725-740; 1978.

The surveillance program started at the Du Pont Company in 1956 is described, and employee morbidity and mortality statistics are presented for the period 1956-1974. With certain exceptions, age-specific cancer incidence and mortality ratios were lower than those found in the Third National Cancer Survey. For men, the incidence ratio for all forms of cancer was 79 and the mortality ratio was 82. The corresponding ratios for women were 111 and 82. Mortality differences between Du Pont and the US decreased with increasing age. The incidence of malignant skin melanoma in men was significantly higher than expected, based on NCI data (Standard Incidence Ratio, SIR = 123, $P < 0.05$). The magnitude of the excess was slightly higher in salaried men (SIR = 127) than in wage role employees (SIR = 121). A significant excess of intestinal cancer occurred in salaried men employees (SIR = 117, $p < 0.05$). Among women, cervical cancer incidence was significantly higher than expected (SIR = 233, p

< 0.05), and it was especially high among the wage roll employees (SIR = 296). Breast cancer incidence was significantly higher in salaried employees (SIR = 136, $p < 0.05$), but was lower than expected in wage roll employees (SIR = 81, $P < 0.05$). Among salaried women employees, there were 11 cases of leukemia, which was significantly greater than the expected 6.1 ($p < 0.05$). Primary sites for which the incidence rates were especially low were for men: the thyroid, larynx, prostate, and buccal cavity, and pharynx; and for women, the lung. Cancer morbidity and mortality rates for the entire company cannot be relied on to indicate an increased risk of cancer resulting from exposure to a particular carcinogenic substance in the company. (16 refs)

78-6552 **Nasopharyngeal Carcinoma Patients in Massachusetts.** (Eng) Goodman, M. L. (Dept. Otolaryngology and Pathology, Massachusetts Eye and Ear Infirmary, James Homer Wright Pathological Lab., Massachusetts General Hosp. and Harvard Medical Sch., Boston, MA); Levine, P. H.; Miller, D. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1591-1598; 1978.

Possible etiological mechanisms of genetic susceptibility and Epstein-Barr virus (EBV) infections in nasopharyngeal carcinoma (NPC) were investigated. Thirty patients treated for NPC from 1972 to 1975 and controls matched by age (within 5 yr), race, sex, and level of education were evaluated for antibodies to EBV viral capsid antigen (VCA) and early antigen (EA). In addition, patients and controls were questioned about their personal habits and dietary practices. The two groups did not differ in the proportion of individuals reporting the use of coffee, tea, alcoholic beverages or cigarettes, although the NPC patients were heavier smokers than the controls (47% vs 22%, respectively, smoked 2 packs or more per day). VCA antibody determinations varied from 1:160 to 1:2560 for NPC patients and 1:40 to 1:1280 for controls, with respective median values of 736 and 370. EA antibody levels were also higher in NPC patients, 13/26 patients having titers of 160 and above, compared with 2/27 controls. Elevated VCA and EA antibodies, low socioeconomic status and heavy smoking habits were the most frequent consistent findings in the NPC group. (4 refs)

78-6553 **Incidence Rates and Risk Factors of Benign Breast Neoplasms.** (Eng) Cole, P. (Dept. Epidemiology, Harvard Sch. Public Health, 677 Huntington Ave., Boston, MA, 02115); Elwood, J. M.; Kaplan, S. D. *Am J Epidemiol* 108(2): 112-120; 1978.

A case-control study of benign breast disease was conducted in the Greater Boston area during 1968-1969. The cases were 938 women initially diagnosed as having fibrocystic disease (FD), fibroadenoma (FA), or a mixed lesion (ML) during the 18-mo study period. Controls were a random sample of the entire reference population. Mailed questionnaires were completed by 678/938 benign breast disease women and 1,807/2,062 controls contacted. Information on factors known to be related to breast cancer was obtained, to allow the similarities of the two conditions to be assessed. The age-standardized incidence rate for FD and FA was 89.4 and 32.8 per 100,000 woman-yr, respectively. The incidence of FD rose to age 45 and then declined sharply, that of FA peaked during the 20's, and that of ML had a mode at age 30-34. Among young women, the highest rates occurred in married nulliparae, but this was not so at higher ages. The incidence of FD and FA did not follow a pattern consistent with parity or age at first birth. The risk of FA increased with late age at menarche and that of FD increased with later natural menopause and with artificial menopause at any age. The risk of FD, but not of FA, also increased in women with arthritis and women in higher socioeconomic groups. The relative incidence of all three benign breast diseases decreased steadily with increasing obesity. The epidemiologic pattern of FD and FA did not correspond to that of breast cancer, which is directly related to obesity but not to age at menarche or to artificial menopause. Previous studies have indicated that the risk of breast cancer in benign disease may be limited to conditions such as severe ductular atypia or carcinoma in situ. (25 refs)

78-6554 Radiation, Smoking and Height Relationships to Lung Cancer in Uranium Miners. (Eng) Archer, V. E. (Natl. Inst. Occupational Safety and Health, Center Disease Control, US Dept. Health, Education and Welfare, 390 Wakara Way, Salt Lake City, UT, 84108). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 1689-1712; 1978.

A prospective mortality study using a life table method was done on 3,366 white underground uranium miners and 1,231 surface workers. Cumulative radon daughter exposure values were calculated for each miner from the date of his first mining to each sequential month of observation through September 1969. Miners were classified into tobacco groups on the basis of cigarette use only. Observed deaths exceeded those expected from respiratory cancer (146 vs 30), pneumoconiosis and related diseases, and accidents related to work. The overall rate of death from respiratory cancer among the miners was 4.9 times greater than that expected from general population data and 6.0 times greater than that found among the control group of surface workers. Exposure-response rela-

tionships with radiation varied with the height of the workers, cigarette smoking, and latent period between start of mining and disease. Of the four factors involved in malignant and nonmalignant respiratory diseases (high free silica, cigarette smoking, and alpha radiation), radiation was the most important. The clinical observation that short uranium miners have more lung disease than larger miners was substantiated by the data gathered in this study. Miners are engaged in physical labor, and tall men have larger lungs so it is postulated that short miners receive a higher dose of noxious agents per unit of lung tissue when performing heavy work. It is concluded that standards for airborne noxious agents should consider the amount of physical labor involved and the size of individual workers. (20 refs)

78-6555 A Comparison Model Approach to the Estimation of Tumor Incidence and Growth: Investigation of a Model of Cancer Latency. (Eng) Tolley, H. (Dept. Statistics, Texas A & M Univ., College Station, TX 77843); Burdick, D.; Manton, K. G.; Stallard, E. *Biometrika* 34(3): 377-389; 1978.

The nature of cancer, with its lengthy latent period, makes collection of data on the entire process of carcinogenesis difficult, as the early rate and pattern of tumor growth are unserved. Stochastic compartment modeling is proposed to solve the problems of estimating the transition probabilities of a partially observed stochastic process. The goal of the model is to synthesize the results of various studies into a coherent model of cancer latency and general population incidence of cancer. The rate coefficients for a time-dependent multiple compartment system were estimated for cancer-susceptible and nonsusceptible portions of a population, with individual rate coefficients being identified as explicit functions of age and tumor growth times. Under the model, tumor latency is relatively lengthy and if the population being studied is elderly, many tumors may not become clinically manifest due to mortality from other diseases. Susceptibility affects the model by fixing the proportion of persons who may get cancers. The model, with parameters representing tumor incidence, time to tumor death given onset, genetic susceptibility to tumor growth, and the effects of competing forces of mortality, was fitted to data on 1969 deaths due to stomach cancer for US men aged ≥ 25 yr. Two forms, one with waiting time distribution for occupants of the latent state and one with a single latency time, achieved excellent fits to the data. (33 refs)

78-6556 Associations of Cancer Mortality with Halomethanes in Drinking Water. (Eng) Cantor, K. (Health Effects Res. Lab., U.S. Environmental Protection Agency, Cincinnati, OH, 45268); Hoover, R.; Mason, T. McCabe, L. J. *J Natl Cancer Inst* 61(4): 979-985; 1978.

Age-standardized cancer mortality rates by site and sex f

tes in 76 US counties were compared with levels of chloroform, bromine-containing trihalomethane(s) (BTHM), and trihalomethane(s) (THM) in the drinking water. The 76 counties are > 50% urban, and over half the population is served by a sampled water supply. Using several demographic, socioeconomic, and industrial variables, a predicted mortality rate was computed and compared with the actual rate. The resulting residual mortality rate for each county was compared with the level of one of the three THM indicators using the bivariate correlation coefficient. Counties were grouped according to the overall percentage of the county served by the sampled water supply (low, intermediate, or high) and by geographic location. Bladder cancer residual mortality in both sexes showed a strong correlation with THM. This correlation persisted in spite of controls for sex, age, class, ethnic group, urban vs rural residence, presence of specific high-risk industries, and lung cancer rates (possible surrogate for levels of cigarette smoking). Brain cancer was not correlated in both sexes, but the associations were not strong, and there was regional variation. Other sites that gave indications of an association were: CHCl₃ levels and male kidney cancer, BTHM and female lung cancer, and THM levels and non-Hodgkin's lymphoma in men. Stomach cancer in women showed a negative correlation. (23 refs)

78-6557 **A Critique of the Statistical Evidence Associating Estrogens with Endometrial Cancer.** (Eng) Weinstein, A. R. (Dept. Medicine, Yale Univ. Sch. Medicine, New Haven, CT, 06510); Horwitz, R. I. *Cancer Res* 38(11, 12): 4001-4005; 1978.

Several types of bias that may have influenced the estrogen-endometrial cancer (E-EMC) association are described. In postmenopausal women with EMC, detection bias is produced because of the dilation and curettage or other diagnostic tests ordered when bleeding occurs as a side effect of E therapy. These tests allow the detection of EMC's that might otherwise be silent and undetected in women who do not take E. The problems of detection bias have been ignored in most case-control studies and they are difficult to deal with, as the bias occurs before cases and controls enter the study. To reduce this bias, the conventional method of epidemiological selection must be altered. Instead of two individual groups being chosen, both cases and controls can be selected from a single sampling frame consisting of people referred for performance of the particular diagnostic test used to identify the target disease. The bias can be reduced further by stratifying patients according to the cogent clinical reason that led to their referral to the hospital for the diagnostic test. Other sources of bias for the E-EMC association include the inadvertent prescription of E's for women with unrecognized EMC, the absence of double-blind interviewing techniques, and arbitrary definitions of exposure to E's. The current controversy about E's and EMC is a result of the inadequacy of conventional methods used for epidemiological research in chronic disease. (18 refs)

78-6558 **Geographic Patterns of Prostate Cancer in the United States.** (Eng) Blair, A. (Environmental Epidemiology Branch, Div. Cancer Cause and Prevention, NCI, NIH, Bethesda, MD, 20014); Fraumeni, J. F. *J Natl Cancer Inst* 61(6): 1379-1384; 1978.

Age-adjusted rates of mortality from prostate cancer during 1950-1969 were correlated by race with demographic, industrial, and agricultural data from 3,056 US counties. Prostate cancer was rare in young whites and nonwhites, increased progressively after 35 yr of age, and reached a plateau after 75 yr of age. Among whites, mortality rates in the midwest and northcentral regions were significantly higher than the total US rate, but the relative difference between the lowest and highest regional rates was only 8%, which is considerably less variation than that seen among whites with other tumors. Mortality among nonwhites was 50% higher than that among whites in all parts of the country where blacks comprise most of the nonwhite population. The rising rate associated with population density among nonwhites, but not whites, suggests that environmental exposures related to urban living may account for the predisposition of American blacks to prostate cancer. Mortality was elevated in counties with a high percentage of residents of Scandinavian descent, consistent with the high incidence reported for Sweden, Norway, and Denmark. Mortality was also elevated in counties with metal-using and textile industries and in regions with high consumption of high-fat foods. (46 refs)

78-6559 **Colon Cancer in a Low-Risk Population.** (Eng) Lyon, J. L. (Dept. Family and Community Medicine, Univ. Utah Coll. Medicine, Salt Lake City, UT, 84132); Sorenson, A. W. *Am J Clin Nutr* 31(10): S227-S230; 1978.

Colon cancer (CC) incidence from 1966 to 1970 was studied for the state of Utah, which had a CC mortality 34% below the US average during 1950-1969. The 10,641 cancer cases over this 5-yr period were compared to the age incidences of the Third National Cancer Survey (TNCS), broken down as to religious group (6,412 were Mormons), and computed as standard incidence ratios (SIR). The overall cancer SIR for the Mormon population was 75.6; for the non-Mormons, 93.9. For CC specifically, the corresponding SIR was 62.9 and 81.6, respectively. Mormons had a significantly lower risk of cancer of esophagus and lung than non-Mormons, due presumably to their religious ban on tobacco and alcohol. Furthermore, Mormon women had cancer incidence rates significantly lower than TNCS values at the colon-correlated sites of corpus uteri, breast, and ovary. Because per capita beef consumption in Utah is greater than the national average, and the anatomic distribution of colon cancer was not different between religious groups, Utah Mormons do not seem to adhere to a vegetarian diet. This conclusion, coupled with the low incidence of colon cancer, does not support the current hypothesis relating meats, fats, and dietary fiber to colonic cancer. (18 refs)

- 78-6560 Mortality in Unionized Truck Drivers.** (Eng) Luepker, R. V. (Lab. Physiological Hygiene, Stadium Gate 27, Univ. Minnesota, Minneapolis, MN, 55455); Smith, M. L. *J Occup Med* 20(10): 677-682; 1978.

Mortality rates among 184,435 male members of the Central States Teamsters Union during May-July 1976 were determined by surveying death certificates issued during that period and then comparing them with the rates for the general US population (GP). In this period, 249 deaths occurred, with the two leading causes being cardiovascular disease (104) and cancer (72). Except for respiratory tract cancer and motor vehicle accident deaths, mortality rates were lower than those in the GP and comparable to those in other employed populations. The mortality rate for all cancers paralleled and was not significantly different from that found in the GP. Respiratory tract cancers, mainly cancer of the lung, were the leading cause of death in the cancer category, accounting for 48% of the cases. Respiratory tract cancer mortality among the teamsters was in slight excess at ages 40-49 and 60-69 and significantly higher at ages 50-59 than that in the GP. It was not possible to separate the effect of cigarette smoking from environmental exposure to explain the excess respiratory tract cancer mortality. (12 refs)

- 78-6561 Occupational Cancer in Chromium Factory Workers.** (Jpn) Sano, T. (Dept. Pathology Occupational Diseases, Inst. Science Labor, Kawasaki-shi, Kanagawa Prefecture 213, Japan); Ebihara, I. *Jpn J Chest Dis* 37(2): 91-101; 1978.

Inflammation and cancer of the skin and mucosa of respiratory tract organs and, also, organs of the digestive tract have been found in chromium factory workers. These lesions are caused by the interaction of Cr and nickel, carcinogenic and inflammatory agents that infiltrate into the body during Cr production, with manganese and vanadium, inflammatory agents. Trivalent Cr plays a leading role in chronic inflammation and carcinogenesis. As of August 1975, there were 17 deaths due to lung cancer, 21 due to cancer of the respiratory organs, and 11 due to cancer of other organs (3 stomach cancers, 4 liver cancers, 1 neck sarcoma, 1 acute lymphatic leukemia, 1 reticulum cell sarcoma, and 1 renal cancer) among former Cr factory workers. Continued inflammatory stimulation induces repeated degeneration and regeneration of cells, leading to instability of the intranuclear gene and, hence, formation of an acquired factor for carcinogenesis. (27 refs)

- 78-6562 Cancer Epidemiology. Series No. 7: Solid Lymphomas in New South Wales.** (Eng) Ford, J. (Central Cancer Registry, Health Commission of NSW, GPO Box 4235, Sydney, New South Wales 2001, Australia). *Cancer Forum* 14: 111-114; 1978.

The incidence of solid lymphomas in New South Wales 1972 was investigated using data from the New South Wales Statutory Cancer Notification Form. Solid lymphomas include all cancers classified as lymphosarcomas, reticulosarcomas, Hodgkin's disease, giant follicular lymphoma, mycosis fungoides, and other malignant lymphomas, including those originating in organs other than lymph nodes. In 1972, the incidence of new cases was 246 for men (5% of cancers) and 212 for women (4% of new cancers). The incidence of solid lymphomas is greatest in the older age group (70 yr and over). There were 288 deaths from this cancer in 1972: 163 in men and 125 in women. There is no evidence that the lymphomas are transmitted by close personal contact. (no refs)

- 78-6563 High Incidence of Metalworkers among Patients with Testicular Cancer.** (Eng) Rhomberg, W. (Dept. Radiotherapy and Special Oncology, Sch. Medicine, Hannover, W. Germany); Schmoll, H. J. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer, by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404-1737-1743; 1978.

Forty-five patients who had undergone radiation therapy for a malignant testicular tumor in 1969-1974 were interviewed to obtain occupational and medical histories. The original diagnoses were as follows: 27 seminomas, 8 teratocarcinomas, 5 embryonal cell carcinomas, and 5 mixed forms. Nine of the patients were skilled metalworkers, and an additional seven patients had formerly been employed in this industry. The group of metalworkers included locksmiths, moldmakers, turners, car mechanics, electromechanics, and tool makers. Occupational exposure to metal compounds or metal dust ranged from 4 to 24 yr. Of the 19 nonmetalworkers, 11 had a history of exposure to metals (for 2-26 yr) that related directly or indirectly to their occupation, and other patients were exposed to metals as a result of a hobby. Only 9/45 patients showed no overt exposure to metals lasting for several years. (11 refs)

- 78-6564 Soft-Tissue Effects Following ²²⁴Ra Injection into Humans.** (Eng) Spiess, H. (Kinderpoliklinik, Universitat Munchen, Munich, W. Germany); Cech, A.; Mays, C. W. *Health Phys* 35(1): 61-81; 1978.

The soft-tissue effects of iv injections of radium-224 for treatment of tuberculosis, ankylosing spondylitis, and several other diseases were investigated in 204 patients treated as juveniles (1-20 yr old) and 612 treated as adults. Dosage and injection span were known in all patients. The average follow-up time from the first injection was 17 yr for the adults and

for the juveniles. Eighteen men injected as adults and two juveniles developed neoplasms, compared with 25 expected. Seventeen women injected as adults and three as juveniles developed cancer, exceeding the eight expected cases. However, when the male and female cases were combined, the 38 observed agreed closely with the 33 expected, suggesting that most of the soft-tissue neoplasms appearing so far are naturally occurring rather than related to ^{224}Ra therapy. Certain types of malignancy did occur at a rate above the expected value: leukemia (2 observed vs 0.8 expected), breast cancer (5 vs 1.2-1.8 expected), and renal cell carcinoma (3 vs 0.6-1.1 expected), which suggests that they might have been radiation-induced. More may appear later because of the latent period characteristic of radiation-induced cancer in humans. (41 refs)

78-6565 Soft Tissue Sarcomas and Intramuscular Injections: An Epidemiological Survey. (Eng) McIlroy, M. B. (Univ. Dept. Therapeutics, City Hosp., Nottingham NG5 1BB, England); Langman, M. J. *Br Med J* 3(41): 864-865; 1978.

Records of patients presenting with soft tissue sarcomas during 1974-1975 in the Trent, England, area were analyzed to determine whether the tumor could have been the result of previous intramuscular injections of iron preparations. Of 46 traceable patients with tumors on the leg, 12 had tumors at sites where injections might have been given, but only 2 cases arose specifically in the buttock. Records of all 46 patients were investigated, and no history of allergy, iron deficiency, or previous treatment by injections of any sort was noted. Likewise, of 19 traceable patients with sarcomas on the arm, 5 had tumors in an area where injections might have been given, but all 19 records proved negative for the types of information sought. These findings, therefore, do not support the suggestion of a causal relationship between sarcoma formation and intramuscular injections of iron or other preparations. (5 refs)

78-6566 Frequent Elevation of Tissue Polypeptide Antigen in the Sera of Workers Exposed to Bladder Carcinogens. (Eng) Kumar, S. (Christie Hosp. and Holt Radium Inst., Withington, Manchester M20 9BX, England); Mason, P.; Brenchley, P.; Taylor, G.; Bjorklund, B.; Eklund, M. *Int J Cancer* 22(5): 542-545; 1978.

Elevated levels of tissue polypeptide antigen (TPA), a substance purified from human tumors and placenta, have been found in sera from cancer patients. The diagnostic, prognostic, or epidemiological significance of sequential determinations of serum TPA levels was evaluated by comparing the results of these determinations in 108 male chemical workers exposed for at least 6 mo to 1-naphthylamine, 2-naphthylamine, benzidine, or a combination of these chemicals; 26 bladder cancer patients; and 63 healthy men. TPA was estimated by a hemagglutination-inhibition technique. The proportion

of individuals with a raised TPA level among the workers and patients was significantly higher (40/108 and 11/26) than that among the normal men (6/63). However, there was no significant difference in the proportion of positives between the workers and patients. It is suggested that a raised serum TPA level may be a very early indicator of hyperplastic bladder epithelium. (18 refs)

78-6567 Management of Urinary Bladder Cancer. (Eng) Fichardt, T. (Dept. Radiotherapy, H. F. Verwoerd Hosp., Pretoria, S. Africa); Sandison, A. G. *S Afr Med J* 54(18): 731-740; 1978.

From 1960 to 1976, 231 patients (198 men, 33 women) with primary cancer of the bladder were referred to a South African Hospital, and the etiology and clinicopathology of this disease are surveyed based on analyses of these patients and the literature. There were 115 transitional cell carcinomas, 87 anaplastic carcinomas, 15 squamous cell carcinomas, 12 papillary carcinomas, and 2 adenocarcinomas among the study group. Most (90%) of the patients presented with hematuria alone; the others presented with hematuria plus frequency and dysuria, incontinence, and bladder calculi. The duration of symptoms ranged from 2 wk to 2 yr, but most lasted 3-6 mo before diagnosis. The lesions were mostly single, situated mainly on the lateral and posterior walls and on the trigonum. Predisposing factors are numerous and varied, but cigarette smoking poses the greatest hazard for the general population. Cigarette smoking blocks the normal metabolism of tryptophan, leading to an accumulation of carcinogenic metabolites of tryptophan in the urine. The basic abnormality is an inability to convert 3-hydroxyanthranilic acid to nicotinamide. Other possible etiologic agents include the aromatic amines used in the dyestuff, rubber, and plastic industries; pesticides, especially those containing naphthylamines; dyes used in the manufacture of foods, cosmetics and drugs; bilharziasis; and coffee-drinking. Bladder calculi and bladder papillomas may also predispose to bladder cancer. (53 refs)

78-6568 Cancer in Cameroon: A Relative Frequency Study. (Eng) Jensen, O. M. (Unit Epidemiology and Biostatistics, International Agency Res. Cancer, 150 Cours Albert-Thomas, 69008 Lyon, France); Tuyns, A. J.; Ravis, P. *Rev Epidemiol Sante Publique* 26(2): 147-159; 1978.

Cancer was diagnosed and confirmed histologically in 1,390 males and 1,418 females in Cameroon, Africa, during 1969-1973. An attempt was made to correlate the cancer pattern with ethnic differences in the state. In both sexes, the pattern was dominated by a few tumor sites: men--skin 30%, malignant lymphomas 13%, and primary liver cancer 11%; women--skin 30%, uterine cervix 16%, and breast 10%. Ethnic differences were more apparent among men than women,

and they were correlated with neoplasms of the pharynx and buccal cavity, lung, connective tissue, and genitourinary tract. A close examination of tumor time trends showed that liver and respiratory tract tumors increased more rapidly than the total number of cancers. On the other hand, the same absolute number (a decreasing proportion) of skin cancers was recorded during each year of the study. These results suggest that the time trends reflect rapid changes in society and in medical facilities rather than actual changes in the frequencies of these tumors. For this reason, the results should be interpreted cautiously. (28 refs)

- 78-6569 Sperm Basic Proteins and Cervical Carcinogenesis (2 Letters to Editor).** (Eng) Reid, B. L. (Dept. Obstetrics and Gynaecology, Univ. Sydney, Sydney, N.S.W., 2006 Australia); French, P. W.; Hagan, B. E.; Coppleson, M. *Lancet* 2(8093): 788-789; 1978.

Two hundred and three men, only 3 of whom were students, were examined in a previous study to determine the correlation between two sperm basic proteins, histone and protamine, and social class. Recasting of the data in nonparametric terms using the analysis of variance by ranks of Kruskal-Wallis showed a correlation that was even more significant than that found originally. Differences in the incidence of cervical cancer by social class are marked and are among the greatest of all differences known. It is possible that the importance of the environment as a constant correlate of social class difference may operate through a subtle direct effect on cellular nucleic acids. A poor environment may impoverish the genome. However, it is important to determine the effects of increased sexual activity, multiple partners, genital herpes, and intercourse only with men who wear condoms or have been vasectomized on cervical cancer incidence before coming to any definite conclusions concerning etiology or prophylaxis. (8 refs)

- 78-6570 Cancer in Eastern Caribbean Islands.** (Eng) Rao, E. V. (Radiotherapy Dept., Public General Hosp., Kingston, Jamaica, W. Indies). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(Part 1), 2404 pp.; 2229-2240; 1978.

The incidence of cancer in the Eastern Caribbean Islands, based on data from the Eastern Caribbean Cancer and Clinical and Pathological Registries, was determined and compared with that in Africa, as 89% of the island residents are Negro. Since most of the islands did not keep cancer statistics, the data pertain mainly to Barbados, which contains more than half the total island populations. In 1971, there were 359 cases of cancer in Barbados, 221 in women and 138

in men. The most common types were in women, 76 of cervical carcinoma and 54 cases of breast cancer. There were no cases of male breast cancer. In men, cancer of stomach (18 cases), esophagus (12), and prostate (20) were most prevalent. Skin cancer showed a high rate in both men and women, 28 cases in men and 23 in women. The tumor incidence in other eastern Caribbean Islands (St. Vincent, St. Lucia, Dominica, etc.) was similar to that seen in Barbados except for a high incidence of bone tumors in St. Vincent. Hepatomas, lymphosarcomas, or leukemias, which are common cancers in Africa, were observed among the population of the Eastern Caribbean Islands. This may be due to changes in geography, environment, and/or social customs. (11 refs)

- 78-6571 Relation of Dysplasia and Carcinoma of the Uterine Cervix to Age at Onset of Sexual Activity and Number of Coital Partners.** (Eng) Berget, A. (Jul 4, DK-2820 Gentofte, Denmark). *Dan Med Bull* 25: 172-176; 1978.

Several risk factors in the development of dysplasia (D) or cervical carcinoma (CC) were investigated using subjects who took part in a CC screening test in a Danish county. The study comprised 67 women with a positive test (40 with D and 27 with CC) and 488 controls with a negative test, all aged 30-55. The risk of developing D or CC increased with increasing age at onset of sexual activity, evaluated from age at first coitus and at first pregnancy. The risk also increased with number of sexual partners, and this risk applied after corrections for social status and age at onset of sexual activity were made. The lower the social status the earlier the onset of sexual activity and, in contrast to controls, women with CC or D had an increasing number of sexual partners with decreasing social status. Thus, the risk of the two conditions increased with decreasing social status. Compared with the middle stratum, the risk was almost five times higher in the low and four times lower in the high stratum. Age at coitus and number of sexual partners were considered of practical use for identifying high-risk groups. In contrast to socioeconomic classification of women by their early education, used here, seems applicable to population screening. The correlations between sexual behavior and D or CC support the theory that the lesions are caused by an infectious agent. (29 refs)

- 78-6572 Diet, Transit Time, Stool Weight, and Cancer in Two Scandinavian Populations.** (Eng) MacLennan, R. (International Agency Res. Cancer, Cours Albert Thomas, 69372 Lyon, Cedex 2, France); Jensen, O. M.; Mosbech, J.; Vuori, H. *Am J Clin Nutr* 31: S239-S242; 1978.

Diet, stool transit time, and stool weight were compared in males, 55-64 yr of age, from urban Copenhagen, Denmark (CD), and rural Kuopio, Finland (KF), to investigate

fold differences in age-adjusted colon cancer incidence and in the two Scandinavian populations. Interviews and a food and drink diary kept for four consecutive days showed that the high incidence CD subjects consumed significantly more white wheat breads, total meats, pork, and beer, and less potatoes and milk. Although constipation and delay in stool passing were more common in CD, frequency of defecation and transit time were similar in both populations, with transit time showing a bimodal distribution. Stool weight was significantly higher in low incidence KF in autumn. Simple correlation coefficients of transit time and stool weight with bowel habits and diet components were calculated, and most correlations were low. In the 5/84 significant relationships that did exist between various foods and transit time or stool weight, all were negative, and the maximum absolute value of the coefficient was 0.28. Although the fiber hypothesis of colon cancer etiology was supported by computed fiber values (30.9 g/day in KF versus 17.2 in CD), the corresponding shorter mean transit time expected in KF was not noted. (0 refs)

78-6573 **Age at Menarche and Menopause of Uterine Cancer Patients.** (Eng) de Graaff, J. (Dept. Obstetrics and Gynecology, Vrije Universiteit, Amsterdam, Netherlands); Stolte, L. A. *Eur J Obstet Gynecol Reprod Biol* 44: 187-193; 1978.

The ages at menarche and at menopause of 868 cervical and endometrial cancer patients diagnosed during 1950-1955 and 1960-1965 were investigated. The mean age at menarche for all the cancer patients occurred in accordance with the literature, at an earlier age during 1960-1965 (13.88 yr, cervical cancer group; 14.03 yr, endometrial cancer group) than during 1950-1955 (14.08 and 14.24 yr, respectively). Analysis of the ages at menarche in relation to year of birth showed that there is no correlation between this factor and cervical endometrial cancer. However, menopause occurred later in the endometrial cancer group (49.52 yr for 1950-1955, 49.86 yr for 1960-1965) than in the cervical cancer group (47.93 yr for 1950-1955, 48.27 yr for 1960-1965). Menopause, therefore, may be a constitutional factor in the development of endometrial and/or cervical cancer. (19 refs)

78-6574 **A Note on the Use of Nitrate Fertilizers in a High-Risk Geographical Area for Stomach Cancer.** (Eng) Zaldivar, R. (International Res. Group Cancer Epidemiology and Human Ecology, 5055 N.W. 7th St., Suite 106, Miami, FL, 33126). *Z Krebsforsch* 92(2): 215-216; 1978.

In the 11 Chilean provinces that are at high risk for stomach cancer (age-adjusted mortality rates/100,000 men, 13.3-84.1), the amount of nitrate fertilizers (NaNO_3 , KNO_3) used, expressed in tons of N, was 238,950 tons. In the four low-risk provinces (mortality rates 39.1-46.2/100,000 men),

only 2,550 tons were used. The use of these fertilizers has been related to stomach cancer mortality. (6 refs)

78-6575 **Family Study of Gastric Carcinoma; With Special Reference to Histological Types.** (Eng) Lehtola, J. (Dept. Internal Medicine, Univ. Central Hosp., Oulu, Finland). *Scand J Gastroenterol Suppl* 13(50): 1-54; 1978.

The occurrence of gastric carcinoma (GC) was determined in 2,243 first-degree relatives of 341 patients (186 men and 155 women) with GC diagnosed in Uusimaa Province, Finland, during 1962-1963. Biopsy specimens of the probands were classified histologically as intestinal (139 cases), diffuse (67 cases), and undifferentiated (68 cases). In 67 cases, the type of GC could not be determined. The male/female ratio in the proband group was nearly identical to that for GC patients in the entire country (1.20 vs 1.28), but the mean age of the group was slightly lower than of other Finnish GC groups (62.6 vs 65.9 yr). The social group distribution of the patients was only slightly different from that of the entire Finnish population. The prevalence of GC among the 2,243 relatives was compared with that among 2,031 relatives of proband controls matched for sex and place and date of birth. Statistical analysis of the results revealed a relative GC risk of 1.50 in relatives of the probands vs controls ($p < 0.05$). When analyzed by cancer type, the relative risk was 1.41 for intestinal GC but 7.0 for diffuse GC ($p < 0.005$). No increased familial predisposition was noted in the probands with undifferentiated GC. When the relative with GC was the mother, the relative risks were 3.67 for the intestinal type cancer and 7.0 for the diffuse type. When several female relatives had GC, the relative risk increased to 8.0. Male cases showed little excess risk over control values. Further studies are warranted of the apparent familial clustering of diffuse GC, particularly among women. (177 refs)

78-6576 **The Incidence of Carcinoma in the Gastric Remnant after Resection for Benign Ulcer Disease.** (Eng) Ewerth, S. (Dept. Surgery, Karolinska Inst., Serafimer Hosp., Stockholm, Sweden); Bergstrand, O.; Hellers, G.; Ost, A. *Acta Chir Scand Suppl* (482): 2-5; 1978.

The incidence of cancer in the gastric remnant in 569 Swedish patients who had undergone gastric resection for benign ulcer disease during 1950-1953 was assessed. The operative technique was very consistent. There were 15 postoperative deaths. Tracing revealed that 223 patients had died, but in only 130 of these patients follow-up was adequate. Two cases with carcinoma in the gastric remnant, one found at autopsy and one radiologically verified, were found. Another patient with carcinoma of the distal esophagus and cardia was not included. Of the 253 patients who were alive when traced, 111 underwent endoscopy and biopsy. In one patient there was a severe dysplasia, but no invasive carcinomas were

found. The incidence of carcinoma in the gastric remnant found in this study was 0.8%, a value lower than that found 10-20 yr ago. The reason for this difference is not known, but it may be related to varying lengths of follow-up time or the marked overall decrease in the incidence of gastric cancer in Sweden. (16 refs)

78-6577 The Incidence of Carcinoma of the Gallbladder and Bile Ducts in Sweden 1958 to 1972. (Eng)

Broden, G. (Dept. Surgery, Karolinska Inst., Serafimer Hosp., Stockholm, Sweden); Ahlberg, J.; Bengtsson, L.; Hellers, G. *Acta Chir Scand Suppl* (482): 24-25; 1978.

The incidence of gallbladder and bile duct carcinomas in Sweden during 1958-1972 was investigated. The 5,913 reported cases were divided into three groups, each covering a 5-yr period. For each period, a mean annual incidence was calculated and related to sex and age. The absolute incidence of gallbladder and bile duct carcinomas increased during the study period. The relative incidence, however, showed a significant increase only among older patients, and especially older women. The risk of developing carcinoma of the gallbladder, which is impossible to cure, may be considered an additional indication for cholecystectomy in gallstone disease. (7 refs)

78-6578 The Changing Incidence of Malignant Melanoma of the Skin in Sweden, 1959-1968. (Eng)

Maliec, E. (Dept. Plastic Surgery, Karolinska sjukhuset, Univ. Stockholm, Stockholm, Sweden); Eklund, G. *Scand J Plast Reconstr Surg* 12(1): 19-27; 1978.

The 3,289 cases of malignant melanoma of the skin recorded in the Swedish Cancer Registry between 1959 and 1968 were analyzed with regard to sex, age, anatomical site, month of registration, and mortality. During this period, the annual

incidence rate (per 100,000) rose from 2.8 to 5.2 for men from 3.3 to 6.0 for women. The percentage annual increase for all sites was 7.0% for both men and women. The incidence rate was higher for women than for men, with a geometric annual mean of 4.4 and 4.0/100,000, respectively. The trunk was the commonest site in men, with a geometric annual mean incidence of 1.66/100,000. For women, the leg was the commonest site, with a corresponding incidence rate of 1.64. The highest increase occurred in women aged 30-39 and in men between the aged 40-64 yr. Mortality rate for malignant melanoma remained fairly constant, showing an annual change of -2% for men and +1% for women. For women, the incidence of melanoma on the legs was higher during the summer months than during the winter months. This could be a direct consequence of short-term exposure to UV radiation. (51 refs)

See also:

*(Rev.): 78-6002, 78-6004, 78-6009, 78-6010, 78-6017, 78-6018, 78-6019, 78-6021, 78-6027, 78-6028, 78-6029, 78-6030, 78-6032, 78-6033, 78-6034, 78-6035, 78-6037, 78-6038, 78-6039, 78-6041, 78-6043, 78-6044, 78-6049, 78-6050, 78-6066, 78-6067, 78-6068, 78-6076, 78-6077

*(Chem.): 78-6131, 78-6135, 78-6136, 78-6148, 78-6153, 78-6156, 78-6183, 78-6187, 78-6243, 78-6271, 78-6288, 78-6300, 78-6370, 78-6374, 78-6376, 78-6380

*(Phys.): 78-6370, 78-6374, 78-6376, 78-6380

MISCELLANEOUS

5579 **Units of Transcription for Cytoplasmic RNA in Mouse Myeloma Cells.** (Eng) Giorno, R. (Dept. Physics and Genetics, Univ. Colorado Medical Center, Denver, CO, 80262); Sauerbier, W. *Proc Natl Acad Sci USA* 75(9): 4374-4378; 1978.

Light mapping was used to estimate the sizes of transcription units for messenger RNA's (mRNA's) in mouse MPC myeloma cells. Cells were irradiated with varying doses of UV light, incubated with ³H-uridine (20-40 μCi/ml) for 1 hr in the presence of actinomycin D (0.08 μg/ml), and extracted. After the mRNA was purified from the cells, approximately 10 predominant mRNA species were resolved above a background of polydisperse mRNA by a 7 M urea/3.5% polyacrylamide slab gel system. UV-inactivation rates for the production of these major mRNA's were determined and converted to transcriptional distances in DNA base pairs. Target size analysis of the predominant mRNA's showed that they originated within transcription units that were at least 2 and possibly up to 14 times the size of the mature mRNA species. This may be a minimum estimate, since the target size for inhibiting mRNA production may have included promoter-distal sequences if they were not necessary for processing. Precursors longer than the observed target sizes may be found by complementary DNA-RNA hybridization. (34 refs)

5580 **Non-polyadenylated Messenger Ribonucleic Acids with Affinity for Polyadenyl-Sephadex** in Friend Murine Leukemia Cells. (Eng) Katinakis, P. (Dept. Biochemistry, Univ. Glasgow, Glasgow G12 8QQ, Scotland); Burdon, R. H. *Biochem Soc Trans* 6(4): 757-758; 1978.

A polyribosomal RNA fraction from Friend murine leukemia cells that has a high affinity for polyadenylic acid [poly(A)]-Sephadex was characterized. This poly(A)-binding RNA, which was isolated from polyribosomes on poly(A)-Sephadex, did not appear to be of ribosomal origin, its synthesis was unaffected by an agent that blocks ribosomal RNA synthesis. The RNA was released from polyribosomes by EDTA in a manner expected of messenger RNA (mRNA) and, after deproteinization, it sedimented in formaldehyde gradients at around 20S. Poly(A)-binding RNA appeared to lack poly(A) sequences longer than seven nucleotides, as judged by its inability to bind to polyuridylic acid [poly(U)]-Sephadex or by ³H-poly(U) hybridization. Binding of intact RNA to poly(A)-Sephadex seemed to be due to a poly(U)-rich region that also contained other nucleotides, notably guanylate residues. When introduced into a cell-free translation system, poly(A)-binding RNA stimulated the incorporation of several labeled amino acids into protein, in a manner linearly related to the amount of added RNA. In experiments in which DNA complementary to Friend cell polyadenylated mRNA was prepared using reverse transcrip-

tase and then used against poly(A)-binding RNA, little sequence homology was found. Thus, poly(A)-binding RNA is a discrete class of nonadenylated mRNA, and it has no direct relationship to polyadenylated mRNA. (16 refs)

78-6581 **Experimental Transplantable Tumors of the Nervous System and Their Application (Meeting Abstract).** (Eng) Mennel, H. D. (Dept. Neuropathology, Univ. Freiburg, Freiburg, W. Germany); Szymas, J. S.; Tamura, M.; Zulch, K. J.; Pakula, H. *J Neuropathol Exp Neurol* 37(5): 659; 1978. (no refs)

78-6582 **DNA Modifying Enzymes of *Agrobacterium tumefaciens*: Effect of DNA Topoisomerase, Restriction Endonuclease, and Unique DNA Endonuclease on Plasmid and Plant DNA.** (Eng) LeBon, J. M. (Dept. Biochemistry, Georgetown Univ. Medical Center, Washington, DC, 20007); Kado, C. I.; Rosenthal, L. J.; Chirikjian, J. G. *Proc Natl Acad Sci USA* 75(9): 4097-4101; 1978.

Three nucleic acid enzymes of *Agrobacterium tumefaciens*, which causes crown gall disease in dicotyledonous plants, were purified and characterized. These enzymes may play a role in the infection process by catalytically activating tumorigenic genes present in the bacterium or, alternatively, by modifying plant host DNA. The first enzyme, *Atu* DNA topoisomerase, relaxed only negatively twisted DNA. The second, *Atu* I, was a type II restriction endonuclease that generated DNA digestion patterns similar to those generated by *Eco*RII with several DNA substrates. The third activity, endonuclease A, preferentially catalyzed the conversion of superhelical DNA's to their corresponding linear forms. Polyacrylamide gel electrophoresis indicated that *Atu* DNA topoisomerase unwound plasmid pCK135 DNA (obtained from the virulent strain 1D135 of *A. tumefaciens*) and had an effect on plant DNA. *Atu* I altered both plant and plasmid DNA's, but endonuclease A only altered plasmid DNA. The mobility of plasmid and plant DNA's was also altered after treatment with combinations of these enzymes, suggesting a possible role for these enzymes in vivo. (28 refs)

78-6583 **Highly Conserved DNA of Ti Plasmids Overlaps T-DNA Maintained in Plant Tumours.** (Eng) Chilton, M. D. (Dept. Microbiology and Immunology, Univ. Washington, Seattle, WA, 98195); Drummond, M. H.; Merlo, D. J.; Sciaky, D. *Nature* 275(5676): 147-149; 1978.

An attempt was made to show that part of the tumor-inducing (Ti) plasmid maintained in tumor cells (T-DNA) is highly conserved among diverse Ti plasmids. It had been suggested that since common tumor-inducing functions are a feature of every Ti plasmid, a region of homologous DNA should

be detectable in all types. Plasmid pTi-B6-806 was mapped and the T-DNA, the region of this plasmid that is maintained in the E9 tobacco tumor, was located. DNA hybridization studies were conducted to test this region of pTi-B6-806 for homology with six representative Ti plasmids. PTi-B6-806 DNA fragments to be used as probes were isolated after preparative agarose gel electrophoresis of plasmid DNA digests and labelled by the nick translation reaction. Four of the six probes hybridized with all of the Ti plasmids examined and, therefore, they must constitute a region of DNA common to all. This region extends on the map from *Sma*I fragment 17 to *Hpa*I fragment 14, a region of 5.5×10^6 daltons or less. However, the results also show that a significant portion of the T-DNA of pTi-B6-806 comprises DNA sequences not common to all Ti plasmids. (26 refs)

78-6584 Homologous DNA Sequences in Different Ti-Plasmids Are Essential for Oncogenicity.

(Eng) Depicker, A. (Laboratorium voor Genetica, Rijksuniversiteit Gent, Ledeganckstraat 35, B-9000 Ghent, Belgium); Van Montagu, M.; Schell, J. *Nature* 275(5676): 150-153; 1978.

In view of the fact that different types of Ti plasmids only have a limited degree of DNA sequence homology but produce transformed plant cells with similar phenotypes, the hypothesis that at least part of the common DNA sequences would be directly and/or indirectly involved in oncogenicity was investigated. Homology between octopine and nopaline Ti plasmids was demonstrated with the use of plasmid DNA from wild-type strains of *Agrobacterium tumefaciens* (strain ACH5 was used as an example of an octopine Ti plasmid and strains C58 and K14 were used as examples of nopaline Ti plasmids). Hybridization experiments employing labeled DNA fragments showed that a specific DNA segment is common to both octopine and nopaline Ti plasmids. At least part of this common segment is involved in the oncogenic properties of the Ti plasmids, as integration of a complete P-type plasmid pRP4 into the pTi-C58 plasmid resulted in a loss of oncogenicity. Dissociation and subsequent hybridization experiments confirmed the fact the integration of pRP4 had indeed occurred in the common DNA sequence. Genetic evidence is given that the segment overlaps with the Ti plasmid DNA found in transformed plant cells. (24 refs)

78-6585 GC/MS Determination of Cytokinins in *Agrobacterium Tumefaciens* Cultures. (Eng)

Claeys, M. (Laboratorium voor Organische Chemie, R.U.G., Krijgslaan 271 [S4], B-9000 Ghent, Belgium); Messens, E.; Van Montagu, M.; Schell, J. *Fresenius Z Anal Chem* 290(2): 125-126; 1978.

Selected ion monitoring (SIM) was used to determine the cytokinins 6-(3-methyl-2-butenylamino)purine and 6-(4-hydroxy-3-methyl-trans-2-butenylamino)purine in oncogenic and nononcogenic *Agrobacterium tumefaciens* strains. The cytokinin levels were higher in the oncogenic strains. (4 refs)

78-6586 Disappearance of Patulin During Alcoholic Fermentation of Apple Juice. (Eng) Stinson, E.

(Eastern Regional Res. Center, Federal Res., Science Education Admin., U.S. Dept. Agriculture, Philadelphia, PA, 19118); Osman, S. F.; Huhtanen, C. N.; Bills, D. D. *Environ Microbiol* 36(4): 620-622; 1978.

Patulin (PL) is a carcinogenic mycotoxin produced by spoilage organisms, including *Penicillium expansum*, which infects apples. Its persistence in apple juice during fermentation was evaluated with the use of eight commercial yeast and three typical American fermentation procedures. In apple juice containing 15 mg/liter added PL, the level was reduced to $< 50 \mu\text{g/liter}$ in all but two cases (with 2 yeasts, 50 and 70 $\mu\text{g/liter}$ were found). In all cases, however, the level was reduced by $> 99\%$ during alcoholic fermentation. In unfermented samples, the level of added PL declined only 10% when the juice was stored for 2 wk, a time equivalent to that required for fermentation. French ciders have relatively high PL levels, and four factors related to harvesting and fermentation procedures are given that may account for the result. (17 refs)

78-6587 Fecal Profiles of Anaerobic Microflora of Large Bowel Cancer Patients and Patients with Nonhereditary Large Bowel Polyps. (Eng) Mastromarino, A.

(Div. Nutritional Biochemistry, Naylor Dana Inst., American Health Foundation, Valhalla, NY, 10595); Reddy, B. Wynder, E. L. *Cancer Res* 38(12): 4458-4462; 1978.

Differences in fecal bacteria profiles were studied in 7 normal controls, 13 patients with colorectal carcinoma, and 7 patients with nonhereditary large bowel polyps. Fresh fecal specimens were processed and analyzed under anaerobic conditions for bacterial growth and 7 α -dehydroxylase (7 α -cholesterol dehydrogenase (CD), and β -glucuronidase (CD) activities. Definitive identification of about 50 isolated anaerobes was based on the Gram reaction, morphology, spore test, gas-liquid chromatography, and other biochemical determinations. There were no significant differences in fecal anaerobic microflora or total aerobic counts, which suggests that taxonomic grouping of fecal bacteria is an inadequate measure of the relative risk of developing large bowel cancer. Results of the enzyme studies showed that 7 α -D and β -G activities were higher in samples from colorectal carcinoma patients and intestinal polyp patients, compared with healthy subjects, but no difference was noted in β -G activity between the three groups. It is postulated that since all subjects were on a Western diet, composition of the diet is a major factor in microbial β -G activity. (50 refs)

78-6588 Fiber in the Diet and Carcinoma. (Fre) Lambert, R.

(Service de Gastroenterologie, Hopital Edouard Herriot, F69003 Lyon, France). *Bull Cancer (Paris)* 65(1): 71-72; 1978.

The proposed relationship between dietary fiber and cancer of the colon is discussed. Dietary fiber has been defined

vegetable residue remaining after acid and alkaline hydrolysis, but this definition is not considered adequate or precise. Dietary fat and intestinal bacteria may be linked to fiber as factors in carcinogenesis. (no refs)

- 589 **Molecular Biology of Metabolic Disease: Defects in the Regulation of Enzymic Activity.** (Eng) Galton, D. J. (Diabetes and Lipid Res. Lab., St. Bartholomew's Hosp., W. Smithfield, London EC1A 7BE, England). *Enzyme* 23(4): 274-288; 1978.

The Jacob-Monod model for regulating enzyme activity was used to analyze the loss of allosteric inhibition of phosphofructokinase by citrate in lipomatosis, disorder of covalent modification of triglyceride lipase in triglyceride-storage disorder, and a defect in the repression of 3-hydroxy-3-methylglutaryl coenzyme A reductase in a new variant of familial hypercholesterolemia. The enzyme regulatory defects were compared with inborn errors of metabolism. (26 refs)

- 590 **Characterization of 7 α -Dehydroxylase in *Clostridium leptum*.** (Eng) Stellwag, E. J. (Medical College of Virginia, Virginia Commonwealth Univ., Richmond, VA 23298); Hylemon, P. B. *Am J Clin Nutr* 31(10): 1524-1527; 1978.

The distribution and enzymatic characterization of 7 α -dehydroxylase (7 α -DH) were studied in whole cells of *Clostridium leptum* as compared to other anaerobic intestinal bacteria. 7 α -DH was assayed both quantitatively and qualitatively in several species of bacteria by culturing with 7 α -chenodeoxycholic acid. 7 α -DH activity was detected in *C. leptum*, but not in 10 other *Bacteroides* species. Enzyme activity was seen over a broad pH range in *C. leptum*, with an optimum at 6.5-7.5. Addition of 0.1 mM sodium cholate to the cultures increased 7 α -DH activity six- to eight-fold during the stationary and declining growth phases. 2,4-Dinitrophenol (0.01, 0.1, or 1 mM), carbonylcyanide-m-chlorobenzylhydrazine (1 mM), and dicyclohexylcarbodiimide (0.2 mM) strongly inhibited 7 α -DH activity, whereas 2-N-heptyl-6-hydroxyquinoline-N-oxide (1 mM) did not. It is hypothesized that 7 α -DH is found only in a restricted population of intestinal bacteria, regulated by primary bile acid concentration; under certain conditions such as diet or disease including colon cancer, total fecal bile acids and 7 α -DH activity are both noted to be high. (20 refs)

- 591 **Increased Sialylation of Surface Glycopeptides of Human Trophoblast Compared with Fetal Cells from the Same Conceptus.** (Eng) Whyte, A. (Dept. Paediatrics, Univ. Cambridge, Cambridge, England); Loke, Y. *Exp Med* 148(4): 1087-1092; 1978.

The surface glycopeptides of human trophoblastic cells were compared with those of other fetal cells from the same embryos using double-labeling methods with isotopes of

L-fucose and D-glucosamine. (The human trophoblast shows some of the features commonly associated with neoplasia, including cellular proliferation and localized invasion.) A faster-eluting neuraminidase-sensitive fraction was observed upon Sephadex chromatography of the trophoblasts when D-glucosamine was used as the precursor. Labeling with fucose did not appear to result in the differences, which suggests that the glycopeptides characteristic of trophoblasts contained glucosamine-derived metabolic products, including sialic acid but excluding fucose. This increased sialylation is similar to, but not identical with, modifications observed in neoplastic cells. It is postulated that two species of glycopeptides may be involved in atypical cellular behavior: one contains sialic acid and other sugars excluding fucose and is associated with localized cellular growth and invasion, and the other contains both sialic acid and fucose and is characteristic of neoplastic cells. (14 refs)

- 78-6592 **Retinol Induces Density-dependent Growth Inhibition and Changes in Glycolipids and LETS.** (Eng) Patt, L. M. (Biochemical Oncology, Fred Hutchinson Cancer Res. Center, Seattle, WA, 98104); Itaya, K.; Hakomori, S. *Nature* 273(5661): 379-381; 1978.

The effect of retinol on cell growth behavior, glycolipid synthesis, and surface-exposed profile of glycoproteins in NIL and NILpy hamster fibroblasts and BALB/c 3T3 mouse cells was examined. Culturing of these cells in irradiated fetal calf serum containing retinol (20 nanomoles/ml) enhanced contact orientation and density-dependent inhibition of cell growth. Associated changes in cell-surface membrane GM₁ level, ganglioside contact response, and in large, external, transformation-sensitive protein (LETS protein) were also observed. Culturing of these cells in UV-irradiated vitamin A-free medium reduced contact orientation and density-dependent inhibition of cell growth. The ability of vitamin A compounds to prevent carcinogenesis could be related to the changes in surface membrane glycolipids and glycoproteins described. However, the glycosyltransferase levels for the synthesis of hematoside in NIL cells grown in medium supplemented with retinol did not differ significantly from those in cells grown in normal medium. Thus, the biochemical basis for the effect on glycolipid synthesis and surface-labeled glycoproteins could be a more complex matter. (17 refs)

- 78-6593 **Dimeric and High Molecular Weight Forms of the Large External Transformation-sensitive Protein on the Surface of Chick Embryo Fibroblasts.** (Eng) McConnell, M. R. (Dept. Pharmacology, Harvard Medical Sch., Boston, MA, 02115); Blumberg, P. M.; Rossow, P. W. *J Biol Chem* 253(20): 7522-7530; 1978.

The subunit structure of the large external transformation-sensitive (LETS) protein of chick embryo fibroblasts was examined. When cells were labeled by lactoperoxidase-catalyzed iodination and analyzed on sodium dodecyl sulfate-polyacrylamide gels in the absence of a reducing agent, no more than 1% of the ¹²⁵I-labeled LETS protein migrated

as a monomer (mol wt: 230,000-240,000). About 20%-23% of LETS migrated as a dimer. The remaining LETS was present as a high-mol-wt (HMW) complex retained at the top of the gel. These proportions did not appear to be an artifact resulting from disulfide bond formation induced by the lactoperoxidase labeling technique. The possibility that the LETS HMW complex was actually LETS dimer that was retained at the top of the gel by trapping was also ruled out. Pulse-labeling experiments demonstrated the conversion of LETS dimer to a HMW complex. The kinetics were biphasic, with rapid conversion of most of the LETS dimer followed by slow conversion of the remainder. The proportion of LETS present as a dimer predicted from the kinetic parameters agreed well with the experimental value determined by lactoperoxidase labeling. LETS dimer and the HMW complex were shed into the medium at similar slow rates; culture conditions seemed to have little influence on the subunit structure of LETS. The significance of the LETS subunit structure for its biological activity remains to be determined. (45 refs)

78-6594 Calcitonin Heterogeneity in Lung Cancer and Medullary Thyroid Cancer. (Eng) Becker, K. L. (Veterans Admin. Hosp., Washington, DC); Snider, R. H.; Silva, O. L.; Moore, C. F. *Acta Endocrinol (Kbh)* 89(1): 89-99; 1978.

To determine whether there is an appreciable difference between the immunoreactive forms (iCT) of human calcitonin in bronchogenic cancer (BGC) and medullary thyroid cancer (MTC), the immunochemical heterogeneity of this hormone was compared in patients with these neoplasms. The sera of 7 MCT patients and 15 BGC patients were analyzed by radioimmunoassay, gel filtration, and region-specific antisera. The serum of the MCT patients contained seven iCT fractions ranging in mol wt from $\geq 30,000$ to 2,500. In contrast, $> 80\%$ of the BGC patients had predominantly high-mol-wt iCT fractions ($\geq 30,000$ and 14,600) in their sera. Following in vitro incubation of the BGC serum at 25 C for 24 hr, the high-mol-wt fractions were converted to a heterogeneous pattern similar to that of MCT serum. Furthermore, in vitro incubation in normal human serum of the high-mol-wt fractions from MTC serum resulted in their degradation to fractions of lower mol wt. In view of the larger amounts of high-mol-wt fractions in the BGC sera and the greater recognition of these fractions by midportion antisera than by carboxyl terminal antisera, it was possible to differentiate, preoperatively, between the hypercalcitonemia of the two patient groups by comparing the midportion of carboxyl terminal ratio of serum iCT levels. Only 1/15 BGC patients had a ratio similar to that encountered with the MTC patients. (21 refs)

78-6595 Spontaneous Hepatomas in Mice Inbred from Ha:ICR Swiss Stock of Sex, Cedar Shaving in Bedding, and Immunization with Fetal Liver or Hepatoma Cells. (Eng) Jacobs, B. B. (American Medical Center Cancer Res. Center and Hosp., Lakewood, CO, 80214); Dieter, D. K. *J Natl Cancer Inst* 61(6): 1531-1534; 1978.

Spontaneous tumor development was compared in Ha: Swiss mice inbred (IB) by mating brother x sister for 2 generations (designated SWJ/Jac) and noninbred (NIB) of the same strain. A total of 350 NIB females (mean age 12 mo) were autopsied, and the following tumors were noted: hepatomas (5), hemangioendotheliomas (4), lymphoid tumors (75), breast tumors (15), lung tumors (104), pituitary tumors (58), thyroid tumors (6), and adrenal tumors (1). When 130 female SWJ/Jac mice (33 retired breeders and 97 virgins) were autopsied at age 12-24 mo, tumor incidence was as follows: hepatomas (9), hemangioendotheliomas (1), lymphoid tumors (36), breast tumors (5), lung tumors (1) and pituitary tumors (1). Tumors found in 86 SWJ/Jac mice autopsied at a mean age of 21.1 mo included hepatomas (1), hemangioendotheliomas (5), lymphoid tumors (9), and pituitary tumors (17). No other tumors were seen. The hepatomas in this group were often multifocal; their high incidence was due to the presence of cedar shavings in the animals' bedding. Treatment of 3-mo-old inbred males with crude extract of spontaneous hepatomas or fetal liver did not protect them from hepatoma development. An interesting result of this experiment was the propensity of inbred males to develop hepatomas (62.8% of animals). (12 refs)

78-6596 Studies on the Culture of Human Cancer Cell Lines and Their Production of Carcinoembryonic Antigen (CEA). (Jpn) Kaneko, Y. (First Dept. Surgery, Kagoshima Univ. Sch. Medicine, Kagoshima, Japan). *Igaku Kenkyu* 48(4): 298-321; 1978.

Three rectal adenocarcinoma cell lines (CaR-1, CaR-2, and CaR-3) were derived from poorly differentiated human rectal adenocarcinoma specimens, and their production of carcinoembryonic antigen (CEA) was investigated. The cells formed a monolayered sheet and attached firmly to the inner surface of culture vessels. Morphologically, they showed few tubular patterns and suggested signet ring cells. Electron microscopy revealed epitheliallike cells with microvilli, desmosomes, interdigitations, and mucinous granules. Doubling time of the three cell lines was 46, 48, and 120 hr, respectively, and their respective modal chromosome numbers were 64, 63, and 64. Histological findings of developed tumors in nude mice, following serial heterologous transplantation, revealed similar patterns as those in primary tumors, i.e., poorly differentiated adenocarcinoma. Metastatic lymph nodes were then explanted to nude mice, and resulting tumors were cultured in vitro. Inoculum from these tumors did not differ from cells taken directly from the original tumor. Small amounts of CEA were demonstrated in the culture medium of lines CaR-1 and CaR-2 and large amounts in line CaR-3. Synthesis of CEA did not change for almost 10-30 serial passages. Changing the culture medium and adding cyclic AMP and dibutyryl cyclic AMP to the medium of CaR-1 and CaR-2 cells did not affect CEA production. Electron microscopy of the high-CEA-producing CaR-3 cells showed well-developed microvilli, glycocalyx, and microvesicles. (87 refs)

597 Growth and Transplantability of Clonally Related Tumorigenic Murine Cell Lines. (Eng) Ker, J. R. (Univ. Warwick, Dept. Biological Sciences, Coventry CV4 7AL, England). *Br J Cancer* 38(4): 513-520; 1978.

In vivo and in vitro growth properties of eight cloned lines derived from a transplantable adenovirus 12-induced CBA mouse tumor cell line, were investigated. All lines were hyperdiploid, with modal numbers of 43 to 48 chromosomes. Metacentric markers were usually present in two or three copies per cell. Polyploidy, assessed separately, was 3%-10% in all clones. Although colony morphology differed markedly between clones, cells within a clone were homomorphic. In vitro, clone doubling times ranged from 24 to 45.4 hr and saturation densities from 31 to 111×10^4 /cm². In their response to the cytotoxic effects of concanavalin A at different concentrations, the clones demonstrated three modes of growth that were unrelated to their origin in vivo. Tumorigenicity, measured by the average number of cells required to produce tumors in 50% of injectable sites (TD₅₀), ranged from 2.5×10^4 cells to 6.3×10^5 cells. Mean latent period to palpable tumor development was 6.8 wk, and the volume doubling of established tumors was 109.4 hr. A high correlation was noted between doubling times in vivo and in vitro, but no correlation was demonstrated between in vitro growth characteristics or morphology and tumorigenicity in vivo. (22 refs)

598 The Growth of the EMT6 Tumour in the Lungs of BALB C Mice Following Intravenous Inoculation of Tumour Cells from Culture. (Eng) Twentyman, P. (Dept. Radiology, Stanford Univ. Medical Center, Stanford, CA, 94305). *Cell Tissue Kinet* 11(1): 57-68; 1978.

Growth of the EMT6 tumor in the lungs of Balb C mice was studied following iv inoculation of different numbers of tumor cells. Cell suspensions were prepared from explants of lungs and the number of in vitro colony forming cells was determined by plating into petri dishes. Following iv injection of 10^4 cells, the population doubling time of clonogenic tumor cells appearing in the cell suspension was 17.2 hr until such that the total tumor cell population per set of lungs reached 10^6 cells (10-12 days). Applying correction factors for changes in ability to extract cells from the lungs into the suspension at various times and also for possible changes in plating efficiency in vitro, the doubling time ranged from 20 to 24 hr. After reaching a total size of 10^6 tumor cells, the population doubling time of clonogenic tumor cells in the lungs increased. During the later stages of tumor growth a correlation was seen between total lung tumor wt and number of clonogenic cells present. For the final 3-4 days of initial period of rapid tumor growth, it is possible to obtain a hemocytometer count of tumor cells in the lung suspension. By plating out various numbers of tumor cells in vitro control lungs and lungs from treated animals, the effect of various treatments on the clonogenic capacity of cells from microscopic tumor foci can be determined. (16 refs)

78-6599 Long-Term Organ Culture of Human Uterine Endocervix. (Eng) Schurch, W. (Dept. Pathology, Univ. Maryland Sch. Medicine, Baltimore, MD, 21201); McDowell, E. M.; Trump, B. F. *Cancer Res* 38(11, part 1): 3723-3733; 1978.

A method for maintaining explants of human uterine endocervix in organ culture for 24 wk is described. The explants were obtained from 18 patients who had hysterectomies for various uterine diseases (usually myoma) and from 2 autopsies. Explants were cultured at 36.5 C according to a long-term organ culture technique using CMRL-1066 medium. The epithelium was viable for the entire time of culture, as demonstrated by ultrastructural morphology and mitotic activity. The morphology of the epithelium was near normal for 2-4 wk. In the following weeks, a slow transformation of the columnar epithelium into a metaplastic epithelium was observed, with a loss of mucus-secreting columnar cells and ciliated cells, and the appearance of layers of cuboidal and flattened epidermoid cells was observed deep in the clefts even after 24 wk in culture. Metaplasia began at the top of the papillae and ridges and extended downward into the clefts. However, in a few areas deep in the clefts, columnar cells retained evidence of mucus secretion during the duration of culture. Transformation of columnar mucous cells into cuboidal and flattened cells with epidermoid characteristics signified the development of epidermoid metaplasia. This observation was supported by ultrastructural studies, which showed mucus secretion and early keratinization in the metaplastic cells. Mitotic figures were seen under the transmission electron microscope for up to 24 wk, and all dividing cells contained mucous granules. Long-term culture of the human endocervix may represent a useful in vitro system for testing metabolic and morphological responses to potential chemical and biological carcinogens. (29 refs)

78-6600 In Vitro Culture of Human Leukemia Cells. (Eng) Dicke, K. A. (Univ. Texas System Cancer Center, M. D. Anderson Hosp. and Tumor Inst., 6723 Bertner, Houston, TX, 77030); Spitzer, G.; Verma, D. S.; McCredie, K. B. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 311-313; 1978.

A technique was developed for the selective in vitro growth of colonies of leukemia cells. It consists of a liquid phase of 15 hr, during which phytohemagglutinin (PHA) is used as a stimulus, before a semisolid phase of 7 days. The PHA colony-formation assay was applied to cells from patients with chronic and acute myeloid leukemia, and the results were found to correlate with clinical outcome of disease in the former and tendency to early relapse in the latter. (5 refs)

ಪ್ರತಿಭಾಷಿತರಾದವರನ್ನು ಸೇರಿಸಿ

Author Index

- Abdelaal, A. S., 78-6381
 Abraham, S. K., 78-6122
 Acheson, N. H., 78-6483
 Acs, G., 78-6205
 Adachi, T., 78-6535
 Adams, J. F., 78-6163
 Adams, R. A., 78-6321
 Ade, N., 78-6390
 Ageenko, A. I., 78-6468
 Agrawal, P., 78-6233
 Agrba, V. Z., 78-6461
 Ahearn, M. J., 78-6508
 Ahlberg, J., 78-6577
 Aisenberg, A. C., 78-6219
 Al-Saleem, T. I., 78-6487
 Al'tshtein, A. D., 78-6086
 Albanus, L., 78-6152
 Alekseeva, G. V., 78-6286
 Alfheim, I., 78-6142
 Allaudeen, H. S., 78-6453
 Allison, D. P., 78-6473
 Almqvist, A., 78-6507
 Aloni, Y., 78-6485
 Ames, B. N., 78-6005
 Amiel, J. L., 78-6074
 Amin, S., 78-6323
 Ander, P., 78-6132
 Anders, A., 78-6059
 Anders, F., 78-6059
 Anderson, D., 78-6165, 78-6260
 Anderson, H. A., 78-6023, 78-6169
 Anderson, L. M., 78-6249, 78-6250
 Anderson, R. S., 78-6296
 Andler, W., 78-6384
 Andon, B. M., 78-6164
 Anghileri, L. J., 78-6061
 Anzil, A. P., 78-6230
 Applebaum, E. L., 78-6515
 Arakelians, A. B., 78-6541
 Aravindakshan, M., 78-6108
 Arbeter, A., 78-6452
 Archer, V. E., 78-6554
 Arrendale, R. F., 78-6337
 Ash, J. F., 78-6396
 Ashby, J., 78-6100, 78-6165, 78-6260
 Asteris, G. T., 78-6376
 August, J. T., 78-6437
 Aungst, C. W., 78-6352
 Aurelian, L., 78-6449
 Baba, S., 78-6207, 78-6208
 Bacchini, P., 78-6524
 Bachman, D. S., 78-6365
 Baird, R. B., 78-6262
 Baird, S., 78-6419
 Bajaj, M. H., 78-6269
 Baker, A. L., 78-6354
 Baker, D. E., 78-6329
 Bakshi, K., 78-6144
 Bakuleva, L. P., 78-6541
 Ball, L. M., 78-6331
 Banerjee, A., 78-6195
 Barale, R., 78-6116
 Barnes, J. R., 78-6270
 Barrett, L. A., 78-6528
 Barthold, S. W., 78-6446
 Bartholomew, J. C., 78-6329
 Basrur, P. K., 78-6172
 Bassin, R. H., 78-6433
 Bates, R. C., 78-6447
 Battle, W. M., 78-6534
 Batzinger, R. P., 78-6328, 78-6333
 Bauchinger, M., 78-6136
 Baughman, B. B., 78-6083
 Baumanis, E. A., 78-6318
 Baxt, W. G., 78-6404
 Beales, P., 78-6062
 Beamer, J. L., 78-6369
 Becci, P. J., 78-6528
 Becker, K. L., 78-6594
 Bedenko, V., 78-6323
 Bend, J. R., 78-6331
 Bendinelli, M., 78-6430
 Benedict, W. F., 78-6195
 Bengtsson, L., 78-6577
 Benitz, K. F., 78-6294
 Benjamin, T., 78-6190, 78-6479
 Benner, J. F., 78-6170
 Bennett, M., 78-6427
 Benson, A. M., 78-6333
 Berget, A., 78-6571
 Bergstrand, O., 78-6576
 Bern, H. A., 78-6346
 Bertino, J. R., 78-6453
 Bertoglio, J., 78-6498
 Bertoni, F., 78-6524
 Beug, H., 78-6390
 Bex, F., 78-6088
 Bickers, D. R., 78-6336
 Bills, D. D., 78-6586
 Birnbaum, W., 78-6383
 Bishop, J. M., 78-6387, 78-6391
 Bjorklund, B., 78-6566
 Blaineau, C., 78-6431
 Blair, A., 78-6558
 Blair, P. B., 78-6409
 Blake, J. A., 78-6039
 Blakeslee, J. R., 78-6242, 78-6440
 Blanc, A. P., 78-6209
 Blinzinger, K., 78-6230
 Blohme, G., 78-6533
 Blokhin, V. A., 78-6143
 Blum, E., 78-6261
 Blumberg, P. M., 78-6593
 Bochkov, N. P., 78-6226
 Bockman, R. S., 78-6488
 Boczkowski, K., 78-6069
 Boh, E., 78-6334
 Bohnke, H., 78-6300, 78-6301
 Bojan, F., 78-6184
 Bolognesi, D. P., 78-6094
 Bone, D. R., 78-6448
 Bonse, G., 78-6167
 Borejko, J., 78-6378
 Borgstrom, G. H., 78-6507
 Borzsonyi, M., 78-6241
 Boulos, B. M., 78-6117
 Bourgaux, P., 78-6462
 Boutin, E., 78-6348
 Bower, R. K., 78-6400
 Bowman, M. C., 78-6119
 Boyd, H., 78-6276
 Bradley, M. O., 78-6366
 Bradshaw, J. J., 78-6340
 Brambilla, G., 78-6053
 Brandt, L., 78-6507
 Brankow, D. W., 78-6283
 Bratosin, S., 78-6485
 Breitburd, F., 78-6091
 Brencley, P., 78-6566
 Bridges, J. W., 78-6267
 Brinkerhoff, G., 78-6021
 Broden, G., 78-6577
 Brodsky, A., 78-6048
 Bronson, D., 78-6523
 Brooks, F. P., 78-6534
 Brown, A. M., 78-6450
 Brown, C. E., 78-6373
 Brown, D. P., 78-6014
 Brown, G. B., 78-6249, 78-6250
 Brown, J. K., 78-6070
 Brown, J. M., 78-6050
 Brown, M., 78-6448
 Brown, N. O., 78-6511
 Brown, R. A., 78-6179
 Brown, T. D., 78-6456
 Broxmeyer, H. E., 78-6488
 Brunden, M. N., 78-6162
 Brunnemann, K. D., 78-6009, 78-6010
 Brunson, K. W., 78-6060
 Brusick, D., 78-6144
 Buchan, P., 78-6343
 Buchanan, J. M., 78-6401
 Buck, C. A., 78-6056
 Budinger, J. M., 78-6249, 78-6250
 Budy, A. M., 78-6546
 Bueding, E., 78-6257, 78-6328, 78-6333
 Buikis, I. M., 78-6318
 Bukaeva, I. A., 78-6461
 Bulbrook, R. D., 78-6544
 Burdick, D., 78-6555
 Burdon, R. H., 78-6580
 Burke, V. T., 78-6334
 Burki, K., 78-6222
 Burkitt, D. P., 78-6038, 78-6043
 Burny, A., 78-6088
 Busch, H., 78-6072
 Busk, L., 78-6152
 Butschak, G., 78-6020
 Butterworth, K. R., 78-6149
 Caldwell, G. G., 78-6459
 Callahan, R., 78-6413
 Calnek, B. W., 78-6406
 Campbell, R. L., 78-6106
 Cannon, W. C., 78-6127, 78-6369
 Cantin, J., 78-6374
 Cantor, K. P., 78-6556
 Carbone, G., 78-6493
 Carcassonne, Y., 78-6209

- Cardesi, E., 78-6547
 Carlson, B. R., 78-6530
 Carlson, D. P., 78-6463
 Carmona, L. G., 78-6262
 Carrano, A. V., 78-6050
 Carroll, R. C., 78-6396
 Carson, H. C., 78-6006
 Carter, D. M., 78-6356
 Carter, R. L., 78-6065
 Cassens, R. G., 78-6203
 Castellazzi, M., 78-6155
 Cato, E. P., 78-6033
 Cavanna, M., 78-6053
 Cha, Y. N., 78-6257, 78-6333
 Chabner, B., 78-6024
 Chaganti, R. S., 78-6506
 Chamberlain, W. J., 78-6171
 Chang, C., 78-6228
 Chang, C. C., 78-6008, 78-6155
 Chantrenne, H., 78-6088
 Chardonnet, Y., 78-6475
 Chatteraj, S. C., 78-6351
 Chaturapit, S., 78-6312
 Chauhan, P. S., 78-6108
 Chen, C. B., 78-6175, 78-6176, 78-6237
 Chesnokova, V. M., 78-6278
 Chhabra, R. S., 78-6227
 Chien, Y. H., 78-6423
 Chilton, M. D., 78-6583
 Chin, A. E., 78-6217
 Chirikjian, J. G., 78-6582
 Chortyk, O. T., 78-6337
 Christiansen, G., 78-6107
 Christman, J. K., 78-6205, 78-6310
 Chu, F. C., 78-6374
 Chun, R. W., 78-6517
 Chute, R. N., 78-6373
 Chuvirov, G. N., 78-6461
 Claeys, M., 78-6585
 Clark, A. M., 78-6238
 Clark, C. G., 78-6180
 Clarke, C. H., 78-6159
 Clary, A. J., 78-6369
 Clauss, U., 78-6429
 Clemmesen, J., 78-6029
 Cleuter, Y., 78-6088
 Cohen, M., 78-6024
 Coiffier, B., 78-6211
 Colburn, W. A., 78-6281
 Cole, P., 78-6553
 Cole, T., 78-6428
 Combes, R. D., 78-6153
 Connelly, R., 78-6459
 Cooper, G. M., 78-6393, 78-6403
 Cooper, H. K., 78-6221
 Cooper, M. D., 78-6078
 Copeland, D. D., 78-6408
 Coplan, M., 78-6261
 Copp, R. P., 78-6310
 Coppleson, M., 78-6569
 Cordell, B., 78-6391
 Cornelius, E. A., 78-6491
 Correa, P., 78-6204
 Coulston, F., 78-6294
 Craddock, V. M., 78-6216
 Craig, D. K., 78-6369
 Craig, E. A., 78-6463
 Crenshaw, J. W., 78-6220
 Crist, W. M., 78-6078
 Croissant, O., 78-6090, 78-6091
 Crombie, I., 78-6448
 Crump, K. S., 78-6111
 Cruse, J. P., 78-6180
 Curphey, T. J., 78-6200
 Cuthbert, J. W., 78-6022
 Cutler, R. G., 78-6422
 Cuzin, F., 78-6478
 Czernobilsky, B., 78-6344
 D'Andrea, A. D., 78-6124
 Dagen, J. E., 78-6499
 Dahl, G. A., 78-6185
 Daniel, F. B., 78-6055
 Das, B., 78-6271
 Davidson, N., 78-6423
 Dawson, P. C., 78-6121
 De Flora, S., 78-6130
 de Graaff, J., 78-6573
 De Groot, L. J., 78-6376
 de la Chapelle, A., 78-6507
 De Larco, J. E., 78-6481
 De Marini, D. M., 78-6150
 de Noronha, F., 78-6443
 De Serres, F. J., 78-6001, 78-6100
 Decker, J. R., 78-6369
 Decleve, A., 78-6418
 Deichmann, W. B., 78-6261
 Deitch, A. D., 78-6426
 Dekegel, D., 78-6088
 Deknudt, G., 78-6147
 Delbecchi, L., 78-6462
 Demerjian, Z. N., 78-6219
 Den Otter, W., 78-6082
 Denda, A., 78-6212
 Dennis, E. H., 78-6280
 DeOme, K. B., 78-6527
 Depass, L. R., 78-6192
 Depicker, A., 78-6584
 DePierre, J. W., 78-6320
 Dethlefsen, L. A., 78-6050
 Dewar, A. E., 78-6505
 DeWitte, W. J., 78-6103
 Diad'kova, A. M., 78-6084
 Diamandopoulos, G. T., 78-6085
 Dicke, K. A., 78-6600
 Dieter, D. K., 78-6595
 Dikshtein, E. A., 78-6532
 Dingemans, K. P., 78-6526
 DiPaolo, J. A., 78-6245
 Dipple, A., 78-6313
 Do Carmo Fajardo Diestel, M., 78-6542
 Doerfler, W., 78-6470
 Dohmae, N., 78-6514
 Doi, K., 78-6207
 Doll, R., 78-6031
 Dong, M., 78-6176
 Dore, J. F., 78-6498
 Dos Reis Peroni, R., 78-6542
 Douglas, G. R., 78-6324
 Dresch, F., 78-6136
 Drewinko, B., 78-6508
 Drummond, M. H., 78-6583
 Dube, S. K., 78-6428
 Dullens, H. F., 78-6082
 Dumont-Herskowitz, R. A., 78-6540
 Duran-Reynals, M. L., 78-6421
 Eastcott, J. W., 78-6427
 Ebihara, I., 78-6561
 Eckelmann, A., 78-6145
 Eckhart, W., 78-6496
 Eklund, G., 78-6566, 78-6578
 Elder, J. H., 78-6436, 78-6439
 Eliakim, M., 78-6513
 Eliasson, R., 78-6482
 Eliseev, V. V., 78-6372
 Elwood, J. M., 78-6553
 Englert, N., 78-6136
 Enzell, C., 78-6177
 Epler, J. L., 78-6102
 Erfle, V., 78-6438
 Erickson, C. A., 78-6480
 Erickson, L. C., 78-6366
 Eriksson, K. E., 78-6132
 Eriksson, L. C., 78-6320
 Erkhov, V. S., 78-6468
 Essex, M., 78-6443, 78-6444
 Evans, I. M., 78-6522
 Ewerth, S., 78-6576
 Fabrizio, P., 78-6459
 Falk, J., 78-6518
 Fanshier, L., 78-6387
 Faras, A., 78-6474
 Farber, E., 78-6492
 Farber, T. M., 78-6123
 Faucounau, N., 78-6248
 Favor, J. B., 78-6220
 Favre, M., 78-6090, 78-6091
 Feinstein, A. R., 78-6151, 78-6557
 Felman, P., 78-6211
 Ferrucci, J. T., 78-6219
 Fiala, E. S., 78-6107
 Fiala, S., 78-6274, 78-6275
 Fialkow, P. J., 78-6509
 Fichardt, T., 78-6567
 Fidler, I. J., 78-6060
 Fiere, D., 78-6211
 Filipe, M. I., 78-6536
 Finegold, S. M., 78-6035
 Fischbein, A., 78-6529
 Fischer, K. M., 78-6432
 Fisenne, I. M., 78-6367
 Fisher, B. K., 78-6519
 Fishman, J., 78-6349
 Fleig, I., 78-6187, 78-6252
 Fleischer, D., 78-6179
 Flesch, H. G., 78-6432
 Ford, J., 78-6562
 Forrester, P. I., 78-6265
 Forster, C. J., 78-6518
 Francke, B., 78-6448
 Fraumeni, J. F., 78-6558
 Fredlund, P., 78-6537
 Fredrickson, T. N., 78-6141
 Freeman, C. G., 78-6236
 French, P. W., 78-6569
 Friedman, H., 78-6430
 Friedman, L., 78-6123
 Frolova, E. I., 78-6466

- Fry, J. R., 78-6267
 Fujie, S., 78-6173
 Fujii, S., 78-6208
 Fujimoto, Y., 78-6521
 Fujinaga, K., 78-6467
 Fukami, J., 78-6305
 Fukasawa, S., 78-6272
 Fukushima, D., 78-6349
 Fumero, S., 78-6116
 Furukawa, M., 78-6194
 Fyrand, O., 78-6046
 Galbadini, J. J., 78-6219
 Galton, D. J., 78-6589
 Gastaut, J. A., 78-6209
 Gateff, E., 78-6075
 Gaunt, I. F., 78-6149
 Gautsch, J. W., 78-6436, 78-6439
 Gaynor, A. L., 78-6356
 Gazdar, A. F., 78-6434
 Gazitt, Y., 78-6426
 Geder, L., 78-6499
 Gensheimer, G. G., 78-6495
 German, J., 78-6506
 Gersbach, H., 78-6308, 78-6325
 Gerspach, A., 78-6564
 Gettemy, D. J., 78-6279
 Ghent, W., 78-6518
 Ghysdael, J., 78-6088
 Gibson, W., 78-6484
 Gillam, M. A., 78-6554
 Gillespie, G. Y., 78-6502
 Gillespie, J. J., 78-6539
 Gillette, J. R., 78-6291
 Gilmore, C. E., 78-6123
 Giorno, R., 78-6579
 Gisselbrecht, S., 78-6431
 Gissmann, L., 78-6476, 78-6477
 Gist-Brocades, N. V., 78-6256
 Glaser, M., 78-6500
 Glatt, H. R., 78-6166
 Glavin, F., 78-6528
 Glazko, T. T., 78-6278
 Glinski, W., 78-6090
 Globar, G., 78-6544
 Globar, G. A., 78-6126
 Glover, T. W., 78-6155
 Glowinski, I. B., 78-6264
 Goel, K. A., 78-6233
 Goggelmann, W., 78-6167
 Goldberg, A. K., 78-6398
 Goldberg, A. R., 78-6397
 Goldenberg, C. J., 78-6463
 Goldfarb, R. H., 78-6395
 Goldin, B., 78-6348
 Goldman, R., 78-6435
 Goldstein, H. M., 78-6549
 Goldstein, J. C., 78-6089
 Gomez, G. A., 78-6494
 Goodall, J., 78-6518
 Goodin, L. J., 78-6121
 Goodman, M. L., 78-6552
 Gorbach, S. L., 78-6348
 Gordonson, J., 78-6383
 Gori, G. B., 78-6036
 Goss, P., 78-6362
 Goto, M., 78-6272
 Graf, T., 78-6390
 Grant, C. E., 78-6324
 Grantham, P. H., 78-6190
 Grasso, P., 78-6149
 Greaser, M. L., 78-6203
 Green, A. E., 78-6044
 Green, N. R., 78-6290
 Green, S., 78-6138
 Green, T., 78-6112
 Greenberger, J., 78-6457
 Greenland, T., 78-6475
 Greenwood, F. C., 78-6544
 Greim, H., 78-6167
 Grierson, W., 78-6012
 Grigor'ev, A. S., 78-6461
 Grimelius, L., 78-6533
 Grimmer, G., 78-6300, 78-6301
 Groneberg, J., 78-6470
 Groopman, J. E., 78-6219
 Gruener, N., 78-6105
 Guess, H. A., 78-6111
 Guirgis, H., 78-6349, 78-6538
 Gusina, N. B., 78-6183
 Guzman, R. C., 78-6527
 Gyles, N. R., 78-6400
 Haddon, W. F., 78-6101
 Haebeler, A. F., 78-6171
 Hagan, B. E., 78-6569
 Hakomori, S., 78-6592
 Halberg, F., 78-6156
 Hall, K. Y., 78-6055
 Han, T., 78-6494
 Hancock, R. L., 78-6265
 Hand, P. H., 78-6413
 Hankin, J. H., 78-6037
 Hansen, C. B., 78-6502
 Hansford-Miller, F., 78-6019
 Hanson, G. A., 78-6363
 Hansson, G., 78-6379
 Haran-Ghera, N., 78-6314
 Hardigree, A. A., 78-6102
 Hardy, W. D., 78-6443
 Hargis, A. M., 78-6516
 Harris, C. C., 78-6528
 Harris, H., 78-6161
 Harris, R., 78-6538
 Hart, R., 78-6242
 Hart, R. W., 78-6055
 Harvey, H. J., 78-6511
 Harvey, R. C., 78-6359
 Hasan, F. M., 78-6168
 Haseltine, W. A., 78-6124
 Hashida, C., 78-6254
 Hashimoto, Y., 78-6246
 Haskins, J. E., 78-6262
 Hathway, D. E., 78-6112
 Hattori, N., 78-6531
 Hauenstein, E., 78-6221
 Haugen, A., 78-6231
 Hauschild, F., 78-6145
 Havers, W., 78-6384
 Hayashi, K., 78-6182
 Hayatsu, H., 78-6160
 Hayes, A., 78-6511
 Hayes, R. B., 78-6131
 Hayhoe, F. G., 78-6457
 Hayward, J. L., 78-6544
 Heath, C. W., 78-6459
 Heaton, K. W., 78-6178
 Hecht, S. S., 78-6009, 78-6175, 78-6176
 78-6237, 78-6323
 Hedden, L., 78-6118
 Hedinger, R. A., 78-6044
 Hehlmann, R., 78-6438
 Heidelberg, C., 78-6283
 Hellers, G., 78-6576, 78-6577
 Henderson, A. R., 78-6216
 Hendy, R. J., 78-6149
 Henle, W., 78-6452
 Henry-Amar, M., 78-6068
 Henry, M. C., 78-6335
 Hens, L., 78-6140
 Henschler, D., 78-6167
 Hensley, G. T., 78-6276
 Hensse, S., 78-6489
 Herman, K., 78-6184
 Hettrick, J. M., 78-6106
 Hewetson, J., 78-6452
 Heyns, K., 78-6225
 Higman, H. C., 78-6337
 Hill, D. L., 78-6115
 Hill, P. A., 78-6522
 Hilleman, M. R., 78-6442
 Hillyard, C. J., 78-6522
 Hinson, J. A., 78-6291
 Hinuma, Y., 78-6454
 Hirano, K., 78-6206
 Hiremath, S. P., 78-6299
 Hirsch, M. S., 78-6219
 Hochberg, M. C., 78-6197
 Hoevels, J., 78-6537
 Hoffmann, D., 78-6009, 78-6010
 78-6175, 78-6176, 78-6237, 78-6323
 Hoffmann, G. R., 78-6359, 78-6360
 Hogg, N., 78-6435
 Holben, J. A., 78-6416
 Holdeman, L. V., 78-6033
 Holder, G. M., 78-6312
 Holtzman, S., 78-6342
 Homburger, F., 78-6321
 Honda, M., 78-6201
 Hooper, A. A., 78-6293
 Hooson, J., 78-6149
 Hoover, R., 78-6556
 Hopfer, S. M., 78-6141
 Horowitz, M., 78-6485
 Horrobin, D. F., 78-6451
 Hortling, L., 78-6388, 78-6389
 Horvat, D. J., 78-6135
 Horwitz, K. B., 78-6350
 Horwitz, R. L., 78-6557
 Hoshi, A., 78-6232
 Hoshino, H., 78-6224
 Huang, C. C., 78-6194
 Huang, P. C., 78-6364
 Huebner, R. J., 78-6417
 Huggins, C. B., 78-6317
 Huhtanen, C. N., 78-6586
 Hunter, T., 78-6484, 78-6496
 Hutchinson, M. A., 78-6496
 Huvos, A., 78-6374
 Hylemon, P. B., 78-6590

- Iakovleva, L. A., 78-6461
 Igarashi, K., 78-6464, 78-6465
 Iitaka, Y., 78-6198
 Ikeda, H., 78-6425
 Ikeda, K., 78-6148
 Ikuta, K., 78-6407
 Imai, S., 78-6414
 Inbar, M., 78-6458
 Inoguchi, T., 78-6535
 Insler, V., 78-6344
 Invernizzi, G., 78-6493
 Ishii, D. N., 78-6309
 Isomoto, H., 78-6535
 Israel, E., 78-6394
 Itaya, K., 78-6592
 Ivanetich, K. M., 78-6340
 Ivankovic, S., 78-6210
 Jablonska, S., 78-6090, 78-6091
 Jacobs, B. B., 78-6595
 Jacobsen, N., 78-6142
 Jacobson, G., 78-6383
 Jacobson, R. J., 78-6509
 Jacquignon, P. C., 78-6166
 Jaggi, W., 78-6345
 James, K., 78-6081
 James, L. A., 78-6554
 Jarzabek-Chorzelska, M., 78-6090
 Jeleniewska, A., 78-6378
 Jenkins, R. L., 78-6262
 Jensen, F. C., 78-6436
 Jensen, O. M., 78-6568, 78-6572
 Jewell, W. R., 78-6504
 Johnson, E. L., 78-6186
 Johnson, R. G., 78-6121
 Johnston, T. P., 78-6115
 Jones, L. A., 78-6346
 Jones, R. F., 78-6411
 Jonsen, J., 78-6142
 Jorgenson, T. A., 78-6138
 Kada, T., 78-6128, 78-6206
 Kaddargi, S. S., 78-6299
 Kadlubar, F. F., 78-6247
 Kado, C. I., 78-6582
 Kaiser, H. E., 78-6528
 Kaledin, V. I., 78-6278, 78-6286
 Kallio, A., 78-6388
 Kameda, S., 78-6531
 Kamine, J., 78-6401
 Kaneko, S., 78-6207
 Kaneko, Y., 78-6596
 Kanematsu, K., 78-6128
 Kang, C. Y., 78-6405
 Kanoh, T., 78-6514
 Kaplan, H. S., 78-6418
 Kaplan, S. D., 78-6553
 Kappas, A., 78-6336
 Karagianes, M. T., 78-6369
 Karmali, R. A., 78-6451
 Karpas, A., 78-6457
 Karrh, B. W., 78-6551
 Kasamatsu, T., 78-6449
 Kashyap, A. K., 78-6137
 Katinakis, P. K., 78-6580
 Kato, Y., 78-6531
 Katyal, S. L., 78-6199
 Katz, A. D., 78-6523
 Kauffman, S. L., 78-6385
 Kawabata, H., 78-6212
 Kawachi, T., 78-6201, 78-6232, 78-6263
 Kawana, T., 78-6449
 Kawashima, R., 78-6259
 Kawashima, S., 78-6346
 Kazemi, H., 78-6168
 Kazumi, T., 78-6207, 78-6208
 Kelleher, P. C., 78-6277
 Keller, R., 78-6080
 Kellerer, A. M., 78-6370
 Kellermann, G., 78-6292
 Kellermann, G. H., 78-6322
 Kelly, P. A., 78-6473
 Kelsey, C. A., 78-6047
 Kennel, S. J., 78-6473
 Kesavan, P. C., 78-6122
 Kettering, W. G., 78-6274
 Kettmann, R., 78-6088
 Khan, M. A., 78-6123
 Khoudokormoff, B., 78-6256
 Khrustalev, S. A., 78-6202
 Kihlman, B. A., 78-6120
 Killough, G. G., 78-6386
 Kilton, L. J., 78-6434
 Kim, J. H., 78-6374
 Kindblom, L. G., 78-6533
 Kirkland, D. J., 78-6151
 Kirsch-Volders, M., 78-6140
 Kirsch, W. M., 78-6486
 Kirven, R., 78-6360
 Kleihues, P., 78-6221
 Klein, G., 78-6314, 78-6456
 Kniazev, P. G., 78-6399
 Knights, E. M., 78-6377
 Knudsen, I., 78-6007
 Kobayashi, K., 78-6531
 Kogan, A. Kh., 78-6332
 Kohn, K. W., 78-6366
 Kohn, R. R., 78-6489
 Kokatnur, M. G., 78-6204
 Kokosha, L. V., 78-6461
 Kolar, G. F., 78-6221
 Kolar, M. C., 78-6132
 Kolesnichenko, T. S., 78-6183
 Komarov, H., 78-6344
 Komorowski, R. A., 78-6363
 Kondo, H., 78-6212
 Kondo, K., 78-6425
 Konishi, Y., 78-6212
 Korobitsyn, L. P., 78-6399
 Korsak, R. K., 78-6545
 Koshimizu, K., 78-6239
 Kosuge, T., 78-6198, 78-6201, 78-6246
 Kouyoumian, J., 78-6089
 Kramers, P. G., 78-6234
 Krass, P. M., 78-6278
 Krepinsky, J., 78-6229
 Krieg, C. J., 78-6428, 78-6429
 Kringstad, K., 78-6132
 Krishna, G., 78-6109
 Krishnan, E. C., 78-6504
 Krueger, G. R., 78-6432
 Krzysek, R., 78-6474
 Kudisch, M., 78-6351
 Kuff, E. L., 78-6472
 Kulakis, C., 78-6107
 Kumagai, M., 78-6531
 Kumaokas, S., 78-6544
 Kumar, S., 78-6566
 Kumar, V., 78-6427
 Kurland, J. I., 78-6488
 Kurokawa, T., 78-6464, 78-6465
 Kurth, R., 78-6402
 Kyner, D. A., 78-6205
 Laamanen, I., 78-6253
 Laerum, O. D., 78-6231
 Laishes, B. A., 78-6492
 Lalovic, D. I., 78-6154
 Lambert, R., 78-6588
 Lampe, J., 78-6020
 Lancaster, W. D., 78-6445
 Lane, M. A., 78-6409
 Lange, B., 78-6452
 Langer, A. M., 78-6529
 Langlands, A. O., 78-6505
 Langman, M. J., 78-6565
 Lankas, G. R., 78-6295
 Lapin, B. A., 78-6461
 Larson, V. M., 78-6442
 Larsson, L. I., 78-6533
 Laub, O., 78-6485
 LaVoie, E., 78-6323
 Lawler, S. D., 78-6151
 Lawton, A. R., 78-6078
 Laychock, A., 78-6499
 Le, M., 78-6068
 Lebedev, V. N., 78-6461
 Lebherz, W. B., 78-6104
 LeBon, J. M., 78-6582
 Leclercq, M., 78-6088
 Lee, A. C., 78-6516
 Lee, C. G., 78-6228
 Lee, L. S., 78-6306
 Lehman, E. D., 78-6442
 Lehtola, J., 78-6575
 Lekevichus, R., 78-6139
 Lemen, R. A., 78-6014
 Lerner, R. A., 78-6097, 78-6436
 78-6439
 Leunen, J., 78-6088
 Levey, G. S., 78-6282
 Levine, P. H., 78-6459, 78-6552
 Levitz, M., 78-6307
 Levy, J. P., 78-6431
 Lewin, M. R., 78-6180
 Lieberman, M., 78-6418
 Lieberman, M. W., 78-6191, 78-6357
 Liem, H. H., 78-6319
 Lijinsky, W., 78-6240
 Lilja, H. S., 78-6200
 Lilly, F., 78-6421
 Lilly, L. J., 78-6214
 Lindberg, S., 78-6379
 Lindquist, N. G., 78-6177
 Lindsay, R. J., 78-6238
 Lippman, M., 78-6024
 Lippman, M. E., 78-6355
 Lipson, R. M., 78-6354
 Litvitskii, P. F., 78-6332
 Lofroth, G., 78-6129
 Loke, Y. W., 78-6591

Lombardi, B., 78-6199
 Lombardi, P., 78-6348
 Long, J. C., 78-6219
 Longnecker, D. S., 78-6200
 Lopez, J. C., 78-6158
 Lopez, D. M., 78-6497
 Loprieno, N., 78-6116
 Low, H., 78-6018
 Lueders, K. K., 78-6472
 Luepker, R. V., 78-6560
 Lum, K., 78-6527
 Lumb, G., 78-6016
 Lundqvist, G., 78-6533
 Lutz, W. K., 78-6345
 Luyten-Kellermann, M., 78-6292
 Lynch, H., 78-6538
 Lynch, H. T., 78-6066, 78-6349
 Lynch, J., 78-6538
 Lynch, J. F., 78-6349
 Lynch, P., 78-6538
 Lynch, P. M., 78-6066
 Lyon, J. L., 78-6559
 Lyon, M., 78-6475
 Macaulay, R. A., 78-6505
 MacDonald, D. G., 78-6316
 Macdonald, I. A., 78-6341
 Macdonald, J. S., 78-6355, 78-6509
 MacDonald, W. E., 78-6261
 MacEwen, E. G., 78-6511
 MacFarlane, J., 78-6179
 MacGillivray, A. J., 78-6456
 Machida, I., 78-6244
 MacKay, B., 78-6539
 MacLennan, R., 78-6572
 Maeda, K., 78-6380
 Magnier, M., 78-6156
 Maguire, J. H., 78-6331
 Maharam, L. C., 78-6280
 Mahdavi, V., 78-6339
 Mahony, D. E., 78-6341
 Maier, P., 78-6227
 Majumdar, S. K., 78-6280
 Mak, S., 78-6469
 Malec, E., 78-6578
 Maloney, K., 78-6349
 Mammerickx, M., 78-6088
 Mandeville, R., 78-6394
 Mantel, N., 78-6243
 Manton, K. G., 78-6555
 Maraud, R., 78-6248
 Markarian, D. S., 78-6461
 Markova, T. P., 78-6461
 Marks, P. A., 78-6426
 Martin, P. M., 78-6350
 Martz, F., 78-6257
 Mason, T. J., 78-6556
 Mastromarino, A. J., 78-6587
 Matheson, D., 78-6144
 Mathews, J. G., 78-6220
 Matka, M., 78-6266
 Matsubara, N., 78-6189
 Matsukura, N., 78-6201
 Matsumoto, T., 78-6239
 Matsunaga, E., 78-6071
 Matsuo, T., 78-6455
 Matsuo, H., 78-6380
 Matsushima, T., 78-6173, 78-6263
 Matsuzawa, A., 78-6425
 Matteucci, D., 78-6430
 Maugh, T. H., 78-6004
 Maurer, P. H., 78-6416
 Mayer, A., 78-6421
 Mays, C. W., 78-6564
 Mazzaresse, R., 78-6323
 McCabe, L. J., 78-6556
 McCallum, H. M., 78-6381
 McCann, J., 78-6005
 McCaw, B. A., 78-6313
 McChesney, E., 78-6294
 McClure, S., 78-6172
 McConnell, M. R., 78-6593
 McCoy, G. D., 78-6237
 McCredie, K. B., 78-6600
 McDevitt, H. O., 78-6098
 McDonald, J. J., 78-6249
 McDowell, E. M., 78-6528, 78-6599
 McGarry, M. P., 78-6424
 McGrath, C. M., 78-6411
 McGuire, W. L., 78-6350
 McIlmurray, M. B., 78-6565
 McMillin-Helsel, C., 78-6393
 McQueen, E. G., 78-6026
 Medina, D., 78-6315
 Medrano, J., 78-6384
 Meier, H., 78-6417
 Meinhardt, T. J., 78-6014
 Meinkoth, J. L., 78-6404
 Melamed, M. R., 78-6374
 Melzer, M. S., 78-6054
 Menczer, J., 78-6344
 Menezes, J., 78-6451
 Mennel, H. D., 78-6581
 Menzies-Gow, N., 78-6353
 Merezko, V. A., 78-6532
 Merkat, I. R., 78-6506
 Merlo, D. J., 78-6583
 Merryman, C. F., 78-6416
 Meruelo, D., 78-6098
 Meschini, A., 78-6493
 Messens, E., 78-6585
 Messripour, M., 78-6125
 Meyer, G., 78-6092
 Mi, M. P., 78-6546
 Mickey, M. R., 78-6459
 Miima, J. G., 78-6532
 Mikami, T., 78-6521
 Miller, A. B., 78-6067
 Miller, D., 78-6552
 Miller, E. C., 78-6185, 78-6247
 Miller, J. A., 78-6185, 78-6247
 Miller, W. R., 78-6343
 Mills, K. T., 78-6346
 Milo, G. E., 78-6242, 78-6245, 78-6440
 Minowada, J., 78-6494
 Mitelman, F., 78-6507
 Miyahara, H., 78-6380
 Miyamoto, M. J., 78-6527
 Mizuno, F., 78-6455
 Mizuno, M., 78-6425
 Mizusaki, S., 78-6239
 Mohammadi, J., 78-6125
 Moloney, W. C., 78-6219
 Mondal, S., 78-6283
 Mondino, A., 78-6116
 Montelaro, R. C., 78-6094
 Montell, L., 78-6368
 Mooney, S., 78-6511
 Moore, C. F., 78-6594
 Moore, D. H., 78-6416
 Moore, M. A., 78-6488
 Moore, W. E., 78-6033
 Mordelet-Dambrine, M., 78-6156
 Moreland, F. M., 78-6123
 Morgan, R. W., 78-6359, 78-6360
 78-6543
 Moriwaki, K., 78-6425
 Moron, M. S., 78-6320
 Morse, E. E., 78-6141
 Mosbech, J., 78-6572
 Moshakis, V., 78-6293
 Mosher, D. F., 78-6057
 Moszczynski, P., 78-6258
 Mountain, I. M., 78-6249
 Mughal, S., 78-6536
 Mulder, G. J., 78-6291
 Muller-Eberhard, U., 78-6319
 Muller, H., 78-6477
 Muller, R. E., 78-6351
 Murakami, M., 78-6305
 Murasko, D. M., 78-6501
 Murata, I., 78-6289
 Murnik, M. R., 78-6268
 Murray, D., 78-6179
 Murray, M. L., 78-6204
 Murthy, D. K., 78-6255
 Musto, J., 78-6451
 Nadasdi, L., 78-6241
 Nadworny, H. A., 78-6277
 Nagao, M., 78-6198, 78-6201, 78-6232
 Nagao, T., 78-6263
 Nagy, A., 78-6184
 Nakai, H., 78-6203
 Nakai, S., 78-6244
 Nakamura, T., 78-6289
 Nakasuji, K., 78-6289
 Nakoinz, I., 78-6419
 Nambaru, S., 78-6251
 Nandi, S., 78-6050
 Naparstek, J., 78-6513
 Naroditskii, B. S., 78-6093
 Narui, K., 78-6206
 Naseem, S. M., 78-6169
 Nash, G., 78-6168
 Nasrallah, A. G., 78-6424
 Natarajan, A. T., 78-6358
 National Cancer Institute, 78-6114
 Natoli, J., 78-6280
 Naujack, K. W., 78-6301
 Nazarenko, S. A., 78-6226
 Nebert, D. W., 78-6319
 Neeffe, J. R., 78-6495
 Nehez, M., 78-6157
 Neiman, P. E., 78-6393
 Nelson, D. S., 78-6490
 Nelson, M., 78-6490
 Nelson, R. L., 78-6161
 Nes, I. F., 78-6371
 Nestmann, E. R., 78-6304

- Neubauer, R. H., 78-6450
 Neumann, H., 78-6457
 Newberne, P., 78-6016
 Newell, D. J., 78-6550
 Newell, G. W., 78-6138
 Newman, M. S., 78-6015
 Newton, M. F., 78-6214
 Nias, A. H., 78-6381
 Nicolson, G. L., 78-6060
 Nigro, N. D., 78-6106
 Nii, S., 78-6407
 Niiyama, Y., 78-6464
 Niman, H. L., 78-6441
 Nishimura, K., 78-6531
 Niskanen, A., 78-6181
 Nissen-Meyer, J., 78-6371
 Nomura, A., 78-6126
 Nonoyama, M., 78-6455
 Notani, N. K., 78-6269
 Nurmiäho, E. L., 78-6388
 O'Berg, M. T., 78-6551
 O'Connor, J., 78-6349
 Obalek, S., 78-6090
 Obe, G., 78-6188, 78-6358
 Odaka, T., 78-6425
 Oehme, J., 78-6196
 Oesch, F., 78-6166, 78-6302
 Ogawa, H., 78-6412
 Ohbuchi, M., 78-6380
 Ohgi, T., 78-6521
 Ohmori, M., 78-6471
 Ohno, S., 78-6314
 Okada, H., 78-6503
 Okada, K., 78-6521
 Okada, N., 78-6503
 Okamoto, T., 78-6198, 78-6246
 Okenquist, S., 78-6403
 Oker-Blom, N., 78-6388
 Olbe, L., 78-6533
 Oldham, P. D., 78-6550
 Olofsson, H., 78-6368
 Olson, C., 78-6445, 78-6446
 Olsson, S., 78-6177
 Ono, T., 78-6422
 Onuma, M., 78-6521
 Ornaf, R. M., 78-6176
 Oroszlan, S., 78-6443
 Orth, G., 78-6090, 78-6091
 Ortiz-Muniz, G., 78-6497
 Osato, T., 78-6455
 Osawa, M., 78-6251
 Osborn, R. C., 78-6527
 Oskarsson, M. K., 78-6439
 Osman, S. F., 78-6586
 Ost, A., 78-6576
 Ostertag, W., 78-6428, 78-6429
 Ostrow, P. T., 78-6365
 Ou, S. Y., 78-6328, 78-6333
 Oud, J. L., 78-6298
 Owens, I. S., 78-6319
 Pai, K. J., 78-6321
 Painter, R. B., 78-6361
 Pakula, H., 78-6581
 Paldy, A., 78-6157
 Paloyan, D., 78-6354
 Panda, B. B., 78-6284
 Pandey, K. N., 78-6170
 Paonessa, D. F., 78-6089
 Parks, W., 78-6497
 Parmiani, G., 78-6493
 Parodi, S., 78-6053
 Parsons, P. G., 78-6362
 Partington, M. W., 78-6518
 Parulekar, S. S., 78-6089
 Patel, P., 78-6451
 Patnaik, A. K., 78-6511
 Paton, D., 78-6260
 Patt, L. M., 78-6592
 Patton, J. T., 78-6447
 Pauley, R. J., 78-6315, 78-6410
 Pavlenko-Mikhailov, Iu. N., 78-6382
 Payne, B. J., 78-6133
 Pearson, G. R., 78-6460
 Pedrinan, L., 78-6310
 Peebles, P. T., 78-6437
 Pegg, A. E., 78-6213
 Pell, S., 78-6551
 Pels, E., 78-6082
 Pelus, L.M., 78-6488
 Penn, L., 78-6076
 Perevozchikov, A. P., 78-6399
 Perry, P. M., 78-6367
 Pesch, H. J., 78-6375
 Peters, E. L., 78-6123
 Peters, P. W., 78-6298
 Petersson, B. G., 78-6537
 Peto, R., 78-6110
 Petrilli, F. L., 78-6130
 Petrucci, P. P., 78-6355
 Petzold, G. L., 78-6162
 Pezzuto, J. M., 78-6326
 Pfeiffer, E. H., 78-6330
 Pfister, H., 78-6476
 Phillips, E., 78-6377
 Pienta, R. J., 78-6104
 Pilotti, A., 78-6177
 Pinter, A., 78-6241
 Plaude, O. O., 78-6318
 Pliss, G. B., 78-6003, 78-6372
 Pogosiants, E. E., 78-6073
 Pohl, L. R., 78-6109
 Poiley, J. A., 78-6104
 Pool, B. L., 78-6223
 Poole, D., 78-6118
 Port, C. D., 78-6335
 Porter, M. W., 78-6373
 Portetelle, D., 78-6088
 Pottathil, R., 78-6417
 Powers, G. J., 78-6127
 Pragani, B., 78-6275
 Pragnell, I. B., 78-6428, 78-6429
 Prasad, U., 78-6064
 Prehn, R. T., 78-6501
 Pritchard, C., 78-6447
 Prosser, J. S., 78-6235
 Provana, A., 78-6547
 Prpic-Majic, D., 78-6135
 Purnell, D. M., 78-6525
 Purohit, M. G., 78-6299
 Pyysalo, H., 78-6181
 Qualtiere, L. F., 78-6460
 Quick, C. A., 78-6474
 Quigley, J. P., 78-6395
 Rabin, H., 78-6450
 Radtke, H. E., 78-6264
 Ragnhult, I., 78-6379
 Rajalakshmi, S., 78-6215, 78-6229
 Ralph, P., 78-6419
 Rambousek, V., 78-6266
 Ramel, C., 78-6113, 78-6132
 Rannug, U., 78-6113, 78-6132
 Rao, E. V., 78-6570
 Rao, P. M., 78-6215, 78-6229
 Rao, T. K., 78-6102
 Rapoport, A., 78-6089
 Rapp, F., 78-6499
 Rappaport, S. M., 78-6279
 Raptis, L., 78-6462
 Raschke, W. C., 78-6419
 Rashad, M. N., 78-6545, 78-6546
 Raskas, H. J., 78-6463
 Rasmuson, A., 78-6368
 Rasmuson, B., 78-6368
 Rasmussen, J. E., 78-6017
 Rassoulzadegan, M., 78-6478
 Ratovitskii, E. A., 78-6399
 Ravisse, P., 78-6568
 Rawlings, V., 78-6037
 Rawson, R. W., 78-6030
 Reddy, B. S., 78-6297, 78-6587
 Reddy, P. V., 78-6255
 Reddy, T. P., 78-6134
 Rehfeld, J. F., 78-6533
 Reich, E., 78-6338
 Reichard, P., 78-6482
 Reid, B. L., 78-6569
 Reid, J. B., 78-6327
 Reid, L. B., 78-6506
 Rein, A., 78-6433
 Reinhardt, C. F., 78-6270
 Resnick, P., 78-6427
 Reuben, R. C., 78-6426
 Reuber, M. D., 78-6011, 78-6275
 Reynolds, R. K., 78-6420
 Rhomberg, W., 78-6563
 Richards, M. J., 78-6517
 Rickwood, D., 78-6456
 Rifkind, R. A., 78-6426
 Riley, V., 78-6058
 Rinsky, R. A., 78-6014
 Ristow, H., 78-6188
 Ritter, J., 78-6196
 Roberts, J. J., 78-6313
 Roesler, G., 78-6429
 Rohner, T. J., 78-6499
 Roninson, I. B., 78-6093
 Roos, E., 78-6526
 Roper, H., 78-6225
 Rosen, J. M., 78-6410
 Rosenfeld, R. S., 78-6349
 Rosenthal, K. S., 78-6458
 Rosenthal, L. J., 78-6582
 Rossow, P. W., 78-6593
 Rostom, A. Y., 78-6385
 Roudebush, C. P., 78-6376
 Rushbrook, C. J., 78-6138
 Russell, S. W., 78-6502
 Russo, A., 78-6317

Aust, J. H., 78-6016
 Ayman, D. S., 78-6350
 Bzesa, G., 78-6090
 Babbarwal, P. S., 78-6170
 Badaie, Y., 78-6206
 Bade, J., 78-6520
 Baeki, T., 78-6244
 Bafaii, H. S., 78-6540
 Bahu, R. K., 78-6284
 Bakamoto, S., 78-6449
 Baknyn, A. V., 78-6143
 Baksun, J. M., 78-6519
 Balerno, R. A., 78-6442
 Balm, R., 78-6510
 Balwa, J., 78-6013
 Bancho-Garnier, H., 78-6068
 Banders, C. L., 78-6127
 Bandison, A. G., 78-6567
 Bandstrom, A., 78-6288
 Banford, E. J., 78-6499
 Bangulia, I. A., 78-6461
 Bano, T., 78-6561
 Baprin, A. N., 78-6218
 Bargent, E. N., 78-6383
 Barma, D. S., 78-6215, 78-6229
 Basada, R., 78-6464
 Bato, T., 78-6380
 Bauerbier, W., 78-6579
 Savage, J. R., 78-6290
 Bave-Soderberg, J., 78-6533
 Sawamura, N., 78-6263
 Schaffhausen, B., 78-6479
 Schain, M., 78-6537
 Schat, K. A., 78-6406
 Schein, P. S., 78-6355, 78-6509
 Schell, J., 78-6584, 78-6585
 Schidlovsky, G., 78-6413
 Schindler, J., 78-6436
 Schlatter, C., 78-6345
 Schlom, J., 78-6413
 Schmeltz, I., 78-6009, 78-6010
 Schmid, E., 78-6136
 Schmoll, H. J., 78-6563
 Schneider, C. J., 78-6146
 Schneider, R., 78-6548
 Schnitzler, E. R., 78-6517
 Schoeny, R. S., 78-6158
 Schoon, I. M., 78-6533
 Schor, N. A., 78-6334
 Schrauzer, G. N., 78-6146
 Schuler, D., 78-6051
 Schurch, W., 78-6528, 78-6599
 Schuttman, W., 78-6049
 Schwind, H., 78-6166
 Sciaky, D., 78-6583
 Sciotto, C. G., 78-6489
 Scolnick, E. M., 78-6423, 78-6472
 Sears, E., 78-6518
 Sebahoun, G., 78-6209
 Seide-Kehoe, R., 78-6205
 Seidegard, J., 78-6320
 Seif, R., 78-6478
 Seiler, J. P., 78-6303
 Seino, Y., 78-6198, 78-6201, 78-6232
 78-6263
 Seits, I. F., 78-6399

Sekikawa, K., 78-6467
 Selikoff, I. J., 78-6023, 78-6169
 78-6529
 Selypes, A., 78-6157
 Senior, B., 78-6540
 Severson, R. F., 78-6337
 Shankel, D. M., 78-6159
 Sharma, C. B., 78-6284
 Sharma, O. K., 78-6347
 Shaub, M., 78-6383
 Sheiness, D., 78-6387
 Shellabarger, C. J., 78-6342
 Shenbom, M., 78-6344
 Sheridan, R. B., 78-6364
 Sheridan, W., 78-6222
 Sherman, H., 78-6270
 Sherr, C. J., 78-6444
 Shevchenko, N. L., 78-6532
 Shih, T. W., 78-6115
 Shih, T. Y., 78-6423
 Shimada, T., 78-6514
 Shimizu, H., 78-6182, 78-6254
 Shimojo, H., 78-6467
 Shinohara, M., 78-6535
 Shinozuka, H., 78-6199
 Shipley, E. A., 78-6039
 Shirai, A., 78-6263
 Shioki, K., 78-6467
 Shudo, K., 78-6198, 78-6246
 Shulman, L. E., 78-6197
 Shultz, T. D., 78-6174
 Siegel, B. V., 78-6077
 Sigel, M. M., 78-6497
 Siiteri, P. K., 78-6028
 Silva, O. L., 78-6594
 Silver, J., 78-6479
 Simmon, V. F., 78-6118
 Simonowitz, D., 78-6354
 Simpson, J. L., 78-6506
 Simpson, N. E., 78-6518
 Singer, S. J., 78-6396
 Singh, P. K., 78-6271
 Singh, S. P., 78-6137
 Slapikoff, S. A., 78-6164
 Sliski, A., 78-6444
 Smerdon, M. J., 78-6357
 Smiley, J. R., 78-6469
 Smith, B. R., 78-6331
 Smith, C. E., 78-6174
 Smith, C. J., 78-6277
 Smith, G. H., 78-6315, 78-6415
 Smith, K. C., 78-6052
 Smith, M. L., 78-6560
 Smith, R. B., 78-6153
 Snider, R. H., 78-6594
 Snook, M. E., 78-6171, 78-6337
 Soboll, H., 78-6470
 Socher, S. H., 78-6315, 78-6410
 Sohler, R., 78-6475
 Sorenson, A. W., 78-6559
 Sorsa, M., 78-6253
 Soto, E. A., 78-6321
 Spector, D. H., 78-6392
 Spector, W. G., 78-6079
 Spiess, H., 78-6564
 Spiller, G. A., 78-6039, 78-6040

Spira, J., 78-6314
 Spitzer, G., 78-6600
 Sporn, M. B., 78-6273
 Squire, R. A., 78-6273
 Stallard, E., 78-6555
 Stambolis, C., 78-6384
 Starek, A., 78-6258
 Stavrou, D., 78-6230
 Steel, G. G., 78-6385
 Stellwag, E. J., 78-6590
 Stemmerman, G., 78-6544
 Stemmermann, G. N., 78-6126
 Stephenson, J. R., 78-6420
 Sternson, L. A., 78-6103
 Steuart, C. D., 78-6174
 Stevens, J. G., 78-6103
 Stich, H. F., 78-6099
 Stinson, E. E., 78-6586
 Stinson, S. F., 78-6273
 Stockbrugger, R., 78-6533
 Stoll, R., 78-6248
 Stollmann, B., 78-6384
 Stolte, L. A., 78-6573
 Stone, J. P., 78-6342
 Stortebeker, T. P., 78-6034
 Stossel, T. P., 78-6219
 Stout, E. R., 78-6447
 Stowers, D. J., 78-6525
 Strand, M., 78-6437
 Strel'tsova, V. N., 78-6382
 Strickland, S., 78-6339
 Strominger, J. L., 78-6458
 Struck, R. F., 78-6115
 Stuart, A. E., 78-6505
 Stukiene, R., 78-6139
 Stula, E. F., 78-6270
 Stupfel, M., 78-6156
 Sturelid, S., 78-6120
 Styles, J. A., 78-6165, 78-6260
 Subramanyam, S., 78-6255
 Sugimoto, C., 78-6514
 Sugimoto, T., 78-6531
 Sugimura, T., 78-6173, 78-6198
 78-6201, 78-6232, 78-6246, 78-6263
 Sugino, Y., 78-6464, 78-6465
 Sugioka, G., 78-6531
 Sugiura, K., 78-6272
 Summitt, R. L., 78-6506
 Sunagawa, M., 78-6212
 Sundaram, K., 78-6108
 Sunderman, F. W., 78-6141
 Suppnick, J. D., 78-6106
 Surjan, A., 78-6241
 Susanne, C., 78-6140
 Sutter, D., 78-6470
 Sutter, V. L., 78-6035
 Suzangar, M., 78-6125
 Suzuki, K., 78-6531
 Suzuki, Y., 78-6529
 Svahlin, H., 78-6368
 Sviridova, I. K., 78-6468
 Swenberg, J. A., 78-6162
 Swetly, P., 78-6428, 78-6429
 Szuts, T., 78-6177
 Szymas, J. S., 78-6581
 Tabuchi, K., 78-6486

- Takahashi, S., 78-6212
 Takamori, K., 78-6535
 Takayama, S., 78-6173
 Takeda, K., 78-6198, 78-6246
 Takemura, N., 78-6182, 78-6254
 Talalay, P., 78-6333
 Tamura, M., 78-6581
 Tanaka, A., 78-6251, 78-6455
 Tanaka, H., 78-6412
 Tanooka, H., 78-6224
 Tardiff, R., 78-6118
 Tardiff, R. G., 78-6158
 Taylor, G., 78-6566
 Taylor, H. W., 78-6240
 Taylor, S., 78-6522
 te Velde, E. R., 78-6095
 Teerenhovi, L., 78-6507
 Teichmann, B., 78-6145
 Teller, M. N., 78-6250
 Teramoto, Y. A., 78-6413
 Terasaki, P. I., 78-6459
 Thatcher, J. D., 78-6280
 The, T. H., 78-6095
 Thielmann, H. W., 78-6308, 78-6325
 Thiess, A. M., 78-6187, 78-6252
 Thilly, W. G., 78-6164
 Thomas, D. B., 78-6027
 Thomas, H. F., 78-6032
 Thomassen, R. W., 78-6516
 Thumfart, W., 78-6375
 Till, J. E., 78-6386
 Timanovskaya, V. V., 78-6461
 Timonen, T., 78-6507
 Tischler, A. S., 78-6063
 Tisell, L. E., 78-6379
 Tishler, P. V., 78-6169
 Tlsty, T. D., 78-6191
 Tobioka, H., 78-6259
 Todaro, G., 78-6413
 Todaro, G. J., 78-6444, 78-6481
 Tohya, M., 78-6380
 Tokieda, T., 78-6251
 Tolley, H. D., 78-6555
 Tomita, H., 78-6239
 Toniolo, A., 78-6430
 Torok, G., 78-6241
 Troll, W., 78-6307
 Trosko, J. E., 78-6008, 78-6155
 Trout, E. C., 78-6274
 Trump, B. F., 78-6528, 78-6599
 Trzos, R. J., 78-6162
 Tsakeres, F., 78-6473
 Tso, T. C., 78-6176
 Tsuji, K., 78-6198, 78-6201
 Tsukamoto, K., 78-6464
 Tubiana, M., 78-6025
 Turner, J. V., 78-6236
 Tuszyński, G. P., 78-6056
 Tuyns, A. J., 78-6568
 Twentyman, P. R., 78-6598
 Uchino, H., 78-6514
 Ueda, N., 78-6317
 Ullberg, S., 78-6177
 Umezawa, K., 78-6173
 Vaheri, A., 78-6057
 Vaidyanath, K., 78-6134
 Vainio, H., 78-6253
 Vakil, D. V., 78-6543
 Van Buskirk, J. J., 78-6486
 van de Ven, W. J., 78-6420
 Van Dongen, C. G., 78-6321
 Van Loveren, H., 78-6082
 Van Montagu, M., 78-6584, 78-6585
 Vande Woude, G. F., 78-6439
 Varmus, H. E., 78-6391, 78-6392
 Veljkovic, V., 78-6154
 Venitt, S., 78-6151
 Venkatesan, N., 78-6195
 Verma, D. S., 78-6600
 Verma, I. M., 78-6423
 Verschaeve, L., 78-6140
 Vesely, D. L., 78-6282
 Viglianti, G. A., 78-6280
 Vijanskaja, T., 78-6139
 Visek, W. J., 78-6042
 Vlodavskii, E. A., 78-6532
 Vogler, L. B., 78-6078
 Vogt, P. K., 78-6396
 Volkert, M. R., 78-6045
 Volkova, A. I., 78-6286
 von Wright, A., 78-6181
 Voogd, C. E., 78-6285
 Vu Van, H., 78-6211
 Vuopio, P., 78-6507
 Vuori, H., 78-6572
 Wade, C. G., 78-6329
 Wagner, B., 78-6016
 Wagner, S., 78-6243
 Wainberg, M. A., 78-6394
 Wakabayashi, K., 78-6201
 Walker, A., 78-6381
 Walker, A. R., 78-6041
 Walker, J. R., 78-6597
 Waller, G., 78-6375
 Wang, D. Y., 78-6544
 Wang, L. H., 78-6087
 Ward, A. D., 78-6236
 Warren, L., 78-6056
 Warren, S., 78-6373
 Watanabe, K., 78-6193, 78-6297
 Watanabe, M., 78-6311
 Waterhouse, J. A., 78-6002
 Webb, G. R., 78-6341
 Weber, W. W., 78-6264
 Wei, L., 78-6099
 Weidenbecher, M., 78-6375
 Weimann, B. J., 78-6428, 78-6429
 Weinberg, A. G., 78-6530
 Weinberger, M. A., 78-6123
 Weinstein, I. B., 78-6096, 78-6306
 Weisburger, E. K., 78-6190
 Weisburger, J. H., 78-6107
 Weiss, S. R., 78-6391
 Westermarck, H., 78-6388
 Whalen, C. E., 78-6310
 Wheldon, T. E., 78-6381
 White, C. M., 78-6235
 White, D. A., 78-6146
 Whiting, R., 78-6099
 Whyte, A., 78-6591
 Wiebkin, P., 78-6267
 Wiener, F., 78-6314
 Wiggans, R. G., 78-6509
 Williams, G. M., 78-6193
 Williamson, R. C., 78-6178
 Wilson, E. L., 78-6338
 Wilson, P., 78-6566
 Winton, W., 78-6102
 Wise, A., 78-6125
 Wolf, B. A., 78-6397, 78-6398
 Wong, T. C., 78-6405
 Woodard, H. Q., 78-6374
 Woods, F., 78-6261
 Woolford, G., 78-6203
 Woolley, P. V., 78-6355, 78-6509
 Wotiz, H. H., 78-6351
 Wozniak, K. D., 78-6512
 Wyke, J. A., 78-6402
 Wynder, E. L., 78-6009, 78-6237
 78-6587
 Yager, J. D., 78-6200
 Yagi, M. J., 78-6409
 Yahagi, T., 78-6198, 78-6201, 78-6232
 78-6263
 Yakovenko, K. N., 78-6226
 Yamaguchi, K., 78-6198
 Yamaha, T., 78-6251
 Yamamoto, K., 78-6455
 Yamamoto, N., 78-6454
 Yamamoto, T., 78-6198
 Yamanaka, S., 78-6272
 Yamauchi, Y., 78-6535
 Yanovich, S., 78-6458
 Yasuda, I., 78-6407
 Yohn, D. S., 78-6242
 Yonemichi, H., 78-6521
 Yoshida, D., 78-6239
 Yoshida, M., 78-6207
 Yoshida, Y., 78-6207
 Yoshikawa, K., 78-6244
 Yoshino, G., 78-6207, 78-6208
 Youness, E., 78-6508
 Young, J. A., 78-6102
 Young, R. J., 78-6014
 Young, S., 78-6313
 Zabos, P., 78-6205
 Zajdela, F., 78-6166
 Zaldivar, R., 78-6574
 Zalmanzon, E. S., 78-6466
 Zapp, J. A., 78-6270
 Zetterberg, G., 78-6287
 Zhdanov, V. M., 78-6086
 Zilkley, B., 78-6172
 Zimmer, S. G., 78-6463
 Zornoza, J., 78-6549
 Zulch, K. J., 78-6581
 Zusman, R. M., 78-6219
 Zverina, Z., 78-6266

Subject Index

tion
 Uterine Neoplasms
 Endometriosis, 78-6541
 Naphthalene, 5-Nitro-
 Ames Test
 Mutagenic Activity, 78-6254
 Cholangioma
 Carcinogenic Potential, 78-6254
 Ear Neoplasms
 Carcinoma, Epidermoid, 78-6254
 Intestinal Neoplasms
 Adenocarcinoma, 78-6254
 Kidney Neoplasms
 Carcinogenic Potential, 78-6254
 Liver Neoplasms
 Carcinogenic Potential, 78-6254
 Mammary Neoplasms, Experimental
 Carcinoma, Ductal, 78-6254
 Rhabdomyosarcoma
 Carcinogenic Potential, 78-6254
 Aldehyde
 Chromatids
 Lymphocytes, 78-6188
 Chromosomes
 Mutagenic Activity, 78-6188
 DNA
 Chromosomes, 78-6188
 Cross-Linking Activity, 78-6188
 Aldehyde, Bromo-
 Ethane, 1,2-Dibromo-
 Carcinogenic Metabolite, 78-6115
 Aldehyde, Chloro-
 Deoxyribonucleosides
 Imidazo-Cyclization Products, 78-6112
 DNA
 Liver, Rat, 78-6112
 Amide, *N*-(5-*p*-Acetamidophenylthiophen-2-yl)-
 Ames Test
 Mutagenic Activity, 78-6260
 Cells, Cultured
 Cell Transformation, Neoplastic, 78-6260
 Amide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 Chromatin
 DNA Repair, 78-6191
 DNA Repair
 Deoxyribonuclease, 78-6191
Escherichia coli
 DNA Repair, 78-6308
 Fibroblasts
 DNA Repair, 78-6191
 12-*O*-Tetradecanoylphorbol-13-acetate
 DNA Repair, 78-6308
 Virus, Feline Sarcoma
 Reverse Transcriptase, 78-6440
 Virus Replication, 78-6440
 Virus, SV40
 Cell Transformation, Neoplastic, 78-6242
 Xeroderma Pigmentosum
 DNA Repair, Review, 78-6055

Acetamide, *N*-Fluoren-2-yl-
 Carcinogenic Metabolite
 Quantitation Method, 78-6190
 Cholesterol
 Metabolism, Liver, 78-6192
 Chromosome Aberrations
 Micronucleus Test, 78-6162
 Glutamyl Transpeptidase
 Hepatocyte, Rat, 78-6492
 Hepatocytes
 Cytotoxicity, Fibroblasts, 78-6267
 Liver Neoplasms
 Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6193
 Precancerous Conditions, 78-6492
 Quinone Reductases
 Liver, Rat, 78-6334
 Sulfate Derivative
 Proteins, Binding, 78-6291
 Transplantation, Homologous
 Hepatocyte, Rat, 78-6492
 Acetamide, *N*-Fluoren-4-yl-
 Hepatocytes
 Cytotoxicity, Fibroblasts, 78-6267
 Acetamide, *N*-(5-Phenylthiophen-2-yl)-
 Ames Test
 Mutagenic Activity, 78-6260
 Cells, Cultured
 Cell Transformation, Neoplastic, 78-6260
 Acetamide, Thio-
 DNA Repair
 Liver, Rat, 78-6216
 Estradiol, 3-Benzate
 Ovalbumin, 78-6347
 Progesterone, 78-6347
 Ovalbumin
 Plasma Levels, Chick, 78-6347
 Progesterone
 Plasma Levels, Chick, 78-6347
 Acetanilide
 Lymphocytes
 Aryl Hydrocarbon Hydroxylases, 78-6292
 Plasma
 Half-Life, 78-6292
 Acetanilide, 4'-Hydroxy-
 Microsomes, Liver
 Metabolism, 78-6291
 Acetic Acid, (4-Chloro-2-methylphenoxy)-
 see Acetic Acid, ((4-Chloro-*o*-tolyl)oxy)-
 Acetic Acid, ((4-Chloro-*o*-tolyl)oxy)-
 Saccharomyces cerevisiae
 Mutagenic Activity, 78-6287
 Acetic Acid, (2,4-Dichlorophenoxy)-
 Cells, Cultured
 Metabolism, Lung, 78-6305
 Mutagenic Activity
 Blue-Green Algae, 78-6271
 Saccharomyces cerevisiae
 Mutagenic Activity, 78-6287
 Sarcoma
 Occupational Hazard, 78-6288

- Acetic Acid, (Ethylenebis(oxyethylenenitrilo))tetra-**
Lymphoma
Rosette Formation, AKR Mouse, 78-6503
Thymoma
Rosette Formation, AKR Mouse, 78-6503
- Acetic Acid, (2,4,5-Trichlorophenoxy)-**
Cells, Cultured
Metabolism, Lung, 78-6305
Saccharomyces cerevisiae
Mutagenic Activity, 78-6287
Sarcoma
Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-, 78-6288
Occupational Hazard, 78-6288
- p*-Acetophenetidide, *N*-Hydroxy-**
Glucuronate Derivative
Proteins, Binding, 78-6291
Microsomes, Liver
Metabolism, 78-6291
Quinone, *N*-Acetylimido-
Sulfate Derivative, 78-6291
Sulfate Derivative
Proteins, Binding, 78-6291
- p*-Acetophenetidide**
Immunosuppression
Carcinogenic Activity, Review, 78-6051
Kidney Neoplasms
Carcinoma, Transitional Cell, 78-6293
Case Report, 78-6293
Lymphocytes
Aryl Hydrocarbon Hydroxylases, 78-6292
Microsomes, Liver
Metabolism, 78-6291
Plasma
Half-Life, 78-6292
- Acetophenone**
Smoke Condensate
Isolation and Characterization, 78-6171
- Acetophenone, 2-Amino-**
Leukemia
Species Difference, 78-6202
- Acetylglucosaminidase**
Benzene
Enzymatic Activity, Rat, 78-6258
Granulocytes, 78-6258
Lymphocytes, 78-6258
- Acetyltransferases**
Benzenamine, 4,4'-Methylenebis-(2-chloro)-
Acetylation Kinetics, 78-6264
Benzidine
Acetylation Kinetics, 78-6264
Fluoren-2-amine
Acetylation Kinetics, 78-6264
Isonicotinic Acid Hydrazide
Acetylation Kinetics, 78-6264
1-Naphthylamine
Acetylation Kinetics, 78-6264
2-Naphthylamine
Acetylation Kinetics, 78-6264
Sulfanilamide, *N*-(4,6-Dimethyl-2-pyrimidinyl)-
Acetylation Kinetics, 78-6264
- Achlorhydria**
Stomach Neoplasms
Gastritis, 78-6533
- Acridinium, 3,6-Diamino-10-methyl-, Chloride**
Escherichia coli
Mutagenic Activity, 78-6159
Mutagenic Activity
Galactose Resistance, 78-6159
Salmonella typhimurium
Mutagenic Activity, 78-6159
- Acriflavine**
see Acridinium, 3,6-Diamino-10-methyl-, Chloride
- Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-**
Enzyme Activation
Microsomes, Testis, 78-6227
Escherichia coli
Enzyme Activation, 78-6227
Oxidoreductases
Microsomes, Testis, 78-6227
Saccharomyces cerevisiae
Growth, Radiation-Sensitive Mutants, 78-6244
- Acrylonitrile**
Occupational Hazard
Carcinogenic Activity, Review, 78-6021
Exposure Standards, Review, 78-6021
- Actinomycin D**
Estradiol, 3-Benzoyl
Ovalbumin, 78-6347
Progesterone, 78-6347
Ovalbumin
Plasma Levels, Chick, 78-6347
Progesterone
Plasma Levels, Chick, 78-6347
Saccharomyces cerevisiae
Mutagenic Activity, 78-6304
- Adenine**
Aflatoxin B1
DNA Cleavage, 78-6124
- Adenine, 3-Methyl-**
Dimethylamine, *N*-Nitroso-
DNA, 78-6215
Urea, Methyl Nitroso-, 78-6215
Urea, Methyl Nitroso-
Chromatin, Liver, 78-6215
- Adenocarcinoma**
Bladder Neoplasms
Epidemiology, 78-6567
Cecal Neoplasms
Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-
78-6286
Cholanthrene, 3-Methyl-, 78-6321
Colonic Neoplasms
4-Biphenylamine, 3,2'-Dimethyl-, 78-6297
Cholanthrene, 3-Methyl-, 78-6321
Colitis, Ulcerative, 78-6537
Genetics, 78-6538
Ultrastructural Study, 78-6536
Intestinal Neoplasms
Acenaphthene, 5-Nitro-, 78-6254
Radiation, Ionizing, 78-6385
Schistosoma japonicum, 78-6535
Kidney Neoplasms
Radium, 78-6564
Ultrastructural Study, 78-6532
Lung Neoplasms
Asbestos, 78-6168
Epidemiology, 78-6547

nocarcinoma (cont'd)

- Mammary Neoplasms, Experimental
 - 4-Biphenylamine, 3,2'-Dimethyl-, 78-6297
 - Phagocytosis, 78-6526
 - Radiation, Ionizing, 78-6050
 - Selenium, 78-6146
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-6342
- Nose Neoplasms
 - Reverse Transcriptase, 78-6521
 - Ultrastructural Study, Sheep, 78-6521
 - Virus-Like Particles, 78-6521
- Prostatic Neoplasms
 - Virus, Cytomegalo, 78-6499
- Radiation, Ionizing
 - 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385
- Rectal Neoplasms
 - Carcinoembryonic Antigen, 78-6596
 - Radiation, Ionizing, 78-6385
 - Transplantation, Heterologous, 78-6596
 - Ultrastructural Study, 78-6596
- Stomach Neoplasms
 - Agammaglobulinemia, 78-6534
 - Anemia, Pernicious, 78-6534
 - Gastritis, 78-6534
 - Immunologic Deficiency Syndromes, 78-6534
 - Plants, Edible, 78-6210
- Thyroid Neoplasms
 - Epidemiology, 78-6523
 - Radiation, Ionizing, 78-6378, 78-6523
 - Radiotherapy, 78-6380
- Uterine Neoplasms
 - Epidemiology, Review, 78-6026

nocarcinoma, Papillary

- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6315
- Thyroid Neoplasms
 - Radiation, Ionizing, 78-6363, 78-6377

enofibroma

- Breast Neoplasms
 - Antibodies, Viral, 78-6412
 - Precancerous Conditions, 78-6553
- Mammary Neoplasms, Experimental
 - Radiation, Ionizing, 78-6050, 78-6342
 - Xanthine, 3-Hydroxy-, 78-6250

enoma

- Cecal Neoplasms
 - Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-78-6286
- Colonic Neoplasms
 - Ultrastructural Study, 78-6536
- Gynecologic Neoplasms
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-6027
 - Transplacental Carcinogenesis, Review, 78-6027
- Liver Neoplasms
 - Carcinoembryonic Antigen, 78-6352
 - Contraceptives, Oral, 78-6352, 78-6354
 - Estrogens, 78-6352
- Lung Neoplasms
 - Carbamic Acid, Ethyl Ester, 78-6185
 - Carbamic Acid, Vinyl Ester, 78-6185
 - Xanthine, 3-Hydroxy-, 78-6249
- Mammary Neoplasms, Experimental
 - Xanthine, 3-Hydroxy-, 78-6250
- Pancreatic Neoplasms
 - Streptozotocin, 78-6208

Adenoma (cont'd)

- Parathyroid Neoplasms
 - Epidemiology, 78-6379
 - Radiation, Ionizing, 78-6376, 78-6377, 78-6379
- Thyroid Neoplasms
 - Imidazole-2-thiol, 1-Methyl-, 78-6248
 - Radiation, Ionizing, 78-6523

Adenosine Cyclic 3',5' Monophosphate

- Erythroleukemia
 - Caffeine, 78-6426
 - Cell Differentiation, 78-6426
 - 2-Imidazolidinone, 4-Methoxy-, 78-6426
 - Theophylline, 78-6426
 - Virus, Friend Murine Leukemia, 78-6426
 - Xanthine, 3-Isobutyl-1-methyl-, 78-6426
- Rectal Neoplasms
 - Carcinoembryonic Antigen, 78-6596
- Retinoic Acid
 - Plasminogen Activators, 78-6338
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Plasminogen Activators, 78-6338
- Virus, Rous Sarcoma
 - Plasminogen Activators, 78-6338

Adrenal Gland Neoplasms

- Epidemiology
 - Inbred Mice, 78-6595
- Kidney Neoplasms
 - Radiotherapy, 78-6384
- Radiation, Ionizing
 - In Utero Exposure, Rat, 78-6382

Adriamycin

- Antineoplastic Agents
 - Ames Test, 78-6232
- Escherichia coli*
 - DNA Repair, 78-6308

Aflatoxin B1

- Adenine
 - DNA Cleavage, 78-6124
- Aflatoxin M1
 - Metabolism, Rat, 78-6125
- Cell Transformation, Neoplastic
 - Diploid Cells, 78-6245
- Cells, Cultured
 - Colony Formation, Agar, 78-6245
- DNA Repair
 - Liver, Rat, 78-6216
- Guanine
 - DNA Cleavage, 78-6124
- Mesenchymoma
 - Cells, Cultured, 78-6245
- Microsomes, Liver
 - DNA Degradation, 78-6124
- Piperidine
 - DNA Cleavage, 78-6124
- Virus, Feline Sarcoma
 - Reverse Transcriptase, 78-6440
 - Virus Replication, 78-6440
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6242

Aflatoxin M1

- Aflatoxin B1
 - Metabolism, Rat, 78-6125
- Starch
 - Hepatotoxicity, 78-6125
 - Metabolism, Rat, 78-6125

Aflatoxin M1 (cont'd)

- Sucrose
- Metabolism, Rat, 78-6125

Agammaglobulinemia

- Stomach Neoplasms
- Adenocarcinoma, 78-6534

Agrobacterium tumefaciens

- Cytokinins
- Isolation and Characterization, 78-6585
- DNA, Bacterial
- Carcinogenic Activity, 78-6584
- DNA-DNA Hybridization, 78-6583, 78-6584
- Plasmids, 78-6583, 78-6584
- Endonucleases
- DNA, Mobility, 78-6582
- Isolation and Characterization, 78-6582
- Isomerases
- Isolation and Characterization, 78-6582
- Plant Tumors
- DNA, Bacterial, 78-6583
- Plasmids
- Cytokinins, 78-6585
- Purine, 5-(4-Hydroxy-3-methyl-*trans*-2-butenylamino)-
- Isolation and Characterization, 78-6585
- Purine, 6-(3-Methyl-2-butenylamino)-
- Isolation and Characterization, 78-6585

Air Pollution

- Activity Cycles
- Health Effects, 78-6156
- Carbon Radioisotopes
- Dose-Response Study, 78-6386
- Nuclear Power, 78-6386
- Carcinogen, Chemical
- Dose-Response Study, 78-6110
- Circadian Rhythm
- Health Effects, 78-6156
- Ethylene, Chloro-
- Dose-Response Study, 78-6110
- Lead
- Body Burden, Child, 78-6136
- Chromosome Aberrations, 78-6136
- Cytogenetic/Biochemical Effect, 78-6135
- Lung Neoplasms
- Carbon Dioxide, 78-6019
- Fossil Fuels, Review, 78-6019

Alanine Aminotransferase

- Benzidine, 3,3'-Dichloro-
- Enzymatic Activity, Dog, 78-6270
- Chloroform
- Hepatotoxicity, Rat, 78-6109

Alanine, 3-(*p*-(Bis(2-chloroethyl)amino)phenyl)-

- Immunosuppression
- Carcinogenic Activity, Review, 78-6051

Alanine, Phenyl-

- Ames Test
- Pyrolysis Product, 78-6201

Alcohol Oxidoreductases

- Hypercholesteremia
- Enzymatic Activity, 78-6589
- Methane, Azoxy-
- Pyrazole, 78-6107
- Methanol, (Methyl-*ONN*-azoxy)-
- Pyrazole, 78-6107

Aldrin

- Cells, Cultured
- Metabolism, Lung, 78-6305

Alkaline Phosphatase

- Aniline, 4,4'-Methylenedi-
- Enzymatic Activity, Dog, 78-6261
- Leukemia, Lymphoblastic
- Enzymatic Activity, 78-6457
- Leukemia, Myeloblastic
- Enzymatic Activity, 78-6457
- Teratoid Tumor
- Retinoic Acid, 78-6339

Alloxan

- Leukemia
- Fish, 78-6233
- Leukocyte Count
- Fish, 78-6233

Alpha Fetoproteins

- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
- Concanavalin A Reactivity, 78-6277
- Serum Levels, 78-6277
- Hepatology
- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6277
- Precancerous Conditions, 78-6277
- Radioimmunoassay, 78-6531

Alpha Particles

- Lung, Dispersion
- Mathematical Model, 78-6370
- Plutonium Oxide
- Lung, Dispersion, 78-6370

Alum

- see Sulfuric Acid, Aluminum Potassium Salt, Dodecahydrate

Aluminum Hydroxide

- Colonic Neoplasms
- Epidemiology, 78-6126
- Peptic Ulcer, 78-6126

Ames Test

- Acenaphthene, 5-Nitro-
- Mutagenic Activity, 78-6254
- Acetamide, *N*-(5-*p*-Acetamidophenylthiophen-2-yl)-
- Mutagenic Activity, 78-6260
- Acetamide, *N*-(5-Phenylthiophen-2-yl)-
- Mutagenic Activity, 78-6260
- Adriamycin
- Antineoplastic Agents, 78-6232
- Alanine, Phenyl-
- Pyrolysis Product, 78-6201
- Amino Acids
- Pyrolysis Product, 78-6201
- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
- S-9 Mix, 78-6263
- Azo Compounds
- Riboflavine, 78-6263
- Benzamide, 3,5-Dinitro-
- Mutagenic Activity, 78-6285
- Benzamide, 2-Methyl-3,5-dinitro-
- Mutagenic Activity, 78-6285
- Benzene, 4-Allyl-1,2-(methylenedioxy)-
- Microsomes, Liver, 78-6290
- Benzenesulfonic Acid, 4,4'-Oxybis-, Dihydrazide-
- S-9 Fraction, 78-6182
- Benzidine, 2,2',5,5'-Tetrachloro-

s Test (cont'd)

- Mutagenic Activity, 78-6254
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 - Mutagenic Activity, 78-6280
- Benzo(a)pyrene
 - Glutathione, 78-6333
 - Microsomes, Liver, 78-6302
 - Mutagenic Activity, 78-6328
 - Phenol, (1,1-Dimethylethyl)-4-methoxy-, 78-6333
 - Quinoline, 6-Ethoxy-1,2-dihydro-2,2,4-trimethyl-78-6333
- Benzo(f)quinoline
 - Amino Derivatives, 78-6239
 - Microsomes, Liver, 78-6239
- Benzo(f)quinoline, 2-Amino-
 - Microsomes, Liver, 78-6239
 - Norharman, 78-6239
- Benzo(h)quinoline
 - Amino Derivatives, 78-6239
 - Microsomes, Liver, 78-6239
- Benzo(h)quinoline, 2-Amino-
 - Microsomes, Liver, 78-6239
 - Norharman, 78-6239
- Benzopurpurine 4B
 - Riboflavine, 78-6263
- 4-Biphenylamine
 - Thiophene Analogs, 78-6260
- 1,4-Butanediol, Dimethylsulfonate
 - Antineoplastic Agents, 78-6232
 - Mutagenic Activity, 78-6163
- 1-Butanone, 4-Hydroxy-1-(3-pyridyl)-
 - Mutagenic Activity, 78-6175
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Mutagenic Activity, 78-6206
- C.I. Acid Red 26, Disodium Salt
 - Riboflavine, 78-6263
- C.I. Direct Rod 28, Disodium Salt
 - Riboflavine, 78-6263
- Carbamic Acid, Vinyl Ester
 - Cytochrome P-450 Inhibitors, 78-6185
 - Mutagenic Activity, 78-6185
- Carbazilquinone
 - Antineoplastic Agents, 78-6232
- Carbon
 - Water, Recycled, 78-6105
- Carcinogen, Chemical
 - Aroclor 1254, 78-6166
 - Cyclohexene Oxide, 78-6166
 - Mutagenic Activity, 78-6100, 78-6206
 - Mutagenic Activity, Review, 78-6005
- Chlorazol Violet N
 - Riboflavine, 78-6263
- Chlorine
 - Bleaching Effluents, Kraft Mills, 78-6132
- Chromic Acid
 - Erythrocytes, 78-6130
 - Microsomes, Liver, 78-6130
- Chromic Acid, Dipotassium Salt
 - Erythrocytes, 78-6130
 - Microsomes, Liver, 78-6130
- Chromic Acid, Lead Salt
 - Erythrocytes, 78-6130
 - Microsomes, Liver, 78-6130
- Chromic Acid, Zinc Salt
 - Ascorbic Acid, 78-6130
 - Erythrocytes, 78-6130
 - Glutathione, 78-6130

Ames Test (cont'd)

- Microsomes, Liver, 78-6130
- Mutagenic Activity, 78-6130
- Permanganic Acid, Potassium Salt, 78-6130
- Sulfurous Acid, Disodium Salt, 78-6130
- Chrysene, 1,2-Dihydro-1,2-dihydroxy-5-methyl-
 - Carcinogenic Metabolite, 78-6323
- Chrysene, 7,8-Dihydro-7,8-dihydroxy-5-methyl-
 - Carcinogenic Metabolite, 78-6323
- Chrysene, 5-Methyl-
 - Carcinogenic Metabolite, 78-6323
- Cosmetics
 - Mutagenic Activity, 78-6150
- p*-Cresol, 2,6-Di-*tert*-butyl-
 - Mutagenic Activity, 78-6206
- Cyclophosphamide
 - Antineoplastic Agents, 78-6232
- Daunomycin
 - Antineoplastic Agents, 78-6232
- Dichromic Acid, Disodium Salt
 - Erythrocytes, 78-6130
 - Microsomes, Liver, 78-6130
- Dimethylamine
 - Microsomes, Liver, 78-6290
- Dimethylamine, *N*-Nitroso-
 - S-9 Mix, 78-6263
- Endosulfan
 - Mutagenic Activity, 78-6163
- Ethane, 1,2-Dichloro-
 - Glutathione, 78-6113
 - NADP, 78-6113
- Ethanol, 2-Hydrazino-
 - S-9 Fraction, 78-6182
- Fluorosulfuric Acid, Methyl Ester
 - Dose-Response Study, 78-6165
- Formic Acid, Ethylenemethylhydrazide
 - Microsomes, Liver, 78-6181
- Formic Acid, 1-Methyl Hydrazide
 - Microsomes, Liver, 78-6181
- 2-Furanol 5-(3-Pyridyl)tetrahydro-
 - Mutagenic Activity, 78-6175
- Glutamic Acid
 - Pyrolysis Products, 78-6198
- Hair Dyes
 - S-9 Fraction, 78-6269
- Histones
 - Pyrolysis Product, 78-6201
- Hydrazine, 2,4-Dinitrophenyl-
 - S-9 Fraction, 78-6182
- Hydrazine, Methyl-
 - Microsomes, Liver, 78-6181
- Hydrazine, (4-Methylphenyl)-
 - S-9 Fraction, 78-6182
- Hydrazine, Monohydrate
 - S-9 Fraction, 78-6182
- Hydrazine, Phenethyl-, Sulfate
 - S-9 Fraction, 78-6182
- Hydrazine, Phenyl-, Monohydrochloride
 - S-9 Fraction, 78-6182
- Hydrazine, Sulfate
 - S-9 Fraction, 78-6182
- ICR 372
 - Genetic Duplication, 78-6359, 78-6360
- Isonicotinic Acid Hydrazide
 - Mutagenic Activity, 78-6163
- Isoquinoline
 - Amino Derivatives, 78-6239

Ames Test (cont'd)

- Microsomes, Liver, 78-6239
- Methane, Azoxy-
 - Mutagenic Activity, 78-6106
- Methanesulfonic Acid, (Antipyrinylmethylamino)-,
 - Monosodium Salt
 - Mutagenic Activity, 78-6163
- Methanesulfonic Acid, Ethyl Ester
 - Genetic Duplication, 78-6359, 78-6360
- Microsomes, Liver
 - Bleaching Effluents, Kraft Mills, 78-6132
- Mitomycin C
 - Antineoplastic Agents, 78-6232
- Muramidase
 - Pyrolysis Product, 78-6201
- 2-Naphthol, 1-(Phenylazo)-
 - Mutagenic Activity, 78-6152
- 2-Naphthol, 1-(α -Tolylazo)-
 - Mutagenic Activity, 78-6152
- 2-Naphthol, 1-(2,4-Xylylazo)-
 - Mutagenic Activity, 78-6152
- 2-Naphthylamine, 1-(Phenylazo)-
 - Mutagenic Activity, 78-6152
- 2-Naphthylamine, 1-(α -Tolylazo)-
 - Mutagenic Activity, 78-6152
- Nicotine, 1'-Nitroso-1'-demethyl-
 - Carcinogenic Metabolite, 78-6175
- Nitroso Compounds
 - Antineoplastic Agents, 78-6232
- Nitrous Acid
 - Genetic Duplication, 78-6359
- Oxalic Acid, Butylhydrazino-
 - S-9 Fraction, 78-6182
- Ozone
 - Water, 78-6105
- Phenanthridine
 - Amino Derivatives, 78-6239
 - Microsomes, Liver, 78-6239
- Phenol, 2-Methoxy-4-(2-propenyl)-
 - Metabolites, 78-6290
 - Microsomes, Liver, 78-6290
- p*-Phenyldiamine
 - Hair Dyes, 78-6269
- Piperazine, 1,4-Bis(3-bromo-1-oxopropyl)-
 - Antineoplastic Agents, 78-6232
- Piperidine
 - Microsomes, Liver, 78-6290
- Potassium Hypochlorite
 - Bleaching Effluents, Kraft Mills, 78-6132
- Proline
 - Microsomes, Liver, 78-6290
- Propane, 2,2'-Oxybis(2-chloro)-
 - Enzyme Activation, 78-6118
- 1-Propanol, 2,3-Dibromo-, Phosphate
 - Mutagenic Activity, 78-6144
- 1-Propanone, 3-(Dimethylamino)-1-(3',4'-methylenedioxyphenyl)-
 - Microsomes, Liver, 78-6290
- 1-Propanone, 3-Piperidyl-1-(3',4'-methylenedioxyphenyl)-
 - Microsomes, Liver, 78-6290
- 1-Propanone, 3-Pyrrolidinyl-1-(3',4'-methylenedioxyphenyl)-
 - Microsomes, Liver, 78-6290
- Pyridine, 2-Amino-5-phenyl-
 - Pyrolysis Product, 78-6201
- Pyridine, 3-(1-Pyrrolin-2-yl)-
 - Mutagenic Activity, 78-6175

Ames Test (cont'd)

- Pyrrolidine
 - Microsomes, Liver, 78-6290
- Quinoline
 - Amino Derivatives, 78-6239
 - Microsomes, Liver, 78-6239
- Radiation, Ionizing
 - Genetic Duplication, 78-6359, 78-6360
- Shelltox
 - Mutagenic Activity, 78-6163
- Smoke Condensate
 - Carcinogenic Potential, 78-6172
- Spermidine
 - Nitrosation, 78-6204
 - Nitrous Acid, Sodium Salt, 78-6204
- Synfuel A-3
 - Microsomes, Liver, 78-6102
 - Mutagenic Activity, 78-6102
- Thiophene, 5-(*p*-Acetamidophenyl)-2-amino-, Hydrochloride
 - Mutagenic Activity, 78-6260
- Thiophene, 2-Amino-5-phenyl-, Hydrochloride
 - Mutagenic Activity, 78-6260
- Triazene, 3-Acetyl-3-methyl-1-(2,4,6-tribromophenyl)-
 - Mutagenic Activity, 78-6223
- Triazene, 3,3-Dimethyl-1-(*p*-chlorophenyl)-
 - Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
 - 78-6223
 - Microsomes, Liver, 78-6223
- Triazene, 3,3-Dimethyl-1-phenyl-
 - Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
 - 78-6223
 - Microsomes, Liver, 78-6223
- Tryptophan
 - Pyrolysis Product, 78-6201
- Ultraviolet Rays
 - Genetic Duplication, 78-6359, 78-6360
- Water Pollutants
 - Aroclor 1254, 78-6158
 - Microsomes, Liver, 78-6158

Amino Acids

- Ames Test
 - Pyrolysis Product, 78-6201

Ammonia

- Colonic Neoplasms
 - Dietary Fiber, 78-6042
 - Metabolism, Review, 78-6042
- Dietary Fiber
 - Carcinogenic Activity, 78-6042
 - Metabolism, Review, 78-6042

Androgens

- Bacteria
 - Metabolism, Feces, 78-6348

Androst-4-ene-3,17-dione

- Bacteria
 - Metabolism, Feces, 78-6348
- Breast Neoplasms
 - Ethnic Groups, 78-6544
- Menopause
 - Ethnic Groups, 78-6544
- Phenol, (1,1-Dimethylethyl)-4-methoxy-
 - Transferases, 78-6333
- Uterine Neoplasms
 - Liver Diseases, 78-6028
 - Obesity, 78-6028

Androstan-17-one, 3 α -Hydroxy-
Hydroxysteroid Dehydrogenases
Vegetarianism, 78-6341
Intestinal Neoplasms
Hydroxysteroid Dehydrogenases, 78-6341

Androstan-3-one, 17 β -Hydroxy-
Mammary Neoplasms, Experimental
1-Piperazineethanol, 4-(3-(2-Chlorophenothiazin-10-yl)propyl)-, 78-6343

rostanediol
Mammary Neoplasms, Experimental
1-Piperazineethanol, 4-(3-(2-Chlorophenothiazin-10-yl)propyl)-, 78-6343

emia, Aplastic
Chromosomes
Hereditary Diseases, Review, 78-6074

emia, Pernicious
Stomach Neoplasms
Adenocarcinoma, 78-6534

giosarcoma
Ethane, 1,2-Dichloro-
Carcinogenic Activity, Mouse, Rat, 78-6114
Eye Neoplasms
Epidemiology, Dog, 78-6516

iline
Chlorine
Chlorination Products, 78-6262
Lethal Dose 50
Toxicity, Rat, 78-6266

iline, *p*-Bromo-
Lethal Dose 50
Toxicity, Rat, 78-6266

iline, *p*-Chloro-
Lethal Dose 50
Toxicity, Rat, 78-6266

iline, *N,N*-Dimethyl-
Chlorine
Chlorination Products, 78-6262

iline, *N,N*-Dimethyl-*p*-phenylazo-
Ames Test
S-9 Mix, 78-6263
***p*-Cresol, 2,6-Di-*tert*-butyl-**
Cytochrome P-450, 78-6218
Free Radicals, 78-6218
Hepatocarcinogenicity, 78-6218

Liver Neoplasms
Cytochrome P-450, 78-6218
Microsomes, Liver
Free Radicals, 78-6218
RNA, Transfer, Methyltransferases
Liver, Rat, 78-6265

iline, 4-Fluoro-
Lethal Dose 50
Toxicity, Rat, 78-6266

iline, *p*-Iodo-
Lethal Dose 50
Toxicity, Rat, 78-6266

iline, 4,4'-Methylenedi-
Alkaline Phosphatase
Enzymatic Activity, Dog, 78-6261
Bladder Neoplasms

Aniline, 4,4'-Methylenedi- (cont'd)
Carcinogenic Potential, Dog, 78-6261
Liver Neoplasms
Carcinogenic Potential, Dog, 78-6261

Aniline, *p*-Nitro-
Lethal Dose 50
Toxicity, Rat, 78-6266

Aniline, *p*-(Phenylazo)-
Carcinoma
Immunosuppression, 78-6278
Transplantation Immunology, 78-6278
Lymphocytes
Immunosuppression, 78-6278
RNA, Transfer, Methyltransferases
Liver, Rat, 78-6265

Anthracene
Aryl Hydrocarbon Hydroxylases
Skin, 78-6336

Anthranilic Acid, 3-Hydroxy-
see Benzoic Acid, 2-Amino-3-hydroxy-

Anti-Antibodies
Cervix Neoplasms
Virus, Herpes Simplex 2, 78-6449
Diethylamine, *N*-Nitroso-
Virus, Murine Leukemia, 78-6417
Virus, Radiation Leukemia, 78-6417
Lung Neoplasms
Diethylamine, *N*-Nitroso-, 78-6417
Virus, Gross Murine Leukemia
Antigen-Antibody Reactions, 78-6431
Syncytia Inhibition Test, 78-6431
Virus, Moloney Murine Leukemia
Antigen-Antibody Reactions, 78-6431
Syncytia Inhibition Test, 78-6431
Virus, Rauscher Murine Leukemia
Antigen-Antibody Reactions, 78-6431
Syncytia Inhibition Test, 78-6431

Antibodies
Estradiol
Pregnancy, Rat, 78-6351
Estriol
Pregnancy, Rat, 78-6351

Antibodies, Viral
Breast Neoplasms
A-Type Particles, 78-6412
Adenofibroma, 78-6412
Virus, Murine Mammary Tumor, 78-6412
Hodgkin's Disease
Virus, Epstein-Barr, 78-6452
Nasopharyngeal Neoplasms
Virus, Epstein-Barr, 78-6552
Prostatic Neoplasms
Virus, Cytomegalo, 78-6499
Virus, Bovine Leukemia
Immune Response, Review, 78-6088

Antibody Formation
Fibrosarcoma
Histocompatibility Antigens, 78-6493
Leukemia, Lymphoblastic
Cell Differentiation, Review, 78-6078
Virus, Feline Leukemia
Cat, Guinea Pig, 78-6442
Viral Proteins, 78-6442

Antibody Formation (cont'd)

- Virus, Friend Murine Leukemia
- Immunosuppression, 78-6427, 78-6430
- Spleen Cells, 78-6430

Antigen-Antibody Complex

- Neoplasms
- Immunoglobulins, Fc, 78-6504
- Prostatic Neoplasms
- Virus, Cytomegalo, 78-6499

Antigen-Antibody Reactions

- Burkitt's Lymphoma
- Immune Serums, 78-6498
- Cervix Neoplasms
- Virus, Herpes Simplex 2, 78-6449
- Fibrosarcoma
- Histocompatibility Antigens, 78-6493
- Leukemia
- B-Lymphocytes, 78-6498
- Leukemia, Lymphoblastic
- Antigens, Neoplasm, 78-6498
- Immune Serums, 78-6498
- Leukemia, Lymphocytic
- Virus, SV40, 78-6085
- Leukemia, Myeloblastic
- Antigens, Neoplasm, 78-6498
- Immune Serums, 78-6498
- Leukemia, Myelocytic
- Antigens, Neoplasm, 78-6498
- Immune Serums, 78-6498
- Lymphosarcoma
- Antigens, Neoplasm, 78-6498
- Immune Serums, 78-6498
- Virus, SV40, 78-6085
- Sarcoma
- Virus, SV40, 78-6085
- Sarcoma, Osteogenic
- Virus, SV40, 78-6085
- Sarcoma, Reticulum Cell
- Virus, SV40, 78-6085
- Virus, Avian Sarcoma
- Glucosamine, 78-6394
- Virus, Epstein-Barr
- Antigens, Viral, 78-6450
- Cell Membrane, 78-6460
- Virus, Feline Leukemia
- Antigens, Viral, 78-6443
- Virus, Feline Sarcoma
- Virus, Feline Leukemia, 78-6444
- Virus, Gross Murine Leukemia
- Anti-Antibodies, 78-6431
- Virus, Herpes Pan
- Antigens, Viral, 78-6450
- Virus, Herpes Papio
- Antigens, Viral, 78-6450
- Virus, Herpes Pongo
- Antigens, Viral, 78-6450
- Virus, Moloney Murine Leukemia
- Anti-Antibodies, 78-6431
- Virus, Murine Mammary Tumor
- Antigens, Viral, 78-6409
- Neoplasm Transplantation, 78-6409
- Virus, Rauscher Murine Leukemia
- Anti-Antibodies, 78-6431
- Virus, Rous Sarcoma
- Antigens, Viral, 78-6401

Antigenic Determinants

- Leydig Cell Tumor
- A-Type Particles, 78-6415
- 12-*O*-Tetradecanoylphorbol-13-acetate
- Plasminogen Activators, 78-6310
- Virus, Epstein-Barr
- Virus, Herpes Pan, 78-6450
- Virus, Herpes Papio, 78-6450
- Virus, Herpes Pongo, 78-6450
- Virus, Feline Sarcoma
- Cell Transformation, Neoplastic, 78-6444
- Virus, Feline Leukemia, 78-6444
- Virus, Murine Mammary Tumor
- A-Type Particles, 78-6415
- Viral Proteins, 78-6413, 78-6415
- Virus, Rous Sarcoma
- Cell Transformation, Neoplastic, 78-6401

Antigens

- Carrageen
- Hypersensitivity, Delayed, 78-6490
- Hodgkin's Disease
- Immunity, Cellular, 78-6487
- 2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-
- Hypersensitivity, Delayed, 78-6490
- Lymphoma
- Immunity, Cellular, 78-6487
- Plasmacytoma
- Mouse, Review, 78-6013
- Radiation, Ionizing
- Hypersensitivity, Delayed, 78-6490
- Reserpine
- Hypersensitivity, Delayed, 78-6490
- Virus, Epstein-Barr
- Antibody-Dependent Cytotoxicity, 78-6460
- Cell Membrane, 78-6460
- Papain, 78-6460
- Staphylococcal Protein A, 78-6460
- Virus, Polyoma
- Cell Transformation, Neoplastic, 78-6496
- Deletion Mutants, 78-6496
- Isolation and Characterization, 78-6496
- Temperature Sensitive Mutants, 78-6496

Antigens, Neoplasm

- Cell Transformation, Neoplastic
- Immune Response, Review, 78-6077
- Immunologic Technics, Review, 78-6077
- Leukemia, Lymphoblastic
- Antigen-Antibody Reactions, 78-6498
- Leukemia, Myeloblastic
- Antigen-Antibody Reactions, 78-6498
- Leukemia, Myelocytic
- Antigen-Antibody Reactions, 78-6498
- Lymphosarcoma
- Antigen-Antibody Reactions, 78-6498
- Neoplasm Metastasis
- Tumor-Associated Products, Review, 78-6065
- Sarcoma
- T-Lymphocytes, 78-6502

Antigens, Viral

- Brain Neoplasms
- Virus, Papova, 78-6486
- Virus, SV40, 78-6486
- Burkitt's Lymphoma
- Virus, Epstein-Barr, 78-6459
- Ependymoma
- Virus, SV40, 78-6486

ens, Viral (cont'd)

Fibroma

Virus, Bovine Papilloma, 78-6446

Fibrosarcoma

Virus, Bovine Papilloma, 78-6446

Hodgkin's Disease

Virus, Epstein-Barr, 78-6459

Leukemia

Virus, Feline Leukemia, 78-6443

Leukemia, Lymphoblastic

Virus, Epstein-Barr, 78-6457

Leukemia, Myeloblastic

Virus, Epstein-Barr, 78-6457

Lymphoma

Virus, Feline Leukemia, 78-6443

Neoplasms, Multiple Primary

Virus, Epstein-Barr, 78-6459

Papilloma

Virus, Bovine Papilloma, 78-6446

Virus, Adeno

Virus Subgroup, 78-6467

Virus, Adeno 3

Cells, Cultured, 78-6464

Immunofluorescence, 78-6467

Virus, Adeno 7

Immunofluorescence, 78-6467

Virus, Adeno 12

Immunofluorescence, 78-6467

Virus, Adeno 31

Immunofluorescence, 78-6467

Virus, Bovine Papilloma

Cell Membrane, 78-6446

Trans-Species Infection, 78-6446

Virus, Epstein-Barr

Antigen-Antibody Reactions, 78-6450

Chromatin, 78-6456

DNA Replication, 78-6454

Histones, 78-6456

Hybrid Cells, 78-6455

IgG Absorption Assay, 78-6456

Lymphocyte Transformation, 78-6454

Lymphocytes, 78-6454

Phosphonoacetic Acid, 78-6450

Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-, 78-6451

Uridine, 2'-Deoxy-5-iodo-, 78-6450

Virus, Feline Leukemia

Antigen-Antibody Reactions, 78-6443

Cell Membrane, 78-6443

Virus, Feline Sarcoma

Radioimmunoassay, 78-6444

Virus, Gross Murine Leukemia

Cell Transformation, Neoplastic, 78-6432

Virus, Herpes Pan

Antigen-Antibody Reactions, 78-6450

Phosphonoacetic Acid, 78-6450

Uridine, 2'-Deoxy-5-iodo-, 78-6450

Virus, Herpes Papio

Antigen-Antibody Reactions, 78-6450

Phosphonoacetic Acid, 78-6450

Uridine, 2'-Deoxy-5-iodo-, 78-6450

Virus, Herpes Pongo

Antigen-Antibody Reactions, 78-6450

Phosphonoacetic Acid, 78-6450

Uridine, 2'-Deoxy-5-iodo-, 78-6450

Virus, Moloney Murine Leukemia

Cell Transformation, Neoplastic, 78-6432

Antigens, Viral (cont'd)

Virus, Murine Mammary Tumor

Antigen-Antibody Reactions, 78-6409

Cell Migration Inhibition, 78-6497

Horizontal Transmission, 78-6414

Lymphocyte Transformation, 78-6497

Neoplasm Transplantation, 78-6409

Virus, Polyoma

Carcinogenic Activity, Review, 78-6092

Deletion Mutants, 78-6479

Isolation and Characterization, 78-6479

Temperature Sensitive Mutants, 78-6479

Virus, Rous Sarcoma

Antigen-Antibody Reactions, 78-6401

Cell Membrane, 78-6402

Cell Transformation, Neoplastic, 78-6402

RNA, Messenger, 78-6401

Temperature Sensitive Mutants, 78-6402

Virus, SV40

Immunity, Cellular, 78-6500

Immunization, 78-6500

Antilymphocyte Serum

Virus, SV40

T-Lymphocytes, 78-6500

Antimony

Bacillus subtilis

Mutagenic Activity, 78-6128

Antineoplastic Agents

Adriamycin

Ames Test, 78-6232

1,4-Butanediol, Dimethylsulfonate

Ames Test, 78-6232

Carbazilquinone

Ames Test, 78-6232

Cyclophosphamide

Ames Test, 78-6232

Daunomycin

Ames Test, 78-6232

Mitomycin C

Ames Test, 78-6232

Nitroso Compounds

Ames Test, 78-6232

Piperazine, 1,4-Bis(3-bromo-1-oxopropyl)-

Ames Test, 78-6232

Antioxidants

Colonic Neoplasms

Epidemiology, Review, 78-6039

Antipyrine

Lymphocytes

Aryl Hydrocarbon Hydroxylases, 78-6292

Plasma

Half-Life, 78-6292

Antipyrine, 4-(Dimethylamino)-

Chromosome Aberrations

Lymphocytes, Rat, 78-6214

Anus Neoplasms

Carcinoma, Transitional Cell

Ultrastructural Study, 78-6539

Areca

Cell Transformation, Neoplastic

Tobacco Extract, 78-6173

Cells, Cultured

Tobacco Extract, 78-6173

Aroclor 1016

Aryl Hydrocarbon Hydroxylases
Lepidoptera, 78-6296

Aroclor 1242

Aryl Hydrocarbon Hydroxylases
Lepidoptera, 78-6296

Aroclor 1254

Aryl Hydrocarbon Hydroxylases
Lepidoptera, 78-6296

Benzo(a)pyrene

Enzyme Activation, 78-6324

Carcinogen, Chemical

Ames Test, 78-6166

Mutagenic Activity, 78-6166

Chrysene, 5-Methyl-

Carcinogenic Metabolite, 78-6323

Dimethylamine, N-Nitroso-

Enzyme Activation, 78-6324

Hydro-Lyases

Enzymatic Activity, 78-6257

Pyrrolidine, 1-Nitroso-

Carcinogenic Metabolite, 78-6237

Water Pollutants

Ames Test, 78-6158

Arthritis, Rheumatoid

Leukemia, Myeloblastic

Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
78-6211

Aryl Hydrocarbon Hydroxylases**Acetanilide**

Lymphocytes, 78-6292

***p*-Acetophenetidide**

Lymphocytes, 78-6292

Anthracene

Skin, 78-6336

Antipyrine

Lymphocytes, 78-6292

Aroclor 1016

Lepidoptera, 78-6296

Aroclor 1242

Lepidoptera, 78-6296

Aroclor 1254

Lepidoptera, 78-6296

Asbestos

Cholanthrene, 3-Methyl-, 78-6169

Dibenz(a,h)anthracene, 78-6169

Lymphocytes, 78-6169

Occupational Hazard, 78-6169

Barbituric Acid, 5-Ethyl-5-phenyl-

Strain Difference, Mouse, 78-6319

Benz(a)anthracene

Lymphocytes, 78-6322

Plant Agglutinins, 78-6322

Benzo(a)pyrene

Cholanthrene, 3-Methyl-, 78-6326

Coumarin, 3, 3'-Methylenebis(4-hydroxy)-, 78-6334

Glutathione, 78-6326

Skin, 78-6336

5,6-Benzoflavone

Strain Difference, Mouse, 78-6319

Carcinoma, Bronchogenic

Benz(a)anthracene, 78-6322

Cholanthrene, 3-Methyl-, 78-6322

Lymphocytes, 78-6322

Smoking, 78-6322

Aryl Hydrocarbon Hydroxylases (cont'd)**Cholanthrene, 3-Methyl-**

Coumarin, 3, 3'-Methylenebis(4-hydroxy)-, 78-6334

Lymphocytes, 78-6322

Plant Agglutinins, 78-6322

Strain Difference, Mouse, 78-6319

Coal Tar

Skin, 78-6336

Hemopexin

Strain Difference, Mouse, 78-6319

Hepatoma

Benz(a)anthracene, 78-6319

Laryngeal Neoplasms

Benz(a)anthracene, 78-6322

Cholanthrene, 3-Methyl-, 78-6322

Lymphocytes, 78-6322

Polychlorobiphenyl Compounds

Lepidoptera, 78-6296

Pyrene

Skin, 78-6336

Skin Diseases

Coal Tar, 78-6336

Smoking

Lymphocytes, 78-6174

Theophylline

Lymphocytes, 78-6292

Asbestos**Aryl Hydrocarbon Hydroxylases**

Lymphocytes, 78-6169

Cholanthrene, 3-Methyl-

Aryl Hydrocarbon Hydroxylases, 78-6169

Dibenz(a,h)anthracene

Aryl Hydrocarbon Hydroxylases, 78-6169

Lung Neoplasms

Adenocarcinoma, 78-6168

Carcinoma, Epidermoid, 78-6529

Epidemiology, 78-6168

Epidemiology, Review, 78-6002

Mesothelioma, 78-6168

Occupational Hazard, 78-6002, 78-6168

Ultrastructural Study, 78-6529

Occupational Hazard

Aryl Hydrocarbon Hydroxylases, 78-6169

Precancerous Conditions, 78-6383

Pleural Diseases

Epidemiology, Review, 78-6023

Occupational Hazard, Review, 78-6023

X-Ray Study, Review, 78-6023

Pleural Neoplasms

Precancerous Conditions, 78-6383

Precancerous Conditions

Pleural Plaques, 78-6383

Prostatic Neoplasms

Occupational Hazard, 78-6168

Smoking

Co-carcinogenic Effect, Review, 78-6002, 78-6032

Asbestosis**Pleural Diseases**

Occupational Hazard, Review, 78-6023

Ascorbic Acid

Chromic Acid, Zinc Salt

Ames Test, 78-6130

Spermidine

Nitrosation, 78-6204

cytoma

- Brain Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6318
 - Virus, B77, 78-6408

xia Telangiectasia

- DNA Repair
 - Endonucleases, 78-6364
- Fibroblasts
 - DNA Repair, 78-6364
- Radiation, Ionizing
 - DNA Repair, 78-6364

phy

- Caffeine
 - Testis, Rat, 78-6123
- Nitrous Acid, Sodium Salt
 - Testis, Rat, 78-6123
- Theobromine
 - Testis, Rat, 78-6123
- Theophylline
 - Testis, Rat, 78-6123

stralia Antigen

- Hepatoma
 - Liver Cirrhosis, 78-6531
 - Radioimmunoassay, 78-6531

omobile Exhaust

- Carcinogen, Chemical
 - Isolation and Characterization, 78-6300
- Polycyclic Hydrocarbons
 - Carcinogenic Activity, 78-6300
- Skin Neoplasms
 - Polycyclic Hydrocarbons, 78-6300

aserine

- see Serine, Diazoacetate (Ester)

athioprine

- Leukemia, Myeloblastic
 - Case Report, 78-6209

o Compounds

- Riboflavine
 - Ames Test, 78-6263

uleno(1,2,3-cd)phenalene

- Benzenecarboxiperoxoic Acid, 3-Chloro-
 - Carcinogenic Metabolite, 78-6289
- 1*H*-Benzo(cd)fluoranthene, 1-Oxo-
 - Carcinogenic Metabolite, 78-6289
- 11*H*-Benzo(cd)fluoranthene, 11-Oxo-
 - Carcinogenic Metabolite, 78-6289

acillus subtilis

- Antimony
 - Mutagenic Activity, 78-6128
- Benzimidazole, 2-(4-Thiazolyl)-
 - Mutagenic Activity, 78-6256
- Benzoic Acid
 - Mutagenic Activity, 78-6256
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Mutagenic Activity, 78-6206
- Carcinogen, Chemical
 - Mutagenic Activity, 78-6206
- Cesium Chloride
 - Mutagenic Activity, 78-6128
- Chloroplatinic Acid
 - Mutagenic Activity, 78-6128
- Ethanol, 2,2',2''-Nitrilotris-
 - Mutagenic Activity, 78-6224

Bacillus subtilis (cont'd)

- Germanium Tetrachloride
 - Mutagenic Activity, 78-6128
- Osmium Tetroxide
 - Mutagenic Activity, 78-6128
- Pimaricin
 - Mutagenic Activity, 78-6256
- Propionic Acid, Calcium Salt
 - Mutagenic Activity, 78-6256
- Rhodium Chloride
 - Mutagenic Activity, 78-6128
- Tellurium
 - Mutagenic Activity, 78-6128
- Vanadium
 - Mutagenic Activity, 78-6128

Bacteria

- Androgens
 - Metabolism, Feces, 78-6348
- Androst-4-ene-3,17-dione
 - Metabolism, Feces, 78-6348
- C.I. Food Brown 1
 - DNA Repair, 78-6153
 - Mutagenic Activity, 78-6153
- Colonic Neoplasms
 - Dietary Fats, 78-6033
 - Epidemiology, Review, 78-6033, 78-6043
 - Epinephrine, 78-6033
- DNA Polymerase
 - DNA Repair, Review, 78-6055
- Estradiol
 - Metabolism, Feces, 78-6348
- Estrogens
 - Metabolism, Feces, 78-6348
- Estrone
 - Metabolism, Feces, 78-6348
- Fluorescein, 2',4',5',7'-Tetrabromo-, Disodium Salt
 - DNA Repair, 78-6153
 - Mutagenic Activity, 78-6153
- Hydroxysteroids
 - Metabolism, Feces, 78-6348
- Intestinal Neoplasms
 - Carcinoma, 78-6587
 - Enzymatic Activity, 78-6587
- Intestinal Polyps
 - Enzymatic Activity, 78-6587
 - Oxidoreductases, 78-6587
- Ultraviolet Rays
 - DNA Repair, Review, 78-6052

Bacteriophages

- Sulfurous Acid, Monosodium Salt
 - Cytosine, 78-6160
 - Mutagenic Activity, 78-6160
 - Transaminated Residues, 78-6160

Bacteroides

- 5 β -Cholan-24-oic Acid, 3 α ,7 α -Dihydroxy-
 - Oxidoreductases, 78-6590

Barbituric Acid, 5-Ethyl-5-phenyl-

- Aryl Hydrocarbon Hydroxylases
 - Strain Difference, Mouse, 78-6319
- Benzo(a)pyrene
 - Carcinogenic Metabolite, 78-6326
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Carcinogenic Metabolite, 78-6326
- Chloroform
 - Metabolism, 78-6109

- Barbituric Acid, 5-Ethyl-5-phenyl- (cont'd)**
 Cyclopentadiene, Tetrachloro-
 Mutagenic Activity, 78-6167
 Dimethylamine, *N*-Nitroso-
 Metabolism, Liver, 78-6217
 Ethane, 1,2-Dibromo-
 Metabolism, Rat, 78-6115
 Hemopexin
 Plasma Levels, Mouse, 78-6319
 Hydro-Lyases
 Enzymatic Activity, 78-6257, 78-6320
 Metabolism, Lung, 78-6331
 Hydroxylases
 Enzymatic Activity, 78-6320
 Liver Neoplasms
 Acetamide, *N*-Fluoren-2-yl-, 78-6193
 Precancerous Conditions, 78-6193
 Methane, Azoxy-
 Formaldehyde, 78-6106
 Microsomes, Liver
 Hydro-Lyases, 78-6320
 Hydroxylases, 78-6320
 Transferases
 Metabolism, Lung, 78-6331
- Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt**
 Phenanthrene, 9,10-Dihydro-9,10-dihydroxy-
 Carcinogenic Metabolite, 78-6312
 Triazene, 3,3-Dimethyl-1-(*p*-chlorophenyl)-
 Ames Test, 78-6223
 Triazene, 3,3-Dimethyl-1-phenyl-
 Ames Test, 78-6223
- Barium**
 Chlorophyll
 Mutagenic Activity, 78-6134
- Bence Jones Protein**
 Multiple Myeloma
 Case Report, 78-6514
 Immunoglobulins, Heavy Chain, 78-6514
- Benz(a)anthracene**
 Carcinoma, Bronchogenic
 Aryl Hydrocarbon Hydroxylases, 78-6322
 Ethane, 1,2-Dibromo-
 Metabolism, Rat, 78-6115
 Hepatoma
 Aryl Hydrocarbon Hydroxylases, 78-6319
 Hemopexin, 78-6319
 Laryngeal Neoplasms
 Aryl Hydrocarbon Hydroxylases, 78-6322
 Lymphocytes
 Aryl Hydrocarbon Hydroxylases, 78-6322
 Plant Agglutinins
 Aryl Hydrocarbon Hydroxylases, 78-6322
 Quinone Reductases
 Liver, Rat, 78-6334
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 7-Bromomethyl-**
 Fibroblasts
 DNA Repair, 78-6313
 Xeroderma Pigmentosum
 Cell Survival, 78-6313
 DNA Repair, 78-6313
- Benz(a)anthracene, 5,6-Dihydro-7,12-dimethyl-5,6-epoxy-**
Escherichia coli
- Benz(a)anthracene, 5,6-Dihydro-7,12-dimethyl-5,6-epoxy-**
 (cont'd)
 DNA Repair, 78-6325
- Benz(a)anthracene, 5,6-Dihydro-5,6-epoxy-7-methyl-**
Escherichia coli
 DNA Repair, 78-6325
- Benz(a)anthracene, 7,12-Dimethyl-**
 Brain Neoplasms
 Astrocytoma, 78-6318
 Glioblastoma Multiforme, 78-6318
 Monoamine Oxygenase Inhibitors, 78-6318
 Serotonin, 78-6318
 Chromosome Aberrations
 Micronucleus Test, 78-6162
Escherichia coli
 Carcinogenic Metabolite, Review, 78-6015
- Leukemia**
 Chromosome Aberrations, 78-6314
 T-Lymphocytes, 78-6314
 2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
 78-6317
- Mammary Neoplasms, Experimental**
 Adenocarcinoma, Papillary, 78-6315
 Dietary Fats, Review, 78-6037
 Estradiol, 78-6351
 Estriol, 78-6351
 Prolactin, 78-6343
 Testosterone, 78-6343
 Virus-Like Particles, 78-6315
 Virus, Murine Mammary Tumor, 78-6315
- Mouth Neoplasms**
 Dose-Response Study, Hamster, 78-6316
 Localization, Cheek Pouch, 78-6316
- Skin Neoplasms**
 Cortisol, 78-6307
 Cortisone, 78-6307
 Croton Oil, 78-6307
 Dexamethasone, 78-6307
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 1-Fluoro-7-methyl-**
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 2-Fluoro-7-methyl-**
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 3-Fluoro-7-methyl-**
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 4-Fluoro-7-methyl-**
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 7-Methyl-**
 Structure-Activity Relationship
 Carcinogenic Activity, Review, 78-6015
- Benz(a)anthracene, 7,8,12-Trimethyl-**
 Leukemia
 2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
 78-6317
 2-Naphthol, 1-((4-(*o*-Tolylazo)-*o*-tolyl)azo)-, 78-6317
 Mammary Neoplasms, Experimental
 2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
 78-6317

a)anthracene, 7,8,12-Trimethyl- (cont'd)
 2-Naphthol, 1-((4-(*o*-Tolylazo)-*o*-tolyl)azo)-, 78-6317
2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
 Oxidoreductases, 78-6317

e)indole, 3-(β -Nitrovinyl)-
***p*-Benzoquinone**
 Benzocarbazole Derivatives, Synthesis, 78-6299
2-Butynedioic Acid, Dimethyl Ester
 Benzocarbazole Derivatives, Synthesis, 78-6299
Ethenetetra carbonitrile
 Benzocarbazole Derivatives, Synthesis, 78-6299
Maleimide, *N*-Phenyl-
 Benzocarbazole Derivatives, Synthesis, 78-6299
1,4-Naphthoquinone
 Benzocarbazole Derivatives, Synthesis, 78-6299

amide, 3,5-Dinitro-
Ames Test
 Mutagenic Activity, 78-6285
Endonucleases
 Mutagenic Activity, 78-6285
Escherichia coli
 Mutagenic Activity, 78-6285
Ultraviolet Rays
 Endonucleases, 78-6285

amide, 2-Methyl-3,5-dinitro-
Ames Test
 Mutagenic Activity, 78-6285
Endonucleases
 Mutagenic Activity, 78-6285
Escherichia coli
 Mutagenic Activity, 78-6285
Ultraviolet Rays
 Endonucleases, 78-6285

enamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
Alpha Fetoproteins
 Concanavalin A Reactivity, 78-6277
 Serum Levels, 78-6277
Glutamyl Transpeptidase
 Shunt, Portacaval, 78-6275
Hepatocytes
 Cytotoxicity, Fibroblasts, 78-6267
Hepatoma
 Alpha Fetoproteins, 78-6277
 Glutamyl Transpeptidase, 78-6274, 78-6275
 Glutathione, 78-6274, 78-6275
 Hypophysectomy, 78-6274
 Shunt, Portacaval, 78-6275
Liver Neoplasms
 Cholangioma, 78-6276
 Nucleotides, Cyclic, 78-6276
 Precancerous Conditions, 78-6276

zenamine, 2-Methyl-4-((2-methylphenyl)azo)-
Carcinoma
 Immunosuppression, 78-6278
 Transplantation Immunology, 78-6278
Cecal Neoplasms
 Adenocarcinoma, 78-6286
 Adenoma, 78-6286
 Carcinogenic Activity, Mouse, 78-6286
Glucocorticoids
 Immunosuppression, 78-6278
Liver Neoplasms
 Carcinogenic Activity, Mouse, 78-6286

Benzenamine, 4,4'-Methylenebis-(2-chloro)-
 Acetylation Kinetics
 Species Difference, 78-6264
 Acetyltransferases
 Acetylation Kinetics, 78-6264
 Aerosols
 Generation Technique, 78-6279

Benzene
 Acetylglucosaminidase
 Enzymatic Activity, Rat, 78-6258
 Granulocytes, 78-6258
 Lymphocytes, 78-6258
 Leukemia
 Occupational Hazard, Review, 78-6014

Benzene, 4-Allyl-1,2-(methylenedioxy)-
 Microsomes, Liver
 Ames Test, 78-6290
 Mutagenic Activity, 78-6290

Benzene, *p*-Bromo-(epoxyethyl)-
Escherichia coli
 Mutagenic Activity, 78-6272

Benzene, 1-Chloro-2,4-dinitro-
 Hydro-Lyases
 Metabolism, Lung, 78-6331
 Phenol, (1,1-Dimethylethyl)-4-methoxy-
 Transferases, 78-6333
 Transferases
 Metabolism, Lung, 78-6331

Benzene, *m*-Chloro-(epoxyethyl)-
Escherichia coli
 Mutagenic Activity, 78-6272

Benzene, 1-(Chloromethyl)-4-nitro-
 Phenol, (1,1-Dimethylethyl)-4-methoxy-
 Transferases, 78-6333

Benzene, 1,2-Dichloro-4-nitro-
 Phenol, (1,1-Dimethylethyl)-4-methoxy-
 Transferases, 78-6333

Benzene, 3,4-Dimethyl-(epoxyethyl)-
Escherichia coli
 Mutagenic Activity, 78-6272

Benzene, (Epoxyethyl)-
Drosophila melanogaster
 Mutagenic Activity, 78-6253
Escherichia coli
 Lethal Dose 50, 78-6272
 Mutagenic Activity, 78-6272
 Lethal Dose 50
 Substituted Derivatives, 78-6272
 Phenol, (1,1-Dimethylethyl)-4-methoxy-
 Hydro-Lyases, 78-6257
 Substituted Derivatives
 Structure-Activity Relationship, 78-6272

Benzene, (Epoxyethyl)-*m*-methoxy-
Escherichia coli
 Mutagenic Activity, 78-6272

Benzene, (Epoxyethyl)-*m*-methyl-
Escherichia coli
 Mutagenic Activity, 78-6272

Benzene, (Epoxyethyl)-*p*-methyl-
Escherichia coli
 Mutagenic Activity, 78-6272

- α -Benzene Hexachloride**
see Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer
- γ -Benzene Hexachloride**
see Cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer
- Benzene, Hexachloro-**
Cells, Cultured
Metabolism, Lung, 78-6305
- Benzene, 4,5-Methylenetris-**
Smoke Condensate
Isolation and Characterization, 78-6337
- Benzenecarboxeperoxoic Acid, 3-Chloro-**
Azuleno(1,2,3-cd)phenalene
Carcinogenic Metabolite, 78-6289
- Benzenediazonium, 4-Sulfo-, Hydroxide, Inner Salt**
Hydroxylases
Localization, Liver, 78-6320
- Benzenediazosulfonic Acid, *p*-(Dimethylamino)-, Sodium Salt**
Chromosomes
Mitosis, 78-6284
Mutagenic Activity, 78-6284
- Benzenesulfonic Acid, 4,4'-Oxybis-, Dihydrazide-**
Ames Test
S-9 Fraction, 78-6182
- Benzidine**
Acetylation Kinetics
Species Difference, 78-6264
Acetyltransferases
Acetylation Kinetics, 78-6264
Aerosols
Generation Technique, 78-6279
Cells, Cultured
Thiophene Analogs, 78-6260
Chlorine
Chlorination Products, 78-6262
Cholesterol
Metabolism, Liver, 78-6192
Chromosome Aberrations
Methanesulfonic Acid, Ethyl Ester, 78-6162
Mutagenic Activity
Thiophene Analogs, 78-6260
Occupational Hazard
Isoantigens, 78-6566
- Benzidine, 3,3'-Dichloro-**
Aerosols
Generation Technique, 78-6279
Alanine Aminotransferase
Enzymatic Activity, Dog, 78-6270
Bladder Neoplasms
Carcinoma, Transitional Cell, 78-6270
Dose-Response Study, Dog, 78-6270
Neoplasms, Multiple Primary, 78-6270
Hepatosoma
Dose-Response Study, Dog, 78-6270
Neoplasms, Multiple Primary, 78-6270
Liver Neoplasms
Neoplasm Metastasis, 78-6270
- Benzidine, 2,2',5,5'-Tetrachloro-**
Ames Test
Mutagenic Activity, 78-6254
Lymphoma
Carcinogenic Potential, 78-6254
Mammary Neoplasms, Experimental
- Benzidine, 2,2',5,5'-Tetrachloro- (cont'd)**
Carcinoma, 78-6254
Fibrosarcoma, 78-6254
- Benzimidazole, 2-(4-Thiazolyl)-**
Bacillus subtilis
Mutagenic Activity, 78-6256
Nitrous Acid
Mutagenic Activity, 78-6256
- 2-Benzimidazolecarbamic Acid, 5-Benzoylmethyl Ester**
Phenol, (1,1-Dimethylethyl)-4-methoxy-
Mutagenic Activity, 78-6328
- 2-Benzimidazolecarbamic Acid, Methyl Ester**
Erythrocytes
Micronucleus Formation, 78-6303
Tubulin
Dose-Response Study, Mouse, 78-6303
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide**
Ames Test
Mutagenic Activity, 78-6280
Bladder
Absorption, Rat, 78-6281
Bladder Neoplasms
Carcinogenic Potential, Monkey, 78-6294
Cholanthrene, 3-Methyl-
Co-carcinogenic Activity, Mouse Embryo Cells
78-6283
Chromosomes
Bone Marrow, Mouse, 78-6280
Mutagenic Activity, 78-6280
Guanyl Cyclase
Bladder, Rat, 78-6282
Enzymatic Activity, 78-6282
Plasma Levels
Dose-Response Study, Rat, 78-6281
Urine
Metabolism, Monkey, 78-6294
- Benzo(a)pyrene**
Ames Test
Mutagenic Activity, 78-6328
Aroclor 1254
Enzyme Activation, 78-6324
Aryl Hydrocarbon Hydroxylases
Skin, 78-6336
Barbituric Acid, 5-Ethyl-5-phenyl-
Carcinogenic Metabolite, 78-6326
Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
Carcinogenic Metabolite, 78-6326
7,8-Benzoflavone
Carcinogenic Metabolite, 78-6326
Cell Transformation, Neoplastic
Collisional Quenchers, 78-6329
Chlorine
Co-carcinogenic Activity, 78-6330
Cholanthrene, 3-Methyl-
Aryl Hydrocarbon Hydroxylases, 78-6326
Carcinogenic Metabolite, 78-6326
Coumarin, 3, 3'-Methylenebis(4-hydroxy)-
Aryl Hydrocarbon Hydroxylases, 78-6334
Hepatotoxicity, Rat, 78-6334
DNA Repair
Dose-Response Study, 78-6324
Enzyme Activation, 78-6324
Epoxide Hydratases
Inactivating Mechanisms, 78-6302

Benzo(a)pyrene (cont'd)

- Fibroblasts
 - Fluorescence Quenching, 78-6329
- Fistula
 - Intestinal Absorption, 78-6327
- Glutathione
 - Ames Test, 78-6333
 - Aryl Hydrocarbon Hydroxylases, 78-6326
- Glutathione Transferases
 - Inactivating Mechanisms, 78-6302
- Hypertension
 - Carcinogenic Activity, 78-6332
- Intestinal Absorption
 - Metabolism, Rat, 78-6327
- Liver
 - Nucleic Acids, Binding, 78-6326
- Lymphatic System
 - Metabolism, Rat, 78-6327
- Methane, Diiodo-
 - Fluorescence Quenching, 78-6329
- Microsomes, Liver
 - Ames Test, 78-6302
 - Inactivating Mechanisms, 78-6302
 - Nucleic Acids, Binding, 78-6326
- Mutagenic Activity
 - Urine, Mouse, 78-6328
- Octane, 1-Iodo-
 - Fluorescence Quenching, 78-6329
- Phenol, (1,1-Dimethylethyl)-4-methoxy-
 - Ames Test, 78-6333
 - Carcinogenic Metabolite, 78-6333
 - Mutagenic Activity, 78-6328
- Quinoline, 6-Ethoxy-1,2-dihydro-2,2,4-trimethyl-
 - Ames Test, 78-6333
 - Carcinogenic Metabolite, 78-6333
 - Mutagenic Activity, 78-6328
- Quinone Reductases
 - Liver, Rat, 78-6334
- Respiratory Tract Neoplasms
 - Anesthesia, 78-6335
 - Brevital, 78-6335
 - Carcinoma, Epidermoid, 78-6335
 - Ethane, 1,1'-Oxybis-, 78-6335
 - Ether, 2,2-Dichloro-1,1-difluoroethyl Methyl 78-6335
 - Iron Oxide, 78-6335
- Skin Neoplasms
 - Chlorine, 78-6330
- Virus, Feline Sarcoma
 - Reverse Transcriptase, 78-6440
- Virus Replication, 78-6440
- Virus, Moloney Murine Sarcoma
 - Fibroblasts, 78-6329
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6242

Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Carcinogenic Metabolite, 78-6326

7,8-Benzoflavone

- Carcinogenic Metabolite, 78-6326

Cholanthrene, 3-Methyl-

- Carcinogenic Metabolite, 78-6326

Glutathione

- Carcinogenic Metabolite, 78-6326

Liver

- Nucleic Acids, Binding, 78-6326

Microsomes, Liver**Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy- (cont'd)**

- Nucleic Acids, Binding, 78-6326

Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxo-7,8,9,10-tetrahydro-**Benzo(a)pyrene**

- Carcinogenic Metabolite, 78-6326

Cell Transformation, Neoplastic

- Collisional Quenchers, 78-6329

Fibroblasts

- Fluorescence Quenching, 78-6329

Methane, Diiodo-

- Fluorescence Quenching, 78-6329

Octane, 1-Iodo-

- Fluorescence Quenching, 78-6329

Virus, Moloney Murine Sarcoma

- Fibroblasts, 78-6329

Benzo(a)pyrene 4,5-Oxide*Escherichia coli*

- DNA Repair, 78-6325

Hydro-Lyases

- Metabolism, Lung, 78-6331

Transferases

- Metabolism, Lung, 78-6331

Benzo(b)perylene**Smoke Condensate**

- Isolation and Characterization, 78-6337

1H-Benzo(cd)fluoranthene, 1-Oxo-**Azulenol(1,2,3-cd)phenalene**

- Carcinogenic Metabolite, 78-6289

11H-Benzo(cd)fluoranthene, 11-Oxo-**Azulenol(1,2,3-cd)phenalene**

- Carcinogenic Metabolite, 78-6289

Benzo(f)quinoline**Ames Test**

- Amino Derivatives, 78-6239

Microsomes, Liver

- Ames Test, 78-6239

Benzo(f)quinoline, 2-Amino-**Microsomes, Liver**

- Ames Test, 78-6239

Norharman

- Ames Test, 78-6239

Benzo(h)quinoline**Ames Test**

- Amino Derivatives, 78-6239

Microsomes, Liver

- Ames Test, 78-6239

Benzo(h)quinoline, 2-Amino-**Microsomes, Liver**

- Ames Test, 78-6239

Norharman

- Ames Test, 78-6239

5,6-Benzoflavone**Aryl Hydrocarbon Hydroxylases**

- Strain Difference, Mouse, 78-6319

Hemoexin

- Plasma Levels, Mouse, 78-6319

7,8-Benzoflavone**Benzo(a)pyrene**

- Carcinogenic Metabolite, 78-6326

Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-

- Carcinogenic Metabolite, 78-6326

Methane, Azoxy-

7,8-Benzoflavone (cont'd)

Formaldehyde, 78-6106

Benzoic Acid

Bacillus subtilis

Mutagenic Activity, 78-6256

Nitrous Acid

Mutagenic Activity, 78-6256

Benzoic Acid, 2-Amino-3-hydroxy-

Kidney

Metabolism, Mouse, 78-6311

Lymphoid Tissue

Kidney, 78-6311

Smoking

Metabolism, 78-6567

Urogenital Neoplasms

Carcinogenic Potential, 78-6311

Benzopurpurine 4B

Riboflavine

Ames Test, 78-6263

p-Benzoquinone

Benz(e)indole, 3-(β -Nitrovinyl)-

Benzocarbazole Derivatives, Synthesis, 78-6299

p-Benzoquinone, 2,3,5-Tris(1-aziridinyl)-

Bone Marrow Cells

Chromosome Aberrations, 78-6298

Chromosome Aberrations

Micronucleus Test, 78-6298

Spermatozoa, 78-6298

Benzylamine, N-Methyl-N-nitroso-

Esophageal Neoplasms

Carcinoma, Epidermoid, 78-6273

Carcinoma, Papillary, 78-6273

Neoplasm Invasiveness, 78-6273

Neoplasms, Multiple Primary, 78-6273

Papilloma, 78-6273

Beryllium Oxide

Lung Neoplasms

Inhalation Study, Rat, 78-6127

Plutonium Oxide, 78-6127

Plutonium Oxide

Alveolar Clearance, 78-6127

Bile Acids and Salts

Dietary Fats

Metabolism, Review, 78-6036

Bile Duct Neoplasms

Age Factors

Epidemiology, Sweden, 78-6577

1,1'-Biphenyl, 2,2',4,4',5,5'-Hexachloro-

Cells, Cultured

Metabolism, Lung, 78-6305

4-Biphenylamine

Ames Test

Thiophene Analogs, 78-6260

Cell Transformation, Neoplastic

Thiophene Analogs, 78-6260

Chromosome Aberrations

Micronucleus Test, 78-6162

4-Biphenylamine, 3,2'-Dimethyl-

Colonic Neoplasms

Adenocarcinoma, 78-6297

Carcinogenic Activity, Rat, 78-6297

4-Biphenylamine, 3,2'-Dimethyl- (cont'd)

Glucuronidase, 78-6297

Intestinal Microflora, 78-6297

Duodenal Neoplasms

Carcinogenic Activity, Rat, 78-6297

Ear Neoplasms

Carcinogenic Activity, Rat, 78-6297

Mammary Neoplasms, Experimental

Adenocarcinoma, 78-6297

Carcinogenic Activity, Rat, 78-6297

Glucuronidase, 78-6297

Intestinal Microflora, 78-6297

Salivary Gland Neoplasms

Carcinogenic Activity, Rat, 78-6297

Skin Neoplasms

Carcinogenic Activity, Rat, 78-6297

4,4'Bipyridinium, 1,1'-Dimethyl-, Dichloride

Carbamic Acid, Ethyl Ester

Co-carcinogenic Effect, 78-6184

Lung Neoplasms

Carbamic Acid, Ethyl Ester, 78-6184

Bis(2-chloroisopropyl)ether

see Propane, 2,2'-Oxybis(2-chloro)-

Bladder

1,2-Benzisothiazolin-3-one, 1,1-Dioxide

Absorption, Rat, 78-6281

Nicotine

Body Burden, Mouse, 78-6177

Bladder Neoplasms

Adenocarcinoma

Epidemiology, 78-6567

Aniline, 4,4'-Methylenedi-

Carcinogenic Potential, Dog, 78-6261

Benzidine, 3,3'-Dichloro-

Dose-Response Study, Dog, 78-6270

1,2-Benzisothiazolin-3-one, 1,1-Dioxide

Carcinogenic Potential, Monkey, 78-6294

Carcinogen, Chemical

Epidemiology, Review, 78-6002

Carcinoma

Epidemiology, 78-6567

Carcinoma, Epidermoid

Epidemiology, 78-6567

Carcinoma, Papillary

Epidemiology, 78-6567

Carcinoma, Transitional Cell

Benzidine, 3,3'-Dichloro-, 78-6270

Epidemiology, 78-6567

Coffee

Epidemiology, 78-6567

Epidemiology

South Africa, 78-6567

Hydroxylamine, N-1-Naphthyl-

Carcinogenic Activity, 78-6247

Deoxyguanosine Adducts, 78-6247

Isoantigens

Precancerous Conditions, 78-6566

Lymphoma

Water Pollution, 78-6556

Neoplasms, Multiple Primary

Benzidine, 3,3'-Dichloro-, 78-6270

Occupational Hazard

Epidemiology, 78-6567

Epidemiology, Review, 78-6002

Isoantigens, 78-6566

Head Neoplasms (cont'd)

- Schistosomiasis
 - Epidemiology, 78-6567
- Smoking
 - Epidemiology, 78-6567
 - Tryptophan, 78-6567
- Streptomycin
 - Caffeine
 - Chromosome Aberrations, 78-6120
 - Chromosome Aberrations
 - Cell Cycle Kinetics, 78-6358
 - Endonucleases
 - Chromosome Aberrations, 78-6358
 - Escherichia coli*
 - DNA Repair, 78-6308

Marrow Cells

- p*-Benzoquinone, 2,3,5-Tris(1-aziridinyl)-
 - Chromosome Aberrations, 78-6298
- α -Cresol, 4,6-Dinitro-, Sodium Salt
 - Chromosome Aberrations, 78-6157
- Mercury
 - Chromosome Aberrations, 78-6139
- Mercury, Chloroethyl-
 - Chromosome Aberrations, 78-6139

Neoplasms

- Caribbean Islands
 - Epidemiology, 78-6570
- Radiation, Ionizing
 - In Utero Exposure, Rat, 78-6382

Skin

- Melanoma
 - Neoplasm Metastasis, Review, 78-6060

Skin Neoplasms

- Astrocytoma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6318
 - Virus, B77, 78-6408
- Benz(a)anthracene, 7,12-Dimethyl-
 - Monoamine Oxygenase Inhibitors, 78-6318
 - Serotonin, 78-6318
- Bromine
 - Water Pollution, 78-6556
- Ependymoma
 - Virus, SV40, 78-6486
- Ethnic Groups
 - Epidemiology, Hawaii, 78-6545
- Glioblastoma Multiforme
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6318
- Urea, Ethyl Nitroso-
 - Precancerous Conditions, 78-6231
 - Transplacental Carcinogenesis, 78-6231
- Virus, B77
 - Age Factors, Rat, 78-6408
 - Ultrastructural Study, Rat, 78-6408
- Virus, Papova
 - Antigens, Viral, 78-6486
- Virus, SV40
 - Antigens, Viral, 78-6486

Soft Tissue Neoplasms

- Adenofibroma
 - Antibodies, Viral, 78-6412
 - Precancerous Conditions, 78-6553
- Androst-4-ene-3,17-dione
 - Ethnic Groups, 78-6544
- Antibodies, Viral

Breast Neoplasms (cont'd)

- A-Type Particles, 78-6412
- Caribbean Islands
 - Epidemiology, 78-6570
- Contraceptives, Oral
 - Epidemiology, Review, 78-6026, 78-6027
- Cystosarcoma Phyllodes
 - Case Report, 78-6510
 - Histological Study, 78-6510
- Cysts
 - Precancerous Conditions, 78-6553
- Diet
 - Epidemiology, Review, 78-6037
- Dietary Fats
 - Epidemiology, Review, 78-6036, 78-6067
- Epidemiology
 - Utah, 78-6559
- Estradiol
 - Epidemiology, Review, 78-6037
 - Ethnic Groups, 78-6544
- Estriol
 - Epidemiology, Review, 78-6037
- Estrogens
 - Epidemiology, Review, 78-6027
 - Receptors, Hormone, Review, 78-6024
- Estrone
 - Epidemiology, Review, 78-6037
 - Ethnic Groups, 78-6544
- Ethnic Groups
 - Epidemiology, Cameroon, 78-6568
 - Epidemiology, Hawaii, 78-6546
- Genetics
 - Blood Groups, Review, 78-6068
 - Epidemiology, 78-6543
 - Epidemiology, Review, 78-6067, 78-6068, 78-6073
 - Estradiol, 78-6349
 - Estriol, 78-6349
 - Estrone, 78-6349
 - FSH, 78-6349
 - LH, 78-6349
 - Prolactin, 78-6349
- Hormones
 - Epidemiology, 78-6349
 - Epidemiology, Review, 78-6067
- Ketosteroids
 - Ethnic Groups, 78-6544
- Mammography
 - Dose Fractionation, Review, 78-6050
- Melatonin
 - Pineal Gland Function, Review, 78-6024
 - Receptors, Hormone, Review, 78-6024
- Menopause
 - Epidemiology, 78-6543
- Obesity
 - Hormones, Review, 78-6037
 - Precancerous Conditions, 78-6553
- Occupational Hazard
 - Epidemiology, 78-6551
- Oncogenic Viruses
 - Vertical Transmission, Review, 78-6068
- Parity
 - Precancerous Conditions, 78-6553
- Precancerous Conditions
 - Age Factors, 78-6553
 - Epidemiology, 78-6553
- Progesterone
 - Ethnic Groups, 78-6544

Breast Neoplasms (cont'd)

- Prolactin
 - Dietary Fats, Review, 78-6037
- Radiation, Ionizing
 - Age Factors, Review, 78-6050
 - Epidemiology, Review, 78-6050
- Radium
 - Epidemiology, 78-6564
- Tranquilizing Agents
 - Epidemiology, 78-6243
- Virus, Murine Mammary Tumor
 - Antibodies, Viral, 78-6412

Brevital

- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-6335

Bromine

- Brain Neoplasms
 - Water Pollution, 78-6556
- Lung Neoplasms
 - Water Pollution, 78-6556
- Lymphoma
 - Water Pollution, 78-6556
- Water Pollution
 - Epidemiology, 78-6556

Bronchial Neoplasms

- Petroleum
 - Carcinogenic Activity, Review, 78-6002

Brown FK

- see C.I. Food Brown 1

Burkitt's Lymphoma

- DNA Polymerase
 - Isolation and Characterization, 78-6453
 - Phosphonoacetic Acid, 78-6453
- Ethnic Groups
 - Epidemiology, Cameroon, 78-6568
- Histocompatibility Antigens
 - Genetics, 78-6459
- Immune Serums
 - Antigen-Antibody Reactions, 78-6498
- Oncogenic Viruses
 - Epidemiology, Review, 78-6031
- Virus, Epstein-Barr
 - Antigens, Viral, 78-6459
 - Carcinogenic Activity, Review, 78-6084
 - DNA Polymerase, 78-6453
 - Epidemiology, Review, 78-6089
 - Genetics, 78-6459
- Virus, Herpes Simplex
 - DNA Polymerase, 78-6453

1,3-Butadiene

- Leukemia
 - Occupational Hazard, Review, 78-6014

Butane, 1,2,3,4-Diepoxy-

- Escherichia coli*
 - DNA Repair, 78-6325

1,4-Butanediol, Dimethylsulfonate

- Ames Test
 - Mutagenic Activity, 78-6163
- Antineoplastic Agents
 - Ames Test, 78-6232
- Immunosuppression
 - Carcinogenic Activity, Review, 78-6051
- Virus, Friend Murine Leukemia

1,4-Butanediol, Dimethylsulfonate (cont'd)

- Hematopoietic Stem Cells, 78-6424

1-Butanone, 4-(*N*-Carbethoxy-*N*-nitrosamino)-1-(3-pyridyl)-

- 2-Furanol 5-(3-Pyridyl)tetrahydro-
 - Hydroxylation, 78-6175

1-Butanone, 4-Hydroxy-1-(3-pyridyl)-

- Ames Test
 - Mutagenic Activity, 78-6175
- Nicotine, 1'-Nitroso-1'-demethyl-
 - Hydroxylation, 78-6175

1-Butanone, 4-(*N*-Methyl-*N*-nitrosamino)-1-(3-pyridyl)-

- Nicotine
 - Synthesis, 78-6176

Butylated Hydroxyanisole

- see Phenol, (1,1-Dimethylethyl)-4-methoxy-

2-Butynedioic Acid, Dimethyl Ester

- Benz(e)indole, 3-(β -Nitrovinyl)-
 - Benzocarbazole Derivatives, Synthesis, 78-6299

Butyraldehyde, 4-Hydroxy-, 2,4-Dinitrophenylhydrazon

- Pyrrolidine, 1-Nitroso-
 - Carcinogenic Metabolite, 78-6237

Butyraldehyde, 4-(*N*-Methyl-*N*-nitrosamino)-4-(3-pyridyl)-

- Nicotine
 - Synthesis, 78-6176

Butyric Acid, 2-Amino-4-(ethylthio)-

- Ames Test
 - Mutagenic Activity, 78-6206
- Bacillus subtilis*
 - Mutagenic Activity, 78-6206
- Chromosome Aberrations
 - Micronucleus Test, 78-6162
- Concanavalin A
 - DNA Replication, 78-6205
 - Lymphocyte Transformation, 78-6205
- DNA Repair
 - Liver, Rat, 78-6216
- DNA Replication
 - Pancreas, Rat, 78-6212
- Escherichia coli*
 - Mutagenic Activity, 78-6206
- Lymphocytes
 - Cell Cycle Kinetics, 78-6205
 - RNA, Transfer, Methyltransferases, 78-6205
- Plant Agglutinins
 - DNA Replication, 78-6205
 - Lymphocyte Transformation, 78-6205
- RNA, Transfer, Methyltransferases
 - Liver, Rat, 78-6265

Butyric Acid, 4-(*p*-Bis(2-Chloroethyl)aminophenyl)-

- Immunosuppression
 - Carcinogenic Activity, Review, 78-6051
- Leukemia, Myeloblastic
 - Arthritis, Rheumatoid, 78-6211
 - Case Report, 78-6209, 78-6211
 - Hodgkin's Disease, 78-6196
 - Nephrosis, 78-6211
- Lymphoma
 - Case Report, 78-6209

Butyric Acid, 4-Hydroxy-4-(3-pyridyl)-

- Nicotine, 1'-Nitroso-1'-demethyl-
 - Urinary Metabolite, 78-6175

ric Acid, 4-Oxo-4-(3-pyridyl)-
 Nicotine, 1'-Nitroso-1'-demethyl-
 Urinary Metabolite, 78-6175
ric Acid, Sodium Salt
 Erythroleukemia
 Cell Differentiation, 78-6426
Acid Red 26, Disodium Salt
 Riboflavine
 Ames Test, 78-6263
Direct Rod 28, Disodium Salt
 Riboflavine
 Ames Test, 78-6263
Food Brown 1
 Bacteria
 DNA Repair, 78-6153
 Mutagenic Activity, 78-6153
 Dyes
 Mutagenic Activity, 78-6153
Food Brown 3, Disodium Salt
 Hemoglobins
 Toxicity, Pig, 78-6149
 Leukocytes
 Toxicity, Pig, 78-6149
ium
 Chlorophyll
 Mutagenic Activity, 78-6134
 Stomach Neoplasms
 Food Contamination, 78-6148
ine
 Atrophy
 Testis, Rat, 78-6123
 Bleomycin
 Chromosome Aberrations, 78-6120
 Cholesterol
 Plasma Levels, 78-6123
 Chromatids
 Vicia faba, 78-6120
 Chromosome Aberrations
 Vicia faba, 78-6120
 Erythroleukemia
 Adenosine Cyclic 3',5' Monophosphate, 78-6426
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Chromosome Aberrations, 78-6120
 Hyperplasia
 Leydig Cells, 78-6123
 Methanesulfonic Acid, Ethyl Ester
 Chromosome Aberrations, 78-6120
 Nitrous Acid, Sodium Salt
 Spermatogenesis, 78-6123
 3,6-Pyridazinedione, 1,2-Dihydro-
 Chromosome Aberrations, 78-6120
 Quinacrine Mustard
 Chromosome Aberrations, 78-6120
 Radiation, Ionizing
 Chromosome Aberrations, 78-6120
 Theophylline
 Mutagenic Activity, 78-6122
 Ultraviolet Rays
 DNA Repair, 78-6121
 Saccharomyces cerevisiae, 78-6121
 Urea, Methyl Nitroso-
 Chromosome Aberrations, 78-6120

Calcitonin

Carcinoma, Bronchogenic
 Immune Serums, 78-6594
 Radioimmunoassay, 78-6594
 Thyroid Neoplasms
 Genetics, 78-6522
 Immune Serums, 78-6594
 Radioimmunoassay, 78-6594

Calcium

Cell Membrane Permeability
 Cell Transformation, Neoplastic, Review, 78-6061
 Lead
 Chromosome Aberrations, 78-6147
 Mitochondria
 Cell Transformation, Neoplastic, Review, 78-6061
 Zinc
 Chromosome Aberrations, 78-6147

Captan

see 4-Cyclohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-

Carbamic Acid, Ethyl Ester

4,4'-Bipyridinium, 1,1'-Dimethyl-, Dichloride
 Co-carcinogenic Effect, 78-6184
 Carbamic Acid, Vinyl Ester
 Carcinogenic Metabolite, 78-6185
 Chromosome Aberrations
 Micronucleus Test, 78-6162
 p-Cresol, 2,6-Di-*tert*-butyl-
 Co-carcinogenic Effect, 78-6184
 DNA Replication
 Lung, Mouse Embryo, 78-6183
 Lung Neoplasms
 Adenoma, 78-6185
 4,4'-Bipyridinium, 1,1'-Dimethyl-, Dichloride
 78-6184
 Carcinogenic Activity, Mouse, 78-6185
 p-Cresol, 2,6-Di-*tert*-butyl-, 78-6184
 Nucleic Acids
 Liver, Binding, 78-6185
 Proteins
 Liver, Binding, 78-6185
 Skin Neoplasms
 Carcinogenic Activity, Mouse, 78-6185
 Papilloma, 78-6185

Carbamic Acid, Methyl-, 1-Naphthyl Ester

Cells, Cultured
 Metabolism, Lung, 78-6305

Carbamic Acid, *N*-Methyl-*N*-nitroso-, Ethyl Ester

DNA Replication
 Pancreas, Rat, 78-6212

Carbamic Acid, Vinyl Ester

Ames Test
 Cytochrome P-450 Inhibitors, 78-6185
 Mutagenic Activity, 78-6185
 Carbamic Acid, Ethyl Ester
 Carcinogenic Metabolite, 78-6185
 Lung Neoplasms
 Adenoma, 78-6185
 Carcinogenic Activity, Mouse, 78-6185
 Skin Neoplasms
 Carcinogenic Activity, Mouse, 78-6185
 Papilloma, 78-6185

Carbamoyl Chloride, Diethyl-

Occupational Hazard

- Carbamoyl Chloride, Diethyl- (cont'd)**
Chromosome Aberrations, 78-6187
- Carbamoyl Chloride, Dimethyl-**
Occupational Hazard
Chromosome Aberrations, 78-6187
- Carbanilic Acid, *m*-Chloro-, Isopropyl Ester**
Food Contamination
Carcinogenic Potential, 78-6186
Maximum Permissible Exposure Level, 78-6186
- Carbazilquinone**
Antineoplastic Agents
Ames Test, 78-6232
- Carbendazim**
see 2-Benzimidazolecarbamic Acid, Methyl Ester
- Carbohydrates**
Glycoproteins
Cell Transformation, Neoplastic, Review, 78-6056
- Carbon**
Ames Test
Water, Recycled, 78-6105
- Carbon Dioxide**
Lung Neoplasms
Air Pollution, 78-6019
Epidemiology, Review, 78-6019
- Carbon Radioisotopes**
Air Pollution
Dose-Response Study, 78-6386
Nuclear Power, 78-6386
Neoplasms, Radiation-Induced
Dose-Response Study, 78-6386
- Carbon Tetrachloride**
DNA Repair
Liver, Rat, 78-6216
RNA, Transfer, Methyltransferases
Liver, Rat, 78-6265
- Carbonyl Cyanide, 3-Chlorophenylhydrazino-**
Clostridium leptum
Oxidoreductases, 78-6590
- Carboxy-Lyases**
Virus, Murine Leukemia
Cell Transformation, Neoplastic, 78-6434
Virus, Helper, 78-6434
Virus, Murine Sarcoma
Cell Transformation, Neoplastic, 78-6434
Enzymatic Activity, 78-6434
Virus Replication, 78-6434
- Carcinoembryonic Antigen**
Cell Transformation, Neoplastic
Chromatin, 78-6072
Intestinal Neoplasms
Immune Response, Review, 78-6083
Liver Neoplasms
Adenoma, 78-6352
Rectal Neoplasms
Adenocarcinoma, 78-6596
Adenosine Cyclic 3',5' Monophosphate, 78-6596
Cells, Cultured, 78-6596
Dibutyl Cyclic AMP, 78-6596
- Carcinogen, Chemical**
Air Pollution
Dose-Response Study, 78-6110
- Carcinogen, Chemical (cont'd)**
Ames Test
Mutagenic Activity, 78-6100, 78-6206
Mutagenic Activity, Review, 78-6005
Aroclor 1254
Ames Test, 78-6166
Mutagenic Activity, 78-6166
Automobile Exhaust
Isolation and Characterization, 78-6300
Bacillus subtilis
Mutagenic Activity, 78-6206
Bladder Neoplasms
Epidemiology, Review, 78-6002
Carcinogenic Metabolite
DNA, Binding, Review, 78-6003
Enzyme Activation, Review, 78-6003
Carcinogenic/Mutagenic Activity
Modification, Review, 78-6001
Cell Membrane Permeability
Cell Transformation, Neoplastic, Review, 78-6061
Cell Transformation, Neoplastic
Bioassay, Hamster Embryo Cells, 78-6104
Dose-Response Study, Review, 78-6004
Precancerous Conditions, Review, 78-6084
Cells, Cultured
Screening Test, Review, 78-6007
Coal Tar
Occupational Hazard, Review, 78-6017
Colonic Neoplasms
Dietary Fiber, 78-6040
Cyclohexene Oxide
Ames Test, 78-6166
Mutagenic Activity, 78-6166
Cytochrome P-450
Enzyme Activation, 78-6340
Dietary Fiber
Metabolism, Review, 78-6040
DNA
Carcinogenic Activity, 78-6154
DNA Repair
Alkaline Elution Assay, Review, 78-6053
Autoradiography, Review, 78-6053
Bioassay, 78-6099
Carcinogenic Activity, 78-6099
Carcinogenic Activity, Review, 78-6054
Screening Test, Review, 78-6053
Dose-Response Study
Risk Factors, 78-6110
Environmental Hazard
Quantitation Method, 78-6119
Escherichia coli
Mutagenic Activity, 78-6206
Genetics
DNA Repair, Review, 78-6055
Glutathione
Carcinogenic Activity, Review, 78-6004
Hepatitis
Virus-Like Particles, Review, 78-6096
Hydro-Lyases
Carcinogenic Metabolite, 78-6166
Intestinal Neoplasms
Bacteria, Review, 78-6035
Melanoma
Mathematical Model, Review, 78-6044
Microsomes, Liver
Enzyme Activation, 78-6340
Mitochondria

inogen, Chemical (cont'd)

- Oxidative Phosphorylation, Review, 78-6061
- Mutagenic Activity
 - Assay System, 78-6100
- Occupational Hazard
 - Epidemiology, 78-6551
 - Women, Review, 78-6022
- Pentafluoropropionyl Derivatives
 - Quantitation Method, 78-6119
- Precancerous Conditions
 - Carcinogenic Activity, Review, 78-6003
- Retinol
 - Carcinogenic Activity, Review, 78-6004
- Ultraviolet Rays
 - Spectrum Analysis, 78-6154
- Virus, RNA Tumor
 - Co-carcinogenic Effect, Review, 78-6096
- Vitamin E
 - Carcinogenic Activity, Review, 78-6004

inogen, Environmental

- Nasopharyngeal Neoplasms
 - Epidemiology, 78-6064
- Neoplasms
 - Epidemiology, Review, 78-6030
- Nitrosamines
 - Quantitation Method, 78-6006
- Nitrous Acid
 - Quantitation Method, 78-6006
- Polycyclic Hydrocarbons
 - Quantitation Method, 78-6301
- Prostatic Neoplasms
 - Epidemiology, 78-6558

inoid Tumor

- Lung Neoplasms
 - APUD Cells, Review, 78-6063

inoma

- Aniline, *p*-(Phenylazo)-
 - Immunosuppression, 78-6278
 - Transplantation Immunology, 78-6278
- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-
 - Immunosuppression, 78-6278
 - Transplantation Immunology, 78-6278
- Bladder Neoplasms
 - Epidemiology, 78-6567
- Colonic Neoplasms
 - Colitis, Ulcerative, 78-6537
- Hypertension
 - Neoplasm Transplantation, 78-6332
- Immunosuppression
 - Transplantation Immunology, 78-6278
- Intestinal Neoplasms
 - Bacteria, 78-6587
 - Glucuronidase, 78-6587
 - Hydroxylases, 78-6587
 - Oxidoreductases, 78-6587
- Kidney Neoplasms
 - Radiotherapy, 78-6384
- Lung Neoplasms
 - APUD Cells, Review, 78-6063
 - Epidemiology, 78-6547
- Mammary Neoplasms, Experimental
 - Benidine, 2,2',5,5'-Tetrachloro-, 78-6254
 - Xanthine, 3-Hydroxy-, 78-6250
- Nasopharyngeal Neoplasms
 - Ultrastructural Study, 78-6064
- Respiratory Tract Neoplasms

Carcinoma (cont'd)

- Occupational Hazard, Review, 78-6049
- Skin Neoplasms
 - Virus, Papilloma, 78-6091
- Stomach Neoplasms
 - Gastrectomy, 78-6576
 - Genetics, 78-6575
- Thyroid Neoplasms
 - Epidemiology, 78-6523
 - Genetics, 78-6522
 - Neoplasm Metastasis, 78-6363, 78-6379, 78-6522
 - Radiation, Ionizing, 78-6363, 78-6523
 - Ultrastructural Study, 78-6505
- Virus, RD-114
 - RNA, Viral, 78-6441

Carcinoma, Bronchiolar

- Lung Neoplasms
 - Epidemiology, 78-6547

Carcinoma, Bronchogenic

- Benz(a)anthracene
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Calcitonin
 - Immune Serums, 78-6594
 - Radioimmunoassay, 78-6594
- Cholanthrene, 3-Methyl-
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Lymphocytes
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Smoking
 - Aryl Hydrocarbon Hydroxylases, 78-6322

Carcinoma, Cloacogenic

- see* Carcinoma, Transitional Cell

Carcinoma, Ductal

- Mammary Neoplasms, Experimental
 - Acenaphthene, 5-Nitro-, 78-6254

Carcinoma, Epidermoid

- Bladder Neoplasms
 - Epidemiology, 78-6567
- Ear Neoplasms
 - Acenaphthene, 5-Nitro-, 78-6254
- Esophageal Neoplasms
 - Animal Model, Rat, 78-6273
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273
 - Epidemiology, 78-6549
 - Ethyl Alcohol, 78-6549
 - Smoking, 78-6549
- Eye Neoplasms
 - Epidemiology, Dog, 78-6516
- Head and Neck Neoplasms
 - Epidemiology, 78-6549
 - Ethyl Alcohol, 78-6549
 - Smoking, 78-6549
- Laryngeal Neoplasms
 - Epidemiology, 78-6380
- Lung Neoplasms
 - Asbestos, 78-6529
 - Occupational Hazard, 78-6529
- Mouth Neoplasms
 - Epidemiology, 78-6375
 - Ethyl Alcohol, 78-6375
 - Smoking, 78-6375
 - Wounds and Injuries, 78-6375
- Pharyngeal Neoplasms
 - Epidemiology, 78-6380
- Respiratory Tract Neoplasms

Carcinoma, Epidermoid (cont'd)

- Benzo(a)pyrene, 78-6335
- Occupational Hazard, Review, 78-6049

Spinal Neoplasms

- Meningomyelocele, 78-6519

Stomach Neoplasms

- Piperazine, 1,4-Dinitroso-, 78-6241
- Plants, Edible, 78-6210

Carcinoma In Situ

- Lung Neoplasms
- Precancerous Conditions, 78-6528

Carcinoma, Papillary

- Bladder Neoplasms
- Epidemiology, 78-6567
- Esophageal Neoplasms
- Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273
- Thyroid Neoplasms
- Radiation, Ionizing, 78-6376, 78-6378

Carcinoma, Transitional Cell

- Anus Neoplasms
- Ultrastructural Study, 78-6539
- Bladder Neoplasms
- Benidine, 3,3'-Dichloro-, 78-6270
- Epidemiology, 78-6567
- Kidney Neoplasms
- p*-Acetophenetidine, 78-6293
- Codeine, 78-6293

Carcinosarcoma

- Hypertension
- Neoplasm Metastasis, 78-6332

Carrageen

- Antigens
- Hypersensitivity, Delayed, 78-6490
- Fibrosarcoma
- Transplantation Immunology, 78-6490
- Macrophages
- Transplantation Immunology, 78-6490

Casein

- RNA, Messenger
- Mammary Development, 78-6410

Cecal Neoplasms

- Adenocarcinoma
- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-78-6286
- Cholanthrene, 3-Methyl-, 78-6321
- Adenoma
- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-78-6286
- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-Carcinogenic Activity, Mouse, 78-6286
- Neoplasms, Multiple Primary
- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385
- Radiation, Ionizing
- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385

Cell Adhesion

- Neoplasms
- Immunoglobulins, Fc, 78-6504
- Virus, Polyoma
- Cell Transformation, Neoplastic, 78-6480
- Ultrastructural Study, BHK Cells, 78-6480

Cell Agglutination

- Virus, Adeno 12
- Concanavalin A, 78-6470
- Morphological Revertants, 78-6470

Cell Differentiation

- Disgerminoma
- Sex Chromosomes, 78-6506
- Erythroleukemia
- Adenosine Cyclic 3',5' Monophosphate, 78-6426
- Butyric Acid, Sodium Salt, 78-6426
- Cell Cycle Kinetics, 78-6426
- Hexamethylene Bisacetamide, 78-6426
- Methane Sulfinylbis-, 78-6426
- Virus, Friend Murine Leukemia, 78-6426
- Neoplasms
- Drosophila melanogaster*, 78-6075
- Neoplasms, Experimental
- Virus, SV40, 78-6085
- Neuraminidase
- Trophoblast, 78-6591
- Teratoid Tumor
- Retinoic Acid, 78-6339
- Turner's Syndrome
- Case Report, 78-6506
- Testis, 78-6506
- Virus, Avian Erythroblastosis
- Cell Transformation, Neoplastic, 78-6390
- Virus, Friend Spleen Focus-Forming
- A-Type Particles, 78-6428
- Virus-Like Particles, 78-6428
- Virus, Murine Leukemia
- Age Factors, 78-6422

Cell Division

- Fibrosarcoma
- Peptides, 78-6481
- Manganese Chloride
- Mutagenic Activity, Algae, 78-6137

Cell Fusion

- Virus, Gross Murine Leukemia
- Glycoproteins, 78-6431
- Virus, Moloney Murine Leukemia
- Glycoproteins, 78-6431
- Virus, Rauscher Murine Leukemia
- Glycoproteins, 78-6431

Cell Membrane

- Glycopeptides
- Trophoblast, 78-6591
- Glycoproteins
- Cell Transformation, Neoplastic, Review, 78-6056
- 78-6057
- Fibroblasts, 78-6593
- Isolation and Characterization, Review, 78-6056
- Virus, Bovine Papilloma
- Antigens, Viral, 78-6446
- Virus, Epstein-Barr
- Antigen-Antibody Reactions, 78-6460
- Antigens, 78-6460
- Virus, Feline Leukemia
- Antigens, Viral, 78-6443
- Virus, Rous Sarcoma
- Antigens, Viral, 78-6402

Cell Membrane Permeability

- Calcium
- Cell Transformation, Neoplastic, Review, 78-6061
- Carcinogen, Chemical

Membrane Permeability (cont'd)

- Cell Transformation, Neoplastic, Review, 78-6061
- Oncogenic Viruses
- Cell Transformation, Neoplastic, Review, 78-6061

Movement**Macrophages**

- Immune Response, Review, 78-6079

Transformation, Neoplastic**Acetamide, *N*-(5-*p*-Acetamidophenylthiophen-2-yl)-**

- Cells, Cultured, 78-6260

Acetamide, *N*-(5-Phenylthiophen-2-yl)-

- Cells, Cultured, 78-6260

Aflatoxin B1

- Diploid Cells, 78-6245

Antigens, Neoplasm

- Immune Response, Review, 78-6077
- Immunologic Technics, Review, 78-6077

Areca

- Tobacco Extract, 78-6173

Benzo(a)pyrene

- Collisional Quenchers, 78-6329

Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-

- Collisional Quenchers, 78-6329

4-Biphenylamine

- Thiophene Analogs, 78-6260

Carcinogen, Chemical

- Bioassay, Hamster Embryo Cells, 78-6104
- Dose-Response Study, Review, 78-6004
- Precancerous Conditions, Review, 78-6084

Cervix Uteri

- Organ Culture, 78-6599

Cesium Radioisotopes

- Mathematical Model, 78-6048

Chromatin

- Carcinoembryonic Antigen, 78-6072
- Gene Control Mechanisms, 78-6072
- Proteins, 78-6072
- RNA, Messenger, 78-6072

Chromosome Aberrations

- Review, 78-6070

Cyclophosphamide

- Cells, Cultured, 78-6195
- Oxidoreductases, 78-6195

DNA Repair

- Theoretical Model, Review, 78-6008

Fluorosulfuric Acid, Methyl Ester

- Dose-Response Study, 78-6165

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Diploid Cells, 78-6245

T-Lymphocytes

- Immune Response, Review, 78-6077
- Immunologic Technics, Review, 78-6077
- Lymphocytotoxicity, Review, 78-6077

Macrophages

- Immune Response, Review, 78-6080
- Phagocytosis, Review, 78-6079

Methanesulfonic Acid, Ethyl Ester

- Diploid Cells, 78-6245

Mutation

- Theoretical Model, Review, 78-6008

Oncogenic Viruses

- Precancerous Conditions, Review, 78-6084

1,2-Oxathiolane, 2,2-Dioxide

- Diploid Cells, 78-6245

2-Oxetanone**Cell Transformation, Neoplastic (cont'd)**

- Diploid Cells, 78-6245

Quinoline, 4-Nitro-, 1-Oxide

- Diploid Cells, 78-6245

Radiation

- Precancerous Conditions, Review, 78-6084

Radiation, Ionizing

- Dose-Response Study, Review, 78-6004

Ruthenium

- Mathematical Model, 78-6048
- Radioisotopes, 78-6048

12-*O*-Tetradecanoylphorbol-13-acetate

- Plasminogen Activators, 78-6310

Thiophene, 5-(*p*-Acetamidophenyl)-2-amino-, Hydrochloride

- Cells, Cultured, 78-6260

Thiophene, 2-Amino-5-phenyl-, Hydrochloride

- Cells, Cultured, 78-6260

Urea, Ethyl Nitroso-

- Neurons, 78-6231
- Ultrastructural Study, Fetal Rat Brain, 78-6231

Virus, Abelson Murine Leukemia

- Macrophages, 78-6419

Virus, Adeno

- Genes, Viral, Review, 78-6086

Virus, Adeno 3

- Cells, Cultured, 78-6464
- DNA, Viral, 78-6464

Virus, Adeno 5

- Chromatin, 78-6466
- DNA, Viral, 78-6466

Virus, Adeno 12

- Morphological Revertants, 78-6470

Virus, Avian Erythroblastosis

- Cell Differentiation, 78-6390
- Hemoglobins, 78-6390

Virus, Avian Leukosis

- Genes, Viral, 78-6387
- Nucleotide Sequence, 78-6387
- Virus, Helper, 78-6388

Virus, Avian Reticuloendotheliosis

- Nucleoproteins, 78-6405

Virus, Avian Sarcoma

- Glucosamine, 78-6394
- Peptides, 78-6481

Virus, B77

- Fibroblasts, 78-6398
- Plasminogen Activators, 78-6398

Virus, Feline Sarcoma

- Antigenic Determinants, 78-6444

Virus, Friend Murine Leukemia

- Erythropoietin, 78-6424
- Hematopoietic Stem Cells, 78-6424
- RNA, Messenger, 78-6580

Virus, Gross Murine Leukemia

- Antigens, Viral, 78-6432

Virus, Marek's Disease Herpes

- Virus Replication, 78-6393

Virus, Moloney Murine Leukemia

- Antigens, Viral, 78-6432
- B-Lymphocytes, 78-6432
- T-Lymphocytes, 78-6432

Virus, Moloney Murine Sarcoma

- Viral Proteins, 78-6437

Virus, Murine Leukemia

- Carboxy-Lyases, 78-6434
- Fibroblasts, 78-6435

Cell Transformation, Neoplastic (cont'd)

- Virus, Murine Sarcoma
 - Carboxy-Lyases, 78-6434
 - Fibroblasts, 78-6435
 - Peptides, 78-6481
- Virus, Polyoma
 - Antigens, 78-6496
 - Cell Adhesion, 78-6480
 - Fibroblasts, 78-6480
 - Genes, Viral, Review, 78-6086
 - Peptides, 78-6481
 - Temperature Sensitive Mutants, 78-6478
- Virus, RNA Tumor
 - Genes, Viral, Review, 78-6086
- Virus, Rous Sarcoma
 - Antigenic Determinants, 78-6401
 - Antigens, Viral, 78-6402
 - Fibroblasts, 78-6398
 - Glycolysis, 78-6396
 - Plasminogen Activators, 78-6338, 78-6395, 78-6397, 78-6398
 - Temperature Sensitive Mutants, 78-6396, 78-6402
 - Virus Replication, 78-6393
- Virus, SV40
 - Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-, 78-6242
 - Aflatoxin B1, 78-6242
 - Benzo(a)pyrene, 78-6242
 - Diethylamine, *N*-Nitroso-, 78-6242
 - Genes, Viral, Review, 78-6086
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-6242
 - Methanesulfonic Acid, Methyl Ester, 78-6242
 - 2-Naphthylamine, 78-6242
 - 1,2-Oxathiolane, 2,2-Dioxide, 78-6242
 - 2-Oxetanone, 78-6242
 - Peptides, 78-6481
 - Plasminogen Activators, 78-6310
 - Quinoline, 4-Nitro-, 1-Oxide, 78-6242
 - Virus, Shope Rabbit Papilloma, 78-6475

Cells, Cultured

- Acetamide, *N*-(5-*p*-Acetamidophenylthiophen-2-yl)-
 - Cell Transformation, Neoplastic, 78-6260
- Acetamide, *N*-(5-Phenylthiophen-2-yl)-
 - Cell Transformation, Neoplastic, 78-6260
- Acetic Acid, (2,4-Dichlorophenoxy)-
 - Metabolism, Lung, 78-6305
- Acetic Acid, (2,4,5-Trichlorophenoxy)-
 - Metabolism, Lung, 78-6305
- Aflatoxin B1
 - Colony Formation, Agar, 78-6245
- Aldrin
 - Metabolism, Lung, 78-6305
- Areca
 - Tobacco Extract, 78-6173
- Benzene, Hexachloro-
 - Metabolism, Lung, 78-6305
- Benzidine
 - Thiophene Analogs, 78-6260
- 1,1'-Biphenyl, 2,2',4,4',5,5'-Hexachloro-
 - Metabolism, Lung, 78-6305
- Carbamic Acid, Methyl-, 1-Naphthyl Ester
 - Metabolism, Lung, 78-6305
- Carcinogen, Chemical
 - Screening Test, Review, 78-6007
- Cyclophosphamide
 - Cell Transformation, Neoplastic, 78-6195
 - Chromosome Aberrations, 78-6195
- Dieldrin

Cells, Cultured (cont'd)

- Metabolism, Lung, 78-6305
- Dimethylamine, *N*-Nitroso-
 - Methylation, 78-6217
- Ethane, 1,1,1-Trichloro-2,2-bis(*p*-chlorophenyl)-
 - Metabolism, Lung, 78-6305
- Formamidine, *N*-(4-Chloro-*o*-tolyl)-*N,N*-dimethyl-
 - Metabolism, Lung, 78-6305
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Colony Formation, Agar, 78-6245
- Hepatoma
 - Hemopexin, 78-6319
- Infectious Mononucleosis
 - Mutation, 78-6164
- Lung Neoplasms
 - Neoplasm Transplantation, 78-6598
- Malathion
 - Metabolism, Lung, 78-6305
- Mesenchymoma
 - Aflatoxin B1, 78-6245
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-, 78-6245
 - Methanesulfonic Acid, Ethyl Ester, 78-6245
 - 1,2-Oxathiolane, 2,2-Dioxide, 78-6245
 - 2-Oxetanone, 78-6245
 - Quinoline, 4-Nitro-, 1-Oxide, 78-6245
- Methanesulfonic Acid, Ethyl Ester
 - Colony Formation, Agar, 78-6245
- 1,2-Oxathiolane, 2,2-Dioxide
 - Colony Formation, Agar, 78-6245
- 2-Oxetanone
 - Colony Formation, Agar, 78-6245
- Phosphorothioic Acid, *O,O*-Diethyl *O*-(*p*-Nitrophenyl) Ester
 - Metabolism, Lung, 78-6305
- Quinoline, 4-Nitro-, 1-Oxide
 - Colony Formation, Agar, 78-6245
- Rectal Neoplasms
 - Carcinoembryonic Antigen, 78-6596
- Retinol
 - Contact Inhibition, 78-6592
 - Glycolipids, 78-6592
 - Glycoproteins, 78-6592
 - Surface Properties, 78-6592
- Spherocytosis, Hereditary
 - Mutation, 78-6164
- Thiophene, 5-(*p*-Acetamidophenyl)-2-amino-, Hydrochloride
 - Cell Transformation, Neoplastic, 78-6260
- Thiophene, 2-Amino-5-phenyl-, Hydrochloride
 - Cell Transformation, Neoplastic, 78-6260
- Virus, Adeno 3
 - Antigens, Viral, 78-6464
 - Cell Transformation, Neoplastic, 78-6464
- Virus, Shope Rabbit Papilloma
 - Virus Interactions, 78-6475

Cervix Neoplasms

- Caribbean Islands
 - Epidemiology, 78-6570
- Ethnic Groups
 - Epidemiology, 78-6029
 - Epidemiology, Cameroon, 78-6568
 - Epidemiology, Hawaii, 78-6546
- Histones
 - Sperm Proteins, 78-6569
- Menarche
 - Epidemiology, 78-6573
- Menopause

vix Neoplasms (cont'd)

- Epidemiology, 78-6573
 - Myoma
 - Organ Culture, 78-6599
 - Occupational Hazard
 - Epidemiology, 78-6551
 - Oncogenic Viruses
 - Epidemiology, Review, 78-6031
 - Parity
 - Epidemiology, 78-6542
 - Precancerous Conditions
 - Epidemiology, Denmark, 78-6571
 - Protamines
 - Sperm Proteins, 78-6569
 - Sex Behavior
 - Epidemiology, 78-6029, 78-6542
 - Epidemiology, Denmark, 78-6571
 - Socioeconomic Factors
 - Epidemiology, 78-6569
 - Sperm Proteins
 - Epidemiology, 78-6569
 - Virus, Herpes Simplex 2
 - Anti-Antibodies, 78-6449
 - Antigen-Antibody Reactions, 78-6449
 - Carcinogenic Activity, Review, 78-6084
 - Seroepidemiology, 78-6449
 - Seroepidemiology, Review, 78-6095
- vix Uteri**
- Cell Transformation, Neoplastic
 - Organ Culture, 78-6599
 - Metaplasia
 - Organ Culture, 78-6599

sium Chloride

- Bacillus subtilis*
 - Mutagenic Activity, 78-6128

sium Radioisotopes

- Cell Transformation, Neoplastic
 - Mathematical Model, 78-6048

lorambucil

- see* Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-

loramphenicol

- Ultraviolet Rays
 - DNA Repair, Review, 78-6052

lorazol Violet N

- Riboflavine
 - Ames Test, 78-6263

lorine

- Ames Test
 - Bleaching Effluents, Kraft Mills, 78-6132
- Aniline
 - Chlorination Products, 78-6262
- Aniline, *N,N*-Dimethyl-
 - Chlorination Products, 78-6262
- Benidine
 - Chlorination Products, 78-6262
- Benzo(a)pyrene
 - Co-carcinogenic Activity, 78-6330
- Skin Neoplasms
 - Benzo(a)pyrene, 78-6330

loroform

- Alanine Aminotransferase
 - Hepatotoxicity, Rat, 78-6109
- Barbituric Acid, 5-Ethyl-5-phenyl-

Chloroform (cont'd)

- Metabolism, 78-6109
- Deuterium
 - Hepatotoxicity, Rat, 78-6109
 - Metabolism, 78-6109
- Hexamethylenetetramine
 - Nitrosation, 78-6225
- Kidney Neoplasms
 - Water Pollution, 78-6556
- Necrosis
 - Hepatotoxicity, Rat, 78-6109
- Phosgene
 - Cytochrome P-450, 78-6109
 - Metabolism, 78-6109
- Water Pollution
 - Epidemiology, 78-6556

***m*-Chloroperbenzoic Acid**

- see* Benzenecarboperoxoic Acid, 3-Chloro-

Chloroplatinic Acid

- Bacillus subtilis*
 - Mutagenic Activity, 78-6128

Chloroquine

- see* Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)

Chocolate Brown HT

- see* C.I. Food Brown 3, Disodium Salt

5 β -Cholan-24-oic Acid, 3 α ,7 α -Dihydroxy-

- Bacteroides
 - Oxidoreductases, 78-6590
- Clostridium leptum*
 - Oxidoreductases, 78-6590

Cholangioma

- Acenaphthene, 5-Nitro-
 - Carcinogenic Potential, 78-6254
- Liver Neoplasms
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6276

Cholangitis

- Lymphosarcoma
 - Case Report, 78-6513

Cholanthrene, 3-Methyl-

- Aryl Hydrocarbon Hydroxylases
 - Strain Difference, Mouse, 78-6319
- Asbestos
 - Aryl Hydrocarbon Hydroxylases, 78-6169
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 - Co-carcinogenic Activity, Mouse Embryo Cells 78-6283
- Benzo(a)pyrene
 - Aryl Hydrocarbon Hydroxylases, 78-6326
 - Carcinogenic Metabolite, 78-6326
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Carcinogenic Metabolite, 78-6326
- Carcinoma, Bronchogenic
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Cecal Neoplasms
 - Adenocarcinoma, 78-6321
- Colonic Neoplasms
 - Adenocarcinoma, 78-6321
- Coumarin, 3, 3'-Methylenebis(4-hydroxy)-
 - Aryl Hydrocarbon Hydroxylases, 78-6334
 - Hepatotoxicity, Rat, 78-6334
- Dimethylamine, *N*-Nitroso-

- Cholanthrene, 3-Methyl- (cont'd)**
 Metabolism, Liver, 78-6217
- Fibrosarcoma**
 Histocompatibility Antigens, 78-6493
 Transplantation Immunology, 78-6490
- Hemopexin**
 Plasma Levels, Mouse, 78-6319
- Hydro-Lyases**
 Enzymatic Activity, 78-6257, 78-6320
- Laryngeal Neoplasms**
 Aryl Hydrocarbon Hydroxylases, 78-6322
- Lymphocytes**
 Aryl Hydrocarbon Hydroxylases, 78-6322
- Lymphoma**
 Dose-Response Study, 78-6321
- Mammary Neoplasms, Experimental**
 Dose-Response Study, 78-6321
 Neoplasm Metastasis, 78-6321
 Neoplasms, Multiple Primary, 78-6321
 Reserpine, 78-6321
- Methane, Azoxy-**
 Formaldehyde, 78-6106
- Microsomes, Liver**
 Enzymatic Activity, 78-6320
 Hydro-Lyases, 78-6320
 Hydroxylases, 78-6320
- Phenanthrene, 9,10-Dihydro-9,10-dihydroxy-**
 Carcinogenic Metabolite, 78-6312
- Plant Agglutinins**
 Aryl Hydrocarbon Hydroxylases, 78-6322
- Reserpine**
 Co-carcinogenic Effect, Hamster, 78-6321
- RNA, Transfer, Methyltransferases**
 Liver, Rat, 78-6265
- Sarcoma**
 Dose-Response Study, 78-6321
- 12-*O*-Tetradecanoylphorbol-13-acetate**
 Co-carcinogenic Activity, Mouse Embryo Cells
 78-6283
- Cholestan-3 β -ol, 5 α ,6 α -Epoxy-**
 Chromosome Aberrations
 Fibroblasts, 78-6362
- Hyperthermia**
 Chromosome Aberrations, 78-6362
- Melanoma**
 Chromosome Aberrations, 78-6362
- Ultraviolet Rays**
 Chromosome Aberrations, 78-6362
 DNA Repair, 78-6362
- Cholesteatoma**
 Ear Neoplasms
 Histological Study, 78-6520
 Precancerous Conditions, 78-6520
 Precancerous Conditions, Review, 78-6062
- Cholesterol**
 Acetamide, *N*-Fluoren-2-yl-
 Metabolism, Liver, 78-6192
- Benzidine**
 Metabolism, Liver, 78-6192
- Caffeine**
 Plasma Levels, 78-6123
- Dietary Fats**
 Metabolism, Review, 78-6036
- Theobromine**
 Plasma Levels, 78-6123
- Cholesterol- α -oxide**
 see Cholestan-3 β -ol, 5 α ,6 α -Epoxy-
- Cholic Acid, Monosodium Salt**
Clostridium leptum
 Oxidoreductases, 78-6590
- Choline**
 Hepatoma
 Serine, Diazoacetate (Ester), 78-6199
- Pancreatic Neoplasms**
 Serine, Diazoacetate (Ester), 78-6199
- Serine, Diazoacetate (Ester)**
 Organ Specificity, 78-6199
- Choroid Plexus Papilloma**
 see Ependymoma
- Chromate**
 Microsomes, Liver
 Metabolism, 78-6129
 Mutagenic Activity, 78-6129
- NADP**
 Metabolism, 78-6129
- Respiratory Tract Neoplasms**
 Epidemiology, 78-6131
 Occupational Hazard, 78-6131
- Salmonella typhimurium*
 Mutagenic Activity, 78-6129
- Chromatids**
 Acetaldehyde
 Lymphocytes, 78-6188
- Caffeine**
Vicia faba, 78-6120
- Cyclophosphamide**
 Diffusion Chamber Cultures, 78-6194
 Oxidoreductases, 78-6195
- Ultraviolet Rays**
 Leukocytes, 78-6356
 Psoralen, 8-Methoxy-, 78-6356
 Psoralen, 4,5',8-Trimethyl-, 78-6356
- Chromatin**
 Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 DNA Repair, 78-6191
- Cell Transformation, Neoplastic**
 Carcinoembryonic Antigen, 78-6072
 Gene Control Mechanisms, 78-6072
 Proteins, 78-6072
 RNA, Messenger, 78-6072
- Dimethylamine, *N*-Nitroso-**
 Purine, 2-Amino-6-methoxy-, 78-6213
- Purine, 2-Amino-6-methoxy-**
 Enzymatic Removal, 78-6213
- Ultraviolet Rays**
 DNA Repair, 78-6357
- Urea, Methyl Nitroso-**
 Purine, 2-Amino-6-methoxy-, 78-6213
- Virus, Adeno 5**
 Cell Transformation, Neoplastic, 78-6466
- Virus, Epstein-Barr**
 Antigens, Viral, 78-6456
- Chromic Acid**
 Erythrocytes
 Ames Test, 78-6130
- Microsomes, Liver**
 Ames Test, 78-6130

omonic Acid, Dipotassium Salt

- Erythrocytes
 - Ames Test, 78-6130
- Microsomes, Liver
 - Ames Test, 78-6130

omonic Acid, Lead Salt

- Erythrocytes
 - Ames Test, 78-6130
- Microsomes, Liver
 - Ames Test, 78-6130
- Saccharomyces cerevisiae*
 - Mutagenic Activity, 78-6304

omonic Acid, Zinc Salt

- Ames Test
 - Mutagenic Activity, 78-6130
- Ascorbic Acid
 - Ames Test, 78-6130
- Erythrocytes
 - Ames Test, 78-6130
- Glutathione
 - Ames Test, 78-6130
- Microsomes, Liver
 - Ames Test, 78-6130
- Permanganic Acid, Potassium Salt
 - Ames Test, 78-6130
- Sulfurous Acid, Disodium Salt
 - Ames Test, 78-6130

romium

- Head and Neck Neoplasms
 - Occupational Hazard, 78-6561
 - Sarcoma, 78-6561
- Kidney Neoplasms
 - Occupational Hazard, 78-6561
- Leukemia, Lymphoblastic
 - Occupational Hazard, 78-6561
- Manganese
 - Co-carcinogenic Activity, 78-6561
- Nickel
 - Co-carcinogenic Activity, 78-6561
- Occupational Hazard
 - Epidemiology, Japan, 78-6561
- Respiratory Tract Neoplasms
 - Epidemiology, Japan, 78-6561
- Sarcoma, Reticulum Cell
 - Occupational Hazard, 78-6561
- Vanadium
 - Co-carcinogenic Activity, 78-6561

romosome Aberrations

- Acetamide, *N*-Fluoren-2-yl-
 - Micronucleus Test, 78-6162
- Antipyrine, 4-(Dimethylamino)-
 - Lymphocytes, Rat, 78-6214
- Benz(a)anthracene, 7,12-Dimethyl-
 - Micronucleus Test, 78-6162
- Benzidine
 - Methanesulfonic Acid, Ethyl Ester, 78-6162
- p*-Benzoquinone, 2,3,5-Tris(1-aziridinyl)-
 - Bone Marrow Cells, 78-6298
 - Micronucleus Test, 78-6298
 - Spermatozoa, 78-6298
- 4-Biphenylamine
 - Micronucleus Test, 78-6162
- Bleomycin
 - Caffeine, 78-6120
 - Cell Cycle Kinetics, 78-6358

Chromosome Aberrations (cont'd)

- Endonucleases, 78-6358
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Micronucleus Test, 78-6162
- Caffeine
 - Vicia faba*, 78-6120
- Carbamic Acid, Ethyl Ester
 - Micronucleus Test, 78-6162
- Carbamoyl Chloride, Diethyl-
 - Occupational Hazard, 78-6187
- Carbamoyl Chloride, Dimethyl-
 - Occupational Hazard, 78-6187
- Cell Transformation, Neoplastic
 - Review, 78-6070
- Cholestan-3 β -ol, 5 α ,6 α -Epoxy-
 - Fibroblasts, 78-6362
 - Hyperthermia, 78-6362
- α -Cresol, 4,6-Dinitro-, Sodium Salt
 - Bone Marrow Cells, 78-6157
 - Dose-Response Study, 78-6157
 - Leukocytes, 78-6157
- Cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer
 - Lymphocytes, Rat, 78-6214
- Cyclophosphamide
 - Cells, Cultured, 78-6195
 - Diffusion Chamber Cultures, 78-6194
 - Micronucleus Test, 78-6162
 - Oxidoreductases, 78-6195
- Diethylamine, *N*-Nitroso-
 - Micronucleus Test, 78-6162
- Dimethylamine, *N*-Nitroso-
 - Lymphocytes, Rat, 78-6214
 - Micronucleus Test, 78-6162
- DNA
 - Cell Cycle Kinetics, 78-6358
- Endonucleases
 - Cell Cycle Kinetics, 78-6358
 - Hamster Ovary Cells, 78-6358
 - Virus, Sendai, 78-6358
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Caffeine, 78-6120
- Heliotrine
 - Piperonyl Butoxide, 78-6238
- Hydrazine, 1,2-Dimethyl-
 - Micronucleus Test, 78-6162
- Imidazole-1-ethanol, 2-Methyl-5-nitro-
 - Micronucleus Test, 78-6162
- Lead
 - Air Pollution, 78-6136
 - Calcium, 78-6147
 - Lymphocytes, 78-6147
- Leukemia
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6314
 - Review, 78-6070
 - Trisomy 15, Mouse, 78-6314
- Leukemia, Myeloblastic
 - Chromosomes, Human, 6-12, 78-6509
 - Chromosomes, Human, 13-15, 78-6507
 - Chromosomes, Human, 16-18, 78-6507
 - Geographic Distribution, 78-6507
- Melanoma
 - Cholestan-3 β -ol, 5 α ,6 α -Epoxy-, 78-6362
 - Ultraviolet Rays, 78-6362
- Mercury
 - Bone Marrow Cells, 78-6139
- Mercury, Chloroethyl-
 - Bone Marrow Cells, 78-6139

Chromosome Aberrations (cont'd)

- Methanesulfonic Acid, Ethyl Ester
 - Caffeine, 78-6120
 - Micronucleus Test, 78-6162
- Methanesulfonic Acid, Methyl Ester
 - Cell Cycle Kinetics, 78-6358
 - Endonucleases, 78-6358
- Mitomycin C
 - Cell Cycle Kinetics, 78-6358
 - Endonucleases, 78-6358
- 2-Naphthylamine
 - Micronucleus Test, 78-6162
- Nitrous Acid, Sodium Salt
 - Lymphocytes, Rat, 78-6214
- Phosphide Sulfide, Tris(1-aziridinyl)-
 - Lymphocytes, 78-6226
- Phosphide Oxide, 1,4-Piperazinediylbis(bis(1-aziridinyl))-
 - 78-6226
- Phosphide Oxide, 1,4-Piperazinediylbis(bis(1-aziridinyl))-
 - Lymphocytes, 78-6226
- Phosphoric Triamide, Hexamethyl-
 - Micronucleus Test, 78-6162
- 3,6-Pyridazinedione, 1,2-Dihydro-
 - Caffeine, 78-6120
- Quinacrine Mustard
 - Caffeine, 78-6120
- Quinoline, 4-Nitro-, 1-Oxide
 - Micronucleus Test, 78-6162
- Radiation, Ionizing
 - Caffeine, 78-6120
 - Cell Cycle Kinetics, 78-6358
 - Endonucleases, 78-6358
 - Imidazole-1-ethanol, 2-Methyl-5-nitro-, 78-6235
 - Piperonyl Butoxide, 78-6238
- Styrene
 - Occupational Hazard, 78-6252
- 1-Triazene, 3,3-Dimethyl-1-phenyl-
 - Lymphocytes, Rat, 78-6214
- Triazene, 3,3-Dimethyl-1-(*m*-pyridyl)-
 - Lymphocytes, Rat, 78-6214
- s*-Triazine, 2,4,6-Tris(1-aziridinyl)-
 - Colchicine, 78-6222
 - Embryo, Mouse, 78-6222
 - Mutagenic Activity, 78-6222
- Ultraviolet Rays
 - Cell Cycle Kinetics, 78-6358
 - Cholestan-3 β -ol, 5 α ,6 α -Epoxy-, 78-6362
 - Endonucleases, 78-6358
 - Fibroblasts, 78-6362
 - Hyperthermia, 78-6362
- Urea, Methyl Nitroso-
 - Caffeine, 78-6120
- Zinc
 - Calcium, 78-6147
 - Lymphocytes, 78-6147

Chromosome Abnormalities

- Macrophages
 - Mitosis, Review, 78-6079
- Salicylanilide, 3,3',5,5',6-Pentachloro-2'-hydroxy-
 - Allium cepa*, 78-6255
 - Mitosis, 78-6255
- Smoke Condensate
 - Ultrastructural Study, Plants, 78-6170

Chromosomes

- Acetaldehyde

Chromosomes (cont'd)

- DNA, 78-6188
- Mutagenic Activity, 78-6188
- Anemia, Aplastic
 - Hereditary Diseases, Review, 78-6074
- Benzenediazosulfonic Acid, *p*-(Dimethylamino)-, Sodium Salt
 - Mitosis, 78-6284
 - Mutagenic Activity, 78-6284
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 - Bone Marrow, Mouse, 78-6280
 - Mutagenic Activity, 78-6280
- Fensulfothion
 - Mitosis, 78-6284
 - Mutagenic Activity, 78-6284
- Hair Dyes
 - Occupational Hazard, 78-6151
- Mercury, (Acetato)phenyl-
 - Occupational Hazard, 78-6140
- Neoplasms
 - Hereditary Diseases, Review, 78-6074
- Neoplasms, Experimental
 - Clone Cells, 78-6597
- p*-Phenylenediamine, *N,N*-Dimethyl-
 - Mitosis, 78-6284
 - Mutagenic Activity, 78-6284
- p*-Toluidine, α,α,α -Trifluoro-2,6-dinitro-*N,N*-dipropyl-*Drosophila melanogaster*, 78-6268
- Turner's Syndrome
 - Genetics, Review, 78-6069
- Xeroderma Pigmentosum
 - Hereditary Diseases, Review, 78-6074

Chromosomes, Human, 6-12

- Leukemia, Myeloblastic
 - Chromosome Aberrations, 78-6509

Chromosomes, Human, 13-15

- Leukemia, Myeloblastic
 - Chromosome Aberrations, 78-6507
- Virus, Epstein-Barr
 - Hybrid Cells, 78-6455

Chromosomes, Human, 16-18

- Leukemia, Myeloblastic
 - Chromosome Aberrations, 78-6507

Chromosomes, Human, 21-22

- Leukemia, Myelocytic
 - Lymphocyte Culture Test, Mixed, 78-6494

Chrysene, 2-Amino-

- Hepatocytes
 - Cytotoxicity, Fibroblasts, 78-6267

Chrysene, 6-Amino-

- Carcinogenic Metabolite
 - Quantitation Method, 78-6190
- Hepatocytes
 - Cytotoxicity, Fibroblasts, 78-6267

Chrysene, 1,2-Dihydro-1,2-dihydroxy-5-methyl-

- Ames Test
 - Carcinogenic Metabolite, 78-6323

Chrysene, 7,8-Dihydro-7,8-dihydroxy-5-methyl-

- Ames Test
 - Carcinogenic Metabolite, 78-6323

Chrysene 5,6-Dihydro-5,6-epoxy-

- Escherichia coli*
 - DNA Repair, 78-6325

rysene, 5-Methyl-
 Ames Test
 Carcinogenic Metabolite, 78-6323
 Aroclor 1254
 Carcinogenic Metabolite, 78-6323
 ric Acid
 Lipomatosis
 Phosphofructokinase, 78-6589
 rulline, *N*-Methyl-*N*-nitroso-
 DNA
 Alkaline Sucrose Gradients, 78-6200
 Pancreas, Liver, Rat, 78-6200
 ostridiopeptidase A
 Neoplasm Metastasis
 Tumor-Associated Products, Review, 78-6065
 ostridium leptum
 Carbonyl Cyanide, 3-Chlorophenylhydrazino-
 Oxidoreductases, 78-6590
 5 β -Cholan-24-oic Acid, 3 α ,7 α -Dihydroxy-
 Oxidoreductases, 78-6590
 Cholic Acid, Monosodium Salt
 Oxidoreductases, 78-6590
 Cyclohexanamine, *N,N*-Methanetetraylbis-
 Oxidoreductases, 78-6590
 Oxidoreductases
 Enzymatic Activity, 78-6590
 Phenol, 2,4-Dinitro-
 Oxidoreductases, 78-6590
 4-Quinolol, 2-Heptyl-, 1-Oxide
 Oxidoreductases, 78-6590
 al Tar
 Aryl Hydrocarbon Hydroxylases
 Skin, 78-6336
 Carcinogen, Chemical
 Occupational Hazard, Review, 78-6017
 Lymphoma
 Occupational Hazard, Review, 78-6017
 Skin Diseases
 Aryl Hydrocarbon Hydroxylases, 78-6336
 Skin Neoplasms
 Occupational Hazard, Review, 78-6017
 obalt Chloride
 Methane, Azoxy-
 Formaldehyde, 78-6106
 odeine
 Kidney Neoplasms
 Carcinoma, Transitional Cell, 78-6293
 olchicine
 s-Triazine, 2,4,6-Tris(1-aziridinyl)-
 Chromosome Aberrations, 78-6222
 olitis, Ulcerative
 Colonic Neoplasms
 Adenocarcinoma, 78-6537
 Carcinoma, 78-6537
 Case Report, 78-6537
 Intestinal Neoplasms
 Precancerous Conditions, Review, 78-6083
 ollagen
 Teratoid Tumor
 Retinoic Acid, 78-6339
 olonic Neoplasms
 Adenocarcinoma

Colonic Neoplasms (cont'd)
 4-Biphenylamine, 3,2'-Dimethyl-, 78-6297
 Cholanthrene, 3-Methyl-, 78-6321
 Colitis, Ulcerative, 78-6537
 Genetics, 78-6538
 Ultrastructural Study, 78-6536
 Adenoma
 Ultrastructural Study, 78-6536
 Aluminum Hydroxide
 Epidemiology, 78-6126
 Ammonia
 Dietary Fiber, 78-6042
 Metabolism, Review, 78-6042
 Antioxidants
 Epidemiology, Review, 78-6039
 Bacteria
 Epidemiology, Review, 78-6033, 78-6043
 4-Biphenylamine, 3,2'-Dimethyl-
 Carcinogenic Activity, Rat, 78-6297
 Glucuronidase, 78-6297
 Intestinal Microflora, 78-6297
 Blacks
 Epidemiology, South Africa, Review, 78-6041
 Carcinogen, Chemical
 Dietary Fiber, 78-6040
 Carcinoma
 Colitis, Ulcerative, 78-6537
 Colitis, Ulcerative
 Case Report, 78-6537
 Cycasin
 Dietary Fiber, 78-6178
 Diet
 Epidemiology, 78-6588
 Epidemiology, Denmark, Finland, 78-6572
 Epidemiology, Review, 78-6043
 Dietary Fats
 Bacteria, 78-6033
 Epidemiology, Review, 78-6039
 Dietary Fiber
 Blacks, 78-6041
 Epidemiology, Review, 78-6039
 Epidemiology
 Utah, 78-6559
 Epinephrine
 Bacteria, 78-6033
 Ethnic Groups
 Epidemiology, Review, 78-6043
 Genetics
 Precancerous Conditions, 78-6538
 Hydrazine, 1,1-Dimethyl
 Diet, 78-6179
 Dietary Fiber, 78-6179, 78-6180
 Intestinal Bacteria
 Epidemiology, Review, 78-6034
 Intestinal Polyps
 Precancerous Conditions, 78-6536
 Precancerous Conditions, Review, 78-6034
 Leukemia, Myeloblastic
 Drug Therapy, 78-6509
 Radiotherapy, 78-6509
 Streptozotocin, 78-6509
 Uracil, 5-Fluoro-, 78-6509
 Urea, 1-(2-Chloroethyl)-3-(4-methylcyclohexyl)-1-
 nitroso-, 78-6509
 Mormons
 Epidemiology, 78-6559
 Mycotoxins

Colonic Neoplasms (cont'd)

- Epidemiology, Review, 78-6034
- Neoplasms, Multiple Primary
 - Genetics, 78-6538
- Peptic Ulcer
 - Aluminum Hydroxide, 78-6126
- Polycyclic Hydrocarbons
 - Intestinal Bacteria, 78-6034
- Polyps
 - Precancerous Conditions, 78-6538
- Precancerous Conditions
 - Ultrastructural Study, 78-6536
- Retinol
 - Epidemiology, Review, 78-6039

Complement

- T-Lymphocytes
 - Immunity, Cellular, 78-6495
- Lymphoma
 - Rosette Formation, AKR Mouse, 78-6503
 - Virus, Murine Leukemia, 78-6503
- Thymoma
 - Rosette Formation, AKR Mouse, 78-6503
 - Virus, Murine Leukemia, 78-6503

Concanavalin A

- Butyric Acid, 2-Amino-4-(ethylthio)-
 - DNA Replication, 78-6205
 - Lymphocyte Transformation, 78-6205
- Neoplasms, Experimental
 - Growth, 78-6597
- Virus, Adeno 12
 - Cell Agglutination, 78-6470
- Virus, RD-114
 - T-Lymphocytes, 78-6441

Congo Red

- see C.I. Direct Rod 28, Disodium Salt

Contact Inhibition

- Growth Substances
 - Retinol, 78-6592
- Retinol
 - Cells, Cultured, 78-6592
- Ultraviolet Rays
 - Retinol, 78-6592

Contraceptives, Oral

- Breast Neoplasms
 - Epidemiology, Review, 78-6026, 78-6027
- Gynecologic Neoplasms
 - Epidemiology, Review, 78-6027
 - Precancerous Conditions, Review, 78-6027
- Hepatoma
 - Case Report, 78-6353
 - Epidemiology, Review, 78-6026, 78-6027
 - Estradiol, 17-Ethynyl-, 78-6353
 - Neoplasm Metastasis, 78-6353
 - Norethisterone Acetate, 78-6353
 - Precancerous Conditions, Review, 78-6027
- Liver Neoplasms
 - Adenoma, 78-6352, 78-6354
 - Case Report, 78-6352, 78-6354, 78-6355
 - Ethinodiol Diacetate, 78-6354
 - Hyperplasia, 78-6355
 - Mestranol, 78-6354, 78-6355
 - Norethynodrel, 78-6355
 - Receptors, Hormone, 78-6355
- Neoplasm Metastasis
 - Case Report, 78-6353

Contraceptives, Oral (cont'd)

- Uterine Neoplasms
 - Epidemiology, Review, 78-6026

Copper

- Chlorophyll
 - Mutagenic Activity, 78-6134

Cortisol

- Skin Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6307

Cortisone

- Skin Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6307

Cosmetics

- Ames Test
 - Mutagenic Activity, 78-6150
- Saccharomyces cerevisiae*
 - Mutagenic Activity, 78-6150

Coumarin, 3, 3'-Methylenebis(4-hydroxy)-

- Benzo(a)pyrene
 - Aryl Hydrocarbon Hydroxylases, 78-6334
 - Hepatotoxicity, Rat, 78-6334
- Cholanthrene, 3-Methyl-
 - Aryl Hydrocarbon Hydroxylases, 78-6334
 - Hepatotoxicity, Rat, 78-6334
- Quinone Reductases
 - Liver, Rat, 78-6334

Coumestrol

- Estradiol
 - Receptors, Hormone, 78-6350

p-Cresol, 2,6-Di-tert-butyl-

- Ames Test
 - Mutagenic Activity, 78-6206
- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 - Cytochrome P-450, 78-6218
- Free Radicals, 78-6218
- Hepatocarcinogenicity, 78-6218
- Carbamic Acid, Ethyl Ester
 - Co-carcinogenic Effect, 78-6184
- Lung Neoplasms
 - Carbamic Acid, Ethyl Ester, 78-6184

o-Cresol, 4,6-Dinitro-, Sodium Salt

- Chromosome Aberrations
 - Bone Marrow Cells, 78-6157
- Dose-Response Study, 78-6157
- Leukocytes, 78-6157

Croton Oil

- Skin Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6307

Cyanogen Bromide

- Virus, Moloney Murine Sarcoma
 - Viral Proteins, 78-6439

Cycasin

- Colonic Neoplasms
 - Dietary Fiber, 78-6178

Cyclohexanamine, *N,N*-Methanetetraylbis-

- Clostridium leptum*
 - Oxidoreductases, 78-6590

Cyclohexane, 1,2,3,4,5,6-Hexachloro-, α -Isomer

- Mutagenic Activity
 - Blue-Green Algae, 78-6271

cyclohexane, 1,2,3,4,5,6-Hexachloro-, γ -Isomer
Chromosome Aberrations
Lymphocytes, Rat, 78-6214

Cyclohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-
Escherichia coli
Mutagenic Activity, 78-6159
Mutagenic Activity
Galactose Resistance, 78-6159
Salmonella typhimurium
Mutagenic Activity, 78-6159

cyclohexene Oxide
Carcinogen, Chemical
Ames Test, 78-6166
Mutagenic Activity, 78-6166
Phenanthrene, 9,10-Dihydro-9,10-epoxy-
Carcinogenic Metabolite, 78-6312

cycloheximide
Virus, Rous Sarcoma
Glycolysis, 78-6396

cyclopenta(cd)pyrene
Smoke Condensate
Isolation and Characterization, 78-6337

cyclopentadiene, Hexachloro-
Escherichia coli
Mutagenic Activity, 78-6167
Microsomes, Liver
Mutagenic Activity, 78-6167

cyclopentadiene, Tetrachloro-
Barbituric Acid, 5-Ethyl-5-phenyl-
Mutagenic Activity, 78-6167
Escherichia coli
Mutagenic Activity, 78-6167
Microsomes, Liver
Mutagenic Activity, 78-6167

cyclophosphamide
Antineoplastic Agents
Ames Test, 78-6232
Cell Transformation, Neoplastic
Cells, Cultured, 78-6195
Chromatids
Diffusion Chamber Cultures, 78-6194
Chromosome Aberrations
Cells, Cultured, 78-6195
Diffusion Chamber Cultures, 78-6194
Micronucleus Test, 78-6162
Glucosephosphate
Carcinogenic/Mutagenic Activity, 78-6195
Hepatocytes
Cytotoxicity, Fibroblasts, 78-6267
Immunosuppression
Carcinogenic Activity, Review, 78-6051
Leukemia, Myeloblastic
Case Report, 78-6197
Hodgkin's Disease, 78-6196
Sjogren's Syndrome, 78-6197
NADP
Carcinogenic/Mutagenic Activity, 78-6195
Oxidoreductases
Cell Transformation, Neoplastic, 78-6195
Chromatids, 78-6195
Chromosome Aberrations, 78-6195
Virus, Marek's Disease Herpes
Viral Vaccines, 78-6406

Cystadenoma, Papillary
Nose Neoplasms
Reverse Transcriptase, 78-6521
Ultrastructural Study, Sheep, 78-6521
Virus-Like Particles, 78-6521

Cystosarcoma Phyllodes
Breast Neoplasms
Case Report, 78-6510
Histological Study, 78-6510

Cysts
Breast Neoplasms
Precancerous Conditions, 78-6553

Cytochrome P-450
Aniline, *N,N*-Dimethyl-*p*-phenylazo-
p-Cresol, 2,6-Di-*tert*-butyl-, 78-6218
Carcinogen, Chemical
Enzyme Activation, 78-6340
Dimethylamine, *N*-Nitroso-
Metabolism, Liver, 78-6217
Hydrocarbons, Halogenated
Carcinogenic Metabolite, Review, 78-6020
Liver Neoplasms
Aniline, *N,N*-Dimethyl-*p*-phenylazo-, 78-6218
Diethylamine, *N*-Nitroso-, 78-6218
Precancerous Conditions, 78-6218
Methane, Azoxy-
N-Demethylation, 78-6106
Mutagens
Enzyme Activation, 78-6340
Mycotoxins
Carcinogenic Metabolite, Review, 78-6020
Nitrosamines
Carcinogenic Metabolite, Review, 78-6020
Phosgene
Chloroform, 78-6109
Polycyclic Hydrocarbons
Carcinogenic Metabolite, Review, 78-6020

Cytokinins
Agrobacterium tumefaciens
Isolation and Characterization, 78-6585
Plasmids, 78-6585

Cytosine
Sulfurous Acid, Monosodium Salt
Bacteriophages, 78-6160

Daunomycin
Antineoplastic Agents
Ames Test, 78-6232

Deoxyribonuclease
Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
DNA Repair, 78-6191
Urea, Methyl Nitroso-
Chromatin, Liver, 78-6215

Deoxyribonucleosides
Acetaldehyde, Chloro-
Imidazo-Cyclization Products, 78-6112
Ethylene, Chloro-
Imidazo-Cyclization Products, 78-6112

Deuterium
Chloroform
Hepatotoxicity, Rat, 78-6109
Metabolism, 78-6109

- Dexamethasone**
 Skin Neoplasms
 Benz(a)anthracene, 7,12-Dimethyl-, 78-6307
 Protease Inhibitors, 78-6307
 Virus, Murine Mammary Tumor
 Glycoproteins, 78-6411
 RNA, Viral, 78-6411
- Dexon**
see Benzenediazosulfonic Acid, *p*-(Dimethylamino)-, Sodium Salt
- Diabetes Mellitus**
 Uterine Neoplasms
 Epidemiology, Review, 78-6066
- 4,4'-Diaminodiphenylmethane**
see Aniline, 4,4'-Methylenedi-
- Diazepam**
 Phenol, (1,1-Dimethylethyl)-4-methoxy-
 Mutagenic Activity, 78-6328
- Diazobenzene Sulfonate**
see Benzenediazonium, 4-Sulfo-, Hydroxide, Inner Salt
- Dibenzo(a,h)anthracene**
 Asbestos
 Aryl Hydrocarbon Hydroxylases, 78-6169
- Dibenzo(p)dioxin, 1,2,7,8-Tetrachloro-**
 Hydro-Lyases
 Metabolism, Lung, 78-6331
 Transferases
 Metabolism, Lung, 78-6331
- Dibenzo-p-dioxin, 2,3,7,8-Tetrachloro-**
 Dimethylamine, *N*-Nitroso-
 Metabolism, Liver, 78-6217
 Sarcoma
 Acetic Acid, (2,4,5-Trichlorophenoxy)-, 78-6288
- Dibutyl Cyclic AMP**
 Rectal Neoplasms
 Carcinoembryonic Antigen, 78-6596
 Teratoid Tumor
 Plasminogen Activators, 78-6339
- Dichromic Acid, Disodium Salt**
 Erythrocytes
 Ames Test, 78-6130
 Microsomes, Liver
 Ames Test, 78-6130
- Dicumarol**
see Coumarin, 3, 3'-Methylenebis(4-hydroxy)-
- Dieldrin**
 Cells, Cultured
 Metabolism, Lung, 78-6305
 Dose-Response Study
 Risk Factors, 78-6111
- Diet**
 Breast Neoplasms
 Epidemiology, Review, 78-6037
 Colonic Neoplasms
 Epidemiology, 78-6588
 Epidemiology, Denmark, Finland, 78-6572
 Epidemiology, Review, 78-6043
 Hydrazine, 1,1-Dimethyl, 78-6179
 Intestinal Neoplasms
 Bacteria, Review, 78-6035
 Epidemiology, Review, 78-6083
- Diet (cont'd)**
 Neoplasms
 Epidemiology, Review, 78-6032
- Dietary Fats**
 Bile Acids and Salts
 Metabolism, Review, 78-6036
 Breast Neoplasms
 Epidemiology, Review, 78-6036, 78-6067
 Cholesterol
 Metabolism, Review, 78-6036
 Colonic Neoplasms
 Bacteria, 78-6033
 Epidemiology, Review, 78-6039
 Intestinal Neoplasms
 Epidemiology, Review, 78-6036
 Methane, Azoxy-
 Metabolism, 78-6106
 Prostatic Neoplasms
 Epidemiology, 78-6558
 Steroids
 Metabolism, Review, 78-6036
 Uterine Neoplasms
 Epidemiology, Review, 78-6036
- Dietary Fiber**
 Ammonia
 Carcinogenic Activity, 78-6042
 Metabolism, Review, 78-6042
 Carcinogen, Chemical
 Metabolism, Review, 78-6040
 Colonic Neoplasms
 Ammonia, 78-6042
 Blacks, 78-6041
 Carcinogen, Chemical, 78-6040
 Cycasin, 78-6178
 Epidemiology, Review, 78-6039
 Hydrazine, 1,1-Dimethyl, 78-6179, 78-6180
 Intestinal Neoplasms
 Bacteria, Review, 78-6038
 Epidemiology, Review, 78-6038
 Toxins
 Metabolism, Review, 78-6040
- Diethylamine, 2,2'-Dichloro-N-methyl-**
 Leukemia, Myeloblastic
 Drug Therapy, 78-6219
- Diethylamine, N-Nitroso-**
 Chromosome Aberrations
 Micronucleus Test, 78-6162
 DNA Repair
 Liver, Rat, 78-6216
 Glutamyl Transpeptidase
 Hepatocyte, Rat, 78-6492
 Liver Neoplasms
 Cytochrome P-450, 78-6218
 Precancerous Conditions, 78-6492
 Lung Neoplasms
 Anti-Antibodies, 78-6417
 Immunization, 78-6417
 Microsomes, Liver
 Free Radicals, 78-6218
 RNA, Transfer, Methyltransferases
 Liver, Rat, 78-6265
 Transplantation, Homologous
 Hepatocyte, Rat, 78-6492
 Virus, Murine Leukemia
 Anti-Antibodies, 78-6417

- ylamine, *N*-Nitroso- (cont'd)**
 Co-carcinogenic Effect, 78-6417
- Virus, Radiation Leukemia**
 Anti-Antibodies, 78-6417
 Co-carcinogenic Effect, 78-6417
- Virus, SV40**
 Cell Transformation, Neoplastic, 78-6242
- ive System Neoplasms**
 Petroleum
 Epidemiology, Review, 78-6018
- Vanadium**
 Occupational Hazard, 78-6561
- thyl Acetylenedicarboxylic Acid**
 see 2-Butynedioic Acid, Dimethyl Ester
- thylamine**
 Microsomes, Liver
 Ames Test, 78-6290
- thylamine, *N*-Nitroso-**
 Ames Test
 S-9 Mix, 78-6263
- Aroclor 1254**
 Enzyme Activation, 78-6324
- Barbituric Acid, 5-Ethyl-5-phenyl-**
 Metabolism, Liver, 78-6217
- Cells, Cultured**
 Methylation, 78-6217
- Cholanthrene, 3-Methyl-**
 Metabolism, Liver, 78-6217
- Chromosome Aberrations**
 Lymphocytes, Rat, 78-6214
 Micronucleus Test, 78-6162
- Cytochrome P-450**
 Metabolism, Liver, 78-6217
- Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-**
 Metabolism, Liver, 78-6217
- DNA**
 Adenine, 3-Methyl-, 78-6215
 Guanine, 7-Methyl-, 78-6215
 Purine, 2-Amino-6-methoxy-, 78-6215
- DNA Repair**
 Dose-Response Study, 78-6324
 Enzyme Activation, 78-6324
- DNA Replication**
 Pancreas, Rat, 78-6212
- Dose-Response Study**
 Risk Factors, 78-6111
- Guanine, 7-Methyl-**
 Chromatin, Liver, 78-6215
- Hepatocytes**
 Cytotoxicity, Fibroblasts, 78-6267
- Nucleic Acids**
 Methylation, 78-6217
- Oxidoreductases**
 Metabolism, Liver, 78-6217
- Purine, 2-Amino-6-methoxy-**
 Chromatin, 78-6213
 DNA, 78-6213
- RNA, Transfer, Methyltransferases**
 Liver, Rat, 78-6265
- Urea, Methyl Nitroso-**
 Adenine, 3-Methyl-, 78-6215
 Purine, 2-Amino-6-methoxy-, 78-6229
- ne**
 see Phosphinde Oxide, 1,4-Piperazinediylbis(bis(1-aziridinyl))-
- Dipyrido(1,2- α 3',2'-*d*)imidazole, 2-Amino-**
 Glutamic Acid
 Mutagens, 78-6198
- Dipyrido(1,2- α 3',2'-*d*)imidazole, 2-Amino-6-methyl-**
 Glutamic Acid
 Mutagens, 78-6198
- Disgerminoma**
 Genetics
 Case Report, 78-6506
 Ovarian Neoplasms
 Genetics, Review, 78-6069
 Sex Chromosomes
 Cell Differentiation, 78-6506
 Testicular Neoplasms
 Metals, 78-6563
- Disulfide, Bis(diethylthiocarbamoyl)-**
 Methane, Azoxy-
 Formaldehyde, 78-6106
- DNA**
 Acetaldehyde
 Chromosomes, 78-6188
 Cross-Linking Activity, 78-6188
 Acetaldehyde, Chloro-
 Liver, Rat, 78-6112
 Carcinogen, Chemical
 Carcinogenic Activity, 78-6154
 Chromosome Aberrations
 Cell Cycle Kinetics, 78-6358
 Citrulline, *N*-Methyl-*N*-nitroso-
 Alkaline Sucrose Gradients, 78-6200
 Pancreas, Liver, Rat, 78-6200
 Dimethylamine, *N*-Nitroso-
 Adenine, 3-Methyl-, 78-6215
 Guanine, 7-Methyl-, 78-6215
 Purine, 2-Amino-6-methoxy-, 78-6213, 78-6215
 Ethylene, Chloro-
 Liver, Rat, 78-6112
 Fluorescence
 Alkaline Elution Assay, 78-6366
 Radiation Effects, 78-6366
 Hydroxylamine, *N*-1-Naphthyl-
 Deoxyguanosine Adducts, 78-6247
 Mitochondria
 Cell Transformation, Neoplastic, Review, 78-6061
 Neutral Red
 Alkaline Sucrose Gradients, 78-6200
 Pancreas, Liver, Rat, 78-6200
 Plasmacytoma
 Endonucleases, 78-6371
 Protease Inhibitors
 Uterus, Mouse, 78-6307
 Purine, 2-Amino-6-methoxy-
 Enzymatic Removal, 78-6213
 Serine, Diazoacetate (Ester)
 Alkaline Sucrose Gradients, 78-6200
 Pancreas, Liver, Rat, 78-6200
 Triazene, 3,3-Dimethyl-1-phenyl-
 Transplacental Alkylation, Rat, 78-6221
 Urea, Methyl Nitroso-
 Chromatin, Liver, 78-6229
 Purine, 2-Amino-6-methoxy-, 78-6213
 Virus, Rous Sarcoma
 Transduction, Genetic, 78-6404
- DNA, Bacterial**
Agrobacterium tumefaciens

DNA, Bacterial (cont'd)

- Carcinogenic Activity, 78-6584
- DNA-DNA Hybridization, 78-6583, 78-6584
- Plasmids, 78-6583, 78-6584
- Plant Tumors
 - Agrobacterium tumefaciens*, 78-6583
 - Plasmids, 78-6583

DNA Polymerase

- Bacteria
 - DNA Repair, Review, 78-6055
- Burkitt's Lymphoma
 - Isolation and Characterization, 78-6453
 - Phosphonoacetic Acid, 78-6453
 - Virus, Epstein-Barr, 78-6453
 - Virus, Herpes Simplex, 78-6453
- DNA Repair
 - Escherichia coli*, 78-6325
- Virus, Adeno 3
 - DNA, Viral, 78-6465
- Virus, Bovine Parvo
 - Enzymatic Activity, 78-6447
- Virus, Kilham Rat
 - Enzymatic Activity, 78-6447
- Virus, Parvo
 - Enzymatic Activity, 78-6447

DNA Repair

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Chromatin, 78-6191
 - Deoxyribonuclease, 78-6191
 - Escherichia coli*, 78-6308
 - Fibroblasts, 78-6191
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-6308
- Acetamide, Thio-
 - Liver, Rat, 78-6216
- Adriamycin
 - Escherichia coli*, 78-6308
- Aflatoxin B1
 - Liver, Rat, 78-6216
- Ataxia Telangiectasia
 - Endonucleases, 78-6364
 - Fibroblasts, 78-6364
- Benz(a)anthracene, 7-Bromomethyl-
 - Fibroblasts, 78-6313
 - Xeroderma Pigmentosum, 78-6313
- Benz(a)anthracene, 5,6-Dihydro-7,12-dimethyl-5,6-epoxy-
 - Escherichia coli*, 78-6325
- Benz(a)anthracene, 5,6-Dihydro-5,6-epoxy-7-methyl-
 - Escherichia coli*, 78-6325
- Benzo(a)pyrene
 - Dose-Response Study, 78-6324
 - Enzyme Activation, 78-6324
- Benzo(a)pyrene 4,5-Oxide
 - Escherichia coli*, 78-6325
- Bleomycin
 - Escherichia coli*, 78-6308
- Butane, 1,2:3,4-Diepoxy-
 - Escherichia coli*, 78-6325
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Liver, Rat, 78-6216
- C.I. Food Brown 1
 - Bacteria, 78-6153
- Carbon Tetrachloride
 - Liver, Rat, 78-6216
- Carcinogen, Chemical
 - Alkaline Elution Assay, Review, 78-6053
 - Autoradiography, Review, 78-6053

DNA Repair (cont'd)

- Bioassay, 78-6099
- Carcinogenic Activity, 78-6099
- Carcinogenic Activity, Review, 78-6054
- Screening Test, Review, 78-6053
- Cell Transformation, Neoplastic
 - Theoretical Model, Review, 78-6008
- Chrysene 5,6-Dihydro-5,6-epoxy-
 - Escherichia coli*, 78-6325
- Diethylamine, *N*-Nitroso-
 - Liver, Rat, 78-6216
- Dimethylamine, *N*-Nitroso-
 - Dose-Response Study, 78-6324
 - Enzyme Activation, 78-6324
- DNA Polymerase
 - Escherichia coli*, 78-6325
- Endonucleases
 - Escherichia coli*, 78-6325
- Fluorescein, 2',4',5',7'-Tetrabromo-, Disodium Salt
 - Bacteria, 78-6153
- Fluorescence
 - Hamster V79 Cells, 78-6366
- Methanesulfonic Acid, Ethyl Ester
 - Liver, Rat, 78-6216
- Mutagens
 - Bioassay, 78-6099
- Phenanthrene 9,10-Dihydro-9,10-epoxy-
 - Escherichia coli*, 78-6325
- Phorbol 12,13-Didecanoate
 - Escherichia coli*, 78-6308
- Platinum, Diaminedichloro-, *cis*
 - Escherichia coli*, 78-6308
- Radiation, Ionizing
 - Ataxia Telangiectasia, 78-6364
 - Harman, 78-6155
 - Norharman, 78-6155
- Retrosine
 - Liver, Rat, 78-6216
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - 6,7-Epoxides, 78-6325
 - Escherichia coli*, 78-6308, 78-6325
- Ultraviolet Rays
 - Caffeine, 78-6121
 - Cholestan-3 β -ol, 5 α ,6 α -Epoxy-, 78-6362
 - Chromatin, 78-6357
 - Fibroblasts, 78-6313
 - Harman, 78-6155
 - Norharman, 78-6155
 - Nucleosome Rearrangement, 78-6357
 - Xeroderma Pigmentosum, 78-6313
- Urea, 1,3-Bis(2-chloroethyl)-1-nitroso-
 - Escherichia coli*, 78-6308
- Urea, Methyl Nitroso-
 - Escherichia coli*, 78-6308
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-6308

DNA Replication

- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Concanavalin A, 78-6205
 - Pancreas, Rat, 78-6212
 - Plant Agglutinins, 78-6205
- Carbamic Acid, Ethyl Ester
 - Lung, Mouse Embryo, 78-6183
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, Ethyl Ester
 - Pancreas, Rat, 78-6212
- Dimethylamine, *N*-Nitroso-
 - Pancreas, Rat, 78-6212
- 2-Propanol, 1,1'-Iminodi-*N*-nitroso-

DNA Replication (cont'd)

- Pancreas, Rat, 78-6212
- Quinoline, 4-(Hydroxyamino)-, 1-Oxide
Pancreas, Rat, 78-6212
- Serine, Diazoacetate (Ester)
Pancreas, Rat, 78-6212
- Streptozotocin
Pancreas, Rat, 78-6212
- Ultraviolet Rays
Dose-Response Study, 78-6361
Hamster V79 Cells, 78-6361
- Virus, Adeno 2
DNA, Viral, 78-6462
- Virus, Epstein-Barr
Antigens, Viral, 78-6454
- Virus, Herpes Simplex 1
Proteins, 78-6448
RNA, 78-6448
Temperature Sensitive Mutants, 78-6448
- Virus, Polyoma
RNA, 78-6482

DNA, Viral

- Virus, Adeno 2
DNA Replication, 78-6462
Endonucleases, 78-6462
Thymidine Incorporation, 78-6462
Virus Replication, Review, 78-6093
- Virus, Adeno 3
Cell Transformation, Neoplastic, 78-6464
Cleavage Sites, 78-6465
DNA Polymerase, 78-6465
Endonucleases, 78-6464, 78-6465
- Virus, Adeno 5
Cell Transformation, Neoplastic, 78-6466
Renaturation Kinetics, 78-6466
- Virus, Adeno 12
DNA-RNA Hybridization, 78-6469
Endonucleases, 78-6469
Morphological Revertants, 78-6470
RNA Replication, 78-6469
- Virus, Bovine Papilloma
DNA-DNA Hybridization, 78-6445
Nucleotide Sequence, 78-6445
- Virus, Papilloma
Mastomys natalensis, 78-6477
Nucleic Acid Conformation, 78-6477
- Virus, Polyoma
Carcinogenic Activity, Review, 78-6092
- Virus, Rous Sarcoma
DNA-RNA Hybridization, 78-6393
Transformation, Genetic, 78-6393, 78-6403
- Virus, SV40
RNA, Messenger, 78-6485
RNA Replication, 78-6485
Virus Replication, Review, 78-6093
- Warts
Virus, Papilloma, 78-6476

Drosophila melanogaster

- Benzene, (Epoxyethyl)-
Mutagenic Activity, 78-6253
- Heliotrine
Mutagenic Activity, 78-6238
- Mercury
Mutagenic Activity, 78-6139
- Mercury, Chloroethyl-
Mutagenic Activity, 78-6139

Drosophila melanogaster (cont'd)

- Mutagens
Eye Pigmentation, 78-6368
Nitroimidazoles, 78-6234
- Neoplasms
Cell Differentiation, 78-6075
- Radiation, Ionizing
Eye Pigmentation, 78-6368
Mutagenic Activity, 78-6238
- Spices
Mutagenic Activity, 78-6122
- Styrene
Mutagenic Activity, 78-6253
- p*-Toluidine, α, α, α -Trifluoro-2,6-dinitro-*N,N*-dipropyl-
Chromosomes, 78-6268
Mutagenic Activity, 78-6268

Drug Therapy

- Leukemia, Myeloblastic
Case Report, 78-6196, 78-6509
Colonic Neoplasms, 78-6509
Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6219
Glomerulonephritis, 78-6209
Hodgkin's Disease, 78-6196, 78-6219
Prednisone, 78-6219
p-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
78-6219
Vincristine, 78-6219
- Lymphoma
Glomerulonephritis, 78-6209
- Uterine Neoplasms
Estrogens, 78-6557

DT-Diaphorases

see Quinone Reductases

Duodenal Neoplasms

- 4-Biphenylamine, 3,2'-Dimethyl-
Carcinogenic Activity, Rat, 78-6297

Dyes

- C.I. Food Brown 1
Mutagenic Activity, 78-6153

Ear Neoplasms

- 4-Biphenylamine, 3,2'-Dimethyl-
Carcinogenic Activity, Rat, 78-6297
- Carcinoma, Epidermoid
Acenaphthene, 5-Nitro-, 78-6254
- Cholesteatoma
Histological Study, 78-6520
Precancerous Conditions, 78-6520
Precancerous Conditions, Review, 78-6062

Endometriosis

- Uterine Neoplasms
Abortion, 78-6541

Endonucleases

- Agrobacterium tumefaciens*
DNA, Mobility, 78-6582
Isolation and Characterization, 78-6582
- Ataxia Telangiectasia
DNA Repair, 78-6364
- Benzamide, 3,5-Dinitro-
Mutagenic Activity, 78-6285
Ultraviolet Rays, 78-6285
- Benzamide, 2-Methyl-3,5-dinitro-
Mutagenic Activity, 78-6285
Ultraviolet Rays, 78-6285
- Bleomycin

Endonucleases (cont'd)

- Chromosome Aberrations, 78-6358
- Chromosome Aberrations
 - Cell Cycle Kinetics, 78-6358
 - Hamster Ovary Cells, 78-6358
- DNA Repair
 - Escherichia coli*, 78-6325
- Methanesulfonic Acid, Methyl Ester
 - Chromosome Aberrations, 78-6358
- Mitomycin C
 - Chromosome Aberrations, 78-6358
- Plasmacytoma
 - DNA, 78-6371
 - Osmium Tetroxide, 78-6371
 - Potassium Chloride, 78-6371
 - Ultraviolet Rays, 78-6371
- Radiation, Ionizing
 - Chromosome Aberrations, 78-6358
- Ultraviolet Rays
 - Chromosome Aberrations, 78-6358
- Virus, Adeno 2
 - DNA, Viral, 78-6462
- Virus, Adeno 3
 - DNA, Viral, 78-6464, 78-6465
- Virus, Adeno 12
 - DNA, Viral, 78-6469
- Virus, Sendai
 - Chromosome Aberrations, 78-6358
- Virus, SV40
 - RNA, Messenger, 78-6485

Endosulfan

- Ames Test
- Mutagenic Activity, 78-6163

Environmental Hazard

- Carcinogen, Chemical
 - Quantitation Method, 78-6119
- Neoplasms
 - Water Supply, Fluoridated, 78-6550
- Pesticides
 - Quantitation Method, 78-6101
- Polycyclic Hydrocarbons
 - Quantitation Method, Review, 78-6010
- Water Supply, Fluoridated
 - Epidemiology, 78-6550

Enzymes

- Neoplasms, Experimental
 - Neoplasm Metastasis, Review, 78-6060

Ependymoma

- Brain Neoplasms
 - Virus, SV40, 78-6486
- Glioblastoma Multiforme
 - Radiotherapy, 78-6365
- Virus, SV40
 - Antigens, Viral, 78-6486

Epichlorohydrine

- see Propane, 1-Chloro-2,3-epoxy-

Epidermal Growth Factors

- see Peptides

Epinephrine

- Colonic Neoplasms
 - Bacteria, 78-6033

Epoxide Hydratases

- Benzo(a)pyrene

Epoxide Hydratases (cont'd)

- Inactivating Mechanisms, 78-6302
- Mutagens
 - Inactivating Mechanisms, 78-6302

Erythrocytes

- 2-Benzimidazolecarbamate, Methyl Ester
 - Micronucleus Formation, 78-6303
- Chromic Acid
 - Ames Test, 78-6130
- Chromic Acid, Dipotassium Salt
 - Ames Test, 78-6130
- Chromic Acid, Lead Salt
 - Ames Test, 78-6130
- Chromic Acid, Zinc Salt
 - Ames Test, 78-6130
- Dichromic Acid, Disodium Salt
 - Ames Test, 78-6130
- 2-Imidazolidinethione, *N*-Nitroso-
 - Micronucleus Formation, 78-6303
- Virus, Bovine Papilloma
 - Hemagglutination, 78-6445

Erythrocytosis

- see Polycythemia

Erythroleukemia

- Adenosine Cyclic 3',5' Monophosphate
 - Caffeine, 78-6426
- Cell Differentiation, 78-6426
- 2-Imidazolidinone, 4-Methoxy-, 78-6426
- Theophylline, 78-6426
- Xanthine, 3-Isobutyl-1-methyl-, 78-6426
- Cell Differentiation
 - Butyric Acid, Sodium Salt, 78-6426
 - Cell Cycle Kinetics, 78-6426
 - Hexamethylene Bisacetamide, 78-6426
 - Methane Sulfinylbis-, 78-6426
- Virus, Friend Murine Leukemia
 - Adenosine Cyclic 3',5' Monophosphate, 78-6426
 - Cell Differentiation, 78-6426
- Virus, Friend Spleen Focus-Forming
 - Virus-Like Particles, 78-6429

Erythropoietin

- Virus, Friend Murine Leukemia
 - Cell Transformation, Neoplastic, 78-6424

Escherichia coli

- Acetamide, *N*-(Acetyloxy)-*N*-fluorene-2-yl-
 - DNA Repair, 78-6308
- Acridinium, 3,6-Diamino-10-methyl-, Chloride
 - Mutagenic Activity, 78-6159
- Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-
 - Enzyme Activation, 78-6227
- Adriamycin
 - DNA Repair, 78-6308
- Benz(a)anthracene, 5,6-Dihydro-7,12-dimethyl-5,6-epoxy-
 - DNA Repair, 78-6325
- Benz(a)anthracene, 5,6-Dihydro-5,6-epoxy-7-methyl-
 - DNA Repair, 78-6325
- Benz(a)anthracene, 7,12-Dimethyl-
 - Carcinogenic Metabolite, Review, 78-6015
- Benzamide, 3,5-Dinitro-
 - Mutagenic Activity, 78-6285
- Benzamide, 2-Methyl-3,5-dinitro-
 - Mutagenic Activity, 78-6285
- Benzene, *p*-Bromo-(epoxyethyl)-
 - Mutagenic Activity, 78-6272
- Benzene, *m*-Chloro-(epoxyethyl)-

Escherichia coli (cont'd)

- Mutagenic Activity, 78-6272
- Benzene, 3,4-Dimethyl-(epoxyethyl)-
 - Mutagenic Activity, 78-6272
- Benzene, (Epoxyethyl)-
 - Lethal Dose 50, 78-6272
 - Mutagenic Activity, 78-6272
- Benzene, (Epoxyethyl)-*m*-methoxy-
 - Mutagenic Activity, 78-6272
- Benzene, (Epoxyethyl)-*m*-methyl-
 - Mutagenic Activity, 78-6272
- Benzene, (Epoxyethyl)-*p*-methyl-
 - Mutagenic Activity, 78-6272
- Benzo(a)pyrene 4,5-Oxide
 - DNA Repair, 78-6325
- Bleomycin
 - DNA Repair, 78-6308
- Butane, 1,2:3,4-Diepoxy-
 - DNA Repair, 78-6325
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Mutagenic Activity, 78-6206
- Carcinogen, Chemical
 - Mutagenic Activity, 78-6206
- Chrysene 5,6-Dihydro-5,6-epoxy-
 - DNA Repair, 78-6325
- 4-Cyclohexene-1,2-dicarboximide, *N*-(Trichloromethyl)thio-
 - Mutagenic Activity, 78-6159
- Cyclopentadiene, Hexachloro-
 - Mutagenic Activity, 78-6167
- Cyclopentadiene, Tetrachloro-
 - Mutagenic Activity, 78-6167
- DNA Repair
 - DNA Polymerase, 78-6325
 - Endonucleases, 78-6325
- 2-Furaldehyde, 5-Nitro-, Semicarbazone
 - Enzyme Activation, 78-6227
- Phenanthrene 9,10-Dihydro-9,10-epoxy-
 - DNA Repair, 78-6325
- Phorbol 12,13-Didecanoate
 - DNA Repair, 78-6308
- Platinum, Diaminedichloro-, *cis*
 - DNA Repair, 78-6308
- Quinacrine
 - Mutagenic Activity, 78-6159
- Sulfuric Acid, Diethyl Ester
 - Mutagenic Activity, 78-6159
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - DNA Repair, 78-6308, 78-6325
- Ultraviolet Rays
 - DNA Repair, Review, 78-6045
 - Mutagenic Activity, Review, 78-6045
- Urea, 1,3-Bis(2-chloroethyl)-1-nitroso-
 - DNA Repair, 78-6308
- Urea, Methyl Nitroso-
 - DNA Repair, 78-6308

Esophageal Neoplasms

- Benzylamine, *N*-Methyl-*N*-nitroso-
 - Neoplasm Invasiveness, 78-6273
- Carcinoma, Epidermoid
 - Animal Model, Rat, 78-6273
- Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273
- Epidemiology, 78-6549
- Ethyl Alcohol, 78-6549
- Smoking, 78-6549
- Carcinoma, Papillary
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273

Esophageal Neoplasms (cont'd)

- Caribbean Islands
 - Epidemiology, 78-6570
- Epidemiology
 - Utah, 78-6559
- Neoplasms, Multiple Primary
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273
 - Epidemiology, 78-6549
- Papilloma
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273
- Piperidine, 4-Chloro-1-nitroso-
 - Carcinogenic Activity, Rat, 78-6240
- Radiotherapy
 - Epidemiology, 78-6380

Estradiol

- Antibodies
 - Pregnancy, Rat, 78-6351
- Bacteria
 - Metabolism, Feces, 78-6348
- Breast Neoplasms
 - Epidemiology, Review, 78-6037
 - Ethnic Groups, 78-6544
 - Genetics, 78-6349
- Coumestrol
 - Receptors, Hormone, 78-6350
- Isoflavone, 7-Hydroxy-4'-methoxy-
 - Receptors, Hormone, 78-6350
- Isoflavone, 4',5,7-Trihydroxy-
 - Receptors, Hormone, 78-6350
- Liver Neoplasms
 - Receptors, Hormone, 78-6355
- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6351
 - Neonatal Hormone Treatment, 78-6346
 - Pituitary Hormones, 78-6346
 - Receptors, Hormone, 78-6351
- Menopause
 - Ethnic Groups, 78-6544
- Pituitary Gland
 - Histological Study, Mouse, 78-6346
- Plasminogen Activators
 - Uterus, Mouse, 78-6307
- Zearalenol
 - Receptors, Hormone, 78-6350
- Zearalenone
 - Receptors, Hormone, 78-6350

Estradiol, 3-Benzozate

- Acetamide, Thio-
 - Ovalbumin, 78-6347
 - Progesterone, 78-6347
- Actinomycin D
 - Ovalbumin, 78-6347
 - Progesterone, 78-6347

Estradiol, 17-Ethynyl-

- DNA, Binding
 - Liver, Rat, 78-6345
- Hepatoma
 - Contraceptives, Oral, 78-6353

Estriol

- Antibodies
 - Pregnancy, Rat, 78-6351
- Breast Neoplasms
 - Epidemiology, Review, 78-6037
 - Genetics, 78-6349
- Mammary Neoplasms, Experimental

- Estriol (cont'd)**
Benz(a)anthracene, 7,12-Dimethyl-, 78-6351
Receptors, Hormone, 78-6351
- Estriol Cyclopentyl Ether**
Estrone, 2-Hydroxy-
Pregnancy, Rat, 78-6351
- Estriol, 16-Epi-**
Mammary Neoplasms, Experimental
Receptors, Hormone, 78-6351
- Estrogens**
Bacteria
Metabolism, Feces, 78-6348
Breast Neoplasms
Epidemiology, Review, 78-6027
Receptors, Hormone, Review, 78-6024
Liver Neoplasms
Adenoma, 78-6352
Case Report, 78-6352
Uterine Neoplasms
Drug Therapy, 78-6557
Epidemiology, 78-6557
Epidemiology, Review, 78-6028, 78-6066
Genetics, Review, 78-6066
Statistical Analysis, 78-6557
- Estrone**
Bacteria
Metabolism, Feces, 78-6348
Breast Neoplasms
Epidemiology, Review, 78-6037
Ethnic Groups, 78-6544
Genetics, 78-6349
DNA, Binding
Liver, Rat, 78-6345
Menopause
Ethnic Groups, 78-6544
Uterine Neoplasms
Liver Diseases, 78-6028
Obesity, 78-6028
- Estrone, 2-Hydroxy-**
Estriol Cyclopentyl Ether
Pregnancy, Rat, 78-6351
- Ethane, 1,2-Dibromo-**
Acetaldehyde, Bromo-
Carcinogenic Metabolite, 78-6115
Barbituric Acid, 5-Ethyl-5-phenyl-
Metabolism, Rat, 78-6115
Benz(a)anthracene
Metabolism, Rat, 78-6115
Microsomes, Liver
Metabolism, Rat, 78-6115
Nucleic Acids
Proteins, 78-6115
Proteins
Binding, Liver, Kidney, Rat, 78-6115
Transferases
Binding, Liver, Kidney, Rat, 78-6115
Enzymatic Activity, 78-6115
- Ethane, 1,2-Dichloro-**
Angiosarcoma
Carcinogenic Activity, Mouse, Rat, 78-6114
Enzyme Activation
Mutagenic Activity, 78-6113
Fibroma
Carcinogenic Activity, Mouse, Rat, 78-6114
- Ethane, 1,2-Dichloro- (cont'd)**
Glutathione
Ames Test, 78-6113
Mutagenic Activity, 78-6113
Mammary Neoplasms, Experimental
Carcinogenic Activity, Mouse, Rat, 78-6114
NADP
Ames Test, 78-6113
Mutagenic Activity, 78-6113
Respiratory Tract Neoplasms
Carcinogenic Activity, Mouse, Rat, 78-6114
Stomach Neoplasms
Carcinogenic Activity, Mouse, Rat, 78-6114
Uterine Neoplasms
Carcinogenic Activity, Mouse, Rat, 78-6114
- Ethane, 1,1'-Oxybis-**
Respiratory Tract Neoplasms
Benzo(a)pyrene, 78-6335
- Ethane, 1,1,1-Trichloro-2,2-bis(p-chlorophenyl)-**
Antineoplastic Activity
Review, 78-6012
Cells, Cultured
Metabolism, Lung, 78-6305
Dose-Response Study
Risk Factors, 78-6111
- Ethanol, 2-Hydrazino-**
Ames Test
S-9 Fraction, 78-6182
- Ethanol, 2,2',2''-Nitrilotris-**
Bacillus subtilis
Mutagenic Activity, 78-6224
Ethanol, *N*-Nitrosoiminodi-
Mutagenic Metabolite, 78-6224
Lymphoma
Carcinogenic Activity, Mouse, 78-6224
Microsomes, Liver
Mutagenic Activity, 78-6224
Nitrous Acid, Sodium Salt
Mutagenic Activity, 78-6224
Sex Factors
Carcinogenic Activity, Mouse, 78-6224
- Ethanol, *N*-Nitrosoiminodi-**
Ethanol, 2,2',2''-Nitrilotris-
Mutagenic Metabolite, 78-6224
- Ethenetetracarbonitrile**
Benz(e)indole, 3-(β -Nitrovinyl)-
Benzocarbazole Derivatives, Synthesis, 78-6299
- Ether, Bis(chloromethyl)-**
Occupational Hazard
Carcinogenic Activity, Review, 78-6009
Quantitation Method, Review, 78-6009
- Ether, 2,2-Dichloro-1,1-difluoroethyl Methyl**
Respiratory Tract Neoplasms
Benzo(a)pyrene, 78-6335
- Ethinylestradiol**
see Estradiol, 17-Ethynyl-
- Ethionine**
see Butyric Acid, 2-Amino-4-(ethylthio)-
- Ethyl Alcohol**
Esophageal Neoplasms
Carcinoma, Epidermoid, 78-6549
Head and Neck Neoplasms

Ethyl Alcohol (cont'd)

- Carcinoma, Epidermoid, 78-6549
- Methanesulfonic Acid, Ethyl Ester
 - Mutagenic Activity, 78-6108
- Mouth Neoplasms
 - Carcinoma, Epidermoid, 78-6375
- Mutagenic Activity
 - Mouse, Rat, 78-6108
- Neoplasms
 - Epidemiology, Review, 78-6032
- Radiation, Ionizing
 - Mutagenic Activity, 78-6108

Ethylene, Chloro-

- Air Pollution
 - Dose-Response Study, 78-6110
- Deoxyribonucleosides
 - Imidazo-Cyclization Products, 78-6112
- DNA
 - Liver, Rat, 78-6112
- Dose-Response Study
 - Risk Factors, 78-6110, 78-6111

Ethylene, Trichloro-

- Yeasts
 - Mutagenic Activity, 78-6116

Ethylidene Gyromitrin

- see Formic Acid, Ethylenemethylhydrazide

Ethynodiol Diacetate

- Liver Neoplasms
 - Contraceptives, Oral, 78-6354

Eye Neoplasms

- Angiosarcoma
 - Epidemiology, Dog, 78-6516
- Carcinoma, Epidermoid
 - Epidemiology, Dog, 78-6516
- Hemangioma
 - Epidemiology, Dog, 78-6516
- Ultraviolet Rays
 - Epidemiology, Dog, 78-6516

Fensulfthion

- Chromosomes
 - Mitosis, 78-6284
 - Mutagenic Activity, 78-6284

Fibroblasts

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - DNA Repair, 78-6191
- Ataxia Telangiectasia
 - DNA Repair, 78-6364
- Benz(a)anthracene, 7-Bromomethyl-
 - DNA Repair, 78-6313
- Benzo(a)pyrene
 - Fluorescence Quenching, 78-6329
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
 - Fluorescence Quenching, 78-6329
- Chick Embryo
 - LETs Protein, Isolation and Characterization 78-6593
- Cholestan-3 β -ol, 5 α ,6 α -Epoxy-
 - Chromosome Aberrations, 78-6362
- Glycoproteins
 - Cell Membrane, 78-6593
 - Chick Embryo, 78-6593
- Macrophages
 - Iododeoxyuridine Incorporation, 78-6435

Fibroblasts (cont'd)

- Methanesulfonic Acid, Ethyl Ester
 - Mutagenic Activity, 78-6161
- Ultraviolet Rays
 - Chromosome Aberrations, 78-6362
 - DNA Repair, 78-6313
- Virus, B77
 - Cell Transformation, Neoplastic, 78-6398
- Virus, Influenza
 - Virus Replication, 78-6435
- Virus, Moloney Murine Sarcoma
 - Benzo(a)pyrene, 78-6329
 - Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-, 78-6329
- Virus, Murine Leukemia
 - Cell Transformation, Neoplastic, 78-6435
- Virus, Murine Sarcoma
 - Cell Transformation, Neoplastic, 78-6435
- Virus, Polyoma
 - Cell Transformation, Neoplastic, 78-6480
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-6398
- Virus, Sendai
 - Virus Replication, 78-6435

Fibroma

- Ethane, 1,2-Dichloro-
 - Carcinogenic Activity, Mouse, Rat, 78-6114
- Ovarian Neoplasms
 - Genetics, 78-6540
- Virus, Bovine Papilloma
 - Antigens, Viral, 78-6446

Fibrosarcoma

- Carrageen
 - Transplantation Immunology, 78-6490
- Cholanthrene, 3-Methyl-
 - Histocompatibility Antigens, 78-6493
 - Transplantation Immunology, 78-6490
- Glycoproteins
 - Isolation and Characterization, Review, 78-6057
- Gold, ((1,2-Dicarboxyethyl)thio)-, Disodium Salt
 - Dose-Response Study, Rat, 78-6133
- Histocompatibility Antigens
 - Antibody Formation, 78-6493
 - Antigen-Antibody Reactions, 78-6493
 - Immunity, Cellular, 78-6493
 - Immunization, 78-6493
 - Immunogenetics, 78-6493
- 2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-
 - Transplantation Immunology, 78-6490
- Lip Neoplasms
 - Epidemiology, 78-6380
- Mammary Neoplasms, Experimental
 - Benzidine, 2,2',5,5'-Tetrachloro-, 78-6254
- Methacrylic Acid, Ethyl Ester
 - Plastic Implants, 78-6372
 - Tryptophan, 78-6372
- Methacrylic Acid, Methyl Ester
 - Plastic Implants, 78-6372
 - Tryptophan, 78-6372
- Nickel
 - Industrial Dusts, 78-6143
- Peptides
 - Cell Division, 78-6481
 - Isolation and Characterization, 78-6481
- Radiation, Ionizing
 - Transplantation Immunology, 78-6490

- Fibrosarcoma (cont'd)**
 Reserpine
 Transplantation Immunology, 78-6490
 Transplantation Immunology
 Macrophages, 78-6490
 Virus, Bovine Papilloma
 Antigens, Viral, 78-6446
- Flagyl**
 see Imidazole-1-ethanol, 2-Methyl-5-nitro-
- Fluoren-2-amine**
 Acetylation Kinetics
 Species Difference, 78-6264
 Acetyltransferases
 Acetylation Kinetics, 78-6264
- Fluorescein, 2',4',5',7'-Tetrabromo-, Disodium Salt**
 Bacteria
 DNA Repair, 78-6153
 Mutagenic Activity, 78-6153
- Fluorescence**
 DNA
 Alkaline Elution Assay, 78-6366
 Radiation Effects, 78-6366
 DNA Repair
 Hamster V79 Cells, 78-6366
- Fluorosulfuric Acid, Methyl Ester**
 Ames Test
 Dose-Response Study, 78-6165
 Cell Transformation, Neoplastic
 Dose-Response Study, 78-6165
- Food Additives**
 Carcinogenic Potential
 Review, 78-6012
- Food Contamination**
 Carbanilic Acid, *m*-Chloro-, Isopropyl Ester
 Carcinogenic Potential, 78-6186
 Maximum Permissible Exposure Level, 78-6186
 Mycotoxins
 Quantitation Method, 78-6101
 Polycyclic Hydrocarbons
 Quantitation Method, 78-6301
 Stomach Neoplasms
 Cadmium, 78-6148
 Manganese, 78-6148
 Zinc, 78-6148
- Formaldehyde**
 Hexamethylenetetramine
 Nitrosation, 78-6225
 Methane, Azoxy-
 Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6106
 7,8-Benzoflavone, 78-6106
 Cholanthrene, 3-Methyl-, 78-6106
 Cobalt Chloride, 78-6106
 Disulfide, Bis(diethylthiocarbamoyl)-, 78-6106
 Metabolism, 78-6106
 Valeric Acid, 2,2-Diphenyl-, 2-(Diethylamino)ethyl
 Ester HCl, 78-6106
- Formamidine, *N*-(4-Chloro-*o*-tolyl)-*N,N*-dimethyl-**
 Cells, Cultured
 Metabolism, Lung, 78-6305
- Formic Acid, Ethylenedimethylhydrazide**
 Microsomes, Liver
 Ames Test, 78-6181
- Formic Acid, Ethylenedimethylhydrazide (cont'd)**
Salmonella typhimurium
 Bacteriolysis, Mouse, 78-6181
- Formic Acid, 1-Methyl Hydrazide**
 Microsomes, Liver
 Ames Test, 78-6181
- Formononetin**
 see Isoflavone, 7-Hydroxy-4'-methoxy-
- Freund's Adjuvant**
 Virus, Feline Leukemia
 Viral Vaccines, 78-6442
- FSH**
 Breast Neoplasms
 Genetics, 78-6349
 Uterine Neoplasms
 Leiomyoma, 78-6541
 Menstruation, 78-6541
- 2-Furaldehyde, 5-Nitro-, Semicarbazone**
 Enzyme Activation
 Microsomes, Testis, 78-6227
Escherichia coli
 Enzyme Activation, 78-6227
 Oxidoreductases
 Microsomes, Testis, 78-6227
- 2-Furanol 5-(3-Pyridyl)tetrahydro-**
 Ames Test
 Mutagenic Activity, 78-6175
 1-Butanone, 4-(*N*-Carbethoxy-*N*-nitrosamino)-1-(3-pyridyl)-
 Hydroxylation, 78-6175
 Nicotine, 1'-Nitroso-1'-demethyl-
 Hydroxylation, 78-6175
- 2-Furanol, Tetrahydro-**
 Pyrrolidine, 1-Nitroso-
 Hydroxylation, 78-6237
- 2-Furanone, 5-(3-Pyridyl)tetrahydro-**
 Nicotine, 1'-Nitroso-1'-demethyl-
 Urinary Metabolite, 78-6175
- Gallbladder Neoplasms**
 Age Factors
 Epidemiology, Sweden, 78-6577
 Cholecystectomy
 Epidemiology, Sweden, 78-6577
- Gastrectomy**
 Stomach Neoplasms
 Carcinoma, 78-6576
 Epidemiology, 78-6576
 Precancerous Conditions, 78-6576
- Gastritis**
 Stomach Neoplasms
 Achlorhydria, 78-6533
 Adenocarcinoma, 78-6534
 Case Report, 78-6533
 Polyps, 78-6533
- Genetics**
 Breast Neoplasms
 Blood Groups, Review, 78-6068
 Epidemiology, 78-6543
 Epidemiology, Review, 78-6067, 78-6068, 78-6073
 Estradiol, 78-6349
 Estriol, 78-6349

Genetics (cont'd)

- Estrone, 78-6349
- FSH, 78-6349
- LH, 78-6349
- Prolactin, 78-6349

Burkitt's Lymphoma

- Histocompatibility Antigens, 78-6459
- Virus, Epstein-Barr, 78-6459

Carcinogen, Chemical

- DNA Repair, Review, 78-6055

Colonic Neoplasms

- Adenocarcinoma, 78-6538
- Neoplasms, Multiple Primary, 78-6538
- Precancerous Conditions, 78-6538

Disgerminoma

- Case Report, 78-6506

Hodgkin's Disease

- Histocompatibility Antigens, 78-6459
- Virus, Epstein-Barr, 78-6459

Hypercholesteremia

- Enzymatic Activity, 78-6589

Intestinal Polyps

- Ultrastructural Study, 78-6536

Kidney Neoplasms

- Epidemiology, Review, 78-6073
- Fish, Review, 78-6059

Leukemia

- Epidemiology, Review, 78-6073

Lymphoma

- Graft Rejection, 78-6491
- Homologous Wasting Disease, 78-6491

Melanoma

- Fish, Review, 78-6059

Neoplasms

- DNA Repair, Review, 78-6055
- Epidemiology, Review, 78-6073
- Neoplasms, Embryonal and Mixed
- Virus, Rous Sarcoma, 78-6400

Neoplasms, Experimental

- Animal Model, Fish, 78-6059

Neoplasms, Multiple Primary

- Virus, Epstein-Barr, 78-6459

Neoplastic Endocrine-Like Syndromes

- Case Report, 78-6518
- Histocompatibility Antigens, 78-6518

Neuroblastoma

- Fish, Review, 78-6059

Ovarian Neoplasms

- Case Report, 78-6540
- Fibroma, 78-6540

Retinoblastoma

- Epidemiology, Review, 78-6071
- Immunity, Cellular, Review, 78-6071

Sarcoma, Reticulum Cell

- Fish, Review, 78-6059

Stomach Neoplasms

- Carcinoma, 78-6575
- Epidemiology, Finland, 78-6575

Thyroid Neoplasms

- Calcitonin, 78-6522
- Carcinoma, 78-6522
- Case Report, 78-6522
- Fish, Review, 78-6059

Ultraviolet Rays

- DNA Repair, Review, 78-6055
- Virus, Rous Sarcoma
- Immune Response, 78-6400

Genetics (cont'd)

- Neoplasm Regression, Spontaneous, 78-6400

Genistein

- see Isoflavone, 4',5,7-Trihydroxy-

Germanium Tetrachloride

- Bacillus subtilis*
- Mutagenic Activity, 78-6128

Glioblastoma Multiforme

- Brain Neoplasms
- Benz(a)anthracene, 7,12-Dimethyl-, 78-6318
- Ependymoma
- Radiotherapy, 78-6365
- Radiotherapy
- Case Report, 78-6365

Glioma

- Glycoproteins
- Isolation and Characterization, Review, 78-6057
- Neoplasm Transplantation
- Cell Line, 78-6581
- Neoplasms, Experimental
- Cell Line, 78-6581
- Urea, Methyl Nitroso-
- Virus-Like Particles, 78-6230
- Virus-Like Particles
- Ultrastructural Study, C-Type Particles, 78-6230

Globin

- Virus, Murine Leukemia
- Age Factors, 78-6422
- Brain, Liver, Mouse, 78-6422
- RNA, 78-6422

Glomerulonephritis

- Leukemia, Myeloblastic
- Drug Therapy, 78-6209
- Lymphoma
- Drug Therapy, 78-6209

Glucocorticoids

- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-
- Immunosuppression, 78-6278

Gluconic Acid, Iron Salt

- Liver Neoplasms
- Siderosis, 78-6193

Glucosamine

- Neuraminidase
- Trophoblast, 78-6591
- Virus, Avian Sarcoma
- Antigen-Antibody Reactions, 78-6394
- Cell Transformation, Neoplastic, 78-6394
- Reverse Transcriptase, 78-6394
- RNA, Viral, 78-6394

Glucose

- Islet Cell Tumor
- Insulin, 78-6207
- Streptozotocin, 78-6207

Glucosephosphate

- Cyclophosphamide
- Carcinogenic/Mutagenic Activity, 78-6195

Glucosephosphate Dehydrogenase

- Leukemia, Myeloblastic
- Enzymatic Activity, 78-6509

Glucuronidase

- Colonic Neoplasms

- Glucuronidase (cont'd)**
 4-Biphenylamine, 3,2'-Dimethyl-, 78-6297
 Intestinal Neoplasms
 Carcinoma, 78-6587
 Intestinal Polyps
 Enzymatic Activity, 78-6587
 Mammary Neoplasms, Experimental
 4-Biphenylamine, 3,2'-Dimethyl-, 78-6297
- Glutamic Acid**
 Ames Test
 Pyrolysis Products, 78-6198
 Dipyrdo(1,2- α :3',2'- d)imidazole, 2-Amino-
 Mutagens, 78-6198
 Dipyrdo(1,2- α :3',2'- d)imidazole, 2-Amino-6-methyl-
 Mutagens, 78-6198
 Mutagens
 Pyrolysis Products, 78-6198
- Glutamic Acid, Monosodium Salt**
 Dose-Response Study
 Mutagenic Activity, Rat, 78-6138
- Glutamyl Transpeptidase**
 Acetamide, *N*-Fluorenyl-2-yl-
 Hepatocyte, Rat, 78-6492
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 Shunt, Portacaval, 78-6275
 Diethylamine, *N*-Nitroso-
 Hepatocyte, Rat, 78-6492
 Hepatoma
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6274, 78-6275
 Glutathione, 78-6274
- Glutathione**
 Benzo(a)pyrene
 Ames Test, 78-6333
 Aryl Hydrocarbon Hydroxylases, 78-6326
 Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 Carcinogenic Metabolite, 78-6326
 Carcinogen, Chemical
 Carcinogenic Activity, Review, 78-6004
 Chromic Acid, Zinc Salt
 Ames Test, 78-6130
 Ethane, 1,2-Dichloro-
 Ames Test, 78-6113
 Mutagenic Activity, 78-6113
 Hepatoma
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6274, 78-6275
 Glutamyl Transpeptidase, 78-6274
 Precancerous Conditions, 78-6274
- Glutathione Transferases**
 Benzo(a)pyrene
 Inactivating Mechanisms, 78-6302
 Mutagens
 Inactivating Mechanisms, 78-6302
- Glycolipids**
 Retinol
 Cells, Cultured, 78-6592
- Glycolysis**
 Virus, Rous Sarcoma
 Cell Transformation, Neoplastic, 78-6396
 Cycloheximide, 78-6396
 Oligomycin, 78-6396
 Temperature Sensitive Mutants, 78-6396
- Glycopeptides**
 Cell Membrane
 Trophoblast, 78-6591
- Glycoproteins**
 Carbohydrates
 Cell Transformation, Neoplastic, Review, 78-6056
 Cell Membrane
 Cell Transformation, Neoplastic, Review, 78-6056
 78-6057
 Fibroblasts, 78-6593
 Isolation and Characterization, Review, 78-6056
 Fibroblasts
 Chick Embryo, 78-6593
 Fibrosarcoma
 Isolation and Characterization, Review, 78-6057
 Glioma
 Isolation and Characterization, Review, 78-6057
 Retinol
 Cells, Cultured, 78-6592
 Rhabdomyosarcoma
 Isolation and Characterization, Review, 78-6057
 Sarcoma, Osteogenic
 Isolation and Characterization, Review, 78-6057
 Virus, Adeno
 Cell Transformation, Neoplastic, Review, 78-6057
 Virus, Gross Murine Leukemia
 Cell Fusion, 78-6431
 Virus, Moloney Murine Leukemia
 Cell Fusion, 78-6431
 Virus, Murine Mammary Tumor
 Dexamethasone, 78-6411
 Hormones, 78-6411
 Virus, Rauscher Murine Leukemia
 Cell Fusion, 78-6431
- Gnidilatin**
 HeLa Cells
 Epidermal Growth Factor, Binding, 78-6306
- Gnilatimacrin**
 HeLa Cells
 Epidermal Growth Factor, Binding, 78-6306
- Goiter, Exophthalmic**
 Thyroid Neoplasms
 Epidemiology, 78-6523
- Gold, ((1,2-Dicarboxyethyl)thio)-, Disodium Salt**
 Fibrosarcoma
 Dose-Response Study, Rat, 78-6133
 Hematoma
 Dose-Response Study, Rat, 78-6133
 Rhabdomyosarcoma
 Dose-Response Study, Rat, 78-6133
- Gold Sodium Thiomalate**
 see Gold, ((1,2-Dicarboxyethyl)thio)-, Disodium Salt
- Gonadotropins, Equine**
 Granulosa Cell Tumor
 Carcinogenic Activity, Mouse, 78-6344
 Lymphosarcoma
 Carcinogenic Activity, Mouse, 78-6344
- Graft vs Host Reaction**
 Virus, Adeno 7
 Immune Response, 78-6468
 Precancerous Conditions, 78-6468
- Graft Rejection**
 Lymphoid Tissue

raft Rejection (cont'd)

Transplantation, Homologous, 78-6491

Lymphoma

Genetics, 78-6491

Histocompatibility Antigens, 78-6491

anulocytes**Benzene**

Acetylglucosaminidase, 78-6258

anulosa Cell Tumor

Gonadotropins, Equine

Carcinogenic Activity, Mouse, 78-6344

rowth**Lung Neoplasms**

EMT6 Mouse Tumor, 78-6598

Neoplasms

Mathematical Model, 78-6555

Neoplasms, Experimental

Clone Cells, 78-6597

Concanavalin A, 78-6597

rowth Substances**Retinol**

Contact Inhibition, 78-6592

uanidine, 1-Methyl-3-nitro-1-nitroso-**Caffeine**

Chromosome Aberrations, 78-6120

Cell Transformation, Neoplastic

Diploid Cells, 78-6245

Cells, Cultured

Colony Formation, Agar, 78-6245

Guanosine

RNA, Methylation, 78-6228

Lymphoblasts, Diploid

Mutagenic Activity, 78-6164

Mesenchymoma

Cells, Cultured, 78-6245

Methanesulfonic Acid, Ethyl Ester

Mutagenic Activity, 78-6161

12-O-Tetradecanoylphorbol-13-acetate

Mutagenic Activity, 78-6295

Virus, SV40

Cell Transformation, Neoplastic, 78-6242

Yeasts

RNA, Methylation, 78-6228

uanine**Aflatoxin B1**

DNA Cleavage, 78-6124

uanine, 7-Methyl-**Dimethylamine, N-Nitroso-**

Chromatin, Liver, 78-6215

DNA, 78-6215

Triazene, 3,3-Dimethyl-1-phenyl-

Liver, Brain, Rat, 78-6221

uanosine**Guanidine, 1-Methyl-3-nitro-1-nitroso-**

RNA, Methylation, 78-6228

Methanesulfonic Acid, Methyl Ester

RNA, Methylation, 78-6228

Sulfuric Acid, Dimethyl Ester

RNA, Methylation, 78-6228

Urea, Methyl Nitroso-

RNA, Methylation, 78-6228

Guanosine Triphosphate**Virus, Polyoma****Guanosine Triphosphate (cont'd)**

Viral Proteins, 78-6484

Guanyl Cyclase

1,2-Benzisothiazolin-3-one, 1,1-Dioxide

Bladder, Rat, 78-6282

Enzymatic Activity, 78-6282

Gynecologic Neoplasms**Adenoma**

4,4'-Stilbenediol, α,α' -Diethyl-, 78-6027

Transplacental Carcinogenesis, Review, 78-6027

Contraceptives, Oral

Epidemiology, Review, 78-6027

Precancerous Conditions, Review, 78-6027

Epidemiology

California, 78-6548

Utah, 78-6559

Hormones

Epidemiology, Review, 78-6067

Hair Dyes**Ames Test**

S-9 Fraction, 78-6269

Occupational Hazard

Chromosomes, 78-6151

p-Phenyldiamine

Ames Test, 78-6269

Harman**Radiation, Ionizing**

DNA Repair, 78-6155

Mutagenic Activity, 78-6155

Ultraviolet Rays

DNA Repair, 78-6155

Mutagenic Activity, 78-6155

Head and Neck Neoplasms**Carcinoma, Epidermoid**

Epidemiology, 78-6549

Ethyl Alcohol, 78-6549

Smoking, 78-6549

Chromium

Occupational Hazard, 78-6561

Neoplasms, Multiple Primary

Epidemiology, 78-6549

Sarcoma

Chromium, 78-6561

Synovioma

Case Report, 78-6524

Histochemical Study, 78-6524

Ultrastructural Study, 78-6524

Heavy Chain Disease**Lymphoma**

Epidemiology, 78-6487

HeLa Cells**Gnidilatin**

Epidermal Growth Factor, Binding, 78-6306

Gnilatimacrin

Epidermal Growth Factor, Binding, 78-6306

Mezerein

Epidermal Growth Factor, Binding, 78-6306

Phorbol 12,13-Dibenzoate

Epidermal Growth Factor, Binding, 78-6306

Phorbol 12,13-Didecanoate

Epidermal Growth Factor, Binding, 78-6306

12-O-Tetradecanoylphorbol-13-acetate

Epidermal Growth Factor, Binding, 78-6306

Heliotrine

- Drosophila melanogaster*
 - Mutagenic Activity, 78-6238
- Piperonyl Butoxide
 - Chromosome Aberrations, 78-6238
 - Mutagenic Activity, 78-6238

Hemadsorption

- Neoplasms
 - Immunoglobulins, Fc, 78-6504

Hemagglutination

- Virus, Bovine Papilloma
 - Erythrocytes, 78-6445

Hemangioendothelioma

- Epidemiology
 - Inbred Mice, 78-6595

Hemangioma

- Eye Neoplasms
 - Epidemiology, Dog, 78-6516

Hematoma

- Gold, ((1,2-Dicarboxyethyl)thio)-, Disodium Salt
 - Dose-Response Study, Rat, 78-6133

Hematopoietic Stem Cells

- Macrophages
 - Colony Stimulating Factor, 78-6488
- Monocytes
 - Colony Stimulating Factor, 78-6488
- Prostaglandins E
 - Colony Stimulating Factor, 78-6488
 - Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-, 78-6488
 - Macrophages, 78-6488
 - Monocytes, 78-6488
- Virus, Friend Murine Leukemia
 - 1,4-Butanediol, Dimethylsulfonate, 78-6424
 - Cell Transformation, Neoplastic, 78-6424
 - Virus Replication, 78-6424

Hemoglobins

- C.I. Food Brown 3, Disodium Salt
 - Toxicity, Pig, 78-6149
- Smoking
 - Hematopoietic Effects, 78-6174
- Virus, Avian Erythroblastosis
 - Cell Transformation, Neoplastic, 78-6390
 - Temperature Sensitive Mutants, 78-6390

Hemopexin

- Aryl Hydrocarbon Hydroxylases
 - Strain Difference, Mouse, 78-6319
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Plasma Levels, Mouse, 78-6319
- 5,6-Benzoflavone
 - Plasma Levels, Mouse, 78-6319
- Cholanthrene, 3-Methyl-
 - Plasma Levels, Mouse, 78-6319
- Hepatoma
 - Benz(a)anthracene, 78-6319
 - Cells, Cultured, 78-6319

Hepatitis

- Leukemia, Myeloblastic
 - Case Report, 78-6219

Hepatoma

- Alpha Fetoproteins
 - Precancerous Conditions, 78-6277

Hepatoma (cont'd)

- Radioimmunoassay, 78-6531
- Australia Antigen
 - Radioimmunoassay, 78-6531
- Benz(a)anthracene
 - Aryl Hydrocarbon Hydroxylases, 78-6319
 - Hemopexin, 78-6319
- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 - Alpha Fetoproteins, 78-6277
 - Glutamyl Transpeptidase, 78-6274, 78-6275
 - Glutathione, 78-6274, 78-6275
 - Hypophysectomy, 78-6274
 - Shunt, Portacaval, 78-6275
- Benzidine, 3,3'-Dichloro-
 - Dose-Response Study, Dog, 78-6270
- Carcinogen, Chemical
 - Virus-Like Particles, Review, 78-6096
- Contraceptives, Oral
 - Case Report, 78-6353
 - Epidemiology, Review, 78-6026, 78-6027
 - Neoplasm Metastasis, 78-6353
 - Precancerous Conditions, Review, 78-6027
- Epidemiology
 - Inbred Mice, 78-6595
- Estradiol, 17-Ethynyl-
 - Contraceptives, Oral, 78-6353
- Ethnic Groups
 - Epidemiology, Cameroon, 78-6568
- Glutamyl Transpeptidase
 - Glutathione, 78-6274
- Glutathione
 - Precancerous Conditions, 78-6274
- Hemopexin
 - Cells, Cultured, 78-6319
- Hypertension
 - Neoplasm Metastasis, 78-6332
- Liver Cirrhosis
 - Australia Antigen, 78-6531
 - Epidemiology, Japan, 78-6531
- Neoplasms, Multiple Primary
 - Benzidine, 3,3'-Dichloro-, 78-6270
 - Serine, Diazoacetate (Ester), 78-6199
- Norethisterone Acetate
 - Contraceptives, Oral, 78-6353
- Pesticides
 - Carcinogenic Activity, Review, 78-6011
- Proteins
 - Gene Control Mechanisms, 78-6072
- Serine, Diazoacetate (Ester)
 - Choline, 78-6199
- Virus, RNA Tumor
 - Virus-Like Particles, Review, 78-6096
- Xanthine, 3-Hydroxy-
 - Transplacental Carcinogenesis, 78-6249

Hereditary Diseases

- Neoplasms
 - DNA Repair, Review, 78-6055
 - Epidemiology, Review, 78-6030

Hexadecane, 2,6,10,14-Tetramethyl-

- Plasmacytoma
 - Mouse, Review, 78-6013

Hexamethylene Bisacetamide

- Erythroleukemia
 - Cell Differentiation, 78-6426

examethylenetetramine

- Chloroform
 - Nitrosation, 78-6225
- Formaldehyde
 - Nitrosation, 78-6225
- Methylamine
 - Nitrosation, 78-6225
- Nitric Acid, Sodium Salt
 - Nitrosation, 78-6225
- Trimethylamine, *N*-Oxide
 - Nitrosation, 78-6225

1,5-Hexatriene, 1,6-Diphenyl-

- Virus, Epstein-Barr
 - Fluorescence Polarization, 78-6458
 - Liposomes, 78-6458
- Virus, Semliki Forest
 - Fluorescence Polarization, 78-6458

exestrol

- see* Phenol, 4,4'-(1,2-Diethylene)di-, *meso*-

stocompatibility Antigens

- Burkitt's Lymphoma
 - Genetics, 78-6459
- Fibrosarcoma
 - Antibody Formation, 78-6493
 - Antigen-Antibody Reactions, 78-6493
 - Cholanthrene, 3-Methyl-, 78-6493
 - Immunity, Cellular, 78-6493
 - Immunization, 78-6493
 - Immunogenetics, 78-6493
- Hodgkin's Disease
 - Genetics, 78-6459
- Leukemia, Lymphocytic
 - Virus, SV40, 78-6085
- T-Lymphocytes
 - Immunity, Cellular, 78-6495
- Lymphoma
 - Graft Rejection, 78-6491
 - Homologous Wasting Disease, 78-6491
- Lymphosarcoma
 - Virus, SV40, 78-6085
- Neoplasms
 - Hereditary Diseases, Review, 78-6074
- Neoplastic Endocrine-Like Syndromes
 - Genetics, 78-6518
- Sarcoma
 - Virus, SV40, 78-6085
- Sarcoma, Osteogenic
 - Virus, SV40, 78-6085
- Sarcoma, Reticulum Cell
 - Virus, SV40, 78-6085
- Virus, Friend Murine Leukemia
 - Immune Response, Review, 78-6098
 - Immunity, Cellular, Review, 78-6098
- Virus, Murine Leukemia
 - Immune Response, Review, 78-6098
 - Immunity, Cellular, Review, 78-6098
- Virus, Radiation Leukemia
 - Immune Response, Review, 78-6098
 - Immunity, Cellular, Review, 78-6098

stones

- Ames Test
 - Pyrolysis Product, 78-6201
- Cervix Neoplasms
 - Sperm Proteins, 78-6569
- Virus, Epstein-Barr

Histones (cont'd)

- Antigens, Viral, 78-6456

Hodgkin's Disease

- Antigens
 - Immunity, Cellular, 78-6487
- Epidemiology
 - Australia, 78-6562
- Histocompatibility Antigens
 - Genetics, 78-6459
- IgG
 - Serum Levels, 78-6487
- IgM
 - Serum Levels, 78-6487
- Leukemia, Myeloblastic
 - Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-78-6196
 - Case Report, 78-6219
 - Cyclophosphamide, 78-6196
 - Drug Therapy, 78-6196, 78-6219
 - Radiotherapy, 78-6196
 - p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-78-6196
 - s*-Triazine, 2,4,6-Tris(1-aziridinyl)-, 78-6196
 - Vinblastine, 78-6196
- Liver Cirrhosis
 - Case Report, 78-6513
- Oncogenic Viruses
 - Epidemiology, Review, 78-6031
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-6452
 - Antigens, Viral, 78-6459
 - Genetics, 78-6459
 - Lymphocytes, 78-6452
 - B-Lymphocytes, 78-6461
 - Virus Activation, Child, 78-6452
 - Virus Excretion, 78-6452
 - Virus Replication, 78-6461

Homologous Wasting Disease

- Lymphoid Tissue
 - Transplantation, Homologous, 78-6491
- Lymphoma
 - Genetics, 78-6491
 - Histocompatibility Antigens, 78-6491

Hormones

- Breast Neoplasms
 - Epidemiology, 78-6349
 - Epidemiology, Review, 78-6067
- Gynecologic Neoplasms
 - Epidemiology, Review, 78-6067
- Neoplasm Metastasis
 - Tumor-Associated Products, Review, 78-6065
- Virus, Murine Mammary Tumor
 - Co-carcinogenic Effect, 78-6411
 - Glycoproteins, 78-6411
 - Precancerous Conditions, 78-6411
 - RNA, Viral, 78-6411

Host vs Graft Reaction

- see* Graft Rejection

Hybrid Cells

- Teratoid Tumor
 - Gene Control Mechanisms, 78-6072
- Virus, Epstein-Barr
 - Antigens, Viral, 78-6455
 - Chromosomes, Human, 13-15, 78-6455

- Hycanthone**
Phenol, (1,1-Dimethylethyl)-4-methoxy-
Mutagenic Activity, 78-6328
- Hydrazine**
Occupational Hazard
Carcinogenic Activity, Review, 78-6009
Quantitation Method, Review, 78-6009
- Hydrazine, 1,1-Dimethyl**
Colonic Neoplasms
Diet, 78-6179
Dietary Fiber, 78-6179, 78-6180
- Hydrazine, 1,2-Dimethyl-**
Chromosome Aberrations
Micronucleus Test, 78-6162
- Hydrazine, 2,4-Dinitrophenyl-**
Ames Test
S-9 Fraction, 78-6182
Pyrrolidine, 1-Nitroso-
Hydroxylation, 78-6237
- Hydrazine, Methyl-**
Microsomes, Liver
Ames Test, 78-6181
- Hydrazine, (4-Methylphenyl)-**
Ames Test
S-9 Fraction, 78-6182
- Hydrazine, Monohydrate**
Ames Test
S-9 Fraction, 78-6182
- Hydrazine, Phenethyl-, Sulfate**
Ames Test
S-9 Fraction, 78-6182
- Hydrazine, Phenyl-, Monohydrochloride**
Ames Test
S-9 Fraction, 78-6182
- Hydrazine, Sulfate**
Ames Test
S-9 Fraction, 78-6182
- Hydro-Lyases**
Aroclor 1254
Enzymatic Activity, 78-6257
Barbituric Acid, 5-Ethyl-5-phenyl-
Enzymatic Activity, 78-6257, 78-6320
Metabolism, Lung, 78-6331
Microsomes, Liver, 78-6320
Benzene, 1-Chloro-2,4-dinitro-
Metabolism, Lung, 78-6331
Benzo(a)pyrene 4,5-Oxide
Metabolism, Lung, 78-6331
Carcinogen, Chemical
Carcinogenic Metabolite, 78-6166
Cholanthrene, 3-Methyl-
Enzymatic Activity, 78-6257, 78-6320
Microsomes, Liver, 78-6320
Dibenzo(p)dioxin, 1,2,7,8-Tetrachloro-
Metabolism, Lung, 78-6331
Phenol, (1,1-Dimethylethyl)-4-methoxy-
Benzene, (Epoxyethyl)-, 78-6257
Enzymatic Activity, 78-6257
Microsomes, Liver, 78-6257
Pregnenolone 16 α -Carbonitrile
Metabolism, Lung, 78-6331
- Hydrocarbons, Chlorinated**
Liver Neoplasms
Occupational Hazard, Review, 78-6014
Occupational Hazard
Carcinogenic Activity, Review, 78-6009
Quantitation Method, Review, 78-6009
- Hydrocarbons, Halogenated**
Cytochrome P-450
Carcinogenic Metabolite, Review, 78-6020
- 2-Hydroxyethylhydrazine**
see Ethanol, 2-Hydrazino-
- Hydroxylamine**
Saccharomyces cerevisiae
Growth, Radiation-Sensitive Mutants, 78-6244
Sulfurous Acid, Monosodium Salt
Mutagenic Activity, 78-6160
- Hydroxylamine, O-Methyl-**
Sulfurous Acid, Monosodium Salt
Mutagenic Activity, 78-6160
- Hydroxylamine, N-1-Naphthyl-**
Bladder Neoplasms
Carcinogenic Activity, 78-6247
Deoxyguanosine Adducts, 78-6247
DNA
Deoxyguanosine Adducts, 78-6247
2-Naphthol, 1-Amino-
Solvolysis Products, 78-6247
4-Naphthol, 1-Amino-
Solvolysis Products, 78-6247
- Hydroxylases**
Barbituric Acid, 5-Ethyl-5-phenyl-
Enzymatic Activity, 78-6320
Microsomes, Liver, 78-6320
Benzenediazonium, 4-Sulfo-, Hydroxide, Inner Salt
Localization, Liver, 78-6320
Cholanthrene, 3-Methyl-
Microsomes, Liver, 78-6320
Intestinal Neoplasms
Carcinoma, 78-6587
Intestinal Polyps
Enzymatic Activity, 78-6587
Peptide Hydrolases
Localization, Liver, 78-6320
- Hydroxysteroid Dehydrogenases**
5 α -Androstan-17-one, 3 α -Hydroxy-
Vegetarianism, 78-6341
Intestinal Neoplasms
5 α -Androstan-17-one, 3 α -Hydroxy-, 78-6341
Enzymatic Activity, Feces, 78-6341
Vegetarianism
Enzymatic Activity, Feces, 78-6341
- Hydroxysteroids**
Bacteria
Metabolism, Feces, 78-6348
- Hypercholesteremia**
Alcohol Oxidoreductases
Enzymatic Activity, 78-6589
Genetics
Enzymatic Activity, 78-6589
- Hyperparathyroidism**
Thyroid Neoplasms
Radiation, Ionizing, 78-6379

- Hyperplasia**
Caffeine
Leydig Cells, 78-6123
- Liver Neoplasms**
Contraceptives, Oral, 78-6355
4,4'-Stilbenediol, α,α' -Diethyl-, 78-6355
- Lung Neoplasms**
Precancerous Conditions, 78-6528
- Mammary Neoplasms, Experimental**
Age Factors, Rat, 78-6525
Estrus, 78-6525
Mitosis, 78-6525
Precancerous Conditions, Rat, 78-6525
- Prostatic Neoplasms**
Virus, Cytomegalo, 78-6499
- Smoke Condensate**
Epidermis, Mouse, 78-6172
- Theobromine**
Leydig Cells, 78-6123
- Thyroid Neoplasms**
Imidazole-2-thiol, 1-Methyl-, 78-6248
- Hypersensitivity, Delayed**
Carrageen
Antigens, 78-6490
- 2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-**
Antigens, 78-6490
- Radiation, Ionizing**
Antigens, 78-6490
- Reserpine**
Antigens, 78-6490
- Hypertension**
Benzo(a)pyrene
Carcinogenic Activity, 78-6332
- Carcinoma**
Neoplasm Transplantation, 78-6332
- Carcinosarcoma**
Neoplasm Metastasis, 78-6332
- Hepatoma**
Neoplasm Metastasis, 78-6332
- Sarcoma**
Neoplasm Transplantation, 78-6332
- Uterine Neoplasms**
Epidemiology, Review, 78-6066
- Hypothermia**
Cholestan-3 β -ol, 5 α ,6 α -Epoxy-
Chromosome Aberrations, 78-6362
- Mammary Neoplasms, Experimental**
Neoplasm Metastasis, 78-6381
- Ultraviolet Rays**
Chromosome Aberrations, 78-6362
- Hypothyroidism**
Thyroid Neoplasms
Imidazole-2-thiol, 1-Methyl-, 78-6248
- Hypoglycemia**
Islet Cell Tumor
Streptozotocin, 78-6207
- Xanthine Phosphoribosyltransferase**
Methanesulfonic Acid, Butyl Ester
Mutagenic Activity, 78-6164
- 372**
Ames Test
Genetic Duplication, 78-6359, 78-6360
- Salmonella typhimurium*
Genetic Duplication, 78-6359, 78-6360
- IgG**
Hodgkin's Disease
Serum Levels, 78-6487
- Lymphoma**
Immunologic Deficiency Syndromes, 78-6487
- IgM**
Hodgkin's Disease
Serum Levels, 78-6487
- Leukemia, Lymphoblastic**
Cytoplasm, Review, 78-6078
- Lymphoma**
Immunologic Deficiency Syndromes, 78-6487
- Imidazo(1,2-a)pyrimidine**
Salmonella typhimurium
Mutagenic Activity, Derivatives, 78-6236
- 1H-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-**
Adenocarcinoma
Radiation, Ionizing, 78-6385
- Cecal Neoplasms**
Neoplasms, Multiple Primary, 78-6385
Radiation, Ionizing, 78-6385
- Intestinal Neoplasms**
Radiation, Ionizing, 78-6385
- Rectal Neoplasms**
Radiation, Ionizing, 78-6385
- Imidazole-1-ethanol, 2-Methyl-5-nitro-**
Chromosome Aberrations
Micronucleus Test, 78-6162
- Phenol, (1,1-Dimethylethyl)-4-methoxy-**
Mutagenic Activity, 78-6328
- Radiation, Ionizing**
Chromosome Aberrations, 78-6235
- Imidazole-2-thiol, 1-Methyl-**
Thyroid Neoplasms
Adenoma, 78-6248
Histological Study, Rat, 78-6248
Hyperplasia, 78-6248
Hyperthyroidism, 78-6248
Precancerous Conditions, 78-6248
- 2-Imidazolidinethione, N-Nitroso-**
Erythrocytes
Micronucleus Formation, 78-6303
- Tubulin**
Dose-Response Study, Mouse, 78-6303
- 2-Imidazolidinone, 4-Methoxy-**
Erythroleukemia
Adenosine Cyclic 3',5' Monophosphate, 78-6426
- 2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-**
Antigens
Hypersensitivity, Delayed, 78-6490
- Fibrosarcoma**
Transplantation Immunology, 78-6490
- Immune Serums**
Burkitt's Lymphoma
Antigen-Antibody Reactions, 78-6498
- Carcinoma, Bronchogenic**
Calcitonin, 78-6594
- Leukemia, Lymphoblastic**
Antigen-Antibody Reactions, 78-6498
Virus, Murine Leukemia, 78-6489
- Leukemia, Myeloblastic**
Antigen-Antibody Reactions, 78-6498
- Leukemia, Myelocytic**

Immune Serums (cont'd)

- Antigen-Antibody Reactions, 78-6498
- Lymphosarcoma
 - Antigen-Antibody Reactions, 78-6498
- Thyroid Neoplasms
 - Calcitonin, 78-6594
- Virus, Murine Leukemia
 - Age Factors, AKR Mouse, 78-6489
 - Thymectomy, AKR Mouse, 78-6489
 - Virus, Gross Murine Leukemia, 78-6489
- Virus, Murine Mammary Tumor
 - Lymphocyte Transformation, 78-6497

Immunity, Cellular

- Fibrosarcoma
 - Histocompatibility Antigens, 78-6493
- Hodgkin's Disease
 - Antigens, 78-6487
- T-Lymphocytes
 - Complement, 78-6495
 - Cytotoxic Effector Cells, 78-6495
 - Histocompatibility Antigens, 78-6495
 - Isoantigens, 78-6495
 - Receptors, Antigen, 78-6495
- Lymphoma
 - Antigens, 78-6487
- Prostatic Neoplasms
 - Virus, Cytomegalo, 78-6499
- Virus, SV40
 - Antigens, Viral, 78-6500
 - T-Lymphocytes, 78-6500

Immunity, Passive

- Sarcoma
 - T-Lymphocytes, 78-6502

Immunization

- Fibrosarcoma
 - Histocompatibility Antigens, 78-6493
- Lung Neoplasms
 - Diethylamine, *N*-Nitroso-, 78-6417
- Virus, Moloney Murine Leukemia
 - Dose-Response Study, 78-6501
 - Ultraviolet Rays, 78-6501
- Virus, Moloney Murine Sarcoma, 78-6501
- Virus, SV40
 - Antigens, Viral, 78-6500

Immunoglobulins, Fc

- Neoplasms
 - Antigen-Antibody Complex, 78-6504
 - Cell Adhesion, 78-6504
 - Hemadsorption, 78-6504
 - Rosette Formation, 78-6504

Immunoglobulins, Heavy Chain

- Multiple Myeloma
 - Bence Jones Protein, 78-6514
 - Cytoplasm, 78-6514

Immunoglobulins, Surface

- Leukemia, Lymphoblastic
 - Cell Differentiation, Review, 78-6078
- Thyroid Neoplasms
 - Lymphoma, 78-6505
- Virus, Moloney Murine Leukemia
 - B-Lymphocytes, 78-6432

Immunologic Deficiency Syndromes

- Leukemia
 - Epidemiology, Review, 78-6031

Immunologic Deficiency Syndromes (cont'd)

- Lymphoma
 - Epidemiology, Review, 78-6031
 - IgG, 78-6487
 - IgM, 78-6487
- Neoplasms
 - Epidemiology, Review, 78-6030
- Stomach Neoplasms
 - Adenocarcinoma, 78-6534
 - Case Report, 78-6534

Immunosuppression

- p*-Acetophenetidine
 - Carcinogenic Activity, Review, 78-6051
- Alanine, 3-(*p*-(Bis(2-chloroethyl)amino)phenyl)-
 - Carcinogenic Activity, Review, 78-6051
- Aniline, *p*-(Phenylazo)-
 - Lymphocytes, 78-6278
- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-
 - Glucocorticoids, 78-6278
- 1,4-Butanediol, Dimethylsulfonate
 - Carcinogenic Activity, Review, 78-6051
- Butyric Acid, 4-(*p*-Bis(2-Chloroethyl)aminophenyl)-
 - Carcinogenic Activity, Review, 78-6051
- Carcinoma
 - Aniline, *p*-(Phenylazo)-, 78-6278
 - Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-, 78-6278
 - Transplantation Immunology, 78-6278
- Cyclophosphamide
 - Carcinogenic Activity, Review, 78-6051
- Norethisterone, 18-Methyl-
 - Carcinogenic Activity, Review, 78-6051
- Radiation, Ionizing
 - Carcinogenic Activity, Review, 78-6051
- 4,4'-Stilbenediol, α, α' -Diethyl-
 - Carcinogenic Activity, Review, 78-6051
- as*-Triazine-3,5(2*H*,4*H*)-dione, 2- β -*D*-Ribofuranosyl-
 - Carcinogenic Activity, Review, 78-6051
- Virus, Friend Murine Leukemia
 - Antibody Formation, 78-6427, 78-6430
 - Fv-3 Gene, 78-6427
 - Genetics, 78-6427
 - Lymphocytes, 78-6427
- Virus, Marek's Disease Herpes
 - Viral Vaccines, 78-6406

Indan, 5-Hydroxyamino-

- Isocyanic Acid, Methyl Ester
 - Quantitation Method, 78-6103
- Liver
 - Quantitation Method, 78-6103

1,3-Indandione

- Smoke Condensate
 - Isolation and Characterization, 78-6171
 - Tumor-Inhibiting Fraction, 78-6171

Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-

- Prostaglandins E
 - Colony Stimulating Factor, 78-6488
 - Hematopoietic Stem Cells, 78-6488

Indole-3-acrylic Acid

- Leukemia
 - Species Difference, 78-6202

Indomethacin

- see* Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl

Infectious Mononucleosis

- Cells, Cultured
- Mutation, 78-6164
- Virus, Epstein-Barr
- Epidemiology, Review, 78-6089

Inflammation

- Macrophages
- Mitosis, Review, 78-6079
- Mouth Neoplasms
- Dentures, 78-6375

Ingenol Dibenzoate

- Nerve Growth Factor
- Ganglia, Chick Embryo, 78-6309

Insecticides

- Leukemia, Myeloblastic
- Lymphoma, 78-6508

Insulin

- Islet Cell Tumor
- Glucose, 78-6207
- Nicotinamide, 78-6207
- Picolinamide, 78-6207
- Streptozotocin, 78-6207

Interferon

- Macrophages
- Immune Response, Review, 78-6077
- Virus, Friend Spleen Focus-Forming
- A-Type Particles, 78-6429
- Virus Replication, 78-6429

Intestinal Neoplasms

- Adenocarcinoma
- Acenaphthene, 5-Nitro-, 78-6254
- Radiation, Ionizing, 78-6385
- Schistosoma japonicum*, 78-6535
- 5 α -Androstan-17-one, 3 α -Hydroxy-
- Hydroxysteroid Dehydrogenases, 78-6341

- Bacteria
- Enzymatic Activity, 78-6587

- Carcinoembryonic Antigen
- Immune Response, Review, 78-6083

- Carcinogen, Chemical
- Bacteria, Review, 78-6035

- Carcinoma
- Bacteria, 78-6587
- Glucuronidase, 78-6587
- Hydroxylases, 78-6587
- Oxidoreductases, 78-6587

- Colitis, Ulcerative
- Precancerous Conditions, Review, 78-6083

- Diet
- Bacteria, Review, 78-6035
- Epidemiology, Review, 78-6083

- Dietary Fats
- Epidemiology, Review, 78-6036

- Dietary Fiber
- Bacteria, Review, 78-6038
- Epidemiology, Review, 78-6038

- Hydroxysteroid Dehydrogenases
- Enzymatic Activity, Feces, 78-6341

- Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
- Colostomy, Rat, 78-6189

- Neoplasms, Multiple Primary
- Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
- 78-6189

- Occupational Hazard

Intestinal Neoplasms (cont'd)

- Epidemiology, 78-6551
- Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-
- Nitrous Acid, Sodium Salt, 78-6241

Polyps

- Bacteria, Review, 78-6035
- Precancerous Conditions, Review, 78-6083

Radiation, Ionizing

- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385

- In Utero Exposure, Rat, 78-6382

Schistosoma japonicum

- Epidemiology, 78-6535
- Socioeconomic Factors
- Epidemiology, Review, 78-6083

Intestinal Polyps**Bacteria**

- Enzymatic Activity, 78-6587
- Colonic Neoplasms
- Precancerous Conditions, 78-6536
- Precancerous Conditions, Review, 78-6034

Genetics

- Ultrastructural Study, 78-6536

Glucuronidase

- Enzymatic Activity, 78-6587

Hydroxylases

- Enzymatic Activity, 78-6587

Oxidoreductases

- Bacteria, 78-6587

Intestines

- Palmitic Acid, Morpholino-
- Body Burden, Rat, 78-6251

Iodine Radioisotopes

- Thyroid Neoplasms
- Epidemiology, 78-6376

Iron

- Chlorophyll
- Mutagenic Activity, 78-6134
- Sarcoma
- Intramuscular Injection, 78-6565

Iron Oxide

- Respiratory Tract Neoplasms
- Benzo(a)pyrene, 78-6335

Islet Cell Tumor

- Glucose
- Insulin, 78-6207
- Nicotinamide
- Insulin, 78-6207
- Pancreatic Neoplasms
- Nicotinamide, 78-6207
- Picolinamide, 78-6207
- Streptozotocin, 78-6207, 78-6208
- Picolinamide
- Insulin, 78-6207
- Streptozotocin
- Glucose, 78-6207
- Hypoglycemia, 78-6207
- Insulin, 78-6207

Isoantigens

- Benzydine
- Occupational Hazard, 78-6566
- Bladder Neoplasms
- Occupational Hazard, 78-6566
- Precancerous Conditions, 78-6566

Isoantigens (cont'd)

- T-Lymphocytes
 - Immunity, Cellular, 78-6495
- 1-Naphthylamine
 - Occupational Hazard, 78-6566
- 2-Naphthylamine
 - Occupational Hazard, 78-6566

Isocyanic Acid, Methyl Ester

- Indan, 5-Hydroxyamino-
 - Quantitation Method, 78-6103

Isoflavone, 7-Hydroxy-4'-methoxy-

- Estradiol
 - Receptors, Hormone, 78-6350

Isoflavone, 4',5,7-Trihydroxy-

- Estradiol
 - Receptors, Hormone, 78-6350

Isomerases

- Agrobacterium tumefaciens*
 - Isolation and Characterization, 78-6582

Isonicotinic Acid Hydrazide

- Acetylation Kinetics
 - Species Difference, 78-6264
- Acetyltransferases
 - Acetylation Kinetics, 78-6264
- Mutagenic Activity
 - Ames Test, 78-6163

Isoquinoline

- Ames Test
 - Amino Derivatives, 78-6239
- Microsomes, Liver
 - Ames Test, 78-6239

Isothiocyanic Acid, 1-Naphthyl Ester

- RNA, Transfer, Methyltransferases
 - Liver, Rat, 78-6265

Isothiocyanic Acid, 2-Naphthyl Ester

- RNA, Transfer, Methyltransferases
 - Liver, Rat, 78-6265

Karyotyping

- Mercury, (Acetato)phenyl-
 - Lymphocytes, 78-6140

Ketosteroids

- Breast Neoplasms
 - Ethnic Groups, 78-6544
- Menopause
 - Ethnic Groups, 78-6544

Kidney

- Benzoic Acid, 2-Amino-3-hydroxy-
 - Lymphoid Tissue, 78-6311
 - Metabolism, Mouse, 78-6311

Kidney Neoplasms

- Acenaphthene, 5-Nitro-
 - Carcinogenic Potential, 78-6254
- p*-Acetophenetidine
 - Case Report, 78-6293
- Adenocarcinoma
 - Radium, 78-6564
 - Ultrastructural Study, 78-6532
- Adrenal Gland Neoplasms
 - Radiotherapy, 78-6384
- Carcinoma
 - Radiotherapy, 78-6384

Kidney Neoplasms (cont'd)

- Carcinoma, Transitional Cell
 - p*-Acetophenetidine, 78-6293
 - Codeine, 78-6293
- Chloroform
 - Water Pollution, 78-6556
- Chromium
 - Occupational Hazard, 78-6561
- Genetics
 - Epidemiology, Review, 78-6073
 - Fish, Review, 78-6059
- Radiotherapy
 - Case Report, 78-6384
- Radium
 - Epidemiology, 78-6564
- Streptozotocin
 - Nicotinamide, 78-6208
- Vanadium
 - Occupational Hazard, 78-6561
- Virus, Avian Leukosis
 - Virus, Helper, 78-6389

Krezonite

- see *o*-Cresol, 4,6-Dinitro-, Sodium Salt

Laryngeal Neoplasms

- Benz(a)anthracene
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Carcinoma, Epidermoid
 - Epidemiology, 78-6380
- Cholanthrene, 3-Methyl-
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Lymphocytes
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Papilloma
 - Virus, Papilloma, 78-6474
- Radiotherapy
 - Epidemiology, 78-6380
- Virus, Papilloma
 - DNA-DNA Hybridization, 78-6474
 - Isolation and Characterization, 78-6474

Lead

- Air Pollution
 - Body Burden, Child, 78-6136
 - Chromosome Aberrations, 78-6136
 - Cytogenetic/Biochemical Effect, 78-6135
- Calcium
 - Chromosome Aberrations, 78-6147
- Chlorophyll
 - Mutagenic Activity, 78-6134
- Lymphocytes
 - Chromosome Aberrations, 78-6147
- Respiratory Tract Neoplasms
 - Radioisotopes, 78-6049

Leiomyoma

- Uterine Neoplasms
 - FSH, 78-6541
 - LH, 78-6541
 - Somatotropin, 78-6541

Leukemia

- Acetophenone, 2-Amino-
 - Species Difference, 78-6202
- Alloxan
 - Fish, 78-6233
- Benz(a)anthracene, 7,12-Dimethyl-
 - Chromosome Aberrations, 78-6314
 - 2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-

emia (cont'd)

- 2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
78-6317
- Benz(a)anthracene, 7,8,12-Trimethyl-
2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
78-6317
- 2-Naphthol, 1-((4-(*o*-Tolylazo)-*o*-tolyl)azo)-, 78-6317
- Benzene
Occupational Hazard, Review, 78-6014
- 1,3-Butadiene
Occupational Hazard, Review, 78-6014
- Cat
Epidemiology, California, 78-6548
- Chromosome Aberrations
Review, 78-6070
- Trisomy 15, Mouse, 78-6314
- Genetics
Epidemiology, Review, 78-6073
- Immunologic Deficiency Syndromes
Epidemiology, Review, 78-6031
- Indole-3-acrylic Acid
Species Difference, 78-6202
- B-Lymphocytes
Antigen-Antibody Reactions, 78-6498
- T-Lymphocytes
Benz(a)anthracene, 7,12-Dimethyl-, 78-6314
- Lymphocytotoxicity, 78-6495
- Occupational Hazard
Epidemiology, Review, 78-6025
- Phagocytosis
Kupffer Cells, 78-6526
- Radiation, Ionizing
Epidemiology, Review, 78-6025
- In Utero Exposure, Rat, 78-6382
- Parabiosis, 78-6373
- Radiotherapy, Review, 78-6025
- Radium
Epidemiology, 78-6564
- Styrene
Occupational Hazard, Review, 78-6014
- Thymus Gland
Phagocytosis, 78-6526
- Virus, Avian Erythroblastosis
Temperature Sensitive Mutants, 78-6390
- Virus, Feline Leukemia
Antigens, Viral, 78-6443
- emia L1210**
Virus, C-Type RNA Tumor
Viral Proteins, 78-6473
- Virus-Like Particles, 78-6473
- Virus-Like Particles
Isolation and Characterization, 78-6473

emia, Lymphoblastic

- Alkaline Phosphatase
Enzymatic Activity, 78-6457
- Antibody Formation
Cell Differentiation, Review, 78-6078
- Antigens, Neoplasm
Antigen-Antibody Reactions, 78-6498
- Chromium
Occupational Hazard, 78-6561
- IgM
Cytoplasm, Review, 78-6078
- Immune Serums
Antigen-Antibody Reactions, 78-6498
- Immunoglobulins, Surface

Leukemia, Lymphoblastic (cont'd)

- Cell Differentiation, Review, 78-6078
- B-Lymphocytes
Precursor Cells, Review, 78-6078
- T-Lymphocytes
Lymphocyte Culture Test, Mixed, 78-6494
- Null Cells
Lymphocyte Culture Test, Mixed, 78-6494
- Virus, Epstein-Barr
Antigens, Viral, 78-6457
- Virus, Murine Leukemia
Age Factors, AKR Mouse, 78-6489
- Immune Serums, 78-6489
- T-Lymphocytes, 78-6489
- Thymectomy, AKR Mouse, 78-6489

Leukemia, Lymphocytic

- Virus, Epstein-Barr
Virus Replication, 78-6461
- Virus, SV40
Antigen-Antibody Reactions, 78-6085
- Histocompatibility Antigens, 78-6085

Leukemia, Myeloblastic

- Alkaline Phosphatase
Enzymatic Activity, 78-6457
- Antigens, Neoplasm
Antigen-Antibody Reactions, 78-6498
- Azathioprine
Case Report, 78-6209
- Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
Arthritis, Rheumatoid, 78-6211
- Case Report, 78-6209, 78-6211
- Nephrosis, 78-6211
- Chromosome Aberrations
Chromosomes, Human, 6-12, 78-6509
- Chromosomes, Human, 13-15, 78-6507
- Chromosomes, Human, 16-18, 78-6507
- Geographic Distribution, 78-6507
- Colonic Neoplasms
Drug Therapy, 78-6509
- Radiotherapy, 78-6509
- Streptozotocin, 78-6509
- Uracil, 5-Fluoro-, 78-6509
- Urea, 1-(2-Chloroethyl)-3-(4-methylcyclohexyl)-1-nitroso-, 78-6509
- Cyclophosphamide
Case Report, 78-6197
- Diethylamine, 2,2'-Dichloro-*N*-methyl-
Drug Therapy, 78-6219
- Drug Therapy
Case Report, 78-6196, 78-6509
- Glomerulonephritis
Drug Therapy, 78-6209
- Glucosephosphate Dehydrogenase
Enzymatic Activity, 78-6509
- Hepatitis
Case Report, 78-6219
- Hodgkin's Disease
Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
78-6196
- Case Report, 78-6219
- Cyclophosphamide, 78-6196
- Drug Therapy, 78-6196, 78-6219
- Radiotherapy, 78-6196
- p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
78-6196
- s*-Triazine, 2,4,6-Tris(1-aziridinyl)-, 78-6196

Leukemia, Myeloblastic (cont'd)

- Vinblastine, 78-6196
- Immune Serums
 - Antigen-Antibody Reactions, 78-6498
- T-Lymphocytes
 - Lymphocyte Culture Test, Mixed, 78-6494
- Lymphoma
 - Case Report, 78-6508
 - Insecticides, 78-6508
- Plant Agglutinins
 - Colony Formation, 78-6600
 - Immune Response, 78-6600
- Prednisone
 - Drug Therapy, 78-6219
- Purine-6-thiol
 - Case Report, 78-6209
- Radiotherapy
 - Case Report, 78-6509
- Sjogren's Syndrome
 - Cyclophosphamide, 78-6197
- p*-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
 - Drug Therapy, 78-6219
- Vincristine
 - Drug Therapy, 78-6219
- Virus, Epstein-Barr
 - Antigens, Viral, 78-6457

Leukemia, Myelocytic

- Antigens, Neoplasm
 - Antigen-Antibody Reactions, 78-6498
- Chromosomes, Human, 21-22
 - Lymphocyte Culture Test, Mixed, 78-6494
- Immune Serums
 - Antigen-Antibody Reactions, 78-6498
- T-Lymphocytes
 - Blast Crisis, 78-6494
 - Lymphocyte Culture Test, Mixed, 78-6494
- Plant Agglutinins
 - Colony Formation, 78-6600
 - Immune Response, 78-6600
- Virus, Epstein-Barr
 - B-Lymphocytes, 78-6461
 - Virus Replication, 78-6461

Leukocytes

- C.I. Food Brown 3, Disodium Salt
 - Toxicity, Pig, 78-6149
- α -Cresol, 4,6-Dinitro-, Sodium Salt
 - Chromosome Aberrations, 78-6157
- Ultraviolet Rays
 - Chromatids, 78-6356

Leydig Cell Tumor

- Antigenic Determinants
 - A-Type Particles, 78-6415
- Virus, Murine Mammary Tumor
 - Viral Proteins, 78-6415

Leydig Cells

- Caffeine
 - Hyperplasia, 78-6123
- Theobromine
 - Hyperplasia, 78-6123

LH

- Breast Neoplasms
 - Genetics, 78-6349
- Uterine Neoplasms
 - Leiomyoma, 78-6541
 - Menstruation, 78-6541

Lip Neoplasms

- Fibrosarcoma
 - Epidemiology, 78-6380
- Radiotherapy
 - Epidemiology, 78-6380

Lipase

- Metabolic Diseases
 - Enzymatic Activity, 78-6589

Lipomatosis

- Citric Acid
 - Phosphofructokinase, 78-6589
- Phosphofructokinase
 - Enzymatic Activity, 78-6589

Lipopolysaccharides

- Virus, Abelson Murine Leukemia
 - Macrophages, 78-6419

Liposomes

- Virus, Epstein-Barr
 - 1,3,5-Hexatriene, 1,6-Diphenyl-, 78-6458
 - Virus-Cell Interactions, 78-6458

Liver

- Benzo(a)pyrene
 - Nucleic Acids, Binding, 78-6326
- Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 - Nucleic Acids, Binding, 78-6326
- Indan, 5-Hydroxyamino-
 - Quantitation Method, 78-6103
- Lymphosarcoma
 - Neoplasm Metastasis, Review, 78-6060

Liver Cirrhosis

- Cerebrohepatorenal Syndrome
 - Case Report, 78-6530
- Hepatoma
 - Australia Antigen, 78-6531
 - Epidemiology, Japan, 78-6531
- Hodgkin's Disease
 - Case Report, 78-6513
- Sarcoma, Reticulum Cell
 - Case Report, 78-6513

Liver Diseases

- Uterine Neoplasms
 - Androst-4-ene-3,17-dione, 78-6028
 - Epidemiology, Review, 78-6028
 - Estrone, 78-6028

Liver Neoplasms

- Acenaphthene, 5-Nitro-
 - Carcinogenic Potential, 78-6254
- Acetamide, *N*-Fluoren-2-yl-
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6193
 - Precancerous Conditions, 78-6492
- Adenoma
 - Carcinoembryonic Antigen, 78-6352
 - Contraceptives, Oral, 78-6352, 78-6354
 - Estrogens, 78-6352
- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 - Cytochrome P-450, 78-6218
- Aniline, 4,4'-Methylenedi-
 - Carcinogenic Potential, Dog, 78-6261
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Precancerous Conditions, 78-6193
- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 - Nucleotides, Cyclic, 78-6276
 - Precancerous Conditions, 78-6276

Neoplasms (cont'd)

- Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-
 - Carcinogenic Activity, Mouse, 78-6286
- Benzidine, 3,3'-Dichloro-
 - Neoplasm Metastasis, 78-6270
- Cholangioma
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6276
- Contraceptives, Oral
 - Case Report, 78-6352, 78-6354, 78-6355
 - Ethinodiol Diacetate, 78-6354
 - Mestranol, 78-6354, 78-6355
 - Norethynodrel, 78-6355
 - Receptors, Hormone, 78-6355
- Cytochrome P-450
 - Precancerous Conditions, 78-6218
- Diethylamine, *N*-Nitroso-
 - Cytochrome P-450, 78-6218
 - Precancerous Conditions, 78-6492
- Estradiol
 - Receptors, Hormone, 78-6355
- Estrogens
 - Case Report, 78-6352
- Hydrocarbons, Chlorinated
 - Occupational Hazard, Review, 78-6014
- Hyperplasia
 - Contraceptives, Oral, 78-6355
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-6355
- Pesticides
 - Carcinogenic Activity, Review, 78-6011
- Piperidine, 4-Chloro-1-nitroso-
 - Carcinogenic Activity, Rat, 78-6240
- Precancerous Conditions
 - Cerebrohepatorenal Syndrome, 78-6530
 - Transplantation, Homologous, 78-6492
- Progesterone
 - Receptors, Hormone, 78-6355
- Siderosis
 - Gluconic Acid, Iron Salt, 78-6193
 - Precancerous Conditions, 78-6193
 - 8-Quinololinol, 78-6193
- Spironolactone
 - Carcinogenic Potential, Review, 78-6016
- Virus, Avian Leukosis
 - Isolation and Characterization, 78-6388
- Virus, Helper, 78-6389

Neoplasms

- Adenocarcinoma
 - Asbestos, 78-6168
 - Epidemiology, 78-6547
- Adenoma
 - Carbamic Acid, Ethyl Ester, 78-6185
 - Carbamic Acid, Vinyl Ester, 78-6185
 - Xanthine, 3-Hydroxy-, 78-6249
- Air Pollution
 - Fossil Fuels, Review, 78-6019
- Asbestos
 - Epidemiology, 78-6168
 - Epidemiology, Review, 78-6002

Lung Neoplasms (cont'd)

- Occupational Hazard, 78-6002, 78-6168
- Ultrastructural Study, 78-6529
- Beryllium Oxide
 - Inhalation Study, Rat, 78-6127
- Bromine
 - Water Pollution, 78-6556
- Carbamic Acid, Ethyl Ester
 - 4,4'-Bipyridinium, 1,1'-Dimethyl-, Dichloride, 78-6184
 - Carcinogenic Activity, Mouse, 78-6185
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-6184
- Carbamic Acid, Vinyl Ester
 - Carcinogenic Activity, Mouse, 78-6185
- Carbon Dioxide
 - Air Pollution, 78-6019
 - Epidemiology, Review, 78-6019
- Carcinoid Tumor
 - APUD Cells, Review, 78-6063
- Carcinoma
 - APUD Cells, Review, 78-6063
 - Epidemiology, 78-6547
- Carcinoma, Bronchiolar
 - Epidemiology, 78-6547
- Carcinoma, Epidermoid
 - Asbestos, 78-6529
 - Occupational Hazard, 78-6529
- Carcinoma In Situ
 - Precancerous Conditions, 78-6528
- Cell Cycle Kinetics
 - EMT6 Mouse Tumor, 78-6598
- Diethylamine, *N*-Nitroso-
 - Anti-Antibodies, 78-6417
 - Immunization, 78-6417
- Epidemiology
 - California, 78-6548
 - Inbred Mice, 78-6595
 - Utah, 78-6559
- Ethnic Groups
 - Epidemiology, Hawaii, 78-6546
- Growth
 - EMT6 Mouse Tumor, 78-6598
- Hyperplasia
 - Precancerous Conditions, 78-6528
- Mesothelioma
 - Asbestos, 78-6168
- Metaplasia
 - Precancerous Conditions, 78-6528
- Neoplasm Transplantation
 - Cells, Cultured, 78-6598
- Nickel
 - Industrial Dusts, 78-6143
 - Inhalation Study, Rat, 78-6143
- Occupational Hazard
 - Case Report, 78-6529
- Plutonium Oxide
 - Beryllium Oxide, 78-6127
 - Inhalation Study, Rat, 78-6127
- Precancerous Conditions
 - Ultrastructural Study, 78-6528
- Radon
 - Occupational Hazard, 78-6554
- Silicosis
 - Epidemiology, 78-6547
 - Histological Study, 78-6547
- Smoking
 - Epidemiology, Review, 78-6031

Lung Neoplasms (cont'd)**Uranium**

- Body Height, 78-6554
- Epidemiology, 78-6554
- Occupational Hazard, 78-6554
- Smoking, 78-6554

Virus, Epstein-Barr

- B-Lymphocytes, 78-6461
- Virus Replication, 78-6461

Xanthine, 3-Hydroxy-

- Transplacental Carcinogenesis, 78-6249

Lymph Nodes**Plutonium Oxide**

- Body Burden, 78-6369

Sarcoma

- T-Lymphocytes, 78-6502
- Virus, Moloney Murine Sarcoma, 78-6502

Lymphatic System**Benzo(a)pyrene**

- Metabolism, Rat, 78-6327

Lymphocyte Depletion

- Virus, Moloney Murine Leukemia
- T-Lymphocytes, 78-6432

Lymphocyte Transformation**Butyric Acid, 2-Amino-4-(ethylthio)-**

- Concanavalin A, 78-6205
- Plant Agglutinins, 78-6205

Virus, Epstein-Barr

- Antigens, Viral, 78-6454

Virus, Murine Mammary Tumor

- Antigens, Viral, 78-6497
- Immune Response, 78-6497
- Immune Serums, 78-6497

Lymphocytes**Acetaldehyde**

- Chromatids, 78-6188

Acetanilide

- Aryl Hydrocarbon Hydroxylases, 78-6292

p-Acetophenetidine

- Aryl Hydrocarbon Hydroxylases, 78-6292

Aniline, p-(Phenylazo)-

- Immunosuppression, 78-6278

Antipyrine

- Aryl Hydrocarbon Hydroxylases, 78-6292

Asbestos

- Aryl Hydrocarbon Hydroxylases, 78-6169

Benz(a)anthracene

- Aryl Hydrocarbon Hydroxylases, 78-6322

Benzene

- Acetylglucosaminidase, 78-6258

Butyric Acid, 2-Amino-4-(ethylthio)-

- Cell Cycle Kinetics, 78-6205
- RNA, Transfer, Methyltransferases, 78-6205

Carcinoma, Bronchogenic

- Aryl Hydrocarbon Hydroxylases, 78-6322

Cholanthrene, 3-Methyl-

- Aryl Hydrocarbon Hydroxylases, 78-6322

Hodgkin's Disease

- Virus, Epstein-Barr, 78-6452

Laryngeal Neoplasms

- Aryl Hydrocarbon Hydroxylases, 78-6322

Lead

- Chromosome Aberrations, 78-6147

Macrophages

- Cytotoxic Activity, Review, 78-6080

Lymphocytes (cont'd)**Mercury, (Acetato)phenyl-**

- Karyotyping, 78-6140

Phosphide Sulfide, Tris(1-aziridinyl)-

- Chromosome Aberrations, 78-6226

Phosphinde Oxide, 1,4-Piperazinediylbis(bis(1-aziridinyl))-

- Chromosome Aberrations, 78-6226

Smoking

- Aryl Hydrocarbon Hydroxylases, 78-6174

Theophylline

- Aryl Hydrocarbon Hydroxylases, 78-6292

Virus, Epstein-Barr

- Antigens, Viral, 78-6454
- Fetal Blood, 78-6454

Virus, Friend Murine Leukemia

- Immunosuppression, 78-6427

Zinc

- Chromosome Aberrations, 78-6147

B-Lymphocytes**Hodgkin's Disease**

- Virus, Epstein-Barr, 78-6461

Leukemia

- Antigen-Antibody Reactions, 78-6498

Leukemia, Lymphoblastic

- Precursor Cells, Review, 78-6078

Leukemia, Myelocytic

- Virus, Epstein-Barr, 78-6461

Lung Neoplasms

- Virus, Epstein-Barr, 78-6461

Virus, Epstein-Barr

- Fluorescence Polarization, 78-6458

Virus, Marek's Disease Herpes

- Transplantation Immunology, 78-6406

Virus, Moloney Murine Leukemia

- Cell Transformation, Neoplastic, 78-6432
- Immunoglobulins, Surface, 78-6432

T-Lymphocytes**Cell Transformation; Neoplastic**

- Immune Response, Review, 78-6077
- Immunologic Technics, Review, 78-6077
- Lymphocytotoxicity, Review, 78-6077

Immunity, Cellular

- Complement, 78-6495
- Cytotoxic Effector Cells, 78-6495
- Histocompatibility Antigens, 78-6495
- Isoantigens, 78-6495
- Receptors, Antigen, 78-6495

Leukemia

- Benz(a)anthracene, 7,12-Dimethyl-, 78-6314
- Lymphocytotoxicity, 78-6495

Leukemia, Lymphoblastic

- Lymphocyte Culture Test, Mixed, 78-6494
- Virus, Murine Leukemia, 78-6489

Leukemia, Myeloblastic

- Lymphocyte Culture Test, Mixed, 78-6494

Leukemia, Myelocytic

- Blast Crisis, 78-6494
- Lymphocyte Culture Test, Mixed, 78-6494

Macrophages

- Hypersensitivity, Delayed, Review, 78-6079

Sarcoma

- Antigens, Neoplasm, 78-6502
- Immunity, Passive, 78-6502
- Killer Cells, 78-6502
- Lymph Nodes, 78-6502

lymphocytes (cont'd)

- Lymphocytotoxicity, 78-6502
- Virus, Moloney Murine Sarcoma, 78-6502
- Sarcoma, Mast Cell
 - Lymphocytotoxicity, 78-6495
- Virus, Epstein-Barr
 - Fluorescence Polarization, 78-6458
- Virus, Marek's Disease Herpes
 - Transplantation Immunology, 78-6406
- Virus, Moloney Murine Leukemia
 - Cell Transformation, Neoplastic, 78-6432
 - Lymphocyte Depletion, 78-6432
- Virus, RD-114
 - Concanavalin A, 78-6441
- Virus, SV40
 - Antilymphocyte Serum, 78-6500
 - Immunity, Cellular, 78-6500

lymphocytosis

- Virus, Bovine Leukemia
 - Epidemiology, Review, 78-6088

lymphoid Tissue

- Benzoic Acid, 2-Amino-3-hydroxy-
 - Kidney, 78-6311
- Graft Rejection
 - Transplantation, Homologous, 78-6491
- Homologous Wasting Disease
 - Transplantation, Homologous, 78-6491
- Virus, Adeno 7
 - Precancerous Conditions, 78-6468

lymphoma (General and Unspecified)

- Acetic Acid, (Ethylenebis(oxyethylenenitrilo))tetra-
 - Rosette Formation, AKR Mouse, 78-6503
- Age Factors
 - Epidemiology, 78-6562
- Antigens
 - Immunity, Cellular, 78-6487
- Benzidine, 2,2',5,5'-Tetrachloro-
 - Carcinogenic Potential, 78-6254
- Bladder Neoplasms
 - Water Pollution, 78-6556
- Bromine
 - Water Pollution, 78-6556
- Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
 - Case Report, 78-6209
- Cat
 - Epidemiology, California, 78-6548
- Cholanthrene, 3-Methyl-
 - Dose-Response Study, 78-6321
- Coal Tar
 - Occupational Hazard, Review, 78-6017
- Complement
 - Rosette Formation, AKR Mouse, 78-6503
- Epidemiology
 - Australia, 78-6562
 - Inbred Mice, 78-6595
- Ethanol, 2,2',2''-Nitrilotris-
 - Carcinogenic Activity, Mouse, 78-6224
- Glomerulonephritis
 - Drug Therapy, 78-6209
- Graft Rejection
 - Genetics, 78-6491
- Heavy Chain Disease
 - Epidemiology, 78-6487
- Histocompatibility Antigens
 - Graft Rejection, 78-6491
 - Homologous Wasting Disease, 78-6491

Lymphoma (General and Unspecified) (cont'd)

- Homologous Wasting Disease
 - Genetics, 78-6491
- Immunologic Deficiency Syndromes
 - Epidemiology, Review, 78-6031
 - IgG, 78-6487
 - IgM, 78-6487
- Leukemia, Myeloblastic
 - Case Report, 78-6508
 - Insecticides, 78-6508
- Neoplasms, Radiation-Induced
 - Virus, Radiation Leukemia, 78-6418
- Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-
 - Nitrous Acid, Sodium Salt, 78-6241
- 1-Propanol, 2,3-Dibromo-, Phosphate
 - Mutagenic Activity, 78-6144
- Thyroid Neoplasms
 - Immunoglobulins, Surface, 78-6505
 - Ultrastructural Study, 78-6505
- Vibrio cholerae*
 - Epidemiology, 78-6487
- Virus, Abelson Murine Leukemia
 - Macrophages, 78-6419
- Virus, Feline Leukemia
 - Antigens, Viral, 78-6443
 - RNA, Viral, 78-6441
- Virus, Gross Murine Leukemia
 - Thymus Gland, Mouse, 78-6432
 - Ultrastructural Study, 78-6432
- Virus, Marek's Disease Herpes
 - Virus-Like Particles, 78-6407
- Virus, Moloney Murine Leukemia
 - Thymus Gland, Mouse, 78-6432
 - Ultrastructural Study, 78-6432
- Virus, Murine Leukemia
 - Complement, 78-6503
 - Virus Replication, 78-6418
- Virus, Radiation Leukemia
 - Virus Replication, 78-6418
- Virus, RD-114
 - RNA, Viral, 78-6441

Lymphoma, Giant Follicular

- Epidemiology
 - Australia, 78-6562

Lymphosarcoma

- Antigens, Neoplasm
 - Antigen-Antibody Reactions, 78-6498
- Cholangitis
 - Case Report, 78-6513
- Ethnic Groups
 - Epidemiology, Cameroon, 78-6568
- Gonadotropins, Equine
 - Carcinogenic Activity, Mouse, 78-6344
- Immune Serums
 - Antigen-Antibody Reactions, 78-6498
- Liver
 - Neoplasm Metastasis, Review, 78-6060
- Phagocytosis
 - Kupffer Cells, 78-6526
- Radiation, Ionizing
 - Parabiosis, 78-6373
- Thymus Neoplasms
 - Virus, Murine Leukemia, 78-6421
- Virus, Bovine Leukemia
 - Epidemiology, Review, 78-6088
 - Horizontal/Vertical Transmission, Review, 78-6088

Lymphosarcoma (cont'd)

Virus, SV40

Antigen-Antibody Reactions, 78-6085

Histocompatibility Antigens, 78-6085

Lysine12-*O*-Tetradecanoylphorbol-13-acetate

Peptide Hydrolases, 78-6307

Macrophages

Carrageen

Transplantation Immunology, 78-6490

Cell Movement

Immune Response, Review, 78-6079

Cell Transformation, Neoplastic

Immune Response, Review, 78-6080

Phagocytosis, Review, 78-6079

Chromosome Abnormalities

Mitosis, Review, 78-6079

Fibroblasts

Iododeoxyuridine Incorporation, 78-6435

Fibrosarcoma

Transplantation Immunology, 78-6490

Hematopoietic Stem Cells

Colony Stimulating Factor, 78-6488

Inflammation

Mitosis, Review, 78-6079

Interferon

Immune Response, Review, 78-6077

Lymphocytes

Cytotoxic Activity, Review, 78-6080

T-Lymphocytes

Hypersensitivity, Delayed, Review, 78-6079

Lymphoma

Virus, Abelson Murine Leukemia, 78-6419

Neoplasm Metastasis

Immune Response, Review, 78-6077

Neoplasms

Chemotaxis, Review, 78-6081

Immunosuppression, Review, 78-6081

Neoplasms, Experimental

Antitumor Activity, Review, 78-6082

Immune Response, Review, 78-6082

Prostaglandins E

Hematopoietic Stem Cells, 78-6488

Radiation, Ionizing

Cytotoxic Activity, Review, 78-6080

Reserpine

Transplantation Immunology, 78-6490

Starch

Cytostatic Activity, 78-6435

Thymoma

Virus, Abelson Murine Leukemia, 78-6419

Virus, Abelson Murine Leukemia

Cell Transformation, Neoplastic, 78-6419

Lipopolysaccharides, 78-6419

Virus, Helper, 78-6419

Virus Replication, 78-6419

Virus, Influenza

Cytostatic Activity, 78-6435

Virus, Murine Leukemia

Cytostatic Activity, 78-6435

Virus, Murine Sarcoma

Cytostatic Activity, 78-6435

Virus, Sendai

Cytostatic Activity, 78-6435

Malabsorption Syndromes

Stomach Neoplasms

Malabsorption Syndromes (cont'd)

Vitamin B12, 78-6534

Malathion

Cells, Cultured

Metabolism, Lung, 78-6305

Maleic Acid*Salmonella typhimurium*

Genetic Duplication, 78-6359

Maleic Hydrazide

see 3,6-Pyridazinedione, 1,2-Dihydro-

Maleimide, *N*-Phenyl-Benz(e)indole, 3-(β -Nitrovinyl)-

Benzocarbazole Derivatives, Synthesis, 78-6299

Mammary Neoplasms, Experimental

Adenocarcinoma

4-Biphenylamine, 3,2'-Dimethyl-, 78-6297

Phagocytosis, 78-6526

Radiation, Ionizing, 78-6050

Selenium, 78-6146

4,4'-Stilbenediol, α,α' -Diethyl-, 78-6342

Adenocarcinoma, Papillary

Benz(a)anthracene, 7,12-Dimethyl-, 78-6315

Adenofibroma

Radiation, Ionizing, 78-6050, 78-6342

Xanthine, 3-Hydroxy-, 78-6250

Adenoma

Xanthine, 3-Hydroxy-, 78-6250

Benz(a)anthracene, 7,12-Dimethyl-

Dietary Fats, Review, 78-6037

Estradiol, 78-6351

Estriol, 78-6351

Prolactin, 78-6343

Testosterone, 78-6343

Virus-Like Particles, 78-6315

Benz(a)anthracene, 7,8,12-Trimethyl-

2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
78-63172-Naphthol, 1-((4-(α -Tolylazo)- α -tolyl)azo)-, 78-6317

4-Biphenylamine, 3,2'-Dimethyl-

Carcinogenic Activity, Rat, 78-6297

Glucuronidase, 78-6297

Intestinal Microflora, 78-6297

Carcinoma

Benzidine, 2,2',5,5'-Tetrachloro-, 78-6254

Xanthine, 3-Hydroxy-, 78-6250

Carcinoma, Ductal

Acenaphthene, 5-Nitro-, 78-6254

Cholanthrene, 3-Methyl-

Dose-Response Study, 78-6321

Neoplasm Metastasis, 78-6321

Reserpine, 78-6321

Epidemiology

Inbred Mice, 78-6595

Estradiol

Neonatal Hormone Treatment, 78-6346

Pituitary Hormones, 78-6346

Ethane, 1,2-Dichloro-

Carcinogenic Activity, Mouse, Rat, 78-6114

Fibrosarcoma

Benzidine, 2,2',5,5'-Tetrachloro-, 78-6254

Hyperplasia

Age Factors, Rat, 78-6525

Estrus, 78-6525

Mitosis, 78-6525

Precancerous Conditions, Rat, 78-6525

Primary Neoplasms, Experimental (cont'd)

- Hyperthermia**
 - Neoplasm Metastasis, 78-6381
- Neoplasms, Multiple Primary**
 - Cholanthrene, 3-Methyl-, 78-6321
- Parity**
 - Precancerous Conditions, 78-6527
- Phagocytosis**
 - Kupffer Cells, 78-6526
- 1-Piperazineethanol, 4-(3-(2-Chlorophenothiazin-10-yl)propyl)-**
 - 5 α -Androstan-3-one, 17 β -Hydroxy-, 78-6343
 - Androstenediol, 78-6343
 - Prolactin, 78-6343
 - Testosterone, 78-6343
- Precancerous Conditions**
 - Transplantation, Homologous, 78-6527
- Progesterone**
 - Neonatal Hormone Treatment, 78-6346
 - Pituitary Hormones, 78-6346
- Prolactin**
 - Castration, 78-6343
 - Dietary Fats, Review, 78-6037
 - Neonatal Hormone Treatment, 78-6346
 - Pituitary Hormones, 78-6346
- Radiation, Ionizing**
 - Dose Fractionation, Review, 78-6050
 - In Utero Exposure, Rat, 78-6382
- Receptors, Hormone**
 - Estradiol, 78-6351
 - Estriol, 78-6351
 - Estriol, 16-Epi-, 78-6351
- Selenium**
 - Antineoplastic Activity, 78-6146
- Spironolactone**
 - Carcinogenic Potential, Review, 78-6016
- 4,4'-Stilbenediol, α,α' -Diethyl-**
 - Carcinogenic Potential, 78-6315
 - Radiation, Ionizing, 78-6342
- Testosterone**
 - Castration, 78-6343
 - Metabolism, 78-6343
- Virus, Murine Mammary Tumor**
 - A-Type Particles, 78-6415
 - B-Type Particles, 78-6413
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6315
 - DNA-RNA Hybridization, 78-6315
 - Horizontal Transmission, 78-6416
 - Pituitary Implants, 78-6411
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-6315
 - Strain Difference, 78-6416
 - Vertical Transmission, 78-6416
- Xanthine, 3-Hydroxy-**
 - Dose-Response Study, Rat, 78-6250
 - Prolactin, 78-6250
 - Transplacental Carcinogenesis, 78-6250

Andelic Acid

- Styrene**
 - Metabolism, 78-6252

Manganese

- Chromium**
 - Co-carcinogenic Activity, 78-6561
- Stomach Neoplasms**
 - Food Contamination, 78-6148

Manganese Chloride

- Cell Division**

Manganese Chloride (cont'd)

- Mutagenic Activity, Algae, 78-6137**
- Dose-Response Study**
 - Mutagenic Activity, Algae, 78-6137
- Plectonema boryanum*
 - Streptomycin Resistance, 78-6137

Manganese Sulfate

- Dose-Response Study**
- Mutagenic Activity, Rat, 78-6138**

Mebendazole

- see 2-Benzimidazolecarbamic Acid, 5-Benzoylmethyl Ester**

Medulloblastoma

- Neoplasm Metastasis**
- Case Report, 78-6517**

Melanin

- Nicotine**
 - Body Burden, Mouse, 78-6177

Melanoma

- Brain**
 - Neoplasm Metastasis, Review, 78-6060
- Carcinogen, Chemical**
 - Mathematical Model, Review, 78-6044
- Cholestan-3 β -ol, 5 α ,6 α -Epoxy-**
 - Chromosome Aberrations, 78-6362
- Genetics**
 - Fish, Review, 78-6059
- Lung**
 - Neoplasm Metastasis, Review, 78-6060
- Neoplasm Metastasis**
 - Case Report, 78-6512
- Nevus**
 - Wounds and Injuries, 78-6512
- Occupational Hazard**
 - Epidemiology, 78-6551
- Ovary**
 - Neoplasm Metastasis, Review, 78-6060
- Phagocytosis**
 - Kupffer Cells, 78-6526
- Site Distribution**
 - Epidemiology, Sweden, 78-6578
- Skin Neoplasms**
 - Ultraviolet Rays, 78-6046
- Ultraviolet Rays**
 - Chromosome Aberrations, 78-6362
 - Dose-Response Study, Review, 78-6044
 - Epidemiology, Review, 78-6044
 - Epidemiology, Sweden, 78-6578
 - Mathematical Model, Review, 78-6044
- Virus, Murine Sarcoma-Leukemia**
 - DNA-RNA Hybridization, 78-6438
- Virus, RNA Tumor**
 - DNA-RNA Hybridization, 78-6438
 - Reverse Transcriptase, 78-6438
 - RNA, Viral, 78-6438
 - Virus-Like Particles, 78-6438
- Wounds and Injuries**
 - Case Report, 78-6512

Melatonin

- Breast Neoplasms**
- Pineal Gland Function, Review, 78-6024**
- Receptors, Hormone, Review, 78-6024**

Menarche

- Cervix Neoplasms**

Menarche (cont'd)

- Epidemiology, 78-6573
- Uterine Neoplasms
- Epidemiology, 78-6573

Meningomyelocele

- Spinal Neoplasms
- Carcinoma, Epidermoid, 78-6519
- Case Report, 78-6519

Menopause

- Breast Neoplasms
- Epidemiology, 78-6543
- Cervix Neoplasms
- Epidemiology, 78-6573
- Uterine Neoplasms
- Epidemiology, 78-6573

Menstruation

- Uterine Neoplasms
- FSH, 78-6541
- LH, 78-6541
- Somatotropin, 78-6541

6-Mercaptopurine

- see Purine-6-thiol

Mercury

- Bone Marrow Cells
- Chromosome Aberrations, 78-6139
- Chlorophyll
- Mutagenic Activity, 78-6134
- Drosophila melanogaster*
- Mutagenic Activity, 78-6139

Mercury, (Acetato)phenyl-

- Lymphocytes
- Karyotyping, 78-6140
- Occupational Hazard
- Chromosomes, 78-6140

Mercury, Chloroethyl-

- Bone Marrow Cells
- Chromosome Aberrations, 78-6139
- Drosophila melanogaster*
- Mutagenic Activity, 78-6139

Mesenchymoma

- Aflatoxin B1
- Cells, Cultured, 78-6245
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
- Cells, Cultured, 78-6245
- Methanesulfonic Acid, Ethyl Ester
- Cells, Cultured, 78-6245
- 1,2-Oxathiolane, 2,2-Dioxide
- Cells, Cultured, 78-6245
- 2-Oxetanone
- Cells, Cultured, 78-6245
- Quinoline, 4-Nitro-, 1-Oxide
- Cells, Cultured, 78-6245

Mesothelioma

- Lung Neoplasms
- Asbestos, 78-6168

Mestranol

- Liver Neoplasms
- Contraceptives, Oral, 78-6354, 78-6355

Metabolic Diseases

- Lipase
- Enzymatic Activity, 78-6589
- Triclycerides

Metabolic Diseases (cont'd)

- Enzymatic Activity, 78-6589

Metals

- Radiation, Ionizing
- Mutagenic Activity, 78-6134
- Testicular Neoplasms
- Disgerminoma, 78-6563
- Epidemiology, 78-6563
- Neoplasms, Embryonal and Mixed, 78-6563
- Occupational Hazard, 78-6563
- Teratoid Tumor, 78-6563

Metaplasia

- Cervix Uteri
- Organ Culture, 78-6599
- Lung Neoplasms
- Precancerous Conditions, 78-6528

Methacrylic Acid, Ethyl Ester

- Fibrosarcoma
- Plastic Implants, 78-6372
- Tryptophan, 78-6372

Methacrylic Acid, Methyl Ester

- Fibrosarcoma
- Plastic Implants, 78-6372
- Tryptophan, 78-6372

Methampyrone

- see Methanesulfonic Acid, (Antipyrinylmethylamino)-, Monosodium S
- Salt

Methane, Azoxy-

- Ames Test
- Mutagenic Activity, 78-6106
- Cytochrome P-450
- N-Demethylation, 78-6106
- Dietary Fats
- Metabolism, 78-6106
- Formaldehyde
- Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6106
- 7,8-Benzoflavone, 78-6106
- Cholanthrene, 3-Methyl-, 78-6106
- Cobalt Chloride, 78-6106
- Disulfide, Bis(diethylthiocarbamoyl)-, 78-6106
- Metabolism, 78-6106
- Valeric Acid, 2,2-Diphenyl-, 2-(Diethylamino)ethyl Ester HCl, 78-6106
- Methanol, (Methyl-ONN-azoxy)-
- Metabolism, Rat, 78-6107
- Microsomes, Liver
- N-Demethylation, 78-6106
- Metabolism, 78-6106
- Mutagenic Activity, 78-6106
- Pyrazole
- Alcohol Oxidoreductases, 78-6107
- Metabolism, Rat, 78-6107

Methane, Diiodo-

- Benzo(a)pyrene
- Fluorescence Quenching, 78-6329
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
- Fluorescence Quenching, 78-6329

Methane Sulfinylbis-

- Erythroleukemia
- Cell Differentiation, 78-6426
- Virus, Friend Spleen Focus-Forming

ane Sulfinylbis- (cont'd)
 Virus-Like Particles, 78-6428

anesulfonic Acid, (Antipyrinylmethylamino)-, Monosodium Salt
 Ames Test
 Mutagenic Activity, 78-6163

anesulfonic Acid, Butyl Ester
 Hypoxanthine Phosphoribosyltransferase
 Mutagenic Activity, 78-6164
 Lymphoblasts, Diploid
 Mutagenic Activity, 78-6164

anesulfonic Acid, Ethyl Ester
 Ames Test
 Genetic Duplication, 78-6359, 78-6360

Benizidine
 Chromosome Aberrations, 78-6162

Caffeine
 Chromosome Aberrations, 78-6120

Cell Transformation, Neoplastic
 Diploid Cells, 78-6245

Cells, Cultured
 Colony Formation, Agar, 78-6245

Chromosome Aberrations
 Micronucleus Test, 78-6162

DNA Repair
 Liver, Rat, 78-6216

Ethyl Alcohol
 Mutagenic Activity, 78-6108

Fibroblasts
 Mutagenic Activity, 78-6161

Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Mutagenic Activity, 78-6161

Mesenchymoma
 Cells, Cultured, 78-6245

Peptide Hydrolases
 Mutation, 78-6161

Phosphoglucomutase
 Mutation, 78-6161

Saccharomyces cerevisiae
 Growth, Radiation-Sensitive Mutants, 78-6244

Salmonella typhimurium
 Genetic Duplication, 78-6359, 78-6360

anesulfonic Acid, Methyl Ester
 Chromosome Aberrations
 Cell Cycle Kinetics, 78-6358

Endonucleases
 Chromosome Aberrations, 78-6358

Guanosine
 RNA, Methylation, 78-6228

Virus, SV40
 Cell Transformation, Neoplastic, 78-6242

Yeasts
 RNA, Methylation, 78-6228

anol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
 Blood Vessels
 Carcinogenic Metabolite, 78-6189

Intestinal Neoplasms
 Colostomy, Rat, 78-6189
 Neoplasms, Multiple Primary, 78-6189

12-*O*-Tetradecanoylphorbol-13-acetate
 Mutagenic Activity, 78-6295

anol, (Methyl-*ONN*-azoxy)-
 Methane, Azoxy-
 Metabolism, Rat, 78-6107

Methanol, (Methyl-*ONN*-azoxy)- (cont'd)
 Pyrazole
 Alcohol Oxidoreductases, 78-6107

Methoxyamine
 see Hydroxylamine, *O*-Methyl-

Methoxyflurane
 see Ether, 2,2-Dichloro-1,1-difluoroethyl Methyl

2-Methyl-4-(dimethylamino)azobenzene
 see *m*-Toluidine, *N,N*-Dimethyl-4-(phenylazo)-

Methyl Fluorosulfonate
 see Fluorosulfuric Acid, Methyl Ester

Methylamine
 Hexamethylenetetramine
 Nitrosation, 78-6225

4,4'-Methylenebis(2-chloroaniline)
 see Benzenamine, 4,4'-Methylenebis(2-chloro)-

Metrifonate
 Sodium Chloride
 Carcinogenic Potential, Rat, 78-6145

Mezerein
 HeLa Cells
 Epidermal Growth Factor, Binding, 78-6306
 Nerve Growth Factor
 Ganglia, Chick Embryo, 78-6309

Microsomes, Liver
 Acetanilide, 4'-Hydroxy-
 Metabolism, 78-6291
p-Acetophenetidine, *N*-Hydroxy-
 Metabolism, 78-6291
p-Acetophenetidine
 Metabolism, 78-6291

Aflatoxin B1
 DNA Degradation, 78-6124

Ames Test
 Bleaching Effluents, Kraft Mills, 78-6132

Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 Free Radicals, 78-6218

Barbituric Acid, 5-Ethyl-5-phenyl-
 Hydro-Lyases, 78-6320
 Hydroxylases, 78-6320

Benzene, 4-Allyl-1,2-(methylenedioxy)-
 Ames Test, 78-6290
 Mutagenic Activity, 78-6290

Benzo(a)pyrene
 Ames Test, 78-6302
 Inactivating Mechanisms, 78-6302
 Nucleic Acids, Binding, 78-6326

Benzo(a)pyrene, 7,8-Dihydro-7,8-dihydroxy-
 Nucleic Acids, Binding, 78-6326

Benzo(f)quinoline
 Ames Test, 78-6239

Benzo(f)quinoline, 2-Amino-
 Ames Test, 78-6239

Benzo(h)quinoline
 Ames Test, 78-6239

Benzo(h)quinoline, 2-Amino-
 Ames Test, 78-6239

Carcinogen, Chemical
 Enzyme Activation, 78-6340

Cholanthrene, 3-Methyl-
 Enzymatic Activity, 78-6320
 Hydro-Lyases, 78-6320
 Hydroxylases, 78-6320

Microsomes, Liver (cont'd)

- Chromate
 - Metabolism, 78-6129
 - Mutagenic Activity, 78-6129
- Chromic Acid
 - Ames Test, 78-6130
- Chromic Acid, Dipotassium Salt
 - Ames Test, 78-6130
- Chromic Acid, Lead Salt
 - Ames Test, 78-6130
- Chromic Acid, Zinc Salt
 - Ames Test, 78-6130
- Cyclopentadiene, Hexachloro-
 - Mutagenic Activity, 78-6167
- Cyclopentadiene, Tetrachloro-
 - Mutagenic Activity, 78-6167
- Dichromic Acid, Disodium Salt
 - Ames Test, 78-6130
- Diethylamine, *N*-Nitroso-
 - Free Radicals, 78-6218
- Dimethylamine
 - Ames Test, 78-6290
- Ethane, 1,2-Dibromo-
 - Metabolism, Rat, 78-6115
- Ethanol, 2,2',2''-Nitrilotris-
 - Mutagenic Activity, 78-6224
- Formic Acid, Ethylenemethylhydrazide
 - Ames Test, 78-6181
- Formic Acid, 1-Methyl Hydrazide
 - Ames Test, 78-6181
- Hydrazine, Methyl-
 - Ames Test, 78-6181
- Isoquinoline
 - Ames Test, 78-6239
- Methane, Azoxy-
 - N*-Demethylation, 78-6106
 - Metabolism, 78-6106
 - Mutagenic Activity, 78-6106
- Mutagens
 - Enzyme Activation, 78-6340
- Phenanthrene, 1,2-Dihydro-1,2-dihydroxy-
 - Carcinogenic Metabolite, 78-6312
- Phenanthrene, 9,10-Dihydro-9,10-dihydroxy-
 - Carcinogenic Metabolite, 78-6312
- Phenanthrene, 9,10-Dihydro-9,10-epoxy-
 - Carcinogenic Metabolite, 78-6312
- Phenanthridine
 - Ames Test, 78-6239
- Phenol, (1,1-Dimethylethyl)-4-methoxy-
 - Hydro-Lyases, 78-6257
- Phenol, 2-Methoxy-4-(2-propenyl)-
 - Ames Test, 78-6290
 - Mutagenic Activity, 78-6290
- Piperidine
 - Ames Test, 78-6290
- Proline
 - Ames Test, 78-6290
- 1-Propanone, 3-(Dimethylamino)-1-(3',4'-methylenedioxyphenyl)-
 - Ames Test, 78-6290
- 1-Propanone, 3-Piperidyl-1-(3',4'-methylenedioxyphenyl)-
 - Ames Test, 78-6290
 - Mutagenic Activity, 78-6290
- 1-Propanone, 3-Pyrrolidinyl-1-(3',4'-methylenedioxyphenyl)-
 - Ames Test, 78-6290
- 5*H*-Pyrido(4,3-*b*)indole, 3-Amino-1-methyl-

Microsomes, Liver (cont'd)

- DNA, Binding, 78-6246
- Pyrrolidine
 - Ames Test, 78-6290
- Pyrrolidine, 1-Nitroso-
 - Carcinogenic Metabolite, 78-6237
- Quinoline
 - Ames Test, 78-6239
- Synfuel A-3
 - Ames Test, 78-6102
- Triazene, 3,3-Dimethyl-1-(*p*-chlorophenyl)-
 - Ames Test, 78-6223
- Triazene, 3,3-Dimethyl-1-phenyl-
 - Ames Test, 78-6223
- Water Pollutants
 - Ames Test, 78-6158
- Misonidazole
 - see 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-
- Mitochondria
 - Calcium
 - Cell Transformation, Neoplastic, Review, 78-6061
 - Carcinogen, Chemical
 - Oxidative Phosphorylation, Review, 78-6061
 - DNA
 - Cell Transformation, Neoplastic, Review, 78-6061
 - Oncogenic Viruses
 - Oxidative Phosphorylation, Review, 78-6061
- Mitomycin C
 - Antineoplastic Agents
 - Ames Test, 78-6232
 - Chromosome Aberrations
 - Cell Cycle Kinetics, 78-6358
 - Endonucleases
 - Chromosome Aberrations, 78-6358
 - Hepatocytes
 - Cytotoxicity, Fibroblasts, 78-6267
- Mitosis
 - Benzenediazosulfonic Acid, *p*-(Dimethylamino)-, Sodium Salt
 - Chromosomes, 78-6284
 - Fensulfothion
 - Chromosomes, 78-6284
 - Mammary Neoplasms, Experimental
 - Hyperplasia, 78-6525
 - p*-Phenylenediamine, *N,N*-Dimethyl-
 - Chromosomes, 78-6284
 - Salicylanilide, 3,3',5,5',6-Pentachloro-2'-hydroxy-
 - Allium cepa*, 78-6255
 - Chromosome Abnormalities, 78-6255
- Monocytes
 - Hematopoietic Stem Cells
 - Colony Stimulating Factor, 78-6488
 - Prostaglandins E
 - Hematopoietic Stem Cells, 78-6488
- Morpholine, Hydrochloride
 - Feces
 - Metabolism, Rat, 78-6251
 - Urine
 - Metabolism, Rat, 78-6251
- Mouth Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-
 - Dose-Response Study, Hamster, 78-6316
 - Localization, Cheek Pouch, 78-6316
 - Carcinoma, Epidermoid

ath Neoplasms (cont'd)

- Epidemiology, 78-6375
- Ethyl Alcohol, 78-6375
- Smoking, 78-6375
- Wounds and Injuries, 78-6375

Dog

- Epidemiology, California, 78-6548

Inflammation

- Dentures, 78-6375

Petroleum

- Epidemiology, Review, 78-6018
- Wounds and Injuries
- Epidemiology, 78-6375

Multiple Myeloma**Bence Jones Protein**

- Case Report, 78-6514

Immunoglobulins, Heavy Chain

- Bence Jones Protein, 78-6514
- Cytoplasm, 78-6514

Pyridase**Ames Test**

- Pyrolysis Product, 78-6201

Tagens**Bleaching Effluents, Kraft Mills**

- Isolation and Characterization, 78-6132

Cytochrome P-450

- Enzyme Activation, 78-6340

DNA Repair

- Bioassay, 78-6099

Drosophila melanogaster

- Eye Pigmentation, 78-6368
- Nitroimidazoles, 78-6234

Epoxide Hydratases

- Inactivating Mechanisms, 78-6302

Glutamic Acid

- Dipyrido(1,2- α :3',2'- d)imidazole, 2-Amino-, 78-6198
- Dipyrido(1,2- α :3',2'- d)imidazole, 2-Amino-6-methyl-, 78-6198

- Pyrolysis Products, 78-6198

Glutathione Transferases

- Inactivating Mechanisms, 78-6302

Microsomes, Liver

- Enzyme Activation, 78-6340

Occupational Hazard

- Women, Review, 78-6022

1-Propanol, 2,3-Dibromo-, Phosphate

- Urine, Rat, 78-6144

Scabies Fungoides**Epidemiology**

- Australia, 78-6562

Scotoxins**Colonic Neoplasms**

- Epidemiology, Review, 78-6034

Cytochrome P-450

- Carcinogenic Metabolite, Review, 78-6020

Food Contamination

- Quantitation Method, 78-6101

Gyna**Cervix Neoplasms**

- Organ Culture, 78-6599

Pyosmine

- see Pyridine, 3-(1-Pyrrolin-2-yl)-

NADP**Chromate**

- Metabolism, 78-6129

Cyclophosphamide

- Carcinogenic/Mutagenic Activity, 78-6195

Ethane, 1,2-Dichloro-

- Ames Test, 78-6113
- Mutagenic Activity, 78-6113

2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-

- Benz(a)anthracene, 7,8,12-Trimethyl-, Oxidoreductases, 78-6317

Leukemia

- Benz(a)anthracene, 7,12-Dimethyl-, 78-6317
- Benz(a)anthracene, 7,8,12-Trimethyl-, 78-6317
- Mammary Neoplasms, Experimental
- Benz(a)anthracene, 7,8,12-Trimethyl-, 78-6317

2-Naphthol, 1-Amino-

- Hydroxylamine, *N*-1-Naphthyl-, Solvolysis Products, 78-6247

4-Naphthol, 1-Amino-

- Hydroxylamine, *N*-1-Naphthyl-, Solvolysis Products, 78-6247

2-Naphthol, 1-(Phenylazo)-

- Ames Test
- Mutagenic Activity, 78-6152

2-Naphthol, 1-(α -Tolylazo)-

- Ames Test
- Mutagenic Activity, 78-6152

2-Naphthol, 1-((4-(α -Tolylazo)- α -tolyl)azo)-

- Leukemia
- Benz(a)anthracene, 7,8,12-Trimethyl-, 78-6317
- Mammary Neoplasms, Experimental
- Benz(a)anthracene, 7,8,12-Trimethyl-, 78-6317

2-Naphthol, 1-(2,4-Xylylazo)-

- Ames Test
- Mutagenic Activity, 78-6152

1,4-Naphthoquinone

- Benz(e)indole, 3-(β -Nitrovinyl)-, Benzocarbazole Derivatives, Synthesis, 78-6299

1-Naphthylamine

- Acetylation Kinetics
- Species Difference, 78-6264
- Acetyltransferases
- Acetylation Kinetics, 78-6264
- Occupational Hazard
- Isoantigens, 78-6566

2-Naphthylamine

- Acetylation Kinetics
- Species Difference, 78-6264
- Acetyltransferases
- Acetylation Kinetics, 78-6264
- Chromosome Aberrations
- Micronucleus Test, 78-6162
- Occupational Hazard
- Isoantigens, 78-6566
- Saccharomyces cerevisiae*
- Growth, Radiation-Sensitive Mutants, 78-6244
- Virus, SV40
- Cell Transformation, Neoplastic, 78-6242

2-Naphthylamine, 1-(Phenylazo)-

- Ames Test
- Mutagenic Activity, 78-6152

2-Naphthylamine, 1-(o-Tolylazo)-
Ames Test

Mutagenic Activity, 78-6152

Nasopharyngeal Neoplasms

Carcinogen, Environmental
Epidemiology, 78-6064

Carcinoma

Ultrastructural Study, 78-6064

Precancerous Conditions

Ultrastructural Study, 78-6064

Smoking

Epidemiology, 78-6552

Socioeconomic Factors

Epidemiology, 78-6552

Virus, Epstein-Barr

Antibodies, Viral, 78-6552

Epidemiology, 78-6552

Epidemiology, Review, 78-6089

Natamycin

see Pimaricin

Necrosis

Chloroform

Hepatotoxicity, Rat, 78-6109

Neoplasm Metastasis

Antigens, Neoplasm

Tumor-Associated Products, Review, 78-6065

Carcinosarcoma

Hypertension, 78-6332

Clostridiopeptidase A

Tumor-Associated Products, Review, 78-6065

Contraceptives, Oral

Case Report, 78-6353

Hepatoma

Contraceptives, Oral, 78-6353

Hypertension, 78-6332

Hormones

Tumor-Associated Products, Review, 78-6065

Liver Neoplasms

Benzidine, 3,3'-Dichloro-, 78-6270

Macrophages

Immune Response, Review, 78-6077

Mammary Neoplasms, Experimental

Cholanthrene, 3-Methyl-, 78-6321

Hyperthermia, 78-6381

Medulloblastoma

Case Report, 78-6517

Melanoma

Case Report, 78-6512

Neoplasms, Experimental

Blood Vessels, Review, 78-6060

Prostaglandins

Tumor-Associated Products, Review, 78-6065

Thyroid Neoplasms

Carcinoma, 78-6363, 78-6379, 78-6522

Radiation, Ionizing, 78-6363, 78-6376

Neoplasm Recurrence, Local

Thyroid Neoplasms

Radiation, Ionizing, 78-6376

Neoplasm Regression, Spontaneous

Virus, Rous Sarcoma

Genetics, 78-6400

Neoplasm Transplantation

Carcinoma

Hypertension, 78-6332

Neoplasm Transplantation (cont'd)

Glioma

Cell Line, 78-6581

Lung Neoplasms

Cells, Cultured, 78-6598

Neoplasms, Experimental

Clone Cells, 78-6597

Organ Transplant

Epidemiology, Review, 78-6076

Sarcoma

Hypertension, 78-6332

Virus, Murine Mammary Tumor

Antigen-Antibody Reactions, 78-6409

Antigens, Viral, 78-6409

Virus-Like Particles, 78-6409

Virus Replication, 78-6409

Neoplasms (General and Unspecified)

Carcinogen, Environmental

Epidemiology, Review, 78-6030

Chromosomes

Hereditary Diseases, Review, 78-6074

Diet

Epidemiology, Review, 78-6032

Drosophila melanogaster

Cell Differentiation, 78-6075

Environmental Hazard

Water Supply, Fluoridated, 78-6550

Ethyl Alcohol

Epidemiology, Review, 78-6032

Genetics

DNA Repair, Review, 78-6055

Epidemiology, Review, 78-6073

Growth

Mathematical Model, 78-6555

Hereditary Diseases

DNA Repair, Review, 78-6055

Epidemiology, Review, 78-6030

Histocompatibility Antigens

Hereditary Diseases, Review, 78-6074

Immunoglobulins, Fc

Antigen-Antibody Complex, 78-6504

Cell Adhesion, 78-6504

Hemadsorption, 78-6504

Rosette Formation, 78-6504

Immunologic Deficiency Syndromes

Epidemiology, Review, 78-6030

Macrophages

Chemotaxis, Review, 78-6081

Immunosuppression, Review, 78-6081

Organ Transplant

Epidemiology, Review, 78-6076

Histological Study, Review, 78-6076

Immunosuppression, Review, 78-6076

Smoking

Epidemiology, Review, 78-6032

Stress

Hormones, Review, 78-6058

Immune Response, Review, 78-6058

Transplantation, Homologous

Kidney, Heart, Review, 78-6076

Tumor Latency

Mathematical Model, 78-6555

Neoplasms, Embryonal and Mixed

Testicular Neoplasms

Metals, 78-6563

Virus, Rous Sarcoma

Neoplasms, Embryonal and Mixed (cont'd)

Genetics, 78-6400

Neoplasms, Experimental**Chromosomes**

Clone Cells, 78-6597

Concanavalin A

Growth, 78-6597

Enzymes

Neoplasm Metastasis, Review, 78-6060

Genetics

Animal Model, Fish, 78-6059

Glioma

Cell Line, 78-6581

Growth

Clone Cells, 78-6597

Immune Response

Neoplasm Metastasis, Review, 78-6060

Macrophages

Antitumor Activity, Review, 78-6082

Immune Response, Review, 78-6082

Neoplasm Metastasis

Blood Vessels, Review, 78-6060

Neoplasm Transplantation

Clone Cells, 78-6597

Piperidine, 1-Nitroso-

Carcinogenic Activity, Rat, 78-6240

Virus, Adeno 12

Clone Cells, 78-6597

Ultrastructural Study, 78-6471

Virus-Like Particles, 78-6471

Virus, RNA Tumor

Co-carcinogenic Effect, Review, 78-6096

Virus, SV40

Cell Differentiation, 78-6085

Neoplasms, Multiple Primary**Bladder Neoplasms**

Benzidine, 3,3'-Dichloro-, 78-6270

Cecal Neoplasms

1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385

Colonic Neoplasms

Genetics, 78-6538

Esophageal Neoplasms

Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273

Epidemiology, 78-6549

Head and Neck Neoplasms

Epidemiology, 78-6549

Hepatoma

Benzidine, 3,3'-Dichloro-, 78-6270

Serine, Diazoacetate (Ester), 78-6199

Intestinal Neoplasms

Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester) 78-6189

Mammary Neoplasms, Experimental

Cholanthrene, 3-Methyl-, 78-6321

Neoplastic Endocrine-Like Syndromes

Case Report, 78-6518

Parathyroid Neoplasms

Radiation, Ionizing, 78-6377, 78-6379

Thyroid Neoplasms

Radiation, Ionizing, 78-6377, 78-6379

Virus, Epstein-Barr

Antigens, Viral, 78-6459

Genetics, 78-6459

Neoplasms, Radiation-Induced

Age Factors

Neoplasms, Radiation-Induced (cont'd)

Epidemiology, Review, 78-6031

Carbon Radioisotopes

Dose-Response Study, 78-6386

Lymphoma

Virus, Radiation Leukemia, 78-6418

Neoplastic Endocrine-Like Syndromes

Genetics

Case Report, 78-6518

Histocompatibility Antigens

Genetics, 78-6518

Neoplasms, Multiple Primary

Case Report, 78-6518

Nephrosis

Leukemia, Myeloblastic

Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)- 78-6211

Nerve Growth Factor

Ingenol Dibenzoate

Ganglia, Chick Embryo, 78-6309

Mezerein

Ganglia, Chick Embryo, 78-6309

Phorbol 12,13-Didecanoate

Ganglia, Chick Embryo, 78-6309

12-*O*-Tetradecanoylphorbol-13-acetate

Cell Survival, 78-6309

Ganglia, Chick Embryo, 78-6309

Neuraminidase

Cell Differentiation

Trophoblast, 78-6591

Glucosamine

Trophoblast, 78-6591

Neurilemmoma

Trigeminal Nerve

Case Report, 78-6515

Neuroblastoma

Genetics

Fish, Review, 78-6059

Neuroleptics

see Tranquilizing Agents

Neurons

Propionic Acid, 3,3'-Thiodi-

Carcinogenic Activity, 78-6117

Tissue Culture, 78-6117

Urea, Ethyl Nitroso-

Cell Transformation, Neoplastic, 78-6231

Neutral Red

DNA

Alkaline Sucrose Gradients, 78-6200

Pancreas, Liver, Rat, 78-6200

Nevus

Melanoma

Wounds and Injuries, 78-6512

Nickel

Chromium

Co-carcinogenic Activity, 78-6561

Fibrosarcoma

Industrial Dusts, 78-6143

Lung Neoplasms

Industrial Dusts, 78-6143

Inhalation Study, Rat, 78-6143

Mammae

Nickel (cont'd)

Metabolism, Mouse, 78-6142

Nickel Sulfide

Blood Cell Volume, Rat, 78-6141

Placenta

Metabolism, Mouse, 78-6142

Polycythemia

Blood Cell Volume, Rat, 78-6141

Radioisotopes

Body Burden, Mouse, 78-6142

Metabolism, Mouse, 78-6142

Tissue Distribution, 78-6142

Sarcoma, Reticulum Cell

Industrial Dusts, 78-6143

Nickel Selenide**Polycythemia**

Blood Cell Volume, Rat, 78-6141

Nickel Sulfide**Nickel**

Blood Cell Volume, Rat, 78-6141

Polycythemia

Strain Difference, Rat, 78-6141

Nicotinamide**Islet Cell Tumor**

Insulin, 78-6207

Kidney Neoplasms

Streptozotocin, 78-6208

Pancreatic Neoplasms

Islet Cell Tumor, 78-6207

Streptozotocin, 78-6208

Nicotine**Bladder**

Body Burden, Mouse, 78-6177

1-Butanone, 4-(*N*-Methyl-*N*-nitrosamino)-1-(3-pyridyl)-

Synthesis, 78-6176

Butyraldehyde, 4-(*N*-Methyl-*N*-nitrosamino)-4-(3-pyridyl)-

Synthesis, 78-6176

Maternal-Fetal Exchange

Metabolism, Mouse, 78-6177

Melanin

Body Burden, Mouse, 78-6177

Respiratory System

Body Burden, Mouse, 78-6177

Nicotine, 1'-Nitroso-1'-demethyl-**Ames Test**

Carcinogenic Metabolite, 78-6175

1-Butanone, 4-Hydroxy-1-(3-pyridyl)-

Hydroxylation, 78-6175

Butyric Acid, 4-Hydroxy-4-(3-pyridyl)-

Urinary Metabolite, 78-6175

Butyric Acid, 4-Oxo-4-(3-pyridyl)-

Urinary Metabolite, 78-6175

2-Furanol 5-(3-Pyridyl)tetrahydro-

Hydroxylation, 78-6175

2-Furanone, 5-(3-Pyridyl)tetrahydro-

Urinary Metabolite, 78-6175

Pyridine, 3-(1-Pyrrolin-2-yl)-

Hydroxylation, 78-6175

Tobacco

E and Z Isomers, 78-6176

Nitric Acid, Potassium Salt**Stomach Neoplasms**

Epidemiology, Chile, 78-6574

Nitric Acid, Potassium Salt (cont'd)

Fertilizers, 78-6574

Nitric Acid, Sodium Salt

Hexamethylenetetramine

Nitrosation, 78-6225

Stomach Neoplasms

Epidemiology, Chile, 78-6574

Fertilizers, 78-6574

Nitrosamines**Carcinogen, Environmental**

Quantitation Method, 78-6006

Cytochrome P-450

Carcinogenic Metabolite, Review, 78-6020

Occupational Hazard

Carcinogenic Activity, Review, 78-6009

Quantitation Method, Review, 78-6009

Stomach Neoplasms

Potato Sap, 78-6210

Nitroso Compounds**Antineoplastic Agents**

Ames Test, 78-6232

***N*-Nitrosoethylenethiourea**

see 2-Imidazolidinethione, *N*-Nitroso-

Nitrous Acid**Ames Test**

Genetic Duplication, 78-6359

Benzimidazole, 2-(4-Thiazolyl)-

Mutagenic Activity, 78-6256

Benzoic Acid

Mutagenic Activity, 78-6256

Carcinogen, Environmental

Quantitation Method, 78-6006

Pimaricin

Mutagenic Activity, 78-6256

Propionic Acid, Calcium Salt

Mutagenic Activity, 78-6256

Salmonella typhimurium

Genetic Duplication, 78-6359

Nitrous Acid, Sodium Salt**Atrophy**

Testis, Rat, 78-6123

Caffeine

Spermatogenesis, 78-6123

Chromosome Aberrations

Lymphocytes, Rat, 78-6214

Ethanol, 2,2',2''-Nitrilotris-

Mutagenic Activity, 78-6224

Intestinal Neoplasms

Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-, 78-6241

Lymphoma

Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-, 78-6241

Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-
Carcinogenic Metabolite, 78-6241

Spermidine

Ames Test, 78-6204

Mutagenic Activity, 78-6204

Stomach Neoplasms

Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-, 78-6241

Theobromine

Spermatogenesis, 78-6123

Theophylline

us Acid, Sodium Salt (cont'd)
 Spermatogenesis, 78-6123

Tryptophan
 Nitrosation, Meat, 78-6203

Tryptophan, Nitroso-
 Hydrogen-Ion Concentration, 78-6203
 Tryptophan, 78-6203

thisterone Acetate
 Hepatoma
 Contraceptives, Oral, 78-6353

thisterone, 18-Methyl-
 Immunosuppression
 Carcinogenic Activity, Review, 78-6051

thynodrel
 Liver Neoplasms
 Contraceptives, Oral, 78-6355

urman
 Benzo(f)quinoline, 2-Amino-
 Ames Test, 78-6239
 Benzo(h)quinoline, 2-Amino-
 Ames Test, 78-6239

Radiation, Ionizing
 DNA Repair, 78-6155
 Mutagenic Activity, 78-6155

Ultraviolet Rays
 DNA Repair, 78-6155
 Mutagenic Activity, 78-6155

e Neoplasms
 Adenocarcinoma
 Reverse Transcriptase, 78-6521
 Ultrastructural Study, Sheep, 78-6521
 Virus-Like Particles, 78-6521

Cystadenoma, Papillary
 Reverse Transcriptase, 78-6521
 Ultrastructural Study, Sheep, 78-6521
 Virus-Like Particles, 78-6521

Piperidine, 4-Chloro-1-nitroso-
 Carcinogenic Activity, Rat, 78-6240

Virus, Herpes
 Virus-Like Particles, 78-6521

Virus, Vaccinia
 Virus-Like Particles, 78-6521

Virus, Visna
 Virus-Like Particles, 78-6521

aldin
 see Methanesulfonic Acid, (Antipyrinylmethylamino)-,
 Monosodium S
 Salt

leic Acids
 Carbamic Acid, Ethyl Ester
 Liver, Binding, 78-6185

Dimethylamine, N-Nitroso-
 Methylation, 78-6217

Ethane, 1,2-Dibromo-
 Proteins, 78-6115

leoproteins
 Virus, Avian Reticuloendotheliosis
 Cell Transformation, Neoplastic, 78-6405
 DNA-RNA Hybridization, 78-6405
 Isolation and Characterization, 78-6405

leotides
 Virus, Polyoma
 RNA, 78-6482

Nucleotides, Cyclic
 Liver Neoplasms
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6276

Obesity
 Breast Neoplasms
 Hormones, Review, 78-6037
 Precancerous Conditions, 78-6553

Uterine Neoplasms
 Androst-4-ene-3,17-dione, 78-6028
 Epidemiology, Review, 78-6028, 78-6066
 Estrone, 78-6028

Occupational Hazard
 Acrylonitrile
 Carcinogenic Activity, Review, 78-6021
 Exposure Standards, Review, 78-6021

Asbestos
 Aryl Hydrocarbon Hydroxylases, 78-6169
 Precancerous Conditions, 78-6383

Benzidine
 Isoantigens, 78-6566

Bladder Neoplasms
 Epidemiology, 78-6567
 Epidemiology, Review, 78-6002
 Isoantigens, 78-6566

Breast Neoplasms
 Epidemiology, 78-6551

Carbamoyl Chloride, Diethyl-
 Chromosome Aberrations, 78-6187

Carbamoyl Chloride, Dimethyl-
 Chromosome Aberrations, 78-6187

Carcinogen, Chemical
 Epidemiology, 78-6551
 Women, Review, 78-6022

Cervix Neoplasms
 Epidemiology, 78-6551

Chromium
 Epidemiology, Japan, 78-6561

Digestive System Neoplasms
 Vanadium, 78-6561

Ether, Bis(chloromethyl)-
 Carcinogenic Activity, Review, 78-6009
 Quantitation Method, Review, 78-6009

Hair Dyes
 Chromosomes, 78-6151

Head and Neck Neoplasms
 Chromium, 78-6561

Hydrazine
 Carcinogenic Activity, Review, 78-6009
 Quantitation Method, Review, 78-6009

Hydrocarbons, Chlorinated
 Carcinogenic Activity, Review, 78-6009
 Quantitation Method, Review, 78-6009

Intestinal Neoplasms
 Epidemiology, 78-6551

Kidney Neoplasms
 Chromium, 78-6561
 Vanadium, 78-6561

Leukemia
 Epidemiology, Review, 78-6025

Leukemia, Lymphoblastic
 Chromium, 78-6561

Lung Neoplasms
 Asbestos, 78-6002, 78-6168
 Carcinoma, Epidermoid, 78-6529
 Case Report, 78-6529

Occupational Hazard (cont'd)

- Radon, 78-6554
 - Uranium, 78-6554
 - Melanoma
 - Epidemiology, 78-6551
 - Mercury, (Acetato)phenyl-Chromosomes, 78-6140
 - Mutagens
 - Women, Review, 78-6022
 - 1-Naphthylamine
 - Isoantigens, 78-6566
 - 2-Naphthylamine
 - Isoantigens, 78-6566
 - Nitrosamines
 - Carcinogenic Activity, Review, 78-6009
 - Quantitation Method, Review, 78-6009
 - Petroleum
 - Epidemiology, Review, 78-6018
 - Prostatic Neoplasms
 - Asbestos, 78-6168
 - Radiation, Ionizing
 - Dosimetry, Review, 78-6047
 - Epidemiology, Review, 78-6025, 78-6047
 - Respiratory Tract Neoplasms
 - Chromate, 78-6131
 - Epidemiology, 78-6560
 - Vanadium, 78-6561
 - Sarcoma
 - Acetic Acid, (2,4-Dichlorophenoxy)-, 78-6288
 - Acetic Acid, (2,4,5-Trichlorophenoxy)-, 78-6288
 - Epidemiology, 78-6288
 - Sarcoma, Reticulum Cell
 - Chromium, 78-6561
 - Styrene
 - Chromosome Aberrations, 78-6252
 - Teratogens
 - Women, Review, 78-6022
 - Testicular Neoplasms
 - Metals, 78-6563
- Octane, 1-Iodo-**
Benzo(a)pyrene
 - Fluorescence Quenching, 78-6329
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
Fluorescence Quenching, 78-6329
- Oil Orange SS**
see 2-Naphthol, 1-(*o*-Tolylazo)-
- Oligomycin**
Virus, Rous Sarcoma
 - Glycolysis, 78-6396
- Phenol, 2,4-Dinitro-, 78-6396

Oncogenic Viruses

- Breast Neoplasms
 - Vertical Transmission, Review, 78-6068
- Burkitt's Lymphoma
 - Epidemiology, Review, 78-6031
- Cell Membrane Permeability
 - Cell Transformation, Neoplastic, Review, 78-6061
- Cell Transformation, Neoplastic
 - Precancerous Conditions, Review, 78-6084
- Cervix Neoplasms
 - Epidemiology, Review, 78-6031
- Hodgkin's Disease
 - Epidemiology, Review, 78-6031
- Mitochondria

Oncogenic Viruses (cont'd)

- Oxidative Phosphorylation, Review, 78-6061
 - Sarcoma, Reticulum Cell
 - Epidemiology, Review, 78-6031
- Osmium Tetroxide**
Bacillus subtilis
 - Mutagenic Activity, 78-6128
- Plasmacytoma
 - Endonucleases, 78-6371
- Ovalbumin**
Acetamide, Thio-
 - Estradiol, 3-Benzoate, 78-6347
 - Plasma Levels, Chick, 78-6347
- Actinomycin D
 - Estradiol, 3-Benzoate, 78-6347
 - Plasma Levels, Chick, 78-6347
- Ovarian Neoplasms**
Disgerminoma
 - Genetics, Review, 78-6069
- Fibroma
 - Genetics, 78-6540
- Genetics
 - Case Report, 78-6540
- Radiation, Ionizing
 - In Utero Exposure, Rat, 78-6382
- Ovary**
Melanoma
 - Neoplasm Metastasis, Review, 78-6060
- Oxalic Acid, Butylhydrazino-**
Ames Test
 - S-9 Fraction, 78-6182
- 1,2-Oxathiolane, 2,2-Dioxide**
Cell Transformation, Neoplastic
 - Diploid Cells, 78-6245
- Cells, Cultured
 - Colony Formation, Agar, 78-6245
- Mesenchymoma
 - Cells, Cultured, 78-6245
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6242
- 2-Oxetanone**
Cell Transformation, Neoplastic
 - Diploid Cells, 78-6245
- Cells, Cultured
 - Colony Formation, Agar, 78-6245
- Mesenchymoma
 - Cells, Cultured, 78-6245
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6242
- Oxidoreductases**
Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-
 - Microsomes, Testis, 78-6227
- Benz(a)anthracene, 7,8,12-Trimethyl-
2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-
78-6317
- Carbonyl Cyanide, 3-Chlorophenylhydrazino-
Clostridium leptum, 78-6590
- 5 β -Cholan-24-oic Acid, 3 α ,7 α -Dihydroxy-
Bacteroides, 78-6590
Clostridium leptum, 78-6590
- Cholic Acid, Monosodium Salt
Clostridium leptum, 78-6590
Clostridium leptum

Oxidoreductases (cont'd)

- Enzymatic Activity, 78-6590
- Cyclohexanamine, *N,N*-Methanetetraylbis-
Clostridium leptum, 78-6590
- Cyclophosphamide
 - Cell Transformation, Neoplastic, 78-6195
 - Chromatids, 78-6195
 - Chromosome Aberrations, 78-6195
- Dimethylamine, *N*-Nitroso-
Metabolism, Liver, 78-6217
- 2-Furaldehyde, 5-Nitro-, Semicarbazone
Microsomes, Testis, 78-6227
- Intestinal Neoplasms
 - Carcinoma, 78-6587
- Intestinal Polyps
 - Bacteria, 78-6587
- Phenol, 2,4-Dinitro-
Clostridium leptum, 78-6590
- 4-Quinolol, 2-Heptyl-, 1-Oxide
Clostridium leptum, 78-6590

Oxyclozanide

- see Salicylanilide, 3,3',5,5',6-Pentachloro-2'-hydroxy-

Ozone

- Ames Test
 - Water, 78-6105

Palmitic Acid, Morpholino-

- Feces
 - Metabolism, Rat, 78-6251
- Intestines
 - Body Burden, Rat, 78-6251
- Urine
 - Metabolism, Rat, 78-6251

Pancreatic Neoplasms

- Adenoma
 - Streptozotocin, 78-6208
- Islet Cell Tumor
 - Nicotinamide, 78-6207
 - Picolinamide, 78-6207
 - Streptozotocin, 78-6207, 78-6208
- Petroleum
 - Epidemiology, Review, 78-6018
- Serine, Diazoacetate (Ester)
 - Choline, 78-6199
 - Precancerous Conditions, 78-6199
- Streptozotocin
 - Dose-Response Study, Rat, 78-6208
 - Nicotinamide, 78-6208
 - Picolinamide, 78-6208

Papain

- Virus, Epstein-Barr
 - Antigens, 78-6460

Papilloma

- Esophageal Neoplasms
 - Benzylamine, *N*-Methyl-*N*-nitroso-, 78-6273
- Laryngeal Neoplasms
 - Virus, Papilloma, 78-6474
- Skin Neoplasms
 - Carbamic Acid, Ethyl Ester, 78-6185
 - Carbamic Acid, Vinyl Ester, 78-6185
- Stomach Neoplasms
 - Piperazine, 1,4-Dinitroso-, 78-6241
 - Plants, Edible, 78-6210
- Virus, Bovine Papilloma
 - Antigens, Viral, 78-6446

Papilloma (cont'd)

- Isolation and Characterization, 78-6445

Paraquat

- see 4,4'-Bipyridinium, 1,1'-Dimethyl-, Dichloride

Parathyroid Neoplasms

- Adenoma
 - Epidemiology, 78-6379
 - Radiation, Ionizing, 78-6376, 78-6377, 78-6379
- Neoplasms, Multiple Primary
 - Radiation, Ionizing, 78-6377, 78-6379

Parity

- Breast Neoplasms
 - Precancerous Conditions, 78-6553
- Mammary Neoplasms, Experimental
 - Precancerous Conditions, 78-6527

Patulin

- Beverages
 - Concentration Levels, 78-6586
 - Fermentation, 78-6586

Pentadecane, 2,6,10,14-Tetramethyl-

- Plasmacytoma
 - Mouse, Review, 78-6013

Peptic Ulcer

- Colonic Neoplasms
 - Aluminum Hydroxide, 78-6126

Peptide Hydrolases

- Hydroxylases
 - Localization, Liver, 78-6320
- Methanesulfonic Acid, Ethyl Ester
 - Mutation, 78-6161
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Lysine, 78-6307
 - Skin, Mouse, 78-6307
- Virus, Murine Mammary Tumor
 - Viral Proteins, 78-6415

Peptides

- Fibrosarcoma
 - Cell Division, 78-6481
 - Isolation and Characterization, 78-6481
- Sarcoma
 - Virus, Murine Sarcoma, 78-6481
- Virus, Avian Sarcoma
 - Cell Transformation, Neoplastic, 78-6481
- Virus, Murine Sarcoma
 - Cell Transformation, Neoplastic, 78-6481
 - Isolation and Characterization, 78-6481
- Virus, Polyoma
 - Cell Transformation, Neoplastic, 78-6481
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6481

Permanganic Acid, Potassium Salt

- Chromic Acid, Zinc Salt
 - Ames Test, 78-6130

Pesticides

- Carcinogenic Potential
 - Review, 78-6012
- Environmental Hazard
 - Quantitation Method, 78-6101
- Hepatoma
 - Carcinogenic Activity, Review, 78-6011
- Liver Neoplasms
 - Carcinogenic Activity, Review, 78-6011

Petroleum

- Bronchial Neoplasms
 - Carcinogenic Activity, Review, 78-6002
- Digestive System Neoplasms
 - Epidemiology, Review, 78-6018
- Extender Oils
 - Carcinogenic Potential, Review, 78-6018
- Mouth Neoplasms
 - Epidemiology, Review, 78-6018
- Occupational Hazard
 - Epidemiology, Review, 78-6018
- Pancreatic Neoplasms
 - Epidemiology, Review, 78-6018
- Prostatic Neoplasms
 - Carcinogenic Activity, Review, 78-6002
- Respiratory Tract Neoplasms
 - Epidemiology, Review, 78-6018
- Skin Neoplasms
 - Carcinogenic Activity, Review, 78-6002
- Stomach Neoplasms
 - Carcinogenic Activity, Review, 78-6002

Phagocytosis

- Leukemia
 - Kupffer Cells, 78-6526
 - Thymus Gland, 78-6526
- Lymphosarcoma
 - Kupffer Cells, 78-6526
- Mammary Neoplasms, Experimental
 - Adenocarcinoma, 78-6526
 - Kupffer Cells, 78-6526
- Melanoma
 - Kupffer Cells, 78-6526

Pharyngeal Neoplasms

- Carcinoma, Epidermoid
 - Epidemiology, 78-6380
- Radiotherapy
 - Epidemiology, 78-6380

Phenacetin

- see p*-Acetophenetidine

Phenanthrene

- Phenanthrene, 1,2-Dihydro-1,2-dihydroxy-
 - Carcinogenic Metabolite, 78-6312
- Phenanthrene, 9,10-Dihydro-9,10-dihydroxy-
 - Carcinogenic Metabolite, 78-6312
- Phenanthrene, 9,10-Dihydro-9,10-epoxy-
 - Carcinogenic Metabolite, 78-6312

Phenanthrene, 1,2-Dihydro-1,2-dihydroxy-

- Microsomes, Liver
 - Carcinogenic Metabolite, 78-6312
- Phenanthrene
 - Carcinogenic Metabolite, 78-6312

Phenanthrene, 9,10-Dihydro-9,10-dihydroxy-

- Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
 - Carcinogenic Metabolite, 78-6312

Cholanthrene, 3-Methyl-

- Carcinogenic Metabolite, 78-6312
- Microsomes, Liver
 - Carcinogenic Metabolite, 78-6312

Phenanthrene

- Carcinogenic Metabolite, 78-6312

Phenanthrene, 9,10-Dihydro-9,10-epoxy-

- Cyclohexene Oxide
 - Carcinogenic Metabolite, 78-6312
- Escherichia coli*

Phenanthrene, 9,10-Dihydro-9,10-epoxy- (cont'd)

- DNA Repair, 78-6325
- Microsomes, Liver
 - Carcinogenic Metabolite, 78-6312
- Phenanthrene
 - Carcinogenic Metabolite, 78-6312

Phenanthridine

- Ames Test
 - Amino Derivatives, 78-6239
- Microsomes, Liver
 - Ames Test, 78-6239

Phenol, 4,4'-(1,2-Diethylene)di-, *meso*-, Dicaprylate

- Phenol, 4,4'-(1,2-Diethylene)di-, *meso*-
 - Residues, Cattle, 78-6259

Phenol, 4,4'-(1,2-Diethylene)di-, *meso*-

- Phenol, 4,4'-(1,2-Diethylene)di-, *meso*-, Dicaprylate
 - Residues, Cattle, 78-6259
- Residues, Cattle
 - Quantitation Method, 78-6259

Phenol, (1,1-Dimethylethyl)-4-methoxy-

- Androst-4-ene-3,17-dione
 - Transferases, 78-6333
- Benzene, 1-Chloro-2,4-dinitro-
 - Transferases, 78-6333
- Benzene, 1-(Chloromethyl)-4-nitro-
 - Transferases, 78-6333
- Benzene, 1,2-Dichloro-4-nitro-
 - Transferases, 78-6333
- Benzene, (Epoxyethyl)-
 - Hydro-Lyases, 78-6257
- 2-Benzimidazolecarbamate, 5-Benzoylmethyl Ester
 - Mutagenic Activity, 78-6328
- Benzo(a)pyrene
 - Ames Test, 78-6333
 - Carcinogenic Metabolite, 78-6333
 - Mutagenic Activity, 78-6328
- Diazepam
 - Mutagenic Activity, 78-6328
- Hycanthone
 - Mutagenic Activity, 78-6328
- Hydro-Lyases
 - Enzymatic Activity, 78-6257
 - Microsomes, Liver, 78-6257
- Imidazole-1-ethanol, 2-Methyl-5-nitro-
 - Mutagenic Activity, 78-6328
- Schistosomicides
 - Mutagenic Activity, 78-6328
- Transferases
 - Liver, Mouse, Rat, 78-6333

Phenol, 2,4-Dinitro-

- Clostridium leptum*
 - Oxidoreductases, 78-6590
- Virus, Rous Sarcoma
 - Oligomycin, 78-6396

Phenol, 2-Methoxy-4-(2-propenyl)-

- Ames Test
 - Metabolites, 78-6290
- Microsomes, Liver
 - Ames Test, 78-6290
 - Mutagenic Activity, 78-6290

p-Phenyldiamine

- Hair Dyes
 - Ames Test, 78-6269

mylenediamine, *N,N*-Dimethyl-
 Chromosomes
 Mitosis, 78-6284
 Mutagenic Activity, 78-6284
 ine Red 20-7600
 see Fluorescein, 2',4',5',7'-Tetrabromo-, Disodium Salt
 ol 12,13-Dibenzoate
 HeLa Cells
 Epidermal Growth Factor, Binding, 78-6306
 ol 12,13-Didecanoate
Escherichia coli
 DNA Repair, 78-6308
 HeLa Cells
 Epidermal Growth Factor, Binding, 78-6306
 Nerve Growth Factor
 Ganglia, Chick Embryo, 78-6309
 gene
 Chloroform
 Cytochrome P-450, 78-6109
 Metabolism, 78-6109
 phide Sulfide, Tris(1-aziridinyl)-
 Lymphocytes
 Chromosome Aberrations, 78-6226
 Phosphinde Oxide, 1,4-Piperazinediylbis(bis(1-aziridinyl))-
 Chromosome Aberrations, 78-6226
 phinde Oxide, 1,4-Piperazinediylbis(bis(1-aziridinyl))-
 Lymphocytes
 Chromosome Aberrations, 78-6226
 Phosphide Sulfide, Tris(1-aziridinyl)-
 Chromosome Aberrations, 78-6226
 pho fructokinase
 Lipomatosis
 Citric Acid, 78-6589
 Enzymatic Activity, 78-6589
 phoglucomutase
 Methanesulfonic Acid, Ethyl Ester
 Mutation, 78-6161
 phonoacetic Acid
 Burkitt's Lymphoma
 DNA Polymerase, 78-6453
 Virus, Epstein-Barr
 Antigens, Viral, 78-6450
 Virus, Herpes Pan
 Antigens, Viral, 78-6450
 Virus, Herpes Papio
 Antigens, Viral, 78-6450
 Virus, Herpes Pongo
 Antigens, Viral, 78-6450
 sphoric Triamide, Hexamethyl-
 Chromosome Aberrations
 Micronucleus Test, 78-6162
 sphorothioic Acid, *O,O*-Diethyl *O*-(*p*-Nitrophenyl) Ester
 Cells, Cultured
 Metabolism, Lung, 78-6305
 olinamide
 Islet Cell Tumor
 Insulin, 78-6207
 Pancreatic Neoplasms
 Islet Cell Tumor, 78-6207
 Streptozotocin, 78-6208

Pimaricin
 Bacillus subtilis
 Mutagenic Activity, 78-6256
 Nitrous Acid
 Mutagenic Activity, 78-6256
 Piperazine, 1,4-Bis(3-bromo-1-oxopropyl)-
 Antineoplastic Agents
 Ames Test, 78-6232
 Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-
 Intestinal Neoplasms
 Nitrous Acid, Sodium Salt, 78-6241
 Lymphoma
 Nitrous Acid, Sodium Salt, 78-6241
 Nitrous Acid, Sodium Salt
 Carcinogenic Metabolite, 78-6241
 Piperazine, 1,4-Dinitroso-
 Carcinogenic Metabolite, 78-6241
 Stomach Neoplasms
 Nitrous Acid, Sodium Salt, 78-6241
 Piperazine, 1,4-Dinitroso-
 Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-
 Carcinogenic Metabolite, 78-6241
 Stomach Neoplasms
 Carcinoma, Epidermoid, 78-6241
 Papilloma, 78-6241
 1-Piperazineethanol, 4-(3-(2-Chlorophenothiazin-10-yl)propyl)-
 Mammary Neoplasms, Experimental
 5 α -Androstan-3-one, 17 β -Hydroxy-, 78-6343
 Androstenediol, 78-6343
 Prolactin, 78-6343
 Testosterone, 78-6343
 Piperidine
 Aflatoxin B1
 DNA Cleavage, 78-6124
 Microsomes, Liver
 Ames Test, 78-6290
 Piperidine, 4-Chloro-1-nitroso-
 Esophageal Neoplasms
 Carcinogenic Activity, Rat, 78-6240
 Liver Neoplasms
 Carcinogenic Activity, Rat, 78-6240
 Nose Neoplasms
 Carcinogenic Activity, Rat, 78-6240
 Stomach Neoplasms
 Carcinogenic Activity, Rat, 78-6240
 Piperidine, 1-Nitroso-
 Neoplasms, Experimental
 Carcinogenic Activity, Rat, 78-6240
 Piperonyl Butoxide
 Heliotrine
 Chromosome Aberrations, 78-6238
 Mutagenic Activity, 78-6238
 Radiation, Ionizing
 Chromosome Aberrations, 78-6238
 Mutagenic Activity, 78-6238
 Pipobrom
 see Piperazine, 1,4-Bis(3-bromo-1-oxopropyl)-
 Pituitary Gland
 Estradiol
 Histological Study, Mouse, 78-6346
 Progesterone
 Histological Study, Mouse, 78-6346

Pituitary Gland (cont'd)

- Prolactin
 - Histological Study, Mouse, 78-6346

Pituitary Hormones

- Ethnic Groups
 - Menopause, 78-6544
 - Plasma/Urinary Levels, 78-6544
- Mammary Neoplasms, Experimental
 - Estradiol, 78-6346
 - Progesterone, 78-6346
 - Prolactin, 78-6346

Pituitary Neoplasms

- Epidemiology
 - Inbred Mice, 78-6595
- Radiation, Ionizing
 - In Utero Exposure, Rat, 78-6382
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Carcinogenic Activity, Rat, 78-6342

Plant Agglutinins

- Benz(a)anthracene
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - DNA Replication, 78-6205
 - Lymphocyte Transformation, 78-6205
- Cholanthrene, 3-Methyl-
 - Aryl Hydrocarbon Hydroxylases, 78-6322
- Leukemia, Myeloblastic
 - Colony Formation, 78-6600
 - Immune Response, 78-6600
- Leukemia, Myelocytic
 - Colony Formation, 78-6600
 - Immune Response, 78-6600

Plant Tumors

- Agrobacterium tumefaciens*
 - DNA, Bacterial, 78-6583
- DNA, Bacterial
 - Plasmids, 78-6583

Plasmacytoma

- Antigens
 - Mouse, Review, 78-6013
- DNA
 - Endonucleases, 78-6371
- Hexadecane, 2,6,10,14-Tetramethyl-
 - Mouse, Review, 78-6013
- Osmium Tetroxide
 - Endonucleases, 78-6371
- Pentadecane, 2,6,10,14-Tetramethyl-
 - Mouse, Review, 78-6013
- Potassium Chloride
 - Endonucleases, 78-6371
- RNA, Messenger
 - Cytoplasm, 78-6579
 - Isolation and Characterization, 78-6579
 - Nucleic Acid Precursors, 78-6579
- Ultraviolet Rays
 - Endonucleases, 78-6371
 - RNA, Messenger, 78-6579
- Virus, Abelson Murine Leukemia
 - Mouse, Review, 78-6013

Plasmids

- Agrobacterium tumefaciens*
 - Cytokinins, 78-6585

Plasminogen Activators

- Estradiol

Plasminogen Activators (cont'd)

- Uterus, Mouse, 78-6307
- Retinoic Acid
 - Adenosine Cyclic 3',5' Monophosphate, 78-6338
 - Fibroblasts, Chick Embryo, 78-6338
 - RNA, Messenger, 78-6338
- Retinol
 - Fibroblasts, Chick Embryo, 78-6338
- Teratoid Tumor
 - Dibutyryl Cyclic AMP, 78-6339
 - Retinoic Acid, 78-6339
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Adenosine Cyclic 3',5' Monophosphate, 78-6338
 - Antigenic Determinants, 78-6310
 - Cell Transformation, Neoplastic, 78-6310
 - Fibroblasts, Chick Embryo, 78-6338
 - Lung, Hamster, 78-6310
 - Retinoic Acid, 78-6338
- Virus, B77
 - Cell Transformation, Neoplastic, 78-6398
- Virus, Rous Sarcoma
 - Adenosine Cyclic 3',5' Monophosphate, 78-6338
 - Cell Transformation, Neoplastic, 78-6338, 78-6395
 - 78-6397, 78-6398
 - Genes, Viral, 78-6397
 - 12-*O*-Tetradecanoylphorbol-13-acetate, 78-6395
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6310

Platinum, Diaminedichloro-, cis

- Escherichia coli*
 - DNA Repair, 78-6308

Plectonema boryanum

- Manganese Chloride
 - Streptomycin Resistance, 78-6137

Pleural Diseases

- Asbestos
 - Epidemiology, Review, 78-6023
 - Occupational Hazard, Review, 78-6023
 - X-Ray Study, Review, 78-6023
- Asbestosis
 - Occupational Hazard, Review, 78-6023
- Fibrosis
 - Occupational Hazard, Review, 78-6023
- Pleural Neoplasms
 - Precancerous Conditions, Review, 78-6023

Pleural Neoplasms

- Asbestos
 - Precancerous Conditions, 78-6383
- Pleural Diseases
 - Precancerous Conditions, Review, 78-6023

Plutonium

- Tissue Distribution
 - Solvent Extraction Method, 78-6367

Plutonium Oxide

- Alpha Particles
 - Lung, Dispersion, 78-6370
- Beryllium Oxide
 - Alveolar Clearance, 78-6127
- Body Burden
 - Inhalation Study, Pig, 78-6369
- Lung
 - Body Burden, 78-6369
- Lung Neoplasms
 - Beryllium Oxide, 78-6127

onium Oxide (cont'd)

Inhalation Study, Rat, 78-6127

Lymph Nodes

Body Burden, 78-6369

onium**Smoking**

Radioisotopes, 78-6049

y A**Virus, Adeno 2**

DNA-RNA Hybridization, 78-6463

Virus, Friend Murine Leukemia

RNA, Messenger, 78-6580

Virus, Polyoma

RNA, Messenger, 78-6484

ychlorobiphenyl Compounds**Aryl Hydrocarbon Hydroxylases**

Lepidoptera, 78-6296

5H-Pyrido(4,3-b)indole, 3-Amino-1-methyl-

DNA, Binding, 78-6246

ycyclic Hydrocarbons**Automobile Exhaust**

Carcinogenic Activity, 78-6300

Carcinogen, Environmental

Quantitation Method, 78-6301

Colonic Neoplasms

Intestinal Bacteria, 78-6034

Cytochrome P-450

Carcinogenic Metabolite, Review, 78-6020

Environmental Hazard

Quantitation Method, Review, 78-6010

Food Contamination

Quantitation Method, 78-6301

Skin Neoplasms

Automobile Exhaust, 78-6300

Smoke Condensate

Quantitation Method, Review, 78-6010

lycythemia**Nickel**

Blood Cell Volume, Rat, 78-6141

Nickel Selenide

Blood Cell Volume, Rat, 78-6141

Nickel Sulfide

Strain Difference, Rat, 78-6141

olyps**Colonic Neoplasms**

Precancerous Conditions, 78-6538

Intestinal Neoplasms

Bacteria, Review, 78-6035

Precancerous Conditions, Review, 78-6083

Stomach Neoplasms

Gastritis, 78-6533

otassium Chloride**Plasmacytoma**

Endonucleases, 78-6371

otassium Hypochlorite**Ames Test**

Bleaching Effluents, Kraft Mills, 78-6132

recancerous Conditions**Asbestos**

Occupational Hazard, 78-6383

Pleural Plaques, 78-6383

Bladder Neoplasms

Isoantigens, 78-6566

Precancerous Conditions (cont'd)**Brain Neoplasms**

Urea, Ethyl Nitroso-, 78-6231

Breast Neoplasms

Adenofibroma, 78-6553

Age Factors, 78-6553

Cysts, 78-6553

Epidemiology, 78-6553

Obesity, 78-6553

Parity, 78-6553

Carcinogen, Chemical

Carcinogenic Activity, Review, 78-6003

Cervix Neoplasms

Epidemiology, Denmark, 78-6571

Colonic Neoplasms

Genetics, 78-6538

Intestinal Polyps, 78-6536

Polyps, 78-6538

Ultrastructural Study, 78-6536

Ear Neoplasms

Cholesteatoma, 78-6520

Hepatoma

Alpha Fetoproteins, 78-6277

Glutathione, 78-6274

Liver Neoplasms

Acetamide, *N*-Fluoren-2-yl-, 78-6492

Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6193

Benzenamine, *N,N*-Dimethyl-4-((3-

methylphenyl)azo)-, 78-6276

Cerebrohepatorenal Syndrome, 78-6530

Cytochrome P-450, 78-6218

Diethylamine, *N*-Nitroso-, 78-6492

Siderosis, 78-6193

Transplantation, Homologous, 78-6492

Lung Neoplasms

Carcinoma In Situ, 78-6528

Hyperplasia, 78-6528

Metaplasia, 78-6528

Ultrastructural Study, 78-6528

Mammary Neoplasms, Experimental

Parity, 78-6527

Transplantation, Homologous, 78-6527

Nasopharyngeal Neoplasms

Ultrastructural Study, 78-6064

Pancreatic Neoplasms

Serine, Diazoacetate (Ester), 78-6199

Pleural Neoplasms

Asbestos, 78-6383

Stomach Neoplasms

Gastrectomy, 78-6576

Thyroid Neoplasms

Imidazole-2-thiol, 1-Methyl-, 78-6248

Virus, Adeno 7

Graft vs Host Reaction, 78-6468

Lymphoid Tissue, 78-6468

Virus, Murine Mammary Tumor

Hormones, 78-6411

RNA, Viral, 78-6411

Prednisone

Leukemia, Myeloblastic

Drug Therapy, 78-6219

Pregnant Mare's Serum

see Gonadotropins, Equine

Pregnenolone 16 α -Carbonitrile

Hydro-Lyases

Metabolism, Lung, 78-6331

- Pregnenolone 16 α -Carbonitrile (cont'd)**
 Transferases
 Metabolism, Lung, 78-6331
- Progesterone**
 Acetamide, Thio-
 Estradiol, 3-Benzate, 78-6347
 Plasma Levels, Chick, 78-6347
 Actinomycin D
 Estradiol, 3-Benzate, 78-6347
 Plasma Levels, Chick, 78-6347
 Breast Neoplasms
 Ethnic Groups, 78-6544
 Liver Neoplasms
 Receptors, Hormone, 78-6355
 Mammary Neoplasms, Experimental
 Neonatal Hormone Treatment, 78-6346
 Pituitary Hormones, 78-6346
 Menopause
 Ethnic Groups, 78-6544
 Pituitary Gland
 Histological Study, Mouse, 78-6346
- Prolactin**
 Breast Neoplasms
 Dietary Fats, Review, 78-6037
 Genetics, 78-6349
 Mammary Neoplasms, Experimental
 Benz(a)anthracene, 7,12-Dimethyl-, 78-6343
 Castration, 78-6343
 Dietary Fats, Review, 78-6037
 Neonatal Hormone Treatment, 78-6346
 1-Piperazineethanol, 4-(3-(2-Chlorophenothiazin-10-yl)propyl)-, 78-6343
 Pituitary Hormones, 78-6346
 Xanthine, 3-Hydroxy-, 78-6250
 Pituitary Gland
 Histological Study, Mouse, 78-6346
 Virus, Murine Mammary Tumor
 Co-carcinogenic Effect, 78-6411
- Proline**
 Microsomes, Liver
 Ames Test, 78-6290
- Propane, 1-Chloro-2,3-epoxy-**
 Yeasts
 Mutagenic Activity, 78-6116
- Propane, 2,2'-Oxybis(2-chloro)-**
 Ames Test
 Enzyme Activation, 78-6118
 Mutagenic Activity
 Body Fluids, Mouse, 78-6118
Saccharomyces cerevisiae
 Mutagenic Activity, 78-6118
- Propane Sultone**
 see 1,2-Oxathiolane, 2,2-Dioxide
- 1-Propanol, 2,3-Dibromo-, Phosphate**
 Ames Test
 Mutagenic Activity, 78-6144
 Lymphoma
 Mutagenic Activity, 78-6144
 Mutagens
 Urine, Rat, 78-6144
- 2-Propanol, 1,1'-Iminodi-N-nitroso-**
 DNA Replication
 Pancreas, Rat, 78-6212
- 1-Propanone, 3-(Dimethylamino)-1-(3',4'-methylenedioxyphenyl)-**
 Microsomes, Liver
 Ames Test, 78-6290
- 1-Propanone, 3-Piperidyl-1-(3',4'-methylenedioxyphenyl)-**
 Microsomes, Liver
 Ames Test, 78-6290
 Mutagenic Activity, 78-6290
- 1-Propanone, 3-Pyrrolidinyl-1-(3',4'-methylenedioxyphenyl)-**
 Microsomes, Liver
 Ames Test, 78-6290
- β -Propiolactone**
 see 2-Oxetanone
- Propionic Acid, Calcium Salt**
Bacillus subtilis
 Mutagenic Activity, 78-6256
 Nitrous Acid
 Mutagenic Activity, 78-6256
- Propionic Acid, 3,3'-Thiodi-**
 Nerve Degeneration
 Spinal Cord, Frog, 78-6117
 Neurons
 Carcinogenic Activity, 78-6117
 Tissue Culture, 78-6117
- Prostaglandins**
 Neoplasm Metastasis
 Tumor-Associated Products, Review, 78-6065
- Prostaglandins E**
 Hematopoietic Stem Cells
 Colony Stimulating Factor, 78-6488
 Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-
 Colony Stimulating Factor, 78-6488
 Hematopoietic Stem Cells, 78-6488
 Macrophages
 Hematopoietic Stem Cells, 78-6488
 Monocytes
 Hematopoietic Stem Cells, 78-6488
- Prostatic Neoplasms**
 Adenocarcinoma
 Virus, Cytomegalo, 78-6499
 Asbestos
 Occupational Hazard, 78-6168
 Carcinogen, Environmental
 Epidemiology, 78-6558
 Caribbean Islands
 Epidemiology, 78-6570
 Dietary Fats
 Epidemiology, 78-6558
 Ethnic Groups
 Epidemiology, 78-6558
 Petroleum
 Carcinogenic Activity, Review, 78-6002
 Virus, Cytomegalo
 Antibodies, Viral, 78-6499
 Antigen-Antibody Complex, 78-6499
 Hyperplasia, 78-6499
 Immunity, Cellular, 78-6499
- Protamines**
 Cervix Neoplasms
 Sperm Proteins, 78-6569

- ins**
 Carbamic Acid, Ethyl Ester
 Liver, Binding, 78-6185
 Cell Transformation, Neoplastic
 Chromatin, 78-6072
 Enzymatic Activity
 Metabolism, Inborn Errors, 78-6589
Ethane, 1,2-Dibromo-
 Binding, Liver, Kidney, Rat, 78-6115
 Nucleic Acids, 78-6115
HepATOMa
 Gene Control Mechanisms, 78-6072
Ultraviolet Rays
 DNA Repair, Review, 78-6052
Virus, Herpes Simplex 1
 DNA Replication, 78-6448
len, 8-Methoxy-
 Ultraviolet Rays
 Chromatids, 78-6356
len, 4,5',8-Trimethyl-
 Ultraviolet Rays
 Chromatids, 78-6356
ne, 2-Amino-6-methoxy-
 Chromatin
 Enzymatic Removal, 78-6213
 Dimethylamine, *N*-Nitroso-
 Chromatin, 78-6213
 DNA, 78-6213, 78-6215
DNA
 Enzymatic Removal, 78-6213
 Enzymatic Removal
 Liver/Kidney Extracts, Rat, 78-6213
 Triazene, 3,3-Dimethyl-1-phenyl-
 Liver, Brain, Rat, 78-6221
Urea, Methyl Nitroso-
 Chromatin, 78-6213
 Chromatin, Liver, 78-6229
 Dimethylamine, *N*-Nitroso-, 78-6229
 DNA, 78-6213
 Xanthine, 78-6229
ne, 5-(4-Hydroxy-3-methyl-*trans*-2-butenylamino)-
Agrobacterium tumefaciens
 Isolation and Characterization, 78-6585
ne, 6-(3-Methyl-2-butenylamino)-
Agrobacterium tumefaciens
 Isolation and Characterization, 78-6585
ne-6-thiol
 Leukemia, Myeloblastic
 Case Report, 78-6209
azole
 Methane, Azoxy-
 Alcohol Oxidoreductases, 78-6107
 Metabolism, Rat, 78-6107
 Methanol, (Methyl-*ONN*-azoxy)-
 Alcohol Oxidoreductases, 78-6107
ene
 Aryl Hydrocarbon Hydroxylases
 Skin, 78-6336
ene, 3,4-Dimethylene
 Smoke Condensate
 Isolation and Characterization, 78-6337
Pyrene, 3,4-Trimethylene
 Smoke Condensate
 Isolation and Characterization, 78-6337
3,6-Pyridazinedione, 1,2-Dihydro-
 Caffeine
 Chromosome Aberrations, 78-6120
Pyridine, 2-Amino-5-phenyl-
 Ames Test
 Pyrolysis Product, 78-6201
Pyridine, 3-(1-Pyrrolin-2-yl)-
 Ames Test
 Mutagenic Activity, 78-6175
 Nicotine, 1'-Nitroso-1'-demethyl-
 Hydroxylation, 78-6175
5*H*-Pyrido(4,3-*b*)indole, 3-Amino-1-methyl-
 Microsomes, Liver
 DNA, Binding, 78-6246
 Polychlorobiphenyl Compounds
 DNA, Binding, 78-6246
Pyrophosphoric Acid, Disodium Salt
 Dose-Response Study
 Mutagenic Activity, Rat, 78-6138
Pyrrolidine
 Microsomes, Liver
 Ames Test, 78-6290
Pyrrolidine, 1-Nitroso-
 Aroclor 1254
 Carcinogenic Metabolite, 78-6237
 Butyraldehyde, 4-Hydroxy-, 2,4-Dinitrophenylhydrazone
 Carcinogenic Metabolite, 78-6237
 2-Furanol, Tetrahydro-
 Hydroxylation, 78-6237
 Hydrazine, 2,4-Dinitrophenyl-
 Hydroxylation, 78-6237
 Microsomes, Liver
 Carcinogenic Metabolite, 78-6237
Quinacrine
Escherichia coli
 Mutagenic Activity, 78-6159
 Mutagenic Activity
 Galactose Resistance, 78-6159
Salmonella typhimurium
 Mutagenic Activity, 78-6159
Quinacrine Mustard
 Caffeine
 Chromosome Aberrations, 78-6120
Quinoline
 Ames Test
 Amino Derivatives, 78-6239
 Microsomes, Liver
 Ames Test, 78-6239
Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutylamino)-
 Virus, Epstein-Barr
 Antigens, Viral, 78-6451
Quinoline, 6-Ethoxy-1,2-dihydro-2,2,4-trimethyl-
 Benzo(a)pyrene
 Ames Test, 78-6333
 Carcinogenic Metabolite, 78-6333
 Mutagenic Activity, 78-6328
Quinoline, 4-(Hydroxyamino)-, 1-Oxide
 DNA Replication

Quinoline, 4-(Hydroxyamino)-, 1-Oxide (cont'd)
Pancreas, Rat, 78-6212

Quinoline, 4-Nitro-
Saccharomyces cerevisiae
Growth, Radiation-Sensitive Mutants, 78-6244

Quinoline, 4-Nitro-, 1-Oxide
Cell Transformation, Neoplastic
Diploid Cells, 78-6245
Cells, Cultured
Colony Formation, Agar, 78-6245
Chromosome Aberrations
Micronucleus Test, 78-6162
Hepatocytes
Cytotoxicity, Fibroblasts, 78-6267
Mesenchymoma
Cells, Cultured, 78-6245
Virus, SV40
Cell Transformation, Neoplastic, 78-6242

8-Quinolinol
Liver Neoplasms
Siderosis, 78-6193

4-Quinolinol, 2-Heptyl-, 1-Oxide
Clostridium leptum
Oxidoreductases, 78-6590

Quinone, N-Acetylimido-
p-Acetophenetidine, *N*-Hydroxy-
Sulfate Derivative, 78-6291

Quinone Reductases
Acetamide, *N*-Fluoren-2-yl-
Liver, Rat, 78-6334
Benz(a)anthracene
Liver, Rat, 78-6334
Benzo(a)pyrene
Liver, Rat, 78-6334
Coumarin, 3, 3'-Methylenebis(4-hydroxy)-
Liver, Rat, 78-6334

Radiation
Cell Transformation, Neoplastic
Precancerous Conditions, Review, 78-6084

Radiation, Ionizing
Adenocarcinoma
1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385
Adrenal Gland Neoplasms
In Utero Exposure, Rat, 78-6382
Ames Test
Genetic Duplication, 78-6359, 78-6360
Antigens
Hypersensitivity, Delayed, 78-6490
Ataxia Telangiectasia
DNA Repair, 78-6364
Bone Neoplasms
In Utero Exposure, Rat, 78-6382
Breast Neoplasms
Age Factors, Review, 78-6050
Epidemiology, Review, 78-6050
Caffeine
Chromosome Aberrations, 78-6120
Cecal Neoplasms
1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385
Cell Transformation, Neoplastic
Dose-Response Study, Review, 78-6004

Radiation, Ionizing (cont'd)
Chromosome Aberrations
Cell Cycle Kinetics, 78-6358
Co-carcinogenic Effect
Mathematical Model, 78-6048
Dose-Response Study
Risk Factors, 78-6111
Drosophila melanogaster
Eye Pigmentation, 78-6368
Mutagenic Activity, 78-6238
Endonucleases
Chromosome Aberrations, 78-6358
Ethyl Alcohol
Mutagenic Activity, 78-6108
Fibrosarcoma
Transplantation Immunology, 78-6490
Harman
DNA Repair, 78-6155
Mutagenic Activity, 78-6155
Imidazole-1-ethanol, 2-Methyl-5-nitro-
Chromosome Aberrations, 78-6235
Immunosuppression
Carcinogenic Activity, Review, 78-6051
Intestinal Neoplasms
Adenocarcinoma, 78-6385
1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385
In Utero Exposure, Rat, 78-6382
Leukemia
Epidemiology, Review, 78-6025
In Utero Exposure, Rat, 78-6382
Parabiosis, 78-6373
Radiotherapy, Review, 78-6025
Lymphosarcoma
Parabiosis, 78-6373
Macrophages
Cytotoxic Activity, Review, 78-6080
Mammary Neoplasms, Experimental
Adenocarcinoma, 78-6050
Adenofibroma, 78-6050, 78-6342
Dose Fractionation, Review, 78-6050
In Utero Exposure, Rat, 78-6382
4,4'-Stilbene diol, α, α' -Diethyl-, 78-6342
Metals
Mutagenic Activity, 78-6134
Norharman
DNA Repair, 78-6155
Mutagenic Activity, 78-6155
Occupational Hazard
Dosimetry, Review, 78-6047
Epidemiology, Review, 78-6025, 78-6047
Ovarian Neoplasms
In Utero Exposure, Rat, 78-6382
Parathyroid Neoplasms
Adenoma, 78-6376, 78-6377, 78-6379
Neoplasms, Multiple Primary, 78-6377, 78-6379
Piperonyl Butoxide
Chromosome Aberrations, 78-6238
Mutagenic Activity, 78-6238
Pituitary Neoplasms
In Utero Exposure, Rat, 78-6382
Rectal Neoplasms
Adenocarcinoma, 78-6385
1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385
Respiratory Tract Neoplasms
Epidemiology, Review, 78-6049

Irradiation, Ionizing (cont'd)

- Salmonella typhimurium*
 - Genetic Duplication, 78-6359, 78-6360

Sarcoma

- Epidemiology, 78-6374

Sarcoma, Osteogenic

- Epidemiology, 78-6374
- Parabiosis, 78-6373

Sarcoma, Reticulum Cell

- Parabiosis, 78-6373

Skin Neoplasms

- In Utero Exposure, Rat, 78-6382

Smoking

- Co-carcinogenic Effect, Review, 78-6049
- Relative Risks, Review, 78-6047

4,4'-Stilbenediol, α,α' -Diethyl-

- Co-carcinogenic Effect, 78-6342

Thyroid Neoplasms

- Adenocarcinoma, 78-6378, 78-6523
- Adenocarcinoma, Papillary, 78-6363, 78-6377
- Adenoma, 78-6523
- Carcinoma, 78-6363, 78-6523
- Carcinoma, Papillary, 78-6376, 78-6378
- Case Report, 78-6363, 78-6378
- Epidemiology, 78-6376, 78-6379, 78-6523
- Hyperparathyroidism, 78-6379
- In Utero Exposure, Rat, 78-6382
- Neoplasm Metastasis, 78-6363, 78-6376
- Neoplasm Recurrence, Local, 78-6376
- Neoplasms, Multiple Primary, 78-6377, 78-6379

Uterine Neoplasms

- In Utero Exposure, Rat, 78-6382

Radioisotopes**Nickel**

- Body Burden, Mouse, 78-6142
- Metabolism, Mouse, 78-6142
- Tissue Distribution, 78-6142

Respiratory Tract Neoplasms

- Lead, 78-6049

Ruthenium

- Cell Transformation, Neoplastic, 78-6048

Smoking

- Polonium, 78-6049

Strontium**Breast Neoplasms**

- Epidemiology, 78-6564

Kidney Neoplasms

- Adenocarcinoma, 78-6564
- Epidemiology, 78-6564

Leukemia

- Epidemiology, 78-6564

Soft Tissue Neoplasms

- Epidemiology, 78-6564
- Radiotherapy, 78-6564

Thallium**Lung Neoplasms**

- Occupational Hazard, 78-6554

Receptors, Hormone**Coumestrol**

- Estradiol, 78-6350

Isoflavone, 7-Hydroxy-4'-methoxy-

- Estradiol, 78-6350

Isoflavone, 4',5,7-Trihydroxy-

- Estradiol, 78-6350

Liver Neoplasms**Receptors, Hormone (cont'd)**

- Contraceptives, Oral, 78-6355

- Estradiol, 78-6355

- Progesterone, 78-6355

Mammary Neoplasms, Experimental

- Estradiol, 78-6351

- Estriol, 78-6351

- Estriol, 16-Epi-, 78-6351

Zearalenol

- Estradiol, 78-6350

Zearalenone

- Estradiol, 78-6350

Rectal Neoplasms**Adenocarcinoma**

- Carcinoembryonic Antigen, 78-6596

- Radiation, Ionizing, 78-6385

- Transplantation, Heterologous, 78-6596

- Ultrastructural Study, 78-6596

Adenosine Cyclic 3',5' Monophosphate

- Carcinoembryonic Antigen, 78-6596

Carcinoembryonic Antigen

- Cells, Cultured, 78-6596

Dibutyl Cyclic AMP

- Carcinoembryonic Antigen, 78-6596

Radiation, Ionizing

- 1*H*-Imidazole-1-ethanol, α -(Methoxymethyl)-2-nitro-, 78-6385

Reserpine**Antigens**

- Hypersensitivity, Delayed, 78-6490

Cholanthrene, 3-Methyl-

- Co-carcinogenic Effect, Hamster, 78-6321

Fibrosarcoma

- Transplantation Immunology, 78-6490

Macrophages

- Transplantation Immunology, 78-6490

Mammary Neoplasms, Experimental

- Cholanthrene, 3-Methyl-, 78-6321

Respiratory System**Nicotine**

- Body Burden, Mouse, 78-6177

Respiratory Tract Neoplasms**Benzo(a)pyrene**

- Anesthesia, 78-6335

- Brevital, 78-6335

- Ethane, 1,1'-Oxybis-, 78-6335

- Ether, 2,2-Dichloro-1,1-difluoroethyl Methyl, 78-6335

- Iron Oxide, 78-6335

Carcinoma

- Occupational Hazard, Review, 78-6049

Carcinoma, Epidermoid

- Benzo(a)pyrene, 78-6335

- Occupational Hazard, Review, 78-6049

Chromate

- Epidemiology, 78-6131

- Occupational Hazard, 78-6131

Chromium

- Epidemiology, Japan, 78-6561

Ethane, 1,2-Dichloro-

- Carcinogenic Activity, Mouse, Rat, 78-6114

Ethnic Groups

- Epidemiology, Cameroon, 78-6568

Lead

- Radioisotopes, 78-6049

Respiratory Tract Neoplasms (cont'd)

- Occupational Hazard
 - Epidemiology, 78-6560
- Petroleum
 - Epidemiology, Review, 78-6018
- Radiation, Ionizing
 - Epidemiology, Review, 78-6049
- Smoking
 - Epidemiology, Review, 78-6049
- Truck Drivers
 - Epidemiology, 78-6560
- Uranium
 - Occupational Hazard, Review, 78-6049
- Vanadium
 - Occupational Hazard, 78-6561

Retinoblastoma

- Genetics
 - Epidemiology, Review, 78-6071
 - Immunity, Cellular, Review, 78-6071

Retinoic Acid

- Adenosine Cyclic 3',5' Monophosphate
 - Plasminogen Activators, 78-6338
- Plasminogen Activators
 - Fibroblasts, Chick Embryo, 78-6338
- RNA, Messenger
 - Plasminogen Activators, 78-6338
- Teratoid Tumor
 - Alkaline Phosphatase, 78-6339
 - Cell Differentiation, 78-6339
 - Collagen, 78-6339
 - Plasminogen Activators, 78-6339
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Plasminogen Activators, 78-6338

Retinol

- Carcinogen, Chemical
 - Carcinogenic Activity, Review, 78-6004
- Cells, Cultured
 - Contact Inhibition, 78-6592
 - Glycolipids, 78-6592
 - Glycoproteins, 78-6592
 - Surface Properties, 78-6592
- Colonic Neoplasms
 - Epidemiology, Review, 78-6039
- Growth Substances
 - Contact Inhibition, 78-6592
- Plasminogen Activators
 - Fibroblasts, Chick Embryo, 78-6338
- Ultraviolet Rays
 - Contact Inhibition, 78-6592

Retrorsine

- DNA Repair
 - Liver, Rat, 78-6216

Reverse Transcriptase

- Melanoma
 - Virus, RNA Tumor, 78-6438
- Nose Neoplasms
 - Adenocarcinoma, 78-6521
 - Cystadenoma, Papillary, 78-6521
- Virus, Avian Sarcoma
 - Glucosamine, 78-6394
- Virus, Bovine Leukemia
 - Isolation and Characterization, Review, 78-6088
- Virus, Feline Sarcoma
 - Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-, 78-6440
 - Aflatoxin B1, 78-6440

Reverse Transcriptase (cont'd)

- Benzo(a)pyrene, 78-6440

Rhabdomyosarcoma

- Acenaphthene, 5-Nitro-
 - Carcinogenic Potential, 78-6254
- Glycoproteins
 - Isolation and Characterization, Review, 78-6057
- Gold, ((1,2-Dicarboxyethyl)thio)-, Disodium Salt
 - Dose-Response Study, Rat, 78-6133

Rhodium Chloride

- Bacillus subtilis*
 - Mutagenic Activity, 78-6128

Riboflavine

- Azo Compounds
 - Ames Test, 78-6263
- Benzopurpurine 4B
 - Ames Test, 78-6263
- C.I. Acid Red 26, Disodium Salt
 - Ames Test, 78-6263
- C.I. Direct Red 28, Disodium Salt
 - Ames Test, 78-6263
- Chlorazol Violet N
 - Ames Test, 78-6263

RNA

- Virus, Herpes Simplex 1
 - DNA Replication, 78-6448
- Virus, Murine Leukemia
 - Globin, 78-6422
- Virus, Polyoma
 - DNA Replication, 78-6482
 - Nucleotides, 78-6482
 - Okazaki Fragments, 78-6482

RNA, Messenger

- Casein
 - Mammary Development, 78-6410
- Cell Transformation, Neoplastic
 - Chromatin, 78-6072
- Plasmacytoma
 - Cytoplasm, 78-6579
 - Isolation and Characterization, 78-6579
 - Nucleic Acid Precursors, 78-6579
 - Ultraviolet Rays, 78-6579
- Retinoic Acid
 - Plasminogen Activators, 78-6338
- Virus, Avian Sarcoma
 - DNA-RNA Hybridization, 78-6391
 - Isolation and Characterization, 78-6391
 - Nucleotide Sequence, 78-6391
- Virus, Friend Murine Leukemia
 - Cell Transformation, Neoplastic, 78-6580
 - DNA-RNA Hybridization, 78-6580
 - Isolation and Characterization, 78-6580
 - Poly A, 78-6580
- Virus, Polyoma
 - Isolation and Characterization, 78-6484
 - Poly A, 78-6484
 - Viral Proteins, 78-6484
- Virus, Rous Sarcoma
 - Antigens, Viral, 78-6401
- Virus, SV40
 - DNA, Viral, 78-6485
 - Endonucleases, 78-6485
 - R-Loop Structures, 78-6485

A Polymerase

- Virus, Rous Sarcoma
 - Isolation and Characterization, 78-6399
 - RNA Replication, 78-6399
 - Virus-Like Particles, 78-6399

A Replication

- Virus, Adeno 2
 - DNA-RNA Hybridization, 78-6463
- Virus, Adeno 12
 - DNA, Viral, 78-6469
- Virus, Rous Sarcoma
 - RNA Polymerase, 78-6399
- Virus, SV40
 - DNA, Viral, 78-6485

A, Transfer, Methyltransferases

- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 - Liver, Rat, 78-6265
- Aniline, *p*-(Phenylazo)-
 - Liver, Rat, 78-6265
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Liver, Rat, 78-6265
 - Lymphocytes, 78-6205
- Carbon Tetrachloride
 - Liver, Rat, 78-6265
- Cholanthrene, 3-Methyl-
 - Liver, Rat, 78-6265
- Diethylamine, *N*-Nitroso-
 - Liver, Rat, 78-6265
- Dimethylamine, *N*-Nitroso-
 - Liver, Rat, 78-6265
- Isothiocyanic Acid, 1-Naphthyl Ester
 - Liver, Rat, 78-6265
- Isothiocyanic Acid, 2-Naphthyl Ester
 - Liver, Rat, 78-6265

A, Viral

- Carcinoma
 - Virus, RD-114, 78-6441
- Lymphoma
 - Virus, Feline Leukemia, 78-6441
 - Virus, RD-114, 78-6441
- Melanoma
 - Virus, RNA Tumor, 78-6438
- Sarcoma
 - Virus, RD-114, 78-6441
- Virus, Abelson Murine Leukemia
 - Viral Proteins, 78-6420
- Virus, Adeno 2
 - DNA-RNA Hybridization, 78-6463
- Virus, Avian Leukosis
 - DNA-RNA Hybridization, 78-6387
 - Nucleotide Sequence, 78-6387
- Virus, Avian Sarcoma
 - DNA-RNA Hybridization, 78-6392
 - Glucosamine, 78-6394
- Virus, Feline Leukemia
 - Tissue Distribution, 78-6441
- Virus, Friend Spleen Focus-Forming
 - Virus-Like Particles, 78-6428
- Virus, Moloney Murine Leukemia
 - Ultrastructural Study, Heteroduplexes, 78-6423
- Virus, Murine Leukemia
 - Age Factors, 78-6422
 - Brain, Liver, Mouse, 78-6422
 - DNA-RNA Hybridization, 78-6422
 - Ultrastructural Study, Heteroduplexes, 78-6423
- Virus, Murine Mammary Tumor

RNA, Viral (cont'd)

- Age Factors, 78-6410
- Dexamethasone, 78-6411
- DNA-RNA Hybridization, 78-6410, 78-6411
- Horizontal Transmission, 78-6411
- Hormones, 78-6411
- Mammary Development, 78-6410
- Precancerous Conditions, 78-6411
- Vertical Transmission, 78-6411
- Virus, Polyoma
 - DNA-RNA Hybridization, 78-6483
 - Nucleotide Sequence, 78-6483
- Virus, RD-114
 - Embryonic Tissue, 78-6441
 - Tissue Distribution, 78-6441
- Virus, RNA Tumor
 - Virus-Like Particles, 78-6472
- Virus, Rous Sarcoma
 - DNA-RNA Hybridization, 78-6404
- Virus, Woolly Monkey
 - Viral Proteins, 78-6420

Ruthenium

- Cell Transformation, Neoplastic
 - Mathematical Model, 78-6048
- Radioisotopes
 - Cell Transformation, Neoplastic, 78-6048

Saccharin

- see 1,2-Benzisothiazolin-3-one, 1,1-Dioxide

Saccharomyces cerevisiae

- Acetic Acid, ((4-Chloro-*o*-tolyl)oxy)-
 - Mutagenic Activity, 78-6287
- Acetic Acid, (2,4-Dichlorophenoxy)-
 - Mutagenic Activity, 78-6287
- Acetic Acid, (2,4,5-Trichlorophenoxy)-
 - Mutagenic Activity, 78-6287
- Acrylamide, 2-(2-Furyl)-3-(5-nitro-2-furyl)-
 - Growth, Radiation-Sensitive Mutants, 78-6244
- Actinomycin D
 - Mutagenic Activity, 78-6304
- Chromic Acid, Lead Salt
 - Mutagenic Activity, 78-6304
- Cosmetics
 - Mutagenic Activity, 78-6150
- Hydroxylamine
 - Growth, Radiation-Sensitive Mutants, 78-6244
- Methanesulfonic Acid, Ethyl Ester
 - Growth, Radiation-Sensitive Mutants, 78-6244
- 2-Naphthylamine
 - Growth, Radiation-Sensitive Mutants, 78-6244
- Propane, 2,2'-Oxybis(2-chloro)-
 - Mutagenic Activity, 78-6118
- Quinoline, 4-Nitro-
 - Growth, Radiation-Sensitive Mutants, 78-6244
- Salicylic Acid
 - Mutagenic Activity, 78-6287
- Ultraviolet Rays
 - Caffeine, 78-6121
- Salicylanilide, 3,3',5,5',6-Pentachloro-2'-hydroxy-**
 - Chromosome Abnormalities
 - Allium cepa*, 78-6255
 - Mitosis, 78-6255
 - Mitosis
 - Allium cepa*, 78-6255
- Salicylic Acid**
 - Saccharomyces cerevisiae*

Salicylic Acid (cont'd)

Mutagenic Activity, 78-6287

Salivary Gland Neoplasms

4-Biphenylamine, 3,2'-Dimethyl-

Carcinogenic Activity, Rat, 78-6297

Salmonella typhimurium

Acridinium, 3,6-Diamino-10-methyl-, Chloride

Mutagenic Activity, 78-6159

Chromate

Mutagenic Activity, 78-6129

4-Cyclohexene-1,2-dicarboximide, *N*-

-(Trichloromethyl)thio-

Mutagenic Activity, 78-6159

Formic Acid, Ethylidenemethylhydrazide

Bacteriolysis, Mouse, 78-6181

ICR 372

Genetic Duplication, 78-6359, 78-6360

Imidazo(1,2-*a*)pyrimidine

Mutagenic Activity, Derivatives, 78-6236

Maleic Acid

Genetic Duplication, 78-6359

Methanesulfonic Acid, Ethyl Ester

Genetic Duplication, 78-6359, 78-6360

Nitrous Acid

Genetic Duplication, 78-6359

Quinacrine

Mutagenic Activity, 78-6159

Radiation, Ionizing

Genetic Duplication, 78-6359, 78-6360

Sulfuric Acid, Diethyl Ester

Mutagenic Activity, 78-6159

Ultraviolet Rays

Genetic Duplication, 78-6359, 78-6360

Sarcoma

Acetic Acid, (2,4-Dichlorophenoxy)-

Occupational Hazard, 78-6288

Acetic Acid, (2,4,5-Trichlorophenoxy)-

Occupational Hazard, 78-6288

Antigens, Neoplasm

T-Lymphocytes, 78-6502

Cholanthrene, 3-Methyl-

Dose-Response Study, 78-6321

Dibenzo-*p*-dioxin, 2,3,7,8-Tetrachloro-

Acetic Acid, (2,4,5-Trichlorophenoxy)-, 78-6288

Head and Neck Neoplasms

Chromium, 78-6561

Hypertension

Neoplasm Transplantation, 78-6332

Iron

Intramuscular Injection, 78-6565

Lung

Neoplasm Metastasis, Review, 78-6060

T-Lymphocytes

Immunity, Passive, 78-6502

Killer Cells, 78-6502

Lymph Nodes, 78-6502

Lymphocytotoxicity, 78-6502

Occupational Hazard

Epidemiology, 78-6288

Radiation, Ionizing

Epidemiology, 78-6374

Soft Tissue Neoplasms

Epidemiology, 78-6565

Epidemiology, Cat, 78-6511

Histological Study, Cat, 78-6511

Virus, Moloney Murine Sarcoma

Sarcoma (cont'd)

Lymph Nodes, 78-6502

T-Lymphocytes, 78-6502

Lymphocytotoxicity, 78-6502

Virus, Murine Sarcoma

Peptides, 78-6481

Virus, RD-114

RNA, Viral, 78-6441

Virus, SV40

Antigen-Antibody Reactions, 78-6085

Histocompatibility Antigens, 78-6085

Sarcoma, Mast Cell

T-Lymphocytes

Lymphocytotoxicity, 78-6495

Sarcoma, Osteogenic

Glycoproteins

Isolation and Characterization, Review, 78-6057

Radiation, Ionizing

Epidemiology, 78-6374

Parabiosis, 78-6373

Virus, SV40

Antigen-Antibody Reactions, 78-6085

Histocompatibility Antigens, 78-6085

Sarcoma, Reticulum Cell

Chromium

Occupational Hazard, 78-6561

Epidemiology

Australia, 78-6562

Ethnic Groups

Epidemiology, Cameroon, 78-6568

Genetics

Fish, Review, 78-6059

Liver Cirrhosis

Case Report, 78-6513

Nickel

Industrial Dusts, 78-6143

Oncogenic Viruses

Epidemiology, Review, 78-6031

Radiation, Ionizing

Parabiosis, 78-6373

Thyroid Neoplasms

Epidemiology, 78-6523

Virus, SV40

Antigen-Antibody Reactions, 78-6085

Histocompatibility Antigens, 78-6085

Schistosoma japonicum

Intestinal Neoplasms

Adenocarcinoma, 78-6535

Epidemiology, 78-6535

Schistosomiasis

Bladder Neoplasms

Epidemiology, 78-6567

Schistosomicides

Phenol, (1,1-Dimethylethyl)-4-methoxy-

Mutagenic Activity, 78-6328

Selenium

Mammary Neoplasms, Experimental

Adenocarcinoma, 78-6146

Antineoplastic Activity, 78-6146

Semicarbazide

Sulfurous Acid, Monosodium Salt

Mutagenic Activity, 78-6160

, Diazoacetate (Ester)
 Choline
 Organ Specificity, 78-6199
 DNA
 Alkaline Sucrose Gradients, 78-6200
 Pancreas, Liver, Rat, 78-6200
 DNA Replication
 Pancreas, Rat, 78-6212
 Hepatoma
 Choline, 78-6199
 Neoplasms, Multiple Primary, 78-6199
 Pancreatic Neoplasms
 Choline, 78-6199
 Precancerous Conditions, 78-6199
 Pituitary
 Brain Neoplasms
 Benz(a)anthracene, 7,12-Dimethyl-, 78-6318
 Chromosomes
 Disgerminoma
 Cell Differentiation, 78-6506
 Epithelial Neoplasms
 Mutagenic Activity, 78-6122
 Estrogens
 Ethnic Groups
 Menopause, 78-6544
 Plasma/Urinary Levels, 78-6544
 Fox
 Ames Test
 Mutagenic Activity, 78-6163
 Glutathione
 Liver Neoplasms
 Gluconic Acid, Iron Salt, 78-6193
 Precancerous Conditions, 78-6193
 8-Quinolinol, 78-6193
 Lung Neoplasms
 Epidemiology, 78-6547
 Histological Study, 78-6547
 Marfan's Syndrome
 Leukemia, Myeloblastic
 Cyclophosphamide, 78-6197
 Diseases
 Coal Tar
 Aryl Hydrocarbon Hydroxylases, 78-6336
 Neoplasms
 Benz(a)anthracene, 7,12-Dimethyl-
 Cortisol, 78-6307
 Cortisone, 78-6307
 Croton Oil, 78-6307
 Dexamethasone, 78-6307
 Benzo(a)pyrene
 Chlorine, 78-6330
 2,2'-Biphenylamine, 3,2'-Dimethyl-
 Carcinogenic Activity, Rat, 78-6297
 Carbamic Acid, Ethyl Ester
 Carcinogenic Activity, Mouse, 78-6185
 Carbamic Acid, Vinyl Ester
 Carcinogenic Activity, Mouse, 78-6185
 Carcinoma
 Virus, Papilloma, 78-6091
 Caribbean Islands
 Epidemiology, 78-6570
 Coal Tar

Skin Neoplasms (cont'd)
 Occupational Hazard, Review, 78-6017
 Dexamethasone
 Protease Inhibitors, 78-6307
 Ethnic Groups
 Epidemiology, Cameroon, 78-6568
 Melanoma
 Ultraviolet Rays, 78-6046
 Papilloma
 Carbamic Acid, Ethyl Ester, 78-6185
 Carbamic Acid, Vinyl Ester, 78-6185
 Petroleum
 Carcinogenic Activity, Review, 78-6002
 Polycyclic Hydrocarbons
 Automobile Exhaust, 78-6300
 Radiation, Ionizing
 In Utero Exposure, Rat, 78-6382
 Ultraviolet Rays
 Precancerous Conditions, Review, 78-6046
 Virus, Papilloma
 Carcinogenic Potential, Review, 78-6090, 78-6091
 Immunosuppression, Review, 78-6090
 Mastomys natalensis, 78-6477
 Virus, Polyoma
 Carcinogenic Activity, Review, 78-6092
 Immunosuppression, Review, 78-6092
 Warts
 Virus, Papilloma, 78-6090, 78-6091

Smoke Condensate
 Acetophenone
 Isolation and Characterization, 78-6171
 Ames Test
 Carcinogenic Potential, 78-6172
 Benzene, 4,5-Methylenetris-
 Isolation and Characterization, 78-6337
 Benzo(b)perylene
 Isolation and Characterization, 78-6337
 Carcinogenic Potential
 Sebaceous Gland Suppression Test, 78-6172
 Chromosome Abnormalities
 Ultrastructural Study, Plants, 78-6170
 Cyclopenta(cd)pyrene
 Isolation and Characterization, 78-6337
 Hyperplasia
 Epidermis, Mouse, 78-6172
 1,3-Indandione
 Isolation and Characterization, 78-6171
 Tumor-Inhibiting Fraction, 78-6171
 Polycyclic Hydrocarbons
 Quantitation Method, Review, 78-6010
 Pyrene, 3,4-Dimethylene
 Isolation and Characterization, 78-6337
 Pyrene, 3,4-Trimethylene
 Isolation and Characterization, 78-6337
 Tumor-Inhibiting Fraction
 Isolation and Characterization, 78-6171

Smoking
 Asbestos
 Co-carcinogenic Effect, Review, 78-6002, 78-6032
 Benzoic Acid, 2-Amino-3-hydroxy-
 Metabolism, 78-6567
 Bladder Neoplasms
 Epidemiology, 78-6567
 Tryptophan, 78-6567
 Carcinoma, Bronchogenic
 Aryl Hydrocarbon Hydroxylases, 78-6322

Smoking (cont'd)

- Esophageal Neoplasms
 - Carcinoma, Epidermoid, 78-6549
- Head and Neck Neoplasms
 - Carcinoma, Epidermoid, 78-6549
- Hemoglobins
 - Hematopoietic Effects, 78-6174
- Leukocyte Count
 - Hematopoietic Effects, 78-6174
- Lung Neoplasms
 - Epidemiology, Review, 78-6031
 - Uranium, 78-6554
- Lymphocytes
 - Aryl Hydrocarbon Hydroxylases, 78-6174
- Mouth Neoplasms
 - Carcinoma, Epidermoid, 78-6375
- Nasopharyngeal Neoplasms
 - Epidemiology, 78-6552
- Neoplasms
 - Epidemiology, Review, 78-6032
- Polonium
 - Radioisotopes, 78-6049
- Radiation, Ionizing
 - Co-carcinogenic Effect, Review, 78-6049
 - Relative Risks, Review, 78-6047
- Respiratory Tract Neoplasms
 - Epidemiology, Review, 78-6049
- Tryptophan
 - Carcinogenic Metabolite, 78-6567
- Uranium
 - Co-carcinogenic Effect, Review, 78-6049

Sodium Bisulfite

- see Sulfurous Acid, Monosodium Salt

Sodium Chloride

- Metrifonate
 - Carcinogenic Potential, Rat, 78-6145

Sodium Erythorbate

- Dose-Response Study
- Mutagenic Activity, Rat, 78-6138

Soft Tissue Neoplasms

- Ethnic Groups
 - Epidemiology, Cameroon, 78-6568
- Radium
 - Epidemiology, 78-6564
 - Radiotherapy, 78-6564
- Sarcoma
 - Epidemiology, 78-6565
 - Epidemiology, Cat, 78-6511
 - Histological Study, Cat, 78-6511

Somatotropin

- Uterine Neoplasms
 - Leiomyoma, 78-6541
 - Menstruation, 78-6541

Spermatogenesis

- Caffeine
 - Nitrous Acid, Sodium Salt, 78-6123
- Theobromine
 - Nitrous Acid, Sodium Salt, 78-6123
- Theophylline
 - Nitrous Acid, Sodium Salt, 78-6123

Spermidine

- Ames Test
- Nitrosation, 78-6204
- Ascorbic Acid

Spermidine (cont'd)

- Nitrosation, 78-6204
- Nitrous Acid, Sodium Salt
- Ames Test, 78-6204
- Mutagenic Activity, 78-6204
- Thiocyanic Acid
- Nitrosation, 78-6204

Spherocytosis, Hereditary

- Cells, Cultured
- Mutation, 78-6164

Spices

- Drosophila melanogaster*
 - Mutagenic Activity, 78-6122
- Sex Chromosomes
 - Mutagenic Activity, 78-6122
- Theophylline
 - Mutagenic Activity, 78-6122

Spinal Neoplasms

- Carcinoma, Epidermoid
 - Meningomyelocoele, 78-6519
- Meningomyelocoele
 - Case Report, 78-6519

Spirolactone

- Liver Neoplasms
 - Carcinogenic Potential, Review, 78-6016
- Mammary Neoplasms, Experimental
 - Carcinogenic Potential, Review, 78-6016
- Species Difference
 - Histological Study, 78-6016
- Testicular Neoplasms
 - Carcinogenic Potential, Review, 78-6016
- Thyroid Neoplasms
 - Carcinogenic Potential, Review, 78-6016

Starch

- Aflatoxin M1
 - Hepatotoxicity, 78-6125
 - Metabolism, Rat, 78-6125
- Macrophages
 - Cytostatic Activity, 78-6435

Steroids

- Dietary Fats
 - Metabolism, Review, 78-6036

4,4'-Stilbenediol, α,α' -Diethyl-

- Gynecologic Neoplasms
 - Adenoma, 78-6027
- Immunosuppression
 - Carcinogenic Activity, Review, 78-6051
- Liver Neoplasms
 - Hyperplasia, 78-6355
- Mammary Neoplasms, Experimental
 - Adenocarcinoma, 78-6342
 - Carcinogenic Potential, 78-6315
 - Radiation, Ionizing, 78-6342
 - Virus, Murine Mammary Tumor, 78-6315
- Pituitary Neoplasms
 - Carcinogenic Activity, Rat, 78-6342
- Radiation, Ionizing
 - Co-carcinogenic Effect, 78-6342

Stomach Neoplasms

- Adenocarcinoma
 - Agammaglobulinemia, 78-6534
 - Anemia, Pernicious, 78-6534
 - Gastritis, 78-6534

nach Neoplasms (cont'd)

- Immunologic Deficiency Syndromes, 78-6534
- Plants, Edible, 78-6210
- Cadmium
 - Food Contamination, 78-6148
- Carcinoma
 - Gastrectomy, 78-6576
 - Genetics, 78-6575
- Carcinoma, Epidermoid
 - Piperazine, 1,4-Dinitroso-, 78-6241
 - Plants, Edible, 78-6210
- Caribbean Islands
 - Epidemiology, 78-6570
- Enterochromaffin Cells
 - Ultrastructural Study, 78-6533
- Epidemiology
 - Mathematical Model, 78-6555
- Ethane, 1,2-Dichloro-
 - Carcinogenic Activity, Mouse, Rat, 78-6114
- Ethnic Groups
 - Epidemiology, Hawaii, 78-6546
- Gastrectomy
 - Epidemiology, 78-6576
 - Precancerous Conditions, 78-6576
- Gastritis
 - Achlorhydria, 78-6533
 - Case Report, 78-6533
- Genetics
 - Epidemiology, Finland, 78-6575
- Immunologic Deficiency Syndromes
 - Case Report, 78-6534
- Manganese
 - Food Contamination, 78-6148
- Nitric Acid, Potassium Salt
 - Epidemiology, Chile, 78-6574
 - Fertilizers, 78-6574
- Nitric Acid, Sodium Salt
 - Epidemiology, Chile, 78-6574
 - Fertilizers, 78-6574
- Nitrosamines
 - Potato Sap, 78-6210
- Papilloma
 - Piperazine, 1,4-Dinitroso-, 78-6241
 - Plants, Edible, 78-6210
- Petroleum
 - Carcinogenic Activity, Review, 78-6002
- Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-
 - Nitrous Acid, Sodium Salt, 78-6241
- Piperidine, 4-Chloro-1-nitroso-
 - Carcinogenic Activity, Rat, 78-6240
- Polyps
 - Gastritis, 78-6533
- Tumor Latency
 - Mathematical Model, 78-6555
- Vitamin B12
 - Malabsorption Syndromes, 78-6534
- Zinc
 - Epidemiology, Japan, 78-6148
 - Food Contamination, 78-6148

ptozotocin

- DNA Replication
 - Pancreas, Rat, 78-6212
- Islet Cell Tumor
 - Glucose, 78-6207
 - Hypoglycemia, 78-6207
 - Insulin, 78-6207
- Kidney Neoplasms

Streptozotocin (cont'd)

- Nicotinamide, 78-6208
- Leukemia, Myeloblastic
 - Colonic Neoplasms, 78-6509
- Pancreatic Neoplasms
 - Adenoma, 78-6208
 - Dose-Response Study, Rat, 78-6208
 - Islet Cell Tumor, 78-6207, 78-6208
 - Nicotinamide, 78-6208
 - Picolinamide, 78-6208

Stress

- Neoplasms
 - Hormones, Review, 78-6058
 - Immune Response, Review, 78-6058

Strontium

- Chlorophyll
 - Mutagenic Activity, 78-6134

Strontium Radioisotopes

- Mammæ
 - Metabolism, Mouse, 78-6142
- Placenta
 - Metabolism, Mouse, 78-6142
- Tissue Distribution
 - Body Burden, Mouse, 78-6142

Styrene

- Drosophila melanogaster*
 - Mutagenic Activity, 78-6253
- Leukemia
 - Occupational Hazard, Review, 78-6014
- Mandelic Acid
 - Metabolism, 78-6252
- Occupational Hazard
 - Chromosome Aberrations, 78-6252

Styrene Oxide

- see Benzene, (Epoxyethyl)-

Sucrose

- Aflatoxin M1
 - Metabolism, Rat, 78-6125

Sudan I

- see 2-Naphthol, 1-(Phenylazo)-

Sudan II

- see 2-Naphthol, 1-(2,4-Xylylazo)-

Sudan III

- see 2-Naphthalenol, 1-((4-(Phenylazo)phenyl)azo)-

Sudan IV

- see 2-Naphthol, 1-((4-(*o*-Tolylazo)-*o*-tolyl)azo)-

Sulfamethazine

- see Sulfanilamide, *N*'-(4,6-Dimethyl-2-pyrimidinyl)-

Sulfanilamide, *N*'-(4,6-Dimethyl-2-pyrimidinyl)-

- Acetylation Kinetics
 - Species Difference, 78-6264
- Acetyltransferases
 - Acetylation Kinetics, 78-6264

Sulfate

- Xanthine, 3-Hydroxy-
 - Metabolism, Liver, 78-6249

Sulfuric Acid, Aluminum Potassium Salt, Dodecahydrate

- see also Alum
- Virus, Feline Leukemia
 - Viral Vaccines, 78-6442

Sulfuric Acid, Diethyl Ester

- Escherichia coli*
 - Mutagenic Activity, 78-6159
- Mutagenic Activity
 - Galactose Resistance, 78-6159
- Salmonella typhimurium*
 - Mutagenic Activity, 78-6159

Sulfuric Acid, Dimethyl Ester

- Guanosine
 - RNA, Methylation, 78-6228
- Yeasts
 - RNA, Methylation, 78-6228

Sulfurous Acid, Disodium Salt

- Chromic Acid, Zinc Salt
 - Ames Test, 78-6130

Sulfurous Acid, Monosodium Salt

- Bacteriophages
 - Cytosine, 78-6160
 - Mutagenic Activity, 78-6160
 - Transaminated Residues, 78-6160
- Hydroxylamine
 - Mutagenic Activity, 78-6160
- Hydroxylamine, *O*-Methyl-
 - Mutagenic Activity, 78-6160
- Semicarbazide
 - Mutagenic Activity, 78-6160

Synfuel A-3

- Ames Test
 - Mutagenic Activity, 78-6102
- Enzyme Activation
 - Mutagenic Activity, 78-6102
- Microsomes, Liver
 - Ames Test, 78-6102

Synovioma

- Head and Neck Neoplasms
 - Case Report, 78-6524
 - Histochemical Study, 78-6524
 - Ultrastructural Study, 78-6524

Tellurium

- Bacillus subtilis*
 - Mutagenic Activity, 78-6128

Teratogens

- Occupational Hazard
 - Women, Review, 78-6022

Teratoid Tumor

- Dibutyl Cyclic AMP
 - Plasminogen Activators, 78-6339
- Hybrid Cells
 - Gene Control Mechanisms, 78-6072
- Retinoic Acid
 - Alkaline Phosphatase, 78-6339
 - Cell Differentiation, 78-6339
 - Collagen, 78-6339
 - Plasminogen Activators, 78-6339
- Testicular Neoplasms
 - Metals, 78-6563

Testicular Neoplasms

- Disgerminoma
 - Metals, 78-6563
- Dog
 - Epidemiology, California, 78-6548
- Metals
 - Epidemiology, 78-6563

Testicular Neoplasms (cont'd)

- Occupational Hazard, 78-6563
- Neoplasms, Embryonal and Mixed
 - Metals, 78-6563
- Spironolactone
 - Carcinogenic Potential, Review, 78-6016
- Teratoid Tumor
 - Metals, 78-6563
- Virus, Avian Leukosis
 - Virus, Helper, 78-6389

Testis

- Turner's Syndrome
 - Cell Differentiation, 78-6506

Testosterone

- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6343
 - Castration, 78-6343
 - Metabolism, 78-6343
 - 1-Piperazineethanol, 4-(3-(2-Chlorophenothiazin-10-yl)propyl)-, 78-6343

12-*O*-Tetradecanoylphorbol-13-acetate

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - DNA Repair, 78-6308
 - Adenosine Cyclic 3',5' Monophosphate
 - Plasminogen Activators, 78-6338
 - Cell Survival
 - Ganglia, Chick Embryo, 78-6309
 - Cholanthrene, 3-Methyl-
 - Co-carcinogenic Activity, Mouse Embryo Cells 78-6283
 - DNA Repair
 - 6,7-Epoxides, 78-6325
 - Escherichia coli*
 - DNA Repair, 78-6308, 78-6325
 - Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Mutagenic Activity, 78-6295
 - HeLa Cells
 - Epidermal Growth Factor, Binding, 78-6306
 - Methanol, (Methyl-*ONN*-azoxy)-, Acetate (Ester)
 - Mutagenic Activity, 78-6295
 - Nerve Growth Factor
 - Cell Survival, 78-6309
 - Ganglia, Chick Embryo, 78-6309
 - Peptide Hydrolases
 - Lysine, 78-6307
 - Skin, Mouse, 78-6307
 - Plasminogen Activators
 - Antigenic Determinants, 78-6310
 - Cell Transformation, Neoplastic, 78-6310
 - Fibroblasts, Chick Embryo, 78-6338
 - Lung, Hamster, 78-6310
 - Retinoic Acid
 - Plasminogen Activators, 78-6338
 - Ultraviolet Rays
 - Co-carcinogenic Activity, Mouse Embryo Cells 78-6283
 - Mutagenic Activity, 78-6295
 - Urea, Methyl Nitroso-
 - DNA Repair, 78-6308
 - Virus, Rous Sarcoma
 - Plasminogen Activators, 78-6395
- Theobromine**
- Atrophy
 - Testis, Rat, 78-6123
 - Cholesterol

romine (cont'd)
 Plasma Levels, 78-6123
 Hyperplasia
 Leydig Cells, 78-6123
 Nitrous Acid, Sodium Salt
 Spermatogenesis, 78-6123
 hylline
 Atrophy
 Testis, Rat, 78-6123
 Caffeine
 Mutagenic Activity, 78-6122
 Erythroleukemia
 Adenosine Cyclic 3',5' Monophosphate, 78-6426
 Lymphocytes
 Aryl Hydrocarbon Hydroxylases, 78-6292
 Nitrous Acid, Sodium Salt
 Spermatogenesis, 78-6123
 Plasma
 Half-Life, 78-6292
 pices
 Mutagenic Activity, 78-6122
 vanic Acid
 spermidine
 Nitrosation, 78-6204
 an
 ee Endosulfan
 ene, 5-(*p*-Acetamidophenyl)-2-amino-, Hydrochloride
 Ames Test
 Mutagenic Activity, 78-6260
 Cells, Cultured
 Cell Transformation, Neoplastic, 78-6260
 ene, 2-Amino-5-phenyl-, Hydrochloride
 Ames Test
 Mutagenic Activity, 78-6260
 Cells, Cultured
 Cell Transformation, Neoplastic, 78-6260
 osphamide
 ee Phosphide Sulfide, Tris(1-aziridinyl)-
 dine, Azido-
 Virus, Friend Spleen Focus-Forming
 A-Type Particles, 78-6429
 Virus Replication, 78-6429
 mu
 Acetic Acid, (Ethylenebis(oxyethylenenitrilo))tetra-
 Rosette Formation, AKR Mouse, 78-6503
 Complement
 Rosette Formation, AKR Mouse, 78-6503
 Virus, Abelson Murine Leukemia
 Macrophages, 78-6419
 Virus, Murine Leukemia
 Complement, 78-6503
 us Gland
 Leukemia
 Phagocytosis, 78-6526
 us Neoplasms
 Lymphosarcoma
 Virus, Murine Leukemia, 78-6421
 Virus, Murine Leukemia
 Mouse, Hybrid, 78-6421
 id Neoplasms
 Adenocarcinoma
 Epidemiology, 78-6523

Thyroid Neoplasms (cont'd)
 Radiation, Ionizing, 78-6378, 78-6523
 Radiotherapy, 78-6380
 Adenocarcinoma, Papillary
 Radiation, Ionizing, 78-6363, 78-6377
 Adenoma
 Imidazole-2-thiol, 1-Methyl-, 78-6248
 Radiation, Ionizing, 78-6523
 Calcitonin
 Immune Serums, 78-6594
 Radioimmunoassay, 78-6594
 Carcinoma
 Epidemiology, 78-6523
 Genetics, 78-6522
 Neoplasm Metastasis, 78-6363, 78-6379, 78-6522
 Radiation, Ionizing, 78-6363, 78-6523
 Ultrastructural Study, 78-6505
 Carcinoma, Papillary
 Radiation, Ionizing, 78-6376, 78-6378
 Epidemiology
 Inbred Mice, 78-6595
 Genetics
 Calcitonin, 78-6522
 Case Report, 78-6522
 Fish, Review, 78-6059
 Goiter, Exophthalmic
 Epidemiology, 78-6523
 Hyperparathyroidism
 Radiation, Ionizing, 78-6379
 Hyperplasia
 Imidazole-2-thiol, 1-Methyl-, 78-6248
 Hyperthyroidism
 Imidazole-2-thiol, 1-Methyl-, 78-6248
 Imidazole-2-thiol, 1-Methyl-
 Histological Study, Rat, 78-6248
 Precancerous Conditions, 78-6248
 Iodine Radioisotopes
 Epidemiology, 78-6376
 Lymphoma
 Immunoglobulins, Surface, 78-6505
 Ultrastructural Study, 78-6505
 Neoplasms, Multiple Primary
 Radiation, Ionizing, 78-6377, 78-6379
 Radiation, Ionizing
 Case Report, 78-6363, 78-6378
 Epidemiology, 78-6376, 78-6379, 78-6523
 In Utero Exposure, Rat, 78-6382
 Neoplasm Metastasis, 78-6363, 78-6376
 Neoplasm Recurrence, Local, 78-6376
 Radiotherapy
 Epidemiology, 78-6380
 Sarcoma, Reticulum Cell
 Epidemiology, 78-6523
 Spironolactone
 Carcinogenic Potential, Review, 78-6016
 Tobacco
 Nicotine, 1'-Nitroso-1'-demethyl-
 E and Z Isomers, 78-6176
p-Toluamide, *N*-Isopropyl- α -(2-methylhydrazino)-
 Leukemia, Myeloblastic
 Drug Therapy, 78-6219
 Hodgkin's Disease, 78-6196
 Toluene-2,4-diamine
 Carcinogenic Metabolite
 Quantitation Method, 78-6190

m-Toluidine, N,N-Dimethyl-4-(phenylazo)-
Hepatocytes
Cytotoxicity, Fibroblasts, 78-6267

p-Toluidine
Lethal Dose 50
Toxicity, Rat, 78-6266

p-Toluidine, α,α,α -Trifluoro-2,6-dinitro-N,N-dipropyl-
Drosophila melanogaster
Chromosomes, 78-6268
Mutagenic Activity, 78-6268

p-Tolyhydrazine
see Hydrazine, (4-Methylphenyl)-

Toxins
Dietary Fiber
Metabolism, Review, 78-6040

Tranquilizing Agents
Breast Neoplasms
Epidemiology, 78-6243

Transfection
see Transformation, Genetic

Transferases
Androst-4-ene-3,17-dione
Phenol, (1,1-Dimethylethyl)-4-methoxy-, 78-6333
Barbituric Acid, 5-Ethyl-5-phenyl-
Metabolism, Lung, 78-6331
Benzene, 1-Chloro-2,4-dinitro-
Metabolism, Lung, 78-6331
Phenol, (1,1-Dimethylethyl)-4-methoxy-, 78-6333
Benzene, 1-(Chloromethyl)-4-nitro-
Phenol, (1,1-Dimethylethyl)-4-methoxy-, 78-6333
Benzene, 1,2-Dichloro-4-nitro-
Phenol, (1,1-Dimethylethyl)-4-methoxy-, 78-6333
Benzo(a)pyrene 4,5-Oxide
Metabolism, Lung, 78-6331
Dibenzo(p)dioxin, 1,2,7,8-Tetrachloro-
Metabolism, Lung, 78-6331
Ethane, 1,2-Dibromo-
Binding, Liver, Kidney, Rat, 78-6115
Enzymatic Activity, 78-6115
Phenol, (1,1-Dimethylethyl)-4-methoxy-
Liver, Mouse, Rat, 78-6333
Pregnenolone 16 α -Carbonitrile
Metabolism, Lung, 78-6331

Transformation, Genetic
Virus, Rous Sarcoma
DNA, Viral, 78-6393, 78-6403
Temperature Sensitive Mutants, 78-6403

Transplantation, Heterologous
Rectal Neoplasms
Adenocarcinoma, 78-6596

Transplantation, Homologous
Acetamide, N-Fluorenyl-2-yl-
Hepatocyte, Rat, 78-6492
Diethylamine, N-Nitroso-
Hepatocyte, Rat, 78-6492
Graft Rejection
Lymphoid Tissue, 78-6491
Homologous Wasting Disease
Lymphoid Tissue, 78-6491
Liver Neoplasms
Precancerous Conditions, 78-6492
Mammary Neoplasms, Experimental

Transplantation, Homologous (cont'd)
Precancerous Conditions, 78-6527
Neoplasms
Kidney, Heart, Review, 78-6076

Transplantation Immunology
Carcinoma
Aniline, p-(Phenylazo)-, 78-6278
Benzenamine, 2-Methyl-4-((2-methylphenyl)azo)-
78-6278
Immunosuppression, 78-6278
Carrageen
Macrophages, 78-6490
Fibrosarcoma
Carrageen, 78-6490
Cholanthrene, 3-Methyl-, 78-6490
2-Imidazolidinone, 1-(5-Nitro-2-thiazolyl)-, 78-6490
Macrophages, 78-6490
Radiation, Ionizing, 78-6490
Reserpine, 78-6490
Reserpine
Macrophages, 78-6490
Virus, Marek's Disease Herpes
B-Lymphocytes, 78-6406
T-Lymphocytes, 78-6406

Trauma
see Wounds and Injuries

Triazene, 3-Acetyl-3-methyl-1-(2,4,6-tribromophenyl)-
Ames Test
Mutagenic Activity, 78-6223

Triazene, 3,3-Dimethyl-1-(p-chlorophenyl)-
Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
Ames Test, 78-6223
Microsomes, Liver
Ames Test, 78-6223
Mutagenic Activity
N-Acetylated Analogs, 78-6223

1-Triazene, 3,3-Dimethyl-1-phenyl-
Chromosome Aberrations
Lymphocytes, Rat, 78-6214

Triazene, 3,3-Dimethyl-1-phenyl-
Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
Ames Test, 78-6223
Carcinogenic Metabolite
Aniline Derivatives, 78-6266

DNA
Transplacental Alkylation, Rat, 78-6221
Guanine, 7-Methyl-
Liver, Brain, Rat, 78-6221
Microsomes, Liver
Ames Test, 78-6223
Mutagenic Activity
N-Acetylated Analogs, 78-6223
Purine, 2-Amino-6-methoxy-
Liver, Brain, Rat, 78-6221

Triazene, 3,3-Dimethyl-1-(m-pyridyl)-
Chromosome Aberrations
Lymphocytes, Rat, 78-6214

α -Triazine-3,5(2H,4H)-dione, 2- β -D-Ribofuranosyl-
Immunosuppression
Carcinogenic Activity, Review, 78-6051

s-Triazine, 2,4,6-Tris(1-aziridinyl)-
Chromosome Aberrations
Embryo, Mouse, 78-6222

azine, 2,4,6-Tris(1-aziridinyl)- (cont'd)

- Mutagenic Activity, 78-6222
- Colchicine
 - Chromosome Aberrations, 78-6222
- Leukemia, Myeloblastic
 - Hodgkin's Disease, 78-6196
- Poecilia reticulata*
 - Fetal Death, 78-6220
 - Mutagenic Activity, 78-6220

iquone

- see *p*-Benzoquinone, 2,3,5-Tris(1-aziridinyl)-

lorfon

- see Metrifonate

lycerides

- Metabolic Diseases
 - Enzymatic Activity, 78-6589

hanolamine

- see Ethanol, 2,2',2''-Nitrilotris-

nylenemelamine

- see *s*-Triazine, 2,4,6-Tris(1-aziridinyl)-

ralin

- see *p*-Toluidine, α,α,α -Trifluoro-2,6-dinitro-*N,N*-dipropy-

rine

- see Piperazine, 1,4-Bis(1-formamido-2,2,2-trichloroethyl)-

ethylamine, *N*-Oxide

- Hexamethylenetetramine
 - Nitrosation, 78-6225

ophan

- Ames Test
 - Pyrolysis Product, 78-6201
- Bladder Neoplasms
 - Smoking, 78-6567
- Fibrosarcoma
 - Methacrylic Acid, Ethyl Ester, 78-6372
 - Methacrylic Acid, Methyl Ester, 78-6372
- Nitrous Acid, Sodium Salt
 - Nitrosation, Meat, 78-6203
- Smoking
 - Carcinogenic Metabolite, 78-6567
- Tryptophan, Nitroso-
 - Nitrous Acid, Sodium Salt, 78-6203
- ophan, Nitroso-
 - Nitrous Acid, Sodium Salt
 - Hydrogen-Ion Concentration, 78-6203
- Tryptophan
 - Nitrous Acid, Sodium Salt, 78-6203

in

- Benzimidazolecarbamic Acid, Methyl Ester
 - Dose-Response Study, Mouse, 78-6303
- Imidazolidinethione, *N*-Nitroso-
 - Dose-Response Study, Mouse, 78-6303

r's Syndrome

- Cell Differentiation
 - Case Report, 78-6506
 - Testis, 78-6506
- Chromosomes
 - Genetics, Review, 78-6069

iolet Rays

- Ames Test
 - Genetic Duplication, 78-6359, 78-6360
- Bacteria

Ultraviolet Rays (cont'd)

- DNA Repair, Review, 78-6052
- Benzamide, 3,5-Dinitro-
 - Endonucleases, 78-6285
- Benzamide, 2-Methyl-3,5-dinitro-
 - Endonucleases, 78-6285
- Caffeine
 - DNA Repair, 78-6121
 - Saccharomyces cerevisiae*, 78-6121
- Carcinogen, Chemical
 - Spectrum Analysis, 78-6154
- Chloramphenicol
 - DNA Repair, Review, 78-6052
- Cholestan-3 β -ol, 5 α ,6 α -Epoxy-
 - Chromosome Aberrations, 78-6362
 - DNA Repair, 78-6362
- Chromatids
 - Leukocytes, 78-6356
- Chromatin
 - DNA Repair, 78-6357
- Chromosome Aberrations
 - Cell Cycle Kinetics, 78-6358
 - Fibroblasts, 78-6362
- DNA Repair
 - Nucleosome Rearrangement, 78-6357
- DNA Replication
 - Dose-Response Study, 78-6361
 - Hamster V79 Cells, 78-6361
- Endonucleases
 - Chromosome Aberrations, 78-6358
- Escherichia coli*
 - DNA Repair, Review, 78-6045
 - Mutagenic Activity, Review, 78-6045
- Eye Neoplasms
 - Epidemiology, Dog, 78-6516
- Fibroblasts
 - DNA Repair, 78-6313
- Genetics
 - DNA Repair, Review, 78-6055
- Harman
 - DNA Repair, 78-6155
 - Mutagenic Activity, 78-6155
- Hyperthermia
 - Chromosome Aberrations, 78-6362
- Melanoma
 - Chromosome Aberrations, 78-6362
 - Dose-Response Study, Review, 78-6044
 - Epidemiology, Review, 78-6044
 - Epidemiology, Sweden, 78-6578
 - Mathematical Model, Review, 78-6044
- Norharman
 - DNA Repair, 78-6155
 - Mutagenic Activity, 78-6155
- Plasmacytoma
 - Endonucleases, 78-6371
 - RNA, Messenger, 78-6579
- Proteins
 - DNA Repair, Review, 78-6052
- Psoralen, 8-Methoxy-
 - Chromatids, 78-6356
- Psoralen, 4,5',8-Trimethyl-
 - Chromatids, 78-6356
- Retinol
 - Contact Inhibition, 78-6592
- Salmonella typhimurium*
 - Genetic Duplication, 78-6359, 78-6360
- Skin Neoplasms

Ultraviolet Rays (cont'd)

- Melanoma, 78-6046
- Precancerous Conditions, Review, 78-6046
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Co-carcinogenic Activity, Mouse Embryo Cells 78-6283
 - Mutagenic Activity, 78-6295
- Virus, Moloney Murine Leukemia
 - Immunization, 78-6501
- Xeroderma Pigmentosum
 - Cell Survival, 78-6313
 - DNA Repair, 78-6313
 - DNA Repair, Review, 78-6055

Uracil, 5-Fluoro-

- Leukemia, Myeloblastic
- Colonic Neoplasms, 78-6509

Uranium

- Lung Neoplasms
 - Body Height, 78-6554
 - Epidemiology, 78-6554
 - Occupational Hazard, 78-6554
 - Smoking, 78-6554
- Respiratory Tract Neoplasms
 - Occupational Hazard, Review, 78-6049
- Smoking
 - Co-carcinogenic Effect, Review, 78-6049

Urea, 1,3-Bis(2-chloroethyl)-1-nitroso-

- Escherichia coli*
- DNA Repair, 78-6308

Urea, 1-(2-Chloroethyl)-3-(4-methylcyclohexyl)-1-nitroso-

- Leukemia, Myeloblastic
- Colonic Neoplasms, 78-6509

Urea, Ethyl Nitroso-

- Brain Neoplasms
 - Precancerous Conditions, 78-6231
 - Transplacental Carcinogenesis, 78-6231
- Cell Transformation, Neoplastic
 - Ultrastructural Study, Fetal Rat Brain, 78-6231
- Neurons
 - Cell Transformation, Neoplastic, 78-6231

Urea, Methyl Nitroso-

- Adenine, 3-Methyl-
 - Chromatin, Liver, 78-6215
- Caffeine
 - Chromosome Aberrations, 78-6120
- Deoxyribonuclease
 - Chromatin, Liver, 78-6215
- Dimethylamine, *N*-Nitroso-
 - Adenine, 3-Methyl-, 78-6215
 - Purine, 2-Amino-6-methoxy-, 78-6229
- DNA
 - Chromatin, Liver, 78-6229
- Escherichia coli*
 - DNA Repair, 78-6308
- Glioma
 - Virus-Like Particles, 78-6230
- Guanosine
 - RNA, Methylation, 78-6228
- Lymphoblasts, Diploid
 - Mutagenic Activity, 78-6164
- Purine, 2-Amino-6-methoxy-
 - Chromatin, 78-6213
 - Chromatin, Liver, 78-6229
 - DNA, 78-6213

Urea, Methyl Nitroso- (cont'd)

- Xanthine, 78-6229
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - DNA Repair, 78-6308
- Yeasts
 - RNA, Methylation, 78-6228

Uridine, 2'-Deoxy-5-iodo-

- Virus, Epstein-Barr
 - Antigens, Viral, 78-6450
- Virus, Herpes Pan
 - Antigens, Viral, 78-6450
- Virus, Herpes Papio
 - Antigens, Viral, 78-6450
- Virus, Herpes Pongo
 - Antigens, Viral, 78-6450

Urogenital Neoplasms

- Benzoic Acid, 2-Amino-3-hydroxy-
 - Carcinogenic Potential, 78-6311
- Epidemiology
 - California, 78-6548
- Ethnic Groups
 - Epidemiology, Cameroon, 78-6568

Uterine Neoplasms

- Adenocarcinoma
 - Epidemiology, Review, 78-6026
- Contraceptives, Oral
 - Epidemiology, Review, 78-6026
- Diabetes Mellitus
 - Epidemiology, Review, 78-6066
- Dietary Fats
 - Epidemiology, Review, 78-6036
- Endometriosis
 - Abortion, 78-6541
- Estrogens
 - Drug Therapy, 78-6557
 - Epidemiology, 78-6557
 - Epidemiology, Review, 78-6028, 78-6066
 - Genetics, Review, 78-6066
 - Statistical Analysis, 78-6557
- Ethane, 1,2-Dichloro-
 - Carcinogenic Activity, Mouse, Rat, 78-6114
- Ethnic Groups
 - Epidemiology, Hawaii, 78-6546
- FSH
 - Menstruation, 78-6541
- Hypertension
 - Epidemiology, Review, 78-6066
- Leiomyoma
 - FSH, 78-6541
 - LH, 78-6541
 - Somatotropin, 78-6541
- LH
 - Menstruation, 78-6541
- Liver Diseases
 - Androst-4-ene-3,17-dione, 78-6028
 - Epidemiology, Review, 78-6028
 - Estrone, 78-6028
- Menarche
 - Epidemiology, 78-6573
- Menopause
 - Epidemiology, 78-6573
- Obesity
 - Androst-4-ene-3,17-dione, 78-6028
 - Epidemiology, Review, 78-6028, 78-6066
 - Estrone, 78-6028
- Pituitary-Adrenal System

Uterine Neoplasms (cont'd)

- Genetics, Review, 78-6066
- Radiation, Ionizing
- In Utero Exposure, Rat, 78-6382
- Somatotropin
- Menstruation, 78-6541

Valeric Acid, 2,2-Diphenyl-, 2-(Diethylamino)ethyl Ester HCl

- Methane, Azoxy-
- Formaldehyde, 78-6106

Vanadium

- Bacillus subtilis*
- Mutagenic Activity, 78-6128
- Chromium
- Co-carcinogenic Activity, 78-6561
- Digestive System Neoplasms
- Occupational Hazard, 78-6561
- Kidney Neoplasms
- Occupational Hazard, 78-6561
- Respiratory Tract Neoplasms
- Occupational Hazard, 78-6561

Vibrio cholerae

- Lymphoma
- Epidemiology, 78-6487

Vinblastine

- Leukemia, Myeloblastic
- Hodgkin's Disease, 78-6196

Vincristine

- Leukemia, Myeloblastic
- Drug Therapy, 78-6219

Vinyl Cyanide

- see Acrylonitrile

Viral Proteins

- Leukemia L1210
- Virus, C-Type RNA Tumor, 78-6473
- Leydig Cell Tumor
- Virus, Murine Mammary Tumor, 78-6415
- Virus, Abelson Murine Leukemia
- Isolation and Characterization, *Xenopus laevis* 78-6420
- RNA, Viral, 78-6420
- Virus, Bovine Leukemia
- Isolation and Characterization, Review, 78-6088
- Virus, C-Type RNA Tumor
- Biosynthesis, Review, 78-6094
- Structure and Morphogenesis, Review, 78-6094
- Virus, Feline Leukemia
- Antibody Formation, 78-6442
- Virus, Friend Murine Leukemia
- Tryptic Peptide Maps, 78-6436
- Virus, Moloney Murine Leukemia
- Tryptic Peptide Maps, 78-6436
- Virus, Moloney Murine Sarcoma
- Cell Transformation, Neoplastic, 78-6437
- Cyanogen Bromide, 78-6439
- Gene Order, 78-6439
- Peptide Mapping, 78-6439
- Tryptic Peptide Maps, 78-6436
- Virus, Moloney Murine Leukemia, 78-6437
- Virus Replication, 78-6437
- Virus, Murine Leukemia
- Fv-1 Gene, 78-6436
- Tryptic Peptide Maps, 78-6436
- Virus, Murine Mammary Tumor
- Antigenic Determinants, 78-6413, 78-6415

Viral Proteins (cont'd)

- Peptide Hydrolases, 78-6415
- Virus, Polyoma
- Guanosine Triphosphate, 78-6484
- Isolation and Characterization, 78-6484
- RNA, Messenger, 78-6484
- Virus, Rauscher Murine Leukemia
- Tryptic Peptide Maps, 78-6436
- Virus, RNA Tumor
- Genes, Viral, Review, 78-6086
- Virus, SV40
- Genes, Viral, Review, 78-6086
- Virus, Woolly Monkey
- Isolation and Characterization, *Xenopus laevis* 78-6420
- RNA, Viral, 78-6420
- Virus, Helper, 78-6420

Viral Vaccines

- Virus, Feline Leukemia
- Freund's Adjuvant, 78-6442
- Sulfuric Acid, Aluminum Potassium Salt, Dodecahydrate, 78-6442
- Virus, Marek's Disease Herpes
- Cyclophosphamide, 78-6406
- Immunosuppression, 78-6406

Virus, Abelson Murine Leukemia

- Lymphoma
- Macrophages, 78-6419
- Macrophages
- Cell Transformation, Neoplastic, 78-6419
- Lipopolysaccharides, 78-6419
- Virus, Helper, 78-6419
- Virus Replication, 78-6419
- Plasmacytoma
- Mouse, Review, 78-6013
- RNA, Viral
- Viral Proteins, 78-6420
- Thymoma
- Macrophages, 78-6419
- Viral Proteins
- Isolation and Characterization, *Xenopus laevis* 78-6420
- Virus, Moloney Murine Leukemia
- Virus, Helper, 78-6419

Virus, Adeno

- Antigens, Viral
- Virus Subgroup, 78-6467
- Cell Transformation, Neoplastic
- Genes, Viral, Review, 78-6086
- Glycoproteins
- Cell Transformation, Neoplastic, Review, 78-6057

Virus, Adeno 2

- DNA, Viral
- DNA Replication, 78-6462
- Endonucleases, 78-6462
- Thymidine Incorporation, 78-6462
- Virus Replication, Review, 78-6093
- Poly A
- DNA-RNA Hybridization, 78-6463
- RNA Replication
- DNA-RNA Hybridization, 78-6463
- RNA, Viral
- DNA-RNA Hybridization, 78-6463

Virus, Adeno 3

- Antigens, Viral

Virus, Adeno 3 (cont'd)

- Cells, Cultured, 78-6464
- Immunofluorescence, 78-6467
- Cells, Cultured
 - Cell Transformation, Neoplastic, 78-6464
- DNA, Viral
 - Cell Transformation, Neoplastic, 78-6464
 - Cleavage Sites, 78-6465
 - DNA Polymerase, 78-6465
 - Endonucleases, 78-6464, 78-6465
- Virus, Adeno 7
 - DNA-DNA Hybridization, 78-6467

Virus, Adeno 5

- Chromatin
 - Cell Transformation, Neoplastic, 78-6466
- DNA, Viral
 - Cell Transformation, Neoplastic, 78-6466
 - Renaturation Kinetics, 78-6466
- Virus, Shope Rabbit Papilloma
 - Virus Replication, 78-6475

Virus, Adeno 7

- Antigens, Viral
 - Immunofluorescence, 78-6467
- Graft vs Host Reaction
 - Immune Response, 78-6468
 - Precancerous Conditions, 78-6468
- Lymphoid Tissue
 - Precancerous Conditions, 78-6468
- Virus, Adeno 3
 - DNA-DNA Hybridization, 78-6467

Virus, Adeno 12

- Antigens, Viral
 - Immunofluorescence, 78-6467
- Cell Agglutination
 - Morphological Revertants, 78-6470
- Cell Transformation, Neoplastic
 - Morphological Revertants, 78-6470
- Concanavalin A
 - Cell Agglutination, 78-6470
- DNA, Viral
 - DNA-RNA Hybridization, 78-6469
 - Endonucleases, 78-6469
 - Morphological Revertants, 78-6470
 - RNA Replication, 78-6469
- Neoplasms, Experimental
 - Clone Cells, 78-6597
 - Ultrastructural Study, 78-6471
 - Virus-Like Particles, 78-6471
- Virus, Adeno 31
 - DNA-DNA Hybridization, 78-6467

Virus, Adeno 31

- Antigens, Viral
 - Immunofluorescence, 78-6467
- Virus, Adeno 12
 - DNA-DNA Hybridization, 78-6467

Virus, Avian Erythroblastosis

- Cell Differentiation
 - Cell Transformation, Neoplastic, 78-6390
- Hemoglobins
 - Cell Transformation, Neoplastic, 78-6390
 - Temperature Sensitive Mutants, 78-6390
- Leukemia
 - Temperature Sensitive Mutants, 78-6390

Virus, Avian Leukosis

- Cell Transformation, Neoplastic
 - Genes, Viral, 78-6387
 - Nucleotide Sequence, 78-6387
- Deletion Mutants
 - Genes, Viral, Review, 78-6087
- Kidney Neoplasms
 - Virus, Helper, 78-6389
- Liver Neoplasms
 - Isolation and Characterization, 78-6388
 - Virus, Helper, 78-6389
- RNA, Viral
 - DNA-RNA Hybridization, 78-6387
 - Nucleotide Sequence, 78-6387
- Testicular Neoplasms
 - Virus, Helper, 78-6389
- Virus, Helper
 - Cell Transformation, Neoplastic, 78-6388
 - Isolation and Characterization, 78-6389
- Virus, Recombinant
 - Genes, Viral, Review, 78-6087

Virus, Avian Reticuloendotheliosis

- Nucleoproteins
 - Cell Transformation, Neoplastic, 78-6405
 - DNA-RNA Hybridization, 78-6405
 - Isolation and Characterization, 78-6405

Virus, Avian Sarcoma

- Deletion Mutants
 - Genes, Viral, Review, 78-6087
- DNA-RNA Hybridization
 - DNA, Vertebrate, 78-6392
- Glucosamine
 - Antigen-Antibody Reactions, 78-6394
 - Cell Transformation, Neoplastic, 78-6394
 - Reverse Transcriptase, 78-6394
 - RNA, Viral, 78-6394
- Nucleotide Sequence
 - DNA, Vertebrate, 78-6392
- Peptides
 - Cell Transformation, Neoplastic, 78-6481
- RNA, Messenger
 - DNA-RNA Hybridization, 78-6391
 - Isolation and Characterization, 78-6391
 - Nucleotide Sequence, 78-6391
- RNA, Viral
 - DNA-RNA Hybridization, 78-6392
- Virus, Recombinant
 - Genes, Viral, Review, 78-6087

Virus, B77

- Brain Neoplasms
 - Age Factors, Rat, 78-6408
 - Astrocytoma, 78-6408
 - Ultrastructural Study, Rat, 78-6408
- Fibroblasts
 - Cell Transformation, Neoplastic, 78-6398
- Plasminogen Activators
 - Cell Transformation, Neoplastic, 78-6398

Virus, Bovine Leukemia

- Antibodies, Viral
 - Immune Response, Review, 78-6088
- Lymphocytosis
 - Epidemiology, Review, 78-6088
- Lymphosarcoma
 - Epidemiology, Review, 78-6088
 - Horizontal/Vertical Transmission, Review, 78-6088

s, Bovine Leukemia (cont'd)

- Reverse Transcriptase
 - Isolation and Characterization, Review, 78-6088
- Viral Proteins
 - Isolation and Characterization, Review, 78-6088

s, Bovine Papilloma

- Antigens, Viral
 - Trans-Species Infection, 78-6446
- Cell Membrane
 - Antigens, Viral, 78-6446
- DNA, Viral
 - DNA-DNA Hybridization, 78-6445
 - Nucleotide Sequence, 78-6445

- Erythrocytes
 - Hemagglutination, 78-6445

- Fibroma
 - Antigens, Viral, 78-6446

- Fibrosarcoma
 - Antigens, Viral, 78-6446

- Papilloma
 - Antigens, Viral, 78-6446
 - Isolation and Characterization, 78-6445

s, Bovine Parvo

- DNA Polymerase
 - Enzymatic Activity, 78-6447

s, C-Type RNA Tumor

- Leukemia L1210
 - Viral Proteins, 78-6473
 - Virus-Like Particles, 78-6473
- Viral Proteins
 - Biosynthesis, Review, 78-6094
 - Structure and Morphogenesis, Review, 78-6094

s, Cytomegalo

- Prostatic Neoplasms
 - Adenocarcinoma, 78-6499
 - Antibodies, Viral, 78-6499
 - Antigen-Antibody Complex, 78-6499
 - Hyperplasia, 78-6499
 - Immunity, Cellular, 78-6499

s, Epstein-Barr

- Antigens
 - Antibody-Dependent Cytotoxicity, 78-6460
 - Papain, 78-6460
 - Staphylococcal Protein A, 78-6460
- Antigens, Viral
 - Antigen-Antibody Reactions, 78-6450
 - DNA Replication, 78-6454
 - Hybrid Cells, 78-6455
 - IgG Absorption Assay, 78-6456
 - Lymphocyte Transformation, 78-6454
 - Lymphocytes, 78-6454
 - Quinoline, 7-Chloro-4-((4-(diethylamino)-1-methylbutyl)amino)-, 78-6451

Burkitt's Lymphoma

- Antigens, Viral, 78-6459
- Carcinogenic Activity, Review, 78-6084
- DNA Polymerase, 78-6453
- Epidemiology, Review, 78-6089
- Genetics, 78-6459

Cell Membrane

- Antigen-Antibody Reactions, 78-6460
- Antigens, 78-6460

Chromatin

- Antigens, Viral, 78-6456
- 1,3,5-Hexatriene, 1,6-Diphenyl-

Virus, Epstein-Barr (cont'd)

- Fluorescence Polarization, 78-6458
- Liposomes, 78-6458
- Histones
 - Antigens, Viral, 78-6456
- Hodgkin's Disease
 - Antibodies, Viral, 78-6452
 - Antigens, Viral, 78-6459
 - Genetics, 78-6459
 - Lymphocytes, 78-6452
 - B-Lymphocytes, 78-6461
 - Virus Activation, Child, 78-6452
 - Virus Excretion, 78-6452
 - Virus Replication, 78-6461
- Hybrid Cells
 - Chromosomes, Human, 13-15, 78-6455
- Infectious Mononucleosis
 - Epidemiology, Review, 78-6089
- Leukemia, Lymphoblastic
 - Antigens, Viral, 78-6457
- Leukemia, Lymphocytic
 - Virus Replication, 78-6461
- Leukemia, Myeloblastic
 - Antigens, Viral, 78-6457
- Leukemia, Myelocytic
 - B-Lymphocytes, 78-6461
 - Virus Replication, 78-6461
- Liposomes
 - Virus-Cell Interactions, 78-6458
- Lung Neoplasms
 - B-Lymphocytes, 78-6461
 - Virus Replication, 78-6461
- Lymphocytes
 - Fetal Blood, 78-6454
- B-Lymphocytes
 - Fluorescence Polarization, 78-6458
- T-Lymphocytes
 - Fluorescence Polarization, 78-6458
- Nasopharyngeal Neoplasms
 - Antibodies, Viral, 78-6552
 - Epidemiology, 78-6552
 - Epidemiology, Review, 78-6089
- Neoplasms, Multiple Primary
 - Antigens, Viral, 78-6459
 - Genetics, 78-6459
- Phosphonoacetic Acid
 - Antigens, Viral, 78-6450
- Uridine, 2'-Deoxy-5-iodo-
 - Antigens, Viral, 78-6450
- Virus, Herpes Pan
 - Antigenic Determinants, 78-6450
- Virus, Herpes Papio
 - Antigenic Determinants, 78-6450
- Virus, Herpes Pongo
 - Antigenic Determinants, 78-6450

Virus, Feline Leukemia

- Antibody Formation
 - Cat, Guinea Pig, 78-6442
- Antigens, Viral
 - Antigen-Antibody Reactions, 78-6443
 - Cell Membrane, 78-6443
- Leukemia
 - Antigens, Viral, 78-6443
- Lymphoma
 - Antigens, Viral, 78-6443
 - RNA, Viral, 78-6441
- RNA, Viral

Virus, Feline Leukemia (cont'd)

- Tissue Distribution, 78-6441
- Sulfuric Acid, Aluminum Potassium Salt, Dodecahydrate
- Viral Vaccines, 78-6442
- Viral Proteins
 - Antibody Formation, 78-6442
- Viral Vaccines
 - Freund's Adjuvant, 78-6442
- Virus, Feline Sarcoma
 - Antigen-Antibody Reactions, 78-6444
 - Antigenic Determinants, 78-6444

Virus, Feline Sarcoma

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Reverse Transcriptase, 78-6440
 - Virus Replication, 78-6440
- Aflatoxin B1
 - Reverse Transcriptase, 78-6440
 - Virus Replication, 78-6440
- Antigenic Determinants
 - Cell Transformation, Neoplastic, 78-6444
- Antigens, Viral
 - Radioimmunoassay, 78-6444
- Benzo(a)pyrene
 - Reverse Transcriptase, 78-6440
 - Virus Replication, 78-6440
- Virus, Feline Leukemia
 - Antigen-Antibody Reactions, 78-6444
 - Antigenic Determinants, 78-6444

Virus, Friend Murine Leukemia

- Antibody Formation
 - Immunosuppression, 78-6427, 78-6430
 - Spleen Cells, 78-6430
- 1,4-Butanediol, Dimethylsulfonate
 - Hematopoietic Stem Cells, 78-6424
- Erythroleukemia
 - Adenosine Cyclic 3',5' Monophosphate, 78-6426
 - Cell Differentiation, 78-6426
- Erythropoietin
 - Cell Transformation, Neoplastic, 78-6424
- Genetics
 - Immune Response, 78-6427
- Hematopoietic Stem Cells
 - Cell Transformation, Neoplastic, 78-6424
 - Virus Replication, 78-6424
- Histocompatibility Antigens
 - Immune Response, Review, 78-6098
 - Immunity, Cellular, Review, 78-6098
- Immunosuppression
 - Fv-3 Gene, 78-6427
 - Genetics, 78-6427
- Lymphocytes
 - Immunosuppression, 78-6427
- RNA, Messenger
 - Cell Transformation, Neoplastic, 78-6580
 - DNA-RNA Hybridization, 78-6580
 - Isolation and Characterization, 78-6580
 - Poly A, 78-6580
- Viral Proteins
 - Tryptic Peptide Maps, 78-6436
- Virus Replication
 - Genotype, 78-6425
 - Mouse, Hybrid, 78-6425

Virus, Friend Spleen Focus-Forming

- Cell Differentiation
 - A-Type Particles, 78-6428
- Erythroleukemia

Virus, Friend Spleen Focus-Forming (cont'd)

- Virus-Like Particles, 78-6429
 - Interferon
 - A-Type Particles, 78-6429
 - Virus Replication, 78-6429
 - Methane, Sulfinylbis-
 - Virus-Like Particles, 78-6428
 - RNA, Viral
 - Virus-Like Particles, 78-6428
 - Thymidine, Azido-
 - A-Type Particles, 78-6429
 - Virus Replication, 78-6429
 - Virus-Like Particles
 - Cell Differentiation, 78-6428
- Virus, Gross Murine Leukemia**
- Anti-Antibodies
 - Antigen-Antibody Reactions, 78-6431
 - Syngenia Inhibition Test, 78-6431
 - Antigens, Viral
 - Cell Transformation, Neoplastic, 78-6432
 - Glycoproteins
 - Cell Fusion, 78-6431
 - Lymphoma
 - Thymus Gland, Mouse, 78-6432
 - Ultrastructural Study, 78-6432
 - Virus, Murine Leukemia
 - Immune Serums, 78-6489

Virus, Helper

- Kidney Neoplasms
 - Virus, Avian Leukosis, 78-6389
- Liver Neoplasms
 - Virus, Avian Leukosis, 78-6389
- Testicular Neoplasms
 - Virus, Avian Leukosis, 78-6389
- Virus, Abelson Murine Leukemia
 - Macrophages, 78-6419
 - Virus, Moloney Murine Leukemia, 78-6419
- Virus, Avian Leukosis
 - Cell Transformation, Neoplastic, 78-6388
 - Isolation and Characterization, 78-6389
- Virus, Murine Leukemia
 - Carboxy-Lyases, 78-6434
- Virus, Woolly Monkey
 - Viral Proteins, 78-6420

Virus, Herpes

- Nose Neoplasms
 - Virus-Like Particles, 78-6521

Virus, Herpes Pan

- Antigens, Viral
 - Antigen-Antibody Reactions, 78-6450
- Phosphonoacetic Acid
 - Antigens, Viral, 78-6450
- Uridine, 2'-Deoxy-5-iodo-
 - Antigens, Viral, 78-6450
- Virus, Epstein-Barr
 - Antigenic Determinants, 78-6450

Virus, Herpes Papio

- Antigens, Viral
 - Antigen-Antibody Reactions, 78-6450
- Phosphonoacetic Acid
 - Antigens, Viral, 78-6450
- Uridine, 2'-Deoxy-5-iodo-
 - Antigens, Viral, 78-6450
- Virus, Epstein-Barr
 - Antigenic Determinants, 78-6450

s, Herpes Pongo
 Antigen, Viral
 Antigen-Antibody Reactions, 78-6450
 Phosphonoacetic Acid
 Antigens, Viral, 78-6450
 Uridine, 2'-Deoxy-5-iodo-
 Antigens, Viral, 78-6450
 Virus, Epstein-Barr
 Antigenic Determinants, 78-6450

s, Herpes Simplex
 Burkitt's Lymphoma
 DNA Polymerase, 78-6453
 Virus, Shope Rabbit Papilloma
 Virus Replication, 78-6475

s, Herpes Simplex 1
 DNA Replication
 Temperature Sensitive Mutants, 78-6448
 Proteins
 DNA Replication, 78-6448
 RNA
 DNA Replication, 78-6448

s, Herpes Simplex 2
 Cervix Neoplasms
 Anti-Antibodies, 78-6449
 Antigen-Antibody Reactions, 78-6449
 Carcinogenic Activity, Review, 78-6084
 Seroepidemiology, 78-6449
 Seroepidemiology, Review, 78-6095

s, Influenza
 Fibroblasts
 Virus Replication, 78-6435
 Macrophages
 Cytostatic Activity, 78-6435

s, Kilham Rat
 DNA Polymerase
 Enzymatic Activity, 78-6447

s-Like Particles
 Erythroleukemia
 Virus, Friend Spleen Focus-Forming, 78-6429
 Glioma
 Ultrastructural Study, C-Type Particles, 78-6230
 Urea, Methyl Nitroso-, 78-6230
 Leukemia L1210
 Isolation and Characterization, 78-6473
 Virus, C-Type RNA Tumor, 78-6473
 Lymphoma
 Virus, Marek's Disease Herpes, 78-6407
 Mammary Neoplasms, Experimental
 Benz(a)anthracene, 7,12-Dimethyl-, 78-6315
 Melanoma
 Virus, RNA Tumor, 78-6438
 Neoplasms, Experimental
 Virus, Adeno 12, 78-6471
 Nose Neoplasms
 Adenocarcinoma, 78-6521
 Cystadenoma, Papillary, 78-6521
 Virus, Herpes, 78-6521
 Virus, Vaccinia, 78-6521
 Virus, Visna, 78-6521
 Virus, Friend Spleen Focus-Forming
 Cell Differentiation, 78-6428
 Methane, Sulfinylbis-, 78-6428
 RNA, Viral, 78-6428
 Virus, Marek's Disease Herpes

Virus-Like Particles (cont'd)
 Ultrastructural Study, 78-6407
 Virus, Moloney Murine Leukemia
 Plaque Assay, 78-6433
 Replication-Defective Mutants, 78-6433
 Virus, Murine Mammary Tumor
 Neoplasm Transplantation, 78-6409
 Virus, Papilloma
 Mastomys natalensis, 78-6477
 Virus, RNA Tumor
 A-Type Particles, 78-6472
 RNA, Viral, 78-6472
 Virus, Rous Sarcoma
 RNA Polymerase, 78-6399

Virus, Marek's Disease Herpes
 B-Lymphocytes
 Transplantation Immunology, 78-6406
 T-Lymphocytes
 Transplantation Immunology, 78-6406
 Lymphoma
 Virus-Like Particles, 78-6407
 Viral Vaccines
 Cyclophosphamide, 78-6406
 Immunosuppression, 78-6406
 Virus-Like Particles
 Ultrastructural Study, 78-6407
 Virus Replication
 Cell Transformation, Neoplastic, 78-6393
 Virus, Rous Sarcoma
 Virus Replication, 78-6393

Virus, Mink Cell Focus-Inducing
 Virus, Moloney Murine Leukemia
 DNA-RNA Hybridization, 78-6423
 Virus, Murine Leukemia
 DNA-RNA Hybridization, 78-6423

Virus, Moloney Murine Leukemia
 Anti-Antibodies
 Antigen-Antibody Reactions, 78-6431
 Syncytia Inhibition Test, 78-6431
 Antigens, Viral
 Cell Transformation, Neoplastic, 78-6432
 Glycoproteins
 Cell Fusion, 78-6431
 Immunization
 Dose-Response Study, 78-6501
 B-Lymphocytes
 Cell Transformation, Neoplastic, 78-6432
 Immunoglobulins, Surface, 78-6432
 T-Lymphocytes
 Cell Transformation, Neoplastic, 78-6432
 Lymphocyte Depletion, 78-6432
 Lymphoma
 Thymus Gland, Mouse, 78-6432
 Ultrastructural Study, 78-6432
 RNA, Viral
 Ultrastructural Study, Heteroduplexes, 78-6423
 Ultraviolet Rays
 Immunization, 78-6501
 Viral Proteins
 Tryptic Peptide Maps, 78-6436
 Virus, Abelson Murine Leukemia
 Virus, Helper, 78-6419
 Virus-Like Particles
 Plaque Assay, 78-6433
 Replication-Defective Mutants, 78-6433
 Virus, Mink Cell Focus-Inducing

- Virus, Moloney Murine Leukemia (cont'd)**
 - DNA-RNA Hybridization, 78-6423
- Virus, Moloney Murine Sarcoma**
 - Carcinogenic Activity, 78-6501
 - Immunization, 78-6501
 - Viral Proteins, 78-6437
- Virus, Murine Leukemia**
 - DNA-RNA Hybridization, 78-6423
- Virus, Moloney Murine Sarcoma**
 - Benzo(a)pyrene
 - Fibroblasts, 78-6329
 - Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
 - Fibroblasts, 78-6329
 - Sarcoma
 - Lymph Nodes, 78-6502
 - T-Lymphocytes, 78-6502
 - Lymphocytotoxicity, 78-6502
 - Viral Proteins
 - Cell Transformation, Neoplastic, 78-6437
 - Cyanogen Bromide, 78-6439
 - Gene Order, 78-6439
 - Peptide Mapping, 78-6439
 - Tryptic Peptide Maps, 78-6436
 - Virus Replication, 78-6437
 - Virus, Moloney Murine Leukemia**
 - Carcinogenic Activity, 78-6501
 - Immunization, 78-6501
 - Viral Proteins, 78-6437
- Virus, Murine Leukemia**
 - Carboxy-Lyases
 - Cell Transformation, Neoplastic, 78-6434
 - Cell Differentiation
 - Age Factors, 78-6422
 - Diethylamine, *N*-Nitroso-
 - Anti-Antibodies, 78-6417
 - Co-carcinogenic Effect, 78-6417
 - Fibroblasts
 - Cell Transformation, Neoplastic, 78-6435
 - Genes, Viral
 - Virus, Recombinant, Review, 78-6097
 - Genotype
 - Mouse, Wild, 78-6425
 - Globin
 - Age Factors, 78-6422
 - Brain, Liver, Mouse, 78-6422
 - RNA, 78-6422
 - Histocompatibility Antigens
 - Immune Response, Review, 78-6098
 - Immunity, Cellular, Review, 78-6098
 - Immune Serums
 - Age Factors, AKR Mouse, 78-6489
 - Thymectomy, AKR Mouse, 78-6489
 - Leukemia, Lymphoblastic
 - Age Factors, AKR Mouse, 78-6489
 - Immune Serums, 78-6489
 - T-Lymphocytes, 78-6489
 - Thymectomy, AKR Mouse, 78-6489
 - Lymphoma
 - Complement, 78-6503
 - Virus Replication, 78-6418
 - Macrophages
 - Cytostatic Activity, 78-6435
 - RNA, Viral
 - Age Factors, 78-6422
 - Brain, Liver, Mouse, 78-6422
- Virus, Murine Leukemia (cont'd)**
 - DNA-RNA Hybridization, 78-6422
 - Ultrastructural Study, Heteroduplexes, 78-6423
- Thymoma**
 - Complement, 78-6503
- Thymus Neoplasms**
 - Lymphosarcoma, 78-6421
 - Mouse, Hybrid, 78-6421
- Viral Proteins**
 - Fv-1 Gene, 78-6436
 - Tryptic Peptide Maps, 78-6436
- Virus, Gross Murine Leukemia**
 - Immune Serums, 78-6489
- Virus, Helper**
 - Carboxy-Lyases, 78-6434
- Virus, Mink Cell Focus-Inducing**
 - DNA-RNA Hybridization, 78-6423
- Virus, Moloney Murine Leukemia**
 - DNA-RNA Hybridization, 78-6423
- Virus Replication**
 - Quantitation Method, 78-6418
- Virus, Murine Mammary Tumor**
 - Antigenic Determinants
 - A-Type Particles, 78-6415
 - Antigens, Viral
 - Antigen-Antibody Reactions, 78-6409
 - Cell Migration Inhibition, 78-6497
 - Horizontal Transmission, 78-6414
 - Lymphocyte Transformation, 78-6497
 - Breast Neoplasms
 - Antibodies, Viral, 78-6412
 - Dexamethasone
 - Glycoproteins, 78-6411
 - RNA, Viral, 78-6411
 - Hormones
 - Co-carcinogenic Effect, 78-6411
 - Glycoproteins, 78-6411
 - Precancerous Conditions, 78-6411
 - RNA, Viral, 78-6411
 - Immune Response
 - Age Factors, 78-6497
 - Immune Serums
 - Lymphocyte Transformation, 78-6497
 - Leydig Cell Tumor
 - Viral Proteins, 78-6415
 - Lymphocyte Transformation
 - Immune Response, 78-6497
 - Mammary Neoplasms, Experimental
 - A-Type Particles, 78-6415
 - B-Type Particles, 78-6413
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6315
 - DNA-RNA Hybridization, 78-6315
 - Horizontal Transmission, 78-6416
 - Pituitary Implants, 78-6411
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-6315
 - Strain Difference, 78-6416
 - Vertical Transmission, 78-6416
 - Milk
 - Horizontal Transmission, 78-6414
 - Isolation and Characterization, 78-6413
 - Neoplasm Transplantation
 - Antigen-Antibody Reactions, 78-6409
 - Antigens, Viral, 78-6409
 - Virus-Like Particles, 78-6409
 - Virus Replication, 78-6409
 - Prolactin
 - Co-carcinogenic Effect, 78-6411

Virus, Murine Mammary Tumor (cont'd)

- RNA, Viral
 - Age Factors, 78-6410
 - DNA-RNA Hybridization, 78-6410, 78-6411
 - Horizontal Transmission, 78-6411
 - Mammary Development, 78-6410
 - Precancerous Conditions, 78-6411
 - Vertical Transmission, 78-6411
- Viral Proteins
 - Antigenic Determinants, 78-6413, 78-6415
 - Peptide Hydrolases, 78-6415

Virus, Murine Sarcoma

- Carboxy-Lyases
 - Cell Transformation, Neoplastic, 78-6434
 - Enzymatic Activity, 78-6434
 - Virus Replication, 78-6434
- Fibroblasts
 - Cell Transformation, Neoplastic, 78-6435
- Macrophages
 - Cytostatic Activity, 78-6435
- Peptides
 - Cell Transformation, Neoplastic, 78-6481
 - Isolation and Characterization, 78-6481
- Sarcoma
 - Peptides, 78-6481

Virus, Murine Sarcoma-Leukemia

- Melanoma
 - DNA-RNA Hybridization, 78-6438

Virus, Papilloma

- DNA, Viral
 - Mastomys natalensis*, 78-6477
 - Nucleic Acid Conformation, 78-6477
- Laryngeal Neoplasms
 - DNA-DNA Hybridization, 78-6474
 - Isolation and Characterization, 78-6474
 - Papilloma, 78-6474
- Mastomys natalensis*
 - Isolation and Characterization, 78-6477
- Skin Neoplasms
 - Carcinogenic Potential, Review, 78-6090, 78-6091
 - Carcinoma, 78-6091
 - Immunosuppression, Review, 78-6090
 - Mastomys natalensis*, 78-6477
 - Warts, 78-6090, 78-6091
- Virus-Like Particles
 - Mastomys natalensis*, 78-6477
- Warts
 - DNA-DNA Hybridization, 78-6474
 - DNA, Viral, 78-6476
 - Isolation and Characterization, 78-6476

Virus, Papova

- Brain Neoplasms
 - Antigens, Viral, 78-6486

Virus, Parvo

- DNA Polymerase
 - Enzymatic Activity, 78-6447

Virus, Polyoma

- Antigens
 - Cell Transformation, Neoplastic, 78-6496
 - Deletion Mutants, 78-6496
 - Isolation and Characterization, 78-6496
 - Temperature Sensitive Mutants, 78-6496
- Antigens, Viral
 - Carcinogenic Activity, Review, 78-6092

Virus, Polyoma (cont'd)

- Deletion Mutants, 78-6479
- Isolation and Characterization, 78-6479
- Temperature Sensitive Mutants, 78-6479
- Cell Adhesion
 - Cell Transformation, Neoplastic, 78-6480
 - Ultrastructural Study, BHK Cells, 78-6480
- Cell Transformation, Neoplastic
 - Genes, Viral, Review, 78-6086
 - Temperature Sensitive Mutants, 78-6478
- DNA, Viral
 - Carcinogenic Activity, Review, 78-6092
- Fibroblasts
 - Cell Transformation, Neoplastic, 78-6480
- Peptides
 - Cell Transformation, Neoplastic, 78-6481
- RNA
 - DNA Replication, 78-6482
 - Nucleotides, 78-6482
 - Okazaki Fragments, 78-6482
- RNA, Messenger
 - Isolation and Characterization, 78-6484
 - Poly A, 78-6484
 - Viral Proteins, 78-6484
- RNA, Viral
 - DNA-RNA Hybridization, 78-6483
 - Nucleotide Sequence, 78-6483
- Skin Neoplasms
 - Carcinogenic Activity, Review, 78-6092
 - Immunosuppression, Review, 78-6092
 - Temperature Sensitive Mutants
 - Phenotype, 78-6478
- Viral Proteins
 - Guanosine Triphosphate, 78-6484
 - Isolation and Characterization, 78-6484

Virus, Radiation Leukemia

- Diethylamine, *N*-Nitroso-
 - Anti-Antibodies, 78-6417
 - Co-carcinogenic Effect, 78-6417
- Histocompatibility Antigens
 - Immune Response, Review, 78-6098
 - Immunity, Cellular, Review, 78-6098
- Lymphoma
 - Virus Replication, 78-6418
- Neoplasms, Radiation-Induced
 - Lymphoma, 78-6418
- Virus Replication
 - Quantitation Method, 78-6418

Virus, Rauscher Murine Leukemia

- Anti-Antibodies
 - Antigen-Antibody Reactions, 78-6431
 - Syncytia Inhibition Test, 78-6431
- Glycoproteins
 - Cell Fusion, 78-6431
- Viral Proteins
 - Tryptic Peptide Maps, 78-6436

Virus, RD-114

- Carcinoma
 - RNA, Viral, 78-6441
- Concanavalin A
 - T-Lymphocytes, 78-6441
- Lymphoma
 - RNA, Viral, 78-6441
- RNA, Viral
 - Embryonic Tissue, 78-6441
 - Tissue Distribution, 78-6441

Virus, RD-114 (cont'd)

- Sarcoma
- RNA, Viral, 78-6441

Virus Replication

- Hodgkin's Disease
 - Virus, Epstein-Barr, 78-6461
- Leukemia, Lymphocytic
 - Virus, Epstein-Barr, 78-6461
- Leukemia, Myelocytic
 - Virus, Epstein-Barr, 78-6461
- Lung Neoplasms
 - Virus, Epstein-Barr, 78-6461
- Lymphoma
 - Virus, Murine Leukemia, 78-6418
 - Virus, Radiation Leukemia, 78-6418
- Virus, Abelson Murine Leukemia
 - Macrophages, 78-6419
- Virus, Adeno 5
 - Virus, Shope Rabbit Papilloma, 78-6475
- Virus, Feline Sarcoma
 - Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-, 78-6440
 - Aflatoxin B1, 78-6440
 - Benzo(a)pyrene, 78-6440
- Virus, Friend Murine Leukemia
 - Genotype, 78-6425
 - Hematopoietic Stem Cells, 78-6424
 - Mouse, Hybrid, 78-6425
- Virus, Friend Spleen Focus-Forming
 - Interferon, 78-6429
 - Thymidine, Azido-, 78-6429
- Virus, Herpes Simplex
 - Virus, Shope Rabbit Papilloma, 78-6475
- Virus, Influenza
 - Fibroblasts, 78-6435
- Virus, Marek's Disease Herpes
 - Cell Transformation, Neoplastic, 78-6393
 - Virus, Rous Sarcoma, 78-6393
- Virus, Moloney Murine Sarcoma
 - Viral Proteins, 78-6437
- Virus, Murine Leukemia
 - Quantitation Method, 78-6418
- Virus, Murine Mammary Tumor
 - Neoplasm Transplantation, 78-6409
- Virus, Murine Sarcoma
 - Carboxy-Lyases, 78-6434
- Virus, Radiation Leukemia
 - Quantitation Method, 78-6418
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-6393
 - Deletion Mutants, 78-6393
- Virus, Sendai
 - Fibroblasts, 78-6435
- Virus, Shope Rabbit Fibroma
 - Virus, Shope Rabbit Papilloma, 78-6475

Virus, RNA Tumor

- A-Type Particles
 - DNA-RNA Hybridization, 78-6472
 - Nucleotide Sequence, 78-6472
- Carcinogen, Chemical
 - Co-carcinogenic Effect, Review, 78-6096
- Cell Transformation, Neoplastic
 - Genes, Viral, Review, 78-6086
- Hepatoma
 - Virus-Like Particles, Review, 78-6096
- Melanoma
 - DNA-RNA Hybridization, 78-6438

Virus, RNA Tumor (cont'd)

- Reverse Transcriptase, 78-6438
 - RNA, Viral, 78-6438
 - Virus-Like Particles, 78-6438
 - Neoplasms, Experimental
 - Co-carcinogenic Effect, Review, 78-6096
 - RNA, Viral
 - Virus-Like Particles, 78-6472
 - Viral Proteins
 - Genes, Viral, Review, 78-6086
 - Virus-Like Particles
 - A-Type Particles, 78-6472
- Virus, Rous Sarcoma**
- Adenosine Cyclic 3',5' Monophosphate
 - Plasminogen Activators, 78-6338
 - Antigens, Viral
 - Antigen-Antibody Reactions, 78-6401
 - Cell Membrane, 78-6402
 - Cell Transformation, Neoplastic, 78-6402
 - RNA, Messenger, 78-6401
 - Temperature Sensitive Mutants, 78-6402
 - Cell Transformation, Neoplastic
 - Antigenic Determinants, 78-6401
 - Temperature Sensitive Mutants, 78-6396, 78-6402
 - Cycloheximide
 - Glycolysis, 78-6396
 - DNA
 - Transduction, Genetic, 78-6404
 - DNA, Viral
 - DNA-RNA Hybridization, 78-6393
 - Transformation, Genetic, 78-6393, 78-6403
 - Fibroblasts
 - Cell Transformation, Neoplastic, 78-6398
 - Genetics
 - Immune Response, 78-6400
 - Neoplasm Regression, Spontaneous, 78-6400
 - Glycolysis
 - Cell Transformation, Neoplastic, 78-6396
 - Temperature Sensitive Mutants, 78-6396
 - Neoplasms, Embryonal and Mixed
 - Genetics, 78-6400
 - Oligomycin
 - Glycolysis, 78-6396
 - Phenol, 2,4-Dinitro-
 - Oligomycin, 78-6396
 - Plasminogen Activators
 - Cell Transformation, Neoplastic, 78-6338, 78-6395
 - 78-6397, 78-6398
 - Genes, Viral, 78-6397
 - RNA Polymerase
 - Isolation and Characterization, 78-6399
 - RNA Replication, 78-6399
 - RNA, Viral
 - DNA-RNA Hybridization, 78-6404
 - 12-*O*-Tetradecanoylphorbol-13-acetate
 - Plasminogen Activators, 78-6395
 - Transformation, Genetic
 - Temperature Sensitive Mutants, 78-6403
 - Virus-Like Particles
 - RNA Polymerase, 78-6399
 - Virus, Marek's Disease Herpes
 - Virus Replication, 78-6393
 - Virus Replication
 - Cell Transformation, Neoplastic, 78-6393
 - Deletion Mutants, 78-6393

s, Rowson-Parr
 see Virus, Friend Murine Leukemia
s, Semliki Forest
 1,3,5-Hexatriene, 1,6-Diphenyl-
 Fluorescence Polarization, 78-6458
s, Sendai
 Endonucleases
 Chromosome Aberrations, 78-6358
 Fibroblasts
 Virus Replication, 78-6435
 Macrophages
 Cytostatic Activity, 78-6435
s, Shope Rabbit Fibroma
 Virus, Shope Rabbit Papilloma
 Virus Interactions, 78-6475
 Virus Replication, 78-6475
s, Shope Rabbit Papilloma
 Cells, Cultured
 Virus Interactions, 78-6475
 Virus, Adeno 5
 Virus Replication, 78-6475
 Virus, Herpes Simplex
 Virus Replication, 78-6475
 Virus, Shope Rabbit Fibroma
 Virus Interactions, 78-6475
 Virus Replication, 78-6475
 Virus, SV40
 Cell Transformation, Neoplastic, 78-6475
s, SV40
 Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 Cell Transformation, Neoplastic, 78-6242
 Aflatoxin B1
 Cell Transformation, Neoplastic, 78-6242
 Age Factors
 Immune Response, Hamster, 78-6085
 Antigens, Viral
 Immunity, Cellular, 78-6500
 Immunization, 78-6500
 Benzo(a)pyrene
 Cell Transformation, Neoplastic, 78-6242
 Brain Neoplasms
 Antigens, Viral, 78-6486
 Ependymoma, 78-6486
 Cell Transformation, Neoplastic
 Genes, Viral, Review, 78-6086
 Diethylamine, *N*-Nitroso-
 Cell Transformation, Neoplastic, 78-6242
 DNA, Viral
 RNA Replication, 78-6485
 Virus Replication, Review, 78-6093
 Ependymoma
 Antigens, Viral, 78-6486
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Cell Transformation, Neoplastic, 78-6242
 Leukemia, Lymphocytic
 Antigen-Antibody Reactions, 78-6085
 Histocompatibility Antigens, 78-6085
 T-Lymphocytes
 Antilymphocyte Serum, 78-6500
 Immunity, Cellular, 78-6500
 Lymphosarcoma
 Antigen-Antibody Reactions, 78-6085
 Histocompatibility Antigens, 78-6085
 Methanesulfonic Acid, Methyl Ester
 Cell Transformation, Neoplastic, 78-6242

Virus, SV40 (cont'd)
 2-Naphthylamine
 Cell Transformation, Neoplastic, 78-6242
 Neoplasms, Experimental
 Cell Differentiation, 78-6085
 1,2-Oxathiolane, 2,2-Dioxide
 Cell Transformation, Neoplastic, 78-6242
 2-Oxetanone
 Cell Transformation, Neoplastic, 78-6242
 Peptides
 Cell Transformation, Neoplastic, 78-6481
 Plasminogen Activators
 Cell Transformation, Neoplastic, 78-6310
 Quinoline, 4-Nitro-, 1-Oxide
 Cell Transformation, Neoplastic, 78-6242
 RNA, Messenger
 DNA, Viral, 78-6485
 Endonucleases, 78-6485
 R-Loop Structures, 78-6485
 Sarcoma
 Antigen-Antibody Reactions, 78-6085
 Histocompatibility Antigens, 78-6085
 Sarcoma, Osteogenic
 Antigen-Antibody Reactions, 78-6085
 Histocompatibility Antigens, 78-6085
 Sarcoma, Reticulum Cell
 Antigen-Antibody Reactions, 78-6085
 Histocompatibility Antigens, 78-6085
 Viral Proteins
 Genes, Viral, Review, 78-6086
 Virus, Shope Rabbit Papilloma
 Cell Transformation, Neoplastic, 78-6475
Virus, Vaccinia
 Nose Neoplasms
 Virus-Like Particles, 78-6521
Virus, Visna
 Nose Neoplasms
 Virus-Like Particles, 78-6521
Virus, Woolly Monkey
 RNA, Viral
 Viral Proteins, 78-6420
 Viral Proteins
 Isolation and Characterization, *Xenopus laevis*
 78-6420
 Virus, Helper
 Viral Proteins, 78-6420
Vitamin B12
 Stomach Neoplasms
 Malabsorption Syndromes, 78-6534
Vitamin E
 Carcinogen, Chemical
 Carcinogenic Activity, Review, 78-6004
Warts
 Skin Neoplasms
 Virus, Papilloma, 78-6090, 78-6091
 Virus, Papilloma
 DNA-DNA Hybridization, 78-6474
 DNA, Viral, 78-6476
 Isolation and Characterization, 78-6476
Water Pollutants
 Aroclor 1254
 Ames Test, 78-6158
 Microsomes, Liver
 Ames Test, 78-6158

Water Pollution

- Bladder Neoplasms
 - Lymphoma, 78-6556
- Brain Neoplasms
 - Bromine, 78-6556
- Bromine
 - Epidemiology, 78-6556
- Chloroform
 - Epidemiology, 78-6556
- Kidney Neoplasms
 - Chloroform, 78-6556
- Lung Neoplasms
 - Bromine, 78-6556
- Lymphoma
 - Bromine, 78-6556

Wounds and Injuries

- Melanoma
 - Case Report, 78-6512
 - Nevus, 78-6512
- Mouth Neoplasms
 - Carcinoma, Epidermoid, 78-6375
 - Epidemiology, 78-6375

Xanthine

- Urea, Methyl Nitroso-
 - Purine, 2-Amino-6-methoxy-, 78-6229

Xanthine, 3-Hydroxy-

- Enzyme Activation
 - Transplacental Carcinogenesis, 78-6249
- Hepatoma
 - Transplacental Carcinogenesis, 78-6249
- Lung Neoplasms
 - Adenoma, 78-6249
 - Transplacental Carcinogenesis, 78-6249
- Mammary Neoplasms, Experimental
 - Adenofibroma, 78-6250
 - Adenoma, 78-6250
 - Carcinoma, 78-6250
 - Dose-Response Study, Rat, 78-6250
 - Prolactin, 78-6250
 - Transplacental Carcinogenesis, 78-6250

Sulfate

- Metabolism, Liver, 78-6249

Xanthine, 3-Isobutyl-1-methyl-

- Erythroleukemia
 - Adenosine Cyclic 3',5' Monophosphate, 78-6426

Xeroderma Pigmentosum

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - DNA Repair, Review, 78-6055
- Benz(a)anthracene, 7-Bromomethyl-
 - Cell Survival, 78-6313
 - DNA Repair, 78-6313
- Chromosomes
 - Hereditary Diseases, Review, 78-6074
- Ultraviolet Rays
 - Cell Survival, 78-6313
 - DNA Repair, 78-6313
 - DNA Repair, Review, 78-6055

Yeasts

- Ethylene, Trichloro-
 - Mutagenic Activity, 78-6116
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - RNA, Methylation, 78-6228
- Methanesulfonic Acid, Methyl Ester
 - RNA, Methylation, 78-6228
- Propane, 1-Chloro-2,3-epoxy-
 - Mutagenic Activity, 78-6116
- Sulfuric Acid, Dimethyl Ester
 - RNA, Methylation, 78-6228
- Urea, Methyl Nitroso-
 - RNA, Methylation, 78-6228

Yellow AB

- see 2-Naphthylamine, 1-(Phenylazo)-

Yellow OB

- see 2-Naphthylamine, 1-(*o*-Tolylazo)-

Zearalenol

- Estradiol
 - Receptors, Hormone, 78-6350

Zearalenone

- Estradiol
 - Receptors, Hormone, 78-6350

Zinc

- Calcium
 - Chromosome Aberrations, 78-6147
- Lymphocytes
 - Chromosome Aberrations, 78-6147
- Stomach Neoplasms
 - Epidemiology, Japan, 78-6148
 - Food Contamination, 78-6148

Chemical Abstracts Service Registry Number Index

- | | | |
|--|---|--|
| 50-00-0, 78-6106, 78-6225 | 53-59-8, 78-6113, 78-6129, 78-6195 | 57-83-0, 78-6346, 78-6347, 78-6355
78-6544 |
| 50-02-2, 78-6307, 78-6411 | 53-70-3, 78-6169 | 57-88-5, 78-6036, 78-6123, 78-6192 |
| 50-06-6, 78-6106, 78-6109, 78-6115
78-6167, 78-6193, 78-6217
78-6257, 78-6319, 78-6320
78-6326, 78-6331 | 53-86-1, 78-6488 | 57-97-6, 78-6015, 78-6037, 78-6162
78-6314, 78-6316, 78-6317
78-6318, 78-6343, 78-6351 |
| 50-07-7, 78-6232, 78-6267, 78-6358 | 53-96-3, 78-6162, 78-6190, 78-6193
78-6267, 78-6291, 78-6334
78-6492 | 58-08-2, 78-6120, 78-6121, 78-6122
78-6123, 78-6426 |
| 50-18-0, 78-6051, 78-6162, 78-6194
78-6195, 78-6196, 78-6197
78-6232, 78-6267, 78-6406 | 54-05-7, 78-6451 | 58-15-1, 78-6214 |
| 50-23-7, 78-6307 | 54-11-5, 78-6176, 78-6177 | 58-22-0, 78-6343 |
| 50-27-1, 78-6037, 78-6349, 78-6351 | 54-25-1, 78-6051 | 58-39-9, 78-6343 |
| 50-28-2, 78-6037, 78-6346, 78-6348
78-6349, 78-6350, 78-6351
78-6355, 78-6544 | 54-42-2, 78-6450 | 58-55-9, 78-6122, 78-6123, 78-6292
78-6426 |
| 50-29-3, 78-6012, 78-6111, 78-6305 | 54-85-3, 78-6163, 78-6264 | 58-89-9, 78-6214 |
| 50-32-8, 78-6242, 78-6302, 78-6324
78-6326, 78-6327, 78-6328
78-6329, 78-6330, 78-6332
78-6333, 78-6334, 78-6335
78-6440 | 54-88-6, 78-6267 | 59-02-9, 78-6004 |
| 50-44-2, 78-6209 | 54-91-1, 78-6232 | 59-87-0, 78-6227 |
| 50-50-0, 78-6347 | 55-17-4, 78-6162, 78-6205, 78-6206
78-6212, 78-6216, 78-6265 | 59-88-1, 78-6182 |
| 50-55-5, 78-6321, 78-6490 | 55-18-5, 78-6162, 78-6216, 78-6218
78-6242, 78-6265, 78-6417
78-6492 | 60-09-3, 78-6265, 78-6278 |
| 50-67-9, 78-6318 | 55-80-1, 78-6274, 78-6275, 78-6276
78-6277 | 60-11-7, 78-6218, 78-6225, 78-6263 |
| 50-71-5, 78-6233 | 55-98-1, 78-6051, 78-6163, 78-6232
78-6424 | 60-29-7, 78-6335 |
| 50-76-0, 78-6304, 78-6347 | 56-23-5, 78-6216, 78-6265 | 60-34-4, 78-6181 |
| 50-81-7, 78-6130, 78-6204 | 56-38-2, 78-6305 | 60-56-0, 78-6248 |
| 50-99-7, 78-6207 | 56-49-5, 78-6106, 78-6169, 78-6217
78-6265, 78-6283, 78-6312
78-6319, 78-6320, 78-6321
78-6322, 78-6326, 78-6334
78-6490, 78-6493 | 60-57-1, 78-6111 |
| 51-03-6, 78-6238 | 56-53-1, 78-6027, 78-6051, 78-6315
78-6342, 78-6355 | 60-80-0, 78-6292 |
| 51-18-3, 78-6196, 78-6222 | 56-55-3, 78-6015, 78-6115, 78-6319
78-6322, 78-6334 | 60-92-4, 78-6338, 78-6426, 78-6596 |
| 51-21-8, 78-6509 | 56-57-5, 78-6162, 78-6242, 78-6245
78-6267 | 61-57-4, 78-6490 |
| 51-28-5, 78-6396, 78-6590 | 56-73-5, 78-6195 | 62-38-4, 78-6140 |
| 51-43-4, 78-6033 | 56-75-7, 78-6052 | 62-44-2, 78-6051, 78-6291, 78-6292
78-6293 |
| 51-75-2, 78-6219 | 56-86-0, 78-6198 | 62-49-7, 78-6199 |
| 51-79-6, 78-6162, 78-6183, 78-6184
78-6185 | 56-87-1, 78-6307 | 62-50-0, 78-6108, 78-6120, 78-6161
78-6162, 78-6216, 78-6244
78-6245, 78-6359, 78-6360 |
| 51-98-9, 78-6353 | 57-14-7, 78-6179, 78-6180 | 62-53-3, 78-6262, 78-6266 |
| 52-01-7, 78-6016 | 57-22-7, 78-6219 | 62-55-5, 78-6216, 78-6347 |
| 52-24-4, 78-6226 | 57-30-7, 78-6223, 78-6312 | 62-68-0, 78-6106 |
| 52-68-6, 78-6145 | 57-50-1, 78-6125 | 62-75-9, 78-6111, 78-6162, 78-6212
78-6213, 78-6214, 78-6215
78-6217, 78-6229, 78-6263
78-6265, 78-6324 |
| 53-03-2, 78-6219 | 57-56-7, 78-6160 | 63-05-8, 78-6028, 78-6333, 78-6348
78-6544 |
| 53-06-5, 78-6307 | 57-57-8, 78-6242, 78-6245 | 63-25-2, 78-6305 |
| 53-16-7, 78-6028, 78-6037, 78-6345
78-6348, 78-6349, 78-6544 | 57-63-6, 78-6345, 78-6353 | 63-91-2, 78-6201 |
| 53-41-8, 78-6341 | 57-68-1, 78-6264 | 64-17-5, 78-6032, 78-6108, 78-6375
78-6549 |

64-67-5, 78-6159	80-51-3, 78-6182	100-75-4, 78-6240
64-86-8, 78-6222	80-62-6, 78-6372	100-97-0, 78-6225
65-85-0, 78-6256	81-07-2, 78-6280, 78-6281, 78-6282 78-6283, 78-6294	101-14-4, 78-6264, 78-6279
66-27-3, 78-6228, 78-6242, 78-6358	83-67-0, 78-6123	101-21-3, 78-6186
66-76-2, 78-6334	83-88-5, 78-6263	101-77-9, 78-6261
66-81-9, 78-6396	83-89-6, 78-6159	102-71-6, 78-6224
67-21-0, 78-6162, 78-6205, 78-6206 78-6212, 78-6216, 78-6265	84-16-2, 78-6259	103-84-4, 78-6292
67-42-5, 78-6503	85-01-8, 78-6312	103-90-2, 78-6291
67-62-9, 78-6160	85-02-9, 78-6239	106-40-1, 78-6266
67-66-3, 78-6109, 78-6225, 78-6556	85-83-6, 78-6317	106-47-8, 78-6266
67-68-5, 78-6426, 78-6428	85-84-7, 78-6152	106-49-0, 78-6266
68-19-9, 78-6534	85-86-9, 78-6317	106-50-3, 78-6269
68-23-5, 78-6355	86-01-1, 78-6484	106-51-4, 78-6299
68-26-8, 78-6004, 78-6039, 78-6338 78-6592	86-40-8, 78-6159	106-89-8, 78-6116
68-76-8, 78-6298	88-10-8, 78-6187	106-93-4, 78-6115
68-89-3, 78-6163	88-32-4, 78-6257, 78-6328, 78-6333	106-99-0, 78-6014
69-05-6, 78-6159	90-64-2, 78-6252	107-06-2, 78-6113, 78-6114
69-72-7, 78-6287	91-22-5, 78-6239	107-13-1, 78-6021
69-89-6, 78-6229	91-53-2, 78-6328, 78-6333	107-20-0, 78-6112
70-18-8, 78-6004, 78-6113, 78-6130 78-6274, 78-6275, 78-6326 78-6333	91-59-8, 78-6162, 78-6242, 78-6244 78-6264, 78-6566	107-27-7, 78-6139
70-25-7, 78-6120, 78-6161, 78-6164 78-6228, 78-6242, 78-6245 78-6295	91-94-1, 78-6270, 78-6279	109-84-2, 78-6182
71-30-7, 78-6160	92-67-1, 78-6162, 78-6260	110-16-7, 78-6359
71-43-2, 78-6014, 78-6258	92-87-5, 78-6162, 78-6192, 78-6260 78-6262, 78-6264, 78-6279 78-6566	110-89-4, 78-6124, 78-6290
72-33-3, 78-6354, 78-6355	93-76-5, 78-6287, 78-6288, 78-6305	111-17-1, 78-6117
73-22-3, 78-6201, 78-6203, 78-6372 78-6567	94-59-7, 78-6290	115-02-6, 78-6199, 78-6200, 78-6212
73-24-5, 78-6124	94-74-6, 78-6287	115-29-7, 78-6163
73-31-4, 78-6024	94-75-7, 78-6271, 78-6287, 78-6288 78-6305	115-90-2, 78-6284
73-40-5, 78-6124	95-80-7, 78-6190	118-00-3, 78-6228
74-89-5, 78-6225	96-09-3, 78-6253, 78-6257, 78-6272	118-74-1, 78-6305
75-01-4, 78-6110, 78-6112	97-00-7, 78-6331, 78-6333	119-26-6, 78-6182, 78-6237
75-07-0, 78-6188	97-53-0, 78-6290	119-65-3, 78-6239
75-11-6, 78-6329	97-56-3, 78-6278, 78-6286	120-12-7, 78-6336
75-44-5, 78-6109	97-63-2, 78-6372	121-00-6, 78-6257, 78-6328, 78-6333
76-38-0, 78-6335	97-77-8, 78-6106	121-69-7, 78-6262
76-57-3, 78-6293	98-86-2, 78-6171	121-75-5, 78-6305
77-47-4, 78-6167	98-92-0, 78-6207, 78-6208	121-81-3, 78-6285
77-78-1, 78-6228	99-54-7, 78-6333	123-33-1, 78-6120
77-92-9, 78-6589	99-98-9, 78-6284	123-75-1, 78-6290
79-01-6, 78-6116	100-01-6, 78-6266	124-20-9, 78-6204
79-44-7, 78-6187	100-14-1, 78-6333	124-38-9, 78-6019
	100-42-5, 78-6014, 78-6252, 78-6253	124-40-3, 78-6290
		126-72-7, 78-6144
		128-37-0, 78-6184, 78-6206, 78-6218
		129-00-0, 78-6336

15-4, 78-6299
79-3, 78-6152
06-2, 78-6159
32-7, 78-6264, 78-6566
56-7, 78-6284
79-4, 78-6241
47-2, 78-6138
85-3, 78-6290
01-6, 78-6285
24-3, 78-6193
79-8, 78-6256
82-3, 78-6051
29-1, 78-6586
78-6, 78-6264
93-8, 78-6308
51-4, 78-6182
54-7, 78-6426
70-6, 78-6337
87-8, 78-6239
27-3, 78-6239
63-3, 78-6155, 78-6239
95-3, 78-6236
20-4, 78-6166, 78-6312
13-1, 78-6107
76-7, 78-6354
81-7, 78-6356
29-6, 78-6193
01-2, 78-6009
04-5, 78-6204
79-4, 78-6338, 78-6339
33-3, 78-6238
03-3, 78-6051, 78-6196, 78-6209
78-6211
80-6, 78-6320
00-2, 78-6305
36-4, 78-6335
34-0, 78-6122, 78-6123, 78-6292
78-6426
84-6, 78-6271
88-8, 78-6590
09-1, 78-6590
74-3, 78-6339, 78-6596
24-6, 78-6488
40-4, 78-6266
20-5, 78-6165

439-14-5, 78-6328
443-48-1, 78-6162, 78-6235, 78-6328
446-72-0, 78-6350
446-86-6, 78-6209
474-25-9, 78-6590
479-13-0, 78-6350
480-54-6, 78-6216
485-72-3, 78-6350
486-84-0, 78-6155
506-68-3, 78-6439
521-18-6, 78-6343
532-12-7, 78-6175
538-75-0, 78-6590
539-44-6, 78-6182
540-37-4, 78-6266
540-73-8, 78-6162
542-88-1, 78-6009
547-81-9, 78-6351
548-26-5, 78-6153
548-93-6, 78-5044, 78-6311, 78-6567
551-06-4, 78-6265
551-93-9, 78-6202
553-24-2, 78-6200
555-60-2, 78-6590
573-58-0, 78-6263
578-76-7, 78-6215, 78-6221
585-08-0, 78-6312, 78-6325
590-96-5, 78-6107
592-62-1, 78-6189, 78-6295
602-87-9, 78-6254
604-59-1, 78-6106, 78-6326
606-23-5, 78-6171
607-30-7, 78-6247
615-53-2, 78-6212
624-83-9, 78-6103
629-27-6, 78-6329
638-36-8, 78-6013
670-54-2, 78-6299
671-16-9, 78-6196, 78-6219
680-31-9, 78-6162
684-93-5, 78-6120, 78-6164, 78-6213
78-6215, 78-6228, 78-6229
78-6230, 78-6308
695-77-2, 78-6167
738-99-8, 78-6226

745-65-3, 78-6488
758-17-8, 78-6181
759-73-9, 78-6231
762-42-5, 78-6299
789-47-9, 78-6267
842-07-9, 78-6152
865-21-4, 78-6196
930-55-2, 78-6237
937-14-4, 78-6289
937-40-6, 78-6273
941-69-5, 78-6299
992-59-6, 78-6263
1116-54-7, 78-6224
1120-71-4, 78-6242, 78-6245
1130-32-1, 78-6426
1155-38-0, 78-6325
1162-65-8, 78-6124, 78-6125, 78-6216
78-6242, 78-6245, 78-6440
1169-79-5, 78-6351
1184-78-7, 78-6225
1204-06-4, 78-6202
1304-56-9, 78-6127
1309-37-1, 78-6335
1328-67-2, 78-6130
1332-21-4, 78-6002, 78-6023, 78-6032
78-6168, 78-6169, 78-6383
78-6529
1401-06-5, 78-6590
1404-19-9, 78-6396
1407-15-4, 78-6232
1434-54-4, 78-6331
1452-77-3, 78-6207, 78-6208
1464-53-5, 78-6325
1509-87-1, 78-6590
1582-09-8, 78-6268
1636-33-5, 78-6265
1637-39-4, 78-6585
1720-32-7, 78-6458
1746-01-6, 78-6217, 78-6288
1881-77-2, 78-6015
1910-42-5, 78-6184
1912-32-9, 78-6164
1921-70-6, 78-6013
2277-92-1, 78-6255
2312-76-7, 78-6157

2518-02-7, 78-6325
 2541-69-7, 78-6015
 2586-60-9, 78-6263
 2606-87-3, 78-6015
 2642-98-0, 78-6190, 78-6267
 2646-17-5, 78-6152
 2834-90-4, 78-6247
 2834-92-6, 78-6247
 3105-97-3, 78-6328
 3118-97-6, 78-6152
 3416-24-8, 78-6394, 78-6591
 3688-53-7, 78-6227, 78-6244
 3697-24-3, 78-6323
 3715-92-2, 78-6303
 3741-15-9, 78-6244
 3761-53-3, 78-6263
 3902-71-4, 78-6356
 4075-81-4, 78-6256
 4213-45-0, 78-6120
 4408-78-0, 78-6450, 78-6453
 4553-89-3, 78-6149
 4637-56-3, 78-6212
 5142-23-4, 78-6215
 5371-52-8, 78-6237
 5916-52-9, 78-6350
 6051-87-2, 78-6319
 6098-44-8, 78-6055, 78-6191, 78-6242
 78-6308, 78-6440
 6164-98-3, 78-6305
 6381-77-7, 78-6138
 6613-85-0, 78-6328
 6795-23-9, 78-6125
 7203-90-9, 78-6223
 7227-91-0, 78-6214, 78-6221, 78-6223
 78-6266
 7439-89-6, 78-6134, 78-6565
 7439-96-5, 78-6148, 78-6561
 7439-97-6, 78-6134, 78-6139
 7440-02-0, 78-6141, 78-6142, 78-6143
 78-6561
 7440-07-5, 78-6367
 7440-24-6, 78-6134
 7440-36-0, 78-6128
 7440-39-3, 78-6134
 7440-43-9, 78-6134

7440-44-0, 78-6105
 7440-47-3, 78-6561
 7440-50-8, 78-6134
 7440-61-1, 78-6049, 78-6554
 7440-62-2, 78-6128, 78-6561
 7440-65-5, 78-6147, 78-6148
 7440-70-2, 78-6061
 7447-40-7, 78-6371
 7631-90-5, 78-6160
 7631-99-4, 78-6225, 78-6574
 7632-00-0, 78-6123, 78-6203, 78-6204
 78-6214, 78-6224, 78-6241
 7646-79-9, 78-6106
 7647-14-5, 78-6145
 7647-17-8, 78-6128
 7664-41-7, 78-6042
 7681-93-8, 78-6256
 7722-64-7, 78-6130
 7726-95-6, 78-6556
 7738-94-5, 78-6130
 7757-79-1, 78-6574
 7757-83-7, 78-6130
 7758-16-9, 78-6138
 7758-97-6, 78-6130, 78-6304
 7773-01-5, 78-6137
 7778-66-7, 78-6132
 7782-39-0, 78-6109
 7782-49-2, 78-6146
 7782-50-5, 78-6132, 78-6262, 78-6330
 7782-77-6, 78-6006, 78-6256, 78-6359
 7784-24-9, 78-6442
 7785-87-7, 78-6138
 7789-00-6, 78-6130
 7803-49-8, 78-6160, 78-6244
 7803-57-8, 78-6182
 8001-28-3, 78-6307
 8001-58-9, 78-6017, 78-6336
 8002-05-9, 78-6002, 78-6018
 8015-95-0, 78-6490
 8062-14-4, 78-6153
 9000-07-1, 78-6490
 9000-71-9, 78-6410
 9001-12-1, 78-6065
 9001-45-0, 78-6297, 78-6587
 9001-62-1, 78-6589

9001-63-2, 78-6201
 9001-67-6, 78-6591
 9001-73-4, 78-6460
 9001-78-9, 78-6261, 78-6339, 78-6457
 9002-62-4, 78-6037, 78-6250, 78-6343
 78-6346, 78-6349, 78-6411
 9002-72-6, 78-6541
 9003-98-9, 78-6191, 78-6215
 9004-10-8, 78-6207
 9005-25-8, 78-6125, 78-6435
 9007-12-9, 78-6494, 78-6522
 9008-11-1, 78-6077, 78-6429
 9035-50-1, 78-6020, 78-6106, 78-6109
 78-6217, 78-6218, 78-6340
 10024-89-2, 78-6251
 10028-15-6, 78-6105
 10028-16-7, 78-6128
 10034-93-2, 78-6182
 10038-98-9, 78-6128
 10043-66-0, 78-6376
 10043-92-2, 78-6554
 10049-07-7, 78-6128
 10588-01-9, 78-6130
 10605-21-7, 78-6303, 78-6306
 11028-71-0, 78-6205, 78-6441, 78-6470
 78-6597
 11056-06-7, 78-6120, 78-6308, 78-6358
 11096-26-7, 78-6424
 11097-69-1, 78-6158, 78-6166, 78-6237
 78-6257, 78-6296, 78-6323
 78-6324
 12001-28-4, 78-6002, 78-6023, 78-6032
 78-6168, 78-6169, 78-6383
 78-6529
 12001-29-5, 78-6002, 78-6023, 78-6032
 78-6168, 78-6169, 78-6383
 78-6529
 12035-72-2, 78-6141
 12059-95-9, 78-6127, 78-6369, 78-6370
 12172-73-5, 78-6002, 78-6023, 78-6032
 78-6168, 78-6169, 78-6383
 78-6529
 12244-57-4, 78-6133
 12268-14-3, 78-6129, 78-6131
 12587-46-1, 78-6370
 12674-11-2, 78-6296
 12798-09-3, 78-6359, 78-6360
 13345-64-7, 78-6317

13394-86-0, 78-6297	16812-54-7, 78-6141	28322-02-3, 78-6267
13479-29-3, 78-6249, 78-6250	16941-12-1, 78-6128	28622-94-8, 78-6325
13551-87-6, 78-6385	17068-78-9, 78-6002, 78-6023, 78-6032 78-6168, 78-6169, 78-6383 78-6529	28822-58-4, 78-6426
13907-45-4, 78-6129, 78-6131	17157-48-1, 78-6115	30516-87-1, 78-6429
13907-47-6, 78-6129, 78-6131	17924-92-4, 78-6350	30746-58-8, 78-6331
13967-48-1, 78-6048	18883-66-4, 78-6207, 78-6208, 78-6212 78-6509	31431-39-7, 78-6328
13981-37-8, 78-6142	19992-69-9, 78-6214	33073-59-5, 78-6509
13981-52-7, 78-6049	20535-83-5, 78-6213, 78-6215, 78-6221 78-6229	33421-40-8, 78-6201
14158-27-1, 78-6142	20816-12-0, 78-6128, 78-6371	34807-41-5, 78-6306, 78-6309
14255-04-0, 78-6049, 78-6134, 78-6135 78-6136, 78-6147	20830-81-3, 78-6232	35065-27-1, 78-6305
14762-75-5, 78-6386	21645-51-2, 78-6126	37574-47-3, 78-6325, 78-6331
14762-78-8, 78-6048	23214-92-8, 78-6232, 78-6308	39362-56-6, 78-6120
14808-79-8, 78-6249	24279-91-2, 78-6232	39638-32-9, 78-6118
14901-08-7, 78-6178	24928-17-4, 78-6306, 78-6309	51778-96-2, 78-6299
15131-84-7, 78-6325	24961-39-5, 78-6313	53469-21-9, 78-6296
15262-20-1, 78-6564	25013-16-5, 78-6257, 78-6328, 78-6333	56875-27-5, 78-6299
15663-27-1, 78-6308	25126-76-5, 78-6343	57303-99-8, 78-6326
16543-55-8, 78-6175, 78-6176	25405-85-0, 78-6306	59963-01-8, 78-6326, 78-6329
16561-29-8, 78-6283, 78-6295, 78-6306 78-6307, 78-6308, 78-6309 78-6310, 78-6325, 78-6338 78-6395	25843-45-2, 78-6106	60195-69-9, 78-6306
16568-02-8, 78-6181	26644-46-2, 78-6241	65445-61-6, 78-6240
16644-15-8, 78-6325	27208-37-3, 78-6337	66107-37-7, 78-6306

THE UNIVERSITY OF CHICAGO

Wiswesser Line Notation Index

.H2.CR-O4, 78-6130 .BA, 78-6134 .BE..O, 78-6127 .BR, 78-6556 .C, 78-6105, 78-6386 .CA, 78-6061 .CD, 78-6134 .CE, 78-6048 .CO..G2, 78-6106 .CR, 78-6561 .CU, 78-6134 .FE, 78-6134, 78-6565 .FE2.O3, 78-6335 .HG, 78-6134, 78-6139 .I, 78-6376 .KA..G, 78-6371 .KA..MN-O4, 78-6130 .KA..N-O3, 78-6574 .KA2.CR-O2-Q2, 78-6130 .MN, 78-6148, 78-6561 .MN..G2, 78-6137 .MN..S-O2-Q2, 78-6138 .NA..G, 78-6145 .NA..N-O2, 78-6123, 78-6203, 78-6204, 78-6214, 78-6224 78-6241 .NA..N-O3, 78-6225, 78-6574 .NA..S-O-Q2, 78-6160 .NA2.CR2-O5-Q3, 78-6130 .NA2.P2-O3-Q4, 78-6138 .NA2.S-O3, 78-6130 .NI, 78-6141, 78-6142, 78-6143, 78-6561 .NI3.S2, 78-6141 .OS..O-4, 78-6128, 78-6371 .PB, 78-6049, 78-6134, 78-6135, 78-6136, 78-6147 .PO, 78-6049 .PU, 78-6367 .PV..O2, 78-6127, 78-6369, 78-6370 .RA, 78-6564 .RH.G3, 78-6128 .RN, 78-6554 .RU, 78-6048 .SB, 78-6128	.SE, 78-6146 .SR, 78-6134, 78-6142 .TE, 78-6128 .UR, 78-6049, 78-6554 .V, 78-6128, 78-6561 .ZN, 78-6147, 78-6148 CS G, 78-6128 E CN, 78-6439 E1YE1O 3PO, 78-6144 E2E, 78-6115 FXFFR CNW ENW DN3&3, 78-6268 G 6-R, 78-6305 G-HG-2, 78-6139 GR C DNU1N1&1, 78-6305 GR CG DG EG FR BG DG EG, 78-6305 GR CMVOY, 78-6186 GVG, 78-6109 GVN1&1, 78-6187 GVN2&2, 78-6187 GXGGG, 78-6216, 78-6265 GXGGG YR DG&R DG, 78-6012, 78-6111, 78-6305 GXGGYQPO&O1&O1, 78-6145 GYGG, 78-6109, 78-6225, 78-6556 YGU1G, 78-6116 G1O1G, 78-6009 G1U1, 78-6110, 78-6112 G2, 78-6132, 78-6262, 78-6330 G2G, 78-6113, 78-6114 G2N1&2G, 78-6219 IU2U2U1 &R &R, 78-6458 IVR, 78-6171 I1I, 78-6329 L B656 HHJ CMVI, 78-6267 L B656 HHJ EMV1, 78-6162, 78-6190, 78-6193, 78-6267 78-6291, 78-6334, 78-6492 L B666J, 78-6312 L B666J HHJ EZ, 78-6264 L B677 MV&T&J CO1 DO1 EO1 JMV1 NO1, 78-6222 L C666J, 78-6336 L D5 C555 A D- EU JUTJ AG AG BG IG JG KG, 78-6305 L D6 B666J, 78-6015, 78-6115, 78-6319, 78-6322, 78-6334 L D6 B666J C H J, 78-6317
--	---

L D6 B666J C J, 78-6015, 78-6037, 78-6162, 78-6314
 78-6316, 78-6317, 78-6318, 78-6343, 78-6351
 L D6 B666J J, 78-6015
 L D6 B666J J MF, 78-6015
 L D6 B666J J1 OF, 78-6015
 L D6 B666J J1 QF, 78-6015
 L D6 B666J J1 RF, 78-6015
 L D6 B666J J1E, 78-6313
 L D6 B6666 2AB TJ, 78-6242, 78-6302, 78-6324, 78-6326
 78-6327, 78-6328, 78-6329, 78-6330, 78-6332
 78-6333, 78-6334, 78-6335, 78-6440
 L D6 B6666 2AB TJ CQ HQ, 78-6326
 L E5 B666 FV OV MUTJ A E, 78-6028, 78-6333, 78-6348
 78-6544
 L E5 B666 FVTTT&J E OQ, 78-6028, 78-6037, 78-6345
 78-6348, 78-6349, 78-6544
 L E5 B666 FVTTT&J E OQ PQ, 78-6351
 L E5 B666 FX OV MUTJ A E KSV1 F-& CT5VOXTJ -
 A&K- B&AE
 78-6016
 L E5 B666 LUTJ A E FY&3Y OQ -B&AEFO, 78-6036
 78-6123, 78-6192
 L E5 B666 OV MUTJ A CQ E FV1Q FQ -B&ACEF
 78-6307
 L E5 B666 OV MUTJ A E FQ -B&AEF, 78-6343
 L E5 B666 OV MUTJ A1 E1 FV1 -B&AEF, 78-6346
 78-6347, 78-6355, 78-6544
 L E5 B666 OVJ A1 E1 FQ -B&AEF, 78-6343
 L E5 B666TJ A1 DQ E1 FY1&2VQ KQ OQ -A&OKO -
 B&AEFM &-NA-
 78-6590
 L E5 B666TJ A1 DQ E1FY2VQ HQ, 78-6590
 L E5 B666TTT&J E FQ F1U1 OQ, 78-6345, 78-6353
 L E5 B666TTT&J E FQ GQ OQ, 78-6037, 78-6349, 78-6351
 L E5 B666TTT&J E FQ OQ, 78-6037, 78-6346, 78-6348
 78-6349, 78-6350, 78-6351, 78-6355, 78-6544
 L E6 B666J C, 78-6323
 L E6 D6656 1A T&&&T&J R, 78-6106, 78-6169, 78-6217
 78-6265, 78-6283, 78-6312, 78-6319, 78-6320
 78-6321, 78-6322, 78-6326, 78-6334, 78-6490
 78-6493
 L G6 D6 B666J, 78-6169
 L5 AHJ-/G 6, 78-6167
 L566 1A LT&&J HNW, 78-6254
 L6TJ AG BG CG DG EG FG *ALPHA, 78-6271
 L6TJ AG BG CG DG EG FG *GAMMA, 78-6214
 L6TJ AMVNNO&2G D1, 78-6509
 L6UTJ A A B1U1Y&UZU1YU1VQ C -T, 78-6338, 78-6339
 L6UTJ A BL/U1Y&U2/ 2Q C C -T, 78-6004, 78-6039
 78-6338, 78-6592

L6V DVJ, 78-6299
 L6V DVJ B- C- E-/- AT3NTJ 3, 78-6298
 L66 BV EVJ, 78-6299
 L66J BMQ, 78-6247
 L66J BNCS, 78-6265
 L66J BNUNR B D& CQ, 78-6152
 L66J BNUNR B DNUNR B&& CQ, 78-6317
 L66J BNUNR B& CQ, 78-6152
 L66J BNUNR B1 D& CQ DSWO HSWO &-NA- 2, 78-6263
 L66J BNUNR& CQ, 78-6152
 L66J BNUNR& CZ, 78-6152
 L66J BOVM1, 78-6305
 L66J BSWQ ENUN 2 CR BQ DQ E1Q &-NA- 2, 78-6149
 L66J BZ, 78-6264, 78-6566
 L66J BZ ESWQ CNUNR D- 2 &-NA- 2, 78-6263
 L66J CZ, 78-6162, 78-6242, 78-6244, 78-6264, 78-6566
 L666 B6 2AB PJ, 78-6336
 NC1U1, 78-6021
 OCN1, 78-6103
 OCO, 78-6019
 ONN1&VO2, 78-6212
 ONN1&1, 78-6111, 78-6162, 78-6212, 78-6213, 78-6214
 78-6215, 78-6217, 78-6229, 78-6263, 78-6265
 78-6324
 ONN1&1R, 78-6273
 ONN2&2, 78-6162, 78-6216, 78-6218, 78-6242, 78-6265
 78-6417, 78-6492
 ONN2GVM2G, 78-6308
 ON1&U&1, 78-6106
 ON1&UN1OV1, 78-6189, 78-6295
 OOO, 78-6105
 OS1&R DOPS&O2&O2, 78-6284
 OS1&1, 78-6426, 78-6428
 PB CR-O2-Q2, 78-6130, 78-6304
 QR BQ DYQ1M1 -LQR BQ DYQ1M1, 78-6033
 QR DMV1, 78-6291
 QR DY2& 2, 78-6259
 QR DY2& 2U, 78-6027, 78-6051, 78-6315, 78-6342, 78-6355
 QVR, 78-6256
 QVR BQ, 78-6287
 QVYZIOV1UNN &10/11, 78-6199, 78-6200, 78-6212
 QVYZIR DN2G2G -L, 78-6051
 QVYZ2S2 -DL, 78-6162, 78-6205, 78-6206, 78-6212, 78-6216
 78-6265
 QVYZ2VQ &-NA- -DL, 78-6138
 QV1N1VQ2O 22, 78-6503

V10R BG DG, 78-6271, 78-6287, 78-6288, 78-6305
 V10R BG DG EG, 78-6287, 78-6288, 78-6305
 V10R DG B, 78-6287
 V1U1VQ -C, 78-6359
 V1XQVQ1VQ, 78-6589
 V2S2VQ, 78-6117
 V3R DN2G2G, 78-6051, 78-6196, 78-6209, 78-6211
 YR&VQ, 78-6252
 INUNO&1, 78-6107
 IYQYQYQYQVO & 2-FE, 78-6193
 2, 78-6032, 78-6108, 78-6375, 78-6549
 2K &Q, 78-6199
 2N2Q2Q, 78-6224
 78-6014, 78-6258
 B3 G6 E666 COT&&T&J F, 78-6325
 B656 EN HMJ FT B656 EN HMJ F, 78-6155
 B6566 B6/CO 4ABBC R BX HO PN DU GHT&&TTJ
 FQ JO1 P
 78-6293
 C3 F6 B666 DOT&T&J, 78-6325
 C566 DO LVOJ BO1, 78-6356
 C6 B5665 2AB S BX IN QN NU JH&&TTTJ FO1 IVH
 KVO1 KQ LOV1 M2 E-NT F6 E596 A BN
 LM&&TTJ NVO1 RQ R2, 78-6219
 C6 B5665 2AB S BX IN QN NU JH&&TTTJ FO1 I
 KVO1 KQ LOV1 M2 E- NT F6 E596 A BN
 LM&&TTJ NVO1 Q Q2, 78-6196
 C666 BKJ B EZ MZ &G &9/21, 78-6159
 C666 BN DNMV INJ B1YQYQYQ1Q L M -BBB
 78-6263
 C666 BN INJ E FZ LN1&1 &GH, 78-6200
 C666 BN ISJ EG B3- AT6N DNTJ D2Q, 78-6343
 C666 BNJ EG IMY&3N2&2 LO1, 78-6159
 C666 BO EV INJ D FZ N G- K-/VM-OT5-16- AN FVN
 IVN LVO PVM SVTJ G J KY N RY 2
 78-6304, 78-6347
 C666 BO EVJ DE FE IR BVO& LE MO NE &-NA- 2
 78-6153
 C755 A EOSO KUTJ AG AG BG FO JG KG LG
 78-6163
 D3 B556 BN EM JV MVTJT&J GO1 H1OVZ KZ L
 78-6232, 78-6267, 78-6358
 D36 I666 B6 2AB U EOT&&&&J, 78-6325, 78-6331
 E3 D5 C555 A D- FO KUTJ AG AG BG JG KG LG
 78-6111
 E3 D6 B6666 2AB U FOTT&&&&J HQ IQ, 78-6326
 78-6329
 F5 C6 B655 DOV GV OO QO RUT&&TTJ LO1, 78-6124
 78-6125, 78-6216, 78-6242, 78-6245, 78-6440

T F6 D5 C666 EM ON&&TTTJ HO1 SOVR CO1 DO1
 EO1& TO1 UVO1, 78-6321, 78-6490
 T3 F5 E666 1A R AX BOTJ IY&3Y J N QQ, 78-6362
 T3-26- BO FVO A&V DU JU LU NU PUTJ H TQ UVQ
 VQ C&Q RO- BT6OTJ CQ
 Z EQ F *, 78-6256
 T3NTJ A- 3PST3NTJ A- 3PS, 78-6226
 T3OTJ B- 2, 78-6325
 T30TJ BR, 78-6253, 78-6257, 78-6272
 T30TJ BIG, 78-6116
 T4OVTJ, 78-6242, 78-6245
 T5M CNJ A1 BSH, 78-6248
 T5MNJ, 78-6107
 T5MTJ, 78-6290
 T5N CSJ DNW B- CT5MVNTJ, 78-6490
 T5NNVJ A BR& DN1&1 E, 78-6214
 T5NNVJ A BR& E, 78-6292
 T5NTJ ANO, 78-6237
 T5OJ BYVZU1- BT5OJ ENW, 78-6227, 78-6244
 T5OSWTJ, 78-6242, 78-6245
 T5OV EHJ CQ DQ EYQ1Q, 78-6130, 78-6204
 T5VNVJ BR, 78-6299
 T50J BNW E1UNMVZ, 78-6227
 T55 AN CUTJ D1OVXQY&&YO1 FQ, 78-6238
 T55-12- 1A Q EN IOVY OVO BUTJ KU2 M NQ N1Q
 78-6216
 T56 BM DN FMVMVJ, 78-6229
 T56 BM DN FMYMVJ GUM, 78-6124
 T56 BM DN FN HNJ ISH, 78-6209
 T56 BM DN FN HNJ IZ, 78-6124
 T56 BM DN FNVNVJ F H, 78-6122, 78-6123, 78-6292
 78-6426
 T56 BM DNJ C- ET5N CSJ, 78-6256
 T56 BMJ D1YZVQ -L, 78-6201, 78-6203, 78-6372, 78-6567
 T56 BMJ D2Z GQ, 78-6318
 T56 BN DN FMYMVJ GUM D- BT50TJ CQ DQ E1Q
 78-6228
 T56 BN DN FNVNVJ B F, 78-6123
 T56 BN DN FNVNVJ B F H, 78-6120, 78-6121, 78-6122
 78-6123, 78-6426
 T56 BO DO CHJ G2U1, 78-6290
 T56 BO DO CHJ G3 H1O2O2O4, 78-6238
 T56 BOV GO IU&TJ FQ, 78-6586
 T56 BSWMVJ, 78-6280, 78-6281, 78-6282, 78-6283, 78-6294
 T56 BVNV GUTJ CSXGGG, 78-6159
 T6KJ A D- 2 &G &G &3/14 &3/17, 78-6184
 T6M DOTJ &GH, 78-6251

T6MPOTJ BO BN2G2G, 78-6051, 78-6162, 78-6194, 78-6195
 78-6196, 78-6197, 78-6232, 78-6267, 78-6406
T6MTJ, 78-6124, 78-6290
T6MVMVJ EF, 78-6509
T6MVMVVJ, 78-6233
T6N CN ENJ B- D- F-/- AT3NTJ 3, 78-6196, 78-6222
T6N CNJ BMSWR DZ& D F, 78-6264
T6N DNTJ ANO DNO, 78-6241
T6NJ C- BT5NTJ A, 78-6176, 78-6177
T6NJ C- BT5NTJ ANO, 78-6175, 78-6176
T6NJ CVZ, 78-6207, 78-6208
T6NJ DVMZ, 78-6163, 78-6264
T6NNVMVJ B- BT5OTJ CQ DQ E1Q -B&CD, 78-6051
T6NTJ ANO, 78-6240
T6NTJ ANO DG, 78-6240
T6NVMVJ EI A- ET5OTJ B1Q CQ -A&C, 78-6450
T6OTJ BO1NUNO&1 CQ DQ EQ F1Q-A&D -D, 78-6178
T6VMMVJ, 78-6120
T6VMMV FHJ F2 FR, 78-6106, 78-6109, 78-6115, 78-6167
 78-6193, 78-6217, 78-6257, 78-6319, 78-6320
 78-6326, 78-6331
T6VMMV FHJ F2 FR &-NA-, 78-6223, 78-6312
T6VMVNV FHJ D FY1UU3 F2U1 &-NA, 78-6335
T6VMVTJ E1YQ- BL6VTJ D F, 78-6396
T66 BM CHJ C C E HO2, 78-6328, 78-6333
T66 BM DN FN HNJ IS- ET5N ONJ DNW, 78-6209
T66 BNJ, 78-6239
T66 BNJ BO EMQ, 78-6212
T66 BNJ BO ENW, 78-6162, 78-6242, 78-6245, 78-6267
T66 BNJ EMY&3N2&2 IG, 78-6451
T66 BNJ ENW, 78-6244
T66 BNJ JQ, 78-6193
T66 BOVJ EQ D- 21, 78-6334
T66 B6 A B- C 1B I BN DN FN HNTJ, 78-6225
T66 CNJ, 78-6239
T67 GNV JN IHJ CG G KR, 78-6328
VHH, 78-6106, 78-6225
VH1, 78-6188
VH1G, 78-6112
WNMYUM&N1&NO, 78-6120, 78-6161, 78-6164, 78-6228
 78-6242, 78-6245, 78-6295
WNR BG ENW, 78-6331, 78-6333
WNR BQ C ENW &-NA-, 78-6157
WNR BQ ENW, 78-6396, 78-6590
WNR DOPS&O2&O2, 78-6305
WNR DYQY1QMVG -DL, 78-6052

WNR D1G, 78-6333
WSO&NUNR DN1&1 &-NA-, 78-6284
WSQQ &ZZ, 78-6182
WS1&O1, 78-6228, 78-6242, 78-6358
WS1&O2, 78-6108, 78-6120, 78-6161, 78-6162, 78-6216
 78-6244, 78-6245, 78-6359, 78-6360
WS1&O2 2U -C, 78-6051, 78-6163, 78-6232, 78-6424
ZH, 78-6042
ZM1, 78-6181
ZM2Q, 78-6182
ZN1&1, 78-6179, 78-6180
ZO1, 78-6160
ZQ, 78-6009, 78-6160, 78-6244
ZR, 78-6262, 78-6266
ZR B DNUNR B, 78-6278, 78-6286
ZR B DR B, 78-6297
ZR BG DR DZ CG, 78-6270, 78-6279
ZR BQ FVQ, 78-5044, 78-6311, 78-6567
ZR BV1, 78-6202
ZR CZ D, 78-6190
ZR DF, 78-6266
ZR DG, 78-6266
ZR DI, 78-6266
ZR DNUNR, 78-6265, 78-6278
ZR DNW, 78-6266
ZR DN1&1, 78-6284
ZR DR, 78-6162, 78-6260
ZR DR DZ, 78-6162, 78-6192, 78-6260, 78-6262, 78-6264
 78-6279, 78-6566
ZR DZ, 78-6269
ZR D1, 78-6266
ZR D1R DZ, 78-6261
ZVMZ, 78-6160
ZVN1&NO, 78-6120, 78-6164, 78-6213, 78-6215, 78-6228
 78-6229, 78-6230, 78-6308
ZVN2&NO, 78-6231
ZVO2, 78-6162, 78-6183, 78-6184, 78-6185
ZVR B CNW ENW, 78-6285
ZV4YZVQ, 78-6307
ZYUS, 78-6216, 78-6347
ZZ &QH, 78-6182
Z1, 78-6225
Z3M4Z, 78-6204
1MM1, 78-6162
1M1, 78-6290

1N1& 3PO1N1& 3PO, 78-6162

1N1&NUNR, 78-6214, 78-6221, 78-6223, 78-6266

1N1&R, 78-6262

1N1&R C DNUNR, 78-6267

1N1&R DNUNR, 78-6218, 78-6225, 78-6263

1N1&R DNUNR C, 78-6274, 78-6275, 78-6276, 78-6277

1OSWO1, 78-6228

1OV1UU1VO1, 78-6299

1UYVO1, 78-6372

1U1R, 78-6014, 78-6252, 78-6253

1U2U1, 78-6014

1VMR, 78-6292

1VO-HG-R, 78-6140

1X&&R BQ, 78-6184, 78-6206, 78-6218

1Y&MVR D1MM1, 78-6196, 78-6219

2N2&YUS&S 2, 78-6106

2OSWO2, 78-6159

2OVYU1, 78-6372

2OV1YVO2&SPS&O1&O1, 78-6305

20R DMV1, 78-6051, 78-6291, 78-6292, 78-6293

3XR&R&VO2N2&2 &GH, 78-6106

THE UNIVERSITY OF CHICAGO

சென்னை 18. 11. 1944

7405
U 16 710.12

Biol.

CARCINOGENESIS ABSTRACTS

DISCERNED COPY
VOL 16, NO 12

JUN 26 1979

VOLUME 16,
ISSUE 12



THE FRANKLIN INSTITUTE PRESS™

1871-1872

CARCINOGENESIS ABSTRACTS

A monthly publication sponsored by the
NATIONAL CANCER INSTITUTE

EDITOR

GEORGE P. STUDZINSKI, M.D., Ph.D.
COLLEGE OF MEDICINE AND DENTISTRY
OF NEW JERSEY, NEWARK

ASSOCIATE EDITOR

JUSSI J. SAUKKONEN, M.D.
JEFFERSON MEDICAL COLLEGE
PHILADELPHIA

NCI STAFF CONSULTANTS

ELIZABETH WEISBURGER, Ph.D.
JOAN W. CHASE, M.S.

Literature Selected, Abstracted, and Indexed
by

Information Services Department, Science Information Services Organization

BRUCE H. KLEINSTEIN, Ph.D., J.D., Assistant Director, Biomedical Projects

RUTHANN E. AUCHINLECK, Managing Editor

The Franklin Research Center
The Benjamin Franklin Parkway, Philadelphia, Pennsylvania 19103

All rights reserved. No part of this journal may be reproduced in any form, for any purpose or by any means, abstracted, or entered into any data base, electronic or otherwise, without specific permission in writing from the publisher.

Copyright © 1979 The Franklin Institute Press

CARCINOGENESIS ABSTRACTS

VOLUME 16, ISSUE 12

CONTENTS

	Cross Reference Abbreviations	Article Numbers	Page
VIEW	(Rev)	78-6601-78-6708	2531
CHEMICAL CARCINOGENESIS	(Chem)	78-6709-78-6802	2557
PHYSICAL CARCINOGENESIS	(Phys)	78-6803-78-6839	2576
AL CARCINOGENESIS'	(Viral)	78-6840-78-6977	2584
MUNOLOGY	(Immun)	78-6978-78-7068	2621
THOGENESIS	(Path)	78-7069-78-7148	2648
DEMOLOGY AND BIOMETRY	(Epid-Biom)	78-7149-78-7185	2667
CELLANEOUS	(Misc)	78-7186-78-7200	2674
THOR INDEX			2677
JECT INDEX			2687
CHEMICAL ABSTRACTS SERVICE REGISTRY NUMBER INDEX			2773
WESSER LINE NOTATION INDEX			2777

GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations or citations of significant carcinogenesis articles collected from the current major biomedical sources of world literature. This service is provided by the National Cancer Institute through a contract with the Franklin Research Center for preparation of the publication, under Contract No. NOI-CP-75885 with the National Cancer Institute, U.S. Department of Health, Education and Welfare. Published and distributed by the Franklin Institute PressSM.

SUBSCRIPTIONS

Published by The Franklin Institute Press, Box 2266, Philadelphia, PA 19103. Subscription rate for Volume 16, consisting of 12 monthly issues and a cumulative index is \$75.00, Domestic and \$90.00, Foreign. Check should be made payable to The Franklin Institute Press.
The Franklin Institute Press Co., Ltd., 3-10, Nihonbashi 2-Chome, Tokyo 103, Japan, is the exclusive distributor of CARCINOGENESIS ABSTRACTS in Japan.

1871-1872

GENERAL INFORMATION

CARCINOGENESIS ABSTRACTS makes available abstracts, annotations, or citations of significant cancer articles collected from the current major biomedical sources of the world literature. This service is provided by the National Cancer Institute through a contract with Franklin Research Center for preparation of the publication.

FREQUENCY OF PUBLICATION: *CARCINOGENESIS ABSTRACTS*, Volume 16, is published in 12 monthly issues with indexes, plus a cumulative index.

ABBREVIATIONS

JOURNAL names are abbreviated according to the *List of Journals Indexed in Index Medicus, Abbreviation Listing*. If the journal is not listed in this, abbreviations are derived from the *International List of Periodical Title Word Abbreviations*.

LANGUAGE of the article is indicated in parentheses after the title and is represented by a three-letter code. The source for these codes is *MARC Manuals Used by the Library of Congress*, pages 183-187.

ABBREVIATIONS used in abstracts:

A	angstrom(s)	mOsm	milliosmolar
ACTH	adrenocorticotrophic hormone	max	maximum
ADP	adenosine diphosphate	mEq	milliequivalent(s)
AMP	adenosine monophosphate	min	minute(s)
ATP	adenosine triphosphate	ml	milliliter(s)
approx	approximately	μl	microliter(s)
av	average	mm	millimeter(s)
BCG	bacillus Calmette-Guerin	mo	month(s)
bid	twice daily	mol wt	molecular weight
C	degree(s) centigrade	N	normal concentration
cal	calorie(s)	NAD	nicotinamide adenine dinucleotide
kcal	kilocalorie(s)	NADH	reduced nicotinamide adenine dinucleotide
cc	cubic centimeter(s)	NADP	nicotinamide adenine dinucleotidephosphate
Ci	curie(s)	NADPH	reduced nicotinamide adenine dinucleotidephosphate
mCi	millicurie(s)	NCI	National Cancer Institute
μCi	microcurie(s)	NIH	National Institutes of Health
cm	centimeter(s)	PAS	periodic acid-Schiff
CNS	central nervous system	po	orally
cpm	counts per minute	ppb	parts per billion
DNA	deoxyribonucleic acid	ppm	parts per million
ED ₅₀	median effective dose	qid	four times daily
EDTA	ethylenediamine tetraacetic acid	qod	every other day
g	gram(s)	QO ₂	oxygen quotient
kg	kilogram(s)	R	roentgen
mg	milligram(s)	RBC	red blood cells (erythrocytes)
μg	microgram(s)	RNA	ribonucleic acid
Hb	hemoglobin	rpm	revolutions per minute
hr	hour(s)	sc	subcutaneous
ia	intra-arterial	sec	second(s)
id	intra-dermal	SGOT	serum glutamic-oxaloacetic transaminase
gA	Immunoglobulin A	SGPT	serum glutamic-pyruvic transaminase
gB	Immunoglobulin B	soln	solution
gG	Immunoglobulin G	TCD	tissue culture dose
gM	Immunoglobulin M	TCD ₅₀	median tissue culture dose
LS	increased life span	tid	three times daily
m	intramuscular	UV	ultraviolet
p	intraperitoneal	WBC	white blood cells (leukocytes)
U	International Unit(s)	wk	week(s)
v	intravenous	wt	weight
km	Michaelis constant	X	times
.D	lethal dose	yr	year(s)
.D ₅₀	median lethal dose		
M	molar		
μM	micromolar		

Handwritten text in a narrow column on the left margin, possibly a page number or reference.

REVIEW

601 Screening Techniques for Identification of Carcinogens: An Overview. (Eng) Magee, P. N. Res. Inst., Temple Univ. Sch. Medicine, Philadelphia, (19140). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 404 pp.; 1935-1942; 1978.

ough none of the current screening techniques for identifying chemical carcinogens are sufficiently far developed to replace conventional long-term animal studies, several show promise. Among the screening procedures reviewed are those for carcinogenesis, precarcinogenesis, in vitro transformation or mutagenesis as end points and those that depend on specific indirect effects of carcinogens. (36 refs.)

602 Mutation and Latency in Carcinogenesis. (Eng) Knudson, A. G. (Inst. Cancer Res., 7701 Burdette Ave., Philadelphia, PA, 19111). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 209-225; 1977.

ent evidence relating somatic mutation and the phenomenon of latency to the origin of cancer is reviewed. A significant fraction of human cancer seems to involve somatic mutation. The chromosomal abnormalities observed in most cancers may in some instances represent this mutation, but in most cases these abnormalities are probably the result rather than the cause. Many environmental carcinogens increase spontaneous mutation rates that give rise to background cancer incidence. Some genetic disorders predispose an individual to environmentally induced carcinogenesis, and others create a high probability of the occurrence of site-specific cancers, even in the apparent absence of environmental carcinogens. The infrequent occurrence of neoplastic transformation of cells in the latter individuals suggests that another factor is required. So also do the phenomena of latency and dependence of cancer incidence. It is proposed that a carcinogen produces a first mutation in some cells, one of which then acquires a second mutation, which might be spontaneous or induced. When the first mutation is inherited, as in dominantly inherited predispositions, an environmental carcinogen would be expected to supply the second mutation. In this case, the latent period between exposure and clinically

apparent tumor should be reduced. The most satisfactory explanation for the age-dependence phenomenon is that the target cells for transformation are embryonal cells that differentiate and cease dividing. Any cancer mutations that occur after differentiation cannot be expressed. The age-specific incidence curves observed for the hereditary forms of these tumors directly reflect the median elapsed time for the occurrence of a second mutation, since the first mutation is present at the time of fertilization. This time is thus similar to the latent period observed following acute exposure to a carcinogen. (40 refs.)

78-6603 Hormonal Control of Tumor Growth. (Eng) Jensen, E. V. (Ben May Lab. Cancer Res., Univ. Chicago, Chicago, IL, 60637). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 209-225; 1977.

The relationship between steroid hormone receptors and the dependence of cancer of the breast and prostate gland on endocrine support is reviewed. Malignant tissues that depend on steroid hormones for their growth and function have a high level of receptor proteins with which the hormones combine to exert their biological effects. Some malignant neoplasms derived from hormone-dependent tissue retain the hormone dependence of the parent tissue, but others escape from hormonal control. Most hormone-dependent breast tumors can be distinguished from autonomous cancers by measuring their estrogen-receptor (estrophilin) levels. Because not all receptor-containing breast cancers are hormone-dependent, it appears that, in the case of neoplastic cells, the coupling between the synthesis of receptor protein and the need for hormonal stimulation in the nucleus is not as precise as it appears to be in normal tissues. Estrophilin levels can predict hormone dependence correctly in 85%-90% of breast cancer patients, but the fundamental criterion of hormone dependence appears to reside in the nucleus, where a simple test for susceptibility of the RNA polymerase system to stimulation by an estrogen-receptor complex might provide a more precise identification of responsive and nonresponsive cancers. Use of the progesterone receptor as a marker for the functionality of estrogen receptor in cancer cells is being evaluated. (33 refs.)

78-6604 Overview, Fact, Myth and Speculation. (Eng) Clayson, D. B. (Eppley Cancer Inst., Univ. Ne-

braska Medical Center, Omaha, NB, 68105). *J Environ Pathol Toxicol* 2(1): 1-8; 1978.

The validity of several assumptions regarding agents that may be considered carcinogenic to humans is questioned. All but 3 of the 20-50 agents known to be effective carcinogens in humans are also carcinogenic in animals. This indicates that between 6%-15% of transspecies extrapolations from humans to animals are discordant. The alternative approach, from the 1,000-1,500 known animal carcinogens to the human situation, is not possible because significant human exposure to many of these carcinogens may not have occurred or because epidemiologic studies may not have been performed. A new definition that would distinguish a true carcinogen from an agent that modifies the expression of another carcinogen is needed. Many of the methods for distinguishing between true carcinogens and modifiers already exist, such as microbial and mammalian cell mutation assays. Problems also exist in extrapolating from high doses, represented by the maximum tolerated dose administered to animals, to the low doses to which humans are exposed. The current view that there is no threshold dose for a carcinogen may also require modification. As more is learned about the mechanism of carcinogenesis, it may be possible to define certain critical rate-limiting steps in the process and to understand more clearly the likely shape of the dose-response curve at levels too low to follow economically by animal bioassay. (20 refs.)

78-6605 Problems of Assessing the Genetic Impact of Mutagens on Man (Meeting Abstract). (Eng) Newcombe, H. B. (Biology and Health Physics Div., Atomic Energy of Canada Ltd., Chalk River, Ontario, Canada). *Can J Genet Cytol* 20(3): 451; 1978. (no refs.)

78-6606 In Vitro Assay for Tumor Promoters and Anti-promoters. (Eng) Chang, C. C. (Dept. Human Development, Coll. Human Medicine, Michigan State Univ., E. Lansing, MI, 48824); Trosko, J. E.; Warren, S. T. *J Environ Pathol Toxicol* 2(1): 43-64; 1978.

The literature relating to the two-stage theory of carcinogenesis is reviewed, and an in vitro assay for tumor promoters and anti-tumor promoters is described. The two-stage theory postulates that carcinogenesis requires an initiation step and a promotion step. It has also been postulated that mutagenesis is responsible for promotion. The two-stage phenomenon has been demonstrated in several in vitro systems designed to measure transformation and mutagenesis in carcinogen-exposed cells that were posttreated with tumor promoters. Tumor promoters and anti-tumor promoters modify the induction of ornithine decarboxylase, the production of plasminogen activator, and the recovery of induced mutations. The recovery of mutagen-induced ouabain-resistant mutations in cultured Chinese hamster cells was increased by

treatment with 12-O-tetradecanoylphorbol-13-acetate (TPA), a tumor-promoter, and decreased by treatment with caffeine, an anti-tumor promoter. This suggests that many induced mutations can be repressed or derepressed by promoters or antipromoters. TPA also increased the recovery of UV-induced 6-thioguanine-resistant mutants. The results support the hypothesis that tumor initiation is due to a mutagenic event, but tumor promotion is the result of an epigenetic process involving cyclic nucleotide modulation of gene expression. (95 refs)

78-6607 Microsomal Enzyme Systems and Drug Toxicities. (Eng) Gillette, J. R. (Chemical Pharmacology Lab., Natl. Heart, Lung and Blood Inst., Bethesda, MD 20014). In: *Rat Hepatic Neoplasia: Workshop on Rat Liver Neoplasia held by William F. Merrell, Co. in Woods Hole, October, 1974*. William F. Merrell, Co. (Woods Hole, MA) 288 pp.; 100-113; 1978.

Toxic compounds that are converted in the body to chemically reactive metabolites that combine irreversibly with macromolecules such as DNA, RNA, protein, and glycogen are discussed. The induction of liver necrosis by halobenzenes illustrates many of the ways by which the severity of drug-induced tissue damage can be affected by changes in the formation or fate of chemically reactive metabolites. Bromobenzene-induced liver necrosis is caused by the covalent binding of a metabolite of bromobenzene formed by a cytochrome P-450 enzyme. The dose of bromobenzene affects the proportion of the dose that is bound to liver microsomes as well as the effects of phenobarbital pretreatment. The reason for the dose effect is related to the fact that liver cells have several ways of metabolizing bromobenzene. Information about the activation of chemicals in biological systems has also derived from studies with carbon tetrachloride. Although many drugs can be converted to chemically reactive metabolites and covalently bound to microsomal protein, it is not clear to what extent the covalent binding can be used to predict the incidence of various kinds of toxicity. (45 refs)

78-6608 Unscheduled DNA Synthesis of Human Cells as a Short-term Assay for Chemical Carcinogens. (Eng) Stich, H. F. (Cancer Res. Centre, Univ. British Columbia, Vancouver, B.C., Canada); San, R. H.; Lam, P. Koropatnick, J.; Lo, L. In: *Human Risk Assessment. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book C), 583 pp.; 1499-1512; 1977.

The use of unscheduled DNA synthesis of human cells as a short-term assay for chemical carcinogens is explored. The extent of DNA fragmentation and repair can be estimated accurately by several methods, but the alkaline sucrose gradient technique and the unscheduled incorporation of

H-thymidine appear to be the only ones suited for large-scale testing programs. Factors affecting DNA repair levels include hyperthermia, genetics, carcinogenic and noncarcinogenic chemicals, refractory period, and species. In tests of a possible correlation between the carcinogenicity of a compound and its ability to trigger DNA repair synthesis, only 1/104 carcinogens screened were not detected by the DNA repair assay. The possibility has been raised that various cell types can be used as target cells. This would offer the opportunity to examine the activation potential of various tissues and to link this property with the action of organotropic carcinogens. The DNA repair bioassay has also been used to study the effect of two chemical carcinogens applied simultaneously or in sequence. When 4-nitroquinoline-1-oxide and N-methyl-N'-nitro-N-nitrosoguanidine were applied sequentially or simultaneously, typically, an additive effect was observed at lower concentrations, an inhibitory effect at higher concentrations. This bioassay can also be used to detect defects at various steps of the repair chain, eg, in xeroderma pigmentosum cells. The DNA fragmentation and DNA repair bioassays should yield information about the validity of applying data from animals to humans or from cultured cells to in vivo conditions. (41 refs)

78-6609 Cancer: Somatic-Genetic Considerations. (Eng) Burnet, F. M. (Dept. Microbiology, Univ. Melbourne, 48 Monomeath Ave., Canterbury, Victoria, Australia). *Adv Cancer Res* 28: 1-29; 1978.

Current knowledge of somatic-genetic factors in the etiology of cancer is reviewed. (90 refs)

78-6610 General Properties of Cancer: Genetic Changes Group Report. (Eng) Harnden, D. G. (Dept. Cancer Studies, Univ. Birmingham, Birmingham B15 2TJ, England); Knudson, A. G.; Miller, R. W.; Brookes, P.; Cairns, J.; Croce, C. M.; Doerfler, W.; Fialkow, P. J.; Green, I.; Magee, P. N.; Muir, C. S.; Paterson, M. C.; Rajewsky, R. F.; Winnacker, E. L. In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 1977: 147-268; 1977.

Evidence surrounding the current genetic theories of cancer biology is reviewed. A number of agents, including environmental exposures of a chemical or irradiation nature in the workplace or elsewhere are known to cause genetic changes in human genomes. Viruses have been specifically implicated as being a transmissible agent in cervical cancer, Burkitt's lymphoma, nasopharyngeal cancer in the Chinese, and Hodgkin's disease early in life. These possible carcinogens may not

always act directly, but rather through intermediates arising from metabolic processes. A few cancers are known to be caused by germinal mutations, but these mutations may also only supply a predisposition to a cancer as in Xeroderma pigmentosum. Somatic mutations as well may be causative in some tumors. Most cancers are known to be karyotypically abnormal, but except for a small number of known markers (eg, the Philadelphia chromosome) these changes are not constant from tumor to tumor and may even occur differently within a single tumor. The valid question arises as to whether the observed chromosome abnormalities are the cause or the result of the tumor. The earliest origins of tumors are being intensely studied, and the glucose-6-phosphodiesterase marker system may be of value in this regard. If genetic changes are crucial to neoplastic change, it is important to learn whether these changes are always the same or whether a subset of the possible changes is sufficient to initiate neoplasia. Study must therefore be conducted in four areas: inherited genes that predispose to malignancy, integration of oncogenic virus genomes, specific chromosome alterations, and mutational events caused by the environment or some other yet unidentified factor. (97 refs)

78-6611 Tumor Initiation and Promotion. (Eng) Scribner, J. D. (Pacific Northwest Res. Foundation, Seattle, WA); Suss, R. *Int Rev Exp Pathol* 18: 137-198; 1978.

The molecular basis for the initiation and promotion of carcinogenesis is reviewed in depth. Cocarcinogenesis and the implications of the two-stage regimen for humans are also covered. (213 refs)

78-6612 Ataxia Telangiectasia: A Genetic Disorder Predisposing to Cancer and Exhibiting Hallmarks of Abnormal DNA Repair In Vitro (Meeting Abstract). (Eng) Paterson, M. C. (Atomic Energy of Canada Ltd., Chalk River Nuclear Labs., Chalk River, Ontario, KOJ 1J0, Canada). *Can J Genet Cytol* 20(3): 435; 1978. (10 refs)

78-6613 Environmentally Induced DNA Damage, Its Faulty Repair, and Malignant Genetic Diseases. (Eng) Paterson, M. C. (Biology and Health Physics Div., Chalk River Nuclear Labs., Chalk River, Ontario KOJ 1J0, Canada). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 39-53; 1977.

The carcinogenic implications of environmentally induced DNA damage and its faulty repair are reviewed. Damage to cellular DNA by physical and chemical carcinogens is a ma-

major factor in environmental carcinogenesis. Living organisms possess several enzymatic mechanisms for repairing this damage. In bacteria, three basic mechanisms have been identified and partially characterized: photoenzymatic repair, excision repair, and recombination repair. Similar mechanisms are also present in mammalian systems, such as human cells in culture. However, it appears that mammalian DNA damaged in vivo is not repaired completely. This is ascribed to (1) nuclear proteins masking a fraction of the damage for repair enzymes and (2) cell differentiation during development, leading to low repair abilities among certain cell types. In general, the lower the proliferative capacity of a cell type, the lower its ability to repair. In addition, damage may be repaired inaccurately, possibly by error-prone mechanisms. Thus, due to incomplete and inaccurate repair, defects may accumulate in DNA, and it is postulated that this accumulation of genetic damage plays a key role in neoplastic transformation induced by physical and chemical carcinogens. This postulate is supported by the association between defective DNA repair and cancer-proneness in humans with certain rare hereditary disorders (eg, xeroderma pigmentosum and ataxia telangiectasia). It is hypothesized that altered DNA predisposes a somatic cell to malignant transformation either by increasing the frequency of mutations and chromosomal changes or by promoting transformation by oncogenic viruses. Although this hypothesis is supported by experimental evidence, the molecular events underlying the emergence of the malignancy remain obscure. (15 refs)

- 78-6614 Xeroderma Pigmentosum and the Role of DNA Repair in Oncogenesis.** (Eng) Giannelli, F. (Paediatric Res. Unit, Prince Philip Res. Labs., Guy's Hosp. Medical Sch., London, SE1 9RT, England). *Bull Cancer (Paris)* 65(3): 323-334; 1978.

Xeroderma pigmentosum (XP) is discussed, with particular reference to its role as a model for actinic carcinogenesis. The light sensitivity of 80%-90% of XP patients can be explained by defects in the excision repair mechanism for reversal of DNA damage (pyrimidine dimers). Measurement of release of supercoils in DNA and the consequent reduction in the density of nucleoids has shown that strand incisions (the first step in excision repair) do not occur at normal rates in XP cells. Restoration of normal rates of DNA repair in XP patients by cell fusion has led to the conclusion that there are at least five complementation groups of the disease (A to E), of which A, C, and D are the most numerous and among which racial differences may exist. Groups A and E are complemented rapidly, group C more slowly, and group D very slowly. Ten to twenty percent of XP patients, known as XP-variants, are not defective in dimer excision but in repair synthesis of excised DNA. XP cells in culture are sensitive to many different chemical carcinogens and mutagens, which indicates that the DNA repair systems defective in XP have broad specificity and that the factor defective in XP recognizes a number of different lesions and initiates their repair by excision. This potential for chemical carcinogenesis may

help explain the presence of DNA abnormalities in cells exposed to sunlight in XP patients. XP should provide a good model for the study of human actinic carcinogenesis, because XP and normal cells seem to differ in their response to light more quantitatively than qualitatively, and the phenomena of XP can be fitted to the mutational, viral, immunologic theories of cancer etiology. (81 refs)

- 78-6615 Chemical-Biological Reactions Common to Teratogenesis and Mutagenesis.** (Eng) Hasegawa, R. D. (Dept. Pharmacology, Vanderbilt Medical Center, Nashville, TN, 37232). *Environ Health Perspect* 24: 87-101; 1978.

A review of the chemical-biological interactions that are common to the mutagenic and teratogenic responses to toxic chemicals is presented. The cytotoxic agents can be grouped as alkylating agents, electrophilic reactants, antimetabolites, intercalating agents, amino acid antagonists, spindle poisons, and additional chemicals that covalently bind to DNA. Alkylating agents and electrophiles react by attacking nucleophilic areas of biological molecules. Antimetabolites inhibit pathways of purine or pyrimidine biosynthesis and formation of thymidylic acid. Nucleic acid replication is inhibited by incorporation of these compounds into nucleic acids as base analogs. Intercalating agents insert between base pairs in DNA and interfere with transcription and replication. Amino acid antagonists inhibit protein and nucleic acid synthesis by interfering with specific amino acids required for protein or nucleic acid synthesis. Specific interference with asparagine, aspartic acid, glutamine, methionine, phenylalanine, and tyrosine utilization has been reported. Spindle poisons condense with microtubular protein and interfere with formation of cellular organelles and the spindle apparatus. Miscellaneous cytotoxins covalently bind with DNA and inhibit DNA synthesis and repair. Representative examples of these classes of cytotoxic agents are given and both specific and general reactions are outlined. (115 refs.)

- 78-6616 Relationship Between Mutagenesis, Carcinogenesis and Teratogenesis (Meeting Abstracts).** (Eng) Magee, P. N. (Fels Res. Inst., Dept. Pathology, Temple Univ. Sch. Medicine, Philadelphia, PA). *Mutat Res* 53: 224-225; 1978. (no refs)

- 78-6617 IARC Monograph Programme on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Preamble.** (Eng) IARC Working Group (Lyon, France). In: *Some N-Nitroso Compounds*. Magee, P. N., Chairman (International Agency for Research on Cancer, Geneva, Switzerland): *IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans*: Vol. 17, 365 pp; 1978.

criteria established by the International Agency for Research on Cancer/World Health Organization ad hoc working group for evaluating the carcinogenic risk of chemicals to humans are presented. (77 refs)

78-6618 Mutations, Chromosome Aberrations and Cancer. (Eng) Bishun, N. (Tissue Culture and Cytogenetics Unit, Res. Dept., Marie Curie Memorial Foundation, The Chart, Oxted, Surrey, England); Smith, N.; Williams, D. *Clin Oncol* 4(3): 251-263; 1978.

The correlation among mutagenesis, clastogenesis, and carcinogenesis produced by many commonly used compounds (drugs, cosmetics, pesticides, food preservatives, industrial chemicals, and oral contraceptives) is summarized. The systematic screening of chemicals for mutagenicity and chromosome-breaking activity has played a significant role in establishing the fundamental points of the chromosome theory of carcinogenesis. Recently, studies have been undertaken to establish correlations between the chromosome damage produced by carcinogenic chemicals and the molecular aberrations that may be involved in the induction of carcinogenesis. A subtle correlation was found between the mutagenic selectivities and carcinogenic potential of N-acetyl-2-aminofluorene and its metabolites. Unfortunately, almost nothing is known of the molecular effects of environmental chemicals on chromosomal material. However, it does not follow that a chemical that is mutagenic and produces chromosomal abnormalities is necessarily carcinogenic, although there may be similarities in those interactions with DNA that lead to chromosomal aberrations, mutagenicity, or carcinogenicity. Thus, far from finding a close correlation between mutagenesis and cancer development, it can be concluded that almost any chemically reactive compound, if tested by a sufficiently wide range of methods, is likely to prove mutagenic in some systems. Laboratory test procedures favored for drug safety evaluation are described. (18 refs)

78-6619 DNA Damage and Repair In Vivo. (Eng) Brash, D. E. (Dept. Biophysics, Ohio State Univ., Columbus, OH, 43210); Hart, R. W. *J Environ Pathol Toxicol* 1(1): 79-114; 1978.

Evidence implicating DNA damage in carcinogenesis is briefly summarized, and in vivo methods for studying DNA adducts and excision repair and strand breaks and strand-break repair are reviewed. Induction of DNA adducts by UV light, rays, and chemical carcinogens is usually monitored through the appearance of identified products in the DNA. Chromatography or immunological detection are the usual identification techniques. DNA-protein and DNA-DNA crosslinks are assayed as changes in DNA extractability, membrane filter adsorption, melting temperature, and maturation rate. Of the five steps involved in excision repair of DNA adducts, base cleavage, excision, polymerization,

and ligation have been studied in vitro and in vivo; incision has only been studied in vitro. Single-strand breaks in cellular DNA in situ can arise from x- or γ -radiation, nicking during excision repair of chemical carcinogens or UV light, or replication past a damaged site. Induction and repair of strand breaks have been studied in vivo by alkaline sucrose gradient sedimentation, DNA polymerase initiation sites, hydroxypapatite chromatography, and S₁ nuclease digestion. Postreplication repair has not been studied in vivo, and photoreactivation of cyclobutane-type pyrimidine dimers in vivo has not been demonstrated. (209 refs)

78-6620 DNA Repair Pathways: In Bacteria and in Mammalian Cells (Meeting Abstract). (Eng) Hanawalt, P. C. (Dept. Biological Sciences, Stanford Univ., Stanford, CA, 94305). *Can J Genet Cytol* 20(3): 433; 1978. (5 refs)

78-6621 Theoretical Mechanisms for Synthesis of Carcinogen-induced Embryonic Proteins: III. The tRNA Methylases: Methylation Mechanism and Function. (Eng) Hancock, R. L. (High River Inst. Theoretical Cancer Study, High River, Alberta, Canada); Hancock, C. D. *Med Hypotheses* 4(5): 497-515; 1978.

It is hypothesized that redundant repressed transfer RNA (tRNA) methylase genes that are active in embryonic states can be derepressed (induced) with carcinogens in adult cells. Supportive evidence, especially from molecular model building, is presented for a mechanism of tRNA methylation. In this mechanism, the adenine moiety of S-adenosyl-L-methionine base-pairs with uracil for all tRNA methylations. These methylations may be required during transcription of a tRNA molecule in order to limit the number of possible conformers that can occur before the molecule develops further. Fragments of tRNA were suitable substrates for enzymatic methylation; however, a subtle treatment such as UV irradiation destroyed the ability of intact tRNA to act as a substrate. It is believed that the tRNA fragments necessarily have little secondary or tertiary structure, which allows the base uracil of tRNA to pair with the adenine portion of S-adenosyl-L-methionine. This theory of the function for methyl groups is discussed in light of the properties of hypomethylated tRNA's. (39 refs)

78-6622 The Role of Specific Nucleic Acid Reactions and Their Repair. (Eng) Brookes, P. (Inst. Cancer Res., Pollards Wood Res. Station, Chalfont St. Giles, Bucks HP8 4SP, England); King, H. W.; Newbold, R.; Osborne, M. R. In: *Characterization and Treatment of Human Tumours. Proceedings of the Seventh International Symposium on the Biological Characterization of Human Tumours held in Budapest, April 13-15, 1977*. Davis, W.; Harrap, K.

R., eds. (Amsterdam: Excerpta Medica): Advances in Tumour Prevention, Detection and Characterization. Vol. 4, 332 pp.; 58-62; 1977.

The mechanisms by which methylating agents and polycyclic hydrocarbons react with DNA are reviewed. Some methylating agents (ie, N-methyl-N-nitrosourea) react with deoxyguanosine to form O⁶-methyldeoxyguanosine. It was suggested that the O⁶-methylguanine moieties in DNA would base-pair with thymine and lead to transition mutations. The extent of tumor formation due to N-methyl-N'-nitro-N-nitrosoguanidine correlates with the extent of O⁶-methylation of guanine. In addition, tissues able to excise O⁶-methylguanine from their DNA have lower tumor incidences following treatment with a carcinogen than tissues that cannot perform this function. According to studies of polycyclic hydrocarbons, an epoxide metabolite of benzo(a)pyrene (BP) and one of 7,12-dimethylbenz(a)anthracene reacts with the 2-amino group of guanine. It is not known why hydrocarbon derivatives react with amino groups that are never attacked by classical alkylating agents. Experiments with the diol epoxide and K-region epoxide of BP indicated that the diol epoxide was the more potent mutagen of the two and that it was able to induce a high frequency of mutations at nontoxic dose levels. The two examples considered suggest that cellular DNA-repair mechanisms may ultimately determine the biological consequences of DNA-carcinogen reactions. (23 refs)

78-6623 Biochemical Aspects of Chemical Carcinogenesis. (Eng) Gielen, J. E. (Laboratoire de Chimie Medicale et de Toxicologie, Institut de Pathologie, Universite De Liege, B4000 Sart Tilman par Liege 1, Belgium). *Bull Cancer (Paris)* 65(3): 249-254; 1978.

A brief review is presented of the carcinogenic abilities of polycyclic aromatic hydrocarbons (PAH), with particular reference to the enzymatic systems necessary to metabolize these carcinogens, the nature of the reactive metabolites, and the cellular targets of the active compounds. PAH are highly lipophilic, and must be made more polar (hydrosoluble) in order to be eliminated from the body. Numerous enzymes and enzyme systems such as the cytochrome P450 monooxygenases, epoxide hydratase, and glutathione-S-transferase have the ability not only to detoxify and eliminate these compounds but also to act as activating agents for them when the reaction product is more active than the parent compound. The activated compounds of PAH are epoxides and dihydrodiol epoxides, which are susceptible to reaction with cellular macromolecules. The relative basal activities (influenced by physiological parameters) of these enzymes determines the intracellular concentration of the carcinogens and are more important therefore than the absolute activity, which can be modified by administration of various chemicals. The compound most likely to be the ultimate carcinogen following benzo(a)pyrene exposure is 7,8-dihydrodiol-9,10-epoxide benzo(a)pyrene. For other unsubstituted PAH, electrophilic benzylic carbonium ions should most easily appear when

located in the so-called bay region of the molecule. Because permanent alteration of the cell results from the action of these carcinogens, it is postulated that nucleophilic centers of macromolecules are subject to electrophilic attack by these compounds. Although complexes between nucleosides and 7,8-dihydrodiol-9,10-epoxide benzo(a)pyrene have been demonstrated in vitro, this phenomenon has not yet been seen in vivo. (45 refs)

78-6624 Characteristics of Excision Repair after High Doses of UV and Chemicals to Human Cells. (Meeting Abstract). (Eng) Setlow, R. B. (Biology Department, Brookhaven Natl. Lab., Upton, NY, 11973). *Can J Genet Cytol* 20(3): 436; 1978. (4 refs)

78-6625 The Importance of the Formation and Persistence of Alkylated Bases in DNA in the Carcinogenicity of N-Nitroso Compounds (Meeting Abstract). (Eng) Pegg, A. E. (Dept. Physiology, Pennsylvania State Univ. Medical Sch., Hershey, PA, 17033). *Can J Genet Cytol* 20(3): 435-436; 1978. (3 refs)

78-6626 Two-Stage (Initiation-Promotion) Carcinogenesis In Vivo and In Vitro. (Eng) Chouroulin, I. (Institut de Recherches Scientifiques sur le Cancer, C.N.R.S., BP8, F 94800, Villejuif, France); Lasne, C. *Cancer (Paris)* 65(3): 255-264; 1978.

Current progress in the in vitro and in vivo investigation of two-stage carcinogenesis is reviewed. In vivo initiation and promotion is each an incomplete carcinogenic action; nevertheless almost all initiators are carcinogens. The promoter, on the other hand, must be irritants capable of inducing hyperplasia. Because the initiation stimulus is transmitted from cell to cell over time, it must involve cellular genetic mechanisms. Promotion, which is reversible, seems to be related to stimulation of mitosis and protein synthesis. For tumor yield should be quantitatively related to the initiator dose, while promoter effectiveness should dictate the speed of tumor development. Whereas these phenomena have been successfully demonstrated using mouse skin, difficulties have been encountered in other systems. A successful in vivo demonstration using rat fibroblasts is reported. Initiators tested were 7,12-dimethylbenz(a)anthracene (DMBA) (1 µg/ml of medium for 24 hr), benzo(a)pyrene (BaP) (1 µg/ml of medium for 6 hr), and benzo(e)pyrene (BeP) (1 µg/ml of medium for 6 hr). Promoters included 12-O-tetradecanoyl-phorbol-13-acetate (TPA) (0.1 µg/ml of medium at each passage) and croton-oil (CrO) (0.01 µg/ml of medium at each passage). Transformation criteria included acceleration of transformation of 20% of cells (by which criterion BaP and BeP followed by CrO were most effective) and acceleration to mutation in injected animals (by which criterion DM

followed by TPA was effective and BaP followed by CrO (most effective). BeP followed by CrO did not enhance carcinogenesis. These results demonstrate effective two-stage carcinogenesis in vitro, and have implications for the occurrence of human lung cancer, where an initiation-promotion model can be hypothesized. (50 refs)

78-6627 **Benzene--Attempts to Establish a Lower Exposure Standard in the United States. A Review.** (Eng) Zenz, C. (2418 Root River Parkway, West Allis, WI, 53227). *Scand J Work Environ Health* 4(2): 103-113; 1978.

The cause-and-effect relationship between benzene exposure and blood abnormalities, including certain leukemias, is reviewed. The National Institute for Occupational Safety and Health has recommended that occupational exposure to benzene be controlled to < 1 ppm in air. The current standard is 10 ppm. (57 refs)

78-6628 **Nitrosamines in the Atmosphere.** (Ger) Lohs, K. (Sitzungsberichte der Akademie der Wissenschaften der DDR, Berlin, E. Germany); Rolle, W.; Gehlert, H. *Akademie der Wissenschaften* (2): 99-105; 1978.

The formation, toxicity, and volatility of dialkyl nitrosamines (DAN) are reviewed. A recent study of the effect of dimethylnitrosamine in rats led to the conclusion that the formation of O-6-methylguanine is more closely associated with the carcinogenic activity of DAN than alkylation of the N-7 of guanine, as was previously thought. Nitrosamine analysis has become much more precise since the development of the thermogravimetric analyzer. Recent work has shown that diethylnitrosamine and diethylammonium nitrate can be rapidly formed in the gas phase, from NO₂ and diethylamine, without the influence of UV light. Review of the literature indicates that at 10⁻⁸ mole of nitrosamine/m³ of inspired air is dangerous to humans and also that industrial sources are releasing amounts of DAN that could quickly lead to dangerous levels. (2 refs)

78-6629 **Mycotoxins.** (Rus) Kravchenko, L. V. (Lab. Enzymology, Inst. Nutrition, Moscow, USSR); Matel'ian, V. A. *Zh Vses Khim O-Va D.I. Mendeleeva* 23(4): 400-405; 1978.

A review is presented summarizing literature pertaining to the physico-chemical properties, toxicity, and carcinogenic activity of a series of most common mycotoxins. Such mycotoxins as aflatoxins, sterigmatocystin, patulin, penicillic acid, ochratoxin, and some metabolites of Alternaria have very strong carcinogenic activity. (198 refs)

78-6630 **Oncogenesis and Vitamin A.** (Rus) Pletsityi, K. D. (Lab. Nonspecific Resistance Organism Immunity, Inst. General Pathology, Pathological Physiology, Moscow, USSR). *Vopr Onkol* 24(10): 85-92; 1978.

A review is presented of the literature concerning the role of vitamin A in tumor resistance. Experimental and clinical data has proved the association between the vitamin A deficiency and the incidence of malignant tumors. Vitamin A deficiency was shown to enhance the carcinogenic effect of benzo(a)pyrene, while administration of vitamin A or its derivatives inhibited the growth of carcinogen-induced tumors. (72 refs)

78-6631 **Inorganic Carcinogens.** (Eng) Rawson, R. W. (Univ. Texas System Cancer Center, Houston, TX). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1893-1906; 1978.

The role of trace metals such as As, Cd, Cr, Ni, Pb, and Be, most of which are nonessential, and asbestos in the genesis and growth of various cancers is reviewed. There is evidence that the presence of an excess amount of certain nonessential elements in an organ or in the whole organism may promote the development of cancers of the lung, gastrointestinal tract, and skin. (66 refs)

78-6632 **Nasopharyngeal Carcinoma.** (Eng) Andersson-Anvret, M. (Dept. Tumor Biology, Karolinska Inst., Stockholm, Sweden); Forsby, N.; Klein, G. *Prog Exp Tumor Res* 21: 100-116; 1978.

The association between Epstein-Barr virus (EBV) and nasopharyngeal carcinoma (NPC) is reviewed. Epidemiological studies suggest that there are intrinsic, probably genetic, cofactors in the etiology of NPC. At least four antigens, or complexes of antigens, have been shown to be EBV-determined: viral capsid antigen (VCA), membrane antigen (MA), early antigens (EA), and complement-fixing soluble antigen (CF/S) and nuclear antigen (EBNA), which are probably identical. Patients with NPC have high antibody titers against VCA, MA, EA (both restricted and diffuse components), and CF/S. EBNA has been found in the epithelial cells of NPC tumors, and there is a complete correlation between the expression of EBNA and detection of EBV DNA in the cells. A comparison of the antibody reactivities indicates that anti-VCA titers are relatively stable over a lifetime, but anti-EA is the most disease-related antibody. It is usually only present in patients with an acute lymphoproliferative, EBV-induced disease. NPC patients have increased IgA antibody levels; however, the EBV-specific IgA antibodies to

VCA and the diffuse component of EA decline to nondetectable levels after treatment of the tumor. The EBV DNA found in NPC tumors is present in the epithelial cells. EBV DNA is only found in the undifferentiated form of the tumor. Different strains of EBV may exist in tumors from different geographical areas. (94 refs)

- 78-6633 **Carcinogenic Mode of Action of Tissue-embedded Foreign Particles.** (Ger) Brand, G. (Dept. Microbiology, Univ. Minnesota Medical Sch., Mayo Box 196, Minneapolis, MN, 55455). *Zentralbl Bakteriol (Orig B)* 166(2/3): 159-168; 1978.

Foreign bodies such as asbestos fibers, schistosoma eggs, or artificial implants are known to cause cancer. Studies of foreign body tumorigenesis in mice have revealed a specific sequence of preneoplastic events: (1) loss of macrophage phagocytic activity; (2) fibrous conversion of the foreign bodies; and (3) during the fifth to sixth month following implantation, contact between the preneoplastic cells and the foreign body surface. The size, shape, and surface properties of the foreign bodies (chemically inert glass or plastic films) were also of decisive importance to the neoplastic process. Specific antigens and indications of immunologic surveillance were not detected. These findings may facilitate the understanding not only of asbestos carcinogenesis and the cocarcinogenic role of smoking, but of the cancer phenomenon in general. (36 refs)

- 78-6634 **Occupational Lung Disease.** (Eng) Jones, R. N. (Tulane Medical Center, New Orleans, LA); Weill, H. *Respiratory Care* 23(10): 989-998; 1978.

Occupational lung disorders are reviewed with regard to their causative agents and the pathologic responses evoked. Dust macules (which are caused by inhaling tin, barium, iron, or coal salts), pulmonary fibrosis, hypersensitivity pneumonia, beryllium lung disease, and obstructive airway diseases are discussed. Asbestos, arsenic, chromates, nickel, radioactive inhalants, the halo bischloromethyl ether, chloromethyl methyl ether, and vinyl chloride monomer may induce respiratory tract tumors. Asbestos may also cause cancer of the larynx and carcinoma of the gastrointestinal tract, and brain tumors and hemangiosarcoma of the liver may result from exposure to vinyl chloride. The diagnosis of occupational lung disease is discussed, as is the setting and enforcement of permissible dust exposure levels. (10 refs)

- 78-6635 **Cancerogenesis, Teratogenesis, Mutagenesis: Correlation Between Animal Experiments and Clinical Findings.** (Ger.) Froberg, H. (Institut für Toxikologie, E. Merck, Frankfurter Strasse 250, 6100 Darmstadt 1, W. Germany). *Arzneim Forsch* 28(11a): 1984-2001; 1978.

Carcinogenicity data from animal tests, epidemiological studies and clinical experience with humans, and mutagenicity tests are reviewed. A good correlation among in vivo carcinogenicity test results, teratogenic effects, and mutagenic effects has been found for alkylating compounds and for compounds that can be metabolized into electrophilic agents. To develop a quick test for carcinogenicity, the mutagenic effects of carcinogens on several bacteria and yeasts have been studied. *Salmonella typhimurium* (Ames test) is considered the most suitable organism for such a quick test. The false-negative rate ranges from 7%-25%. The highest false-negative rate came from a study in which many carcinogenic compounds were tested. In this study, several metals that were not carcinogenic in animals, MoVI, PtII, RhI, and RuII, gave positive results. Thus, the Ames test as currently performed is not suitable for testing the carcinogenicity of metals. With microsomal activation, the Ames test gave a 9% error rate when tested on known N-nitrosamines; for polycyclic aromatic hydrocarbons, the error rate was only 2%. Ames and coworkers have assumed that there is a quantitative relationship between mutagenicity and carcinogenicity, but other workers have not confirmed this relationship. It appears that only substances that can be converted to reactive electrophilic compounds show a positive correlation between mutagenicity and carcinogenicity. Possible reasons for a lack of correlation with other compounds include extreme volatility, lack of activating substances, and rapid reaction of the compound with metabolites present in *S. typhimurium*. (11 refs)

- 78-6636 **Structural Prognostication of Carcinogenicity and Tumor-Enhancing Activity in Various Organic Chemicals.** (Eng) Van Duuren, B. L. (Lab. Organic Chemistry, New York Univ. Medical Center, New York, NY 10016). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2 (part 1), 2404 pp.; 2071-2084; 1978.

Structure-activity relationships of direct-acting alkylating agents, tumor promoters, and cocarcinogens are reviewed. Among the 100 alkylating agents examined were epoxides (mono-, bi- and polyfunctional), β - and γ -lactones, and a variety of chloro ethers. The β -lactones are carcinogenic but the γ -lactones are not. The bifunctional epoxides exhibit carcinogenicity more frequently than the monofunctional analogs. In monofunctional agents, the presence of a reactive adjacent functional group results in carcinogenic activity. Trichloroethylene (TCE) and vinyl chloride (VC) may be metabolized via an epoxide intermediate. TCE is carcinogenic in mice but not in rats, and VC is a known human carcinogen. Because of the wide differences in chemical structure, reactivity, and physical properties of tumor promoters, cocarcinogens, and tumor inhibitors, they probably exert their a-

ities in several different ways. They may simply alter the rate of absorption and disappearance of a carcinogen, or they may alter its metabolic pathway. Tumor-promoting agents, particularly the phorbol esters, are likely to interact at the cell membrane. As a result of structure-activity studies, it has become possible to assign biological activity to certain classes of compounds that have not yet been tested. (41 refs)

- 78-6637 **Nutritional Modulation of Carcinogenesis.** (Eng) Newberne, P. M. (Dept. Nutrition Food Science, Massachusetts Inst. Technology, Cambridge, MA); Rogers, A. E.; Gross, R. L. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer in New York, NY: Vol. 1(part 1): 1193 pp.; 665-691; 1977.

Based on epidemiologic studies that clearly reveal a change in the incidence of some forms of cancer when populations move from one geographic location to another and change in habits, evidence has accumulated to indicate that nutrition is associated with resistance or susceptibility to cancer. The following positive correlations between cancer in five major target organs and environmental risk factors have been conclusively established: (1) esophagus: heavy consumption of cigarettes and alcohol; wheat bread and tea consumption; (2) stomach: low socioeconomic status; consumption of salt-smoked, or fermented foods; decreased consumption of vegetables; (3) colon: increased consumption of meats, saturated fats, and highly refined carbohydrates with low dietary fiber and decreased vitamin A; (4) liver: decreased dietary lipotropes and increased dietary fat; consumption of toxic substances such as alcohol and mycotoxins and modified microsomal enzymes; (5) breast: higher socioeconomic status, with increased consumption of animal fat; obesity, or, more likely, both increased weight and height; potential increases in estrogen synthesis. It is hoped that through continued study, nutrition may soon be used as a major modifier in efforts toward prevention of cancer. (76 refs)

- 78-6638 **Dietary Effects on Chemical Carcinogenesis in the Livers of Rats.** (Eng) Rogers, A. E. (Lab. Experimental Pathology, Dept. Nutrition and Food Science, Massachusetts Inst. Technology, Cambridge, MA, 02139). *Rat Hepatic Neoplasia: Workshop on Rat Liver Neoplasia* edited by William F. Merrell, Co. in *Woods Hole, October, 1974*. William F. Merrell, Co. (Woods Hole, MA): 288 pp.; 1978. 225-265; 1978.

The effects of different types of diets and of specific nutrients and additives on the development of chemically induced hepatic tumors in rats are reviewed. Natural-ingredient diets provide some degree of protection against hepatic carcinogenesis, although the mechanism by which they do so is not known.

Lipotropic factors (choline, methionine, folic acid), with the possible exception of vitamin B₁₂, protect rats against the induction of hepatic tumors by a number of chemicals, and there is evidence of direct biochemical interaction between lipotropes and carcinogens. Vitamin A protects against the induction of gastrointestinal neoplasia by aflatoxin B₁ and other carcinogens, but it does not consistently affect the induction of hepatic tumors by aflatoxin. Dietary effects on the toxicity of carcinogens may or may not correlate with their effects on carcinogenicity. Dietary manipulation offers a valuable tool for exploring the mechanisms of action of chemical carcinogens. (no refs)

- 78-6639 **Experimentally Produced Brain Tumors (Meeting Abstract).** (Eng) Wechsler, W. (Dept. Neuropathology, Univ. Dusseldorf, Dusseldorf, W. Germany). *J Neuropathol Exp Neurol* 37(5): 575; 1978. (no refs)

- 78-6640 **The Thyroid, Prolactin, and Breast Cancer.** (Eng) VanderLaan, W. P. (Scripps Clinic and Res. Foundation, 10666 N. Torrey Pines Road, La Jolla, CA, 92037); Larson, B. A. *Arch Intern Med* 138(11): 1611-1612; 1978.

The role of thyroid dysfunction and prolactin secretion in the etiology of breast cancer is reviewed. Breast cancer growth in rats can be stimulated by iodine deprivation, and rat mammary gland dysplasia may be induced by thyroid-stimulating hormone (TSH) acting directly on the breast. Prolactin is influential in murine mammary tumorigenesis. In breast cancer patients, the findings of altered TSH and prolactin responses to thyrotropin-releasing hormone may be part of a general pituitary dysfunction in breast cancer. (10 refs)

- 78-6641 **Correcting a Bias?** (Eng) Hutchison, G. B. (Harvard Univ. Sch. Public Health, Boston, MA, 02115); Rothman, K. J. *N Engl J Med* 299(20): 1129-1130; 1978.

An attempt to compensate for the presumed greater detection of endometrial cancer among estrogen users by choosing as a comparison group women with benign gynecologic conditions only adds to the accumulating evidence that exogenous estrogens induce endometrial cancer. In other words, it too introduces a selection bias. (4 refs)

- 78-6642 **Menopausal Estrogens and Endometrial Cancer.** (Eng) Weiss, N. S. (Dept. Epidemiology, Univ. Washington and Fred Hutchinson Cancer Res. Center, Seattle, WA, 98195). In: *Estrogens and Cancer. Proceedings*

of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977. Colorado Regional Cancer Center (Denver, CO): 197 pp.; 27-34; 1978.

The relationship between endometrial cancer and the menopausal use of estrogens is discussed. The incidence of endometrial cancer in the United States rose sharply soon after the rapid increase in estrogen use among American menopausal women, whereas the incidence rate in England, where noncontraceptive estrogen use has been much less common, is lower than in the United States and has not increased during the same period. Five recent studies have shown that menopausal women with endometrial cancer had a greater exposure to exogenous estrogens than did controls. The incidence in users was estimated to be 3-10 times that in nonusers. Estrogen use has also been found to predispose to endometrial hyperplasia; women with this condition have an increased risk of carcinoma. Other evidence suggesting a causative role for the menopausal use of estrogens includes: (a) conditions in which endogenous estrogen levels are elevated or are not countered by progesterone secretion that predispose to endometrial cancer, (b) the use of noncontraceptive estrogens in the treatment of gonadal agenesis and breast cancer that leads to an increased incidence of endometrial cancer, and (c) endometrial proliferation as a prominent effect of estrogenic hormones. Other evidence suggests that if low-dose, short-duration estrogen therapy is used, the increased risk of endometrial cancer will be less than the average increase found in most of the recent studies. It is suggested that case control and cohort studies should be employed together in future studies of the relationship between noncontraceptive estrogen use and the development of endometrial cancer. (34 refs)

78-6643 Hormonal Carcinogenesis: A Novel Hypothesis for the Role of Hormones. (Eng) Nandi, S. (Dept. Zoology, Univ. California, Berkeley, CA, 94720). *J Environ Pathol Toxicol* 2(1): 13-20; 1978.

Information related to the role of hormones in carcinogenesis is reviewed as background for a hypothesis on hormonal carcinogenesis. The hypothesis is based on the following findings: (1) hormones regulate cell division in normal cells; (2) successful neoplastic transformation requires hormonal stimulation of cells receiving a carcinogenic insult; and (3) normal cells, when present in a high ratio, inhibit the growth of neoplastic cells. According to the hypothesis, hormones are neither mutagenic nor carcinogenic. They play a dual role in carcinogenesis induced by direct-acting carcinogens, such as viruses, chemicals, and radiation. This dual role is postulated to be (1) the fixation of neoplastically transformed cells, and (2) control of the latent period for the appearance of overt tumors by regulating the growth and ultimate death of normal cells. This would alter the ratio of normal to neoplastic cells in favor of the neoplastic cells, which would, in turn, facilitate the growth of tumor cells. This hypothesis is testable in the mammary tumor system. (40 refs)

78-6644 Use of Hormones and Human Tumorigenesis. (Eng) Lewison, E. F. (Johns Hopkins Hospital, Baltimore, MD). In: *Third International Symposium on Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer, New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 609-616; 1977.

The relationship between the long-term use of exogenous hormones and the development of malignant and benign tumors of the breast, endometrium, vagina, and liver is reviewed. Studies showing that estrogens act directly on normal breast tissue cells by promoting growth and differentiation and that there is a high level of positive estrogen receptor in the cytoplasm of about one-third of all breast cancer patients suggest that estrogen is involved in the etiology of this disease. A significant fact is that >99% of the patients are female. However, epidemiologic studies of the relationship between breast cancer and estrogens or oral contraceptives have yielded inconclusive results. Recent studies have shown conclusively that the prolonged use of estrogens by postmenopausal women is correlated with a marked increase in the risk of developing endometrial cancer. Cancer of the vagina, a relatively rare disease, has been linked to the ingestion during pregnancy of nonsteroid estrogens, including diethylstilbestrol and its various analogs, by the mothers of the patients. Primary liver tumors, liver adenomas, and liver hyperplasia have all been shown to develop with increased incidence in young women taking oral contraceptives. In determining the ultimate risk of developing malignancies in all of these situations, one must take into account the exact type of synthetic estrogen used as well as the duration of use. (24 refs)

78-6645 Pathologic Aspects of Tumors and Tumor-like Lesions in Women Taking Oral Contraceptives and Hormones. (Eng) Fechner, R. E. (Dept. Pathology, Univ. Virginia Sch. Medicine, Charlottesville, VA, 22903). In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977*. Colorado Regional Cancer Center (Denver, CO): 197 pp.; 111-145; 1978.

The pathology of liver, breast, and genital tract tumors and other related lesions in women taking oral contraceptives is reviewed. In addition, studies on the incidences of these tumors are reviewed. The largest group of tumors that appear to be associated with hormones is in the liver. Liver cancer, adenoma is composed of hepatocytes lined by Kupffer cells and with nuclear variation more diverse than that seen in normal liver; it contains no bile ducts or ductules. Bile ducts do occur in focal nodular hyperplasia, and this feature is the crucial diagnostic difference between the two conditions. Gross and microscopic features of these conditions do not differ in hormone users from those in nonusers. However, some evidence suggests that there may be a quantitative increase in tumor vascular changes in women taking oral contraceptives.

ceptives, as compared with those in women not taking hormones. Hepatocellular carcinoma is also discussed. Most investigators are inclined to agree that oral contraceptives could not be incriminated in the pathogenesis of cervical dysplasia on the basis of the evidence thus far available. The endometrium is the uterine tissue most frequently altered morphologically by oral contraceptives. The changes are due mainly to progestin, which may produce a conspicuous decidual reaction, while the glands become atrophic (glandular-stromal dissociation). Endometrial adenocarcinoma has been illustrated in an endometrium that exhibited stromal-glandular dissociation. The histologic appearance of endometrial carcinomas has been unusual in two respects. One study demonstrated an excess of secretory and clear-cell carcinoma in hormone users when compared with young women with endometrial cancer who had never used oral contraceptives. The second alteration is the greater degree of eosinophilia in the neoplastic cells of women who have received hormones. Only a small proportion of uteri removed during therapy after oral contraceptive therapy demonstrate morphologic changes in the myometrium or the myomas harbored therein. Cervical adenosis has become clinically detectable during oral contraceptive therapy and has regressed when the hormones are discontinued. Breast tissue undergoes no histologically distinctive changes during contraceptive hormone therapy. However, the hormones occasionally produce lobular hyperplasia with secretory activity either in normal breast or within adenomas. (184 refs)

78-6646 **Oral Contraceptives and Cancer.** (Eng) Kretzschmar, R. M. (Dept. Obstetrics and Gynecology, Univ. Iowa, Iowa City, IA, 52240). *CA* 28(2): 8-123; 1978.

There is no convincing evidence that oral contraceptives cause cancer, but they have been associated with the occurrence of benign liver tumors. (no refs)

78-6647 **Postmenopausal Estrogens and Endometrial Cancer: A Pathologist's Overview.** (Eng) Sommers, S. C. (Dept. Pathology, Lenox Hill Hosp., New York, NY, 10021). In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977.* Colorado Regional Cancer Center (Denver, CO): 197 pp.; 43-53; 1978.

The relationship between the use of postmenopausal estrogens and the development of endometrial cancer is discussed. Previous epidemiologic-statistical and retrospective case control studies were poorly planned and failed to answer the questions of how often and how significantly perimenopausal estrogen administration may contribute to the development of endometrial carcinoma. Prospective cohort studies are

needed to avoid the acknowledged defects of case control and retrospective studies. It is suggested that, if there is a family history of endometrial or gastrointestinal carcinoma, obesity, diabetes, hypertension, proven endometrial hyperplasia or dysplasia, or an abnormal genotype, estrogens should probably not be given, as they might exaggerate an already abnormal situation and contribute eventually to development of endometrial carcinoma. In other women, estrogens may be prescribed without fear of endometrial carcinogenesis, provided that an initial endometrial specimen does not exhibit hyperplasia or dysplasia, that the dose is in the low (growth stimulating) rather than in the high (growth inhibiting) range, and that the estrogen is not given continuously. If estrogen is administered in a high dose, progestagen should be administered concomitantly about 1 wk in every 4. Follow-up endometrial aspiration or curette biopsies are desirable to appraise the endometrial response at intervals. (63 refs)

78-6648 **Effect of Oral Contraceptive Use on the Risk of Human Breast and Cervical Cancer.** (Eng) Stadel, B. V. (Natl. Inst. Child Health and Human Development, NIH, Bethesda, MD, 20014). In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977.* Colorado Regional Cancer Center (Denver, CO): 197 pp.; 89-102; 1978.

The epidemiology of breast and cervical cancers and the effect of oral contraceptive (OC) use on the risk of these cancers are reviewed. No overall increase in breast cancer risk for OC users has been observed in four case control studies or three cohort studies. However, only one study specifically analyzed possible interactions between OC use and previously reported breast cancer risk factors, such as age of first full-term birth and benign breast disease. This study suggests that OC use may increase breast cancer risk among certain subgroups of women and indicates the need for further study. The available data regarding the effect of OC use on the risk of cervical cancer and its precursors are confusing and inconclusive. Most of these data do not control directly for possible confounding between OC use and either age at first coitus or number of sexual partners. The use of such variables as socioeconomic status, marital status, and parity in an effort to control indirectly for possible confounding between OC use and sexual behavior may be insufficient. Separate evaluation of the effect of OC use on the risk of cervical cancer and its precursors for women with previously reported cervical cancer risk factors would be useful, since OC use might affect risk only among women exposed to a venereally transmitted carcinogen prior to OC use. Virtually all available studies of the effect of OC use on the risk of cervical cancer and its precursors have had cervical dysplasia or carcinoma in situ as their biologic endpoint. Diagnostic criteria for these lesions appear to vary widely, as do estimates of their invasive potential. The current information is insufficient to warrant the conclusion that OC use affects the risk of either breast or cervical cancer, but it does indicate a need for further studies. (34 refs)

- 78-6649 Breast Cancer and Menopausal Estrogens.** (Eng) Ross, R. (Dept. Pathology, Univ. Southern California, Sch. Medicine, Los Angeles, CA, 90033); Mack, T.; Gerkins, V.; Hill, A. In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977.* Colorado Regional Cancer Center (Denver, CO): 197 pp.; 55-65; 1978.

Despite evidence implicating estrogens in mammary tumor development in animals, the role of exogenous menopausal estrogens in human breast cancer (BC) etiology remains unresolved. Of two case-control studies, the first revealed a large protective effect of hormones, but the second showed a two-fold increased risk of BC in hormone users. When the studies were combined and adjusted for age at diagnosis and menopause, estrogens were found to have no effect. In a recent study of 1,891 women using conjugated estrogens and followed for an av of 12 yr, the overall relative risk of BC was 1.3. However, women in this study who took at least 0.625 mg estrogen on other than a daily basis and followed by > 10 yr had a relative risk of 4.7. The methodology for a current case-control study designed to overcome the methodologic errors of previous studies is described. Drug histories of cases and controls matched by age, marital status, and social class are being obtained through personal interviews, medical record abstraction, and pharmacy records, and a detailed menstrual and reproductive history is also being taken. (33 refs)

- 78-6650 Oral Contraceptives and Liver Tumors.** (Eng) Bloustein, P. A. (Dept. Lab. Medicine, Wesley Medical Center, Wichita, KS, 67202). In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977.* Colorado Regional Cancer Center (Denver, CO): 197 pp.; 103-110; 1978.

Reported associations between oral contraceptive use and the development of focal nodular hyperplasia, liver adenoma, and hepatocellular carcinoma are reviewed, and the effects of oral contraceptives on liver metabolism and ultrastructure are summarized. Serum transaminase activity and bromosulphalein retention increased in women who took the first estrogen- and progestagen-containing oral contraceptives in 1960, but the incidence of markedly deranged liver function has decreased as the amount of steroids has been reduced. Liver dysfunction is usually transient, suggesting that the hepatocytes can adapt. The most striking ultrastructural change observed in the hepatocytes with oral contraceptive use was an increase in the amount of smooth endoplasmic reticulum. Some evidence suggests that the development of focal nodular hyperplasia might be related to contraceptive use, although this evidence is not as compelling as it is with other hepatic lesions. In general, the former lesion occurs in women in the child-bearing years, regardless of oral contraceptive use. Vascular lesions were described in focal nodular hyperplasia prior to the introduction of birth control pills and thus cannot be ascribed to the use of these agents. However, hemorrhage within a preexisting lesion of focal nodular

hyperplasia may be related to oral contraceptive use, because only since the advent of the pill have patients with this lesion presented with significant intrahepatic hemorrhage or acute hepatic rupture. Hepatocellular adenoma was an exceptionally uncommon tumor prior to the widespread use of oral contraceptives. The incidence of this lesion has increased in recent years in women taking oral contraceptives, especially those who had taken the hormone for several years. The most severe complication is rupture of the tumor with resultant hemorrhage. There is evidence suggesting that the tumor produces a biologically active substance that causes diarrhea. Liver cell carcinoma has been reported in women using oral contraceptive steroids and estrogen-like drugs, but the number of such cases in no way represents the type of increase in frequency that has been seen in adenomas. (29 refs)

- 78-6651 Treatment of Menopause.** (Eng) Morris, J. M. (Dept. Obstetrics and Gynecology, Yale Univ. Sch. Medicine, New Haven, CT, 06516). In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977.* Colorado Regional Cancer Center (Denver, CO): 197 pp. 13-26; 1978.

The risks and benefits of hormone replacement therapy during menopause are discussed. Although patients taking exogenous estrogens appear to have an increased risk of developing endometrial and breast cancer, there is evidence that the endometrial tumors induced by exogenous estrogen are of lower-grade malignancy than their nonexogenously induced counterparts. In post-menopausal women, estrogen administration can reverse a negative calcium balance, halt bone loss, and arrest vertebral deformity and loss of height. (19 refs)

- 78-6652 Vaginal and Cervical Pathology Associated with Prenatal Exposure to Diethylstilbestrol.** (Eng) Robboy, S. J. (Dept. Pathology, Massachusetts General Hosp., Boston, MA, 02114); Prat, J.; Welch, W. R. In: *Estrogens and Cancer. Proceedings of a meeting held by the Colorado Regional Cancer Center in Denver, Sept. 10, 1977.* Colorado Regional Cancer Center (Denver, CO): 197 pp.; 153-171; 1978.

Intrauterine exposure to diethylstilbestrol (DES) has been linked to clear cell adenocarcinoma of the vagina and cervical nonneoplastic abnormalities of the lower genital tract in women (vaginal adenosis, cervical ectropion, vaginal and cervical ridges) and, more recently, to anomalies of the epididymis and testis in men. The pathologic findings in DES-exposed subjects are reviewed at length. Ongoing animal studies of the mechanisms of DES are summarized, as are clinical studies of the natural history of DES-induced adenosis and the potential risks of consuming meat from DES-fed animals during pregnancy. (52 refs)

6653 **Alcohol and Cancer.** (Eng) Tuyns, A. J. (International Agency Res. Cancer, Lyons, France). *Alcohol Health Res World* 2(4): 20-31; 1978.

role of alcohol in the development of cancer, particularly cancer of the mouth, tongue, larynx, pharynx, esophagus, liver, and lung, is reviewed. Alcohol consumption has also been associated with stomach, pancreatic, and rectal cancer. In some of these cancers, the effects of drinking and smoking are usually combined in an additive or even multiplicative manner. (108 refs)

6654 **Biological Autoxidation. 1. Decontrolled Iron: An Ultimate Carcinogen and Toxicant: An Hypothesis.** (Eng) Kon, S. H. (Henry Dazian Dept. Chemistry, Mount Sinai Hosp., New York, NY, 10029). *Med Hypotheses* 4: 445-471; 1978.

accumulation of endogenous hydrolyzed iron that is out of the control of ferritin is hypothesized to act as an inorganic carcinogen. Uncontrolled ionic iron promotes autoxidation, which crosslinks biomolecules and produces destructive active oxygen. The disease can be initiated by the transient actions of metallic catalysts of oxidation, a foreign body, a metabolic insufficiency, a burst of ionizing radiation, or free radicals. These events give rise to porous residual bodies containing limited-access spaces (LAS), which remain as a permanent tissue imprint. The pores exclude ferritin and digestive and antiautoxidation enzymes but admit ionic iron, which, sheltered from ferritin, forms a decontrolled-iron package (DIP). Thus, a DIP is an accumulation of endogenous hydrolyzed iron in LAS, an open-pore space of a porous body. The DIP uses the energy of ambient pH fluctuations to erupt from the LAS, swamp the available ferritin, poison its surroundings, catalyze autoxidation, and crosslink cell components into additional LAS carriers. There is a striking resemblance between preneoplastic carcinogenesis and DIP growth and expansion. Protein denaturants and reagents, oxidation catalysts, metabolism inhibitors, and foreign bodies such as asbestos, all initiators of carcinogenesis, are also initiators of DIP growth. It is likely that carcinogenesis includes DIP growth as a preneoplastic stage. (121 refs)

6655 **Nitrites: The Newberne Report (Letter to Editor).** (Eng) Lachance, P. A. (Dept. Food Science, Rutgers State Univ. New Jersey, New Brunswick, NJ 08903). *Science* 202(4368): 576; 1978.

The results of a Food and Drug Administration-sponsored study of nitrites were misinterpreted. There was no solid evidence in that study that nitrites are carcinogens. The results do not permit assigning nitrite a proximate carcinogenic status, and it was stated that nitrite appears to be more a promoter of neoplasia than an initiator. (no refs)

78-6656 **N-Nitroso-N-ethylurea.** (Eng) IARC Working Group (Lyon, France). In: *IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man: Some N-Nitroso Compounds*. (Lyon: International Agency for Research on Cancer): 17: 191-215; 1978.

Experimental evidence on the carcinogenicity of N-nitroso-N-ethylurea (NEU) is reviewed. NEU has been used in the laboratory synthesis of diazoethane and has been studied for growth promoting activity in plants, but it has never been used commercially. The compound is carcinogenic in mice, rats, Syrian golden hamsters, rabbits, opossums, pigs, and monkeys. NEU produces benign and malignant tumors, the sites of which vary with the route of administration. Prenatal exposure induces tumors of the nervous system. Treatment of *Salmonella typhimurium*, *Escherichia coli*, and *Drosophila melanogaster* with NEU leads to mutations, and treatment of cultured human fibroblasts with the compound results in both chromatid and chromosome aberrations. Although no human case reports or epidemiological data were available for analysis, it was concluded that NEU should be regarded as if it were carcinogenic to humans. (101 refs)

78-6657 **Is Cigarette the Wrong Scapegoat?** (Swe) Carlens, E. (Lidingo, Sweden). *Lakartidningen* 75(40): 3531-3533; 1978.

Experimental and epidemiological studies indicate that tar containing benzopyrene (BP) is a relatively weak carcinogen. Therefore, such cigarette smoke components as acrolein, formaldehyde, and hydrocyanic acid, which contribute to BP carcinogenesis by destroying the mucociliary barrier of the lungs, are equally important factors in the genesis of lung tumors. (10 refs)

78-6658 **Is Pulmonary Dysfunction the Common Denominator for the Multiple Effects of Cigarette Smoking?** (Eng) Cohen, B. H. (Dept. Epidemiology, Johns Hopkins Univ. Sch. Hygiene and Public Health, Baltimore, MD, 21205). *Lancet* 2(8098): 1024-1027; 1978.

The hypothesis that impaired pulmonary function underlies the diversity, multiplicity, and putative "nonspecificity" of the effects of cigarette smoking was assessed in a brief literature review. The differences in degree as well as type of clinical expression probably derive from the basic role of pulmonary function in mediating the action of the many different constituents of the tobacco itself against the background of various environmental exposures and intrinsic host factors. Which of the specific effects become manifest in different individuals depends on the dynamic interplay of genes, environment, and chance that determines individual vulnerability. (38 refs)

- 78-6659 Organochlorine Pesticides and Liver Cancer Deaths in the United States, 1930-1972.** (Eng) Deichmann, W. B. (Res. and Teaching Center Toxicology, Coral Gables, FL, 33124); MacDonald, W. E. *Exotoxicol Environ Saf* 1(1): 89-110; 1977.

The association between the use of organochlorine pesticides, primarily chlorophenothane (DDT), aldrin, and dieldrin, in the US and liver cancer deaths between 1930 and 1972 was investigated. Intensive and prolonged occupational exposure to these pesticides has been found to result in increased accumulation of the compounds in the body fat, emaciation, hypertension, elevated serum cholesterol, impairment of neuromuscular function, carcinoma, and generalized pathologic conditions of the liver. Other studies, however, have indicated no undesirable effects or loss of working time as a result of pesticide exposure, in spite of high body burdens. The incidence of hepatic cancer has fallen from 8.8 per 100,000 in 1930 to 5.6 per 100,000 in 1972. This steady decline in total liver cancer death rates fails to support any clear-cut evidence of an increase in liver cancer deaths since the introduction of organochlorine pesticides in the US. If any of the organochlorine pesticide compounds do have carcinogenic potential, it will first be noted in persons engaged in the manufacture, formulation, and extensive use of these compounds. (51 refs)

- 78-6660 Carcinogenicity of Saccharin.** (Eng) Reuber, M. D. (Chemical Carcinogenesis Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501). *Environ Health Perspect* 25: 173-200; 1978.

Published and unpublished carcinogenicity studies of saccharin are reviewed. Saccharin is carcinogenic for the urinary bladder in rats and mice, with male rats being more susceptible than females. Rats exposed as fetuses develop neoplasms more readily than rats exposed as weanlings. The lesions in the urinary bladder go through the stages of hyperplasia, hyperplastic nodules, and carcinomas. Neoplasms of the urinary bladder in rats were not caused by stones, parasites, sodium, or impurities. There is a cocarcinogenic effect between saccharin and methylnitrosourea for the urinary bladder. Chronic renal disease appears to result from low doses of saccharin, tumors of the renal pelvis and urinary bladder from higher doses. Benign and malignant neoplasms at all sites are increased significantly in mice and rats ingesting higher doses of saccharin. These neoplasms are present in the reproductive and hematopoietic systems and, to a lesser extent, in the lungs, vascular system, and squamous epithelium. Saccharin initiates skin neoplasms when its application is followed by croton oil. Three epidemiological studies reportedly show no increase in cancer of the urinary bladder following saccharin use. However, a Canadian study found an estimated 60% increase in cancer of the urinary bladder associated with the use of artificial sweeteners in men. (60 refs)

- 78-6661 Viral and Cellular Contributions to Expression of the Transformed State.** (Eng) Pollack, R. I. (Dept. Microbiology, State Univ. New York, Stony Brook, NY, 11594). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 169-179; 1977.

A review is presented of cell growth control in culture and its relationship to the virally induced progression from normal to malignant phenotype. Although present understanding of growth control is limited by the inability to grow many normal cell types in culture, the fibroblast model (although quite complex) has been well studied. Diminished serum requirement, loss of density-dependent growth, and relaxation of the anchorage requirement for G_1 growth are important examples of the phenotypic changes induced by various oncogenic viruses; each is sufficient to trap a normal fibroblast in the G_1 phase of the cell cycle. Tumorigenicity requires a specific group of phenotypic changes induced by viral genes; this tumor syndrome has been elucidated for SV40-transformed rodent fibroblasts injected into athymic nude mice: it consists of loss of cytoarchitectural order, loss of anchorage requirement, and production of plasminogen activator (these are necessary but not sufficient causes for tumorigenicity in this system). Although it is known that expression of the SV40 A gene is necessary for expression of the tumor syndrome, future research is needed to determine which cellular molecules the A gene product interacts with to generate this syndrome. In addition, it is necessary to delineate syndromes of transformation for nonfibroblastic cells, which account for more than 90% of human tumors. More recently, keratinocytes, macrophages, arterial endothelial cells, capillary mural cells, and many hormone-dependent glandular cells have been put into culture and should aid researchers in unraveling the mechanism of expression of the transformed state. (42 refs)

- 78-6662 Food: A Carcinogenic Hazard?** (Eng) Kroes, R. (Lab. Pathology, Natl. Inst. Public Health, P.O. Box 1, Bilthoven, Netherlands). In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 659-664; 1977.

Information on the possible carcinogenic hazard of food is reviewed. Natural components of food that pose a carcinogenic threat include nitrate (in spinach, leeks, and purslane), fats, and proteins. Natural contaminants of food that are carcinogenic include aflatoxin, cycasin, safrole, sterigmatocystin, penicillic acid, patulin, luteoskyrin, cyclochlorotrine, and pyrrolizidine alkaloids. Manmade chemicals that contaminate

foods come from a wide range of sources. The food additives butter yellow, ponceaux SX, and 3R, guinea green, fuchsin, amaranth, furfurylfuramide, and dulcin are now relatively prohibited, based on recent findings that they may be carcinogenic. Contamination can occur from food packaging chemicals; ie, vinyl chloride, a proven carcinogen is now federally regulated. Various agricultural chemicals still in use and under suspicion include dibromoethane, monochloropropane, aramite, chlordecone, lindane, DDT, aldrin, dieldrin, and nitrogen fertilizers. In addition, essentially carcinogenic substances used for disinfection, with promotion, weed and insect control, and chemical synthesis reach food as a result of environmental pollution. The last source of carcinogenic contamination is from food processing: smoke curing introduces polycyclic aromatic hydrocarbons, nitrate and nitrite curing introduces nitrosamines, and processing through the use of fungi introduces mycotoxins into the diet. (25 refs)

78-6663 Potential Cancer Risk in the Graphic Arts and Textiles Industries. (Eng) Kay, K. (Environmental Sciences Lab.-Dept. Community Medicine, Mount Sinai Sch. Medicine, City Univ. New York, Fifth Ave. & 14th St., New York, NY, 10029). In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 857-899; 1977.

Occupational environments in the graphic arts and textiles industries were investigated through survey of numerous establishments. Examination of printing and ink establishments provided information on a number of chemicals. Although the literature reports no carcinogenic effects of carbon black per se, high-boiling oils used with carbon black in inks may act as solvents for cancer-producing chemicals adsorbed on the carbon. Chlorinated hydrocarbon solvents, petroleum hydrocarbon solvents, and high volatility solvents are used widely and may contribute to carcinogenesis although convincing evidence of this is lacking. Among the polymers and solventless inks found in use were formaldehyde resins, halogenated aromatic amines, benzophenone, anthraquinone, and styrene, having chemical relatives that are proven carcinogens. Ballpoint and felt-pen inks employ pigments and dyes comparable to those used in printing inks and incorporate alkyd resins, glycols, allyl alcohol-styrene polymers, formaldehyde-urea-ketone and urea-formaldehyde resins. In the textiles industry, the following carcinogenic dyes have been reported: Direct black 38, Direct brown 95, Direct blue 6, Disperse blue 3, Vat green 1, and Basic green 4, all containing the chromophore, -N=N-. Among flame retardants widely used are tris (2,3-dibromopropyl) phosphate, and three of its isomers. Contaminants have been shown to be mutagenic for histidine-deficient strains of *Salmonella typhimurium*. Furthermore, chemical relatives of other flame retardants, namely, halogenated and organophosphorous compounds, are

known carcinogens. Finally, three fluorescent whitening agents, 3-benzyl-4-methyl-7-hydroxycoumarin, disodium 4,4'-bis-(2,4-dimethoxybenzamido)-2,2'-stilbenedisulfonate, and disodium 4,4'-bis-(4,6-dianilino-s-triazin-2-yl)amino-2,2'-stilbenedisulfonate have been shown to enhance the carcinogenic potency of germicidal UV radiation in mice. (182 refs)

78-6664 Proposal for Abatement of Polycyclic Aromatic Hydrocarbon Levels Through Standardization of Benzo(a)pyrene Level in the Air. (Ger) Tomingas, R. (No affiliation given); Pott, F. *Lufthyg Silikoseforsch* 10: 28-50; 1978.

In view of the carcinogenic effects of benzo(a)pyrene (BP) and other polycyclic aromatic hydrocarbons, the atmospheric BP level should be reduced to 10 nanograms/m³. However, this level should be adopted on an interim basis, because it cannot be regarded as perfectly safe. (61 refs)

78-6665 Rationale for One Double Event in Model for Radiation Carcinogenesis (Letter to Editor). (Eng) Baum, J. W. (Safety and Environmental Protection Div., Brookhaven Natl. Lab., Upton, NY, 11973). *Health Phys* 35(4): 581-582; 1978.

The author responds to criticism of an article on the development of the multiple simultaneous event model for radiation carcinogenesis in which the critics suggest that two double events should have been considered. Two double events had been considered and rejected because of poor fits to the experimental data. A rationale for the absence of two double events is provided by considering the possibility that these events are gene alterations: two genes located near each other at the end of the chromosome may both be lost due to a terminal deletion, but there may be no similar mechanism for a second double event in the same cell or its ancestors. The critics also referred to claims for the predictive power of the model. The model was used to make predictions primarily to design experiments to test it further and improve the model. Claims were not made concerning the probability that the predictions would be validated. (2 refs)

78-6666 Formation of Thymine Containing Dimers in Skin Exposed to Ultraviolet Radiation. (Eng) Johnson, B. E. (Dept. Dermatology, Univ. Dundee, Dundee, Scotland). *Bull Cancer (Paris)* 65(3): 283-297; 1978.

A review is presented of the formation of thymine dimers in skin exposed to UV radiation (UVR). UVR is known to damage nuclear DNA, causing primarily pyrimidine-pyrimidine dimers. Although thymine is always a constituent of the dimer, thymine-cytosine dimers are also readily formed. Thymine dimers revert to the monomeric form after exposure to short

wavelength UVR (240 nanometers [nm]). Photoreactivation (a photoenzymatic process dependent on long wavelength UVR [320/390 nm] or even visible radiation) also repairs dimers, although it is not well documented in mammalian systems. However, excision repair (dark repair) has been demonstrated in humans. Studies with ³H-thymidine labeling and fluorescent antibody binding have provided evidence that DNA is photochemically changed in vivo in skin exposed to UVR, and that this change in DNA involves the formation of thymine dimers. Estimates of the yield of these dimers vary with radiation dose and wavelength, but the documented range is 0.016-2.6% of total DNA activity. In vitro studies have indicated linear increases in dimer yields over the range 50-1,000 Joules (J)/meter² (m²), and an exposure of monolayer cultures to 100 J/m² at 254 or 260 nm yielded a common value of approx 0.45% of total DNA activity as dimers. The most efficient dimer-forming activity appears to occur at approx 280-290 nm, with efficiency dropping off at both limits, perhaps more steeply at the upper end. Evidence concerning the persistence of dimers in vivo is conflicting, but a decrease in dimer incidence with time after irradiation appears to occur. Although thymine-containing dimers are formed in mammalian skin exposed to UVR in vivo, a relationship between this phenomenon and neoplastic transformation in skin remains to be proven. (73 refs)

- 78-6667 **The Controversy over the Health Effects of Radiation.** (Eng) Najarian, T. (Boston Veterans Admin. Hosp., Boston, MA). *Technol Review* 81(2): 74-82; 1978.

Nuclear workers exposed to low levels of ionizing radiation at the Portsmouth Naval Shipyard [av lifetime exposures of 2-10 Rems (Roentgen equivalent man)] and the Hanford Atomic Plant (av lifetime exposures of 1-15 Rems) are developing leukemia and/or multiple myeloma and pancreatic cancer at higher than expected rates. An excess of leukemia deaths has also been reported in men exposed to 1 Rem during a Nevada atom bomb test. The new data are either faulty or low-level radiation is much more harmful than was assumed previously. (11 refs)

- 78-6668 **X-Ray Dose Fractionation and Oncogenic Transformations in Cultured Mouse Embryo Cells (2 Letters to Editor).** (Eng) Peto, R. (Dept. Regius Professor Medicine, Radcliffe Infirmary, Oxford, England); Hall, E. J.; Miller, R. *Nature* 274(5674): 926; 1978.

In a previous study of the x-ray transformation of cultured cells, an inappropriate statistical technique was used to present the data, and, as a result, the accuracy of the data in some cases was overestimated considerably. In a reply, the authors agree with this criticism but they emphasize that the conclusion that split doses produce more transformations than a single exposure of the same total dose still holds. (1 ref.)

- 78-6669 **Molecular Biological Aspects of Oncogenesis Caused by RNA Tumor Viruses.** (Ger) Chanock, P. (Abteilung fur Molekularbiologie, J.-W.-Goethe-Universität, Theodor-Stern-Kai 7, 6000 Frankfurt-Main 70, Germany); Ebener, U.; Steel, L. K.; Laube, H.; Kornhuber, B.; Thorbeck, R. R. *Klin Paediatr* 190(1): 9-27; 1978.

The molecular mechanisms by which RNA tumor viruses (oncornaviruses) transfer their genetic information from genomic RNA (70S RNA) of the virions to cellular DNA, which leads to neoplastic transformation, are discussed. Special attention is focused on the biochemical and serological properties of reverse transcriptase (RT) and the role of the enzyme in the life cycle of RNA tumor viruses. The role of RT in myelofibrosis is also discussed. This RT is antigenically related to the RT of certain primate RNA tumor viruses as to that of human leukemia cells. The role of RT inhibition in viral oncogenesis, particularly that of partially thiolated polycytidylic acid (MPC), which has a high binding affinity to viral RT, is reviewed. The results of the use of MPC in a few leukemia patients are reported. (141 refs.)

- 78-6670 **The Oncogenic Potential of Viruses.** (Eng) Irfinger, W. (Institut für Genetik, Abt. für Virologie, Universität Köln, Weyertal 121, 5000 Cologne 41, Germany). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 87-110; 1977.

The oncogenic potential of viruses is reviewed with respect to the mechanism of malignant transformation. A variety of viruses can transform cells in culture or induce tumors in animals. A transformed or tumor cell can be distinguished from a normal cell, but transformed cells do not always exhibit all the properties of tumor cells. In virus-transformed cells, viral DNA persists frequently in covalent linkage with cellular DNA. Viral DNA often is integrated in the form of fragments. In cells transformed by adenovirus type 12 and simian virus 40, the sites of integration of the viral genome appear to differ from one cell line to another. In transformed cells, specific viral functions are expressed; in the case of homologous cells transformed by RNA tumor viruses, the entire viral genome is transcribed, and infectious virus is produced. Specific viral genes appear to be essential for transformation, in that mutants lacking these genes cannot transform. However, normal cells contain and express these similar genes without being transformed. In fact, nearly all apparently normal cells carry endogenous viruses belonging to the RNA tumor virus group. Several possible mechanisms relating the integration of viral genomes to neoplastic transformation are proposed: (1) fixation of the viral genome in host cells; (2) specific viral gene products (under host control) are responsible for transformation; (3) integration of viral genes causes mutations; (4) chromosome aberrations and

nslocations; (5) repair mechanisms in eukaryotic cells; and "hit-and-run" mechanism. Each of these hypotheses is incompatible with experimental observations, although they may be too limited to explain the actual biologic phenomenon of transformation. In different systems, transformation may be caused at different levels, and different principles may be involved. (32 refs)

78-6671 Viral "Tumor Antigens": A Novel Type of Mammalian Regulator Protein. (Eng) Weil, R. (Dept. Molecular Biology, Univ. Geneva, 30 quai Ernest-Ansermet, CH-1211 Geneva 4, Switzerland). *Biochim Biophys Acta* 516(3): 301-388; 1978.

The biological relevance of polyoma virus (PyV), simian virus (SV40), and the papilloma viruses is reviewed in the general context of cell and tumor biology. The gene products of PyV and SV40, ie, their virus-specific tumor (T) antigens, seem to represent a novel type of mammalian regulator protein. There is a striking contrast between the broad spectrum of biological effects induced by SV40 and PyV and the small size of their genome. The late region of both viruses comprises two (possibly 3) genes that specify the major (VP1) and minor (VP2 and VP3) capsid proteins. The early region, expressed before onset of viral DNA replication, comprises approximately 50% of the viral genome. It contains most of the information for lytic infection and all for abortive infection, induction of mitosis, initiation and maintenance of cell transformation, tumor formation in animals, and, possibly, for the repression of virus-specific transplantation antigens. The experimental evidence is compatible with the hypothesis that the biological effects of the early region are mediated by T antigens, although their mode of action is not known. Additional topics covered include new antigens (transplantation, surface, and T antigens) in virus-induced tumors, transformation of mammalian cells in vitro, cytological and molecular events of lytic and abortive infection with PyV and SV40 in tissue culture cells, the nature and biological significance of the virus-specific imprints in PyV- and SV40-transformed cells, and virus-specific pleiotropic effector genes and their significance for cell transformation and tumor induction. (717 refs).

78-6672 Advances in the Viral Etiology of Leukemia and Lymphoma. (Eng) Gallo, R. C. (Lab. Tumor Cell Biology, NCI, Bethesda, MD, 20014); Meyskens, F. L. *Leukemia Hematol* 15(4): 379-398; 1978.

The emphasis of this review is a critical examination of evidence for the presence of RNA tumor virus information in human leukemia. This evidence comprises the detection of extracytoplasmic particles with properties of known oncornaviruses and of reverse transcriptase (RT) related to murine and certain nonhuman primate enzymes, the rare demonstration of oncornaviruses by electron microscopy, and demon-

stration of extracellular viral particles. It is not known how the viral information enters humans and if it is important for leukemia or any disease. The major difficulty in linking C-type oncornaviruses with human disease is the rarity of isolating replicating biologically active virus from human tissue. This problem is approached by reviewing conditions under which most animal isolates are obtained. Baboon endogenous virus, simian sarcoma virus (SiSV), and gibbon ape leukemia virus may be of special interest, since components related to these viruses have been reported in human leukemic tissues. However, there are also reports of (1) a transient release of virallike particles, (2) isolation of a partially characterized infectious or transforming virus, and (3) replicating infectious virus from human cells. Failure to detect SiSV-related proviral nucleic acid sequences in human leukemic tissues may be due to the facts that SiSV is integrated only in a rare cell, complete SiSV is integrated and then completely or partially excised, or infecting virus is integrated in a tissue remote from the primary phenotypic expression. Several explanations are given for the presence of intracytoplasmic particles, which have several properties similar to those of oncornaviruses, in human leukemia cells. The detection of oncornaviruslike particles, RT and specific antibody to RT on the surface of specific types of leukemic cells may lead to the development of markers for human leukemia. (103 refs)

78-6673 The Structure and Expression of Human Herpesvirus DNAs in Productive Infection and in Transformed Cells. (Eng) Roizman, B. (Committee on Virology, Univ. Chicago, Chicago, IL, 60637); Frenkel, N.; Kieff, E. D.; Spear, P. G. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conference on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1069-1111; 1977.

The structure of herpesvirus DNA and its expression during the reproductive cycle and in transformed cells are reviewed. Evidence is presented for coordinate and sequential synthesis of viral gene products, and mechanisms and requirements for this sequential synthesis are also reviewed. Several lines of evidence indicate that posttranscriptional processing of viral RNA plays a major role in the initiation of viral polypeptide synthesis. Two representatives from the β class of herpes simplex virus (HSV) proteins (thymidine kinase and VP143) and one from the γ class (a glycoprotein) are expressed in some transformed cell lines. In studies of the presence of HSV DNA and RNA in transformed cells, the entire genome was never found. The fraction of the genome retained varied from one cell line to another and in different sublines of the same line. The number of copies of the retained sequences was relatively low. Studies with Epstein-Barr virus (EBV)-infected cells indicate that the majority of reproductively infected cell lines contain most and probably all of the EBV genome. The synthesis of viral DNA in reproductively infected cells is linked to the synthesis of cell DNA. The differences between EBV and

HSV might be a consequence of differences in inherent regulation of viral gene function in HSV- and EBV-infected cells or they might be due to differences in the inherent permissiveness of the cells transformed by these viruses. Cells transformed by HSV are permissive, whereas those transformed by EBV are largely nonpermissive. Since EBV replicates only in B lymphocytes, cell factors present in the permissive host cell but lacking in the more restrictive lymphocytes could account for the difference between HSV and EBV in the retention and expression of viral genes in transformed cells. (164 refs)

- 78-6674 RD-114 Virus: Characterization and Identification.** (Eng) McAllister, R. M. (Children's Hosp. Los Angeles, 4650 Sunset Blvd., Los Angeles, CA, 90027); Gardner, M. B.; Nicolson, M. O.; Gilden, R. V.; Davidson, N. *Prog Exp Tumor Res* 21: 196-215; 1978.

Studies performed to characterize and identify RD-114 virus and to determine its role in feline and human cancer are reviewed. RD-114 virus is a xenotropic, endogenous virus of domestic cats and of several closely related feline species. A relationship between RD-114 and endogenous baboon C-type viruses has led to the suggestion that under natural conditions, C-type viral genes can be transferred between remotely phylogenically related species. Electron microscopy of the high-mol-wt RNA extracted from RD-114 virus showed that it was an inverted dimer containing two monomer subunits with symmetrically placed loops. The two monomers are joined at their nonpoly(adenosine) ends. The dimer linkage and symmetrical loops seen in RD-114 viral RNA probably are a prototype for common structural features of all mammalian C-type viral RNA's. Transfection studies indicated that human rhabdomyosarcoma productively infected with RD-114 virus contained virus-specific DNA that specifies complete information for progeny virus formation. This DNA is integrated into the chromosomal DNA. RD-114 RNA was detected in most malignant feline tissues that transcribe feline leukemia virus (FeLV) RNA, but the converse was not seen. Thus, RD-114 gene expression may be a function of dividing cells and may be activated by FeLV infection. In older cats with naturally occurring cancer, however, transcription and translation of the endogenous RD-114 genome was not detected in the absence of FeLV expression. (78 refs)

- 78-6675 Synthesis and Processing of the Translational Products of RNA Tumor Viruses.** (Eng) Jamjoom, G. A. (Riyadh Univ. Sch. Medicine, Riyadh, Saudi Arabia); Arlinghaus, R. B. *Methods Cancer Res* 15: 3-67; 1978.

The characterization of the translation products of some RNA tumor viruses is reviewed. The virus life cycle, the structure of the viral genome, and the identification and mapping of viral genes are discussed. The major polypeptides of

the mature virions of murine and avian C-type viruses are tabulated. Studies of virus-specific messenger RNA (mRNA) indicate that the mRNA's for the *gag* and *pol* gene products are genome-length RNA species that appear to be identical to the viral genomic RNA. The mRNA's for the *env* and *src* gene products include segments of the genome extending from the *env* or *src* gene, respectively, to the 3'-end. The gene product is required for cell transformation. Modified, uncleaved precursors of the core proteins have been identified on the surface of mouse leukemia cells. The ability of these precursors to interact with nucleic acids and to associate with the cell membrane suggests that they may be capable of interfering with normal processes under conditions in which they accumulate in the cells. Exchange of genetic material among endogenous viruses of the murine system may play a role in the etiology of mouse leukemia. Evidence that the recombinant MCF viruses are the cause of leukemia in AKR mice is suggestive but still circumstantial. Studies of the peptide maps of gp70 molecules from MCF viruses and AKV viruses indicate that recombination does occur and that it occurs in the *env* gene. There is probably more than one way in which cells can become malignant, at least where C-type viruses are concerned. Two different genes, *src* and *env*, appear to be important in virus-induced sarcomas and leukemias, respectively. The gene products are different in structure, and their mode of action may also be different. (242 refs)

- 78-6676 Avian Leukemia Viruses: Interaction with Target Cells In Vivo and In Vitro.** (Eng) G. T. (German Cancer Res. Center, Neuenheimerfeld 2, D-6900 Heidelberg, W. Germany); Beug, H. *Biochim Biophys Acta* 516(3): 269-299; 1978.

The origin, oncogenic spectrum, in vitro target cell specificity, and genetic properties of avian leukemia viruses (ALV) are reviewed. Both nondefective LV, which predominantly induce lymphatic leukemias after long latency periods, and defective LV, which predominantly cause acute erythroid and myeloid leukemias after short latency periods, are considered. Susceptibility and resistance to avian sarcoma virus (ASV) and nondefective LV are determined by dominant cellular genes and by specific properties of the lymphoid target cells. Nongenetic factors that determine susceptibility and resistance include the host immune system, inoculation route, the way in which the virus was passaged or selected, and the type of tumor cells from which the virus was obtained. The oncogenic spectrum and latency period do not allow an accurate distinction between ASV and the different types of ALV. Since none of the ALV apparently possess the *src* gene, ASV, this property is proposed as a main distinguishing criterion. It is suggested that highly leukemogenic, defective LV have evolved from low leukemogenic, nondefective strains in a multistep process involving the acquisition of several properties. A tentative and highly speculative model of leukemogenesis by defective LV is presented. The model is based on three assumptions: the transformation gene of defective LV represents a modified sequence of a cellular gene.

red for normal differentiation; a necessary step in topopoietic differentiation is the interaction of an early differentiation factor with a certain cellular protein, thereby triggering the synthesis or activation of differentiation factors characteristic of later stages; the virus-coded transformation is still capable of binding to the early differentiation factor. The most obvious explanation for the transformation of cell specificity of ALV in hematopoietic cells is the ability that they are able to infect and replicate only in certain types of cells. (150 refs)

77 Oncornavirus Leukemogenesis as a Model for Selective Neoplastic Transformation. (Eng) Gardner, I. L. (Lab. Experimental Oncology, Dept. Pathology, Stanford Univ., Stanford, CA, 94305); Baird, S. In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 135-152; 1977.

Review is presented of the classification, pathogenetic role, and functions of oncornaviruses (ORV). ORV are subdivided into two groups on the basis of whether they carry the early or late gene for the expression of the malignant phenotype in target cells. Specific loci on the chromosomes of rapidly transforming (RT) ORV code for malignant phenotype expression within hours to days after host cells are infected. The genome of the RT avian ORV has been defined by the coding genes: *gag* which codes for a polyprotein precursor; *pol* which codes for nucleic acid polymerases; *env* which codes for a budding site polyprotein; *sarc*, the transforming gene which codes for common oligonucleotides; and *poly* whose function is unknown. Most RT ORV are replication-defective and require the aid of a "helper virus" to produce bud-derivations in host cells. Some ORV are slowly transforming and induce lymphocytic-leukemia weeks to months after infection into target host cells; helper viruses usually fall into this group. These viruses do not carry the *sarc* gene and are not able to transform lymphocytes but may transform fibroblasts. Many model systems emphasize the fact that malignant transformation by ST ORV requires a specific interaction between cell and virus and that replication and transcription of the viral genome is not sufficient to cause malignant transformation in cells other than susceptible thymic lymphocytes. Pathogenesis by ST ORV involves preleukemic and leukemic phases that select for expression of virion proto-oncogenes. Various models are presented that attempt to account for the origin, embryonic function, and leukemogenesis of these viruses including the oncogene, provirus, enhancer, and receptor-mediated transformation models. (24

Central Veterinary Inst., Lelystad, Netherlands); Bloemers, H. P. *Biochim Biophys Acta* 516(3): 249-268; 1978.

Recent developments concerning the genetic functions of murine sarcoma viruses (SV) and other replication-defective (RD) mammalian C-type viruses [Abelson leukemia virus (AbLV), nonmurine mammalian SV, spleen focus-forming virus (SFFV)] are reviewed. These viruses share several characteristics: they are able to transform cells in vitro and are oncogenic in vivo; they are RD and can be rescued from nonproducer cells by helper LV, yielding pseudotype viruses; their genome, which consists of a 30S RNA subunit (approx 6,000 nucleotides), evolved by recombination of LV-specific genetic information and sequences derived from normal rat cells (Kirsten and Harvey SV), normal mouse cells (Moloney SV), or xenotropic murine LV (SFFV). The origin of the non-LV sequences in the genomes of AbLV, feline SV, and woolly monkey SV is unknown. At least one of the functions of the LV sequences in the RD viral genomes may be to provide 3'- and 5'-terminal ends necessary for circularization and integration of the defective genome in the chromosomal DNA of a susceptible host cell. Several findings are given to support this speculation. The gene products of the *gag*, *pol*, or *env* genes of the LV genome are apparently not required for cell transformation by SV because they are partly or completely absent in transformed nonproducer cells infected by different SV. This does not exclude the possibility that unexpressed *gag*, *pol*, or *env* gene sequences are present and play a role in transformation or that a putative LV-specific *onc* gene cooperates somehow with SV-specific sequences. The non-LV sequences in the genome of most RD viruses are not related to each other. Apparently, SV and other RD transforming C-type viruses are vehicles of sequences that are involved in malignant transformation and, possibly, in unknown processes in normal cells. (145 refs)

78-6679 Search for RNA Tumor Virus in Humans. (Eng) Gardner, M. B. (Dept. Pathology, Univ. Southern California Sch. Medicine, Los Angeles, CA, 90033); Rashed, S.; Shimizu, S.; Rongey, R. W.; Henderson, B. E.; McAllister, R. M.; Klement, V.; Charman, H. P.; Gilden, R. V.; Heberling, R. L.; Huebner, R. J. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation.* Hiatt, H. H.; Watson, J. D.; Winston, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory: Origins of Human Cancer. Vol. 4 (Book B), 699 pp.; 1235-1251; 1977.

A number of unsuccessful attempts to identify RNA tumor viruses in human material, using techniques that have proved successful in identifying RNA tumor viruses in animal tumor systems, are summarized. Virus particles could not be identified by electron microscopy, infectious viruses could not be isolated from cultured human tissues, and virus antigens or antibodies could not be identified by radioimmunoassay in a large number of human specimens. These specimens were tissues and sera from patients with a variety of cancers and

78-6680 The Genome of the Mammalian Sarcoma Viruses. (Eng) Van Zaane, D. (Dept. Virology,

diseases (sarcomas, carcinomas, lymphomas, leukemias, lupus erythematosus, xeroderma pigmentosum, and others). The fact that no antibodies to the well-characterized RNA tumor viruses of animal origin were detected in the many hundreds of human sera tested (not even in sera from individuals known to have been exposed to such viruses, eg, feline leukemia virus) is strong evidence against the horizontal infection of humans with any of the known mammalian C-type viruses. These results, however, do not rule out the possibility that noninfectious endogenous RNA tumor viral genomes are present in human cells. (55 refs)

- 78-6680 Limited In Vitro Replicative Lifespan of Cultured Human Epithelial Cells: A Survey of Recent Literature.** (Eng) Ruben, R. L. (Dept. Anatomy, Hahnemann Medical Coll., 230 Broad St., Philadelphia, PA, 19102). *Growth* 42(10): 357-368; 1978.

The literature concerning limits to the in vitro replicative life span of normal, neoplastic, and virus-transformed human epithelial cells is reviewed. Normal diploid cells routinely undergo protracted crisis in vitro, leading to the eventual loss of the cultures. Epithelial cell tumors, such as melanomas, oat cell carcinomas, and colorectal adenocarcinomas, have met with greater culturing success. However, all cell lines surviving long-term culture have had an aneuploid karyotype. Neoplastic and virus-transformed cells either degenerate in culture if diploid or survive the degenerative phase if aneuploid. Since most cells of autochthonous tumors in vivo are diploid, aneuploidy and immortality may be associated with primary or secondary transformation in vitro rather than with malignancy in vivo. In view of this evidence, the concept that neoplastic or virus-transformed cells possess unlimited potential for division should be reevaluated. (66 refs)

- 78-6681 Patterns of Gene Expression and the Cellular Origins of Human Leukaemias.** (Eng) Greaves, M. (Membrane Immunology Lab., Imperial Cancer Res. Fund, London, England); Janossy, G. *Biochim Biophys Acta* 516(2): 193-230; 1978.

Detailed phenotypic profiles on human lymphoid leukemias and on permanent cell lines established from the leukemic cells were studied to determine whether there are any consistent antigenic properties of these leukemias and whether there are any antigens of the tumor cells that could be recognized by the host. The currently available data suggest that: (1) most, if not all, of human hematopoietic malignancies are monoclonal derivatives of cells resident at the time of leukemia initiation in various linkage and differentiation compartments from the common lymphomyeloid progenitors; (2) with improved banding techniques, a high frequency of non-random chromosomal alterations are detectable in leukemia; (3) leukemic cells express a relatively stable morphological,

cytochemical, enzymatic, and antigenic phenotype that is probably unique for and qualitatively normal for the cell lineage in which the disease arises; and (4) there is no unequivocal evidence at present for leukemia-specific membrane antigen or other alterations. These interpretations challenge the rationale of immunotherapy aimed at deliberate autoimmunity in leukemia and suggest that consistent and qualitatively unique membrane changes in neoplasia may not exist. (22 refs)

- 78-6682 Stress and Disease: The Missing Link. A Vasoconstrictor Theory. III. Stress, Vasospasm and General Disease.** (Eng) Boyd, G. W. (Univ. Dept. Medicine, 4 Collins St., Hobart 7000, Australia). *Med Hypotheses* 4(5): 432-444; 1978.

Repeated ischemic cell damage from vasospasm may be a factor in the development of autoimmune disease and cancer. DNA may be altered directly or through errors in DNA replication. The similarities between viral transformation and neoplasia have led to the proposition that much cancer might be explained if, as a species, humans have evolved by the gradual build-up of viruses. (66 refs)

- 78-6683 Herpesviruses, Veneral Disease, and Cancer.** (Eng) Rapp, F. (Dept. Microbiology, Milton S. Eisenhower Medical Center, Pennsylvania State Univ., Hershey, PA, 17033). *Am Sci* 66(6): 670-674; 1978.

The probability is significant that certain human tumors may, in part, be caused by herpesviruses (HV). Epstein-Barr virus is uniformly associated with Burkitt's lymphoma, nasopharyngeal carcinoma in the southern Chinese, and a significant percentage of non-African lymphomas. Herpes simplex virus type 2, which causes venereal diseases, is associated with cervical cancer. All HV cause cells to synthesize DNA and undergo mitosis, and they can transform normal cells from many species to cells with malignant properties. (10 refs)

- 78-6684 Epidermodysplasia Verruciformis. An Autosomal Recessive Disease Characterized by Viral Warts and Skin Cancer. A Model for Viral Oncogenesis.** (Eng) Lutzner, M. A. (Dermatology Branch, Room 231, Building 10, NCI, NIH, Bethesda, MD, 20014). *Bull Cancer (Paris)* 65(2): 169-182; 1978.

The case reports of 147 epidermodysplasia verruciformis (EV) patients comprising 125 families were reviewed. EV is an autosomal recessive disease that usually begins in infancy or childhood, with an average age of onset of 9. It was found that EV has no racial predisposition and no geographic preference, and it occurs with the same frequency in both sexes.

warts are most common, but pityriasis-like warts occur in approx 75% of patients. The warts are disseminated and chronic. Cancer may develop as early as age 13; the average age at onset is 31. About one-third of the patients develop cancer of the skin, usually in exposed areas. The cancer is often squamous cell carcinoma. The average time from wart to cancer is 24 yr, but cancer has been developed in as little as 2 yr. Malignant transformation is usually in carcinoma-in situ of the Bowenoid type or squamous cell carcinoma. Two deaths occurred by local invasion; one metastasis was recorded. One patient died of Burkitt's lymphoma and another was reported to have disseminated reticulosis. Approx 8% of the patients are mentally retarded. Humoral immunity is normal, but cell-mediated immunity is depressed in most patients. The gene defect in this disease has not yet been uncovered. (135 refs)

685 Hepatocellular Carcinoma and the Hepatitis B Virus: Evidence for a Causal Association. (Eng) Weiss, R. A. (Lab. Epidemiology, New York Blood Center, New York, NY). *Prog Med Virol* 24: 40-69; 1978.

Evidence is presented in support of the hypothesis that hepatitis B virus (HBV) is a primary etiological agent in the development of hepatocellular carcinoma (HCC). HCC is widespread in parts of Africa and southeast Asia, but relatively uncommon in the US and other Western countries. Patients in the high-incidence areas are quite young, and they are predominantly male. The hypothesis is supported by the similarity in the geographic distribution of HCC and HBV, evidenced by radioimmunoassay of the HBV surface antigen (HBsAg). Moreover, there is an unusually high frequency of HBsAg and antibody to the hepatitis B core antigen (anti-HBc) in HCC patients; the amount of free circulating surface antigen seems to be relatively low. Based on available information, the relative risk of past HBV infection in HCC patients is 1.0-1.5 in high-incidence areas and 10-15 in low-incidence areas. HBsAg prevalence in high-incidence areas appears to be inversely related to patient age, and mother-child transmission is suspected. Although HCC is strongly associated with the HBsAg carrier state, genetic and environmental factors (eg, malaria, malnutrition) may also contribute to susceptibility to HCC. (114 refs)

686 Epigenetic and Viral Aspects of Cancer: Group Report. (Eng) Weiss, R. A. (Imperial Cancer Research Fund Labs., Lincoln's Inn Fields, P.O. Box 123, London WC2A 3PX, England); Baserga, R. L.; Basilico, C.; Bernhard, H.; Bustan, H.; Gardner, R. L.; Green, H.; Illmensee, K.; Jisch, R.; Kaplan, H.; Koprowski, H.; Lilly, F.; McGuire, E.; Pollack, R. E.; Sato, G. H.; Weissman, I. L. In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a Meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 269-288; 1977.

Epigenetic and viral hypotheses of carcinogenesis are summarized. Teratocarcinoma development usually involves transformation from a malignant pluripotent stem cell to non-malignant, differentiated progeny. This reversibility of the neoplastic state is an interesting concept for further research, especially since several human tumors of fetal origin, including retinoblastoma and Wilms' tumor, frequently regress postnatally. There is evidence that hormones play a role in tumor development, perhaps by activating an oncogene. Recent developments in culture methods are clarifying hormone requirements and interactions in neoplastic cell lines. Transformation by oncogenic DNA viruses appears to require the synthesis of specific viral gene products, with integration of the viral genome or genome fragments being essential for persistence of the transformed state. Papovaviruses, whose early genome region codes for both large T (tumor) and small t antigens, cause parotid gland tumors in animals and benign warts in humans. Most transforming RNA viruses (retroviruses) carry the oncogene *sarc*, but their replication is dependent on helper viruses that possess all virion genes. Although simian sarcoma helper virus and baboon endogenous virus have been isolated from humans, they have not been implicated in human disease. Papilloma virus, the cause of warts, is the only virus known to cause tumors in humans. (13 refs)

78-6687 Tumor Phenotype and Progression: Group Report. (Eng) Eichmann, K. P. (Institut für Immunologie und Genetik, Deutsches Krebsforschungszentrum, Im Neuenheimer Feld 280, 6900 Heidelberg, W. Germany); Askonas, B. A.; Franke, W. W.; Howard, J. C.; Lennox, E. S.; Mitchison, N. A.; Nicolson, G. L.; Ohno, S.; Rajewsky, K. In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 289-304; 1977.

Information concerning the phenotype of malignant cells and the immune reaction to malignant cells is reviewed. Topics discussed include the morphologic and biochemical properties of the tumor cell membrane, the three-dimensional arrangement of the malignant cells, characteristics of tumor cells with a high rate of metastasis, genetic characteristics of malignant cells, tumor cell antigens, and the biological function of antigens. A review of host responses to tumors includes immune surveillance, the role of immunity in metastasis, the association between the presence of histocompatibility antigens and susceptibility to tumors, manipulation of the immune system (ie, enhancement of tumor immunogenicity, stimulation of lymphocyte compartments, and idiotypic manipulation), nonspecific defense mechanisms, and tumor-associated antigens. (22 refs)

78-6688 Immune Surveillance. (Eng) Howard, J. C. (Dept. Immunology, ARC Inst. Animal Physi-

ology, Brabraham, Cambridge CB2 4AT, England). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a Meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977*. Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 227-245; 1977.

The "strong" form of the immunosurveillance theory is discussed. This theory differs from weaker versions in asserting that the "true" tumor incidence in animals endowed with an immune system is higher than the observed incidence and that the difference is accounted for by the activity of the immune system in recognizing new antigens on incipient neoplastic cells and disposing of the cells that bear them. Two statements must be true for the strong theory of immune surveillance to be valid: (1) immunodepression will lead to an increase in the incidence of tumors in any age class; and (2) incipient neoplasias will be immunogenic. The concept has not stood up to experimental test. A generalized increase in spontaneous tumor incidence is not seen in experimental or natural states of immunodepression. Even highly immunogenic tumors are no more easily induced in immunodepressed animals than in normal controls. Only virus-induced cancers show an increased incidence in immunodepressed animals, just as such animals are most susceptible to nononcogenic viral infections. There is no evidence that the incidence of overt cancer is limited by the activity of the immune system except in the case of horizontally transmitted oncogenic viruses. It is worth wondering why immune surveillance is not effective in other situations in which tumors are highly antigenic. (25 refs)

78-6689 Some New Data on the Pathogenesis of Chronic Lymphoid Leukemia (Meeting Abstract). (Rus) Polianskaia, A. M. (No affiliation given). *Vopr Onkol* 24(9): 121; 1978. (no refs)

78-6690 Mechanisms of Tumor Immunity. (Eng) Cohen, S. (Dept. Pathology, Univ. Connecticut Health Center, Farmington, CT, 06032); Cohen, M. *Am J Pathol* 93(2): 449-457; 1978.

Ways in which the immune response system is able to effect the proliferation of tumor cells are reviewed. The immune mechanisms of tumor-cell killing are divided into two groups: cell-mediated and non-cell-mediated mechanisms. Effector cells in tumor cell-mediated destruction include lymphocytes, which secrete soluble mediators, participate in antibody-dependent reactions, and can become cytotoxic themselves; macrophages, which may be rendered cytotoxic by an arming factor from activated T lymphocytes; and granulocytes, which participate in antibody-dependent cytotoxic reactions. Noncellular mechanisms of tumor-cell destruction include lymphotoxins, produced by lymphocytes stimulated with specific antigen or with nonspecific mitogens that can kill or damage various cell lines in vitro; and complement, which

can induce holes or lesions in the cell membrane of tumor cells. Nontoxic effects of the immune system involve modification of cell mobility (important in metastasis and loss of contact inhibition). Although many natural protective measures are built into the immune system, genetic and predisposing factors can affect tumor resistance. (27 refs)

78-6691 Evolutionary Development of IgE and the Role of Anaphylactic-Type Reactions in Resistance to Solid Tumors. (Eng) Lynch, N. R. (Institut de Recherches Scientifiques sur le Cancer, B.P. N 8, 94800 Villejuif, France); Salomon, J. C.; Turner, K. J. *Cancer Immunol Immunother* 4(4): 223-228; 1978.

The evolutionary development of IgE and the role of anaphylactic reactions in resistance to solid tumors are reviewed. The necessary components for allergic or anaphylactic-type reactions are found in many species, including marsupials, which have long diverged from the main mammalian line. This finding suggests that the existence of such reactions has been evolutionarily conserved, and thus may have a regulatory or protective role in the organism. Evidence suggests that anaphylactic-type reactions are a constituent of anti-tumor mechanisms is presented. These reactions can increase the access of immune effectors into solid tumors, though progressively growing tumors appear to have a self-protective mechanism inhibiting such reactions. It has not yet been established that the anaphylactic system directly modulates the activity of lymphoid cells, although involvement in the expression of delayed hypersensitivity seems certain. The possibility that anaphylactic-type reactions contribute to parasite rejection is an attractive but controversial idea. (79 refs)

78-6692 Immunology of Rat Hepatic Neoplasia. (Eng) Baldwin, R. W. (Cancer Res. Campaign Lab., Univ. Nottingham, Nottingham, England). In: *Rat Hepatic Neoplasia: Workshop on Rat Liver Neoplasia held by William F. Merrell, Co. in Woods Hole, October, 1974*. William F. Merrell, Co. (Woods Hole, MA): 288 pp.; 228-240; 1978.

The expression of tumor-associated rejection antigens, tumor-specific cell-surface antigens, and tumor-associated embryonic antigens in carcinogen-induced rat liver tumors is discussed. The most significant characteristic of carcinogen-induced tumors is expression of the tumor-specific cell-surface antigens that may function as tumor-rejection antigens. This type of neoantigen is not always expressed by transformed cells and cannot be used as a criterion of malignant change. However, there are few exceptions to the general finding that each tumor has a characteristic antigen, and characterization of this specificity may be useful in evaluating the role of carcinogenic agents in transformation. Embryonic antigen expression may be a concomitant effect of neoplastic change; thus, typing of these antigens could provide a suitable preliminary screening assay for carcinogen-induced transformation. The role of these antigens in tumor rejection is unresolved, but it is unlikely that they are responsible for

ity elicited against transplanted hepatic tumors. (31 refs)

693 **On the Pathogenesis of Cutaneous Malignant Lymphoma (CML).** (Ger) Steigleder, G. K. Universitat Hautklinik Koln, Joseph-Stelzmann-Strasse 9, Koln 41, W. Germany). *Z Hautkr* 53(20): 719-726;

Previously published suggestion concerning the pathogenesis of cutaneous malignant lymphoma (CUML) is discussed. This suggestion still appears valid in view of current observations. Both T- and B-cell lymphomas of the skin have been reported. The suggested model of CUML development involves proliferation of lymphocytes stimulated by persistent injury or chronic exposure to UV light; if a lymphocyte contains neoplastic information, a clone of neoplastic cells will arise. The stimulated lymphocytes secrete lymphokinin, which influences other cells (eg, by decreasing the movement of macrophages). Because T-lymphocytes are more common than B-lymphocytes, neoplastic transformation of T-lymphocytes is more probable. Mycosis fungoides (MF) can be regarded as analogous to eczema, while B-cell CUML can be considered a type of exanthema. The involvement of allergic allergy is consistent with the association of MF with occupations such as construction work and others leading to sun exposure. In this model, infectious mononucleosis is considered a pseudolymphoma. (43 refs)

694 **Antigenic Changes in the Transformed Cell.** (Eng) Lennox, E. S. (MRC Lab. Molecular Biology, Hills Road, Cambridge CB2 2QH, England); Sikora, I. *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a Meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, W. Germany): 314 pp.; 197-207; 1977.

The effect of the immune system on antigenic changes in the surface components of transformed cells was studied. Tumor antigens are defined by various modes of immune response: (1) in vivo by induction of transplantation immunity or delayed-type hypersensitivity and in vitro by serological or cell-mediated reactions against tumor cells as targets. By comparing these separate modes of response, various antigens can be defined on the same tumor. The following antigens can be detected on tumor cells, some unique, some shared with normal cells: (1) normal transplantation antigens (the targets of immune rejection reactions when tissue is transplanted between individuals); (2) tumor-specific transplantation antigens (TSTA: defined in protection assays in which animals immunized with the tumor are subsequently challenged with the same or other tumors, with studies revealing that certain induced tumors carry unique TSTA); (3) virus-determined antigens (those of DNA and RNA tumor viruses; they can serve as transplantation antigens as well as immunogens for antibody formation and thus may be targets of humoral as well as cell-mediated attack); (4) embryonic

antigens (defined by antisera prepared in syngeneic animals by injection of preparations of mid-term embryos or by sera from multiparous animals; they can detect antigens on tumor cells transformed by viruses or induced by carcinogens); and (5) differentiation antigens (those present on most cells of one particular individual). TSTA appear to be the most likely candidates for recognition in immune surveillance, and they may play a major role in the interaction between the immune system and cells undergoing transformation. (12 refs)

78-6695 **Immunodeficiency and the Pathogenesis of Lymphoma and Leukemia.** (Eng) Louie, S. (Tufts Univ. Sch. Medicine, Boston, MA, 02111); Schwartz, R. S. *Semin Hematol* 15(2): 117-138; 1978.

The possible pathogenic connection between immunologic abnormalities and malignant lymphoproliferative diseases is examined in light of clinical situations associated with a high risk of lymphoma. Several congenital immunodeficiency diseases are associated with a high risk of malignant lymphoma: ataxia telangiectasia, Wiscott-Aldrich syndrome, common variable immunodeficiency, and Bloom's syndrome. A characteristic of all these diseases is the presence of abnormalities outside the immune system, and the genetic disturbance in these conditions can manifest itself by structural changes in chromosomes. These diseases may be regarded as premalignant states characterized by chromosome instability and an associated immunologic disturbance. The risk of developing malignant lymphoma is also increased in recipients of renal allografts and in untreated patients with certain immunoinflammatory diseases (autoimmune hemolytic anemia, Sjogren's syndrome). The risk of developing acute myelogenous leukemia is increased in patients treated with immunosuppressive agents for immunoinflammatory diseases and in cancer patients treated with either aggressive chemotherapy or chemotherapy combined with radiotherapy. It is suggested that some 'spontaneous' lymphomas may arise as the result of the enhanced proliferation of mutant lymphocytes brought about by the selective pressure of impaired immunoregulation. (164 refs)

78-6696 **Geographical Distribution of Gastrointestinal Cancer and Breast Cancer and Its Relation to Selenium Deficiency.** (Eng) Jansson, B. (Natl. Large Bowel Cancer Project, Univ. Texas System Cancer Center, M. D. Anderson Hosp. and Tumor Inst., Houston, TX, 77030); Mahaly, M. A.; Seibert, G. B. *In: Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 1161-1178; 1977.

Evidence indicating that selenium may act as an inhibitor of cancer of the colon, rectum, and breast is reviewed. Studies in the US and New Zealand showed that low rates of colon, rectal, and breast cancer were associated with high Se concen-

trations in the soil and vice versa. A New Zealand veterinarian observed that the incidence of intestinal adenocarcinomas and other neoplasms in sheep was reduced after the introduction of Se drenching. Similar findings were obtained in studies of rats fed carcinogens with and without Se. Another study showed that spontaneous mammary tumors were inhibited by Se in mice. An unexpected finding was that the incidence age was lowered for tumor-bearing mice fed Se compared with untreated control tumor-bearers. Se deficiency reduces the fertility of animals and may also have the same effect in humans. Thus, the finding of a negative correlation between birth rates and colon, rectal and breast cancer rates in a study of the natality and mortality statistics for 48 continental states further supports the hypothesis that Se or some other agent acts as an inhibitor of these cancers. (38 refs)

- 78-6697 **Causes of Cancer in Man.** (Eng) Miller, R. W. (NCI, NIH, Bethesda, MD, 20014). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a Meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 25-38; 1977.

The causes of cancer in humans are reviewed, with emphasis on physical and chemical carcinogens. Among the physical carcinogens are two forms of radiation, ionizing and UV. Ionizing radiation can penetrate deeply and induce various forms of cancer, whereas UV light is not penetrating and induces only skin cancer. Asbestos acts as a physical carcinogen in that its physical properties, ie, the length and diameter of the fibers, apparently account for its carcinogenicity in various organs. Chemical carcinogens induce respiratory tract, urinary bladder, blood, and liver malignancies, especially in heavily exposed industrial workers. A subtle influence of chemical carcinogenesis early in life is indicated by the retention of old-country cancer rates in migrants whose children have the cancer rates of the new country and by the distinctive patterns of excessive mortality from specific cancers observed in the 3,056 US counties between 1950-1969. Some of these patterns appear to be related to industries within the counties. Despite extensive laboratory evidence for viral oncogenesis, no virally induced human cancers have been demonstrated. Host factors, including single genes, chromosomal aberrations, and immunodeficiency, may place individuals, families, or ethnic groups at a very high or low risk of certain cancers. (33 refs)

- 78-6698 **Variations in Cancer Death Rates.** (Eng) Lew, E. A. (Dept. Epidemiology and Statistics, American Cancer Society, Inc., New York, NY). In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 1127-1138; 1977.

Current observations of the variations in cancer death rates among various populations are reviewed briefly, with a warning directed against premature overzealous interpretation of apparent differences. Although US death rates increased 5.2% in the first 7 mo of 1975, the age-adjusted death rate actually increased only approx 1% over the entire yr. It has been an observable phenomenon, however, that sharp increases are noted in cancer death rates over the first half of any year. Comparisons among nations must be made with care, particularly when 10-fold differences in population exist, eg, the US and Sweden. Moreover, methods of death certification and registration vary from country to country, automatically introducing bias. It is suggested that it may be at least as useful to study mortality rates from different parts of the US as to compare different countries. Several special studies investigating social factors are discussed; in general it has been found that death rates from most cancers are higher for the lower socioeconomic classes (as defined by occupation, education, and/or economic status), although the opposite is true for prostate cancer and leukemias in men and breast cancer in women. Studies of native white vs foreign-born white vs nonwhite populations have also been useful in determining several interesting relationships, such as the increased digestive tract cancer and decreased prostate cancer mortality in these non-native populations. Better medical care facilities will continue to improve both tumor classification and the definition of various cancer incidence and mortalities. (no refs)

- 78-6699 **Our Daily Habits and Cancer.** (Swe) Werko, G. (Farmaceutisk Divisionen, Astra Lakemedel AB, Sweden). *Lakartidningen* 75(10): 919-920; 1978.

The risk of cancer is increased by tobacco, alcohol, and dietary habits. Positive correlations have been found between alcohol consumption and laryngeal and buccal cavity cancer and between high-fat diets and leukemia or colon and breast cancer. (11 refs)

- 78-6700 **Benign and Malignant Breast Tumors: Epidemiological Similarities.** (Eng) Sartwell, P. E. (Dept. Epidemiology, Johns Hopkins Univ. Sch. Hygiene and Public Health, Baltimore, MD, 21205); Arthes, F. Tonascia, J. A. *Int J Epidemiol* 7(3): 217-221; 1978.

Risk factors were estimated in 783 cases of cystic disease of the breast (CD), 155 cases of breast fibroadenoma (FA), 10 cases of miscellaneous breast conditions, and 284 cases of breast cancer. FA occurred in the youngest patients, the highest percentage (39.4%) occurring in patients aged 15-24. The cancer patients were the oldest, 67.6% of the cases occurring in women aged 45-64 yr. Relative risks were calculated for each of the four breast conditions, with respect to the possible risk factors that could be categorized dichotomously. These were nulliparity, first pregnancy at age 25 or older, being premenopausal at the time of discovery of the lesion, and marriage at age 20 or over. Each patient was compared

her own control matched for age, race, ever-married, time of hospitalization, and pay status. Risks associated with being nulliparous were all above unity, but only that the cancer patients was statistically significant ($p < 0.05$). First pregnancy at an older age, all the risks were above unity, and those for CD and cancer were significantly elevated. In all four disease classes, the patients had fewer pregnancies than their controls; the differences were significant only for CD and FA. CD patients had experienced both natural and artificial menopause less commonly than their controls; breast cancer patients had experienced only artificial menopause, more commonly than their controls. This general similarity between benign breast disease and breast cancer in certain reproductive and menstrual features plus previous evidence that breast cancer is more frequent in women who have previously had benign breast disease suggest a similarity in etiologic factors. (9 refs)

701 The "Urban Factor" and Lung Cancer: Cigarette Smoking or Air Pollution? (Eng) Carnow, J. (Dept. Preventive Medicine, Univ. Illinois, Chicago, IL 60680). *Environ Health Perspect* 22: 17-21; 1978.

Evidence that urban pollution is a significant etiologic factor in lung cancer was provided by four groups of studies. Studies of urban-rural death rates revealed a doubling of lung cancer mortality in large cities compared with rural areas, which could not be explained by a difference in smoking habits. Studies of lung cancer death rates in migrants also suggested a relationship to urban pollution. Among studies that used regression analysis in an attempt to identify urban factors responsible for urban-rural differences in death rates from lung carcinoma, one indicated that the products of solid fuel combustion may be an important etiologic factor and another showed a significant relationship between the death rates and atmospheric concentrations of (a) pyrene (BP) levels. After standardization for such variables as smoking habits, age, sex, and duration of residence, sampling studies in the US and other countries found that lung cancer incidence was significantly associated with urbanization and air pollution. Data on lung cancer deaths among coke oven workers were used to calculate a dose-response relationship between BP and lung cancer in an urban community. The risk of excess lung cancer is due to continuous exposure to BP is higher than previously assumed. (24 refs)

702 Identification of Carcinogenic Hazards from Aromatic Amine Exposures and Measures for Control. (Eng) Parkes, H. G. (Health Res. Unit, British Rubber Manufacturers' Assoc., Scala House, Holloway Circus, London, W.1, England). In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, NY). Vol. 2 (part 1), 2404 pp.; 1913-1920; 1978.

Variation in the duration, intensity, and conditions of occupational exposure to carcinogens such as aromatic amines will determine whether the resultant cancer incidence is 100% or negligible. Suspicions may be aroused by chemical similarities between established and suspect carcinogens and they may be reinforced by evidence of experimental mutagenicity, teratogenicity, or carcinogenicity, but positive epidemiological identification is the final confirmation of a carcinogenic hazard. (12 refs)

78-6703 Cell Cycle Dependency of Transformation. (Eng) Baserga, R. L. (Dept. Pathology, Fels Res. Inst., Temple Univ. Sch. Medicine, Philadelphia, PA, 19140). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977*. Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 125-133; 1977.

A review of the major studies concerning the cell-cycle dependency of transformation, coupled with more recent work showing a close relationship between transformation and mutagenesis, suggest that G_1 cells are more susceptible to transformation than are G_0 cells because they have a higher relative transcriptional activity, and transcriptionally active chromatin is more accessible to mutagens and carcinogens than is transcriptionally inactive chromatin. It has also been shown that S-stage cells are susceptible to transformation for the same reason G_1 cells: incorporation of the transforming agent into the accessible DNA. It may be predicted from these conclusions that G_0 cells are strongly resistant to transformation by chemical carcinogens, and research is discussed that supports this contention. (18 refs)

78-6704 DNA Polymerase and Mutation. (Eng) Bridges, B. (Univ. Sussex, Sussex, England). *Nature* 275(5681): 591-592; 1978.

Several theories on the interaction between DNA damage and DNA polymerases in the generation of permanently altered nucleotide sequences and, as a result, cancer and certain heritable diseases are presented. (4 refs)

78-6705 Cellular Susceptibility to Neoplasia and Its Reversibility. (Eng) Gardner, R. L. (Dept. Zoology, Univ. Oxford, South Parks Road, Oxford OX1 3PS, England). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a Meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977*. Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 111-124; 1977.

Cell susceptibility to neoplastic transformation and its reversibility are reviewed. The susceptibility has been proven to be

dependent on both intrinsic and extrinsic elements. Intrinsically, genetic manifestation of susceptibility is the result of polymorphic loci coding for enzymes involved in DNA repair or metabolism of carcinogens and proteins affecting the activity of tumor viruses. Among the extrinsic factors that have been implicated in neoplasia because they alter the cellular environment are chemical carcinogens, 'foreign bodies', prolonged endocrine imbalances, and other elements that impose stress on tissue homeostasis. It is suggested that all these extrinsic mechanisms act by stimulating cell division. A number of methods have been used to suppress neoplasticity; such as incorporation of a thymidine analog into DNA, use of temperature-sensitive mutant cells that are neoplastic only at the permissive temperature, or hybridization of malignant cells with nonmalignant cells that have lost a single chromosome. Differentiation of tumor cells to a terminal, postmitotic state can result in loss of neoplasticity. Furthermore, a model of plant and animal teratoma cells and nuclei isolated from adenocarcinoma cells suggests that neoplasia may in some instances also be reversed. (33 refs)

78-6706 Characterization of Hepatic Nodules. (Eng)

Becker, F. F. (Dept. Anatomic and Experimental Pathology, M. D. Anderson Hosp. and Tumor Inst., 6723 Bertner Ave., Houston, TX, 77030). In: *Rat Hepatic Neoplasia: Workshop on Rat Liver Neoplasia held by William F. Merrell, Co. in Woods Hole, October, 1974*. William F. Merrell, Co. (Woods Hole, MA): 288 pp.; 42-57; 1978.

Research on premalignant and malignant hepatic nodules is reviewed. Identification of premalignant lesions in animal tissues exposed chronically to chemical carcinogens would provide insight into the possibility of a chemical etiology for similar lesions in humans, allow analyses to be focused on one small region rather than the entire organ, and would help in understanding of characteristics of malignant cells. Evidence indicating that hepatic nodules are premalignant lesions is presented. Study of hepatic nodules allows carcinogen-altered tissue to be examined at various stages in its progression toward malignancy and nodules of different stages to be compared with each other and with nonnodular parenchymal areas. Techniques allowing cytogenetic analysis of the chromosome composition of solid hepatic tissues have been developed recently. Problems in transplanting of hepatic nodules successfully are reviewed. The hepatic nodule is the best available tissue to which new analytical techniques can be applied and upon which new oncological concepts can be tested. (35 refs)

78-6707 Sequential Cellular Alterations During Hepatocarcinogenesis. (Eng) Bannasch, P. (Abteilung

fur Cytopathologie, Institut fur Experimentelle Pathologie Deutschen Krebsforschungszentrum, Heidelberg, W. Germany). In: *Rat Hepatic Neoplasia: Workshop on Rat Liver Neoplasia held by William F. Merrell, Co. in Woods Hole, October, 1974*. William F. Merrell, Co. (Woods Hole, MA): 288 pp.; 58-99; 1978.

The sequential alterations observed in liver cells under the influence of hepatocarcinogens are discussed. Reversible alterations are characterized by a predominantly centrilobular loss of glycogen and by a concomitant dispersal of the basophilic bodies. The outstanding feature of persistent alterations is an excessive storage of glycogen, which appears to be due to the direct action of the carcinogen, resulting in an irreversible cell lesion. At the end of the preneoplastic phase in most animals, so-called hyperplastic nodules develop. These nodules appear to originate from areas storing glycogen in excess, and most nodules are probably a mixture of precancerous, cancerous, and diverse intermediary cells. It is highly probable, however, that the hyperplastic nodule is only one of at least two histologically different formations in which neoplastic transformation can take place. Nearly the entire liver parenchyma appears to undergo irreversible cellular alterations, with cellular enlargement being due mainly to excessive glycogen storage and/or a proliferation of the membranes of the agranular endoplasmic reticulum. The data suggest that the glycogen-free basophilic tumor cell is the result of a long process that begins with an excessive storage of glycogen and perhaps passes through a stage of membrane proliferation or fat accumulation. (67 refs)

78-6708 Characteristics of Malignant Phenotypes

Familial Polyposis. (Eng) Wells, S. A. (Duke

Univ. Medical Center, Durham, NC, 27710); Johnson, A. F.; Gregory, R. O. In: *International Advances in Surgical Oncology*. Murphy, G. P., ed. (New York: Alan R. Liss, Inc. Vol 1, 288 pp.; 29-52; 1978.

The clinical characteristics of various gastrointestinal polyposis syndromes, particularly familial polyposis coli (FPC), and their relationship to colorectal cancer are reviewed. FPC, which is transmitted as an autosomal dominant trait, is characterized by development in adolescence or early adult life of adenomatous polyps on the mucous membrane of the colon and rectum. The incidence of colon malignancy in FPC is thought to approach 100%. Proper therapy is contingent on distinguishing the disease from inflammatory polyps, lymphoid polyps, hamartomatous polyps, multiple neurofibromatosis (Von Recklinghausen's disease), lipomatous polyps, hyperplasia or metaplasia, and pneumatosis cystoides intestinalis. There are no serum, plasma, blood cell, or urinary markers associated with the disease. Surgery is the treatment of choice, the most common procedures being total proctocolectomy with abdominal ileostomy, total colectomy with ileoproctostomy (TC and I) (preferred), or anal ileostomy. Regression of rectal polyps occurs in approx 10%-15% of FPC patients undergoing surgery, but this phenomenon is not permanent. Other syndromes that share two common traits with FPC (autosomal dominant inheritance and propensity to develop gastrointestinal carcinoma) include Gardner's syndrome, Turcot's syndrome, familial 'minor' adenomatous polyposis coli, juvenile polyposis coli, and Peutz-Jeghers syndrome. The etiology of all these syndromes may involve genetic, environmental, and immunologic factors. (91 refs)

CHEMICAL CARCINOGENESIS

709 **Gas Standards for Air Pollution Analysis.** (Eng) Taylor, J. K. (Natl. Bureau Standards, Gaithersburg, MD); Hughes, E. E.; Dorko, W. D.; Cadoff, B. C.; Schmidt, W. P.; Marinenko, R. J.; Friend, D. G.; Meeks, S.; De, E.; Johns, R. H.; Deardorff, E. R.; Friend, D. G.; Er, B.; Heinrich, K.; Myklebust, R. *NBS Tech Note* 80-104; 1978.

Research was conducted to develop or improve methods for deriving primary gas standards for air pollution analyses by gravimetric, volumetric, and manometric techniques. Another objective was to provide the traceability of all air pollution measurements, including those concerned with the occupational environment, ambient air, and source emissions, to a common set of analytical standards. (8 refs)

710 **Applications of Infrared Measurements to Air Pollution Problems.** (Eng) Deslattes, R. (Natl. Bureau Standards, Gaithersburg, MD); Howard, C. J.; Radtke, H. E.; Lafferty, W. J.; Freund, S.; Layer, H.; Schenck, R.; Rougen, J.; Maki, A.; Beers, Y.; Evenson, K.; Wells, J.; Evove, J.; Sweger, D.; Hughes, E.; Okabe, H.; Streit, G. *NBS Tech Note* (963): 22-36; 1978.

Techniques and instrumentation were developed to utilize infrared laser devices in measuring air pollutants, particularly NO_2 and O_3 , and molecules that play a role in pollution, OH and HO_2 . The latter two molecules are radical species in the chemistry of smog formation and atmospheric ozone destruction by anthropogenic pollutants. (8 refs)

711 **In Vivo Covalent Binding of Chemicals to DNA as a Short-Term Test for Carcinogenicity** (Abstract). (Eng) Lutz, W. K. (Inst. Toxicology, Univ. Zurich, CH-8603 Schwerzenbach, Switzerland); Matter, C. *Experientia* 34(7): 927; 1978. (no refs)

712 **Effects of Vinyl Chloride in Man. A Cytogenetic Follow-up Study.** (Eng) Hansteen, I. L. (Lab. Cytogenetics, Telemark Central Hosp., Olavsgt. 26, 3900 Porsgrunn, Norway); Hillestad, L.; Thiis-Evensen, E.; Heldaas, S. *Scand J Clin Lab Invest* 51(2): 271-278; 1978.

A second cytogenetic study was made of 37/39 workers at a vinyl chloride plant 2.5 yr after a first study, during which time the workers had minimal exposure to vinyl chlo-

ride monomer (VCM). The second study, performed with 32 matched controls, was conducted to determine whether the chromosome damage found in the first study was due to VCM. Of the original 39 workers, 14 had been chosen at random, 13 because they had heavy exposure to VCM for years, and 12 because of abnormal clinical findings that later proved to be normal. Breaks, gaps, and stable rearrangements were scored as 100 metaphases/person from 48-hr lymphocyte cultures. In the first study, the mean number of chromosome breaks was significantly higher for the workers (3.41%) than for the controls (1.79%). No significant difference between these groups was found in the second study. In addition, there was no significant difference in mean number of breaks between the first control group and both groups of the second study. The results of the two studies provide evidence that VCM was the cause of the chromosome damage found in the first study. In the second study, the mean number of sister chromatid exchanges per cell was the same (7.6) for both workers and controls. (19 refs)

78-6713 **Hepatic Angiosarcoma due to Vinyl Chloride. Report of 4 Cases.** (Fre) Roche, J. (Clinique d'Hépatologie-Gastroenterologie, C.H.U. de Grenoble, Hôpital des Sablons, F 38700 La Tronche, France); Fournet, J.; Hostein, J.; Panh, M.; Bonnet-Eymard, J. B. *Gastroenterol Clin Biol* 2(8/9): 669-678; 1978.

Angiosarcomas of the liver developed in four men (aged 63, 48, 38, and 42 yr) who had worked in the production of vinyl chloride (VC) for 12, 26, 10, and 12 yr, respectively. Initial symptoms were abdominal pain in three patients and back pain (due to metastases) in one. The most constant laboratory indication was elevated serum alkaline phosphatase [55-77 milli(m)IU]; gamma-glutamyltranspeptidases were markedly elevated (217 and 298 mIU) in the two patients in which they were measured. α -Fetoprotein was normal, and SGOT and SGPT were close to normal. Two patients had hepatomegaly, which was the principal clinical finding in the literature on other VC-induced angiosarcomas. Arteriography revealed the angiosarcoma in the four patients, although $^{99\text{m}}\text{Tc}$ liver scans in two patients were also indicative of liver pathology. Gastrointestinal tract hemorrhage occurred in two patients and hemorrhage into the peritoneum in two. Survival from onset of symptoms was 2-13 mo. Clinical, biochemical, and histological features of previously reported patients with VC-induced angiosarcomas are reviewed. (50 refs)

78-6714 **Occurrence of Hepatocellular Tumors and Hemangiosarcoma in Rats Inhaled 1,2-Di-**

bromoethane Under the Influence of Disulfiram Treatment (Meeting Abstract). (Eng) Wong, L. C. (Pharmacology and Toxicology, Midwest Res. Inst., Kansas City, MO, 64110); Winston, J. M.; Hong, C. B.; Lee, C. C.; Bhandari, J. C. *Pharmacologist* 20(3): 174; 1978. (no refs)

78-6715 Chromosomal Analyses in Vinyl Chloride-exposed Workers. (Eng) Purchase, I. F. (Central Toxicology Lab., Central Medical Group and Mond Div., Imperial Chemical Industries Ltd., Alderley Park, Macclesfield, Cheshire, England); Richardson, C. R.; Anderson, D.; Paddle, G. M.; Adams, W. G. *Mutat Res* 57(3): 325-334; 1978.

The chromosomal morphology of cultured peripheral lymphocytes from 81 men [57 employed in plants manufacturing vinyl chloride (VC) or polyvinyl chloride (PVC), 19 on-site controls, and 5 off-site controls] was studied. There was a significant increase in chromosomal abnormalities among the VC/PVC-exposed workers compared with the controls. Total B cells (those with a chromatid break or gap or a chromosome gap), total Cu cells (those with larger unstable chromosomal abnormalities), and total Cs cells (those with larger unstable chromosomal abnormalities) showed the greater increases in autoclave operators, with smaller increases in the other five job categories. The increase in chromosomal aberrations was correlated with length of exposure and with a history of exposure to short-term excursion levels (approx 15 ppm) of VC during the year prior to sampling (1973-1974). There were no correlations between results of liver function tests and chromosome aberrations. Quantity of tobacco smoked 18 mo after blood sampling was correlated with total C cells, as was smoker vs nonsmoker status at 18 mo postsampling. It was not possible to determine whether smoking history, length of employment, or exposure to short-term excursion levels of VC was the most important variable in determining C-cell abnormalities. (15 refs)

78-6716 Use of Terbium(III) Fluorescence Enhancement to Selectively Monitor DNA and RNA Guanine Residues and Their Alteration by Chemical Modification. (Eng) Ringer, D. P. (Biomedical Div., Samuel Roberts Noble Foundation, Ardmore, OK, 73401); Burchett, S.; Kizer, D. E. *Biochemistry* 17(22): 4818-4825; 1978.

The abilities of DNA and ribosomal RNA (rRNA) from rat liver to enhance terbium(III) (Tb^{3+}) fluorescence were studied under aqueous conditions at pH 6 and at room temperature. DNA and rRNA excited by 290-nanometer light enhanced Tb^{3+} fluorescence by 1,800-fold and 13,000-fold, respectively. Of various ribose and deoxyribose nucleotides examined, only those containing guanine, guanosine 5'-monophosphate (Guo-5'-P) and deoxy-Guo-5'-P, produced significant enhancement of fluorescence. Similarly, only po-

ly(G) of the homopolymers of the four common ribonucleotides was a strong enhancer to Tb^{3+} fluorescence. The Guo-5'-P moiety in the polymer state was at least five to six times more efficient in increasing fluorescence than the free nucleotide. No enhancement was produced with 7-methyl-Guo-5'-P or Guo-5'-P modified at the 1 and N² positions by kethox and at the 8 position by N-acetoxy-2-acetylaminofluorene (NA-AAF). Chemical modification of rat liver DNA and rRNA in vitro with NA-AAF resulted in a 70% and 30% respectively, reduction in their ability to enhance Tb^{3+} fluorescence. Thus, Tb^{3+} may be useful as a guanine-specific fluorescent probe for RNA and DNA. Also, since modifications at several different positions on guanine caused loss of enhancing ability, the terbium probe may be useful in detecting guanine modifications. (40 refs)

78-6717 Chemical and Toxicological Studies on Bracken Fern, *Pteridium aquilinum* Var. *latiusculum*. I. Introduction, Extraction and Fractionation of Constituents, and Toxicological Studies Including Carcinogenicity Tests. (Eng) Yoshihira, K. (Natl. Inst. Hygienic Science, Kamiyoga-1-Chome, Setagaya-ku, Tokyo, Japan); Fukuo, M.; Kuroyanagi, M.; Natori, S.; Umeda, M.; Morohoshi, T.; Enomoto, M.; Saito, M. *Chem Pharm Bull (Tokyo)* 26(12): 2346-2364; 1978.

Twenty-four pterosins and related compounds isolated from Bracken Fern, *Pteridium aquilinum* var. *latiusculum*, were tested for toxicity in HeLa cells and experimental animals. Young fronds and rhizomes were collected from various sites, extracted with organic solvents and/or water, and, in some cases, further purified by column and thin-layer chromatography. Typical isolates were a group of C₁₄ or sesquiterpenoids with a 1-indanone nucleus, termed pterosins, and their related glucosides and pterosides. Several compounds produced morphologic changes in HeLa cells within test samples for 3 days. In animal toxicity tests, the compounds were administered to DDD mice, Wistar male rats, and male Hartley guinea pigs by feeding (5 or 15 g/day), injection (0.2 or 0.3 ml/20 g), or gastric intubation (0.4 ml/g). Four of the pterosins were toxic for the mice as were several crude bracken fractions. A frequent finding was injury to actively dividing cells, including involution of thymus and cytotoxic changes in the lymph follicles and crypts of the small intestine. Long-term carcinogenesis feeding experiments were undertaken in male Wistar and Fischer rats, with 4-340 mg/day of fern isolates added to their diet for 35-205 days. Crude extracts of fronds and rhizomes induced tumors preferentially in Fischer rats, particularly in the ileum, but purified fractions and extracts did not. Ileal tumors included adenomas and papillary carcinomas. (49 refs)

78-6718 Comparison of the Oncogenicities of Four Structurally Isomeric N-Hydroxyanthrins

Teller, M. N. (Donald S. Walker Lab., Sloan-Kettering Cancer Res., 145 Boston Post Road, Rye, NY, 10580); Jinger, J. M.; Zvilichovsky, G.; Watson, A. A.; McDon-J. J.; Stohrer, G.; Brown, G. B. *Cancer Res* 38(7): 2042; 1978.

oncogenicities of all of the four possible isomeric hydroxyxanthines were compared in rats. 3-Hydroxyxanthine induced sarcomas in 88%-100% of the rats when administered sc at a dose of 1 mg 3x/wk for 8 wk; at a dose of 0.1 mg (same route and schedule), both sarcomas and fibromas appeared in 25%-58% of the rats given the isomer. Total 3-hydroxyxanthine doses ranging from 12-66 mg/rat induced sc tumors in 25%-100% of the rats, some of which also developed hepatocellular carcinomas. The oncogenicity of 7-hydroxyxanthine approached that of the 3-isomer, particularly in the induction of liver tumors. At 1 mg given 3x/wk for 8 wk, the 7-isomer induced both sc and liver tumors in 40% of the rats compared with 89% for the 3-isomer. Liver tumors were induced in 45% and 65% of the rats given 1.0-mg ip injections (3 x/wk for 8 wk) of the 7- and 3-isomers, respectively. The difference was not significant. 9-Hydroxyxanthine showed no oncogenicity, which is the result of the stability of its sulfate ester. 1-Hydroxyxanthine was not oncogenic in assay; it has induced tumors at doses > 1.0 mg, but there is no explanation for its oncogenicity by any correlation of chemical properties with those of the 3- and 7-isomers. (25 refs)

19 **The Tumor Promoting Action of Diterpene Esters In Vivo Correlates with Their Potency to Release Prostaglandins from Macrophages In Vitro (Meeting Abstract).** (Eng) Brune, K. (Dept. Pharmacology, Birmum, CH-4056 Basel, Switzerland); Hecker, E. *Experientia* 34(7): 922; 1978. (no refs)

720 **Snuff and Cancer of the Oral Cavity.** (Swe) Axelsson, T. (Institutionen for oral kirurgi, Lunds universitet, S-214 21 Malmo, Sweden); Mornstad, H.; Sundin, B. *Lakartidningen* 75(22): 2224-2226; 1978.

The relationship between snuff and oral cancer was studied in a series of 375 oral cancer patients found in Sweden during 1971. Forty-one patients admitted using snuff at the time of diagnosis, 8 had been snuff-dippers, 18 denied ever using snuff, and 109 admitted using tobacco in other ways; there were no data on tobacco use for 199 patients. A relationship between snuff and cancer was documented in 14 patients and found probable in 19. In 67.3% of the verified snuff-dippers, the cancer was localized to regions of the oral cavity where the snuff had been placed. Most patients, both in the entire group and the snuff-dippers, were in the age bracket 40-80 yr. The percentage of snuff-related tumors increased with advancing age. The findings indicate that snuff-dipping

increases the cancer risk at the site where the snuff is placed five- to sixfold. (7 refs)

78-6721 **The Role of Aflatoxin in Primary Liver Cancer (Meeting Abstract).** (Eng) Pang, R. T. (Dept. Internal Medicine, Univ. Indonesia, Jalan Tanah, Abang Enpat, Indonesia). In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 39; 1977. (no refs)

78-6722 **Liver Carcinogenesis in Rats after Aflatoxin B₁ Administration. A Light- and Electron-Microscopic Study.** (Eng) Novi, A. M. (Pathologisches Institut, Universitat Dusseldorf, Moorenstrasse 5, D-4000 Dusseldorf, W. Germany). *Curr Top Pathol* 65: 115-163; 1977.

Changes induced in the liver of female Wistar II rats by aflatoxin B₁ (AFB₁; 25 µg/day, 5 days/wk for 8 wk, by gastric intubation) were examined by light (LM) and electron microscopy (EM). Animals were sacrificed weekly during the first 8 wk and then at various intervals up to 66 wk. By 16 wk, only an increased number of megalocytes was detected by LM. EM revealed profound changes in the liver parenchyma. Depending on their location within the liver lobule, hepatocytes underwent two main patterns of modification. While periportal hepatocytes rapidly degenerated, more peripherally located parenchymal cells first showed a high proliferation of the endoplasmic reticulum, similar to that seen in hepatocytes engaged in drug metabolism; then they underwent degeneration in a pattern similar to that seen in the periportal cells. The endoplasmic reticulum was affected first, followed by depletion of glycogen and alteration of Golgi zones and then by damage to mitochondria and the cell membrane. Between 17 and 37 wk, LM revealed the appearance of small periportal foci of newly born hepatocytes that subsequently degenerated and a progressive disappearance of megalocytes. Sublobular changes resembling those seen previously were seen by EM. After 42 wk, LM showed a diffuse proliferation of periportal hepatocytes, which resulted in the appearance of islands of neoplastic cells. Degenerative changes in the new hepatocytes consisted either in addensations of their basophilic cytoplasm toward the periphery of the cell or in vacuolation of their cytoplasm. EM revealed changes corresponding to those observed at the cellular level. Thus, the lag time preceding the appearance of recognizable cancer is characterized by a constant spectrum of ultramicroscopic changes. Since the cells showed identical phenotypical changes after AFB₁ was discontinued, it is postulated that the cells have an endogenous mechanism for the storage and release of AFB₁. (49 refs)

78-6723 **Ultrastructure of Hepatic Neoplasia.** (Eng) Butler, W. H. (Imperial Chemical Industries, Ltd.,

s Fed 2-Acetylaminofluorene Followed by Phenobarbital. (g) Pugh, T. D. (Dept. Pathology, Univ. Wisconsin Sch. Medicine, Madison, WI, 53706); Goldfarb, S. *Cancer Res* 38(12): 4450-4457; 1978.

attempt was made to characterize the hyperplastic hepatocellular islands that develop during chemical carcinogenesis and to determine whether hepatocellular carcinomas develop from progenitor cells through a sequential process. Buffalo rats 30 days old were fed a 30% protein diet containing 0.02% 2-acetylaminofluorene (2-AAF) for 4 wk followed by 0.05% phenobarbital in the same diet for 39 wk, a combination that induced hepatocellular neoplasms in 5/6 and 195 phenotypically altered islands. To determine relations between cell replication rates and other traits of islands, ³H-thymidine-labeling indices and the activities of three enzymes were studied in serially cut cryostat sections. The enzymes were ATPase, glucose-6-phosphatase (G6P), and γ -glutamyl transpeptidase (GGT). Among the islands, 89% were GGT+, 71% were ATP-, and only 35% were G6P+; those islands with variations in all three enzymes had significantly higher ³H-thymidine-labeling indices than islands with only one or two variations. The acquisition of GGTase activity was the best marker for hyperplastic islands, although decreased G6P activity characterized a small subpopulation of islands with higher rates of cell replication. It is concluded that the increasing phenotypic diversity of islands associated with increasing rates of cell replication supports the clonal hypothesis of carcinogenesis. (15 refs)

78-6730 Mechanism of In Vitro Mutagenic Activation and Covalent Binding of N-Hydroxy-2-Acetylaminofluorene in Isolated Liver Cell Nuclei from Rat and Mouse. (Eng) Sakai, S. (Biochemical Pharmacology Section, Lab. Chemical Pharmacology, Developmental Therapeutics Program, Div. Cancer Treatment, NIH, Bethesda, MD, 20014); Reinhold, C. E.; Thorgeirsson, S. S. *Cancer Res* 38(7): 2820-2826; 1978.

in vitro mutagenic activation (MA) of N-hydroxy-2-acetylaminofluorene (N-OH-AAF) in *Salmonella typhimurium* strain TA1538 by isolated rat and mouse liver cell nuclei and the in vitro covalent binding of N-OH-AAF to nuclear nucleic acids and proteins were studied. Mutagenic MA of N-OH-AAF and 2-acetylaminofluorene by liver cell nuclei was observed. The MA of N-OH-AAF by mouse nuclei was 6-7 times greater than that by rat nuclei (2,300 vs 350 revertants/10 μ g N-OH-AAF). The mutagenicity of N-OH-AAF was increased twofold when NADPH was included with the nuclei; this increase was inhibited by α -naphthoflavone. The carboxylesterase:amidase inhibitor paraoxon (10^{-5} M) completely inhibited the MA of N-OH-AAF in the 100,000 g supernatant from rat liver (RLS), whereas 50% inhibition of N-OH-AAF mutagenesis was observed when adenosine 3'-phosphate 5'-phosphosulfate was added to the

RLS. The addition of RLS to mouse liver nuclei resulted in a decreased mutagenicity of N-OH-AAF that could be further inhibited by paraoxon. On the other hand, the addition of this supernatant to rat liver nuclei resulted in an increase in N-OH-AAF mutagenicity that was insensitive to inhibition by paraoxon. Vitamin E and butylated hydroxytoluene (10^{-2} M) inhibited the MA of N-OH-AAF by 20%-30% in the mouse nuclear preparation, whereas vitamin C at the same concentration increased N-OH-AAF mutagenesis two- to threefold. Addition of cysteamine and methionine (10^{-2} M) to the mouse nuclear preparation resulted in 60% and 30% inhibition, respectively, of N-OH-AAF mutagenesis. The first step in the in vitro MA of N-OH-AAF by isolated rat and mouse liver cell nuclei and/or RLS and the in vitro covalent binding of N-OH-AAF to nuclear nucleic acids and protein occur primarily via deacetylation of the carcinogen by either the membrane-bound amidase or the N,O-acyltransferase in the supernatant fraction. (48 refs)

78-6731 Metabolism of 2-Acetylaminofluorene in Primary Cultures of Rat Hepatocytes (Meeting Abstract). (Eng) Spilman, S. D. (Dept. Environmental Toxicology, Univ. California, Davis, CA, 95616); Byard, J. L. *Pharmacologist* 20(3): 175; 1978. (1 ref)

78-6732 Influence of Chemical Carcinogens on the Physiological Effects of Adrenomimetics. (Rus) Gurkalo, V. K. (Lab. Chemical Carcinogens, N. N. Petrov Res. Inst. Oncology, Leningrad, USSR); Pliss, G. B. *Vopr Onkol* 24(4): 49-53; 1978.

The carcinogen-induced activation of α - and β -adrenoreceptors was studied in isolated rat and guinea pig atria. Exposure of the atria to the hepatocarcinogens 2-acetylaminofluorene (AAF: 1 μ g/ml) or diethylnitrosamine (DENA: 0.01 μ g/ml) increased the rate and amplitude of spontaneous contractions. At large concentrations, AAF and DENA (10-50 and 1-10 μ g/ml, respectively) inhibited the spontaneous contractions completely. AAF selectively inhibited the inotropic effect of norepinephrine in rat atria, but not in guinea pig atria. DENA had a similar effect on rat and guinea pig atria. In rat atria, sensitivity to noradrenaline decreased with increase in AAF concentrations. (15 refs)

78-6733 Nitrates and Nitrites in the Human Diet (Letter to Editor). (Eng) Hartman, P. E. (Mergenthaler Lab. Biology, Johns Hopkins Univ., Baltimore, MD, 21218). *Science* 202(4365): 260; 1978.

Of the gastric nitrite in healthy individuals, 28% is probably derived from cured meats as opposed to the < 20% quoted previously. Cured meats are suspect teratogens, and strong correlations have been demonstrated between nitrite ingestion, gastric nitrite, and stomach cancer. Heavy fertilization

may be increasing the nitrate content of spinach and tomatoes. (10 refs)

- 78-6734 Chronic Inhalation Toxicity of 2-Nitropropane in Rats (Meeting Abstract).** (Eng) Griffin, T. B. (No affiliation given); Benitz, K. F.; Coulston, F.; Rosenblum, I. *Pharmacologist* 20(3): 145; 1978. (no refs)

- 78-6735 Studies on the Immunological Response in Rats with Experimental Urinary Bladder Carcinoma.** (Jpn) Nishijima, T. (Dept. Urology, Osaka City Univ. Medical Sch., Osaka, Japan). *Jpn J Urol* 69(8): 1044-1054; 1978.

Experimental bladder tumors were induced in Wister rats by po administration of 0.025% N-butyl-N-(4-hydroxybutyl)nitrosamine (BBN), and changes in the bladder histopathology and in the mitogen response of lymphocytes with time were investigated. There were no significant differences in body wt between the experimental and control groups. However, there was a significant increase in bladder wt and a significant decrease in thymus wt ($p < 0.01$) in the experimental group. At 12 wk after BBN administration, hyperplasia began to develop in the bladder mucous epithelium; there was gradual growth in a papillary form and, after 24 wk, transitional cell carcinomas were found in all rats. The stimulation index values of the lymphocytes from the cervical and mesenteric lymph nodes began to decrease significantly ($p < 0.005$) from the time hematuria was observed macroscopically (about 21 wk after BBN treatment), but there were no significant differences between the responses to phytohemagglutinin (PHA) and concanavalin A (Con A). The SI values of the thymic lymphocytes decreased significantly ($p < 0.01$) from the time hyperplasia was found in the bladder mucous membrane (about 15 wk), before the significant reduction in the mitogen responses of lymph node lymphocytes. The decrease was much greater with Con A than with PHA stimulation. (55 refs)

- 78-6736 Malignancies Associated with Immunosuppressive or Cytotoxic Therapy.** (Eng) Penn, I. (Dept. Surgery, Univ. Colorado Medical Center, 4200 E. Ninth Ave., Denver, CO, 80220). *Surgery* 83(5): 492-502; 1978.

The increased risk of malignancy in five groups of patients receiving immunosuppressive or cytotoxic therapy is discussed. There is a 36% incidence of neoplasia in transplant recipients who received organs from donors with cancer. Transplant patients have a 6% risk of developing de novo tumors at some time after transplantation; this frequency is 100 times greater than that for the general population in the same age range. Patients who undergo transplantation as treatment for malignancies have variable recurrence rates: 62.5% following liver replacement for hepatic malignancies, and 31% after bone marrow grafting for acute leukemia. The

treatment of renal malignancies by kidney transplantation has a 53% recurrence rate if the interval between treatment of the cancer and transplantation is ≤ 1 yr; no recurrence have been noted with longer waiting periods. Neoplasms have also occurred in nontransplant patients treated with immunosuppressive agents. Cancer patients treated with chemotherapy may subsequently develop second or third tumors. A statistically significant increase in the incidence of acute myeloid leukemia has been noted in patients given cytotoxic therapy for multiple myeloma, Hodgkin's disease, and ovarian cancer. Thus, immunosuppressive and cancer chemotherapeutic agents should be used with caution, and the long-term use of immunosuppressive agents in nontransplant patients with nonmalignant disease should be restricted to those in whom all other forms of therapy have failed. (10 refs)

- 78-6737 Depression by Ethionine of Phosphorylation and Oxidation in Hepatic Mitochondria.** (Eng) Higgins, E. S. (Dept. Biochemistry, Medical Coll. Virginia, Virginia Commonwealth Univ., Richmond, VA, 23298); Friend, W. H.; Rogers, K. S. *Experientia* 34(5): 578-579; 1978.

The immediate effect of ethionine on the ATP generating capacity of male Sprague-Dawley rat hepatic mitochondria was investigated in vivo and in vitro. The induction of hepatic steatosis and the suppression of hepatic ATP levels, protein synthesis, and gluconeogenesis subsequent to ethionine administration may be consequences of its interference with the mitochondrial phosphorylation of ADP. The mitochondrial dysfunction is not a direct action of ethionine on the organelle. (19 refs)

- 78-6738 Saccharin, Cancer, and Calories (2 Letters to the Editor).** (Eng) Campbell, T. C. (Federation of American Societies Experimental Biology, Bethesda, MD 20014). *Science* 202(4365): 260-262; 1978.

In comments on a report based on estimates of the Food and Drug Administration (FDA) linking bladder cancer to saccharin ingestion, species differences in response to carcinogens are discussed. The responses can vary by three or four orders of magnitude. The assumption that noncaloric sweeteners reduce calorie consumption is also questioned. In reply, the FDA's risk estimate is also questioned and several examples of how uses of saccharin do decrease calorie intake are cited. (6 refs)

- 78-6739 Histological Conformity of Implantation Tumors Produced by Kidney Cell Lines Derived from Dimethylnitrosamine-treated Rats with Dimethylnitrosamine-induced Renal Mesenchymal Tumors.** (Eng) Hard, G. C. (Fels Res. Inst., Temple Univ. Sch. Medicine)

Philadelphia, PA, 19140). *Cancer Res* 38(7): 1974-1978; 1978.

Histology of five implantation tumors induced in Wistar by the deposition of cultured cell lines derived from diethylnitrosamine (DMN)-treated rats is described and compared with the morphology of the predominant kidney plasma induced in vivo by a single high dose (60 mg/kg) of DMN. The cell lines leading to growth upon implantation of long-established, continuously growing cultures obtained from either a DMN-induced renal mesenchymal tumor or from rats treated shortly (2 hr) before with a carcinogenic dose (60 mg/kg) of DMN. The latter cultures expressed morphological transformation at subcultures 5 or 6. All of the implantation tumors were of the mesenchymal type, comprising a range of cell forms that included fibroblastlike spinules, smooth muscle fibers, and giant cells, which resembled common aspects of the parent mesenchymal tumors induced in the rat kidney by DMN. The deposition of cells renally illustrated the survival of remnants of preexisting neoplasms as epithelial profiles scattered through the proliferating malignant tissue, a feature most characteristic of the parent tumor. The results confirmed the malignant nature of the various cell lines tested, and they were consistent with the premise that the in vivo-in vitro system selected cells in culture that represented the same target population from which the renal mesenchymal tumors were derived in vivo. (16 refs)

7740 Macroautoradiographic Study of the Distribution of ^{14}C -Dimethylnitrosamine in Mice and Rats. (Eng) Takahashi, G. (Dept. Pathology, Chest Disease Res. Inst., Kyoto Univ., Kyoto, Japan). *Bull Chest Dis Inst Kyoto Univ* 11(1/2): 81-85; 1978.

Dimethylnitrosamine (DMN: 5 μCi) was administered intravenously to pregnant ICR mice on day 18 of gestation, and the distribution of radioactivity in the mice and their fetuses was determined by macroautoradiography of serial sagittal sections. Intense radioactivity was noted 30 min after intravenous administration in the liver, kidneys, bone marrow, glandular organs, skin appendages, pancreas, mucous membranes of the oral cavity, intestines, and glandular stomach. This pattern was maintained for up to 2 hr. The distribution of radioactivity was similar 30 min after per os administration of the carcinogen with intense labeling noted primarily in the liver, kidney, spleen, and pancreas. Labeled DMN or its metabolites passed through the placenta and were evenly distributed in fetal liver, intestines, bone marrow, and thymus. Because the in vivo carcinogenic activity of DEN is due to its demethylated metabolites, a comparison of radioactive label concentration in tissue susceptibility to DMN has limited value. (18 refs)

7741 Occurrence of Methylguanidine and Agmatine in Foods. (Eng) Kawabata, T. (Dept. Biomed-

ical Res. Food, Natl. Inst. Health, Tokyo, Japan); Ohshima, H.; Ino, M. *IARC Sci Publ* (19): 415-423; 1978.

Various fresh and processed foods were analyzed by a new series of methods for the presence of two mutagenic guanidino compounds, nitrosated methylguanidine (MG) and nitrosated agmatine (AG). The methods consisted of trichloroacetic acid extraction, separation of the extract into fractions by ion-exchange chromatography, colorimetric analysis of each fraction, and identification of the compounds by gas chromatography-mass spectrometry. Sixty-five fresh and smoked or cured fish and meat samples were analyzed. MG was not detected or was negligible in all samples except in three smoke-dried fish products: skipjack (178 mg/kg), chubmackerel-horse mackerel mixture (56 mg/kg), and sardines (37.8 mg/kg). These seasoning products, which are declining in use, are only used in small quantities and cannot be related to the high gastric cancer incidence in Japan. Relatively high AG levels (mg/kg) were found in fresh abalone and top shell (40-240) and in some processed fish, including fresh Alaskan pollack roe (72.5), salt-dried herring (217), salted squid (650), dried filefish (112), and smoked scallops (44.8). It is suggested that the high MG levels found in earlier food studies were due to the use of metallic salts, which catalyzed the oxidation of creatine or creatinine to give MG as an artifact. (16 refs)

78-6742 Ornithine Decarboxylase and Polyamines: Relationship to N-Nitrosamine Carcinogenesis. (Eng) Farwell, D. C. (Univ. New Hampshire, Durham, NH). *Diss Abstr Int B* 39(2): 706; 1978. (no refs)

78-6743 Effects of Carcinogenic and Non-carcinogenic Chemicals on Plasma Esterases in BALB/c Mice. (Eng) Tyndall, R. L. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Clapp, N. K.; Davidson, K. A.; Colyer, S. P.; Burtis, C. A. *Chem Biol Interact* 23(2): 159-169; 1978.

Esterase profiles of plasma from female BALB/c mice treated with a variety of carcinogenic, weakly carcinogenic, and non-carcinogenic chemicals were analyzed to determine whether these chemicals produce esterase alterations similar to those of diethylnitrosamine and butylated hydroxytoluene and if such alterations are more sensitive indicators of these chemicals than standard clinical chemistry tests. Mice that received diethylnitrosamine (40 mg/liter), dinitrosopiperazine (100 mg/liter), dipropylnitrosamine (100 mg/liter), dimethylhydrazine (50, 100, or 800 mg/liter), urethane (10,000 mg/liter), or dimethyldinitrosopiperazine (50, 100, or 800 mg/liter) in their drinking water for 7 days had similarly altered plasma esterase profiles. These consisted of increased activity in four esterase bands. The increased activity persisted in some bands after cessation of carcinogen exposure. Exposure to high concentrations of the weakly carcinogenic or

noncarcinogenic compounds nitrosohydroxyproline (100 mg/liter), nitrosomethoxymethylamine (100 mg/liter), 1-nitroso-4-methylpiperazine (100 mg/liter), nitroso-2,6-dimethylpiperidine (100 mg/liter), or ethyl methanesulfonate (200 mg/liter) caused no obvious plasma esterase alterations. Carbon tetrachloride ingestion (800 mg/liter) increased the activity in one esterase band and decreased the activity in a second band. Analysis of serum from test mice for levels of SGOT, alkaline phosphatase, lactate dehydrogenase-lactate substrate, and *D*- γ -glutamyl transpeptidase did not differentiate between mice treated with selected carcinogens and those treated with noncarcinogens and/or carbon tetrachloride. (33 refs)

- 78-6744 Low Level of Unscheduled [³H]Thymidine Incorporation into Hyperplastic Nodule and Hepatoma Cells after Exposure to Dimethylnitrosamine In Vivo.** (Eng) Ogawa, K. (Dept. Pathology, Sapporo Medical Coll., S.1, W.17, Chuo-ku, Sapporo 060, Japan); Enomoto, K.; Yoshida, Y.; Onoe, T. *Chem Biol Interact* 23(2): 259-262; 1978.

In order to investigate why hyperplastic liver nodules that arise prior to and with hepatoma development are resistant to the cytotoxic and growth inhibitory effects of various hepatocarcinogens, the effect of *in vivo* exposure to dimethylnitrosamine (DMN) was investigated in male Wistar rats bearing these lesions. The lesions had been induced by continuous administration of diethylnitrosamine in the drinking water (6 mg/kg/day) for 4 wk or by 3'-methyl-4-(dimethylamino)azobenzene in the diet (0.06%) for 12 wk. They were then maintained carcinogen-free for 28 and 8 wk, respectively. Rats were treated with 30 mg/kg DMN ip; 4.5 hr later, chilled Eagle's minimal essential medium was perfused through the portal vein. DNA synthesis was studied in excised livers that had been incubated with ³H-thymidine. Without prior DMN treatment, fairly large numbers of grains were localized on the cytoplasm and nuclei of both normal liver and hyperplastic nodules, probably due to incorporation of tritiated moieties from catabolized ³H-thymidine into macromolecules other than nuclear DNA. After DMN treatment, normal liver cells showed increased number of grains on the nuclei. The two hyperplastic nodules and one hepatoma examined showed no increase in ³H-thymidine incorporation into nuclei after DMN treatment, whereas surrounding liver showed a distinct increase of grains on the nuclei, as did normal liver cells. These results show that *in vivo* DMN treatment decreases unscheduled DNA synthesis in hyperplastic nodules and hepatomas, suggesting that these lesions may be capable of escaping from carcinogen-induced DNA damage *in vivo*. (16 refs)

- 78-6745 Transformation of Mouse Mammary Glands in Whole Organ Culture by Carcinogenic Aromatic Amines and Amides (Meeting Abstract).** (Eng) Tonelli, Q.

J. (Inst. for Cancer Res., Fox Chase Cancer Center, Philadelphia, PA, 19111); Sorof, S. *J Cell Biol* 79(2, part 2): 73a; 1978. (no refs)

- 78-6746 Deuterium Isotope Effect on Rat Liver DNA Damage In Vivo by Dimethylnitrosamine.** (Eng) Oshiro, Y. (Searle Labs., Box 5110, Chicago, IL, 60680); Sarma, D. S.; Zielinski, W. L. *Chem Biol Interact* 23(2): 263-266; 1978.

The deuterated analog (DMN-d₆) of dimethylnitrosamine (DMN) proved to be less effective than DMN in inducing *in vivo* DNA damage after ip administration to rats in doses from 0.37 to 3.20 mg/kg. Hepatic cell DNA damage after labeling with ³H-thymidine was measured by alkaline sucrose gradient sedimentation. At all dose levels, DMN-d₆ treatment resulted in a greater percentage of labeled DNA sedimenting at a mol wt > 13 x 10⁶ than DMN treatment. The distribution of slower-sedimenting DNA was always lower after DMN-d₆ than after DMN treatment. The sedimentation as heavy DNA, compared with control DNA, was taken to be a direct indication of DNA damage. The differences in the ability of DMN-d₆ and DMN to cause *in vivo* DNA damage were apparent at 10 min after administration, but they became less apparent with longer periods between treatment and sacrifice. Because DMN has to be metabolized to chemically modify DNA, it is hypothesized that the differential mutagenic capability of DMN-d₆ is due to the slower activation of this perdeuterated analog. (16 refs)

- 78-6747 Blood, Tissue and Subcellular Distribution of Radioactivity Following Administration of ¹⁴C-Dimethylnitrosamine (DMN) or ¹⁴C-Dimethylamine (DMA) in Guinea Pigs (Meeting Abstract).** (Eng) Chaudhari, A. (Pharmacology Dept., Wayne State Univ., Detroit, MI 48202); Dutta, S. *Pharmacologist* 20(3): 175; 1978. (no refs)

- 78-6748 The Significance of Some Modifying Factors During Transplacental Blastomogenesis by Ethylnitrosourea in Rabbits.** (Eng) Dimant, I. N. (Uzbek Res. Inst. Oncology and Radiology, 700 000 Tashkent USSR); Beniashvili, D. Sh. *Neoplasma* 25(4): 453-460; 1978.

The effects of three treatments, organospecific immunity, chronic irritation of the peripheral nerve, and methylnitrosourea (MNU), on the development of ethylnitrosourea (ENU)-induced transplacental blastomogenesis were studied in chinchilla rabbits. Groups of 36 weaned rabbits, whose mothers had been treated during pregnancy with ENU, either served as controls or underwent one of the three treatments. The immunized animals (Group 1) were treated three times with antigens from the homologous tissue of the sciatic nerve

Group 2, the left sciatic nerve was chronically irritated by applying a nylon squeezing free ligature, the ends of which were attached to the femoral muscle. Contraction of this muscle produced irritation of the nerve. The MNU-treated animals (Group 3) received a weekly iv dose of 10 mg/kg for 5 mo. All animals were observed for 4 yr. The total number of tumors developing in survivors, 27 Group 1, 26 Group 2, 10 Group 3, and 36 control rabbits, was 17, 12, 4, and 10, respectively. The total number of peripheral nervous system tumors in the respective groups was 8, 7, 2, and 4. It appears that the first two treatments increased the frequency of tumor development, particularly that of peripheral nerve tumors. MNU apparently inhibited ENU-induced tumorigenesis. (25 refs)

78-6749 Studies of Experimental Brain Tumorigenesis Induced by Transplacental Administration of Diethylnitrosourea in Rats. (Jpn) Inui, S. (Second Dept. Surgery, Nara Medical Univ., Nara, Japan). *J Nara Med Assoc* 9(1): 288-306; 1978.

Wistar rats were given a single transplacental dose of ethylnitrosourea (ENU: 10-75 mg/kg), 4-hydroxyaminoquinoline N-oxide (4-HAQO: 7 mg/kg), or N-bis(2-hydroxypropyl)nitrosamine (DHPN: 2,000 mg/kg) on gestation days 18 or 20, and DNA synthesis and histopathological changes in the fetal brain were analyzed. Four hours after administration on day 18, ENU, 4-HAQO, and DHPN inhibited DNA synthesis by 87.7%, 85.6%, and 87.3%, respectively, with the inhibition due to ENU being dose-dependent. In ENU- and 4-HAQO-treated rats, the inhibition persisted for 48 hr, followed by a stimulation of synthesis at 72 hr after administration. Brain tumors were induced by ENU but not by 4-HAQO and DHPN. The incidence in males and females ranged from 36.4% to 66.7% at 23 wk and from 90.3% to 100% at 30 wk. In order of incidence, the tumors were oligodendrogliomas, astrocytomas, mixed gliomas, anaplastic gliomas, medulloblastomas, and trigeminal neurinomas. Atypical cell foci were observed as early as 10 wk, and the incidence of these foci was higher than that of the tumors at 23 wk and lower at 30 wk. The labeling index of the oligodendrogliomas, astrocytomas, anaplastic gliomas, neurinomas, and atypical cell foci was 1.92, 3.16, 7.52, 3.49, and 0.075, respectively. The atypical cell foci may represent preneoplastic lesions. (41 refs)

78-6750 Influence of Repeated Liver Regeneration on Hepatic Carcinogenesis by Diethylnitrosamine in Mice. (Eng) Pound, A. W. (Dept. Pathology, Univ. Queensland, Brisbane, Australia); McGuire, L. J. *Br J Cancer* 37(4): 595-602; 1978.

The effect of repeated doses of CCl₄ ip at 5-wk intervals on the tumorigenic response of diethylnitrosamine (DEN)-treated random-bred male Quackenbush mice was determined.

The mice were divided into six groups: 1 (a), 0.25 ml/kg CCl₄ ip; (b), 24 hr later, 80 mg/kg DEN ip and, 5 wk later, repeated doses of CCl₄; 2, regimen 1b; 3, regimen 1a + DEN; 4, DEN; 5, repeated CCl₄; 6, regimen 1a. No tumors were noted in mice given single or multiple injections of CCl₄. DEN + repeated CCl₄ injections increased the tumor yield in the liver over that occurring in animals treated with DEN alone. A single hepatonecrotic dose of CCl₄ 24 hr before DEN treatment also increased the number of tumors produced. The effects of the pre-DEN CCl₄ dose and the post-DEN CCl₄ doses were additive and were therefore independent of each other. The neoplasms produced were hepatocellular tumors and cholangiomas. Kidney tumors (papillary adenomas and adenocarcinomas) were also noted in the CCl₄ + DEN-treated mice. These findings indicate that the liver in the regenerative phase after a hepatonecrotic CCl₄ dose is more susceptible than normal liver to the initiating action of DEN and that repeated doses of CCl₄ after DEN provide a promoting stimulus. (18 refs)

78-6751 Effects of the Carcinogen, N-Methyl-N'-Nitro-N-Nitroguanidine (MNNG), on Differentiation of Human Prostate Epithelium in Organ Culture (Meeting Abstract). (Eng) Sanefuji, H. (Dept. Pathology, Univ. Maryland Sch. Medicine, Baltimore, MD); Heatfield, B. M.; Trump, B. F. *J Cell Biol* 79(2, part 2): 71a; 1978. (no refs)

78-6752 Binding of 1-Methyl-1-nitrosourea (MNU) and 1-Methyl-3-nitro-1-nitrosoguanidine (MNNG) to L1210 Cell Histones (Meeting Abstract). (Eng) Pinsky, S. D. (V. Lombardi Cancer Res. Center, Georgetown Univ., Washington, DC); Tew, K. D.; Woolley, P. V. *J Cell Biol* 79(2, part 2): 113a; 1978. (no refs)

78-6753 Mycosis Fungoides, Nitrogen Mustard and Skin Cancer. (Eng) Du Vivier, A. (St. Mary's Hosp., London W2, England); Vonderheid, E. C.; Van Scott, E. J.; Urbach, F. *Br J Dermatol* 99(1): 61-63; 1978.

The development of epithelial malignancies prior to or after treatment with topically applied nitrogen mustard (mechlorethamine, HN₂) was determined in 202 patients (135 men, 67 women) with mycosis fungoides and its Sezary variant. The records of the patients, who had been followed for up to 9 yr, were reviewed. Epithelial cancers developed in 24 patients, 3 of whom were excluded from further study because of previous treatment with arsenic. The remaining 21 patients (12 men, 9 women) had a mean age at onset of lesions of 64 yr. The mean time between initiation of HN₂ therapy and detection of epithelial neoplasms other than actinic keratoses was 4.0 yr. Of eight patients who had lesions removed prior to HN₂ therapy (Group I), five developed additional lesions after HN₂ treatment [4 squamous cell carcinomas (SCC), 4

keratoacanthomas, and 1 basal cell carcinoma]. Three of these patients had multiple tumors. The other 13 patients (Group II), who had no evidence of epithelial lesions initially, subsequently developed actinic keratoses (4 patients) and 7 SCC, 3 keratoacanthomas, and 2 basal cell carcinomas (9 patients, 3 of whom developed multiple tumors). Two of the SCC were found in locations protected from solar radiation, which offers convincing evidence that topically applied HN₂ is a carcinogen or cocarcinogen. These lesions, which appeared 6-7 yr after the start of treatment, corresponded to the use of approx 14 g HN₂/4 yr (SCC on scrotum) and 16 g/7 yr (SCC on buttock). (7 refs.)

78-6754 In Vivo Synthesis of 1-Naphthyl N-Methylnitrosocarbamate (Nitrosocarbaryl) in the Rat and Guinea Pig (Meeting Abstract). (Eng) Rickard, R. W. (Graduate Center Toxicology, Univ. Kentucky, Lexington, KY, 40506); Dorrough, H. W. *Pharmacologist* 20(3): 146; 1978. (no refs)

78-6755 Inhibition of the Alkylation of Nucleic Acids and of the Metabolism of 1,2-Dimethylhydrazine by Aminoacetonitrile. (Eng) Pegg, A. E. (Dept. Physiology, Milton S. Hershey Medical Center, Pennsylvania State Univ., Hershey, PA, 17033). *Chem Biol Interact* 23(2): 273-279; 1978.

Administration of aminoacetonitrile (AAN) ip to female Sprague-Dawley rats before ip injection of ¹⁴C-1,2-dimethylhydrazine (DMH) altered the latter's metabolism to ¹⁴CO₂ and ¹⁴C-azomethane. The amount of ¹⁴CO₂ expired by treated rats was determined by drawing air through their cages, collecting it in gas-filled wash bottles, and measuring the radioactivity present in the bottles. When 100 mg/kg AAN was administered before 2 mg/kg ¹⁴C-DMH, CO₂ production 8 hr later was reduced from 28% to 4%, and azomethane increased from 1.3% to approx 50% of the administered DMH dose. Results were similar when the ¹⁴C-DMH dose was reduced to 0.5 mg/kg. When ¹⁴C-DMH dose was increased to 50 or 200 mg/kg, azomethane expiration rose to 33% and 41% of the administered dose, respectively, and ¹⁴CO₂ exhalation dropped to 11% and 6%. Alkylation of rat DNA at the 7-methylguanine position after administration of 2 mg/kg DMH was inhibited 95% in the liver, 89% in kidney, and 57% in colon by pretreatment with 100 mg/kg AAN. O⁶-methylguanine production was inhibited to a similar extent. The difference in the degree of inhibition of colon DNA suggests that at least one critical step in the conversion of DMH to a methylating agent for the colon occurs outside the liver. (26 refs)

78-6756 Chemically Induced Gangliogliomas in the Opossum (*Didelphis virginiana* Kerr) (Meeting

Abstract). (Eng) Jurgelski, W. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC); Hudson, M.; Henry, J. M.; Falk, H. L. *J Neuropathol Exp Neurol* 37(5): 638; 1978. (no refs)

78-6757 Molecular Basis for the Neoplastic Transformation of Neuroectodermal Cells by Nitrosoureas. (Eng) Chang, M. J. (Ohio State Univ., Columbus, OH, 43210). *Diss Abstr Int B* 39(2): 701; 1978. (no refs)

78-6758 Long-Term Effects of Phenobarbitone-Na Male Fischer Rats. (Eng) Butler, W. H. (Safety Medicines Dept., Pharmaceuticals Div., ICI, Ltd., Mereside, Alderley Park, Macclesfield, Cheshire, SK10 4TG, England). *Br J Cancer* 37(3): 418-423; 1978.

Preliminary results are given of an experiment in which male inbred Fischer rats were fed sodium phenobarbitone at a level of 500 ppm in the diet for 1 wk followed by 1,000 ppm for 103 wk. Of 33 rats surviving ≥ 80 wk, 11 had small foci of nodular hyperplasia; however, 1 killed at 102 wk had a larger lesion (0.75 cm) with no evidence of local invasion or metastases. All livers had evidence of parenchymal cell damage. No nodular hyperplasia was found in the controls. It is concluded that this compound is not carcinogenic in Fischer rats. (9 refs)

78-6759 Evidence from *Haemophilus influenzae* for Error-Prone Repair Systems with Different Mutation Specificities (Meeting Abstract). (Eng) Kimball, R. (Biology Div., Oak Ridge Natl. Lab., Oak Ridge, TN, 37830); Boling, M. E.; Perdue, S. W. *Mutat Res* 53(2): 209-210; 1978. (no refs)

78-6760 Effect of Pyrazole on Tumor Induction by Methylazoxymethanol (MAM) Acetate: Relationship to Metabolism of MAM (Meeting Abstract). (Eng) Zidek, M. S. (Sloan-Kettering Inst., New York, NY, 10021); Tan, Q. H. *Pharmacologist* 20(3): 174; 1978. (2 refs)

78-6761 Relative Sensitivities of Forward and Reverse Mutation Assays in *Salmonella typhimurium*. (Eng) Skopek, T. R. (Toxicology Group, Dept. Nutrition and Food Science, Massachusetts Inst. Technology, Cambridge, MA, 02139); Liber, H. L.; Kaden, D. A.; Thilly, W. G. *Proc Natl Acad Sci USA* 75(9): 4465-4469; 1978.

The relative sensitivities of the forward and reverse mutation assays in *Salmonella typhimurium* were compared with the

of 16 mutagens of both base-substitution and frameshift types. Forward mutation of 8-azaguanine resistance was measured with strains TM35 and TM677; reverse mutation of histidine prototrophy was measured with the Ames tester strains TA1535, TA1537, TA1538, TA98, and TA100. The spontaneous revertant mutant fractions ranged from 1×10^{-8} (TA1538) to 22×10^{-8} (TA100). The mean spontaneous forward mutant fraction was 1.7×10^{-5} for TM35 and 1×10^{-5} for TM677. The two approaches were found to be sensitive for all 16 mutagens; ie, induction of significant mutation occurred at similar concentrations in the forward mutation assay and in the most sensitive of the five Ames tester strains. The combined assays provide an improved means of quantitating chemomutagenic potential. (9 refs)

78-6762 **Mutagenic Properties of Quinoline.** (Eng) Hollstein, M. C. (Univ. California, Berkeley, CA). *Abstr Int B* 39(2): 673; 1978. (no refs)

78-6763 **Macroautoradiographic Assays of ^{14}C -Labeled 4-Nitroquinoline 1-Oxide in Adult and Fetal Liver.** (Eng) Takahashi, G. (Dept. Pathology, Chest Disease Res. Inst., Kyoto Univ., Kyoto, Japan); Kinoshita, K.; Uehira, K. *Bull Chest Dis Res Inst Kyoto Univ* 11(1/2): 90; 1978.

The distribution of ^{14}C -labeled 4-nitroquinoline 1-oxide (4NQO) and its metabolites in adult, pregnant, and fetal mice was studied by macroautoradiography. 4NQO (7.5 μCi , 230 μg) was administered iv to pregnant mice on days 16 or 19 of gestation or injected iv or sc in young adult mice. All animals were killed 1, 3, 5, or 24 hr later. Concentrations of radioactive substances were nearly equal in fetal tissue, placenta, and maternal blood 1 hr after iv administration. Although fetal organ distribution of radioactivity was uniform on day 16 of gestation, the 19 kidneys, urinary bladder, and intestinal tract were intensely labeled on day 19. In normal adult mice, radioactivity 1 hr after iv 4NQO administration was concentrated most intensely in the gallbladder, small intestine, lungs, kidneys, and bladder. There was moderate activity in the liver, heart, muscle, salivary glands, and blood. Radioactive substances tended to deposit at the injection site after sc administration, but they also accumulated in the gallbladder, intestine, bladder, and kidneys. No direct relationship between organ susceptibility to 4NQO and the concentration of radioactivity was observed. (10 refs)

78-6764 **Effect of a Single Injection of 4-Dimethylaminoazobenzene on Some Properties of Rat Liver Cell Mitochondria.** (Rus) Medvedeva, N. D. (Lab. Tumor Cytology, Inst. Cytology, Acad. Sciences, USSR, Leningrad, USSR); Khodosova, I. A. *Tsitologiya* 20(10): 1102-1106; 1978.

The effects of a single ip injection (250 mg/kg) of the hepatocarcinogen 4-dimethylaminoazobenzene (DAB) and its non-carcinogenic analog, 4-diethylaminoazobenzene (DEAB), on rat liver mitochondria were studied. DAB had no noticeable effect on mitochondrial and cytoplasmic malate dehydrogenase activity. Both substances caused a significant inhibition of mitochondrial cytochrome oxidase activity on the first day (to 51.3% by DAB and to 42.2% by DEAB), but there was no significant difference between the effects of the two substances. This activity returned to normal levels on day 4. The swelling-contraction properties of isolated mitochondria were reduced in the presence of succinate and ATP, even 18 days after a single DAB injection. However, DAB had no effect on mitochondria from the kidney, which is not its target organ. The changes are interpreted as the result of the action of the carcinogen on the mitochondrial membranes. (15 refs)

78-6765 **Need for Ascertaining Environmental History in Leukemia and Malformations in Children.** (Swe) Haglund, B. (Halsovardsenheten, Bassjukhuset, S-541 00 Skovde, Sweden); Kjellman, B. *Lakartidningen* 75(39): 3436-3437; 1978.

The necessity and importance of ascertaining the environmental history of children with leukemia and malformations are illustrated by case reports. A woman exposed occupationally to xylene vapors and freon in an unventilated room gave birth to a boy who was subsequently exposed to styrene, acetone, and xylene while his father was building a plastic boat. The boy developed acute lymphocytic leukemia at age 2 yr. One of the mother's colleagues gave birth to a girl with bilateral talipes equinovarus, the other to a girl with gastroschisis and exstrophy of the bladder. A 10-yr-old boy developed acute leukemia a few months after he was exposed to solvents, mainly trichloroethylene, while building plastic models. A 6-yr-old girl exposed frequently to solvents and living in a house in which the radioactivity level was about four times higher than the background level developed leukemia. The girl's grandfather, who had been occupationally exposed to solvents as an automobile mechanic, died from chronic lymphocytic leukemia. (13 refs)

78-6766 **A Study of Cyclic Nucleotide Metabolism and the Histology of Rat Liver During 3'-Methyl-4-dimethylaminoazobenzene Carcinogenesis. II. Cyclic AMP Metabolism.** (Eng) Boyd, H. (Dept. Pharmacology, Univ. Miami Sch. Medicine, P. O. Box 520875, Miami, FL, 33152); McAfee, D. A.; Rubin, J. J. *Tissue Cell* 10(3): 477-494; 1978.

Adenyl cyclase (AC) activity and cyclic AMP (cAMP) metabolism were studied in Sprague-Dawley rat livers undergoing carcinogenesis induced by dietary 3'-methyl-4-dimethylaminoazobenzene (0.06%). All treated rats developed

cholangiocarcinomas by 10 wk. AC activity, assessed from the crude membrane fraction, increased slowly for 3 wk and then rose to a peak of 565% of control values at 9 wk post-treatment. Stimulation with isoproterenol increased this peak to 640%; guanylimidodiphosphate (Gpp(NH)p) and sodium fluoride also stimulated AC activity, but glucagon had no significant effect. Enzyme activity was higher in tumor-bearing animals than in controls, and it was higher in areas of tumor involvement than in noninvolved tissue from the same animal. The patterns of responsiveness to the AC stimulators indicated that there were differences not only in the onset of hormonal stimulation of the enzyme and the transduction mechanism, but also a difference in the time at which these changes were no longer seen. The former preceded tumor induction; the latter accompanied tumor induction and persisted in tumors and uninvolved tissue. Cyclic AMP levels in treated animals showed a consistent and significant elevation (120%-250%) after wk 3, but they were strikingly increased in tumor tissue (up to 1500%). Particulate phosphodiesterase activities increased in the first 2 wk after treatment, then declined in tumor-bearing livers to levels similar to those in non-tumor-bearing livers. (64 refs)

- 78-6767 A Study of Cyclic Nucleotide Metabolism and the Histology of Rat Liver During 3'-Methyl-4-dimethylaminoazobenzene Carcinogenesis. III. Cyclic GMP Metabolism.** (Eng) Boyd, H. (Dept. Pharmacology, Univ. Miami Sch. Medicine, P.O. Box 520875, Miami, FL, 33152); McAfee, D. A.; Laumen, G.; Rubin, J. J. *Tissue Cell* 10(3): 495-504; 1978.

Cyclic guanosine monophosphate (cGMP) metabolism and guanylate cyclase (GC) activity were studied during the development of liver cholangiocarcinomas in Sprague-Dawley rats fed a diet containing 0.06% 3'-methyl-4-dimethylaminoazobenzene. Tumor incidence at 10 wk was 100%. Changes in GC activity were apparent in soluble (SF) and particulate fractions (PF), with and without added Triton X-100 (0.5%). PF activity without Triton increased slightly in the first 3 wk post-treatment, decreased to 50% of control values at 3-5 wk, and then increased to a sharp peak (300% of control values) at 10 wk. SF activity varied in a roughly inverse manner: it showed no appreciable changes for 3 wk, increased to 180% of control values during wk 3-5, then returned to normal before increasing to a sharp peak at wk 10. This peak correlated with the peak tumor incidence. The activity of SF- and Triton-activated PF enzyme increased in tumor-bearing animals, but even more so in tumor tissues. Cyclic GMP levels in uninvolved tissue dropped to 30%-40% of control values during the first 2 wk, but they increased at 4 wk and reached 160% of control values by 10 wk. SF cyclic GMP-phosphodiesterase was consistently elevated (150% of control values) in treated animals after 1 wk, as were PF activities at 1 μ M substrate. Higher substrate concentrations, decreased PF enzyme activity. The ratio of cGMP/cAMP decreased in the following order: < in tumors < uninvolved tissue < controls. (33 refs)

- 78-6768 Work-related Origin of Cancer in the Nose, Mouth, Throat, and Larynx.** (Ger) Kup, H. (Sitzungsberichte der Akademie der Wissenschaften und der Literatur, Berlin, E. Germany). *Akademie der Wissenschaften* 2(2): 20-25; 1978.

The rarity of cancer of the ear, nose, and throat in relation to the total cancer morbidity makes it difficult to single out work-related causes. Among a series of 15 patients seen by the author, there were 12 cases of laryngeal carcinoma, 2 cases of epipharyngeal cancer, and 1 case of nasal carcinoma. Only three of these patients were nonsmokers. The only possible association with occupation could be made by six laryngeal carcinoma patients: four had occupational exposure to naphthalene, and two worked in the tire manufacturing industry. It is suggested that most of these cancers result from non-work-related causes. (11 refs)

- 78-6769 Pathways of DNA Repair Operating in Yeast Cells Treated with the Pyrrolo(1,4)benzodiazepine Antitumor Antibiotics.** (Eng) Hannan, M. A. (Div. Medicinal Chemistry and Pharmacognosy, Coll. Pharmacy, Univ. Kentucky, Lexington, KY, 40506); Hurley, L. H. *J Antibiot (Tokyo)* 31(9): 911-913; 1978.

Two principal pathways for the repair of potentially mutagenic lesions produced by anthramycin, sibiromycin and tomaymycin in *Saccharomyces cerevisiae* were identified. These pathways utilize the *RAD 1* gene product concerned with the excision of UV-induced dimers and the *RAD 52* gene product concerned with the repair of ionizing radiation-induced DNA strand breaks. (9 refs)

- 78-6770 Application of Ames' Test to Dyes.** (Ita) Vicini, M. (Laboratorio Chimico Farmaceutico E. G. G. nelli S.p.A., Milan, Italy); Nosotti, A. *Boll Chim Fis* 117(7): 402-415; 1978.

Quinoline yellow, tartrazine, patent blue V, food yellow 4, carmoisine, Allura Red AC and Encyanine were tested for mutagenicity in *Salmonella typhimurium* using the Ames test. None of the substances tested induced mutations. (5 refs)

- 78-6771 A Basis for Short-term In Vivo Carcinogenesis Bioassay (Meeting Abstract).** (Eng) Cueto, J. (Carcinogenesis Testing Program, NCI, Bethesda, MD, 20014); Douglas, J. F. *Pharmacologist* 20(3): 175; 1978. (1 ref)

- 78-6772 The Formation of Dihydrodiols by the Chemical or Enzymic Oxidation of Benz(a)anthracene**

7,12-Dimethylbenz(a)anthracene. (Eng) Tierney, B. (Cancer Res. Inst., Inst. Cancer Res., Royal Cancer Hosp., Fulham Road, London SW3 6JB, England); Hewer, A. D.; MacNicoll, A. D.; Giovanni Gervasi, P.; Rattle, H.; Grover, P. L.; Sims, P. *Chem Biol Interact* 23(2): 257; 1978.

A quick simple method of obtaining small amounts of trans-dihydrodiols (trans-DHD) from benz(a)anthracene (BA) and 7,12-dimethylbenz(a)anthracene (DMBA) is described. The characterized trans-DHD of these hydrocarbons were used as reference compounds in the high-pressure liquid chromatography (HPLC) analysis of DHD mixtures formed by the metabolism of the parent hydrocarbons by rat liver microsomal preparations, and the formation of several previously unreported metabolic non-K-region trans-DHD was detected. The non-K-region DHD trans-1,2-dihydro-1,2-dihydroxy-BA and trans-3,4-dihydro-3,4-dihydroxy-BA together with small amounts of the trans-8,9- and trans-10,11-DHD, were formed when BA was oxidized in a reaction mixture containing ascorbic acid, ferrous sulfate, and EDTA. When oxidized in a similar system, DMBA yielded the K-region DHD trans-5,6-dihydro-5,6-dihydroxy-DMBA and the non-K-region DHD trans-1,2-dihydro-1,2-dihydroxy-DMBA, trans-3,4-dihydro-3,4-dihydroxy-DMBA, trans-8,9-dihydro-8,9-dihydroxy-DMBA, trans-10,11-dihydro-10,11-dihydroxy-DMBA, and a trace of the 1,2-DHD. The structures and stereochemistry of the DHD were established by comparisons of their UV spectra and HPLC characteristics with those of authentic compounds or, when no authentic compounds were available, by UV, nuclear magnetic resonance, and mass spectral analysis. The metabolic DHD were identified by comparisons of their chromatographic and fluorescence spectral characteristics with compounds of known structures. The principal metabolic DHD formed from BA and DMBA were the trans-5,6- and trans-8,9-DHD. The 1,2- and 10,11-DHD were identified as minor products of the metabolism of BA, and tentative identification of the trans-3,4-DHD as a metabolite was made from fluorescence and chromatographic data. The minor metabolic DHD formed from DMBA were the trans-3,4-DHD and the trans-10,11-DHD, but the trans-1,2-DHD was not detected. (23 refs)

78-6773 Biochemical Studies on Cultured Epithelium-like Cells. (Eng) Weisburger, E. K. (Carcinogen Metabolism and Toxicology Branch, NCI, Bethesda, MD, 20895); Idoine, J.; Elliot, J. M. In: *Rat Liver Neoplasia; Workshop on Rat Liver Neoplasia held by William F. Merrell, Co. Woods Hole, October, 1974*. William F. Merrell, Co. Woods Hole, MA: 288 pp.; 220-227; 1978.

Spontaneous transformation and aryl hydrocarbon hydroxylase (AHH) activity were monitored in eight epithelial cell lines derived from the livers of 7- to 10-day-old Fischer rats. In an initial assay, the specific AHH activity was 8.3-13.4 units/mg protein/min incubation at 37 C. Of four lines kept

in continuous culture, the AHH activity of one was high at the end of 6 mo, the activity of another showed only trace levels, and the activities of the other two remained constant. No tumors developed in animals inoculated with cells from two of the four lines, whereas one animal each developed a tumor (out of each experimental group of 8 rats) following inoculation with cells from the other two lines. AHH activity increased following treatment of cultures with benz(a)anthracene, 7,12-dimethylbenz(a)anthracene, 3-methylcholanthrene, and, to a lesser extent, with phenobarbital. The data indicate that there are biochemical differences in epithelial cell lines derived from the same host tissue and that the level of AHH activity cannot be used as a criterion for the transformed or normal status of cells in culture. (9 refs)

78-6774 Development and Characterization of Mammary Tumors and Mammary Dysplasias Induced with 7,12-Dimethylbenz(a)anthracene in Lewis Rats. (Eng) Haslam, S. N. (Univ. California, Berkeley, Berkeley, CA). *Diss Abstr Int B* 39(2): 503; 1978. (no refs)

78-6775 Induced Cancer of the Mammary Gland in Rats. (Rus) Kisliakova, N. D. (Hormonal Lab., Res. Inst. Oncology, Rostov-on-Don, USSR); Verevkina, G. A.; Ogorodnikova, L. S.; Kucherova, T. I. *Vopr Onkol* 24(8): 93-97; 1978.

A new method for the mass induction of experimental cancer of the mammary gland in rats is described. To stimulate proliferation in the mammary gland epithelium, random-bred albino rats were fed fresh human placenta (5-6 g/day), which contains large amounts of estrogens and progesterone, for 1.5 mo. One week after the start of this treatment, the animals received dimethylbenz(a)anthracene (DMBA: 10 mg, intragastrically). Ten days later rats received a second dose of DMBA and, 1 mo later, a third dose. Mammary gland tumors were induced in 22/26 rats (1 rat developed a uterine tumor). Tumors were first detected within 1.5 mo of administration of the last DMBA dose. Histologically, the tumors were similar to glandular human breast carcinoma. (7 refs)

78-6776 Developmental Stage of the Rat Mammary Gland as Determinant of its Susceptibility to 7,12-Dimethylbenz(a)anthracene. (Eng) Russo, I. H. (Experimental Pathology Lab., Dept. Biology, Michigan Cancer Foundation, 110 E. Warren Ave., Detroit, MI, 48201); Russo, J. *J Natl Cancer Inst* 61(6): 1439-1449; 1978.

Mammary gland development was compared in 80 normal Sprague-Dawley (SP) virgin rats, aged 2-112 days, and 60 SD virgin rats that, at age 55 days, were treated with 7,12-dimethylbenz(a)anthracene (DMBA: 10 mg/100 g intragastrically). Animals of both groups were killed weekly and their

mammary glands were examined in whole-mount preparations. During the first wk of postnatal life, the rat mammary gland was composed of a single lactiferous duct with 3-5 secondary branches termed terminal end-buds (TEB). The TEB grew in size and number for 21 days, then differentiated mainly into alveolar buds (AB). The AB increased steadily until age 70-84 days and differentiated into lobules with the onset of the estrous cycle at 35-42 days. Beginning at age 14 days, undifferentiated TEB's decreased in size and diameter to become terminal ducts (TD), a process that persisted until age 84 days. Therefore, in 84-day-old animals, most TEB's had differentiated into AB or TD. DMBA was administered to the rats at a time when differentiation into AB should have been rapidly progressing and TEB's decreasing. However, the number of TEB's remained constant or decreased only slightly from 5.4 to 4.6 TEB/mm². Between days 14 and 21, TEB's began increasing markedly in size (103.0 to 226.3 μ) and were termed intraductal proliferations (IDP). Mitotic activity was high in IDP's, which evolved into microtumors (400 to 750 μ) with the characteristics of adenocarcinoma. In the DMBA-treated animals, decreases in AB and lobules were correlated inversely with the increase in TEB and TD. Although these types of differentiations have not been proven conclusively in humans, there is evidence for an analogous mechanism in the development of human breast carcinoma. (32 refs)

78-6777 The Effects of UV Light and 7,12-Dimethylbenz(a)anthracene (DMBA) on C57BL Mouse Melanocytes (Meeting Abstract). (Eng) Blog, F. B. (Dept. Dermatology, Erasmus Univ., Rotterdam, Netherlands); Szabo, G. *J Cell Biol* 79(2, part 2): 34a; 1978. (no refs)

78-6778 Alteration of Pituitary Hormone Secretion in Rats During the Induction of Mammary Tumors by Dimethylbenzanthracene. (Fre) El Abed, A. (Laboratoire des Hormones polypeptidiques, C.N.R.S., 91190 Gif-sur-Yvette, France); Valens, M.; Langlois, Y.; Kerdellhue, B. *C R Acad Sci [D]* (Paris) 285(16): 1483-1486; 1977.

The secretion of pituitary hormones in various phases of the estrous cycle was studied in female Sprague-Dawley rats during the induction of mammary tumors by 9,10-dimethyl-1,2-benzanthracene (DMBA). Palpable mammary tumors usually appeared in the rats during the 14th cycle after the administration of 15 mg DMBA by gastric tube. Serum concentrations of luteinizing hormone (LH), follicle-stimulating hormone (FSH), prolactin (PRL), thyroid-stimulating hormone (TSH), and gonadotrophic hormone (GH) were measured on the morning and afternoon of the proestrus and estrus phases of the 2nd-14th cycles after drug administration. An immediate effect of the carcinogen was a decrease in the preovulation peaks of LH and FSH, observed in the afternoon of proestrus; at the same time, concentrations of PRL remained elevated. Concentrations of GH and TSH

were usually lower in the DMBA-treated rats. Apparent pituitary hormone secretion is altered in rats susceptible to mammary tumor formation by DMBA. (17 refs)

78-6779 Role of Pineal Gland in Aetiology and Treatment of Breast Cancer (Letter to Editor). (Er) Tapp, E. (Group Lab., Preston Royal Infirmary, Preston Lancashire, England). *Lancet* 2(8097): 1001-1002; 1978.

Hypersecretion of the pineal gland may be implicated in the etiology of breast cancer. Melatonin increases the secretion of prolactin, a substance implicated in the development of breast cancer. Dimethylbenzanthracene-induced mammary tumors in rats show increased malignancy if melatonin secretion is given. Moreover, suppression of the pineal gland under conditions of continuous daylight reduced the incidence and degree of malignancy of breast tumors. (5 refs)

78-6780 Morphological Changes in the Development of 7,12-Dimethylbenz(a)anthracene-induced Mammary Tumors. (Jpn) Kobuchi, K. (Dept. Surgery, Okayama Univ. Medical Sch., Shikata-cho, Okayama 700 Japan). *Okayama Igakkai Zasshi* 90(3/4): 313-334; 1978.

7,12-Dimethylbenz(a)anthracene (DMBA: 20 mg) was administered by gastric intubation to 55-day-old Sprague-Dawley rats, and morphological changes occurring during mammary gland carcinogenesis were followed. Rats (2-3) were sacrificed daily during the first week after DMBA administration, and then weekly up to 20 wk. At each stage, mammary glands were examined by the whole-mount method, optical microscopy, and electron microscopy. On day 4 after DMBA administration, mammary epithelial cells at the peripheral portion of the duct showed dilatation of the endoplasmic reticulum and Golgi apparatus. By 2 wk, hypersecretion was well-established. At 4 wk, the epithelial cells were atrophied and their intracellular organelles were regressed. However, myoepithelial cells displayed cytoplasmic extensions that surrounded mammary epithelial cells. The stroma around the ductules showed marked proliferation. At 7 wk, hyperplastic alveolar nodules were found. Among some rat stroma growing around the ductules became fibroadenomatous. At 9 wk, tumors (tubular adenocarcinomas) were palpable. Electron microscopy of the carcinoma demonstrated an increased nucleocytoplasmic ratio, nuclear irregularities, hypertrophy of the nuclear membrane, and enlargement of the intracellular space. (31 refs)

78-6781 Measurement of Aryl Hydrocarbon Hydroxylase and NADH-dependent Cytochrome c Reductase Activities in Mitogen-activated Human Lymphocytes. (Eng) Kouri, R. E. (Dept. Biochemical Oncology, Microbiological Associates, Bethesda, MD, 20014); Imblum

L.; Prough, R. A. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for Detection and Prevention of Cancer. (New York, NY): 1978. 2(part 1), 2404 pp.; 1659-1676; 1978.

Factors that influence mitogen activation and induction of hydrocarbon hydroxylase (AHH) activity in cultured human lymphocytes were examined. The most effective mitogens for lymphocyte activation were phytohemagglutinin (PHA), PHA plus pokeweed mitogen, leucoagglutinin. The AHH activity of cultured human lymphocytes was influenced by factors such as type and lot of fetal calf serum, type and concentration of mitogen, kind of inducer, initial cell concentration, and degree of mitogen activation. The degree of 3-methylcholanthrene (3-MC)-induced (and noninduced) AHH activity paralleled the degree of mitogen activation. At concentration of 2-5 μ M, benz(a)anthracene and 5,6-benzoflavone efficiently induced AHH activity to levels similar to that induced by 3-MC. Presentation of AHH activity based on level of cytochrome *c* reductase activity rather than on the number of cells or total DNA resulted in consistently lower interassay variation. Two other factors that helped yield a lower interassay variation were the use of lot No. 88973 fetal calf serum, which resulted in basal AHH levels 5-15 times above zero time fluorescence values, and the use of an initial cell concentration of 1.5×10^6 cells/ml, which routinely resulted in peak AHH activity for both 3-MC-induced and control cultures at 72 hr. (42 refs)

6782 **Induction of Tumor-Cytostatic Macrophages by Phorbol-12,13-Diester (Meeting Abstract).** (Eng) Grimm, W. (Institut f. Immunologie der Universitat, Heidelberg, W. Germany); Gerns, D. *Z Immunitaetsforsch* 5(1): 19; 1978. (no refs)

6783 **Electrochemical Investigations on Several Dimethylbenzo(a)pyrenes.** (Eng) Park, S. M. (Dept. Chemistry, Univ. New Mexico, Albuquerque, NM, 87131); Michnovicz, J. J.; Daub, G. H. *Anal Biochem* 90(1): 4-388; 1978.

Electrochemical studies were made of five dimethylbenzo(a)pyrene (DMBP) derivatives, 1,3-, 1,6-, 2,3-, 3,6-, and 5-DMBP, all of which have stronger carcinogenicity than benzo(a)pyrene (BP). A three-electrode cell housing a platinum disk working electrode, a Pt wire auxiliary electrode,

and a silver wire pseudoreference electrode were used to record the cyclic voltammetry and chronopotentiometry in (1:1) acetonitrile-benzene solns with tetra-n-butylammonium perchlorate as a supporting electrolyte. The oxidation and reduction of DMBP's were demonstrated to be reversible one-electron processes, forming the cation and anion radicals, respectively, as the electrode reaction products. Cyclic voltammograms indicated that the cation radicals are very stable. The half-lives calculated from pseudo-first-order decay rate constants were estimated to be 150, 11.4, and 7.7 sec for the 1,3-, 2,3-, and 3,6-DMBP's, respectively. The one-electron electrooxidations may be due to the stabilization of the cation radicals by methyl substituents and to the dryness of the solvent. The cation radicals of the DMBP's were more stable than those of benzo(a)pyrene (previous study). Studies of reactions of cation radicals with nucleophiles such as water and pyridine showed that blocking at the 6-position lowered the reactivity of the cation radical toward water, a mild nucleophile. Pyridine was too strong to be affected by 6-position blocking. (27 refs)

78-6784 **Benzo(a)Pyrene 7,8-Dihydrodiol-9,10-oxide Modification of DNA: Relation to Chromatin Structure and Reconstitution.** (Eng) Yamasaki, H. (Dept. Medicine, Columbia Univ. Coll. Physicians and Surgeons, 701 W. 168th St., New York, NY, 10032); Roush, T. W.; Weinstein, I. B. *Chem Biol Interact* 23(2): 201-213; 1978.

DNA was modified by in vitro reaction with the 7,8-dihydrodiol-9,10-oxide of benzo(a)pyrene (BPDE), and the effects of this modification on the in vitro reconstitution of nucleosome structures were examined. In addition, intact chromatin (CT) was reacted with BPDE, and the distribution of the covalently bound BP residues was studied. Purified duck reticulocyte DNA modified by incubation with BPDE was more susceptible to partial digestion by the single-strand-specific endonuclease *S*₁ than unmodified DNA, suggesting slight denaturation of the helix at sites of modification. CT that was reconstituted using this carcinogen-modified DNA and unmodified CT-associated proteins showed the same kinetics and extent of digestion by *Staphylococcal* nuclease and similar nucleosome profiles on sucrose density gradient centrifugation as those obtained with native CT or CT reconstituted with unmodified DNA. Polyacrylamide gel electrophoresis of DNA fragments obtained from nuclease digests of the reconstituted CT's suggested that the CT containing carcinogen-modified DNA had the same subnucleosome structure as that reconstituted with unmodified DNA. After intact duck reticulocyte CT was reacted directly with BPDE, nuclease digestion studies indicated that 65% of the carcinogen was bound to the 'open' regions of CT and 35% to 'closed' regions (ie, regions highly susceptible to digestion by *Staphylococcal* nuclease vs regions relatively resistant to digestion). The results indicate that although covalent binding of a BP derivative to DNA produces local

distortions in conformation of the helix, this modification does not appear to interfere with the ability of the DNA to associate with histones to form nucleosome structures. Although DNA in the open regions of CT is more susceptible to reaction with the BP derivative, there is appreciable reaction with the DNA associated with histones. (46 refs)

- 78-6785 Uptake, Binding to Cytoplasmic Protein, Translocation to the Nucleus and Morphological Transformation of Human Cells by Benzo(a)pyrene and 7,12-Dimethylbenzanthracene.** (Eng) Ekelman, K. B. (Ohio State Univ., Columbus, OH, 43210). *Diss Abstr Int B* 39(2): 499-500; 1978. (no refs)

- 78-6786 Pathologic Changes Induced in Respiratory Tract Mucosa by Polycyclic Hydrocarbons of Differing Carcinogenic Activity.** (Eng) Topping, D. C. (Natl. Inst. Environmental Health Sciences, Research Triangle Park, Durham, NC, 27709); Pal, B. C.; Martin, D. H.; Nelson, F. R.; Nettesheim, P. *Am J Pathol* 93(2): 311-324; 1978.

The toxic effects of benzo(e)pyrene (BeP), pyrene, anthracene, benz(a)anthracene (BA), dibenz(a,c)anthracene (DBA), benzo(a)pyrene (BP), and dimethylbenz(a)anthracene (DMBA) were investigated in the respiratory mucosa of male Fischer F-344 rats. These compounds were chosen because they comprise a spectrum of polycyclic hydrocarbons (PCH) ranging from noncarcinogens to initiators, weak carcinogens, and strong carcinogens. Heterotopic tracheal grafts were maintained under the dorsal skin of syngeneic recipients. The grafts were exposed to the test substance from an intraluminal beeswax pellet, which released the material slowly over 8 wk. The histopathologic effects were periodically assessed semiquantitatively. All PCH exhibited varying degrees of toxicity for the respiratory epithelium and submucosa. BeP showed the least toxicity, followed by pyrene and anthracene. BA and DBA caused marked epithelial and submucosal changes: epithelial hyperplasia, undifferentiated epithelium, and squamous metaplasia and marked mononuclear infiltration in the subepithelial connective tissue. With BP, the epithelial and submucosal changes were similar but stronger. DMBA was the most toxic substance, causing epithelial necrosis followed by generalized keratinizing squamous metaplasia; the subepithelial changes consisted of an early acellular exudate and, at 8 wk, marked condensation and hyalinization of the lamina propria. The toxic response pattern of the tracheal mucosa to carcinogens was characterized by the chronicity of epithelial and connective tissue damage, as opposed to the short-lived hyperplastic and inflamma-

tory response elicited by the noncarcinogens and initiators. (21 refs)

- 78-6787 Histochemical Patterns of a Few Key Carbohydrate Metabolic Enzymes During Hepatocarcinogenesis (Meeting Abstract).** (Ger) Hacker, H. J. (Hamburg, W. Germany); Bannasch, P. *Zentralbl Allg Pathol* 122(3): 289; 1978. (no refs)

- 78-6788 Inducibility of Aryl Hydrocarbon Hydroxylase in Cultured Human Lymphocytes.** (Eng) Fleischer, K. A. (Dept. Tropical Medicine, Univ. Liverpool, Box 147, Liverpool L69 3BX, England); Price Evans, D.; Canning, M. V. *J Med Genet* 15(3): 182-188; 1978.

The method for estimating aryl hydrocarbon hydroxylase (AHH) activity in cultured human lymphocytes was modified in an attempt to obtain reproducible values for AHH induction. However, none of the modifications produced the level of reproducibility required for quantitative genetic studies. (18 refs)

- 78-6789 Vaginal and Cervical Squamous Cell Dysplasia in Women Exposed to Diethylstilbestrol (DES) in Utero.** (Eng) Burke, L. (Beth Israel Hosp., 330 Brook Ave., Boston, MA, 02215); Antonioli, D.; Rosen, S. A. *Obstet Gynecol* 132(5): 537-544; 1978.

Pelvic, colposcopic, and histologic examinations were performed on 250 patients with a known or suspected history of in utero exposure to diethylstilbestrol (DES). (Exposure was confirmed in 205 cases.) Colposcopically abnormal areas were graded for severity according to predetermined criteria and biopsy evaluation was made for adenosis in vaginal squamous metaplasia, glandular or squamous metaplasia, nonglycogenated mucosa, and dysplasia in both vaginal and cervical specimens. No glandular or squamous carcinomas were noted, but a nonneoplastic abnormality was found in biopsy specimens of 82% of the women. Gross abnormalities included cervical columnar epithelium in 147 women and a cervicovaginal hood in 147. Adenosis was present in vaginal walls, fornices, and/or hood in 185 cases. Colposcopically abnormal areas were present in vaginas and hoods of 211 patients. Fifteen patients had squamous dysplasia, the degree of which was mild in 11 and moderate in 4 cases. The dysplasia involved the cervix in 14 patients and the vagina in 4. Patients with cervical mosaic and vaginal epithelium had dysplasia on biopsy much more frequently than patients with similar colposcopic appearances in the vagina. Thus, colposcopic data on cervical dysplasia can

applied to the evaluation of dysplasia in vaginal adenosis without modification. (16 refs)

78-6790 **Comparative Sensitivity of Animals of Various Species to the Carcinogenic Effect of Tryptophan Metabolites.** (Rus) Khrustalev, S. A. (Lab. Systemic Food Diseases, Cancer Res. Center, Moscow, USSR). *Patol Biol Eksp Ter* (4): 61-63; 1978.

The carcinogenic activity of tryptophan metabolites in guinea pigs, Syrian hamsters, and random-bred rats was compared. Animals of each species were divided into two groups: Group 1 was inoculated sc with 2-aminoacetophenone (2-AAP) and Group 2 was inoculated sc with 3-indolylacrylic acid (3-IAA) 2- to 3-day intervals. Both groups received inoculations for 5 mo; the total dose of each agent was 500-740 mg for guinea pigs, 200 mg for hamsters, and 200 mg for rats. The tumorigenic activity of 2-AAP and 3-IAA was most pronounced in the hamsters (there were 6 cases of lymphosarcoma, 2 cases of myeloid leukemia, 2 cases of lymphoid leukemia, and 1 case of plasmacytic leukemia of hemopoietic cells). Among the rats there were seven cases of lymphosarcoma, and among the guinea pigs there was one case of lymphoid leukemia. Both metabolites also induced various benign and malignant tumors in the hamsters (6 insulinomas of the pancreas, 1 adenoma of the pancreas, 2 liver cholangiomas, 1 papilloma of the vagina, 3 fibrosarcomas and fibromas, 1 melanoma, and 1 papillary carcinoma of the urinary bladder). Both metabolites also induced various tumors in rats (2 fibromas, 1 fibrosarcoma, 1 cystic epithelioma, 1 ovarian folliculoma, 1 uterine myxofibroma, 6 mammary and carcinomas). (7 refs)

78-6791 **Role of Hormones in Mammary Neoplasia.** (Eng) Nandi, S. (Dept. Zoology, Univ. California, Berkeley, CA, 94720). *Cancer Res* 38(11, part 2): 4046-4049; 1978.

A hypothesis for the role of hormones in mammary neoplasia as well as neoplasia in other endocrine organs and target tissues is presented. It is assumed that hormones are not carcinogenic by themselves but that they modify the host and/or target tissues during the events initiated by carcinogens such as viruses, chemicals, or radiation. The hypothesis is based on the observations that hormones are involved in regulating cell division in normal mammary cells, the emergence of overt mammary tumors requires hormonal stimulation of cells receiving a carcinogenic stimulus, normal cells have finite divisional life but neoplastic cells are immortal, and normal mammary cells might inhibit the growth of small numbers of preneoplastic and neoplastic cells. The hypothesis is that hormones have at least two functions in mammary carcinogenesis initiated by direct-acting carcinogens. First, hormones are necessary for DNA synthesis and mitosis of the initial transformed cells and, consequently, for their conver-

sion into fixed transformed cells with heritable characteristics. Second, hormones stimulate the rapid growth of normal cells with finite divisional life, leading to a reduction in their number in localized areas surrounding the transformed cells or in the tissue as a whole. This releases the transformed cells from the growth inhibitory influence of normal cells, allowing the emergence of overt tumors. (48 refs)

78-6792 **Two In Vivo Assays for Mutagenicity Compared: The Sperm Abnormality and Micronucleus Assays (Meeting Abstract).** (Eng) Bruce, W. R. (Dept. Medical Biophysics, Univ. Toronto, Toronto, Canada); Heddle, J. A. *Mutat Res* 53(2): 159-160; 1978. (no refs)

78-6793 **The Formation of a Large Molecular Weight Form of ACTH by Chemically-induced Rat and Mouse Lung Tumors (Meeting Abstract).** (Eng) Khan, M. N. (Molecular Pharmacology, Biochemistry Section, Lab. Pharmacology, NIEHS, Research Triangle Park, NC, 27709); Ghosh, A.; Linnoila, I.; Nettesheim, P.; DiAugustine, R. P. *Environ Health Perspect* 20: 232-233; 1977. (1 ref)

78-6794 **Dose-Effect Studies on Estrogen Induced Mammary Cancers in Mice.** (Eng) Rudali, G. (Equipe de Recherche 190 du C.N.R.S., Fondation Curie, 26, Rue d'Ulm, 75231 Paris, Cedex 05, France); Jullien, P.; Vives, C.; Apiou, F. *Biomedicine* 29(2): 45-46; 1978.

Castrated male (C3H x RIII)F1 mice were treated with graded doses of estradiol-17 β and estrone in a life-span experiment to determine to what extent the carcinogenic activity of estrogens in the mammary gland was dose-related. The mice were castrated at age 25 days; on the same day, paraffin pellets containing estradiol-17 β were implanted sc for continuous resorption. These pellets contained 1, 2.5, 5, 10, or 100 μ g of hormone. Other castrated mice received estrone po in the food at concentrations of 0.02, 0.2, or 2 ppm, corresponding to daily intakes of 0.06, 0.6 and 6 μ g, respectively. In addition, castrated females were ovariectomized and treated with the lower doses of the estrogens to see if they were sufficient to induce a vaginal estrus. Although the lower doses of both hormones induced vaginal estrus, they did not produce mammary tumors in the males. With the higher doses, however, the mammary tumor incidence was 95.8%-100%. These findings can be compared with clinical evidence in humans that continuous ingestion of low doses

of estrogens does not enhance breast cancer development. (44 refs)

78-6795 Liver Tumours Associated with Oral Contraceptive Agents (Meeting Abstract). (Eng) Britton, W. J. (A. W. Morrow Dept. Gastroenterology, Royal Prince Alfred Hosp., Sydney, Australia); Gallagher, N. D.; Little, J. M. *Aust NZ J Med* 8(5): 559; 1978. (no refs)

78-6796 Tumor Promoter Inhibits Production of Liver Retinol Binding Protein (Meeting Abstract). (Eng) Borek, C. (Coll. Physicians Surgeons, Columbia Univ., New York, NY); Smith, J. E. *J Cell Biol* 79(2, Part 2): 78a; 1978. (no refs)

78-6797 Factors Affecting the Induction of Chromosomal Aberrations by Cadmium in Chinese Hamster Cells (Meeting Abstract). (Eng) Deaven, L. L. (Cellular and Molecular Biology Group, Los Alamos Scientific Lab., Univ. California, Los Alamos, NM, 87545); Campbell, E. W. *J Cell Biol* 79(2, part 2): 131a; 1978. (no refs)

78-6798 Hormonal Control of Preneoplastic Epidermal Cells During Carcinogenesis in Mice (Meeting Abstract). (Eng) Lupulescu, A. (Dept. Dermatology, Wayne State Univ., Detroit, MI); Rogers, J.; Birmingham, D. *J Cell Biol* 79(2, part 2): 187a; 1978. (no refs)

78-6799 The Influence of Nutrition on the Growth of Experimental Kidney Tumors in Rats (Meeting Abstract). (Eng) Hartig, F. (Mannheim, W. Germany); Hebold, G. *Vet Pathol* 15(4): 574; 1978. (no refs)

78-6800 Mesotheliomas Induced by Sterigmatocystin in Wistar Rats. (Eng) Terao, K. (Res. Inst. Chemobiodynamics, Chiba Univ., Inohana 1-8-1, Chiba 280, Japan). *Gann* 69(2): 237-247; 1978.

The direct carcinogenic potency of sterigmatocystin (SMC)

to the peritoneum after repeated ip injections was investigated in male Wistar rats. Three groups of rats received ip injections of SMC, O-acetylsterigmatocystin (AcO-SMC), dihydrosterigmatocystin (dihydro-SMC) 1 x/wk for 23 weeks and they were observed for 80 wk. Only SMC produced mesothelioma in 20/40 rats given a total dose of 20-25 mg. No mesotheliomas were found in rats injected with AcO-SMC or dihydro-SMC. Both epithelioid and mesenchymal mesotheliomas were induced by SMC. Hepatocellular carcinomas were found in one rat treated with SMC and five given AcO-SMC. These results suggest that SMC is a potent carcinogen when administered ip. Although both dihydro-SMC and SMC formed fine, needlelike crystals when injected ip, dihydro-SMC did not produce any mesotheliomas. Dihydro-SMC has no double bond in the terminal furan ring; this double bond has been postulated as an important determinant of the acute toxicity and carcinogenicity of aflatoxin B₁ in rats. It is suggested, therefore, that the carcinogenicity of SMC to the peritoneum may be related more closely to the presence of this double bond than to its physical form in the peritoneal cavity. (28 refs)

78-6801 Induction by Degraded Carrageenan of Colorectal Tumors in Rats. (Eng) Wakabayashi, T. (Res. Inst., Nippon Chemiphar Co., Ltd., Misato-shi, Saitama, Japan); Inagaki, T.; Fujimoto, Y.; Fukuda, Y. *Cancer Lett* 4(3): 171-176; 1978.

The carcinogenicity of degraded carrageenan derived from the red seaweed *Eucheuma spinosum* was examined in Sprague-Dawley rats. In the first experiment, rats were divided into four groups of 30 males and 30 females each: Group 1, 2, and 3 received 10%, 5%, and 1% carrageenan in the diet, respectively, for up to 24 mo, and Group 4 served as a control. The rats were sacrificed at various intervals up to 24 mo after the start of the experiment. A total of 59/60 Group 1 and 12/60 Group 2 rats developed colorectal squamous metaplasia; 19/60 and 53/60 rats, respectively, developed colorectal tumors. The tumors were squamous cell carcinomas, adenocarcinomas and adenomas; two Group 1 rats and one Group 2 rat had metastases of squamous cell carcinomas to the retroperitoneal lymph nodes. In a second experiment, 20 animals of each sex were given a 5% aqueous solution of degraded carrageenan in the drinking water for 15 mo. Metaplasia was noted in all 40 rats, and 11 developed tumors. In addition to the tumor types found in the first experiment, one myosarcoma was also observed. In the third experiment, rats were divided into three groups of 15 males and 15 females each. Group 1 was given 5 g/kg and Group 2 1 g/kg of degraded carrageenan by stomach tube for 15 mo. Metaplasia was noted in all Group 1 rats and in 11/30 Group 2 rats. Adenocarcinomas and adenomas were noted in eight Group 1 animals. No colorectal metaplasia or tumors were noted in control rats in all three experiments. These findings indicate that degraded carrageenan is carcinogenic to the colorectal epithelium of the rat. (9 refs)

802 The Use of Short-Term Tests to Measure the Preventive Action of Reducing Agents on Formation and Activation of Carcinogenic Nitroso Compounds. (g) Lo, L. W. (Cancer Res. Centre, Univ. British Columbia, Vancouver V6T 1W5, Canada); Stich, H. F. *Mutat* 57(1): 57-67; 1978.

effect of reducing agents on the nitrosation of methylhydrazine (MG) and on the in vitro activation of dimethylnitrosamine (DMN) was examined in cultured human fibroblasts. The reducing agents were sodium ascorbate, cysteine, glutathione, and propyl gallate. The addition of sodium ascorbate to an NaNO_2 and MG mixture resulted in reduced DNA repair synthesis. The inhibitory action of sodium ascorbate on nitrosation of MG was also observed when the extent of chromosome aberrations was measured and when the colony-forming capacity of human fibroblasts was estimated. A repair synthesis of cultured human fibroblasts incubated with a rat liver microsomal activation system and DMN was greatly reduced in the presence of the reducing agents. The inhibitory effect could have been exerted during activation of DMN, or it could also be due to the trapping of free radicals formed during activation. The results do not distinguish between these two possibilities. Alkaline sucrose gradient analysis showed that sodium ascorbate inhibited the shift to smaller DNA fragments that was observed when cultured cells were exposed to liver microsome-activated DMN. These results indicate that DNA repair synthesis is a suitable short-term test for studying the numerous interactions between several carcinogenic agents and for estimating the capacity of inhibitory agents to affect formation and activation of chemical carcinogens. (45 refs)

See also:

- *(Rev.): 78-6601, 78-6603, 78-6604, 78-6605, 78-6606, 78-6607, 78-6608, 78-6609, 78-6611, 78-6612, 78-6613, 78-6614, 78-6615, 78-6616, 78-6617, 78-6618, 78-6619, 78-6621, 78-6622, 78-6623, 78-6624, 78-6625, 78-6626, 78-6627, 78-6628, 78-6629, 78-6630, 78-6631, 78-6634, 78-6635, 78-6636, 78-6637, 78-6638, 78-6639, 78-6640, 78-6641, 78-6642, 78-6643, 78-6644, 78-6645, 78-6646, 78-6647, 78-6648, 78-6649, 78-6650, 78-6651, 78-6652, 78-6653, 78-6654, 78-6655, 78-6656, 78-6657, 78-6658, 78-6659, 78-6660, 78-6662, 78-6663, 78-6664, 78-6697, 78-6701, 78-6702, 78-6703, 78-6706, 78-6707.
- *(Phys.): 78-6805, 78-6808, 78-6810, 78-6815, 78-6817, 78-6836.
- *(Viral): 78-6862, 78-6868, 78-6899, 78-6908, 78-6933, 78-6969, 78-6970.
- *(Immun.): 78-6984, 78-6994, 78-6998, 78-7011, 78-7033, 78-7041, 78-7042, 78-7043, 78-7045, 78-7046, 78-7048, 78-7049, 78-7051, 78-7053, 78-7059, 78-7061, 78-7063, 78-7064.
- *(Path.): 78-7075, 78-7076, 78-7088, 78-7095, 78-7107, 78-7110, 78-7113, 78-7114, 78-7122, 78-7125, 78-7126, 78-7133, 78-7141.
- *(Epid.-Biom.): 78-7149, 78-7150, 78-7151, 78-7156, 78-7162, 78-7163, 78-7172, 78-7173, 78-7174, 78-7176, 78-7181, 78-7183, 78-7184.

PHYSICAL CARCINOGENESIS

- 78-6803 **Malignant Melanoma of Skin and Sunspot Activity (Letter to Editor).** (Eng) Wigle, D. T. (Bureau Epidemiology, Lab. Centre Disease Control, Health and Welfare Canada, Ottawa, Canada K1A 0L2). *Lancet* 2(8079): 38; 1978.

Studies in the US and Canada showed that the incidence of malignant melanoma during 1935-75 varied cyclically, with the peaks correlating closely with periods of peak sunspot activity. The increased flux of cosmic rays during peak sunspot activity may destroy stratospheric ozone, so that more UV radiation reaches the earth's surface. (no refs)

- 78-6804 **Re: "Mathematical Models of Age and Ultraviolet Effects on the Incidence of Skin Cancer among Whites in the United States" (2 Letters to Editor).** (Eng) Lee, J. A. (Dept. Epidemiology, Sch. Public Health, Univ. Washington, Seattle, WA, 98195); Fears, T. R.; Scotto, J.; Schneiderman, M. A. *Am J Epidemiol* 107(3): 259-262; 1978.

Discrepancies between the regression coefficients for the incidence of nonmelanoma skin cancer and malignant melanoma with increasing age have implied that both these conditions are related to annual dose rather than cumulative lifetime UV exposure. It is suggested, however, that the similarity of the regression coefficient relating latitude and the coefficient relating age to the incidence of nonmelanoma skin cancer indicates that these tumors are produced by the cumulative effect of UV damage; this cumulative relationship may also hold for melanoma. In a rebuttal, it is held that since a flat age/nonmelanoma incidence is not found, neither melanoma nor nonmelanoma incidences are directly related to total UV exposure. It is suggested that nonmelanoma incidence is related to the cumulative UV dose and that melanoma incidence is related to the annual UV dose. (14 refs)

- 78-6805 **Interaction of Near-Ultraviolet Radiation and Irradiated L-Tryptophan in the Production of Mutations in *Escherichia coli*.** (Eng) Webb, R. B. (Div. Biological and Medical Res., Argonne Natl. Lab., Argonne, IL, 60439); Yoakum, G. H.; Perou, J.; Eisenstark, A. *Mutat Res* 51(3): 427-432; 1978.

The interaction between near-UV radiation and preirradiated tryptophan (TP) in *Escherichia coli* growing in chemostat cultures was studied. Near-UV radiation alone produced T5-resistant mutants in *E. coli* WP2 culture at a rate of 25.3×10^{-8} mutant/cell/day/watt m^{-2} . It produced 59.5×10^{-8} mu-

tant/cell/day/watt m^{-2} in the excision-deficient strain WP2. When TP was present in the culture during near-UV treatment, the rate of mutagenesis was increased two-fold for strain WP2 and seven-fold for strain WP2s. In the absence of near-UV radiation, TP increased the mutation rate by a factor of 1.3 in strain WP2 and by a factor of 1.7 in strain WP2s. The presence of unirradiated TP (5-10 mg/liter) reduced the rate of near-UV mutagenesis. The data suggest that the nonmutagenic repair of lesions produced by the interaction of components of TP and near-UV radiation requires the *uvrA* + endonuclease. (20 refs.)

- 78-6806 **Effects of Dietary Constituents on Ultraviolet Light-mediated Carcinogenesis.** (Eng) Black, J. S. (Photobiology Lab., Veterans Admin. Hosp., Houston, TX, 77030); Chan, J. T.; Brown, G. E. *Cancer Res* 38(13): 1384-1387; 1978.

The effects of dietary supplements of antioxidants and enzyme inducers on UV-mediated carcinogenesis in female hairless mice were investigated. The animals received either a (1) regular, balanced, laboratory meal, (2) regular meal supplemented with 0.05% (wt/wt) phenobarbital, (3) regular meal with disulfiram (2 mg/g meal), (4) regular meal with 0.5% (wt/wt) butylated hydroxytoluene (BHT), or (5) regular meal with 0.1% (wt/wt) reduced glutathione. At the end of 17 wk, the animals in Group 3 were transferred to a regular meal because of wt loss. After 10 days, the mice received UV radiation treatments 5 days/wk through week 17. The initial suberythemal dose was 0.84 joule/cm²/day; it was increased every 2 wk by 0.34 joule/cm² until a daily dose of 2.1 joules/cm² was given. An accumulative dose of 152 joules/cm² was administered. The reduced glutathione diet had no effect on tumor incidence, but the BHT, phenobarbital, and disulfiram diets all significantly suppressed the initiation and development of actinic lesions and tumors. Analysis of the percentage of animals bearing lesions and tumors, percentage of animals bearing tumors, and numbers of tumors/animal indicated that disulfiram provided the greatest protection. Tumor inhibition thus appears to be due to the induction of systemic physiological responses. (39 refs)

- 78-6807 **Inhibition of UV-induced DNA Repair at Different Steps by Quinacrine and Anthralin.** (Eng) Kulkarni, M. S. (Lab. Molecular Biology, Univ. Alabama at Birmingham, AL, 35294); Yelding, K. L. *Biochem Biophys Res Commun* 83(4): 1531-1537; 1978.

The effects of anthralin (AT) and quinacrine (QA) on DNA

er synthesis were evaluated in L1210 cells, a mouse lymphoma line. The cells were labeled with ^3H -thymidine, irradiated with UV light, and incubated for 15 or 45 min with or without QA or AT (both, 50 μM). The appearance and disappearance of single-strand breaks was followed by neutral sedimentation in neutral sucrose gradients. QA reduced the reduction in nucleoid sedimentation rate resulting from UV irradiation. After 15 min of post-UV incubation with QA, these nucleoids still sedimented like those from nonirradiated cells. With 45 min post-UV incubation, however, the sedimentation profile resembled that of irradiated cells. After 15 min of post-UV incubation without QA. These findings suggest that QA inhibits the UV endonuclease step in DNA repair. AT apparently increased the extent of single-strand breaks seen at 15 min post-UV incubation. There was no evidence for repair after 45 min post-UV incubation with AT, as indicated by an increased nucleoid sedimentation rate. However, AT enhanced the initial appearance of nicked DNA, followed by a return to normal, but QA delayed the nicking process. Although both agents inhibit repair synthesis, they produce different biological effects. (25 refs)

6808 Mutagenicity and Carcinogenicity of 8-MOP/UV A in Cell Cultures. (Eng) Burger, P. M. (Inst. Dermatology, Univ. Medical Centre, State Univ. Leiden, Leiden, Netherlands); Simons, J. W. *Bull Cancer (Paris)* 65(3): 281-282; 1978.

ants deficient for the enzyme hypoxanthine-guanine-phosphoribosyltransferase (HGPRT) were selected in V-79 Chinese hamster cells and in human skin fibroblasts (HSF) after application of the photochemotherapeutic (PUVA-therapy) regimen of 8-methoxypsoralen (8-MOP) (10 $\mu\text{g}/\text{ml}$) and UV irradiation (0-600 J/m ^{2}). In the V-79 cells, there was a linear relationship between UV dose and mutation frequency. However, with respect to varying doses of 8-MOP, a constant number of induced mutants (5.5×10^{-8} per μg 8-MOP/ml per J/m ^{2} per unit dose) was detected. HSF also yielded a constant number of induced mutants/unit dose of 8-MOP (3.3×10^{-8} per unit dose for dividing cells and 0.6×10^{-8} /unit dose for nondividing cells). If these results are extrapolated to human skin cells, it is predicted that 7.4×10^{-5} mutants/unit dose for dividing cells and 1.4×10^{-5} mutants/unit dose for nondividing, (or a total of 1.7×10^{-5} mutants induced per unit dose of PUVA therapy) may be expected to arise. A carcinogenic effect (the induction of cell transformation in C3H/3T3 mouse cells) was also observed. (12 refs)

6809 8-Methoxypsoralen as a Frameshift Mutagen in Bacteria and Chinese Hamster Cells in the Skin—Implications for Genetic Risk in Man (Meeting Abstract). (Eng) Bridges, B. A. (MRC Cell Mutation Unit, Univ. Sussex, Falmer, Brighton, England); Mottershead, R. A.; Arlett, C. F. *Mutat Res* 53(2): 156; 1978. (no refs)

78-6810 Characterization of Sister Chromatid Exchange Induction by 8-Methoxypsoralen plus Near UV Light. (Eng) Latt, S. A. (Genetics Div., Childrens Hosp. Medical Center, 300 Longwood Ave., Boston, MA, 02115); Loveday, K. S. *Cytogenet Cell Genet* 21(4): 184-200; 1978.

Factors influencing the induction of sister chromatid exchanges (SCE's) in Chinese hamster ovary (CHO) cells by 8-methoxypsoralen (8-MOP) + near-UV light were investigated. Exposure of CHO monolayers to approx 6×10^{-6} M 8-MOP or UV light (up to 2×10^4 ergs/mm ^{2} , primarily at 365 nanometers) prior to bromodeoxyuridine (BUDR) treatment had little effect on SCE frequency. However, irradiation of cells shortly after 8-MOP addition resulted in a marked increase in SCE's that was dependent on 8-MOP dose and UV dose, but not on the interval between the two. SCE induction by 8-MOP + UV light was compatible with cell survival over multiple cycles. Study of cells that had undergone three cycles of BUDR incorporation indicated that 8-MOP + UV light increased SCE formation during the third cycle. Thus, cells surviving treatment included those damaged and responding with an increase in SCE's. Both the frequency and location of SCE's induced at different times within the DNA synthesis period varied in a manner that indicated that exchange induction was restricted to regions that replicated during or after DNA damage. (56 refs)

78-6811 Inflammatory Reactions Induced by Ultraviolet Irradiation. (Eng) Greaves, M. W. (Inst. Dermatology and St. John's Hosp. Diseases of the Skin, Homerton Grove, London E9 6BX, England); Hensby, C. N.; Black, A. K.; Plummer, N. A.; Finchman, N.; Warin, A. P.; Camp, R. *Bull Cancer (Paris)* 65(3): 299-303; 1978.

Human skin irradiated with different wavelengths of the UV spectrum in the presence and absence of anti-inflammatory drugs was examined pharmacologically. In all experiments, human abdominal skin was irradiated; it was examined visually for erythema, and exudate from irradiated cells was collected at different times after irradiation and analyzed by bioassay using the cascade superfusion technique, by radioimmunoassay, and by gas-liquid chromatography-mass spectrometry. Unirradiated skin from the same subject was used as the control. Blood flow (BF) changes were also measured by photoelectric plethysmography. After irradiation with six times the minimal erythema dose (MED) of UVC [< 290 nanometers (nm) in wavelength], the levels of arachidonic acid (ARA) and prostaglandins (PG)E $_2$ and F $_2$ were all increased (1.5-3.0 times control), reaching a max at 18-24 hr, but returned to control levels by 48 hr, by which time the erythema had also subsided. When the PG inhibitor indomethacin (IM) was administered in divided doses beginning 48 hr before irradiation, the levels of PGE $_2$ and PGF $_2$ were both held to the control level, but ARA levels were enhanced in both irradiated and nonirradiated skin. IM caused a reduction in BF that correlated with a reduction in visual erythema. Irradiation with 3 MED of UVB (290-320 nm) also

produced peaks in levels of ARA, PGE₂, PGF₂ and other ARA metabolites at 24 hr (5 to 20 times control), which returned to control values by 48 hr. Erythema did not decrease at 48 hr, however, and IM treatment produced only a small change in BF in irradiated skin while apparently completely suppressing PG activity. UVA (320-400 nm) treatment of patients after administration of 8-methoxypsoralen produced no changes in PGE₂ or PGF₂ levels in irradiated areas, but ARA levels were again increased at 24 hr. It is suggested that increased levels of PGE₂ protect epidermal cells from mutagenic action of UVB by increasing cellular levels of cyclic AMP and thus reducing the vulnerable pool of actively dividing cells. (16 refs)

78-6812 Detection of DNA Repair in Nuclei Isolated from Human Cultured Cells (Meeting Abstract). (Eng) Robison, S. H. (Dept. Pathology, Thomas Jefferson Univ., Philadelphia, PA); Fansler, B. S. *J Cell Biol* 79(2, part 2): 137a; 1978. (no refs)

78-6813 Quantitative Changes in Respiratory Enzyme Activity in Premalignant Lesions and Experimentally Irradiated Skin. (Eng) Pearse, A. D. (Dept. Medicine, Welsh Natl. Sch. Medicine, Heath Park, Cardiff, Wales); Marks, R. *Bull Cancer (Paris)* 65(3): 351-356; 1978.

The morphological and cell kinetic findings were compared with metabolic changes within the epidermis of patients exposed to solar radiation to investigate the earliest changes in UV-induced neoplasia. Quantitative enzyme histochemical techniques were employed to examine lesional and paralesional skin from four patients with solar keratoses and one with Bowen's disease. Buttock skin was also sampled from these patients to compare the lesional sun-exposed skin with non-sun-exposed skin. Succinic dehydrogenase (SDH) activity was evenly distributed in buttock skin, and there was very little change in total sun-exposed epidermal SDH activity, but it was significantly reduced in the granular cell layer in lesional and paralesional skin of all patients with solar keratoses. Glucose-6-phosphate dehydrogenase (G6PDH) epidermal total activities were increased in both lesional and paralesional skin compared with buttock epidermis in all patients. In all skin samples G6PDH activity was preferentially increased in the granular cell layer. Lactate dehydrogenase (LDH) activity did not show a difference in localization in any tissues studied. When buttock skin of five healthy subjects was irradiated 10x during a 14-day period with 1-2 minimum erythematous doses of 290-400 nanometer, UV irradiation, SDH activity dropped approx 50% in the granular skin layer, but was unchanged in basal layer and total activities. Additionally, G6PDH activity increased significantly in the total epidermis and basal cell layer and somewhat in the granular cell layer after irradiation. It is suggested that these findings may serve as a model for persistent UV light exposure leading to epidermal damage in man. (2 refs)

78-6814 Experimentally Induced Intestinal Metaplasia in Wistar Rats by X-Ray Irradiation. (E) Watanabe, H. (Dept. Cancer Res., Res. Inst. Nuclear Medicine and Biology, Hiroshima Univ., Kasumi 1-2-3, Hiroshima, 734, Japan). *Gastroenterology* 75(5): 796-799; 1978.

The induction of intestinal metaplasia with Paneth cells and intestinal marker enzymes in Wistar rats by local x-irradiation is described. The gastric region of 5-wk-old female Wistar rats was irradiated daily with 500 rads of x-rays up to a total of six times. Goblet cells and marker enzymes of small intestine, such as lactase, trehalase, and maltase, appeared in the pyloric region of the glandular stomach of rats from the first week after final irradiation. Intestinal-type crypts without Paneth cells were observed at 8 wk, sucrose activity at 26 wk, and intestinal metaplasia with Paneth cells at 71 wk. The number of goblet cells, intestinal type crypts and Paneth cells increased with age. Gastric adenocarcinoma did not develop after irradiation. These studies suggest that a small dose of carcinogen might cause only intestinal metaplasia and that increasing the dose level might result in carcinoma associated with metaplasia or, in the case of higher doses, carcinoma alone. (26 refs)

78-6815 Acute Myelomonocytic Leukemia after Irradiation and Chemotherapy for Ewing's Sarcoma. (Eng) Smithson, W. A. (Dept. Pediatrics, Mayo Clinic, Rochester, MN); Burgert, E. O.; Childs, D. S.; Hoagland, H. *Mayo Clin Proc* 53(11): 757-759; 1978.

A 16-yr-old boy was diagnosed with acute myelomonocytic leukemia 16 mo after the start of irradiation and chemotherapy for Ewing's sarcoma of the upper right leg. The primary lesion was treated with a daily midplane dose of 200 rads x-rays for 25 sessions over 29 days (total, 5,000 rads). Beginning at the same time, vincristine (VCR: 2 mg/m², max dose 2.0 mg) and cyclophosphamide (CP: 500 mg/m²) were given weekly for 10 wk, followed by a 4-wk rest period. Then doxorubicin (0.015 mg/kg/day) was given for 5 days, followed by resumption of the weekly VCR and CP therapy for 5 wk. Four cycles of triple-agent chemotherapy were administered. Fourteen months after the initial presentation, a portion of the right lung involved by metastatic disease was resected and the lungs were treated with a total of 1,800 rads of x-rays in 12 fractions over 16 days. Simultaneously, doxorubicin (DR: 30 mg/m²/day) was given for 3 days and repeated once in 3 wk. Two months after the lung resection, the Hb level decreased and undifferentiated blasts appeared in the peripheral blood; the total WBC count was 2,500/mm³ with 1% blasts. DR therapy was continued for 3 mo, at which time the percentage of blasts increased to 38% of a total WBC count of 40,000/mm³. At death, there was no clinical evidence for active Ewing's sarcoma. Although nothing in this case report suggests the need to modify the current approach to Ewing's sarcoma, it does signal the need for surveillance for second neoplasms of all types. (21 refs)

78-6816 **Otolaryngologic Sequelae of Radiation Therapy.** (Eng) Kupperman, G. L. (Univ. Massachusetts, Amherst, MA); Kleiner, H.; Stageman, D. L.; Martinelli, D. *Am J Otolaryngol* 57(11): 510-515; 1978.

The ears, nose, throat, and hearing of 100 persons who had been exposed to head, neck, or chest irradiation at least 10 years previously were examined in an attempt to document the long-term effects of radiation treatment. There was no evidence of clinical carcinoma in the ear, nose, or throat structures of any of the patients. Temporomandibular-joint diseases were found in three patients without the common historical physical findings usually found. One patient had bilateral enlargement of the submaxillary glands and two had unilateral enlargement. Few cases of conductive hearing loss or middle-ear disease were observed. Nine of 27 patients with sensorineural hearing loss gave no history of excessive noise exposure or ototoxic drug therapy. The uncommonly high number of patients with this type of hearing loss indicates there may be some relationship to previous irradiation. (16 refs)

78-6817 **Effect of Some Nonradioactive and Radioactive Chemicals on Spleen Structure.** (Rus) Anomareva, T. V. (Res. Inst. Radiation Hygiene, Leninabad, USSR); Merkushev, G. N. *Arkh Anat Gistol Embriol* (4): 47-52; 1978.

To evaluate the effect of common environmental pollutants on spleen structure, mice and rats received a series of radioactive (^{137}Cs , ^{226}Ra , ^{89}Sr , ^{65}Zn) and nonradioactive [$\text{Pb}(\text{NO}_3)_2$, BeCl_2] nuclides, hexamethylenediamine (HMDA), and chloromethylmercury (CMM) in their drinking water for 1-2 yr. The compounds were given in concentrations (slightly greater than av and max permissible doses). Chronic exposure to ^{226}Ra and ^{225}Ra (total absorbed dose 0.3 and 5.4 rads, respectively), HMDA, and CMM enhanced cell proliferation in the red pulp. Structural rearrangement of the spleen tissue was accompanied by hyperplasia of the lymphoid tissue. Hyperplasia of the hematopoietic stem cells and enhanced plasma cell proliferation were detected after exposure of the rodents to ^{226}Ra , ^{89}Sr , ^{65}Zn , and BeCl_2 . At the end of experiment, the animals had a marked decrease in immune reactivity that coincided with aplasia of the lymphatic component of the spleen. (17 refs)

78-6818 **On the Mechanism of "Dissolution" in Liquids of PuO_2 by Alpha Decay.** (Eng) Fleischer, R. (General Electric Res. and Development Center, Schenectady, NY, 12301); Raabe, O. G. *Health Phys* 35(4):545-548; 1978.

Subparticle formation by alpha decay of plutonium-239 in plutonium oxide particles was studied by the nuclear track technique to locate the sites of deposition of fissionable atoms

or groups of atoms. Fission was induced on a high-purity quartz detector by thermal neutrons, and the individual tracks were revealed by chemical etching. Analysis of the results indicated that in a vacuum, at the most, a single ^{239}Pu atom was ejected with the uranium-235 recoil nucleus. Therefore, direct spraying-out of multiatom fragments does not occur. The most plausible model of the previously observed subparticle formation in water is that a damaged region is produced in the original particle and that the damage is subsequently released by exposure to water. (10 refs)

78-6819 **A Morphometric Study of Inhaled $^{239}\text{PuO}_2$ in Rodents (Meeting Abstract).** (Eng) Rhoads, K. (Battelle, Pacific Northwest Lab., Richland, WA, 99352); Sanders, C. L. *Pharmacologist* 20(3): 276; 1978. (no refs)

78-6820 **Toxicology of Inhaled Transuranic Oxides in Rodents (Meeting Abstract).** (Eng) Sanders, C. L. (Battelle, Pacific Northwest Lab., Richland, WA, 99352); Mahaffey, J. A. *Pharmacologist* 20(3): 175; 1978. (no refs)

78-6821 **Estimation of Risk for Lung Cancer Induced by Inhaled Alpha Emitting Radionuclides (Meeting Abstract).** (Eng) Mahaffey, J. A. (Battelle, Pacific Northwest Lab., Richland, WA, 99352); Sanders, C. L. *Pharmacologist* 20(3): 175; 1978. (no refs)

78-6822 **Factors Affecting the Mobility of Plutonium-238 Dioxide in the Rat.** (Eng) Stradling, G. N. (Natl. Radiological Protection Board, Harwell, Didcot, Oxon, England); Ham, G. J.; Smith, H.; Cooper, J.; Breadmore, S. E. *Int J Radiat Biol* 34(1): 37-47; 1978.

A major factor influencing the movement of plutonium-238 from lungs to blood after the administration of oxide suspensions to female inbred rats via intubation was the presence of 0.001- μm -diameter particles. In a polydisperse suspension of particles, this fraction increased with time, probably because of fragmentation of larger particles induced by alpha decay. The rate of this process could account for the greater transportability in vivo of plutonium-238 relative to plutonium-239 when the oxides were inhaled. In blood, 0.001- μm -diameter plutonium-238 oxide particles underwent a rapid reaction to form a low-mol-wt species before plutonium was complexed with transferrin and citrate ions. The filtration of this species through the kidneys may explain the observed enhanced urinary excretion of plutonium relative to administered plutonium citrate. The mechanism of urinary excretion and the relationship between cumulative urinary excretion and body content for plutonium-238 were similar to those

previously observed for plutonium-239, even though different methods of preparation of the oxides were used (calcination of the oxalate, $^{238}\text{PuO}_2$; condensation of the metal vapor, $^{239}\text{PuO}_2$). (25 refs.)

78-6823 Studies on the Uptake of Plutonium from Major Indian Soils and Its Distribution in Plants.

(Eng) Vyas, B. N. (Biology and Agricultural Div., Bhabha Atomic Res. Centre, Trombay, Bombay 400 085, India); Mistry, K. B. *J Environ Qual* 7(4): 533-536; 1978.

Studies of the plant uptake of plutonium from three major Indian soil types showed that the highest amount of Pu was absorbed from acidic laterite soil, followed by alluvial soil and medium-black alkaline soil. Pu concentration ratios in beans and maize ranged from 1.5 to 9.6×10^{-4} for aerial tissues and from 13.9 to 15.3×10^{-4} for roots. In mature bean plants, the Pu distribution showed a marked acropetal gradient. (19 refs)

78-6824 Size Fractionation Methods: Measuring Plutonium in Respirable Dust. (Eng) Hayden,

J. A. (Rockwell International, Rocky Flats Plant, P.O. Box 464, Golden, CO, 80401). *Science* 202(4369): 753-754; 1978.

Four soil fractionation methods were used to separate three size fractions of two soil samples in order to evaluate the inhalation hazard of soil contaminated with plutonium. Increasing amounts of soil mass and Pu activity (0.5%-38% and 7%-84%, respectively) remained in the soil fraction $<10 \mu\text{m}$. The two samples showed similar trends in the changing Pu content of this fraction, the change being highly dependent on the fractionation method used. (5 refs)

78-6825 Influence of Injected Mass of Plutonium on Its Biological Distribution. (Eng) Guilmette, R. A.

(Inhalation Toxicology Res. Inst., P.O. Box 5890, Albuquerque, NM, 87115); Lindenbaum, A.; Friedman, A.; Sullivan, J.; Moretti, E.; Russell, J. J. *Health Phys* 35(4): 529-536; 1978.

The distribution and retention in mouse tissues of an injected mass of plutonium were investigated using four α -emitting Pu isotopes of differing specific activities: ^{236}Pu ($532 \mu\text{Ci}/\mu\text{g}$ Pu), ^{239}Pu ($0.0613 \mu\text{Ci}/\mu\text{g}$), ^{242}Pu ($0.0039 \mu\text{Ci}/\mu\text{g}$), and ^{244}Pu ($1.93 \times 10^{-5} \mu\text{Ci}/\mu\text{g}$). Female B6CF₁/Anl mice 88 days old, were injected iv with $1 \mu\text{Ci}/\text{kg}$ of ^{236}Pu , ^{239}Pu , ^{242}Pu , or $0.3 \mu\text{Ci}/\text{kg}$ of ^{244}Pu . Groups of five animals receiving each isotope were sacrificed 4-96 days after injection, and the following tissues were analyzed radiochemically by α -liquid scintillation counting for Pu content: liver, spleen, kidneys, lungs, lymph nodes, thigh muscle, ovaries, femurs, and tibias (marrow samples from both tibias were analyzed). In all the tissues analyzed, the early distribution and retention of ^{236}Pu ,

which was injected at a mass level (2 nanograms/kg) equivalent to 20% of the max permissible body burden (MPBB) of ^{239}Pu , did not differ from those of ^{239}Pu , even though injected masses differed by $>8,000$. Retention of the isotope of lower specific activity, ^{242}Pu and ^{244}Pu , which were injected at mass equivalents of 28,000 and 99,000 times the MPBB of ^{239}Pu , respectively, was generally three to six times higher than retention of ^{236}Pu and ^{239}Pu in the soft tissues. These results suggest the occurrence of an aggregation of ^{242}Pu and ^{244}Pu in vivo. They also suggest that the use of metabolic data obtained in animal studies with high levels of injected ^{239}Pu allows valid extrapolation downward to at least the level of an MPBB exposure. (13 refs)

78-6826 Radiation Leukemia in C57BL/6 Mice. Correlation of Altered Expression of Terminal

Deoxynucleotidyl Transferase to Induction of Leukemia

(Eng) Pazmino, N. H. (Cancer Biology Program, National Cancer Institute, Frederick Cancer Res. Center, Frederick, MD, 21501); Mullan, R.; Ihle, J. N. *J Exp Med* 148(5): 1338-1350; 1978.

In a study of the effects of irradiation on early thymic differentiation, terminal deoxynucleotidyl transferase (TdT) activity was measured in C57BL/c mice irradiated with total body x-rays 1, 3, or 4 x/wk, beginning at age 1 mo. Early radiation schedule resulted in a 60%-80% loss of thymocytes at 1 wk post-treatment, but the subsequent repopulation of the thymus was dose dependent. Peak TdT I and II activities (expressed as percentage of nonirradiated controls) in thymus and bone marrow decreased dramatically after a single 175-rad irradiation, but they returned to normal values by 4 wk. Recovery of thymus TdT activity after two or three 175-rad doses took longer, and in the bone marrow there was only 50% recovery. Four doses reduced activity to undetectable levels, and no TdT was detected in bone marrow for at least 18 wk. Radiation-induced thymic lymphomas in C57BL/c mice were TdT-positive; high levels of thymus TdT activity were detectable only in overtly leukemic mice, regardless of the time after irradiation. Bone marrow-reconstitution studies immediately after the last radiation abrogated the leukemic effect, and TdT levels were normal 6 wk after reconstitution. Studies of the effects of 4 x 175 rads on mice of various ages showed that susceptibility to leukemia induction and ability to recover normal TdT activity were age-dependent. TdT activity was induced in bone marrow B cells from thymectomized mice given a nonleukemic radiation dose (1×350 rads) but not in B cells from mice given 4 x 175 rads. This indicates that a leukemogenic radiation dose eliminates or alters bone marrow stem cell population inducible for TdT. Since the loss of the TdT stem cell was thymus-independent, the primary effect of irradiation occurs in the bone marrow. (10 refs)

78-6827 Chronic Myelogenous Leukemia Development After Thymic Irradiation. (Eng) Shimaoka,

Medicine B, Roswell Park Memorial Inst., Buffalo, Sokal, J. E. In: *Third International Symposium on the Prevention and Detection of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): (part 1): 1193 pp; 819-830; 1977.

patients with Philadelphia (Ph') chromosome-positive chronic myelogenous leukemia (CML) and a history of thymic irradiation are presented. The first patient underwent thoracic irradiation for breathing difficulties and a widened mediastinum on chest x-ray at 3 mo of age. Three of 100 rads in air were given every other day, for a total of 100 rads. He developed leukemia 18 yr later; chromosome analysis revealed 46XY with a single Ph' chromosome in all metaphases prior to the blastic stage, after which 47XY with two Ph' chromosomes gradually became the sole karyotype. The second patient was given three doses/wk of 132 rads in air to the thorax and an enlarged thymus at the age of 2 mo, for a total dose of 100 rads. Leukemia developed at the age of 22 yr; blast karyotyping revealed a mode of 45 chromosomes (a deletion of one of the Group C chromosomes) and a single Ph' chromosome in all metaphases. It is concluded that both patients had radiation-induced CML. (39 refs)

28 **Behaviour of Murine Lymphosarcoma Cells During Prolonged Cultivation In Vivo.** (Eng) Kovacs, V. (Inst. Biophysics, Czechoslovak Acad. Sci., Brno 12, Czechoslovakia); Drasil, V. *Folia Biol (Praha)* 24(4): 253-263; 1978.

Transplantable lymphosarcoma growing as a free suspension in the peritoneal cavity of isologous hosts was isolated after fractionated irradiation of C57BL mice (4 x 160 R at 2-wk intervals). During a 14-yr cultivation period, the modal number of tumor chromosomes declined from 41-42 to 39, the mean survival time of tumor hosts decreased, and the resistance of the tumor cells increased slightly. Strain specificity remained unchanged. (25 refs)

29 **Estimation of Life Shortening Resulting from Radiogenic Cancer per rem of Absorbed Dose.** Gotchy, R. L. (Radiological Assessment Branch, U.S. Environmental Protection Agency, Washington, DC, 20555). *Health Phys* 35(4): 563-565; 1978.

The amount of life shortening (LS) from the cancer that may be expected from a radiation dose equivalent to 1 rem (roentgen-equivalent-man) was estimated. Equations are given for estimating the LS for a given age cohort. The av LS for each age cohort was estimated by summing all the years of life lost for each age cohort over the entire population. The total LS associated with all cancer types for total body irradiation was derived from the sums of av LS by cancer

type. The total LS per million man-rem is 3,200 yr for the general public. The av loss of life expectancy per individual is 1.2 days/rem. An upper-bound estimate of the av LS of radiation workers per 5-rem dose equivalent was calculated using the 20- to 70-yr population. The av loss of life expectancy for this group for all types of cancer was calculated as 0.63 day/man-rem. It is believed, however, that the actual age distribution of nuclear workers may be skewed toward the younger years. Therefore, the LS of occupationally exposed radiation workers in the nuclear industry is expected to lie between 0.63 and 1.2 days/rem. (4 refs)

78-6830 **Late Effects after Incorporation of the Short-lived α -Emitters ^{224}Ra and ^{227}Th in Mice.** (Eng) Muller, W. A. (Institut für Biologie Gesellschaft für Strahlen- und Umweltforschung mbH, D-8042 Neuherberg, W. Germany); Gossner, W.; Hug, O.; Luz, A. *Health Phys* 35(1): 33-35; 1978.

Dose dependence and temporal distribution of osteosarcomas (OS's) and other malignancies were investigated following single and repeated ip injections of ^{224}Ra of varying activity and single injections of ^{227}Th in NMRI mice. The highest concentration of ^{224}Ra occurred in the long bones, followed by the vertebral column, especially the lumbar spine. After a single injection of ^{224}Ra (1.0-50 $\mu\text{Ci/kg}$, 30-1,500 rads), the OS incidence ranged from 7% to 22% (at 360 rads) in females vs 1.3% in controls. Males had a max incidence of 8.5% at 750 rads (dose range, 5-50 $\mu\text{Ci/kg}$, 150-1,500 rads) vs 0% in controls. A single injection of ^{227}Th in females produced a max incidence of 60% at 1,000 rads. Fractionated application of ^{224}Ra resulted in a higher overall incidence of OS than that from a single injection of the same mean skeletal dose. The highest tumor incidence (> 90%) was observed with the lowest activity per injection, 0.5 $\mu\text{Ci/kg}$ 2 x/wk for 36 wk (total, 1,080 rads). Tumor incidence was also increased when the same mean skeletal dose was distributed over a longer treatment period. When 0.5 $\mu\text{Ci/kg}$ was given 2 x/wk for 36 wk, a 92% OS incidence resulted at 17 mo, but 0.5 $\mu\text{Ci/kg}$ 2 x/wk for 12 wk resulted in only a 46% incidence at 23 mo. OS's occurring with single-dose treatments at the higher doses were located mainly in the head bones; these treatments also increased the incidence of ossified fibromas. Fractionated doses and lower single doses generally produced osteosarcomas in sites other than the head. An earlier occurrence of leukemia was seen in some ^{227}Th -treated groups and in one group given repeated ^{224}Ra injections. (34 refs)

78-6831 **Toxic Autonomous Thyroid Nodule Following Neck Irradiation.** (Eng) Scott, R. S. (Medical Unit, Princess Margaret Hosp., Christchurch, New Zealand); Brownlie, B. E.; Bremner, J. M. *Aust NZ J Med* 8(5): 521-522; 1978.

A 37-yr-old man with history of neck irradiation in childhood

presented with thyrotoxicosis due to an autonomous thyroid nodule. Adjacent to the nodule was a radiation scar from radiotherapy administered 35 yr previously. The long latent period is not unusual, in that thyroid abnormalities have been observed > 30 yr after radiation. Hemithyroidectomy was performed, and histological examination excluded malignant change. (6 refs)

- 78-6832 Radiation Induced Cancer in the Head and Neck Region.** (Eng) Matsumura, Y. (Div. Head and Neck Surgery, Natl. Kyushu Cancer Center, Fukuoka, Japan); Nomura, Y.; Eguchi, S.; Horie, A. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 807-818; 1977.

Twelve patients with radiation-induced cancer, 8 initially treated for benign conditions (Group A) and 4 initially treated for malignant tumors (Group B), were analyzed. The primary tumors in the latter group were reticulum cell sarcoma of the maxilla, squamous cell carcinoma of the larynx (2), and nasopharyngeal carcinoma. At the time of therapy, the average of the Group A patients was 19 yr, that of the Group B patients 44 yr. However, the mean latent interval for Group A was 33 yr, vs only 13 yr for Group B; no cancer developed < 7 yr after the initiation of radiation therapy. Although the radiation dose was not known for some patients, it was estimated that those in Group A received 2,000-5,000 rads, those in Group B 6,000-8,000 rads. Damage to the head and neck skin in Group A patients included fibrosis, stigma, irregular pigmentation, and leukoderma, but Group B patients had only a slight dermatosis. The difference in damage is suggested to be due to the radiation dose, source, and field. Histologically, the radiation-induced cancers were generally anaplastic. They spread diffusely and were of uniform thickness. The radiation-induced cancers are proposed to arise by the two-stage mechanism. The possibility of cancer developing in irradiated patients after a certain latent interval is distinct; these patients should be monitored carefully. (23 refs)

- 78-6833 Contamination of the Home Environment by Patients Treated with Iodine-131: Initial Results.** (Eng) Jacobson, A. P. (Dept. Environmental and Industrial Health, Univ. Michigan Sch. Public Health, Ann Arbor, MI, 48109); Plato, P. A.; Toerock, D. *Am J Public Health* 68(3): 225-230; 1978.

Twin-sodium iodide-radiation detectors were used to analyze iodine-131 transfer from thyroid patients to their families. The study included seven families with 17 persons, 11 of whom were under age 16. Direct radiation exposure of family members due to the proximity of the radioactive patients

ranged from 0.17 to 126 mR/day (natural background radiation is 0.35 mR/day). The range of max activity of iodine-131 in the thyroids was 92-110,000 picocuries, and it resulted in thyroid dose equivalents of 4-1,330 milliroentgen-equivalent-man. Based on recent estimates of thyroid cancer, the latter dose equivalent could double the risk of thyroid malignancy in children over what is expected normally. Such a risk implies the addition of 10 induced cases to the 10 natural occurring cases per million people per year. (20 refs)

- 78-6834 Fate of Inhaled Transuranic Oxides (Meeting Abstract).** (Eng) Adee, R. R. (Battelle, Pacific Northwest Lab., Richland, WA, 99352); Sanders, C. L.; Halsey, J. G. *Pharmacologist* 20(3): 276; 1978. (no refs)

- 78-6835 Development of Osteosarcoma in Rats After Irradiation.** (Eng) Solheim, O. P. (Norsk Hydro Inst. Cancer Res., Oslo, Norway). *Acta Radiol [Theor (Stockh)]* 16(5): 433-446; 1977.

Tissue abnormalities preceding the development of osteogenic sarcoma were determined in inbred male rats exposed to ^{90}Sr (in equilibrium with its daughter isotope ^{90}Y) radiation with activities of 0, 0.05, 0.2, 0.67, or 1.32 $\mu\text{Ci/cm}$. The animals were exposed to the radiation for various times ranging from 4 to 360 days and then sacrificed at times ranging from 4 to 540 days. In the medullary cavity, the main effect of the irradiation was a slow and dose-dependent replacement of the hematopoietic elements by fat. Neither extensive fibrosis nor irregular nontumorous bone was found in the marrow cavity of any animal exposed to continuous irradiation. After termination of irradiation, the hematopoietic tissue was reformed. In the cortical bone, the number of osteocytes decreased at a dose-dependent rate; this decrease continued after the radiation was discontinued. Osteoclasts were found on the periosteal surface of all bones exposed for 14-45 days but they were rarely found on the endosteal surface. The endosteal surfaces were always lined by resting osteoblasts after exposure to at least 0.42 Gray unit/day (all groups above 0.05 $\mu\text{Ci/cm}$). A total of 22 tumors were noted: 4 in the 0.2-, 12 in the 0.67-, and 12 in the 1.32- $\mu\text{Ci/cm}$ groups. Nineteen of these tumors were sarcomas and 10 were osteogenic sarcomas. Malignant proliferation began after approx 180 days of irradiation. Osteogenic sarcomas had their centers at the endosteal surface. The adjacent bone marrow was nearly aplastic, but the endosteum was thickened by fibroblasts and osteoblasts. The number of osteocytes in the adjacent bone was greatly reduced. It is suggested that radiation injury accumulated in osteogenic cells until some mechanism involved in the control of proliferation and differentiation failed. (7 refs)

- 78-6836 Preliminary Studies of the In Vitro Cellular Effects of Asbestos and Fine Glass Dusts.** (Eng) Sincok, A. M. (Dept. Human Genetics and Biometry, Univ.

London, London NW1 2HE, England). In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Ten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4 (Book B), pp. 941-954; 1977.

These hamster cells that had been cloned from a single line cell and checked for a total modal variation of were incubated with various samples of asbestos (0.1 ml) for 48 hr and then subjected to genetic analysis. With of the asbestos samples (S.F.A. chrysotile and U.I.C.C. dolomite), there was a high level of aberration, with > 50% of cells analyzed being karyotypically abnormal. The other asbestos samples also showed significant levels of genetic abnormality. No significant genetic damage was observed in cells treated with coarse fiberglass. A fine glass fiber sample (code 100), which contained fibers within the diameter length ranges of asbestos, also produced a significant level of aberration in treated cells. Both chemical leaching and mechanical milling of asbestos fibers significantly reduced the ability of asbestos to induce in vitro genetic damage. These results indicate the importance of physical morphology in the induction of genetic damage by asbestos. They indicate that this system could be used to grade fine dusts on the basis of induced genetic aberration. (10 refs)

837 Malignant Mesothelioma Arising after Direct Application of Asbestos and Fiber Glass to the Pericardium. (Eng) Churg, A. (Dept. Pathology, Sch. Medicine, Univ. California, San Francisco, CA, 94143); Warnock, J.; Bensch, K. G. *Am Rev Respir Dis* 118(2): 419-424;

A 57-year-old man developed a malignant mesothelioma 15 yr after he had been treated for angina pectoris by dusting of the pericardial cavity with a mixture of fibrous dusts. Electron diffraction and microprobe analysis indicated that approximately 67% of the fibers were tremolite and anthophyllite asbestos and the remainder were fiber glass. This is believed to be the first human malignant mesothelioma associated with direct mesothelial contact with fibrous dusts. (14 refs)

838 Mortality after Long Exposure to Cumingtonite-Grunerite. (Eng) McDonald, J. C. (TUC Health Research Inst. Occupational Health, London Sch. Hygiene

and Tropical Medicine, Keppel St., Gower St., London WC1, England); Gibbs, G. W.; Liddell, F. D.; McDonald, A. D. *Am Rev Respir Dis* 118(2): 271-277; 1978.

The causes of death of 657/660 former cumingtonite-grunerite workers were investigated to determine if exposure to these minerals carried an increased risk of respiratory tract cancer. Excess mortality with a clear dust-exposure relationship was found for pneumoconiosis and respiratory tuberculosis, but there was no convincing evidence of an increase in respiratory cancer. (9 refs)

78-6839 Inhalation Studies with Polyurethane Foam Dust in Relation to Respiratory Tract Carcinogenesis. (Eng) Thyssen, J. (Institut für Toxikologie der BAYER AG, 5600 Wuppertal 1, W. Germany); Kimmerle, G.; Dickhaus, S.; Emminger, E.; Mohr, U. *J Environ Pathol Toxicol* 1(4): 501-508; 1978.

Male and female Sprague-Dawley rats inhaled freshly generated polyurethane foam dusts (8.65 mg/m³ air) for 6 hr/day, 5 days/wk, over a 12-wk period. The exposure had no apparent effect on appearance, behavior, body wt, or life-span, and there were no indications of a carcinogenic effect of the dusts on the respiratory tract. (11 refs)

See also:

*(Rev.): 78-6609, 78-6612, 78-6614, 78-6620, 78-6624, 78-6633, 78-6634, 78-6665, 78-6666, 78-6667, 78-6668, 78-6697, 78-6704.

*(Chem.): 78-6753, 78-6759, 78-6765, 78-6777, 78-6792.

*(Viral): 78-6941, 78-6947.

*(Immun.): 78-6979, 78-7061.

*(Path.): 78-7072, 78-7090, 78-7091, 78-7092, 78-7102, 78-7121.

*(Epid.-Biom.): 78-7155, 78-7170, 78-7171, 78-7179, 78-7182.

VIRAL CARCINOGENESIS

- 78-6840 Cell-free Translation of Avian Sarcoma Virus RNA: Suppression of the *gag* Termination Codon Does Not Augment Synthesis of the Joint *gag/pol* Product.** (Eng) Weiss, S. R. (Dept. Microbiology, Univ. California, San Francisco, CA, 94143); Hackett, P. B.; Oppermann, H.; Ullrich, A.; Levintow, L.; Bishop, J. M. *Cell* 15(2): 607-614; 1978.

The effect of yeast suppressor transfer RNA's (tRNA's) on the cell-free translation of two viral gene products (Pr76*gag* and P180*gag/pol*) from the 38S messenger RNA (mRNA) of avian sarcoma virus (ASV) was studied. Although the synthesis of P180*gag/pol* was not enhanced by the suppressor tRNA's, and the use of amber suppressor tRNA reduced the synthesis of Pr76*gag* and resulted in the synthesis of a new 80,000-dalton polypeptide (P80) that contained the antigenic determinants of *gag*. Studies using chemical inhibitors of the initiation of translation demonstrated that P80 was generated by the addition of amino acids to the carboxy terminus of Pr76. The data indicate that the ASV *gag* gene is terminated by an amber codon, UAG, and that suppression of this termination signal does not lead to uninterrupted translation from *pol*. A second termination signal about 120 nucleotides from the UAG termination appears to define the length of P80. Thus, at least two suppression events would be necessary to synthesize P180. This argues against the hypothesis that suppression of nonsense codon(s) at the end of the *gag* gene is responsible for the synthesis of P180 in the ASV-infected cell. (28 refs)

- 78-6841 Characterization of Transformation-Sensitive Membrane-Proteins in Chick Embryo Fibroblasts Transformed with Avian Sarcoma Virus.** (Eng) Isaka, T. (Dept. Viral Oncology, Cancer Inst., Kami-Ikebukuro, Toshima-ku, Tokyo 170, Japan); Ikawa, Y.; Yoshida, M. *Virology* 90(2): 330-343; 1978.

A membrane polypeptide of 90,000 daltons (P90) found in increased amounts of chick embryo fibroblasts transformed by avian sarcoma virus was characterized, and its accumulation in transformed cells was studied. Compared with normal cells, there was a marked increase in the amount of P90, some increase in P79, and a decrease in P200 in transformed cells. Autoradiography showed that P90 had exactly the same isoelectric point as a polypeptide of 91,000 daltons (P91). Comparison of the peptide maps of P90 and P91 showed that the two polypeptides had very similar primary structures. P91 was shown not to be a glycoprotein, and it appeared to be the glycosylated form of P90. However, there was no conversion of P91 to P90 by deglycosylation. It is concluded that P91 is formed by glycosylation of the growing polypeptide

P90 on polysomes, that P90 is formed in the absence of glycosylation, and that the completed polypeptides are not interconvertible. (26 refs)

- 78-6842 A Simple Method for Rapid Purification of Avian Sarcoma Virus Supercoiled DNA by Selective Precipitation of Infected Cell Chromatin.** (Eng) Goto, R. V. (Dept Microbiology, Coll. Physicians and Surgeons, Columbia Univ., New York, NY, 10032). *Anal Biochem* 90(1): 256-261; 1978.

A simple and effective method of partially purifying avian sarcoma virus (ASV) supercoiled (form I) DNA from infected quail tumor cells (QT6) consists of lysing infected nuclei in water and precipitating the resulting chromatin with sodium chloride. This method offers some advantages over the Hirt procedure in that low-mol-wt viral DNA can be extracted rapidly and only 1%-2% cellular DNA is found in the viral fraction, even when the total number of cells to be processed is as high as 3.4×10^9 . The total amount of ASV-specific DNA recovered was 70%-100% of that recovered by the Hirt method. Similar results were obtained with simian virus 40 (SV40)-infected cells were processed by two methods, with the chromatin method extracting about 85% of the SV40 DNA extracted by the Hirt procedure. The efficiency of the two methods was compared by monitoring the recovery of ASV DNA after final purification. Approximately 3×10^9 QT6 cells were labeled with ^3H -thymidine immediately after ASV infection. The nuclei were prepared and divided into two parts, and each half was subjected to one of the purification procedures. Complete purification of ASV form I DNA, as monitored by alkali sucrose gradient analysis, required only two steps by the chromatin method, whereas at least three or four steps, including two CsCl-propidium iodide equilibrium gradient centrifugation steps were required to eliminate contaminating cellular DNA in the Hirt supernatant. Because of the extremely low levels of host cell DNA in the postchromatin fraction, the method is particularly suitable for purifying low-mol-wt viral DNA's, such as Rous tumor virus and SV40 DNA's. (6 refs)

- 78-6843 *env* Gene of Chicken RNA Tumor Viruses: Content of Conservation in Cellular and Viral Genomes.** (Eng) Fujita, D. J. (Cancer Res. Lab., Univ. Western Ontario, London, N6A 5B7 Ontario, Canada); Tavaré, Varmus, H. E.; Bishop, J. M. *J Virol* 27(3): 465-474; 1978.

Molecular hybridization studies with DNA complementary to *env* gene sequences of chicken leukosis-sarcoma v

s (cDNA) were performed with a variety of avian s to examine the natural distribution and possible ori- of viral *env* genes. The results indicate that nucleotide nces closely related to cDNA are present in chicken but not in other avian DNA's tested or in calf thymus salmon sperm DNA's. These results are in marked con- to experience with the transforming gene (*src*) of avian ma viruses, for which homologs have been demonstrat- the DNA's of all avian species tested and also in the 's of other vertebrates. The *env* gene sequences were to be present in the DNAs of both chick helper fac- sitive and chick helper factor-negative (chf-) chicken at a level of one copy per haploid genome. The presence v gene sequences in chf- DNA suggests that the chf- type does not result from a major deletion of *env* gene nces, and it is consistent with the existence of mech- is for the regulation of *env* gene expression. Thermal uration studies were used to further examine the ology between the *env* genes of representatives of viral roups A to E. The results indicate that the hybrids ed between cDNA and RNA's of subgroup A to E es are closely matched. This suggests that a substantial on of the *env* genes of viruses of subgroups A to E con- of shared, closely related sequences and that these genes bly have a common evolutionary origin. (36 refs.)

844 Isolation and Characterization of the Envelope Glycoprotein of an Avian Osteopetrosis Virus: t of Host Cell on Antigenic Reactivity. (Eng) Van El- L. J. (Rockefeller Univ., 1230 York Ave., New York, 10021); Smith, R. E. *Virology* 90(1): 80-89; 1978.

physical and antigenic properties of the surface glyco- in (gp85) of avian myeloblastosis-associated virus [V-2(O)] were compared with those of avian myeloblasto- rus (AMV) gp85. Sodium dodecyl sulfate-polyacryla- gel electrophoresis and guanidine hydrochloride col- chromatography indicated that the gp85's of the two es were qualitatively similar, but that MAV-2(O) con- d approx 3.46 times more gp85 than AMV. When the of MAV-2(O) grown in chick embryo fibroblasts (CEF) examined by competition radioimmunoassay (RIA) us- rabbit anti-AMV gp85 serum, it exhibited 100- to -fold less reactivity than the AMV glycoprotein. In con- MAV-2(O) isolated directly from the serum of osteope- chickens competed efficiently. If MAV-2(O)-infected n of osteopetrotic chickens was replicated in CEF for 1 t subsequently lost its ability to compete in RIA with V. AMV itself did not compete effectively when grown EF, but it did so when grown in myeloblasts. MAV-2(O) ed from serum or CEF competed equally in RIA's using erum against B77 virus gp85, and glycoproteins from -grown MAV-2(O) and serum AMV showed equal ivity to the B77 antiserum. These results suggest MAV-2(O) and AMV gp85 show similar host- ndent glycosylation patterns in vivo and different erns when the viruses are grown in cultured CEF. (refs)

78-6845 Fractionation of Proteins from Rous Sarcoma Virus and Avian Myeloblastosis Virus by Isoelectric Focusing. (Eng) Mach, O. (Inst Molecular Genetics, Czechoslovak Acad of Sciences, Prague 6, Czechoslovakia); Sovova, V.; Cerna, H. *Folia Biol (Praha)* 24(4): 233-241; 1978.

The technique of isoelectric focusing in pH gradients was used to isolate proteins from Rous sarcoma virus and avian myeloblastosis virus. The purified major group-specific (gs) protein p27 (pI = 9.1) and gp86 (pI = 5.3) were obtained after disruption of virus with 1% nonionogenic detergent in the presence of 6M urea. Proteins p10, p15, and p19 were present in the same pH range (pI = 6.8). A strongly basic, immunologically active protein, presumably p12, was found in the alkaline region at pH 10.8. These proteins fully retained their immunological activity. On the other hand, in the acidic region between pH 4 and 5, strong precipitates were regularly found. These precipitates were complexes that were formed by interaction of the acid components of ampholines with the viral proteins during isoelectric focusing. Almost all viral proteins were present, differing only in quantity. The complexes were stable in 1% nonionogenic detergent with 6M urea. They were dissociated with 1% sodium dodecyl sulfate and 5M urea, and they had no immunological activity. The methods of virus disruption and the possibility of formation of the ampholine-viral protein complexes are discussed. (23 refs)

78-6846 Assay for Early Cytoplasmic Effects of the *src* Gene Product of Rous Sarcoma Virus. (Eng) McClain, D. A. (Rockefeller Univ., New York, NY, 10021); Maness, P. F.; Edelman, G. M. *Proc Natl Acad Sci USA* 75(6): 2750-2754; 1978.

The *src* gene product of Rous sarcoma virus (RSV) was as- sayed by microinjection of cytoplasmic extracts of chick em- bryo fibroblasts transformed by the Prague strain of RSV (subgroup A) into normal mouse 3T3 fibroblasts and then examining the actin-containing structures in these cells by immunofluorescence microscopy. Microinjection of these cytoplasmic extracts, but not extracts of normal cells, caused a dissolution of the 3T3 cell microfilament bundles that was max within 30 min. The activity could still be measured after a tenfold dilution of the cytoplasmic extracts (14 mg/ml original protein concentration). The activity was trypsin-sen- sitive and it was destroyed by boiling, but it was not RNase-sensitive. Protein synthesis was not required for the dissolution. Microinjected cytoplasts prepared from normal 3T3 cells also dissolved the microfilament bundles, indicating that the cell nucleus was not required for expression of activi- ty. Extracts made from fibroblasts transformed by RSV hav- ing a temperature-sensitive mutation in the *src* gene were also temperature-sensitive in this assay. Thus, the activity of extracts from cells infected with *src* mutant virus, but not from cells infected with wild- type virus, was destroyed either by in vitro incubation of the extract at the nonpermissive temperature before

injection or by incubation of recipient cells at the nonpermissive temperature after injection. It is concluded that the microinjection assay can detect a cytoplasmic activity coded for by the *src* gene RSV and that an early direct or indirect target of the *src* gene product is the cytoskeleton and cell motility system. This result is discussed in relation to the hypothesis that submembranous arrays of microfilaments, microtubules, and their associated proteins interact with cell-surface receptors to form a surface-modulating assembly that functions as a key regulator of cell growth. (13 refs)

- 78-6847 Linkage of the Endogenous Avian Leukosis Virus Genome of Virus-producing Chicken Cells to Inhibitory Cellular DNA Sequences.** (Eng) Cooper, G. M. (Sidney Farber Cancer Inst., Dept. Pathology, Harvard Medical Sch., Boston, MA, 02115); Silverman, L. *Cell* 15(2): 573-577; 1978.

To account for the inefficient transcription and lack of infectivity of endogenous Rous-associated virus-O (RAV-O) DNA of uninfected virus-producing (V+) cells, it was proposed that the endogenous RAV-O genome was linked to a cis-acting control element of cellular DNA that inhibited its transcription. The prediction of this model that shearing the DNA of uninfected V+ cells might convert the endogenous RAV-O genome to an infectious form by separating it from the control element was tested. The DNA of uninfected V+ chicken cells was infectious at a low level. In addition, the infectivity of RAV-O DNA of uninfected V+ cells was increased approx sixfold by shearing the DNA to the approx size of the viral genome. In contrast, shearing the DNA of RAV-O-infected cells reduced its infectivity 5- to 20-fold. Thus, the infectivity of endogenous RAV-O DNA relative to the infectivity of DNA of RAV-O infected cells was increased approx 60-fold by shearing. These results suggest that the endogenous RAV-O genome of V+ chicken cells is linked to cellular DNA sequences that inhibit its infectivity in transfection assays, as predicted by the cis-acting control element model. Alternative interpretations, however, such as the possibility that shearing leads indirectly to internal rearrangements in the viral genome, cannot be excluded. Since the infectivity of sheared DNA of uninfected cells was approx 0.3% of the infectivity of sheared DNA of RAV-O-infected cells, the size of the control element was roughly estimated at 0.1%-1% in the size of the RAV-O genome, or 10-100 nucleotide base pairs. These findings suggest that the endogenous viral genome of uninfected V+ chicken cells is linked to specific inhibitory cellular DNA sequences, and they provide further evidence consistent with the model of endogenous viral gene regulation. (17 refs)

- 78-6848 Evidence That the Transforming Gene of Avian Sarcoma Virus Encodes a Protein Kinase Associated with a Phosphoprotein.** (Eng) Levinson, A. D. (Dept. Microbiology, Univ. California, San Francisco, CA,

94143); Oppermann, H.; Levintow, L.; Varmus, H. E.; Bishop, J. M. *Cell* 15(2): 561-572; 1978.

Immunoprecipitation with rabbit tumor antisera to avian sarcoma virus (ASV) was used to isolate a phosphorylated protein of mol wt 60,000 daltons (pp60). This protein appeared to be encoded by the *src* gene responsible for the induction and maintenance of neoplastic transformation. Phosphorylation of pp60 could be accomplished in vitro with extracts of ASV-infected cells. With a temperature-sensitive conditionally lethal mutation in *src*, phosphorylation of the pp60 protein was temperature-sensitive. Since the mutant *src* was not expressed at the restrictive temperature, phosphorylation of pp60 must be required for its function as the putative effector of *src*. Immunoprecipitates prepared with extracts of ASV-infected cells and the rabbit tumor antisera contained a protein kinase activity that catalyzed phosphorylation of the heavy chain of immunoglobulin molecules, using either ATP or guanosine triphosphate as phosphate donor. It appears that the protein kinase may also be encoded in *src* and that the enzymatic activity may be an intrinsic property of pp60. Phosphorylation of pp60 in cellular extracts was inhibited by calcium ionophores, whereas the immunoprecipitable kinase activity was not, suggesting that the kinase responsible for pp60 phosphorylation may be distinct from that encoded in *src*. The data suggest that phosphorylation of specific cellular targets may account for transformation of the host cell by *src*. (36 refs)

- 78-6849 Ascorbic Acid Retards RNA-Tumor Virus Infection of Avian Cells in Culture (Meeting Abstract).** (Eng) Bissell, M. J. (Lab. Chemical Biodynamics, LBL, Univ. California, Berkeley, CA, 94720); Hatie, C.; Farnsworth, D.; Schwarz, R.; Soo, W. J. *J Cell Biol* 79(2, part 1): 401a; 1978. (no refs)

- 78-6850 Limited Host Range of Avian Tumor Viruses in Pigeon Cells.** (Eng) Meyers, P. (Dept. Microbiology, Mayo Clinic and Mayo Medical Sch., Rochester, MN 55901); Ritts, G. D.; Heise, J. M.; Qualtiere, L. F. *Virology* 90(1): 162-165; 1978.

Show Racer (SHR) and White Corneau (WC) pigeon embryo fibroblasts (PEF) were found to be virtually resistant to infection with Rous sarcoma virus (RSV) of subgroups A, B, C, D, and E. Focus assays were performed on five SHR PEF and six WC PEF cultures, using nine RSV strains, representing all five subgroups. PEF were absolutely resistant to subgroups A and E and 2/3 subgroup B viruses. Schmidt-Rupprecht (SR)-RSV, subgroup B, plated to some degree on many PEF, but its max plating efficiency (EOP) was 0.0002 relative to standard chicken embryo fibroblasts (CEF). Subgroup C viruses plated poorly, but SR-RSV, subgroup D, plated relatively high titers in several PEF, the max EOP being 0.001 relative to CEF. Although SHR is an outbred and WC is an inbred strain, there was little difference in PEF resistance to RSV. (16 refs)

1 Binding of Tryptophanyl-tRNA to the Reverse Transcriptase of Replication-defective Avian Viruses. (Eng) Panet, A. (Dept. Virology, Hebrew Hadassah Medical Sch., Jerusalem, Israel); Weil, G.; R. *J Virol* 28(2): 434-443; 1978.

NA polymerase activity and the binding to tryptophan transfer RNA of the reverse transcriptase (TR) from five avian sarcoma virus mutants, two (LA 335 and LA 336) carrying an early replication lesion and three (LA 343, and LA 672) defective in a late replication lesion, were examined. Compared with the activities of the wild-type parents, both RT activities were found to be reduced in LA 335 and LA 336. When the other mutant viruses were grown at the permissive temperature, the RT activities were nearly as heat-stable as those of the wild-type parents. At the nonpermissive temperature, however, both RT activities were reduced in LA 338, LA 343, and LA 672. The relative amounts of RT antigen in these virions was determined by a competition immunoassay, it was found that the mutants contained reduced amounts of RT protein per total protein when harvested at the nonpermissive temperature. Even the infectious virus mutants produced at the permissive temperature always contained less enzyme protein than the wild-type parents. (27 refs)

52 Characterization of Rous Sarcoma Virus *src* Gene Products Synthesized In Vitro. (Eng) Beecher, K. (Tumor Virology Lab., Salk Inst., San Diego, CA, CA); Hunter, T. *J Virol* 28(2): 551-566; 1978.

Cell-free translation products of Rous sarcoma virus (RSV) virion RNA were further characterized by parallel denaturation of the heat-denatured virion RNA's of Prague group B (PR-B), Schmidt-Ruppin subgroup D (SR-D), transformation-defective (td) SR-D RSV in the messenger RNA-dependent reticulocyte lysate system. Two-dimensional analysis of the tryptic peptides of the 17,000 (17K), 25K, and 60K proteins synthesized by PR-B and SR-D RSV, but not td SR-D RNA, revealed that they have overlapping amino acid sequences. However, these products showed strain-specific differences, manifested by their electrophoretic mobility in acrylamide gels and by their tryptic peptides. Virion RNA's coding for the 17K, 25K, and 60K proteins were polyadenylated and they sedimented with maximum S values at about 23S, 19S-20S, and 18S, respectively. Transformation-specific proteins with mol wts of 39K and 33K were also synthesized. These proteins were related to the three proteins, and were synthesized from polyadenylated RSV RNA of approx 21S-22S. RNase T₁-resistant oligonucleotides were analyzed in parallel, and the *src*-specific oligonucleotides were found to be first present in equimolar amounts in gradient fractions sedimenting at 21S-22S. The results suggest that synthesis of the 60K protein is initiated near the 3' end of the *src* gene and that the 39K, 33K, and 17K proteins are initiated internally in the

src gene. All these proteins appear to be initiated independently, but they may have a common termination site. (30 refs)

78-6853 Study of Group-specific Proteins in Mammalian Cells Transformed by Rous Sarcoma Virus.

(Eng) Sovova, V. (Inst. Molecular Genetics, Czechoslovak Acad. Sciences, Praha 6, Czechoslovakia); Bosch, V.; Cerna, H. *Folia Biol (Praha)* 24(3): 199-205; 1978.

Lysates of rat, mouse, and hamster cells transformed by Rous sarcoma virus (RSV) and labeled with ³⁵S-methionine were examined for the presence of group-specific (gs) proteins by means of antisera, and the size of these molecules was determined by gel electrophoresis. Gs proteins were not detected in any of the immunoprecipitates analyzed. Only in RSH-mix cells (Syrian hamster embryo fibroblasts transformed in vitro by the Schmidt-Ruppin strain of RSV) was there a quantitative difference in the amount of a 70,000-dalton protein. Autoradiograms of simultaneously tested immunoprecipitates of lysates of RSV-transformed chicken cells clearly showed the presence of p27 and, after a short-term labeling period, the gs precursor molecule pr76. In previous experiments using complement fixation, gs proteins were detected in two of the RSV-transformed mammalian cell lines. The most probable explanation for this discrepancy is that the precursor molecule is attached to subcellular structures in mammalian tumor cells. (21 refs.)

78-6854 Analysis of Rous Sarcoma Virus (RSV) RNA Structure by Means of Specific Nucleases.

(Eng) Darlix, J. L. (Departement de Biologie Molculaire, Universite de Geneve, CH-1211-Geneve 4, Switzerland); Spahr, P. F.; Bromley, P. A. *Virology* 90(2): 317-329; 1978.

The effects of *Escherichia coli* RNases III and IV and *Aspergillus oryzae* T₁ ribonuclease and S1 nuclease on Rous sarcoma virus (RSV) 70S and 35S RNA were analyzed by the fingerprinting technique. The viral RNA was very susceptible to both RNases III and IV, with five regions being highly susceptible. Of these five regions, two, represented by T₁ oligonucleotides 13 and 2, are directly related to the initiation sites for reverse transcriptase. A large number of RSV RNA regions excluded from Sephadex G-75 were resistant to degradation by T₁ RNase, with 35S RNA being much less resistant than 70S RNA. Regions of 70S RNA with oligonucleotides 8, 15, and 9 were very resistant to S₁ nuclease; similar results were found with 35S RNA. Large T₁ oligonucleotides were not very susceptible to RNases III and IV, with the exception of oligonucleotides 2 and 18. The results indicate that RSV RNA is a highly structured molecule in which the two 35S subunits are intimately associated via a large number of specific interactions to form the 70S complex. (37 refs)

- 78-6855 Fluorescent Probe Studies of Normal, Persistently Infected, Rous Sarcoma Virus-transformed, and Trypsinized Rat Cells.** (Eng) Burleson, G. R. (Lobound Lab., Dept. Microbiology, Univ. Notre Dame, Notre Dame, IN, 76556); Kulpa, C. F.; Edwards, H. E.; Thomas, J. K. *Exp Cell Res* 116(2): 291-300; 1978.

The fluorescent probes pyrene, pyrene butyric acid, and N-phenyl-1-naphthylamine were used to study membranes of normal rat cells, Rous sarcoma virus (RSV)-transformed cells, cells treated with a proteolytic enzyme, and cells persistently infected with lymphocytic choriomeningitis virus. The lifetimes of excited pyrene and pyrene butyric acid showed only minor changes among normal, transformed, trypsinized or persistently infected cells. However, pyrene, but not pyrene butyric acid, lifetimes were shorter in cell membranes than in homogeneous solvents. The quenching of excited pyrene in cells by quencher molecules was slower than corresponding reactions in homogeneous solutions, indicating that the probe was screened from the quencher by the membrane. Quenching reactions with the pyrene butyric acid probe were similar in cells and homogeneous solvents, which indicated that pyrene and pyrene butyric acid reside in different lipid regions of the membrane. Transformed and trypsinized cells showed increased membrane fluidity, compared with normal and persistently infected cells. Membrane fluidity was determined from the excimer/monomer fluorescence ratios of pyrene and by the polarization of N-phenyl-1-naphthylamine fluorescence. Several techniques distinguished between normal and transformed or trypsinized cells; however, the only parameter unique to RSV transformation was a blue shift of the fluorescence maxima of N-phenyl-1-naphthylamine. This shift, which reflected a less polar environment of N-phenyl-1-naphthylamine in virus-transformed cells, was reversed by trypsinization. Therefore, the shift was apparently due to the presence of a trypsin-sensitive protein unique to the transformed cells. (31 refs)

- 78-6856 Characterization of the Avian Sarcoma Virus Protein p60src.** (Eng) Brugge, J. S. (Dept. Pathology, Univ. Colorado Medical Center, Denver, CO, 80262); Steinbaugh, P. J.; Erikson, R. L. *Virology* 91(1): 130-140; 1978.

The intracellular distribution of the avian sarcoma virus (ASV) protein p60src was investigated in transformed chicken and hamster cells by radioimmunoprecipitation of fractionated cells and by immunofluorescent staining of fixed cells with monospecific antiserum to p60src. The src protein was found exclusively in the cytoplasmic fraction in both a soluble and an insoluble form. Furthermore, the antigenicity of p60src was extremely heat-labile in vitro relative to Pr76, the precursor to the internal structural proteins of ASV. Antibody against p60src in serum from rabbits bearing tumors induced by the Schmidt-Ruppin (subgroup D) strain of ASV did not recognize p60src from cells transformed by other ASV strains, including Prague C, B77, and Bryan (RAV-50).

Reevaluation of the precipitation of p60src from NY68-transformed cells led to the conclusion that there is no reduction in the synthesis or antigenicity of p60src when cells are cultured under conditions nonpermissive for transformation by the mutant virus. (15 refs)

- 78-6857 Large-Scale Production, Concentration and Purification of Rous Sarcoma Virus in Tissue Culture.** (Eng) Dostalova, V. (Inst. Molecular Genetics, Czechoslovak Acad. Sciences, Prague 6, Czechoslovakia); Hložek, I.; Mach, O.; Lacinova, J. *Folia Biol (Praha)* 24(4): 225-232; 1978.

A method is described for the production of large amounts of purified virus from tissue cultures of chicken cells transformed by the Prague strain of Rous sarcoma virus, subgroup C. The method is based on the use of roller culture bottle precipitation with polyethylene glycol 6000, and purification by zonal centrifugation. Mean yields in cells cultured for 2 wk were 4.9 mg virus/liter of culture medium. (21 refs)

- 78-6858 Long-term Karyological Study of RVP, Tumour Cells Grown In Vitro.** (Eng) Sainarova, I. (Inst. Molecular Genetics, Czechoslovak Acad. Sciences, Prague 6, Czechoslovakia); Svoboda, J. *Folia Biol (Praha)* 24(4): 242-252; 1978.

The RVP₃ cell line established from an in vivo passaged RVP tumor, which originated in a C57BL/6 mouse infected with the Prague strain of Rous sarcoma virus (RSV), was analyzed at different passage levels and characterized virologically and cytogenetically. The RVP₃ cell line has 2-5 medium-sized large metacentric chromosomes; the majority of the mitoses contain about 1 subtelocentric chromosome; and in addition every mitosis contains chromosomal structures corresponding to the group (termed m-ms) that comprises both minute chromatin bodies and microchromosomes. In three RVP₃ monocellular clones with a similar karyotype, the number of m-ms increased proportionally to the total number of chromosomes in individual metaphases. It was established that the RVP₃ cell population does not contain any transmissible agent producing chromosomal changes in mouse cells and that RVP₃ cells are highly malignant. No RSV could be rescued from the clone tested. The possibility that m-ms's are generated by chromosome excision followed by replication and reintegration is discussed. The accompanying recombination events could lead to deletion or loss of proviral sequences adjacent to the region of the chromosome breaks. (11 refs)

- 78-6859 Differentiated Expression of Virus Genome in Mammalian Cells Transformed by Rous Sarcoma Virus.** (Rus) Tatosyan, A. G. (Lab. Leukemia Virology

Res. Center, Moscow, USSR); Kiselev, F. L. *Vopr* 24(10): 9-13; 1978.

ly was designed to attempt to verify the hypothesis that specific RNA in rat XC and TWERC cells murine cells, and hamster XPO cells transformed by Rous virus (RSV), undergoes transcription from different parts of integrated viral genome. Incubation of the RNA product of RSV with RNA isolated from transfected cells revealed a high level of molecular hybridization (80%) with RNA from cells producing gS-antigenic XC cells and helper-dependent TWERC cells) hybridized with a low level of hybridization (10%-13%) with RSV-3 cells. (14 refs)

60 **Demonstration of Marek's Disease Tumor-associated Surface Antigen in Chickens Infected with Noncancerous Marek's Disease Virus and Herpesvirus** (Eng) Schat, K. A. (Dept. Avian and Aquatic Medicine, New York State Coll. Veterinary Medicine, Ithaca, NY, 14853); Calnek, B. W. *J Natl Cancer Inst* 55:855-857; 1978.

presence of Marek's disease tumor-associated surface antigen (MATSA) was demonstrated on spleen cells from P-line chickens inoculated 5-6 days earlier with herpesvirus of turkeys (HVT) and SB-1 (a noncancerous Marek's disease virus). Fluorescence, which was restricted to the cell membrane, was more intense on cells from SB-1-infected chickens than those from HVT-infected chickens. Antisera against MATSA expressed on five Marek's disease lymphoblastoid cell lines were able to recognize the MATSA present on infected spleen cells. No viral membrane antigens and a low incidence of viral internal antigens could be demonstrated. (16 refs.)

61 **The Herpesvirus of Marek's Disease and Human Cancer.** (Eng) Makari, J. G. (Makari Res. Center, Englewood, NJ, 07631). In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer in New York, NY: Vol. 1(part 1): 1193 pp.; 263-278; 1977.

Previously found common antigenicity between glycoprotein (gp) antigens from human cancers and those from spleen cells of chickens with Marek's disease was explored further by a highly sensitive, modified Schutz-Dale technique. A search was made for serum antibodies (ab) to Marek's disease herpesvirus (MDHV) in 141 subjects with and without cancer and for MDHV antigens in the deoxyribose-purified (DPF) of normal and cancerous tissues. Serum antibodies prepared from MDHV were found in 63% of

65 subjects with nonmalignant diseases, 74% of 69 subjects with carcinoma (ca), and 4/7 subjects with melanoma and sarcoma. Serum antibodies to gp antigens prepared from herpes simplex virus types 1 and 2 and Epstein-Barr virus were found in only 4% of 52 subjects with nonmalignant diseases, 9% of 54 subjects with ca, and none of the 5 subjects with melanoma and sarcoma. All 8 subjects with early colon and breast cancer had serum antibodies against MDHV, vs 56% of 18 subjects with late disease. MDHV antigens were found in the DPF of all nine cancer types tested, but not in normal lung DPF or in DPF from fibroid uterus. These antigens were not detected with comparable gp antigens extracted from mouse hepatitis virus and influenza virus or with six control tissue antigens prepared from human and chicken tissues. Certain types of cancer were associated with antigens prepared from various strains of MDHV: ie, colon cancer was associated with the GA strain, breast, ovary, and lung cancer were associated with the CR-64 strain, and liver cancer and lymphoma were associated with the GA-2 strain. The associations were found in both the serum antibody and DPF studies. This was not the case with the gp antigens from the other herpesviruses investigated. (27 refs)

78-6862 **Murine Mammary Tumor Virus Expression During Mammary Tumorigenesis in BALB/c Mice (Meeting Abstract).** (Eng) Pauley, R. J. (Dept. Cell Biology, Baylor Coll. Medicine, Houston, TX, 77030); Socher, S. H. *J Cell Biol* 79(2, part 2): 358a; 1978. (no refs)

78-6863 **Murine Mammary Tumor Virus (MuMTV) Infection of an Epithelial Cell Line Established from C57BL/6 Mouse Mammary Glands.** (Eng) Vaidya, A. B. (Dept. Microbiology and Immunology, Hahnemann Medical Coll., Philadelphia, PA, 19102); Lasfargues, E. Y.; Sheffield, J. B.; Coutinho, W. G. *Virology* 90(1): 12-22; 1978.

An epithelioid cell line (C57MG) derived from the mammary glands of a 23-wk-old C57BL/6 breeder mouse was found to be susceptible to infection by murine mammary tumor virus (MuMTV). After infection of 10^6 C57MG cells/ml with approximately 10^{11} virus particles/ml, there were no changes in all morphological or growth characteristics, but annealing experiments with cellular DNA and complementary MuMTV DNA indicated that the infected C57MG cells had acquired three times as many MuMTV-related DNA sequences as uninfected cells. When high molecular weight cellular DNA was separated from unintegrated relatively low-molecular weight DNA and then hybridized again with complementary MuMTV DNA, the threefold increase was again demonstrated in the high-molecular weight fraction of infected cells vs the noninfected cells. This indicated a high level of MuMTV DNA integration in infected C57MG cells. Infected cells synthesized approximately 150 copies of MuMTV RNA per cell, and the concentration was enhanced by the addition of dexamethasone to the culture medium. Uninfected cells had no observable MuMTV RNA

under any conditions. Infected cells released 80 MuMTV virions/cell/24 hr before dexamethasone stimulation and 240 virions/cell/24 hr after stimulation. Electron microscopy indicated that the virions were indistinguishable from standard MuMTV. Thermal denaturation profiles were inadequate in determining whether the MuMTV produced by infected C57MG cells was of infecting virus or endogenous C57 virus origin. This in vitro infection of in vivo target cells for MuMTV provides an additional useful research tool. (27 refs)

78-6864 The Use of Molecular Hybridization to Track the Mode of Transmission and Distribution of Murine Mammary Tumor Viruses. A Model for Etiologic Studies of Human Breast Cancer. (Eng) Schlom, J. (NCI, NIH, Bethesda, MD, 20014); Colcher, D.; Drohan, W.; Kettmann, R. *Prog Exp Tumor Res* 21: 140-158; 1978.

Molecular hybridization techniques were used to determine whether mouse mammary tumor virus (MMTV) can be introduced into a given mouse via the germ cell line or via a non-germ line mechanism. Labeled 60S-70S RNA from a C3H mouse mammary tumor virus [MMTV(C3H)] was hybridized to an excess of DNA from livers of normal C3H mice, and the unhybridized RNA fraction was isolated. This recycled RNA hybridized to the DNA of C3H mammary tumors, but not to the DNA of C3H livers. Thus, it represents the portion of the MMTV(C3H) genome that is transmitted via a non-germ line mechanism. These sequences were present as provirus in the DNA's of mammary tumors of C3H, RIII, and GR mice, but they were not detected in the DNA's of mammary tumors of C3HfC57BL or BALB/c mice or in the DNA's of normal livers of several mouse strains. Thermal analysis of hybrids also demonstrated qualitative differences in MMTV proviral sequences in different mouse strains. The use of ³H-MMTV complementary DNA indicated quantitative differences in MMTV proviral sequences in different mouse strains and between normal and neoplastic tissues of a given mouse. The complexities of the murine model must therefore be considered in studies dealing with a viral involvement in human breast cancer. (23 refs)

78-6865 Translation of Mouse Mammary Tumor Virus RNA: Precursor Polypeptides are Phosphorylated during Processing. (Eng) Nüsse, R. (Dept. Virology, Antoni van Leeuwenhoekhuis, Plesmanlaan 122, Amsterdam, Netherlands); Asselbergs, F. A.; Salden, M. H.; Michalides, R. J.; Bloemendal, H. *Virology* 91(1): 106-115; 1978.

The synthesis of viral polypeptides of murine mammary tumor virus (MMTV) was studied by pulse-labeling of MMTV-producing cells and by translating MMTV virion RNA in vitro, in *Xenopus laevis* oocytes. Virus-related polypeptides were detected by immunoprecipitation with monospecific antisera against the major viral proteins gp49 and p24 and analysis of the immunoprecipitates

on polyacrylamide gels. In pulse-labeled MMTV-producing cells (Mm5mt/cl), a precursor 73,000-dalton polypeptide was immunoprecipitated by anti-p2 serum (Pr73gag). Pr73gag comigrated with the 73,000 dalton glycosylated precursor for the envelope proteins (Pr73gag) immunoprecipitated by anti-gp4 serum. Pr73gag was, during chase, converted into 76,000-dalton polypeptide that also reacted with the anti-p2 serum (Pr76gag). After prolonged incubation, the mature internal protein p24 was synthesized. Pulse-labeling with ³²P and subsequent chasing revealed that phosphate was incorporated into Pr76gag but not into Pr73gag. Isolated virion 70S RNA of MMTV, microinjected into *Xenopus* oocytes, gave rise to the synthesis of Pr73gag, Pr76gag, and p24, all immunoprecipitated by anti-p24 serum, and the viral core proteins p14 and p10, precipitated by polyvalent anti-MMTV serum. 70S RNA did not instruct synthesis of the viral envelope glycoproteins. (27 refs)

78-6866 Genetic Control of Endogenous Murine Mammary Tumour Viruses Reinvestigated. (Eng) Bentvelzen, P. (Radiobiological Inst. TNO, 151 Lange Kleiweg, Rijswijk, Netherlands); Brinkhof, J.; Haaijman, J. J. *Br J Cancer* 14(10): 1137-1147; 1978.

Milk samples of progeny mice from crosses between mouse strains that differ in the degree of release of endogenous murine mammary tumor virus (MuMTV) were tested for the presence of MuMTV antigens by the Sepharose bead immunofluorescence assay. The results indicated that, compared with nonexpressor BALB/c mouse, a single dominant gene controls the release of MuMTV in C3Hf mouse. However, the low-cancer strain C57BL carries a dominant, epistatic gene that can inhibit virus release in hybrids with C3Hf to a large extent. The high-cancer strain GRS harbors a dominant gene that, in F1 hybrids of all strains tested, causes the release of large quantities of a virus (MuMTV-P) that induces tumors at an early age. Other genetic factors control the release of a lesser oncogenic virus in the GRS strain. The animals tested for MuMTV antigens were subsequently force-bred for mammary carcinoma production. Although there was a good correlation between virus release at a young age and tumor development at a late age, a substantial number of virus-negative mice developed mammary tumors. (3 refs)

78-6867 Expression of Mammary Tumor Virus Antigens in the Male Secondary Genital Organs of the GRS/A Mouse Strain. (Eng) Imai, S. (2nd Dept. Pathology, Nara Medical Univ., Nara, Japan); Imai, N.; Morimoto, J.; Tsubura, Y. *J Nara Med Assoc* 29(3): 503-508; 1978.

Four immunological methods (immunodiffusion, microimmunodiffusion, immunofluorescence absorption, and radioimmunoassay) were compared for their ability to detect mammary tumor virus (MTV) antigen, and a pilot study was

out to correlate the expression of MTV antigen in genital organs to the Mtv-2 gene in GRS/A strain mice. Immunassay was the most sensitive method for detecting MTV antigen. Studies of tissue extracts of the epididymis and the coagulating gland suggested that MTV antigen expression in these organs might be under the control of the 2 gene locus. (20 refs)

868 Involvement of Mouse Mammary Tumor Virus in Spontaneous and Hormone-induced Mammary Tumors in Low-Mammary-Tumor Mouse Strains.

Michalides, R. (Div. Virology, Antoni van Leeuwenhoek Institute, Netherlands Cancer Inst., Amsterdam, Netherlands); Deemter, L.; Nusse, R.; Ropcke, G.; Boot, L. *J Virol* 55:551-559; 1978.

The possible involvement of mouse mammary tumor virus (MTV) in spontaneous and hormone-induced mammary tumors in low-mammary-tumor mouse strains was studied by comparing the amounts of MTV DNA and MTV RNA in mammary tumors, mammary glands, and other organs of mice with and without hormonal treatments. Hormone treatment was performed as follows: four hypophyseal isografts were implanted, two into each kidney, of C3Hf females at the age of 6-8 wk; one hypophysis was implanted into the left side of castrated O20 males at the age of 6-8 wk. Estrone (1 mg/liter) was added to the drinking water throughout the whole life-span of the mouse, and 3 mg of progesterone was given sc. The virus-induced mammary tumors contained 10³ MTV DNA copies in their cellular genome than the amount present in the genome of cells of noninfected tissues. The former also contained larger amounts of MTV RNA than those found in the lactating mammary glands of these animals. Hormone-induced mammary tumors contained the same number of MTV DNA copies in their cellular genome as the other tissues of the animals. Hormone treatment resulted in an increased expression of MTV RNA in the lactating mammary glands, but this enhanced level of MTV expression was not maintained in the hormone-induced mammary tumors. Spontaneous mammary tumors of BALB/c mice did not contain enhanced amounts of MTV DNA or MTV RNA. (27 refs.)

869 An Inquiry into the Involvement of Nonhuman Primate Breast Cancer Viruses in the Human Disease: Experiments in Malignant Breast Ductal Stem Cells.

McGrath, C. M. (Dept. Biology, Michigan Cancer Foundation, Detroit, MI, 48201); Soule, H. D. In: *Mechanisms of Carcinogenesis. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, W. W.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory); Origins of Human Cancer. Vol. 4 (Book 99 pp.; 1287-1303; 1977.

An attempt was made to determine if mouse mammary tu-

mor virus (MMTV)- or Mason-Pfizer monkey virus (MPMV)-related information is expressed in malignant breast ductal cells (MCF-7) and is present as a provirus in malignant cell DNA. Line MCF-7 was derived from cells cloned from pleural effusions of a scirrhous carcinoma, which contained cells with prominent histologic and hormonal markers. MMTV-related RNA was detected upon hybridization (hyb) with MMTV complementary DNA (cDNA) in 0.5 M Na⁺ and under stringent conditions of hyb (hydrolysis with a single-strand-specific nuclease), suggesting extensive sequence homology between MMTV-related RNA in MCF-7 cells and murine breast virus genomic RNA. MMTV-related RNA was not detected in two nonbreast malignant cell lines under identical hyb conditions; ie, MMTV cDNA does not complex nonspecifically with any human cell RNA at the RNA concentrations used. In addition, antigens cross-reactive with MMTV virion antigens were detected in the MCF-7 cells, suggesting that MMTV-related RNA is informational in the malignant breast stem cell. However, MMTV-related DNA sequences were not detected in MCF-7 DNA under hyb conditions capable of detecting one av copy per cell. MPMV-related RNA was not detected in MCF-7 cells using MPMV cDNA representing a minimum of 60% of MPMV RNA, and the major internal protein of MPMV (p27) was not detected by immunofluorescence assay or radioimmunoassay. MPMV sequences were not found in MCF-7 DNA at a significantly higher frequency than that found in normal human cells, which suggests that the normal breast precursor of MCF-7 was not infected by MPMV, at least not in the same manner as that in other human cells. Although the data do not rule out the possibility of information related to MMTV or MPMV in DNA of the human breast carcinoma cells, the inability to detect a major portion of the genome related to either virus in DNA of MCF-7 cells provides a rationale for isolating viruses directly from breast carcinoma cells for use as probes in etiologic studies of breast cancer. (54 refs)

78-6870 The Relative Hydrophobicity of Oncornaviral Structural Proteins. (Eng) Marcus, S. L.

(Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Smith, S. W.; Racevskis, J.; Sarkar, N. H. *Virology* 86(2): 398-412; 1978.

The interaction of the major structural polypeptides of murine mammary tumor virus (MuMTV) and Rauscher murine leukemia virus (R-MuLV) with alkyl-agarose derivatives containing hydrocarbon arms of various lengths was examined. Both R-MuLV and MuMTV polypeptides were selectively adsorbed to octylimino (C₈)-agarose and decylimino (C₁₀)-agarose columns. The proteins were eluted with buffers containing 8.5 M ethylene glycol. In certain cases, the polypeptides could be removed from the columns only by the addition of detergent to the ethylene glyco-containing buffers. By examining the chromatographic behavior of the polypeptides, and the conditions required for their elution, it was possible to determine the relative potential for the hydro-

phobic interaction (degree of hydrophobicity) of each viral protein. The most hydrophobic MuMTV polypeptide was the glycoprotein gp34, p15(E) and the gag precursor prp67 appeared to be the most hydrophobic R-MuLV proteins. The least hydrophobic MuMTV polypeptides were p23 and p16, and those of R-MuLV proteins were the two major internal proteins, p10 and p12. Four degrees of apparent hydrophobicity, from least hydrophobic to strongly hydrophobic, could be discerned by this procedure. The major core proteins (MuMTV p23, R-MuLV p30) of both viruses were slightly hydrophobic, but their major glycoproteins were moderately hydrophobic. The degree of hydrophobicity of oncornaviral polypeptides, as determined by hydrophobic chromatography, to the different subviral components with which the proteins were associated. Hydrophobic chromatography may provide a new parameter for the characterization of oncornaviral polypeptides. (52 refs)

- 78-6871 Source and Hormone-Dependence of GIX-gp70 in Mouse Serum.** (Eng) Obata, Y. (Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Stockert, E.; Yamaguchi, M.; Boyse, E. A. *J Exp Med* 148(3): 793-798; 1978.

Strain 129 mice (not overt producers of virus) were examined to determine the tissue origin and hormone dependence of serum GIX-gp70 (a 70,000 dalton polypeptide distinguished by the type-specific antigen GIX). The source of GIX-gp70 could not be traced to any organ or tissue that is known to be GIX-positive by serological tests. Hematopoietic tissues were excluded as the source of serum GIX-gp70 by tests with reciprocal radiation chimeras made from 129 and 129-GIX-donors and recipients. Thymus and spleen were excluded because excision of these organs did not affect levels of GIX-gp70 in the serum. The serum of young adult 129 males contained approx four times as much GIX-gp70 as adult 129 females and the levels rose with increasing age in both sexes. Castration of 129 males reduced the level of serum GIX-gp70 to that of females, and the level was fully restored by testosterone. Thus, the epididymis and seminal fluid, although rich in GIX-gp70, do not contribute significant amounts of GIX-gp70 to the serum. The level of GIX-gp70 in the serum of testosterone-treated females, although more than double that of untreated females, did not reach the level of normal males, under the conditions tested. This may signify that GIX-gp70 production in males is subject to imprinting by testosterone in early life. Evidently, the main source of GIX-gp70 is a tissue or organ that is common to males and females, is directly or indirectly responsive to testosterone, and has not so far been identified serologically as GIX-positive. (16 refs.)

- 78-6872 Xenotropic MuLV are Subject to Restriction by the Fv-1 Locus of the Mouse (Meeting Abstract).** (Eng) Gautsch, J. W. (Dept. Cellular and Develop-

mental Immunology, Res. Inst. Scripps Clinic, La Jolla, CA)
Jensen, F. C.; Elder, J. H.; Lerner, R. A. *J Cell Biol* 79(part 2): 406a; 1978. (no refs)

- 78-6873 Wild Rat Type C Virus: Isolation and Characterization.** (Eng) Rasheed, S. (Dept. Pathology, Univ. Southern California Sch. Medicine, Los Angeles, CA 90033); Charman, H. P.; Gardner, M. B. *Virology* 89(2): 605-609; 1978.

C-type virus was isolated from a wild male rat (*Rattus rattus*) and compared with the C-type viruses from two strains of laboratory rat. The wild rat was injected sc with 200 mg of 3-methylcholanthrene, and the resulting fibrosarcoma, which developed at the inoculation site within 13 mo, was directly cultured. The wild-rat tumor cells were tested for virus production at various passages by the supernatant RNA-dependent DNA polymerase activity assay. The results of virus purification, neutralization, and interference studies showed that the isolated virus exhibits the general properties of rat leukemia virus (RLV) including the RLV-specific p30 antigen. It showed low infectivity for rat cells and no other species. It was antigenically similar to other RLV strains of laboratory rat origin. It is concluded that a C-type virus genome is potentially activable in fetal rats never subjected to laboratory inbreeding. (12 refs)

- 78-6874 Physicochemical, Biological and Serological Properties of a Leukemogenic Virus Isolated from Cultured RadLV-induced Lymphomas of C57BL/Ka Mice.** (Eng) Declève, A. (Cancer Biology Res. Lab., Dept. Radiology, Stanford Univ. Sch. Medicine, Stanford, CA 94305); Lieberman, M.; Ihle, J. N.; Rosenthal, P. N.; Lunn, M. L.; Kaplan, H. S. *Virology* 90(1): 23-35; 1978.

A radiation leukemia virus (RadLV/VL₁) produced by C57BL/Ka cells, which were derived from a RadLV-induced C57BL/Ka mouse thymic lymphoma, was characterized by physicochemical studies with ³H-uridine-, ³H-thymidine- and ³H-leucine-labeled RadLV/VL₁, as well as inhibition of the viral reverse transcriptase (RT) by antiserum against Moloney murine leukemia virus RT, indicated that the isolate was a C-type RNA virus. The virus was not neutralized by antibodies prepared against other ecotropic or xenotropic murine C-type viruses, used singly or in combination, but was neutralized by rabbit and mouse anti-RadLV serum. The inability of RadLV and RadLV/VL₁ to induce cytopathic changes or transformed cell foci in mink lung cell culture even after addition of xenotropic BL/Ka(X) virus, indicated that the in vitro and in vivo host-age patterns of these viruses could not be attributed to the presence of competent or defective mink cell focus-forming virus. In co-infection experiments using live or UV-inactivated viruses, none of the viruses tested was able to interfere with RadLV and RadLV/VL₁. However, purified RadLV/VL₁ protein gp70

ificantly interfered with the infectivity of RadLV and VL/VL₃, indicating that the major envelope glycoprotein of C57BL/Ka thymotropic and leukemogenic viruses is distinct from other tested viruses. This was supported by the fact that in competitive radioimmunoassay for RadLV/VL₃, that used an anti-RadLV/VL₃ antiserum, none of the viruses competed equivalently to RadLV/VL₃ gp71. Other results for the p12 and p30 component of RadLV/VL₃ did not show this specificity. (20 refs)

78-6875 The Pattern of Involvement of Murine Lymphoid Tissues by Primary Lymphomas of Different Cell Lineage. (Eng) Baird, S. (Veterans Admin. Hosp., La Jolla Village Drive, La Jolla, CA, 92161); Raschke, J. *J. Natl. Cancer Inst. Pathol* 9(1): 47-50; 1978.

The metastatic patterns of several different lymphomas were compared with the recirculation pattern of normal T lymphocytes. T and non-T lymphomas were induced in newborn B/c mice by injections of Abelson leukemia virus preparations. Many primary T-cell lymphomas migrated in the lymphoid system in areas occupied by normal recirculating lymphocytes (eg, the periarteriolar white sheath of the spleen and thymus). Other lymphomas lacking T-cell markers did not show this pattern of migration. This lends support to the use of classifying human lymphoid neoplasms as either T- or B-derived on the basis of their surface antigen markers and morphology in lymph nodes. The data indicate that murine lymphomas are good models for studying the metastatic pattern of some human lymphomas under prospective controlled conditions. (3 refs)

78-6876 Genetic and Immunological Controls of Murine Leukemia Viruses in AKR Mice. (Eng) Nowinski, R. C. (Fred Hutchinson Cancer Res. Center, 1124 Columbia St., Seattle, WA, 98104); Baron, J. M. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1977*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 1978; 198; 1977.

Low-leukemic AKR mice, low-leukemic C57L mice, and their hybrids were examined for the production of murine leukemia virus (MuLV) p30 antigen, the production of infectious non-oncogenic MuLV (ieMuLV), and the development of immunity against various proteins of the endogenous virion. C57L mice did not produce ieMuLV, and p30 expression in their spleens was low. In contrast, AKR mice produced high titers of ieMuLV, and p30 expression in the spleens was high. F₁ hybrids produced high titers of ieMuLV and expressed high p30 levels in their spleens. Backcross C57L x AKR x C57L F₁ mice genetically segregated for the expression of infectious MuLV and p30: 68% produced ieMuLV, and did not contain detectable virus. This ratio was statisti-

cally compatible with a two-gene model for the control of MuLV. Virus-producing backcross mice expressed high levels of p30; most of the nonproducers expressed low p30 levels, but a minor population expressed high p30 levels. The latter represent a phenotype distinct from that of the AKR and C57L parents. AKR and C57L strains produced low titers of natural antibody (ab) against eMuLV; their F₁ hybrids produced anti-MuLV ab in a higher titer than either of the parents. Progeny of the backcross mice that contained ieMuLV produced anti-MuLV ab's. Thus, the persistent production of ab against high-titered MuLV in these mice did not result in immunological tolerance toward viral antigens. Mice that did not contain ieMuLV could be separated into ab-producing and ab-nonproducing classes. The absence of a detectable ab to MuLV in a mouse was invariably associated with a virus-free phenotype. Overt production of endogenous eMuLV probably served as a major immunogenic stimulus for the production of anti-MuLV ab in these mice. (73 refs)

78-6877 Acquisition of Oncogenicity by Endogenous Mouse Type C Viruses: Effects of Variations in *env* and *gag* Genes. (Eng) Devare, S. G. (Lab. Cellular and Molecular Biology, NCI, Bethesda, MD, 20014); Rapp, U. R.; Todaro, G. J.; Stephenson, J. R. *J. Virol* 28(2): 457-465; 1978.

Competition immunoassays were used to compare the structural proteins of different groups of highly leukemogenic viruses: dual-tropic isolates derived from the thymuses of preleukemic or leukemic AKR mice and strains generated by in vitro or in vivo passage of a weakly infectious endogenous virus of C3H mouse cells. The results show that three AKR-derived isolates and two C3H passage-derived isolates possess *gag* gene-coded proteins immunologically indistinguishable from those of AKR murine leukemia virus. This observation confirms the results of a previous study in which the major structural proteins of several AKR-derived mink cell focus-inducing isolates were found to be similar to those of AKR-MuLV. In contrast, *gag* gene-coded proteins of two other leukemogenic isolates of C3H origin, including one ecotropic and one dual-tropic virus, were indistinguishable from those of Moloney MuLV. All of the highly leukemogenic viruses, including those of AKR and C3H origin, as well as C3H-derived virus, FC clone 3, possessed common envelope glycoprotein antigenic determinants of a class not exhibited by the nononcogenic ecotropic viruses from which they were derived. The results suggest that oncogenic variants of endogenous ecotropic mouse C-type viruses are derived by genetic recombination. This recombinational event appears to involve the acquisition, by different ecotropic viruses, of a common class of endogenous virus-coded envelope glycoprotein determinants that may be required, but not necessarily sufficient, for leukemogenicity. (19 refs)

78-6878 A Target Cell Assay for Friend Virus. (Eng) Steinheider, G. (Albert Einstein Coll. Medicine,

1300 Morris Park Ave., Bronx, NY, 10461); Steeves, R. *Leuk Res* 2(3): 197-200; 1978.

An assay was developed for the in vitro infection and subsequent quantification of the hemopoietic target cells from spleen and bone marrow of DBA/2 mice that can replicate Friend virus (FV). The presence of macroscopic spleen colonies in recipient CDF₁ mice was used to quantify the number of infected FV-releasing cells. DBA/2 mice were bled on two consecutive days and sacrificed. Bone marrow and spleen cells were isolated and cultured with 5×10^4 spleen focus-forming units/ml of FV at 37 C for 1 hr and then injected into CDF₁ mice. After 8 days, the CDF₁ spleens were removed and spleen colonies counted. Inoculation with different dilutions of infected DBA/2 cells revealed a linear relationship between the quantity of infected cells transferred and the number of resulting spleen colonies. Lethal irradiation (5,000 R) of 50% of the infected DBA/2 donor cells before injection yielded 73.8 ± 2.1 colonies from the intact cells but none from the irradiated cells, showing that the production of virus and the generation of spleen colonies requires a living cell and ruling against the passive transfer of cell-adherent virus only. When the numbers of infected spleen and bone marrow cells from bled and unbled DBA/2 donor mice that resulted in spleen colonies in recipient mice were compared, it was concluded that bone marrow had more target cells than spleen and that the hemopoietic stimulus of bleeding was greater in the spleen than in the bone marrow. This assay may prove useful in finding and improving the conditions for true oncogenic transformation by FV in vitro, which was not observed in this experiment. (18 refs)

78-6879 Antibody-induced Loss of Friend Virus Leukemia Cell Surface Antigens Occurs During Progression of Erythroleukemia In F₁ Mice. (Eng) Doig, D. (Hamilton, MT, 59840); Chesebro, B. *J Exp Med* 148(5): 1109-1121; 1978.

The ability of anti-Friend leukemia virus (FLV) serum + rabbit complement (C') to lyse leukemic spleen cells from (B10A x A/WynSn)F₁ and (B10.A x A.BY)F₁ mice was studied. Leukemic cells from 27/28 mice showed high specific lysis (20%-45%) when tested 8-9 days after inoculation of the mice with B-tropic FLV (1,500 focus-forming units iv), but cells of only 5/31 mice had > 20% lysis 30-90 days post-inoculation. Quantitation of FV-induced cell surface antigens (CSA) by absorption of cytotoxic antibodies demonstrated that the antigen levels were 4.5-fold higher in early (8-9 days) leukemic cells than in later cells. No difference in absorption of antibody to H-2d histocompatibility antigens was noted when (C57BL/10 x BALB.B)F₁ anti-(B10.D2 x BALB/c)F₁ (anti-H-2d) serum was absorbed with (B10.A x A/WynSn)F₁ early or late or leukemic spleen cells or with normal cells. There were also no changes in H-2K or H-2D antigens. Membrane immunofluorescence studies confirmed the loss of FV-induced CSA in the later stages of leukemia. Acetone-fixed cells

showed no difference between early or late F₁ leukemic spleen cells when tested for the presence of cytoplasmic antigen with conjugated goat anti-Moloney LV. The decrease in FV-induced CSA could be reversed after adoptive transfer of leukemic cells to lethally irradiated (900 rads) unimmunized congenic hybrid mice, but not after transfer to immunized recipients. When late F₁ leukemic spleen cells were cultured in vitro in the absence of anti-FV antibody, no increase in FV-induced CSA was noted. Conversely, no decrease in FV-induced CSA occurred in early leukemic spleen cells cultured with high concentrations of cytotoxic anti-F serum. The observed decreased antigen expression may represent a tumor cell adaptation to host immune pressures generated during the course of FV leukemia. (19 refs)

78-6880 Biosynthesis and Metabolism of Viral Proteins Expressed on the Surface of Murine Leukemia Virus-infected Cells. (Eng) Ledbetter, J. A. (Fred Hutchinson Cancer Res. Ctr., 1124 Columbia St., Seattle, WA, 98104); Nowinski, R. C.; Eisenman, R. N. *Virology* 91(1): 116-129; 1978.

Viral proteins expressed on the surface of the Friend virus-induced leukemia E(female)G2 were detected by immune precipitation assays from detergent lysates of cells that were labeled by the ¹²⁵I-lactoperoxidase method. These proteins included gp70 and two glycosylated polyproteins that contained antigens coded by the *gag* gene. The 95,000-dalton polyprotein (gpP95*gag*) reacted with antisera against p30, p12, and p10; the 85,000-dalton polyprotein (gpP85*gag*) reacted with antisera against p30 and p12, but not against p10. Neither polyprotein reacted with reverse transcriptase antiserum. GpP95*gag* and gpP85*gag* contained glucosamine only, but gp70 contained both glucosamine and fucose. Tryptic peptide maps of the iodinated polyproteins indicated that gpP95*gag* and gpP85*gag* were highly related. Viral polyproteins with the same electrophoretic mobilities as the cell surface gpP95*gag* and gpP85*gag* were also detected by immune precipitation of lysates from cells that were metabolically labeled with radioactive methionine or arginine. In kinetic studies, gpP95*gag* was labeled rapidly following a 10-min pulse, in contrast, gpP85*gag* was not detected until late in the chase period, suggesting that this glycosylated polyprotein was a processed product of gpP95*gag*. Mapping of arginine-labeled tryptic peptides of different *gag* polyproteins supported this assumption. In fact, peptide mapping with isolated viral proteins demonstrated that gpP95*gag* and gpP85*gag* contained peptides of p30, p15/p12, and p10. The presence of p10 peptides in both polyproteins was unanticipated, since the gpP85*gag* did not react in immune precipitation assays with anti-p10 serum. By peptide mapping, both of the glycosylated polyproteins also appeared highly related to Pr75*gag*, the intracellular precursor to virion *gag* proteins. The data suggest that gpP85*gag* is a glycosylated cell surface form of the primary *gag* precursor Pr75*gag*, and that processing of gpP95*gag* results in the production of gpP85*gag*. (35 refs)

881 **The Integrated Genome of Murine Leukemia Virus.** (Eng) Steffen, D. (Dept. Biology, Center Res., Massachusetts Inst. Technology, Cambridge, 02139); Weinberg, R. A. *Cell* 15(3): 1003-1010; 1978.

Southern gel filter transfer technique was used to characterize the integrated genome of Moloney murine leukemia virus (M-MuLV) and the genomes of the endogenous mouse viruses. A study of 10 clones of rat NRK cells independently infected by M-MuLV indicated a minimum of 15 integration sites into which the M-MuLV provirus could be inserted. No common integration site was observed among these clones. Cells productively infected by M-MuLV acquired multiple proviruses, but infected cells unable to produce virus contained only one M-MuLV provirus. Once established, the integrated genomes were stable for at least two years after reinfection. A spectrum of *Eco*RI-cleaved mouse DNA fragments containing endogenous MuLV genomes was determined through the use of M-MuLV probes. DNA's of different inbred laboratory mouse strains yielded similar provirus patterns, with each strain showing minor characteristic differences. In some instances, mouse cells infected by M-MuLV yielded additional proviruses beyond those seen in uninfected cells. DNA's from three different M-MuLV-induced lymphomas indicated, as in rat cells, multiple possible integration sites. (20 refs)

882 **T1 Oligonucleotide Maps of Moloney and HIX Murine Leukemia Viruses.** (Eng) Faller, D. V. (Dept. and Center for Cancer Res., Massachusetts Inst. Technology, Cambridge, MA, 02139); Hopkins, N. *Virology* 90(2): 265-273; 1978.

Genomes of a HIX virus, its putative ecotropic parent (Moloney IC), and another clonal isolate of Moloney virus (Moloney clone-1) were studied by RNase T1 oligonucleotide analysis. HIX and Moloney IC shared 42 out of 51-53 of their cleavable T1 oligonucleotides, and the two clonal isolates of Moloney virus shared 46 T1 oligonucleotides. Each virus possessed some oligonucleotides not found in the others. Based on oligonucleotide mapping, it was concluded that: oligonucleotides shared by HIX and Moloney viruses probably lie in the same relative positions on their genomes; the oligonucleotides found in Moloney IC but not HIX, and vice versa, are clustered in corresponding regions in the 3' halves of the Moloney IC and HIX maps; and the oligonucleotides that differ between the two clonal isolates of Moloney virus are scattered along the 5' half of the Moloney IC oligonucleotide map. (26 refs)

883 **Host Response Against Oncorna Virus Induced Lymphomas in Mice.** (Eng) Becker, S. (Dept. of Cancer Biology, Karolinska Inst., S-104 01 Stockholm 60, Sweden); Klein, E. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Sym-*

posium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977. Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 61-68; 1978.

The host response against oncornavirus-induced lymphomas in mice was first demonstrated by transplantation immunity and then by the presence of antibodies reacting with the cell surface. With the development of in vitro methods and the discovery of various lymphocyte subpopulations, attention was focused on cell-mediated killing effects. Three cell types with killing potential can be distinguished: lymphocytes mediating antibody-dependent killing, sensitized T cells, and natural killer (NK) cells. The reactivity to cell-mediated cytotoxicity in vitro do not, however, correlate. Two serologically cross-reacting lymphoma culture lines (YAC-1 and RBL-5) differ in their sensitivity to cell-mediated cytotoxicity by both NK and sensitized T cells. A cell line selected from YAC for low amounts of Moloney cell-surface antigen recognized by mouse antibodies (YAC-IR) does not show decreased sensitivity to the killing effect of either NK or sensitized T cells. Histocompatibility restriction prevails in short-term cytotoxic tests with sensitized effectors. In cold target inhibition tests with A x C57BL F₁ (H-2a, H-2b); but not C57BL (H-2b), effectors, anti-RBL-5 (H-2b) cytotoxicity was inhibited by YAC-1 (H-2a), indicating the recognition of both H-2a and H-2b target cells. When F₁ effector spleen cells were cultured in vitro in the presence of RBL-5 cells, effector cells against RBL-5 were selected with regard to histocompatibility restriction, as indicated by the fact that YAC-1 cells were not inhibitory. (27 refs)

78-6884 **Biological, Immunological, and Biochemical Evidence that HIX Virus is a Recombinant Between Moloney Leukemia Virus and a Murine Xenotropic C Type Virus.** (Eng) Fischinger, P. J. (Lab. Viral Carcinogenesis, NCI, Bethesda, MD, 20014); Frankel, A. E.; Elder, J. H.; Lerner, R. A.; Ihle, J. N.; Bolognesi, D. P. *Virology* 90(2): 241-254; 1978.

Various characteristics of HIX virus were studied to determine its origin and relationship to other murine C-type oncornaviruses. Cloned amphotropic HIX virus adapted to grow in human RD cells (HIX-RD) could enter mouse cells but essentially lost the capacity to propagate in them. The neutralizing antibody-accessible determinants on HIX and HIX-RD resembled those of Moloney murine leukemia virus (M-MuLV) and, to a lesser extent, those of Friend or Rauscher MuLV and two murine xenotropic viruses (MuX). The entire *gag* region structural proteins of HIX appeared to be very similar or identical to those of M-MuLV. HIX and HIX-RD gp70 polypeptides were not mixtures of MuX and M-MuLV gp70's but exhibited type-specific determinants different from known MuLV's. Tryptic peptide maps of the gp70's of HIX-type viruses indicated that HIX virus is the product of recombination between ecotropic and xenotropic

viruses within the envelope gene. There appeared to be little if any HIX-specific complementary DNA sequences that were not a part of or closely related to M-MuLV RNA. (46 refs)

- 78-6885 Mammalian Retrovirus-associated RNase H Is Virus Coded.** (Eng) Lai, M. T. (Tumor Virology Lab., Salk Inst., San Diego, CA, 92112); Verma, I. M.; Tronick, S. R.; Aaronson, S. A. *J Virol* 27(3): 823-825; 1978.

The virus-coded nature of the RNase H activity of a temperature-sensitive Rauscher murine leukemia virus (R-MuLV) mutant with thermolabile polymerase activity (*ts29*) was studied. Both the DNA polymerase and RNase H activities of the mutant viral enzyme were two to three times more thermolabile than the corresponding activities from the wild-type R-MuLV. A wild-type (phenotypic) revertant of *ts29* was isolated in which the kinetics of heat inactivation of the DNA polymerase and RNase H were comparable to that of the wild-type viral reverse transcriptase. The restriction to infection by two other R-MuLV mutants (*ts17* and *ts19*, which are also defective in early postpenetration functions) did not appear to be associated with a measurable function of the viral enzyme. The thermolability of the RNase H associated with *ts29* reverse transcriptase demonstrates that this activity, as well as that of the DNA polymerase, is a product of the mammalian C-type viral genome. The two activities appear to be localized on the same polypeptide chain. (18 refs.)

- 78-6886 Characteristics of Cellular RNA in Viral Leukemia.** (Rus) Butenko, Z. A. (Inst. Problem Oncology, Kiev, USSR); Smirnova, I. A. *Vopr Onkol* 24(10): 18-23; 1978.

This study was designed as an attempt to evaluate characteristic features of polyadenylate rich (poly-A) sequences in RNA from Rauscher leukemia virus (RLV)-transformed cells. RNA specimens were isolated from the spleens of BALB/c mice inoculated with RLV and of leukemia-free BALB/c mice. There was no difference in content of cytoplasmic poly-A rich RNA between the leukemic and normal cells, while the content of nuclear poly-A rich RNA in leukemic cells was slightly greater than that in normal cells. Electrophoretic analysis of nuclear poly-A rich RNA showed the presence of additional 30-35 S fraction in leukemic cells, which was absent in normal cells. (29 refs)

- 78-6887 Biosynthesis of Rauscher Leukemia Viral Proteins.** (Eng) Arlinghaus, R. B. (Biology Dept., Univ. Texas System Cancer Center, M. D. Anderson Hosp. and Tumor Inst., Houston, TX, 77030); Jamjoom, G. A.; Arcement, L. J.; Naso, R. B.; Karshin, W. L. In: *Third Inter-*

national Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 293-309; 1977.

The synthesis of murine leukemia viral proteins was investigated by immune precipitation of ³⁵S-methionine pulse-labeled and chased cultures of Rauscher murine leukemia virus (RLV)-infected JLS-V16 cells. The RLV structural proteins were found to be synthesized in infected cells by way of high-mol-wt precursor polyproteins. In a working model for the biosynthesis of RLV proteins, Pr1a + b, Pr2a + b, and Pr3 [approx 200,000 daltons (200K), 90K, and 80K, respectively] are primary gene products derived by translation of 35S viral RNA. Pr3 arises by a translational control mechanism in which ribosomes detach from the 35S viral messenger RNA after translating Pr3. However, the ribosomes can proceed to translate the polymerase gene in 1/10-20 times to yield Pr1a + b. Pr1a + b thus is cleaved to form reverse transcriptase (RT)-specific proteins [Pr RT1 (145K), 2 (135K), and 3 (85K)] and, possibly, Pr3. Viral proteins p30, p15, p12, and p10 and RT are contained in Pr1a + b, gp69/71 and p12E are contained in Pr2a + b. Pr1b may be derived from Pr1a by proteolytic cleavage or some other posttranslational alteration. Pr2a and Pr2b are assumed to have the same core polypeptide that is glycosylated to different extents during or after translation to produce Pr2a or Pr2b. p30, p15, p12, and p10 are mainly generated by a series of cleavages of Pr3, producing Pr4, Pr5, and Pr6 (65K, 55K, and 45K) in that order. p10 is produced by cleavage of Pr4 to Pr5, p15 by cleavage of Pr5 to Pr6, and p30 and p12 by cleavage of Pr6. Treatment of infected cells with a proteolytic enzyme inhibitor such as tolylsulfonylphenylalanyl chloromethyl ketone caused a build-up of Pr1a + b and prevented the formation of mature structural proteins and RT. The main feature of this model is that a translational control mechanism exists that results in synthesis of a much lower amount of RT than viral structural proteins. (33 refs)

- 78-6888 Inhibition by RNA of RNase H Activity Associated with Reverse Transcriptase in Rauscher Murine Leukemia Virus Cores.** (Eng) Sarngadharan, M. G. (Dept. Cell Biology, Litton Bionetics, Inc., Bethesda, MD, 20014); Kalyanaraman, V. S.; Gallo, R. C. *J Virol* 27(3): 568-575; 1978.

Virion cores of Rauscher murine leukemia virus (R-MuLV) were examined to determine if the absence of detectable RNase H activity in these fractions was a result of some selective inhibition. Earlier findings that R-MuLV cores, when isolated by a density gradient, show no detectable RNase H activity were confirmed. The absence of RNase H activity in the core fractions was shown to be due to a selective inhibition of this activity by viral RNA. Direct evidence that viral RNA-mediated inhibition of the RNase H activity was the

cause of the apparent absence of RNase H activity in the isolated core fraction was obtained by studying the effect of viral and other RNAs on RNase H purified from isolated viral cores. Irrespective of the substrate used for measurement, the enzyme was generally inhibited by a variety of natural and synthetic RNAs. Although viral RNA was a strong inhibitor, polyguanylic acid was by far the most potent inhibitor of the reverse transcriptase-associated RNase H. The RNase H activity associated with a highly purified preparation of simian sarcoma virus reverse transcriptase was also similarly inhibited by RNA, suggesting that this property may be general for mammalian RNA tumor viruses. A rabbit anti-reverse transcriptase IgG inhibited both the reverse transcriptase and RNase H activities of the enzyme purified from cores. Overall, the results are consistent with the idea that RNase H and reverse transcriptase are two activities present on the same protein molecule. (21 refs.)

78-6889 A Study of the Synthesis and Processing of Rauscher Leukemia Virus-specific Polypeptides in Infected Cells and Their Assembly into Virus Particles. (Eng) Jamjoom, G. A. (Univ. Texas Health Science Center, Houston Graduate Sch. Biomedical Sciences, Houston, TX). *Diss Abstr Int [B]* 39(2): 666; 1978. (no refs)

78-6890 Physical Association of Murine Leukemia Virus Envelope Glycoprotein and Mouse Major Histocompatibility Antigens on Leukemia Cell Surfaces and their Recognition as Target Antigens by Cytotoxic T-Lymphocytes (Meeting Abstract). (Eng) Zarling, D. A. (Univ. Wisconsin, Madison, WI, 53706); Watson, A.; Duke, R. E.; Keshet, I.; Bach, F. H. *J Cell Biol* 79(2, part 2): 401a; 1978. (no refs)

78-6891 Persistence of Sindbis Virus in Oncornavirus-producing Cultures. (Rus) Pogodina, V. V. (Inst. Poliomyelitis and Viral Encephalitis, Moscow, USSR); Fokina, G. I.; Kiseleva, L. L.; Graevskaia, N. A.; Sito, A. F.; Ivanov, A. P. *Vopr Virusol* (2): 196-201; 1978.

Two continuous cell lines (PKR-1 and PKR-2) were derived from a primary culture of BALB/c embryo cells infected with Rauscher leukemia virus (RLV). PKR-1 and PKR-2 were infected with Sindbis virus (SV) and/or repeatedly infected with RLV. The cultures, maintained for 18 mo, retained their ability to produce C-type oncornavirus. Marked reverse transcriptase activity was detected in line PKR-1 from passages 19 to 68 and in PKR-2 from passages 26 to 70. The lines contained the murine oncornavirus-specific antigen gs-1. Characteristic features of SV persistence were development of homologous interference with SV without interferon participation and progressive repression of arbor virus and oncornavirus function. (8 refs)

78-6892 B-MuX: A Unique Murine C-type Virus Containing the "env" Gene of Xenotropic Viruses and the "gag" Gene of the Ecotropic Virus. (Eng) Ihle, J. N. (Cancer Biology Program, NCI Frederick Cancer Res. Center, Frederick, MD, 21501); Fischinger, P.; Bolognesi, D.; Elder, J.; Gautsch, J. W. *Virology* 90(2): 255-264; 1978.

A unique xenotropiclike virus (B-MuX) isolated by clonal purification from iododeoxyuridine-induced Kirsten sarcoma virus-transformed BALB/c nonproducer cells was characterized. B-MuX had the host-range properties of other murine C-type xenotropic viruses. It reciprocally shared interference properties with BALB:virus-2 (BALB-1), and the neutralization spectra suggested that the two viruses had very similar, if not identical, neutralizing antibody-eliciting determinants on their gp70 polypeptides. B-MuX contained an *env* gene similar to that of the endogenous BALB-2 xenotropic virus and *gag* gene products similar to those of the endogenous ecotropic virus. The p30 of B-MuX was structurally similar to that of the endogenous ecotropic virus and differed from the p30 of the BALB/c xenotropic virus, as determined by tryptic peptide analysis. The results suggest that B-MuX is a new type of endogenous xenotropic virus or a recombinant between the endogenous xenotropic and ecotropic viruses. The *gag* gene region does not contribute to this virological property. (34 refs)

78-6893 Surface Morphology of Normal and Virus-transformed Cells In Vitro. (Rus) Samil'chuk, E. I. (Lab. Experimental Oncology, Inst. Experimental Pathology and Therapy, Acad. Medical Sciences, USSR, Sukhumi, USSR); Timanovskaia, V. V.; Kukava, G. G. *Tsitologiia* 20(10): 1205-1208; 1978.

The morphology of the mink lung cell line Mv1Lu and its Kirsten sarcoma virus-transformed counterpart, Ki-Mv1Lu, was studied in vitro. In monolayer culture, the morphology of the normal cells differed sharply from that of the transformed cells. The Mv1Lu monolayer consisted of uniform, flat, polygonal cells with surface microvilli. At low cell density, foci of dense growth consisting of cells morphologically identical to those of the monolayer were seen. The transformed cells were characterized by morphological heterogeneity; the cells were polygonal, flat, spherical, and spindle-shaped. Microvilli, blebs, and filopodia were seen on the surface. The morphology of the Ki-Mv1Lu cells was independent of cell density. There was, however, nothing in the cell-surface morphology that was absolutely specific for the transformed cells. (10 refs)

78-6894 Comparison of the Genomic Organization of Kirsten and Harvey Sarcoma Viruses. (Eng) Shih, T. Y. (Lab. Tumor Virus Genetics, NCI, Bethesda, MD, 20014); Williams, D. R.; Weeks, M. O.; Maryak, J. M.; Vass, W. C.; Scolnick, E. M. *J Virol* 27(1): 45-55; 1978.

The genomic RNA's of Kirsten murine sarcoma virus (Ki-MuSV), Harvey murine sarcoma virus (Ha-MuSV), and the replication-defective endogenous rat virus were compared, and their structure and function were correlated. Genome organization and sequence homology were studied by fingerprinting large RNase T1-resistant oligonucleotides (ON) and by cross-protecting homologous ON against RNase A and T1 digestion with complementary DNA (cDNA) prepared from each of the other viral RNA's. Ki-MuSV and Ha-MuSV RNA shared a series of rat-derived ON beginning approx 1 kilobase (kb) from the 3' end and extending to within 1.5 kb at the 5' end of Ki-MuSV RNA. The total map distance covered was about 5.5 kb. The eight ON covering the 1.5 kb assigned to the 5' end of Ki-MuSV RNA were not found in Ha-MuSV RNA. However, by protection of Ki-MuSV RNA with cDNA from the rat virus, 5/8 ON could be designated as rat genetic information. Since Ha-MuSV is approx 6.5 kb and Ki-MuSV is 8.0 kb in size, the major difference between them is the 1.5 kb at the 5' end of Ki-MuSV RNA not present in Ha-MuSV RNA. When the viral RNA's were translated in vitro in cell-free systems, Ki-MuSV yielded a major product of 50,000 daltons, Ha-MuSV RNA a major product of 22,000 daltons. These polypeptides are clearly distinguishable from each other upon tryptic mapping and apparently are translated from the full-genome-length viral RNA, suggesting that they are products of the 5' portions of the viral RNA. They may provide the necessary protein markers for identifying in vivo virus-coded proteins. (23 refs)

78-6895 In Vitro Isolation of Stable Rat Sarcoma Viruses. (Eng) Rasheed, S. (Dept. Pathology, Univ. Southern California Sch. Medicine, Los Angeles, CA, 90023); Gardner, M. B.; Huebner, R. J. *Proc Natl Acad Sci USA* 75(6): 2972-2976; 1978.

At low passage, a Sprague-Dawley (SD-1) rat embryo culture released an endogenous ecotropic C-type leukemia virus (SD-RaLV) and, after about 20 further passages, it underwent spontaneous transformation. The SD-RaLV released from the transformed cells did not cause rapid transformation of other rat embryo cells. However, when the transformed cells were cocultivated repeatedly with three different chemically transformed and serially transplanted rat tumor cell lines (4-nitroquinoline oxide-induced sarcoma, dimethylhydrazine-induced carcinoma, and N,N'-2,7-fluorenylene bis-2,2,2-trifluoroacetamide-induced hepatoma), rapidly fibroblast-transforming "sarcoma" viruses (RaSV) were recovered after each attempt. RaSV was not recovered from one of these tumor cell lines before transplantation, and focus-forming virus could not be rescued from these same tumor cells C-type by cocultivation with other cells releasing heterologous C-type viruses. Foci were induced on normal rat kidney (NRK) and several other rat embryo cell strains within 7-15 days, and both productive and nonproductive NRK clones were derived. The productive clones were positive for rat-specific p30 antigen and the RaSVs released were

serially transmitted to other rat embryo cells. RaSV genome was rescued from the nonproductive clones by superinfection with SD-RaLV, wild rat C-type virus, and several heterologous C-type viruses. It is concluded that naturally occurring transformation-specific (*src*) genes are being recovered in vitro in the form of stable sarcoma viruses from cells into which no sarcoma virus had apparently been introduced. The viruses differ from the Kirsten and Harvey strains of murine sarcoma virus in that they apparently contain no murine leukemia virus sequences and are of purely rat origin. (41 refs)

78-6896 Characterisation of a Tumor-Specific Antigen on the Surface of Feline Lymphosarcoma Cells. (Eng) Snyder, H. W. (Lab. Viral Oncology and Veterinary Oncology, Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Hardy, W. D.; Zuckerman, E. E.; Fleissner, E. *Nature* 275(5681): 656-658; 1978.

A 70,000-mol-wt protein that exhibited antigenic determinants in common with the feline oncornavirus-associated cell membrane antigen (FOCMA) was identified on the surface of feline lymphosarcoma (LSA) cells and cultured transformed feline lymphoblastoid cells. Cell surface molecules carrying FOCMA determinants were characterized by enzymatic radioiodination of viable cells, precipitation with typing serum from detergent lysates, and analysis of proteins in immunoprecipitates by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). In LSA cells, viral antibodies in FOCMA typing serum always precipitated a single protein species, regardless of whether the cell showed concomitant expression of feline leukemia virus (FeLV) structural antigens. In contrast, immunoprecipitation of FeLV gp70 molecules with FeLV-neutralizing serum was restricted to FeLV-positive cells. Absorption experiments proved that the 70,000-mol-wt FOCMA molecule in LSA cells was serologically distinguishable from surface antigens related to FeLV structural proteins. FOCMA-carrying FeSV-transformed nonproducer mink cells gave two protein species approx 85,000 and 65,000 mol-wt upon SDS-PAGE after precipitation by FOCMA antiserum. (18 refs)

78-6897 The Retrovirus Particles in Human Myeloma Cells RPMI8226: Morphological, Biochemical, Immunological and Infective Transmission Studies. (Eng) Grofova, M. (Cancer Res. Inst., Slovak Acad. Sciences, 88 32 Bratislava, Czechoslovakia); Popovic, M.; Ogura, H.; Matoska, J.; Lizonova, A.; Nilsson, K.; Kuzela, S. *Neoplasma* 25(4): 423-437; 1978.

The morphological, biochemical, and immunological properties of a retrovirus (RPMI8226V) isolated from RPMI8226 human myeloma cells were investigated, and propagation of the virus in various animal and human cells was studied. Ultrastructural observations revealed that RPMI8226V particles have a C-type morphology. They were classified into

three categories: immature and mature particles indistinguishable from typical C-type mammalian oncornaviruses and typical C-type particles with an intermediate membrane. Budding particles in secondarily infected human neoplastic cells contained doughnut-shaped nucleoids characteristic of C-type particles. ³H-uridine-labeled RPMI8226V particles had a buoyant density of 1.17 g/ml in a sucrose gradient containing high mol wt RNA, and the distribution of viral structural proteins upon sodium dodecyl sulfate-polyacrylamide gel electrophoresis was characteristic of that of oncornaviruses. The mol wt of the internal structural proteins ranged from 10,000 to 30,000 daltons. The virus contained an Mg-dependent reverse transcriptase (RT). The cellular homogenate and ultracentrifugate from RPMI8226 cultures did not react with antibodies against murine leukemia virus, feline leukemia virus, RD114, Mason-Pfizer monkey virus, simian sarcoma virus, and avian leukemia/sarcoma virus. The only reaction occurred with anti-bovine leukemia virus (BLV) antibodies. However, anti-BLV serum inhibiting the RT activity of BLV to 60% did not cross-react with the RT of RPMI8226V. In contrast to BLV concentrates, neither XC or KC cells showed syncytia formation by RPMI8226V. RPMI8226V replication was restricted to human tumor and normal human glialike cells. The data indicate that RPMI8226V may be of bovine origin. (40 refs)

78-6898 Induction of Deoxythymidine Kinase Activity in Several Mammalian Cell Lines after Infection with Six Different Strains of Equine Herpesvirus Type 1 (EHV-1). (Eng) Allen, G. P. (Dept. Microbiology, Univ. Mississippi Medical Center, Jackson, MS, 39216); McGowan, J.; Bryans, J. T.; Randall, C. C.; Gentry, G. A. *Virology* 90(2): 51-359; 1978.

The ability of six strains of equine herpesvirus type 1 (EHV-1) to induce deoxythymidine kinase (dTK) activities in mouse 3T3 cells, horse embryo fibroblasts (KyED), equine transitional cell carcinoma cells (HT), and owl monkey kidney cells (OMK) was studied. Increased deoxythymidine triphosphate (dTTP)-resistant dTK activity was observed after infection of all cell types with any of the six strains of EHV-1. Increased levels of dTTP-resistant dTK were also detected in liver homogenates derived from Syrian hamsters infected with EHV-1. This activity appeared to be virus-coded. EHV-1 appeared to induce a dTK activity in HT and KyED cells that comigrated in polyacrylamide gels with the cytosol dTK of the host cell. Replication of wild-type EHV-1 in 3T3 cells was inhibited by arabinosylthymine (ara-T), but replication of the virus in OMK, HT, and KyED cells was considerably more resistant to inhibition by ara-T. The data indicate that infection of cells with any strain of EHV-1 results in the induction of a new, virus-coded dTK activity. (40 refs)

78-6899 Transformation of Warts to Malignancy in Alimentary Carcinoma in Cattle. (Eng) Jarrett, W.

F. (Dept. Veterinary Pathology, Univ. Glasgow, Bearsden Road, Glasgow G61 1QH, Scotland). *Bull Cancer (Paris)* 65(2): 191-194; 1978.

The naturally occurring high incidence of alimentary carcinoma in beef cattle in the highland areas of Britain offers an excellent opportunity to study the interactions between a papilloma virus and an environmental chemical, possibly derived from bracken fern, that has putative mutagenic and immunosuppressive properties. This development of carcinoma is associated with the presence of bracken fern on high-incidence farms, and many of the carcinomas arise in preexisting virus-induced papillomas. Papillomas are much commoner and occur in greater numbers in animals of all ages on the cancer farms than in cattle on lowland farms. Carcinomas have only been seen in cattle >6 yr old. The virus has been isolated, visualized, and purified; it produces papillomas when inoculated into the alimentary tract and skin. A total of 165 carcinomas were found in 80 cows with squamous carcinoma of the alimentary tract. They were distributed as follows: palate 4%, pharynx 8%, tongue 7%, esophagus and groove 41%, and rumen 30%. Papillomas or warts of the squamous epithelium lining the upper alimentary tract were found in 96% of these cows. There was a range of tumors from benign papillomas to papillomas with areas of malignant transformation to frank ulcerating and invasive carcinoma. (5 refs)

78-6900 Some Physical Properties of a Human Laryngeal Papillomavirus. (Ger) Liebermann, H. (Friedrich-Loeffler-Institut für Tierseuchenforschung, DDR-2201 Insel Riems, über Greifswald, E. Germany); Solisch, P.; Hahnefeld, H. *Arch Geschwulstforsch* 48(6): 552-558; 1978.

Ultracentrifugation of a human laryngeal papillomavirus produced particles sedimenting at 130S and 214S. Electron microscopy revealed the presence of 37-nanometer (nm) and 49-nm particles, with the latter having a buoyant density in CsCl gradients of 1.34 g/ml. This virus resembles a polyoma virus. (19 refs)

78-6901 In Vitro Transcription of Polyoma DNA. (Eng) Monjardino, J. (Div. Cell Studies, Royal Free Hosp. Sch. Medicine, 8 Hunter St., London WC1N 1BP, England); James, A. *Biochem Biophys Res Commun* 84(3): 618-626; 1978.

In vitro transcription of supercoiled DNA from nondefective polyoma virus (PV) by *Escherichia coli* RNA polymerase was successfully carried out, and the resulting complementary RNA (cRNA) was seen to be double stranded to a large extent, longer than the coding strand, and initiated from a limited number of sites. The incubation mixture for RNA synthesis contained 20 µg/ml PV DNA I, 75 µg/ml *E. coli*

RNA polymerase, and nucleotides including ^3H -GTP (0.15 Ci/millimole). Results of chromatographic separation of the double-stranded RNA product on CF11 cellulose and pancreatic RNA digestion, followed by cRNA annealing, proved that the transcription was at least partially symmetric. Electron microscope visualization of the transcription complexes showed long RNA chains up to 3x the length of the template strand. In the presence of rifampicin, transcription was usually initiated from 2-3 initiation sites, proceeding at a rate of 16 nucleotides/sec under suboptimal conditions. When 32 transcription complexes were lined up on linear DNA molecules, it was evident that the RNA could not have originated from a common initiation site. Further, cRNA chains in some complexes differed in length by more than a transcript of the entire DNA template. These findings support the hypothesis that transcription took place from both DNA strands. (16 refs)

78-6902 Expression of Polyoma Virus In Heterokaryons Between Embryonal Carcinoma Cells and Differentiated Cells. (Eng) Boccara, M. (Departement de Biologie Moleculaire, Institut Pasteur, 25 rue du Dr. Roux, 75015 Paris, France); Kelly, F. *Virology* 90(1): 147-150; 1978.

Polyoma virus (PV)-resistant PCC3/A/1 embryonal carcinoma cells were fused with two types of susceptible cells: differentiated teratoma-derived PCD3 cells, which are fully permissive for PV, and BHK21 cells, which can express only the early viral genes. Cells from one parent were infected with 20 plaque forming units of PV, fused 4 hr later with the other ^3H -thymidine labeled parent using β -propiolactone-inactivated Sendai virus, and examined by autoradiography 48 hr post-infection for the presence of V antigen or radioactively labeled nuclei. When infected PCC3/A/1 cells were fused with PCD3 cells, 20% of the heterokaryons were V antigen-positive. Both parent cells were necessary for viral expression, suggesting that the virus entered the PCC3/A/1 cells but was expressed only upon fusion with permissive cells. Fusion of infected PCD3 cells with PCC3/A/1 cells resulted in 60% antigen-positive heterokaryons, a percentage comparable to that of infected PCD3 homokaryons or isolated PCD3 cells. Nonpermissive preinfected BHK21 cells produced 5% antigen-positive heterokaryons when fused with PCC3/A/1 cells, compared with no apparent staining in unfused, infected BHK21 cells or BHK21 homokaryons. These experiments indicate that blocked response to PV in pluripotent teratocarcinoma cells occurs after penetration of the viral particle but before the synthesis of the T antigen, and that susceptibility to infection in the differentiated cells is dominant over resistance in the embryonal carcinoma cells. (10 refs)

78-6903 Isolation of Polyoma Viruses Lacking Endonuclease *Hind*III Sites. (Eng) Anderson, D. M. (Roche Inst. Molecular Biology, Nutley, NJ, 07110); Bendig, M. M.; Folk, W. R. *Virology* 89(2): 637-642; 1978.

Spontaneous mutants of polyoma virus that have specific alterations at one of two endonuclease *Hind*III sites in the polyoma genome (one in the early region, the other in the late region) were isolated and partially characterized. The viruses were divided into two groups on the basis of their genome structure and complementation requirements: Group I viruses did not require complementing genomes in order to form plaques, but Group II viruses did. Infection by group I viruses produced homogeneous populations of DNA, but infection by Group II viruses always produced two DNA species, regardless of the number of attempts to plaque-purify them. In almost every instance, a DNA without one *Hind*III site was accompanied by a DNA having an alteration at the other *Hind*III site. Many of the viruses in both groups synthesized viral DNA at 32 and 39 C, although several Group II mutants did not synthesize form I DNA as well at 39 C as at 32 C, indicating thermolability of early gene products. Several of these mutants will be useful in probing the function of the polyoma genome. Group I mutants may express altered early functions. In addition, most of these viruses are capable of proceeding through a normal lytic infection and may therefore, be more useful as cloning vehicles than wild-type polyoma DNA. Group II mutants, altered in the early or late region, are difficult to grow in large amounts, and the individual species of the pairs of mutants must be separated before the extent and effects of the alterations can be defined. (28 refs)

78-6904 Polyoma Virus High Molecular Weight Nuclear RNA Codes for Capsid Protein VP2 In Vitro. (Eng) Kamen, R. I. (Transcription Lab., Imperial Cancer Res. Fund, London WC2A 3PX, England); Wheeler, T. Smith, A. E. *Virology* 89(2): 461-474; 1978.

An attempt was made to determine whether polyoma virus (Py) nuclear RNA molecules, which contain messenger RNA-like sequences but which are assumed not yet extensively spliced, can direct the synthesis of viral proteins in vitro. RNA extracted from the nuclei of 3T6 mouse cells late after infection by Py was translated in vitro to give one predominant polypeptide. This polypeptide was identified as virion protein VP2 by polyacrylamide gel electrophoresis, specific immunoprecipitation, and tryptic peptide analysis. The active nuclear RNA species sedimented between 15S and 50S on sucrose gradients after denaturation. The bulk of the active viral nuclear RNA did not bind to oligo(dT)-cellulose and synthesis of Py VP2 was inhibited by m⁷Gp under conditions in which translation of Py-complementary RNA and EMC RNA are not inhibited. It is concluded that VP2 coding activity is not only a function of mature cytoplasmic 19S RNA, but also of Py nuclear RNA, which is probably capped but not polyadenylated. Determination of the size of the active RNA molecules indicated that all size classes of nuclear RNA have VP2 coding activity and that this activity is unlikely to result from contamination of the high mol wt RNA with aggregated 19S RNA. (49 refs)

6905 **Evaluation of Normal and Neoplastic Human Tissue for BK Virus.** (Eng) Israel, M. A. (Lab. Virology, Natl. Inst. Allergy and Infectious Diseases, Bethesda, MD, 20014); Martin, M. A.; Takemoto, K. K.; Wley, P. M.; Aaronson, S. A.; Solomon, D.; Khoury, G. *Virology* 90(2): 187-196; 1978.

The role of the human papovavirus BK virus (BKV) in human neoplasia was evaluated using nucleic acid hybridization and indirect immunofluorescence techniques. None of 12 normal organ preparations, 5 human tumors, or 16 human tumor cell lines was positive for sequences homologous to BKV DNA. Detectable levels of antibody specific for BKV (tumor)-antigen were not found in any of 55 human tumor cell lines or in serum samples from any of 67 cancer patients. On the basis of these findings, it is unlikely that BKV is commonly associated with human cancer. (37 refs)

6906 **Cell Surface Morphology and Adhesive Properties of Normal and Virally Transformed Cells Treated with Tunicamycin, an Inhibitor of Protein Glycosylation.** (Eng) Duksin, D. (Dept. Biochemistry, Biological Structure and Medicine, Univ. Washington, Seattle, WA, 98195); Holbrook, K.; Williams, K.; Bornstein, P. *Exp Cell Res* 116(1): 153-165; 1978.

The effects of tunicamycin (0.1 or 0.2 $\mu\text{g/ml}$) on the biochemical and functional surface properties of mouse 3T3 cells, simian virus 40-transformed 3T3 cells, and polyoma virus-transformed 3T3 cells were studied. The cells of all lines became rounded, leaving long attenuated cell processes attached to the substrate. In addition, the nontransformed 3T3 cells showed extensive ruffling of the cell surface and dilation and loss of ribosomes from the rough endoplasmic reticulum. There was a marked reduction in all cell lines in a cell surface protein that was similar or identical to fibronectin. Tunicamycin treatment also resulted in a reduction in the length of cell adhesion to the substrate, a decrease in plating efficiency, and a marked inhibition in the cell spreading rate on the surface of the tissue culture plate; the latter two changes were more pronounced in the transformed cells than the nontransformed cells. The data suggest that cell surface glycoproteins, including fibronectin, play a role in determining the surface morphology and adhesive properties of cells. (8 refs)

6907 **Mobility of Histones on the Chromosomes of Simian Virus 40.** (Eng) Beard, P. (Swiss Inst. Experimental Cancer Res., CH-1066 Epalinges, Switzerland). *Cell* 15(3): 955-967; 1978.

near simian virus 40 (SV40) chromosomes were prepared by *EcoRI* nuclease cleavage of the circular SV40 chromosomes released from virions with dithiothreitol at pH 9.8. Chromatin-DNA hybrids were constructed with segments of

^3H -labeled, naked SV40 DNA covalently joined via the *EcoRI*-generated cohesive ends to segments of linear SV40 chromosome. Upon incubation of chromatin-DNA hybrids at 37 C and moderate ionic strength, histones migrated onto the labeled DNA while retaining the nucleosome structure. This was shown by (1) the pattern of micrococcal nuclease digestion of labeled DNA, (2), by nitrocellulose filter binding of labeled DNA after redigestion of the chromatin-DNA hybrids with *EcoRI*, and (3), by examination of chromatin-DNA hybrids in the electron microscope. Migration was slow, being apparent after several hours. Parallel experiments in which naked DNA and chromosomes were mixed without joining showed no transfer of nucleosomal histone between DNA molecules. The kinetics of *EcoRI* cleavage of the DNA in virion-derived SV40 chromosomes are also consistent with the notion that nucleosomal histones, in the absence of other proteins, can move on DNA. (59 refs)

78-6908 **Influence of Agents Reducing Cell-Substrate Adhesion on Protein and Glycoprotein Patterns of Transformed Mouse Fibroblasts in Culture.** (Rus) Liubimov, A. V. (Lab. Mechanisms Carcinogenesis, Oncology Res. Center, Acad. Medical Sciences, USSR, Moscow, USSR). *Tsitologiya* 20(10): 1179-1185; 1978.

The effect of substances that reduce cell-substrate adhesion (trypsin, EDTA, and urea) on the protein and glycoprotein patterns of methylcholanthrene-transformed mouse fibroblasts (L cells), the serum-free derivative of L cells (LSF), and mouse fibroblasts transformed by simian virus 40 (M-22 cells) was studied by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Incubation with trypsin (10-1,000 $\mu\text{g/ml}$ for 10 min at 37 C) caused no marked changes in the protein and glycoprotein patterns of L cells, but urea (1 mole, 2 hr) and EDTA (200 $\mu\text{g/ml}$, 2 hr) removed a protein with a mol wt of 240,000. In LSF, trypsin (10 $\mu\text{g/ml}$, 10 min, 37 C) reduced the level of a 271,000-dalton protein considerably. In M-22 cells, trypsin (1-1,000 $\mu\text{g/ml}$) caused a protein with a mol wt of 270,000 to disappear. Urea extracted a 270,000-dalton protein nearly completely and a 139,000-dalton protein considerably. EDTA had no marked effect on protein and glycoprotein patterns, but it removed a protein with a mol wt of 270,000. The findings suggest that deficient contact reactions of transformed cells may be due to changes in the pattern and properties of cell components that are sensitive to alterations of adhesion. (7 refs)

78-6909 **Augmented Immunogenicity of Tumor Cell Membranes Produced by Infection with Influenza Virus as Compared to Moloney Sarcoma Virus.** (Eng) Takeichi, N. (Lab. Pathology, Cancer Inst., Hokkaido Univ. Sch. Medicine, Sapporo, Japan); Austin, F. C.; Oikawa, T.; Boone, C. W. *Cancer Res* 38(12): 4580-4584; 1978.

The immunogenicity of tumor cell membranes from Moloney

murine sarcoma virus (MSV)- and influenza virus-infected cells was tested. The tumor-associated transplantation antigens (TATA) of crude membrane extracts from simian virus 40 (SV40)-transformed BALB/3T3 tumor cells lytically infected with influenza virus were much more immunogenic than the TATA of extracts from uninfected cells, as measured by their ability to induce heightened resistance to tumor graft challenge or by heightened lymphocyte-mediated cytotoxicity against tumor cells in vitro. When intact tumor cells (as opposed to membrane extracts) were productively infected with MSV, they became so immunogenic they would only grow in x-irradiated syngeneic animals. However, crude membrane extracts from the MSV-infected tumor cells showed no increase in TATA activity analogous to that seen after infection with influenza virus. Thus, influenza virus augmentation of tumor membrane TATA may be different from the oncornavirus augmentation of intact tumor cell TATA reported previously. It appears that MSV and, possibly, other oncornaviruses cannot be used to augment the TATA activity of tumor cell membranes in the same way that other surface-budding viruses can. (25 refs)

- 78-6910 Simian Virus 40 A Gene Function: DNA Content Analysis of Chinese Hamster Cells Transformed by an Early Temperature-sensitive Virus Mutant.** (Eng) Robinson, C. C. (Dept. Pathology, Univ. Colorado Medical Center, 4200 E. Ninth Ave., Denver, CO, 80262); Lehman, J. M. *Proc Natl Acad Sci USA* 75(9): 4389-4393; 1978.

Replication of Chinese hamster embryo cell lines (ChH WT-2, ChH A58, and ChH A58-1) transformed by wild-type simian virus 40 (WT SV40) and the SV40 temperature-sensitive A gene mutant (*tsA58*) was examined by flow microfluorometry and autoradiography of ³H-thymidine-labeled cells to determine whether DNA synthesis in transformed cells is initiated by the A gene. At the permissive temperature (37 C), the *tsA58*-transformed cells resembled the WT transformants in cell morphology, colony formation, high saturation density, and rapid replication. They seemed to revert to a normal phenotype at the nonpermissive temperature (40.5 C). However, flow microfluorometric analysis of the cell DNA content at 40.5 C showed that under conditions in which normal embryo cells enter a resting state, the *tsA58* transformants continued to replicate. The low density of the cultures was maintained by a balance among cell proliferation, cell death, and sloughing of dead cells into the culture medium, and it did not represent a return to normal growth. Thus, loss of the phenotypic characteristics of transformation cannot be explained by arrest of the cells in the G₁ (G₀) phase of the cell cycle. It is suggested that the initiator hypothesis for the function of the A gene product in the maintenance of transformation must be reevaluated. (34 refs)

- 78-6911 Suppressive Effects on Simian Virus 40-induced Oncogenesis of Several Immunosuppressive**

Agents and Hormonal Modifications Applied During the Latent Period. (Eng) Ohtaki, S. (Dept. Pathology, Nalt. Inst. Health, Tokyo Gakuen 4-7-1, Musashi-Murayama, Tokyo 190-12, Japan). *Cancer Res* 38(12): 4698-4710; 1978.

Host factors affecting tumorigenesis at different stages, especially before and after the appearance of tumor cells, were studied in two experiments: (1) simian virus 40 (SV40)-induced oncogenesis in Syrian golden hamsters, to investigate the latent period; and (2) transplantation of homologous, primary SV40 tumor cells into hamster, to investigate a late stage of tumorigenesis. The mean latent period for SV40 tumor induction was about 6 wk shorter in female hamsters than in males ($p < 0.02$). The mean latent period in castrated females was 12.6 wk longer than that in intact females. In contrast, tumor development was not significantly affected by orchietomy. Estrone treatment (0.2-0.3 mg, sc) at birth delayed tumor development significantly in both sexes. Tumor development was significantly accelerated by testosterone treatment (0.5 mg, sc) of males at age 15 wk, the mean latency period being decreased by 11 wk. By each of these hormonal modifications, growth of transplanted SV40 tumors was influenced in a different way from that of tumor induction. Immunosuppressive treatments, such as thymectomy or administration of antilymphocyte sera, cyclophosphamide, or cortisone acetate, delayed and decreased tumor development when applied in the latent period, and the degree or pattern of this effect varied from one procedure to another, depending on the sex and age of the animals. In contrast, tumor growth was markedly accelerated in thymectomized or cortisone-treated hosts of either sex. The different and almost reverse effect of the same procedures in the two phases of tumorigenesis indicates that two distinct mechanisms may be operating during these phases. The different effect may be due to the absence of "mature" neoplastic cells in the latent period, except for a few weeks before the appearance of a palpable tumor. These results suggest that the latent period may be needed for completion of SV40-induced neoplastic conversion of cells. (47 refs)

- 78-6912 Simian Virus 40 Production After Viral Uncoating in the CV-1 Cell Nucleus.** (Eng) Gershey, L. (Rockefeller Univ., New York, NY, 10021); Diacumakos, E. G. *J Virol* 28(1): 415-416; 1978.

A normal yield of infectious simian virus 40 was produced by a culture of African green monkey kidney CV-1 cells after a few cells were inoculated with an av of one particle of plaque-purified virus per nucleus by microinjection. (7 refs)

- 78-6913 Simian Virus 40 DNA Replication in Isolated Replicating Viral Chromosomes.** (Eng) Su, R. (Dept. Microbiology, Univ. Kansas, Lawrence, KS, 66044); DePamphilis, M. L. *J Virol* 28(1): 53-65; 1978.

Three soluble subnuclear systems (nuclear extracts, nucle-

protein complexes, and replicating chromosomes prepared from simian virus 40-infected CV-1 cells) were studied to determine the minimum conditions necessary for DNA replication in vitro. DNA synthesis was measured with and without CV-1 cell cytosol in the reaction mixture. With nuclear extracts, cytosol stimulated total DNA synthesis (2-3 times), elongation of nascent DNA chains, maturation and joining of Okazaki pieces (transient intermediates), and the conversion of replicating viral DNA into covalently closed, superhelical DNA. The response of nucleoprotein complexes was similar but often 10%-30% lower. In contrast, isolated replicating chromosomes in the presence of cytosol appeared only to complete and join Okazaki pieces on the DNA templates. They failed to extend nascent DNA chains prelabeled in vivo with ^3H -thymidine or labeled in vitro with α - ^{32}P -deoxycytosine triphosphates (α - ^{32}P -dCTP). Consequently, the conversion of viral replicating DNA into mature DNA was given to eight times less than that observed in nuclear extracts. In the absence of cytosol, Okazaki pieces were still labeled with α - ^{32}P -dCTP in vitro, but completely failed to join the growing nascent DNA chains, demonstrating the requirement of cytosol factors at this stage in DNA elongation. Treatment of nucleoprotein complexes with 0.6-2.0 M NaCl resulted in complete loss of endogenous DNA synthesis and lack of response to cytosol. These results demonstrate that continuation of endogenous DNA replication in vitro requires both the soluble cytosol fraction and a complex nucleoprotein template whose ability to continue DNA synthesis depends on its concentration and ionic environment during its preparation. (31 refs)

78-6914 Non-Coordinate Expression of SV40 Transformation in Cell Hybrids (Meeting Abstract). (Eng) Howell, N. (Sidney Farber Cancer Inst., Boston, MA, 02115); Sager, R. *J Cell Biol* 79(2, part 2): 396a; 1978. (no refs)

78-6915 SV40-transformed Hamster Cells Resistant to 100-250 $\mu\text{g}/\text{ml}$ of Ethidium Bromide. (Eng) Volf, G. (Max-Planck-Institut für Virusforschung, 7400 Tübingen, W. Germany); Tejmar, L.; Borell, S.; Klietmann, W. *Cell Sci* 33: 157-169; 1978.

Simian virus 40 (SV40)-transformed hamster cells selected for resistance to ethidium bromide (EB, 100-250 $\mu\text{g}/\text{ml}$) were characterized. EB resistance was maintained during propagation in EB-containing medium and thus seemed to be genetically determined. The EB-resistant cells were considerably less tumorigenic in Syrian golden hamsters than wild-type cells. They had a normal ultrastructure, except for a highly condensed conformation of the mitochondrial cristae. Unlike wild-type cells, the resistant cells showed no significant inhibition of mitochondrial DNA synthesis in the presence of 16 $\mu\text{g}/\text{ml}$ EB. The polypeptides of the mitochondrial fractions of wild-type and resistant cells were significantly different:

polypeptide 14, which was present in wild-type cells, was absent or greatly reduced in the mitochondria of the resistant cells, and the relative proportions of polypeptides 1:2 and 7:8:9 were different. The mechanism of EB resistance in mammalian cells has not been determined. (27 refs)

78-6916 Simian Virus 40 T Antigen Binds to Host Cell Chromosomes. (Eng) D'Alisa, R. M. (Rockefeller Univ., New York, NY, 10021); Gershey, E. L. *Nature* 274(5667): 164-166; 1978.

The binding of simian virus 40 (SV40) T antigen to mammalian chromosomes was studied using a fibroblastic cell strain derived from a male Indian deer, *Muntiacus muntjak*, an animal with only seven chromosomes, which can be unequivocally identified. Cultured muntjac cells were infected with SV40 strain VA 4554 at multiplicities between 200 and 500 plaque-forming units/cell. Seventy-two hours after infection, the chromosomes were swollen in 0.075 M KCl for 12 min, fixed in 3:1 methanol:acetic acid, air-dried, and stained for T antigen by indirect immunofluorescence. The preparations were photographed on an inverted microscope. Alternatively, cells were grown in glass slides and arrested with colcemid before KCl treatment and fixation. No differences in chromosome morphology were detected with the two methods. T antigen was associated with all chromosomes, including the small Y, of infected cells. The binding appeared to be specific for the fluorescent chromosomes of infected cells, since uninfected cells did not contain fluorescent chromosomes and (2) presumably uninfected cells within the SV40-treated population also did not fluoresce. The similarity of the T-antigen binding patterns to Q- and G-banding patterns suggests that the antigen binds to initiation sites for host cell DNA replication. (27 refs)

78-6917 Definition of the Boundaries of the Origin of DNA Replication in Simian Virus 40. (Eng) Subramanian, K. N. (Dept. Microbiology, Univ. Illinois Medical Center, Chicago, IL, 60612); Shenk, T. *Nucleic Acids Res* 5(10): 3635-3642; 1978.

Using DNA sequencing techniques, the exact location of the deletion in *dl* 892, a viable deletion mutant of simian virus 40 (SV40) reported to map very near the unique replication origin of SV40, was determined. Autoradiographic studies showed that *dl* 892 lacks a segment of 19 nucleotides (located between positions 187 and 205 in the *Eco* RII-G sequence, both inclusive) present in wild-type SV40 DNA. In view of the fact that segments of DNA occurring distal position 186 could be deleted without affecting the replication of the viral DNA (and the ability of the virus to grow without a helper), one of the boundaries for the replication origin located distal to the *Hind* (11 + 111) -A-C junction could be placed at position 186. Previous studies indicate that sequences to the left of map position 0.660 are not needed for initiation of

SV40 DNA replication. Therefore, the other boundary of the replication origin has been placed at map position 0.660 corresponding to the *Hae* III restriction site at nucleotide 102 in the *Eco* RII-G sequence. Thus, according to the present estimate, the SV40 replication origin consists of a max of 85 nucleotides located between positions 102 and 185 in the sequence of the *Eco* RII-G fragment derived previously. This sequence includes the 27-nucleotide perfect palindrome, the 17-nucleotide "true" palindrome (containing a single imperfection in its sequence), the 17-nucleotide A-T-rich sequence, and the 15-nucleotide imperfect palindrome, all shown to be present in the previously derived *Eco* RII-G sequence. These and previous results have enabled the boundaries of the replication origin to be narrowed down to 85 nucleotides within the sequence of SV40. (10 refs)

78-6918 The Detection and Characterization of Multiple Forms of SV40 Nucleoprotein Complexes.

(Eng) Garber, E. A. (Dept. Biochemical Sciences, Princeton Univ., Princeton, NJ, 08540); Seidman, M. M.; Levine, A. J. *Virology* 90(2): 305-316; 1978.

Using a newly developed method for extracting multiple forms of simian virus 40 (SV40) nucleoprotein complexes from the nuclei of virus-infected cells, the production and maturation of the SV40 chromosome in vivo were studied. Four different forms of SV40 DNA-protein complexes were identified: a heterogeneous (75S to 200S) replicative intermediate; a 75S minichromosome; a 200S previrion particle; and a mature virion (250S). After replicating in the heterogeneous 75S to 200S complexes, the newly replicating chromosome appeared to be converted to the 75S form, which matured into the 200S previrion structure. By 45 to 60 min, mature virions were observed. The sequence of appearance of these complexes was suggestive of a precursor-product relationship. The data demonstrated that 90% of the viral DNA was encapsulated by 4 hr after its synthesis. (25 refs)

78-6919 Identification of Oncogenic Chromosomes in Transformed Human Cells. (Eng) Croce, C. M.

(Wistar Inst. Anatomy and Biology, 36th and Spruce Sts., Philadelphia, PA, 19104). In: *Neoplastic Transformation: Mechanisms and Consequences. Proceedings of a meeting held by the Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences in Berlin, May 9-13, 1977.* Dahlem Workshop on Neoplastic Transformation: Mechanisms and Consequences (Berlin, W. Germany): 314 pp.; 69-85; 1977.

To determine whether viral DNA sequences are associated with specific chromosomes in transformed cells and whether integrated viral DNA sequences are responsible for expression of the transformed phenotype and malignancy, a large number of somatic cell hybrids between rodent cells and dif-

ferent simian virus 40 (SV40)-transformed human cells were produced and investigated. Three independent SV40 T-antigen-positive hybrid clones between CT1D mouse cells and LN-SV SV40-transformed human cells were subcloned, and the subclones were analyzed for the expression of SV40 tumor antigen and the presence or absence of human chromosomes 7. Somatic cell hybrids between thymidine kinase-deficient IT-22 mouse or A3 hamster cells and GN54VA transformed human fibroblasts were also studied. The results showed that the SV40 genome can be integrated in at least two different human chromosomes: 7 and 17. These and further studies demonstrated that it was possible to map the integration sites of oncogenic viruses in virus-transformed human cells and to identify the human chromosomes responsible for the expression of malignancy by somatic cell hybridization. Since these chromosomes can be isolated on a normal inbred mouse strain background, it should also be possible to elicit humoral and cellular immunoresponses to human tumor-associated antigens using hybrid cells. (26 refs)

78-6920 The Sequences between 0.59 and 0.54 Map Units on SV40 DNA Code for the Unique Function of Small t Antigen. (Eng) Paucha, E. (Translation Lab Imperial Cancer Res. Fund, London WC2A 3PX, England); Smith, A. E. *Cell* 15(3): 1011-1020; 1978.

The hypothesis that the carboxy terminal portion of simian virus 40 (SV40) small t antigen is encoded by a sequence of nucleotides from the region between 0.59-0.54 map units on SV40 DNA was tested by characterizing the putative shortened forms or fragments of small t produced by SV40 mutants (dl 884, dl 885, dl 890) with deletions in this genomic region. Attempts to isolate the putative fragments of small t from mutant-infected cells, or from cell-free systems primed with messenger RNA (mRNA) from mutant-infected cells, resulted in only low yields of the fragments. Experiments using purified SV40 mRNA in low background cell-free systems, in which large T and small t could be detected without immunoprecipitation, suggested that these low yields were due to the reduced amounts of mRNA coding for the shortened forms of small t present in the mutant-infected cells. Larger amounts of putative fragments of small t were produced by translation of deletion mutant complementary RNA (synthesized in vitro using purified deletion mutant DNA and *Escherichia coli* RNA polymerase). Fingerprint analysis of the proteins produced showed that they contained most, if not all, of the methionine peptides common to small t and large T. Furthermore, the fragments of small t produced in response to dl 884 and dl 890 lacked two methionine peptides that were present in small t but not in large T. The results provide direct evidence that the region between 0.59-0.54 map units on SV40 DNA codes for polypeptide sequences that are unique to small t, and they establish that the nucleotide sequences from this region are both a coding sequence (for small t) and an intervening sequence (for large T). (38 refs)

78-6921 Selective Degradation of Newly Synthesized Nonmessenger Simian Virus 40 Transcripts.

(Eng) Chiu, N. H. (Lab. Biology Viruses, Natl. Inst. Allergy and Infectious Diseases, NIH, Bethesda, MD, 20014); Radonovich, M. F.; Thoren, M. M.; Salzman, N. P. *J Virol* 18(2): 590-599; 1978.

Under pulse-chase conditions with glucosamine, the processing, transport, and degradation of a relatively homogeneous population of nuclear viral RNA molecules (nvRNA) labeled during a brief pulse were examined 45 hr postinfection. By pretreating the simian virus 40-infected BSC-1 cells with glucosamine, ³H-uridine labeling of cellular and viral RNA could be halted immediately by the addition of cold uridine. Both messenger and nonmessenger regions of the late strand were transcribed during the 5-min labeling period and, after various chase periods, nvRNA's that sedimented at 16S, 17.5S, and 19S were observed. The fraction of the late strand that did not give rise to stable late messenger RNA (50% of nvRNA) was degraded rapidly, with a half-life of 8 min. A 40% fraction was transported to the cytoplasm, and the remaining 10% was sedimented from the nucleus at 2S-4S. These 2S-4S nvRNA's had a half-life of 3 hr, and hybridization studies suggested that they were partly coded for by the late-strand nonmessenger region and were derived from the initial nuclear processing step. Another part was coded for by the late-strand messenger region and may result from cleavage of 19S RNA into 17.5 and 16 S RNA's. Transport of nvRNA into the cytoplasm was detected after a 5-min pulse and a 7-min chase. Accumulation in the cytoplasm was maximal after a 30-min to 1-hr chase. At least two viral cytoplasmic species were noted: 19S (half-life 2 hr), which is likely transported directly from the nucleus, and 16S (half-life 5 hr), which may be formed by cleavage of the larger species in the nucleus and/or cytoplasm. (31 refs)

78-6922 Does Simian Virus 40 DNA Integrate into Cellular DNA During Productive Infection? (Eng)

Rigby, P. W. (Dept. Biochemistry, Imperial Coll. Science and Technology, London SW7 2AZ, England); Berg, P. *J Virol* 28(2): 475-489; 1978.

The possibility that simian virus 40 (SV40) DNA becomes covalently integrated into the chromosomal DNA of productively infected cells was investigated by isolating and purifying SV40 DNA late after infection of African green monkey kidney cells. This fraction contained large amounts of SV40 DNA sequences, but their behavior in electron microscope and restriction endonuclease mapping experiments indicated that most, if not all, of the viral DNA was not covalently joined to cellular DNA. Instead, it was organized into large, tandem, head-to-tail polymeric structures of wholly viral origin. The results showed that heteroduplexes between high-mol-wt SV40 DNA and a mutant SV40 DNA probe were not formed unless the former was digested with *EcoRI* or *HpaII*, enzymes that cleave SV40 DNA once. The electrophoretic mobility of the high-mol-wt SV40 DNA was not

perceptibly altered by digestion with several restriction endonucleases that cleave cellular DNA but not SV40 DNA. However, digestion with *EcoRI*, *HpaII*, *KpnI*, or *TaqI* converted all the SV40 DNA into unit-length, linear molecules. The high-mol-wt SV40 DNA was highly infectious in the standard plaque assay, and this infectivity was increased by digestion with *EcoRI* or *HpaII*. Because the polymers appeared upon extensive replication of the viral DNA and homopolymers were produced after coinfections with two physically distinguishable SV40 genomes, it is concluded that they are synthesized by a replicational mechanism. Although the data cannot exclude a low level of SV40 DNA integration during productive infection, sedimentation criteria used previously to demonstrate integration are misleading because novel forms of viral DNA, which copurify with cellular DNA, mimic integrated genomes. (46 refs)

78-6923 Presence of Histone H1 in the Chromosome of SV40 Virions Isolated from the Nuclei of Infected Cells. (Rus) Nedospasov, S. A. (Inst. Molecular Biology, Moscow, USSR); Bakaev, V. V.; Georgiev, G. P. *Dokl Akad Nauk SSSR* 241(2): 485-487; 1978.

The isolation of virions of simian virus 40 (SV40) from infected CV-1 monkey cells is described. Cells were incubated with ³H-thymidine (2 µCi/ml) and ¹⁴C-amino acids (5 µCi/ml) for 72 hr, washed, and centrifuged; the supernatant was then centrifuged in a sucrose gradient. The DNA radioactivity curve contained two distinct peaks, one 60S and the other at 200S. The peaks of protein radioactivity coincided with the peaks of DNA radioactivity. Electron microscope analysis of material from each peak showed that the light peak contained particles with a diameter (approx 250 Å) that corresponded to that of the native compact form of free SV40 minichromosomes. The heavy peak contained typical mature SV40 virions (420-450 Å in diameter) with clearly visible capsomers. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis detected all five histones (including histone H1) in both the free minichromosomes and the virion particles. The ratio of histone H1 to the other histones was stoichiometric (1 molecule of H1/nucleosome). These findings indicate that the mature virus particle contains a chromosome that is extremely similar to the chromosome of the eukaryotes. (13 refs)

78-6924 Expression of Simian Virus 40 Early Genes in Transformed Rat Cells is Correlated with Maintenance of the Transformed Phenotype. (Eng) Gaudray, P. (Centre de Biochimie, Université de Nice, Parc Valrose, 06034 Nice, France); Rassoulzadegan, M.; Cuzin, F. *Proc Natl Acad Sci USA* 75(10): 4987-4991; 1978.

Experiments that show a clear temperature-dependence of the synthesis of all the immunoprecipitable tumor (T) antigenic polypeptides in the temperature-sensitive *A-N* (*tsA-N*)

simian virus 40 (SV40) transformants, but not in *tsA-A* SV40 transformants, are presented. Early viral polypeptides synthesized in SV-transformed FR 3T3 rat fibroblasts were identified by immunoprecipitation using anti-T antigen immune serum. Four polypeptide classes were identified, none of which was detectable in extracts of nontransformed cells or precipitated from transformed cell extracts by nonimmune serum. Their apparent mol wts were 92,000, 63,000, 56,000, and 19,000 daltons. A similar pattern was observed in extracts from lytically infected cells, but there was a reproducible quantitative difference between the relative amounts of immunoprecipitable polypeptide species in lytic and transformed cell extracts: the middle T 56,000- and 63,000-dalton species amounted to 5%-8% of the total T antigen-associated radioactivity in lytic cell extracts, compared with 20%-30% in transformed cells. It appears unlikely that these components result from random proteolysis or any other chemical modification in extracts. Previously reported evidence suggests that these middle T polypeptides may represent virus-coded polypeptides, produced by translation of independent spliced messenger RNA's or by post-translational processing of a larger viral polypeptide (large T); alternatively, they may be cellular gene products in a specific and stoichiometric association with viral proteins. In ³⁵S-methionine-labeled *tsA30-A* transformants, which maintain the transformed phenotype at high temperature, only minor quantitative variations in the pattern of T antigen synthesis were observed when the cultures were shifted from 33 C to 40.5 C. In contrast, the labeling rate of the four virus-specific polypeptides decreased by 90% or more at the high temperature in the *tsA-N* transformants. In all cases, a coordinated variation of the radioactivity associated with the different polypeptide classes was observed. These results suggest that the synthesis and/or processing of early viral proteins is controlled by different mechanisms in various types of SV40 transformants, and that it may be under the positive control a virus-coded protein in transformed N cells. (32 refs)

78-6925 Neoplastic Transformation of Permissive Monkey Cells by Subgenomic Fragments of SV40 DNA and Rescue of SV40 Virions from Transformed Cells by Cell Fusion. (Eng) Moyer, R. C. (Thorman Cancer Res. Lab., Trinity Univ., San Antonio, TX, 78284); Moyer, M. P.; Gerodetti, M. H. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2 Part 1), 2404 pp.; 1443-1464; 1978.

The *EcoRI*/*Hap A* fragment of simian virus 40 (SV40) DNA, which comprises 74% of the viral genome including the early region, and the *EcoRI*/*Hap B* fragment, which comprises 26% of the genome including most of the late regions, were tested for their ability to transform permissive TC7/CV-1 monkey cells. The two fragments were separated by electro-

phoresis and added to cell cultures in the presence of diethylaminoethyl-dextran. Results of indirect immunofluorescence tests revealed that either fragment transformed the cell without evidence of cytopathology or V antigen. Cells transformed by the 74% fragment displayed SV40 T antigen and were tumorigenic in newborn hamsters, but cells transformed by the 26% fragment appeared to express a new antigen and were not tumorigenic. Fusion of cells transformed by both types of fragments resulted in SV40 cytopathology and antigen production, which indicated that the two defective provirus complements were rescued to produce infectious virions. An "alien DNA" hypothesis is proposed to account for virus-induced and other forms of neoplasia. (61 refs)

78-6926 Growth Control in Simian Virus 40-transformed Rat Cells: Temperature-independent Expression of the Transformed Phenotype in *tsA* Transformants Derived by Agar Selection. (Eng) Rassoulzadegan, M. (Centre de Biochimie, Universite de Nice, Parc Valrose, 06034 Nice, France); Perbal, B.; Cuzin, F. *J Virol* 28(1): 1-5; 1978.

Fischer rat fibroblasts (FR 3T3) were infected with a temperature-sensitive simian virus 40 (SV40) mutant (*tsA30*) using transformed lines isolated from foci (*tsA30-N*) or from colonies grown in soft agar (*tsA30-A*). All (4/4) of the *tsA-A* transformants expressed the transformed phenotype at 40.5 C, but 2/3 *tsA-N* lines were temperature dependent for all the transformation characteristics studied. The pattern of temperature-dependent vs temperature-independent maintenance in these transformants thus appears similar to that found previously for transformed lines isolated after infection with polyoma virus *tsA* mutants. Temperature-shift experiments fused to CV-1 monkey cells indicated that all the *tsA-A* transformants carry mutant endogenous viral genomes. SV40 tumor antigen could be detected by immunofluorescence in all of the transformants at 33 C. Two distinct patterns of immunofluorescent staining were seen: all transformants of *tsA-A* showed homogeneous nuclear staining, but all the *tsA-N* showed a heterogeneous pattern. These results suggest that distinct modes of regulation of tumor antigen synthesis might be associated with the A and N mechanisms of transformation. (18 refs)

78-6927 Mapping of Early SV40-Specific Functions by Microinjection of Different Early Viral DNA Fragments. (Eng) Mueller, C. (Institut für Molekularbiologie und Biochemie, Freien Universität Berlin, W. Berlin 33, Arnimallee 22, Germany); Graessmann, A.; Graessmann, M. *Cell* 15(2): 579-585; 1978.

The ability of simian virus 40 (SV40) DNA fragments containing 60%-100% of the early genome region to induce early virus-specific functions in monkey and mouse kidney cells was studied. The following early virus-specific events were

ected 24 hr after microinjection of SV40 DNA I or its in vitro transcript, cRNA complementary to the early viral DNA strand: intranuclear tumor (T) antigen and U antigen accumulation; stimulation of cellular DNA synthesis; initiation of late SV40 gene expression; and induction of adenovirus 2 fiber protein synthesis. Proteins with T antigenicity were efficiently synthesized in cells after microinjection of any of the four SV40 DNA fragments sharing a common end of about 0.73 map unit and containing 58%-100% of the early genome. Cells injected with fragment Bgl I/Bam HI-A, which contains 100% of the early genome, were T-antigen-negative. Only the full-length early fragment Hpa II/Bam HI-A induced the remaining early SV40-specific functions. The data indicate that only an intact early viral protein can stimulate both viral and cellular DNA synthesis, DNA sequences at map position 0.67 being required for intracellular early viral transcription. (41 refs)

78-6928 Transforming Viruses Directly Reduce the Cellular Growth Requirement for a Platelet-Derived Growth Factor. (Eng) Scher, C. D. (Dept. Pediatrics, Sidney Farber Cancer Inst., Boston, MA, 02115); Pledger, W. J.; Martin, P.; Antoniades, H.; Stiles, C. D. *J Cell Physiol* 137(3, part 1): 371-380; 1978.

The ability of early passage mouse embryo fibroblasts, mouse 3T3 cells, early passage diploid human fibroblasts, and 3T3 cells and human fibroblasts transformed by simian virus 40 (SV40) or retroviruses to grow in medium supplemented with defibrinogenated platelet-poor plasma (PPP) was studied. The untransformed cells grew to higher cell densities in medium supplemented with serum than in medium supplemented with PPP. However, the human fibroblasts displayed this differential growth response only in the presence of hypophysiologic concentrations of calcium. When heated (100 C) extracts of human platelets were added to PPP-supplemented medium, growth of human and mouse embryo fibroblasts was the same as that in serum-supplemented medium. Fibroblasts transformed with SV40 or retroviruses grew to the same densities in medium supplemented with serum of PPP. Infection of BALB/c-3T3 cells with SV40 rapidly induced them to grow in PPP medium, whereas 3T3 cells infected but not transformed by retroviruses did not replicate in PPP medium. Cells that did not proliferate well in PPP medium did not form tumors when inoculated into athymic nude mice, whereas many of the lines that grew well in PPP medium were tumorigenic in nude mice.

78-6929 Another Chromosomal Assignment for a Simian Virus 40 Integration Site in Human Cells. (Eng) Kucherlapati, R. (Dept. Biochemical Sciences, Princeton Univ., Princeton, NJ, 08540); Hwang, S. P.; Shimizu, N.; McDougall, J. K.; Botchan, M. R. *Proc Natl Acad Sci USA* 75(9): 4460-4464; 1978.

Hybrid cell lines between B82, a murine cell line deficient in thymidine kinase (TK), and GM637, a simian virus 40 (SV40)-transformed human fibroblast cell line, were analyzed for SV40 T antigen (T-ag), V antigen (V-ag), and integrated viral DNA. Total high-mol wt DNA from GM637 was separated from free viral DNA by sucrose gradient fractionation and agarose gel electrophoresis. This high-mol-wt DNA was hydrolyzed with restriction endonuclease *BalI*, and SV40 DNA was detected by blot hybridization. Only one band containing SV40 DNA was consistently noted in GM637 and eight other hybrid cell DNA preparations, all of which were T-ag-positive. Cells lacking this viral DNA band were all T-ag-negative. Indirect immunofluorescence studies failed to show a correlation between the presence of T-ag and the human TK gene, which is located on chromosome 17, indicating that SV40 was not integrated on human chromosome 17. Likewise, none of the 16 gene markers tested, representing 14 different chromosomes, correlated with the presence of T-ag. Chromosome analyses of 20 B82/GM637 hybrid cell lines showed that, with one exception, the only chromosome present in all T-ag+ cell hybrids was chromosome 8. All hybrids that were T-ag+ had a single band of DNA that hybridized with SV40 DNA and migrated to the same position in all hybrids, indicating that the position of the viral sequences in the hybrids was stable. These data and those of earlier studies indicate that SV40 can integrate at one of several sites in the human genome and that this process is not chromosome specific. (30 refs)

78-6930 Viable Deletion Mutants in the Simian Virus 40 Early Region. (Eng) Feunteun, J. (Institut de Recherches Scientifiques sur le Cancer, B.P. no. 8, 94800-Villejuif, France); Kress, M.; Gardes, M.; Monier, R. *Proc Natl Acad Sci USA* 75(9): 4455-4459; 1978.

Simian virus 40 (SV40) mutants that have lost the unique site for the restriction enzyme *Taq* I at 0.565 were developed in monkey CV-1 cells in order to isolate *hr-t*-like SV40 mutants. Purified *taq* I linear SV40 DNA (50 µg/ml) was exposed to λ 5'-exonuclease (20 µg/ml) at 0 C for 30 min in 67 mM glycinate buffer, pH 9.4/2.5 mM MgCl₂, and then used for plaque assay on CV-1 cells. Five *Taq* I-resistant clones were purified, and DNA from each was prepared for restriction enzyme analysis using *Hind*III, *Hinf*I, *Alu* I, and *Hae* III. Two deletion classes were seen, both located at 0.565 on the genome. One deletion was approx 20-30 base pairs long, the other approx 250-310 base pairs long. To characterize the T antigens present in these deletion mutants, CV-1 cells infected with the mutants were labeled for 3 hr with ³⁵S-methionine and subjected to immunoprecipitation followed by gel electrophoresis. No difference was noted between any of the mutants and wild-type large T antigen. Wild-type small t antigen (t-ag), however, was missing from the mutants; in four mutants it was replaced by new polypeptides of mol wt 18,500, 17,000, 16,500, and 12,500 in greatly reduced quantities. In the fifth mutant, no t-ag of any type was detectable. The low yield of t-ag antigen was not due to a rapid turnover, and in

mixed infections the respective yields of wild-type and mutant t-ag were as expected. Thus, the deletions affect a region that is part of the t-ag coding sequences, but not those of large T antigen. Like *hr-t* polyoma mutants, the transforming ability of these cells, as measured by growth induction in agar, was clearly altered. A second complementation group in the SV40 early region is therefore defined by virtue of the fact that these mutants can complement a *tsA* mutant for transformation. (27 refs)

78-6931 Are the SV40 Induced Antigens and the Morphology of SV40 Transformed Cells Correlated (Meeting Abstract)? (Eng) Weller, N. K. (Dept. Molecular, Cellular and Developmental Biology, Univ. Colorado, Boulder, CO). *J Cell Biol* 79(2, part 2): 405a; 1978. (no refs)

78-6932 Regulation of SV40 T Antigen in SV40 Transformed Mouse Cells (Meeting Abstract). (Eng) Axelrod, H. R. (Dept. Biochemistry, Princeton Univ., Princeton, NJ); Levine, A. J. *J Cell Biol* 79(2, part 2): 401a; 1978. (2 refs)

78-6933 Induction of Infectious Virus DNA and Virus Particles by Mitomycin C in SV40-Transformed Mouse Cells. (Eng) Lavialle, C. (Institut de Recherches Scientifiques sur le Cancer, B.P. No. 8, 94800-Villejuif, France); Suarez, H. G.; Morris, A. G.; Cassingena, R. *J Gen Virol* 41(1): 201-205; 1978.

The efficiency of mitomycin C (MC) in inducing simian virus 40 (SV40) in nonpermissive mouse cells was investigated. Following infection of 3T3-4E Swiss mouse cells with SV40 (multiplicity of infection, 100 plaque-forming units/cell), an SV40-transformed cell line (SV/3T3-4E) was isolated and four independent clones were derived. The parental line and three of the clones (1, 2, and 4) spontaneously produced small amounts of infectious virus DNA but treatment of these cells with MC (0.3-3.0 $\mu\text{g}/\text{ml}$) enhanced the yield of virus DNA 10- to 100-fold. Infectious virus particles, never detected in untreated cultures, were occasionally induced by MC in the parental line and in clones 1, 2 and 4. A correlation was found between the level of SV40 DNA inducibility of a given clone and its capacity to produce virions in detectable amounts after MC treatment. It is concluded that SV/3T3-4E cells contain small amounts of free infectious SV40 DNA and that most cells in the parental line are capable of producing SV40 DNA spontaneously. (16 refs)

78-6934 Genetics of SV40 T-Antigen Expression: Studies of Twins, Heritable Syndromes and Cancer Families. (Eng) Blattner, W. A. (Environmental Branch,

NCI, Bethesda, MD, 20014); Lubiniecki, A. S.; Mulvihill, J. J.; Lalley, P.; Fraumeni, J. F. *Int J Cancer* 22(3): 231-238 1978.

The susceptibility of human skin fibroblasts to simian virus 40 (SV40) infection has been suggested as a marker of high risk for leukemia and, possibly, other cancers. To search for a pattern of genetic transmission, fibroblasts from 9 pairs of identical twins (in 7 pairs, one co-twin had acute leukemia in 1, both had aplastic anemia) and from 129 members of 21 cancer-prone families (including 16 members with cancer) were tested for SV40 tumor (T) antigen expression (SVTAE) in a 3-day immunofluorescence assay. The variance of T-antigen values was significantly less in identical than in fraternal twins or nontwin sibs (taken from cancer-prone or control families), suggesting a heritable component in the regulation of SV40 infection. In the families, SV40 T-antigen values of parents and children, as determined in the fibroblast immunofluorescence assay, were compared to models of Mendelian inheritance. At least three modes of inheritance, autosomal dominant, recessive, and x-linked, were observed. There was a tendency for offspring values to be distributed unimodally with respect to parental values. With the exception of Fanconi's anemia, the values for patients with cancer or predisposing syndromes were not elevated uniformly. The findings suggest that the genetic regulation of T-antigen expression is complex and, since no one pattern clearly predominates, probably accounts for a lack of disease effect of SVTAE. Thus, the utility of SVTAE as a marker of cancer risk appears limited. (35 refs)

78-6935 Quantitative Correlation Between Simian Virus 40 T-Antigen Synthesis and Late Viral Gene Expression in Permissive and Nonpermissive Cells. (Eng) Graessmann, A. (Institut fuer Molekularbiologie und Biochemie, Freien Universitaet Berlin, W. Berlin 33, W. Germany); Graessmann, M.; Guhl, E.; Mueller, C. *J Cell Biol* 77(1): R1-R8; 1978.

The intranuclear synthesis and accumulation of simian virus 40 (SV40) tumor (T) antigen was investigated by immunofluorescence and cytofluorometry in permissive CV1 monkey cells and nonpermissive 3T3 mouse cells at different times after infection. CV1 cells accumulated T antigen continuously over a 48-hr period after infection, whereas in 3T3 cells the T-antigen content remained constant and at a comparatively low level. Only those CV1 cells that had attained a threshold concentration of intranuclear T antigen [28 arbitrary units (AU)] synthesized viral capsid proteins (V antigen). Nearly 100% of nonpermissive 3T3 cells infected with 500 plaque-forming units of SV40 became T-antigen positive at 65 hr postinfection, but V antigen was not detected at any time. The minimum T-antigen threshold value (52 AU) for V-antigen synthesis in the nonpermissive cells was attained by microinjecting the cells with large amounts of SV40 DNA. The 3T3 cells may synthesize T antigen less efficiently and/or degrade it at a higher rate than monkey cells. Since the in-

nuclear presence of at least 250-500 viral particles or viral NA molecules overcomes nonpermissiveness, an early event in the infectious cycle (adsorption, penetration, intracellular release) must be involved in the nonpermissive type of response in infected mouse cells. (23 refs)

78-6936 Studies of Simian Sarcoma and Simian Sarcoma-associated Virus. II. Isolation of the Major Viral Glycoprotein, Properties of This Component and Specific Antiserum. (Eng) Thiel, H. J. (Dept. Surgery, Duke Univ. Medical Center, Durham, NC, 27710); Beug, H.; Graf, T.; Schwarz, H.; Schafer, W.; Bergholz, C.; Deinhardt, V. *Virology* 90(2): 360-365; 1978.

The characteristics of the major virus glycoprotein (gp) of the simian sarcoma virus complex (SSV) were studied. The glycoprotein, designated gp70, was isolated in highly pure form from virus-free supernatants of tissue culture medium using immunoadsorbents. SSV gp70 easily degraded to a 100,000-dalton product designated gp45. Both gp preparations appeared to be serologically and immunologically homogeneous. Antiserum prepared in rabbits to the purified SSV gp70 had a high neutralizing titer against SSV and was strongly cytotoxic for SSV-producing cells. The serum seemed to react mainly in a type-specific fashion with minor amounts of group and no interspecies reactivities. The interspecies reactivity of gp45 was somewhat reduced in comparison with gp70. (14 refs)

78-6937 Activation of Endogenous Type C Virus in BALB/c Mouse Cells by Herpesvirus DNA (Oncornavirus Activation/Transfection). (Eng) Boyd, A. L. (Viral Oncology Program, Frederick Cancer Res. Center, Frederick, MD, 21501); Derge, J. G.; Hamper, B. *Proc Natl Acad Sci USA* 75(9): 4558-4562; 1978.

Several virion and nonvirion DNA's were tested for their ability to activate endogenous C-type virus in BALB/c-derived mouse cells using the calcium precipitation technique. The DNA's from all herpesviruses (HV's) tested activated xenotropic C-type virus synthesis, whereas under the same conditions the DNA's from vaccinia virus and simian virus 40 were negative. Activation of C-type virus by HV DNA's was essentially linear up to about 0.2 µg DNA, but a decrease was usually observed at DNA concentrations ≥1 µg. Activation was also observed with UV-irradiated HV's, but not with irradiated vaccinia virus. Activation of C-type virus by DNA from herpes simplex virus (HSV) was observed after shearing or sonication of the DNA to an av size of 10⁶ daltons, but it was not observed with DNA sonicated to an av size of 1 x 10⁶ daltons. Alkali denaturation of DNA from HSV or treatment of the DNA with DNase, but not with DNase, destroyed its ability to activate C-type virus, as did crosslinking of the DNA with 4,5',8-trimethylpsoralen and light. The results suggest that HSV may function as a cocarcinogen. (20 refs)

78-6938 Physical Mapping of Herpes Simplex Virus-induced Polypeptides. (Eng) Marsden, H. S. (MRC Virology Unit, Inst. Virology, Univ. Glasgow, Glasgow G11 5JR, Scotland); Stow, N. D.; Preston, V. G.; Timbury, M.C.; Wilkie, N. M. *J Virol* 28(2): 624-642; 1978.

Correlation of the types of polypeptides induced by 29 herpes simplex virus type 1/type 2 intertypic recombinants with the crossover points in the recombinant DNAs enabled the map positions of many polypeptides to be deduced. The location on the physical map of 25 ³⁵S-methionine-labeled polypeptides, 11 phosphorylated polypeptides, and 4 glycosylated polypeptides was identified. Together with data from a previous study on the mapping of five immediate-early polypeptides, the results show that representatives of four groups of protein-immediate-early, late, phosphorylated, and glycosylated-map in both long and short regions. The functional organization of the herpes simplex virus genome does not therefore restrict any of these four groups to either the long or the short region. (19 refs)

78-6939 Antibodies to Herpes Simplex Virus in Cervical Cancer. (Jpn) Ishiguro, T. (Dept. Obstetrics and Gynecology, Kansai Medical Univ., Moriguchi-shi, Osaka Prefecture 570, Japan); Ozaki, Y. *Adv Obstet Gynecol (Jpn)* 30(1): 9-13; 1978.

Levels of antibodies to herpes simplex virus types 1 and 2 (HSV-1 and HSV-2) were measured in 79 Japanese patients with cervical cancer, 11 with dysplasia, and 155 controls (including 145 healthy women and 10 with ovarian cancer). There was no significant difference in mean antibody titers among these groups. Furthermore, 18.2% of the dysplasia patients, 16.7% of those with carcinoma in situ (CIS), 11.1% of those with early invasive cancer (EIC), 3.2% of those with invasive cancer (IC), 0% of those with advanced cancer (AC), and 8.3% of healthy controls had no antibodies to either virus. Statistically, these data were not significant (SNS). The prevalence of single infection with HSV-1, as defined by an HSV-2/HSV-1 ratio of ≤65, was 36.4% in dysplasia, 50.0% in CIS, 27.8% in EIC, 19.4% in IC, 27.8% in AC, and 31.7% in healthy controls (data SNS). The prevalence of HSV-2-positive women, as defined by an HSV-2/HSV-1 ratio of ≥93, was 9.1% in dysplasia, 0% in CIS, 16.7% in EIC, 11.1% in AC, and 9.0% in healthy controls (data SNS); however, the prevalence in IC was 19.4%, a rate significantly higher than that in healthy controls (p < 0.05). These results suggest that HSV-1 is not associated with cervical cancer or dysplasia and that HSV-2 is involved with the progression but probably not the initiation of cervical cancer. (13 refs)

78-6940 Preliminary Studies of a Non-Virionic Antigen Associated with Herpes Simplex Virus 1 and 2 (HSV₁-HSV₂). (Eng) Accinni, R. (Clinica Medica II dell'Universita di Milano, Milano, Italy); Ferrara, R.;

Tarro, G.; Sisillo, E.; Privitera, L.; Biancardi, C.; Bartorelli, A. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 199-210; 1977.

Nonvirion antigens of herpes virus (HSV₁) were studied radioimmunologically, by gel filtration on Sephadex G-200, and by polyacrylamide gel electrophoresis (PAGE). In the radioimmunological study, soluble extracts of normal or infected cells incubated with pathological sera were found to possess different immunological properties. Gel filtration demonstrated two peaks in pellet extracts from infected cells that were absent in analogous extracts from normal cell pellets. PAGE demonstrated that soluble extracts from suspensions of infected cells contained three radioactive bands that were absent in extracts from normal cells. (7 refs)

78-6941 Ultraviolet Reactivation of Herpes Simplex Virus Is Mutagenic and Inducible in Mammalian Cells. (Eng) Bandyopadhyay DasGupta, U. (Dept. Therapeutic Radiology, Yale Univ. Sch. Medicine, 333 Cedar St., New Haven, CT, 06510); Summers, W. C. *Proc Natl Acad Sci USA* 75(5): 2378-2381; 1978.

Experiments were performed using the thymidine kinase (TK) locus of herpes simplex virus type 1 (HSV-1), strain Cl 101, to determine whether UV reactivation (Weigle reactivation) in mammalian cells is error-prone, thus leading to increased mutagenesis. The survival of UV-irradiated HSV-1 on UV-irradiated Vero cells was higher than that on unirradiated cells. A time lapse of 17 hr between irradiation of the host cells and infection with virus was needed to achieve max reactivation. In parallel experiments in which the frequencies of occurrence of the forward mutation in the TK gene of the virus were measured, growth of HSV-1 on UV-irradiated cells yielded progeny virus that had higher frequencies of TK- mutants than did progeny from infections of control cells. The time course of development of this mutagenic effect was the same as that for the development of UV-reactivation capacity. Furthermore, development of UV reactivation could be blocked by inhibition of protein synthesis. These findings suggest that an error-prone inducible UV-reactivation phenomenon exists in mammalian cells. (19 refs)

78-6942 Studies on the Synthesis and Structure of Herpes Simplex Virus RNA. (Eng) Stringer, J. R. (Univ. California, Irvine, Irvine, CA). *Diss Abstr Int [B]* 39(2): 511; 1978. (no refs)

78-6943 Temperature-sensitive Mutants of Herpes Simplex Virus Type 1. Defective in Transcription and Post-transcriptional Functions Required for Viral DNA Studies. (Eng) Parris, D. S. (Dept. Microbiology, Univ. Tennessee, Knoxville, TN, 37196); Courtney, R. J.; Schaffner, P. A. *Virology* 90(2): 177-186; 1978.

DNA- temperature-sensitive (*ts*) mutants of herpes simplex virus (HSV) representing four complementation groups (A, B, C, and D) were examined in temperature shift-down experiments to determine whether viral DNA synthesis is inhibited at the level of transcription or translation at the nonpermissive temperature. Synthesis of viral DNA after temperature shift-down appeared to require *de novo* transcription in the case of the *ts*A mutants, *ts*B2, and *ts*C4, but not in the case of *ts*D9. For *ts*B2 and *ts*C4, approx 2 hr was required for recovery of DNA synthesis capabilities following shift-down. Viral transcripts specifying proteins required for DNA synthesis were synthesized 2 hr postinfection in *ts*D9-infected cells at the nonpermissive temperature. Polypeptide synthesis by all four mutants following shift-down in the presence of an inhibitor was similar to that observed in mutant-infected cells maintained at the nonpermissive temperature in the absence of an inhibitor. Restoration of the wild-type pattern of viral polypeptide synthesis following shift-down in the presence of an inhibitor was greater for the groups A, C, and D mutants than from the group B mutants. (14 refs)

78-6944 Transformation of Hamster Embryo Fibroblasts by a Specific Fragment of the Herpes Simplex Virus Genome. (Eng) Camacho, A. (Dept. Microbiology, Univ. Chicago, Chicago, IL, 60637); Spear, P. G. *Cell* 15(3): 993-1002; 1978.

Isolated restriction endonuclease fragments of the herpes simplex virus type 1 (HSV-1) genome were introduced into hamster embryo cells to identify DNA sequences capable of transforming the cells with respect to acquisition of properties correlated with tumorigenicity. One of the fragments generated by cleavage of HSV-1 DNA with the restriction endonuclease *Xba*I induced transformation at a frequency of about 10 colonies per quantity of fragment recovered from 1 µg of uncut DNA; fractions containing the other *Xba*I fragments failed to induce transformation reproducibly, although occasional colonies were detected. The fragment with transforming activity (*Xba*I-F) is 15.5 x 10⁶ daltons in mol wt and is located between 0.30 and 0.45 map units on the HSV-1 genome. The *Xba*I-F transformants were selected for their ability to replicate in low concentrations of serum; in addition, they were found to attain high saturation densities in the presence of 10% serum and to form colonies in semisolid medium. Moreover, the transformed cells produced at least one viral gene product (a membrane glycoprotein) encoded in the fragment used for transformation, indicating not only that viral DNA was incorporated in the cells but that viral genes were expressed. (37 refs)

6945 **Genital Herpesvirus in the Etiology of Human Cervix Uteri Cancer.** (Eng) Kessler, I. I. (Dept. Epidemiology, Johns Hopkins Univ., Baltimore, MD, 21205). In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): 1977. 1(part 1): 1193 pp.; 211-229; 1977.

Investigations of the possible role of herpesviruses in cervical cancer (CC) were made. The first was a retrospective, case-control study of 350 Yugoslavian women with histologically confirmed squamous carcinoma of the uterine cervix and 350 matched controls. Herpes simplex virus type 2 (HSV-2) antibody titers were higher among the CC patients than the controls. The relative risk of CC in women positive for HSV-2 was significantly increased, even after adjustments were made for the effects of 34 epidemiologic factors associated with CC. In addition, compared with controls, women who developed CC were more likely to smoke and drink alcoholic beverages, they had higher blood pressures, their parents tended to be older at their birth, they had more multiple marriages and an earlier start of sexual activity, consistently reduced coital frequency, and they were more likely to have abnormal menstrual histories. There was an unexpected two- to threefold greater mortality, especially at younger ages, of the first husbands of the CC patients, compared with the control first husbands. In a second study, still in progress, large groups of women are being observed: one married men who, at some other time, had wives who developed CC and a second married to men without such histories. To date, abnormal pap tests have been found in 13.6% of the "case" wives vs 7.2% of the control wives. CC or carcinoma in situ has been diagnosed in 3.5% of the case wives and 1.2% of the control wives. These data support an etiological role for HSV-2 in CC and they suggest that hormonal factors may also be involved. (23 refs)

6946 **Evaluation of the Effects of Phosphonoacetic Acid and 2-Deoxy-D-glucose on Herpesvirus Saimiri and Epstein-Barr Virus.** (Eng) Ablashi, D. (CI-Frederick Cancer Res. Center, Fort Detrick, Frederick, MD, 21501); Armstrong, G.; Fellows, C.; Easton, J.; Carson, G.; Twardzik, D. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 245-261; 1977.

The effect of phosphonoacetic acid (PAA) and 2-deoxy-D-glucose on *Herpesvirus saimiri* (HVS) and on Epstein-Barr virus (EBV) was studied in vitro. PAA significantly and reversibly inhibited the replication of HVS in African green monkey kidney (Vero) cells when it was added immedi-

ately after a 2-hr absorption period, but not later. The compound had no effect on the virus itself, on viral absorption, or on the induction of early antigen, but it did inhibit the production of late and membrane antigens almost completely. The failure of more template to reverse the inhibition of Vero cell DNA polymerases by PAA suggests that PAA does not interact with the uncoated DNA of input HVS in Vero cells. Rather, PAA may block a cellular process essential for HVS replication. PAA had no effect on the induction of endogenous or exogenous EBV early antigen, and it did not inhibit the appearance of EBV nuclear antigen in actively replicating Raji cells. Thus, the compound may have no effect on the replication of EBV DNA in Raji cells. This DNA is either integrated with the cell genome or in a free circular form. However, under certain conditions, PAA inhibited the expression of EBV membrane and viral capsid antigens. If the resistance of HVS to PAA, beginning 2 hr after the end of absorption, is similar to that of EBV DNA in Raji cells, infecting HVS DNA may assume a relatively stable and resistant form within the infected cell. It is not known if this would involve integration of input HVS DNA with the cellular genome. 2-Deoxy-D-glucose had no effect on HVS. (30 refs)

78-6947 **Host Cell Reactivation of Epstein-Barr Virus in Normal and Repair-defective Leukocytes.** (Eng) Henderson, E. E. (Dept. Microbiology and Immunology, Temple Univ. Sch. Medicine, Philadelphia, PA, 19140). *Cancer Res* 38(10): 3256-3263; 1978.

Epstein-Barr virus (EBV)-induced transformation efficiencies were quantitated for peripheral blood WBC from two patients with xeroderma pigmentosum (XP) and a patient with Cockayne's syndrome. No differences in transformation efficiencies were observed among XP, Cockayne's syndrome, or normal adult WBC, as judged by transformed centers or limiting dilution assays. However, when either the WBC or the infecting EBV was UV-irradiated prior to virus exposure, XP WBC showed a greater decrease in ability to undergo in vitro transformation. The doses of the transforming functions of the B95-8 strain of EBV lethal to 37% of the animals were 70, 725, and 750 ergs/mm² when measured in XP, XP heterozygotic, and normal adult WBC, respectively. Transforming activity of UV-irradiated EBV could be partially restored by illuminating infected XP WBC with visible light 36 hr after virus exposure. Illumination with visible light had little effect on transformation of neonatal or normal adult WBC by UV-irradiated EBV. Colony formation in soft agarose by the resulting lymphoblastoid cell lines established from primary XP or Cockayne's syndrome WBC showed increased sensitivity to UV irradiation when compared with lymphoblastoid cell lines established from heterozygotic or normal individuals. These results indicate that EBV is similar to other herpesviruses in being subject to host cell reactivation and that the EBV-induced transformation does not require or alter these functions. (39 refs.)

78-6948 Detection of Epstein-Barr Virus in Biopsied Malignant Lymphoma Cell and Its Continuous Culture. (Eng) Suzuki, T. (First Dept. Pathology, Niigata Univ. Sch. Medicine, 1-Bancho, Asahimachi-dori, Niigata 951, Japan); Oboshi, S.; Ishihara, N.; Emura, I.; Ohnishi, Y.; Sairenji, T.; Hinuma, Y. *Gann* 69(2): 213-221; 1978.

The detection of the Epstein-Barr (EB) virus genome in the tumor cells from a patient with non-Burkitt-type B-cell lymphoma and the establishment of tumor cell lines (Fujimaki-I and -II) from these cells are described. More than 80% of the tumor cells obtained from a pericardial effusion of the patient were positive for EB nuclear antigen (EBNA), and herpes-type virus particles were detected in the tumor cell nuclei. Tumor cells from a biopsy specimen of the original sc tumor grew continuously in vitro without any cell death (Fujimaki-I) and tumor cells from the pericardial effusion proliferated without a growth lag phase or cell death (Fujimaki-II). The cytological characteristics of the Fujimaki-II cells were similar to those of the tumor cells in vivo. Seventy percent of these cells were positive for EBNA and 4% were positive for EB virus capsid antigen. The modal chromosome number of the cells in the twenty-fourth generation was 46, and no marker chromosomes were found. IgG was monoclonally demonstrated in the culture medium. Fatal generalized toxoplasmosis resulted when the original tumor tissue was inoculated ip, sc, or under the renal capsule in athymic nude mice. Fujimaki-II cells heterotransplanted sc or ip into mice did not induce tumors, and the virus released by these cells did not transform cord blood lymphocytes. These data suggest that the Fujimaki cells did not produce detectable amounts of infectious virus. This may be the first case of non-Burkitt type lymphoma in which the EB virus genome was directly detected in the tumor cells. (22 refs)

78-6949 A Study on Epstein Barr Virus Receptor Activity in Cell Free Extracts of Human Lymphoblastoid Cells. (Eng) Ng, M. H. (Dept. Microbiology, Queen Mary Hosp. Compound, Univ. Hong Kong, Hong Kong, China); Lui, S. W.; Chan, K. H. *Arch Virol* 58(3): 221-233; 1978.

The detection of Epstein Barr virus (EBV) receptor activity in the surface extract of Raji cells and to a lesser extent in that of BJAB cells both are B lymphoblastoid cells susceptible to EBV infection) is reported, and results of a preliminary attempt to purify the Raji cell receptors are also presented. Receptor activity was not detected in extracts of P₃HR-1 cells and human diploid fibroblasts, neither of which are known to be susceptible to EBV infection. Receptor activity in the Raji cell extract was found to be associated with membranous structures, which may have rendered the activity resistant to treatment with trypsin and sonication. The receptor activity was abrogated if the extract was exposed to neutral detergent. Binding of receptor activity was observed when Raji cell extract was chromatographed on a column of immobilized EBV

components. Subsequent electrophoretic analysis showed however, that this procedure did not result in an appreciable purification of the receptor activity. Neutral detergent-treated extract was similarly chromatographed. The resulting eluates did not contain detectable receptor activity but were heterogeneous in protein content compared with the original extract. It is uncertain if these EBV binding proteins are involved in the receptor activity of the extract. (17 refs)

78-6950 Expression of Altered Ribonucleotide Reductase Activity Associated with the Replication of the Epstein-Barr Virus. (Eng) Henry, B. E. (Dept. Microbiology, Univ. Mississippi Medical Center, Jackson, MS 39216); Glaser, R.; Hewetson, J.; O'Callaghan, D. J. *Virology* 89(1): 262-271; 1978.

Extracts of control and 5-iododeoxyuridine (IUdR)-induced epithelial/Burkitt hybrid cells (D98/HR-1) and Raji cells were assayed for ribonucleotide reductase (RR) activity in the presence and absence of hydroxyurea (HU). The expression of the Epstein-Barr virus (EBV) genome in these cells can be regulated by induction with IUdR or superinfection with the virus. HU inhibited the enzyme activity of control D98/HR-1 and Raji cells by > 70% at both low (2×10^{-4} M) and high (5×10^{-4} M) concentrations. However the reductase activities of IUdR-induced D98/HR-1 cells, Raji cells superinfected with EBV, and IUdR-induced Raji cells were resistant to both levels of HU; enzyme activities of 85% to > 100% of control values were obtained in all cases. Under conditions that allowed only partial expression of the EBV genome (before removal of IUdR; early time after superinfection), significant levels (70%-85%) of HU-resistant RR activity were obtained in the presence of 2×10^{-4} M HU, but only 30%-45% of control reductase activity was observed at the higher concentration. Mixing experiments employing combinations of various D98/HR-1 control and induced cell extracts indicated that the HU-resistant reductase activity of induced D98/HR-1 cells was due to altered enzyme activity and not to nonenzymatic factors. Experiments in which HU was preincubated with control and induced cell extracts showed that the HU-resistant RR in extracts of induced D98/HR-1 cells was not due to inactivation of the inhibitor by an enzyme or factor present in these cells. These findings, together with similar findings for equine herpesvirus type 1 and herpes simplex virus types 1 and 2, suggest that alteration of RR activity may be a key event in herpesvirus replication. (22 refs.)

78-6951 Epidemiological Evidence for Causal Relationship Between Epstein-Barr Virus and Burkitt Lymphoma from Ugandan Prospective Study. (Eng) de The G. (International Agency Res. Cancer, WHO, 150 cours Albert Thomas, 69372 Lyon, France); Geser, A.; Day, N. E.; Tukei, P. M.; Williams, E. H.; Beri, D. P.; Smith, P. G.

A. G.; Bornkamm, G. W.; Feorino, P.; Henle, W. *Nature* 267(5673): 756-761; 1978.

From 1972-1977, a large, prospective, seroepidemiological study was undertaken in Uganda to evaluate evidence that Epstein-Barr virus (EBV) is a necessary factor for the development of Burkitt's lymphoma (BL). From 1972 to 1974, 147 samples were taken from 42,000 children up to age 8. Of these, 31 cases of BL had been diagnosed in the area, 14 of which were in children originally bled. (Only 3/31 patients were missed at original bleeding; the rest were from a region covered by the field team or were too old for inclusion.) In 14 children, the interval between initial serum collection and BL onset was 7-54 mo. Significantly higher titers of antibodies (ab's) to virus capsid antigen (VCA) were seen in BL sera than in matched controls. These titers were indicative of the high ab levels associated with BL and present long before clinical onset. There were no marked differences in pre- and post-BL ab titers to herpes simplex and cytomegalovirus, eliminating the possibility that this was due to a long-standing defect in cell-mediated immunity. Six of seven patients with detectable EBV DNA in tumor tissue had prediagnosis VCA ab titers high or higher than those in controls. Thus, the association of BL with pre-diagnosis VCA titers may be confined to cases characterized by EBV DNA in the tumor cells. There may be two types of BL with different etiologies, one associated with EBV and predominantly in tropical areas and one not associated with EBV and found in temperate areas. The results strongly support a causal association between EBV infection and BL. However, even in a region where infection with EBV is nearly universal, only 1 child in 1,000 develops BL; therefore, other factors are required. The potential oncogenicity of EBV is realized only in exceptional circumstances. (45 refs.)

52 A New Manifestation of Infection with Epstein-Barr Virus. (Eng) Smith, H. (Dept. Infectious Diseases and Immunological Medicine, Northwick Hospital, Harrow HA1 3UJ, England); Denman, A. M. *British Medical Journal* 2(6132): 248; 1978.

A 47-year-old woman with a specific T-lymphocyte defect suffered a prolonged bizarre illness produced by Epstein-Barr virus (EBV). The illness was intermediate in severity and outlasted the common self-limiting infectious mononucleosis and the more aggressive malignant condition associated with this virus. Antibodies to EB virus capsid antigen (detected by immunofluorescence) showed a progressive rise in titer. It is suggested that the patient suffered from a specific defect in cell-mediated immunity to EBV, shown morphologically in her failure to produce atypical mononuclear cells and functionally in her inability to mount a specific reaction against cells infected by this agent. The patient's history of recurrent varicella indicated that there was a constitutional inability to react to DNA containing herpesviruses. The patient responded to treatment with corticosteroids. (4 refs)

78-6953 Study of Epstein-Barr Virus-determined Nuclear Antigen (EBNA) by Chromatography on Fixed Cell Nuclei. (Eng) Hirsch, I. (Dept. Experimental Virology, Inst. Sera and Vaccines, 101 03 Prague, Czechoslovakia); Suchankova, A.; Zavadova, H.; Vonka, V. *International Journal of Cancer* 22(5): 535-541; 1978.

A low concentration of NaCl (50 mM) and pH 6.0 were found to be optimal conditions for the in vitro conversion of Epstein-Barr virus nuclear antigen (EBNA)-negative nuclei to EBNA-positive forms by the addition of complement-fixing (CF) antigen from the Raji human lymphoblastoid cell lines. The conversion was sensitive to treatment with DNase but not RNase, which suggests that EBV-CF antigen binds specifically to nuclear DNA. Identical conditions were optimal for both the elution of EBNA from in vitro-converted nuclei and the adsorption and elution of EBV-CF antigen in DNA-cellulose chromatography. This indicates that the DNA-binding properties of EBNA antigen can be studied in a microchromatographic system employing fixed nuclei. The advantages of this method are its simplicity, capacity for analyzing many samples per experiment, and requirements of very small amounts of material. Elution patterns for EBNA were significantly different when nuclei from different cell lines (Raji, Ramos, and P3HR-1) were converted to EBNA positivity. EBNA was eluted from converted nuclei of EBV genome-positive P3HR-1 cells at an almost 0.1 M higher concentration of NaCl than was necessary for a similar degree of elution from nuclei of EBV genome-negative Ramos cells. (22 refs)

78-6954 DNA of Epstein-Barr Virus. IV. Linkage Map of Restriction Enzyme Fragments of the B95-8 and W91 Strains of Epstein-Barr Virus. (Eng) Given, D. (Dept. Medicine, Univ. Chicago, Chicago, IL, 60637); Kieff, E. *Journal of Virology* 28(2): 524-542; 1978.

The arrangement of restriction enzyme sites *EcoRI*, *Hsu I*, and *Sal I* in the DNA of the B95-8 and W91 isolates of Epstein-Barr virus (EBV) was determined from the size of the single-enzyme-cleaved fragments and from blot hybridizations that identify which fragments cut from the DNA with one enzyme contain nucleotide sequences in common with fragments cut from the DNA with a second enzyme. The data indicate that there is a consistent and unique sequence arrangement of approx 95 to 100 x 10⁶ daltons of B95-8 strain EBV DNA. Both termini were variable in length, but the opposite ends of the molecule contained homologous sequences. The high degree of homology between the opposite ends of the molecule and the similarity in size of the "additional" 500-base pair segments suggest that there are identical repeating units at both ends of the DNA. The size of almost all *EcoRI*, *Hsu I*, and *Sal I* fragments of W91 DNA was identical to that of the B95-8 DNA fragments except that W91 DNA contained approx 7 x 10⁶ daltons of "extra" DNA in the *Hsu I*-*EcoRI*-C overlap region of B95-8 DNA. Furthermore, the restriction endonuclease fragments of the W91

and B95-8 DNA's, which are of similar mol wt, had homologous nucleotide sequences. Moreover, the W91 fragments contained only sequences from a single region of the B95-8 genome. Two lines of evidence indicate that the extra DNA in W91 *Eco*RI C and *Hsu*I D₁ and D₂ fragments is viral DNA and not cellular. (48 refs)

78-6955 Electron Microscopic Studies of Tumor Viruses. II. Entry and Uncoating of Epstein-Barr Virus. (Eng) Miyamoto, K. (Pathology Div., Natl. Cancer Center Res. Inst., Tsukiji 5-1-1, Chuo-ku, Tokyo 104, Japan); Sairenji, T.; Hinuma, Y. *Microbiol Immunol* 22(8): 463-474; 1978.

The entry of Epstein-Barr virus into human lymphoblastoid cells (Daudi) was studied ultrastructurally. Two types of virions were observed on the cell surface: the zone separating the capsid from the dense core in one case was electron-translucent, but this zone in the other case was more dense because of the presence of finely granular material. These virions were referred to as light and dense particles, respectively. Virus attached to the cell surface at time 0 consisted of 56% dense particles, 41% light particles, and 3% empty particles (ie, devoid of cores). In samples taken at time 0, no free virus was found in the cell cytoplasm, although a few (<1%) of the cells showed degenerative changes. When the cells were warmed to 37 C, two distinct types of interaction between the viral envelope and cell membrane ensued, these changes being observed most frequently in samples taken at 30-60 min. The first change was simultaneous dissolution of both the viral envelope and cell membrane at the point of attachment, presumably resulting in passage of viral capsids into the cytoplasm. The other change was dissolution only of the cell membrane at the site of attachment, resulting in penetration of the enveloped virus. Envelope dissolution appeared to occur subsequently in the cytoplasm, with release of capsids. Fusion of the viral envelope with the cell membrane was not observed. Enveloped viruses may differ in the mechanism by which they gain entry and in their ability to cause cell fusion at early stages of infection. (21 refs)

78-6956 Acute Lymphocytic Leukemia in the Course of a Primary Herpes Infection with Hepatitis and Bone Marrow Aplasia in a Two-Year-Old Child. (Fre) De Lumley, L. (Travail du Service de Pédiatrie I, Centre Hospitalier Universitaire Dupuytren, 87000 Limoges, France); Malinvaud, G.; Boulesteix, J.; Gaillard, S.; Colombeau, M. C. *Pédiatrie* 33(3): 291-298; 1978.

Acute lymphocytic leukemia (ALL) developed in a 2-yr-old boy admitted to the hospital with viral hepatitis. Prior to admission, the child was given sulfamethoxazole, trimethoprim, and pipebuzone for fever and stomatitis. Acute hepatitis was diagnosed on the basis of fever, icterus, pruritis, and hepatomegaly. Blood studies revealed anemia, leukopenia

(10% polynuclear neutrophils, 90% lymphocytes), and thrombocytopenia. Viral serology was negative; however, there was a high level of antiherpes antibodies. The patient was treated with cefalotine, (125 mg/kg/day), gentamycin (4 mg/kg/day), prednisone (2 mg/kg/day), and RBC and WB transfusions. The cefalotin was replaced by colistin (1 million units/day) when a dermatitis reaction occurred. The patient improved and was well enough to leave the hospital on the 31st day with regression of hepatic and inflammatory symptoms. Four months after the initial admission, he was readmitted with clinical symptoms of a tumor, including hepatosplenomegaly and widespread adenopathy. A bone marrow puncture revealed 82% small lymphoblasts. Despite treatment with combination chemotherapy for ALL, the child died 17 mo after the first admission. The question arises as to whether the herpesvirus infection was severe as a result of a preleukemic condition or whether the herpesvirus was responsible for the ALL. (26 refs)

78-6957 Establishment of a Human Hepatoma Cell Line Which Produces Hepatitis B Surface Antigen (HBsAg). (Eng) Alexander, J. (Virus Cancer Res. Unit., Polio-myelitis Res. Foundation, Johannesburg, South Africa); Bey, E.; Macnab, G. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 321-330; 1977.

Biochemical and immunological studies were made of a cell line established from a primary liver carcinoma of a 24-yr-old man. One aim of the studies was to examine the cells for the presence of virus particles and viral antigens. The cells, which had been in continuous culture for 30 mo, had a male human chromosome pattern. The modal chromosome number was 56.6, and markers were present. The enzymes conformed to the human pattern, and glucose-6-phosphate dehydrogenase type A isozyme was found. Electron microscope studies showed that the cells contained glycogen granules and that microvilli and smooth endoplasmic reticulum were common. There was no evidence of virus particles, and there were no signs of lesions commonly associated with hepatitis-infected liver cells. The cells produced hepatitis B surface antigen (HBsAg), which was detected predominantly in the culture supernatant. The antigen levels increased with time, and antigen was produced at approx 500 µg/day/10⁶ cells. HBsAg activity in the culture supernatant was associated with amorphous particles ranging from 16 to 24 nanometers in diameter. The particles had a buoyant density in CsCl of 1.18-1.2 g/cm³. (23 refs)

78-6958 Hepatocellular Carcinoma Associated with HBs Antigen: A Study of 25 Cases from Vietnam. (Fre) Doury, J. C. (Laboratoire d'anatomie pathologique, In-

de Medecine tropicale du service de sante des armees, du Pharo, 13998 Marseille-Armees, France); Roche, J. B. *Rev Epidemiol Sante Publique* 26(2): 161-170; 1978.

orcein staining and indirect immunofluorescence on thin sections, hepatitis B surface antigen was demonstrated in 64% of 25 hepatocellular carcinomas (including 1 bone metastasis) in Vietnamese patients. Literature findings suggesting a causal role for hepatitis B virus in primary liver cancer are discussed. (29 refs)

5959 **A Virus Similar to Human Hepatitis B Virus Associated with Hepatitis and Hepatoma in Woodchucks.** (Eng) Summers, J. (Inst. Cancer Res., Fox Chase Cancer Center, Philadelphia, PA, 19111); Smolec, J. Snyder, R. *Proc Natl Acad Sci USA* 75(9): 4533-4537; 1978.

ence indicates that human hepatitis B virus (HBV) belongs to a novel class of DNA viruses that may be causative agents of hepatitis in species other than humans. Approximately 10% of the serum samples from woodchucks at a research laboratory contained DNA polymerase (Pase)-containing particles in amounts comparable to those found in hepatitis B surface antigen (HBsAg)-positive sera. Three animals (CW307, CW326, and CW316) subsequently died and were necropsied further to determine the possible relationship between the particles and hepatic lesions. CW307 and CW326 had hepatocellular carcinomas and evidence of chronic active hepatitis, and their sera contained detectable levels of endogenous DNA Pase, but serum from CW316, with no apparent hepatic lesions, had no detectable DNA Pase-containing particles. Large numbers of spherical and tubular forms approximately 30-40 nm in diameter were found in the serum of CW307 and CW326. They strongly resembled HBsAg, but floated at a slightly lower density in CsCl gradients. A second population of particles banding at 1.215 g/cm³ contained a DNA polymerase (Pase) activity. The DNA was similar in size and structure to HBV DNA. In addition, each DNA molecule appeared to contain a substantial single-stranded region, as in HBV DNA. The action of the internal DNA Pase on the single-stranded region accounts for the endogenous DNA Pase activity observed in the serum of some woodchucks. In the CsCl gradient, fractions of the gradient that had endogenous DNA Pase activity contained double-layered particles, 55 nm in diameter, that resembled Dane particles, thought to be the complete virus of hepatitis B. It is likely that these double-layered particles contain the DNA and DNA Pase activity. The double-layered particles found in the diseased woodchucks were similar but not identical to those associated with hepatitis B. Preliminary experiments indicate that the exposed antigenic sites on the small particles do not cross-react with HBsAg. (48 refs)

5960 **An Electroncytochemical Study of 5'-Nucleotide Phosphodiesterase in Hepatoma, Cirrhosis**

and Hepatitis. (Eng) Tsou, K. C. (Harrison Dept. Surgical Res., Sch. Medicine, Univ. Pennsylvania, Philadelphia, PA, 19104); Lo, K. W.; Rhoads, J. E.; Mackie, J. A. *In: Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 331-342; 1977.

Polyacrylamide gel electrophoresis analyses have recently shown that there is an isozyme of 5'-nucleotide phosphodiesterase (5'-NPDase) in the sera of patients with hepatitis B and hepatoma and in some with cirrhosis. To determine the significance of this enzyme in linking the etiology of hepatoma with hepatitis B-related virus, hepatoma tissues from two patients were studied ultrastructurally and electron cytochemically. The sera of these two patients were positive for 5'-NPDase and negative for hepatitis B surface antigen (HBsAg). α -Fetoprotein levels were 10 and 400 nanograms/ml, respectively. Both tumor samples were positive for viruslike particles resembling Dane particles of hepatitis B serum. These particles were also seen in cirrhotic areas of the liver, but most were found enclosed in cytoplasmic vesicles. One interesting feature of the cytoplasmic organelles was an abundance of circular vesicular membranes that often surrounded the mitochondria. This feature was also seen in the cirrhotic areas of both patients. An electron cytochemical method demonstrated the presence of deposits of 5'-NPDase activity on particles obtained by precipitation of hepatitis B serum with anti-HBsAg. These data suggest that there may be a group of human hepatomas etiologically linked to hepatitis B-associated virus and that some cases of cirrhosis may be a reflection of the host-immune response to viral proliferation. A possible role for 5'-NPDase in hepatitis B infection is suggested. (48 refs)

78-6961 **Liver Cell Dysplasia and Hepatitis B Surface and Core Antigens in Cirrhosis and Hepatocellular Carcinoma of Autopsy Cases.** (Eng) Sakurai, M. (1st Dept. Pathology, Osaka Univ. Medical Sch., Osaka, Japan). *Acta Pathol Jpn* 28(5): 705-719; 1978.

The relationships between cirrhosis, hepatoma, liver cell dysplasia, serum and cellular levels of hepatitis B surface and core antigens (HBsAg and HBcAg), and serum levels of antibody to HBsAg (anti-HBs) were studied based on examination of tissues from 180 autopsy cases with cirrhosis and/or hepatoma. Dysplasia was not associated with hepatoma in the absence of cirrhosis and was significantly more common in cirrhotic livers with tumors than in cirrhotic livers without tumors. The association between cirrhosis and dysplasia did not vary with age or sex. There was a significant correlation between high HBsAg levels and cirrhosis plus dysplasia, and orcein-positive cells were found significantly more often in cirrhotic livers than in noncirrhotic livers. The morphological characteristics of the orcein-positive liver cells could not

be correlated with dysplasia or hepatoma. HBcAg levels did not correlate significantly with dysplasia or hepatoma, but they did correlate with active cirrhosis, marked liver cell degeneration and necrosis, and membranous diffuse-type HBsAg in liver cells. (21 refs)

78-6962 Detection of Adenovirus Integration Sites in Host Cell Chromosomes. (Eng) McDougall, J. K. (Cold Spring Harbor Lab., Cold Spring Harbor, NY, 11524); Dunn, A. R.; Gallimore, P. H. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 129-137; 1977.

The location and function of integrated adenovirus (Ad) DNA sequences in rodent and human cells were determined by several approaches. Somatic cell hybrids were derived by fusion of virus-transformed rat cells and mouse cells deficient in thymidine kinase. The hybrid lines were isolated by a double selective system. Chromosomes (Chr) in the hybrids bearing mouse centromeres were identified by in situ hybridization with mouse satellite complementary RNA as the probe. The only rat Chr positively identified in the hybrid lines were 1, 2, 11, 12, 17, and 19; however, none of these Chr was consistently present in every hybrid line. All hybrids were screened for virus messenger RNA (mRNA) by in situ hybridization using ¹²⁵I-labeled Ad DNA. None of the six rat Chr identified was consistently present in viral mRNA or T-antigen-positive lines, and none of the 15 rat isozymes tested, apart from thymidine kinase, was always present in hybrid cells expressing viral functions. Thus, cytogenetic analysis and rat isozyme segregation data indicated that many hybrids contained new Chr translocations not present in the parental lines and that none of the marker Chr in the transformed cells appeared in the hybrid lines. The situation with an Ad-transformed human cell line was similar, in that these cells were heterogeneous in Chr constitution, with an average count in the triploid range and with only a small amount of viral DNA/diploid quantity of human DNA. Thus, Ad DNA is integrated into only a few Chr of rat cells, but it was not possible to define a specific Chr associated with the expression of Ad T antigen. (26 refs)

78-6963 Synthesis of a Structural Adenovirus Polypeptide in the Absence of Viral DNA Replication. (Eng) Persson, H. (Dept. Microbiology, Biomedical Center, Box 581, S-751 23 Uppsala, Sweden); Pettersson, U.; Mathews, M. B. *Virology* 90(1): 67-79; 1978.

The kinetics and control of adenovirus type 2 (Ad2) polypeptide IX (ppIX) synthesis were studied in HeLa cells pulse-labeled with ³⁵S-methionine at different times beginning at 2 hr postinfection. Cell extracts were immunoprecipitated

with antisera against Ad2 early polypeptides or structural components, and the immunoprecipitates were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Polypeptide IX, a structural protein, was first detectable 6 hr postinfection, 4 hr before ppII and III, and 6 hr before ppIV, the major structural components. When cells were treated with cytosine arabinoside for 1 hr after Ad2 infection, ppIX antiserum precipitated a 12,000-mol-wt peptide that specifically comigrated with ppIX on SDS-PAGE thus demonstrating the synthesis of ppIX in the absence of viral RNA replication. Cycloheximide which blocks protein synthesis as well as viral replication, did not inhibit ppIX synthesis or accumulation of its messenger RNA (mRNA), although the mRNA's of the other structural components remained undetectable. RNA from cycloheximide-treated, virus-infected cells also directed the synthesis of a 12,000-mol-wt polypeptide that comigrated with ppIX and was precipitated by antiserum against ppIX. Size fractionation of the RNA indicated that it sedimented at the same rate, 9S, as the late mRNA for ppIX. Because transcription of this polypeptide occurs at a different time and is under different regulation than the other late structural genes, ppIX may have functions in addition to its structural role in the virus particle. (48 refs)

78-6964 The Spliced Structures of Adenovirus 2 Fiber Message and the Other Late mRNAs. (Eng) Chow, L. T. (Cold Spring Harbor Lab., P.O. Box 100, Cold Spring Harbor, NY, 11524); Broker, T. R. *Cell* 15(2): 497-510; 1978.

Several rightward-transcribed (R)-strand transcripts in the adenovirus 2 (Ad2) map interval between coordinates 27 and 50 were mapped. In common with eight previously reported Ad2 messenger RNA's (mRNA's), four RNA species were identified with the same tripartite leader sequence derived from coordinates 16.6, 19.6, and 26.6. This leader was joined to messages with 5' ends located at coordinates 29.1, 30.5, 33.9, and 42.8, respectively. The RNA transcripts extending from 29.1 to 39.0 covered most or all of the virus-associated RNA genes (29.0-30.1). The late RNA's could be grouped into several families of transcripts that differed substantially in the coordinate at which the leader sequences were attached to the main body of message. The number of RNA species transcribed from the Ad2 chromosome between coordinates 29 and 50 exceeded the number of identified Ad2 proteins. A subpopulation of the polysomal mRNA's for the fiber protein, the most distal gene (86.3-91.2) in the rightward transcription unit, had a fourth leader component "y" (78.6-79.1) in addition to the common tripartite leader sequences. Less frequent fiber RNA species had a fifth "x" (79.6-77.3) and/or sixth "z" (84.7-85.1) leader segment. mRNA transcribed at late times from the L strand between coordinates 15.0 and 11.0 had a single component leader located 200 bases from the main body of the gene (coordinates 16.1-15.7). This leader originated from a site close to the first leader component (16.5-16.6) of the R-strand messages. (38 refs)

6965 **Enucleated Cells as an Approach to the Study of Cytoplasmic Events in Adenovirus Replication (Meeting Abstract).** (Eng) Gayathri, A. K. (Dept. Human Genetics, Yale Univ., New Haven, CT); Hodge, L. D. *Cell Biol* 79(2, part 2): 402a; 1978. (no refs)

6966 **A Supplementary Adenoviral Leader Sequence and its Role in Messenger Translation.** (Eng) ...nn, A. R. (Cold Spring Harbor Lab., P.O. Box 100, Cold Spring Harbor, NY, 11524); Mathews, M. B.; Chow, L. T.; ...brook, J. *Cell* 15(2): 511-526; 1978.

The population of adenovirus 2-simian virus 40 (Ad2-SV40) hybrid messenger RNA's (mRNA's) generated late in the infection of HeLa cells with the Ad2-SV40 hybrid Ad2+ND1 dp2 was studied biochemically and by electron microscope heteroduplex analysis. During the course of lytic infection with the hybrid, a viral mRNA was synthesized whose 5' end included the start of the Ad2 fiber gene and whose 3' end lay in SV40 sequences. Two forms of this mRNA existed in the cytoplasm of HeLa cells at late times of infection. Both forms contained the conventional tripartite leader present on "late" Ad2 mRNA's, but they differed in the presence or absence of a fourth leader sequence consisting of 180 nucleotides of Ad2 sequences located 2,000 nucleotides upstream from the gene. Both forms of the hybrid mRNA were isolated by fractionation of total poly(A)-containing Ad2+ND1 dp2 late cytoplasmic RNA in methylmercuric hydroxide-agarose gels. It was established that the presence or absence of the fourth leader sequence had little or no influence on the ability of the messenger to direct the synthesis of the 23,000- to 24,000-dalton protein(s) in a cell-free translation system derived from rabbit reticulocytes. (23 refs)

6967 **Activation of RNA Particles from Human Tumors by Human Adenovirus.** (Eng) Winters, W. (Univ. Texas Health Science Center, San Antonio, TX); ...ri, A.; Sykes, J. A.; O'Toole, C. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 279-292; 1977.

Cultures of various human tumor and normal cells were grown in standard medium and inoculated with 5-500 plaque-forming units/cell of stock adenovirus 5 (Ad5). Cells of 15/20 sarcomas of different histologic types, 9/10 melanomas, and those of all breast, cervical, and squamous cell carcinomas examined readily supported complete Ad5 replication. The cells contained membrane-enclosed particles, and culture supernatants contained RNA particles with specific DNA polymerase activities. The normal embryonic, fetal, newborn, and adult cells also supported Ad replication.

Cell-free fluids of untreated or Ad-infected permissive human tumor and normal cell cultures had no DNA polymerase activities. Cell-free fluids of nonpermissive human tumor cells contained DNA polymerase activities at 3-9 days following Ad5 infection. The activation of detectable DNA polymerase-containing particles was dose- and time-dependent. Additional studies designed to label specifically the DNA and RNA in replicate cultures revealed that cell-free supernatants contained ³H-uridine-labeled particles with densities of 1.14-1.18 g/ml in equilibrium sucrose density gradients. DNA polymerase activities were detected within this range of density in fractions taken from these gradients. Heat-inactivated and antiserum-neutralized Ad5 did not infect or activate human tumor cells. The results suggest that infectious viruses with which humans normally have contact could, under proper host conditions, play a role in activating latent viruses associated with human tumors. (23 refs)

78-6968 **Defective Virions of Simian Adenovirus 7 Produced by One Passage at High Multiplicity of Infection.** (Eng) Price, J. A. (Dept. Microbiology, Univ. Massachusetts, Amherst, MA, 01003); Norkin, L. C. *Virology* 91(1): 194-197; 1978.

An assay for the rapid microtitration of simian adenovirus 7 (SA7) particles capable of inducing an adenovirus cytopathic effect (rounded, swollen cells) is described. Utilizing this assay for cytopathic particles, it was found that only one passage of standard SA7 at a high multiplicity of infection (moi) resulted in virus stocks containing an excess of noninfectious cytopathic particles. These stocks also contained an excess of defective virions that retained the ability to enhance the replication of human adenovirus type 5 during mixed infection of monkey cells. SA7 stocks produced on two simian virus 40-transformed simian cell lines contained larger fractions of these defective virions than stocks produced on normal cell lines. Autointerference activity was not observed in SA7 stocks containing an excess of defective cytopathic particles, and there were no detectable levels of shortened viral genomes. The results indicate that SA7 stocks containing an excess of defective particles are readily produced during only one passage at high moi. There need not be a drop in viral yields during subsequent passages to indicate that this phenomenon is occurring. (20 refs)

78-6969 **Phosphorylation of Chromatin- and Ribosome-associated Proteins in Cells Transformed by Adenovirus, Murine Sarcoma Virus, and Methylcholanthrene.** (Eng) Segawa, K. (Inst. Medical Science, Univ. Tokyo, Minato-ku, Tokyo 108, Japan); Oda, K.; Yuasa, Y.; Shiroki, K.; Shimojo, H. *J Virol* 27(3): 800-808; 1978.

The patterns of phosphorylation of chromatin- and ribosome-associated proteins in cells transformed by adenovirus

type 12 (Ad12), murine sarcoma virus (MSV), or methylcholanthrene (MC) were studied. A highly phosphorylated 90,000-dalton (90K) protein was detected in chromatin and ribosomes from Ad12-transformed WJ1, GY1, CY1, and WY3 cells; MSV-transformed H-NRK, K-BALB/3T3, and XC cells; MCA-transformed TCL15 mouse cells; and simian virus 40-transformed T22 monkey cells. The phosphorylated 90K protein was not detected in untransformed NRB, 3Y1, NRK, MV1Lu, or C3H.10T1/2 cells or in S+L-Mi mink cells transformed by Moloney MSV. The S+L-Mi cells showed rapid reversion to a normal morphology after several subcultures while continuing to contain rescuable MSV. The pattern of phosphorylation of proteins other than 90K in these cells resembled that of transformed rather than untransformed cells. Some 90K proteins became detectable in some untransformed cells that were stimulated to grow to the high density characteristic of transformed cells. Peptide map analysis indicated that the 90K protein of the transformed cells was similar irrespective of species of cell or transforming agent. The data suggest that the alteration in the pattern of endogenous protein phosphorylation in the chromatin and ribosomes is closely related to the transformed state of cells. (20 refs.)

- 78-6970 Characterization of Adenovirus Type 12 Tumour Antigen Produced in Chick Fibroblasts.** (Eng) Pusztai, R. (Inst. Microbiology, Univ. Medical Sch., H-6720 Szeged, Hungary); Tarodi, B.; Beladi, I. *J Gen Virol* 41(2): 433-436; 1978.

The type-specific tumor (T) antigen produced in adenovirus 12 (Ad12)-infected nonpermissive chick fibroblasts and permissive HEp-2 cells was characterized by immunoprecipitation with normal hamster serum and anti-T serum and sodium dodecyl sulfate-polyacrylamide gel electrophoresis. The anti-T serum precipitated a polypeptide with an apparent mol wt of 50,000 and a smaller polypeptide of mol wt approx 11,000 from the infected HEp-2 and chick cells extracts. There was a considerably lower amount of the larger polypeptide in the chick cells than in the HEp-2 cells. These polypeptides were also obtained in the presence of 50 µg/ml cytosine arabinoside, but they were not observed when normal hamster serum was used or when mock-infected chick fibroblast extract was precipitated by anti-T serum. It is concluded that the size of the major component of Ad12-specific T antigen is 50,000 daltons. (8 refs)

- 78-6971 Poliovirus Single-stranded RNA and Double-stranded RNA: Differential Infectivity in Enucleate Cells.** (Eng) Detjen, B. M. (Dept. Biology, Washington Univ., St. Louis, MO, 63130); Lucas, J.; Wimmer, E. *J Virol* 27(3): 582-586; 1978.

The ability of single-stranded poliovirus virion RNA (PV-RNA) and double-stranded poliovirus replicative form

(RF) RNA to replicate in enucleated mouse L cells was investigated. Monolayers containing >99% enucleated L cells were prepared by cytochalasin B (10 µg/ml) treatment in conjunction with centrifugation. Control nucleate cells were prepared by similar treatment with cytochalasin B and omission of the centrifugation step. Virion RNA replicated successfully in the absence of the cell nucleus, but RF RNA did not produce any detectable progeny in enucleated cells. The results show that the nucleus is involved in RF RNA infection, presumably in the early stages. (47 refs)

- 78-6972 Poly(C)- and Poly(G)-rich Tracts in the RNA of Oncornaviruses.** (Eng) Phillips, L. A. (Lab. Viral Carcinogenesis, NCI, Bethesda, MD, 20014); Pang, R. H. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 311-319; 1977.

Evidence is given for the presence of poly(C)- and poly(G)-rich tracts in both untreated and RNase T₁- and RNase C-treated [dimethyl-³H] viral RNA of avian, murine, hamster, feline, and primate oncornaviruses (OV's) as detected by polynucleotide agarose affinity columns. This finding demonstrates the universality of these tracts in the RNA of leukemia and sarcoma viruses. The tracts may play a genomic stabilizing role in OV RNA, and they may also play a role in the transcription of OV RNA. Poly(C) may be involved in the replication of some other viruses, as encephalomyocarditis virus-infected cells contain a viral specific polymerase that can use poly(C) homopolymers as templates, and poly(C) homopolymers can also serve as templates for Q β replicase. If it can be determined that OV RNA contains specific polynucleotide sequences or tracts that are not present in nonviral messenger RNA (mRNA) of human cells, it may be possible to make sequence-specific drugs against the unique viral-specific polynucleotide sequences or tracts to block transcription of viral RNA into viral DNA by reverse transcriptase and to block translation of viral RNA into viral proteins by the cellular ribosomes, while not affecting the translation of cellular nonviral mRNA into cellular proteins. Thus, a possible molecular approach to cancer therapy may become a reality. (27 refs)

- 78-6973 DNA Polymerase Reactions in Human Bladder Tumor Cells in Culture: Characterization of the RNA-associated Product.** (Eng) Davis, A. R. (Dept. Microbiology and Immunology, Univ. California Sch. Medicine, Los Angeles, CA, 90024); Vayuvegula, B. S.; Nayak, D. P. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Preven-

and Detection of Cancer (New York, NY): Vol. 1(part 193 pp.; 17-26; 1977.

RNAse-sensitive endogenous DNA polymerase activity in the cytoplasmic pellet fraction of cultured T-24 derived from a human transitional cell carcinoma of the bladder was partially characterized. Pretreatment of the cytoplasmic pellet with RNAse reduced the activity threefold, and the reaction was inhibited 80% by actinomycin D (AD). The reaction kinetics were linear for 5 min only. Analysis of the DNA product of 5-min reactions, prepared in the presence of Cs₂SO₄, by equilibrium-density gradient centrifugation revealed a nucleic acid product migrating as 10%-47% RNA (density 1.63) and 30%-70% DNA (density 1.39-1.42). Less than 1% of the product prepared under the same conditions from T-24 nuclei or mitochondria or from the cytoplasmic fraction of WI-38 cells, a normal human diploid cell line, migrated as RNA. The label in the RNA region of the gradients no longer migrated as RNA after treatment with the total product with high- and low-salt RNAse or S-1 nuclease. Under conditions in which hybrids are maintained with high-salt RNAse or S-1 nuclease, small amounts of material were seen migrating in the hybrid region of the gradients. The ³H-DNA recovered from the RNA region of the gradients was partially resistant to S-1 nuclease. Thus, part of the RNA-associated DNA product was a RNA-DNA hybrid and part was single-stranded DNA. Known RNA tumor viruses catalyze the transcription of RNA to DNA via reverse transcriptase, with early products being RNA-DNA hybrids. The DNA polymerase reaction product migrating as DNA and other properties of the reaction indicate that reverse transcriptase is present in the T-24 cells. (31 refs)

6974 **Ultrastructural Study of Virus-like Particles in Chinese Hamster Lung Cells.** (Eng) Calafat, J. (Dept. Virology, The Netherlands Cancer Institute, 108 Saristraat, Amsterdam, Netherlands); Hilken, J. G. *J Gen Virol* 41(2): 417-420; 1978.

Virus-like particles found in the E36 cell line derived from Chinese hamster lung cells are described. Two kinds of particles were found: viruslike particles in the cytoplasm associated with the centrioles, and extracellular viruslike particles resembling the immature and mature forms of the oncornaviruses but different from the classical B- and C-type particles. Apparently, the intracytoplasmic, budding, immature, and mature particles noted were different stages in the morphogenesis of the same virus. No major C- or B-type virus proteins were detected, indicating that there was no immunological relationship between the virus characterized and the oncornavirus from mammals or mouse B-type virus. (12 refs)

6975 **Transformation of Normal Bone-Marrow Cells by a Leukaemic Cell Line Associated with a**

Presumptive New Human Virus. (Eng) Karpas, A. (Dept. Haematological Medicine, Univ. Cambridge Clinical Sch., Hills Road, Cambridge, England); Wreghitt, T. G.; Nagington, J. *Lancet* 2(8098): 1016-1019; 1978.

Experiments were carried out to determine whether the intracytoplasmic leukoviruslike bodies found in the cells and culture medium of a T-cell lymphoblast line established from a leukemic patient could transform normal human bone marrow cells in vitro. The leukemic cell line (241 cells) was initiated from the buffy coat of a man with terminal acute monocytic leukemia superimposed on a longstanding non-Hodgkin's lymphoma. When the cells were lethally irradiated and added to normal bone marrow cells from a female donor, the latter were transformed into continuous replication, and they produced the same types of viruslike particles. The new cell line (P2-241), which has been in continuous culture for 4 mo, consists of female B lymphoblasts quite distinct from the male T-cell line 241. Sectioned cells of the nonirradiated line 241 and marrow line P2-241 contained two types of round viruslike particle in the Golgi region, one approx 50 nanometers (nm) and the other 70 nm in diameter. Formation of 50-nm particles by intracytoplasmic budding was observed. It appears that marrow cells are transformed in vitro to continuous replication by the 241 cell line. (10 refs)

78-6976 **Reverse Transcriptase Activity in Extracts of Human Prostatic Tissues.** (Eng) Job, L. (Dept. Medical Viral Oncology, Roswell Park Memorial Inst., Buffalo, NY); Carter, W. A.; Arya, S. K. *Oncology* 35(5): 202-205; 1978.

Three types of human prostatic tissue (normal, hyperplastic, and adenocarcinoma) were examined for particle-associated reverse transcriptase (RT) activity. The activity was isolated by banding the tissue extract in an equilibrium sucrose density gradient followed by phosphocellulose chromatography of the lysate of material banding at 1.14-1.20 g/ml. It was characterized by its utilization of poly(Cm), which is a specific template for oncornaviral RT, and by its inhibition by selective inhibitors of viral RT, such as poly(Am) and poly(I). The activity did not utilize synthetic DNA templates, such as poly(dA):oligo(dT) and poly[d(A-T)], which is consistent with the interpretation that the DNA polymerase activity isolated from prostatic tissue is a RT. Of the tissues examined, 1/4 normal, 5/6 hyperplastic, and 2/2 adenocarcinoma specimens contained viruslike RT activity. These results suggest that oncornaviruses may be associated with the human prostate. (18 refs)

78-6977 **Inclusion Bodies Containing Virus-like Particles in Acute Lymphocytic Leukaemia of Childhood.** (Eng) Smith, H. (Dept. Pathology, Royal Brisbane Hosp., Queensland 4029, Australia). *Leuk Res* 2(2): 133-140; 1978.

Lymphoid cells from 28 children with acute lymphocytic leukemia (ALL) and from three control groups (children with acute nonlymphocytic leukemia or nonleukemic malignancies and normal children) were examined for the presence of distinctive cytoplasmic inclusions. Light microscopy revealed such inclusions (0.7-7.6 μm) in 12/28 ALL patients. The inclusions tended to be close to the nucleus or were attached to it by a stalk. Inclusions were not seen in nonlymphoid cells; they were more frequent in leukemic than in nonleukemic lymphocytes, and more frequent in marrow than in blood cells. In all other patients with inclusions, they were found only in leukemic cells. Inclusions were more frequent in necrobiotic cells than in viable lymphoid cells in blood but not marrow of one patient. Electron microscopy showed that the inclusions are enclosed particles with a membrane usually 10-30 nanometers (nm) thick but in some cases up to 100 nm thick. Budding from the inside of the enclosing membrane was suggested by connection between it and some particles and by clustering of particles around some villi. Inclusions similar to those in ALL were seen rarely in normal lymphocytes and in acute nonlymphocytic leukemic cells, but they were seen often in nonleukemic malignancies. Profusely populated groups (≥ 20 particles in a profile) and villous intrusions of the enclosing membrane occurred only in the ALL

patients. The particles do not resemble known viruses, but some characteristics suggest that they have a viral nature. (23 refs)

See also:

*(Rev.): 78-6610, 78-6616, 78-6620, 78-6632, 78-6639, 78-6661, 78-6669, 78-6670, 78-6671, 78-6672, 78-6673, 78-6674, 78-6675, 78-6676, 78-6677, 78-6678, 78-6679, 78-6680, 78-6682, 78-6683, 78-6684, 78-6685, 78-6686, 78-6697, 78-6978, 78-6982, 78-6983, 78-6984, 78-6985, 78-6989, 78-7005, 78-7006, 78-7008, 78-7015, 78-7016, 78-7017, 78-7018, 78-7021, 78-7023, 78-7024, 78-7029, 78-7034, 78-7036, 78-7038, 78-7045, 78-7050, 78-7065, 78-7068.

*(Path.): 78-7073, 78-7082, 78-7084, 78-7101, 78-7118, 78-7124.

IMMUNOLOGY

5978 **A Serum Inhibitor of Cell Mediated Immunological Functions in Healthy Mice and Possible Relations with C-Type Viral Expression and Spontaneous Plasmia.** (Eng) Krueger, J. G. (Thorman Cancer Res. Lab., City Univ., San Antonio, TX, 78284); Moyer, R. C. *In: 3rd International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1 (part 1) 193 pp.; 409-427; 1977.

... suggesting the presence of an inhibitor of cell-mediated immunological functions in the serum of healthy DBA/2J mice possessing latent C-type leukemia virus are presented. The phytohemagglutinin responsiveness (PHAR) of peritoneal lymphocytes in a whole blood culture was measured. ³H-thymidine uptake in nine inbred strains of mice. High PHAR was seen in different strains, even though lymphocytes were taken from 14-wk-old sex-matched mice. High responders showed no C-type virus expression; mice of the low responder strains showed C-type virus expression throughout their lives. The low PHAR of these strains was investigated further using low-responder DBA/2J mice plus high-responder C57BL/6J mice for comparison. Low responsiveness of DBA/2J mice was not related to dose-response factor, to a time shift in response to PHA, to age. Splenic lymphocytes showed good PHAR, ruling out the possibility of an intrinsic cellular defect. Low responder serum reduced the PHAR of syngeneic splenic lymphocytes, and splenic lymphocytes failed to incorporate PHA actively in the presence of this serum. C57BL/6J mice showed no significant inhibiting effect on allogeneic syngeneic lymphocytes. C57BL/6J serum increased the response of C57BL/6J splenic lymphocytes to tumor-associated antigens from a syngeneic mammary carcinoma by 100%. DBA/2J serum prevented the response of DBA/2J splenic lymphocytes to syngeneic tumor-associated antigens. Ability of lymphocytes to perform cell-mediated functions may be inhibited by factors in the serum of healthy mice. (22 refs)

5979 **Influence of a Mixture of Chemical Protectors on the Lymphoid Regeneration of Bone Marrow and Thymus in Irradiated Mice.** (Eng) Delrez, M. (Laboratoire d'Anatomie Pathologique, Institut de Pathologie, Université de Liege, B-4000 Liege, Belgium). *Experientia* 34(9): 1122-1222; 1978.

... effect of chemical radioprotectors on the lymphoid peak in bone marrow and on the thymic cell population was deter-

mined in female C57BL mice. At 25 min before x-irradiation (400 or 650 R), the mice received 15 mg glutathione po; 10 min later, they received 8.5 mg cysteine ip followed 5 min later by ip injection of a mixture of 1.4 mg 2-β-aminoethylthiouronium, 3 mg cysteamine, and 0.65 mg 5-hydroxytryptamine. Administration of the chemical protectors without subsequent irradiation had little effect on the bone marrow but induced a significant decrease in thymocyte number. Administration of the protectors before irradiation permitted a better bone marrow repopulation and reduced late thymic atrophy (a condition that precedes the appearance of thymic lymphomas), compared with the effects of the radiation in nonprotected animals. The lymphoid peak was considerably increased in the radioprotected animals. Thus, under conditions that were previously shown to reduce the development of lymphomas, a lymphoid rebound appeared in the bone marrow. (8 refs)

78-6980 **Electronic Cell Volume Analysis of Growth and Rejection of EL-4 Ascites Tumor Cells.** (Eng) Zucker, R. M. (Papanicolaou Cancer Res. Inst., P.O. Box 236188, Miami, FL, 33123); Harari, J.; Finley, J.; Kambour, M. *Am J Vet Res* 39(10): 1710-1712; 1978.

The in vivo growth of EL-4 leukemia cells in the peritoneal cavities of syngeneic (C57BL/6) and allogeneic (DBA/2) mice was monitored daily by electronic cell volume and cell number analysis, cell morphology, and labeling index. In C57BL/6 mice, tumor cell number increased from 2×10^5 to 8×10^5 , cell volume decreased from 440 to 360 μm^3 , and the labeling index decreased from 57% to 20% from days 5 to 9 after inoculation. The volume decrease was correlated with an increased cell cycle time and a decreased proliferating fraction. During the later stages (10-15 days) of EL-4 ascites tumor growth in these mice, the cell number remained constant but the cells continued to grow and divide. In DBA/2 mice, rejection of tumor cells began on day 7 after inoculation; cell number decreased from 4.6×10^5 to 2.5×10^5 cells, modal population volume from 370 to 355 μm^3 . The degree of rejection was determined by electronic index; these variables were not uniform among the mice. Concomitant with the decrease in cell number and labeling index seen between days 7 and 8, the number of lymphocytes, macrophages, and dead tumor cells increased. The size of the lymphocytes was 240 μm^3 , and that of the macrophages and dead tumor cells was 640-1,280 μm^3 . Lymphocyte volume decreased during the rejection process until it was 110 μm^3 at complete rejection. These results demonstrate that electronic cell volume analysis can be used to study the influx of different cell types during the rejection as well as the growth of tumor cells. (12 refs)

78-6981 Complete Tumor Inhibition by a Physiological Factor in Cell-Free Tumour Ascites. (Eng)

Kapp, J. F. (Schering AG, Research Labs., Mullerstrasse 170, 1000 Berlin 65, W. Germany). In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society. Symposium held in Edinburgh, September 12-14th, 1977.* Univ. Edinburgh. (Edinburgh, Scotland): pp. 285-291; 1977.

Anti-inflammatory activity was demonstrated in mouse and rat cell-free tumor ascites fluids. Ascites from Ehrlich (including diploid and tetraploid), L1210, Krebs, and Yoshida mouse tumors and from Yoshida hepatoma, Yoshida sarcoma, and Zajdela hepatoma rat tumors were tested in the carrageenan NMRJ mouse paw edema test. Mouse tumor ascites inhibited the inflammation reaction from 40.5% to 58.5%, rat tumor ascites from 28% to 45%, but normal sera from either species were inactive. There were no inflammatory cells in the site of the inflammatory stimulus. With respect to antitumor activity, tumor growth was inhibited completely when 1 ml tumor ascites was injected ip 24 hr before ip injection of 1×10^5 tumor cells. An inoculum of 3×10^5 DBA/1J mammary adenocarcinoma cells injected sc into NMRJ mice and grown to 5 mm in diameter underwent continuous regression when 1 ml L1210 ascites fluid was injected. Likewise, human HeLa cells (1×10^5) injected ip into nu/nu mice were inhibited by injection of the mice with 1 ml Yoshida ascites 24 hr previously, demonstrating interspecies protection. Because sensitized lymphocytes from spleens and lymph nodes did not demonstrate cytotoxic activity in a ^{51}Cr -release assay, inhibition of tumor growth apparently occurs by stimulation of nonspecific host defense mechanisms. (9 refs)

78-6982 Anti-viral Activity Induced by Culturing Lymphocytes with Tumor-derived or Virus-transformed Cells. Identification of the Anti-viral Activity as Interferon and Characterization of the Human Effector Lymphocyte Subpopulation. (Eng) Trinchieri, G. (Swiss Inst. Experimental Cancer Res., Epalinges, CH 1066, Switzerland); Santoli, D.; Dee, R. R.; Knowles, B. B. *J Exp Med* 147(5): 1299-1313; 1978.

The viral inhibitor(s) released in the supernates of mixed cultures of mouse or human lymphocytes and certain tumor-derived or virus-transformed mouse or human cell cultures was characterized. The inhibitor thus induced proved to be a protein that was active against a variety of unrelated viruses and was not toxic for cells. It did not directly inactivate viruses but inhibited viral replication via an intracellular mechanism that involved RNA and protein synthesis by the cells. The inhibitor was identified as interferon, which was contained in at least two molecular species, of approx 25,000 and 45,000 daltons, respectively. The supernates of the mixed cultures also displayed an anticellular activity and an ability to inhibit DNA synthesis and cell multiplication. The antiviral and anticellular activities were positively correlated in supernates from various mixed cultures and in partially purified

preparations. The human cell population involved in interferon production was composed primarily of Fc receptor-positive, surface immunoglobulin negative, non-T-cell lymphocytes. The ability of certain cell lines to induce interferon was significantly associated with tumor origin or with in vitro transformation by certain viruses (Epstein-Barr virus, murine sarcoma virus). (56 refs)

78-6983 Anti-viral Activity Induced by Culturing Lymphocytes with Tumor-derived or Virus-transformed Cells. Enhancement of Human Natural Killer Cell Activity by Interferon and Antagonistic Inhibition of Susceptibility of Target Cells to Lysis. (Eng) Trinchieri, G. (Swiss Inst. Experimental Cancer Res., Epalinges, CH 1066, Switzerland); Santoli, D. *J Exp Med* 147(5): 1314-1333; 1978.

The effects of the interferon produced in mixed cultures of human lymphocytes and human tumor-derived or virus-transformed lines on the two cell populations were studied. Interferon preparation increased by severalfold the nonspecific cytotoxicity of human lymphocytes, the effector cells being cytotoxic for any target cell line tested and the enhanced cytotoxicity lasting for 2-3 days. The activity of spontaneous (natural) killer lymphocytes was not separable from the cytotoxic activity induced by the interferon preparations, suggesting that the interferon enhanced the activity of the natural killer cells rather than generating a new population of effector cells. In the mixed cell cultures in which interferon was produced, the interferon mediated an enhanced cytotoxicity representing up to 70%-90% of the observed cytotoxicity. The effect of interferon on the target cells was antagonistic to its effect on the lymphocytes, the susceptibility of various cells to cell-mediated lysis being decreased and in some cases almost completely suppressed after pretreatment with interferon. Interferon rendered target cells resistant to natural killer cells via an intracellular mechanism requiring RNA and protein synthesis. Virus-infected cells and most tumor cells were not similarly protected by interferon. It is suggested that interferon might render the natural killer cell system an inducible selective defense mechanism against tumor and virus-infected cells. This might occur by stimulation of efficient nonspecific cytotoxic cells and by protection of normal cells from lysis. (56 refs)

78-6984 Increases and Perturbations in Thymocytes Forming Spontaneous Rosettes During Viral and Chemical Oncogenesis in Hamsters. (Eng) Leffell, M. S. (Clinical Micro Lab., North Carolina Memorial Hosp., Univ. North Carolina, Chapel Hill, NC, 27514); Coggin, J. H. *Clin Immunol Immunopathol* 11(3): 275-287; 1978.

Because spontaneous (E) rosette formation with sheep RBC has been linked to diseases that involve deficient cellular immunity including malignancies, thymocytes from strain LVG Syrian golden hamsters were tested for their ability to form

rosettes with sheep RBC. Preliminary studies revealed an unusual seasonal variation in the percentage of rosette-forming cells (RFC) (17.2% formation in winter, compared with 10.5% in later summer and fall). Further studies showed that, regardless of the seasonal variation, the number of RFC increased during viral and chemical carcinogenesis. Infection of neonate hamsters with simian virus 40 or adenovirus 13 resulted in a 50% or greater increase in RFC in thymuses of individual hamsters assayed as early as a 2 wk postinfection. Treatment of weanling hamsters with dimethylbenzanthracene resulted in significant increases in thymocyte RFC at the time of primary sarcoma appearance. The carcinogen-induced RFC increases appeared to be limited to a small subset of thymocytes that reacted in dextran-containing medium, compared with a larger population that bound sheep RBC in the presence of heat-inactivated, sheep RBC-absorbed serum. In the thymocyte subset, the increases in RFC were most marked after primary tumor induction. Passively transferred tumor cells had only a slight effect on RFC values. (19 refs)

78-6985 Immune Response of the Syrian Golden Hamster to SV40 Induced and Transplantable Tumors. (Eng) Goldrosen, M. H. (Dept. Surgical Oncology, Tumor Immunology Group, Roswell Park Memorial Inst., Buffalo, NY, 14263); Dent, P. B. *Fed Proc* 37(7): 2092-2095; 1978.

The effector mechanisms that are operative against the growth of simian virus 40 (SV40)-induced tumors were investigated. Cell-mediated immunity of progressor, postexcision, and immune peripheral blood lymphocytes was evaluated in vivo by the lymphocytotoxicity assay. At a ratio of 200:1, three types of peripheral blood lymphocytes caused a significant reduction in the number of H50 (SV40-transformed) target cells but did not reduce the number of hamster fibroblasts. No significant cytotoxicity was observed with these effector populations on BHK-21 cells, which do not carry SV40-associated antigens. Thus, cytolytic effector cells can be detected at all phases of progressive tumor growth and at all stages of tumorigenesis evaluated. However, the presence of these cells may not reflect functional in vivo tumor rejection. The effects of the three sera on in vitro lymphocytotoxicity were also investigated. The only situation in which serum blocking factors were noted was in animals that experienced a recurrence of a previously excised tumor. Serum blocking factors were not present in the early and middle stages of progressive tumor growth; therefore, they could not be an important factor in the critical early stages of tumor growth. Serum taken from hamsters shortly after inoculation with H50 cells but before the clinical appearance of the tumor demonstrated characteristics similar to those of serum taken from immunized hamsters. These sera augmented the cytotoxic reactivity of tumor immune lymphocytes in vitro and did not induce normal lymphocytes to specifically kill tumor cells. These findings suggest that progressive primary tumor growth may be due to a lack of serum arming and potentiating

activity rather than the presence of serum blocking factors. On the other hand, serum blocking factors may correlate with the presence of recurrent or metastatic disease rather than with progressive tumor growth alone. (25 refs)

78-6986 Studies of Effector Cell, Antibody, and Target Cell Interactions in Natural Cell-mediated Cytotoxicity. (Eng) Koide, Y. (Dept. Surgery, UCLA Sch. of Medicine, Univ. California, Los Angeles, CA, 90024); Kwok, R.; Takasugi, M. *Int J Cancer* 22(5): 546-551; 1978.

IgG antibodies bound to effector lymphocytes obtained from human peripheral blood through Fc receptors were observed to determine the specificity of natural cell-mediated cytotoxicity (NMC) against cultured target cells, including sheep RBC (SRBC), antilymphocyte serum-treated lymphocytes, and a lymphoblastoid cell line. When effector cells, which possessed natural antibodies specific for antigens on cultured cells but lacked IgG antibodies specific for antigens on SRBC, were incubated in IgG anti-SRBC, they became specifically cytotoxic to SRBC. As a result of this gained specific cytotoxicity, NMC and antibody-dependent CMC was diminished, suggesting that natural antibodies were displaced and Fc receptors on effector cells occupied by IgG anti-SRBC. It is concluded that treatment of effector cells with serum may result in increased or decreased cytotoxicity, depending upon the serum composition of specific antibodies. This type of modulation has important implications for antibody regulation of CMC (blocking and unblocking). (23 refs)

78-6987 Spontaneous and Antibody-dependent Cell-mediated Cytotoxicity by Human T Cell Subpopulations. (Eng) Gupta, S. (Memorial Sloan-Kettering Cancer Center, New York, NY, 10021); Fernandes, G.; Nair, M.; Good, R. A. *Proc Natl Acad Sci USA* 75(10): 5137-5141; 1978.

Human peripheral blood non-T cells, T cells, and their subpopulations (T_{μ} , T_{γ} , T_{ϕ} , T_{γ} -depleted cells, and T_{μ} -depleted cells) were assayed for their ability to mediate spontaneous lymphocyte-mediated cytotoxicity (SLMC) against K562 tumor cells and antibody-dependent cellular cytotoxicity (ADCC) against chicken RBC coated with antibody. (T_{μ} = T cells with receptors for IgM; T_{γ} = T cells with receptors for IgG; T_{ϕ} = T cells without receptors for IgM or IgG.) Non-T cells and purified T cells mediated both ADCC and SLMC. Depletion of T_{μ} cells did not influence these activities, whereas depletion of T_{γ} cells resulted in almost complete loss of both activities. Isolated T_{μ} cells and T_{ϕ} cells did not demonstrate significant ADCC or SLMC activity, and T_{γ} cells demonstrated strong SLMC activity but did not demonstrate significant ADCC activity. The data demonstrate that SLMC and ADCC activities in T lymphocytes are mediated by T_{γ} cell subpopulations. The two cytotoxic reactions may be mediated by two distinct subsets of T_{γ} cells or by a single effector cell type using two different mechanisms. (54 refs)

- 78-6988 Cytotoxic Activity of Guinea-Pig Lymph-Node Cells Stimulated In Vitro.** (Eng) Altman, A. (Dept. Cellular and Developmental Immunology, Scripps Clinic and Res. Foundation, 10666 N. Torrey Pines Road, La Jolla, CA, 92037); Rapp, H. J. *Int J Cancer* 22(5): 595-601; 1978.

Several stimulators of lymphocyte proliferation were tested for their capacity to induce strain 13 guinea pig lymph node cells (LNC) to become cytotoxic in vitro. The stimulating agents were 5% fetal calf serum; 5×10^5 M 2-mercaptoethanol; 1.6×10^5 virally induced, syngeneic, 543 sarcoma cells or various amounts of allogeneic (strain 2) mineral oil-induced peritoneal exudate cells, all treated with 60 μ g/ml mitomycin C; and 2.5 μ g/ml phytohemagglutinin (PHA) or concanavalin A. LNC cultured in the presence of these agents were cytotoxic to normal syngeneic fibroblasts and to syngeneic, (13F, 543), allogeneic (L1), or xenogeneic (human myeloid line K562) tumor cells, but not the PHA-induced strain 2 lymphoblasts. LNC had weak or no cytotoxic activity when freshly harvested from guinea pigs, indicating that acquisition of cytotoxicity was a result of events occurring in culture. Potentiation of cytotoxicity was accompanied by a marked proliferation of the cultured LNC; the combination of several stimulatory agents had an additive effect on the generation of cytotoxicity. (20 refs)

- 78-6989 Production of Migration Inhibitory Factor (MIF) by Human Leukocytes Following Exposure to Epstein-Barr Virus.** (Eng) Wilson, G. L. (Dept. Microbiology, Univ. Maryland, College Park, MD, 20742); Young, B. G. *Cell Immunol* 38(1): 147-156; 1978.

The ability of Epstein-Barr virus (EBV) to induce a cell-mediated immune response (CMI) in cultures of human WBC was investigated. Partially purified EBV, obtained from culture fluids of AV-1 cells, was inactivated by exposure to UV radiation. Inactivated virus was mixed with peripheral WBC from Hodgkin's disease (HD), infectious mononucleosis (IM), and malignant lymphoma patients as well as from normal individuals. Production of migration inhibitory factor (MIF), as measured by guinea pig macrophage migration inhibition (MMI), was used as an indicator of CMI response. Significant differences in MIF response were observed subsequent to exposure of the cells to EBV. WBC from patients in each of the disease categories tested exhibited greater MIF production than WBC from normal controls. Furthermore, there was a significant difference between the response of IM-derived cells and the response of HD- and lymphoma-derived cells. Serum from each subject was examined for immunoglobulin specific for EBV capsid antigen (anti-VCA). Although the majority of individuals within the disease categories tested had elevated anti-EBV serum titers, no correlation could be made between elevated anti-VCA titer and levels of MIF production. (22 refs)

- 78-6990 Effect of Host Inflammatory Response on the Tumorigenicity of a Mouse Melanoma (Meeting**

Abstract). (Eng) Freedman, V. H. (Rockefeller Univ., New York, NY); Calvelli, T. A.; Silagi, S.; Silverstein, S. C. *J Cell Biol* 79(2, part 2): 91a; 1978. (no refs)

- 78-6991 Do Tumors Escape Surveillance by Depression of Macrophage Inflammation?** (Eng) Norman, S. J. (Dept. Pathology, Univ. Florida Coll. Medicine, Gainesville, FL, 32610); Scharadt, M.; Sorkin, E. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 247-257; 1977.

The hypothesis that growing tumors compromise monocyte chemotaxis and decrease macrophage (MP) accumulation in inflammatory foci, thereby altering the ratio of MP to tumor cells, decreasing the probability for close contact of these cells, and escaping growth control by MP, was investigated. Nitrocellulose filter disks were inserted into the connective tissue of DA rats with and without transplanted 7,12-dimethylbenz(a)anthracene (DMBA)-induced tumors. Twelve days after transplantation, both groups showed a biphasic pattern of MP accumulation, but in tumor-bearers both the total number of cells and their rate of accumulation were diminished markedly. This depression in MP accumulation was approx 65% and it lasted for at least 5 days, despite the persistence of the inflammatory stimulus. Thus, tumor-bearing induced a generalized defect in MP inflammation by altering MP accumulation not only in the peritoneal cavity but also in inflamed connective tissue. The defect required the continued presence of the tumor, as it was rapidly corrected by surgical excision of the primary tumor. However, the defect usually occurred late in tumor growth, required a threshold number of tumor cells often exceeding 100 million, and did not exist when the animal possessed latent tumor cells that remained viable following surgical resection of the primary tumor. Thus, it is concluded that the defect in monocyte inflammation arises as a result of neoplastic progression and that it is not a factor that limits surveillance or contributes to the emergence of nascent cancer cells. (27 refs)

- 78-6992 Inflammation and Activation of Macrophage** (Eng) Fauve, R. M. (Unite d'Immunophysiologie Cellulaire, Institut Pasteur, 25 Rue du Dr. Roux, 7501 Paris, France); Hevin, M. B.; Dauge, M. C. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 258-263; 1977.

C57Bl mice were highly resistant to challenge with virulent *Listeria monocytogenes*, *Schistosoma mansoni*, and the Lewis lung carcinoma (LLC) when inflammatory reactions were induced previously at a distance from the site of infection by injecting the dorsal area sc with magnesium silicate embedded in calcium phosphate gel. When *L. monocytogenes* was

ected iv (10 LD₅₀), mice with the inflammation had 10³ and less organisms in spleen and liver on the third day than control mice. Similarly, mice with an inflammatory reaction are free of worms 45 days postinfection with 100 furcocercariae of *S. mansoni*. When 10⁶ LLC cells were injected into the rear left footpad 4 days before or 4 days after induction of the inflammatory reaction, tumor growth was delayed. No metastases were found in the lungs of treated mice 20 days after LLC inoculation. When mice received 10⁶ LLC cells in the sternal area one day before inflammation induction, metastases were delayed and there were many WBC and macrophages among the tumor cells. These results suggest that some sort of serum mediator acted on the pathogens, since these were never in physical contact with the inflammatory reaction. (10 refs)

6-6993 **Regulation of Macrophage Tumoricidal Function: A Role for Prostaglandins of the E Series.** (Eng) Schultz, R. M. (Lab. RNA Tumor Viruses, NCI, Bethesda, MD, 20014); Pavlidis, N. A.; Stylos, W. A.; Chirigos, M. A. *Science* 202(4365): 320-321; 1978.

The ability of several agents to interfere with macrophage activation by interferon (IF) preparations in vitro was measured using macrophages from male CD2F₁ mice. IF (1,000 units/ml culture medium) was added to the macrophage cultures for 16 hr. Agents tested for inhibitory action were added in 100-μl portions. Treated macrophages were tested for cytotoxicity to MBL-2 lymphoblastic leukemia cells. IF-activated macrophages suppressed leukemia cell growth at 48 hr by 67%. When prostaglandin E₁ or E₂ was added simultaneously with IF to resting macrophages or for 2 hr after treatment, the tumoricidal activity of the macrophages was inhibited. A prostaglandin concentration of > 10⁻⁷ M was necessary for macrophage suppression. Prostaglandin F₂ was ineffective at all concentrations tested when added simultaneously or after IF treatment. Dibutyl cyclic AMP and hydrocortisone also decreased macrophage tumoricidal activity. Subsequent treatment of IF-activated macrophages with dibutyl guanosine 3',5'-monophosphate (GMP), indomethacin, or colchicine did not influence macrophage tumoricidal function. The regulation of macrophage function by IF and prostaglandin E may occur through a balance in cyclic AMP- and cyclic GMP-associated signals. Direct evidence for this conclusion awaits corroboration of levels of cyclic nucleotides in the macrophages with the observed effects of the agents tested. (22 refs)

6-6994 **Striking Differences in Interaction Between Macrophages and Two Closely Related Tumor Lines.** (Eng) Kaplan, G. (Dept. Morphology, Inst. Medical Biology, Tromsø, Norway); Seljelid, R. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 347-363; 1977.

A murine methylcholanthrene-induced solid sarcoma (MSS), its ascites form (MAA), and a solid tumor line derived from it (MAS) were used to study the in vitro interaction of syngeneic mouse macrophages with the tumor cells from intraperitoneal transfers. After sc or ip injection into C₃H mice or their F₁ hybrids, MSS cells yielded solid tumors which metastasized efficiently. MAS cells injected sc yielded solid tumors and injected ip yielded nonmetastasizing ascites when injected ip. Ascites from the mice which developed them were grown in culture. Macrophages from the MAA cultures became activated as indicated by their morphology and ability to ingest complement-coated particles. The macrophages proliferated and inhibited tumor cell growth; macrophage proliferation then decreased as the tumor cells died. The presence of tumor cells was apparently required for macrophage activation, and the macrophages appeared to kill the tumor cells by nonphagocytic means. The MAS cells showed less macrophage activation, and tumor cell killing was not apparent. The data show a correlation between the ability of tumor cells to give rise to solid metastasizing tumors and the extent to which the tumor could be inhibited and killed in vitro by syngeneic macrophages; ie, the more aggressive tumor in vivo is less susceptible to macrophage activation and killing in vitro. (30 refs)

78-6995 **Antitumour Effects of Macrophages in BCG Contact Therapy of Rat Tumours.** (Eng) Hopper, D. G. (Cancer Res. Campaign Labs., Univ. Nottingham, Nottingham, England); Pimm, M. V. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* Univ. Edinburgh. (Edinburgh, Scotland): pp. 182-187; 1977.

Wistar/Nottingham rats were inoculated sc with 2 x 10⁶ sarcoma Mc7 or Mc40A cells or hepatoma D23 cells, alone or in admixture with 1 mg BCG, and the cellular contents of the resulting tumors were investigated. Sarcoma cells alone produced progressively growing tumors, and the sarcoma cell: macrophage (MP) ratio was 1:0.2 for the Mc7 cells at 22 days and 1:0.4 for the Mc40A cells at 17 days. BCG + sarcoma cells, however, produced small nodular resector tumors, and the ratios were 1:1.3 for the Mc7 cells at 22 days and 1:1.0 for the Mc40A cells at 17 days. Lymphocyte levels were similar in regressor and progressor tumors. Hepatoma D23 was not as susceptible to BCG contact therapy as the sarcomas. BCG + hepatoma cells produced tumors whose growth was progressive although retarded compared to that of tumors developing from injection of hepatoma cells alone. The hepatoma cell:MP ratio in the BCG-progressor tumors was 1:0.3 at 6 days; that in the progressor tumors developing from tumor cells injected alone was 1:0.1. These results suggest that for BCG to prevent the growth of 2 x 10⁶ tumor cells, sufficient MP must be present at the tumor site. This hypothesis is strengthened by the results of experiments in which the hepatoma cell-BCG inocula was enriched with 50% MP. An increase in MP numbers to the extent that the

hepatoma cell:MP ratio was 1:1 rendered the tumor cells susceptible to BCG contact therapy; ie, it prevented their growth completely. (33 refs)

- 78-6996 Stimulation of Tumor Growth in Mice by High Doses of BCG.** (Eng) Kaledin, V. I. (Inst. Cytology and Genetics, Siberian Dept. USSR Acad Sciences, Novosibirsk, 90, USSR); Kurunov, Y. N.; Matienko, N. A.; Nikolov, V. P. *J Natl Cancer Inst* 61(6): 1393-1396; 1978.

The effect of BCG vaccination on the growth of Krebs-2 carcinoma was studied in virgin female C57BL/6 (susceptible to BCG infection), CC57BR/M (resistant to BCG), and C3Hf (resistant to BCG, mammary tumor virus-free) mice. Animals were vaccinated at 2 mo of age with 0.05 mg live BCG. Thirty days later, 1 or 2 x 10⁵ Krebs-2 carcinoma cells were injected im and the animals killed 15 days later. The number of living BCG in C57BL/6 mice was 30 times higher than that in C3Hf mice, and the tumors in vaccinated C57BL/6 mice were 1.5 times heavier than those in control C57BL/6 mice. The tumor wt of BCG-vaccinated and control C3Hf mice did not differ, indicating that tumor growth was stimulated only in the BCG-sensitive strain. Various doses of BCG were given to C57BL/6 and CC57BR/M mice 14 days prior to Krebs-2 transplantation. When the BCG dose was sufficiently high (1.0 mg), tumor growth was stimulated in both mouse groups. Dead mycobacteria and inert particles (India ink) promoted tumor growth, suggesting that this effect may be due to the presence of large quantities of corpuscular particles which load the macrophages and render them incapable of guarding against the transplanted tumor cells. When Krebs-2 carcinoma was transplanted in CC57BR/M mice 80 days after BCG vaccination (when no viable BCG were detected in the spleen), tumor growth was inhibited 41.5%. These results indicate that the effects of BCG are based on the diversion of effector cells (macrophages) and on the activation of pools of these cells. High doses of BCG may be dangerous in human cancer immunotherapy, and patients predisposed to cancer should be revaccinated with BCG to maintain the effector cell pool at a high level of activation. (18 refs)

- 78-6997 The Effect of the Macrophage Cytotoxic Agent Iota Carrageenan on the Growth of Two Transplantable Murine Ascites Tumours.** (Eng) Pugh-Humphreys, R. G. (Dept. Zoology, Univ. Aberdeen, Tillydrone Ave., Aberdeen AB9 2TN, Scotland); Richardson, C. A.; Wilson, A. R.; Cruickshank, W. J. *In: The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 142-154; 1977.

The role of macrophages (MP) in the growth of ascites tumor allografts in C57BL10 mice was evaluated with the use of a

commercial preparation of the MP cytotoxic agent iota carrageenan (IC: Auby Gel X52). Groups of 10 mice were inoculated ip with 10 µg IC 24 hr before they were inoculated ip with threshold or subthreshold doses of Landschutz ascite carcinoma (LAC) or EL4 ascites lymphoma cells. In preliminary experiments, light and electron microscope studies of aspirated lymphoreticular cells at varying times after ip injection of IC alone revealed that IC was selectively cytotoxic for peritoneal MP but not for lymphocytes or polymorphonuclear WBC. Injection of the tumor cells alone elicited an inflammatory response within the peritoneal cavity, and host lymphoreticular cells, particularly MP and neutrophils, congregated around the tumor cells. In IC-treated mice, only the polymorphonuclear WBC appeared to congregate around the tumor cells. IC had a dose-dependent cytotoxic effect on LAC cells. Doses of 10 and 25 µg had no significant effect on tumor growth, but 50-200 µg reduced the percentage of mice that developed tumors and 500 µg inhibited tumor development completely. In contrast, IC had no deleterious effect on the viability of EL4 cells, even at high concentrations. IC potentiated the growth and development of EL4 lymphomas in the mice while simultaneously killing MP. Thus, MP play an important role in regulating the growth of the EL4 lymphoma in C57BL10 mice. The nature of the cytotoxic effect of IC on the LAC cells is being studied. Selective killing of MP may abrogate an important growth-controlling mechanism within the tumor-bearing host. (45 refs)

- 78-6998 The Effects of Altered Phagocytic Activity on Growth of Primary and Metastatic Tumour** (Eng) Sadler, T. E. (Urology and Transplant Unit, Dept. Surgery, Royal Postgraduate Medical Sch., Hammersmith Hosp., London, W12 0HS, England); Jones, P. D. E.; Castro, J. E. *In: The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 155-163; 1977.

Silica (Si), cortisone acetate (CA), trypan blue (TB), diethyl stilbestrol (DES), and *Corynebacterium parvum* were used in macrophage phagocytosis studies that determined an inverse relationship between phagocytic index and metastatic behavior of tumors, but no apparent relationship between primary tumor growth (as measured by size) and phagocytosis. Si given iv (0.1 ml) and ip (0.5 ml) 3 hr later on days 4 and 9 after sc implantation of the Lewis lung carcinoma in the lower flank of C57BL/10 ScSn mice depressed the phagocytic index and significantly increased pulmonary metastases above control values. Similar results were obtained with CA (0.1 ml sc given on days 4 and 11 after tumor transplantation. DES (0.1 ml sc) given on days 0-4 after tumor inoculation and TB (0.4 ml ip 24 hr and 0.1 ml ip 3 hr prior to tumor implantation had little effect on the phagocytic index values, neither compound significantly altered the number of metastases. *C. parvum* (0.2 ml iv) given at tumor implantation caused a significant rise in the phagocytic index and a significant reduction in the number of metastases and overall tumor growth. Si

and DES also decreased tumor growth, but TB had no parent effect. (30 refs)

78-6999 Depression of Mononuclear Phagocyte Function in C57Bl Mice Bearing Lewis Lung Carcinoma. (Eng) Otu, A. A. (Dept. Surgery, St. Margaret's Hospital, Calabar, Nigeria); Russell, R. J.; Wilkinson, P. C.; Hevin, M-B. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 281-284; 1977.

To study the influence of Lewis lung carcinoma on macrophage function in C57Bl mice, the effects of growing tumors, tumor cell extracts, and sera from tumor-bearing animals on clearance of colloidal carbon from the circulation, chemotaxis of peritoneal macrophages in vitro, and the in vitro macrophage colony formation capacity of bone marrow derived cells were investigated. During the first 72 hr after tumor implantation, there was a pronounced depression in all these functions. For 2 wk thereafter, all functions were normal or above normal, but all were again severely depressed when the tumors metastasized and the mice had entered the terminal stage. Sera or tumor supernatant from tumor-bearing mice depressed carbon clearance in normal mice and inhibited the in vitro locomotion of peritoneal macrophages in normal mice. The data suggest that the tumor cells used a product which circulated in the blood and which was inhibitory to macrophages. (14 refs)

78-7000 Promotion of L1210 Tumour Growth by Macrophages. (Eng) Huget, R. P. (Dept. Microbiology, Lab. Immunology, Univ. Ulm, 7900 Ulm, W. Germany); Bitter-Suermann, D.; Opitz, H. G. *Z Immunitätsforsch* 154(5): 441; 1978.

To study the role of macrophages in tumor immunity, 10^4 L1210 mouse lymphoma cells derived from the peritoneal cavity or from in vitro cultures were injected sc into DBA/2 mice; and tumor development was monitored. Mice injected with peritoneal exudate cells developed tumors within 9 days. Tumor incidence was reduced and tumor development delayed by several days when in vitro derived cells were injected or when adherent cells were removed from the peritoneally derived cells prior to injection. Addition of adherent cells in high numbers to in vitro derived L1210 cells accelerated tumor growth, as did the addition of syngeneic liver cells. Differences in tumor development among the various groups were statistically significant ($p < 0.05$). The tumor-promoting effect of the adherent cells in the recipient was apparently not due to an increased proliferative capacity of the tumor cells. The data suggest that, under certain conditions, a cell population with the functional properties of macrophages is able to promote tumor development,

probably due to its nonspecific effect on the microenvironment of the growing tumor. (14 refs)

78-7001 Toxic Effects of Tumour Cells on Macrophages. (Eng) Fauve, R. M. (Unité d'Immunophysiologie cellulaire, Institut Pasteur, 25 rue du Dr. Roux, 75015 Paris, France); Hevin, M-B. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 264-270; 1977.

The substance(s) responsible for the toxic effects of Lewis tumor cells on peritoneal macrophages from C57Bl/6 and NCS mice was studied. Bradykinin in concentrations of 10^{-7} to 10^{-8} M increased the spreading of peritoneal exudate macrophages in vitro. In contrast, the spreading of macrophages was decreased in the presence of $0.1 \mu\text{g/ml}$ or more (up to $100 \mu\text{g/ml}$) of the active fraction obtained from Lewis tumor cell supernatant (AP). No decrease was observed when the macrophages were exposed to the same concentrations of the 10^3 - to 10^4 -mol wt fraction from fetal mouse kidney cells. Also, no decrease was observed when the peritoneal cells were incubated with bradykinin and then with AP. No increase in spreading was observed when the peritoneal cells were incubated first with AP and then with bradykinin. The data suggest that tumor culture supernatants inhibit the fixation of bradykinin to macrophage membranes. (27 refs)

78-7002 The Role of Complement Cleavage Products in Killing of Tumor Cells by Macrophages. (Eng) Schorlemmer, H. U. (Inst. Medical Microbiology, Johannes-Gutenberg-Univ., 6500 Mainz, W. Germany); Hadding, U.; Bitter-Suermann, D.; Allison, A. C. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 68-77; 1977.

Macrophages (MP) recovered from the peritoneal cavities of Swiss mice were cultured with pure guinea pig complement cleavage product C3b ($20 \mu\text{g/ml}$) or lipopolysaccharide (LPS: $25 \mu\text{g/ml}$), and the effect of these substances on the capacity of MP to lyse tumor cells was determined. ^{51}Cr -labeled L 5178 YE lymphoma cells were added to the MP 24 or 48 hr after culture initiation, and both cell types were cultured together for a further 16-18 hr. Unlike unstimulated control cultures, MP incubated with C3b or LPS lysed all or nearly all of the tumor cells. The MP also lysed P-815 mastocytoma cells and human lymphoblastic cells. The mastocytoma cells appeared to be the most sensitive to the lytic effect. This effect was C3b dose- and time-dependent, and it was considerably reduced in the presence of serum. The tachyphylaxis and cross-desensitization observed with reference samples of C3a identified the active agent in the media

of stimulated MP as C3a. Further evidence that the lytic activity is due to C3a comes from experiments using specific antiserum against C3a, which is presumably released from C3b-stimulated MP. These results show that C3b is a potent naturally occurring stimulator of MP activities, including tumor cell lysis. (11 refs)

- 78-7003 Tumor-induced Immunosuppression.** (Eng) Friedman, H. (Dept. Microbiology and Immunology, Albert Einstein Medical Center, Philadelphia, PA, 19141); Specter, S.; Watanabe, M.; Pan, S. H. *Am J Pathol* 93(2): 499-513; 1978.

Three tumor systems (lymphoma cell line FBL-3, mastocytoma cell line P815X, and plasmacytoma cell lines MOPC 104E, MOPC 31C, MOPC 173, and TEPC 15) were studied for their ability to modify humoral immunity to sheep RBC (SRBC) in vitro and in vivo. For the in vivo studies, BALB/c or DBA/2 mice infected with the Friend leukemia virus (FLV) complex or inoculated with tumor cells were challenged by ip or iv immunization with 4×10^6 SRBC. For the in vitro studies, 5×10^6 spleen cells from tumor-bearing or normal animals were cultured and immunized with 0.1 ml of a 0.1% suspension of SRBC (2×10^6 cells). FLV and all tumors depressed the hemolytic antibody plaque-forming cell response in susceptible mice, and spleen cells from normal animals were suppressed by intact tumor cells as well as by cell-free extracts from the plasmacytomas and mastocytomas. Cocultivation of FBL-3 lymphoma cells, which were initially induced by FLV in neonatal C57BL/6 mice, also resulted in immunosuppression. However, cell-free extracts from these immunosuppressive lymphomas were nonsuppressive. Thus, the lymphoma model did not support the concept that immunosuppression always occurs as a consequence of soluble factors. The mechanisms of suppression, although not fully defined, were different for each system investigated. (32 refs)

- 78-7004 Correlation Between Immunosuppression to Tumorigenesis. I. Humoral Immunosuppressing Factor.** (Eng) Hakim, A. A. (Dept. Surgery, Div. Surgical Oncology, Coll. Medicine, Univ. Illinois Medical Center, 180 Longwood Drive, Kankakee, IL, 60901). *Eur J Cancer* 14(11): 1249-1260; 1978.

The biological activities of agents released as a result of tumor cell proliferation were identified. Factors that alter cellular immunity and macrophage and lymphoid cell functions can be obtained from sera of animals bearing proliferating tumors. Inbred female Fischer rats and CFW mice bearing syngeneic mammary adenocarcinomas were used. The cytotoxicity of peritoneal macrophages and splenocytes against tumor cells and the effect of serum on the incorporation of ^3H -thymidine into normal splenocyte DNA in response to phytohemagglutinin (PHA) were assessed at various times during tumor growth. Serum enhanced the response of nor-

mal splenocytes to PHA if drawn before the 6th wk following sc injection of tumor cells; serum drawn after 7 wk suppressed this response. Macrophages and splenocytes from normal animals had no effect, those from animals bearing tumors enhanced tumor cell proliferation, and those from animals immunized against the tumor were cytotoxic against syngeneic tumor cells. Cell-free supernatants from mixed cultures of sensitized splenocytes, ie, splenocytes from immunized animals (at all ratios) and from animals bearing progressively growing tumors (at low ratios), with syngeneic tumor cells stimulated the response of normal syngeneic splenocytes to PHA. Since supernatants from cultures of splenocytes alone had no effect and supernatants from cultures of tumor cells alone had statistically significant suppressing activity, it is suggested that the suppressing agent(s) is released by the tumor cells. (58 refs)

- 78-7005 Adjuvant-induced Thymus-derived Suppressor Cells of Cell-mediated Tumour Immunity.** (Eng) Glaser, M. (Lab. Cell Biology, NCI, Bethesda, MD, 20014). *Nature* 275(5681): 654-656; 1978.

Inoculation of mice with complete Freund's adjuvant (CFA) markedly inhibited the ability of immune spleen cells to prevent simian virus 40 (SV40)-induced tumor growth in the Winn tumor cell neutralization assay and to lyse SV40-transformed cells in the ^{51}Cr -release assay. Female (BALB/c $\times$ C57BL/6)F₁ (CBF₁) mice were inoculated ip with 0.5 ml CFA and, immediately afterward, inoculated sc with 2×10^7 C57SV cells (an SV40-transformed cell line of BALB/c mice). T-cell-enriched fractions of spleen cells were purified and used in the ^{51}Cr -release assay against C57SV cells. A marked depression of cell-mediated cytotoxicity was observed. Challenge, sc, of another group of mice with 1×10^5 MKSA cells (an SV40-transformed line of BALB/c mice) 15 days after CFA immunization or mixing T-cell-enriched spleen cells from a third group of mice with 1×10^5 MKSA cells and injecting this mixture sc into normal CBF₁ mice resulted in death of most of these mice from progressive tumor growth. No tumors or death occurred in untreated control animals. Similarly, in the Winn assay, inhibition of tumor cell neutralization activity of lymphocytes from CFA-treated mice resulted in tumor growth and death of all animals as opposed to untreated mice, in which tumor cell neutralization resulted in complete protection against tumor growth. Thus, CBF₁ mice challenged with syngeneic SV40-transformed cells and CFA failed to develop effector cells; instead, they developed suppressor T cells that inhibited in vivo effector T-cell functions. Adult thymectomy reversed the suppression caused by CFA, which indicates that the intact adult thymus was necessary for the induction of suppressor cells. The results suggest that there is a delicate balance between the antitumor effects of adjuvants and their immunosuppressive effects. (17 refs)

- 78-7006 Enhancement of Avian Sarcoma Virus-induced Tumor Growth after Pretreatment with BCG.**

(Eng) Wainberg, M. A. (Lady Davis Inst. Medical Res., Jewish General Hosp., Montreal, Quebec, Canada); Israel, E. *Affect Immun* 22(2): 328-333; 1978.

The immunoprophylactic effects of BCG and the methanol extraction residue (MER) of BCG were investigated in chickens inoculated with avian sarcoma viruses (ASV). The chickens were inoculated in the wing web with 5 mg BCG or 0.5 mg MER 7 days before they were inoculated with 10^3 focus-forming units of virus. Of 17 chickens inoculated with BCG and ASV at the same site, 15 developed progressively growing neoplasms that eventually led to death. Injection of BCG into the wing web contralateral to that used for virus inoculation or injection of MER into the same wing web did not affect tumor development. Administration of BCG also gave rise to heightened levels of tumor-associated, cell-mediated immunity, both in chickens that received a subsequent injection of Prague strain A of Rous sarcoma virus and in normal, BCG-inoculated controls. BCG immunostimulation was apparently restricted to the cellular immune system, since an indirect immunofluorescence assay revealed no effect of BCG injection of the immunofluorescence staining capacity of virus-induced, tumor-bearing hosts. It is concluded that BCG prophylaxis is stimulatory to both Rous sarcoma development and to antitumor immunity in tumor-bearing hosts. (20 refs)

78-7007 **Characteristic Features of Immunologically-induced Changes in the Tumorigenesis in Mice of Various Strains.** (Rus) Volegov, A. I. (Lab. Pathophysiology, A. Hertsen Res. Oncological Inst., Moscow, USSR). *Biull Eksp Biol Med* 86(11): 572-574; 1978.

Tumor induction was compared in 3 to 4-mo-old C3HA, C3H/He, BALB/c, C57BL/6, and random-bred mice. Mice were immunized with BCG (0.01 or 1 mg in 0.2 ml of physiological saline) or with complete Freund adjuvant (CFA; 0.02 ml). Thirteen days after CFA administration and 30 days after BCG vaccination, animals were inoculated im with 20-methylcholanthrene (20-MC; 1 mg in 0.1 ml of vegetable oil). In C3H/He mice vaccinated with BCG, tumors started to develop significantly later than in non-treated mice; in C57BL/6 mice, BCG vaccination decreased the latent period of tumor development. CFA administration decreased the latent period of tumor development in C3H/He and C57BL/6 mice, inhibited tumor development in BALB/c and random-bred mice, and did not affect the tumor resistance of C3HA mice. These findings might indicate a role for hereditary factors in tumorigenesis. (12 refs)

78-7008 **Development of Immunocompetence in Hamsters and the Influence of DNA Oncogenic Viruses.** (Eng) Friedman, H. (Dept. Microbiology and Immunology, Albert Einstein Medical Center, Philadelphia, PA, 19141). *Fed Proc* 37(7): 2069-2071; 1978.

The effect of simian virus 40 (SV40) and adenovirus 12 (Ad12) infection on the development of immunocompetence in hamsters was investigated. Immunocompetence to sheep RBC developed rapidly in newborn hamsters. Animals inoculated on days 2-3 after birth developed few plaque-forming cells (PFC), but inoculation on days 5-7, 10-12, and 20-40 resulted in progressively greater responses (to $35-45 \times 10^3$ PFC/spleen). Inoculation of hamsters with SV40 at various times during the first 40 days of life had only a slight effect on PFC response 2-3 mo later. Hamsters given SV40 during the first week of life showed a slight depression in the antibody response to sheep RBC when challenged later in life. Tumors only occurred in animals treated with SV40 during the first 2 wk of life. Maturation of the immune response to sheep RBC was impaired in hamsters inoculated at birth with SV40. Injection of Ad12 into neonatal animals had similar effects. A slight to moderate effect on immune responsiveness to *Escherichia coli* lipopolysaccharide occurred after neonatal infection with SV40 or Ad12. Max effects were noted with animals 1-2 wk old. Cell-mediated immune responses to alloantigens, mycobacteria, and tumor virus-associated antigens were assessed in normal and virus-inoculated hamsters. Inoculation of normal hamsters with BCG as early as 2 wk of age resulted in marked cellular hypersensitivity: in vivo and in vitro reactions to challenge with purified protein derivative of tuberculin (PPD) were noted. Hamsters inoculated at birth with either SV40 or Ad12 also developed sensitivity to PPD after in vivo injection of BCG, but the in vitro response was weaker. The responsiveness of hamsters to 10% saline extracts of SV40 or Ad12 tumor extracts was directly related to prior inoculation at birth with virus. Sensitization became evident around 2 wk of age, reached a max around 3-4 wk, and decreased considerably by 60-100 days. This model may be useful in examining the interactions between host immune defense mechanisms and oncogenic DNA viruses that do not directly affect lymphoid cells. (25 refs)

78-7009 **Natural Killer Cells. In Vitro and In Vivo.** (Eng) Henney, C. S. (Fred Hutchinson Cancer Res. Center, 1124 Columbia St., Seattle, WA, 98104); Tracey, D.; Durdik, J. M.; Klimpel, G. *Am J Pathol* 93(2): 459-467; 1978.

Lymphoid cell populations from normal animals (rats, mice, and humans) can lyse tumor cells but not normal cells in in vitro ^{51}Cr release assays. The cells responsible for this cytotoxic activity have been characterized in the mouse and termed natural killer (NK) cells. In an attempt to discover whether NK cells are clonally distributed, they were absorbed onto monolayers of susceptible cells and then tested against other susceptible cells to see if the NK activity had been removed. The reactivity of NK cells from BCG-treated C57BL/6 mice against L5178 lymphoma cells was selectively removed by incubation of the effector cells on an L5178 cell monolayer but to a much lesser degree on a monolayer of normal DBA/2 spleen cells, which are insensitive to NK-mediated lysis. Adsorption on the L5178 cell monolayer also removed NK reactivity against susceptible human

Chang cells. On Chang cell monolayers, NK reactivity toward both L5178 and Chang cells was totally removed by adsorption. These experiments provided no evidence for clonal distribution of NK reactivities, but they indicated that target cell susceptibility to NK action may be determined by a shared antigenic specificity. Further BCG prophylaxis assays indicate that C57BL/6 NK cells were more efficient than growth-inhibitory adherent cells in vivo in retarding EL-4 ascites tumor growth. Although preliminary, these studies show that NK cells are an important host defense against progression to a neoplastic state. (25 refs)

78-7010 Modulation of Sensitivity to Natural Killer Cell Lysis after In Vitro Explantation of a Mouse Lymphoma. (Eng.) Becker, S. (Dept. Obstetrics and Gynecology, Sch. Medicine, Univ. North Carolina at Chapel Hill, Chapel Hill, NC, 27514); Kiessling, R.; Lee, N.; Klein, G. *J Natl Cancer Inst* 61(6): 1495-1498; 1978.

The kinetics of modulation to increased sensitivity to natural killer (NK) cell lysis was studied in explant cultures of a YAC ascites lymphoma induced in A/Sn mice by Moloney murine leukemia virus (M-MuLV). The sensitivity of all cultures to the same CBA spleen effector population was measured in a 6-hr ^{51}Cr -release assay. The lytic sensitivity was not significantly increased after 2 hr or 1 or 5 days of culture. The sensitivity was clearly increased by 3 wk, but max sensitivity was attained only after 1-2 mo of culturing. With the more sensitive competition assay, a change in sensitivity to NK lysis could be demonstrated as early as 2-24 hr of culture. Expression of the M-MuLV-determined, cell-surface antigen, measured by quantitative absorption with intact cells, increased in parallel with NK sensitivity, whereas the H-2a alloantigen concentration decreased during in vitro culture. These findings plus those from previous work indicate that establishment of an in vitro YAC line leads to profound membrane changes in the YAC cell. These changes may condition the cell to respond with an increased NK lytic susceptibility, even if the virally determined antigenic sites by themselves are not responsible for NK lysis. (19 refs)

78-7011 Alterations of the Antibody Response Following in Utero Exposure to Diethylstilbestrol. (Eng.) Luster, M. I. (Environmental Biology and Chemistry Branch, Natl. Inst. Environmental Health Sciences, Research Triangle Park, NC, 27709); Faith, R. E.; McLachlan, J. A. *Bull Environ Contam Toxicol* 20(4): 433-437; 1978.

The effects of a single administration of diethylstilbestrol (DES) to mice in the prenatal period on their immune response to sheep RBC (SRBC) and *Escherichia coli* 0127 lipopolysaccharide (LPS), thymic-dependent and thymic-independent antigens, respectively, were determined. Pregnant Swiss-Webster mice were injected with DES (1.0 mg sc) on day 16 of gestation. Offspring were immunized ip with either

0.2 ml of a 10% suspension of SRBC or 0.1 ml of a 0.1-mg/ml soln of *E. coli* 0127 LPS at age 7 wk. In utero exposure to DES did not alter the antibody response to SRBC. Lipid plaque-forming cells (PFC's) were decreased in DES-treated females and significantly increased in DES-treated males compared with controls. The enhanced action of DES on direct (IgM) LPS PFC in male mice could also be demonstrated during a secondary response (mice boosted with 0.01 mg LI 3 wk after the primary immunization), but antibody suppression was not observed in females. It is possible that exposure to DES during certain periods of fetal development stimulates or selectively programs a particular population of lymphocytes, particularly those involved with T-independent antigen responses, resulting in a relatively long-term alteration in the antibody response. (26 refs)

78-7012 Surface Membrane Changes of T Cells Induced by Syngeneic Tumour Cells. I. Formation and Uptake of Complexes of Ig and Tumour Antigens by T Cells. (Eng.) Dular, U. (Dept. Medicine, Immunology, Manitoba Inst. Cell Biology, Univ. Manitoba, Winnipeg, Manitoba R3E 0V9, Canada); Chow, D. A.; Paraskevas, F. *Int J Cancer* 22(5): 611-620; 1978.

The formation and uptake of complexes of immunoglobulin (Ig) and tumor antigens by spleen T cells were studied in mice inoculated ip with syngeneic tumor cells (L5178Y lymphoblastic leukemia and AKR thymoma). A dose of 10^7 cells from either source produced a significant increase in Ig-bearing spleen cells 6 hr later. Incubation of normal spleen cells in serum taken 6 hr after tumor cell inoculation reproduced in vitro the changes observed in the spleen cell population of the serum donors, demonstrating the presence of a cytophilic Ig that was taken up by T cells. Serum collected after tumor cell inoculation also contained Ig-antigen complexing factor (IACF) in the 4S fraction. Papain-digested tumor cell preparations, which contained specific tumor antigen activity, formed complexes with Ig in the presence of this factor. These complexes were cytophilic for a subpopulation of T cells, which thereby became Ig-positive. It is suggested that the cytophilic Ig present in the serum 6 hr after the injection of large numbers of tumor cells may also represent complexes of Ig and tumor antigens, the formation of which is mediated by IACF. (24 refs)

78-7013 Surface Membrane Changes of T Cells Induced by Syngeneic Tumour Cells. II. T-Cell Defects Induced by Small Tumour Cell Inocula or Tumour Cell Antigens. (Eng.) Chow, D. A. (Dept. Medicine and Immunology, Manitoba Inst. Cell Biology, Univ. Manitoba, Winnipeg, Manitoba R3E 0V9, Canada); Paraskevas, F.; Dular, U. *Int J Cancer* 22(5): 621-629; 1978.

T-cell defects induced by the injection of syngeneic L5178Y lymphoblastic leukemia cells in vivo were studied using

BA/2 mice. Injection of a large number of tumor cells was followed within 6 hr by the uptake of cytophilic immunoglobulin (probably complexes) by a subpopulation of T lymphocytes. This "6-hr T-cell response" was abrogated when small number of tumor cells (100) small amounts of an antigen-containing preparation from tumor cells, or repeated immunogenic doses of disaggregated human immunoglobulin are injected prior to the immunogenic challenge. Abrogation of the 6-hr T-cell response resulted in specific antitumor immunity, as measured by in vitro growth inhibition. It also resulted in loss of the enhancing effect of antibody formation against sheep RBC that is normally detected in a B-cell cooperative system when T cells are used 6 weeks after they have been primed with sheep RBC. It is suggested that this T-cell defect may represent a mechanism whereby tumor cells, in the early stages of their growth, interfere with the inductive stages of the immune response for a sufficient period of time to allow growth of the tumor beyond immune control. (25 refs)

78-7014 Resistance to Transplanted Cancer in Mice Increased by Live *Brucella* Vaccine. (Eng) Dard, L. (Centre Regional Anticancereux, Pontchaillou. F 35031, Rennes Cedex, France); Le Garrec, Y.; David, C.; Boujas, L. *Br J Cancer* 38(3): 464-467; 1978.

The effect of a chronic state of infection on the resistance of male (C57B16 x DBA2)F₁ mice to challenge by EL4 lymphoma cells was determined. Mice were vaccinated by iv injection of 5 x 10² or 5 x 10⁶ live *Brucella abortus*. One to 6 weeks later, half the animals in each group were killed; the remainder received 10³ EL4 lymphoma cells ip. Study of the lymphoid organs of the animals that were killed revealed that after injection of the lower dose, the number of bacteria increased until the third week. With the larger dose, however, the max number of bacteria was found as early as the first week. In all cases, the infection subsided by 6 wk. With the lower dose, the survival of lymphoma-bearing mice was slightly but significantly increased from weeks 3 to 5. Infection with the larger dose provoked a longer survival, particularly during weeks 1 and 2 after inoculation of the vaccine. Resistance to lymphoma was better correlated with the number of infectious organisms than with the degree of hyperplasia of the liver and spleen. It is suggested that *Brucella* may secrete a substance that inhibits cell division or that immune mechanisms against the bacteria may also act against the cancer cells. (23 refs.)

78-7015 Antigenic Content of an In Vitro Cellular Immunity Reactive Cell Surface Extract of Raji Cells. (Eng) Ng, M. H. (Medical Health Dept., Inst. Radiology Oncology, Queen Elizabeth Hosp., Kowloon, Hong Kong); Ng, W. S.; Ho, H. C. *Arch Environ Contam Toxicol* 8(3): 213-220; 1978.

The antigenic content of a Raji cell surface extract (TS) was studied by sequential absorption of rabbit antiserum to this extract with different cells. Anti-TS was reacted repeatedly with human blood cells followed by HeLa cells and human diploid fibroblasts. The residual reactivity was studied by stepwise absorption with an Epstein-Barr virus (EBV)-negative B lymphoblastoid cell line, BJAB, and then with B1-19, a clone derived from EBV-transformed BJAB cells. These processes delineated three kinds of reactivities in anti-TS having specificities for antigens on B blasts, EBV-determined cell surface antigen(s) and antigens that are exclusive to Raji cells. These results are compatible with the heterogeneous protein content of TS and with the results of earlier in vitro cell-mediated immunity studies using a similar preparation of Raji cell TS. (17 refs)

78-7016 High Lipid Fluidity of Leukemic Cell Membranes Through Host Immune Response and Shedding of Tumor Antigen-enriched Rigid Plasma Membrane Regions. (Eng) Van Blitterswijk, W. J. (Div. Cell Biology, Antoni van Leeuwenhoek-Huis, Netherlands Cancer Inst., Amsterdam, Netherlands); Van Der Sluis, P. J.; Hilgers, J.; Hilkmann, H. A.; Feltkamp, C. A.; Emmelot, P. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 341-344; 1978.

Thymus-derived leukemia (GRSL) in GR/A mice, maintained in ascites form by ip transplantation, possess mammary tumor virus (MTV)-induced cell-surface antigens (CSA). Seven days after transplantation, excess CSA are present in the ascites fluid in three forms: soluble, complexed, and on membrane vesicles. The microviscosity (fluidity) of phospholipids in intact cells and membrane fractions was determined by fluorescence polarization (P) using 1,6-diphenyl-1,3,5-hexatriene as a probe. Within a few transplantations in ascites form, the GRSL cells became fluid in normal nonirradiated hosts, but not in irradiated hosts. If cells from irradiated hosts were transplanted in nonirradiated hosts, the P value dropped, but the reverse did not occur to any marked degree. The increased fluidity of the cells in the ascites of normal hosts may reflect immunoselection and the shedding of CSA-enriched rigid vesicles: ie, abortive virus formation. The high fluidity of transplanted ascites GRSL cells may result in a decreased antibody-induced redistribution of antigens and a decreased sensitivity to immune killing. (7 refs)

78-7017 Modification of In Vitro Cell Mediated Immune Responses by Protein Components of a Leukemogenic Virus. (Eng) Fowler, A. K. (NCI, Frederick Cancer Res. Center, NIH, Frederick, MD, 21501); Twardzik, D. R.; Weislow, O. S.; Reed, C. D.; Hellman, A. In: *Advances*

in *Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977*. Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 234; 1978.

Freeze-thaw supernatants from banded Rauscher murine leukemia virus were chromatographed and the effluent fractions were tested for lymphocyte stimulatory activity and for inhibition of lymphocyte transformation induced by mitogens. Two samples from the first peak stimulated lymphocyte transformation but had no effect on mitogen-induced transformation. Material from the second peak exerted no stimulatory or inhibitory effect. A trypsin-sensitive component of the third peak suppressed mitogen-induced transformation and lipopolysaccharide stimulation. (no refs)

78-7018 Oncogenicity and Immunogenicity Associated with Membranes Isolated from Cell-free Ascites Fluid of Lymphoma-bearing Mice. (Eng) Raz, A. (Basic Res. Program, NCI, Frederick, MD, 21501); Barzilai, R.; Spira, G.; Inbar, M. *Cancer Res* 38(8): 2480-2485; 1978.

The immunogenicity and oncogenicity of membranes isolated from the cell-free ascites fluid of Moloney virus-induced lymphoma (YAC lymphoma)-bearing mice were determined. The supernatant obtained after the second centrifugation of the ascites fluid was separated into two fractions by ultracentrifugation: a supernatant containing soluble membrane fragments (AFM-s) and a pellet of insoluble membrane vesicles (AFM-p). Radioimmunoassay for murine leukemia virus p30 antigen (30,000-mol wt protein) indicated that the specific activity of this antigen is 20 times higher in the AFM-p fraction than in the YAC cell homogenate. Electron microscopy of the AFM-p fraction showed that viruslike (C-type) particles were associated with the plasma membrane vesicles. In vivo experiments with strain A mice indicated that the growth of an ascites tumor is associated with two biological activities, immunity and oncogenicity, and that both are present in the cell-free ascites fluid. The two activities can be disassociated by high-speed centrifugation or UV irradiation. In all cases, oncogenicity, but not immunogenicity, was abolished by UV irradiation. The immune activity is suggested to be associated with plasma membrane vesicles that contain a high specific activity of tumor-specific surface antigens. The oncogenic activity is associated with a tumor promoter factor that increases the efficiency of either intact lymphoma cells or the C-type particles. The tumor promoter factor is associated with the insoluble membrane fraction, since all oncogenic activity was associated with AFM-p. The immune activity is associated with both the soluble and insoluble fractions, since immunity was induced by both AFM-p and AFM-s. The possibility that shedding of plasma membrane vesicles and viruslike particles is associated with the same mechanism on the lymphoma cell surface membrane is explored. (38 refs)

78-7019 Surface Antigens in Acute Myeloblastic Leukemia: A Study Using Heterologous Antisera (Eng) Tupchong, L. (Nuffield Dept. Clinical Medicine, Radcliffe Infirmary, Oxford OX2 6HE, England); MacLennan, I. C. *Br J Cancer* 38(4): 481-495; 1978.

Attempts were made to confirm the presence of leukemia-specific antigens by producing rabbit antiserum against human leukemic myeloblasts by a variety of techniques. Ninety-five rabbits were immunized against myeloblasts from 19 acute myeloblastic leukemia patients by techniques designed to reduce the level of antibody against normal tissue antigens. Specificities of the antisera produced were determined by a K-cell-mediated cytotoxicity test and by a noncytotoxic radioimmune anti-immunoglobulin (¹²⁵I)-iodinated horse anti-rabbit binding assay. The antisera showed equal affinity to leukemic myeloblast preparations and normal splenic-lymphocyte preparations. When subsequently absorbed against pooled cadaveric human spleen cells to remove antibody against polymorphic antigens and tested against two leukemic myeloblastic preparations, 17/95 antisera had activity against at least one of the preparations. Several of the 17 absorbed sera also showed activity against phytohemagglutinin (PHA)-stimulated lymphoblasts. Although these antisera showed little activity against marrow per se, when they were absorbed twice against equal volumes of normal marrow, there was complete loss of activity against myeloblasts and PHA-stimulated lymphocytes, leading to the conclusion that specificity in these antisera was directed only against normal tissue antigens. It is concluded that although these sera contained activity against antigens common to leukemic myeloblasts and a minority population of normal marrow cells, they had no detectable activity against leukemia-specific antigens. (46 refs)

78-7020 Influence of Micrococcus, BCG and Related Polysaccharides on the Proliferation of the L1210 Leukaemia. (Eng) Verloes, R. (Laboratorium voor Chemie der Proteinen, Instituut voor Moleculaire Biologie, Paardenstraat 65; B-1640 Sint-Genesius-Rode, Belgium); Atassi, G.; Dumont, P.; Kanarek, L. *Br J Cancer* 38(5): 599-605; 1978.

The effects of *Micrococcus lysodeikticus* (ML), BCG, and series of structurally related complement-activating polysaccharides on the nonspecific and specific immune resistance of CDF₁ mice against L1210 lymphoid leukemia were compared. The mice were vaccinated ip with 10⁷ irradiated (8 kilorads) cultured leukemia cells and 1 mg BCG, ML, zymosan A, chitin, dextran sulfate, or irradiated cells alone 1 mo before ip challenge with 10⁶ leukemia cells. They had increased mean survival times (MST) of 165%, 67%, 159%, 98%, 76%, and 53%, respectively, over controls. These experiments were also conducted via the iv route, but the ip route was more effective in increasing MST over controls and inducing survivors to day 90 after tumor grafting. The addition of serial booster injections of 1 mg bacteria or polysac-

arides after tumor grafting increased the MST further and produced a higher percentage of long-term survivors. Treatments before or after tumor grafting with single or multiple doses of bacteria or polysaccharides alone was not able to increase MST over control values. Furthermore, serial injections of these complement activators + irradiated leukemia cells post-tumor grafting were also ineffective. (26 refs)

78-7021 Leukemia in AKR Mice. II. Two Modes of Suppression of In Vitro Antibody Formation by Leukemic Cells. (Eng) Roman, J. M. (Dept. Biological Sciences, Purdue Univ., Lafayette, IN, 47907); Golub, E. S. *Cell Immunol* 40(2): 316-325; 1978.

The in vitro suppression of the anti-sheep RBC (anti-SRBC) response of normal AKR mouse spleen cells by leukemic spleen and thymus cells from spontaneous AKR leukemias was further investigated. Leukemic cells that suppressed only syngeneic or semiallogeneic cells were termed restricted (R), and those that suppressed both syngeneic and allogeneic cells were called nonrestricted (NR). The suppression mediated by leukemic cells (representing approx 80% of spontaneous tumors) was overcome by the presence of irradiated allogeneic cells, allogeneic supernates, or mitogens. NR suppressors, however, exerted suppression in the presence of these substances. No differences in age, sex, or patterns of organ enlargement were observed between mice bearing R or NR tumors; in fact, individual leukemic AKR mice were sometimes found to have one lymphoid organ with R cells and another with NR cells. The combinations of allogeneic cells used for generating the supernates that overcame R suppression may differ from each other in background, histocompatibility antigen, or total genetic makeup. The activity of these supernates was nondialyzable and was partially sensitive to heat inactivation. When leukemic or normal cells were incubated with supernate and then washed, normal cells were still suppressible, but the leukemic cells were no longer suppressive, indicating that the supernates act on the leukemic cells rather than the normal cells. Suppression mediated by B10.A Haasch murine leukemia virus-induced leukemic cells was similar to that seen in AKR leukemic cells; both required cell contact, and 80% of the mice exhibited R suppression. This may indicate a mode of suppression common to many lymphoid leukemias, particularly those requiring congenital or long-term virus infection. The two types of suppression may reflect a modulation of leukemic cell-surface properties or a selection of leukemic cell populations resulting from specific host antitumor defense mechanisms. (22 refs)

78-7022 Leukemia in AKR Mice. III. Size Distribution of Suppressor T-Cells in AKR Leukemia and Neonatal Mice. (Eng) Mulder, A. M. (Dept. Biological Sciences, Purdue Univ., Lafayette, IN, 47907); Durdik, J. M.; Roth, P.; Golub, E. S. *Cell Immunol* 40(2): 326-335; 1978.

The size distribution of suppressor T cells of leukemic and normal AKR mice and neonatal (C57BL/6J x DBA/2J)F₁ (BDF₁) mice was compared by velocity sedimentation. All fractions of leukemic or neonatal thymus cells suppressed the immune responses of normal cells in vitro, but all fractions of normal AKR thymus cells enhanced these responses. Spleen cells showed a different pattern of suppression. Both neonatal and leukemic spleens contained suppressor cells, but these were found only among cells sedimenting at 3.3 mm/hr or faster; ie, large cells. Cells sedimenting slower than 3.3 mm/hr neither suppressed nor enhanced the response of normal cells. In contrast, small cells in the normal spleen contained cells that enhanced the responses of normal cells. Large and small suppressor thymus cells gave similar titration curves, indicating that suppression by the small cell fraction is not due to small numbers of large cells contaminating the population and that cells in both fractions have comparable specific activity. The suppression by lymph node cells was similar to that by spleen cells: only large cells suppressed. In contrast, an sc growing lymphoma exhibited a suppression pattern similar to that of the thymus. These data are consistent with the idea that the leukemia arising in the thymus exists as both large and small cells that are able to suppress normal responses but that only the large cells are exported to lymph nodes and spleen or that both large and small cells are exported but the small cells become large cells in the peripheral organs. Both neonatal and leukemic suppression was abrogated by x-irradiation or mitomycin C treatment. Both types of suppression required cell contact. The similarities between suppressor cells in neonatal and leukemic mice may indicate that the immunological events in spontaneous leukemia of AKR mice are to some extent a replay of the early events in normal T-cell differentiation. (26 refs)

78-7023 Protection of Newly-Hatched Chickens from Marek's Disease (JMV) by Normal Spleen Cells from Older Animals. (Eng) Lam, K. M. (Dept. Microbiology and Immunology, Temple Univ. Sch. Medicine, Philadelphia, PA, 19140); Linna, T. J. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers, J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 111-113; 1978.

The transfer of spleen cells from intact, nonimmunized adult chickens (≥ 8 wk old) that are naturally resistant to JMV to newly hatched chickens protected the recipients from JMV-induced leukemia. The results of various donor treatments (thymectomy, bursectomy, cyclophosphamide, or x-radiation, or their combinations) and spleen cell treatments (to remove macrophages or T cells) showed that non-T, non-B, nonmacrophage lymphoid cells are involved in the protection. (5 refs)

78-7024 Immunoprevention of Naturally Occurring Endogenous Murine Type-C RNA Viruses. (Eng)

Kende, M. (Lab. RNA Tumor Viruses, NCI, Frederick, MD, 21501); Stephenson, J. R.; Kelloff, G. J.; Al-Ghazzouli, I. K.; Dinowitz, M. *Nature* 273(5661): 383-385; 1978.

The induction of immunity to endogenous C-type viruses and the expression of these viruses were examined in BALB/c, NIH Swiss, and C57BL mice. Rauscher leukemia virus (R-MuLV) and Gross leukemia virus (G-MuLV) immunogens were injected sc on day 0 (200 µg), ip on days 14 and 28 (100 µg), and ip at 2-4 wk intervals (50 µg) for a total of 12 immunizations. All mice were killed 3 wk after the last immunization. All three strains had a virus-specific cytotoxic humoral immune response to both the R-MuLV and G-MuLV immunogens. Sera from C57BL mice were the most reactive, and sera from BALB/c were the least reactive. A predominant type-specific activity was seen with both immunogens, but a greater degree of cross-reactivity was seen with the former. At the time of killing, lymphocytes from all strains were nonreactive in cell-mediated cytotoxicity assays against the target cell used. Both immunogens induced an immunity to naturally occurring ecotropic virus in BALB/c mice: infectious ecotropic virus was detected in only 5/54 immunized animals in XC assays performed at the third passage vs 23/29 controls. At the first passage, only 3/54 animals yielded isolatable virus, and the mean titers were much less than those of controls. Both MuLV p30 and p12 expression in the spleen was closely correlated with the ability to isolate virus. Like BALB/c mice, immunized C57BL mice yielded few ecotropic virus isolates, but the incidence of isolates from control C57BL mice was lower than that from BALB/c mice. Investigations of class II and class III xenotropic virus activity at the third passage indicated that the immunized animals did not show marked reduction in endogenous xenotropic virus activity. However, the R-MuLV-immunized C57BL mice showed a trend of decreased activity in both infectivity and antigen assays that was significant with class II virus. (30 refs)

78-7025 Isolation of Leukaemia Specific Antigenic Macromolecular Moiety from Dinitrophenylated

Normal Human Leukocytes. (Eng) Karande, A. A. (Cancer Res. Inst., Tata Memorial Centre, Bombay 400 012, India); Sahasrabudhe, M. B. *Indian Acad Sci Proc [B]* 87(1): 1-8; 1978.

In an attempt to isolate tumor-specific antigen from non-malignant tissues, conformational changes were introduced in the cell-surface macromolecules of normal human WBC, and changes in the antigenicity of the cells were studied. Normal WBC suspensions were dinitrophenylated by treating them with 1-fluoro-2,4-dinitrobenzene (10⁴ molecules/cell). Agglutination titers of 256 to 2,048 against leukemic cells were obtained with rabbit antiserum against the dinitrophenylated normal WBC. Affinity chromatography on an antibody-sepharose column yielded three fractions, two of

which contained leukemia-specific antigenic moieties. Strong specific antileukemic agglutinating activity was obtained when the latter two fractions were injected in an oil-in-water emulsion into mice. The antileukemic agglutinating activity was weak when the pooled fractions were given in saline. The antigenic fractions had mol wts of 48,000 and 21,000, respectively. The use of the oil-in-water emulsion provided the necessary core for the antigenic proteins to orient themselves and perhaps acquire the same hydrophobic and hydrophilic orientation as that seen on intact cell membranes. (27 refs)

78-7026 Cell Sorter Analysis of Leukaemia-associated Antigens on Human Myeloid Precursors. (Eng)

Janossy, G. (Dept. Immunology, Royal Free Hosp., Pond St. London NW3, England); Francis, G. E.; Capellaro, D.; Goldstone, A. H.; Greaves, M. F. *Nature* 276(5684): 176-178; 1978.

The presence of leukemia-associated antigens on human bone marrow cells was studied by separating them in a fluorescence-activated cell sorter (FACS) according to their binding of anti-Ia-like or anti-acute lymphoblastic leukemia (ALL) antibodies. After most mature granulocytic and erythroid cells were eliminated, adult marrow samples contained 10%-14% cells reactive with anti-Ia serum (Ia⁺ cells). These Ia⁺ cells included larger cells, mainly myeloblasts and promyelocytes, and smaller SmIg⁺ (surface membrane immunoglobulin-positive) B lymphocytes. The Ia⁻ cells were myelocytes, maturing granulocytic and erythroid cells, a few lymphocytes, and <5% myeloblasts and promyelocytes. Colony and cluster-forming cells were present in both Ia⁺ and Ia⁻ fractions. Clone-forming cells were found almost exclusively among the intermediate-to-large cells, suggesting that small Ia⁺, ALL⁺ non-T, non-B cells did not form myeloid colonies. This was confirmed by staining fetal marrow cells with anti-ALL serum, separating them on the FACS, and culturing them for 7 days. The ALL⁺ and ALL⁻ fractions gave, respectively, 3.2 and 626 clones per 10⁴ cells. These results suggest that a proportion of human myeloid precursors are Ia⁺ and that a few are very strongly Ia⁺. The results also suggest that the two Ia⁺ non-T, non-B cell types (the smaller ALL⁺ cell and the larger ALL⁻ cell) might represent the target cells for leukemia transformation in the common (non-T, non-B) form of ALL and in acute myeloblastic leukemia, respectively. (27 refs)

78-7027 Cell-mediated Anti-tumor Response in the RL(male) 1 System. I. T Nature, Nature, H-2

DInvolvement and Antigenic Specificity of the Effector Cells. (Eng) Duprez, V. (Laboratoire d'Immunologie et de Virologie des Tumeurs, INSERM U 151, Hôpital Cochin, 27, Rue du Faubourg Saint Jacques, F-75674 Paris Cedex 14, France); Gomard, E.; Levy, J. P. *Eur J Immunol* 8(9): 650-655; 1978.

the effector cells involved in the cell-mediated antitumor response in the RL male 1 lymphoma cell system were characterized. After in vivo priming and in vitro secondary stimulation by radio-induced RL(male) 1 cells (RLC), syngeneic BALB/c splenic lymphocytes showed a specific but very weak cytolytic activity against target RLC. Under the same experimental conditions, spleen cells from (B6 x BALB/c)F₁ were at least 100-fold more efficient. In both cases the effector cells were cytolytic T lymphocytes (CTL), and normal M2 antigens of the tumor cell surface were involved in the effector-to-target cell interaction. The immune CTL were highly specific for target RLC; of cells from 13 other transplantable lymphomas as well as leukemic cells from spontaneously leukemic AKR mice, only P815 mastocytoma cells showed weak cross-reactivity. The exact nature of the antigen recognized by anti-RLC CTL remains unknown; the hematopoietic histocompatibility antigens which could have accounted for the F₁ antiparent reaction appear irrelevant. (28 refs)

78-7028 Increased Incidence of Spontaneous Lymphomas in Thymectomized Mice. (Eng) Cornelius, E. A. (Section Nuclear Medicine, Dept. Diagnostic Radiology, Yale Univ. Sch. Medicine, New Haven, CT). In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 1975-351; 1977.

The influence of thymectomy on the incidence of spontaneous lymphomas was examined in (C57BL/1 x A)F₁ [(BA)F₁] mice and in (SJL/J x C57BL/1)F₁ [(SB)F₁] mice. Mice were thymectomized (Tx) within 24 hr of birth. Tx(BA)F₁ mice demonstrated impaired cellular immunity, histologic changes similar to the graft-vs-host reaction, and a markedly increased incidence of lymphomas compared with normal control mice of the same age. This difference was greatest before old age, at which time tumors became more frequent in controls. Presumptive evidence of a subcellular oncogenic agent in the tumors of Tx mice was obtained. Most tumors arising in Tx(BA)F₁ mice were rejected on transplantation into normal syngeneic controls, suggesting that the impaired cellular immunity of the Tx mice permitted the growth of tumors that were more strongly antigenic than those that occurred in normal mice. The incidence of nonlymphoid tumors was not increased in Tx(BA)F₁ mice. In (SB)F₁ mice, thymectomy was associated with a moderately increased incidence of lymphomas for a 4-mo period. Tx(SB)F₁ mice, unlike Tx(BA)F₁ mice, demonstrated no lymphoproliferative changes and no graft-vs-host like lesions. (11 refs)

78-7029 T-Cell Lymphoma Induction by Radiation Leukemia Virus in Athymic Nude Mice. (Eng) Ricciardi-Castagnoli, P. (Cancer Biology Res. Lab., Dept. Radiology, Stanford Univ. Medical Center, Stanford, CA, 94305); Lieberman, M.; Finn, O.; Kaplan, H. S. *J Exp Med* 148(5): 1292-1310; 1978.

Extrathymic lymphoblastic leukemias developed in 3/5 athymic nude BALB/c mice and 1/8 athymic nude NIH Swiss mice inoculated with radiation-induced leukemia virus (RadLV), and a permanent cell line (BALB/Nu₁) was established from a BALB/c lymphoma. BALB/Nu₁ was highly tumorigenic when injected ip into BALB/c and NIH/Swiss nude mice, and electron microscopy of tumor cells showed features of a typical stimulated lymphocyte and numerous C-type viral particles, designated RadLV/Nu₁ virus. Reverse transcriptase activity of this virus coincided with that of ³H-labeled material in the 1.15-1.1 g/cm³ gradient density region. On polyacrylamide gel electrophoresis, RadLV/Nu₁ and RadLV/VL₃ viruses coelectrophoresed in most protein regions except the major envelope protein (gp71). Radioimmunoassay indicated that RadLV/Nu₁ possessed surface antigens common to or cross-reacting with the envelope antigens of RadLV/VL₃ and, to a lesser extent, xenotropic and ecotropic murine leukemia viruses. Cross-reactivity between the gp71 determinants of RadLV/Nu₁ and RadLV/VL₃ was demonstrated by membrane immunofluorescence test. RadLV/Nu₁ had thymotropic and leukemogenic activity in C57BL/Ka mice, but not in NIH/Swiss nude (nu/nu) or (nu/+). In vitro RadLV/Nu₁ was B-tropic, poorly fibrotropic, and limited to xenotropic activity. In several areas, therefore, RadLV/Nu₁ appeared to be closely similar to its parent virus RadLV. Terminal deoxynucleotidyl transferase activity in the lymphoma cells of BALB/c nude and NIH/Swiss nude mice was tenfold higher than in normal spleen cells of control nude mice. When tested for several T-cell antigens, these lymphoma cells were seen to be Thy-1.2+, Ly-2.2+, Ly-1.2-, and TL-. Immunoglobulin determinants were not noted. This profile resembles that of mature peripheral T cells, although the antigens were present in much lower quantities. (32 refs)

78-7030 Atypical Lymphoid Leukemia in Ataxia Telangiectasia. (Eng) Levitt, R. (Section Publications, Mayo Clinic, 200 First St. SW, Rochester, MN, 55901); Pierre, R. V.; White, W. L.; Siekert, R. G. *Blood* 52(5): 1003-1011; 1978.

The case reports of two sisters (23- and 25-yr-old) with ataxia telangiectasia (AT) are presented. The elder sister developed an atypical subacute lymphocytic leukemia characterized by large atypical lymphocytes and prolymphocytes, and the absence of organomegaly. Results of cytogenetic banding disclosed that both patients had identical abnormalities associated with chromosome 14 (t(14q11:14q34)) with the leukemic sister showing additional stable aberrations. The presence of the marker in the leukemic cells of the elder sister supports the theory that her leukemic cells arose from a preexisting AT lymphocyte clone. Furthermore, it is suggested that the

presence of an abnormal lymphocyte clone, depressed cellular immunity, or susceptibility to chromosomal damage or rearrangement may be responsible for the increased incidence of malignancy in AT patients and AT carriers. (37 refs)

- 78-7031 Crossed Tumor Antigenicity Between the Experimental Tumors Ca M, Ca H, Ca O and Ca M₄.** (Spa) Cuadrado, S. P. (Departamento de Immunopatologia, Instituto Nacional de Oncologia, Madrid-3, Spain); Moreno Koch, M. C.; Tome Paule, C. *Rev Esp Oncol* 25(1): 45-47; 1978.

The cross antigenicity between the experimental tumors Ca M, Ca H, Ca O and Ca M₄ was studied in C₃H/he mice. Eighty animals first received a Ca M tumor transplant. After the surgical removal of this tumor, four subgroups of animals received transplants of Ca M (autograft), Ca H, Ca O or Ca M₄ tumors. Another group received a primary Ca H transplant; its two subgroups received secondary Ca H and Ca M grafts. The third group received primary Ca O transplants; the secondary transplants were Ca M and Ca O in the subgroups. The fourth group was grafted first with Ca M₄, and then Ca M and Ca M₄ were grafted in the subgroups. The immune status produced by the primary Ca M and Ca M₄ tumors offered protection against subsequent transplants of both Ca M and Ca M₄ (crossed antigenicity), but not against Ca O and Ca H transplants. The immunity produced by Ca O offered, however, protection against a subsequent Ca M transplant. The Ca H tumor failed to produce any immune protection against other tumors. The absence of crossed antigenicity between tumors in which viral particles are present suggests that these viral particles are not involved in the pathogenesis of the tumors. (no refs)

- 78-7032 HLA and Glioma.** (Eng) de Moerloose, P. (Transplantation Immunology Unit, Div. Immunology and Allergy, Hopital Cantonal, 1211 Geneva 4, Switzerland); Jeannet, M.; Martin-Achard, A.; de Tribolet, N.; Seiler, R.; Guanella, N. *Tissue Antigens Histocompat Immunogenet* 12(2): 146-148; 1978.

Eighty glioma patients were typed for several histocompatibility antigens (HLA-A, -B, and -DRw), with age at time of diagnosis, sex, blood, group, and degree of malignancy being considered. Phenotypic frequencies of Bw35 and Drw1 antigens were increased in patients compared with controls, but when p values were corrected by the number of antigens tested (35), the results were not significant. (7 refs)

- 78-7033 Immunotherapy of Chemically-induced Tumors in the Hamster Cheek Pouch with Dinitrochlorobenzene.** (Eng) Marshack, M. (Dept. Oral Pathology, Loyola Univ. Sch. Dentistry, Maywood, IL, 60153); Toto, P.; Kerman, R. *J Dent Res* 57(4): 625-630; 1978.

The effect of dinitrochlorobenzene (DNCB) sensitization in the buccal pouch of hamsters prior to or after the induction of cheek pouch tumors with 7,12-dimethylbenz(a)anthracene (DMBA) was studied. For tumor induction, the buccal pouches were painted with 0.5% DMBA in mineral oil 2x/wk for 14 wk. For DNCB sensitization, two pellets of cotton, each containing 1 ml of 0.1% DNCB soln, were inserted in each pouch and left to remain until expelled by the hamster (usually after 10 min). Hamsters were given two DNCB treatments, 48 hr apart; 14 days later, the pouches were challenged with a 0.01% DNCB soln. Hamster cheek pouches treated with DMBA only developed advanced invasive squamous cell carcinomas. Infiltration of the connective tissue surrounding the tumors was slight, with an av of 1 lymphocytes and < 1 plasma cell/50 μ m². When hamsters were treated with DMBA and then DNCB, benign noninvasive papillary tumors resulted. Hamsters pretreated with DNCB followed by DMBA (starting at the same time as the challenge dose of DNCB was being responded to) developed squamous cell carcinoma in situ. However, with repeated (1-4) DNCB rechallenges, 5/16 pouches developed noninvasive papillary tumors characterized by hyperplasia with dyskeratosis and 11 pouches showed no tumors. A lymphocytic and plasma cell-mediated response was seen in all DNCB-treated groups, and the lymphocyte response was greatest (av of 50 lymphocytes/50 μ m²) in hamsters treated with DNCB following tumor induction. Lymphocytes not only invaded the lamina propria but also the epithelium. These lymphocytes may be reacting to tumor-associated antigens. (11 refs)

- 78-7034 Antibody to Epstein-Barr Virus in Patients with Carcinoma of the Nasopharynx.** (Eng) Inuyama, Y. (Dept. Oto-rhino-laryngology, Sch. Medicine, Keio Univ. Tokyo, Japan); Asaoka, K.; Nakajima, Y.; Horiuchi, M.; Takasaki, K.; Ozu, R. *Keio J Med* 26(2): 79-89; 1977.

Antibody titers against the viral capsid antigen (VCA) of Epstein-Barr virus (EBV) were studied in 59 patients with a malignant tumor of the nasopharynx (51 carcinomas, 8 lymphomas), 117 patients with a tumor of other sites of the head and neck, and 263 patients without malignant disease (controls). The geometric mean titer was 1:794 and the antibody-positive rate 82% in the patients with nasopharyngeal tumors. The corresponding values were 1:190 and 37% in patients with other head and neck tumors and 1:120 and 20% in the controls. Riddit analysis indicated that the differences among the groups were statistically significant. The anti-VCA titer was determined repeatedly in patients with nasopharyngeal carcinoma. It was found that in cases of recurrence or metastasis, the titer was elevated markedly, but in patients with a short- or long-term remission, it was decreased. However, radiotherapy appeared to increase the anti-VCA titer. This could result from either antigen stimulation due to the destruction of cancer cells or the production of more antibodies by B cells following relief from the suppression of T cells. Anti-VCA titers might be useful in the

gnosis of nasopharyngeal carcinoma. The findings suggest that EBV is either carcinogenic or is a passenger virus in nasopharyngeal carcinoma. (16 refs)

78-7035 Protective Effect of Thymectomy in a High Cancer Risk Population. (Eng) Papatestas, A. (Dept. Surgery, Mount Sinai Sch. Medicine, New York, NY 10029); Kark, A. E.; Jenkins, G.; Aufses, A. H. *In: Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 6-May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 353-365; 1977.

The cancer risk in myasthenia gravis (MG) was investigated in relation to thymic pathology, the differences in cancer incidence between patients with and without thymectomy (Tx), and the role of Tx in altering the risk of oncogenesis. The records of 2,000 patients registered at a New York hospital and a London hospital were reviewed; 189 of these patients had undergone Tx. A total of 149 neoplasms were found in the entire population. In women, a significant increase in relative risk (RR) was present prior to Tx both in those with thymomatous MG and those with associated thymomas. Following Tx, the relative risk decreased and there was no difference from the expected numbers. In men, the RR prior to Tx was also higher than that following Tx, but there were no significant differences in RR. In both sexes, patients with thymomas and those with many germinal centers had a higher RR than patients without thymomas. The post-Tx decrease in RR was more marked in patients who were operated early after the onset of disease and in those who had reached remission. The observed numbers of neoplasms in the period after the onset of remission were lower than expected, but in all other periods observed numbers were higher than expected. Thus, Tx may have a protective effect against the high cancer risk associated with thymic abnormalities. (25 refs)

78-7036 Immunological Studies in Epidermodysplasia Verruciformis. (Eng) Jablonska, S. (Dept. Dermatology, Warsaw Sch. Medicine, ul. Koszykowa 82a, 02-008 Warsaw, Poland); Orth, G.; Jarzabek-Chorzelska, M.; Rzesza, G.; Obalek, S.; Glinski, W.; Favre, M.; Croissant, O. *Bull Cancer (Paris)* 65(2): 183-190; 1978.

Immunofluorescence (IF) and cell-mediated immunity (CMI) studies were made of 14 cases of Epidermodysplasia verruciformis (EV), 10 familial and 4 nonfamilial. Numerous malignancies plus one occurrence of Bowen's disease were found in three nonfamilial cases; malignancies were also found in three nonfamilial cases. Attempts were made to determine (1) the antigenic differences of viruses inducing EV by IF; (2) the significance of CMI for progression of the disease and possible malignant transformation; (3) whether

the types of human papillomavirus (HPV) inducing EV differ in their oncogenic potential; and (4) any possible correlation of the type of HPV with CMI. Two different types of HPV, HPV-3 and HPV-4, were found in the EV cases. There was a relationship between the type of HPV and the clinical picture of EV as well as the occurrence of malignant transformation. HPV-4 was more oncogenic: of the 12 cases typed, transformation occurred in 1/7 with HPV-3 and in 5/5 with HPV-4. CMI was depressed in most of the active cases and preserved in regressing and abortive cases. However, a low CMI was found in EV cases infected with HPV-3 and in persistent flat warts, also due to HPV-3, that did not undergo malignant transformation. In contrast, in a case of EV due to HPV-4, malignant transformation occurred in spite of a preserved, although lowered, CMI. HPV-1, which is responsible for plantar warts, and HPV-2, responsible for common warts, have no evident oncogenic potential. (29 refs)

78-7037 Common Human Melanoma Membrane Antigens Detected by Mixed Hemadsorption Microassay with Serum from a Patient Undergoing Immunotherapy with Autologous Tumor Cells. (Eng) Liao, S. K. (Ontario Cancer Foundation Hamilton Clinic, c/o Henderson General Hosp., 711 Concession St., Hamilton, Ontario, L8V 1C3, Canada); Leong, S. P.; Sutherland, C. M.; Dent, P. B.; Kwong, P. C.; Krementz, E. T. *Cancer Res* 38(12): 4395-4400; 1978.

The postimmune serum (PIS) and preimmune plasma of a patient with malignant melanoma who received 28 immunizations with autologous, irradiated, cultured melanoma cells plus BCG were tested for their reactivity in mixed hemadsorption microassays against several cell lines. No reactivity against any of the five allogeneic melanoma cell lines or five nonmelanoma lines tested was detected in the preimmune plasma, but strong reactivity against all five melanoma lines in varying titers (1:2,560 to 1:40,960) and against 2/5 nonmelanoma lines in lower titers (1:320 and 1:1,280) were found in the unabsorbed PIS. Absorption with human AB RBC, pooled peripheral blood WBC, and spleen homogenates of normal adults, did not alter this reactivity. Further absorption with cells of a human laryngeal carcinoma (KB) line reduced reactivity against the five melanoma lines 4- to 16-fold (1:160 to 1:5,120), but a strong residual common reactivity against these lines still remained. This absorbed serum no longer reacted with any of the five nonmelanoma lines. Removal of reactivity against fetal calf serum (FCS) by absorption with insolubilized FCS reduced reactivity against the melanoma cell lines to the same extent, but failed to remove reactivity against the nonmelanoma cell lines completely. Further absorption of PIS with human fetal homogenate or BCG had no effect on its reactivity against the melanoma lines. Blood group isoantigens, histocompatibility locus antigens, and FCS were ruled out as causes of the observed seropositivity against the melanoma cell lines in the final two stages of sequential absorptions. These data provide further evidence for the existence of common melanoma-associated antigens. (32 refs)

- 78-7038 Genetic Control of Resistance to Tumors.** (Eng) Phillips-Quagliata, J. M. (Dept. Pathology, New York Univ. Medical Center, 550 First Ave., New York, NY, 10016); Walker, M. C. *Am J Pathol* 93(2): 487-498; 1978.

To determine whether genes mapping within the major murine histocompatibility complex (H-2) control resistance to a BALB/c plasmacytoma (MPC-11), hybrid mice from over 20 parental strains were injected sc in the right dorsal flank with 3×10^5 viable MPC-11 cells (a dose fatal to 100% of BALB/c controls). F₁ hybrids between BALB/c and other BALB congenic resistant (CR) strains were as susceptible to MPC-11 as homozygous BALB/c mice, which demonstrated that H-2 heterozygosity was not sufficient for resistance. F₁ hybrids between BALB/c and all but two recombinant inbred strains were resistant. In (BALB/c x B10 CR)F₁ hybrids, resistance was dependent on the presence of a single dominant autosomal gene or gene complex that segregates independently of H-2 and that has slight differential effects on various H-2 haplotypes. The resistant mice became susceptible to MPC-11 cells injected after 450 rads γ irradiation, but they remained resistant after iv silica injection. Studies of hybrids between B6 and BALB/c mice suggested that their tumor resistance was decreased by the expression of Gross viral antigens of some other viral determinant. The mechanism for this is not understood, but it might involve tolerance, suppression, or enhancing antibody. (39 refs)

- 78-7039 Resonance Raman-Spectroscopic Studies of the Hapten Features Involved in the Binding of 2,4-Dinitrophenyl Haptens by the Mouse Myeloma Proteins MOPC 315 and MOPC 460.** (Eng) Kumar, K. (Dept. Chemistry, St. Mary's Univ., Halifax, N.S., Canada); Phelps, D. J.; Carey, P. R.; Young, N. M. *Biochem J* 175(2): 727-735; 1978.

The resonance Raman spectra of four dinitrophenyl (Dnp) haptens were characterized, and the binding of the haptens to mouse myeloma proteins MOPC 315 IgA and MOPC 460 IgA was studied by resonance Raman spectroscopy. The interactions between a given hapten and MOPC 315 IgA were often quite distinct from those between the same hapten and MOPC 460 IgA. More importantly, the nature of the R side chain in a Dnp-NHR compound appeared to modify the contacts between the chromophore and the protein. Thus, with the haptens studied, there was no unique set of contacts between the Dnp group and the binding site. The contacts expected between ϵ -2,4-dinitrophenyl-L-lysine and the site on MOPC 315 IgA were not detected. However, the contacts between this hapten and the site on MOPC 460 IgA were close to those predicted by a model for MOPC 315 IgA. (25 refs)

- 78-7040 Symmetry of Binding Sites of a Mouse IgA Myeloma Protein (MOPC 315).** (Eng) Eisen-

berg, R. (Dept. Medicine, Univ. North Carolina, Chapel Hill, NC, 29514); Plotz, P. *Biochemistry* 17(22): 4801-4807; 1978.

The mechanism of monovalency of the 7S subunit of a mouse IgA myeloma protein (MOPC 315) against a large antigen was investigated using affinity labels. In an equilibrium Farr assay, the subunit was shown to have only one valence for a univalent 40,000-mol wt antigen (dinitrophenyl-dextran). In four experiments, an average valence of 1.1 and a K_a of 1.4×10^6 were obtained. This monovalency did not appear to be an artifact of the Farr assay used, as a small hapten (dinitrophenyl- ϵ -aminocaproic acid) tested in the same assay gave an average valence of 1.8 and a K_a of 1.4×10^6 . Blocking binding sites with a small affinity labeling hapten, bromoacetyl-dinitrophenyl-lysine, quantitatively reduced the binding of large molecules in a manner strongly favoring the model of a symmetric 7S molecule, in which the binding of a large ligand to one site inhibits an adjacent site from binding a second large ligand. The results of dinitrophenyl-lysine-agarose column chromatography strongly favored the steric hindrance model of MOPC 315 monovalency. Such a model implies that the Fab binding sites are symmetrically arranged in the 7S subunit, but that they are too close together to permit two large antigens to bind simultaneously. (32 refs)

- 78-7041 Autoimmunity and Tumor Resistance. 3-Methylcholanthrene Tumorigenesis in New Zealand Black Mice.** (Eng) Morton, J. I. (Div. Immunology, Dept. Medicine, Univ. Oregon Health Sciences Center, Portland, OR, 97201); Siegel, B. V.; Moore, R. D. *Am J Pathol* 93(2): 469-485; 1978.

The susceptibility of New Zealand Black (NZB) mice to epidermal tumors induced by 3-methylcholanthrene (3-MCA) was examined as a function of age, sex, immune and autoimmune status, immunosuppressive treatment, and hepatic aryl hydrocarbon hydroxylase (AHH) levels. Results of a comparison of the frequency, type, and distribution by sex of 3-MCA-induced tumors in NZB, BALB/c, C57BL/6, and SJL/J mice given a single sc injection of 150 μ g 3-MCA at age 4 mo revealed that NZB mice had a longer tumor latency period and a lower tumor incidence than the other strains. Although most of the tumors in the other groups were sarcomas, the NZB group had an almost equal number of sarcomas (21) and squamous cell carcinomas (18). In NZB mice injected with 3-MC at different ages, 4-mo-old females responded with the greatest incidence and most rapid development of tumors. Treatment with antithymocyte serum decreased the final incidence of tumors while x-irradiation with 500 rads accelerated the onset of tumors in 4-mo-old NZB mice. The development of glomerulonephritis and hematopoietic carcinomas was independent of 3-MCA injection and the presence of skin tumors. Liver AHH specific activity following ip injection of 100 mg/kg 3-MCA was highest in female NZB mice, and the activity increased gradually with age. (30 refs)

78-7042 Circulating Immune Complexes in Rats Bearing Chemically Induced Tumors. I. Sequential Determination During Growth of Tumours at Various Body Sites. (Eng) Hoffken, K. (Cancer Res. Campaign Lab., Univ. Nottingham, Univ. Park, Nottingham NG7 2RD, England); Hoffken, M. R.; McLaughlin, P. J.; Moore, V. E.; Baldwin, R. *Int J Cancer* 21(4): 496-504; 1978.

transplanted 3-methylcholanthrene-induced sarcoma (Mc7) and a 4-dimethylaminoazobenzene-induced hepatoma (D23) were used to determine whether the binding of circulating immune complexes to the complement component C3 can be employed to monitor tumor growth in rats. Circulating immune complexes were measured in sequential monitoring studies in tumor-bearing WAB/Not rats. The assay employed was based on the radioimmunoprecipitation of immune factors with ^{125}I -Clq or on an indirect test whereby the inhibition of binding of ^{125}I -Clq to IgG aggregates by tumor-bearer sera were measured. After an initial peak coincident with a tumor diameter of about 0.5 cm, the Clq-binding activity of sarcoma-bearer sera decreased until they ultimately bound less Clq than normal rat serum. Sc implantation or growth of D23 hepatoma resulted in a change in Clq binding by tumor-bearer serum similar to that observed with sc inoculation of Mc7 sarcoma. The percentage of specific inhibition of tumor-bearer serum decreased at late stages of tumor growth, but it was reversed in rats whose tumors were excised 7 days after implantation. Antibody reactivity increased with increasing tumor burden, and the specific humoral activity was lost following resection. Following iv tumor cell injection, immune-complex levels increased until the terminal stages of the disease. Measurement of Clq-binding serum factors represents a useful method for monitoring the growth and burden of experimental animal tumors. (33 refs)

78-7043 Induction of a Tumor with Greatly Increased Metastatic Growth Potential by Injection of Cells from a Low-Metastatic H-2 Heterozygous Tumor Cell Line into an H-2 Incompatible Parental Strain. (Eng) Kerst, R. S. (Natl. Cancer Inst. Canada Res. Group, Dept. Pathology, Queen's Univ., Kingston, Ontario K7L 3N6, Canada); Twiddy, R. R.; Robertson, D. M. *Int J Cancer* 22(5): 533-594; 1978.

In H-2-heterozygous sarcoma, MDAY, originally induced by 3-methylcholanthrene in an (A x DBA/2) F_1 (H-2a x H-2d; DBA/2 F_1) hybrid was adapted for growth in the H-2-incompatible DBA/2 parent strain by serial ip transplantation of ascites tumor cells. Upon serial passage, the cells gradually lost their K-2Kk antigenicity. When these H-2Kk-negative, H-2d-positive variant (MDAY-D2) cells were injected sc into DBA/2 AD F_1 recipients, they gave rise to highly metastatic tumors. Widespread metastases to liver, spleen, lungs, and kidneys were evident within 12-18 days. There was total replacement of liver and spleen within 2-3 wk. The tendency to metastasize and loss of the K-2Kk haplotype appeared stable and irreversible. In contrast, MDAY cells either did not

metastasize or did so only to the liver in AD F_1 mice. The MDAY-D2 tumor originally appeared to be an H-2-loss variant tumor of an originally heterozygous tumor growing in AD F_1 x hosts. However, several results cast doubt on its variant nature; for example, MDAY and MDAY-D2 differed in their expression of a wide variety of cell-surface antigens and receptors. If MDAY-D2 is not a variant but is instead a newly induced DBA/2 tumor, the most likely causative agent might be an oncogenic virus harbored by the MDAY cells that is released upon cell death in vivo subsequent to an allogeneic reaction. Although its precise origin is not clear, MDAY-D2 has metastasizing properties that make it desirable for studying the biology of metastasis. (26 refs)

78-7044 T-T Cell Collaboration in Rejection of a Syngeneic SV40-induced Sarcoma in Mice. (Eng) Glaser, M. (Lab. Cell Biology, NCI, Bethesda, MD, 20014); Law, L. W. *Nature* 273(5661): 385-387; 1978.

The concept of T-T cell interactions during cell-mediated immune responses to tumor-associated antigens in vivo was examined. Female BALB/c mice were inoculated sc with mKSA-Tu5 tumor culture cells that had lost their oncogenic potential but had maintained tumor-specific transplantation antigen activity. Additional mice were inoculated ip with mitomycin C-treated (50 $\mu\text{g}/\text{ml}$) Meth-A cells, a methylcholanthrene-induced tumor with strong tumor-specific transplantation antigen activity. In the first experiment, various effector:target cell ratios of lymph node and thymus cells from normal BALB/c mice or mice immunized against mKSA or Meth-A cells plus mKSA or Meth-A cells were injected sc into normal BALB/c mice. No protection against mKSA cells was seen with 25:1 or lower ratios of anti-Meth-A lymph node:Meth-A cells. Thymus cells did not protect against either tumor at any cell number tested. Studies of the ability of immune lymph node and thymus cells to collaborate in the rejection of mKSA cells indicated that injection of immune lymph node or thymus cells alone did not prevent tumor growth, but together they completely inhibited tumor growth. Untreated anti-mKSA immune lymph node cells that could prevent tumor growth lost their activity after treatment with specific anti-Thy-1.2 serum and complement; this indicated that the cell type in the immune lymph nodes needed to neutralize mKSA is T-cell-dependent. Untreated immune lymph node or thymus cells at suboptimal concentrations did not neutralize mKSA when injected alone, but they did prevent tumor growth when injected together. This collaborative effect could be abrogated by pretreatment of either cell type with anti-Thy-1.2 and complement, but not complement alone, before mixing, this indicated that T cells in both lymph nodes and thymus were needed for collaboration. (12 refs)

78-7045 Effect of Generalized Graft-Versus-Host-Reaction on B- and T-Lymphocytes and a Benzpy-

rene-induced Murine Sarcoma. (Eng) Witting, C. (Pathologisches Institut, Westring 17, D-4400 Munster, W. Germany); Hultsch, E. *Z Krebsforsch* 92(3): 255-265; 1978.

The effects of a generalized graft-vs-host reaction (GVHR) on peripheral blood lymphocytes and benzo(a)pyrene (BP)-induced sarcomas were studied in F₁ hybrids of A/Jax female mice and BALB/c males. The GVHR was induced in 3-mo-old hybrid mice by the ip transplantation of A/Jax spleen cells (100×10^6). Twelve months later, BP ($4 \mu\text{g/g}$) was applied sc 1x/wk for 4 wk. Tumor incidence 12 wk after the last application was 74.5% in 200 GVHR mice, compared with 49.5% in 200 control mice. The tumors in the GVHR group were significantly larger and heavier than those in the controls, and metastases were observed in 21% of the mice vs 0% of the controls. In both groups, >95% of the tumors were myosarcomas. The percentage of T lymphocytes declined from 56%-65% in the controls to 36%-45% in the GVHR animals; the percentage of B lymphocytes increased from 21%-30% to 46%-55%. Examination of the marginal zone of the tumors revealed a marked decrease in T cells and total round cell count. There were no changes in B cells in this zone. It is concluded that the GVHR severely impairs cells, probably T lymphocytes or their subgroups, that play an essential role in tumor defense mechanisms. (26 refs)

78-7046 Effect of Splenectomy on B- and T-Lymphocytes and a Benzpyrene-induced Murine Sarcoma. (Eng) Witting, C. (Pathologisches Institut, Westring 17, D-4400 Munster, W. Germany); Hultsch, E. *Z Krebsforsch* 92(3): 267-274; 1978.

The effects of splenectomy (SP) on peripheral blood lymphocytes and benzo(a)pyrene (BP)-induced sarcomas were studied in female hybrids from A/Jax female mice and BALB/c males. The hybrids were splenectomized 3 wk after birth; beginning 10 wk later, the mice received three single doses of $4 \mu\text{g/g}$ BP at weekly intervals. Tumor incidence 12 wk after the last dose was 28.5% in 200 SP animals, and 49.5% in 200 BP controls. Histologically, all tumors of SP mice and 96.9% of those in control mice were myosarcomas, but the tumors were significantly smaller in the SP group. There was a significant decrease in B lymphocytes and an increase in T lymphocytes in the SP mice, compared with BP controls and untreated controls. In the marginal zone of the tumors, there was a decrease in B cells and an increase in T cells but no change in total round cell count. Studies of spleen lymphocytes in 100 untreated hybrid mice revealed a very high proportion of B lymphocytes. The antineoplastic effects of SP may have been due to the B-cell reduction, which allowed T cells to increase proportionally and thus improve host defense mechanisms. (17 refs)

78-7047 Combined Effects of Immune Depression and Carcinogenesis. (Eng) Trainin, N. (Dept. Cell

Biology, Weizmann Inst. Science, Rehovot, Israel); Small M.; Gabizon, A. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY) Vol. 2(part 1), 2404 pp.; 1621-1630; 1978.

The influence of various lymphoid subpopulations on the behavior of transplantable tumors (a simian virus 40-induced and a methylcholanthrene-induced fibrosarcoma; the 3 LI Lewis lung carcinoma) was investigated in inbred mice. When spleen cells from tumor-bearing T mice (thymectomized, irradiated, and injected with thymocytes) were transferred with tumor cells into syngeneic recipients, the incidence of tumor metastases was increased. Spleen cells from B mice (thymectomized, irradiated, and injected with bone marrow cells) did not exhibit this activity. Thymocytes from nonsensitized animals also accelerated the growth of a primary tumor. Fractionation of the spleen cell population from tumor-bearing mice by velocity sedimentation yielded one fraction of tumor-sensitized cells that inhibited the growth of the same tumor and a second fraction that enhanced tumor growth upon transfer. These experiments suggest that subpopulations of lymphocytes with differing characteristics coexist within the spleen and that these subpopulations may exhibit opposing reactivities upon exposure to syngeneic tumor cells. Thus, immunosuppressive agents could bring about contradictory results, depending on whether the tumor-inhibiting or tumor-enhancing subpopulation is impaired. (11 refs)

78-7048 Chemical Modification and Immunogenicity of Membrane Fractions from Mouse Tumour Cells. (Eng) Staab, H. J. (Friedrich Miescher Laboratorium der Max Planck Gesellschaft, Spemannstrasse 37-39, 7400 Tübingen, W. Germany); Anderer, F. A. *Br J Cancer* 38(4): 496-502; 1978.

The effects of several chemicals on the immunogenicity of a crude membrane fraction isolated from mouse STU-D17 tumor cells (STU mouse embryo cells transformed by Rous sarcoma virus) were tested in syngeneic mice. Groups of 10 STU mice were immunized with membrane equivalents corresponding to 10^4 , 10^3 , 10^2 , and 10^1 cells of modified crude membrane fractions of tumor cells or of the crude membrane fractions from DMS-modified tumor cells. A challenge dose of 10^5 viable tumor cells was given sc 14 days after immunization. Compared to 0% tumor-free control animals, immunization with unmodified membrane fraction produced 20% tumor-free animals after 90 days. Immunogenicity was appreciably elevated when the crude membrane fraction was modified by dimethyl sulfate (DMS: 50 mM) and partially elevated when modified by acetic anhydride (50 mM) or glutardialdehyde (200 mM) + [1-ethyl-3(3-dimethylaminopropyl)carbodiimide (20 mM) + methylamine (3 M)]. The membrane

ection treated with DMS produced up to 60% tumor-free animals, and the mean survival time of animals developing tumors showed the greatest increase over controls. However, the membrane fraction from DMS-modified tumor cells failed to produce an immune response, a result that is difficult to explain. For all chemical modifiers, membrane equivalents of 10^2 - 10^4 tumor cells were most effective. Neuraminidase had no apparent effect on the immunogenicity of the tumor cells. The immunogenetic enhancement was specific, as immunization with STU 51A/232B or fetal tissue membrane fractions, even when methylated, had no influence on tumor growth or animal survival. (21 refs)

78-7049 Angioimmunoblastic Lymphadenopathy, Immunoblastic Sarcoma of B Cells. Animal Model: Autoimmune-prone Inbred Armenian Hamster. (Eng) Yernool, G. (Sidney Farber Cancer Inst., 44 Binney St., Boston, MA, 02115); Gagnon, H. J.; Battaglini, A. *Am J Pathol* (1): 209-212; 1978.

An animal model of angioimmunoblastic lymphadenopathy (AIL) and immunoblastic sarcoma of B cells (IBS) was developed using autoimmune-prone inbred Armenian hamsters treated with nitroso compounds. When the 16th- to 17th-generation animals were 482 days old, 25 were treated with methyl nitrosourea (MNU: 70 mg/kg). Within the period 88-390 days after treatment, 12 animals developed massive hepatomegaly and mesenteric lymph node tumors with or without Peyer's patches, pulmonary lesions, and bilateral renal medullary hyperplasia. The 9-yr tumor-free status of the colony was interrupted when 12 untreated young adults developed lesions identical to those noted in the older treated animals. The characteristic presentation of the massive mesenteric lymph node tumors following MNU treatment suggests that these latent B-cell lesions may be selectively responsive to a new and persisting antigen, with the development of AIL when cells are diploid (polyclonal) and the development of IBS when cells are monoclonal. The MNU-associated, spontaneous Armenian hamster AIL and the transplantable IBS feature the characteristic triad of lesions associated with clinical AIL, ie, the multicellularity of proliferating B cells, an extensively arborizing vasculature, and deposits of eosinophilic material. (14 refs)

78-7050 Marek's Disease: Effect of Virus Pathogenicity and Genetic Susceptibility on Responses of Peripheral Blood Lymphocytes to Concanavalin-A. (Eng) Schat, K. A. (Dept. Avian and Aquatic Animal Medicine, New York State Coll. Veterinary Medicine, Cornell Univ., Ithaca, NY, 14853); Schultz, R. D.; Calnek, B. W. In: *Advances in Comparative Leukemia Research 1977. Proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, The Netherlands on 22-26 August, 1977.* Bentvelzen, P.; Hilgers,

J.; Yohn, D. S., eds. (New York: Elsevier/North Holland Biomedical Press): 503 pp.; 183-186; 1978.

The response of peripheral blood lymphocytes (PBL) to concanavalin A (Con A) was compared using genetic susceptibility to Marek's disease (MD) and virus pathogenicity as variables. A transient depression of the response of PBL to Con A occurred 7 days postinjection of MD-resistant N-line and MD-susceptible P-line chickens with the nononcogenic (SB-1) and oncogenic (GA-5) isolates of MD virus. A permanent depression, starting 2 wk postinjection, developed only in susceptible P-line chickens inoculated with GA-5. (7 refs)

78-7051 Lack of Correlation Between In Vivo Rejection of Syngeneic Fibrosarcomas and In Vitro Non-specific Macrophage Cytotoxicity. (Eng) Evans, R. (Dept. Tumour Immunology, Chester Beatty Res. Inst., Inst. Cancer Res., Clifton Ave, Belmont, Sutton, Surrey SM2 5PX, England); Booth, C. G.; Spencer, F. *Br J Cancer* 38(5): 583-590; 1978.

The relationship between in vitro antitumor activity and in vivo tumor rejection was studied using C57BL mice immunized against two syngeneic benzpyrene-induced fibrosarcomas (FS1 and FS6). After the mice were immunized by sc injection of 10^6 tumor cells and subsequent tumor excision, ip injections of graded doses of FS1 or FS6 cells resulted in a totally specific rejection (in terms of survival). Similarly, the peritoneal inflammatory response (in terms of cell content) was specific; on the peak days of cell production (5-7 days postchallenge) the macrophage content was approx 34%-43%. Conversely, macrophage monolayers prepared from the peritoneal exudates of immunized mice were generally nonspecifically cytotoxic, though occasional specific cytotoxicity was detected. In a 48 hr assay, peritoneal cells from immunized mice lysed ^{125} I-labeled tumor cells in a specific manner, and the loss of lytic potential after pretreatment of cells with anti- θ serum indicated that the reaction was T lymphocyte-mediated. Winn tests further showed that immune peritoneal exudate cells protected normal mice against specific tumor cells, but had no effect on unrelated tumor appearance or growth. It is concluded that the specific rejection of FS1 and FS6 was mediated by T lymphocytes, with possible assistance from macrophages. (16 refs)

78-7052 Host Genetic Factors Influencing the Occurrence of Leukemoid Reaction in BALB/cMk Mice Bearing Transplanted Tumors. (Eng) Kodama, T. (Lab. Pathology, Cancer Inst., Hokkaido Univ. Sch. Medicine, Sapporo 060, Japan); Kawamura, T.; Kobayashi, H. *Cancer Res* 38(12): 4662-4664; 1978.

Hybridization experiments involving BALB/cMk and C57BL/6 mice were carried out to study host genetic factors influencing the occurrence of the leukemoid reaction in

BALB/cMk mice bearing transplanted syngeneic fibrosarcomas. All BALB/cMk mice and all BALB/cMk x C57BL/6 F₁ (CB6F₁) and C57BL/6 x BALB/cMk F₁ (B6CF₁) hybrids, but none of the C57BL/6 mice, were tumor growth-positive (G+) and leukemoid reaction-positive (R+) after fragments of the BALB/cMk tumor CMT-6 were implanted under their back skin. Segregants that were G-,R- were found among the F₂ and C57BL/6-backcross mice. The ratio of G+,R+ to G-,R- mice was 22:8 in the F₂ mice and 14:12 in the C57BL/6-backcross progeny. Neither G+,R- mice nor G-,R+ mice were found. Mice in the parental C57BL/6, F₂, and C57BL/6-backcross groups that were G-,R- were again made tumor-bearing with the C57BL/6 BMT-3 tumor and tested for tumor growth and the leukemoid reaction. G+ mice were observed in all C57BL/6, F₂, and C57BL/6-backcross mice. No R+ mice were found. No BALB/cMk mice, but all C57BL/6 mice with BMT-3 tumor implants were G+. Neither strain with BMT-3 implants was R+. All CB6F₁ and B6CF₁ mice were G+,R+. Segregants that were G-,R- and those that were G+,R- were found among the F₂ and BALB/cMk-backcross mice. Although the ratio of G+:G- mice was 19/7 in F₂ mice, a 12:7 R+:R- ratio was obtained in F₂ mice, seven G+,R- mice, were found. In the BALB/cMk-backcross progeny, the ratio of G+,R+ to G-,R- mice was 14:11. G+,R+ mice were observed in all BALB/cMk, F₂, and BALB/cMk-backcross mice again made tumor-bearing with CMT-6 tumors. These results, the segregation of a positive or negative leukemoid reaction in BALB/cMk mice, were in good agreement with a single dominant gene model and did not fit a two-gene interpretation. The results suggest that occurrence of the leukemoid reaction is regulated genetically in BALB/cMk mice bearing transplanted tumors. (15 refs)

78-7053 Characteristics and In Vitro Growth Influencing Effects of the Spleen Cell Population in a Methylcholanthrene Induced Mouse Sarcoma System. (Eng) Cornain, S. (Dept. Tumor Biology, Karolinska Institutet, S-10401, Stockholm 60, Sweden); Klein, E. *Z Immunitaetsforsch* 154(2): 101-114; 1978.

The effects of immunization on the growth of two transplantable methylcholanthrene-induced fibrosarcomas (MC57 X and MC57 Y) were studied in syngeneic C57Bl mice and their (ABY x C57Bl)F₁ hybrids. For immunization, three injections of 10⁶ x-irradiated tumor cells were given sc at 10-day intervals. Ten days after the last injection, the mice were challenged sc with viable tumor cells (10³, 10⁴, 10⁵, 10⁶, 5 x 10⁶, or 5 x 10⁷ cells); 24 hr before challenge, the mice received 400 R total-body radiation. Preimmunization resulted in complete protection against tumor X, but it enhanced the growth of tumor Y. The enhancement was specific, since the effect was absent in mice pretreated with X cells. The spleens of tumor-bearing animals were enlarged, and they contained a high percentage of blasts; the proportion of complement 3 (C3)- and/or Fc-receptor cells was increased even in the T-cell enriched population. These changes were also observed

to a lesser extent in the lymph nodes. The in vitro growth of tumor cells was inhibited by C3-receptor cells from the spleens of control and tumor-bearing mice and by T-cell fractions from tumor-bearer spleens. The T cells that inhibited target cell growth were detected 14 days after tumor challenge; cells obtained 7 days postinoculation, ie, before tumor growth, enhanced target cell growth. The inhibitory effect of the T cells was nonspecific. The results of the in vitro test reflected only the tumor status of the host and did not relate to preimmunization. (21 refs)

78-7054 Cooperation Between Immune T Cells and Normal Macrophages in the Expression of Tumor Immunity in Chickens. (Eng) Palladino, M. A. (Dept. Pathology, New York Univ. Sch. Medicine, 550 First Ave., New York, NY, 10016); Thorbecke, G. J. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 86-96; 1977.

The possibility that the adoptive tumor-specific immunity expressed in the Winn assay by spleen cells from normal and bursectomized (Bu-x) chickens is mediated by a collaboration between immune T cells and normal recipient monocytes was investigated. Local tumor-specific immunity was transferred into 2- to 4-wk-old recipients with immune T cells from B2/B2 chickens that had rejected B2/B2 fibrosarcoma transplants with the aid of *Corynebacterium parvum*. Immunity was induced and transferred in the total absence of B cells in donors and recipients. Immunity was transferred in normal but not in irradiated or silica-pretreated recipients. Irradiated recipients were reconstituted by (1) iv injection of Bu-x spleen or bone marrow cells and (2) local addition of Bu-x spleen or purified monocytes to the tumor + immune T-cell inoculum. The induction of a delayed hypersensitivity reaction to human gamma-globulin (HGG) in a tumor cell injection site also resulted in tumor immunity. HGG immune spleen cells failed to inhibit tumor cell growth locally unless HGG was also injected in the same site. This immunity, mediated by an adoptive local delayed reaction, could also not be demonstrated in irradiated recipients and thus mimicked tumor-specific immunity in requiring cooperation between immune T cells, antigen, and macrophages. Activation of macrophages through local delayed reactions is concluded to be of paramount importance in obtaining tumor growth inhibition in Winn tests in chickens. (26 refs)

78-7055 Reduced Spontaneous Antitumor Resistance of the Elderly Rat is Restored by *Corynebacterium parvum*. (Eng) Keller, R. (Immunobiology Res. Group, Univ. Zurich, Schonleinstrasse 22, CH-8032 Zurich, Switzerland). *Br J Cancer* 38(4): 557-560; 1978.

The inoculation of DA rats with *Corynebacterium parvum*

increased the in vivo resistance of rats (in terms of survival) aged 30 days or 3-4, 12, or 16-18 mo to the later inoculation of 7,12-dimethylbenzanthracene (DMBA)-induced fibrosarcoma ascites tumor cells. The survival of rats inoculated with syngeneic DMBA-induced tumor cells alone decreased with age and was independent of body wt. However, when the rats were inoculated ip with 3 mg heat-killed strain 2683 *parvum* 1 wk before tumor challenge, local tumor resistance was increased similarly in rats aged 30 days and 3-4 mo; both age groups, *C. parvum* treatment was well-supported and survival time after tumor cell injection was prolonged significantly. Pretreatment with *C. parvum* was much more toxic for rats aged 12 and 16-18 mo and caused the loss of 10%-30% of these animals within 10 days. In the survivors, resistance to tumor challenge varied. In 5/6 experiments, the pretreatment caused a distinct enhancement of local tumor resistance that was sometimes more pronounced than that in younger animals; in one experiment, only minor simulatory effects were noted. (30 refs)

78-7056 In Vivo Anti-tumour Activity of *C. parvum* Stimulated Peritoneal Exudate Cells. (Eng) McBride, W. H. (Dept. Bacteriology, Univ. Edinburgh Medical Sch., Edinburgh, Scotland); Peters, L. J.; Mason, K. A.; Morrow, G.; Milas, L. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 1973-181; 1977.

The antitumor activity of *Corynebacterium parvum*-stimulated peritoneal exudate cells (CP-PEC) was investigated in 3Hf/Bu/E mice in a Winn-type sc transfer assay. The cells inhibited tumor growth when they were admixed with 10^3 or 10^4 fibrosarcoma cells (Fsa) and injected into the mice. When the ratio of CP-PEC:Fsa was 10:1 or higher, there was a significant delay before tumors were detected and a lower final incidence of tumor takes. Normal cells or plastic microspheres interfered with the action of the CP-PEC, presumably by interposing themselves between the target and effector cells. In other experiments, mice were inoculated with 4×10^5 Fsa sc and with three doses of CP ip. There was little activity in the PEC from normal tumor-bearing mice not treated with CP. In contrast, the level of activity in PEC from CP treated normal and tumor-bearing mice was considerable, even at an effector:target cell ratio of 2.5:1. This activity level was maintained in the tumor bearers for some time after injection of CP, but it declined in the normal mice. This suggests that there is some interplay between the stimulus provided by the tumor and that provided by CP, possibly involving a specifically immune T-cell population. There was an unexpected observation of depressed reactivity in the splenic T-cell pool from CP-treated tumor bearers, a finding that contrasts with the enhanced activity of splenic T cells upon iv transfer. This conflict may be due to the different abilities of T-cell subsets to migrate to the tumor site. (19 refs)

78-7057 A Study on Cells Participating in the Antitumour Effect of *Corynebacterium Parvum*. (Eng) Stiffel, C. (Dept. Immunogenetics (U 125 INSERM and ER 70 CNRS), Fondation Curie, Institut du Radium, 26 rue D'Ulm, 75 231 Paris Cedex 05, France); Mazurek, C.; Chalvet, H.; Biozzi, G. In: *The Macrophage and Cancer. Proceedings of the European Reticuloendothelial Society Symposium held in Edinburgh, September 12-14, 1977.* European Reticuloendothelial Society (Edinburgh, Scotland): 448 pp.; 164-172; 1977.

The effects of x-rays and silica on the activity of *Corynebacterium parvum* were studied in two quantitative systems: (1) phagocytosis and antibody synthesis in C3H mice and (2) mortality induced by a transplantable syngeneic mammary carcinoma. Ip injections of 5 mg silica (a specific macrophage toxin) or 500 rads whole-body irradiation inhibited *C. parvum*-induced stimulation of the phagocytic activity of liver and spleen macrophages when given two days before, but not two days after, *C. parvum* administration. The same irradiation dose totally abolished the antibody response to 10^8 sheep red blood cells (SRBC) injected 3 or 10 days later, but partial antibody synthesis occurred when the antigen was injected 20 days after irradiation and agglutination titers were normal in mice immunized at 40 days. Silica had no effect on the antibody response to SRBC or on the adjuvant effect that *C. parvum* supplied to the reaction. Silica injected 2 days before or 4 days after *C. parvum* abolished the bacteria's normal antitumor effect (ie, survival) against the transplanted mammary carcinoma. Irradiation also abolished this effect when administered 3 or 10 days before to *C. parvum*. The antitumor activity of *C. parvum* was restored totally when it was injected 40 days after irradiation. Both silica and x-rays had a similar effect on *C. parvum* antitumor capabilities and macrophage phagocytosis, but only x-rays inhibited antibody synthesis in response to SRBC. (19 refs)

78-7058 Tumor-directed Antibodies in Cancer: A Study of Immune Complexes Eluted from Kidneys of Humans and Experimental Animals with Malignancies. (Eng) Slovin, S. F. (Columbia Univ., New York, NY, 10027). *Diss Abstr Int B* 39(4): 1700-1701; 1978. (no refs)

78-7059 Reduction of Respiratory Tract Binding of Benzo(a)pyrene in Mice by Immunization. (Eng) Moolten, F. (Dept. Microbiology, Boston Univ. Sch. Medicine, Boston, MA, 02118); Capparell, N.; Boger, E. *J Natl Cancer Inst* 61(5): 1347-1349; 1978.

The feasibility of immunizing the respiratory tracts of mice with 5-fluoro-12-methylbenzanthryl-7-acetic acid (FMBAAA)-bovine serum albumin (BSA) conjugates was investigated. Male inbred A/J mice were immunized ip with the FMBAAA-BSA conjugate (0.2 mg antigen) in two doses 2 wk apart. Intranasal immunization (30 μ g antigen in 5 ml

buffer soln was pipetted onto each nostril) was begun 1 wk after the second ip injection and continued 3x/wk for six instillations. Five days after the last dose of antigen, each mouse received two instillations of benzo(a)pyrene (BP) in each nostril 2 hr apart [total dose 92 nanograms (ng)]. The mice were killed 20 hr later, and BP was measured in the nose and tracheal tissues. Total BP bound to these respiratory tissues was 0.0334 ng in animals that received FMBAAA-BSA conjugate, 0.0502 ng in unimmunized mice, 0.0527 ng in BSA-treated mice, and 0.0507 ng in mice treated with an unconjugated mixture of FMBAAA and BSA. γ -Globulin fractions eluted from tracheas of immunized, but not unimmunized, mice exhibited significant BP-binding capacity, indicating that the reduction of bound BP in immunized mice was caused by the interception of BP by antibodies in respiratory secretions. The one-third reduction of BP in the respiratory tracts of the FMBAAA-BSA-immunized mice given 92 ng of BP is equivalent to a reduction in exposure of 30 ng. The data suggest the potential capacity of respiratory tract immunization to cope with significant human exposures to BP and other polycyclic aromatic hydrocarbon carcinogens. (5 refs)

78-7060 Gamma Heavy Chain Disease in Man: Translation and Partial Purification of mRNA Coding for the Deleted Protein. (Eng) Alexander, A. (Dept. Biochemical Sciences, Princeton Univ., Princeton, NJ, 08540); Barritault, D.; Buxbaum, J. *Proc Natl Acad Sci USA* 75(10): 4774-4778; 1978.

Long-term-cultured lymphoid cells from a patient with heavy chain disease continue to produce a protein antigenically identical to the deleted heavy chain disease protein found in the patient's serum. The size of the messenger RNA (mRNA) coding for this protein was determined, and evidence for the presence or absence of translatable light (L)-chain mRNA was sought. The primary in vitro translation product was 1,500-2,000 daltons larger than the polypeptide portion of the cytoplasmic or secreted protein. Methionine was present at the amino terminus of the translation product. The mRNA sedimented at 15.5S on sucrose gradients and, therefore, appears to be smaller than the 17S mRNA coding for normal-sized mouse γ chains. It contained a base sequence that coded for a hydrophobic amino-terminal peptide not found in the cytoplasmic protein. There was no evidence for the synthesis of translatable L-chain mRNA by these cells. The data suggest that the protein resulted from a primary somatic genetic event that gave rise to a cell product bearing a normal aminoterminal sensitive to limited proteolytic digestion. The serum protein thus appears to begin in the hinge region but, in fact, contains a normal heavy chain initiation site. (28 refs)

78-7061 Cancer Related to Asbestos Exposure: Immunological Studies of Patients at Risk. (Eng) Ka-

gan, E. (Dept. Immunology, Natl. Res. Inst. Occupational Diseases, South African Medical Res. Council, Johannesburg, S. Africa); Solomon, A.; Cochrane, J. C.; Beissner, I.; Gluckman, J.; Kuba, P.; Webster, I. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held at the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976*. International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1631-1638; 1978.

Several immunological parameters were evaluated in two groups of asbestos-exposed patients who were at possible risk of developing asbestos-related cancer. Group 1 consisted of 20 patients with asbestos-related pleural thickening; Group 2 consisted of 26 patients with parenchymal asbestosis. Comparatively minor immunological abnormalities were noted among Group 1 patients, and they were generally restricted to isolated disturbances of humoral immunity. All control subjects and all except 2 Group 1 patients exhibited strong cutaneous delayed hypersensitivity reactions to 2,4-dinitrochlorobenzene (DNCB), but 17/26 Group 2 patients manifested cutaneous anergy, indicating defective expression of cell-mediated immunity. In vitro assays also showed significantly impaired cellular immunity in Group 2. Levels of secretory IgA and serum concentrations of IgA, IgG, IgM, and IgE were significantly increased in Group 2, but serum IgA and IgG levels were significantly reduced in Group 1. The percentage and absolute numbers of T lymphocytes were significantly lower in Group 2 compared with controls. Cold-reactive serum lymphocytotoxins were detected in 55% of Group 1 and 81% of Group 2 patients vs 22% of controls. These antibodies showed autoreactivity in 25% of both groups vs 7% of controls. The imbalance between cellular and humoral immunity seen in these patients may be a manifestation of abnormal T-cell/B-cell interaction and may have relevance to the possible causal role of asbestos in the induction of malignancy. (28 refs)

78-7062 Early Increase of Serum Alpha-Fetoprotein in Spontaneous Hepatocarcinogenesis in Mice. (Eng) Jalanko, H. (Dept. Serology and Bacteriology, University of Helsinki, 00290 Helsinki 29, Finland); Virtanen, I.; Engvall, E.; Ruoslahti, E. *Int J Cancer* 21(4): 453-459; 1978.

The concentration of serum alpha-fetoprotein (AFP) was followed by competitive double antibody radioimmunoassay in C3H mice having a high incidence of spontaneous hepatocellular carcinoma. No general elevation of serum AFP levels with age was seen in mice without tumors. With a single exception, mice bearing hepatocellular carcinomas had increased serum AFP levels. In some mice this increase followed a biphasic course. Mice killed within 1 mo of the time when an elevation of serum AFP was first observed had small tumors or no detectable tumor. Premalignant lesions were present in the livers of 11/16 mice that had elevated AFP b

cancer, but only 1/14 mice with normal AFP had such lesions. Autologous antibodies to AFP were produced in a group of C3H mice by immunization with rat AFP. These AFP antibodies reduced the amount of serum AFP but had no effect on the incidence of spontaneous hepatomas. The results suggest that spontaneous hepatocarcinogenesis proceeds through almost the same premalignant lesions as chemically induced carcinogenesis and that an increase in AFP production occurs early during this process, often preceding macroscopic lesions. Repeated testing for AFP in individuals at risk of developing liver cancer might reveal tumors at a curable stage. (24 refs)

7063 A Microassay for the Detection of Tumour-specific Complement-dependent Serum Cytotoxicity Against a Chemically Induced Rat Hepatoma. (Eng) Price, M. R. (Cancer Res. Campaign Labs., Univ. Nottingham, University Park, Nottingham NG7 2RD, England). *Transplantation* 25(4): 224-226; 1978.

A short-term ^{51}Cr -release microassay was developed for the detection of tumor-specific complement-dependent serum cytotoxicity against a transplanted 4-dimethylaminoazobenzene-induced rat hepatoma (D23). The method involves the sequential sampling of supernatants in the wells of microtiter plates containing serum (100 μl), hepatoma cells (10^4 cells/0.1 ml), and guinea pig serum (100 μl) as complement. In a representative test, the complement-dependent cytotoxicity of hepatoma D23-bearer serum and normal rat control serum was determined after 2-, 4-, and 8-hr incubations. The complement-dependent cytotoxicity of tumor bearer serum was present at the three time intervals, even at dilutions of 1/80 or 4 hr of incubation. Using this assay, the serum from rats bearing ip grafts of hepatoma D23 exhibited tumor-specific complement-dependent cytotoxicity for hepatoma D23 cells. Serum from rats bearing sc tumor grafts also exhibited tumor-specific complement-dependent cytotoxicity, in contrast to previous reports. Attempts to use rabbit serum as a source of complement met with variable success. The application of this technique economizes on the time, labor, and reagents required to perform the test in individual plates. (11 refs)

7064 Circulating Immune Complexes in Rats Bearing Chemically Induced Tumours. II. Characterization of Sera from Different Stages of Tumour Growth. (Eng) Witten, K. (West Deutsches Tumorzentrum, Innere Universitätsklinik, Hufelandstrasse 55, D-4300 Essen 1, W. Germany); Price, M. R.; Moore, V. E.; Baldwin, R. W. *Int J Cancer* 22(5): 576-582; 1978.

Sera of WAB/Not rats bearing ip implants of a 4-dimethylaminoazobenzene-induced hepatoma were fractionated

by Sephadex G200 and diethylaminoethylcellulose ion-exchange column chromatography and then examined for their capacity to bind radioiodinated complement C1q to immune complexes, to bind soluble tumor-specific antigen, and to react with antigens on the surface of hepatoma cells. Sera were obtained from rats bearing implants for 3 and 9 days, since these samples display significantly higher and lower capacities, respectively, to bind C1q compared with normal rat serum. Immune complexes were present at both early and late stages of tumor growth; however, inherent limitations in the C1q test prevented their detection in unfractionated sera from late tumor bearers. Fractionated sera from early and late tumor-growth stages displayed significantly greater C1q-binding activity than corresponding normal rat serum fractions in the region 7S-19S, but in unfractionated sera, only samples from rats at early tumor stages exhibited this activity. The results suggest that the detection of immune complexes in unfractionated late tumor-bearer serum by ^{125}I -C1q fixation is masked by the presence of excess tumor-specific antibodies and by a shift from complement-fixing to noncomplement-fixing antibodies. (25 refs)

78-7065 Compared Humoral and Cellular Immune Response to Herpes Simplex (HSV-2) Antigens in Women with Cervical Carcinoma. (Eng) Thiry, L. (Institut Pasteur du Brabant, Brussels, Belgium); Sprecher-Goldberger, S.; Hannecart-Pokorni, E.; Gould, I.; Bossens, M. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 231-243; 1977.

The specific immune status of patients with cervical dysplasia and cervical carcinoma was characterized by determining their specific humoral and cell-mediated responses and comparing the results with those of a control group. The test subjects comprised (1) 30 with a normal cervix, (2) 33 with dysplasia, (3) 32 with cervical carcinoma, (4) 11 cancer patients undergoing radiation therapy, and (5) 35 patients that had been treated (24, surgically). Neutralizing antibodies (ab) to herpes simplex virus type 2 (HSV-2) were studied by neutralization kinetics before and after adsorption of serum samples with *Staphylococcus Cowan A*. Ab remaining in the supernatant were non-IgG ab. These ab were found in 6%, 42%, 66%, 27%, and 6% of the respective five groups. Heparinized blood was obtained from most of these patients. Thymidine incorporation into washed nonpurified lymphocytes was studied in the presence of killed HSV-2 and HSV-1, as well as in the presence of mitomycin C-treated hamster cells transformed by HSV-2 or simian virus 40 or nontransformed cells. Specific stimulation by the HSV-2-transformed cells was obtained in 0%, 24%, 36%, 82%, and 11% of the five groups. Differences among the various groups were less clear-cut when HSV-2 virions were

used as antigens to stimulate the lymphocytes (24%, 37%, 45%, 40%, and 10%, respectively, had lymphocytes responding to the virions). The lymphocyte responses were rarely associated with the non-IgG ab. Some in vivo mechanism must modulate the immune reactions in such a way that the simultaneous presence of non-IgG ab to HSV-2 and lymphocyte responsiveness to this virus is excluded. There was also a negative correlation between the humoral and cell-mediated responses. (16 refs)

- 78-7066 Carcino-embryonic Antigen (CEA) and Alpha-Fetoprotein (AFP) in Cancer-prone Families.** (Eng) Guirgis, H. (Creighton Univ., Omaha, NB); Lynch, H.; Keller, R.; Tomasi, T.; Vandervoorde, J.; Lynch, J.; Rankin, L.; Maloney, K.; Harris, R. In: *Prevention and Detection of Cancer. Proceedings of the Third International Symposium on Detection and Prevention of Cancer Held by the International Study Group for the Detection and Prevention of Cancer in New York, April 26 - May 1, 1976.* International Study Group for the Detection and Prevention of Cancer. (New York, NY): Vol. 2(part 1), 2404 pp.; 1563-1571; 1978.

Carcinoembryonic antigen (CEA) and/or α -fetoprotein (AFP) levels were measured in 698 individuals (521 relatives and 177 spouses) from 11 cancer-prone families. CEA was determined by the Hansen method and AFP by a fluorescence technique. The families were divided into the following categories: (1) breast cancer kindreds (6 families); (2) cancer family syndrome (CFS) kindreds (3 families); (3) malignant melanoma kindred (1 family); and (4) Hodgkin's disease kindred (1 family). Individuals were classified as smokers or nonsmokers. Mean plasma CEA titers/family were 1.11-3.08 ng/ml with standard deviations in the range 0.46-9.13 ng/ml. There were significant intrafamily variations and even greater interfamily variations. Significant elevations of CEA were detected in the spouses of CFS relatives and in nonsmoking spouses of patients from breast cancer-prone families. In the two families tested for AFP, 50% of the individuals were positive for AFP on their lymphocytes. No correlation was found between elevated plasma CEA titers and the AFP test, but there was a significant correlation between a positive AFP test and the suppression of stimulated lymphocytes by the patient's sera. The significance of these results remains to be determined. (17 refs)

- 78-7067 Incidence of Spontaneous Tumours in Neonatally Thymectomized Rats.** (Eng) Dawson, P. J. (Dept. Pathology, Univ. Chicago, Chicago, IL, 60637); Fieldsteel, A. H.; McCusker, J. *Br J Cancer* 38(3): 476-478; 1978.

The spontaneous tumor incidence in neonatally thymectomized (Thy-x) Wistar/Lewis and Buffalo rats was compared with that in intact rats. In addition, 214/1,007 Thy-x Wistar/Lewis rats also were treated with antithymocyte serum (ATS) and/or 250-450 rads of whole-body x-radiation. The most common ATS dose used was 0.5-2.4 ml ip weekly for 3-6 mo beginning at 2-3 mo of age. A total of 33/51 Thy-x Buffalo rats also were irradiated, and 16 of these were given 0.5 ml ATS ip weekly for 6 mo beginning at 2-7 mo of age. Three intact Buffalo and five intact Wistar/Lewis rats received 0.5 ml ATS weekly for 3 mo beginning at 2-7 mo of age. Buffalo rats (24) also were irradiated after they were inoculated with *Mycobacterium leprae* into the foot pads. Tumors were noted in 1/23 male and 10/28 female Thy-x and 2/4 male and 17/32 female intact Buffalo rats. For Wistar rats, the respective figures were 43/517 and 66/490 for the Thy-x animals and 4/65 and 6/43 for intact animals. Breast tumors were the most common tumors, followed by leukemia and fibrosarcomas. It was concluded that the incidence of spontaneous tumors in the two strains was not influenced by prolonged immunosuppression. (11 refs.)

- 78-7068 Immune Response of Prostatic Cancer Patients to Cytomegalovirus-infected and -transformed Human Cells.** (Eng) Laychock, A. M. (Milton S. Hershey Medical Center, Pennsylvania State Univ. Coll. Medicine, Hershey, PA); Gedér, L.; Sanford, E. J.; Rapp, F. *Cancer* 42(4): 1766-1771; 1978.

The possibility of a relationship between cytomegalovirus (CMV) infection and prostatic cancer was investigated. The indirect immunofluorescence (IF) test was used to determine the prevalence of humoral immunity to CMV-induced antigens in 35 patients with prostate cancer, 22 with benign prostatic hyperplasia, 17 with transitional cell carcinoma (TCC) of the bladder, and 18 nonurogenital cancer controls. Sera from the four groups were screened for antibodies to the CMV-transformed cells CMV-Mj-HEL-2 and CMV-Mj-HEL-2,T-1. A high percentage of sera from all test groups reacted when tested against fixed and unfixed preparations of CMV-Mj-HEL-2,T-1 cells at passages 46 and 56, respectively. There were no major differences among the groups when the sera were tested against unfixed cells. With fixed cell preparations, approx 80% of serum specimens from prostatic cancer and bladder cancer patients demonstrated fluorescence. In IF tests against cells infected with strain CMV-Mj, the prostatic cancer group had the fewest patients with titers $< 1:4$ (3%) but the most patients with titers $\geq 1:64$ (66%). Sixty-five percent of the patients with TCC also possessed antibody titers $\geq 1:64$. Thirty-two percent of the patients with benign hyperplasia had antibody titers $\leq 1:4$, and only 32% reacted with CMV-Mj-infected cells in serum dilutions $\geq 1:64$. In the controls, 39% had titers $< 1:4$, 44% had titers between 1:4 and 1:64, and only 22% had titers $\geq 1:64$. A significantly higher number of prostatic cancer patients reacted with the CMV-Mj-infected cells at a serum dilution

4 or higher than did patients with benign hyperplasia or controls. There was no significant difference in reactivity between the prostatic cancer patients and the TCC patients, though a higher ratio of the prostatic cancer group had V-Mj antibodies. When screened against CMV-transfected human cell lines, the reactivity of the sera followed rate of expression of CMV-related antigens of cell lines in these tests. The data enhance the possibility of finding etiologic association between CMV and prostatic cancer. (refs)

See also:

*(Rev.): 78-6671, 78-6681, 78-6682, 78-6684, 78-6687, 78-6688, 78-6689, 78-6690, 78-6691, 78-6692, 78-6693, 78-6694, 78-6695.

*(Chem.): 78-6735, 78-6736, 78-6782.

*(Viral): 78-6860, 78-6871, 78-6875, 78-6879, 78-6883, 78-6884, 78-6890, 78-6909, 78-6911, 78-6919, 78-6929, 78-6939, 78-6956.

*(Path.): 78-7071, 78-7101, 78-7110, 78-7136.

*(Epid.-Biom.): 78-7165, 78-7169.

PATHOGENESIS

- 78-7069 Chromosomal Evidence for the Transplantability of the Shay Chloroleukemia.** (Eng) Derelan-ko, M. J. (Dept. Biology, 952 Brown Bldg., Washington Sq., New York, NY, 10003); Moorthy, A. S.; LoBue, J.; Meagher, R. C.; Gordon, A. S.; Mitra, J.; Fredrickson, T. N. *J Surg Oncol* 10(5): 441-445; 1978.

Female Long-Evans rats were injected iv with Shay chloroleukemia (SCL) cells, and the resulting tumor cells were studied to determine whether or not SCL is a true transplantable leukemia resulting from colonization of donor leukemic cells. During blast crisis in the recipient rats, the leukemic blast cells were mitotically active in the peripheral circulation, and a large number of mitotic figures were found. Karyotype analyses of the SCL tumor cells used to induce the leukemia demonstrated the presence of X and Y chromosomes. Similarly, all 180 metaphase figures of leukemic blast cells from the peripheral circulation of female rats inoculated with SCL showed the X and Y chromosomes. The data demonstrate that transmission of SCL is a result of donor cell colonization and that SCL can be considered a true transplantable leukemia. Viruses described in SCL by other workers are probably not involved in the transmission of this leukemia. (20 refs)

- 78-7070 Chromosomes and Causation of Human Cancer and Leukemia: XXVIII. Value of Detailed Chromosome Studies on Large Numbers of Cells in CML.** (Eng) Sonta, S. (Roswell Park Memorial Inst., 666 Elm St., Buffalo, NY, 14263); Sandberg, A. A. *Am J Hematol* 3: 121-126; 1977.

Bone marrow aspirates, peripheral blood, and, in one case, splenic tissue from six patients with Philadelphia chromosome (Ph¹)-positive chronic myelocytic leukemia (CML) were subjected to chromosomal analyses. Clones with additional chromosome abnormalities (besides a Ph¹ and a missing Y chromosome) were shown to exist in the uncomplicated, chronic phase of CML when a large number of cells (110 to 500) was examined. These abnormalities were not evident during routine cytogenetic study (10 to 50 cells). The abnormalities were an extra Ph¹ chromosome, missing chromosomes 16 and 17, an isochromosome of the long arm of No. 17, and an abnormal chromosome 2. In 5/6 cases, aneuploid clones observed during the examination of 110 to 500 cells during the chronic phase became the dominant ones during the blastic phase. The data indicate that in some cases of CML, a variety of karyotypically abnormal cells is already present in the marrow during the chronic phase and that the quantitative shift in favor of these abnormal cells as a result of their increased proliferative capacity constitutes the

chromosomal aspects of the blastic phase, rather than the de novo genesis of cells with chromosome changes previously not seen in the marrow. (8 refs.)

- 78-7071 An Autopsy Case of Acute Myeloblastic Leukemia with Persistent High Anti-EBV-antibody Titer.** (Jpn) Musashi, M. (Third Dept. Internal Medicine Hokkaido Univ. Sch. Medicine, Hokkaido, Japan); Maekawa, I.; Sakurada, K.; Fujimoto, N.; Noguchi, H.; Shiraiishi, T. *Jpn J Clin Hematol* 19(8): 1079-1085; 1978.

A case of acute myeloblastic leukemia (AML) with persistent high anti-Epstein Barr virus (EBV)-antibody titer is presented. A 62-yr-old woman presented with fatigue, palpitation, and shortness of breath. Physical examination revealed anemia, but there was no hepatosplenomegaly or superficial lymphadenopathy. Examination of peripheral blood showed moderate anemia, thrombocytopenia, and no atypical cells. However, 4.5% of atypical cells were observed in a bone-marrow specimen; these cells were positive for peroxidase activity and Sudan Black B stain and nucleolus staining showed M-type cells. These results prompted the diagnosis of AML. Chromosome study of the marrow cells revealed karyotype (46xx, 5q-). Two mo later, a tumor was found at the middle nasal meatus that proved to be a well-differentiated hemangioma. Examination of the patient's anti-EBV titer revealed a titer of 1:1280 for antibody to viral capsid antigen (anti-VCA) of 1:80 for antibody to erythrocyte antigen (anti-EA). Anti-VCA and anti-EA persisted at high levels (1:640 and 1:320, respectively) throughout the clinical course. After the appearance of the nasal tumor, the number of leukemic cells increased in the peripheral blood and bone marrow, and a marked hepatosplenomegaly developed. Autopsy revealed leukemic cell proliferation and infiltration in the bone marrow, liver, and spleen. The occurrence of AML with high anti-EBV is rare, and the relationship between AML and EBV infection found in the present case is discussed. (27 refs)

- 78-7072 Leukemia in Radiation-treated Patients: Cytogenetic Studies in Eight Cases.** (Eng) Goh, K. (Medical Oncology-Hematology Unit, Monroe Community Hosp., Rochester, NY, 14603); Bauman, A.; Bakemeier, R.; Lee, H.; Woll, J. E. *Am J Med Sci* 276(2): 189-195; 1978.

The case reports and cytogenetic studies are presented of eight patients (6 women, 2 men; age range 17-66 yr) who developed leukemia several years after diagnosis and treatment for an unrelated primary malignancy. These patients originally had a multiple myeloma, a Wilms' tumor, Hodg-

s disease, a seminoma, a breast cancer, or lymphomas (3 refs). All patients received extensive radiation therapy (100-18,329 rads); two were also treated with nitrogen mustard and one each with chlorambucil, cyclophosphamide, and doxorubicin. Cytogenetic studies were performed on all patients by direct methods; and chromosome abnormalities were seen in every patient. Only 42% of the karyotyped metaphases were diploid, and it was quite common to find less than the full diploid complement of chromosomes. The most consistent abnormal chromosomes, present in 91/516 metaphases analyzed, was a small G chromosome replacing the normal G. Other chromosome abnormalities included ring chromosomes, large acrocentric chromosomes, submetacentric chromosomes, pulverized chromosomes, pseudodiploid metaphases, and chromosome breaks. It is suggested that in these patients there was a causal relationship between previous radiation treatment and subsequent development of leukemia. (19 refs)

78-7073 Lymphoblastoid Cell Lines from Patients with Chronic Lymphocytic Leukemia: Identification of Tumor Origin by Idiotypic Analysis. (Eng) Hurley, J. N. (Rockefeller Univ., New York, NY, 10021); Fu, S. M.; Kunkel, H. G.; McKenna, G.; Scharff, M. D. *Proc Natl Acad Sci USA* 75(11): 5706-5710; 1978.

A large number of continuous lymphoblastoid cell lines were derived from the peripheral blood mononuclear cells of two male chronic lymphocytic leukemia patients with monoclonal IgM bands in the serum. Anti-idiotypic antisera obtained against the isolated IgM band proteins of both patients showed that the serum IgM band, the membrane IgM on the leukemic lymphocytes, and the intracellular IgM in the plasma cells were idiotypically identical. The idiotypic determinants thus constitute tumor-specific antigens. Surface and intracellular immunofluorescence studies using the idiotypic antisera showed that certain cell lines from each patient were derived from the leukemic cells and others were derived from individual normal B lymphocytes. The leukemic lines contained different percentages of lymphoid cells with the idiotype-specific membrane immunoglobulin (Ig) and different percentages of plasma cells with intracellular Ig of the same specificity. Hemagglutination-inhibition analyses of cell line supernatants also demonstrated specific Ig synthesis. Aside from Ig specificity, there were no differences between the leukemic lines and those derived from normal cells. After approximately 6 mo of continuous culture, one leukemic cell line was cloned in soft agarose. The cloning efficiency was approximately 10%. Immunofluorescence studies of seven clones showed that all were of leukemic origin and that they had higher percentages of cells positive for surface IgM than the parental line. (23 refs)

78-7074 A Case of AML Preceded by Fibrinous Bronchitis During the Preleukemic Phase. (Jpn) Nun-

oue, N. (Second Dept. Internal Medicine, Showa Univ. Sch. Medicine, Japan); Kitamura, K.; Tomoyasu, S.; Asano, Y.; Yokoyama, S.; Tsuruoka, N.; Shimizu, M.; Sano, K. *Jpn J Clin Hematol* 19(9): 1226-1233; 1978.

A case of acute myelogenous leukemia (AML) preceded by the preleukemic phase, which developed following admission for fibrinous bronchitis, is reported. A 41-yr-old man presented with fever and recurrent attacks of dyspnea with expulsion of bronchial casts that lasted for several months and was diagnosed as fibrinous bronchitis. About a month after admission, a few abnormal leukocytes appeared in his blood smear. Bone marrow aspirates revealed: NCC $17.3 \times 10^4/\text{cm}^3$, minimal bactericidal level 45.4%, and prothrombin 12.8%. A diagnosis of AML was confirmed. Review of the blood count from the onset of fibrinous bronchitis to the clinical manifestation of AML revealed progressive anemia with anisopoikilocytosis, and an episode of thrombocytosis, monocytosis, and reticulocytosis of unknown cause. The hematologic abnormalities that predated the development of AML were considered to be preleukemic in nature. (32 refs)

78-7075 Quantitative Morphological Evaluation of Basal Lamina Changes During Epithelial Carcinogenesis (Meeting Abstract). (Eng) White, F. H. (Dept. Oral Pathology, Univ. Sheffield, Sheffield, England); Gohari, K. *Clin Sci Mol Med* 55(3): 15P; 1978. (no refs)

78-7076 Evidence for the Progressive Nature of Neoplastic Transformation In Vitro. (Eng) Barrett, J. C. (Div. Biophysics, Sch. Hygiene and Public Health, Johns Hopkins Univ., Baltimore, MD, 21205); Ts'o, P. O. *Proc Natl Acad Sci USA* 75(8): 3761-3765; 1978.

Evidence is presented that neoplastic transformation in vitro is progressive, involving multiple developmental stages. Syrian hamster embryo cell cultures were treated with benzo(a)pyrene (BP) and monitored for the acquisition of in vitro phenotypes associated with neoplasia. Morphological transformation and enhanced fibrinolytic activity were observed within 2 wk and anchorage-independent growth was observed after 6-15 wk. Development of the potential to grow in soft agar did not appear to be related to morphological transformation. One of four spontaneously established Syrian hamster embryo cell lines produced tumors in vivo. None of eight cultures treated with BP for 24 hr produced tumors, but all produced fibrosarcomas in 100% of injected animals after they had acquired anchorage independence in vitro. The data suggest that multiple cellular changes are required for the expression of anchorage-independent growth and that neoplastic transformation in vitro progresses through qualitatively different stages. The results strongly justify the study of oncogenesis in cell cultures as a model for neoplastic transformation in vivo. (24 refs.)

- 78-7077 Genetics of the Large, External, Transformation-sensitive (LETS) Protein: Assignment of a Gene Coding for Expression of LETS to Human Chromosome 8.** (Eng) Owerbach, D. (Roswell Park Memorial Inst., New York State Dept. Health, Buffalo, NY, 14263); Doyle, D.; Shows, T. B. *Proc Natl Acad Sci USA* 75(11): 5640-5644; 1978.

The genetic basis of the large, external, transformation-sensitive (LETS) protein (fibronectin) was analyzed with the use of human-mouse somatic cell hybrids formed from human WI-38 fibroblasts and a mouse L-cell line not expressing LETS. External membrane proteins of these hybrids, which had reduced numbers of human but not mouse chromosomes, were labeled by lactoperoxidase-catalyzed iodination. Cell-surface proteins were identified after sodium dodecyl sulfate/polyacrylamide gel electrophoresis by autoradiography of the dried gel. The LETS protein was identified in parental human cells, and it segregated in the hybrids. The LETS protein segregated concordantly with the chromosome 8 enzyme marker glutathione reductase and human chromosome 8. In addition, dependence on cellular density and trypsin sensitivity were similar for LETS in the hybrid cells and in the human fibroblasts. These findings demonstrate that a gene, *LETS*, encoded on chromosome 8, is responsible for LETS protein expression in humans. Because LETS has been implicated in tumorigenicity and cellular transformation, it is of interest that rearrangement or modifications in the number of chromosome 8 have been associated with certain forms of cancer. (45 refs)

- 78-7078 Hemodynamic Characteristics in Microcirculatory Blood Channels During Early Tumor Growth.** (Eng) Endrich, B. (Dept. AMES-Bioengineering, M-005, Univ. California at San Diego, La Jolla, CA, 92093); Intaglietta, M.; Reinhold, H. S.; Gross, J. F. *Cancer Res* 39(1): 17-23; 1979.

RBC velocity was determined in the microcirculation of the BA 1112 rat sarcoma and in regions located outside the tumor (repairing tissue), which were used as a control and were assumed to be representative of the inflammatory-reparative reaction of a healing wound. Studies were carried out in female Wistar rats of an inbred strain fitted with modified Algire window chambers utilizing television techniques in situ. Observations and measurements were started 14 days after implantation of tumor or adjacent tissue in the chamber and continued for 18 days. RBC velocity in repairing tissue was of the order of 0.80 ± 0.15 (S.E.) mm/sec in terminal arterioles, 0.2 ± 0.09 mm/sec in postcapillary venules, and 0.2 ± 0.1 mm/sec in collecting venules and remained constant during the observation period. By comparison, RBC velocity in terminal arterioles of the tumor tissue increased from 0.72 ± 0.08 to 0.95 ± 0.09 mm/sec and decreased slightly in postcapillary and collecting venules. Vasomotion in terminal arterioles of the malignant tissue could be demonstrated only until day 22 after implantation. Since vasomo-

tion was observed only during the initial phase of tumor growth and the arteriolar RBC velocity increased during the same period, the findings indicate that blood is supplied to mesenchymal tumors by means of dilated arterioles. In adjacent repairing tissue (control), the RBC velocity of postcapillary and collecting venular remained constant. By comparison, RBC velocity decreased in venular outflow vessels of the tumor, suggesting that venous drainage is facilitated during progressive spread of the BA 1112 sarcoma. (22 refs)

- 78-7079 Cell Movement and the Mechanism of Invasiveness: A Survey of the Behaviour of Some Normal and Malignant Cells Implanted into the Developing Chick Wing Bud.** (Eng) Tickle, C. (Dept. Biology as Applied to Medicine, Middlesex Hosp. Medical Sch., Cleveland St. London W1P 6DB, England); Crawley, A.; Goodman, M. *J Cell Sci* 31: 293-322; 1978.

Normal and malignant cells were implanted into the loose mesenchyme of developing chick wing buds to investigate the mechanisms of tumor invasiveness. Two to 3 days later, the distribution of the grafted cells was examined histologically. Cells from mouse lung tumor, rat bladder tumor, and human breast tumor did not invade the mesenchyme, but trophoblastic, sarcoma 180, and neuroblastoma cells and cultured hamster fibroblasts (BHK, PyBHK, Nil 8, HSV Nil 8) did. Cells from embryonic pigmented retina and heart ventricle were invasive. Thus, cell movement may not be a common feature of all invasive tumors. Lack of contact inhibition of movement, although probably involved in the invasiveness of sarcoma 180 cells, did not appear to be necessary for invasion. Cells that exhibited contact inhibition of movement (BHK and PyBHK) were also invasive. Both normal and transformed cells (BHK and PyBHK; Nil 8 and HSV Nil 8) moved into the mesenchyme. Other invading cells, such as trophoblastic, neuroblastoma, and (to a small extent) HSV Nil 8 cells, destroyed the adjacent host tissue, and this may be important in their invasiveness. Trophoblastic cells and the fibroblasts were often elongated along the basement membrane at the ectoderm/mesenchyme border and also closely apposed to the endothelial linings of the blood vessels. Sarcoma 180 and neuroblastoma cells clustered around nerves. The embryonic tissues and neuroblastoma cells were often associated with blood vessels. A striking finding was that carcinoma cells were frequently found positioned within the wing ectoderm on the basement membrane. This affinity of carcinoma cells for epithelium rather than the mesenchyme suggests that the invasiveness of carcinomas involves the same mechanisms as the epithelial invasion of the mesenchyme that occurs during gland development. (77 refs.)

- 78-7080 Banding Studies on Chromosomes in Diffuse "Histiocytic" Lymphomas: Correlation of 14q+ Marker Chromosome with Cytology.** (Eng) Fukuhara, S. (Dept. Medicine, Univ. Chicago Hosp., 950 E. 59th St.

ago, IL, 60637); Rowley, J. D.; Variakojis, D.; Sweet, J. *Blood* 52(5): 989-1002; 1978.

Chromosomes were studied in involved lymph nodes of nine patients with diffuse histiocytic lymphoma. Two of the patients had stage II disease; their tumors were fibrotic and had mitotic cells. One patient was in stage III, and the remaining patients had stage IV disease. The modal chromosome number of abnormal cells from these last seven patients was diploid in two, hyperdiploid in four, and near-triploid in one. Complete banding studies, including peripheral blood and bone marrow, of six cases, and partial analysis of the ninth indicate that (1) every patient had a distinct cell line with common markers, with a few cells showing minor variations; (2) although certain chromosomes (Nos. 1, 2, 3, 9, 12, 14) were structurally affected more often than others, no markers with the same banding pattern were noted among patients; and (3) the cytologic type of lymphoma could be correlated with the karyotype in all seven patients. When the Rues and Collins classification was used, three patients whose tumors were composed predominantly of large noncleaved cells showed a 14q translocation leading to the formation of a 14q+ marker chromosome. This marker was not observed in four patients whose tumors had a majority of cleaved cells. These preliminary results suggest that there are consistent chromosome changes associated with specific subtypes of lymphoproliferative disorders analogous to the Philadelphia chromosome in chronic myelogenous leukemia. (32 refs)

78-7081 Effect of Microtubule Inhibitors on Malignant Invasion In Vitro. (Eng) Mareel, M. M. (Dept. Experimental Cancerology, Clinic Radiotherapy and Nuclear Medicine, Univ. Hosp., De Pintelaan 135, B-9000 Ghent, Belgium); De Brabander, M. J. *J Natl Cancer Inst* 70: 787-792; 1978.

Malignant C3H/3T3 mouse cells MO₁ invaded embryonic heart fragments in an organotypic coculture system in semisolid medium. Malignant invasion was obvious from an irregular boundary between the MO₁ nodule and the heart tissue, the penetration of strands and solitary MO₁ cells into the heart, and destruction of the heart. In this system, at a concentration of 1 µg/ml, the microtubule inhibitors colchicine, demecolcine, vincristine sulfate, vinblastine sulfate, or mefloquine [5-(2-thienylcarbonyl)-1H-benzimidazol-2-yl]carbamate (Nocodazole) totally inhibited malignant invasion. At this concentration the drugs were also mitostatic, which was apparent from the absence of mitoses and from the absence of postmetaphase figures. At a mitostatic dose of 10 µg/ml, 5-fluorouracil (5-FU), cytosine arabinoside, or bleomycin did not interfere with malignant invasion. Combined treatment of the cocultures with 5-FU (10 µg/ml) plus Nocodazole (1 µg/ml) completely inhibited invasion. These cocultures also showed the effective inhibition of mitosis by 5-FU, because Nocodazole-induced chromosome breaks were absent. The reversibility of the anti-invasive effect of a 4-day treatment with Nocodazole

(1 µg/ml) was demonstrated in shaker cocultures with the use of fluid medium. These in vitro experiments indicate that cytoplasmic microtubules are involved in malignant invasion and that cell division and invasion constitute separate characteristics of malignant cells. (32 refs)

78-7082 Evidence that the Malignant Lymphoma of Sjogren's Syndrome is a Monoclonal B-Cell Neoplasm. (Eng) Zulman, J. (Immunology and Arthritis Section, Veterans Admin. Medical Center, 4150 Clement St., San Francisco, CA, 94121); Jaffe, R.; Talal, N. *N Engl J Med* 299(22): 1215-1220; 1978.

Nine patients who developed malignant lymphoma (ML) following Sjogren's syndrome (SS) and antecedent benign salivary gland lesions were studied histopathologically and immunohistologically. Paraffin-embedded tissue blocks of malignant lymphoma were examined in all cases, and SS salivary gland lymphoid lesions in four. The unlabeled antibody-enzyme method (peroxidase antihorseradish peroxidase) was used for immunohistologic analyses. There were six female patients, the average age was 48 yr and the average duration from onset of SS to diagnosis of ML was 10 yr. Two patients received no irradiation or immunosuppressive therapy before ML development. Histologically, the tumors varied within themselves in cellular composition, content of epithelioid histiocytes, presence of plasmacytoid features, and numbers of plasma cells, but 6/9 patients had cytoplasmic immunoglobulin within the ML cells, confirming their B-cell nature. All six of these patients had IgMκ chains exclusively, but only one had rheumatoid arthritis, whereas all three patients who did not have IgMκ chains had rheumatoid arthritis. No tumor cells contained lysozyme, and none gave evidence of phagocytosis. These lymphomas could have arisen coincidentally with Sjogren's syndrome or could represent a stage of B-cell dedifferentiation in which the concentration of intracytoplasmic immunoglobulin falls below a detectable threshold. (23 refs)

78-7083 Familial Hodgkin's Disease. (Eng) Hull, P. J. (Royal Infirmary, Manchester M13 9WL, England); Delamore, I. W. *Postgrad Med J* 54(636): 676-677; 1978.

The occurrence of Hodgkin's disease in a 23-yr-old man, his 22-yr-old sister, and their 13-yr-old cousin is reported. All had mediastinal involvement, and, in each case, the histology was that of nodular sclerosing Hodgkin's disease. (8 refs)

78-7084 Membranous Proliferative Glomerulonephritis and Castleman's Lymphoma. (Fre) Genin, R. (Clinique Medicale, Hopital du Bocage, F. 21034 Dijon)

Cedex, France); Justrabo, E.; Devillebichot, C.; Michiels, R.; Riflé, G. *J Urol Nephrol (Paris)* 84(9): 667-672; 1978.

Castleman's pseudotumor, localized retroperitoneally and associated with membranoproliferative glomerulonephritis, was diagnosed in a 56-yr-old man. This association has never been reported before. A polyclonal plasmacyte population with IgA, IgG, and IgM was found in the tissue surrounding the lymph nodes involved. Disappearance of proteinuria and regression of the nephrotic syndrome were seen after the pseudotumor was removed. Castleman's pseudotumor is very rare; only 200 cases have been reported hitherto. It affects men and women alike, mostly during the second and third decades of life. While a fortuitous association of the two diseases cannot be ruled out, the glomerular affection may be due to the presence of angiofollicular hyperplasia, as are the frequent immunological and hematological manifestations. Thus, the nephropathy may be due to the production of circulating immunocomplexes. According to another hypothesis, the two diseases may have a common etiological factor. Castleman's pseudotumor was found in some cases following vaccination or the resorption of suture material. Bacterial or viral etiology has not been proved. The patient presented here had a facial wound about 2 mo before the onset of glomerulonephritis, which may have been the port of entry of an antigen. (9 refs)

78-7085 Development of the Granulomatous Response in Murine Toxocariasis. (Eng) Kayes, S. G. (Veterans Admin. Hosp., 1310 24th Ave., South, Nashville, TN, 37203); Oaks, J. A. *Am J Pathol* 93(2): 277-294; 1978.

To characterize the cellular evolution of muscle-associated granuloma in murine toxocariasis, *Toxocara canis* eggs were collected from the stools of naturally infected puppies, cultured, checked for the presence of stage 2 (L2) infective larvae, and administered to male HM/ICR mice (1,000 eggs/mouse) by stomach intubation. Necropsies were performed between 11-14, 28-35, and 70-77 days postinfection, and the carcass musculature was examined by light and electron microscopy following appropriate histologic preparation. By day 11, the initial granulomas consisted primarily of eosinophils. By day 14, many of the eosinophils appeared to be dying and undergoing disintegration; the resulting cellular debris was sometimes seen inside macrophages. Two distinct lesions were observed during the 28-35 day interval: the first, representing a continuation of the evolving mononuclear progression seen at day 14 and containing predominantly macrophages, was outlined by fibroblasts and enveloped in a distinct collagenous capsule; the second lesion day contained foci of polymorphs and mononuclear cells similar to those seen at 11-14 days. By 77 days, the granulomas had developed into fibrotically encapsulated epithelioid lesions surrounding the larva. The results indicate that infection by this parasitic larvae nematode progresses from an inflammatory reaction to a loose eosinophil-containing exudate, and, finally, to a

compacted, encapsulated granuloma, a situation likened to hepatic granuloma and schistosome egg granuloma. (36 refs)

78-7086 Studies on a Japanese Burkitt Lymphoma. Part 2: Serial Transplantation of a Japanese Burkitt Lymphoma Cell Line in Hamsters. (Jpn) Kishimoto, I. (Dept. Internal Medicine, Okayama Univ. Medical School, Okayama, Japan). *Jpn J Clin Hematol* 19(9): 1186-1191; 1978.

An Epstein-Barr virus genome-free Japanese Burkitt lymphoma line (JBL) was transplanted ip for 10 serial passages in newborn Syrian hamsters treated with antilymphocyte serum. All the hamsters developed invasive tumors and massive ascites. The hamster-passaged line maintained a characteristic starry-sky pattern closely resembling that of human biopsy material. Some of the animals developed not only bone marrow invasion but also a leukemic blood picture. Since the JBL line was converted to an ascitic form, its serial passage was greatly facilitated. (13 refs)

78-7087 Possible Role of Connective Tissue in Epidermal Neoplasia. (Eng) Mitrani, E. (Dept. Medicine, Welsh Natl. Sch. Medicine, Heath Park, Cardiff, Wales). *Br J Dermatol* 99(3): 233-244; 1978.

Interactions between epidermal cells were defined within a proposed mathematical model of mammalian skin. Examination was made of how epidermis retains its mass both normally and after various disturbances and how this mechanism may fail, as in neoplasia. The model assumes that the lower the epidermal layer (ie, nearer the dermis), the higher the probability of a cell in that layer to enter mitosis. Computer tests of the model suggested that in highly proliferating epidermis, competition for cell space in the basal layer may be sufficient to generate considerable forces in the papillary dermis. According to data on human and pig epidermal hyperplasia, basal cells are submitted to considerable lateral forces and these and not dermal hyperplasia or dermal edema are the forces responsible for the increasingly folded dermoepidermal junction. In the condition of a persistently high mitotic rate, the model remains stable only if new connective tissue (collagen) synthesis is not induced by the developing papillary tension. New collagen tissue formation relieves the stress only temporarily; then the epidermis is in a situation similar to the one prevailing before synthesis began, except that it has induced a given amount of dermal growth. Thus, a self-expanding system emerges. Clearly, collagen synthesis plays an important role in epithelial neoplasia. (38 refs)

78-7088 A Causal Relation Between Bowen's Disease and a Lewisite Scar. (Ger) Krause, H. (Abteilung Dermatologie der Medizinischen Fakultät, Technische Universität München, Germany).

Hochschule, Goethestrasse 27-29, D-5100 Aachen, W. Germany); Grussendorf, E. I. *Hautarzt* 29(9): 490-493; 1978.

A 22-yr-old woman developed Bowen's disease of the right lower leg 8 yr after accidental contact of that area with Lewisite (mustard gas). The tumor excision was followed by a local recurrence 2 yr later. (25 refs)

78-7089 Basal Cell Epithelioma in a Vaccination Scar. (Eng) Hazelrigg, D. E. (Section Dermatology, Veterans Affairs Medical Clinic, 421 Chestnut St., Evansville, IN, 47713). *J Dermatol* 17(9): 723-725; 1978.

A 60-yr-old woman developed a superficial basal cell epithelioma (BCE) in a smallpox vaccination scar. She had been vaccinated at 12 yr of age. Two BCE's had been removed from the posterior neck and forehead 2 and 1 yr before the present presented with the lesion described. The latter had regressed gradually over 6 yr. (12 refs)

78-7090 DNA Repair Deficiency in Lymphocytes from Patients with Actinic Keratosis. (Eng) Darub, J. M. (Dept. Biochemistry and Dermatology, University of Glasgow, Glasgow, G12 8QQ, Scotland); Mackie, R.; and J. D. *Bull Cancer (Paris)* 65(3): 357-362; 1978.

DNA repair activity in peripheral blood lymphocytes was measured in 18 patients with actinic keratosis (AK) and 18 matched controls by comparing the incorporation of ³H-thymidine into UV-irradiated cells with that into nonirradiated cells. The incorporation was followed by autoradiography or scintillation spectrometry. The rate of incorporation of ³H-thymidine (5.0 µCi/ml, 18.5 Ci/mmole) into hydroxyurea-treated (1.5 x 10⁻³M) AK lymphocytes measured at different times after irradiation of 20 Joules (J)/meter²(m²) was constantly lower than that into normal cells at all times examined. When UV dose response was studied by measuring total incorporation of ³H-thymidine in lymphocytes at different UV doses 4 hr after irradiation, the repair activity of AK cells was approx 50% that of controls at all UV doses examined. For both cell populations, the repair curve plateaued between 10 and 20 J/m². Autoradiographic analysis of AK lymphocytes after irradiation at 5.0 and 20 J/m² determined the av UV-induced DNA repair synthesis to be 47.7% and 48.9% of normal lymphocytes, respectively. Measurement of the differential of total incorporation of ³H-thymidine into hydroxyurea-treated and untreated patient and control lymphocytes after 20 J/m² irradiation revealed a DNA repair rate of 41%-56% that of normal controls. The results of using two different analytical methods to examine DNA repair are quite similar and show that the rates of repair in AK cells are all in the range 38-63% that of normal cells. (12 refs)

78-7091 DNA Repair in Conditions Associated with Malignancy: Aging and Actinic Keratosis. (Eng) Ringborg, U. (Radiumhemmet, Karolinska Hosp., S-104 01 Stockholm 60, Sweden); Lambert, B.; Swanbeck, G. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): pp. 1193; 39-52; 1977.

UV light-induced DNA excision repair was studied in two conditions associated with malignancy, aging and actinic keratosis, in an attempt to clarify the relationship between a decreased capacity for DNA repair and malignant disease. WBC from 48 healthy subjects aged 13-94 yr were irradiated with 3.2, 6.4, 9.6, or 19.2 joules/m² UV light and incubated with ³H-thymidine in the presence of hydroxyurea. There were great variations in max incorporation of ³H-thymidine among individuals, and they were considered biological in origin. For the three higher UV doses, DNA repair synthesis was significantly lower in subjects 60-94 yr old than in subjects 13-59 yr old. The av DNA repair capacity decreased by about 30% between 20 and 90 yr of age. WBC from 10 patients with actinic keratoses (5 men, 5 women aged 53-81) and from 10 healthy controls of the same age range were subjected to the same treatment as above. The capacity for DNA repair synthesis was 30% lower in the actinic keratosis patients than in the controls. The possibility that the reduced capacity for DNA repair in aging and actinic keratosis predisposes an individual to malignant disease is discussed. (14 refs)

78-7092 Posttraumatic Epithelioma. (Ita) Fallica, L. (Clinica Dermosifilopatica, Università di Catania, Catania, Italy); Guarneri, B. *G Ital Dermatol* 113(7/8): 477-479; 1978.

Three patients (two men, one woman, aged 70, 62, and 56 yr, respectively) developed basal-cell carcinoma of the skin after mechanical trauma of the same site. The latency from the trauma to the manifestation of the carcinoma was 7-15 mo. Literature data, as well as the rarity of posttraumatic carcinomas, suggest that the trauma only accelerates the development of the neoplasia as a function of a quantitative modification of the cells that are qualitatively already anaplastic, ie, that trauma acts only as a promoting factor. (13 refs)

78-7093 Hydatidiform Mole as a Complication of Artificial Insemination. (Eng) Stone, S. C. (Dept. Obstetrics and Gynecology, Univ. California at Irvine, 101 City Drive S., Orange, CA, 92668). *Am J Obstet Gynecol* 132(6): 696-697; 1978.

A hydatidiform mole developed in a 22-yr-old woman who

had undergone artificial insemination with fresh donor sperm. A spontaneous abortion occurred at 8 wk gestation, and scanty tissue of molar appearance was passed. The pathology report confirmed the diagnosis of molar pregnancy. (2 refs)

78-7094 Naevoid Basal Cell Carcinoma Syndrome and Charcot-Marie-Tooth Disease: Two Autosomal Dominant Disorders Segregating in a Family. (Eng) Heimler, A. (Div. Human Genetics, Dept. Pediatrics, Long Island Jewish-Hillside Medical Center, New Hyde Park, NY, 11040); Friedman, E.; Rosenthal, A. D. *J Med Genet* 15(4): 288-291; 1978.

A family is described in which 16 members (three generations) have Charcot-Marie-Tooth disease; at least six members also have nevoid basal cell carcinoma syndrome (NBCS), and one member with both disorders has two daughters with NBCS. These two disorders have not been reported previously in the same patient or within one family. A diagnosis of NBCS should be considered in any patient with recurrent dentigerous cysts at an early age. (8 refs)

78-7095 Pathogenesis of Prolactin-secreting Pituitary Adenomas. (Eng) Sherman, B. M. (Clinical Res. Center, Univ. Hosps., Iowa City, IA, 52242); Schlechte, J.; Halmi, N. S.; Chapler, F. K.; Harris, C. E.; Duello, T. M.; VanGilder, J.; Granner, D. K. *Lancet* 2(8098): 1019-1021; 1978.

In a comprehensive multidisciplinary study of pituitary tumors, 42 women (aged 18-54) with amenorrhea and hyperprolactinemia [25-833 nanograms (ng)/ml] related to a pituitary adenoma were studied. Seventy-four percent of these patients developed amenorrhea and/or galactorrhea in immediate association with the use or discontinuation of oral contraceptives (24 women) or postpartum (7 women). All underwent transsphenoidal surgery for resection of the tumor, and there was enough adenomatous tissue for immunocytochemical studies in 35 specimens. Prolactin was localized unequivocally in 31 specimens by immunoperoxidase staining with anti-rat prolactin serum. The mean serum prolactin concentration did not differ in the groups with and without galactorrhea, but it was significantly greater in the 15 patients with adenomas 10 mm or more in diameter (226 ng/ml) than in those with lesions < 10 mm (94 ng/ml). The authors suggest that estrogens, which modulate prolactin secretion in normal humans and animals, can induce the growth and expression of otherwise silent pituitary lesions and that this should be considered a risk of oral contraceptive use. (17 refs)

78-7096 Meningiomas in Five Members of a Family over Two Generations, in One Member Simultane-

ously with Acoustic Neurinomas. (Eng) Delleman, J. V. (Ophthalmogenetic Dept., Netherlands Ophthalmic Research Inst., Wilhelmina Gasthuis, Amsterdam, Netherlands); IJong, J. G.; Bleeker, G. M. *Neurology (Minneapolis)* 28(6): 567-570; 1978.

The familial occurrence of meningioma in five members of a family over a two-generation span is reported. The proband was a 33-yr-old man who had undergone enucleation of the left eye at age 22 yr; a meningioma psammomatous of the optic nerve sheath, with infiltration of the surrounding tissue was found. Two recurrences were noted over the next 5 yr. A sister had a spinal meningioma removed at age 25 and a hyperostotic meningioma removed at age 31. The proband's father and uncle also had meningiomas. None of these family members had evidence of neurofibromatosis. The autopsy report of the paternal aunt who had died at the age of 43 indicated that she had bilateral acoustic neurinomas and multiple meningiomas. A sister of the proband has multiple café-au-lait spots. It is suggested that all these cases are manifestations of neurofibromatosis. (55 refs)

78-7097 Possible Maternal Effect on Severity of Neurofibromatosis. (Eng) Miller, M. (Dept. Medicine and Pediatrics, Univ. Washington Sch. Medicine, Seattle, WA, 98105); Hall, J. G. *Lancet* 2(8099): 1071-1072; 1978.

A review of medical records of patients under 18 yr old seen for signs or symptoms suggestive of neurofibromatosis revealed 62 patients (from 54 families) with the disease who were then assessed for severity of disease and whether the individual was the result of a new mutation or born to an affected father or mother. Patients were considered to have this autosomal dominant disease if they demonstrated café-au-lait spots > 1.5 cm in diameter. Severity of disease was graded on a 0-3 scale based on physical and mental parameters. The expected numbers of new mutations and patients born to affected mothers or fathers (based on gene probabilities and relative fertilities) was compared with the observed cases; 16 patients had new mutations, compared with an expected number of 31, 15 were born to an affected father compared with an expected 11, and 31 were born to an affected mother, compared with an expected 20. The mean severity of disease was highest in those born to affected mothers (2.4), least in those born to affected fathers (0.8), and intermediate in new mutations (1.6). Cases due to new mutations or born to affected fathers were relatively evenly distributed among the four grades of severity, but significantly more of those born to affected mothers had severe complications. The 21 cases with severe complications born of affected mothers tended to present the disease early in childhood. Furthermore, in the six instances in which two affected sibs were born to an affected mother, all suffered the most severe form of complications. It is suggested that the severity observed in cases born to affected mothers is due to some factor in the intrauterine environment produced by the affected mother.

interacts only with fetuses that carry the neurofibrinogen gene. This interaction may be humorally mediated (10 refs)

7098 Ulcer of the Tongue after CCNU Therapy for a Spinal-Cord Tumor. (Eng) Walker, M. D. Brain Tumor Study Group, NCI, Bethesda, MD, 20014; Hardison, E. P.; Scully, R. E. *N Engl J Med* 299(19): 1010-1067; 1978.

A 7-yr-old woman with a history of hepatitis and pelvic tumor 1 yr before admission presented with pain in the arm. She was diagnosed with an intradural-extramedullary, mixed astrocytoma/ependymoma-type glioma of the spinal cord. Over the next 3 yr, she received 12 doses of 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea because of the growth of malignant glial cells in cerebrospinal fluid cultures. Upon second admission, she developed general debility secondary to anaplastic cell carcinoma, with metastases to cervical lymph nodes, lungs, and pleuras. Anatomical diagnosis also revealed myelinolysis, quiescent myelopathy and cavernous hemangiomas of the liver. (5 refs)

7099 Enzymes of Neutrophils from Patients with Precancerous States of the Larynx. (Eng) Lisiewicz, J. (Dept. Hematology, Inst. Internal Medicine, Cracow Medical Academy, Cracow, Poland); Gierlek, T.; Pilch, J.; Piasa, B.; Namyslowski, G. *Rev Esp Oncol* 25(1): 7-11; 1978.

Levels of β -glucuronidase, alkaline and acid phosphatases, N-acetyl- β -glucosaminidase (NAGA), myeloperoxidase, glycogen, and lipids in the peripheral blood neutrophils of 20 healthy men and 24 men with precancerous states of the larynx (leukoplakia, papillomas, or pachydermia) were compared. The patients and healthy controls did not differ in terms of absolute total neutrophil counts or absolute counts of neutrophils staining for β -glucuronidase, acid phosphatase, NAGA, myeloperoxidase, PAS, or lipids. However, mean values of neutrophilic leukocytosis, the absolute count of alkaline phosphatase-positive neutrophils, and alkaline phosphatase activity were significantly higher in the patients than in the controls. In contrast, intracellular β -glucuronidase activity and lipid content were significantly lower in the patients. (11 refs)

7100 Mucoepidermoid Tumor of the Epipharynx. (Ger) Nowak, R. (Klinik und Poliklinik für Hals-, Nasen- und Ohrenkrankheiten, Wilhelm-Pieck-Universität, Doberanerstrasse 139, DDR-25 Rostock 1, E. Germany); Putzke, H. P. *Arch Otorhinolaryngol (NY)* 221(3): 223-233; 1978.

A mucoepidermoid tumor of the epipharynx was diagnosed by biopsy in a 39-yr-old woman. She developed a local recur-

rence about 1 yr after tumor resection. Histopathological investigation revealed that the primary tumor consisted mainly of undifferentiated cells, plus glandular duct and acinar epithelial cells and epidermoid cells at various stages of maturity. The findings suggest that these tumors originate from pluripotent cells of the transition between the acinus and duct, rather than from major glandular and salivary ducts. They are classified as being of low-grade malignancy. (12 refs)

78-7101 Intracisternal Tubular Inclusions in Nasopharyngeal Carcinoma. (Eng) Prasad, U. (E.N.T. Dept., Faculty Medicine, Univ. Malaya, Kuala Lumpur, Malaysia); Gogusev, J. *J Laryngol Otol* 92(11): 979-989; 1978.

Twenty biopsy specimens from patients with nasopharyngeal carcinoma were examined ultrastructurally for the presence of intracisternal tubular inclusions (ITI). These inclusions were observed in the cytoplasm in eight cases. The tubules were arranged in reticular arrays. Dense particles with an outer diameter of 200 Å were interconnected by light bands, and these aggregates were bound by a membrane that was continuous with the granular endoplasmic reticulum (GER). In five other cases, a slightly different type of cytoplasmic inclusion was found. The inclusion was large and almost spherical, with filamentous aggregates criss-crossing in a haphazard way. The aggregates seemed to be connected to the GER. ITI may be the result of an immunological reaction initiated by injured cells infected with Epstein-Barr virus, which acts as an antigen. (37 refs)

78-7102 Radiation-induced Carcinoma of the Thyroid with Severe Thyrotoxicosis. (Eng) Citrin, D. L. (Dept. Human Oncology, 701C Univ. Hosps., 1300 Univ. Ave., Madison, WI, 53706); Pederson, J.; Rose, D. P.; Carbone, P. P. *Wisconsin Medical J* 77(10): 597-599; 1978.

The case of a 65-yr-old woman with radiation-induced metastatic thyroid cancer and severe thyrotoxicosis is presented. The patient had received radiation therapy to the neck and upper chest at age 8 yr. During the year prior to presentation, there was a rapid increase in the size of a neck mass that had been present for many years. The patient had taken thyroxine intermittently during these years but had taken no thyroid medication for at least 3 yr prior to presentation. The patient presented with complete dysphagia. Hyperkinesia, tremor, and atrial fibrillation suggested hyperthyroidism; this was supported by thyroid function tests, but the patient was unable to swallow the ^{131}I for a radioiodine uptake test. Biopsy of the neck mass revealed a mixed follicular and papillary thyroid carcinoma. The firm, nontender mass measured at least 10 x 10 cm and formed a firm collar around the trachea, which could not be clearly identified. Postmortem examination showed that the left lobe of the thyroid gland measured

10 x 5 cm and contained necrotic areas. The right lobe was also involved, with invasion of the capsule and neck muscles. Histologic examination revealed a well-differentiated follicular and papillary carcinoma with colloid present in the follicles. There was no evidence of hyperplastic normal thyroid tissue. The lungs were filled with metastatic thyroid nodules up to 2 cm in diameter. In previously irradiated patients, hyperthyroidism does not exclude the possibility of thyroid cancer. (6 refs)

78-7103 Morphological Aspects of Parathyroid Transplantation. Clinical Relevancy of Induced Invasive Tissue Growth. (Ger) Klempa, I. (Abteilung für Allgemein- und Abdominalchirurgie, Zentrum der Chirurgie der Universität, Theodor-Stern-Kai 7, D-6000 Frankfurt/Main, W. Germany); Steinau, U.; Frei, U.; Koch, K. M.; Usadel, K. H.; Rottger, P. *Chirurg* 49(11): 704-710; 1978.

A 40-yr-old man with hypercalcemia and increased parathyroid hormone levels due to terminal renal insufficiency underwent total parathyroidectomy following which one of the parathyroid glands (showing typical adenomatous hyperplasia without signs of potential or manifest malignancy) was grafted in the forearm. The autograft started growing rapidly 5 mo after grafting, and the serum calcium and parathyroid hormone levels increased. An autonomous tumor-like adenomatous formation measuring 2.5-3 cm in diameter and weighing 3.1 g was excised 9 mo after the implantation. The parathyroid tissue excised showed solid, relatively cell-rich growth interspersed with giant nuclei. The tumor invaded the vessels of the normal muscular tissue. The findings suggest that induced invasive tissue growth is possible following parathyroid autografting and that only parathyroid tissue without adenomatous hyperplasia should be used for this purpose. (16 refs)

78-7104 Growth of Murine Mammary Tumor Cells in an Artificial Capillary System: Ultrastructural Study. (Fre) Mouriquand, J. (Laboratoire d'Histologie, Cytologie et Cyto-genetique, Faculte de Medecine de Grenoble, 38700 La Tronche, France); Mouriquand, C.; Mermet, M. A.; Louis, J.; Petitpas, E. *Biol Cell (Paris)* 31(1): 27-30; 1978.

A method is described for growing murine mammary tumor cells on an artificial capillary system. The cells were from spontaneous mammary tumors in the PS strain of mice. Actually, the cells had reverted to epithelial morphology since they were in the 162nd passage. The artificial capillary system used copolymers of acrylic or polysulfone permeable to substances of mol wt up to 50,000 or 100,000, respectively. The cells were grown in a medium containing 10% decomplexed calf serum, penicillin and streptomycin. Light and electron microscopy was used to monitor growth of the cells in the wk 3-5. Epithelial clumps could be seen between the

capillaries or clustering against them. Murine mammary tumor virus (MTV) particles were numerous, with A-type particles in the cytoplasm and B-type particle in the vacuoles in the intercellular spaces and free in the culture medium. Intracytoplasmic ductlike vacuoles similar to those described in human breast cancer cells were observed, and it is believed that these have not previously been reported in murine mammary tumor cells. (43 refs)

78-7105 Cytology of Secretions and Aspiration Biopsy. Diagnostic Applications in Mammary Diseases. (Ita) Magni, E. (Servizio di anatomia e istologia patologica, Ospedale Civile S. Maria delle Croci, Ravenna, Italy); Montagna, G.; Tabarri, B. *Pathologica* 70(1007/1008): 495-502; 1978.

Secretion cytology was performed in 1,780 women with mammary secretion, and needle biopsy was performed in 73 women with palpable breast nodes. The secretion cytology was indicative of pathological changes: the cytological changes were rated as grade III in 24 cases; papilloma was diagnosed histologically in 19/24 cases, papillomatosis in 5 and intraductal carcinoma in 1. The cytological findings were rated as grade IV-V in 15 cases: intraductal carcinoma was found histologically in 6 cases and invasive carcinoma in 9. The needle biopsy, performed in 395 women with solid nodes and in 357 patients with cystic nodes, failed to reveal tumor cells in any of the cysts. The reliability of needle biopsy was studied in 184 histologically verified cases: the histology revealed malignant tumors in 75 cases; the cytological findings were concordant with the histology in 62 cases, false negative in 5, and 8 biopsies were unevaluable. In 109 histologically negative cases, the biopsy was correctly negative in 85 cases, false positive in 4, and 20 biopsy specimens were unevaluable. The findings indicate the great diagnostic value of secretion and biopsy cytology in the diagnosis of breast diseases. (10 refs)

78-7106 Prevalence of Hypertension and Diabetes in Breast Cancer: A Case-Control Study in 179 Patients and Age-matched, Non-hospitalized Controls. (Eng) Adami, H. O. (Dept. Surgery, Univ. Hosp., S-750 14 Uppsala, Sweden); Rimsten, A. *Clin Oncol* 4(3): 243-249; 1978.

The prevalence of hypertension and diabetes was investigated in 179 consecutive breast cancer patients (mean age, 63 yr) seen in four Swedish counties during a 5-mo period and 17 age-matched controls. Pertinent information was gathered by questionnaire, hospital records, and by interview. The blood pressure of all subjects was measured, and blood samples were taken for hormone analyses. Thirty-three patients and 28 controls were premenopausal. Hypertension was found in 33.5% of the patients and 32.4% of the controls. The distribution of systolic and diastolic blood pressure in the patients was in close accordance with that in the controls, even when the women were subdivided into pre- and postmenopausal

categories. Age was significantly correlated with diastolic and systolic blood pressure. The use of antihypertensive drugs was similar in the two groups. The prevalence of diabetes was also similar: 10 patients (9 under treatment) and 8 controls (all under treatment). Apparently, there is no association between breast cancer and hypertension or diabetes. (27 refs)

78-7107 Fibrocystic Breast Disease in Oral-Contraceptive Users. A Histopathological Evaluation of Epithelial Atypia. (Eng) LiVolsi, V. A. (Dept. Pathology, Yale Univ. Sch. Medicine, New Haven, CT, 06510); Stadel, V.; Kelsey, J. L.; Holford, T. R.; White, C. *N Engl J Med* 9(8): 381-385; 1978.

A case-control study was performed to determine whether the association between oral contraceptive (OC) use and fibrocystic disease (FD) varies according to the extent of epithelial atypia associated with the latter. The patients, 205 premenopausal women aged 20-44 yr, and individually matched controls had been admitted as in- or outpatients to two Connecticut hospitals during 1971-1973. The frequency of FD was not differentially associated with prior OC use when cases were subdivided according to histologic patterns. However, there were variations in the frequency of FD among long-term pill users compared with nonusers according to the degree of epithelial cytologic atypia being considered. The results support the hypothesis that the use of OC for 2-4 yr or more is associated with a decreased frequency of FD with only minimal epithelial atypia, and not with the more florid forms of this condition. These findings suggest that a spectrum of cystic disease exists and that the long-term use of OC protects against the forms of FD that are not firmly associated with an increased risk of breast cancer, but not against the premalignant forms. (40 refs.)

78-7108 Volume and Hyperplastic Lesions of Breast of Japanese Women in Hawaii and Japan. (Eng) Sasano, N. (Dept. Pathology, Tohoku Univ. Sch. Medicine, Sendai, Japan); Tateno, H.; Stemmermann, G. N. *Prev Med* 2(2): 196-204; 1978.

Breast volume and the frequency of ductal hyperplasia were determined in autopsy subjects in Japan and Hawaii. The volume of the total breast (TBV) and of the mammary gland was calculated from specimen radiographs of serially sliced breasts enucleated at necropsy. TBV had a relatively good correlation with the Quetelet index (body wt/height squared) except among Japanese women <45 yr of age. Glandular volume had no significant correlation with the Quetelet index except in Japanese women ≥45 yr. In Hawaiian women, the TBV decreased until 39 yr of age and thereafter showed little change. In Japan, the TBV showed a conspicuous decrease with old age, possibly due to a loss of adipose tissue. In premenopausal women, mean values of glandular breast

volume decreased up to 49 yr of age in parallel with the TBV. The volume density increased slightly from puberty to 39 yr; it then decreased and stabilized at a lower level after 40. Women with premenopausal oophorectomy had volume density values within the standard deviation, but they had a smaller mean value. Women with postmenopausal oophorectomy had generally low values. Epithelial hyperplasia was much more common in Hawaiian than Japanese women after age 30. The Hawaiian women were divided into native-born issei and Hawaiian-born nisei. A total of 19/37 nisei showed epithelial hyperplasia; this incidence (51.4%) was far higher than that in the issei (14.5%) and in Japanese women (18.7%). In Hawaiian women aged ≥45 yr, the volume of the mammary gland, corrected by the Quetelet index, was larger when epithelial hyperplasia was present than when it was absent. Women with premenopausal oophorectomy had no epithelial hyperplasia. (4 refs)

78-7109 Metastases In Vitro: Assays for Cell-release and Invasion (Meeting Abstract). (Eng) Li, A. P. (Cancer Res. and Treatment Center, Univ. New Mexico, Albuquerque, NM, 87131). *Proc Am Assoc Cancer Res* 19: 55; 1978. (no refs)

78-7110 Transformed Cell Lines Susceptible or Resistant to In Vivo Surveillance Against Tumorigenesis. (Eng) Patek, P. Q. (Developmental Biology Lab., Salk Inst., P.O. Box 1809, San Diego, CA, 92112); Collins, J. L.; Cohn, M. *Nature* 276(5687): 510-511; 1978.

Steps involved in the transformation of a normal cell to a malignant one were investigated by transforming normal continuous cell lines in vitro and selecting for transformants by their growth in agarose. The tumorigenicity of the cloned, transformed cell lines was tested in normal and immunodeficient (ID) mice. Based on tumorigenic potential, the cells were divided into three classes: O, growth in agarose only; I, growth in agarose + tumor formation in ID mice; C, growth in agarose + tumor formation in normal and ID mice. Normal fibroblast cells (N) derived from BALB/c, BALB.B, and C57BL/6 embryo cultures were transformed virally (simian virus 40 or Abelson murine leukemia virus) or chemically (20-methylcholanthrene, N-methyl-N'-nitro-N-nitrosoguanidine, 11,12-dihydro-3-methylcholanthrene-11,12-epoxide or 5,6-dihydrobenz(a)anthracene-5,6-epoxide). Of 14 agarose-positive cell clones derived from viral transformation, 2 were classified O, 9 I, and 3 C. Of 21 clones derived from chemical transformation, 2 were classified O, 4 I, and 15 C. Overall, 31/35 agarose-positive clones formed tumors in ID mice; ie, there was an 89% correlation between in vitro growth in agarose and in vivo tumorigenicity. The proportion of I tumorigenic transformed lines was 9/12 for the virally transformed lines and 4/19 for the chemically transformed lines. Expressed in terms of surveillance efficiency, 75% of

the former and 21% of the latter were susceptible to a surveillance mechanism that was impaired in ID mice. It is not known whether all C lines are derived from I intermediates; however, clones that have undergone the mutation N to I require another mutation, I to C, to grow as a tumor in normal mice. (7 refs)

- 78-7111 **Lymphoma after S.L.E. (Letter to Editor).** (Eng) Gibbs, A. R. (Dept. Pathology, Welsh Natl. Sch. Medicine, Cardiff CF4 4XN, Wales); Seal, R. M. *Lancet* 2(8098): 1046; 1978.

Primary pulmonary malignant lymphoma developed in a 57-yr-old woman 20 yr after the onset of systemic lupus erythematosus, the first published case of this association. Microscopy revealed a nodular, poorly differentiated-type non-Hodgkin's lymphoma. (no refs)

- 78-7112 **Needle Tract Implantation in a Case of Needle Biopsy for Lung Cancer.** (Eng) Ichiba, M. (Dept. Surgery, Tokyo Medical Coll., 6-7-1, Nishishinjuku, Shinjuku-ku, Tokyo 160, Japan); Aida, M.; Konaka, C.; Hayata, Y. *Jpn J Clin Oncol* 8(1): 71-73; 1978.

A pulmonary adenocarcinoma was implanted in the tract of a small-bore needle biopsy in a 60-yr-old man. It was diagnosed 19 mo after the original biopsy. This was the first case of tumor cell implantation encountered among a series of 582 biopsies and it is the eighth case reported in the literature. (13 refs.)

- 78-7113 **Etiology of Malignant Pericardial Mesothelioma.** (Ger) Eck, H. (Pathologisches Institut, Städtisches Krankenhaus Höchst, Gotenstr. 6-8, 6230 Frankfurt 80, W. Germany); Berg-Schlosser, V. *Dtsch Med Wochenschr* 103(44): 1751-1753; 1978.

Malignant mesothelioma of the pericardium was diagnosed in a 45-yr-old man and in a 77-yr-old woman. Mild lung fibrosis and a slightly but significantly increased number of asbestos needles in the pulmonary parenchyma were found in the 45-yr-old man who, being a scrap dealer, may have been exposed to asbestos occupationally; however, other unknown etiological factors must also have been involved. No asbestos bodies were found in the 77-yr-old woman, but hemosiderin, obviously originating from phagocytized erythrocytes, was found in the tumor cells. This phagocytic activity of the mesothelial cells seems to be a proof of the mesothelial origin of such tumors. (9 refs)

- 78-7114 **Epithelial Dysplasia in Cancerous and Noncancerous Esophagi.** (Eng) Sato, E. (Second Dept.

Pathology, Kagoshima Univ. Faculty Medicine, Kagoshima 890, Japan); Tokunaga, M.; Sakae, K.; Mukada, T.; Sasano, N. *Tohoku J Exp Med* 124(2): 117-128; 1978.

The prevalence, severity, distribution, and histology of esophageal dysplasia were studied in 86 grossly normal human specimens obtained at necropsy and in 100 specimens of esophageal carcinoma obtained at surgery. Fifty-four of the necropsy specimens showed dysplasia, 32 from men and 22 from women. Seventeen men and three women showed moderate or severe dysplasia or carcinoma in situ (CIS). The incidence and severity of dysplasia seemed to increase with alcohol and cigarette use, but the results were inconclusive. The incidence and severity also increased with age. Chronic inflammation, glycogenic acanthosis, and hyperplasia of the duct epithelium were observed frequently, but they were not significantly related to the incidence of dysplasia. Most of the surgical specimens showed a more advanced dysplasia. Those that showed extensive dysplasia tended to have a mosaic pattern of CIS and dysplasia. Carcinoma and extensive dysplasia was found in 24 instances, carcinoma and multifocal dysplasia in 46. Of the 24 cases that did not have treatment prior to surgery, 15 showed dysplasia coexisting with the carcinoma, and 19 showed multicentric foci of carcinoma. The invasive carcinoma sometimes did not show a sharp demarcation from the surrounding mucosa, especially in cases involving well-differentiated squamous cell carcinoma. The results indicate that dysplasia plays a significant role as a precursor lesion in esophageal carcinogenesis. (19 refs)

- 78-7115 **Conversion of a Poorly Differentiated Human Adenocarcinoma to Ascites Form with Invasion and Metastasis in Nude Mice: Brief Communication.** (Eng) Takahashi, S. (First Dept. Surgery, Nara Medical Univ., 840 Shijo-cho, Kashihara, Nara 634, Japan); Konishi, Y.; Nakatani, K.; Inui, S.; Kojima, K.; Shiratori, T. *J Natl Cancer Inst* 60(4): 925-929; 1978.

Cells from a poorly differentiated Borrmann Type III adenocarcinoma of the stomach of a 70-yr-old man were converted to the ascites form in nude BALB/c mice. The tumor cells were passaged sc in nude mice following removal from the patient. At passage 9, the histology of the cells was similar to that of the original tumor; 10^7 cells were then injected ip into nude mice. The injection resulted in ascites carcinoma with invasion (carcinomatous peritonitis) and liver metastasis. Serial transfers of 1.5×10^5 dispersed cancer cells into the peritoneal cavity of nude mice converted this adenocarcinoma to an ascites form. Hemorrhagic ascites accumulated within 3 wk at the first passage and 4-6 wk in serial passages. Carcinomatous peritonitis occurred consistently in the diaphragm, mesenteries, omentum, and pancreas; metastases were seen in the liver and spleen. Iv injection of 6×10^5 cells at ascites passage 3 into nude mice produced metastatic lesions in the lung and heart. The histology of the invasive and metastatic lesions was similar to that of the original tumor. (8 refs)

8-7116 **Relationship Between Gastric Polyps and Gastric Mucosa.** (Cze) Mertl, L. (I. vnitřní klinika katedry dětského lékařství, Fakultní nemocnice Pod Biskupem, V Uvalu 84, Prague 5, Czechoslovakia); Gregor, O. *Československá Lek 24(9): 842-847; 1978.*

The gastric mucosa and gastric secretion were studied in 74 patients with gastric polyps. Malignant (carcinomatous) polyps were found in 19 patients, adenomatous polyps in 25, hyperplastic polyps in 24, eosinophilic granulomas in 2, fibromas in 2, hemangioma and neurinoma in 1 each. The gastric mucosa was normal in 7 patients with benign polyps; superficial gastritis was found in 10 patients with benign and 2 with malignant polyps, atrophic gastritis in 31 patients with benign and 10 patients with malignant polyps, and atrophic gastritis with metaplasia in 7 patients with benign and malignant polyps each. Histamine-resistant achlorhydria was found in 38 patients with benign polyps and in 18 patients with malignant polyps. The findings indicate that chronic inflammation is a primary process in the pathogenesis of mucosal polyps, the polyp being a secondary, local manifestation of gastric mucosa disease. Chronic inflammation also appears to lead to the development of hyperplastic adenomas. The findings suggest a relationship between alterations in the gastric mucosa and tumor growth in the stomach. (24 refs)

8-7117 **'Small Intestinal Type' and 'Colonic Type' Intestinal Metaplasia of the Human Stomach, and Their Relationship to the Histogenic Types of Gastric Adenocarcinomas.** (Eng) Teglbjaerg, P. S. (Inst. Pathology, Odense Univ. Hosp., DK 5000 Odense, Denmark); Nielsen, H. O. *Acta Pathol Microbiol Scand [A] 86(5): 351-355; 1978.*

The frequency of different histochemical types of intestinal metaplasia (IM) was compared in patients with gastric carcinomas of presumed intestinal histogenesis and patients with tumors of nonintestinal histogenesis. Operative specimens were collected from 27 consecutive patients undergoing surgery for stomach cancer. Seven tumors were classified as intestinal cell carcinomas, 6 as pylorocardiac gland carcinomas, 3 as mucous cell carcinomas, and 1 was unclassifiable. Small intestinal type IM, identified by the staining of goblet cells with alcian blue pH 2.6-PAS, was present in all 24 stomachs with IM. Colonic type IM, identified by the reaction of goblet cells with periodate-borohydride/KOH/PAS, was found in scattered IM foci in 11 cases. The frequency of the latter type was significantly greater in stomachs with tumors of intestinal histogenesis than in those with tumors of nonintestinal histogenesis (ie, intestinal cell carcinomas vs pylorocardiac carcinomas). It is suggested that colonic type IM may have a premalignant potential and that the small intestinal type is merely of a reactive character. (18 refs)

78-7118 **Genetic and Communicable Effects on Carcinoembryonic Antigen Expressivity in the Cancer**

Family Syndrome. (Eng) Guirgis, H. A. (Dept. Preventive Medicine/Public Health, Creighton Univ. Sch. Medicine, Omaha, NB, 68178); Lynch, H. T.; Harris, R. E.; Vandevoorde, J. P. *Cancer Res 38(8): 2523-2528; 1978.*

The potential of plasma carcinoembryonic antigen (CEA) as a discriminant of genetic cancer risk was tested in 272 members of six pedigrees manifesting the cancer family syndrome (CFS) and 191 normal controls. In addition, the role of age and duration of cigarette smoking in CEA expression was quantitated, and connubial effects on CEA levels in this syndrome were determined. The CEA distributions per se were transformed to square root CEA (srCEA) to correct for skewness and kurtosis. Duration of smoking and age showed a positive correlation with srCEA level in both family members and controls. The mean srCEA did not differ between family members and controls of the same smoking status. However, when pedigree data were classified by closeness of relationship to cancer patients, there was a significant linear increase in mean srCEA with increasing genetic cancer risk. Unrelated spouses had mean srCEA levels similar to those for the corresponding risk class of their direct-line mates, and the intraclass correlations of srCEA between direct-line relatives and their spouses approached significance when both spouses were concordant for smoking status. It is suggested that there is a genetic-connubial effect on plasma CEA levels that presumably is due to a common environmental agent acting with the degree of genetic predisposition to cancer. An integrative target cell theory is proposed to explain CEA expressivity in the CFS. The model involves the interaction of the CFS genotype with physical factors (cigarette smoking) and a communicable agent (oncogenic virus). (36 refs)

78-7119 **Peutz-Jeghers Syndrome. Intussusception and Associated Malignancies.** (Fre) Besson, A. (Service de Chirurgie A, 1011 CHUV, Lausanne, Switzerland); Delacretaz, F.; Pescia, G.; Vecerina, S.; Saegesser, F. *Chirurgie 104(2): 117-130; 1978.*

The case histories of six patients representing three families with Peutz-Jeghers syndrome (PJS) are presented. Two of the patients developed cancer and eventually died with multiple metastases. The first patient, a 34-yr-old man, had multiple polyps of the stomach and small and large intestines throughout his life, and he had undergone resection of the jejunum and ileum. At the age of 23, a polyp in the sigmoid colon was invaded by an undifferentiated adenocarcinoma. Ten years later, the patient died of complications following surgery for rectal polyps. Hepatic metastases were observed at this time. The other patient, the mother of two sons with the syndrome, died at the age of 48 with multiple metastases from adenocarcinoma of the uterus. She had the cutaneous symptoms of PJS, and x-rays had shown polyps in the ileum and rectum, but the latter had never given rise to clinical symptoms. The literature reports a 5% incidence of gastrointestinal cancers associated with PJS; 1/6 women with the syndrome develop an ovarian tumor, frequently before age 15. (41 refs)

- 78-7120 Chromosomes and B and T Cells in Mycosis Fungoides.** (Eng) Goh, K. (Medical-Oncology Unit, Monroe Community Hosp., 435 E. Henrietta Road, Rochester, NY, 14603); Reddy, M. M.; Joishy, S. K. *Am J Med Sci* 276(2): 197-204; 1978.

Cytogenetic studies were performed using peripheral blood WBC from four male patients (aged 59-78 yr) with mycosis fungoides (MF). The WBC cultures were stimulated with phytohemagglutinin or pokeweed mitogen. Pseudodiploid cells were found in all patients, and they varied between 24% and 42% in each culture. The most consistent chromosomal pattern among the pseudodiploid metaphases was a larger chromosome number 2, the long arm of which appeared longer than that of its homolog. Chromosomal fragments were seen in various proportions in cultures from three patients. The absolute number of lymphocytes from these three patients was lower than that in control cultures. The absolute number of T lymphocytes was also reduced, but only one patient showed a lower proportion of T lymphocytes. Two patients showed a higher frequency but lower absolute number of B cells compared with controls; the frequency and absolute number of B cells were within normal limits in the third patient. The results support the concept that MF is a neoplastic disease, and they suggest that MF and Sezary syndrome are two separate entities. (16 refs)

- 78-7121 Pigmented Xerodermoid: First Report of a Family.** (Eng) Hofmann, H. (Universitäts-Hautklinik Heidelberg, Vosstrasse 2, D-69 Heidelberg, W. Germany); Schnyder, U. W. *Bull Cancer (Paris)* 65(3): 347-350; 1978.

A family concentration of pigmented xerodermoid, a xeroderma pigmentosum variant, is presented. In a second generation of 13 children, 2 men and 2 women patients had typical clinical signs of xeroderma pigmentosum, including increased pigmentation, strong sunburn reactions, an inordinate number of freckles, solar keratoses, and multiple squamous cell carcinoma of the skin in later life; 3/13 children died in early infancy, and 3 of the 6 remaining siblings died of internal malignancies. Their deceased parents had apparently been normal, as were 22 offspring of this second generation, except for marked sun sensitivity and blistering in 2 sons of one patient. One of these men subsequently died of mammary carcinoma. Of 24 members of the fourth generation, 2 had extensive body freckles, but at an age range of 7-25 no other skin aberrations were so far reported. DNA repair mechanisms were investigated in UV-irradiated lymphocytes and epidermal cells of one of the second generation patients. In these cells, the base-excision repair mechanism was normal but the DNA synthesis time was abnormally long, indicative of pigmented xerodermoid. This appears to be the first report of a family concentration of this condition. (8 refs)

- 78-7122 Morphology and Metastatic Nature of Induced Hepatic Nodular Lesions in C57BL x C3H F₁**

Mice. (Eng) Vesselinovitch, S. D. (Dept. Pathology, Pritzker Sch. Medicine, Univ. Chicago, Chicago, IL, 60637); Mihailovich, N.; Rao, K. V. *Cancer Res* 38(7): 2003-2010; 1978.

The metastatic capabilities of well-defined nodular hepatic lesions in mice induced by benzo(a)pyrene (single treatment of 75-150 µg/g), ethylnitrosourea (single treatment of 60-120 µg/g), benzidine-2HCl (continuous treatment of 100 ppm in the diet), and diethylnitrosamine (4 treatments totaling 6-12 µg/g) were evaluated to determine if it is possible to diagnose cancer in the mouse solely by histological examination of the primary lesion. Coded liver and lung tissues from 1,264 treated (C57BL/6J x C3HeB/FeJ)F₁ mice were assessed independently for the presence of primary nodular lesions and metastases, respectively. Primary lesions were classified according to their size, cell morphology, and growth patterns into hyperplastic, adenomatous, and trabecular nodules. None of the 126 mice bearing hyperplastic nodules had pulmonary metastases. Four of 291 mice with adenomatous nodular lesions showed metastases. However, of 733 mice with the trabecular type of nodular lesions, alone or in combination with other lesions, 266 showed pulmonary metastases. The metastases were first detected in mice dying between 51 and 60 wk of age (5%), and the rate increased as a function of age at death, reaching an incidence of 51% in mice surviving > 81 wk. It was concluded that nodules showing trabecular and the more anaplastic solid sheet type of growths represented bona fide hepatocellular carcinomas in the mouse. However, before bioassay data of mouse carcinogenesis are interpreted, the nodular lesions should be classified, the experimental conditions fully assessed, and factors modifying hepatocarcinogenesis considered. (23 refs)

- 78-7123 Frequency of Occurrence of Hepatocellular Carcinoma in Patients with Porphyria Cutanea Tarda During Long-Term Observation (Meeting Abstract).** (Eng) Kordac, V. (First Medical Clinic, Charles Univ., Prague, Czechoslovakia). *J Clin Chem Clin Biochem* 16(1): 67; 1978. (no refs)

- 78-7124 Primary Liver Carcinoma: The Possible Role of Sex Hormones (2 Letters to Editor).** (Dut) Zuidema, P. J. (Amsterdam, Netherlands); Houwert, G. J.; Brandt, K. H.; Bronkhorst, F. B. *Ned Tijdschr Geneesk* 122(42): 1634; 1978.

Pathologically increased serum alpha-fetoprotein levels were found in 6/14 patients with primary liver carcinoma. Two other patients with mixed hepatocellular-cholangiolar liver carcinoma were negative. (4 refs)

- 78-7125 Focal Nodular Hyperplasia of the Liver and Oral Contraceptives.** (Eng) Kinch, R. (Depts.

Obstetrics and Gynaecology, Montreal General Hosp., 1650 Cedar Ave., Montreal, Quebec, Canada H3G 1A4); Lough, J. *Am J Obstet Gynecol* 132(7): 717-727; 1978.

Six cases of focal nodular hyperplasia discovered since 1973 at Montreal General Hospital are presented. Two cases are reported in depth; four were published previously and are summarized. The age of the patients ranged from 21 to 45 yr (four were > 40 yr), and all six had taken oral contraceptives for 4-12 yr. Four had taken preparations containing mestranol and one had taken 50 µg ethinyl estradiol + 2.5 norgestrel for 6 yr; the product was unknown in the sixth patient. Each patient had suffered long-term (2-3 yr) upper epigastric discomfort and pain that was mostly located in the area over the hepatic lesion. Upper gastrointestinal series were negative and liver enzyme tests, when performed, were normal. All tumors had the same gross appearance: 3- to 6-cm nodular masses whose resected surface was yellowish and multinodular, with a central fibrous scar radiating into the parenchyma (similar to focal postnecrotic cirrhosis). The normal lobular architecture was replaced by nodules of regenerating parenchyma, with the hepatocytes being arranged in a pseudoacinar pattern. Ductular proliferation was seen along the margins in the vascular stroma. Focal nodular hyperplasia and hepatic adenoma are suggested to be two distinct processes. It is postulated that the estrogen component of the oral contraceptives causes vascular lesions characterized by myointimal hyperplasia and thrombosis, leading to infarct, necrosis, and subsequent nodular hepatic regeneration. Gray-scale ultrasonography is an effective screening method that also aids in a definitive diagnosis. (18 refs)

78-7126 Liver Cell Adenoma Associated with Oral Contraceptive Hormone Therapy. (Eng) Nicolaides, N. J. (133 Leichhardt St., Brisbane, Queensland 4000, Australia). *Med J Aust* 2(6): 274-276; 1978.

The case of a 40-yr-old woman who developed a hepatic adenoma in association with 7 yr of oral contraceptive therapy (norethisterone/ethinyl estradiol) is presented. Upon removal of the tumor, widespread interparenchymal hemorrhage was discovered. Catastrophic intraperitoneal hemorrhage could well have eventuated had the capsule been breached, and needle biopsy of such a lesion may have proved disastrous. (29 refs)

78-7127 A Close Association of Hepatocellular Carcinoma with Cirrhotic Changes Among 275,857 Pathological Autopsy Cases in Japan During 16 Years from 1958 to 1973. (Eng) Miyaji, T. (Japanese Inst. Pathology, Osaka 530, Japan). *Kurume Med J* 24(Suppl): S63-S79; 1977.

The association between hepatocellular carcinoma (HCC) and cirrhotic changes was studied based on 275,857 autopsy cases registered in Japan during 1958 through 1973. The most

frequent malignancy was gastric carcinoma, followed by leukemia, bronchogenic carcinoma, and HCC. During the 16-yr period, 7,002 HCC's and 15,136 cases of cirrhotic changes not associated with HCC or cholangiocellular carcinoma were registered. A gradual increase in HCC's, especially in men, was noted. Seventy percent of the HCC's in men and 53.3% of those in women were associated with cirrhosis, as were 20.7% of cholangiocellular carcinomas in men and 10.7% of those in women. Liver fibrosis was associated with HCC in 1.6% of affected men and 2.0% of the women and with cholangiocellular carcinoma in 8.7% of affected men and 22.0% of the women. Of the cirrroses associated with HCC, 85.8% of those in men and 80.8% of those in women were either unspecified (US) cirrhosis or the Nagayo-Miyake B-type (posthepatic or macronodular) cirrhosis. Among the cirrroses not associated with carcinoma, 50.0% in men and 44.7% in women were of the B and US types. A'-type cirrhosis (transitional between postnecrotic and B type) and B'-type cirrhosis (transitional between fibrosis and B type) were associated with HCC in 29.7% and 23.2% of men, respectively, and in 20.7% and 21.5% of women. The number of postnecrotic and A'-type cirrroses associated with HCC tended to increase over the 16-yr interval. The peak of HCC in men and women was at 55 yr and that of B type and US cirrhosis was at 60 yr. (66 refs)

78-7128 The Histiocytic Origin of the Multinucleated Giant Cells in Myeloma Kidney. (Eng) Factor, S. M. (Dept. Pathology, Albert Einstein Coll. Medicine, Bronx, NY); Winn, R. M.; Biempica, L. *Hum Pathol* 9(1): 114-120; 1978.

The cellular origin of the syncytial multinucleated giant cells in a patient with myeloma kidney disease was determined by light and electron microscopy (EM) and by immunofluorescence. The patient was a 71-yr-old black man with uremia and dysphagia. Even though renal tissue was not fixed for > 20 hr after the patient's death, the multinucleated giant cells appeared well-preserved, contrasting with the marked degeneration of the tubular epithelium (TE). There was a consistent correlation between the presence of giant cells around tubular casts and breaks in the basement membrane. Although casts were present within Bowman's space, they were never associated with giant cells, possibly because the parietal glomerular basement membrane is relatively resistant to penetration by inflammatory cells. One possible source of misinterpretation of a direct transition between the giant cells and the TE was noted by light and EM. Degenerated TE was frequently found in close apposition with giant cells or, in some instances, within spaces enclosed by giant cell cytoplasmic processes. However, EM showed that there was no fusion of cell membranes. There was also a clear difference in staining reactions and a sharp delimitation of cell contours between the giant cells and the TE when the renal tissue was stained for reticulin fibers. Immunofluorescence studies suggested that the multinucleated giant cells phagocytose fragments of proteinaceous casts. The histiocytic origin of the

giant cells was confirmed directly by EM; a lock and key configuration, numerous mitochondria, large numbers of membrane-bound structures, dense bodies, numerous cytoplasmic vesicles, and luminal projections of the plasma membrane were present. It is concluded that the giant cells arise in the interstitial tissues, migrate through breaks in the tubular basement membrane, and engulf and surround intratubular protein casts. (23 refs)

- 78-7129 An Unusual Osteosclerotic Metastasis to the Calcaneus in Basal-Cell Adenocarcinoma of the Kidney in a Child.** (Cze) Mrzena, V. (Ortopedicka klinika, Fakulta detskeho lekarstvi KU, Karlovo nam. 1, 128 00 Prague 2, Czechoslovakia); Belikova, H. *Acta Chir Orthop Traumatol Cech* 45(3): 214-216; 1978.

Osteosclerotic metastasis to the calcaneus was the first symptom of a kidney tumor in a 13-yr-old boy. Basal-cell adenocarcinoma of the kidney was found at surgery. (5 refs)

- 78-7130 Filiform Polyposis.** (Eng) Zegel, H. G. (Dept. Radiology, Hosp. Univ. Pennsylvania, 3400 Spruce St., Philadelphia, PA, 19104); Laufer, I. *Radiology* 127(3): 615-619; 1978.

Filiform polyposis presents a characteristic radiographic appearance of long, filamentous filling defects in an otherwise normal colon. It has been documented in patients with a prior history of ulcerative or granulomatous colitis. In addition, a solitary filiform polyp was discovered in a patient with previously undiagnosed inflammatory bowel disease, and in a patient with Crohn's disease involving the small bowel. Filiform/polyposis may represent a nonspecific sequela of diffuse mucosal inflammation, and it should not be mistaken for a neoplastic form of polyposis. (5 refs)

- 78-7131 Cell Proliferation of the Colorectal Mucosa in Normal Subjects and Patients with Preneoplastic or Neoplastic Pathology of the Colon and Rectum.** (Fre) Maskens, A. P. (Unite de Recherche Cancerologique, Clinique Saint-Michel, Rue de Lantsheere 19, B-1040 Brussels, Belgium). *Acta Gastroenterol Belg* 41(5/6): 226-240; 1978.

Proliferation of cells in the colorectal mucosa was studied by radioactive thymidine uptake in samples from 25 patients (26-71 yr of age; av age 54 yr) with intestinal polyps (14 adenomatous, 9 hyperplastic, 2 villous) and in samples of normal mucosa of 5 tumor-free subjects and 11 patients with colorectal cancer. Distribution of radioactivity was graphed for the lower, middle, and upper thirds of the glandular epithelium in normal and pathological mucosa. In mucosa from normal patients, 66.3% of the radioactivity was recovered in

the lower third, 33.5% in the middle third, and 0.3% in the upper third. In mucosa samples of normal appearance from the cancer patients, 57.7% of the radioactivity was recovered in the lower third, 39.4% in the middle third, and 3.0% in the upper third. In mucosa samples from hyperplastic polyps, 40% of the radioactivity was recovered in the lower third, 93.5% in the lower two-thirds, and 7.5% in the upper third. The location of radioactivity was difficult to calculate exactly in samples from adenomatous polyps due to the morphology and size of the glands; however, av uptake was elevated (23.1% compared with 8.3%) toward the surface regions. Only two samples from villous polyps were analyzed for radioactivity, and the fragile villi made counting difficult. (32 refs)

- 78-7132 The Background Processes in the Development of Endometriosis and Endometrioid Cancer of the Ovaries.** (Rus) Kolosov, A. E. (N. N. Petrov Res. Inst. Oncology, USSR Ministry Health, Leningrad, USSR). *Vopr Onkol* 24(3): 44-50; 1978.

Surgical specimens from 87 patients with ovarian endometriosis and 23 patients with endometrioid carcinoma of the ovaries were analyzed morphologically to determine the changes or background processes leading to these conditions. The group included 61 pre- and 49 menopausal patients. The background lesions included microfollicular cysts (54 patients) and hyperplasia of the ovarian stroma (73 patients: 22 with diffuse, 37 with focal, and 14 with mixed hyperplasia). The premenopausal patients had predominantly follicular cysts and focal hyperplasia, but the menopausal patients had diffuse hyperplasia. Hyperplasia of the endometrium was diagnosed in 41 patients. (16 refs)

- 78-7133 Laurence-Moon-Bardet-Biedl Syndrome and Endometrial Carcinoma.** (Ger) Peiker, G. (Frauenklinik, Friedrich-Schiller-Universitat, Bachstr. 18, DDR-69 Jena, E. Germany); Bohm, W.; Carol, W. *Zentralbl Gynaekol* 100(17): 1126-1130; 1978.

Laurence-Moon-Bardet-Biedl syndrome associated with Stein-Leventhal syndrome and polycystic ovaries was diagnosed in an 18-yr-old obese woman with irregular monophasic menstrual cycles. Endometrial adenocarcinoma was found 5 yr later. Compared with normal values, the total estrogen excretion in the urine was increased (43 $\mu\text{g}/24$ hr), and the 17-ketosteroid excretion was low (2 mg/24 hr). The genesis of the endometrial adenocarcinoma can be explained by the extraglandular (peripheral) aromatization of androgens into estrogens, especially by the transformation of androstenedione into estrone and the absence of estradiol and progesterone during the long-term exposure of the endometrium to estrone. (9 refs)

78-7134 Choriocarcinoma with Metastasis in a Postmenopausal Woman. (Eng) Dougherty, C. M. (Louisiana State Univ. Sch. Medicine, LSU Unit, 5825 Airline Highway, Baton Rouge, LA, 70805); Cunningham, C.; Mickal, A. *Am J Obstet Gynecol* 132(6): 700-701; 1978.

The case of a black woman who had a 34-yr latent period between her last pregnancy (at age 52) and subsequent metastatic choriocarcinoma (ChC) is described. Amenorrhea began after her last pregnancy. Molar remnants, including hydropic villi, retained within the uterus for possibly 34 yr furnished proof of primary trophoblastic neoplasia. Viable, growing ChC in the lungs at autopsy proved the malignant potential of the tumor. (4 refs)

78-7135 Primary Adenocarcinoma of the Vagina: A Case Report with Light and Electron Microscopic Study. (Eng) Navas Palacios, J. J. (Dept. Anatomic Pathology, Ciudad Sanitaria de la Seguridad Social "1 de Octubre," Carretera de Andalucía, km. 5.4, Madrid 26, Spain); de Agustin de Agustin, P. P.; Martinez Tello, F. J. *Acta Cytol (Baltimore)* 22(5): 358-361; 1978.

The case of a 46-yr-old woman with primary vaginal adenocarcinoma unrelated to hormone therapy is reported, and the results of histological and ultrastructural studies are presented. The patient presented with vaginal bleeding after intercourse, and, on examination, an exophytic necrotic mass was found to fill most of the vagina. After removal of the vaginal content, the cervix appeared unremarkable; a 6-cm implantation area was noted on the posterior vaginal wall and fornix. The presence of a primary site other than the vagina was ruled out by lower gastrointestinal tract and chest x-rays, laparoscopy, and rectosigmoidoscopy. The pathologic diagnosis of primary adenocarcinoma of the endometrioid subgroup was made. The ultrastructural features were different from those of clear cell-type vaginal adenocarcinoma. Glycogen aggregates were absent, and the rough endoplasmic reticulum was sparse and did not form stacks. Endocervical type adenocarcinoma was ruled out because of the absence of characteristic mucinous secretory granules. The ultrastructure of the mesonephric nests was different from that of tumors with a true mesonephric origin. The ultrastructural features of renal cell carcinoma were also absent in this case. Because of the ultrastructural features of normal endometrium and endometrial adenocarcinoma are similar to the observations in the present case, it is concluded that the adenocarcinoma was of müllerian origin. (27 refs)

78-7136 Viral Particles in Cervical Condylomatous Lesions. (Eng) Della Torre, G. (Divisione di Oncologia Sperimentale A., Istituto Nazionale Tumori, Via G. Venezian 1, 20133 Milan, Italy); Pilotti, S.; de Palo, G.; Rilke, F. *Tumori* 64(5): 549-553; 1978.

Koilocytosis is a pathologic alteration of squamous epithelial cells of the uterine cervix that is characterized by small nuclei with an indented outline and clumped chromatin surrounded by a halo of transparent cytoplasm. Biopsy specimens from eight patients with condylomatous lesions of the uterine cervix were examined histologically and ultrastructurally. Cervical smears showed koilocytotic atypia in varying numbers of cells in all cases, and malignant epithelial tumor cells were found in one case. In seven cases, the colposcopic pattern was that of early condylomatosis; one case was colposcopically negative except for a small leukoplakia. Ultrastructural examination demonstrated many intranuclear viral particles morphologically identical to papovavirions in the koilocytotic epithelial cells of four patients. The particles were also identical to those of the papilloma virus found in genital condylomata acuminata, suggesting that cervical condylomatous lesions are related etiologically to condylomata acuminata. (8 refs)

78-7137 Early Vulvar Carcinoma. (Eng) Kunschner, A. (Dept. Obstetrics and Gynecology, Magee-Womens Hosp., Univ. Pittsburgh, Pittsburgh, PA, 15213); Kanbour, A. I.; David, B. *Am J Obstet Gynecol* 132(6): 599-606; 1978.

Fifty-eight cases of early vulvar neoplasia recorded in Tumor Registry at Magee-Womens Hospital during 1947-1976 were reviewed. There were 23 cases of squamous cell carcinoma in situ; 16 cases of carcinoma in situ with microinvasion, and 19 cases of occult carcinoma. All these cases were treated surgically with the modality varying from local excision to radical vulvectomy with inguinal lymph node dissection. Regardless of the surgical therapy, none of the patients died of the vulvar malignancy. As part of routine gynecologic surveillance, more critical attention should be given to examination for early epithelial neoplastic lesions of the vulva. This can be accomplished by simple inspection and the use of special diagnostic aids. Biopsy specimens should be reviewed histologically for depth of invasion and the treatment individualized to the patient, with considerations for conservation, especially in younger women. (16 refs)

78-7138 Neoplastic Potential of the Dysgenetic Gonad. Case Report. (Eng) Fayed, J. A. (Truman Medical Center, 2301 Holmes, Kansas City, MO, 64108); Jonas, H. S.; Masterson, B. J. *Mo Med* 75(11): 564-566; 1978.

The case of a 16-yr-old girl with a 46/XY karyotype in whom bilateral gonadoblastoma developed in the gonadal streaks is presented. Gonadal dysgenesis was diagnosed on the basis of the patient's karyotype and lack of secondary sex characteristics. An infantile uterus was found at laparotomy, and streak gonads were identified in the position of normal ovaries. Biopsy disclosed gonadoblastoma, and a total hysterectomy and bilateral salpingogonadectomy were performed. This case il-

illustrates the malignant potential of the dysgenetic gonad. (10 refs)

- 78-7139 Carcinoma In Situ of the Bladder. Report of 10 Cases.** (Fre) Steg, A. (Clinique Urologique, Hopital Cochin, 27 rue du Faubourg Saint-Jacques, 75674 Paris Cedex 14, France). *Ann Urol (Paris)* 12(2): 73-82; 1978.

The symptoms, diagnosis, and therapeutic aspects of carcinoma in situ (CIS) of the urinary bladder are described, and the histology and significance of CIS for invasive carcinoma are discussed in connection with 10 cases of CIS (9 men, 1 woman, aged 51-74 yr). No occupational carcinogenic factors were found. Differentiation between CIS and dysplasia is often very difficult. In simple dysplasia, the number of cell layers is increased. In atypical dysplasia of the epithelium, this change is also associated with nuclear anomalies, while the lack of cohesion and maturation, and giant cells are characteristic of CIS. It is a well-established fact that CIS can evolve into invasive carcinoma; such transformation is seen in 20-50% of all cases, and takes place in 1-6 yr after the diagnosis of CIS. Each invasive carcinoma of the urinary bladder in the 10 patients had a CIS stage. (23 refs)

- 78-7140 Natural History of Papillary Lesions of the Urinary Bladder in Schistosomiasis.** (Eng) Kuntz, R. E. (Dept. Parasitology, Southwest Foundation Res. and Education, San Antonio, TX, 78284); Cheever, A. W.; Bryan, G. T.; Moore, J. A.; Huang, T. C. *Cancer Res* 38(11, part 1): 3836-3839; 1978.

Attempts to autologously and heterologously transfer bladder lesions in monkeys (*Cebus apella*) infected with *Schistosoma haematobium* (1,000 or 2,000 cercariae) are reported, along with the natural history of papillary transitional cell lesions in these animals. The animals were examined by laparotomy and cystotomy 94-164 wk postinfection, and six were chosen for autologous transplant studies, three for heterologous transplant studies. These animals were followed for 14-136 wk postcystotomy and then subjected to necropsy. At cystotomy, variable epithelial hyperplasia was observed in the urinary bladder of the nine monkeys. All biopsy specimens showed marked *S. haematobium* infection. At necropsy, involution of the urothelial hyperplasia had occurred in 8/9 monkeys; the only monkey in which the hyperplasia persisted was also the only monkey with marked active infection. No growth of transplanted epithelium was evident by palpation or by gross or microscopic examination. The results indicate that proliferative transitional cell lesions have restricted growth potential. (15 refs)

- 78-7141 Desmosome Ultrastructure and the Biological Behavior of Chemical Carcinogen-induced Uri-**

nary Bladder Carcinomas. (Eng) Pauli, B. U. (Dept. Pathology, Rush Medical Coll., 1753 W. Congress Parkway, Chicago, IL, 60612); Cohen, S. M.; Alroy, J.; Weinstein, R. S. *Cancer Res* 38(10): 3276-3285; 1978.

The relationship between desmosomes and tumor invasiveness was studied in urinary bladder transitional cell carcinomas induced in male Fischer rats by N-[4-(5-nitro-2-furyl)-2-thiazolyl]formamide (0.2% of diet for 26 wk). The number of desmosomes per unit area of plasma membrane was significantly reduced in the carcinomas. However, the percentage of cell surface area occupied by desmosomes were greater in carcinomas than in controls. This was accounted for on the basis of increases in desmosomal size, which result from squamous differentiation within the tumors. Areas of transitional cell differentiation and squamous differentiation demonstrated an equal capacity for invasiveness. Desmosomes were abundant in invading nests of tumor cells. These findings argue against the idea that decreased intercellular adhesion is a prerequisite for tumor invasion since strong interadhesion is probably a function of the area occupied by the intercellular junctions. (48 refs.)

- 78-7142 On the Etiology of Cancer of the Prostate** (Eng) Stein-Werblowsky, R. (Bikur Cholim Hosp., POB 492, Jerusalem, Israel). *Eur Urol* 4(5): 370-373; 1978.

Experiments demonstrating that the penetration of spermatozoa into the prostatic tissue of Wistar rats can cause abscesses, lithiasis, and carcinoma are presented. Nine of 20 animals developed one of these lesions following the injection of 0.1-0.2 ml of autologous sperm or homologous sperm into the anterior prostate at its junction with the neck of the bladder. (11 refs)

- 78-7143 On the Etiology of Testicular Tumors. An Experimental Study.** (Eng) Stein-Werblowsky, R. (Bikur Cholim Hosp., Jerusalem, Israel). *Eur Urol* 4(1): 57-59; 1978.

The etiology of testicular tumors was examined in Wistar rats. One testicle was extirpated and minced, and 0.2 ml of the resulting suspension was injected into the other testicle. Four of 10 rats developed testicular abnormalities within 32-47 days after injection. One rat had a tumor filling the whole abdomen; the testicle was not enlarged and the tumor, which was readily transplantable to homologous recipients, was apparent only on histological examination of the testis. The testicle of another rat was greatly enlarged, and normal tissue was wholly replaced by tumor. No distant metastases were observed, but the animal died of local spread 3 wk after orchietomy. These results demonstrate that testicular tumors can develop rapidly as a result of autofertilization. It

possible that mature, motile spermatozoa, instead of being discharged into the lumen of the seminiferous tubuli, penetrate the cells of the germinal epithelium. These results also point to a possible mechanism common to all carcinogenic processes, that of fertilization in the sense of nuclear fusion. The function of a carcinogenic agent may be the promotion of nuclear fusion. The chromosome abnormalities, such as excess of chromatin material and aneuploidy generally associated with cancer, lend further support to this hypothesis. (11 refs)

78-7144 Measurement of Urinary Epinephrine in Screening for Pheochromocytoma in Multiple Endocrine Neoplasia Type II. (Eng) Hamilton, B. P. (Dept. Medicine, Veterans Admin. Hosp., 3900 Lock Raven Blvd., Baltimore, MD, 21218); Landsberg, L.; Levine, R. J. *Am J Med* 65(6): 1027-1032; 1978.

A sensitive screening test for the pheochromocytoma component of multiple endocrine neoplasia type II (MEN-II), comparable to the calcitonin radioimmunoassay for medullary carcinoma of the thyroid (MCT), has been lacking. A large family with MEN-II was screened for pheochromocytoma by measuring 24-hr urinary catecholamines as norepinephrine and epinephrine fractions. Pheochromocytomas had been confirmed at surgery in six family members and at autopsy in one. All seven have had associated MCT. In an additional family members, only MCT has been detected. In 5/6 patients treated surgically, the urinary epinephrine fraction was the dominant or sole abnormality, and an epinephrine level $> 20 \mu\text{g}/24 \text{ hr}$, was diagnostic of pheochromocytoma. An increased urinary epinephrine fraction appears to be a characteristic feature of MEN-II pheochromocytomas, and measurement of this fraction a sensitive and reliable screening test in afflicted families. (16 refs)

78-7145 Seminoma in a 12-Year-Old Male with 46 XY/45 XO Karyotype. (Eng) Doll, D. C. (Dept. Medicine, West Virginia Univ. Medical Center, Morgantown, WV, 26506); Kandzari, S.; Jenkins, J. J.; Amato, S.; Jones, B. *Cancer* 42(4): 1823-1825; 1978.

The case of a 12-yr-old boy with pure seminoma in an undescended testis and 46 XY/45 XO karyotype is presented. The presence of the undescended testicle may have been a predisposing factor in the development of the malignancy. The abnormal chromosome constitution might have been an isolated finding with no relation to the patient's malignancy. However, it is possible that the abnormal karyotype was a contributing factor in both the undescended testicle and seminoma. (17 refs)

78-7146 Gonadal Tumors in Testicular Dysgenesis Syndrome. (Rus) Bronshtein, M. E. (Lab. Pathologic Anatomy, Inst. Experimental Endocrinology and Chemistry Hormones, Moscow, USSR); Iur'eva, N. P.; Golubeva, I. V. *Probl Endokrinol (Mosk)* 24(2): 30-35; 1978.

Of 84 patients (1.5-39 yr of age) with testicular dysgenesis syndrome, 26 had gonadal tumors. These tumors were diagnosed in only two patients at clinical examination; the rest were diagnosed at laparotomy. Tumor size ranged from 1.5 x 0.8 cm to 12 x 8 x 7 cm. Of the 26 patients, 10 had both testicles located either in the abdomen (9) or in the inguinal canal (1); 6 of the former had tumors of both gonads. Histologically, the tumors were classified as gonadoblastomas (18) or dysgerminomas (8). (11 refs)

78-7147 Malignant Transformation of Osteoid Osteoma. (Ger) Wahl, H. (Orthopädische Klinik, Medizinische Akademie Magdeburg, Schellingstrasse 3/4, DDR-301 Magdeburg, E. Germany); Dominok, G. W. *Zentralbl Chir* 103(22): 1490-1492; 1978.

An osteoid osteoma of the left tibia was diagnosed and partially resected in a 22-yr-old woman 3 yr after onset of the first symptoms. The symptoms (pain, swelling, and local hyperthermia) recurred 3 mo later. An osteosarcoma of the same site was diagnosed following partial resection performed about 2 yr after the first operation. To date, she remains free from any metastasis. (6 refs)

78-7148 Tissue Culture Studies on Human Giant Cell Tumor and Osteosarcoma. (Eng) Ishii, S. (Dept. Orthopaedic Surgery, Hokkaido Univ. Sch. Medicine, Kita-14-jo, Nishi-5-Chome, Sapporo, Hokkaido, Japan); Yamawaki, S.; Sasaki, T.; Usui, M.; Ubayama, Y.; Minami, A.; Yagi, T. *Int Orthop* 2(1): 77-86; 1978.

The cytological aspects of giant cell tumors (GCT) and osteosarcomas (OS) were investigated in vivo and in vitro. Primary cultures of GCT were successful in 33/41 trials, and 34/33 were transferred serially for more than two subculture passages. Two established lines (G-1 and G-2) were maintained for long-term cultivation. Primary OS cultures were successful in only 22/61 trials, and none of the cultures survived > 10 passages. Although round at initial cultivation, the giant cells became fibroblastlike after several hours; after 8-10 mo of cultivation, G-1 and G-2 became epitheliallike. The ultrastructure of the giant cells became increasingly complex with time in culture, displaying numerous microvilli, well-developed organelles, and free ribosomes, but the ultrastructure of the OS cells in primary culture

was essentially the same as that of the in vivo tumor cells. The GCT began with a normal chromosome complement of 46 and a superficially normal diploid karyotype. In contrast, the OS were characterized by aneuploid modal numbers. The number of polyploid and aneuploid cells in the GCT increased with passage number, resulting in a modal chromosome number of 65 at subculture 325. The lactate dehydrogenase (LDH) isozyme patterns of the GCT showed max activity in fractions IV and V; those of the OS showed max activity in fraction III. The LDH isozyme patterns of G-1 and G-2 showed max activity in fractions III and IV, respectively. In later transfer generations, G-1 and G-2 produced, in Wistar/Mk rats, tumors by sc inoculation and tumor death by iv inoculation. Sarcomatous stromal cells in the sc tumors were enmeshed in the hyaline network, resulting in the formation of osteoid or chondroid tissue. The results strongly suggest that the benign GCT may have become malignant during the long in vitro transfer period. (15 refs)

See also:

- *(Rev.): 78-6605, 78-6610, 78-6612, 78-6616, 78-6617, 78-6624, 78-6626, 78-6639, 78-6645, 78-6651, 78-6652, 78-6686, 78-6687, 78-6689, 78-6690, 78-6695, 78-6700, 78-6703, 78-6705, 78-6706, 78-6707, 78-6708.
- *(Chem.): 78-6712, 78-6713, 78-6715, 78-6722, 78-6723, 78-6728, 78-6729, 78-6733, 78-6739, 78-6742, 78-6744, 78-6756, 78-6757, 78-6765, 78-6776, 78-6780, 78-6786, 78-6789, 78-6790, 78-6795, 78-6797, 78-6798, 78-6801.
- *(Phys.): 78-6809, 78-6810, 78-6814, 78-6827, 78-6831, 78-6832, 78-6836.
- *(Viral): 78-6858, 78-6862, 78-6875, 78-6899, 78-6919, 78-6929, 78-6934, 78-6948, 78-6952, 78-6956, 78-6958, 78-6961, 78-6962, 78-6977.
- *(Immun.): 78-6980, 78-7035, 78-7036, 78-7066.
- *(Epid.-Biom.): 78-7152, 78-7153, 78-7154, 78-7161.

EPIDEMIOLOGY AND BIOMETRY

8-7149 **Chronic Dental Infections in the Etiology of Glioblastomas (Meeting Abstract).** (Eng) Storbecker, P. (Akerbyvagen 282, Taby/Stockholm, Sweden). *Neuropathol Exp Neurol* 37(5): 695; 1978. (1 ref)

8-7150 **Proliferative State of Blast Cell Progenitors in Acute Myeloblastic Leukemia (AML).** (Eng) Minden, M. D. (Ontario Cancer Inst., 500 Sherbourne St., Toronto, Ontario M4X 1K9, Canada); Till, J. E.; McCulloch, A. *Blood* 52(3): 592-600; 1978.

Blast cell colony formation by WBC from the peripheral blood of 15 patients with acute myeloblastic leukemia was studied. Colony formation by WBC from these patients was influenced markedly by the concentration of media conditioned by the leukocytes (LCM) in the presence of phytohemagglutinin (PHA). At low PHA LCM concentrations, colonies were compact, immobile, and situated at the bottom of the culture dishes. At higher PHA LCM concentrations, colonies were larger and globular in shape, and they floated in the methylcellulose. The former colony morphology was usually associated with peroxidase-negative cells of blastlike morphology that did not form erythrocyte (E) rosettes. Colonies at high PHA LCM concentrations were composed of small cells (resembling normal lymphocytes) that readily formed E rosettes. The major finding is that progenitors of blast cell colonies are predominantly in a rapid cell cycle, as judged by determinations of the proportion of the cycle in the S phase using tritiated thymidine or hydroxyurea as a DNA synthesis-specific inhibitor. Unless there are very unusual distributions of the cells around the cell cycle, the data indicate that the population of cells detected by this assay is rapidly proliferating. The culture assay for blast cell progenitors detects a minority cell population of morphologically identified blasts; this subpopulation has sufficient proliferative potential for colony formation. The observation that blast progenitors are rapidly proliferating is consistent with the view that they serve to maintain the rest of the blast population. (36 refs.)

78-7151 **The Association between Bronchial Cancer Mortality and the Tobacco Industry in Switzerland between 1970 and 1975.** (Fre) Cardis, F. (Lausanne, Switzerland). *Broncho-Pneumologie* 28(1): 29-30; 1978.

Between 1970 and 1975, bronchial cancer mortality in Swiss women increased 41.5%, vs only 27.8% for Swiss men. During this time, mortality due to other malignant tumors de-

creased 8.91% in men but increased 3.62% in women, and mortality from all malignant tumors increased by 13.34% in men and 4.74% in women. The increase in bronchial cancer mortality in women may have been the result of an increase in the number of women smoking cigarettes. (no refs.)

78-7152 **Nasopharyngeal Cancer in Sabah: Statistics, Observations and Personal Experiences (Meeting Abstract).** (Eng) Kulkarni, M. G. (Queen Elizabeth Hosp., Kota Kinabalu, Sabah, Malaysia). In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 64-65; 1977. (no refs)

78-7153 **Perspective of Cancer in Iran (Meeting Abstract).** (Eng) Habibi, A. (Dept. Pathology, Tehran Univ., 32 Churchill Ave., Tehran, Iran). In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977.* Philippine Cancer Society. (Manila, Philippines): 184 pp.; 85-86; 1977. (no refs)

78-7154 **Celiac Disease. II. Differences in Its Course in Survivors and Deceased Subjects, and Tumor Incidence.** (Cze) Dvorak, M. (Interni oddeleni fakultni polikliniky, Karlovo nam. 32, 121 11, Prague 2, Czechoslovakia); Fric, P.; Lojda, Z. *Cesk Gastroenterol Vyz* 32(4): 217-226; 1978.

Thirty-two adults with celiac disease were followed from 1955 to 1975. Thirteen of the 32 patients died. Cardiopulmonary disease was the cause of death in six cases, suicide in one and malignant tumors in six. The tumors included malignant lymphomas (4), bronchogenic carcinoma (1), and gynecological carcinoma (1). One of the survivors developed jejunal adenocarcinoma, which was successfully resected. The tumors were considered to be a late complication of prolonged celiac disease. Gluten-free diets reduced the cancer risk only partly. (18 refs)

- 78-7155 Lung Cancer Mortality in Uranium Miners (Methodological Aspects) (Letter to Editor).** (Eng) Kunz, E. (Inst. Hygiene and Epidemiology, Centre Radiation Hygiene, 100 42 Prague 10, Czechoslovakia); Sevc, J.; Placek, V. *Health Phys* 35(4): 579-580; 1978.

Data from a 1976 epidemiological study of lung cancer mortality in Czech uranium miners were reevaluated for a group of miners entering employment in 1948-1952. The mortality experience was analyzed in terms of number of cases per unit time at risk, which began with entry into the study group and ended at the cut-off date of the analysis. The results of this recalculation show a coincident picture of response to the increase of cumulative exposure, as in the earlier evaluation. Significant differences between observed and expected values of lung cancer were found in all exposure categories with the exception of the lowest one. The differences in the frequency of additional cases among exposure categories were also significant and, contrary to a study on American uranium miners, confirmed the increase in lung cancer incidence in relation to dose from the first category of exposure upward. The recalculation led to some increase in the estimated number of additional cases in higher exposure categories and a decrease in lower categories compared with the earlier treatment. The distortions caused by the earlier method, however, are not large because the Czech miners entered work and exposure in the same relatively narrow 5-yr period and reached the final exposure levels mostly in the first 5 yr of work. (4 refs)

- 78-7156 Lung Cancer on a Nickel Smelting Island (Meeting Abstract).** (Eng) Reed, D. (California State Health Dept., Berkeley, CA, 94704); Lessard, R.; Maheux, B. *Am J Epidemiol* 108(3): 233; 1978. (no refs)

- 78-7157 Current Vital Statistics: Methods and Interpretation.** (Eng) Adelstein, A. M. (Office Population Censuses and Surveys, London WC2B 6JP, England). *Br Med J* 2(6143): 983-987; 1978.

Statistical analyses of mortality rates in England and Wales show a sharp rise in lung and bronchial cancer deaths among women aged >45 yr and, since 1961, a moderate decline in the rates among men. Differences in smoking habits may account for some of the differences in mortality among men of different social classes. Proportional mortality rates for stomach, intestinal, and rectal cancer are lower than expected among immigrants from India, with Asians having even lower rates than Indian-born Britons. (9 refs)

- 78-7158 Incidence of Bilateral Tumours in a Population-based Series of Breast-Cancer Patients. I. Two Approaches to an Epidemiological Analysis.** (Eng) Pri-

or, P. (Birmingham Regional Cancer Registry, Queen Elizabeth Hosp., Birmingham, England); Waterhouse, J. A. *Br Cancer* 37(4): 620-634; 1978.

The incidence of second primary breast tumors in the contralateral breast was determined among 21,967 primary breast cancer patients registered with the Birmingham Regional Cancer Registry between 1936 and 1964. The results were based on >90,000 woman-years at risk and 399 second primary tumors. When the data were analyzed by a method that included coincidental tumors, the overall risk of a tumor in the contralateral breast was found to be 3.0 times that of a first primary in the general population. The corresponding risks for the three main age groups (at time of diagnosis of first primary tumor) were 5.6 (15-44 yr), 3.7 (45-59 yr), and 1.8 (≥ 60 yr). When coincidental tumors were excluded from the analysis, the relative risk was found to be 2.4 overall and 5.3, 3.0, and 1.0 for the respective age groups. The level of risk was negatively correlated with age at first primary, and the relative risk remained fairly constant over time. (11 refs)

- 78-7159 Carcinoma of the Breast in Women under the Age of 30 (Letter to Editor).** (Eng) Logan, C. J. (Ulster Hosp, Dundonald, Down, England). *Br Med J* 2(6143): 1023; 1978.

Of a group of 213 breast cancer patients, only 3 were <30 yr old, vs 10 of a previously reported group of 195 patients. a high incidence for this age group. Malignancy was discovered in the three women only upon biopsy. None has developed metastases after several years. (1 ref)

- 78-7160 Carcinoma of the Breast in Women under the Age of 30 (Letter to Editor).** (Eng) Jones, V. (Radiotherapy Center, Velindre Hosp., Cardiff, Wales); Slater, A. *Br Med J* 2(6174): 1300; 1978.

Assessment of the age distribution of breast carcinoma patients referred to a radiotherapy center from 1955 to 1977 showed that the age incidence had not changed markedly during this period. The number of women <30 yr old presenting with breast cancer remained at $\leq 1.1\%$. (no refs)

- 78-7161 Tubular Carcinoma of the Breast. A Study of Frequency.** (Eng) Carstens, P. H. (Dept. Pathology, Univ. Louisville Sch. Medicine, P.O. Box 35260, Louisville, KY, 40232). *Am J Clin Pathol* 70(2): 204-210; 1978.

The frequency of tubular carcinoma of the breast among 409 infiltrating breast carcinomas diagnosed during 1974 in the Louisville, Kentucky, area was determined. This carcinoma was histologically defined as an infiltrating breast cancer con-

sisting of tubular structures separated by an abundant amount of hyalinized stroma. A total of 42 tubular carcinoma cases were identified in women aged 26-84 yr. Five patients had bilateral breast cancer, and in three, the contralateral breast also contained tubular carcinoma. Two patients had multifocal tubular carcinoma. Axillary lymph nodes were available for examination with 36/42 breast specimens, and metastatic foci were found in 6. Three histological types of tubular carcinoma were identified: (1) sclerosing type, characterized by a haphazard arrangement of tubules infiltrating into surrounding fat and connective tissue; (2) pure type, consisting of desmoplastic hyalinized stroma surrounding a few tubular structures; and (3) mixed type, in which at least 75% of the tumor was of the tubular variety. Five patients had the first type (none had involved lymph nodes), 9 had the second type (1 with involved nodes), and 28 had the third type (5 with involved nodes). All 42 patients are alive 2 yr after diagnosis; 2 patients have had recurrences. These findings suggest that this type of breast cancer is not as rare as previously reported. (9 refs.)

78-7162 The Hormonal Basis of Breast Cancer: Elevated Plasma Levels of Estrogen, Prolactin, and Progesterone. (Eng) Henderson, B. E. (Dept. Pathology, Univ. Southern California, Los Angeles, CA, 90033); Pike, M. C.; Gerkins, V. R.; Casagrande, J. T. In: *Incidence of Cancer in Humans. Proceedings of the Cold Spring Harbor Conferences on Cell Proliferation*. Hiatt, H. H.; Watson, J. D.; Winsten, J. A., eds. (Cold Spring Harbor: Cold Spring Harbor Laboratory): Origins of Human Cancer. Vol. 4, 602 pp.; 77-86; 1977.

The familial risk of breast cancer was studied in relation to a variety of hormone interactions. There was little difference in the physical characteristics or menstrual histories of 49 daughters of breast cancer patients and 46 daughters of control women. However, a bivariate scatter plot of prolactin and follicular phase total estrogen values showed that levels were clearly elevated in 12/34 daughters of patients vs only 1/30 daughters of controls. Luteal phase hormones were also elevated in the former in one study, but not in a second study, and the levels of plasma estrogen, progesterone, and follicle-stimulating hormone (FSH) were elevated in both studies. Nulliparity and a familial risk of breast cancer were increased in bilateral breast cancer patients diagnosed before age 40, and age at first pregnancy was increased in patients diagnosed between ages 40 and 49. Follicular phase estrone and FSH were significantly elevated in 18 daughters of bilateral breast cancer patients. It is concluded that the familial risk of breast cancer is associated with elevated levels of several hormones, including prolactin, progesterone, and estrogen. However, this 'intrinsic' risk is modified by age at first full-term pregnancy. (41 refs.)

78-7163 Hair Dyes and Human Cancer (Meeting Abstract). (Eng) Hennekens, C. H. (Channing

Lab., Harvard Medical Sch., Boston, MA, 02115); Speizer, F. E.; Rosner, B.; Bain, C.; Belanger, C. *Am J Epidemiol* 108(3): 240-241; 1978. (no refs)

78-7164 Dietary Correlates of Cancer Risk Among Five Ethnic Groups in Hawaii (Meeting Abstract). (Eng) Kolonel, L. N. (Biomedical Sciences D-102, Univ. Hawaii, Honolulu, HI, 96822); Nomura, A. N.; Hankin, J. H. In: *Summary Program and Abstracts of Papers. Proceedings of the 3rd Asian Cancer Conference Held by the Philippine Cancer Society under the Auspices of the Asian Federation of Organizations for Cancer Research and Control in Manila, 26-30 September, 1977*. Philippine Cancer Society. (Manila, Philippines): 184 pp.; 81-82; 1977. (no refs)

78-7165 Seroepidemiologic Assessment of Feline Leukaemia Virus Infection Risk for Man. (Eng) Krakower, J. M. (Lab. RNA Tumor Viruses, NCI, Bethesda, MD, 20014); Aaronson, S. A. *Nature* 273(5662): 463-464; 1978.

The close environmental association of cats and humans and the known infectivity of feline leukemia virus (FeLV) for human cells in tissue culture prompted the examination of human sera for immunological evidence of exposure to this oncogenic agent. Radioimmunoprecipitation assays for antibodies to the major internal protein (p30) and envelope glycoprotein (gp70) of FeLV were performed with sera from 2,074 individuals, including healthy volunteers, subjects with occupational or home exposure to FeLV-infected cats, laboratory workers with occupational exposure to RNA tumor viruses (including FeLV), and patients with various neoplastic and other disorders. No individual had detectable antibodies against either FeLV protein. Sera from patients with hematologic malignancies and from subjects with known or potential home and occupational exposure to FeLV or FeLV-infected animals were further evaluated in competition radioimmunoassays for p30. None of these persons demonstrated detectable FeLV p30 antigen levels. It is concluded that FeLV infection occurs rarely, if at all, in the population studied. (16 refs)

78-7166 Cancer Incidence Amongst Cosmetologists (Meeting Abstract). (Eng) Walrath, J. (Connecticut Cancer Epidemiology Unit, New Haven, CT, 06510). *Am J Epidemiol* 108(3): 234; 1978. (no refs)

78-7167 Leukemia in Olmsted County, Minnesota, 1965-1974. (Eng) Linos, A. (Dept. Medical Statistics and Epidemiology, Mayo Clinic, Rochester, MN);

Kyle, R. A.; Elveback, L. R.; Kurland, L. T. *Mayo Clin Proc* 53(11): 714-718; 1978.

An earlier study on the incidence of leukemia in Olmsted County, Minnesota, was extended to include 1965 through 1974. In this decade, the overall crude annual incidence rate was 9.5/100,000. The male:female ratio was 2.4:1, and the incidence rate in men was significantly higher than that in women, especially in the older age groups. Adjusted for age and sex, the mean annual incidence rate among men (17.3) was significantly higher than rates reported previously from other areas (ranging from 10.7 to 13), and it had increased significantly during 1935-1974. The increase was most prominent in men aged 50 yr and over, and it was detectable only in the subgroup with acute myelocytic leukemia. The observed number of farmers with leukemia (8) was significantly higher than that expected (2.8). (26 refs)

78-7168 **Epidemiology of Brain Tumors in Children.** (Eng) Gold, E. B. (Johns Hopkins Univ., Baltimore, MD). *Diss Abstr Int B* 39(2): 665; 1978. (no refs)

78-7169 **Serological and Epidemiological Studies on the Relationships of Human Leukemia and Bovine Lymphosarcoma (Meeting Abstract).** (Eng) Donham, K. J. (Inst. Agricultural Medicine, Univ. Iowa, Oakdale, IA, 52319); Berg, J. W.; Sawin, R. S. *Am J Epidemiol* 108(3): 241-242; 1978. (no refs)

78-7170 **A Case-Control Study of Intracranial Meningiomas in Women (Meeting Abstract).** (Eng) Preston-Martin, S. (USC Sch. Medicine, Los Angeles, CA, 90033). *Am J Epidemiol* 108(3): 233-234; 1978. (no refs)

78-7171 **Radiation- and Host-Factors in Thyroid Tumors Following Thymus Irradiation (Meeting Abstract).** (Eng) Shore, R. (New York Univ. Medical Center, New York, NY, 10016); Hempelmann, L.; Pasternack, B.; Woodard, E. *Am J Epidemiol* 108(3): 233; 1978. (no refs)

78-7172 **Latency Analysis in an Occupational Study of Stomach Cancer (Meeting Abstract).** (Eng) Blum, S. B. (Univ. North Carolina, Chapel Hill, NC, 27514); Smith, A. H.; Andjelkovich, D.; Tyroler, H. A. *Am J Epidemiol* 108(3): 239; 1978. (no refs)

78-7173 **Mortality Among Laundry and Dry Cleaning Workers (Meeting Abstract).** (Eng) Blair, A.

(Environmental Epidemiology Branch, NCI, Bethesda, MD, 20014); Decoufle, P.; Grauman, D. *Am J Epidemiol* 108(3): 238; 1978. (no refs)

78-7174 **Cancer Mortality and Agricultural Activity: An Association With Cotton Production and Large Farms in the Southeastern United States (Meeting Abstract).** (Eng) Clark, L. (Univ. North Carolina, Chapel Hill, NC, 27514); Shy, C. M.; Portier, K. M.; Most, B. M.; Florin, J. W. *Am J Epidemiol* 108(3): 242; 1978. (no refs)

78-7175 **Cancer Mortality in Selected Louisiana Parishes and Drinking Water Source (Meeting Abstract).** (Eng) Gottlieb, M. (Tulane Univ., New Orleans, LA, 70112). *Am J Epidemiol* 108(3): 232; 1978. (no refs)

78-7176 **Diet and Cancer of the Colon (Meeting Abstract).** (Eng) Graham, S. (SUNY at Buffalo, Buffalo, NY, 14261); Mettlin, C. *Am J Epidemiol* 108(3): 241; 1978. (no refs)

78-7177 **'Adult Type' Cancer in Children and Adolescents.** (Eng) Pratt, C. B. (St. Jude Children's Res. Hosp., P.O. Box 318, Memphis, TN, 38101); Johnson, W. W.; Hustu, H. O. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 961-970; 1977.

The incidence of "adult type" cancer was investigated among 1,009 children with malignant solid neoplasms admitted to a pediatric oncology center between March 1962 and January 1976. Fifty-eight of these patients, aged 3 mo to 19 yr, had tumors usually encountered only in adults. The median age of the 58 patients was 12 yr 9 mo; 22 were Caucasian boys, 18 Caucasian girls, 11 black boys, and 7 black girls. Thirty-seven patients had carcinomas (12 colon, 5 hepatocellular, 3 adrenal, 3 melanoma, 2 pancreas, 2 dysgerminoma, 2 teratocarcinoma, and 1 case each of parotid, thyroid, cholangiohepatocellular, kidney, rectum, seminoma, choriocarcinoma, and adamantinoma). Twenty-one patients had sarcomas (9 liposarcoma, 5 synovial, 2 schwannoma, 2 alveolar soft part, and 1 case each of histiocytoma, angiosarcoma, and mesenchymoma). During the same time period, 960 children were seen with acute lymphocytic leukemia, and 269 with nonlymphocytic leukemia. Among the solid tumor patients, 150 had Hodgkin's disease, 142 neuroblastoma, 127 non-Hodgkin's lymphoma, 99 Wilms' tumor, 99 rhabdomyosarcoma, 74 osteosarcoma, 75 Ewing's sarcoma, and 51 retinoblastoma. There is no indication that more pediatric

patients are being referred with adult-type cancers. However, the admission of nine patients with carcinoma of the bowel within a 15-mo period suggests an increasing frequency of this tumor among pediatric patients. Although almost all tumors that are more commonly seen in adults may occur in children, the occurrence of these rare tumors is of potential importance in regard to the implications of the changing diagnostic categories of patients referred to pediatric oncology centers. (17 refs)

78-7178 Gallbladder Disease and Gallbladder Cancer among American Indians in Tricultural New Mexico. (Eng) Morris, D. L. (Univ. Texas Sch. Public Health, Houston, TX); Buechley, R. W.; Key, C. R.; Morgan, M. V. *Cancer* 42(5): 2472-2477; 1978.

The incidence and mortality rates of gallbladder disease and gallbladder cancer were determined for American Indians (A-I), Spanish-Americans (S-A), and Anglo-Americans (A-A) in New Mexico using the data from the New Mexico Tumor Registry. The results confirmed previous reports of an extremely high incidence of gallbladder disease and cancer in A-I. Cases from the registry were used as numerators, data from the 1970 New Mexico population census as denominators; age adjustment was performed using the US standard population. Male mortality from noncancerous gallbladder disease was 15.3 for A-I and 2.8 for AA; the S-A were intermediate at 4.1. The distribution was similar for women. Computed gallbladder cancer incidence rates indicated a 15-fold increase for A-I vs A-A women, and a similar 6-fold increase in men. Again, S-A had intermediate rates. Standard incidence ratios were calculated for these groups, and indicated the magnitude of the differences between these populations. For A-A men and women, the ratios were 72.7 and 66.7, respectively. Analogous figures for A-I and S-A were 463.6 and 1,004.8, and 200.0 and 500.0. Several risk factors, including parity, obesity, age, cholesterol level, and genetic factors, were explored. Genetic factors appeared to be the most important, with parity a contributing factor. Thus, A-I may have a strong genetic disposition toward gallbladder disease and gallbladder cancer. This hypothesis might also explain the high rates in S-A, a high percentage of which have Indian ancestors. (20 refs)

78-7179 Occupational Dermatoses in the Uranium Industry. (Cze) Sevcova, M. (Kozni oddeleni, ZUNZ UP, 269 00 Pribram-Zdabor, Czechoslovakia). *Prac Lek* 30(4): 138-141; 1978.

The incidence of dermatoses was not increased in uranium industry workers, but the incidence of basal cell carcinoma was 3.6/10,000 (10.2/10,000 for > 5 yr exposure) vs an expected incidence of 0.8/10,000. (13 refs)

78-7180 Effect of a High Beef Diet on Bacterial Flora and Chemical Components of Human Feces: a Summary of Results. (Eng) Hentges, D. J. (Univ. Missouri-Columbia, Columbia, MO); Burton, G. C.; Flynn, M. A.; Franz, J. M.; Gehrke, C. W.; Gerhardt, K. O.; Maier, B. R.; Tsutakawa, R. K.; Wixom, R. L. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976*. International Study Group for the Prevention and Detection of Cancer (New York, NY): Vol. 1(part 1): 1193 pp.; 693-706; 1977.

To examine the effect of a diet high in beef on the bacterial and chemical composition of the feces of humans, 10 male student volunteers were placed on a 4-mo diet series consisting of a control diet, a meatless diet, a high-meat diet, and a second control diet, each of 1-mo duration. The diets were carefully formulated to standardize calories, proteins, fat, carbohydrates, cholesterol, fiber, minerals and vitamins. The fat and fiber content were held constant in all four diets. Total protein was the same in the control and meatless diets but was doubled in the high-meat diet. During the last week of each diet, three fecal samples were collected under oxygen-free conditions from each of the volunteers and analyzed for aerobic and anaerobic bacteria, proximates and minerals, individual amino acids, creatine and creatinine, tryptophan metabolites, polyamines, fatty acids, and neutral and acid steroids. Results showed that there was some increase in the proportion of anaerobic to aerobic bacteria in the feces during the high-meat diet, compared with the meatless diet. There was no increase in the concentration of neutral or acid steroids, and although the high-meat diet contained more cholesterol than did the meatless diet, significantly less cholesterol and fewer cholesterol metabolites were excreted during the high-meat regimen. The results fail to confirm the differences in flora composition and steroid concentration that have previously been observed between populations at high and low risk for colon cancer. (15 refs)

78-7181 Methodologic Issues in Epidemiologic Studies of Endometrial Cancer and Exogenous Estrogen. (Eng) Hulka, B. S. (Dept. Epidemiology and Biostatistics, Sch. Public Health, Univ. North Carolina, Chapel Hill, NC, 27514); Hogue, C. J.; Greenberg, B. G. *Am J Epidemiol* 107(4): 267-276; 1978.

Methodologic issues relevant to epidemiologic studies in general and to studies of the association between endometrial cancer and exogenous estrogen intake are discussed. Problems common to all epidemiologic studies concern pathologic misclassification of disease status; determination of the proper type, source, and amount of data to collect; studies of indications for therapy vs studies of the therapy as the cause of the disease; and differences in medical care in different populations. There are also problems associated with the prolonged follow-up of large numbers of patients and with deter-

mination of the appropriate population source from which to estimate the expected incidence of disease. Cohort studies may be biased by the fact that the two patient groups are not necessarily alike with respect to all possible confounding variables. Case-compeer studies have provided the primary epidemiologic basis for the current concern over increased risk of endometrial cancer associated with exogenous estrogen use. These studies involve matching endometrial cancer patients selected from community hospitals with noncancer or "other cancer" patients at the same hospitals or with ostensibly normal community women. They present problems in determining if the study cases are representative of all cases of the disease and in choosing a suitable comparison group from hospitals, the community, hysterectomized vs nonhysterectomized women, or women undergoing a common diagnostic procedure. The potential risk associated with estrogen use must also be weighed against possible benefits; this topic cannot be addressed by case-compeer studies. (23 refs)

78-7182 Low-Level Radiation and Cancer Deaths. (Eng) Sanders, B. S. (5823 Soledad Road, La Jolla, CA, 92037). *Health Phys* 34(6): 521-538; 1978.

The incidence of cancer among 17,600 men and 3,900 women who worked at the Hanford Works atomic plant in Richland, Washington, and were exposed to low levels of radiation was determined. For exposed men hired between 1944 and 1971, the mean radiation dose increased progressively from 142 millirems (mrem; rem = roentgen-equivalent-man) in 1944 to 3,745 mrem in 1972; the percentage of deaths from cancer was 19.85 for 1944-1959 and 20.27 for 1960-1972. For those hired in 1944 and 1945, the observed dose gradient rose from 142 mrem in 1944 to 3,762 mrem in 1972; the percentage of cancer deaths was 20.67 for 1944-1959 and 19.10 for 1960-1972. From the time of hire to October 1972, the percentage of deaths from cancer was 20.36 for men hired between 1946 and 1971 and 20.05 for those hired in 1944 and 1945. For each year, the mean dose level of those who died from cancer was not significantly different from the mean of those who died from other causes. The mean dose level for the majority of those who died in a specific year was lower than the mean for survivors in the year of death or in the 1 or 2 yr preceding the year of death. Similar findings were observed among the female employees. Although male employees with radiation exposure had a higher proportion of cancer deaths than the general male population in the state of Washington, there was no indication that the radiation was the cause of this difference. Further studies of exposed men vs nonexposed male employees, and out-of-plant controls indicated that the exposed employees had a greater longevity than identified siblings of such employees, nonexposed employees, and matched controls. Thus, all observations to date indicate no adverse health effects among employees that can be attributed to occupational radiation exposure within permissible limits. (17 refs.)

78-7183 Measurement of Environmental Nitrosamines. (Eng) Cohen, J. B. (Natl. Enforcement Investigations Center, Office Enforcement, U.S. Environmental Protection Agency, Denver, CO); Bachman, J. D. *IARC Sci Publ* (19): 357-372; 1978.

The results of a sampling program designed to identify sources of direct emission of nitrosamines (NA's) into the environment and to assess environmental formation of NA's are presented. Samples were taken during summer and winter in the proximity of industrial plants, primarily sources of amines, and analyzed for NA's. There was little indication of the presence or formation of N-nitroso compounds in the air in the vicinity of the sources examined, even when these compounds were observed in plant wastewater. The wastewater samples indicated that N-nitroso compounds are present in chemical, pesticide, and coke plant effluents, some persistently. The in situ sampling did not present any evidence to substantiate secondary production in the vicinity of amine sources. There were no data that would indicate the formation of NA's in the environment, as opposed to their presence as a contaminant. Samples collected along and transverse to major highways showed no contribution of NA's from engine exhaust. The presence of NA's in deionized water (introduced by the mixed-bed resins used for deionization) constitutes a potential, long-term risk for the general population. (9 refs)

78-7184 Dietary Aflatoxins and Human Primary Liver Cancer. (Eng) Peers, F. G. (Tropical Products Inst., 56-62 Gray's Inn Road, London WC1X 8LU, England); Linsell, C. A. *Ann Nutr Aliment* 31(4/5/6): 1005-1018; 1977.

The combined results of two studies, one performed in the Murang'a district of Kenya and the other in Swaziland, show a high degree of positive correlation between calculated ingestion levels of aflatoxin, expressed as nanograms/kilogram/day (x), and the adult incidence rates of hepatocellular carcinoma, expressed as cases/10⁵ adults/year (y). This positive correlation is true for both men and women. Based on an assumed 2-kg/day intake of wet diet at a mean body wt of 70 kg, the calculated relationship for adult women is: $y = 4.14 \log_{10} x - 0.80$. With the added assumption of a daily intake of 2 liters/day of native beer, the regression equation for adult men is: $y = 21.96 \log_{10} x - 11.17$. These regression data are essentially compatible with comparable data recorded by independent workers in Thailand and Mozambique, the latter being the region where the highest rates of liver cancer and the highest aflatoxin ingestion levels have been recorded to date. A highly significant regression line was also calculated using crude liver cancer incidence data and aflatoxin exposure levels from all available studies: $y = 7.60 \log_{10} x - 3.60$. (27 refs)

7185 **Variable Globin Chain Synthesis in Mouse Erythroleukemia Cells.** (Eng) Alter, B. P. (Div. Hematology and Oncology, Children's Hosp. Medical Center, 300 Longwood Ave., Boston, MA, 02115); Goff, S. C. *Leukemia* 52(5): 1047-1057; 1978.

Various mouse erythroleukemia cell (ELC) lines were treated with chemical inducers to determine the proportion of β minor globin (β -MG) synthesis. When cultured with 2% methylsulfoxide, cell lines 745 and 707 from DBA/2 mice synthesized 25%-48% β -MG (75%-52% β -major globin). Induction of the same cells with butyric acid (1 millimole) led to approx 40% β -MG production, but induction with hemin (75 μ M) resulted in > 80% β -MG synthesis. ELC lines derived from other mouse strains show variations in the proportions of globin chains, compared with those from DBA/2 mice, when subjected to similar treatment. The chemical inducing agents did not influence protein stability or the initiation of globin synthesis. Translation of poly(A)-containing cytoplasmic RNA in a wheat germ-free system resulted in the same proportions of β -MG as those seen in intact cells. It is concluded that the induction of globin production depends on the specific interaction of erythroid pre-

cursors with the inducing agents and is due to the regulation of transcription, differential posttranscriptional stabilization of messenger RNA, or the selective proliferation of cells whose specific type of globin production is predetermined. (37 refs)

See also:

- *(Rev.): 78-6634, 78-6637, 78-6641, 78-6642, 78-6644, 78-6646, 78-6648, 78-6649, 78-6651, 78-6652, 78-6653, 78-6659, 78-6662, 78-6664, 78-6667, 78-6680, 78-6684, 78-6685, 78-6696, 78-6697, 78-6698, 78-6699, 78-6700, 78-6701, 78-6703.
- *(Chem.): 78-6709, 78-6710, 78-6712, 78-6720, 78-6765.
- *(Phys.): 78-6803, 78-6804, 78-6829, 78-6833, 78-6838.
- *(Viral): 78-6905, 78-6951, 78-6961.
- *(Immun.): 78-6980, 78-7035, 78-7036, 78-7066.
- *(Path.): 78-7096, 78-7097, 78-7107, 78-7127.

MISCELLANEOUS

- 78-7186 **Compartmental Modeling and Analysis for Carcinogenic Experiments.** (Eng) Whitmore, R. W. (Texas A & M Univ., College Station, TX, 77843). *Diss Abstr Int B* 39(4): 1854; 1978. (no refs)

- 78-7187 **Further Studies on Liver Glyoxalase I and Glyoxalase II. Activity in Mice Bearing Sarcoma 180 and L1210 Leukemia in ascites form.** (Eng) Winter, R. (Inst. Biology and Physiology, Silesian Medical Acad., 41 808 Zabrze, Poland); Piskorska, D.; Jerzykowski, T. *Neoplasma* 25(4): 465-470; 1978.

Glyoxalase I and II activities were determined in the livers and tumor cells of BDF₁ and DBA₂ mice bearing implanted sarcoma 180 and L1210 leukemia in ascites form. The ascites fluids had been injected into the peritoneal cavity. No glyoxalase II activity was detected in the cells of either tumor, but glyoxalase I activity was reduced in comparison with that in normal tissues. Hepatic glyoxalase I and II activities decreased progressively during tumor development. This reduction was as high as 40%-60% of the initial levels. The results corroborate previous findings on the absence or reduction of glyoxalase activity in tumor cells and in the liver during tumor development, and they indicate that a tumor developing ip affects glyoxalase activity in the liver, an organ not directly attacked by the tumor. (23 refs)

- 78-7188 **Impaired Regulation of Activity of Cell Membrane Adenylcyclase and Guanylcyclase as Possible Factor of Carcinogenesis.** (Rus) Khermann, B. (Dept. Internal Medicine, Emil Weil Hosp., Budapest, Hungary). *Vopr Onkol* 24(10): 28-31; 1978.

A discussion is presented of the implications of the recently discovered enzymatic imbalance in the cancer cell. The prevalence of catabolism over anabolism in the tumor cell confirms the hypothesis that the impaired metabolism of the tumor cell can be measured in terms of changes in the balance of activity of adenylcyclase vs activity of guanylcyclase. (14 refs)

- 78-7189 **A Comparison of Polysomal Messenger Ribonucleoprotein Particles from Normal and Neoplastic Rat Liver.** (Eng) Cardelli, J. A. (Dept. Oncology, McArdle Lab. Cancer Res., Medical Sch., Univ. Wisconsin, Madison, WI, 53706); Pitot, H. C. *Cancer Res* 38(7): 2099-2102; 1978.

Free polysomes were isolated from normal and regenerating rat liver and from Morris hepatomas 7777, 7800, 5123C, and

9618A and compared in terms of their messenger ribonucleoprotein particles (MRP). Sucrose gradient analysis ruled out the possibility of any significant messenger RNA (mRNA) degradation in these polysome preparations, since the polysomes were relatively large (8-12 ribosomes/mRNA) and the tissues contained only small amounts of disomes. The EDTA-disrupted polysomes were fractionated on oligodeoxythymidylic acid-cellulose columns. The column-bound polyribadenylic acid-containing MRP were eluted with a formamide buffer, precipitated with ethanol, and subjected to polyacrylamide gel electrophoresis in the presence of sodium dodecyl sulfate. The mRNA-associated proteins from the different tissues were qualitatively similar, but two proteins with a mol wt of 66,000 and the 109,000, which were found as minor proteins in normal liver, appeared on gels as major protein bands when hepatoma MRP were examined. The two proteins were also minor proteins in regenerating and host liver MRP, suggesting that their increase in the hepatomas was not due to tissue growth or to exposure to an altered hormonal environment per se. The possibility exists that the increase in certain MRP is related in some way to the initiation and/or maintenance of the neoplastic state. (15 refs)

- 78-7190 **Analysis of RNA Initiated in Isolated Mouse Myeloma Nuclei Using Purine Nucleoside 5' [γ-S] Triphosphates as Affinity Probes.** (Eng) Smith, M. M. (Dept. Molecular Biology, Univ. Edinburgh, King's Buildings, Mayfield Road, Edinburgh, Scotland EH9 3JR); Reeve, A. E.; Huang, R. C. *Cell* 15(2): 615-626; 1978.

The purine ribonucleoside triphosphate analogs adenosine 5' [γ-S] triphosphate and guanosine 5' [γ-S] triphosphate were used as affinity probes to study RNA chain initiation in isolated nuclei from the mouse myeloma 66.2 cell line. The binding was specific and dependent on inclusion of the sulfur nucleotide analogs in the in vitro synthetic reaction. The affinity-labeled RNA appeared to be initiated in vitro. The nuclei incorporated 1.2-2.4 picomoles of uridine monophosphate/10⁶ nuclei/min, and the rate of synthesis was unaffected by substitution of the nucleotide analogs for the normal nucleotides. The percentage of total RNA synthesized that was incorporated into sequences initiated in vitro was 7.8%-9.6%. The size of the total RNA synthesized ranged from <5S to 45S, and the size of the affinity-labeled sequences ranged from <5S to 28S. The incorporation into RNA initiated in vitro was abolished by α-amanitin in concentrations inhibitory for polymerase II or III. Over 70% of the total incorporation into ribosomal 5S RNA transcripts was into sequences initiated in vitro. This initiation was catalyzed by polymerase III and was specific for guanosine triphosphate as the initiating nucleotide. The RNase T1 fingerprint of the newly initiated 5S RNA indicated that this gene was accurately initiated and elongated in vitro. (52 refs)

78-7191 **Circulating DNA and Oncogenesis.** (Eng) Anker, P. (Div. Oncohaematology, Dept. Medicine, Cantonal Hospital, Geneva, Switzerland); Stroun, M.; Maunula, P. In: *Third International Symposium on the Detection and Prevention of Cancer held by the International Study Group for the Prevention and Detection of Cancer in New York, April 26-May 1, 1976.* International Study Group for the Prevention and Detection of Cancer (New York, NY): 1977. 1(part 1): pp. 1193; 3-16; 1977.

The DNA released from eukaryote cells (reDNA) was characterized, and the possible existence of circulating DNA was explored. The material released from frog auricles and from human blood lymphocytes had the UV absorption characteristics of cellular DNA and was 95% digested by DNase but insensitive to RNase and pronase. The mol wt of the reDNA was not as homogeneous as that of cellular DNA: the latter had one peak over 18S, but reDNA had two main peaks of about 12S and 7S. Separation on CsCl gradients indicated that the AT/GC ratio was the same for both DNA's. The reDNA was not due to a DNA loss from damaged cells, and there was constant renewal of extracellular DNA with each change of medium, suggesting that release is governed by a precise homeostatic mechanism. The DNA was released in a complex containing proteins, among which were enzymes necessary for partial extracellular synthesis of reDNA. Near-neighbor analyses showed that this extracellular synthesis was a true synthesis. The spontaneous transfer of DNA from bacterial to higher organisms that occurs during bacterial infection and its subsequent transcription might be a general biological process that occurs between eukaryote cells in normal conditions, and it suggests the existence of circulating DNA. In view of the fact that DNA extracted from malignant cells can transform normal cells, an oncogenic role of circulating DNA is postulated. It is possible that a circulating DNA may not necessarily be transforming but might be involved in changes associated with malignancy. (29 refs)

78-7192 **In Vitro Cell Proliferation Studies in Hodgkin's Disease.** (Eng) Cowley, J. G. (Natl. Inst. Oncology and Radiology, Havana, Cuba). *Neoplasma* 25(1): 3-90; 1978.

DNA replication in lymph node cells from 30 untreated Hodgkin's disease patients was studied autoradiographically using two different tritiated thymidine pulse-labeling techniques. Nine patients with inflammatory reactive conditions served as controls. The results confirmed previous labeling data, but they showed that DNA synthesis can also be detected among the reticulum cells. Cell proliferation in Hodgkin's and Reed-Sternberg cells appeared to be a characteristic of large mononuclear elements even though some multinuclear elements were detected in DNA synthesis. Large multipolar mitoses were frequent in the Hodgkin's patients, which suggested that the multinuclear cells may have evolved from the mononuclear cells. (17 refs)

78-7193 **Super-sensitive DNA-Repair Test-Liquid for Confirming DNA-Damagicity of Mutagens** (Meeting Abstract). (Eng) Nishioka, H. (Doshisha Univ., Kyoto, Japan). *Mutat Res* 53(2): 242-243; 1978. (1 ref)

78-7194 **Coordinated Regulation of Octopine Degradation and Conjugative Transfer of Ti Plasmids in *Agrobacterium tumefaciens*: Evidence for a Common Regulatory Gene and Separate Operons.** (Eng) Klapwijk, P. M. (Dept. Biochemistry, State Univ. Leiden, 2300 RA Leiden, Netherlands); Scheulderman, T.; Schilperoort, R. A. *J Bacteriol* 136(2): 775-785; 1978.

In an attempt to determine whether octopine utilization is genetically related to conjugative *Agrobacterium tumefaciens* Ti plasmid transfer by the functioning of a shared regulatory mechanism and whether there is a correlation between the genes responsible for octopine metabolism and tumorigenesis, the analog noroctopine was used to isolate *A. tumefaciens* mutants with altered regulatory patterns for Ti plasmid-associated degrading genes. Results of cointegration, conjugation, and octopine oxidase and permease enzyme assays revealed three distinct mutant groups: (1) those with a constitutive level of octopine enzymes and a high degree of spontaneous Ti transfer; (2) those with high enzyme levels but not transfer; and (3) those in which both conjugation and octopine utilization were induced by noroctopine. It was concluded that these results were due to the activity of a common gene. Additionally, when homo-octopine was used to produce mutants defective in octopine metabolism, strain virulence was dependent on the ability to transfer the Ti plasmid. The results indicate that conjugative transfer does not play a significant part in crown gall tumor induction in plants. (38 refs)

78-7195 **Use of the Betabinomial Distribution When Testing for Weak Mutagenic Activity in the Dominant-Lethal Test** (Meeting Abstract). (Eng) Vuataz, L. (Mathematics Section, Nestle Products Technical Assistance Co. Ltd., 1814 La Tour de Peilz, Switzerland); Sotek, J. *Mutat Res* 53(2): 281; 1978. (2 refs)

78-7196 **Mutagenicity Testing by Automated Measurement of Colony Growth and Morphology** (Meeting Abstract). (Eng) Glaser, D. A. (Dept. Molecular Biology and Virus Lab., Univ. California, Berkeley, CA, 94720); Konrad, M. W. *Mutat Res* 53(2): 192; 1978. (no refs)

78-7197 **Mitogens for Murine Embryo Cell Lines.** (Eng) Herschman, H. R. (Sch. Medicine, Univ. California, Los Angeles, CA, 90024); Passovoy, D. S.; Pruss, R. M.; Aharonov, A. *J Supramol Struct* 8(3): 263-268; 1978.

The growth-promoting activities of fetal bovine serum, cortisol, phorbol myristate acetate (PMA), prostaglandin F₂, insulin, epidermal growth factor (EGF), and fibroblast

growth factor (FGF) were evaluated in four murine embryo cell lines (Swiss 3T3, BALB 3T3, M2, and C3H10T1/2). All cell lines showed an increased saturation density when grown in progressively higher serum concentrations, but the M2 cell line demonstrated a somewhat reduced response to serum in comparison with the other three lines. The results were variable: positive results for most mitogens ranged from 50% to 100% increases in cell numbers. PMA was mitogenic for both Swiss 3T3 and M2 cells but not for 10T1/2. Prostaglandin F₂ was mitogenic only for Swiss 3T3 cells, but cortisol was inactive in all four lines. Insulin was stimulatory only for Swiss 3T3 cells, and then only at very high concentrations when compared with other polypeptide mitogens. A 50% increase in saturation density occurred in Swiss 3T3 cells in response to 1 ng/ml of EGF, whereas a similar growth response was elicited only with 1,000-10,000 ng/ml insulin. At 100 ng/ml, EGF resulted in mitogenic responses of 172%, 284%, 127%, and 312% in Swiss 3T3, Mw, C3H10T1/2, and BALB 3T3 cells, respectively. At 10 ng/ml, FGF caused increases of 144%, 121%, 85%, and 122% in the respective cell lines; at 100 ng/ml the corresponding values were 217%, 199%, 122%, and 171%. These experiments show that serum, EGF, and FGF stimulate cell division in all four lines, albeit to varying degrees for the different target cells. (23 refs)

- 78-7198 **Sulfated Mucopolysaccharides from Normal Swiss 3T3 Cell Line and Its Tumorigenic Mutant ST1: Possible Role of Chondroitin Sulfates in Neoplastic Transformation.** (Eng) Dietrich, C. P. (Departamento de Bioquímica e Farmacologia, Escola Paulista de Medicina, Caixa Postal 20372, Universidade de São Paulo, Brazil); Armelin, H. A. *Biochem Biophys Res Commun* 84(3): 794-801; 1978.

The surface sulfate mucopolysaccharides (MPS) of normal Swiss 3T3 cells and a glucocorticoid-sensitive Swiss 3T3 tumor mutant (ST1) were compared. MPS were labeled by incubation with Na₂³⁵SO₄ (10 µCi/ml), cultured, trypsinized, and analysed by agarose gel electrophoresis and enzymatic degradation with appropriate enzymes. Normal 3T3 cells contained primarily chondroitin sulfate B (CSB) (49% in whole cells) and heparin sulfate (HS) (40%), plus small amounts of CSA (11%). The ST1 mutant contained chondroitin sulfate A (CSA) (43%), HS (44%), and small amounts of CSB (13%). After the addition of hydrocortisone to ST1 cell cultures, the cells showed some of the properties of normal cells (cell density, anchorage, and serum dependence), but their sulfated MPS composition remained constant. Degradation of normal and ST1 cells with chondroitinases B and ABC showed that they contained only 4-sulfated disaccharides. Normal Swiss 3T3 and 3T6 cells, both of which undergo spontaneous transformation with time in culture, showed increased CSA and decreased CSB after exhibiting phenotypic characteristics of transformation, including loss of contact inhibition. These results support the hypothesis that glucuronic acid-containing CSA A plays a role in the

stimulation of cell division in neoplastic and embryonic tissues. (19 refs)

- 78-7199 **Establishment and Characterization of a New Human Urinary Bladder Carcinoma Cell Line (PS-1).** (Eng) Sanford, E. J. (Div. Urology, Dept. Surgery, Milton S. Hershey Medical Center, Pennsylvania State Univ., Coll. Medicine, Hershey, PA, 17033); Geder, L.; Dagen, E.; Laychock, A. M.; Ladda, R.; Rohner, T. J. *Invest Urol* 16(3): 246-252; 1978.

An epithelioid cell line (PS-1) was established from a human transitional cell carcinoma of the urinary bladder. Sc injection of the epithelioid cells into weanling athymic nude mice induced solid tumors histologically similar to the original tumor. A cell line was also established from a tumor induced in an athymic nude mouse (PS-1, T-1). Both cell lines exhibited essentially identical growth characteristics and formed monolayer growth of epithelioid cells in culture. Electron microscope studies confirmed the epithelioid morphology. No fibroblastoid elements were observed. Chromosomal analysis revealed heteroploidy with persistent marker chromosomes. All cells contained a Y chromosome. The presence of a tumor-specific antigen(s) in PS-1 cells was suggested by radioimmunoassay and cytotoxicity assays with peripheral allogeneic lymphocytes from other transitional cell cancer patients. Serum from urinary bladder cancer patients reacted with nuclear antigens of the established cells. (22 refs)

- 78-7200 **Serum Ribonucleases in Pancreatic Cancer: Correlation to Tumor Histology.** (Eng) Funakoshi, T. (Third Dept. Internal Medicine, Faculty Medicine, Kyushu Univ., 1-1 Maeda-shi 3-chome, Higashi-ku, Fukuoka City 812, Japan); Wakasugi, H.; Kimura, T.; Matsumoto, M.; Ibayashi, H. *Gastroenterol Jpn* 13(5): 390-394; 1978.

Serum ribonuclease (RNase) levels of 18 normal persons, 18 patients with chronic pancreatitis, and 26 patients with pancreatic cancer were determined using polycytidylic acid as substrate. The RNase levels of patients with four histologic types of pancreatic cancer were compared, ie, in patients with ductal cell adenocarcinoma, and anaplastic cell, acinar cell, and islet cell carcinoma. The normal level of serum RNase was 92 ± 6.3 units (U). In chronic pancreatitis and pancreatic cancer it was 118 ± 10.4 and 249 ± 16.8 U, respectively. Patients with ductal cell adenocarcinoma had the highest serum RNase levels. The phosphocellulose chromatography patterns of the serum RNase and pancreatic cancer tumor extract were compared with those of serum RNase and pancreatic extracts from normal subjects. Control serum and control pancreatic extract both exhibited three main elution patterns, but pancreatic cancer serum and tumor extract exhibited a different elution pattern and high activity. It is suggested that enzyme elevations may be due to the synthesis of a new species of RNase and an increased enzyme synthesis by the proliferating tumor cells. If this theory proves correct, early diagnosis of pancreatic cancer may be accomplished by analyzing serum RNase isozymes. (13 refs)

Author Index

- aronson, S. A., 78-6885, 78-6905
 78-7165
 blashi, D., 78-6946
 bo-Darub, J. M., 78-7090
 ccinni, R., 78-6940
 dami, H. O., 78-7106
 dams, W. G., 78-6715
 dee, R. R., 78-6834
 delstein, A. M., 78-7157
 haronov, A., 78-7197
 ida, M., 78-7112
 l-Ghazzouli, I. K., 78-7024
 lexander, A., 78-7060
 lexander, J., 78-6957
 llen, G. P., 78-6898
 llison, A. C., 78-7002
 lroy, J., 78-7141
 lter, B. P., 78-7185
 ltman, A., 78-6988
 mato, S., 78-7145
 nderer, F. A., 78-7048
 ndererson, D., 78-6715
 ndererson, D. M., 78-6903
 nderesson-Anvret, M., 78-6632
 ndjelkovich, D., 78-7172
 nker, P., 78-7191
 ntoniades, H., 78-6928
 ntonioli, D., 78-6789
 piou, F., 78-6794
 rcement, L. J., 78-6887
 rlett, C. F., 78-6809
 rlinghaus, R. B., 78-6675, 78-6887
 rmelin, H. A., 78-7198
 rrmstrong, G., 78-6946
 rthes, F. G., 78-6700
 rya, S. K., 78-6976
 sano, Y., 78-7074
 saoka, K., 78-7034
 skonas, B. A., 78-6687
 sselbergs, F. A., 78-6865
 tassi, G., 78-7020
 ufes, A. H., 78-7035
 ustin, F. C., 78-6909
 xell, T., 78-6720
 xelrod, H. R., 78-6932
 ach, F. H., 78-6890
 achman, J. D., 78-7183
 laird, S., 78-6677, 78-6875
 akaev, V. V., 78-6923
 akemeier, R., 78-7072
 aldwin, R. W., 78-6692, 78-7042
 78-7064
 allintine, E. J., 78-7163
 annasch, P., 78-6707, 78-6787
 aron, J. M., 78-6876
 arrett, J. C., 78-7076
 arritault, D., 78-7060
 arrow, G., 78-7056
 artorelli, A., 78-6940
 arzilai, R., 78-7018
 aserga, R. L., 78-6686, 78-6703
 asilico, C., 78-6686
 attaglinio, A., 78-7049
 Baum, J. W., 78-6665
 Bauman, A., 78-7072
 Beard, P., 78-6907
 Becker, F. F., 78-6706
 Becker, S., 78-6883, 78-7010
 Beemon, K., 78-6852
 Beers, Y., 78-6710
 Beissner, E. I., 78-7061
 Beladi, I., 78-6970
 Belikova, H., 78-7129
 Bendig, M. M., 78-6903
 Beniashvili, D. S., 78-6748
 Benitz, K. F., 78-6734
 Bennett, P. H., 78-7163
 Bensch, K. G., 78-6837
 Bentvelzen, P., 78-6866
 Berg, J. W., 78-7169
 Berg, P., 78-6922
 Berg-Schlosser, V., 78-7113
 Bergholz, C., 78-6936
 Beri, D. P., 78-6951
 Bernhard, H. P., 78-6686
 Besson, A., 78-7119
 Betz, E. H., 78-6979
 Beug, H., 78-6676, 78-6936
 Bey, E., 78-6957
 Bhandari, J. C., 78-6714
 Biancardi, C., 78-6940
 Biempica, L., 78-7128
 Biozzi, G., 78-7057
 Birmingham, D., 78-6798
 Bishop, J. M., 78-6840, 78-6843
 78-6848
 Bishun, N., 78-6618
 Bissell, M. J., 78-6849
 Bitter-Suermann, D., 78-7002
 Black, A. K., 78-6811
 Black, H. S., 78-6806
 Blair, A., 78-7173
 Blattner, W. A., 78-6934
 Bleeker, G. M., 78-7096
 Bloemendal, H., 78-6865
 Bloemers, H. P., 78-6678
 Blog, F. B., 78-6777
 Bloustein, P. A., 78-6650
 Blum, S. B., 78-7172
 Boccara, M., 78-6902
 Boger, E., 78-7059
 Bohm, W., 78-7133
 Boling, M. E., 78-6759
 Bolognesi, D., 78-6892
 Bolognesi, D. P., 78-6884
 Bonnet-Eymard, J., 78-6713
 Boone, C. W., 78-6909
 Boot, L., 78-6868
 Booth, C. G., 78-7051
 Borek, C., 78-6796
 Borell, S., 78-6915
 Bornkamm, A. G., 78-6951
 Bornstein, P., 78-6906
 Bosch, V., 78-6853
 Bossens, M., 78-7065
 Botchan, M. R., 78-6929
 Boulesteix, J., 78-6956
 Boyd, A. L., 78-6937
 Boyd, G. W., 78-6682
 Boyd, H., 78-6766, 78-6767
 Boyse, E. A., 78-6871
 Bozzetti, F., 78-6728
 Brand, G., 78-6633
 Brandt, K. H., 78-7124
 Brash, D. E., 78-6619
 Breadmore, S. E., 78-6822
 Bremner, J. M., 78-6831
 Bridges, B., 78-6704
 Bridges, B. A., 78-6809
 Brinkhof, J., 78-6866
 Brisou, B., 78-6958
 Britton, W. J., 78-6795
 Broker, T. R., 78-6964
 Bromley, P. A., 78-6854
 Bronkhorst, F. B., 78-7124
 Bronshtein, M. E., 78-7146
 Brookes, P., 78-6610, 78-6622
 Brown, G. B., 78-6718
 Brown, G. E., 78-6806
 Brownlie, B. E., 78-6831
 Bruce, W. R., 78-6792
 Brugge, J. S., 78-6856
 Brune, K., 78-6719
 Bryan, G. T., 78-7140
 Bryans, J. T., 78-6898
 Budinger, J. M., 78-6718
 Buechley, R. W., 78-7178
 Burchett, S., 78-6716
 Burger, P. M., 78-6808
 Burgert, E. O., 78-6815
 Burke, L., 78-6789
 Burleson, G. R., 78-6855
 Burnet, F. M., 78-6609
 Burtis, C. A., 78-6743
 Burton, G. C., 78-7180
 Bustan, H., 78-6686
 Butenko, Z. A., 78-6886
 Butler, W. H., 78-6723, 78-6758
 Buxbaum, J., 78-7060
 Byard, J. L., 78-6731
 Cadoff, B. C., 78-6709
 Cairns, J., 78-6610
 Calafat, J., 78-6974
 Calnek, B. W., 78-6860, 78-7050
 Calvelli, T. A., 78-6990
 Camacho, A., 78-6944
 Camp, R., 78-6811
 Campbell, E. W., 78-6797
 Campbell, T. C., 78-6738
 Canning, M. V., 78-6788
 Capellaro, D., 78-7026
 Capparelli, N., 78-7059
 Carbone, P. P., 78-7102
 Cardelli, J. A., 78-7189
 Cardis, F., 78-7151
 Carey, P. R., 78-7039
 Carlens, E., 78-6657
 Carnow, B. W., 78-6701
 Carol, W., 78-7133

- Carstens, H. B., 78-7161
Carter, W. A., 78-6976
Casagrande, J. T., 78-7162
Cassingena, R., 78-6933
Castro, J. E., 78-6998
Cerna, H., 78-6845, 78-6853
Chalvet, H., 78-7057
Chan, J. T., 78-6806
Chan, K. H., 78-6949
Chandra, P., 78-6669
Chang, C. C., 78-6606
Chang, M. J., 78-6757
Chapler, F. K., 78-7095
Charman, H. P., 78-6679, 78-6873
Chaudhari, A., 78-6747
Cheever, A. W., 78-7140
Chesebro, B., 78-6879
Childs, D. S., 78-6815
Chirigos, M. A., 78-6993
Chiu, N. H., 78-6921
Chouroulinkov, I., 78-6626
Chow, D. A., 78-7012, 78-7013
Chow, L. T., 78-6964, 78-6966
Churg, A., 78-6837
Citrin, D. L., 78-7102
Clapp, N. K., 78-6743
Clark, L., 78-7174
Clayson, D. B., 78-6604
Cochrane, J. C., 78-7061
Coggin, J. H., 78-6984
Cohen, B. H., 78-6658
Cohen, B. L., 78-6738
Cohen, J. B., 78-7183
Cohen, M. C., 78-6690
Cohen, S., 78-6690
Cohen, S. M., 78-7141
Cohn, M., 78-7110
Colcher, D., 78-6864
Collins, J. L., 78-7110
Colombeau, M. C., 78-6956
Colyer, S. P., 78-6743
Cooper, G. M., 78-6847
Cooper, J., 78-6822
Cornain, S., 78-7053
Cornelius, E. A., 78-7028
Costa, A., 78-6728
Coulston, F., 78-6734
Courtney, R. J., 78-6943
Coutinho, W. G., 78-6863
Cowley, J. G., 78-7192
Crawley, A., 78-7079
Croce, C. M., 78-6610, 78-6919
Croissant, O., 78-7036
Cruickshank, W. J., 78-6997
Cueto, C., 78-6771
Cunningham, C., 78-7134
Cuzin, F., 78-6924, 78-6926
D'Alisa, R. M., 78-6916
Dagen, J. E., 78-7199
Darlix, J. L., 78-6854
DasGupta, U. B., 78-6941
Daub, G. H., 78-6783
Dauge, M. C., 78-6992
David, B., 78-7137
David, C., 78-7014
Davidson, K. A., 78-6743
Davidson, N., 78-6674
Davis, A. R., 78-6973
Dawson, P. J., 78-7067
Day, N. E., 78-6951
Dazord, L., 78-7014
de Agustin de Agustin, P. P., 78-7135
De Brabander, M. J., 78-7081
De Jong, J. G., 78-7096
De Lumley, L., 78-6956
de Moerloose, P., 78-7032
de Palo, G., 78-7136
de-The, G., 78-6951
de Tribolet, N., 78-7032
Dean, A. G., 78-6951
Deardorff, E. R., 78-6709
Deaven, L. L., 78-6797
Decleve, A., 78-6874
Decoufle, P., 78-7173
Dec, R. R., 78-6982
Deichmann, W. B., 78-6659
Deinhardt, F., 78-6936
Delacretaz, F., 78-7119
Delamore, I. W., 78-7083
Della Torre, G., 78-7136
Delleman, J. W., 78-7096
Delrez, M., 78-6979
Denman, A. M., 78-6952
Dent, P. B., 78-6985, 78-7037
DePamphilis, M. L., 78-6913
Derelanko, M. J., 78-7069
Derge, J. G., 78-6937
Deslattes, R., 78-6710
Detjen, B. M., 78-6971
Devare, S. G., 78-6877
Devillebichot, C., 78-7084
Devoe, J., 78-6710
Diacumakos, E. G., 78-6912
DiAugustine, R. P., 78-6793
Dickhaus, S., 78-6839
Dietrich, C. P., 78-7198
Dimant, I. N., 78-6748
Dinowitz, M., 78-7024
Doerfler, W., 78-6610, 78-6670
Doig, D., 78-6879
Doll, D. C., 78-7145
Dominok, G. W., 78-7147
Donham, K. J., 78-7169
Dorko, W. D., 78-6709
Dorough, H. W., 78-6754
Dostalova, V., 78-6857
Dougherty, C. M., 78-7134
Douglas, J. F., 78-6771
Doury, J. C., 78-6958
Doyle, D., 78-7077
Drasil, V., 78-6828
Drohan, W., 78-6864
Du Vivier, A., 78-6753
Duello, T. M., 78-7095
Duke, R. E., 78-6890
Duksin, D., 78-6906
Dular, U., 78-7012, 78-7013
Dumont, P., 78-7020
Dunn, A. R., 78-6962, 78-6966
Duprez, V., 78-7027
Durdik, J. M., 78-7009, 78-7022
Dutta, S., 78-6747
Dvorak, M., 78-7154
Easton, J., 78-6946
Ebener, U., 78-6669
Eck, H., 78-7113
Edelman, G. M., 78-6846
Edwards, H. E., 78-6855
Eguchi, S., 78-6832
Eichmann, K. P., 78-6687
Eisenberg, R., 78-7040
Eisenman, R. N., 78-6880
Eisenstark, A., 78-6805
Ekelman, K. B., 78-6785
El Abed, A., 78-6778
Elder, J., 78-6892
Elder, J. H., 78-6872, 78-6884
Elliot, J. M., 78-6773
Elveback, L. R., 78-7167
Emmelot, P., 78-7016
Emminger, E., 78-6839
Emura, I., 78-6948
Endrich, B., 78-7078
Engvall, E., 78-7062
Enomoto, K., 78-6744
Enomoto, M., 78-6717
Erikson, R. L., 78-6856
Evans, R., 78-7051
Evenson, K., 78-6710
Factor, S. M., 78-7128
Faith, R. E., 78-7011
Falk, H. L., 78-6756
Faller, D. V., 78-6882
Fallica, L., 78-7092
Fansler, B. S., 78-6812
Farson, D., 78-6849
Farwell, D. C., 78-6742
Fauve, R. M., 78-6992, 78-7001
Favre, M., 78-7036
Fayez, J. A., 78-7138
Fears, T. R., 78-6804
Fechner, R. E., 78-6645
Fellows, C., 78-6946
Feltkamp, C. A., 78-7016
Fernandes, G., 78-6987
Ferrara, R., 78-6940
Feunteun, J., 78-6930
Fialkow, P. J., 78-6610
Fieldsteel, A. H., 78-7067
Fincham, N., 78-6811
Finley, J., 78-6980
Finn, O., 78-7029
Fischinger, P., 78-6892
Fischinger, P. J., 78-6884
Flad, H. D., 78-7000
Fleischer, R. L., 78-6818
Fleissner, E., 78-6896
Fletcher, K. A., 78-6788
Florin, J. W., 78-7174
Flynn, M. A., 78-7180
Fokina, G. I., 78-6891
Folk, W. R., 78-6903
Forsby, N., 78-6632
Fournet, J., 78-6713
Fowler, A. K., 78-7017
Francis, G. E., 78-7026
Franke, W. W., 78-6687
Frankel, A. E., 78-6884
Franz, J. M., 78-7180
Fraumeni, J. F., 78-6934

- Fredrickson, T. N., 78-7069
 Freedman, V. H., 78-6990
 Frei, U., 78-7103
 Frenkel, N., 78-6673
 Freund, S., 78-6710
 Fric, P., 78-7154
 Friedman, A., 78-6825
 Friedman, E., 78-7094
 Friedman, H., 78-7003, 78-7008
 Friend, D. G., 78-6709, 78-6709
 Friend, W. H., 78-6737
 Friis, R. R., 78-6851
 Froberg, H., 78-6635
 Fu, S. M., 78-7073
 Fujimoto, N., 78-7071
 Fujimoto, Y., 78-6801
 Fujita, D. J., 78-6843
 Fukuda, Y., 78-6801
 Fukuhara, S., 78-7080
 Fukuoka, M., 78-6717
 Funakoshi, A., 78-7200
 Gabizon, A., 78-7047
 Gagnon, H. J., 78-7049
 Gaillard, S., 78-6956
 Galante, E., 78-6728
 Gallagher, N. D., 78-6795
 Gallimore, P. H., 78-6962
 Gallo, R. C., 78-6672, 78-6888
 Garber, E. A., 78-6918
 Gardes, M., 78-6930
 Gardner, M. B., 78-6674, 78-6679
 78-6873, 78-6895
 Gardner, R. L., 78-6686, 78-6705
 Gaudray, P., 78-6924
 Gautsch, J. W., 78-6872, 78-6892
 Gayathri, A. K., 78-6965
 Geder, L., 78-7068, 78-7199
 Gehlert, P., 78-6628
 Gehrke, C. W., 78-7180
 Gemsa, D., 78-6782
 Genin, R., 78-7084
 Jenkins, G., 78-7035
 Gentry, G. A., 78-6898
 Georgiev, G. P., 78-6923
 Gerhardt, K. O., 78-7180
 Gerkins, V., 78-6649
 Gerkins, V. R., 78-7162
 Gerodetti, M. H., 78-6925
 Gershey, E. L., 78-6912, 78-6916
 Geser, A., 78-6951
 Ghosh, A., 78-6793
 Giannelli, F., 78-6614
 Gibbs, A. R., 78-7111
 Gibbs, G. W., 78-6838
 Gielen, J. E., 78-6623
 Gierak, T., 78-7099
 Gilden, R. V., 78-6674, 78-6679
 Gillette, J. R., 78-6607
 Giovanni Gervasi, P., 78-6772
 Given, D., 78-6954
 Glaser, D. A., 78-7196
 Glaser, M., 78-7005, 78-7044
 Glaser, R., 78-6950
 Glinski, W., 78-7036
 Gluckman, J., 78-7061
 Goff, S. C., 78-7185
 Gogusev, J., 78-7101
 Goh, K., 78-7072, 78-7120
 Gohari, K., 78-7075
 Gold, E. B., 78-7168
 Goldfarb, S., 78-6729
 Goldrosen, M. H., 78-6985
 Goldstone, A. H., 78-7026
 Golub, E. S., 78-7021, 78-7022
 Golubeva, I. V., 78-7146
 Gomard, E., 78-7027
 Good, R. A., 78-6987
 Goodman, M., 78-7079
 Gordon, A. S., 78-7069
 Gossner, W., 78-6830
 Gotchy, R. L., 78-6829
 Gothoskar, S. V., 78-6727
 Gottlieb, M., 78-7175
 Gould, I., 78-7065
 Graessmann, A., 78-6927, 78-6935
 Graessmann, M., 78-6927, 78-6935
 Graevskaia, N. A., 78-6891
 Graf, T., 78-6676, 78-6936
 Graham, S., 78-7176
 Granner, D. K., 78-7095
 Grauman, D., 78-7173
 Greaves, M., 78-6681
 Greaves, M. F., 78-7026
 Greaves, M. W., 78-6811
 Green, H., 78-6610, 78-6686
 Greenberg, B. G., 78-7181
 Gregor, O., 78-7116
 Gregory, R. O., 78-6708
 Greifer, B., 78-6709
 Griffin, T. B., 78-6734
 Grimm, W., 78-6782
 Grofova, M., 78-6897
 Gross, J. F., 78-7078
 Gross, R. L., 78-6637
 Grover, P. L., 78-6772
 Grussendorf, E. I., 78-7088
 Guanella, N., 78-7032
 Guarneri, B., 78-7092
 Guhl, E., 78-6935
 Guilmette, R. A., 78-6825
 Guirgis, H., 78-7066
 Guirgis, H. A., 78-7118
 Guntaka, R. V., 78-6842
 Gupta, S., 78-6987
 Gurkalo, V. K., 78-6732
 Haaijman, J. J., 78-6866
 Habibi, A., 78-7153
 Hacker, H. J., 78-6787
 Hackett, P. B., 78-6840
 Hadding, U., 78-7002
 Hadley, J. G., 78-6834
 Haglund, B., 78-6765
 Hahnefeld, H., 78-6900
 Hakim, A. A., 78-7004
 Hall, E. J., 78-6668
 Hall, J. G., 78-7097
 Halmi, N. S., 78-7095
 Ham, G. H., 78-6822
 Hamilton, B. P., 78-7144
 Hampar, B., 78-6937
 Hanawalt, P. C., 78-6620
 Hancock, C. D., 78-6621
 Hancock, R. L., 78-6621
 Hankin, J. H., 78-7164
 Hannan, M. A., 78-6769
 Hannecart-Pokorni, E., 78-7065
 Hansteen, I. L., 78-6712
 Haot, J., 78-6979
 Harari, J., 78-6980
 Harbison, R. D., 78-6615
 Hard, G. C., 78-6739
 Hardy, W. D., 78-6896
 Harnden, D. G., 78-6610
 Harris, C. E., 78-7095
 Harris, R., 78-7066
 Harris, R. E., 78-7118
 Hart, R. W., 78-6619
 Hartig, F., 78-6799
 Hartman, P. E., 78-6733
 Haslam, S. N., 78-6774
 Hatie, C., 78-6849
 Hayata, Y., 78-7112
 Hayden, J. A., 78-6824
 Hazelrigg, D. E., 78-7089
 Heatfield, B. M., 78-6751
 Heberling, R. L., 78-6679
 Hebold, G., 78-6799
 Hecker, E., 78-6719
 Heddle, J. A., 78-6792
 Heimler, A., 78-7094
 Heinrich, K., 78-6709
 Heise, J. M., 78-6850
 Heldaas, S. S., 78-6712
 Hellman, A., 78-7017
 Hempelmann, L., 78-7171
 Henderson, B. E., 78-6679, 78-7162
 Henderson, E. E., 78-6947
 Henle, W., 78-6951
 Henney, C. S., 78-7009
 Henry, B. E., 78-6950
 Henry, J. M., 78-6756
 Hensby, C. N., 78-6811
 Hentges, D. J., 78-7180
 Herschman, H. R., 78-7197
 Hevin, M. B., 78-6992, 78-7001
 Hewer, A., 78-6772
 Hewetson, J., 78-6950
 Higgins, E. S., 78-6737
 Hilgers, J., 78-7016
 Hilken, J. G., 78-6974
 Hilkmann, H. A., 78-7016
 Hill, A., 78-6649
 Hillestad, L., 78-6712
 Hinuma, Y., 78-6948, 78-6955
 Hirsch, I., 78-6953
 Hlozanek, I., 78-6857
 Ho, H. C., 78-7015
 Hoagland, H. C., 78-6815
 Hodge, L. D., 78-6965
 Hoffken, K., 78-7042, 78-7064
 Hofmann, H., 78-7121
 Hogue, C. J., 78-7181
 Holbrook, K., 78-6906
 Holden, K. M., 78-6724
 Holford, T. R., 78-7107
 Hollstein, M. C., 78-6762
 Hong, C. B., 78-6714
 Hopkins, N., 78-6882

- Hopper, D. G., 78-6995
Horie, A., 78-6832
Horiuchi, M., 78-7034
Hostein, J., 78-6713
Hougen, J., 78-6710
Houwert, G. J., 78-7124
Howard, C. J., 78-6710
Howard, J. C., 78-6687, 78-6688
Howell, N., 78-6914
Howley, P. M., 78-6905
Huang, R. C., 78-7190
Huang, T. C., 78-7140
Hudson, P. M., 78-6756
Huebner, R. J., 78-6679, 78-6895
Hug, O., 78-6830
Huget, R. P., 78-7000
Hughes, E., 78-6710
Hughes, E. E., 78-6709
Hulka, B. S., 78-7181
Hull, P. J., 78-7083
Hultsch, E., 78-7045, 78-7046
Hunter, T., 78-6852
Hurley, J. N., 78-7073
Hurley, L. H., 78-6769
Hustu, H. O., 78-7177
Hutchison, G. B., 78-6641
Hwang, S. P., 78-6929
IARC Working Group, 78-6617
78-6656
Ibayashi, H., 78-7200
Ichiba, M., 78-7112
Idoine, J., 78-6773
Ihle, J. N., 78-6826, 78-6874, 78-6884
78-6892
Ikawa, Y., 78-6841
Ikeh, V., 78-6979
Ilmensee, K., 78-6686
Imai, N., 78-6867
Imai, S., 78-6867
Imblum, R. L., 78-6781
Inagaki, T., 78-6801
Inbar, M., 78-7018
Ino, M., 78-6741
Intaglietta, M., 78-7078
Inui, S., 78-6749, 78-7115
Inuyama, Y., 78-7034
Isaka, T., 78-6841
Ishiguro, T., 78-6939
Ishihara, N., 78-6948
Ishii, S., 78-7148
Israel, E., 78-7006
Israel, M. A., 78-6905
Ivanov, A. P., 78-6891
Jablonska, S., 78-7036
Jacobson, A. P., 78-6833
Jaenisch, R., 78-6686
Jaffe, R., 78-7082
Jalanko, H., 78-7062
James, A., 78-6901
Jamjoom, G. A., 78-6675, 78-6887
78-6889
Janossy, G., 78-6681, 78-7026
Jansson, B., 78-6696
Jarrett, W. F., 78-6899
Jarzabek-Chorzelska, M., 78-7036
Jeannet, M., 78-7032
Jenkins, J. J., 78-7145
Jensen, E. V., 78-6603
Jensen, F. C., 78-6872
Jerzykowski, T., 78-7187
Job, L., 78-6976
Johns, R. H., 78-6709
Johnson, A. H., 78-6708
Johnson, B. E., 78-6666
Johnson, W. W., 78-7177
Joishy, S. K., 78-7120
Jonas, H. S., 78-7138
Jones, B., 78-7145
Jones, G., 78-6723
Jones, P. D., 78-6998
Jones, R. N., 78-6634
Jones, V., 78-7160
Jullien, P., 78-6794
Jung, E. G., 78-7121
Juraskova, V., 78-6828
Jurgelski, W., 78-6756
Justrabo, E., 78-7084
Kaden, D. A., 78-6761
Kagan, E., 78-7061
Kaledin, V. I., 78-6996
Kalyanaraman, V. S., 78-6888
Kambour, M., 78-6980
Kamen, R. I., 78-6904
Kanarek, L., 78-7020
Kanbour, A. I., 78-7137
Kandzari, S., 78-7145
Kaplan, G., 78-6994
Kaplan, H., 78-6686
Kaplan, H. S., 78-6874, 78-7029
Kapp, J. F., 78-6981
Karande, A. A., 78-7025
Kark, A. E., 78-7035
Karpas, A., 78-6975
Karshin, W. L., 78-6887
Kawabata, T., 78-6741
Kawamura, T., 78-7052
Kay, K., 78-6663
Kayes, S. G., 78-7085
Keller, R., 78-7055, 78-7066
Kelloff, G. J., 78-7024
Kelly, F., 78-6902
Kelsey, J. L., 78-7107
Kende, M., 78-7024
Kerbel, R. S., 78-7043
Kerdelhue, B., 78-6778
Kerman, R., 78-7033
Keshet, I., 78-6890
Kessler, I. I., 78-6945
Kettmann, R., 78-6864
Key, C. R., 78-7178
Khan, M. N., 78-6793
Kherrmann, B., 78-7188
Khodosova, I. A., 78-6764
Khoury, G., 78-6905
Khrustalev, S. A., 78-6790
Kieff, E., 78-6954
Kieff, E. D., 78-6673
Kiessling, R., 78-7010
Kimball, R. F., 78-6759
Kimmerle, G., 78-6839
Kimura, T., 78-7200
Kinch, R., 78-7125
King, H. W., 78-6622
Kinoshita, K., 78-6763
Kiselev, F. L., 78-6859
Kiseleva, L. L., 78-6891
Kishimoto, H., 78-7086
Kisliakova, N. D., 78-6775
Kitamura, K., 78-7074
Kizer, D. E., 78-6716
Kjellman, B., 78-6765
Klapwijk, P. M., 78-7194
Klein, E., 78-6883, 78-7053
Klein, G., 78-6632, 78-7010
Kleiner, H., 78-6816
Klement, V., 78-6679
Klempa, L., 78-7103
Klietmann, W., 78-6915
Klimpel, G., 78-7009
Knowler, W. C., 78-7163
Knowles, B. B., 78-6982
Knudson, A. G., 78-6602, 78-6610
Kobayashi, H., 78-7052
Kobuchi, K., 78-6780
Koch, K. M., 78-7103
Kodama, T., 78-7052
Koide, Y., 78-6986
Kojima, K., 78-7115
Kolonel, L. N., 78-7164
Kolosov, A. E., 78-7132
Kon, S. H., 78-6654
Konaka, C., 78-7112
Konishi, Y., 78-7115
Konrad, M. W., 78-7196
Koprowski, H., 78-6686
Kordac, V., 78-7123
Kornhuber, B., 78-6669
Koropatnick, J., 78-6608
Kouri, R. E., 78-6781
Krakower, J. M., 78-7165
Krause, H., 78-7088
Kravchenko, L. V., 78-6629
Krementz, E. T., 78-7037
Kress, M., 78-6930
Kretschmar, R. M., 78-6646
Kroes, R., 78-6662
Krueger, J. G., 78-6978
Kuba, P., 78-7061
Kucherlapati, R., 78-6929
Kucherova, T. L., 78-6775
Kukava, G. G., 78-6893
Kulkarni, M. G., 78-7152
Kulkarni, M. S., 78-6807
Kulpa, C. F., 78-6855
Kumar, K., 78-7039
Kunkel, H. G., 78-7073
Kunschner, A., 78-7137
Kuntz, R. E., 78-7140
Kunz, E., 78-7155
Kup, W., 78-6768
Kupperman, G. L., 78-6816
Kurland, L. T., 78-7167
Kuroyanagi, M., 78-6717
Kurunov, Y. N., 78-6996
Kuzela, S., 78-6897
Kwok, R., 78-6986
Kwong, P. C., 78-7037

Kyle, R. A., 78-7167
 Lachance, P. A., 78-6655
 Lacinova, J., 78-6857
 Ladda, R., 78-7199
 Lafferty, W. J., 78-6710
 Lai, M. H., 78-6885
 Lalley, P., 78-6934
 Lam, K. M., 78-7023
 Lam, P., 78-6608
 Lambert, B., 78-7091
 Landsberg, L., 78-7144
 Langlois, Y., 78-6778
 Larson, B. A., 78-6640
 Lasfargues, E. Y., 78-6863
 Lasne, C., 78-6626
 Latt, S. A., 78-6810
 Laube, H., 78-6669
 Laufer, I., 78-7130
 Laumen, G., 78-6767
 Lavialle, C., 78-6933
 Law, L. W., 78-7044
 Laychock, A. M., 78-7068, 78-7199
 Layer, H., 78-6710
 Le Garrec, Y., 78-7014
 Ledbetter, J. A., 78-6880
 Lee, C. C., 78-6714
 Lee, H., 78-7072
 Lee, J. A., 78-6804
 Lee, N., 78-7010
 Leffell, M. S., 78-6984
 Lehman, J. M., 78-6910
 Lennox, E. S., 78-6687, 78-6694
 Leong, S. P., 78-7037
 Lerner, R. A., 78-6872, 78-6884
 Lessard, R., 78-7156
 Levine, A. J., 78-6918, 78-6932
 Levine, R. J., 78-7144
 Levinson, A. D., 78-6848
 Levintow, L., 78-6840, 78-6848
 Levitt, R., 78-7030
 Levy, J. P., 78-7027
 Lew, E. A., 78-6698
 Lewison, E. F., 78-6644
 Li, A. P., 78-7109
 Liao, S. K., 78-7037
 Liber, H. L., 78-6761
 Liddell, F. D., 78-6838
 Lieberman, M., 78-6874, 78-7029
 Liebermann, H., 78-6900
 Lilly, F., 78-6686
 Lindenbaum, A., 78-6825
 Linna, T. J., 78-7023
 Linnoila, I., 78-6793
 Linos, A., 78-7167
 Linsell, C. A., 78-7184
 Lisiewicz, J., 78-7099
 Little, J. M., 78-6795
 Liubimov, A. V., 78-6908
 LiVolsi, V. A., 78-7107
 Lizonova, A., 78-6897
 Lo, K. W., 78-6960
 Lo, L., 78-6608
 Lo, L. W., 78-6802
 LoBue, J., 78-7069
 Loewengart, G., 78-6725
 Logan, C. J., 78-7159
 Lohs, K., 78-6628
 Lojda, Z., 78-7154
 Lough, J., 78-7125
 Louie, S., 78-6695
 Louis, J., 78-7104
 Loveday, K. S., 78-6810
 Lubiniecki, A. S., 78-6934
 Lucas, J., 78-6971
 Lui, S. W., 78-6949
 Lung, M. L., 78-6874
 Lupulescu, A., 78-6798
 Luster, M. I., 78-7011
 Lutz, W. K., 78-6711
 Lutzner, M. A., 78-6684
 Luz, A., 78-6830
 Lynch, H., 78-7066
 Lynch, H. T., 78-7118
 Lynch, J., 78-7066
 Lynch, N. R., 78-6691
 MacDonald, W. E., 78-6659
 Mach, O., 78-6845, 78-6857
 Mack, T., 78-6649
 Mackie, J. A., 78-6960
 Mackie, R., 78-7090
 MacLennan, I. C., 78-7019
 Macnab, G., 78-6957
 MacNicoll, A. D., 78-6772
 Maekawa, I., 78-7071
 Magee, P. N., 78-6601, 78-6610
 78-6616
 Magni, E., 78-7105
 Mahaffey, J. A., 78-6820, 78-6821
 Maheux, B., 78-7156
 Maier, B. R., 78-7180
 Maisin, J. R., 78-6979
 Makari, J. G., 78-6861
 Maki, A., 78-6710
 Malahy, M. A., 78-6696
 Malinvaud, G., 78-6956
 Maloney, K., 78-7066
 Maness, P. F., 78-6846
 Marcus, S. L., 78-6870
 Mareel, M. M., 78-7081
 Marinenko, R. J., 78-6709
 Marks, R., 78-6813
 Marsden, H. S., 78-6938
 Marshack, M., 78-7033
 Martin-Achard, A., 78-7032
 Martin, D. H., 78-6786
 Martin, M. A., 78-6905
 Martin, P., 78-6928
 Martinelli, D., 78-6816
 Martinez Tello, F. J., 78-7135
 Maryak, J. M., 78-6894
 Maskens, A. P., 78-7131
 Mason, K. A., 78-7056
 Masterson, B. J., 78-7138
 Mathews, M. B., 78-6963, 78-6966
 Matienko, N. A., 78-6996
 Matoska, J., 78-6897
 Matsumoto, M., 78-7200
 Matsumura, Y., 78-6832
 Mattelin, G., 78-6979
 Maurice, P., 78-7191
 Mazurek, C., 78-7057
 McAfee, D. A., 78-6766, 78-6767
 McAllister, R. M., 78-6674, 78-6679
 McBride, W. H., 78-7056
 McClain, D. A., 78-6846
 McCusker, J., 78-7067
 McDonald, A. D., 78-6838
 McDonald, J. C., 78-6838
 McDonald, J. J., 78-6718
 McDougall, J. K., 78-6929, 78-6962
 McEwan, R., 78-6826
 McGowan, J. J., 78-6898
 McGrath, C. M., 78-6869
 McGuire, L. J., 78-6750
 McGuire, W. L., 78-6686
 McKenna, G., 78-7073
 McLachlan, J. A., 78-7011
 McLaughlin, P. J., 78-7042
 Meagher, R. C., 78-7069
 Medvedeva, N. D., 78-6764
 Meeks, S., 78-6709
 Menzel, D. B., 78-6724
 Merkushev, G. N., 78-6817
 Mermet, M. A., 78-7104
 Mertl, L., 78-7116
 Mettlin, C., 78-7176
 Meyers, P., 78-6850
 Meyskens, F. L., 78-6672
 Michalides, R., 78-6868
 Michalides, R. J., 78-6865
 Michiels, R., 78-7084
 Michnovicz, J. J., 78-6783
 Mickal, A., 78-7134
 Mihailovich, N., 78-7122
 Milas, L., 78-7056
 Miller, M., 78-7097, 78-7163
 Miller, R., 78-6668
 Miller, R. W., 78-6610, 78-6697
 Minami, A., 78-7148
 Mistry, K. B., 78-6823
 Mitchison, N. A., 78-6687
 Mitra, J., 78-7069
 Mitrani, E., 78-7087
 Miyaji, T., 78-7127
 Miyamoto, K., 78-6955
 Mohr, U., 78-6839
 Monier, R., 78-6930
 Monjardino, J., 78-6901
 Monti, G., 78-7105
 Moolten, F., 78-7059
 Moore, J. A., 78-7140
 Moore, R. D., 78-7041
 Moore, V. E., 78-7042, 78-7064
 Moorthy, A. S., 78-7069
 Moreno Koch, M. C., 78-7031
 Moretti, E., 78-6825
 Morgan, M. V., 78-7178
 Morimoto, J., 78-6867
 Mornstad, H., 78-6720
 Morohoshi, T., 78-6717
 Morris, A. G., 78-6933
 Morris, D. L., 78-7178
 Morris, J. M., 78-6651
 Morton, J. I., 78-7041
 Most, B. M., 78-7174
 Mottershead, R. P., 78-6809
 Mouriquand, C., 78-7104
 Mouriquand, J., 78-7104
 Moyer, M. P., 78-6925

- Moyer, R. C., 78-6925, 78-6978
 Mrzena, V., 78-7129
 Mueller, C., 78-6927, 78-6935
 Muir, C. S., 78-6610
 Mukada, T., 78-7114
 Mulder, A. M., 78-7022
 Muller, W. A., 78-6830
 Mulvihill, J. J., 78-6934
 Musashi, M., 78-7071
 Myklebust, R., 78-6709
 Nagington, J., 78-6975
 Nair, M., 78-6987
 Najarian, T., 78-6667
 Nakajima, Y., 78-7034
 Nakatani, K., 78-7115
 Namyslowski, G., 78-7099
 Nandi, S., 78-6643, 78-6791
 Naso, R. B., 78-6887
 Natori, S., 78-6717
 Navas Palacios, J. J., 78-7135
 Nayak, D. P., 78-6973
 Nedospasov, S. A., 78-6923
 Nelson, F. R., 78-6786
 Neri, A., 78-6967
 Nettesheim, P., 78-6786, 78-6793
 Newberne, P. M., 78-6637
 Newbold, R., 78-6622
 Newcombe, H. B., 78-6605
 Ng, M. H., 78-6949, 78-7015
 Ng, W. S., 78-7015
 Nicolaides, N. J., 78-7126
 Nicolson, G. L., 78-6687
 Nicolson, M. O., 78-6674
 Nielsen, H. O., 78-7117
 Nikolin, V. P., 78-6996
 Nilsson, K., 78-6897
 Nishijima, T., 78-6735
 Nishioka, H., 78-7193
 Noguchi, H., 78-7071
 Nomura, A. N., 78-7164
 Nomura, Y., 78-6832
 Norkin, L. C., 78-6968
 Normann, S. J., 78-6991
 Nosotti, A., 78-6770
 Novi, A. M., 78-6722
 Nowak, R., 78-7100
 Nowinski, R. C., 78-6876, 78-6880
 Nunoue, N., 78-7074
 Nusse, R., 78-6865, 78-6868
 O'Callaghan, D. J., 78-6950
 O'Toole, C., 78-6967
 Oaks, J. A., 78-7085
 Obalek, S., 78-7036
 Obata, Y., 78-6871
 Oboshi, S., 78-6948
 Oda, K., 78-6969
 Ogawa, K., 78-6744
 Ogorodnikova, L. S., 78-6775
 Ogura, H., 78-6897
 Ohnishi, Y., 78-6948
 Ohno, S., 78-6687
 Ohshima, H., 78-6741
 Ohtaki, S., 78-6911
 Oikawa, T., 78-6909
 Okabe, H., 78-6710
 Olin, R. G., 78-7150
 Onoe, T., 78-6744
 Opitz, H. G., 78-7000
 Oppermann, H., 78-6840, 78-6848
 Orth, G., 78-7036
 Osborne, M. R., 78-6622
 Oshiro, Y., 78-6746
 Otu, A. A., 78-6999
 Owerbach, D., 78-7077
 Ozaki, Y., 78-6939
 Ozu, R., 78-7034
 Paddle, G. M., 78-6715
 Pal, B. C., 78-6786
 Palladino, M. A., 78-7054
 Pan, S. H., 78-7003
 Panet, A., 78-6851
 Pang, R. H., 78-6972
 Pang, R. T., 78-6721
 Panh, M., 78-6713
 Papatestas, A. E., 78-7035
 Paraskevas, F., 78-7012, 78-7013
 Park, S. M., 78-6783
 Parkes, H. G., 78-6702
 Parris, D. S., 78-6943
 Passovoy, D. S., 78-7197
 Pasternack, B., 78-7171
 Patek, P. Q., 78-7110
 Paterson, M. C., 78-6610, 78-6612
 78-6613
 Paucha, E., 78-6920
 Pauley, R. J., 78-6862
 Pauli, B. U., 78-7141
 Pavlidis, N. A., 78-6993
 Pazmino, N. H., 78-6826
 Pearse, A. D., 78-6813
 Pearson, G., 78-6946
 Pederson, J., 78-7102
 Peers, F. G., 78-7184
 Pegg, A. E., 78-6625, 78-6755
 Peiker, G., 78-7133
 Penn, I., 78-6736
 Perbal, B., 78-6926
 Perdue, S. W., 78-6759
 Perez Cuadrado, S., 78-7031
 Perou, J., 78-6805
 Persson, H., 78-6963
 Pescia, G., 78-7119
 Peters, L. J., 78-7056
 Petitpas, E., 78-7104
 Peto, R., 78-6668
 Pettersson, U., 78-6963
 Phelps, D. J., 78-7039
 Phillips, L. A., 78-6972
 Phillips-Quagliata, J. M., 78-7038
 Piastucka, B., 78-7099
 Pienta, R. J., 78-6726
 Pierre, R. V., 78-7030
 Pike, M. C., 78-7162
 Pilch, J., 78-7099
 Pilotti, S., 78-7136
 Pimm, M. V., 78-6995
 Pinsky, S. D., 78-6752
 Piskorska, D., 78-7187
 Pitot, H. C., 78-7189
 Pitts, J. D., 78-7090
 Placek, V., 78-7155
 Plato, P. A., 78-6833
 Pledger, W. J., 78-6928
 Pletsityi, K. D., 78-6630
 Pliss, G. B., 78-6732
 Plotz, P., 78-7040
 Plummer, N. A., 78-6811
 Pogodina, V. V., 78-6891
 Polianskaia, A. M., 78-6689
 Pollack, R. E., 78-6661, 78-6686
 Ponomareva, T. V., 78-6817
 Popovic, M., 78-6897
 Portier, K. M., 78-7174
 Pott, F., 78-6664
 Pound, A. W., 78-6750
 Prasad, U., 78-7101
 Prat, J., 78-6652
 Pratt, C. B., 78-7177
 Preston-Martin, S., 78-7170
 Preston, V. G., 78-6938
 Price Evans, D. A., 78-6788
 Price, J. A., 78-6968
 Price, M. R., 78-7042, 78-7063
 78-7064
 Prior, P., 78-7158
 Privitera, L., 78-6940
 Prough, R. A., 78-6781
 Pruss, R. M., 78-7197
 Pugh-Humphreys, R. G., 78-6997
 Pugh, T. D., 78-6729
 Purchase, I. F., 78-6715
 Pusztai, R., 78-6970
 Putzke, H. P., 78-7100
 Qualtiere, L. F., 78-6850
 Raabe, O. G., 78-6818
 Racevskis, J., 78-6870
 Radford, H. E., 78-6710
 Radonovich, M. F., 78-6921
 Rafferty, K. A., 78-6680
 Rajewsky, K., 78-6687
 Rajewsky, M. F., 78-6610
 Ranadive, K. J., 78-6727
 Randall, C. C., 78-6898
 Rankin, L., 78-7066
 Rao, K. V., 78-7122
 Rapp, F., 78-6683, 78-7068
 Rapp, H. J., 78-6988
 Rapp, U. R., 78-6877
 Raschke, W., 78-6875
 Rasheed, S., 78-6679, 78-6873, 78-6895
 Rassoulzadegan, M., 78-6924, 78-6926
 Rattle, H., 78-6772
 Rawson, R. W., 78-6631
 Raz, A., 78-7018
 Reddy, M. M., 78-7120
 Reed, C. D., 78-7017
 Reed, D., 78-7156
 Reeve, A. E., 78-7190
 Reinhold, C. E., 78-6730
 Reinhold, H. S., 78-7078
 Reuber, M. D., 78-6660
 Rhoads, J. E., 78-6960
 Rhoads, K., 78-6819
 Ricciardi-Castagnoli, P., 78-7029
 Richardson, C. A., 78-6997
 Richardson, C. R., 78-6715
 Richardson, E. P., 78-7098
 Rickard, R. W., 78-6754

- Rifle, G., 78-7084
Rigby, P. W., 78-6922
Rilke, F., 78-7136
Rimsten, A., 78-7106
Ringborg, U., 78-7091
Ringer, D. P., 78-6716
Ritts, G. D., 78-6850
Robboy, S. J., 78-6652
Robertson, D. M., 78-7043
Robinson, C. C., 78-6910
Robison, S. H., 78-6812
Roche, J., 78-6713
Roche, J. C., 78-6958
Rogers, A. E., 78-6637, 78-6638
Rogers, J., 78-6798
Rogers, K. S., 78-6737
Rohner, T. J., 78-7199
Roizman, B., 78-6673
Rolle, W., 78-6628
Roman, J. M., 78-7021
Rongey, R. W., 78-6679
Ropcke, G., 78-6868
Rose, D. P., 78-7102
Rosen, S., 78-6789
Rosenblum, I., 78-6734
Rosenthal, A. D., 78-7094
Rosenthal, P. N., 78-6874
Ross, R., 78-6649
Rothman, K. J., 78-6641
Rottger, P., 78-7103
Roush, T. W., 78-6784
Rowley, J. D., 78-7080
Ruben, R. L., 78-6680
Rubin, J. J., 78-6766, 78-6767
Rudali, G., 78-6794
Ruoslahti, E., 78-7062
Russell, J. J., 78-6825
Russell, R. J., 78-6999
Russo, I. H., 78-6776
Russo, J., 78-6776
Rzesa, G., 78-7036
Sadler, T. E., 78-6998
Saegesser, F., 78-7119
Sager, R., 78-6914
Sahasrabudhe, M. B., 78-7025
Sainerova, H., 78-6858
Sairenji, T., 78-6948, 78-6955
Saito, M., 78-6717
Sakae, K., 78-7114
Sakai, S., 78-6730
Sakurada, K., 78-7071
Sakurai, M., 78-6961
Salden, M. H., 78-6865
Salomon, J. C., 78-6691
Salzman, N. P., 78-6921
Sambrook, J., 78-6966
Samil'chuk, E. I., 78-6893
San, R. H., 78-6608
Sandberg, A. A., 78-7070
Sanders, B. S., 78-7182
Sanders, C. L., 78-6819, 78-6820
78-6821, 78-6834
Sanefuji, H., 78-6751
Sanford, E. J., 78-7068, 78-7199
Sano, K., 78-7074
Santoli, D., 78-6982, 78-6983
Sarkar, N. H., 78-6870
Sarma, D. S., 78-6746
Sarngadharan, M. G., 78-6888
Sartwell, P. E., 78-6700
Sasaki, T., 78-7148
Sasano, N., 78-7108, 78-7114
Sato, E., 78-7114
Sato, G. H., 78-6686
Sawin, R. S., 78-7169
Schafer, W., 78-6936
Schaffer, P. A., 78-6943
Scharadt, M., 78-6991
Scharff, M. D., 78-7073
Schat, K. A., 78-6860, 78-7050
Scheide, E., 78-6709
Schenck, P., 78-6710
Scher, C. D., 78-6928
Scheulderman, T., 78-7194
Schilperoort, R. A., 78-7194
Schlatter, C., 78-6711
Schlechte, J., 78-7095
Schlom, J., 78-6864
Schmidt, W. P., 78-6709
Schneiderman, M. A., 78-6804
Schnyder, U. W., 78-7121
Schorlemmer, H. U., 78-7002
Schultz, R. D., 78-7050
Schultz, R. M., 78-6993
Schwartz, R. S., 78-6695
Schwarz, H., 78-6936
Schwarz, R., 78-6849
Scolnick, E. M., 78-6894
Scott, R. S., 78-6831
Scotto, J., 78-6804
Scribner, J. D., 78-6611
Scully, R. E., 78-7098
Seal, R. M., 78-7111
Segawa, K., 78-6969
Seibert, G. B., 78-6696
Seidman, M. M., 78-6918
Seiler, R., 78-7032
Seljelid, R., 78-6994
Setlow, R. B., 78-6624
Sevc, J., 78-7155
Sevcova, M., 78-7179
Sheffield, J. B., 78-6863
Shenk, T., 78-6917
Sherman, B. M., 78-7095
Shih, T. Y., 78-6894
Shimaoka, K., 78-6827
Shimizu, M., 78-7074
Shimizu, N., 78-6929
Shimizu, S., 78-6679
Shimojo, H., 78-6969
Shiraishi, T., 78-7071
Shiratori, T., 78-7115
Shiroki, K., 78-6969
Shore, R., 78-7171
Shows, T. B., 78-7077
Shy, C. M., 78-7174
Siegel, B. V., 78-7041
Siekert, R. G., 78-7030
Sikora, K., 78-6694
Silagi, S., 78-6990
Silverman, L., 78-6847
Silverstein, S. C., 78-6990
Simons, J. W., 78-6808
Sims, P., 78-6772
Sincock, A. M., 78-6836
Sisillo, E., 78-6940
Sito, A. F., 78-6891
Skopek, T. R., 78-6761
Slater, A., 78-7160
Slovin, S. F., 78-7058
Small, M., 78-7047
Smirnova, I. A., 78-6886
Smith, A. E., 78-6904, 78-6920
Smith, A. H., 78-7172
Smith, H., 78-6822, 78-6952, 78-6977
Smith, J. E., 78-6796
Smith, M. M., 78-7190
Smith, N., 78-6618
Smith, P. G., 78-6951
Smith, R. E., 78-6844
Smith, S. W., 78-6870
Smithson, W. A., 78-6815
Smolec, J. M., 78-6959
Snyder, H. W., 78-6896
Snyder, R., 78-6959
Socher, S. H., 78-6862
Sokal, J. E., 78-6827
Solheim, O. P., 78-6835
Solisch, P., 78-6900
Solomon, A., 78-7061
Solomon, D., 78-6905
Sommers, S. C., 78-6647
Sonta, S. I., 78-7070
Soo, W. J., 78-6849
Sorkin, E., 78-6991
Sorof, S., 78-6745
Sotek, J., 78-7195
Soule, H. D., 78-6869
Sovova, V., 78-6845, 78-6853
Spahr, P. F., 78-6854
Spear, P. G., 78-6673, 78-6944
Specter, S., 78-7003
Spencer, F., 78-7051
Spilman, S. D., 78-6731
Spira, G., 78-7018
Sprecher-Goldberger, S., 78-7065
Staab, H. J., 78-7048
Stadel, B. V., 78-6648, 78-7107
Stageman, D. L., 78-6816
Steel, L. K., 78-6669
Steeves, R., 78-6878
Steffen, D., 78-6881
Steg, A., 78-7139
Steigleder, G. K., 78-6693
Stein-Werblowsky, R., 78-7142
78-7143
Steinau, U., 78-7103
Steinbaugh, P. J., 78-6856
Steinheider, G., 78-6878
Stemmermann, G. N., 78-7108
Stephenson, J. R., 78-6877, 78-7024
Stich, H. F., 78-6608, 78-6802
Stiffel, C., 78-7057
Stiles, C. D., 78-6928
Stockert, E., 78-6871
Stohrer, G., 78-6718
Stone, S. C., 78-7093
Stortebecker, P., 78-7149

- Stow, N. D., 78-6938
 Stradling, G. N., 78-6822
 Streit, G. E., 78-6710
 Stringer, J. R., 78-6942
 Stroun, M., 78-7191
 Stylos, W. A., 78-6993
 Su, R. T., 78-6913
 Suarez, H. G., 78-6933
 Subramanian, K. N., 78-6917
 Suchankova, A., 78-6953
 Sullivan, J., 78-6825
 Summers, J., 78-6959
 Summers, W. C., 78-6941
 Sundstrom, B., 78-6720
 Suss, R., 78-6611
 Sutherland, C. M., 78-7037
 Suzuki, T., 78-6948
 Svoboda, J., 78-6858
 Swanbeck, G., 78-7091
 Sweet, D. L., 78-7080
 Sweger, D., 78-6710
 Sykes, J. A., 78-6967
 Szabo, G., 78-6777
 Szmunn, W., 78-6685
 Tabbari, B., 78-7105
 Takahashi, G., 78-6740, 78-6763
 Takahashi, S., 78-7115
 Takasaki, K., 78-7034
 Takasugi, M., 78-6986
 Takeichi, N., 78-6909
 Takemoto, K. K., 78-6905
 Tal, J., 78-6843
 Talal, N., 78-7082
 Tan, Q. H., 78-6760
 Tapp, E., 78-6779
 Tarodi, B., 78-6970
 Tarro, G., 78-6940
 Tateno, H., 78-7108
 Tatosyan, A. G., 78-6859
 Taylor, J. K., 78-6709
 Teglbjaerg, P. S., 78-7117
 Tejmar, L., 78-6915
 Teller, M. N., 78-6718
 Terao, K., 78-6800
 Tew, K. D., 78-6752
 Thiel, H. J., 78-6936
 Thiis-Evensen, E., 78-6712
 Thilly, W. G., 78-6761
 Thiry, L., 78-7065
 Thomas, J. K., 78-6855
 Thorbeck, R. R., 78-6669
 Thorbecke, G. J., 78-7054
 Thoren, M. M., 78-6921
 Thorgeirsson, S. S., 78-6730
 Thyssen, J., 78-6839
 Tickle, C., 78-7079
 Tierney, B., 78-6772
 Timanovskaia, V. V., 78-6893
 Timbury, M. C., 78-6938
 Todaro, G. J., 78-6877
 Toeroek, D., 78-6833
 Tokunaga, M., 78-7114
 Tomasi, T., 78-7066
 Tome Paule, C., 78-7031
 Tomingas, R., 78-6664
 Tomoyasu, S., 78-7074
 Tonascia, J. A., 78-6700
 Tonelli, Q. J., 78-6745
 Topping, D. C., 78-6786
 Toth, P., 78-7022
 Toto, P., 78-7033
 Toujas, L., 78-7014
 Tracey, D., 78-7009
 Trainin, N., 78-7047
 Trinchieri, G., 78-6982, 78-6983
 Tronick, S. R., 78-6885
 Trosko, J. E., 78-6606
 Trump, B. F., 78-6751
 Ts'o, P. O., 78-7076
 Tsou, K. C., 78-6960
 Tsubura, Y., 78-6867
 Tsuruoka, N., 78-7074
 Tsutakawa, R. K., 78-7180
 Tukei, P. M., 78-6951
 Tupchong, L., 78-7019
 Turner, K. J., 78-6691
 Tutel'ian, V. A., 78-6629
 Tuyns, A. J., 78-6653
 Twardzik, D., 78-6946
 Twardzik, D. R., 78-7017
 Twiddy, R. R., 78-7043
 Tyndall, R. L., 78-6743
 Tyroler, H. A., 78-7172
 Ubayama, Y., 78-7148
 Ullrich, A., 78-6840
 Umeda, M., 78-6717
 Urbach, F., 78-6753
 Usadel, K. H., 78-7103
 Usui, M., 78-7148
 Vaidya, A. B., 78-6863
 Valens, M., 78-6778
 Van Blitterswijk, W. J., 78-7016
 Van Deemter, L., 78-6868
 Van Der Sluis, P. J., 78-7016
 Van Duuren, B. L., 78-6636, 78-6725
 Van Eldik, L. J., 78-6844
 Van Scott, E. J., 78-6753
 Van Zaane, D., 78-6678
 VanderLaan, W. P., 78-6640
 Vandervoort, J., 78-7066
 Vandervoort, J. P., 78-7118
 VanGilder, J., 78-7095
 Variakojis, D., 78-7080
 Varmus, H. E., 78-6843, 78-6848
 Vass, W. C., 78-6894
 Vayuegula, B. S., 78-6973
 Vecerina, S., 78-7119
 Verevkin, G. A., 78-6775
 Verloes, R., 78-7020
 Verma, I. M., 78-6885
 Vesselinovitch, S. D., 78-7122
 Viola, M., 78-6770
 Virtanen, I., 78-7062
 Vives, C., 78-6794
 Volegov, A. I., 78-7007
 Vonderheid, E. C., 78-6753
 Vonka, V., 78-6953
 Vuataz, L., 78-7195
 Vyas, B. N., 78-6823
 Wahl, H., 78-7147
 Wainberg, M. A., 78-7006
 Wakabayashi, K., 78-6801
 Wakasugi, H., 78-7200
 Walker, M. C., 78-7038
 Walker, M. D., 78-7098
 Walrath, J., 78-7166
 Walsh, C., 78-6772
 Warin, A. P., 78-6811
 Warnock, M. L., 78-6837
 Warren, S. T., 78-6606
 Watanabe, H., 78-6814
 Watanabe, M., 78-7003
 Waterhouse, J. A., 78-7158
 Watson, A., 78-6890
 Watson, A. A., 78-6718
 Webb, R. B., 78-6805
 Webster, I., 78-7061
 Wechsler, W., 78-6639
 Weeks, M. O., 78-6894
 Weil, G., 78-6851
 Weil, R., 78-6671
 Weill, H., 78-6634
 Weinberg, R. A., 78-6881
 Weinstein, I. B., 78-6784
 Weinstein, R. S., 78-7141
 Weisburger, E. K., 78-6773
 Weislow, O. S., 78-7017
 Weiss, N. S., 78-6642
 Weiss, R. A., 78-6686
 Weiss, S. R., 78-6840
 Weissman, I. L., 78-6677, 78-6686
 Welch, W. R., 78-6652
 Weller, N. K., 78-6931
 Wells, J. S., 78-6710
 Wells, S. A., 78-6708
 Werko, L., 78-6699
 Wheeler, T., 78-6904
 White, C., 78-7107
 White, F. H., 78-7075
 White, R. G., 78-6999
 White, W. L., 78-7030
 Whitmore, R. W., 78-7186
 Wigle, D. T., 78-6803
 Wilkie, N. M., 78-6938
 Wilkinson, P. C., 78-6999
 Williams, D., 78-6618
 Williams, D. R., 78-6894
 Williams, E. H., 78-6951
 Williams, K., 78-6906
 Williams, S. J., 78-6724
 Wilson, A. R., 78-6997
 Wilson, G. L., 78-6989
 Wimmer, E., 78-6971
 Winn, R. M., 78-7128
 Winnacker, E. L., 78-6610
 Winston, J. M., 78-6714
 Winter, R., 78-7187
 Winters, W. D., 78-6967
 Wirth, P. J., 78-6730
 Witting, C., 78-7045, 78-7046
 Wixom, R. L., 78-7180
 Wolf, G., 78-6915
 Woll, J. E., 78-7072
 Wong, L. C., 78-6714
 Woodard, E., 78-7171
 Woolley, P. V., 78-6752
 Wreghitt, T. G., 78-6975

Yagi, T., 78-7148
Yamaguchi, M., 78-6871
Yamasaki, H., 78-6784
Yamawaki, S., 78-7148
Yasuhira, K., 78-6763
Yerganian, G., 78-7049
Yielding, K. L., 78-6807
Yonkum, G. H., 78-6805
Yokoyama, S., 78-7074
Yoshida, M., 78-6841

Yoshida, Y., 78-6744
Yoshihira, K., 78-6717
Young, B. G., 78-6989
Young, N. M., 78-7039
Yuasa, Y., 78-6969
Yurieva, N. P., 78-7146
Zarling, D. A., 78-6890
Zavadova, H., 78-6953
Zedeck, M. S., 78-6760
Zegel, H. G., 78-7130

Zenz, C., 78-6627
Zielinski, W. L., 78-6746
Zucker, R. M., 78-6980
Zuckerman, E. E., 78-6896
Zuidema, P. J., 78-7124
Zulman, J., 78-7082
Zvilichovsky, G., 78-6718

BRITISH MUSEUM LIBRARY

Subject Index

normalities

- Carcinogen, Chemical
 - Teratogenic Effect, Review, 78-6616
- Methane, Chlorodifluoro-
 - Epidemiology, Child, 78-6765
- Radiation
 - Teratogenic Effect, Review, 78-6616
- Xylene
 - Epidemiology, Child, 78-6765

etamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-

- Cell Transformation, Neoplastic
 - Organ Culture, Mammas, 78-6745

DNA

- Liver, Rat, 78-6716

Fluorescence

- Liver, Rat, 78-6716

RNA, Ribosomal

- Liver, Rat, 78-6716

Xeroderma Pigmentosum

- DNA Repair, Review, 78-6624

etamide, *N*-Fluoren-2-yl-

- Cell Transformation, Neoplastic
 - Organ Culture, Mammas, 78-6745

Chromosome Aberrations

- Carcinogenic Potential, Review, 78-6618

Hepatoma

- Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6729
- Hyperplasia, 78-6729
- Precancerous Conditions, 78-6729

Liver

- Metabolism, Rat, 78-6731
- Mutagenic Activation, 78-6730

Mutagenic Activity

- Micronucleus/Sperm Abnormality Assays, 78-6792

Norepinephrine

- Receptors, Adrenergic, 78-6732

Receptors, Adrenergic

- Atria, Rat, Guinea Pig, 78-6732

etamide, *N*-(4-((2-Hydroxy-5-methylphenyl)azo)phenyl)-

Occupational Hazard

- Textile Workers, Review, 78-6663

etic Acid, (Ethylenedinitrilo)tetra-

Proteins

- Cell Transformation, Neoplastic, 78-6908
- Fibroblasts, 78-6908

etic Acid, 5-Fluoro-12-methylbenzanthryl-

Benzo(a)pyrene

- Antibody Formation, 78-7059
- Binding, Respiratory Tract, 78-7059
- Serum Albumin, Bovine, 78-7059
- Respiratory Tract Neoplasms
 - Benzo(a)pyrene, 78-7059

etic Anhydride

Sarcoma

- Transplantation Immunology, 78-7048

Acetohydroxamic Acid, *N*-Fluoren-2-yl-

Ascorbic Acid

- Nucleic Acids, Binding, 78-6730

7,8-Benzoflavone

- Mutagenic Activity, 78-6730

Cell Transformation, Neoplastic

- Organ Culture, Mammas, 78-6745

p-Cresol, 2,6-Di-*tert*-butyl-

- Mutagenic Activity, 78-6730

Enzyme Induction

- Mutagenic Activity, 78-6730

Ethylamine, 2,2'-Dithiobis-

- Nucleic Acids, Binding, 78-6730

Liver

- Mutagenic Activation, 78-6730
- Nucleic Acids, Binding, 78-6730

Methionine

- Nucleic Acids, Binding, 78-6730

Phosphoric Acid, Diethyl-, *p*-Nitrophenyl Ester

- Mutagenic Activity, 78-6730

Salmonella typhimurium

- Mutagenic Activity, 78-6730

Vitamin E

- Mutagenic Activity, 78-6730

Acetone

Leukemia, Lymphoblastic

- Case Report, 78-6765

Acetophenone, 2-Amino-

Fibroma

- Species Difference, 78-6790

Fibrosarcoma

- Species Difference, 78-6790

Gynecologic Neoplasms

- Fibroma, 78-6790
- Granulosa Cell Tumor, 78-6790
- Papilloma, 78-6790
- Species Difference, 78-6790

Leukemia, Lymphocytic

- Species Difference, 78-6790

Leukemia, Myelocytic

- Species Difference, 78-6790

Leukemia, Plasmacytic

- Species Difference, 78-6790

Liver Neoplasms

- Cholangioma, 78-6790
- Species Difference, 78-6790

Lymphosarcoma

- Species Difference, 78-6790

Mammary Neoplasms, Experimental

- Species Difference, 78-6790

Pancreatic Neoplasms

- Adenoma, 78-6790
- Islet Cell Tumor, 78-6790
- Species Difference, 78-6790

Achlorhydria

Stomach Neoplasms

- Polyps, 78-7116

Acrolein

Benzo(a)pyrene

- Co-carcinogenic Effect, Review, 78-6657

Lung Neoplasms

- Smoke Condensate, Review, 78-6657

Actin

- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-6846
 - Immunofluorescence Microscopy, 78-6846
 - Temperature Sensitive Mutants, 78-6846

Actinomycin D

- Bladder Neoplasms
 - DNA Polymerase, 78-6973
- Leukemia
 - Neoplasms, Multiple Primary, 78-7072
- Leukemia, Myeloblastic
 - Drug Therapy, 78-6815

Adenocarcinoma

- Carcinoembryonic Antigen
 - Oncogenic Viruses, 78-7118
 - Smoking, 78-7118
- Cell Differentiation
 - Neoplasm Suppression, Review, 78-6705
- Genetics
 - Carcinoembryonic Antigen, 78-7118
- Gynecologic Neoplasms
 - Contraceptives, Oral, 78-6645
- Intestinal Neoplasms
 - Carrageen, 78-6801
 - Celiac Disease, 78-7154
 - Cells, Cultured, 78-6680
 - Selenium, 78-6696
- Liver Neoplasms
 - Mycotoxins, 78-6723
- Lung Neoplasms
 - Needle Biopsy Implantation, Case Report, 78-7112
- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6775
 - 78-6776, 78-6780
 - Immunity, Cellular, 78-7004
 - Neoplasm Regression, 78-6981
 - Ultrastructural Study, Rat, 78-6780
- Ovarian Neoplasms
 - Cysts, 78-7132
 - Endometrial Hyperplasia, 78-7132
 - Histological Study, 78-7132
 - Hyperplasia, 78-7132
- Prostatic Neoplasms
 - Reverse Transcriptase, 78-6976
- Stomach Neoplasms
 - Histochemical Study, 78-7117
 - Transplantation, Heterologous, 78-7115
- Thyroid Neoplasms
 - Neoplasm Metastasis, 78-7102
 - Radiation, Ionizing, 78-7102
- Uterine Neoplasms
 - Genetics, 78-7118
 - Laurence-Moon-Biedl Syndrome, 78-7133
- Vaginal Neoplasms
 - Case Report, 78-7135
 - Ultrastructural Study, 78-7135

Adenofibroma

- Breast Neoplasms
 - Contraceptives, Oral, 78-6645
 - Precancerous Conditions, 78-6700
- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6780

Adenoma

- Hyperthyroidism
 - Radiation, Ionizing, 78-6831

Adenoma (cont'd)

- Intestinal Neoplasms
 - Bracken Fern, 78-6717
 - Carrageen, 78-6801
- Kidney Neoplasms
 - Streptozotocin, 78-6799
- Liver Neoplasms
 - Contraceptives, Oral, 78-6645, 78-6650, 78-6795
 - 78-7126
- Pancreatic Neoplasms
 - Acetophenone, 2-Amino-, 78-6790
 - Indole-3-acrylic Acid, 78-6790
- Parathyroid Glands
 - Transplantation, Autologous, 78-7103
- Pituitary Neoplasms
 - Contraceptives, Oral, 78-7095
 - Prolactin, 78-7095
- Thyroid Neoplasms
 - Radiation, Ionizing, 78-7171

Adenosine Cyclic 3',5' Monophosphate

- Cholangioma
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6766

Adenosine Triphosphatase

- Hepatoma
 - Hyperplasia, 78-6729

Adenosine Triphosphate

- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 - Mitochondrial Swelling, 78-6764
- Butyric Acid, 2-Amino-4-(ethylthio)-
 - Mitochondria, Liver, 78-6737
- Plasmacytoma
 - RNA Replication, 78-7190

Adenyl Cyclase

- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 - Glucagon, 78-6766
 - Isoproterenol, 78-6766
 - Sodium Fluoride, 78-6766
- Cell Membrane
 - Cell Transformation, Neoplastic, 78-7188
- Cell Transformation, Neoplastic
 - Enzymatic Activity, 78-7188
- Cholangioma
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6766

Adipose Tissue

- Breast Neoplasms
 - Precancerous Conditions, 78-7108

Aflatoxin B1

- Hepatoma
 - Food Contamination, 78-6721
 - Precancerous Conditions, 78-6722
 - Ultrastructural Study, Rat, 78-6722
- Mycotoxins
 - Carcinogenic Activity, Review, 78-6629
- Virus, SV40
 - Virus Replication, Review, 78-6620

Aflatoxin G1

- Hepatoma
 - Food Contamination, 78-6721

Aging

- Keratosis
 - DNA Repair, 78-7091

gmatine

see Guanidine, (4-Aminobutyl)-

grobacterium tumefaciens**Norocitopine**

Metabolism, 78-7194

Octopine

Extrachromosomal Inheritance, 78-7194

Metabolism, 78-7194

Regulatory Gene Mutants, 78-7194

Tumorigenicity, 78-7194

ir Pollution**Benzo(a)pyrene**

Epidemiology, Review, 78-6664

Maximum Permissible Concentrations, Review
78-6664

Free Radicals

Spectrum Analysis, 78-6710

Lung Neoplasms

Epidemiology, Review, 78-6701

Nitrogen Oxide

Spectrum Analysis, 78-6710

Nitrosamines

Carcinogenic Activity, Review, 78-6628

Concentration Levels, 78-7183

Occupational Hazard

Quantitation Method, 78-6709

Ozone

Spectrum Analysis, 78-6710

Polycyclic Hydrocarbons

Epidemiology, Review, 78-6664

Maximum Permissible Concentrations, Review
78-6664

Quantitation Method

Reference Standards, 78-6709

lanine, 3-(3,4-Dihydroxyphenyl)-**Breast Neoplasms, 78-6728**

Case Report, 78-6728

Growth, 78-6728

Prolactin, 78-6728

ldrin**Liver Neoplasms**

Epidemiology, Review, 78-6659

alkaline Phosphatase**Angiosarcoma**

Ethylene, Chloro-, 78-6713

Laryngeal Neoplasms

Neutrophils, 78-7099

alkylating Agents**Brain Neoplasms**

Species Difference, Review, 78-6639

Carcinogenic Metabolite

Mutagenic Activity, Review, 78-6635

Nucleic Acids

Teratogenic/Mutagenic Activity, Review, 78-6615

Structure-Activity Relationship

Carcinogenic Activity, Review, 78-6636

lpha Fetoproteins**Breast Neoplasms**

Genetics, 78-7066

Cholangioma

Serum Levels, 78-7124

Hepatoma

Immune Serums, 78-6960

Serum Levels, 78-7124

Alpha Fetoproteins (cont'd)**Hodgkin's Disease**

Genetics, 78-7066

Liver Neoplasms

Carcinoma, 78-7062

Hepatoma, 78-7062

Melanoma

Genetics, 78-7066

Alpha Particles**Lung Neoplasms**

Risk Estimation, 78-6821

Plutonium Oxide

Dissolution, Water, 78-6818

Uranium

Plutonium Oxide, 78-6818

 α -Amanitine**Plasmacytoma**

RNA Replication, 78-7190

Americium Oxide**Bone and Bones**

Inhalation Study, Rat, 78-6834

Lung Clearance

Inhalation Study, Rat, 78-6834

Lung Neoplasms

Inhalation Study, Rat, 78-6820

Radioisotopes, 78-6820

Ames Test**Carcinogen, Chemical**

Mutagenic Activity, 78-6792

Dyes

Mutagenic Activity, 78-6770

Molybdenum

Carcinogenic Activity, Review, 78-6635

Platinum

Carcinogenic Activity, Review, 78-6635

Psoralen, 8-Methoxy-

Mutagenic Activity, 78-6809

Quinoline

Cholanthrene, 3-Methyl-, 78-6762

Microsomes, Liver, 78-6762

Mutagenic Activity, 78-6762

5-Quinolinol

Mutagenic Activity, 78-6762

8-Quinolinol

Mutagenic Activity, 78-6762

Radiation

Mutagenic Activity, 78-6792

Rhodium

Carcinogenic Activity, Review, 78-6635

Ruthenium

Carcinogenic Activity, Review, 78-6635

Amino Acids**Antagonists**

Teratogenic/Mutagenic Activity, Review, 78-6615

Heavy Chain Disease

Proteins, 78-7060

Virus, Rous Sarcoma

Viral Proteins, 78-6852

Aminotransferases**Contraceptives, Oral**

Liver Function, Review, 78-6650

Anaphylaxis**IgE**

Immune Response, Solid Tumors, 78-6691

Anaphylaxis (cont'd)

- Neoplasms
- Immune Response, Solid Tumors, 78-6691

Anemia, Aplastic

- Benzene
- Epidemiology, Review, 78-6627
- Virus, SV40
- Twins, 78-6934

Aneuploidy

- Chromosome Abnormalities
- Histological Study, Rat, 78-7143
- Testicular Neoplasms
- Histological Study, Rat, 78-7143

Angiosarcoma

- Bone Neoplasms
- Strontium Radioisotopes, 78-6835
- Ethane, 1,2-Dibromo-
- Carcinogenic Activity, Rat, 78-6714
- Disulfide, Bis(diethylthiocarbamoyl)-, 78-6714
- Ethylene, Chloro-
- Alkaline Phosphatase, 78-6713
- Case Report, 78-6713
- Epidemiology, Review, 78-6634
- Glutamyl Transpeptidase, 78-6713
- Occupational Hazard, 78-6713

Aniline, *N,N*-Diethyl-*p*-phenylazo-

- Mitochondria, Liver
- Cytochrome Oxidase, 78-6764
- Malate Dehydrogenase, 78-6764

Aniline, *N,N*-Dimethyl-*p*-phenylazo-

- Adenosine Triphosphate
- Mitochondrial Swelling, 78-6764
- Hepatoma
- Antigen-Antibody Complex, 78-7042, 78-7064
- Complement, 78-7063
- Mitochondria, Liver
- Cytochrome Oxidase, 78-6764
- Malate Dehydrogenase, 78-6764
- Succinic Acid
- Mitochondrial Swelling, 78-6764

Anthracene

- Trachea
- Mucosal Toxicity, Rat, 78-6786

1,8,9-Anthracenetriol

- Leukemia L1210
- Ultraviolet Rays, 78-6807
- Ultraviolet Rays
- DNA Repair, 78-6807

Anthralin

- see 1,8,9-Anthracenetriol

Anthramycin

- Saccharomyces cerevisiae*
- DNA Repair, 78-6769

Anthraquinone

- Occupational Hazard
- Printers, Review, 78-6663

Anti-Antibodies

- Virus, Friend Murine Leukemia
- Antigens, Viral, 78-6879
- Virus, Rauscher Murine Leukemia
- Reverse Transcriptase, 78-6888
- Ribonuclease, 78-6888

Antibiotics, Antineoplastic

- Saccharomyces cerevisiae*
- DNA Repair, 78-6769

Antibodies, Viral

- Bladder Neoplasms
- Virus, Cytomegalo, 78-7068
- Breast Neoplasms
- Virus, Marek's Disease Herpes, 78-6861
- Carcinoma
- Virus, Marek's Disease Herpes, 78-6861
- Cervix Neoplasms
- Antigens, Bacterial, 78-7065
- Virus, Herpes Simplex 1, 78-6939
- Virus, Herpes Simplex 2, 78-6939, 78-7065
- Colonic Neoplasms
- Virus, Marek's Disease Herpes, 78-6861
- Head and Neck Neoplasms
- Virus, Epstein-Barr, 78-7034
- Leukemia, Myeloblastic
- Virus, Epstein-Barr, 78-7071
- Liver Neoplasms
- Virus, Marek's Disease Herpes, 78-6861
- Lung Neoplasms
- Virus, Marek's Disease Herpes, 78-6861
- Lymphoma
- Virus, Marek's Disease Herpes, 78-6861
- Melanoma
- Virus, Marek's Disease Herpes, 78-6861
- Nasopharyngeal Neoplasms
- Virus, Epstein-Barr, 78-7034
- Ovarian Neoplasms
- Virus, Herpes Simplex 1, 78-6939
- Virus, Herpes Simplex 2, 78-6939
- Virus, Marek's Disease Herpes, 78-6861
- Prostatic Hypertrophy
- Virus, Cytomegalo, 78-7068
- Prostatic Neoplasms
- Virus, Cytomegalo, 78-7068
- Sarcoma
- Virus, Marek's Disease Herpes, 78-6861
- Thymoma
- Virus Replication, 78-6874
- Virus, C-Type RNA Tumor
- Radioimmunoassay, Tissue Samples, 78-6679
- Virus, Epstein-Barr
- Immunosuppression, 78-6952
- Virus, Marek's Disease Herpes, 78-6861
- Virus, Herpes Simplex 2
- Antigens, Bacterial, 78-7065
- Virus, Polyoma, BK
- Immune Serums, 78-6905

Antibody Formation

- Benzo(a)pyrene
- 7-Acetic Acid, 5-Fluoro-12-methylbenzanthryl-
- 78-7059
- Corynebacterium parvum*
- Radiation, Ionizing, 78-7057
- Silica, 78-7057
- Leukemia
- Cells, Cultured, 78-7021
- Immunosuppression, 78-7021
- T-Lymphocytes, 78-7022
- Virus, Murine Leukemia, 78-7021, 78-7022
- Virus, Rauscher Murine Leukemia, 78-7021
- Leukemia, Lymphoblastic
- Transplantation Immunology, 78-7013

Antibody Formation (cont'd)

- Leukemia, Lymphocytic
 - Immune Response, Review, 78-6689
- Lymphoma
 - T-Lymphocytes, 78-7022
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Antigens, 78-7011
 - IgG, 78-7011
 - Lipopolysaccharides, 78-7011
 - Transplacental Effects, 78-7011
- Virus, Adeno 12
 - Lipopolysaccharides, 78-7008
- Virus, Gross Murine Leukemia
 - Formaldehyde, 78-7024
- Virus, Murine Leukemia
 - Immune Response, 78-6876
- Virus, Rauscher Murine Leukemia
 - Cells, Cultured, 78-7021
 - Formaldehyde, 78-7024
- Virus, SV40
 - Antigens, 78-7008
 - Lipopolysaccharides, 78-7008

Antibody Specificity

- Leukemia, Lymphocytic
 - IgM, 78-7073
- Leukemia, Myeloblastic
 - Antigens, 78-7019
- Mammary Neoplasms, Experimental
 - Antigen-Antibody Complex, 78-7058
- Multiple Myeloma
 - Virus, Bovine Leukemia, 78-6897
- Neoplasms
 - Antigen-Antibody Complex, 78-7058
- Virus, Epstein-Barr
 - B-Lymphocytes, 78-7015
- Virus, Herpes Simplex 1
 - Antigens, Viral, 78-6940
- Virus, Herpes Simplex 2
 - Antigens, Viral, 78-6940
- Virus, Simian Sarcoma
 - Viral Proteins, 78-6936

Antigen-Antibody Complex

- Hepatoma
 - Aniline, *N,N*-Dimethyl-*p*-phenylazo-, 78-7042
 - 78-7064
 - Complement, 78-7064
 - Complement Binding Assay, 78-7042
 - Growth, 78-7064
 - Transplantation Immunology, 78-7042
- Leukemia, Lymphoblastic
 - T-Lymphocytes, 78-7012, 78-7013
 - Transplantation Immunology, 78-7012
- Mammary Neoplasms, Experimental
 - Antibody Specificity, 78-7058
 - Kidney, 78-7058
 - Virus, Murine Mammary Tumor, 78-7058
- Neoplasms
 - Antibody Specificity, 78-7058
 - Kidney, 78-7058
- Sarcoma
 - Cholanthrene, 3-Methyl-, 78-7042
 - Complement Binding Assay, 78-7042
 - Transplantation Immunology, 78-7042
- Thymoma
 - T-Lymphocytes, 78-7012
 - Transplantation Immunology, 78-7012

Antigen-Antibody Reactions

- Bladder Neoplasms
 - Antigens, Neoplasm, 78-7199
- Burkitt's Lymphoma
 - Virus, Epstein-Barr, 78-6951
- Hepatoma
 - Australia Antigen, 78-6961
- Leukemia, Lymphoblastic
 - Precancerous Conditions, 78-7026
- Leukemia, Myeloblastic
 - Precancerous Conditions, 78-7026
- Liver Cirrhosis
 - Australia Antigen, 78-6961
- Lymphocytes
 - Immunoglobulins, Surface, 78-6986
- Lymphoma
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
 - Virus, Moloney Murine Leukemia, 78-6883
 - Virus, Rauscher Murine Leukemia, 78-6883
- Lymphosarcoma
 - Virus, Feline Leukemia, 78-6896
- Melanoma
 - Antigenic Determinants, 78-7037
 - BCG*, 78-7037
 - Immune Serums, 78-7037
- Osteopetrosis
 - Virus, Avian Myeloblastosis, 78-6844
- Plasmacytoma
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
- Sarcoma, Mast Cell
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
- Stomach Neoplasms
 - Carcinoembryonic Antigen, 78-7058
- Virus, Adeno 12
 - Antigens, Viral, 78-6970
- Virus, Avian Myeloblastosis
 - Glycoproteins, 78-6844
- Virus, Epstein-Barr
 - Precancerous Conditions, 78-6951
- Virus, Feline Leukemia
 - Occupational Hazard, 78-7165
 - Seroepidemiology, Humans, 78-7165
- Virus, Friend Murine Leukemia
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
- Virus, Gross Murine Leukemia
 - Viral Proteins, 78-6880
- Virus, Marek's Disease Herpes
 - Antigens, 78-6860
- Virus, Murine Leukemia
 - Antigens, Viral, 78-6876
- Virus, Murine Mammary Tumor
 - Viral Proteins, 78-6865
- Virus, Rous Sarcoma
 - Viral Proteins, 78-6856
- Virus, Turkey Herpes
 - Antigens, 78-6860

Antigenic Determinants

- Fibrosarcoma
 - Virus, Rat Leukemia, 78-6873
- Leukemia, Lymphoblastic
 - Bone Marrow Cells, 78-7026
 - Hematopoietic Stem Cells, 78-7026
 - B-Lymphocytes, 78-7026

Antigenic Determinants (cont'd)

- Null Cells, 78-7026
- Precancerous Conditions, 78-7026
- Leukemia, Lymphocytic
 - IgM, 78-7073
- Leukemia, Myeloblastic
 - Bone Marrow Cells, 78-7026
 - Hematopoietic Stem Cells, 78-7026
- Lymphoma
 - Immunity, Cellular, 78-7027
 - Killer Cells, 78-7009
 - Virus, Radiation Leukemia, 78-7029
- Lymphosarcoma
 - Virus, Feline Leukemia, 78-6896
- Melanoma
 - Antigen-Antibody Reactions, 78-7037
 - BCG, 78-7037
 - Laryngeal Neoplasms, 78-7037
- Neoplasms, Experimental
 - Transplantation Immunology, 78-7031
- Virus, Epstein-Barr
 - Cell Transformation, Neoplastic, 78-7015
 - Immunity, Cellular, 78-7015
- Virus, Moloney Murine Leukemia
 - Viral Proteins, 78-6877
 - Virus, Recombinant, 78-6884
- Virus, Murine Leukemia
 - Genes, Viral, 78-6892
 - Isolation and Characterization, 78-6892
 - Viral Proteins, 78-6871, 78-6877
- Virus, Radiation Leukemia
 - Viral Proteins, 78-6874
- Virus, Rauscher Murine Leukemia
 - Viral Proteins, 78-6889
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-6853
 - Viral Proteins, 78-6853
- Virus, Simian Sarcoma
 - Viral Proteins, 78-6936

Antigens

- Cell Transformation, Neoplastic
 - Immune Response, Review, 78-6694
- Leukemia
 - Immune Response, Review, 78-6681
 - Phenotype, Review, 78-6681
- Leukemia, Myeloblastic
 - Antibody Specificity, 78-7019
 - Cell Membrane, 78-7019
- Liver Neoplasms
 - Carcinogen, Chemical, 78-6692
 - Isolation and Characterization, Review, 78-6692
- Lymphoma
 - Immune Response, Review, 78-6681
- Myeloma Proteins
 - Binding Sites, 78-7040
- Neoplasms
 - Immune Response, Review, 78-6687
- 4,4'-Stilbenediol, α, α' -Diethyl-
 - Antibody Formation, 78-7011
- Virus, Adeno 12
 - Immunity, Cellular, 78-7008
- Virus, Marek's Disease Herpes
 - Antigen-Antibody Reactions, 78-6860
 - Cell Membrane, 78-6860
 - Spleen Cells, 78-6860
 - Virus, Turkey Herpes, 78-6860
- Virus, SV40

Antigens (cont'd)

- Antibody Formation, 78-7008
- Immunity, Cellular, 78-7008
- Virus, Turkey Herpes
 - Antigen-Antibody Reactions, 78-6860

Antigens, Bacterial

- Cervix Neoplasms
 - Antibodies, Viral, 78-7065
- Virus, Herpes Simplex 2
 - Antibodies, Viral, 78-7065

Antigens, Neoplasm

- Bladder Neoplasms
 - Antigen-Antibody Reactions, 78-7199
 - Carcinoma, Transitional Cell, 78-7199
 - Lymphocytotoxicity, 78-7199
- Immunosuppression
 - Cell Transformation, Neoplastic, Review, 78-6688
- Leukemia
 - Benzene, 1-Fluoro-2,4-dinitro-, 78-7025
 - Leukocytes, 78-7025
- Leukemia, Lymphoblastic
 - Transplantation Immunology, 78-7013
- Lymphocytes
 - Cytotoxicity, 78-6988
- Lymphoma
 - Cross Reactions, 78-7027
- Mammary Neoplasms, Experimental
 - Immune Serums, 78-6978
 - Virus, Murine Leukemia, 78-6978
- Melanoma
 - Hemadsorption Microassay, 78-7037
- Virus, Murine Leukemia
 - Immune Serums, 78-6978
 - Immunity, Cellular, 78-6978

Antigens, Viral

- Carcinoma
 - Virus, Marek's Disease Herpes, 78-6861
- Cell Transformation, Neoplastic
 - Immune Response, Review, 78-6694
- Erythroleukemia
 - Virus, Friend Murine Leukemia, 78-6879
- Histocompatibility Antigens
 - Immune Response, Review, 78-6694
- Leukemia
 - Immune Response, 78-7016
 - Virus, Murine Mammary Tumor, 78-7016
- Lymphoma
 - Cells, Cultured, 78-7010
 - Ultraviolet Rays, 78-7018
 - Virus, Moloney Murine Leukemia, 78-7010, 78-7018
- Melanoma
 - Virus, Marek's Disease Herpes, 78-6861
- Neoplasms
 - Virus, Polyoma, BK, 78-6905
- Plasmacytoma
 - Immune Response, Mouse, 78-7038
 - Virus, Gross Murine Leukemia, 78-7038
- Sarcoma
 - Virus, Marek's Disease Herpes, 78-6861
- Sarcoma, Reticulum Cell
 - Virus, Epstein-Barr, 78-6948
- Teratoid Tumor
 - Virus, Polyoma, 78-6902
 - Virus, SV40, 78-6932
- Virus, Adeno 12
 - Antigen-Antibody Reactions, 78-6970

Antigens, Viral (cont'd)

- Cytosine, 1- β -*D*-Arabinofuranosyl-, 78-6970
- Isolation and Characterization, Chick Fibroblasts 78-6970
- Virus, C-Type RNA Tumor
 - Radioimmunoassay, Tissue Samples, 78-6679
- Virus, Epstein-Barr
 - Complement Fixation, 78-6953
 - DNA, Binding, 78-6953
 - Immunity, Cellular, 78-7015
 - Isolation and Characterization, 78-6953
 - B-Lymphocytes, 78-7015
 - Phosphonoacetic Acid, 78-6946
 - Raji Cells, 78-7015
- Virus, Feline Leukemia
 - Isolation and Characterization, 78-6896
- Virus, Friend Murine Leukemia
 - Anti-Antibodies, 78-6879
 - Antibody-Dependent Cytotoxicity, 78-6879
 - Cell Membrane, 78-6879
 - Immune Response, 78-6879
 - Immunity, Passive, 78-6879
 - Radiation, Ionizing, 78-6879
- Virus, Herpes Saimiri
 - Phosphonoacetic Acid, 78-6946
- Virus, Herpes Simplex 1
 - Antibody Specificity, 78-6940
 - Cell Transformation, Neoplastic, 78-6944
 - DNA, Viral, 78-6944
 - Immune Serums, 78-6940
- Virus, Herpes Simplex 2
 - Antibody Specificity, 78-6940
 - Immune Serums, 78-6940
 - Immunity, Cellular, 78-7065
- Virus, Moloney Murine Leukemia
 - Ascitic Fluid, 78-7018
 - Cell Membrane, 78-7018
- Virus, Murine Leukemia
 - Antigen-Antibody Reactions, 78-6876
 - Mouse, AKR, 78-6876
- Virus, Murine Mammary Tumor
 - Immune Response, 78-7016
 - Milk, Hybrid Mouse, 78-6866
 - Radioimmunoassay, Epididymis, Coagulating Gland 78-6867
 - Vertical Transmission, 78-6866
- Virus, Papilloma
 - Carcinogenic Activity, Review, 78-6671
- Virus, Polyoma
 - Carcinogenic Activity, Review, 78-6671
 - Cell Differentiation, 78-6902
 - Hybrid Cells, 78-6902
- Virus, Polyoma, BK
 - Epidemiology, 78-6905
- Virus, Rauscher Murine Leukemia
 - Immunization, 78-7024
- Virus, RD-114
 - Isolation and Characterization, Review, 78-6674
- Virus, SV40
 - Carcinogenic Activity, Review, 78-6671
 - Cell Cycle Kinetics, 78-6932
 - Cell Fusion, 78-6925
 - Cell Transformation, Neoplastic, 78-6925, 78-6926 78-6930, 78-6931, 78-6932
 - Chromosomes, Binding, 78-6916
 - Deletion Mutants, 78-6920, 78-6930
 - DNA, Viral, 78-6920, 78-6927

Antigens, Viral (cont'd)

- Endonucleases, 78-6925
- Genetics, 78-6934
- Hybrid Cells, 78-6914, 78-6929
- Nucleotide Sequence, 78-6920
- Temperature Sensitive Mutants, 78-6926
- Viral Proteins, 78-6924
- Virus Replication, 78-6935

Antilymphocyte Serum

- Fibrosarcoma
 - T-Lymphocytes, 78-7067
- Leukemia
 - T-Lymphocytes, 78-7067
- Mammary Neoplasms, Experimental
 - T-Lymphocytes, 78-7067
- Burkitt's Lymphoma
 - Transplantation, Heterologous, 78-7086
- Lymphocytes
 - IgG, 78-6986
- T-Lymphocytes
 - Transplantation Immunology, 78-7044
- Neoplasms, Experimental
 - Virus, SV40, 78-6911

Antimetabolites

- Nucleic Acids
 - Teratogenic/Mutagenic Activity, Review, 78-6615

Ara-T

- see* Thymine, 1- β -*D*-Arabinofuranosyl-

Areca

- Fibrosarcoma
 - Carcinogenic Potential, Ingredients, 78-6727
- Mouth Neoplasms
 - Carcinogenic Potential, Ingredients, 78-6727
 - Precancerous Conditions, 78-6727
 - Vitamin B, 78-6727
- Skin Neoplasms
 - Carcinogenic Potential, Ingredients, 78-6727
- Vitamin B
 - Carcinogenic Potential, Ingredients, 78-6727

Arsenic

- Respiratory Tract Neoplasms
 - Epidemiology, Review, 78-6634

Aryl Hydrocarbon Hydroxylases

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Liver, Rat, 78-6773
- Benz(a)anthracene
 - Liver, Rat, 78-6773
 - Lymphocytes, 78-6781
 - Plant Agglutinins, 78-6781
- Benz(a)anthracene, 7,12-Dimethyl-
 - Liver, Rat, 78-6773
- 5,6-Benzoflavone
 - Lymphocytes, 78-6781
 - Plant Agglutinins, 78-6781
- Cell Transformation, Neoplastic
 - Liver, Rat, 78-6773
- Cells, Cultured
 - Liver, Rat, 78-6773
- Cholanthrene, 3-Methyl-
 - Liver, Mouse, 78-7041
 - Liver, Rat, 78-6773
 - Lymphocytes, 78-6781
 - Mitogens, 78-6781
 - Plant Agglutinins, 78-6781

- Aryl Hydrocarbon Hydroxylases (cont'd)**
 Cytochrome Reductases
 Quantitation Method, 78-6781
 DNA
 Quantitation Method, 78-6781
 Lymphocytes
 Enzyme Induction, 78-6788
 Phenotype, 78-6788
 Quinoline
 Mutagenic Activity, 78-6762
- Asbestos**
 Autoantibodies
 Precancerous Conditions, 78-7061
 Chromosome Abnormalities
 Fibers, 78-6836
 Hamster Cells, 78-6836
 Cummingtonite-Grunerite
 Carcinogenic Potential, 78-6838
 Gastrointestinal Neoplasms
 Epidemiology, Review, 78-6634
 Genetics
 Epidemiology, Review, 78-6697
 Immunity, Cellular
 Precancerous Conditions, 78-7061
 Immunoglobulins
 Occupational Hazard, 78-7061
 Iron
 Co-carcinogenic Effect, 78-6654
 Lung Neoplasms
 Epidemiology, Review, 78-6631
 Precancerous Conditions, 78-7061
 B-Lymphocytes
 Precancerous Conditions, 78-7061
 T-Lymphocytes
 Precancerous Conditions, 78-7061
 Mesothelioma
 Case Report, 78-6837
 Dust Application, Surgery, 78-6837
 Pericardium, 78-6837, 78-7113
 Neoplasms, Experimental
 Immune Response, Review, 78-6633
 Precancerous Conditions, Review, 78-6633
 Occupational Hazard
 Immunity, Cellular, 78-7061
 Respiratory Tract Neoplasms
 Cummingtonite-Grunerite, 78-6838
 Epidemiology, 78-6838
 Epidemiology, Review, 78-6634
 Stomach Neoplasms
 Occupational Hazard, 78-7172
- Asbestosis**
 Autoantibodies
 Occupational Hazard, 78-7061
 Benzene, 1-Chloro-2,4-dinitro-
 Hypersensitivity, Delayed, 78-7061
 Immunoglobulins
 Serum Levels, 78-7061
 Occupational Hazard
 B-Lymphocytes, 78-7061
 T-Lymphocytes, 78-7061
- Ascorbic Acid**
 Acetohydroxamic Acid, *N*-Fluoren-2-yl-
 Nucleic Acids, Binding, 78-6730
 Virus, Rous Sarcoma
 Cell Transformation, Neoplastic, 78-6849
 Glucose, 2-Deoxy-, 78-6849
- Ascorbic Acid, Monosodium Salt**
 Dimethylamine, *N*-Nitroso-
 Metabolism, 78-6802
 Guanidine, Methyl-
 Metabolism, 78-6802
- Astrocytoma**
 Brain Neoplasms
 Urea, Ethyl Nitroso-, 78-6749
 Spinal Cord Neoplasms
 Case Report, 78-7098
 Neoplasms, Multiple Primary, 78-7098
- Ataxia Telangiectasia**
 Chromosome Aberrations
 Fibroblasts, Review, 78-6612
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 DNA Repair, Review, 78-6612
 Leukemia, Lymphoblastic
 Case Report, 78-7030
 Chromosome Aberrations, 78-7030
 Chromosomes, Human, 13-15, 78-7030
 Radiation, Ionizing
 DNA Repair, Review, 78-6612
- Australia Antigen**
 Hepatoma
 Antigen-Antibody Reactions, 78-6961
 Cells, Cultured, 78-6957
 Epidemiology, 78-6961
 Immune Serums, 78-6960
 Immunohistochemical Study, 78-6958
 Liver Cirrhosis, 78-6961
 Neoplasm Metastasis, 78-6958
 Precancerous Conditions, 78-6961
 Seroepidemiology, Review, 78-6685
 Ultrastructural Study, 78-6957
 Virus, Hepatitis, 78-6685
 Liver Cirrhosis
 Antigen-Antibody Reactions, 78-6961
 Epidemiology, 78-6961
- Autoantibodies**
 Asbestos
 Precancerous Conditions, 78-7061
 Asbestosis
 Occupational Hazard, 78-7061
- Autoimmune Diseases**
 Cerebral Ischemia, Transient
 Stress, Review, 78-6682
- Bacteria**
 Colonic Neoplasms
 Diet, 78-7180
 Diet
 Gastrointestinal System, 78-7180
 Glioblastoma Multiforme
 Carcinogenic Metabolite, 78-7149
 Tooth Diseases, 78-7149
 Mutagens
 Colony Growth and Morphology, 78-7196
- Barbituric Acid, 5-Ethyl-5-phenyl-**
 Aryl Hydrocarbon Hydroxylases
 Liver, Rat, 78-6773
 Benzene, Bromo-
 Metabolism, Review, 78-6607
 Hepatoma
 Acetamide, *N*-Fluoren-2-yl-, 78-6729
 Liver Neoplasms

Barbituric Acid, 5-Ethyl-5-phenyl- (cont'd)

Dietary Proteins, 78-6742

Skin Neoplasms

Ultraviolet Rays, 78-6806

Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt**Liver Neoplasms**

Carcinogenic Potential, Rat, 78-6758

Hyperplasia, 78-6758

BCG**Carcinoma**

Growth, 78-6996

Macrophages, 78-6996

Transplantation Immunology, 78-6996

Cholanthrene, 3-Methyl-

Immune Response, Mouse, 78-7007

Hepatoma

Macrophages, 78-6995

Neoplasm Regression, 78-6995

Immunity, Cellular

Cell Transformation, Neoplastic, 78-7006

Leukemia L1210

Immune Response, 78-7020

Lymphoma

Killer Cells, 78-7009

Melanoma

Antigen-Antibody Reactions, 78-7037

Antigenic Determinants, 78-7037

Immune Response, Mouse, 78-6990

Sarcoma

Macrophages, 78-6995

Neoplasm Regression, 78-6995

Virus, Adeno 12

Tuberculin, 78-7008

Virus, B77

Cell Transformation, Neoplastic, 78-7006

Virus, Rous Sarcoma

Cell Transformation, Neoplastic, 78-7006

Immunity, Cellular, 78-7006

Methanol Extraction Residue, 78-7006

Virus, SV40

Tuberculin, 78-7008

Benz(a)anthracene**Aryl Hydrocarbon Hydroxylases**

Liver, Rat, 78-6773

Lymphocytes

Aryl Hydrocarbon Hydroxylases, 78-6781

Microsomes, Liver

Carcinogenic Metabolite, 78-6772

Plant Agglutinins

Aryl Hydrocarbon Hydroxylases, 78-6781

Trachea

Mucosal Toxicity, Rat, 78-6786

Tracheal Neoplasms

Precancerous Conditions, 78-6786

Benz(a)anthracene, 1,2-Dihydro-1,2-dihydroxy-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 8,9-Dihydro-8,9-dihydroxy-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 10,11-Dihydro-10,11-dihydroxy-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 1,2-Dihydro-1,2-dihydroxy-7,12-dimethyl-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-7,12-dimethyl-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 5,6-Dihydro-5,6-dihydroxy-7,12-dimethyl-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 8,9-Dihydro-8,9-dihydroxy-7,12-dimethyl-**Microsomes, Liver**

Carcinogenic Metabolite, 78-6772

Benz(a)anthracene, 7,12-Dimethyl-**Aryl Hydrocarbon Hydroxylases**

Liver, Rat, 78-6773

Cell Transformation, Neoplastic

Organ Culture, Mammary, 78-6745

Cells, Cultured

Cell Transformation, Neoplastic, 78-6785

Croton Oil

Two-Stage Carcinogenesis, Review, 78-6626

Fibroblasts

Proteins, Binding, 78-6785

Two-Stage Carcinogenesis, Review, 78-6626

Fibrosarcoma*Corynebacterium parvum*, 78-7055**T-Lymphocytes**

Cell Transformation, Neoplastic, 78-6984

Rosette Formation, 78-6984

Mammary Neoplasms, Experimental

Adenocarcinoma, 78-6775, 78-6776, 78-6780

Adenofibroma, 78-6780

FSH, 78-6778

Gonadotropins, 78-6778

Histological Study, Rat, 78-6775

Hormone Dependency, 78-6774

LH, 78-6778

Melatonin, 78-6779

Neoplasm Transplantation, 78-6774

Placental Extracts, 78-6775

Precancerous Conditions, 78-6774, 78-6776, 78-6780

Prolactin, 78-6778, 78-6779

Thyrotropin, 78-6778

Ultrastructural Study, Rat, 78-6780

Microsomes, Liver

Carcinogenic Metabolite, 78-6772

Mouth Neoplasms

Benzene, 1-Chloro-2,4-dinitro-, 78-7033

Carcinoma, Epidermoid, 78-7033

Papilloma, 78-7033

Precancerous Conditions, 78-7075

Neoplasms, Experimental

Chemotaxis, 78-6991

12-O-Tetradecanoylphorbol-13-acetate

Two-Stage Carcinogenesis, Review, 78-6626

Trachea

Mucosal Toxicity, Rat, 78-6786

Tracheal Neoplasms

Precancerous Conditions, 78-6786

Ultraviolet Rays

Melanocytes, Mouse, 78-6777

Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-

- Cholangioma
 - Adenosine Cyclic 3',5' Monophosphate, 78-6766
 - Adenyl Cyclase, 78-6766
 - Guanosine Cyclic 3',5' Monophosphate, 78-6767
 - Guanyl Cyclase, 78-6767
 - Phosphodiesterases, 78-6766
 - Precancerous Conditions, 78-6767

- Glucagon
 - Adenyl Cyclase, 78-6766

- Hepatoma
 - DNA Replication, 78-6744

- Isoproterenol
 - Adenyl Cyclase, 78-6766

- Liver Neoplasms
 - Cholangioma, 78-6767

- Sodium Fluoride
 - Adenyl Cyclase, 78-6766

Benzene

- Anemia, Aplastic
 - Epidemiology, Review, 78-6627
- Chromosome Aberrations
 - Epidemiology, Review, 78-6627
- Leukemia, Myeloblastic
 - Epidemiology, Review, 78-6627
- Occupational Hazard
 - Epidemiology, Review, 78-6627
 - Maximum Permissible Exposure Level, Review 78-6627

Benzene, Bromo-

- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Metabolism, Review, 78-6607
- Cytochrome P-450
 - Hepatotoxicity, Review, 78-6607
- Microsomes, Liver
 - Metabolism, Review, 78-6607
- Nucleic Acids, Binding
 - Hepatotoxicity, Review, 78-6607

Benzene, 1-Chloro-2,4-dinitro-

- Asbestosis
 - Hypersensitivity, Delayed, 78-7061
- Mouth Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-7033
 - Immunity, Cellular, 78-7033
 - Immunotherapy, 78-7033

Benzene, 1-Fluoro-2,4-dinitro-

- Leukemia
 - Antigens, Neoplasm, 78-7025

Benzidine, Dihydrochloride

- Hepatoma
 - Neoplasm Metastasis, 78-7122
- Liver Neoplasms
 - Benzo(a)pyrene, 78-7122
 - Diethylamine, *N*-Nitroso-, 78-7122
 - Neoplasm Metastasis, 78-7122

2-Benzimidazolecarbamate Acid, 5-(2-Thienylcarbonyl)- Methyl Ester

- Cell Division
 - Cells, Cultured, 78-7081
- Sarcoma
 - Cell Migration Inhibition, 78-7081
- Uracil, 5-Fluoro
 - Cell Migration Inhibition, 78-7081

1,2-Benzisothiazolin-3-one, 1,1-Dioxide

- Bladder Neoplasms
 - Carcinogenic Activity, Mouse, Rat, 78-6660
 - Epidemiology, Review, 78-6660
 - Histological Study, Rat, Review, 78-6660
 - Species Difference, 78-6738
- Gynecologic Neoplasms
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Lung Neoplasms
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Lymphosarcoma
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Mammary Neoplasms, Experimental
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Skin Neoplasms
 - Croton Oil, 78-6660
- Urea, Methyl Nitroso-
 - Co-carcinogenic Effect, Review, 78-6660

Benzo(a)pyrene

- 7-Acetic Acid, 5-Fluoro-12-methylbenzanthryl-
 - Antibody Formation, 78-7059
 - Binding, Respiratory Tract, 78-7059
 - Serum Albumin, Bovine, 78-7059
- Acrolein
 - Co-carcinogenic Effect, Review, 78-6657
- Air Pollution
 - Epidemiology, Review, 78-6664
 - Maximum Permissible Concentrations, Review 78-6664
- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
 - Carcinogenic Metabolite, Review, 78-6623
- Cell Transformation, Neoplastic
 - Anchorage-Independent Growth, 78-7076
 - Cells, Cultured, 78-7076
- Cells, Cultured
 - Cell Transformation, Neoplastic, 78-6785
- Croton Oil
 - Two-Stage Carcinogenesis, Review, 78-6626
- DNA Repair
 - Cell Cycle Kinetics, 78-6785
- Fibrinolysis
 - Cell Transformation, Neoplastic, 78-7076
- Fibroblasts
 - Proteins, Binding, 78-6785
 - Two-Stage Carcinogenesis, Review, 78-6626
- Fibrosarcoma
 - Cells, Cultured, 78-7076
 - Transplantation Immunology, 78-7051
- Formaldehyde
 - Co-carcinogenic Effect, Review, 78-6657
- Hepatoma
 - Neoplasm Metastasis, 78-7122
- Hydrocyanic Acid
 - Co-carcinogenic Effect, Review, 78-6657
- Liver Neoplasms
 - Benzidine, Dihydrochloride, 78-7122
- Lung Neoplasms
 - Dose-Response Study, Review, 78-6701
 - Epidemiology, Review, 78-6701
 - Occupational Hazard, Review, 78-6701
 - Smoke Condensate, Review, 78-6657
- Myosarcoma
 - Graft vs Host Reaction, 78-7045
 - Splenectomy, 78-7046
- Respiratory Tract Neoplasms
 - 7-Acetic Acid, 5-Fluoro-12-methylbenzanthryl-

benzo(a)pyrene (cont'd)

- 7-Acetic Acid, 5-Fluoro-12-methylbenzanthryl-78-7059

Retinol

- Carcinogenic Activity, Review, 78-6630

Sarcoma

- Graft vs Host Reaction, 78-7045
- B-Lymphocytes, 78-7045
- T-Lymphocytes, 78-7045

Serum Albumin, Bovine

- Immunization, 78-7059

Smoking

- Co-carcinogenic Activity, 78-6701

12-*O*-Tetradecanoylphorbol-13-acetate

- Two-Stage Carcinogenesis, Review, 78-6626

Trachea

- Mucosal Toxicity, Rat, 78-6786

Tracheal Neoplasms

- Precancerous Conditions, 78-6786

benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-**Benzo(a)pyrene**

- Carcinogenic Metabolite, Review, 78-6623

Chromatin

- DNA, Binding, 78-6784
- Nucleic Acid Conformation, 78-6784

Endonucleases

- DNA, Binding, 78-6784

Histones

- DNA, Binding, 78-6784

benzo(a)pyrene, 1,3-Dimethyl-**Cation Radicals**

- Electrochemistry, 78-6783

Oxidation-Reduction

- Electrochemistry, 78-6783

benzo(a)pyrene, 1,6-Dimethyl-**Cation Radicals**

- Electrochemistry, 78-6783

Oxidation-Reduction

- Electrochemistry, 78-6783

benzo(a)pyrene, 2,3-Dimethyl-**Cation Radicals**

- Electrochemistry, 78-6783

Oxidation-Reduction

- Electrochemistry, 78-6783

benzo(a)pyrene, 3,6-Dimethyl-**Cation Radicals**

- Electrochemistry, 78-6783

Oxidation-Reduction

- Electrochemistry, 78-6783

benzo(a)pyrene, 4,5-Dimethyl-**Cation Radicals**

- Electrochemistry, 78-6783

Oxidation-Reduction

- Electrochemistry, 78-6783

benzo(b)triphenylene**Trachea**

- Mucosal Toxicity, Rat, 78-6786

benzo(e)pyrene**Trachea**

- Mucosal Toxicity, Rat, 78-6786

5,6-Benzoflavone**Lymphocytes**

- Aryl Hydrocarbon Hydroxylases, 78-6781

5,6-Benzoflavone (cont'd)**Plant Agglutinins**

- Aryl Hydrocarbon Hydroxylases, 78-6781

7,8-Benzoflavone**Acetohydroxamic Acid, *N*-Fluoren-2-yl-**

- Mutagenic Activity, 78-6730

Benzophenone**Occupational Hazard**

- Printers, Review, 78-6663

Beryllium Chloride**Hyperplasia**

- Spleen, Mouse, Rat, 78-6817

Betel

- see Areca

Bile Duct Neoplasms**Cholangioma**

- Epidemiology, Japan, 78-7127

Bladder Neoplasms**Antigens, Neoplasm**

- Antigen-Antibody Reactions, 78-7199

Lymphocytotoxicity, 78-7199**1,2-Benzisothiazolin-3-one, 1,1-Dioxide**

- Carcinogenic Activity, Mouse, Rat, 78-6660

Epidemiology, Review, 78-6660**Histological Study, Rat, Review, 78-6660****Species Difference, 78-6738****1-Butanol, 4-(Butylnitrosamino)-**

- Immune Response, Rat, 78-6735

Precancerous Conditions, 78-6735**Carcinoma In Situ**

- Case Report, 78-7139

Neoplasm Invasiveness, 78-7139**Precancerous Conditions, 78-7139****Carcinoma, Transitional Cell**

- Antigens, Neoplasm, 78-7199

1-Butanol, 4-(Butylnitrosamino)-, 78-6735**Cell Line, Epithelioid, 78-7199****DNA-RNA Hybridization, 78-6973****Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-78-7141****Neoplasm Invasiveness, 78-7141****Reverse Transcriptase, 78-6973****Ribonuclease, 78-6973*****Schistosoma haematobium*, 78-7140****Virus, Cytomegalo, 78-7068****Cells, Cultured**

- Reverse Transcriptase, 78-6973

Desmosomes

- Neoplasm Invasiveness, 78-7141

Ultrastructural Study, 78-7141**DNA Polymerase**

- Actinomycin D, 78-6973

Ribonuclease, 78-6973**Lymphocytes**

- Concanavalin A, 78-6735

Immune Response, Rat, 78-6735**Plant Agglutinins, 78-6735*****Schistosoma haematobium***

- Histological Study, Monkey, 78-7140

Neoplasm Transplantation, 78-7140**Sex Chromosomes**

- Cell Line, Epithelioid, 78-7199

Virus, Cytomegalo

- Antibodies, Viral, 78-7068

- Bleomycin**
 - Sarcoma
 - Cell Migration Inhibition, 78-7081
- Blood Platelets**
 - Fibroblasts
 - Growth Substances, 78-6928
 - Virus, SV40
 - Growth Substances, 78-6928
- Blood Vessels**
 - DNA
 - Carcinogenic Potential, 78-7191
- Bone Marrow**
 - Leukemia, Radiation-Induced
 - DNA Nucleotidyltransferases, 78-6826
- Bone Marrow Cells**
 - Leukemia, Lymphoblastic
 - Antigenic Determinants, 78-7026
 - Leukemia, Myeloblastic
 - Antigenic Determinants, 78-7026
 - Lung Neoplasms
 - Colony Formation, 78-6999
 - Radiation, Ionizing
 - Radiation-Protective Agents, 78-6979
- Bone Neoplasms**
 - Angiosarcoma
 - Strontium Radioisotopes, 78-6835
 - Fibrosarcoma
 - Strontium Radioisotopes, 78-6835
 - Neoplasm Recurrence, Local
 - Case Report, 78-7147
 - Osteoma, Osteoid
 - Neoplasm Recurrence, Local, 78-7147
 - Sarcoma, Osteogenic
 - Neoplasm Recurrence, Local, 78-7147
 - Strontium Radioisotopes, 78-6835
- Bone Resorption**
 - Estrogens
 - Menopause, Review, 78-6651
- Bracken Fern**
 - Digestive System Neoplasms
 - Epidemiology, Cattle, 78-6899
- HeLa Cells**
 - Toxicity, Pterosins, Pterosides, 78-6717
- Intestinal Neoplasms**
 - Adenoma, 78-6717
 - Carcinogenic Activity, Rat, 78-6717
 - Carcinoma, Papillary, 78-6717
- Virus, Papilloma**
 - Co-carcinogenic Effect, 78-6899
- Bradykinin**
 - Lung Neoplasms
 - Macrophages, 78-7001
 - Tumor Cell Supernatant, 78-7001
- Brain Neoplasms**
 - Alkylating Agents
 - Species Difference, Review, 78-6639
 - Astrocytoma
 - Urea, Ethyl Nitroso-, 78-6749
 - Carcinogen, Chemical
 - Epidemiology, 78-7168
 - Ethylene, Chloro-
 - Epidemiology, Review, 78-6634
 - Glioma
 - Urea, Ethyl Nitroso-, 78-6749
 - Hormones
 - Epidemiology, 78-7168
 - Meningioma
 - Epidemiology, 78-7170
 - Oligodendroglioma
 - Urea, Ethyl Nitroso-, 78-6749
 - Oncogenic Viruses
 - Epidemiology, 78-7168
 - Species Difference, Review, 78-6639
 - Polycyclic Hydrocarbons
 - Species Difference, Review, 78-6639
 - 2-Propanol, 1,1'-Iminodi-*N*-nitroso-
 - Transplacental Carcinogenesis, 78-6749
 - Quinoline, 4-(Hydroxyamino)-, 1-Oxide
 - Transplacental Carcinogenesis, 78-6749
 - Transplantation, Heterologous
 - Nude Mouse, Review, 78-6639
 - Urea, Ethyl Nitroso-
 - Transplacental Carcinogenesis, 78-6749
- Brain Neoplasms (cont'd)**
 - Urea, Ethyl Nitroso-, 78-6749
- Breast Diseases**
 - Cysts
 - Contraceptives, Oral, 78-7107
- Breast Neoplasms**
 - Adenocarcinoma
 - Alanine, 3-(3,4-Dihydroxyphenyl)-, 78-6728
 - Adenofibroma
 - Contraceptives, Oral, 78-6645
 - Precancerous Conditions, 78-6700
 - Age Factors
 - Epidemiology, 78-6700, 78-7159, 78-7160
 - Alanine, 3-(3,4-Dihydroxyphenyl)-
 - Case Report, 78-6728
 - Growth, 78-6728
 - Prolactin, 78-6728
 - Carcinoma, Ductal
 - Cytodiagnosis, 78-7105
 - Epidemiology, 78-7161
 - Neoplasm Metastasis, 78-7161
 - Virus, RNA Tumor, 78-6869
 - Contraceptives, Oral
 - Epidemiology, Review, 78-6645, 78-6648
 - Precancerous Conditions, 78-6648, 78-7107
 - Ultrastructural Study, Review, 78-6645
 - Cysts
 - Cytodiagnosis, 78-7105
 - Diabetes Mellitus
 - Epidemiology, Sweden, 78-7106
 - Diet
 - Epidemiology, Review, 78-6637
 - Dietary Fats
 - Epidemiology, Review, 78-6699
 - Epidemiology
 - Iran, 78-7153
 - Estrogens
 - Epidemiology, 78-7162
 - Epidemiology, Review, 78-6644, 78-6649, 78-6651
 - Menopause, 78-6649
 - Menopause, Review, 78-6651
 - Receptors, Hormone, Review, 78-6603
 - Ethnic Groups
 - Epidemiology, Hawaii, 78-7164
 - Ethylene, Trichloro-
 - Epidemiology, 78-7173
 - Genetics

Breast Neoplasms (cont'd)

- Alpha Fetoproteins, 78-7066
- Carcinoembryonic Antigen, 78-7066

Hair Dyes

- Epidemiology, 78-7163, 78-7166

Hormones

- Cell Division, 78-6791
- Plasma Levels, 78-7162

Hyperplasia

- Contraceptives, Oral, 78-6645
- Epidemiology, Japan, Hawaii, 78-7108
- Precancerous Conditions, 78-7108

Hypertension

- Epidemiology, Sweden, 78-7106

Mastitis

- Precancerous Conditions, 78-6700

Melatonin

- Epidemiology, 78-6779
- Prolactin, 78-6779

Myasthenia Gravis

- Thymectomy, 78-7035

Neoplasm Metastasis

- Age Factors, 78-7159

Neoplasms, Multiple Primary

- Epidemiology, 78-7158

Papilloma

- Cytodiagnosis, 78-7105

Parity

- Epidemiology, 78-6700

Precancerous Conditions

- Adipose Tissue, 78-7108
- Epidemiology, 78-6700

Progesterone

- Epidemiology, 78-7162
- Receptors, Hormone, Review, 78-6603

Prolactin

- Epidemiology, 78-7162
- Thyroid Dysfunction, Review, 78-6640

RNA Polymerase

- Estrogens, 78-6603

Selenium

- Antitumor Activity, Review, 78-6696
- Epidemiology, Review, 78-6696

Smoking

- Carcinoembryonic Antigen, 78-7066

Steroids

- Receptors, Hormone, Review, 78-6603

Thyrotropin

- Thyroid Dysfunction, Review, 78-6640

Thyrotropin Releasing Hormone

- Prolactin, 78-6640
- Thyrotropin, 78-6640

Virus, Marek's Disease Herpes

- Antibodies, Viral, 78-6861

Virus, Mason-Pfizer Monkey

- DNA-RNA Hybridization, 78-6869

Virus, Murine Mammary Tumor

- DNA-RNA Hybridization, 78-6869

Bronchial Neoplasms**Smoking**

- Epidemiology, Switzerland, 78-7151

Bronchitis**Leukemia, Myeloblastic**

- Precancerous Conditions, 78-7074

Brucella abortus**Lymphoma*****Brucella abortus* (cont'd)**

- Dose-Response Study, Mouse, 78-7014
- Immune Response, Mouse, 78-7014
- Immunization, 78-7014

Burkitt's Lymphoma**Leukemia**

- Transplantation, Heterologous, 78-7086

Transplantation, Heterologous

- Antilymphocyte Serum, 78-7086
- Histological Study, Hamster, 78-7086
- Neoplasm Invasiveness, 78-7086

Virus, Epstein-Barr

- Antigen-Antibody Reactions, 78-6951
- Seroepidemiology, Uganda, 78-6951

Warts

- Genetics, Review, 78-6684

2,3-Butanediol, 1,4-Dimercapto-**Virus, SV40**

- Chromosomes, 78-6907

1-Butanol, 4-(Butylnitrosamino)-**Bladder Neoplasms**

- Carcinoma, Transitional Cell, 78-6735
- Immune Response, Rat, 78-6735
- Precancerous Conditions, 78-6735

2-Butanone, 1,1-Dihydroxy-3-ethoxy-**DNA**

- Liver, Rat, 78-6716

Fluorescence

- Liver, Rat, 78-6716

RNA, Ribosomal

- Liver, Rat, 78-6716

Butylhydroxytoluene

- see *p*-Cresol, 2,6-Di-*tert*-butyl-

Butyric Acid**Erythroleukemia**

- Globin, 78-7185

Butyric Acid, 2-Amino-4-(ethylthio)-**Mitochondria, Liver**

- Adenosine Triphosphate, 78-6737
- Oxidative Phosphorylation, 78-6737

Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-**Leukemia**

- Neoplasms, Multiple Primary, 78-7072

C.I. Basic Green 4**Occupational Hazard**

- Textile Workers, Review, 78-6663

C.I. Disperse Yellow 3

- see Acetamide, *N*-(4-((2-Hydroxy-5-methylphenyl)azo)phenyl)-

C.I. Vat Green 1**Occupational Hazard**

- Textile Workers, Review, 78-6663

Cadmium**Chromosome Aberrations**

- Cells, Cultured, 78-6797
- Culture Media, 78-6797

Caffeine**Ultraviolet Rays**

- Mutagenic Activity, Review, 78-6606

- Calcitonin**
Skin Neoplasms
Cholanthrene, 3-Methyl-, 78-6798
- Calcium**
Fibroblasts
Growth Substances, 78-6928
- Carbamic Acid, Ethyl Ester**
Esterases
Plasma Levels, Mouse, 78-6743
Mutagenic Activity
Micronucleus/Sperm Abnormality Assays, 78-6792
- Carbamic Acid, N-Methyl-N-nitroso-, 1-Naphthyl Ester**
Haemophilus influenzae
DNA Repair, 78-6759
Radiation, Ionizing
DNA Repair, 78-6759
Stomach
Synthesis, Guinea Pig, Rat, 78-6754
Ultraviolet Rays
DNA Repair, 78-6759
- Carbodiimide, 1-Ethyl-3-(3-dimethylaminopropyl)-**
Sarcoma
Transplantation Immunology, 78-7048
- Carbohydrates**
Virus, Gross Murine Leukemia
Viral Proteins, 78-6880
- Carbon Tetrachloride**
Diethylamine, N-Nitroso-
Co-carcinogenic Effect, 78-6750
Esterases
Plasma Levels, Mouse, 78-6743
Hepatoma
Diethylamine, N-Nitroso-, 78-6750
Microsomes, Liver
Metabolism, Review, 78-6607
- Carboxy-Lyases**
Carcinogen, Chemical
Two-Stage Carcinogenesis, Review, 78-6606
Kidney Neoplasms
Dimethylamine, N-Nitroso-, 78-6742
Polyamines, 78-6742
Liver Neoplasms
Diethylamine, N-Nitroso-, 78-6742
Polyamines, 78-6742
- Carcinoembryonic Antigen**
Adenocarcinoma
Genetics, 78-7118
Oncogenic Viruses, 78-7118
Smoking, 78-7118
Breast Neoplasms
Genetics, 78-7066
Smoking, 78-7066
Cell Transformation, Neoplastic
Immune Response, Review, 78-6694
Hodgkin's Disease
Genetics, 78-7066
Liver Neoplasms
Carcinogen, Chemical, 78-6692
Melanoma
Genetics, 78-7066
Oncogenic Viruses
Genetics, 78-7118
RNA, Transfer, Methyltransferases
- Carcinoembryonic Antigen (cont'd)**
Uracil, 78-6621
Smoking
Genetics, 78-7118
Stomach Neoplasms
Antigen-Antibody Reactions, 78-7058
- Carcinogen, Chemical**
Abnormalities
Teratogenic Effect, Review, 78-6616
Ames Test
Mutagenic Activity, 78-6792
Carboxy-Lyases
Two-Stage Carcinogenesis, Review, 78-6606
Carcinogenic Potential
Bioassay, Review, 78-6601
Cell Cycle Kinetics
Cell Transformation, Neoplastic, Review, 78-6703
Cell Transformation, Neoplastic
Animal Model, Hamster Embryo, 78-6726
Bioassay, Review, 78-6601
DNA, Binding, Review, 78-6604
Dose-Response Study, Review, 78-6604
Mathematical Model, 78-7186
Mutagenic Activity, Review, 78-6616
Chromosome Abnormalities
Carcinogenic Activity, Review, 78-6610
DNA
Mutagenic Activity, Review, 78-6616
DNA Repair
Bioassay, Review, 78-6608
Carcinogenic Activity, Review, 78-6608, 78-6619
Carcinogenic/Mutagenic Activity, 78-6711
Cell Transformation, Neoplastic, Review, 78-6613
In Vivo Measurement Methods, Review, 78-6619
Food Additives
Risk Factors, Review, 78-6662
Food Contamination
Risk Factors, Review, 78-6662
Gastrointestinal Neoplasms
Diet, Review, 78-6638
Genetics
Cell Transformation, Neoplastic, Review, 78-6705
Hormones
Co-carcinogenic Effect, Review, 78-6643
Immunosuppression
Epidemiology, Review, 78-6697
Liver Neoplasms
Antigens, 78-6692
Carcinoembryonic Antigen, 78-6692
Choline, 78-6638
Chromosomes, Review, 78-6706
Diet, Review, 78-6638
Glycogen, 78-6707
Histocompatibility Antigens, 78-6692
Methionine, 78-6638
Precancerous Conditions, Review, 78-6706, 78-6707
Ultrastructural Study, Review, 78-6707
Vitamin B12, 78-6638
Mammary Neoplasms, Experimental
Hormones, 78-6791
Mutagenic Activity
Bioassay, Review, 78-6601
Colony Growth and Morphology, 78-7196
Micronucleus/Sperm Abnormality Assays, 78-6792
Neoplasms, Experimental
Trans-Species Extrapolation, Review, 78-6604
Nucleic Acids

Carcinogen, Chemical (cont'd)

- Mutagenic Activity, Review, 78-6611
- Occupational Hazard
 - Epidemiology, Review, 78-6697
- Plasminogen Activators
 - Two-Stage Carcinogenesis, Review, 78-6606
- Precancerous Conditions
 - Co-carcinogenic Activity, Review, 78-6611
- Respiratory Tract Neoplasms
 - Occupational Hazard, 78-6768
- Retinol
 - Immune Response, Review, 78-6630
- Risk Evaluation
 - Criteria, Review, 78-6617
- Skin Neoplasms
 - Co-carcinogenic Activity, Review, 78-6611
- Structure-Activity Relationship
 - Carcinogenic Activity, Review, 78-6636
- Xeroderma Pigmentosum
 - DNA Repair, Review, 78-6614

Carcinogen, Environmental

- Cell Differentiation
 - Mutation, Review, 78-6602
- Chromosome Abnormalities
 - Cell Transformation, Neoplastic, Review, 78-6602
- DNA
 - Mutagenic Activity, Review, 78-6609
- Genetics
 - Cell Transformation, Neoplastic, Review, 78-6609
 - 78-6610
- Hereditary Diseases
 - Cell Transformation, Neoplastic, Review, 78-6602
 - Mutation, Review, 78-6602
- Latency Period
 - Cell Transformation, Neoplastic, Review, 78-6602
- Precancerous Conditions
 - Co-carcinogenic Activity, Review, 78-6611

Carcinoma

BCG

- Growth, 78-6996
- Macrophages, 78-6996
- Transplantation Immunology, 78-6996
- Carrageen
 - Cytotoxic Activity, 78-6997
- Cervix Neoplasms
 - Virus, Herpes Simplex 2, 78-6939
- Epidemiology
 - Pediatric Patients, 78-7177
- Intestinal Neoplasms
 - Pediatric Patients, 78-7177
- Liver Neoplasms
 - Alpha Fetoproteins, 78-7062
 - Radioimmunoassay, 78-7062
- Lung Neoplasms
 - Foreign Body Reaction, 78-6992
 - Neoplasm Metastasis, 78-6998
 - Phagocytosis, 78-6998
- Nasopharyngeal Neoplasms
 - Cellular Inclusions, 78-7101
 - Virus, Epstein-Barr, 78-7034
- Pancreatic Neoplasms
 - Ribonuclease, 78-7200
- Pharyngeal Neoplasms
 - Neoplasm Recurrence, Local, 78-7100
- Virus, C-Type RNA Tumor
 - Ultrastructural Study, Review, 78-6679

Carcinoma (cont'd)

- Virus, Marek's Disease Herpes
 - Antibodies, Viral, 78-6861
 - Antigens, Viral, 78-6861
- Vulvar Neoplasms
 - Diagnosis, 78-7137
 - Epidemiology, 78-7137

Carcinoma, Basal Cell

- Cicatrix
 - Case Report, 78-7089
- Kidney Neoplasms
 - Case Report, 78-7129
 - Neoplasm Metastasis, 78-7129
- Muscular Atrophy
 - Case Report, 78-7094
 - Genetics, 78-7094
- Nevus
 - Genetics, 78-7094
- Odontogenic Cysts
 - Genetics, 78-7094
- Skin Neoplasms
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
 - Wounds and Injuries, 78-7092
- Smallpox Vaccine
 - Cicatrix, 78-7089
- Uranium
 - Epidemiology, 78-7179
 - Occupational Hazard, 78-7179
- Wounds and Injuries
 - Case Report, 78-7092

Carcinoma, Bronchogenic

- Celiac Disease
 - Epidemiology, 78-7154

Carcinoma, Ductal

- Breast Neoplasms
 - Cytodiagnosis, 78-7105
 - Epidemiology, 78-7161
 - Neoplasm Metastasis, 78-7161
 - Virus, RNA Tumor, 78-6869
- Pancreatic Neoplasms
 - Ribonuclease, 78-7200

Carcinoma, Ehrlich Tumor

- Ascitic Fluid
 - Anti-Inflammatory Activity, 78-6981

Carcinoma, Epidermoid

- Cervix Neoplasms
 - Virus, Herpes Simplex 2, 78-6945
- Digestive System Neoplasms
 - Virus, Papilloma, 78-6899
- Esophageal Neoplasms
 - Precancerous Conditions, 78-7114
- Head and Neck Neoplasms
 - Radiation, Ionizing, 78-6832
- Intestinal Neoplasms
 - Carrageen, 78-6801
- Mouth Neoplasms
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-7033
- Skin Neoplasms
 - Cholanthrene, 3-Methyl-, 78-7041
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
 - Ultraviolet Rays, 78-6813
 - Virus, Papilloma, 78-6684
 - Xeroderma Pigmentosum, 78-7121
- Sulfide, Bis(2-chloroethyl)-
 - Case Report, 78-7088

- Carcinoma, Epidermoid (cont'd)**
 Neoplasm Recurrence, Local, 78-7088
 Tongue Neoplasms
 Case Report, 78-7098
 Neoplasm Metastasis, 78-7098
 Urea, 1-(2-Chloroethyl)-3-cyclohexyl-1-nitroso-
 78-7098
 Virus, Adeno 5
 Reverse Transcriptase, 78-6967
 Virus Activation, 78-6967
 Virus Replication, 78-6967
 Vulvar Neoplasms
 Epidemiology, 78-7137
 Warts
 Genetics, 78-7036
 Virus, Papilloma, 78-7036
- Carcinoma In Situ**
 Bladder Neoplasms
 Case Report, 78-7139
 Neoplasm Invasiveness, 78-7139
 Precancerous Conditions, 78-7139
 Cervix Neoplasms
 Virus, Herpes Simplex 2, 78-6939
 Esophageal Neoplasms
 Precancerous Conditions, 78-7114
 Vulvar Neoplasms
 Epidemiology, 78-7137
- Carcinoma, Mucinous**
 Stomach Neoplasms
 Histochemical Study, 78-7117
- Carcinoma, Oat Cell**
 Cells, Cultured
 Cell Division, Review, 78-6680
- Carcinoma, Papillary**
 Intestinal Neoplasms
 Bracken Fern, 78-6717
- Carcinoma, Transitional Cell**
 Bladder Neoplasms
 Antigens, Neoplasm, 78-7199
 1-Butanol, 4-(Butylnitrosamino)-, 78-6735
 Cell Line, Epithelioid, 78-7199
 DNA-RNA Hybridization, 78-6973
 Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-
 78-7141
 Neoplasm Invasiveness, 78-7141
 Reverse Transcriptase, 78-6973
 Ribonuclease, 78-6973
Schistosoma haematobium, 78-7140
 Virus, Cytomegalo, 78-7068
 Virus, Equine Rhinopneumonitis
 Thymidine Kinase, 78-6898
 Virus Replication, 78-6898
- Carrageen**
 Carcinoma
 Cytotoxic Activity, 78-6997
 Edema
 Tumor Ascitic Fluid, 78-6981
 Intestinal Neoplasms
 Adenocarcinoma, 78-6801
 Adenoma, 78-6801
 Carcinogenic Activity, Rat, 78-6801
 Carcinoma, Epidermoid, 78-6801
 Lymphoma
 Cytotoxic Activity, 78-6997
- Carrageen (cont'd)**
 Growth, 78-6997
 Macrophages
 Cytotoxic Activity, 78-6997
- Carrier Proteins**
 Hepatoma
 Retinol, 78-6796
 12-*O*-Tetradecanoylphorbol-13-acetate, 78-6796
- Celiac Disease**
 Carcinoma, Bronchogenic
 Epidemiology, 78-7154
 Gynecologic Neoplasms
 Epidemiology, 78-7154
 Hodgkin's Disease
 Epidemiology, 78-7154
 Intestinal Neoplasms
 Adenocarcinoma, 78-7154
 Epidemiology, 78-7154
- Cell Adhesion**
 Cholanthrene, 3-Methyl-
 Cell Transformation, Neoplastic, 78-6908
 Virus, Epstein-Barr
 Ultrastructural Study, Daudi Cells, 78-6955
 Virus, Polyoma
 Tunicamycin, 78-6906
 Virus, SV40
 Cell Transformation, Neoplastic, 78-6908
 Tunicamycin, 78-6906
- Cell Differentiation**
 Adenocarcinoma
 Neoplasm Suppression, Review, 78-6705
 Carcinogen, Environmental
 Mutation, Review, 78-6602
 Giant Cell Tumors
 Cells, Cultured, 78-7148
 Guanidine, 1-Methyl-3-nitro-1-nitroso-
 Organ Culture, Prostate, 78-6751
 Homeostasis
 Cell Transformation, Neoplastic, Review, 78-6705
 Intestinal Neoplasms
 Precancerous Conditions, 78-7131
 Leukemia
 Phenotype, Review, 78-6681
 Precancerous Conditions, Review, 78-6681
 Leukemia, Myeloblastic
 T-Lymphocytes, 78-7150
 Lymphoma
 Phenotype, Review, 78-6681
 Precancerous Conditions, Review, 78-6681
 Mammary Neoplasms, Experimental
 Precancerous Conditions, 78-6776
 Nephroblastoma
 Neoplasm Regression, Review, 78-6686
 Retinoblastoma
 Neoplasm Regression, Review, 78-6686
 Stomach Neoplasms
 Transplantation, Heterologous, 78-7115
 Teratoid Tumor
 Neoplasm Regression, Review, 78-6686
 Neoplasm Suppression, Review, 78-6705
 Virus, Polyoma
 Antigens, Viral, 78-6902
- Cell Division**
 2-Benzimidazolecarbamyl Acid, 5-(2-Thienylcarbonyl)-
 Methyl Ester

Cell Division (cont'd)

- Cells, Cultured, 78-7081
- Cells, Cultured
 - Epithelial Cells, Review, 78-6680
- Colchicine
 - Cells, Cultured, 78-7081
- Colchicine, *N*-Deactyl-*N*-methyl-
 - Cells, Cultured, 78-7081
- Cortisol
 - Embryo, Mouse, 78-7197
- Growth Substances
 - Embryo, Mouse, 78-7197
- Hormones
 - Co-carcinogenic Effect, Review, 78-6643
- Insulin
 - Embryo, Mouse, 78-7197
- Oncogenic Viruses
 - Epithelial Cells, Review, 78-6680
- Prostaglandins F
 - Embryo, Mouse, 78-7197
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Embryo, Mouse, 78-7197
- Vinblastine Sulfate
 - Cells, Cultured, 78-7081
- Vincristine Sulfate
 - Cells, Cultured, 78-7081

Cell Fusion

- Virus, SV40
 - Antigens, Viral, 78-6925

Cell Membrane

- Adenyl Cyclase
 - Cell Transformation, Neoplastic, 78-7188
- Cell Transformation, Neoplastic
 - Peptide Hydrolases, 78-6855
- Glycoproteins
 - Hybrid Cells, 78-7077
- Guanyl Cyclase
 - Cell Transformation, Neoplastic, 78-7188
- Leukemia, Myeloblastic
 - Antigens, 78-7019
- Lymphoma
 - Immune Response, 78-7018
 - Virus, Moloney Murine Leukemia, 78-7018
- Neoplasm Metastasis
 - Phenotype, Review, 78-6687
- Neoplasms
 - Surface Properties, Review, 78-6687
- Peptide Hydrolases
 - Fluorescence, 78-6855
- Tunicamycin
 - Glycoproteins, 78-6906
- Virus, Avian Myeloblastosis
 - Glycoproteins, 78-6844
- Virus, Avian Sarcoma
 - Proteins, 78-6841
- Virus, Epstein-Barr
 - Raji Cells, 78-6949
 - Receptor Activity, 78-6949
 - Viral Proteins, 78-6955
- Virus, Friend Murine Leukemia
 - Antigens, Viral, 78-6879
- Virus, Influenza
 - Histocompatibility Antigens, 78-6909
- Virus, Lymphocytic Choriomeningitis
 - Fluorescence, 78-6855
 - 1-Naphthylamine, *N*-Phenyl-, 78-6855

Cell Membrane (cont'd)

- Pyrene, 78-6855
- Pyrenebutyric Acid, 78-6855
- Virus, Marek's Disease Herpes
 - Antigens, 78-6860
- Virus, Moloney Murine Leukemia
 - Antigens, Viral, 78-7018
 - Nucleotidases, 78-7018
- Virus, Moloney Murine Sarcoma
 - Histocompatibility Antigens, 78-6909
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-6855
 - Fluorescence, 78-6855
 - 1-Naphthylamine, *N*-Phenyl-, 78-6855
 - Pyrene, 78-6855
 - Pyrenebutyric Acid, 78-6855
- Virus, SV40
 - Histocompatibility Antigens, 78-6909

Cell Migration Inhibition

- 2-Benzimidazolecarbamic Acid, 5-(2-Thienylcarbonyl)-
 - Methyl Ester
 - Uracil, 5-Fluoro, 78-7081
- Sarcoma
 - 2-Benzimidazolecarbamic Acid, 5-(2-Thienylcarbonyl)- Methyl Ester, 78-7081
 - Bleomycin, 78-7081
 - Cells, Cultured, 78-7081
 - Colchicine, 78-7081
 - Colchicine, *N*-Deactyl-*N*-methyl-, 78-7081
 - Cytosine, 1- β -*D*-Arabinofuranosyl-, 78-7081
 - Uracil, 5-Fluoro, 78-7081
 - Vinblastine Sulfate, 78-7081
 - Vincristine Sulfate, 78-7081

Cell Movement

- Contact Inhibition
 - Neoplasm Invasiveness, 78-7079
- Virus, SV40
 - Histones, 78-6907

Cell Transformation, Neoplastic

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Organ Culture, Mammary, 78-6745
- Acetamide, *N*-Fluoren-2-yl-
 - Organ Culture, Mammary, 78-6745
- Acetic Acid, (Ethylenedinitrilo)tetra-
 - Proteins, 78-6908
- Acetohydroxamic Acid, *N*-Fluoren-2-yl-
 - Organ Culture, Mammary, 78-6745
- Adenyl Cyclase
 - Cell Membrane, 78-7188
 - Enzymatic Activity, 78-7188
- Antigens
 - Immune Response, Review, 78-6694
- Antigens, Viral
 - Immune Response, Review, 78-6694
- Aryl Hydrocarbon Hydroxylases
 - Liver, Rat, 78-6773
- Benz(a)anthracene, 7,12-Dimethyl-
 - Cells, Cultured, 78-6785
 - T-Lymphocytes, 78-6984
 - Organ Culture, Mammary, 78-6745
- Benzo(a)pyrene
 - Anchorage-Independent Growth, 78-7076
 - Cells, Cultured, 78-6785, 78-7076
 - Fibrinolysis, 78-7076
- Carcinoembryonic Antigen
 - Immune Response, Review, 78-6694

Cell Transformation, Neoplastic (cont'd)

- Carcinogen, Chemical
 - Animal Model, Hamster Embryo, 78-6726
 - Bioassay, Review, 78-6601
 - DNA, Binding, Review, 78-6604
 - Dose-Response Study, Review, 78-6604
 - Mathematical Model, 78-7186
 - Mutagenic Activity, Review, 78-6616
- Cell Membrane
 - Peptide Hydrolases, 78-6855
- Cholanthrene, 11,12-Dihydro-11,12-epoxy-3-methyl-
 - Immunosuppression, 78-7110
- Cholanthrene, 3-Methyl-
 - Cell Adhesion, 78-6908
 - Immunosuppression, 78-7110
 - Proteins, Phosphorylation, 78-6969
- Chondroitin
 - Fibroblasts, 78-7198
 - Glucuronates, 78-7198
- Dibenz(a,h)anthracene, 5,6-Dihydro-5,6-epoxy-
 - Immunosuppression, 78-7110
- Fibroblasts
 - Growth Substances, 78-6928
 - Transplantation, Heterologous, 78-7079
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - Immunosuppression, 78-7110
- Guanyl Cyclase
 - Cell Membrane, 78-7188
 - Enzymatic Activity, 78-7188
- Immunity, Cellular
 - BCG, 78-7006
- Mucopolysaccharides
 - Fibroblasts, 78-7198
- Oncogenic Viruses
 - Fibroblasts, Review, 78-6661
- Radiation, Ionizing
 - Dose-Response Study, 78-6668
 - Theoretical Model, 78-6665
- Trypsin
 - Proteins, 78-6908
- Ultraviolet Rays
 - Psoralen, 8-Methoxy-, 78-6808
- Urea
 - Proteins, 78-6908
- Virus, Abelson Murine Leukemia
 - Immunosuppression, 78-7110
 - Replication-Defective Virus, Review, 78-6678
 - Virus, Helper, Review, 78-6678
- Virus, Adeno 2
 - Chromosomes, 78-6962
 - DNA-RNA Hybridization, 78-6962
- Virus, Adeno 12
 - T-Lymphocytes, 78-6984
 - Proteins, Phosphorylation, 78-6969
- Virus, Avian Leukosis
 - Target Cells, Review, 78-6676
- Virus, Avian Sarcoma
 - Proteins, 78-6841
 - Target Cells, Review, 78-6676
- Virus, B77
 - BCG, 78-7006
- Virus, Epstein-Barr
 - Antigenic Determinants, 78-7015
- Virus, Friend Spleen Focus-Forming
 - Replication-Defective Virus, Review, 78-6678
 - Virus, Helper, Review, 78-6678
- Virus, Herpes Simplex 1

Cell Transformation, Neoplastic (cont'd)

- Antigens, Viral, 78-6944
- DNA, Viral, 78-6944
- Endonucleases, 78-6944
- Fibroblasts, 78-6944
- Virus, Kirsten Murine Sarcoma
 - Ultrastructural Study, Mink Lung Cells, 78-6893
- Virus, Murine Mammary Tumor
 - Genetics, 78-6866
- Virus, Murine Sarcoma
 - Proteins, Phosphorylation, 78-6969
 - Replication-Defective Virus, Review, 78-6678
 - Virus, Helper, Review, 78-6678
- Virus, Polyoma
 - Antigenic Determinants, Review, 78-6671
 - Antigens, Review, 78-6671
 - Glycoproteins, 78-6906
 - Tunicamycin, 78-6906
- Virus, Rat Leukemia
 - Carcinogenic Activity, 78-6895
- Virus, Rauscher Murine Leukemia
 - Glycoproteins, 78-6890
 - Histocompatibility Antigens, 78-6890
- Virus, RNA Tumor
 - Fibroblasts, 78-6928
- Virus, Rous Sarcoma
 - Actin, 78-6846
 - Antigenic Determinants, 78-6853
 - Ascorbic Acid, 78-6849
 - BCG, 78-7006
 - Cell Membrane, 78-6855
 - Chromosomes, 78-6858
 - Microtubules, 78-6846
 - Phosphoproteins, 78-6848
 - RNA, Viral, 78-6859
 - Viral Proteins, 78-6856
- Virus, SV40
 - Antigenic Determinants, Review, 78-6671
 - Antigens, Review, 78-6671
 - Antigens, Viral, 78-6925, 78-6926, 78-6930, 78-6931
 - 78-6932
 - Cell Adhesion, 78-6908
 - Cell Cycle Kinetics, 78-6910
 - Deletion Mutants, 78-6930
 - DNA Replication, 78-6910
 - DNA, Viral, 78-6925, 78-6933
 - Ethidium Bromide, 78-6915
 - Fibroblasts, 78-6928
 - Fibroblasts, Rat, 78-6926
 - Fibroblasts, Review, 78-6661
 - A Gene Product, 78-6910
 - Genes, Viral, Review, 78-6661
 - Glycoproteins, 78-6906
 - Histocompatibility Antigens, 78-6909
 - Hybrid Cells, 78-6914, 78-6919
 - Immunosuppression, 78-7110
 - Isolation and Characterization, 78-6924
 - T-Lymphocytes, 78-6984
 - Temperature Sensitive Mutants, 78-6926, 78-6931
 - Tunicamycin, 78-6906
 - Viral Proteins, 78-6924
- Cells, Cultured
 - Adenocarcinoma
 - Cell Division, Review, 78-6680
 - Aryl Hydrocarbon Hydroxylases
 - Liver, Rat, 78-6773
 - Benz(a)anthracene, 7,12-Dimethyl-

Cells, Cultured (cont'd)

- Cell Transformation, Neoplastic, 78-6785
- 2-Benzimidazolecarbamoyl-5-(2-Thienylcarbonyl)-Methyl Ester
 - Cell Division, 78-7081
- Benzo(a)pyrene
 - Cell Transformation, Neoplastic, 78-6785, 78-7076
- Bladder Neoplasms
 - Reverse Transcriptase, 78-6973
- Cadmium
 - Chromosome Aberrations, 78-6797
- Carcinoma, Oat Cell
 - Cell Division, Review, 78-6680
- Cell Division
 - Epithelial Cells, Review, 78-6680
- Colchicine
 - Cell Division, 78-7081
- Colchicine, *N*-Deacetyl-*N*-methyl-
 - Cell Division, 78-7081
- DNA
 - Homeostasis, 78-7191
 - Nucleic Acid Release, 78-7191
- Fibroblasts
 - Growth Substances, 78-6928
- Fibrosarcoma
 - Benzo(a)pyrene, 78-7076
 - Neoplasm Metastasis, 78-7109
- Giant Cell Tumors
 - Cell Differentiation, 78-7148
 - Chromosomes, 78-7148
 - Lactate Dehydrogenase Isoenzymes, 78-7148
- Hepatoma
 - Australia Antigen, 78-6957
 - Chromosomes, 78-6957
 - Glucosephosphate Dehydrogenase, 78-6957
- Intestinal Neoplasms
 - Adenocarcinoma, 78-6680
- Leukemia
 - Antibody Formation, 78-7021
- Lung Neoplasms
 - Neoplasm Metastasis, 78-7109
- T-Lymphocytes
 - Antitumor Activity, K526 Cells, 78-6987
- Lymphoma
 - Antigens, Viral, 78-7010
 - Histocompatibility Antigens, 78-7010
 - Killer Cells, 78-7010
- Lymphosarcoma
 - Chromosomes, 78-6828
 - Radioresistance, 78-6828
 - Transplantation Immunology, 78-6828
- Mammary Neoplasms, Experimental
 - Neoplasm Metastasis, 78-7109
- Melanoma
 - Cell Division, Review, 78-6680
- Mutagens
 - Colony Growth and Morphology, 78-7196
- Neoplasm Metastasis
 - Bromo Cyclic AMP, 78-7109
- Radiation, Ionizing
 - DNA Repair, 78-6812
- Sarcoma
 - Cell Migration Inhibition, 78-7081
- Sarcoma, Osteogenic
 - Chromosomes, 78-7148
 - Lactate Dehydrogenase Isoenzymes, 78-7148
- 12-*O*-Tetradecanoylphorbol-13-acetate

Cells, Cultured (cont'd)

- Mutagenic Activity, Review, 78-6606
- Ultraviolet Rays
 - DNA Repair, 78-6812
- Vinblastine Sulfate
 - Cell Division, 78-7081
- Vincristine Sulfate
 - Cell Division, 78-7081
- Virus, Equine Rhinopneumonitis
 - Virus Replication, 78-6898
- Virus, Leukemia
 - B-Lymphocytes, 78-6975
- Virus, Murine Mammary Tumor
 - Virus-Like Particles, 78-7104
 - Virus Replication, 78-6863
- Virus, Rauscher Murine Leukemia
 - Antibody Formation, 78-7021
- Virus, Rous Sarcoma
 - Chromosomes, 78-6858
- Virus, SV40
 - Virus Replication, 78-6912
- Cellular Inclusions**
 - Leukemia, Lymphoblastic
 - Ultrastructural Study, 78-6977
 - Nasopharyngeal Neoplasms
 - Carcinoma, 78-7101
 - Ultrastructural Study, 78-7101
 - Virus, Epstein-Barr, 78-7101
- Cellulose Tetranitrate**
 - Neoplasms, Experimental
 - Macrophages, 78-6991
- Cerebral Ischemia, Transient**
 - Autoimmune Diseases
 - Stress, Review, 78-6682
 - Neoplasms
 - Stress, Review, 78-6682
- Cerebral Vasospasm**
 - see Cerebral Ischemia, Transient
- Cervix Neoplasms**
 - Antigens, Bacterial
 - Antibodies, Viral, 78-7065
 - Carcinoma
 - Virus, Herpes Simplex 2, 78-6939
 - Carcinoma, Epidermoid
 - Virus, Herpes Simplex 2, 78-6945
 - Carcinoma In Situ
 - Virus, Herpes Simplex 2, 78-6939
 - Condylomata Acuminata
 - Ultrastructural Study, 78-7136
 - Virus-Like Particles, 78-7136
 - Contraceptives, Oral
 - Diagnosis, Review, 78-6648
 - Epidemiology, Review, 78-6648
 - Precancerous Conditions, 78-6648
 - Epidemiology
 - Iran, 78-7153
 - Ethyl Alcohol
 - Epidemiology, Yugoslavia, 78-6945
 - Ethylene, Trichloro-
 - Epidemiology, 78-7173
 - Smoking
 - Epidemiology, Yugoslavia, 78-6945
 - Virus, Herpes Simplex 1
 - Antibodies, Viral, 78-6939
 - Lymphocyte Transformation, 78-7065

- Cervix Neoplasms (cont'd)**
 Virus, Herpes Simplex 2
 Antibodies, Viral, 78-6939, 78-7065
 Carcinogenic Activity, Review, 78-6683
 Epidemiology, Yugoslavia, 78-6945
 Immunity, Cellular, 78-7065
 Lymphocyte Transformation, 78-7065
 Precancerous Conditions, 78-6939, 78-7065
 Seroepidemiology, 78-6939, 78-6945
- Cesium Radioisotopes**
 Hyperplasia
 Spleen, Mouse, Rat, 78-6817
- Charcot-Marie Disease**
 see Muscular Atrophy
- Chemotaxis**
 Lung Neoplasms
 Macrophages, 78-6999
 Neoplasms, Experimental
 Benz(a)anthracene, 7,12-Dimethyl-, 78-6991
 Macrophages, 78-6991
 Monocytes, 78-6991
- Chlorambucil**
 see Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
- Chloroma**
 see Leukemia, Myelocytic
- Cholangioma**
 Alpha Fetoproteins
 Serum Levels, 78-7124
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 Adenosine Cyclic 3',5' Monophosphate, 78-6766
 Adenyl Cyclase, 78-6766
 Guanosine Cyclic 3',5' Monophosphate, 78-6767
 Guanyl Cyclase, 78-6767
 Phosphodiesterases, 78-6766
 Precancerous Conditions, 78-6767
 Bile Duct Neoplasms
 Epidemiology, Japan, 78-7127
 Liver Neoplasms
 Acetophenone, 2-Amino-, 78-6790
 Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6767
 Indole-3-acrylic Acid, 78-6790
- Cholanthrene, 11,12-Dihydro-11,12-epoxy-3-methyl-**
 Immunosuppression
 Cell Transformation, Neoplastic, 78-7110
- Cholanthrene, 3-Methyl-**
 Aryl Hydrocarbon Hydroxylases
 Liver, Mouse, 78-7041
 Liver, Rat, 78-6773
 BCG
 Immune Response, Mouse, 78-7007
 Cell Adhesion
 Cell Transformation, Neoplastic, 78-6908
 Cell Transformation, Neoplastic
 Proteins, Phosphorylation, 78-6969
 Chromatin
 Proteins, Phosphorylation, 78-6969
 Fibrosarcoma
 Immunization, 78-7053
 Virus, Rat Leukemia, 78-6873
 Freund's Adjuvant
 Immune Response, Mouse, 78-7007
 Glomerulonephritis
- Cholanthrene, 3-Methyl- (cont'd)**
 Mouse, New Zealand Black, 78-7041
 Hormones
 Co-carcinogenic Effect, 78-6798
 Immune Response
 Mouse, New Zealand Black, 78-7041
 Immunosuppression
 Cell Transformation, Neoplastic, 78-7110
 L Cells
 Proteins, 78-6908
 Lymphocytes
 Aryl Hydrocarbon Hydroxylases, 78-6781
 Cytochrome Reductases, 78-6781
 T-Lymphocytes
 Antilymphocyte Serum, 78-7041
 Mitogens
 Aryl Hydrocarbon Hydroxylases, 78-6781
 Neoplasms, Experimental
 Histocompatibility Antigens, 78-7044
 Plant Agglutinins
 Aryl Hydrocarbon Hydroxylases, 78-6781
 Quinoline
 Ames Test, 78-6762
 Ribosomes
 Proteins, Phosphorylation, 78-6969
 Sarcoma
 Antigen-Antibody Complex, 78-7042
 Histocompatibility Antigens, 78-7043
 Macrophages, 78-6994
 Neoplasm Metastasis, 78-7043
 Skin Neoplasms
 Antilymphocyte Serum, 78-7041
 Calcitonin, 78-6798
 Carcinoma, Epidermoid, 78-7041
 Cortisol, 78-6798
 Mouse, New Zealand Black, 78-7041
 Precancerous Conditions, 78-6798
 Prostaglandins E, 78-6798
 Radiation, Ionizing, 78-7041
 Sarcoma, 78-7041
 Thyroxine, 78-6798
 Virus, Rat Leukemia
 Virus Activation, 78-6873
- Cholesterol**
 Colonic Neoplasms
 Diet, 78-7180
 Diet
 Metabolism, 78-7180
- Choline**
 Liver Neoplasms
 Carcinogen, Chemical, 78-6638
- Chondroitin**
 Contact Inhibition
 Fibroblasts, 78-7198
 Cortisol
 Fibroblasts, 78-7198
 Fibroblasts
 Cell Transformation, Neoplastic, 78-7198
 Glucuronates
 Cell Transformation, Neoplastic, 78-7198
- Chondroitin Sulfate**
 see Chondroitin
- Chondroitinases**
 Disaccharides
 Fibroblasts, 78-7198

Choriocarcinoma

- Hydatidiform Mole
Case Report, 78-7134

Lung

- Neoplasm Metastasis, 78-7134
- Neoplasm Metastasis
Case Report, 78-7134

Chromate

- Respiratory Tract Neoplasms
Epidemiology, Review, 78-6634

Chromatids

- Ethylene, Chloro-
Occupational Hazard, 78-6712
- Psoralen, 8-Methoxy-
Hamster Ovary Cells, 78-6810
- Ultraviolet Rays, 78-6810
- Ultraviolet Rays
Hamster Ovary Cells, 78-6810

Chromatin

- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
DNA, Binding, 78-6784
Nucleic Acid Conformation, 78-6784
- Cholanthrene, 3-Methyl-
Proteins, Phosphorylation, 78-6969
- Virus, Adeno 12
Proteins, Phosphorylation, 78-6969
- Virus, Murine Sarcoma
Proteins, Phosphorylation, 78-6969

Chromosome Aberrations

- Acetamide, *N*-Fluoren-2-yl-
Carcinogenic Potential, Review, 78-6618
- Ataxia Telangiectasia
Fibroblasts, Review, 78-6612
- Benzene
Epidemiology, Review, 78-6627
- Cadmium
Cells, Cultured, 78-6797
Culture Media, 78-6797
- Contraceptives, Oral
Carcinogenic Potential, Review, 78-6618
- Cosmetics
Carcinogenic Potential, Review, 78-6618
- Dimethylamine, *N*-Nitroso-
Metabolism, 78-6802
- Drugs
Carcinogenic Potential, Review, 78-6618
- Food Additives
Carcinogenic Potential, Review, 78-6618
- Guanidine, Methyl-
Metabolism, 78-6802
- Leukemia
Precancerous Conditions, Review, 78-6681
- Leukemia, Lymphoblastic
Ataxia Telangiectasia, 78-7030
- Lymphoma
Precancerous Conditions, 78-6695
- Mutagens
Carcinogenic Potential, Review, 78-6618
- Pesticides
Carcinogenic Potential, Review, 78-6618
- Sarcoma, Reticulum Cell
Case Report, 78-7080
Chromosomes, Human, 13-15, 78-7080
- Urea, Ethyl Nitroso-

Chromosome Aberrations (cont'd)

- Mutagenic Activity, Review, 78-6656
- Virus, DNA Tumor
Carcinogenic Activity, Review, 78-6670
- Virus, RNA Tumor
Carcinogenic Activity, Review, 78-6670

Chromosome Abnormalities

- Aneuploidy
Histological Study, Rat, 78-7143
- Asbestos
Fibers, 78-6836
Hamster Cells, 78-6836
- Carcinogen, Chemical
Carcinogenic Activity, Review, 78-6610
- Carcinogen, Environmental
Cell Transformation, Neoplastic, Review, 78-6602
- Disgerminoma
Case Report, 78-7138
- Ethylene, Chloro-
Occupational Hazard, 78-6712, 78-6715
Smoking, 78-6715
- Ethylene, Chloro- Polymer
Occupational Hazard, 78-6715
Smoking, 78-6715
- Fiber Glass
Hamster Cells, 78-6836
- Hereditary Diseases
Cell Transformation, Neoplastic, Review, 78-6602
- Leukemia
Radiation, Ionizing, 78-7072
- Leukemia, Myelocytic
Aneuploidy, 78-7070
- Mycosis Fungoides
Chromosomes, Human, 1-3, 78-7120
B-Lymphocytes, 78-7120
T-Lymphocytes, 78-7120
- Oncogenic Viruses
Carcinogenic Activity, Review, 78-6610
- Radiation
Carcinogenic Activity, Review, 78-6610
- Testicular Neoplasms
Disgerminoma, 78-7145
Histological Study, Rat, 78-7143

Chromosomes

- Giant Cell Tumors
Cells, Cultured, 78-7148
- Hepatoma
Cells, Cultured, 78-6957
- Lymphosarcoma
Cells, Cultured, 78-6828
- Sarcoma, Osteogenic
Cells, Cultured, 78-7148
- Virus, Adeno 2
Cell Transformation, Neoplastic, 78-6962
DNA, Viral, 78-6962
RNA, Messenger, 78-6964
- Virus, Rous Sarcoma
Cell Transformation, Neoplastic, 78-6858
Cells, Cultured, 78-6858
Chromatin Bodies, 78-6858
Metaphase, 78-6858
- Virus, SV40
2,3-Butanediol, 1,4-Dimercapto-, 78-6907
DNA Replication, 78-6913
Endonucleases, 78-6907
Histones, 78-6907, 78-6923

Chromosomes (cont'd)

- Hybrid Cells, 78-6914
- Nucleoproteins, 78-6918
- Virus Replication, 78-6918

Chromosomes, Human, 1-3

- Mycosis Fungoides
- Chromosome Abnormalities, 78-7120

Chromosomes, Human, 6-12

- Glycoproteins
- Hybrid Cells, 78-7077
- LETS Protein, 78-7077
- Virus, SV40
- DNA, Viral, 78-6919, 78-6929
- Hybrid Cells, 78-6919, 78-6929

Chromosomes, Human, 13-15

- Leukemia, Lymphoblastic
- Ataxia Telangiectasia, 78-7030
- Sarcoma, Reticulum Cell
- Chromosome Aberrations, 78-7080

Chromosomes, Human, 16-18

- Virus, SV40
- DNA, Viral, 78-6919
- Hybrid Cells, 78-6919

Chromosomes, Human, 21-22

- Leukemia, Myelocytic
- Karyotyping, 78-7070
- Radiation, Ionizing, 78-6827

Cicatrix

- Carcinoma, Basal Cell
- Case Report, 78-7089
- Smallpox Vaccine, 78-7089

Cockayne's Syndrome

- see Dwarfism

Colchicine

- Cell Division
- Cells, Cultured, 78-7081
- Sarcoma
- Cell Migration Inhibition, 78-7081

Colchicine, *N*-Deacetyl-*N*-methyl-

- Cell Division
- Cells, Cultured, 78-7081
- Sarcoma
- Cell Migration Inhibition, 78-7081

Colitis, Ulcerative

- Polyps
- Case Report, 78-7130
- Colon, 78-7130

Collagen

- Skin Neoplasms
- Precancerous Conditions, 78-7087

Colon

- Polyps
- Colitis, Granulomatous, 78-7130
- Colitis, Ulcerative, 78-7130

Colonic Neoplasms

- Adenocarcinoma
- Genetics, 78-7118
- Diet
- Bacteria, 78-7180
- Cholesterol, 78-7180
- Epidemiology, Review, 78-6637

Colonic Neoplasms (cont'd)

- Dietary Fats
- Epidemiology, Review, 78-6699
- Virus, Marek's Disease Herpes
- Antibodies, Viral, 78-6861

Complement

- Fibrosarcoma
- Growth, 78-7053
- Hepatosoma
- Aniline, *N,N*-Dimethyl-*p*-phenylazo-, 78-7063
- Antigen-Antibody Complex, 78-7064
- Lymph Nodes
- Transplantation Immunology, 78-7044
- Macrophages
- Cleavage Product, Isolation and Characterization
- 78-7002
- Cytotoxic Activity, 78-7002

Concanavalin A

- Bladder Neoplasms
- Lymphocytes, 78-6735
- Ethanol, 2-Mercapto-
- Antitumor Activity, 78-6988
- Lymphocytes
- Cytotoxicity, 78-6988
- Virus, Marek's Disease Herpes
- Immune Response, 78-7049
- Lymphocyte Transformation, 78-7049
- T-Lymphocytes, 78-7050
- Virus, Rauscher Murine Leukemia
- Lymphocyte Transformation, 78-7017

Condylomata Acuminata

- Cervix Neoplasms
- Ultrastructural Study, 78-7136
- Virus-Like Particles, 78-7136
- Virus-Like Particles
- Ultrastructural Study, 78-7136
- Virus, Papilloma
- Virus-Like Particles, 78-7136
- Virus, Papova
- Virus-Like Particles, 78-7136

Connective Tissue

- Skin Neoplasms
- Mitosis, 78-7087
- Precancerous Conditions, 78-7087

Contact Inhibition

- Cell Movement
- Neoplasm Invasiveness, 78-7079
- Chondroitin
- Fibroblasts, 78-7198
- Neuroblastoma
- Neoplasm Transplantation, 78-7079
- Sarcoma 180, Crocker
- Neoplasm Transplantation, 78-7079

Contraceptives, Oral

- Aminotransferases
- Liver Function, Review, 78-6650
- Breast Diseases
- Cysts, 78-7107
- Breast Neoplasms
- Adenofibroma, 78-6645
- Epidemiology, Review, 78-6645, 78-6648
- Hyperplasia, 78-6645
- Precancerous Conditions, 78-6648, 78-7107
- Ultrastructural Study, Review, 78-6645

Contraceptives, Oral (cont'd)

- Cervix Neoplasms
 - Diagnosis, Review, 78-6648
 - Epidemiology, Review, 78-6648
 - Precancerous Conditions, 78-6648
- Chromosome Aberrations
 - Carcinogenic Potential, Review, 78-6618
- Gynecologic Neoplasms
 - Adenocarcinoma, 78-6645
 - Epidemiology, Review, 78-6645
 - Myoma, 78-6645
 - Ultrastructural Study, Review, 78-6645
- Hepatoma
 - Case Report, 78-6795
 - Epidemiology, Review, 78-6645, 78-6650
- Hyperplasia
 - Epidemiology, Review, 78-6650
- Liver Neoplasms
 - Adenoma, 78-6645, 78-6650, 78-6795, 78-7126
 - Case Report, 78-6795, 78-7125, 78-7126
 - Epidemiology, Review, 78-6644, 78-6645, 78-6646, 78-6650
 - Estradiol, 17-Ethynyl-, 78-7126
 - Hyperplasia, 78-6645, 78-6650, 78-6795, 78-7125
 - Norethisterone, 78-7126
 - Precancerous Conditions, Review, 78-6646
 - Tumor Hemorrhage, 78-7126
 - Ultrastructural Study, 78-7125
 - Ultrastructural Study, Review, 78-6645
- Phthalide, 4,5,6,7-Tetrabromo-3,3-bis(4-hydroxy-3-sulphophenyl)-
 - Liver Function, Review, 78-6650
- Pituitary Neoplasms
 - Adenoma, 78-7095
 - Prolactin, 78-7095

Articortropin

- Lung Neoplasms
 - Isolation and Characterization, 78-6793

Cortisol

- Cell Division
 - Embryo, Mouse, 78-7197
- Chondroitin
 - Fibroblasts, 78-7198
- Leukemia, Lymphoblastic
 - Macrophages, 78-6993
- Skin Neoplasms
 - Cholanthrene, 3-Methyl-, 78-6798

Cortisone Acetate

- Lung Neoplasms
 - Neoplasm Metastasis, 78-6998
 - Phagocytosis, 78-6998
- Neoplasms, Experimental
 - Virus, SV40, 78-6911

Corynebacterium parvum

- Fibrosarcoma
 - Antitumor Activity, 78-7056
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-7055
 - T-Lymphocytes, 78-7056
 - Macrophage Activation, 78-7056
 - Macrophages, 78-7054
 - Transplantation Immunology, 78-7055
- Lung Neoplasms
 - Neoplasm Metastasis, 78-6998
 - Phagocytosis, 78-6998
- Macrophages

Corynebacterium parvum (cont'd)

- Phagocytosis, 78-7057
- Mammary Neoplasms, Experimental
 - Antitumor Activity, 78-7057
 - Radiation, Ionizing, 78-7057
 - Silica, 78-7057
 - Transplantation Immunology, 78-7057
- Radiation, Ionizing
 - Antibody Formation, 78-7057
 - Antitumor Activity, 78-7057
 - Phagocytosis, 78-7057
- Silica
 - Antibody Formation, 78-7057
 - Antitumor Activity, 78-7057
 - Phagocytosis, 78-7057

Cosmetics

- Chromosome Aberrations
 - Carcinogenic Potential, Review, 78-6618

Coumarin, 3-Benzyl-7-hydroxy-4-methyl-

- Ultraviolet Rays
 - Occupational Hazard, 78-6663
 - Textile Workers, Review, 78-6663

p-Cresol, 2,6-Di-tert-butyl-

- Acetohydroxamic Acid, N-Fluorenyl-
 - Mutagenic Activity, 78-6730
- Esterases
 - Plasma Levels, Mouse, 78-6743
- Skin Neoplasms
 - Ultraviolet Rays, 78-6806

Croton Oil

- Benz(a)anthracene, 7,12-Dimethyl-
 - Two-Stage Carcinogenesis, Review, 78-6626
- Benzo(a)pyrene
 - Two-Stage Carcinogenesis, Review, 78-6626
- Skin Neoplasms
 - 1,2-Benzisothiazolin-3-one, 1,1-Dioxide, 78-6660

Curium Oxide

- Bone and Bones
 - Inhalation Study, Rat, 78-6834
- Lung Clearance
 - Inhalation Study, Rat, 78-6834
- Lung Neoplasms
 - Inhalation Study, Rat, 78-6820
 - Radioisotopes, 78-6820

Cycloheximide

- Virus, Adeno 2
 - Peptides, 78-6963

Cyclophosphamide

- Leukemia
 - Neoplasms, Multiple Primary, 78-7072
- Leukemia, Myeloblastic
 - Drug Therapy, 78-6815
- Neoplasms, Experimental
 - Virus, SV40, 78-6911
- Virus, Marek's Disease Herpes
 - Immunity, Passive, 78-7023

Cysteine

- Dimethylamine, N-Nitroso-
 - Metabolism, 78-6802
- Guanidine, Methyl-
 - Metabolism, 78-6802
- Radiation, Ionizing
 - Lymphocytes, 78-6979

Cysts

- Breast Diseases
 - Contraceptives, Oral, 78-7107
- Breast Neoplasms
 - Cytodiagnosis, 78-7105
- Ovarian Neoplasms
 - Adenocarcinoma, 78-7132
 - Endometriosis, 78-7132

Cytochrome Oxidase

- Aniline, *N,N*-Diethyl-*p*-phenylazo-
 - Mitochondria, Liver, 78-6764
- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 - Mitochondria, Liver, 78-6764

Cytochrome P-450

- Benzene, Bromo-
 - Hepatotoxicity, Review, 78-6607
- Nucleic Acids, Binding
 - Hepatotoxicity, Review, 78-6607

Cytochrome Reductases

- Aryl Hydrocarbon Hydroxylases
 - Quantitation Method, 78-6781
- Cholanthrene, 3-Methyl-
 - Lymphocytes, 78-6781

Cytosine, 1- β -D-Arabinofuranosyl-

- Sarcoma
 - Cell Migration Inhibition, 78-7081
- Virus, Adeno 2
 - Peptides, 78-6963
- Virus, Adeno 12
 - Antigens, Viral, 78-6970

Cytosol

- Virus, SV40
 - DNA Replication, 78-6913

Desmosomes

- Bladder Neoplasms
 - Neoplasm Invasiveness, 78-7141
 - Ultrastructural Study, 78-7141

Deuterium

- Dimethylamine, *N*-Nitroso-
 - DNA Sedimentation, 78-6746

Dexamethasone

- Virus, Murine Mammary Tumor
 - RNA Replication, 78-6863

Diabetes Mellitus

- Breast Neoplasms
 - Epidemiology, Sweden, 78-7106

Dibenz(a,h)anthracene, 5,6-Dihydro-5,6-epoxy-

- Immunosuppression
 - Cell Transformation, Neoplastic, 78-7110

Dibutyl Cyclic AMP

- Leukemia, Lymphoblastic
 - Macrophages, 78-6993

Dieldrin

- Liver Neoplasms
 - Epidemiology, Review, 78-6659

Diet

- Bacteria
 - Gastrointestinal System, 78-7180
- Breast Neoplasms
 - Epidemiology, Review, 78-6637
- Cholesterol

Diet (cont'd)

- Metabolism, 78-7180
- Colonic Neoplasms
 - Bacteria, 78-7180
 - Cholesterol, 78-7180
 - Epidemiology, Review, 78-6637
- Esophageal Neoplasms
 - Epidemiology, Review, 78-6637
- Intestinal Neoplasms
 - Epidemiology, 78-7176
 - Vegetables, 78-7176
- Liver Neoplasms
 - Epidemiology, Review, 78-6637
- Stomach Neoplasms
 - Epidemiology, Review, 78-6637

Dietary Fats

- Breast Neoplasms
 - Epidemiology, Review, 78-6699
- Colonic Neoplasms
 - Epidemiology, Review, 78-6699
- Leukemia
 - Epidemiology, Review, 78-6699

Dietary Proteins

- Kidney Neoplasms
 - Dimethylamine, *N*-Nitroso-, 78-6742
 - Streptozotocin, 78-6799
- Liver Neoplasms
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6742
 - Diethylamine, *N*-Nitroso-, 78-6742

Diethylamine

- Liver
 - Tissue Distribution, Guinea Pig, 78-6747
- Lung
 - Tissue Distribution, Guinea Pig, 78-6747

Diethylamine, 2,2'-Dichloro-*N*-methyl-

- Haemophilus influenzae*
 - DNA Repair, 78-6759
- Leukemia
 - Neoplasms, Multiple Primary, 78-7072
- Skin Neoplasms
 - Carcinoma, Basal Cell, 78-6753
 - Carcinoma, Epidermoid, 78-6753
 - Keratoacanthoma, 78-6753
 - Mycosis Fungoides, 78-6753
 - Neoplasms, Multiple Primary, 78-6753

Diethylamine, *N*-Nitroso-

- Carbon Tetrachloride
 - Co-carcinogenic Effect, 78-6750
- Esterases
 - Plasma Levels, Mouse, 78-6743
- Hepatoma
 - Carbon Tetrachloride, 78-6750
 - DNA Replication, 78-6744
 - Liver Regeneration, 78-6750
 - Neoplasm Metastasis, 78-7122
 - Precancerous Conditions, 78-6744
- Liver Neoplasms
 - Benzidine, Dihydrochloride, 78-7122
 - Carboxy-Lyases, 78-6742
 - Dietary Proteins, 78-6742
- Receptors, Adrenergic
 - Atria, Rat, Guinea Pig, 78-6732

Digestive System Neoplasms

- Bracken Fern

estive System Neoplasms (cont'd)

- Epidemiology, Cattle, 78-6899
- Carcinoma, Epidermoid
 - Virus, Papilloma, 78-6899
- Ethnic Groups
 - Epidemiology, England, Wales, 78-7157
 - Epidemiology, Review, 78-6698
- Ethylene, Trichloro-
 - Epidemiology, 78-7173
- Socioeconomic Factors
 - Epidemiology, Review, 78-6698
- Uterine Neoplasms
 - Case Report, 78-7119
 - Peutz-Jeghers Syndrome, 78-7119
- Virus, Papilloma
 - Epidemiology, Cattle, 78-6899
 - Precancerous Conditions, 78-6899
- methylaniline**
 - Liver
 - Tissue Distribution, Guinea Pig, 78-6747
 - Lung
 - Tissue Distribution, Guinea Pig, 78-6747
- methylaniline, N-Nitroso-**
 - Ascorbic Acid, Monosodium Salt
 - Metabolism, 78-6802
 - Chromosome Aberrations
 - Metabolism, 78-6802
 - Cysteine
 - Metabolism, 78-6802
 - Deuterium
 - DNA Sedimentation, 78-6746
 - DNA Repair
 - Metabolism, 78-6802
 - DNA Sedimentation
 - Liver, Rat, 78-6746
 - Ethanethiol, 2-Amino-
 - Metabolism, 78-6802
 - Gallic Acid, Propyl Ester
 - Metabolism, 78-6802
 - Hepatoma
 - DNA Replication, 78-6744
 - Kidney Neoplasms
 - Carboxy-Lyases, 78-6742
 - Dietary Proteins, 78-6742
 - Fibrosarcoma, 78-6739
 - Histological Study, 78-6739
 - Neoplasm Transplantation, 78-6739
 - Ultrastructural Study, 78-6739
 - Maternal-Fetal Exchange
 - Tissue Distribution, Mouse, 78-6740
 - Purine, 2-Amino-6-methoxy-
 - DNA, Liver, 78-6625
 - Enzymes, 78-6625
 - Tissue Distribution, Mouse
 - Autoradiography, 78-6740
- propylamine, N-Nitroso-**
 - Esterases
 - Plasma Levels, Mouse, 78-6743
- Direct Black 38**
 - Hepatoma
 - Carcinogenic Activity, Rat, 78-6771
 - Liver Neoplasms
 - Precancerous Conditions, Mouse, 78-6771
 - Occupational Hazard
 - Textile Workers, Review, 78-6663

Direct Blue 6

- Hepatoma
 - Carcinogenic Activity, Rat, 78-6771
- Liver Neoplasms
 - Precancerous Conditions, Mouse, 78-6771
- Occupational Hazard
 - Textile Workers, Review, 78-6663

Direct Brown 95

- Hepatoma
 - Carcinogenic Activity, Rat, 78-6771
- Liver Neoplasms
 - Precancerous Conditions, Mouse, 78-6771

Disaccharides

- Chondroitinases
- Fibroblasts, 78-7198

Disgerminoma

- Chromosome Abnormalities
 - Case Report, 78-7138
- Testicular Neoplasms
 - Case Report, 78-7145
 - Chromosome Abnormalities, 78-7145
 - Histological Study, 78-7146

Disulfide, Bis(diethylthiocarbamoyl)-

- Angiosarcoma
 - Ethane, 1,2-Dibromo-, 78-6714
- Hepatoma
 - Ethane, 1,2-Dibromo-, 78-6714
- Skin Neoplasms
 - Ultraviolet Rays, 78-6806

Disulfiram

- see Disulfide, Bis(diethylthiocarbamoyl)-

Diterpene Esters

- Prostaglandins
 - Macrophages, 78-6719

DNA

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Liver, Rat, 78-6716
- Aryl Hydrocarbon Hydroxylases
 - Quantitation Method, 78-6781
- Blood Vessels
 - Carcinogenic Potential, 78-7191
- 2-Butanone, 1,1-Dihydroxy-3-ethoxy-
 - Liver, Rat, 78-6716
- Carcinogen, Chemical
 - Mutagenic Activity, Review, 78-6616
- Carcinogen, Environmental
 - Mutagenic Activity, Review, 78-6609
- Cells, Cultured
 - Nucleic Acid Release, 78-7191
- Homeostasis
 - Cells, Cultured, 78-7191
- Lymphocytes
 - Nucleic Acid Release, 78-7191
- Nucleic Acid Release
 - Frog Auricles, 78-7191
- Terbium
 - Fluorescence, 78-6716

DNA, Bacterial

- Mutagens
 - Colony Growth and Morphology, 78-7196

DNA Nucleotidyltransferases

- Leukemia, Radiation-Induced
 - Bone Marrow, 78-6826

DNA Nucleotidyltransferases (cont'd)

- Enzymatic Activity, Mouse, 78-6826
- Radiation, Ionizing, 78-6826
- Thymus Gland, 78-6826

Lymphoma

- Virus, Radiation Leukemia, 78-7029

DNA Polymerase**Bladder Neoplasms**

- Actinomycin D, 78-6973
- Ribonuclease, 78-6973

Escherichia coli

- DNA Repair, Review, 78-6620

Mutation

- Hereditary Diseases, Review, 78-6704

Ultraviolet Rays

- Mutagenic Activity, Review, 78-6704
- Nucleotide Sequence, Review, 78-6704

Virus, Hepatitis

- Isolation and Characterization, Woodchuck
78-6959

Virus, Rauscher Murine Leukemia

- Temperature Sensitive Mutants, 78-6885
- Thermolability, 78-6885

Virus, Rous Sarcoma

- Temperature Sensitive Mutants, 78-6851

DNA Repair**Anthramycin**

- Saccharomyces cerevisiae*, 78-6769

Antibiotics, Antineoplastic

- Saccharomyces cerevisiae*, 78-6769

Benzo(a)pyrene

- Cell Cycle Kinetics, 78-6785

Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester

- Haemophilus influenzae*, 78-6759

Carcinogen, Chemical

- Bioassay, Review, 78-6608
- Carcinogenic Activity, Review, 78-6608, 78-6619
- Carcinogenic/Mutagenic Activity, 78-6711
- Cell Transformation, Neoplastic, Review, 78-6613
- In Vivo Measurement Methods, Review, 78-6619

Diethylamine, 2,2'-Dichloro-*N*-methyl-

- Haemophilus influenzae*, 78-6759

Dimethylamine, *N*-Nitroso-

- Metabolism, 78-6802

Guanidine, Methyl-

- Metabolism, 78-6802

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Carcinogenic Activity, Review, 78-6608, 78-6622
- Haemophilus influenzae*, 78-6759

Hereditary Diseases

- Cell Transformation, Neoplastic, Review, 78-6613

Keratinosis

- Aging, 78-7091
- Leukocytes, 78-7091
- Lymphocytes, 78-7090
- Ultraviolet Rays, 78-7090
- Urea, Hydroxy-, 78-7090

Leukemia L1210

- Ultraviolet Rays, 78-6807

Methanesulfonic Acid, Ethyl Ester

- Haemophilus influenzae*, 78-6759

Mitomycin C

- Haemophilus influenzae*, 78-6759

Mutagens

- Escherichia coli*, 78-7193
- Quantitation Method, 78-7193

DNA Repair (cont'd)**Polycyclic Hydrocarbons**

- Carcinogenic Activity, Review, 78-6622
- Quinoline, 4-Nitro-, 1-Oxide

- Carcinogenic Activity, Review, 78-6608

Radiation, Ionizing

- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester, 78-6759
- Carcinogenic Activity, Review, 78-6619
- Cell Transformation, Neoplastic, Review, 78-6613
- Cells, Cultured, 78-6812
- Haemophilus influenzae*, 78-6759
- In Vivo Measurement Methods, Review, 78-6619
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-, 78-6812
- Saccharomyces cerevisiae*, 78-6769

Sibromycin

- Saccharomyces cerevisiae*, 78-6769

Skin Neoplasms

- Xeroderma Pigmentosum, 78-7121

Tomaymycin

- Saccharomyces cerevisiae*, 78-6769

Ultraviolet Rays

- 1,8,9-Anthracenetriol, 78-6807
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester, 78-6759
- Carcinogenic Activity, Review, 78-6619
- Cell Transformation, Neoplastic, Review, 78-6613
- Cells, Cultured, 78-6812
- Escherichia coli*, 78-6805
- Haemophilus influenzae*, 78-6759
- In Vivo Measurement Methods, Review, 78-6619
- Keratinosis, 78-7091
- Leukocytes, 78-7091
- Pyrimidine Dimers, Review, 78-6666
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-, 78-6812
- Quinacrine, 78-6807
- Saccharomyces cerevisiae*, 78-6769
- Species Difference, Review, 78-6666
- Tryptophan, 78-6805
- Xeroderma Pigmentosum, 78-7121
- Urea, Methyl Nitroso-
- Carcinogenic Activity, Review, 78-6622
- Virus, Herpes Simplex 1
- Ultraviolet Rays, 78-6941

DNA Replication**Cell Cycle Kinetics**

- Cell Transformation, Neoplastic, Review, 78-6703

Hepatoma

- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6744
- Diethylamine, *N*-Nitroso-, 78-6744
- Dimethylamine, *N*-Nitroso-, 78-6744
- Hyperplasia, 78-6729
- Precancerous Conditions, 78-6744

Hodgkin's Disease

- Autoradiographic Study, 78-7192
- Histological Study, 78-7192
- Lymph Nodes, 78-7192

Leukemia

- Immunosuppression, 78-7022

Lymphocytes

- Interferon, 78-6982

Mammary Neoplasms, Experimental

- Plant Agglutinins, 78-7004

2-Propanol, 1,1'-Iminodi-*N*-nitroso-

- Brain, Rat, 78-6749

Quinoline, 4-(Hydroxyamino)-, 1-Oxide

NA Replication (cont'd)

- Brain, Rat, 78-6749
- Urea, Ethyl Nitroso-
 - Brain, Rat, 78-6749
- Virus, Epstein-Barr
 - Carcinogenic Activity, Review, 78-6683
- Virus, Herpes
 - Carcinogenic Activity, Review, 78-6683
- Virus, Herpes Simplex 1
 - Genes, Viral, 78-6943
 - Temperature Sensitive Mutants, 78-6943
- Virus, Herpes Simplex 2
 - Carcinogenic Activity, Review, 78-6683
- Virus, SV40
 - Cell Nucleus, 78-6913
 - Cell Transformation, Neoplastic, 78-6910
 - Chromosomes, 78-6913
 - Cytosol, 78-6913
 - Deletion Mutants, 78-6917
 - DNA, Viral, 78-6922
 - Endonucleases, 78-6927
 - Ethidium Bromide, 78-6915
 - Nucleoproteins, 78-6913
 - Nucleotide Sequence, 78-6917
 - Temperature Sensitive Mutants, 78-6910

NA, Viral

- Mammary Neoplasms, Experimental
 - Hormones, 78-6868
 - Virus, Murine Mammary Tumor, 78-6868
- Virus, Adeno 2
 - Chromosomes, 78-6962
 - Hybrid Cells, 78-6962
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, Review, 78-6670
- Virus, Avian Sarcoma
 - Chromatin, Precipitation, 78-6842
 - Purification Method, 78-6842
- Virus, Epstein-Barr
 - DNA-DNA Hybridization, 78-6954
 - Endonucleases, 78-6954
 - Nucleotide Sequence, 78-6954
- Virus, Herpes
 - Cell Transformation, Neoplastic, Review, 78-6673
 - Isolation and Characterization, Review, 78-6673
- Virus, Herpes Simplex
 - Psoralen, 4,5'-8-Trimethyl-, 78-6937
 - Ultraviolet Rays, 78-6937
- Virus, Herpes Simplex 1
 - Antigens, Viral, 78-6944
 - C-Type Virus Activation, Mouse, 78-6937
 - Cell Transformation, Neoplastic, 78-6944
- Virus, Herpes Simplex 2
 - C-Type Virus Activation, Mouse, 78-6937
- Virus, Moloney Murine Leukemia
 - DNA-RNA Hybridization, 78-6881
 - Integration Sites, 78-6881
 - Virus Replication, 78-6881
- Virus, Murine Mammary Tumor
 - DNA-DNA Hybridization, 78-6863
 - Virus Replication, 78-6863
- Virus, Polyoma
 - Endonuclease-Resistant Mutants, 78-6903
 - RNA Polymerase, 78-6901
 - RNA Replication, 78-6901
- Virus, Rous Sarcoma
 - Infectivity, Sheared Nucleic Acid, 78-6847
 - Transformation, Genetic, 78-6847

DNA, Viral (cont'd)

- Virus, SV40
 - Antigens, Viral, 78-6920, 78-6927
 - C-Type Virus Activation, Mouse, 78-6937
 - Cell Transformation, Neoplastic, 78-6925, 78-6933
 - Cell Transformation, Neoplastic, Review, 78-6670
 - Chromosomes, Human, 6-12, 78-6919, 78-6929
 - Chromosomes, Human, 16-18, 78-6919
 - DNA-DNA Hybridization, 78-6922
 - DNA Replication, 78-6922
 - Endonucleases, 78-6922, 78-6925
 - Hybrid Cells, 78-6919, 78-6926, 78-6929
 - Mitomycin C, 78-6933
 - Purification Method, 78-6842
 - RNA, Messenger, 78-6920
 - Virus Replication, 78-6927
- Virus, Vaccinia
 - C-Type Virus Activation, Mouse, 78-6937

L-Dopa

see Alanine, 3-(3,4-Dihydroxyphenyl)-

Drug Therapy

- Leukemia, Myeloblastic
 - Actinomycin D, 78-6815
 - Case Report, 78-6815
 - Cyclophosphamide, 78-6815
 - Vincristine, 78-6815
- Neoplasms, Multiple Primary
 - Hodgkin's Disease, 78-6736
 - Leukemia, Myeloblastic, 78-6736
 - Multiple Myeloma, 78-6736
 - Ovarian Neoplasms, 78-6736

Drugs

- Chromosome Aberrations
 - Carcinogenic Potential, Review, 78-6618

Dwarfism

- Virus, Epstein-Barr
 - Lymphocyte Transformation, 78-6947

Dyes

- Ames Test
 - Mutagenic Activity, 78-6770

Ear Neoplasms

- Neurilemmoma
 - Genetics, 78-7096
 - Neoplasms, Multiple Primary, 78-7096
- Radiation, Ionizing
 - Carcinogenic Potential, 78-6816

Edema

- Carrageen
 - Tumor Ascitic Fluid, 78-6981

5,8,11,14-Eicosatetranoic Acid

- Ultraviolet Rays
 - Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-, 78-6811
 - Psoralen, 8-Methoxy-, 78-6811

Endometrial Hyperplasia

- Ovarian Neoplasms
 - Adenocarcinoma, 78-7132
 - Endometriosis, 78-7132

Endometriosis

- Ovarian Neoplasms
 - Cysts, 78-7132
 - Endometrial Hyperplasia, 78-7132

Endometriosis (cont'd)

- Histological Study, 78-7132
- Hyperplasia, 78-7132

Endonucleases

- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
 - DNA, Binding, 78-6784
- Ultraviolet Rays
 - DNA Repair, Review, 78-6620
- Virus, Epstein-Barr
 - DNA, Viral, 78-6954
 - Nucleotide Sequence, 78-6954
- Virus, Herpes Simplex 1
 - Cell Transformation, Neoplastic, 78-6944
- Virus, Rous Sarcoma
 - Escherichia coli*, 78-6854
 - Nucleotide Sequence, 78-6854
 - Reverse Transcriptase, 78-6854
 - RNA, Viral, 78-6854
- Virus, SV40
 - Antigens, Viral, 78-6925
 - Chromosomes, 78-6907
 - Deletion Mutants, 78-6930
 - DNA Replication, 78-6927
 - DNA, Viral, 78-6922, 78-6925
 - Virus Replication, 78-6927

Enteritis, Regional

- Polyps
 - Case Report, 78-7130
 - Stomach, 78-7130

Environmental Hazard

- Mutagens
 - Epidemiology, Review, 78-6605
- Quantitation Method
 - Reference Standards, 78-6709

Enzymes

- Dimethylamine, *N*-Nitroso-
 - Purine, 2-Amino-6-methoxy-, 78-6625
- Purine, 2-Amino-6-methoxy-
 - Nitroso Compounds, 78-6625

Epinephrine

- Pheochromocytoma
 - Genetics, 78-7144
 - Urinary Levels, 78-7144

Erythema

- Ultraviolet Rays
 - Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-, 78-6811
 - Prostaglandins E, 78-6811
 - Prostaglandins F, 78-6811
 - Psoralen, 8-Methoxy-, 78-6811

Erythrocytes

- Leukemia, Myeloblastic
 - Colony Formation, 78-7150
- Sarcoma
 - Hemodynamics, 78-7078
 - Mitosis, 78-7078

Erythroleukemia

- Butyric Acid
 - Globin, 78-7185
- Globin
 - RNA, Messenger, 78-7185
- Heme

Erythroleukemia (cont'd)

- Globin, 78-7185
- Methane, Sulfinylbis-
 - Globin, 78-7185
- Virus, Friend Murine Leukemia
 - Antibody-Dependent Cytotoxicity, 78-6879
 - Antigens, Viral, 78-6879

Escherichia coli

- DNA Polymerase
 - DNA Repair, Review, 78-6620
- Mutagens
 - DNA Repair, 78-7193
- Psoralen, 8-Methoxy-
 - Mutagenic Activity, 78-6809
- Ultraviolet Rays
 - DNA Repair, 78-6805
 - Mutagenic Activity, 78-6805
- Virus, Rous Sarcoma
 - Endonucleases, 78-6854

Esophageal Neoplasms

- Carcinoma, Epidermoid
 - Precancerous Conditions, 78-7114
- Carcinoma In Situ
 - Precancerous Conditions, 78-7114
- Diet
 - Epidemiology, Review, 78-6637
- Epidemiology
 - Iran, 78-7153
- Ethyl Alcohol
 - Precancerous Conditions, 78-7114
- Smoking
 - Precancerous Conditions, 78-7114

Esterases

- Carbamic Acid, Ethyl Ester
 - Plasma Levels, Mouse, 78-6743
- Carbon Tetrachloride
 - Plasma Levels, Mouse, 78-6743
- p*-Cresol, 2,6-Di-*tert*-butyl-
 - Plasma Levels, Mouse, 78-6743
- Diethylamine, *N*-Nitroso-
 - Plasma Levels, Mouse, 78-6743
- Dipropylamine, *N*-Nitroso-
 - Plasma Levels, Mouse, 78-6743
- Hydrazine, 1,1-Dimethyl-
 - Plasma Levels, Mouse, 78-6743
- Hydroxylamine, *O,N*-Dimethyl-*N*-nitroso-
 - Plasma Levels, Mouse, 78-6743
- Methanesulfonic Acid, Ethyl Ester
 - Plasma Levels, Mouse, 78-6743
- Piperazine, 2,5-Dimethyl-1,4-dinitroso-
 - Plasma Levels, Mouse, 78-6743
- Piperazine, 1,4-Dinitroso-
 - Plasma Levels, Mouse, 78-6743
- Piperazine, 1-Methyl-4-nitroso-
 - Plasma Levels, Mouse, 78-6743
- Piperidine, 2,6-Dimethyl-1-nitroso-
 - Plasma Levels, Mouse, 78-6743
- Proline, 4-Hydroxy-1-nitroso-
 - Plasma Levels, Mouse, 78-6743

Estradiol

- Mammary Neoplasms, Experimental
 - Dose-Response Study, Mouse, 78-6794

Estradiol, 17-Ethynyl-

- Liver Neoplasms
 - Contraceptives, Oral, 78-7126

adiol, 17-Ethynyl- (cont'd)
Hyperplasia, 78-7125

ogens
Bone Resorption
Menopause, Review, 78-6651
Breast Neoplasms
Epidemiology, 78-7162
Epidemiology, Review, 78-6644, 78-6649, 78-6651
Menopause, 78-6649
Menopause, Review, 78-6651
Receptors, Hormone, Review, 78-6603
RNA Polymerase, 78-6603
Gynecologic Neoplasms
Epidemiology, Review, 78-6644
Liver Neoplasms
Epidemiology, Review, 78-6650
Uterine Neoplasms
Classification, 78-7181
Epidemiology, 78-6647, 78-7181
Epidemiology, Review, 78-6641, 78-6642, 78-6651
Genetics, Review, 78-6647
Menopause, Review, 78-6642, 78-6647, 78-6651
Precancerous Conditions, Review, 78-6641
Risk Factor Analysis, 78-7181

one
Mammary Neoplasms, Experimental
Dose-Response Study, Mouse, 78-6794
Virus, Murine Mammary Tumor, 78-6868
Neoplasms, Experimental
Virus, SV40, 78-6911

ne, 1,2-Dibromo-
Angiosarcoma
Carcinogenic Activity, Rat, 78-6714
Disulfide, Bis(diethylthiocarbamoyl)-, 78-6714
Hepatoma
Carcinogenic Activity, Rat, 78-6714
Disulfide, Bis(diethylthiocarbamoyl)-, 78-6714

ne, 1,1,1-Trichloro-2,2-bis(p-chlorophenyl)-
Liver Neoplasms
Epidemiology, Review, 78-6659

anethiol, 2-Amino-
Dimethylamine, *N*-Nitroso-
Metabolism, 78-6802
Guanidine, Methyl-
Metabolism, 78-6802

anol, 2-Mercapto-
Concanavalin A
Antitumor Activity, 78-6988
Cytotoxicity
Antitumor Activity, 78-6988
Lymphocytes
Cytotoxicity, 78-6988
Peritoneal Fluid
Antitumor Activity, 78-6988

er, Bis(chloromethyl)-
Phosphonium, Tetrakis(hydroxymethyl)-, Chloride
Chromatographic Study, 78-6725

idium Bromide
Virus, SV40
Cell Transformation, Neoplastic, 78-6915
DNA Replication, 78-6915
Drug-Resistant Mutants, 78-6915
Mitochondria, 78-6915

Ethidium Bromide (cont'd)
Proteins, 78-6915

Ethyl Alcohol
Cervix Neoplasms
Epidemiology, Yugoslavia, 78-6945
Esophageal Neoplasms
Precancerous Conditions, 78-7114
Gastrointestinal Neoplasms
Epidemiology, Review, 78-6653
Laryngeal Neoplasms
Epidemiology, Review, 78-6699
Mouth Neoplasms
Epidemiology, Review, 78-6699
Respiratory Tract Neoplasms
Epidemiology, Review, 78-6653
Smoking
Co-carcinogenic Activity, Review, 78-6653

Ethylamine, 2,2'-Dithiobis-
Acetohydroxamic Acid, *N*-Fluoren-2-yl-
Nucleic Acids, Binding, 78-6730
Radiation, Ionizing
Lymphocytes, 78-6979

Ethylene, Chloro-
Angiosarcoma
Alkaline Phosphatase, 78-6713
Case Report, 78-6713
Epidemiology, Review, 78-6634
Glutamyl Transpeptidase, 78-6713
Occupational Hazard, 78-6713
Brain Neoplasms
Epidemiology, Review, 78-6634
Occupational Hazard
Chromatids, 78-6712
Chromosome Abnormalities, 78-6712, 78-6715
Respiratory Tract Neoplasms
Epidemiology, Review, 78-6634
Smoking
Chromosome Abnormalities, 78-6715
Structure-Activity Relationship
Carcinogenic Activity, Review, 78-6636

Ethylene, Chloro- Polymer
Occupational Hazard
Chromosome Abnormalities, 78-6715
Smoking
Chromosome Abnormalities, 78-6715

Ethylene, Trichloro-
Breast Neoplasms
Epidemiology, 78-7173
Cervix Neoplasms
Epidemiology, 78-7173
Digestive System Neoplasms
Epidemiology, 78-7173
Leukemia
Epidemiology, 78-7173
Occupational Hazard, 78-7173
Leukemia, Lymphoblastic
Case Report, 78-6765
Liver Neoplasms
Epidemiology, 78-7173
Occupational Hazard, 78-7173
Lung Neoplasms
Epidemiology, 78-7173
Occupational Hazard
Epidemiology, 78-7173
Structure-Activity Relationship

Ethylene, Trichloro- (cont'd)
 Carcinogenic Activity, Review, 78-6636

Extrachromosomal Inheritance
Agrobacterium tumefaciens
 Octopine, 78-7194

Eye Neoplasms
 Meningioma
 Genetics, 78-7096
 Radiation, Ionizing
 Carcinogenic Potential, 78-6816

Fiber Glass
 Chromosome Abnormalities
 Hamster Cells, 78-6836

Fibrinolysis
 Benzo(a)pyrene
 Cell Transformation, Neoplastic, 78-7076

Fibroblasts
 Acetic Acid, (Ethylenedinitrilo)tetra-
 Proteins, 78-6908
 Benz(a)anthracene, 7,12-Dimethyl-
 Proteins, Binding, 78-6785
 Two-Stage Carcinogenesis, Review, 78-6626
 Benzo(a)pyrene
 Proteins, Binding, 78-6785
 Two-Stage Carcinogenesis, Review, 78-6626
 Blood Platelets
 Growth Substances, 78-6928
 Calcium
 Growth Substances, 78-6928
 Cell Transformation, Neoplastic
 Transplantation, Heterologous, 78-7079
 Chondroitin
 Cell Transformation, Neoplastic, 78-7198
 Contact Inhibition, 78-7198
 Cortisol
 Chondroitin, 78-7198
 Disaccharides
 Chondroitinases, 78-7198
 Growth Substances
 Cell Transformation, Neoplastic, 78-6928
 Cells, Cultured, 78-6928
 Hybrid Cells
 LETS Protein, 78-7077
 Mucopolysaccharides
 Cell Transformation, Neoplastic, 78-7198
 12-*O*-Tetradecanoylphorbol-13-acetate
 Two-Stage Carcinogenesis, Review, 78-6626
 Transplantation, Heterologous
 Mesenchyme, Chick Wing, 78-7079
 Trypsin
 Proteins, 78-6908
 Urea
 Proteins, 78-6908
 Virus, Equine Rhinopneumonitis
 Thymidine Kinase, 78-6898
 Virus Replication, 78-6898
 Virus, Herpes Simplex 1
 Cell Transformation, Neoplastic, 78-6944
 Virus, Rat Sarcoma
 Carcinogenic Activity, 78-6895
 Virus, RNA Tumor
 Cell Transformation, Neoplastic, 78-6928
 Growth Substances, 78-6928
 Virus, SV40
 Cell Transformation, Neoplastic, 78-6928

Fibroblasts (cont'd)
 Genetics, 78-6934
 Growth Substances, 78-6928
 Proteins, 78-6908

Fibroma
 Acetophenone, 2-Amino-
 Species Difference, 78-6790
 Gynecologic Neoplasms
 Acetophenone, 2-Amino-, 78-6790
 Indole-3-acrylic Acid, 78-6790
 Indole-3-acrylic Acid
 Species Difference, 78-6790
 Radium
 Dose-Response Study, Mouse, 78-6830
 Xanthine, 3-Hydroxy-
 Carcinogenic Activity, Rat, 78-6718

Fibrosarcoma
 T-Lymphocytes
 Antilymphocyte Serum, 78-7067
 Immunosuppression, Rat, 78-7067
 Radiation, Ionizing
 Immunosuppression, Rat, 78-7067
 Acetophenone, 2-Amino-
 Species Difference, 78-6790
 Areca
 Carcinogenic Potential, Ingredients, 78-6727
 Benz(a)anthracene, 7,12-Dimethyl-
Corynebacterium parvum, 78-7055
 Benzo(a)pyrene
 Cells, Cultured, 78-7076
 Transplantation Immunology, 78-7051
 Bone Neoplasms
 Strontium Radioisotopes, 78-6835
 Cells, Cultured
 Neoplasm Metastasis, 78-7109
 Cholanthrene, 3-Methyl-
 Immunization, 78-7053
 Virus, Rat Leukemia, 78-6873
 Complement
 Growth, 78-7053
Corynebacterium parvum
 Antitumor Activity, 78-7056
 T-Lymphocytes, 78-7056
 Macrophage Activation, 78-7056
 Macrophages, 78-7054
 Transplantation Immunology, 78-7055
 Gamma Globulins
 Hypersensitivity, Delayed, 78-7054
 Genetics
 Leukemoid Reaction, 78-7052
 Hypersensitivity, Delayed
 Immunity, Passive, 78-7054
 Immunity, Passive
 Monocytes, 78-7054
 Neoplasm Metastasis, 78-7047
 Immunosuppression
 Spleen Cells, 78-7047
 Indole-3-acrylic Acid
 Species Difference, 78-6790
 Kidney Neoplasms
 Dimethylamine, *N*-Nitroso-, 78-6739
 B-Lymphocytes
 Neoplasm Metastasis, 78-7047
 T-Lymphocytes
 Growth, 78-7053
 Immunity, Cellular, 78-7051

Fibrosarcoma (cont'd)

- Immunity, Passive, 78-7054
- Neoplasm Metastasis, 78-7047
- Transplantation Immunology, 78-7051, 78-7054

Macrophages

- Histocompatibility Antigens, 78-7051
- Immunity, Cellular, 78-7051
- Transplantation Immunology, 78-7051

Neoplasm Transplantation

- Immunosuppression, 78-7047
- Leukemoid Reaction, 78-7052

Radiation, Ionizing

- Cytotoxicity, Spleen Cells, 78-7053
- Immunization, 78-7053

Virus, Rat Leukemia

- Antigenic Determinants, 78-6873
- Isolation and Characterization, 78-6873

Fluorescence

- Acetamide, *N*-(Acetyloxy)-*N*-fluorenyl-2-yl-
Liver, Rat, 78-6716

- 2-Butanone, 1,1-Dihydroxy-3-ethoxy-
Liver, Rat, 78-6716

Terbium

- DNA, 78-6716
- Guanine, 78-6716
- Liver, Rat, 78-6716
- RNA, Ribosomal, 78-6716

Fluorescent Dyes**Ultraviolet Rays**

- Occupational Hazard, 78-6663
- Textile Workers, Review, 78-6663

Food Additives

- Carcinogen, Chemical
Risk Factors, Review, 78-6662
- Chromosome Aberrations
Carcinogenic Potential, Review, 78-6618
- Nitric Acid
Carcinogenic Potential, Review, 78-6662

Food Contamination

- Carcinogen, Chemical
Risk Factors, Review, 78-6662
- Guanidine, (4-Aminobutyl)-
Quantitation Method, 78-6741
- Guanidine, Methyl-
Quantitation Method, 78-6741
- Hepatitis
Aflatoxin B1, 78-6721
- Aflatoxin G1, 78-6721
- Mycotoxins, 78-7184

Insecticides

- Carcinogenic Potential, Review, 78-6662

Meningioma

- Nitrous Acid, Sodium Salt, 78-7170

Mycotoxins

- Carcinogenic Potential, Review, 78-6662

Nitrous Acid

- Carcinogenic Potential, Review, 78-6655, 78-6662

Pesticides

- Carcinogenic Potential, Review, 78-6662

Stomach Neoplasms

- Nitric Acid, 78-6733
- Nitrous Acid, 78-6733

Foreign Bodies**Homeostasis**

- Cell Transformation, Neoplastic, Review, 78-6705

Foreign Bodies (cont'd)**Iron**

- Co-carcinogenic Effect, 78-6654
- Neoplasms, Experimental
Immune Response, Review, 78-6633
- Precancerous Conditions, Review, 78-6633

Foreign Body Reaction**Lung Neoplasms**

- Carcinoma, 78-6992
- Immune Response, 78-6992

Formaldehyde

- Benzo(a)pyrene
Co-carcinogenic Effect, Review, 78-6657
- Lung Neoplasms
Smoke Condensate, Review, 78-6657
- Virus, Gross Murine Leukemia
Antibody Formation, 78-7024
- Immunity, Cellular, 78-7024
- Virus, Rauscher Murine Leukemia
Antibody Formation, 78-7024
- Immunity, Cellular, 78-7024

Formamide, *N*-(4-(5-Nitro-2-furyl)-2-thiazolyl)-

- Bladder Neoplasms
Carcinoma, Transitional Cell, 78-7141

Free Radicals

- Air Pollution
Spectrum Analysis, 78-6710
- Ozone
Smog Formation, 78-6710

Freon

- see Methane, Chlorodifluoro-

Freund's Adjuvant

- Cholanthrene, 3-Methyl-
Immune Response, Mouse, 78-7007
- T-Lymphocytes
Immunosuppression, 78-7005
- Virus, SV40
Immunity, Cellular, 78-7005
- T-Lymphocytes, 78-7005
- Transplantation Immunology, 78-7005

FSH

- Mammary Neoplasms, Experimental
Benz(a)anthracene, 7,12-Dimethyl-, 78-6778

Galactosidases

- Radiation, Ionizing
Enzymatic Activity, 78-6814

Gallbladder Diseases

- Ethnic Groups
Epidemiology, New Mexico, 78-7178

Gallbladder Neoplasms

- Ethnic Groups
Epidemiology, New Mexico, 78-7178
- Genetics
Indians, North American, 78-7178

Gallic Acid, Propyl Ester

- Dimethylamine, *N*-Nitroso-
Metabolism, 78-6802
- Guanidine, Methyl-
Metabolism, 78-6802

Gamma Globulins

- Fibrosarcoma

Gamma Globulins (cont'd)

Hypersensitivity, Delayed, 78-7054

Gastric Mucosa

Stomach Neoplasms

Metaplasia, 78-7117

Gastritis

Stomach Neoplasms

Polyyps, 78-7116

Precancerous Conditions, 78-7116

Gastrointestinal Neoplasms

Asbestos

Epidemiology, Review, 78-6634

Carcinogen, Chemical

Diet, Review, 78-6638

Ethnic Groups

Epidemiology, Hawaii, 78-7164

Ethyl Alcohol

Epidemiology, Review, 78-6653

Mycotoxins

Diet, Review, 78-6638

Retinol, 78-6638

Trace Elements

Epidemiology, Review, 78-6631

Gastrointestinal System

Diet

Bacteria, 78-7180

Genetics

Adenocarcinoma

Carcinoembryonic Antigen, 78-7118

Asbestos

Epidemiology, Review, 78-6697

Breast Neoplasms

Alpha Fetoproteins, 78-7066

Carcinoembryonic Antigen, 78-7066

Carcinogen, Chemical

Cell Transformation, Neoplastic, Review, 78-6705

Carcinogen, Environmental

Cell Transformation, Neoplastic, Review, 78-6609
78-6610

Carcinoma, Basal Cell

Muscular Atrophy, 78-7094

Nevus, 78-7094

Odontogenic Cysts, 78-7094

Carcinoma, Epidermoid

Warts, 78-7036

Colonic Neoplasms

Adenocarcinoma, 78-7118

Ear Neoplasms

Neurilemmoma, 78-7096

Eye Neoplasms

Meningioma, 78-7096

Fibrosarcoma

Leukemoid Reaction, 78-7052

Gallbladder Neoplasms

Indians, North American, 78-7178

Hodgkin's Disease

Alpha Fetoproteins, 78-7066

Carcinoembryonic Antigen, 78-7066

Case Report, 78-7083

Histological Study, 78-7083

Leukemia

Virus, SV40, 78-6934

Lymphoma

Virus, SV40, 78-6934

Melanoma

Genetics (cont'd)

Alpha Fetoproteins, 78-7066

Carcinoembryonic Antigen, 78-7066

Meningioma

Case Report, 78-7096

Neurofibromatosis, 78-7096

Neoplasms

Epidemiology, Review, 78-6697

Neurofibromatosis

Child, 78-7097

Morbidity, 78-7097

Oncogenic Viruses

Carcinoembryonic Antigen, 78-7118

Cell Transformation, Neoplastic, Review, 78-6705

Pheochromocytoma

Case Report, 78-7144

Epinephrine, 78-7144

Neoplastic Endocrine-Like Syndromes, 78-7144

Radiation, Ionizing

Epidemiology, Review, 78-6697

Skin Neoplasms

Xeroderma Pigmentosum, 78-7121

Smoking

Carcinoembryonic Antigen, 78-7118

Spinal Neoplasms

Meningioma, 78-7096

Thyroid Neoplasms

Pheochromocytoma, 78-7144

Ultraviolet Rays

Epidemiology, Review, 78-6697

Uterine Neoplasms

Adenocarcinoma, 78-7118

Virus, Murine Mammary Tumor

Cell Transformation, Neoplastic, 78-6866

Virus, SV40

Antigens, Viral, 78-6934

Fibroblasts, 78-6934

Xeroderma Pigmentosum

Case Report, 78-7121

Giant Cell Tumors

Cells, Cultured

Cell Differentiation, 78-7148

Chromosomes, 78-7148

Lactate Dehydrogenase Isoenzymes, 78-7148

Glioblastoma Multiforme

Bacteria

Carcinogenic Metabolite, 78-7149

Tooth Diseases

Bacteria, 78-7149

Epidemiology, 78-7149

Glioma

Brain Neoplasms

Urea, Ethyl Nitroso-, 78-6749

Histocompatibility Antigens

Antigen Frequency, 78-7032

Globin

Erythroleukemia

Butyric Acid, 78-7185

Heme, 78-7185

Methane, Sulfinylbis-, 78-7185

RNA, Messenger, 78-7185

Glomerulonephritis

Cholanthrene, 3-Methyl-

Mouse, New Zealand Black, 78-7041

Hyperplasia

merulonephritis (cont'd)

Lymph Nodes, 78-7084

Lymphoma

Case Report, Castleman's Tumor, 78-7084

Immunoglobulins, 78-7084

icagon

Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-

Adenyl Cyclase, 78-6766

ucose, 2-Deoxy-

Virus, Epstein-Barr

Virus Replication, 78-6946

Virus, Herpes Saimiri

Virus Replication, 78-6946

Virus, Rous Sarcoma

Ascorbic Acid, 78-6849

ucosephosphatase

Hepatoma

Hyperplasia, 78-6729

ucosephosphate Dehydrogenase

Hepatoma

Cells, Cultured, 78-6957

Liver Neoplasms

Morpholine, *N*-Nitroso-, 78-6787

Precancerous Conditions, 78-6787

Skin Neoplasms

Ultraviolet Rays, 78-6813

Ultraviolet Rays

Enzymatic Activity, 78-6813

cosidases

Radiation, Ionizing

Enzymatic Activity, 78-6814

curonates

Chondroitin

Cell Transformation, Neoplastic, 78-7198

curonidase

Laryngeal Neoplasms

Neutrophils, 78-7099

atamyl Transpeptidase

Angiosarcoma

Ethylene, Chloro-, 78-6713

Hepatoma

Hyperplasia, 78-6729

ataraldehyde

Sarcoma

Transplantation Immunology, 78-7048

atathione

Radiation, Ionizing

Lymphocytes, 78-6979

Skin Neoplasms

Ultraviolet Rays, 78-6806

atathione Reductase

Glycoproteins

LETs Protein, 78-7077

ycerolphosphate Dehydrogenase

Nervous System Neoplasms

Urea, Ethyl Nitroso-, 78-6757

ycinonitrile

Hydrazine, 1,2-Dimethyl-

DNA, Alkylation, 78-6755

Metabolism, Rat, 78-6755

Glycogen

Liver Neoplasms

Carcinogen, Chemical, 78-6707

Glycoproteins

Chromosomes, Human, 6-12

Hybrid Cells, 78-7077

LETs Protein, 78-7077

Glutathione Reductase

LETs Protein, 78-7077

Hybrid Cells

Cell Membrane, 78-7077

Tunicamycin

Cell Membrane, 78-6906

Virus, Avian Myeloblastosis

Antigen-Antibody Reactions, 78-6844

Cell Membrane, 78-6844

Isolation and Characterization, 78-6844

Virus, Polyoma

Cell Transformation, Neoplastic, 78-6906

Tunicamycin, 78-6906

Virus, Rauscher Murine Leukemia

Cell Transformation, Neoplastic, 78-6890

Histocompatibility Antigens, 78-6890

T-Lymphocytes, 78-6890

Virus, SV40

Cell Transformation, Neoplastic, 78-6906

Tunicamycin, 78-6906

Gonadotropins

Mammary Neoplasms, Experimental

Benz(a)anthracene, 7,12-Dimethyl-, 78-6778

Graft vs Host Reaction

T-Lymphocytes

Lymphocyte Depletion, 78-7045

Lymphoma

Thymectomy, 78-7028

Myosarcoma

Benzo(a)pyrene, 78-7045

Sarcoma

Benzo(a)pyrene, 78-7045

B-Lymphocytes, 78-7045

T-Lymphocytes, 78-7045

Neoplasm Metastasis, 78-7045

Granuloma

Toxocara canis

Connective Tissue, Mouse, 78-7085

Ultrastructural Study, Mouse, 78-7085

Granulosa Cell Tumor

Gynecologic Neoplasms

Acetophenone, 2-Amino-, 78-6790

Indole-3-acrylic Acid, 78-6790

Growth

Breast Neoplasms

Alanine, 3-(3,4-Dihydroxyphenyl)-, 78-6728

Carcinoma

BCG, 78-6996

Fibrosarcoma

Complement, 78-7053

T-Lymphocytes, 78-7053

HeLa Cells

Tumor Ascitic Fluid, 78-6981

Hepatoma

Antigen-Antibody Complex, 78-7064

Leukemia L1210

Macrophages, 78-7000

Growth (cont'd)

- Lung Neoplasms
 - Phagocytosis, 78-6998
- Lymphoma
 - Carrageen, 78-6997
- Lymphosarcoma
 - Cell Cycle Kinetics, 78-6980
 - Peritoneal Cavity, Mouse, 78-6980
- Mammary Neoplasms, Experimental
 - Immune Serums, 78-7004
- Melanoma
 - Inflammation, 78-6990
- Neoplasms
 - Phenotype, Review, 78-6687

Growth Substances

- Cell Division
 - Embryo, Mouse, 78-7197
- Fibroblasts
 - Blood Platelets, 78-6928
 - Calcium, 78-6928
 - Cell Transformation, Neoplastic, 78-6928
 - Cells, Cultured, 78-6928
- Virus, RNA Tumor
 - Fibroblasts, 78-6928
- Virus, SV40
 - Blood Platelets, 78-6928
 - Fibroblasts, 78-6928

Guanidine, (4-Aminobutyl)-

- Food Contamination
- Quantitation Method, 78-6741

Guanidine, Methyl-

- Ascorbic Acid, Monosodium Salt
 - Metabolism, 78-6802
- Chromosome Aberrations
 - Metabolism, 78-6802
- Cysteine
 - Metabolism, 78-6802
- DNA Repair
 - Metabolism, 78-6802
- Ethanethiol, 2-Amino-
 - Metabolism, 78-6802
- Food Contamination
 - Quantitation Method, 78-6741
- Gallic Acid, Propyl Ester
 - Metabolism, 78-6802

Guanidine, 1-Methyl-3-nitro-1-nitroso-

- Ataxia Telangiectasia
 - DNA Repair, Review, 78-6612
- Cell Differentiation
 - Organ Culture, Prostate, 78-6751
- DNA Repair
 - Carcinogenic Activity, Review, 78-6608, 78-6622
- Haemophilus influenzae*
 - DNA Repair, 78-6759
- Immunosuppression
 - Cell Transformation, Neoplastic, 78-7110
- Leukemia L1210
 - Histones, Binding, 78-6752

Guanine

- Terbium
 - Fluorescence, 78-6716

Guanine, 7-Methyl-

- Hydrazine, 1,2-Dimethyl-
 - DNA, Alkylation, 78-6755

Guanosine

- Phosphonium, Tetrakis(hydroxymethyl)-, Chloride
 - Phosphine Product, 78-6725

Guanosine Cyclic 3',5' Monophosphate

- Cholangioma
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6767

Guanosine Triphosphate

- Plasmacytoma
 - RNA Replication, 78-7190

Guanyl Cyclase

- Cell Membrane
 - Cell Transformation, Neoplastic, 78-7188
- Cell Transformation, Neoplastic
 - Enzymatic Activity, 78-7188
- Cholangioma
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6767

Gynecologic Neoplasms

- Acetophenone, 2-Amino-
 - Species Difference, 78-6790
- Adenocarcinoma
 - Contraceptives, Oral, 78-6645
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Celiac Disease
 - Epidemiology, 78-7154
- Contraceptives, Oral
 - Epidemiology, Review, 78-6645
 - Ultrastructural Study, Review, 78-6645
- Estrogens
 - Epidemiology, Review, 78-6644
- Fibroma
 - Acetophenone, 2-Amino-, 78-6790
 - Indole-3-acrylic Acid, 78-6790
- Granulosa Cell Tumor
 - Acetophenone, 2-Amino-, 78-6790
 - Indole-3-acrylic Acid, 78-6790
- Hair Dyes
 - Epidemiology, 78-7163
- Indole-3-acrylic Acid
 - Species Difference, 78-6790
- Myoma
 - Contraceptives, Oral, 78-6645
- Papilloma
 - Acetophenone, 2-Amino-, 78-6790
 - Indole-3-acrylic Acid, 78-6790
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Epidemiology, Review, 78-6644
 - Histological Study, 78-6789
 - Precancerous Conditions, 78-6652, 78-6789
 - Transplacental Carcinogenesis, 78-6789
 - Transplacental Carcinogenesis, Review, 78-6652

Haemophilus influenzae

- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester
 - DNA Repair, 78-6759
- Diethylamine, 2,2'-Dichloro-*N*-methyl-
 - DNA Repair, 78-6759
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
 - DNA Repair, 78-6759
- Methanesulfonic Acid, Ethyl Ester
 - DNA Repair, 78-6759
- Mitomycin C
 - DNA Repair, 78-6759
- Radiation, Ionizing

***Haemophilus influenzae* (cont'd)**

DNA Repair, 78-6759

Ultraviolet Rays

DNA Repair, 78-6759

Hair Dyes

Breast Neoplasms

Epidemiology, 78-7163, 78-7166

Gynecologic Neoplasms

Epidemiology, 78-7163

Leukemia

Epidemiology, 78-7166

Occupational Hazard

Epidemiology, 78-7166

Skin Neoplasms

Epidemiology, 78-7163

Thyroid Neoplasms

Epidemiology, 78-7163

Laptens

Myeloma Proteins

Spectrum Analysis, Raman, 78-7039

Head and Neck Neoplasms

Carcinoma, Epidermoid

Radiation, Ionizing, 78-6832

Radiation, Ionizing

Precancerous Conditions, 78-6832

Virus, Epstein-Barr

Antibodies, Viral, 78-7034

Hearing Disorders

Radiation, Ionizing

Epidemiology, 78-6816

Heavy Chain Disease

Immunoglobulins

Gene Deletion, 78-7060

Proteins

Amino Acids, 78-7060

RNA, Messenger

Immunoglobulins, 78-7060

Proteins, 78-7060

HeLa Cells

Bracken Fern

Toxicity, Pterosins, Pterosides, 78-6717

Growth

Tumor Ascitic Fluid, 78-6981

Virus, Adeno 2

Virus Replication, 78-6965

Virus, Adeno 2 - SV40 Hybrid

RNA, Messenger, 78-6966

Hemangioma

Nose Neoplasms

Neoplasms, Multiple Primary, 78-7071

Hemangioma, Cavernous

Liver Neoplasms

Case Report, 78-7098

Neoplasms, Multiple Primary, 78-7098

Hematopoietic Stem Cells

Leukemia, Lymphoblastic

Antigenic Determinants, 78-7026

Leukemia, Myeloblastic

Antigenic Determinants, 78-7026

Virus, Avian Leukosis

Target Cells, Review, 78-6676

Virus, Friend Murine Leukemia

Colony Formation, 78-6878

Hematopoietic Stem Cells (cont'd)

Virus Replication, 78-6878

Heme

Erythroleukemia

Globin, 78-7185

Hemosiderin

Mesothelioma

Pericardium, 78-7113

Hepatitis

Leukemia, Lymphoblastic

Case Report, Child, 78-6956

Virus-Like Particles

Isolation and Characterization, Woodchuck
78-6959**Hepatoma**Acetamide, *N*-Fluoren-2-yl-

Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6729

Hyperplasia, 78-6729

Precancerous Conditions, 78-6729

Aflatoxin B1

Food Contamination, 78-6721

Precancerous Conditions, 78-6722

Ultrastructural Study, Rat, 78-6722

Aflatoxin G1

Food Contamination, 78-6721

Alpha Fetoproteins

Serum Levels, 78-7124

Aniline, *N,N*-Dimethyl-*p*-phenylazo-

Antigen-Antibody Complex, 78-7042, 78-7064

Complement, 78-7063

Antigen-Antibody Complex

Complement Binding Assay, 78-7042

Growth, 78-7064

Transplantation Immunology, 78-7042

Ascitic Fluid

Anti-Inflammatory Activity, 78-6981

Australia Antigen

Antigen-Antibody Reactions, 78-6961

Cells, Cultured, 78-6957

Epidemiology, 78-6961

Immunohistochemical Study, 78-6958

Neoplasm Metastasis, 78-6958

Precancerous Conditions, 78-6961

Seroepidemiology, Review, 78-6685

Ultrastructural Study, 78-6957

BCG

Macrophages, 78-6995

Neoplasm Regression, 78-6995

Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-

DNA Replication, 78-6744

Benzidine, Dihydrochloride

Neoplasm Metastasis, 78-7122

Benzo(a)pyrene

Neoplasm Metastasis, 78-7122

Chromosomes

Cells, Cultured, 78-6957

Complement

Antigen-Antibody Complex, 78-7064

Contraceptives, Oral

Case Report, 78-6795

Epidemiology, Review, 78-6645, 78-6650

Diethylamine, *N*-Nitroso-

Carbon Tetrachloride, 78-6750

DNA Replication, 78-6744

Liver Regeneration, 78-6750

Hepatoma (cont'd)

- Neoplasm Metastasis, 78-7122
- Precancerous Conditions, 78-6744
- Dimethylamine, *N*-Nitroso-
 - DNA Replication, 78-6744
- Direct Black 38
 - Carcinogenic Activity, Rat, 78-6771
- Direct Blue 6
 - Carcinogenic Activity, Rat, 78-6771
- Direct Brown 95
 - Carcinogenic Activity, Rat, 78-6771
- DNA Replication
 - Precancerous Conditions, 78-6744
- Ethane, 1,2-Dibromo-
 - Carcinogenic Activity, Rat, 78-6714
 - Disulfide, Bis(diethylthiocarbamoyl)-, 78-6714
- Fibrosis
 - Epidemiology, Japan, 78-7127
- Glucosephosphate Dehydrogenase
 - Cells, Cultured, 78-6957
- Hyperplasia
 - Adenosine Triphosphatase, 78-6729
 - DNA Replication, 78-6729
 - Glucosephosphatase, 78-6729
 - Glutamyl Transpeptidase, 78-6729
- Immune Serums
 - Alpha Fetoproteins, 78-6960
 - Australia Antigen, 78-6960
 - Phosphodiesterases, 78-6960
 - Virus, Hepatitis, 78-6960
- Liver Cirrhosis
 - Australia Antigen, 78-6961
 - Epidemiology, Japan, 78-7127
 - Phosphodiesterases, 78-6960
 - Precancerous Conditions, 78-7123
 - Virus, Hepatitis, 78-6958
 - Virus-Like Particles, 78-6960
- Liver Neoplasms
 - Alpha Fetoproteins, 78-7062
 - Radioimmunoassay, 78-7062
- Mycotoxins
 - Epidemiology, Kenya, Swaziland, 78-7184
 - Food Contamination, 78-7184
 - Ultrastructural Study, Rat, 78-6723
- Neoplasm Metastasis
 - Histological Study, 78-7122
 - Lung, Mouse, 78-7122
- Phosphodiesterases
 - Ultrastructural/Cytochemical Study, 78-6960
- Polyribosomes
 - RNA, Messenger, 78-7189
- Porphyria
 - Precancerous Conditions, 78-7123
- Retinol
 - Carrier Proteins, 78-6796
- RNA, Messenger
 - Nucleoproteins, 78-7189
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Carrier Proteins, 78-6796
- Urea, Ethyl Nitroso-
 - Neoplasm Metastasis, 78-7122
- Virus, Hepatitis
 - Australia Antigen, 78-6685
 - Epidemiology, Vietnam, 78-6958
 - Phosphodiesterases, 78-6960
 - Seroepidemiology, Review, 78-6685
 - Ultrastructural Study, Woodchuck, 78-6959

Hepatoma (cont'd)

- Vertical Transmission, Review, 78-6685
- Virus-Like Particles, 78-6960
- Virus-Like Particles
 - Isolation and Characterization, Woodchuck, 78-6959
- Xanthine, 3-Hydroxy-
 - Carcinogenic Activity, Rat, 78-6718

Hereditary Diseases

- Age Factors
 - Cell Transformation, Neoplastic, Review, 78-6602
- Carcinogen, Environmental
 - Cell Transformation, Neoplastic, Review, 78-6602
 - Mutation, Review, 78-6602
- Chromosome Abnormalities
 - Cell Transformation, Neoplastic, Review, 78-6602
- DNA Repair
 - Cell Transformation, Neoplastic, Review, 78-6613
- Mutagens
 - Epidemiology, Review, 78-6605

2,5-Hexadienoic Acid, 3-Methoxy-5-methyl-4-oxo-Mycotoxins

- Carcinogenic Activity, Review, 78-6629

1,6-Hexanediamine

- Hyperplasia
 - Spleen, Mouse, Rat, 78-6817

Histocompatibility Antigens

- Antigens, Viral
 - Immune Response, Review, 78-6694
- Fibrosarcoma
 - Macrophages, 78-7051
- Glioma
 - Antigen Frequency, 78-7032
- Liver Neoplasms
 - Carcinogen, Chemical, 78-6692
- T-Lymphocytes
 - Immune Response, 78-7044
- Lymphoma
 - Cells, Cultured, 78-7010
 - Immunity, Cellular, 78-7027
 - Virus, Moloney Murine Leukemia, 78-6883
 - Virus, Rauscher Murine Leukemia, 78-6883
- Neoplasms, Experimental
 - Cholanthrene, 3-Methyl-, 78-7044
 - Transplantation Immunology, 78-7044
- Plasmacytoma
 - Immune Response, Mouse, 78-7038
 - Radiation, Ionizing, 78-7038
 - Silica, 78-7038
- Sarcoma
 - Cholanthrene, 3-Methyl-, 78-7043
 - Neoplasm Metastasis, 78-7043
 - Neoplasm Transplantation, 78-7043
 - Transplantation Immunology, 78-7044
 - Virus, SV40, 78-7044
- Virus, Influenza
 - Cell Membrane, 78-6909
 - Transplantation Immunology, 78-6909
- Virus, Moloney Murine Sarcoma
 - Cell Membrane, 78-6909
 - Transplantation Immunology, 78-6909
- Virus, Rauscher Murine Leukemia
 - Cell Transformation, Neoplastic, 78-6890
 - Glycoproteins, 78-6890
 - T-Lymphocytes, 78-6890

stocompatibility Antigens (cont'd)

- Virus, SV40
 - Cell Membrane, 78-6909
 - Cell Transformation, Neoplastic, 78-6909
 - Transplantation Immunology, 78-6909
- Virus, Influenza, 78-6909
- Virus, Moloney Murine Sarcoma, 78-6909

stones

- Benzo(a)pyrene, 7,8-Dihydroxy-9,10-oxy-7,8,9,10-tetrahydro-
 - DNA, Binding, 78-6784
- Virus, SV40
 - Cell Movement, 78-6907
 - Chromatin-DNA Hybrids, 78-6907
 - Chromosomes, 78-6907, 78-6923
 - Virus-Like Particles, 78-6923

dgkin's Disease

- Celiac Disease
 - Epidemiology, 78-7154
- DNA Replication
 - Autoradiographic Study, 78-7192
 - Histological Study, 78-7192

Genetics

- Alpha Fetoproteins, 78-7066
- Carcinoembryonic Antigen, 78-7066
- Case Report, 78-7083
- Histological Study, 78-7083

Lymph Nodes

- DNA Replication, 78-7192

Neoplasms, Multiple Primary

- Drug Therapy, 78-6736

Virus, Epstein-Barr

- Migration Inhibitory Factor, 78-6989

meostasis

- Cell Differentiation
 - Cell Transformation, Neoplastic, Review, 78-6705
- DNA
 - Cells, Cultured, 78-7191
- Foreign Bodies
 - Cell Transformation, Neoplastic, Review, 78-6705

rmones

- Brain Neoplasms
 - Epidemiology, 78-7168
- Breast Neoplasms
 - Cell Division, 78-6791
 - Plasma Levels, 78-7162
- Carcinogen, Chemical
 - Co-carcinogenic Effect, Review, 78-6643
- Cell Division
 - Co-carcinogenic Effect, Review, 78-6643
- Cholanthrene, 3-Methyl-
 - Co-carcinogenic Effect, 78-6798
- Mammary Neoplasms, Experimental
 - Cell Division, 78-6791
 - Co-carcinogenic Effect, Review, 78-6643
 - DNA, Viral, 78-6868
 - RNA, Viral, 78-6868
- Oncogenic Viruses
 - Co-carcinogenic Effect, Review, 78-6643
- Radiation
 - Co-carcinogenic Effect, Review, 78-6643

ybrid Cells

- Fibroblasts
 - LETS Protein, 78-7077
- Glycoproteins

Hybrid Cells (cont'd)

- Cell Membrane, 78-7077
- Chromosomes, Human, 6-12, 78-7077
- L Cells
 - LETS Protein, 78-7077
- Teratoid Tumor
 - Virus, Polyoma, 78-6902
- Virus, Adeno 2
 - DNA-RNA Hybridization, 78-6962
 - DNA, Viral, 78-6962
 - RNA, Messenger, 78-6962
- Virus, Polyoma
 - Antigens, Viral, 78-6902
- Virus, SV40
 - Antigens, Viral, 78-6914, 78-6929
 - Cell Transformation, Neoplastic, 78-6914, 78-6919
 - Chromosomes, 78-6914
 - Chromosomes, Human, 6-12, 78-6919, 78-6929
 - Chromosomes, Human, 16-18, 78-6919
 - DNA, Viral, 78-6919, 78-6926, 78-6929

Hydatidiform Mole

- Choriocarcinoma
 - Case Report, 78-7134
- Insemination, Artificial
 - Case Report, 78-7093

Hydrazine, 1,1-Dimethyl-

- Esterases
 - Plasma Levels, Mouse, 78-6743

Hydrazine, 1,2-Dimethyl-

- DNA, Alkylation
 - Colon, Liver, Kidney, 78-6755
- Glycinonitrile
 - DNA, Alkylation, 78-6755
 - Metabolism, Rat, 78-6755
- Guanine, 7-Methyl-
 - DNA, Alkylation, 78-6755
- Purine, 2-Amino-6-methoxy-
 - DNA, Alkylation, 78-6755

Hydro-Lyases

- Polycyclic Hydrocarbons
 - Metabolism, Review, 78-6623

Hydrocarbons, Halogenated

- Microsomes, Liver
 - Metabolism, Review, 78-6607
- Nucleic Acids, Binding
 - Hepatotoxicity, Review, 78-6607

Hydrocyanic Acid

- Benzo(a)pyrene
 - Co-carcinogenic Effect, Review, 78-6657
- Lung Neoplasms
 - Smoke Condensate, Review, 78-6657

Hydroxylamine, O,N-Dimethyl-N-nitroso-

- Esterases
 - Plasma Levels, Mouse, 78-6743

Hyperplasia

- Beryllium Chloride
 - Spleen, Mouse, Rat, 78-6817
- Breast Neoplasms
 - Contraceptives, Oral, 78-6645
 - Epidemiology, Japan, Hawaii, 78-7108
 - Precancerous Conditions, 78-7108
- Cesium Radioisotopes
 - Spleen, Mouse, Rat, 78-6817

Hyperplasia (cont'd)

- Contraceptives, Oral
 - Epidemiology, Review, 78-6650
- Glomerulonephritis
 - Lymph Nodes, 78-7084
- Hepatoma
 - Acetamide, *N*-Fluoren-2-yl-, 78-6729
 - Adenosine Triphosphatase, 78-6729
 - DNA Replication, 78-6729
 - Glucosephosphatase, 78-6729
 - Glutamyl Transpeptidase, 78-6729
- 1,6-Hexanediamine
 - Spleen, Mouse, Rat, 78-6817
- Liver Neoplasms
 - Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt 78-6758
 - Contraceptives, Oral, 78-6645, 78-6650, 78-6795 78-7125
 - Estradiol, 17-Ethynyl-, 78-7125
 - Mestranol, 78-7125
 - Precancerous Conditions, Review, 78-6707
- Mercury, Chloromethyl-
 - Spleen, Mouse, Rat, 78-6817
- Nitric Acid, Lead Salt
 - Spleen, Mouse, Rat, 78-6817
- Ovarian Neoplasms
 - Adenocarcinoma, 78-7132
 - Endometriosis, 78-7132
- Radium
 - Spleen, Mouse, Rat, 78-6817
- Skin Neoplasms
 - Precancerous Conditions, 78-7087
- Strontium Radioisotopes
 - Spleen, Mouse, Rat, 78-6817
- Zinc Radioisotopes
 - Spleen, Mouse, Rat, 78-6817

Hypersensitivity

- Melanoma
 - Precancerous Conditions, Review, 78-6693

Hypersensitivity, Delayed

- Asbestosis
 - Benzene, 1-Chloro-2,4-dinitro-, 78-7061
- Fibrosarcoma
 - Gamma Globulins, 78-7054
 - Immunity, Passive, 78-7054

Hypertension

- Breast Neoplasms
 - Epidemiology, Sweden, 78-7106

Hyperthyroidism

- Adenoma
 - Radiation, Ionizing, 78-6831
- Radiation, Ionizing
 - Case Report, 78-6831
- Thyroid Neoplasms
 - Radiation, Ionizing, 78-7102

Hypoxanthine Phosphoribosyltransferase

- Psoralen, 8-Methoxy-
 - Mutagenic Activity, 78-6808
- Ultraviolet Rays
 - Mutagenic Activity, 78-6808

IgA

- Myeloma Proteins
 - Binding Sites, 78-7040
 - 2,4-Dinitrophenyl Haptens, Binding, 78-7039

IgE

- Anaphylaxis
 - Immune Response, Solid Tumors, 78-6691

IgG

- Lymphocytes
 - Antilymphocyte Serum, 78-6986
 - Immunoglobulins, Surface, 78-6986
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Antibody Formation, 78-7011

IgM

- Leukemia, Lymphocytic
 - Antibody Specificity, 78-7073
 - Antigenic Determinants, 78-7073
 - B-Lymphocytes, 78-7073
- Lymphoma
 - Sjogren's Syndrome, 78-7082

Immune Serums

- Hepatoma
 - Alpha Fetoproteins, 78-6960
 - Australia Antigen, 78-6960
 - Phosphodiesterases, 78-6960
 - Virus, Hepatitis, 78-6960
- Leukemia, Myeloblastic
 - Antibody-Dependent Cytotoxicity, 78-7019
- Lung Neoplasms
 - Macrophages, 78-6999
- Mammary Neoplasms, Experimental
 - Antigens, Neoplasm, 78-6978
 - Growth, 78-7004
 - Immunity, Cellular, 78-7004
 - Immunosuppressive Factor, 78-7004
 - Macrophages, 78-7004
 - Monocytes, Spleen, 78-7004
- Melanoma
 - Antigen-Antibody Reactions, 78-7037
- Virus, Herpes Simplex 1
 - Antigens, Viral, 78-6940
- Virus, Herpes Simplex 2
 - Antigens, Viral, 78-6940
- Virus, Murine Leukemia
 - Antigens, Neoplasm, 78-6978
 - Immunity, Cellular, 78-6978
 - Lymphocytes, 78-6978
 - Viral Proteins, 78-6871
- Virus, Polyoma, BK
 - Antibodies, Viral, 78-6905
- Virus, SV40
 - Arming/Potentiating Activity, 78-6985
 - Lymphocytotoxicity, 78-6985

Immunity, Cellular

- Asbestos
 - Occupational Hazard, 78-7061
 - Precancerous Conditions, 78-7061
- BCG
 - Cell Transformation, Neoplastic, 78-7006
- Cervix Neoplasms
 - Virus, Herpes Simplex 2, 78-7065
- Fibrosarcoma
 - T-Lymphocytes, 78-7051
 - Macrophages, 78-7051
- Lymphocytes
 - Effector Cells, 78-6986
 - Immunoglobulins, Surface, 78-6986
 - Interferon, 78-6983
- T-Lymphocytes

Immunity, Cellular (cont'd)

Immunoglobulins, Fc, 78-6987

Lymphoma

Antigenic Determinants, 78-7027
Histocompatibility Antigens, 78-7027
Killer Cells, 78-6883, 78-7009
T-Lymphocytes, 78-6883, 78-7027
Radiation, Ionizing, 78-7027
Virus, Moloney Murine Leukemia, 78-6883, 78-7010
Virus, Rauscher Murine Leukemia, 78-6883

Mammary Neoplasms, Experimental

Adenocarcinoma, 78-7004
Immune Serums, 78-7004
Plant Agglutinins, 78-7004

Mouth Neoplasms

Benzene, 1-Chloro-2,4-dinitro-, 78-7033

Skin Neoplasms

Virus, Papilloma, 78-7036

Virus, Adeno 12

Age Factors, 78-7008
Animal Model, Hamster, 78-7008
Antigens, 78-7008

Virus, Epstein-Barr

Antigenic Determinants, 78-7015
Antigens, Viral, 78-7015
Migration Inhibitory Factor, 78-6989

Virus, Gross Murine Leukemia

Formaldehyde, 78-7024

Virus, Herpes Simplex 2

Antigens, Viral, 78-7065

Virus, Marek's Disease Herpes

Genetic Resistance, 78-7049
Strain Difference, 78-7050

Virus, Murine Leukemia

Antigens, Neoplasm, 78-6978
Immune Serums, 78-6978

Virus, Rauscher Murine Leukemia

Formaldehyde, 78-7024

Virus, Rous Sarcoma

BCG, 78-7006

Virus, SV40

Age Factors, 78-7008
Animal Model, Hamster, 78-7008
Antigens, 78-7008
Freund's Adjuvant, 78-7005
Hamster, 78-6985
Lymphocytes, 78-6985

Warts

Virus, Papilloma, 78-7036

Immunity, Passive**Fibrosarcoma**

Hypersensitivity, Delayed, 78-7054
T-Lymphocytes, 78-7054
Monocytes, 78-7054
Neoplasm Metastasis, 78-7047

Leukemia, Lymphoblastic

Virus, Marek's Disease Herpes, 78-7023

Lung Neoplasms

Neoplasm Metastasis, 78-7047

T-Lymphocytes

Monocytes, 78-7054

Virus, Friend Murine Leukemia

Antigens, Viral, 78-6879

Virus, Marek's Disease Herpes

Cyclophosphamide, 78-7023
Killer Cells, 78-7023
Radiation, Ionizing, 78-7023

Immunity, Passive (cont'd)

Spleen Cells, 78-7023

Immunization

Benzo(a)pyrene
Serum Albumin, Bovine, 78-7059

Fibrosarcoma

Cholanthrene, 3-Methyl-, 78-7053
Radiation, Ionizing, 78-7053

Lymphoma

Brucella abortus, 78-7014

Immunoglobulins**Asbestos**

Occupational Hazard, 78-7061

Asbestosis

Serum Levels, 78-7061

Heavy Chain Disease

Gene Deletion, 78-7060
RNA, Messenger, 78-7060

Leukemia, Lymphoblastic

Transplantation Immunology, 78-7013

Lymphoma

Glomerulonephritis, 78-7084

Immunoglobulins, Fc**Lymphocytes**

Interferon, 78-6982

T-Lymphocytes

Antibody-Dependent Cytotoxicity, 78-6987
Antitumor Activity, K526 Cells, 78-6987
Immunity, Cellular, 78-6987

Immunoglobulins, Heavy Chain**Virus, Rous Sarcoma**

Protein Kinase, 78-6848

Immunoglobulins, Light Chain**Lymphoma**

Sjogren's Syndrome, 78-7082

Salivary Gland Neoplasms

Sjogren's Syndrome, 78-7082

Immunoglobulins, Surface**Leukemia, Lymphoblastic**

T-Lymphocytes, 78-7012

Lymphocytes

Antibody-Dependent Cytotoxicity, 78-6986
Antigen-Antibody Reactions, 78-6986
IgG, 78-6986
Immunity, Cellular, 78-6986

Thymoma

T-Lymphocytes, 78-7012

Immunosuppression**Antigens, Neoplasm**

Cell Transformation, Neoplastic, Review, 78-6688

Carcinogen, Chemical

Epidemiology, Review, 78-6697

Cholanthrene, 11,12-Dihydro-11,12-epoxy-3-methyl-

Cell Transformation, Neoplastic, 78-7110

Cholanthrene, 3-Methyl-

Cell Transformation, Neoplastic, 78-7110

Dibenz(a,h)anthracene, 5,6-Dihydro-5,6-epoxy-

Cell Transformation, Neoplastic, 78-7110

Fibrosarcoma

Neoplasm Transplantation, 78-7047

Spleen Cells, 78-7047

Freund's Adjuvant

T-Lymphocytes, 78-7005

Guanidine, 1-Methyl-3-nitro-1-nitroso-

Immunosuppression (cont'd)

- Cell Transformation, Neoplastic, 78-7110
- Kidney Neoplasms
 - Transplantation, Homologous, 78-6736
- Leukemia
 - Antibody Formation, 78-7021
 - Culture Supernatant, 78-7021
 - DNA Replication, 78-7022
 - T-Lymphocytes, 78-7022
 - Mitomycin C, 78-7022
 - Plant Agglutinins, 78-7021
 - Radiation, Ionizing, 78-7021, 78-7022
 - Spleen/Thymus Cells, 78-7022
 - Surface Properties, 78-7021
 - Transplantation, Homologous, 78-6736
 - Virus, Murine Leukemia, 78-7022
- Liver Neoplasms
 - Transplantation, Homologous, 78-6736
- Lung Neoplasms
 - Neoplasm Transplantation, 78-7047
 - Spleen Cells, 78-7047
- Lymphoma
 - Antigen-Antibody Reactions, 78-7003
 - T-Lymphocytes, 78-7022
 - Neoplasm Transplantation, 78-7028
 - Thymectomy, 78-7028
 - Tumor Extract vs Culture Supernatant, 78-7003
- Oncogenic Viruses
 - Cell Transformation, Neoplastic, Review; 78-6688
- Plasmacytoma
 - Antigen-Antibody Reactions, 78-7003
 - Tumor Extract vs Culture Supernatant, 78-7003
- Sarcoma, Mast Cell
 - Antigen-Antibody Reactions, 78-7003
 - Tumor Extract vs Culture Supernatant, 78-7003
- Virus, Abelson Murine Leukemia
 - Cell Transformation, Neoplastic, 78-7110
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-6952
 - Case Report, 78-6952
 - T-Lymphocytes, 78-6952
- Virus, Friend Murine Leukemia
 - Antigen-Antibody Reactions, 78-7003
 - Tumor Extract vs Culture Supernatant, 78-7003
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-7110

Immunosuppressive Agents

- Leukemia, Myeloblastic
 - Precancerous Conditions, 78-6695
- Lymphoma
 - Precancerous Conditions, 78-6695
- Neoplasms
 - Epidemiology, 78-6736

Indole-3-acetic Acid, 1-(p-Chlorobenzoyl)-5-methoxy-2-methyl-

- Erythema
 - Ultraviolet Rays, 78-6811
- Ultraviolet Rays
 - 5,8,11,14-Eicosatetraenoic Acid, 78-6811
 - Prostaglandins E, 78-6811
 - Prostaglandins F, 78-6811

Indole-3-acrylic Acid

- Fibroma
 - Species Difference, 78-6790
- Fibrosarcoma
 - Species Difference, 78-6790
- Gynecologic Neoplasms

Indole-3-acrylic Acid (cont'd)

- Fibroma, 78-6790
- Granulosa Cell Tumor, 78-6790
- Papilloma, 78-6790
- Species Difference, 78-6790
- Leukemia, Lymphocytic
 - Species Difference, 78-6790
- Leukemia, Myelocytic
 - Species Difference, 78-6790
- Leukemia, Plasmacytic
 - Species Difference, 78-6790
- Liver Neoplasms
 - Cholangioma, 78-6790
 - Species Difference, 78-6790
- Lymphosarcoma
 - Species Difference, 78-6790
- Mammary Neoplasms, Experimental
 - Species Difference, 78-6790
- Pancreatic Neoplasms
 - Adenoma, 78-6790
 - Islet Cell Tumor, 78-6790
 - Species Difference, 78-6790

Infectious Mononucleosis

- Virus, Epstein-Barr
- Migration Inhibitory Factor, 78-6989

Insecticides

- Food Contamination
- Carcinogenic Potential, Review, 78-6662

Insulin

- Cell Division
- Embryo, Mouse, 78-7197

Interferon

- Killer Cells
 - Lymphocyte Culture Test, Mixed, 78-6983
- Leukemia, Lymphoblastic
 - Macrophages, 78-6993
- Lymphocytes
 - Antibody-Dependent Cytotoxicity, 78-6983
 - DNA Replication, 78-6982
 - Immunity, Cellular, 78-6983
 - Immunoglobulins, Fc, 78-6982
 - Isolation and Characterization, 78-6982
 - Killer Cells, 78-6983
 - Lymphocyte Culture Test, Mixed, 78-6982
 - Virus Replication, 78-6982
- Macrophages
 - Prostaglandins E, 78-6993
- Neoplasms
 - Lymphocytes, 78-6983
- Virus, Epstein-Barr
 - Lymphocytes, 78-6982
- Virus, Murine Sarcoma
 - Lymphocytes, 78-6982
- Virus, Newcastle Disease
 - Lymphocytes, 78-6983
- Virus, Vaccinia
 - Lymphocytes, 78-6983
- Virus, Vesicular Stomatitis
 - Lymphocytes, 78-6983

Intestinal Neoplasms

- Adenocarcinoma
 - Carrageen, 78-6801
 - Celiac Disease, 78-7154
 - Cells, Cultured, 78-6680
 - Selenium, 78-6696

Intestinal Neoplasms (cont'd)

- Adenoma
 - Bracken Fern, 78-6717
 - Carrageen, 78-6801
- Bracken Fern
 - Carcinogenic Activity, Rat, 78-6717
- Carcinoma
 - Pediatric Patients, 78-7177
- Carcinoma, Epidermoid
 - Carrageen, 78-6801
- Carcinoma, Papillary
 - Bracken Fern, 78-6717
- Carrageen
 - Carcinogenic Activity, Rat, 78-6801
- Celiac Disease
 - Epidemiology, 78-7154
- Cell Differentiation
 - Precancerous Conditions, 78-7131
- Diet
 - Epidemiology, 78-7176
 - Vegetables, 78-7176
- Intestinal Polyps
 - Diagnosis, Review, 78-6708
 - Genetics, Review, 78-6708
- Peutz-Jeghers Syndrome
 - Genetics, Review, 78-6708
- Polyps
 - Precancerous Conditions, 78-7131
- Precancerous Conditions
 - Thymidine Incorporation, 78-7131
- Selenium
 - Antitumor Activity, Review, 78-6696
 - Epidemiology, Review, 78-6696

Intestinal Polyps

- Intestinal Neoplasms
 - Diagnosis, Review, 78-6708
 - Genetics, Review, 78-6708

Iodine

- Mammary Neoplasms, Experimental
 - Thyroid Dysfunction, Review, 78-6640

Iodine Radioisotopes

- Thyroid Neoplasms
 - Dose-Response Study, 78-6833
 - Mutagenic Activity, 78-6833

Iron

- Asbestos
 - Co-carcinogenic Effect, 78-6654
- Foreign Bodies
 - Co-carcinogenic Effect, 78-6654
- Hydrogen-Ion Concentration
 - Toxicity, 78-6654
- Oxidation-Reduction
 - Carcinogenic Activity, 78-6654
- Oxygen
 - Enzymatic Deactivation, 78-6654
- Phagocytosis
 - Toxicity, 78-6654
- Islet Cell Tumor
 - Pancreatic Neoplasms
 - Acetophenone, 2-Amino-, 78-6790
 - Indole-3-acrylic Acid, 78-6790
 - Ribonuclease, 78-7200

Isoantigens

- T-Lymphocytes

Isoantigens (cont'd)

- Immune Response, 78-7044
- Lymphoma
 - Virus, Radiation Leukemia, 78-7029
- Urea, Ethyl Nitroso-
 - Transplacental Carcinogenesis, 78-6748

Isoproterenol

- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 - Adenyl Cyclase, 78-6766

Karyotyping

- Leukemia, Myelocytic
 - Chromosomes, Human, 21-22, 78-7070
- Sarcoma, Reticulum Cell
 - Lymph Nodes, 78-7080

Keratoacanthoma

- Skin Neoplasms
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753

Keratinosis

- Aging
 - DNA Repair, 78-7091
- Leukocytes
 - DNA Repair, 78-7091
- Lymphocytes
 - DNA Repair, 78-7090
- Skin Neoplasms
 - Ultraviolet Rays, 78-6813
- Ultraviolet Rays
 - DNA Repair, 78-7090, 78-7091
- Urea, Hydroxy-
 - DNA Repair, 78-7090

Kethoxal

- see* 2-Butanone, 1,1-Dihydroxy-3-ethoxy-

Kidney Neoplasms

- Adenoma
 - Streptozotocin, 78-6799
- Carcinoma, Basal Cell
 - Case Report, 78-7129
 - Neoplasm Metastasis, 78-7129
- Dimethylamine, *N*-Nitroso-
 - Carboxy-Lyases, 78-6742
 - Dietary Proteins, 78-6742
 - Histological Study, 78-6739
 - Neoplasm Transplantation, 78-6739
 - Ultrastructural Study, 78-6739
- Fibrosarcoma
 - Dimethylamine, *N*-Nitroso-, 78-6739
- Methanol, (Methyl-*ONN*-azoxy)-
 - Pyrazole, 78-6760
- Neoplasm Metastasis
 - Calcaneus, 78-7129
- Plasmacytoma
 - Case Report, 78-7128
 - Ultrastructural Study, Giant Cells, 78-7128
- Polyamines
 - Carboxy-Lyases, 78-6742
- Streptozotocin
 - Dietary Proteins, 78-6799
- Transplantation, Homologous
 - Immunosuppression, 78-6736
- Urea, Ethyl Nitroso-
 - Transplacental Carcinogenesis, 78-6748

L Cells

- Cholanthrene, 3-Methyl-
 - Proteins, 78-6908

L Cells (cont'd)

- Hybrid Cells
 - LETS Protein, 78-7077
- Virus, Polio
 - RNA Replication, 78-6971

Lactate Dehydrogenase Isoenzymes

- Giant Cell Tumors
 - Cells, Cultured, 78-7148
- Sarcoma, Osteogenic
 - Cells, Cultured, 78-7148

Laryngeal Neoplasms

- Ethyl Alcohol
 - Epidemiology, Review, 78-6699
- Leukoplakia
 - Enzymatic Activity, 78-7099
 - Neutrophils, 78-7099
- Melanoma
 - Antigenic Determinants, 78-7037
- Neutrophils
 - Alkaline Phosphatase, 78-7099
 - Enzymatic Activity, 78-7099
 - Glucuronidase, 78-7099
 - Lipids, 78-7099
 - Precancerous Conditions, 78-7099
- Papilloma
 - Enzymatic Activity, 78-7099
 - Neutrophils, 78-7099
 - Virus-Like Particles, 78-6900
- Precancerous Conditions
 - Enzymatic Activity, 78-7099
- Virus, Papilloma
 - Isolation and Characterization, 78-6900

Laurence-Moon-Biedl Syndrome

- Uterine Neoplasms
 - Adenocarcinoma, 78-7133
 - Case Report, 78-7133

Letterer-Siwe Disease

- Virus-Like Particles
 - Ultrastructural Study, 78-6977

Leukemia

- T-Lymphocytes
 - Antilymphocyte Serum, 78-7067
 - Immunosuppression, Rat, 78-7067
- Radiation, Ionizing
 - Immunosuppression, Rat, 78-7067
- Antibody Formation
 - Cells, Cultured, 78-7021
 - Immunosuppression, 78-7021
- Antigens
 - Immune Response, Review, 78-6681
 - Phenotype, Review, 78-6681
- Antigens, Neoplasm
 - Benzene, 1-Fluoro-2,4-dinitro-, 78-7025
 - Leukocytes, 78-7025
- Antigens, Viral
 - Immune Response, 78-7016
- Burkitt's Lymphoma
 - Transplantation, Heterologous, 78-7086
- Cell Differentiation
 - Phenotype, Review, 78-6681
 - Precancerous Conditions, Review, 78-6681
- Chromosome Aberrations
 - Precancerous Conditions, Review, 78-6681
- Dietary Fats
 - Epidemiology, Review, 78-6699

Leukemia (cont'd)

- Ethylene, Trichloro-
 - Epidemiology, 78-7173
 - Occupational Hazard, 78-7173
- Hair Dyes
 - Epidemiology, 78-7166
- Immunosuppression
 - Culture Supernatant, 78-7021
 - DNA Replication, 78-7022
 - Spleen/Thymus Cells, 78-7022
 - Surface Properties, 78-7021
- Interspecies Transmission
 - Interspecies Transmission, 78-6672
- T-Lymphocytes
 - Antibody Formation, 78-7022
 - Immunosuppression, 78-7022
- Mitomycin C
 - Immunosuppression, 78-7022
- Neoplasms, Multiple Primary
 - Actinomycin D, 78-7072
 - Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-78-7072
 - Cyclophosphamide, 78-7072
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-7072
 - Radiation, Ionizing, 78-7072
- Plant Agglutinins
 - Immunosuppression, 78-7021
- Polyribonucleotides
 - Drug Therapy, Review, 78-6669
- Radiation, Ionizing
 - Case Report, 78-7072
 - Chromosome Abnormalities, 78-7072
 - Epidemiology, Review, 78-6667
 - Immunosuppression, 78-7021, 78-7022
 - Occupational Hazard, 78-6667
- Radium
 - Dose-Response Study, Mouse, 78-6830
- Sex Factors
 - Epidemiology, Minnesota, 78-7167
- Socioeconomic Factors
 - Epidemiology, Review, 78-6698
- Solvents
 - Case Report, 78-6765
 - Epidemiology, Child, 78-6765
- Thorium
 - Dose-Response Study, Mouse, 78-6830
 - Radioisotopes, 78-6830
- Transplantation, Homologous
 - Immunosuppression, 78-6736
- Virus, Baboon
 - Isolation and Characterization, Review, 78-6672
- Virus, C-Type RNA Tumor
 - Ultrastructural Study, Review, 78-6679
 - Virus-Like Particles, Review, 78-6672
- Virus, Feline Leukemia
 - Horizontal Transmission, 78-7165
- Virus, Gibbon Ape Lymphoma
 - Isolation and Characterization, Review, 78-6672
- Virus, Gross Murine Leukemia
 - Viral Proteins, 78-6880
- Virus, Murine Leukemia
 - Antibody Formation, 78-7021, 78-7022
 - Immunosuppression, 78-7022
- Virus, Murine Mammary Tumor
 - Antigens, Viral, 78-7016
- Virus, Rauscher Murine Leukemia
 - Antibody Formation, 78-7021

Leukemia (cont'd)

- Virus, Simian Sarcoma
Isolation and Characterization, Review, 78-6672
- Virus, SV40
Genetics, 78-6934
Twins, 78-6934

Leukemia L1210

- Ascitic Fluid
Anti-Inflammatory Activity, 78-6981
- BCG*
Immune Response, 78-7020
- Guanidine, 1-Methyl-3-nitro-1-nitroso-
Histones, Binding, 78-6752
- Lyases
Enzymatic Activity, Liver, 78-7187
- Macrophages
Growth, 78-7000
- Micrococcus lysodeikticus*
Immune Response, 78-7020
- Polysaccharides
Immune Response, 78-7020
- Radiation, Ionizing
Immunotherapy, 78-7020
- Ultraviolet Rays
1,8,9-Anthracenetriol, 78-6807
DNA Repair, 78-6807
Quinacrine, 78-6807
- Urea, Methyl Nitroso-
Histones, Binding, 78-6752
- Zymosan
Immune Response, 78-7020

Leukemia, Lymphoblastic

- Acetone
Case Report, 78-6765
- Antibody Formation
Transplantation Immunology, 78-7013
- Antigen-Antibody Complex
Transplantation Immunology, 78-7012
- Antigen-Antibody Reactions
Precancerous Conditions, 78-7026
- Antigenic Determinants
Bone Marrow Cells, 78-7026
Hematopoietic Stem Cells, 78-7026
B-Lymphocytes, 78-7026
Null Cells, 78-7026
Precancerous Conditions, 78-7026
- Antigens, Neoplasm
Transplantation Immunology, 78-7013
- Ataxia Telangiectasia
Case Report, 78-7030
Chromosome Aberrations, 78-7030
Chromosomes, Human, 13-15, 78-7030
- Cellular Inclusions
Ultrastructural Study, 78-6977
- Ethylene, Trichloro-
Case Report, 78-6765
- Hepatitis
Case Report, Child, 78-6956
- Immunoglobulins
Transplantation Immunology, 78-7013
- T-Lymphocytes
Antigen-Antibody Complex, 78-7012, 78-7013
Immunoglobulins, Surface, 78-7012
Transplantation Immunology, 78-7013
- Macrophages
Antitumor Activity, 78-6993

Leukemia, Lymphoblastic (cont'd)

- Cortisol, 78-6993
- Dibutyl Cyclic AMP, 78-6993
- Interferon, 78-6993
- Prostaglandins E, 78-6993
- Methane, Chlorodifluoro-
Case Report, 78-6765
In Utero Exposure, 78-6765
- Styrene
Case Report, 78-6765
- Virus, Bovine Leukemia
Epidemiology, 78-7169
- Virus, Herpes
Case Report, Child, 78-6956
Precancerous Conditions, 78-6956
- Virus-Like Particles
Ultrastructural Study, 78-6977
- Virus, Marek's Disease Herpes
Immunity, Passive, 78-7023
- Virus, SV40
Twins, 78-6934
- Xylene
Case Report, 78-6765
In Utero Exposure, 78-6765

Leukemia, Lymphocytic

- Acetophenone, 2-Amino-
Species Difference, 78-6790
- Antibody Formation
Immune Response, Review, 78-6689
- IgM
Antibody Specificity, 78-7073
Antigenic Determinants, 78-7073
- Indole-3-acrylic Acid
Species Difference, 78-6790
- B-Lymphocytes
Clone Cells, 78-7073
IgM, 78-7073
Immune Response, Review, 78-6689

Leukemia, Monoblastic

- Virus, Leukemia
Isolation and Characterization, 78-6975
T-Lymphocytes, 78-6975
Virus-Like Particles, 78-6975

Leukemia, Myeloblastic

- Actinomycin D
Drug Therapy, 78-6815
- Antigen-Antibody Reactions
Precancerous Conditions, 78-7026
- Antigenic Determinants
Bone Marrow Cells, 78-7026
Hematopoietic Stem Cells, 78-7026
- Antigens
Antibody Specificity, 78-7019
Cell Membrane, 78-7019
- Benzene
Epidemiology, Review, 78-6627
- Bronchitis
Precancerous Conditions, 78-7074
- Cyclophosphamide
Drug Therapy, 78-6815
- Drug Therapy
Case Report, 78-6815
- Erythrocytes
Colony Formation, 78-7150
- Immune Serums
Antibody-Dependent Cytotoxicity, 78-7019

Leukemia, Myeloblastic (cont'd)

- Immunosuppressive Agents
 - Precancerous Conditions, 78-6695
- Leukocytes
 - Colony Formation, 78-7150
- T-Lymphocytes
 - Cell Differentiation, 78-7150
- Neoplasms, Multiple Primary
 - Drug Therapy, 78-6736
 - Virus, Epstein-Barr, 78-7071
- Plant Agglutinins
 - Colony Formation, 78-7150
- Precancerous Conditions
 - Case Report, 78-7074
- Radiation, Ionizing
 - Case Report, 78-6815
- Sarcoma, Ewing's
 - Neoplasms, Multiple Primary, 78-6815
- Sex Factors
 - Epidemiology, Minnesota, 78-7167
- Thymidine
 - Cell Cycle Kinetics, 78-7150
- Urea, Hydroxy-
 - Cell Cycle Kinetics, 78-7150
- Vincristine
 - Drug Therapy, 78-6815
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-7071
 - Case Report, 78-7071

Leukemia, Myelocytic

- Acetophenone, 2-Amino-
 - Species Difference, 78-6790
- Chromosome Abnormalities
 - Aneuploidy, 78-7070
- Chromosomes, Human, 21-22
 - Karyotyping, 78-7070
 - Radiation, Ionizing, 78-6827
- Indole-3-acrylic Acid
 - Species Difference, 78-6790
- Radiation, Ionizing
 - Case Report, 78-6827
- Sex Chromosomes
 - Blast Crisis, Shay Tumor, 78-7069
 - Neoplasm Transplantation, 78-7069

Leukemia, Plasmacytic

- Acetophenone, 2-Amino-
 - Species Difference, 78-6790
- Indole-3-acrylic Acid
 - Species Difference, 78-6790

Leukemia, Radiation-Induced

- DNA Nucleotidyltransferases
 - Bone Marrow, 78-6826
 - Enzymatic Activity, Mouse, 78-6826
 - Thymus Gland, 78-6826
- Radiation, Ionizing
 - DNA Nucleotidyltransferases, 78-6826

Leukemia, Subleukemic

- Virus, SV40
 - Twins, 78-6934

Leukocytes

- Keratosis
 - DNA Repair, 78-7091
- Leukemia
 - Antigens, Neoplasm, 78-7025
- Leukemia, Myeloblastic

Leukocytes (cont'd)

- Colony Formation, 78-7150
- Neoplasms
 - Immunity, Cellular, Review, 78-6690
- Ultraviolet Rays
 - DNA Repair, 78-7091
- Virus, Epstein-Barr
 - Migration Inhibitory Factor, 78-6989

Leukoplakia

- Laryngeal Neoplasms
 - Enzymatic Activity, 78-7099
 - Neutrophils, 78-7099

LH

- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6778

Lipids

- Laryngeal Neoplasms
 - Neutrophils, 78-7099

Lipopolysaccharides

- Macrophages
 - Cytotoxic Activity, 78-7002
 - 4,4'-Stilbene diol, α,α' -Diethyl-
 - Antibody Formation, 78-7011
- Virus, Adeno 12
 - Antibody Formation, 78-7008
- Virus, SV40
 - Antibody Formation, 78-7008

Listeria monocytogenes

- Silicic Acid, Magnesium Salt
 - Immune Response, 78-6992
 - Macrophage Activation, 78-6992

Liver

- Acetamide, *N*-Fluoren-2-yl-
 - Metabolism, Rat, 78-6731
 - Mutagenic Activation, 78-6730
- Acetohydroxamic Acid, *N*-Fluoren-2-yl-
 - Mutagenic Activation, 78-6730
 - Nucleic Acids, Binding, 78-6730
- Diethylamine
 - Tissue Distribution, Guinea Pig, 78-6747
- Dimethylamine
 - Tissue Distribution, Guinea Pig, 78-6747

Liver Cirrhosis

- Australia Antigen
 - Antigen-Antibody Reactions, 78-6961
 - Epidemiology, 78-6961
- Hepatoma
 - Australia Antigen, 78-6961
 - Epidemiology, Japan, 78-7127
 - Phosphodiesterases, 78-6960
 - Precancerous Conditions, 78-7123
 - Virus, Hepatitis, 78-6958
 - Virus-Like Particles, 78-6960

Liver Neoplasms

- Acetophenone, 2-Amino-
 - Species Difference, 78-6790
- Adenocarcinoma
 - Mycotoxins, 78-6723
- Adenoma
 - Contraceptives, Oral, 78-6645, 78-6650, 78-6795
 - 78-7126
- Aldrin
 - Epidemiology, Review, 78-6659

Liver Neoplasms (cont'd)

- Antigens
 - Isolation and Characterization, Review, 78-6692
- Barbituric Acid, 5-Ethyl-5-phenyl-
 - Dietary Proteins, 78-6742
- Barbituric Acid, 5-Ethyl-5-phenyl-, Sodium Salt
 - Carcinogenic Potential, Rat, 78-6758
 - Hyperplasia, 78-6758
- Benzidine, Dihydrochloride
 - Neoplasm Metastasis, 78-7122
- Benzo(a)pyrene
 - Benzidine, Dihydrochloride, 78-7122
- Carcinogen, Chemical
 - Antigens, 78-6692
 - Carcinoembryonic Antigen, 78-6692
 - Choline, 78-6638
 - Chromosomes, Review, 78-6706
 - Diet, Review, 78-6638
 - Glycogen, 78-6707
 - Histocompatibility Antigens, 78-6692
 - Methionine, 78-6638
 - Precancerous Conditions, Review, 78-6706, 78-6707
 - Ultrastructural Study, Review, 78-6707
 - Vitamin B12, 78-6638
- Carcinoma
 - Alpha Fetoproteins, 78-7062
 - Radioimmunoassay, 78-7062
- Cholangioma
 - Acetophenone, 2-Amino-, 78-6790
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6767
 - Indole-3-acrylic Acid, 78-6790
- Contraceptives, Oral
 - Case Report, 78-6795, 78-7125, 78-7126
 - Epidemiology, Review, 78-6644, 78-6645, 78-6646
 - 78-6650
 - Estradiol, 17-Ethynyl-, 78-7126
 - Hyperplasia, 78-6795
 - Norethisterone, 78-7126
 - Precancerous Conditions, Review, 78-6646
 - Tumor Hemorrhage, 78-7126
 - Ultrastructural Study, 78-7125
 - Ultrastructural Study, Review, 78-6645
- Dieldrin
 - Epidemiology, Review, 78-6659
- Diet
 - Epidemiology, Review, 78-6637
- Diethylamine, *N*-Nitroso-
 - Benzidine, Dihydrochloride, 78-7122
 - Carboxy-Lyases, 78-6742
 - Dietary Proteins, 78-6742
- Direct Black 38
 - Precancerous Conditions, Mouse, 78-6771
- Direct Blue 6
 - Precancerous Conditions, Mouse, 78-6771
- Direct Brown 95
 - Precancerous Conditions, Mouse, 78-6771
- Estrogens
 - Epidemiology, Review, 78-6650
- Ethane, 1,1,1-Trichloro-2,2-bis(*p*-chlorophenyl)-
 - Epidemiology, Review, 78-6659
- Ethylene, Trichloro-
 - Epidemiology, 78-7173
 - Occupational Hazard, 78-7173
- Glucosephosphate Dehydrogenase
 - Precancerous Conditions, 78-6787
- Hemangioma, Cavernous

Liver Neoplasms (cont'd)

- Case Report, 78-7098
- Neoplasms, Multiple Primary, 78-7098
- Hepatoma
 - Alpha Fetoproteins, 78-7062
 - Radioimmunoassay, 78-7062
- Hyperplasia
 - Contraceptives, Oral, 78-6645, 78-6650, 78-7125
 - Estradiol, 17-Ethynyl-, 78-7125
 - Mestranol, 78-7125
 - Precancerous Conditions, Review, 78-6707
- Indole-3-acrylic Acid
 - Species Difference, 78-6790
- Morpholine, *N*-Nitroso-
 - Glucosephosphate Dehydrogenase, 78-6787
- Mycotoxins
 - Ultrastructural Study, Rat, 78-6723
- Neoplasm Metastasis
 - Histological Study, 78-7122
 - Lung, Mouse, 78-7122
- Polyamines
 - Carboxy-Lyases, 78-6742
- Progestational Hormones
 - Epidemiology, Review, 78-6650
- Propane, 2-Nitro-
 - Inhalation Study, Rat, 78-6734
 - Precancerous Conditions, 78-6734
- Transplantation, Homologous
 - Immunosuppression, 78-6736
 - Precancerous Conditions, Review, 78-6706
- Urea, Ethyl Nitroso-
 - Neoplasm Metastasis, 78-7122
- Virus, Marek's Disease Herpes
 - Antibodies, Viral, 78-6861
- Xanthine, 7-Hydroxy-
 - Carcinogenic Activity, Rat, 78-6718

Lung

- Choriocarcinoma
 - Neoplasm Metastasis, 78-7134
- Diethylamine
 - Tissue Distribution, Guinea Pig, 78-6747
- Dimethylamine
 - Tissue Distribution, Guinea Pig, 78-6747

Lung Neoplasms

- Acrolein
 - Smoke Condensate, Review, 78-6657
- Adenocarcinoma
 - Needle Biopsy Implantation, Case Report, 78-7112
- Air Pollution
 - Epidemiology, Review, 78-6701
- Alpha Particles
 - Risk Estimation, 78-6821
- Americium Oxide
 - Inhalation Study, Rat, 78-6820
 - Radioisotopes, 78-6820
- Asbestos
 - Epidemiology, Review, 78-6631
 - Precancerous Conditions, 78-7061
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Benzo(a)pyrene
 - Dose-Response Study, Review, 78-6701
 - Epidemiology, Review, 78-6701
 - Occupational Hazard, Review, 78-6701
 - Smoke Condensate, Review, 78-6657
- Bone Marrow Cells

Lung Neoplasms (cont'd)

- Colony Formation, 78-6999
- Bradykinin
 - Macrophages, 78-7001
 - Tumor Cell Supernatant, 78-7001
- Carcinoma
 - Foreign Body Reaction, 78-6992
 - Neoplasm Metastasis, 78-6998
 - Phagocytosis, 78-6998
- Cells, Cultured
 - Neoplasm Metastasis, 78-7109
- Corticotropin
 - Isolation and Characterization, 78-6793
- Cortisone Acetate
 - Neoplasm Metastasis, 78-6998
 - Phagocytosis, 78-6998
- Corynebacterium parvum*
 - Neoplasm Metastasis, 78-6998
 - Phagocytosis, 78-6998
- Curium Oxide
 - Inhalation Study, Rat, 78-6820
 - Radioisotopes, 78-6820
- Ethylene, Trichloro-
 - Epidemiology, 78-7173
- Foreign Body Reaction
 - Immune Response, 78-6992
- Formaldehyde
 - Smoke Condensate, Review, 78-6657
- Hydrocyanic Acid
 - Smoke Condensate, Review, 78-6657
- Immune Serums
 - Macrophages, 78-6999
- Immunity, Passive
 - Neoplasm Metastasis, 78-7047
- Immunosuppression
 - Spleen Cells, 78-7047
- B-Lymphocytes
 - Neoplasm Metastasis, 78-7047
- T-Lymphocytes
 - Neoplasm Metastasis, 78-7047
- Lymphoma
 - Lupus Erythematosus, Systemic, 78-7111
- Macrophages
 - Carbon Clearance, 78-6999
 - Chemotaxis, 78-6999
 - Colony Formation, 78-6999
 - Tumor Cell Supernatant, 78-7001
- Neoplasm Transplantation
 - Immunosuppression, 78-7047
- Nickel
 - Epidemiology, 78-7156
 - Occupational Hazard, 78-7156
- Occupational Hazard
 - Epidemiology, 78-7155
- Pesticides
 - Epidemiology, 78-7174
 - Occupational Hazard, 78-7174
- Phagocytosis
 - Growth, 78-6998
 - Neoplasm Metastasis, 78-6998
- Plutonium
 - Risk Estimation, 78-6821
- Plutonium Oxide
 - Inhalation Study, Rat, 78-6819, 78-6820
 - Radioisotopes, 78-6820
- Silica
 - Neoplasm Metastasis, 78-6998

Lung Neoplasms (cont'd)

- Phagocytosis, 78-6998
- Silicic Acid, Magnesium Salt
 - Immune Response, 78-6992
 - Macrophage Activation, 78-6992
- Smoking
 - Epidemiology, 78-7156
- 4,4'-Stilbenediol, α, α' -Diethyl-
 - Phagocytosis, 78-6998
- Trace Elements
 - Epidemiology, Review, 78-6631
- Trypan Blue
 - Phagocytosis, 78-6998
- Uranium
 - Dose-Response Study, 78-7155
 - Epidemiology, 78-7155
- Virus, Marek's Disease Herpes
 - Antibodies, Viral, 78-6861
- Lupus Erythematosus, Systemic**
 - Lung Neoplasms
 - Lymphoma, 78-7111
 - Lymphoma
 - Case Report, 78-7111
 - Virus, C-Type RNA Tumor
 - Ultrastructural Study, Review, 78-6679
- Luteoskyrin**
 - Mycotoxins
 - Carcinogenic Activity, Review, 78-6629
- Lyases**
 - Leukemia L1210
 - Enzymatic Activity, Liver, 78-7187
 - Sarcoma 180, Crocker
 - Enzymatic Activity, Liver, 78-7187
- Lymph Nodes**
 - Complement
 - Transplantation Immunology, 78-7044
 - Glomerulonephritis
 - Hyperplasia, 78-7084
 - Hodgkin's Disease
 - DNA Replication, 78-7192
 - Sarcoma, Reticulum Cell
 - Karyotyping, 78-7080
- Lymphocyte Depletion**
 - Graft vs Host Reaction
 - T-Lymphocytes, 78-7045
 - Lymphoma
 - T-Lymphocytes, 78-7028
 - Myosarcoma
 - B-Lymphocytes, 78-7046
 - Virus, Marek's Disease Herpes
 - T-Lymphocytes, 78-7050
- Lymphocyte Transformation**
 - Cervix Neoplasms
 - Virus, Herpes Simplex 1, 78-7065
 - Virus, Herpes Simplex 2, 78-7065
 - Virus, Epstein-Barr
 - Dwarfism, 78-6947
 - Ultraviolet Rays, 78-6947
 - Xeroderma Pigmentosum, 78-6947
 - Virus, Leukemia
 - B-Lymphocytes, 78-6975
 - Virus, Marek's Disease Herpes
 - Concanavalin A, 78-7049
 - Virus, Rauscher Murine Leukemia

Lymphocyte Transformation (cont'd)

- Concanavalin A, 78-7017
- Plant Agglutinins, 78-7017
- Viral Proteins, 78-7017

Lymphocytes

- Antibody-Dependent Cytotoxicity
 - Effector Cells, 78-6986
- Antigens, Neoplasm
 - Cytotoxicity, 78-6988
- Antilymphocyte Serum
 - IgG, 78-6986
- Aryl Hydrocarbon Hydroxylases
 - Enzyme Induction, 78-6788
 - Phenotype, 78-6788
- Benz(a)anthracene
 - Aryl Hydrocarbon Hydroxylases, 78-6781
- 5,6-Benzoflavone
 - Aryl Hydrocarbon Hydroxylases, 78-6781
- Bladder Neoplasms
 - Concanavalin A, 78-6735
 - Immune Response, Rat, 78-6735
 - Plant Agglutinins, 78-6735
- Cholanthrene, 3-Methyl-
 - Aryl Hydrocarbon Hydroxylases, 78-6781
 - Cytochrome Reductases, 78-6781
- Concanavalin A
 - Cytotoxicity, 78-6988
- DNA
 - Nucleic Acid Release, 78-7191
- Ethanol, 2-Mercapto-
 - Cytotoxicity, 78-6988
- Immunity, Cellular
 - Effector Cells, 78-6986
- Immunoglobulins, Surface
 - Antibody-Dependent Cytotoxicity, 78-6986
 - Antigen-Antibody Reactions, 78-6986
 - IgG, 78-6986
 - Immunity, Cellular, 78-6986
- Interferon
 - Antibody-Dependent Cytotoxicity, 78-6983
 - DNA Replication, 78-6982
 - Immunity, Cellular, 78-6983
 - Immunoglobulins, Fc, 78-6982
 - Isolation and Characterization, 78-6982
 - Killer Cells, 78-6983
 - Lymphocyte Culture Test, Mixed, 78-6982
 - Virus Replication, 78-6982
- Keratosis
 - DNA Repair, 78-7090
- Lymphosarcoma
 - Neoplasm Regression, Spontaneous, 78-6980
 - Peritoneal Cavity, Mouse, 78-6980
- Mitomycin C
 - Cytotoxicity, 78-6988
- Neoplasms
 - Interferon, 78-6983
 - Killer Cells, 78-6983
- Peritoneal Fluid
 - Cytotoxicity, 78-6988
- Plant Agglutinins
 - Cytotoxicity, 78-6988
- Radiation, Ionizing
 - Cysteine, 78-6979
 - Ethylamine, 2,2'-Dithiobis-, 78-6979
 - Glutathione, 78-6979
 - Pseudourea, 2-(2-Aminoethyl)-2-thio-, 78-6979
 - Radiation-Protective Agents, 78-6979

Lymphocytes (cont'd)

- Serotonin, 78-6979
- Virus, Epstein-Barr
 - Interferon, 78-6982
- Virus, Murine Leukemia
 - Immune Serums, 78-6978
 - Plant Agglutinins, 78-6978
- Virus, Murine Sarcoma
 - Interferon, 78-6982
- Virus, Newcastle Disease
 - Interferon, 78-6983
- Virus, SV40
 - Immunity, Cellular, 78-6985
- Virus, Vaccinia
 - Interferon, 78-6983
- Virus, Vesicular Stomatitis
 - Interferon, 78-6983

B-Lymphocytes

- Asbestos
 - Precancerous Conditions, 78-7061
- Asbestosis
 - Occupational Hazard, 78-7061
- Fibrosarcoma
 - Neoplasm Metastasis, 78-7047
- Leukemia, Lymphoblastic
 - Antigenic Determinants, 78-7026
- Leukemia, Lymphocytic
 - Clone Cells, 78-7073
 - IgM, 78-7073
 - Immune Response, Review, 78-6689
- Lung Neoplasms
 - Neoplasm Metastasis, 78-7047
- Lymphoma
 - Neoplasm Metastasis, 78-6875
 - Sjogren's Syndrome, 78-7082
- Mycosis Fungoides
 - Chromosome Abnormalities, 78-7120
 - Plant Agglutinins, 78-7120
- Myosarcoma
 - Lymphocyte Depletion, 78-7046
 - Splenectomy, 78-7046
- Sarcoma
 - Benzo(a)pyrene, 78-7045
 - Graft vs Host Reaction, 78-7045
- Skin Neoplasms
 - Melanoma, 78-6693
- Virus, Epstein-Barr
 - Antibody Specificity, 78-7015
 - Antigens, Viral, 78-7015
- Virus, Leukemia
 - Cells, Cultured, 78-6975
 - Lymphocyte Transformation, 78-6975

T-Lymphocytes

- Fibrosarcoma
 - Antilymphocyte Serum, 78-7067
 - Immunosuppression, Rat, 78-7067
- Leukemia
 - Antilymphocyte Serum, 78-7067
 - Immunosuppression, Rat, 78-7067
- Mammary Neoplasms, Experimental
 - Antilymphocyte Serum, 78-7067
 - Immunosuppression, Rat, 78-7067
- Antilymphocyte Serum
 - Transplantation Immunology, 78-7044
- Asbestos
 - Precancerous Conditions, 78-7061

T-Lymphocytes (cont'd)

- Asbestosis
 - Occupational Hazard, 78-7061
- Benz(a)anthracene, 7,12-Dimethyl-
 - Cell Transformation, Neoplastic, 78-6984
 - Rosette Formation, 78-6984
- Cells, Cultured
 - Antitumor Activity, K526 Cells, 78-6987
- Cholanthrene, 3-Methyl-
 - Antilymphocyte Serum, 78-7041
- Fibrosarcoma
 - Corynebacterium parvum*, 78-7056
 - Growth, 78-7053
 - Immunity, Cellular, 78-7051
 - Immunity, Passive, 78-7054
 - Neoplasm Metastasis, 78-7047
 - Transplantation Immunology, 78-7051, 78-7054
- Freund's Adjuvant
 - Immunosuppression, 78-7005
- Graft vs Host Reaction
 - Lymphocyte Depletion, 78-7045
- Histocompatibility Antigens
 - Immune Response, 78-7044
- Immunoglobulins, Fc
 - Antibody-Dependent Cytotoxicity, 78-6987
 - Antitumor Activity, K526 Cells, 78-6987
 - Immunity, Cellular, 78-6987
- Isoantigens
 - Immune Response, 78-7044
- Leukemia
 - Antibody Formation, 78-7022
 - Immunosuppression, 78-7022
- Leukemia, Lymphoblastic
 - Antigen-Antibody Complex, 78-7012, 78-7013
 - Immunoglobulins, Surface, 78-7012
 - Transplantation Immunology, 78-7013
- Leukemia, Monoblastic
 - Virus, Leukemia, 78-6975
- Leukemia, Myeloblastic
 - Cell Differentiation, 78-7150
- Lung Neoplasms
 - Neoplasm Metastasis, 78-7047
- Lymphoma
 - Antibody Formation, 78-7022
 - Immunity, Cellular, 78-6883, 78-7027
 - Immunosuppression, 78-7022
 - Lymphocyte Depletion, 78-7028
 - Neoplasm Metastasis, 78-6875
 - Virus, Radiation Leukemia, 78-7029
- Monocytes
 - Immunity, Passive, 78-7054
- Mycosis Fungoides
 - Chromosome Abnormalities, 78-7120
 - Plant Agglutinins, 78-7120
- Myosarcoma
 - Splenectomy, 78-7046
- Neoplasms
 - Immunity, Cellular, Review, 78-6690
- Neoplasms, Experimental
 - Immunosurveillance, Review, 78-6688
- Sarcoma
 - Benzo(a)pyrene, 78-7045
 - Graft vs Host Reaction, 78-7045
- Skin Neoplasms
 - Melanoma, 78-6693
- 4,4'-Stilbenediol, α,α' -Diethyl-
 - Immune Response, 78-7011

T-Lymphocytes (cont'd)

- Thymoma
 - Antigen-Antibody Complex, 78-7012
 - Immunoglobulins, Surface, 78-7012
- Virus, Adeno 12
 - Cell Transformation, Neoplastic, 78-6984
 - Rosette Formation, 78-6984
- Virus, Epstein-Barr
 - Immunosuppression, 78-6952
- Virus, Marek's Disease Herpes
 - Concanavalin A, 78-7050
 - Lymphocyte Depletion, 78-7050
- Virus, Murine Leukemia
 - Genes, Viral, 78-6677
 - Precancerous Conditions, Review, 78-6677
- Virus, Rauscher Murine Leukemia
 - Glycoproteins, 78-6890
 - Histocompatibility Antigens, 78-6890
- Virus, SV40
 - Cell Transformation, Neoplastic, 78-6984
 - Freund's Adjuvant, 78-7005
 - Rosette Formation, 78-6984
- Lymphoma (General and Unspecified)**
 - Antigen-Antibody Reactions
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
 - Antigenic Determinants
 - Killer Cells, 78-7009
 - Antigens
 - Immune Response, Review, 78-6681
 - Antigens, Neoplasm
 - Cross Reactions, 78-7027
 - Antigens, Viral
 - Cells, Cultured, 78-7010
- BCG
 - Killer Cells, 78-7009
- Brucella abortus*
 - Dose-Response Study, Mouse, 78-7014
 - Immune Response, Mouse, 78-7014
 - Immunization, 78-7014
- Carrageen
 - Cytotoxic Activity, 78-6997
 - Growth, 78-6997
- Cell Differentiation
 - Phenotype, Review, 78-6681
 - Precancerous Conditions, Review, 78-6681
- Cell Membrane
 - Immune Response, 78-7018
- Cells, Cultured
 - Killer Cells, 78-7010
- Chromosome Aberrations
 - Precancerous Conditions, 78-6695
- Epidemiology
 - Iran, 78-7153
- Glomerulonephritis
 - Case Report, Castleman's Tumor, 78-7084
 - Immunoglobulins, 78-7084
- Graft vs Host Reaction
 - Thymectomy, 78-7028
- Histocompatibility Antigens
 - Cells, Cultured, 78-7010
- Immunity, Cellular
 - Antigenic Determinants, 78-7027
 - Histocompatibility Antigens, 78-7027
 - Killer Cells, 78-6883, 78-7009
- Immunosuppression
 - Neoplasm Transplantation, 78-7028

Lymphoma (General and Unspecified) (cont'd)

- Thymectomy, 78-7028
- Tumor Extract vs Culture Supernatant, 78-7003
- Immunosuppressive Agents
 - Precancerous Conditions, 78-6695
- Lung Neoplasms
 - Lupus Erythematosus, Systemic, 78-7111
- Lupus Erythematosus, Systemic
 - Case Report, 78-7111
- B-Lymphocytes
 - Neoplasm Metastasis, 78-6875
- T-Lymphocytes
 - Antibody Formation, 78-7022
 - Immunity, Cellular, 78-6883, 78-7027
 - Immunosuppression, 78-7022
 - Lymphocyte Depletion, 78-7028
 - Neoplasm Metastasis, 78-6875
- Macrophages
 - Cytotoxic Activity, 78-7002
- Nasopharyngeal Neoplasms
 - Virus, Epstein-Barr, 78-7034
- Nitrous Acid
 - Carcinogenic Potential, Review, 78-6655
- Radiation, Ionizing
 - Immunity, Cellular, 78-7027
- Sjogren's Syndrome
 - Histological Study, 78-7082
 - IgM, 78-7082
 - Immunoglobulins, Light Chain, 78-7082
 - B-Lymphocytes, 78-7082
- Ultraviolet Rays
 - Antigens, Viral, 78-7018
- Virus, Abelson Murine Leukemia
 - Neoplasm Metastasis, 78-6875
- Virus, C-Type RNA Tumor
 - Ultrastructural Study, Review, 78-6679
- Virus, Epstein-Barr
 - Migration Inhibitory Factor, 78-6989
- Virus, Feline Leukemia
 - Horizontal Transmission, 78-7165
- Virus, Marek's Disease Herpes
 - Antibodies, Viral, 78-6861
- Virus, Moloney Murine Leukemia
 - Antigen-Antibody Reactions, 78-6883
 - Antigens, Viral, 78-7010, 78-7018
 - Cell Membrane, 78-7018
 - Histocompatibility Antigens, 78-6883
 - Immunity, Cellular, 78-6883, 78-7010
- Virus, Radiation Leukemia
 - Antigenic Determinants, 78-7029
 - DNA Nucleotidyltransferases, 78-7029
 - Isoantigens, 78-7029
 - T-Lymphocytes, 78-7029
 - Ultrastructural Study, 78-7029
- Virus, Rauscher Murine Leukemia
 - Antigen-Antibody Reactions, 78-6883
 - Histocompatibility Antigens, 78-6883
 - Immunity, Cellular, 78-6883
- Virus, SV40
 - Genetics, 78-6934
 - Twins, 78-6934

Lymphoma, Histiocytic

see Sarcoma, Reticulum Cell

Lymphosarcoma

- Acetophenone, 2-Amino-
 - Species Difference, 78-6790

Lymphosarcoma (cont'd)

- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 - Carcinogenic Activity, Mouse, Rat, 78-6660
- Cells, Cultured
 - Radioresistance, 78-6828
- Chromosomes
 - Cells, Cultured, 78-6828
- Growth
 - Cell Cycle Kinetics, 78-6980
 - Peritoneal Cavity, Mouse, 78-6980
- Indole-3-acrylic Acid
 - Species Difference, 78-6790
- Lymphocytes
 - Neoplasm Regression, Spontaneous, 78-6980
 - Peritoneal Cavity, Mouse, 78-6980
- Macrophages
 - Neoplasm Regression, Spontaneous, 78-6980
- Transplantation Immunology
 - Cells, Cultured, 78-6828
- Virus, Feline Leukemia
 - Antigen-Antibody Reactions, 78-6896
 - Antigenic Determinants, 78-6896

Lysine

- Myeloma Proteins
 - 2,4-Dinitrophenyl Haptens, Binding, 78-7039

Macrophages

- Carcinoma
 - BCG*, 78-6996
- Carrageen
 - Cytotoxic Activity, 78-6997
- Complement
 - Cleavage Product, Isolation and Characterization 78-7002
 - Cytotoxic Activity, 78-7002
- Corynebacterium parvum*
 - Phagocytosis, 78-7057
- Diterpene Esters
 - Prostaglandins, 78-6719
- Fibrosarcoma
 - Corynebacterium parvum*, 78-7054
 - Histocompatibility Antigens, 78-7051
 - Immunity, Cellular, 78-7051
 - Transplantation Immunology, 78-7051
- Hepatoma
 - BCG*, 78-6995
- Interferon
 - Prostaglandins E, 78-6993
- Leukemia L1210
 - Growth, 78-7000
- Leukemia, Lymphoblastic
 - Antitumor Activity, 78-6993
 - Cortisol, 78-6993
 - Dibutyl Cyclic AMP, 78-6993
 - Interferon, 78-6993
 - Prostaglandins E, 78-6993
- Lipopolysaccharides
 - Cytotoxic Activity, 78-7002
- Lung Neoplasms
 - Bradykinin, 78-7001
 - Carbon Clearance, 78-6999
 - Chemotaxis, 78-6999
 - Colony Formation, 78-6999
 - Immune Serums, 78-6999
 - Tumor Cell Supernatant, 78-7001
- Lymphoma
 - Cytotoxic Activity, 78-7002

Macrophages (cont'd)

- Lymphosarcoma
 - Neoplasm Regression, Spontaneous, 78-6980
- Mammary Neoplasms, Experimental
 - Immune Serums, 78-7004
 - Transplantation Immunology, 78-7057
- Neoplasms
 - Antibody-Dependent Cytotoxicity, Review, 78-6690
- Neoplasms, Experimental
 - Cellulose Tetranitrate, 78-6991
 - Chemotaxis, 78-6991
- Plasmacytoma
 - Cytotoxic Activity, 78-7002
- Prostaglandins E
 - Antitumor Activity, 78-6993
- Sarcoma
 - Activation, Ascites Cells, 78-6994
 - BCG, 78-6995
 - Cholanthrene, 3-Methyl-, 78-6994
 - Immune Response, 78-6994
 - Neoplasm Metastasis, 78-6994
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Antitumor Activity, 78-6782
 - Prostaglandins, 78-6719
 - Prostaglandins E, 78-6782

Malate Dehydrogenase

- Aniline, *N,N*-Diethyl-*p*-phenylazo-
 - Mitochondria, Liver, 78-6764
- Aniline, *N,N*-Dimethyl-*p*-phenylazo-
 - Mitochondria, Liver, 78-6764

Mammary Neoplasms, Experimental

- T-Lymphocytes
 - Antilymphocyte Serum, 78-7067
 - Immunosuppression, Rat, 78-7067
- Radiation, Ionizing
 - Immunosuppression, Rat, 78-7067
- Acetophenone, 2-Amino-
 - Species Difference, 78-6790
- Adenocarcinoma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6775
 - 78-6776, 78-6780
 - Immunity, Cellular, 78-7004
 - Neoplasm Regression, 78-6981
 - Ultrastructural Study, Rat, 78-6780
- Adenofibroma
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6780
- Antigen-Antibody Complex
 - Antibody Specificity, 78-7058
 - Kidney, 78-7058
- Antigens, Neoplasm
 - Immune Serums, 78-6978
 - Virus, Murine Leukemia, 78-6978
- Benz(a)anthracene, 7,12-Dimethyl-
 - FSH, 78-6778
 - Gonadotropins, 78-6778
 - Histological Study, Rat, 78-6775
 - Hormone Dependency, 78-6774
 - LH, 78-6778
 - Melatonin, 78-6779
 - Neoplasm Transplantation, 78-6774
 - Placental Extracts, 78-6775
 - Precancerous Conditions, 78-6774, 78-6776, 78-6780
 - Prolactin, 78-6778, 78-6779
 - Thyrotropin, 78-6778
 - Ultrastructural Study, Rat, 78-6780
- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide

Mammary Neoplasms, Experimental (cont'd)

- Carcinogenic Activity, Mouse, Rat, 78-6660
- Carcinogen, Chemical
 - Hormones, 78-6791
- Cell Differentiation
 - Precancerous Conditions, 78-6776
- Cells, Cultured
 - Neoplasm Metastasis, 78-7109
- Corynebacterium parvum*
 - Antitumor Activity, 78-7057
 - Transplantation Immunology, 78-7057
- Estradiol
 - Dose-Response Study, Mouse, 78-6794
- Estrone
 - Dose-Response Study, Mouse, 78-6794
 - Virus, Murine Mammary Tumor, 78-6868
- Hormones
 - Cell Division, 78-6791
 - Co-carcinogenic Effect, Review, 78-6643
 - DNA, Viral, 78-6868
 - RNA, Viral, 78-6868
- Immune Serums
 - Growth, 78-7004
 - Immunity, Cellular, 78-7004
 - Immunosuppressive Factor, 78-7004
 - Macrophages, 78-7004
 - Monocytes, Spleen, 78-7004
- Indole-3-acrylic Acid
 - Species Difference, 78-6790
- Iodine
 - Thyroid Dysfunction, Review, 78-6640
- Macrophages
 - Transplantation Immunology, 78-7057
- Neoplasm Regression
 - Tumor Ascitic Fluid, 78-6981
- Oncogenic Viruses
 - Hormones, 78-6791
- Pituitary Hormones
 - Estrus, 78-6778
- Plant Agglutinins
 - DNA Replication, 78-7004
 - Immunity, Cellular, 78-7004
 - Leukocyte Culture Test, Mixed, 78-7004
- Progesterone
 - Virus, Murine Mammary Tumor, 78-6868
- Prolactin
 - Hypophyseal Isograft, 78-6868
 - Thyroid Dysfunction, Review, 78-6640
 - Virus, Murine Mammary Tumor, 78-6868
- Radiation, Ionizing
 - Corynebacterium parvum*, 78-7057
 - Hormones, 78-6791
- RNA, Viral
 - Precancerous Conditions, 78-6862
- Selenium
 - Antitumor Activity, Review, 78-6696
- Silica
 - Corynebacterium parvum*, 78-7057
- Thyrotropin
 - Thyroid Dysfunction, Review, 78-6640
- Virus, Murine Mammary Tumor
 - Antigen-Antibody Complex, 78-7058
 - DNA-RNA Hybridization, 78-6864
 - DNA, Viral, 78-6868
 - RNA, Viral, 78-6862, 78-6868
 - Virus-Like Particles, 78-7104

stitis
 Breast Neoplasms
 Precancerous Conditions, 78-6700

elanoma
 Antigenic Determinants
 Antigen-Antibody Reactions, 78-7037
 Antigens, Neoplasm
 Hemadsorption Microassay, 78-7037

BCG
 Antigen-Antibody Reactions, 78-7037
 Antigenic Determinants, 78-7037
 Immune Response, Mouse, 78-6990

Cells, Cultured
 Cell Division, Review, 78-6680

Genetics
 Alpha Fetoproteins, 78-7066
 Carcinoembryonic Antigen, 78-7066

Growth
 Inflammation, 78-6990

Hypersensitivity
 Precancerous Conditions, Review, 78-6693

Immune Serums
 Antigen-Antibody Reactions, 78-7037

Laryngeal Neoplasms
 Antigenic Determinants, 78-7037

Skin Neoplasms
 Lymphocyte Transformation, Review, 78-6693
 B-Lymphocytes, 78-6693
 T-Lymphocytes, 78-6693
 Precancerous Conditions, Review, 78-6693

Ultraviolet Rays
 Age Factors, 78-6804
 Epidemiology, 78-6803, 78-6804
 Precancerous Conditions, Review, 78-6693
 Sunspot Activity, 78-6803

Virus, Adeno 5
 Reverse Transcriptase, 78-6967
 Virus Activation, 78-6967
 Virus Replication, 78-6967

Virus, Marek's Disease Herpes
 Antibodies, Viral, 78-6861
 Antigens, Viral, 78-6861

Melatonin
 Breast Neoplasms
 Epidemiology, 78-6779
 Prolactin, 78-6779
 Mammary Neoplasms, Experimental
 Benz(a)anthracene, 7,12-Dimethyl-, 78-6779

Meningioma
 Brain Neoplasms
 Epidemiology, 78-7170
 Eye Neoplasms
 Genetics, 78-7096
 Genetics
 Case Report, 78-7096
 Neurilemmoma
 Case Report, 78-7096
 Neurofibromatosis
 Genetics, 78-7096
 Nitrous Acid, Sodium Salt
 Food Contamination, 78-7170
 Radiation, Ionizing
 Epidemiology, 78-7170
 Spinal Neoplasms
 Genetics, 78-7096
 Tooth Diseases

Meningioma (cont'd)
 Epidemiology, 78-7149
 Wounds and Injuries
 Epidemiology, 78-7170

Menopause
 Breast Neoplasms
 Estrogens, 78-6649

Mercury, Chloromethyl-
 Hyperplasia
 Spleen, Mouse, Rat, 78-6817

Mesothelioma
 Asbestos
 Case Report, 78-6837
 Dust Application, Surgery, 78-6837
 Pericardium, 78-6837
 Pericardium
 Asbestos, 78-7113
 Case Report, 78-7113
 Hemosiderin, 78-7113
 Peritoneal Neoplasms
 Sterigmatocystin, 78-6800
 Sterigmatocystin
 Histological Study, Rat, 78-6800

Mestranol
 Liver Neoplasms
 Hyperplasia, 78-7125

Metaplasia
 Radiation, Ionizing
 Gastric Mucosa, Rat, 78-6814
 Stomach Neoplasms
 Gastric Mucosa, 78-7117
 Precancerous Conditions, 78-7117

Methane, Chlorodifluoro-
 Abnormalities
 Epidemiology, Child, 78-6765
 Leukemia, Lymphoblastic
 Case Report, 78-6765
 In Utero Exposure, 78-6765

Methane, Sulfinylbis-
 Erythroleukemia
 Globin, 78-7185

Methanesulfonic Acid, Ethyl Ester
 Esterases
 Plasma Levels, Mouse, 78-6743
Haemophilus influenzae
 DNA Repair, 78-6759
 Virus, SV40
 Virus Replication, Review, 78-6620

Methanesulfonic Acid, Methyl Ester
 Virus, SV40
 Virus Replication, Review, 78-6620

Methanol, (Methyl-*ONN*-azoxy)-
 Kidney Neoplasms
 Pyrazole, 78-6760
 Oxidoreductases
 Enzyme Activation, 78-6760
 Skin Neoplasms
 Papilloma, 78-6760
 Pyrazole, 78-6760

Methionine
 Acetohydroxamic Acid, *N*-Fluoren-2-yl-
 Nucleic Acids, Binding, 78-6730

- Methionine (cont'd)**
Liver Neoplasms
Carcinogen, Chemical, 78-6638
- Methionine, S-Adenosyl-**
Ultraviolet Rays
RNA, Transfer, Methyltransferases, 78-6621
Uracil
RNA, Transfer, Methyltransferases, 78-6621
- Methylamine**
Sarcoma
Transplantation Immunology, 78-7048
- Micrococcus lysodeikticus***
Leukemia L1210
Immune Response, 78-7020
- Microsomes, Liver**
Benz(a)anthracene
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 1,2-Dihydro-1,2-dihydroxy-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 8,9-Dihydro-8,9-dihydroxy-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 10,11-Dihydro-10,11-dihydroxy-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 1,2-Dihydro-1,2-dihydroxy-7,12-
dimethyl-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 3,4-Dihydro-3,4-dihydroxy-7,12-
dimethyl-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 5,6-Dihydro-5,6-dihydroxy-7,12-
dimethyl-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 8,9-Dihydro-8,9-dihydroxy-7,12-
dimethyl-
Carcinogenic Metabolite, 78-6772
Benz(a)anthracene, 7,12-Dimethyl-
Carcinogenic Metabolite, 78-6772
Benzene, Bromo-
Metabolism, Review, 78-6607
Carbon Tetrachloride
Metabolism, Review, 78-6607
Hydrocarbons, Halogenated
Metabolism, Review, 78-6607
Nitrosamines
Ames Test, Review, 78-6635
Polycyclic Hydrocarbons
Ames Test, Review, 78-6635
Quinoline
Ames Test, 78-6762
- Microtubules**
Virus, Rous Sarcoma
Cell Transformation, Neoplastic, 78-6846
- Migration Inhibitory Factor**
Hodgkin's Disease
Virus, Epstein-Barr, 78-6989
Infectious Mononucleosis
Virus, Epstein-Barr, 78-6989
Lymphoma
Virus, Epstein-Barr, 78-6989
Virus, Epstein-Barr
Immunity, Cellular, 78-6989
Leukocytes, 78-6989
- Migration Inhibitory Factor (cont'd)**
Ultraviolet Rays, 78-6989
- Mitochondria**
Virus, SV40
Ethidium Bromide, 78-6915
- Mitochondria, Liver**
Aniline, *N,N*-Diethyl-*p*-phenylazo-
Cytochrome Oxidase, 78-6764
Malate Dehydrogenase, 78-6764
Aniline, *N,N*-Dimethyl-*p*-phenylazo-
Cytochrome Oxidase, 78-6764
Malate Dehydrogenase, 78-6764
Butyric Acid, 2-Amino-4-(ethylthio)-
Adenosine Triphosphate, 78-6737
Oxidative Phosphorylation, 78-6737
- Mitogens**
Cholanthrene, 3-Methyl-
Aryl Hydrocarbon Hydroxylases, 78-6781
- Mitomycin C**
Haemophilus influenzae
DNA Repair, 78-6759
Leukemia
Immunosuppression, 78-7022
Lymphocytes
Cytotoxicity, 78-6988
Virus, SV40
DNA, Viral, 78-6933
Virus-Like Particles, 78-6933
- Mitosis**
Sarcoma
Erythrocytes, 78-7078
Skin Neoplasms
Connective Tissue, 78-7087
- Molybdenum**
Ames Test
Carcinogenic Activity, Review, 78-6635
- Monocytes**
Fibrosarcoma
Immunity, Passive, 78-7054
T-Lymphocytes
Immunity, Passive, 78-7054
Neoplasms, Experimental
Chemotaxis, 78-6991
- Morpholine, *N*-Nitroso-**
Liver Neoplasms
Glucosephosphate Dehydrogenase, 78-6787
- Mouth Neoplasms**
Areca
Carcinogenic Potential, Ingredients, 78-6727
Precancerous Conditions, 78-6727
Vitamin B, 78-6727
Benz(a)anthracene, 7,12-Dimethyl-
Benzene, 1-Chloro-2,4-dinitro-, 78-7033
Precancerous Conditions, 78-7075
Benzene, 1-Chloro-2,4-dinitro-
Immunity, Cellular, 78-7033
Immunotherapy, 78-7033
Carcinoma, Epidermoid
Benz(a)anthracene, 7,12-Dimethyl-, 78-7033
Ethyl Alcohol
Epidemiology, Review, 78-6699
Papilloma
Benz(a)anthracene, 7,12-Dimethyl-, 78-7033

Mouth Neoplasms (cont'd)

Precancerous Conditions

Basal Lamina, 78-7075

Ultrastructural Study, Hamster, 78-7075

Snuff

Epidemiology, 78-6720

Mucopolysaccharides

Fibroblasts

Cell Transformation, Neoplastic, 78-7198

Multiple Myeloma

Neoplasms, Multiple Primary

Drug Therapy, 78-6736

Radiation, Ionizing

Epidemiology, Review, 78-6667

Occupational Hazard, 78-6667

Virus, Bovine Leukemia

Antibody Specificity, 78-6897

Virus, C-Type RNA Tumor

Isolation and Characterization, 78-6897

Reverse Transcriptase, 78-6897

Virus-Like Particles, 78-6897

Muscular Atrophy

Carcinoma, Basal Cell

Case Report, 78-7094

Genetics, 78-7094

Mustard Gas

see Sulfide, Bis(2-chloroethyl)-

Mutagens

Bacteria

Colony Growth and Morphology, 78-7196

Cells, Cultured

Colony Growth and Morphology, 78-7196

Chromosome Aberrations

Carcinogenic Potential, Review, 78-6618

DNA, Bacterial

Colony Growth and Morphology, 78-7196

DNA Repair

Escherichia coli, 78-7193

Quantitation Method, 78-7193

Environmental Hazard

Epidemiology, Review, 78-6605

Hereditary Diseases

Epidemiology, Review, 78-6605

Salmonella typhimurium

Azaguanine Resistance, 78-6761

Histidine Reversion, 78-6761

Mutation

Betabinomial Distribution

Mouse, 78-7195

DNA Polymerase

Hereditary Diseases, Review, 78-6704

Myasthenia Gravis

Breast Neoplasms

Thymectomy, 78-7035

Neoplasms

Thymectomy, 78-7035

Thymectomy

Immune Response, 78-7035

Thymoma

Thymectomy, 78-7035

Mycosis Fungoides

Chromosome Abnormalities

Chromosomes, Human, 1-3, 78-7120

Mycosis Fungoides (cont'd)

B-Lymphocytes

Chromosome Abnormalities, 78-7120

Plant Agglutinins, 78-7120

T-Lymphocytes

Chromosome Abnormalities, 78-7120

Plant Agglutinins, 78-7120

Skin Neoplasms

Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753

Drug Therapy, 78-6753

Mycotoxins

Aflatoxin B1

Carcinogenic Activity, Review, 78-6629

Food Contamination

Carcinogenic Potential, Review, 78-6662

Gastrointestinal Neoplasms

Diet, Review, 78-6638

Retinol, 78-6638

Hepatoma

Epidemiology, Kenya, Swaziland, 78-7184

Food Contamination, 78-7184

Ultrastructural Study, Rat, 78-6723

2,5-Hexadienoic Acid, 3-Methoxy-5-methyl-4-oxo-

Carcinogenic Activity, Review, 78-6629

Liver Neoplasms

Adenocarcinoma, 78-6723

Ultrastructural Study, Rat, 78-6723

Luteoskyrin

Carcinogenic Activity, Review, 78-6629

Patulin

Carcinogenic Activity, Review, 78-6629

Sterigmatocystin

Carcinogenic Activity, Review, 78-6629

Structure-Activity Relationship

Carcinogenic Activity, Review, 78-6629

Myelofibrosis

Reverse Transcriptase

Enzymatic Activity, Review, 78-6669

Myeloma Proteins

Antigens

Binding Sites, 78-7040

Haptens

Spectrum Analysis, Raman, 78-7039

IgA

Binding Sites, 78-7040

2,4-Dinitrophenyl Haptens, Binding, 78-7039

Lysine

2,4-Dinitrophenyl Haptens, Binding, 78-7039

Myoma

Gynecologic Neoplasms

Contraceptives, Oral, 78-6645

Myosarcoma

Benzo(a)pyrene

Graft vs Host Reaction, 78-7045

Splenectomy, 78-7046

B-Lymphocytes

Lymphocyte Depletion, 78-7046

Splenectomy, 78-7046

T-Lymphocytes

Splenectomy, 78-7046

Splenectomy

Immune Response, 78-7046

2-Naphthylamine

Aromatic Amines

2-Naphthylamine (cont'd)

Carcinogenic Potential, Review, 78-6702

Epidemiology, Review, 78-6702

Occupational Hazard

Epidemiology, Review, 78-6702

1-Naphthylamine, *N*-Phenyl-

Virus, Lymphocytic Choriomeningitis

Cell Membrane, 78-6855

Virus, Rous Sarcoma

Cell Membrane, 78-6855

Nasopharyngeal Neoplasms

Carcinoma

Cellular Inclusions, 78-7101

Virus, Epstein-Barr, 78-7034

Cellular Inclusions

Ultrastructural Study, 78-7101

Ethnic Groups

Epidemiology, Malaysia, 78-7152

Lymphoma

Virus, Epstein-Barr, 78-7034

Virus, Epstein-Barr

Antibodies, Viral, 78-7034

Cellular Inclusions, 78-7101

DNA, Viral, Review, 78-6632

Histological Study, Review, 78-6632

Seroepidemiology, Review, 78-6632

Neoplasm Metastasis

Breast Neoplasms

Age Factors, 78-7159

Carcinoma, Ductal, 78-7161

Cell Membrane

Phenotype, Review, 78-6687

Cells, Cultured

Bromo Cyclic AMP, 78-7109

Choriocarcinoma

Case Report, 78-7134

Lung, 78-7134

Fibrosarcoma

Cells, Cultured, 78-7109

Immunity, Passive, 78-7047

B-Lymphocytes, 78-7047

T-Lymphocytes, 78-7047

Hepatoma

Australia Antigen, 78-6958

Benzidine, Dihydrochloride, 78-7122

Benzo(a)pyrene, 78-7122

Diethylamine, *N*-Nitroso-, 78-7122

Histological Study, 78-7122

Lung, Mouse, 78-7122

Urea, Ethyl Nitroso-, 78-7122

Kidney Neoplasms

Calcaneus, 78-7129

Carcinoma, Basal Cell, 78-7129

Liver Neoplasms

Benzidine, Dihydrochloride, 78-7122

Histological Study, 78-7122

Lung, Mouse, 78-7122

Urea, Ethyl Nitroso-, 78-7122

Lung Neoplasms

Carcinoma, 78-6998

Cells, Cultured, 78-7109

Cortisone Acetate, 78-6998

Corynebacterium parvum, 78-6998

Immunity, Passive, 78-7047

B-Lymphocytes, 78-7047

T-Lymphocytes, 78-7047

Neoplasm Metastasis (cont'd)

Phagocytosis, 78-6998

Silica, 78-6998

Lymphoma

B-Lymphocytes, 78-6875

T-Lymphocytes, 78-6875

Virus, Abelson Murine Leukemia, 78-6875

Mammary Neoplasms, Experimental

Cells, Cultured, 78-7109

Sarcoma

Animal Model, Mouse, 78-7043

Cholanthrene, 3-Methyl-, 78-7043

Graft vs Host Reaction, 78-7045

Histocompatibility Antigens, 78-7043

Macrophages, 78-6994

Stomach Neoplasms

Transplantation, Heterologous, 78-7115

Thyroid Neoplasms

Adenocarcinoma, 78-7102

Tongue Neoplasms

Carcinoma, Epidermoid, 78-7098

Neoplasm Recurrence, Local

Bone Neoplasms

Case Report, 78-7147

Osteoma, Osteoid, 78-7147

Sarcoma, Osteogenic, 78-7147

Carcinoma, Epidermoid

Sulfide, Bis(2-chloroethyl)-, 78-7088

Pharyngeal Neoplasms

Carcinoma, 78-7100

Case Report, 78-7100

Neoplasm Regression, Spontaneous

Lymphosarcoma

Lymphocytes, 78-6980

Macrophages, 78-6980

Neoplasm Transplantation

Bladder Neoplasms

Schistosoma haematobium, 78-7140

Fibrosarcoma

Immunosuppression, 78-7047

Leukemoid Reaction, 78-7052

Kidney Neoplasms

Dimethylamine, *N*-Nitroso-, 78-6739

Leukemia, Myelocytic

Sex Chromosomes, 78-7069

Lung Neoplasms

Immunosuppression, 78-7047

Lymphoma

Immunosuppression, 78-7028

Mammary Neoplasms, Experimental

Benz(a)anthracene, 7,12-Dimethyl-, 78-6774

Neuroblastoma

Contact Inhibition, 78-7079

Neoplasm Invasiveness, 78-7079

Sarcoma

Hemodynamics, 78-7078

Histocompatibility Antigens, 78-7043

Sarcoma 180, Crocker

Contact Inhibition, 78-7079

Neoplasm Invasiveness, 78-7079

Neoplasms (General and Unspecified)

Anaphylaxis

Immune Response, Solid Tumors, 78-6691

Antigen-Antibody Complex

Antibody Specificity, 78-7058

Neoplasms (General and Unspecified) (cont'd)

- Kidney, 78-7058
- Antigens
 - Immune Response, Review, 78-6687
- Cell Membrane
 - Surface Properties, Review, 78-6687
- Cerebral Ischemia, Transient
 - Stress, Review, 78-6682
- Epidemiology
 - Pediatric Patients, 78-7177
- Ethnic Groups
 - Epidemiology, Review, 78-6697
- Genetics
 - Epidemiology, Review, 78-6697
- Growth
 - Phenotype, Review, 78-6687
- Immunosuppressive Agents
 - Epidemiology, 78-6736
- Leukocytes
 - Immunity, Cellular, Review, 78-6690
- Lymphocytes
 - Interferon, 78-6983
 - Killer Cells, 78-6983
- T-Lymphocytes
 - Immunity, Cellular, Review, 78-6690
- Myasthenia Gravis
 - Thymectomy, 78-7035
- Oncogenic Viruses
 - Stress, Review, 78-6682
- Pesticides
 - Epidemiology, 78-7174
 - Occupational Hazard, 78-7174
- Socioeconomic Factors
 - Epidemiology, Review, 78-6698
- Virus, Polyoma, BK
 - Antigens, Viral, 78-6905
 - DNA-DNA Hybridization, 78-6905

Neoplasms, Experimental

- Antigenic Determinants
 - Transplantation Immunology, 78-7031
- Asbestos
 - Immune Response, Review, 78-6633
 - Precancerous Conditions, Review, 78-6633
- Benz(a)anthracene, 7,12-Dimethyl-
 - Chemotaxis, 78-6991
- Carcinogen, Chemical
 - Trans-Species Extrapolation, Review, 78-6604
- Cellulose Tetranitrate
 - Macrophages, 78-6991
- Chemotaxis
 - Macrophages, 78-6991
- Cholanthrene, 3-Methyl-
 - Histocompatibility Antigens, 78-7044
- Foreign Bodies
 - Immune Response, Review, 78-6633
 - Precancerous Conditions, Review, 78-6633
- Histocompatibility Antigens
 - Transplantation Immunology, 78-7044
- T-Lymphocytes
 - Immunosurveillance, Review, 78-6688
- Monocytes
 - Chemotaxis, 78-6991
- Neoplasm Invasiveness
 - Mesenchyme, Chick Wing, 78-7079
- Schistosoma*
 - Immune Response, Review, 78-6633
 - Precancerous Conditions, Review, 78-6633

Neoplasms, Experimental (cont'd)

- Transplantation, Heterologous
 - Neoplasm Invasiveness, 78-7079
- Virus-Like Particles
 - Transplantation Immunology, 78-7031
- Virus, SV40
 - Antilymphocyte Serum, 78-6911
 - Castration, 78-6911
 - Cortisone Acetate, 78-6911
 - Cyclophosphamide, 78-6911
 - Estrone, 78-6911
 - Testosterone, Propionate, 78-6911
 - Transplantation Immunology, 78-6911

Neoplasms, Multiple Primary

- Breast Neoplasms
 - Epidemiology, 78-7158
- Ear Neoplasms
 - Neurilemmoma, 78-7096
- Hodgkin's Disease
 - Drug Therapy, 78-6736
- Leukemia
 - Actinomycin D, 78-7072
 - Butyric Acid, 4-(*p*-Bis(2-chloroethyl)aminophenyl)-
78-7072
 - Cyclophosphamide, 78-7072
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-7072
 - Radiation, Ionizing, 78-7072
- Leukemia, Myeloblastic
 - Drug Therapy, 78-6736
 - Sarcoma, Ewing's, 78-6815
 - Virus, Epstein-Barr, 78-7071
- Liver Neoplasms
 - Hemangioma, Cavernous, 78-7098
- Multiple Myeloma
 - Drug Therapy, 78-6736
- Nose Neoplasms
 - Hemangioma, 78-7071
 - Virus, Epstein-Barr, 78-7071
- Ovarian Neoplasms
 - Drug Therapy, 78-6736
- Skin Neoplasms
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
 - Xeroderma Pigmentosum, 78-7121
- Spinal Cord Neoplasms
 - Astrocytoma, 78-7098

Neoplasms, Radiation-Induced

- Nuclear Reactors
 - Epidemiology, 78-6829, 78-7182
- Occupational Hazard
 - Dose-Response Study, Hanford Atomic Plant
78-7182
 - Epidemiology, 78-6829

Neoplastic Endocrine-Like Syndromes

- Pheochromocytoma
 - Genetics, 78-7144

Nephroblastoma

- Cell Differentiation
 - Neoplasm Regression, Review, 78-6686

Nervous System Neoplasms

- Neuroblastoma
 - Animal Model, Opossum, 78-6756
 - Urea, Ethyl Nitroso-, 78-6756
- Urea, Ethyl Nitroso-
 - Carcinogenic Activity, Review, 78-6656
 - Chronic Nerve Irritation, 78-6748

Nervous System Neoplasms (cont'd)

- Glycerolphosphate Dehydrogenase, 78-6757
- Transplacental Carcinogenesis, 78-6748, 78-6757

Urea, Methyl Nitroso-

- Transplacental Carcinogenesis, 78-6757

Neurilemmoma

- Ear Neoplasms
 - Genetics, 78-7096
 - Neoplasms, Multiple Primary, 78-7096
- Meningioma
 - Case Report, 78-7096

Neuroblastoma

- Neoplasm Invasiveness
 - Mesenchyme, Chick Wing, 78-7079
- Neoplasm Transplantation
 - Contact Inhibition, 78-7079
 - Neoplasm Invasiveness, 78-7079
- Nervous System Neoplasms
 - Animal Model, Opossum, 78-6756
 - Urea, Ethyl Nitroso-, 78-6756
- Virus-Like Particles
 - Ultrastructural Study, 78-6977

Neurofibromatosis

- Genetics
 - Child, 78-7097
 - Morbidity, 78-7097
- Meningioma
 - Genetics, 78-7096

Neutrophils

- Laryngeal Neoplasms
 - Alkaline Phosphatase, 78-7099
 - Enzymatic Activity, 78-7099
 - Glucuronidase, 78-7099
 - Leukoplakia, 78-7099
 - Lipids, 78-7099
 - Papilloma, 78-7099
 - Precancerous Conditions, 78-7099

Nevus

- Carcinoma, Basal Cell
 - Genetics, 78-7094

Nickel

- Lung Neoplasms
 - Epidemiology, 78-7156
 - Occupational Hazard, 78-7156
- Respiratory Tract Neoplasms
 - Epidemiology, Review, 78-6634

Nickel Chloride

- Lung Clearance
 - Kinetics, Rat, 78-6724

Nitric Acid

- Food Additives
 - Carcinogenic Potential, Review, 78-6662
- Stomach Neoplasms
 - Food Contamination, 78-6733

Nitric Acid, Lead Salt

- Hyperplasia
 - Spleen, Mouse, Rat, 78-6817

Nitrocellulose

- see Cellulose Tetranitrate

Nitrogen Mustard

- see Diethylamine, 2,2'-Dichloro-*N*-methyl-

Nitrogen Oxide

- Air Pollution
 - Spectrum Analysis, 78-6710

Nitrosamines

- Air Pollution
 - Carcinogenic Activity, Review, 78-6628
 - Concentration Levels, 78-7183
- Microsomes, Liver
 - Ames Test, Review, 78-6635
- Water, Deionized
 - Risk Factors, 78-7183
- Water Pollution
 - Concentration Levels, 78-7183
 - Emission Sources, 78-7183

Nitroso Compounds

- Purine, 2-Amino-6-methoxy-DNA, Liver, 78-6625
- Enzymes, 78-6625

Nitrosocarbaryl

- see Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester

Nitrous Acid

- Food Contamination
 - Carcinogenic Potential, Review, 78-6655, 78-6662
- Lymphoma
 - Carcinogenic Potential, Review, 78-6655
- Stomach Neoplasms
 - Food Contamination, 78-6733

Nitrous Acid, Sodium Salt

- Meningioma
 - Food Contamination, 78-7170

Norepinephrine

- Acetamide, *N*-Fluoren-2-yl-Receptors, Adrenergic, 78-6732

Norethisterone

- Liver Neoplasms
 - Contraceptives, Oral, 78-7126

Norocetopine

- Agrobacterium tumefaciens*
 - Metabolism, 78-7194

Nose Neoplasms

- Hemangioma
 - Neoplasms, Multiple Primary, 78-7071
- Neoplasms, Multiple Primary
 - Virus, Epstein-Barr, 78-7071
- Virus, Epstein-Barr
 - Case Report, 78-7071

Nuclear Reactors

- Neoplasms, Radiation-Induced
 - Epidemiology, 78-7182

Nucleic Acids

- Alkylating Agents
 - Teratogenic/Mutagenic Activity, Review, 78-6615
- Antimetabolites
 - Teratogenic/Mutagenic Activity, Review, 78-6615
- Carcinogen, Chemical
 - Mutagenic Activity, Review, 78-6611
- Virus, RD-114
 - Isolation and Characterization, Review, 78-6674

Nucleoproteins

- Hepatoma

nucleoproteins (cont'd)

- RNA, Messenger, 78-7189
- RNA, Messenger
 - Liver Regeneration, 78-7189
- Virus, SV40
 - Chromosomes, 78-6918
 - DNA Replication, 78-6913
 - Isolation and Characterization, 78-6918
 - Virus Replication, 78-6918

nucleotidases

- Virus, Moloney Murine Leukemia
 - Cell Membrane, 78-7018

Occupational Hazard

- Acetamide, *N*-(4-((2-Hydroxy-5-methylphenyl)azo)phenyl)-
 - Textile Workers, Review, 78-6663
- Air Pollution
 - Quantitation Method, 78-6709
- Angiosarcoma
 - Ethylene, Chloro-, 78-6713
- Anthraquinone
 - Printers, Review, 78-6663
- Asbestos
 - Immunity, Cellular, 78-7061
 - Immunoglobulins, 78-7061
- Asbestosis
 - Autoantibodies, 78-7061
 - B-Lymphocytes, 78-7061
 - T-Lymphocytes, 78-7061
- Benzene
 - Epidemiology, Review, 78-6627
 - Maximum Permissible Exposure Level, Review 78-6627
- Benzophenone
 - Printers, Review, 78-6663
- C.I. Basic Green 4
 - Textile Workers, Review, 78-6663
- C.I. Vat Green 1
 - Textile Workers, Review, 78-6663
- Carcinogen, Chemical
 - Epidemiology, Review, 78-6697
- Carcinoma, Basal Cell
 - Uranium, 78-7179
- Coumarin, 3-Benzyl-7-hydroxy-4-methyl-
 - Ultraviolet Rays, 78-6663
- Direct Black 38
 - Textile Workers, Review, 78-6663
- Direct Blue 6
 - Textile Workers, Review, 78-6663
- Ethylene, Chloro-
 - Chromatids, 78-6712
 - Chromosome Abnormalities, 78-6712, 78-6715
- Ethylene, Chloro- Polymer
 - Chromosome Abnormalities, 78-6715
- Ethylene, Trichloro-
 - Epidemiology, 78-7173
- Fluorescent Dyes
 - Ultraviolet Rays, 78-6663
- Hair Dyes
 - Epidemiology, 78-7166
- Leukemia
 - Ethylene, Trichloro-, 78-7173
 - Radiation, Ionizing, 78-6667
- Liver Neoplasms
 - Ethylene, Trichloro-, 78-7173
- Lung Neoplasms

Occupational Hazard (cont'd)

- Epidemiology, 78-7155
 - Nickel, 78-7156
 - Pesticides, 78-7174
 - Multiple Myeloma
 - Radiation, Ionizing, 78-6667
 - 2-Naphthylamine
 - Epidemiology, Review, 78-6702
 - Neoplasms
 - Pesticides, 78-7174
 - Neoplasms, Radiation-Induced
 - Dose-Response Study, Hanford Atomic Plant 78-7182
 - Epidemiology, 78-6829
 - Pancreatic Neoplasms
 - Radiation, Ionizing, 78-6667
 - 1-Propanol, 2,3-Dibromo-, Phosphate
 - Textile Workers, Review, 78-6663
 - Radiation, Ionizing
 - Dose-Response Study, Review, 78-6667
 - Epidemiology, Review, 78-6667
 - Respiratory Tract Neoplasms
 - Carcinogen, Chemical, 78-6768
 - Epidemiology, East Germany, 78-6768
 - Epidemiology, Review, 78-6634
 - Precancerous Conditions, Review, 78-6634
 - Stomach Neoplasms
 - Asbestos, 78-7172
 - Epidemiology, 78-7172
 - Latency Period, 78-7172
 - Styrene
 - Printers, Review, 78-6663
 - Virus, Feline Leukemia
 - Antigen-Antibody Reactions, 78-7165
- Octopine**
- Agrobacterium tumefaciens*
 - Extrachromosomal Inheritance, 78-7194
 - Metabolism, 78-7194
 - Regulatory Gene Mutants, 78-7194
 - Tumorigenicity, 78-7194
- Odontogenic Cysts**
- Carcinoma, Basal Cell
 - Genetics, 78-7094
- Oligodendroglioma**
- Brain Neoplasms
 - Urea, Ethyl Nitroso-, 78-6749
- Oncogenic Viruses**
- Adenocarcinoma
 - Carcinoembryonic Antigen, 78-7118
 - Brain Neoplasms
 - Epidemiology, 78-7168
 - Species Difference, Review, 78-6639
 - Carcinoembryonic Antigen
 - Genetics, 78-7118
 - Cell Cycle Kinetics
 - Fibroblasts, Review, 78-6661
 - Cell Division
 - Epithelial Cells, Review, 78-6680
 - Cell Transformation, Neoplastic
 - Fibroblasts, Review, 78-6661
 - Chromosome Abnormalities
 - Carcinogenic Activity, Review, 78-6610
 - Genetics
 - Cell Transformation, Neoplastic, Review, 78-6705
 - Hormones

Oncogenic Viruses (cont'd)

Co-carcinogenic Effect, Review, 78-6643

Immunosuppression

Cell Transformation, Neoplastic, Review, 78-6688

Mammary Neoplasms, Experimental

Hormones, 78-6791

Mutagenic Activity

Colony Growth and Morphology, 78-7196

Neoplasms

Stress, Review, 78-6682

Transformation, Genetic

Stress, Review, 78-6682

Virus-Like Particles

Ultrastructural Study, Hamster Lung Cells, 78-6974

Osteoma, Osteoid**Bone Neoplasms**

Neoplasm Recurrence, Local, 78-7147

Osteopetrosis

Virus, Avian Myeloblastosis

Antigen-Antibody Reactions, 78-6844

Ovarian Neoplasms**Adenocarcinoma**

Cysts, 78-7132

Endometrial Hyperplasia, 78-7132

Histological Study, 78-7132

Hyperplasia, 78-7132

Endometriosis

Cysts, 78-7132

Endometrial Hyperplasia, 78-7132

Histological Study, 78-7132

Hyperplasia, 78-7132

Neoplasms, Multiple Primary

Drug Therapy, 78-6736

Virus, Herpes Simplex 1

Antibodies, Viral, 78-6939

Virus, Herpes Simplex 2

Antibodies, Viral, 78-6939

Virus, Marek's Disease Herpes

Antibodies, Viral, 78-6861

Oxidoreductases

Methanol, (Methyl-*ONN*-azoxy)-

Enzyme Activation, 78-6760

Oxygen**Iron**

Enzymatic Deactivation, 78-6654

Oxygenases

Polycyclic Hydrocarbons

Metabolism, Review, 78-6623

Ozone**Air Pollution**

Spectrum Analysis, 78-6710

Free Radicals

Smog Formation, 78-6710

Pancreatic Neoplasms

Acetophenone, 2-Amino-

Species Difference, 78-6790

Adenoma

Acetophenone, 2-Amino-, 78-6790

Indole-3-acrylic Acid, 78-6790

Carcinoma

Ribonuclease, 78-7200

Carcinoma, Ductal

Ribonuclease, 78-7200

Pancreatic Neoplasms (cont'd)

Indole-3-acrylic Acid

Species Difference, 78-6790

Islet Cell Tumor

Acetophenone, 2-Amino-, 78-6790

Indole-3-acrylic Acid, 78-6790

Ribonuclease, 78-7200

Radiation, Ionizing

Epidemiology, Review, 78-6667

Occupational Hazard, 78-6667

Ribonuclease

Serum Levels, 78-7200

Pancreatitis

Ribonuclease

Serum Levels, 78-7200

Papilloma**Breast Neoplasms**

Cytodiagnosis, 78-7105

Gynecologic Neoplasms

Acetophenone, 2-Amino-, 78-6790

Indole-3-acrylic Acid, 78-6790

Laryngeal Neoplasms

Enzymatic Activity, 78-7099

Neutrophils, 78-7099

Virus-Like Particles, 78-6900

Mouth Neoplasms

Benz(a)anthracene, 7,12-Dimethyl-, 78-7033

Skin Neoplasms

Methanol, (Methyl-*ONN*-azoxy)-, 78-6760

Parathyroid Glands**Adenoma**

Transplantation, Autologous, 78-7103

Parathyroid Neoplasms

Precancerous Conditions

Transplantation, Autologous, 78-7103

Parity

Breast Neoplasms

Epidemiology, 78-6700

Patulin

Mycotoxins

Carcinogenic Activity, Review, 78-6629

Peptide Hydrolases

Cell Membrane

Fluorescence, 78-6855

Cell Transformation, Neoplastic

Cell Membrane, 78-6855

Peptides

Virus, Adeno 2

Cycloheximide, 78-6963

Cytosine, 1- β -*D*-Arabinofuranosyl-, 78-6963

Cytosine, 1- β -*D*-Arabinofuranosyl-, 78-6963

RNA, Messenger, 78-6963

Virus, Herpes Simplex 1

Physical Mapping, 78-6938

Temperature Sensitive Mutants, 78-6943

Virus, Recombinant, 78-6938

Virus, Herpes Simplex 2

Physical Mapping, 78-6938

Virus, Recombinant, 78-6938

Pericardium**Mesothelioma**

Asbestos, 78-6837, 78-7113

Case Report, 78-7113

Pericardium (cont'd)

Hemosiderin, 78-7113

Peritoneal Neoplasms

Mesothelioma

Sterigmatocystin, 78-6800

Pesticides

Chromosome Aberrations

Carcinogenic Potential, Review, 78-6618

Food Contamination

Carcinogenic Potential, Review, 78-6662

Lung Neoplasms

Epidemiology, 78-7174

Occupational Hazard, 78-7174

Neoplasms

Epidemiology, 78-7174

Occupational Hazard, 78-7174

Peutz-Jeghers Syndrome

Digestive System Neoplasms

Uterine Neoplasms, 78-7119

Intestinal Neoplasms

Genetics, Review, 78-6708

Phagocytosis

Corynebacterium parvum

Macrophages, 78-7057

Radiation, Ionizing, 78-7057

Silica, 78-7057

Iron

Toxicity, 78-6654

Lung Neoplasms

Carcinoma, 78-6998

Cortisone Acetate, 78-6998

Corynebacterium parvum, 78-6998

Growth, 78-6998

Neoplasm Metastasis, 78-6998

Silica, 78-6998

4,4'-Stilbenediol, α, α' -Diethyl-, 78-6998

Trypan Blue, 78-6998

Pharyngeal Neoplasms

Carcinoma

Neoplasm Recurrence, Local, 78-7100

Neoplasm Recurrence, Local

Case Report, 78-7100

Radiation, Ionizing

Carcinogenic Potential, 78-6816

Pheochromocytoma

Epinephrine

Genetics, 78-7144

Urinary Levels, 78-7144

Genetics

Case Report, 78-7144

Neoplastic Endocrine-Like Syndromes

Genetics, 78-7144

Thyroid Neoplasms

Genetics, 78-7144

Phosphodiesterases

Cholangioma

Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6766

Hepatoma

Immune Serums, 78-6960

Liver Cirrhosis, 78-6960

Ultrastructural/Cytochemical Study, 78-6960

Virus, Hepatitis, 78-6960

Phosphonium, Tetrakis(hydroxymethyl)-, Chloride

Ether, Bis(chloromethyl)-

Chromatographic Study, 78-6725

Guanosine

Phosphine Product, 78-6725

Skin Neoplasms

Carcinogenic Activity, Mouse, 78-6725

Oxalate Salt, 78-6725

Phosphonoacetic Acid

Virus, Epstein-Barr

Antigens, Viral, 78-6946

Virus Replication, 78-6946

Virus, Herpes Saimiri

Antigens, Viral, 78-6946

Virus Replication, 78-6946

Phosphoproteins

Virus, Rous Sarcoma

Cell Transformation, Neoplastic, 78-6848

Genes, Viral, 78-6848

Protein Kinase, 78-6848

Temperature Sensitive Mutants, 78-6848

Phosphoric Acid, Diethyl-, *p*-Nitrophenyl Ester

Acetohydroxamic Acid, *N*-Fluoren-2-yl-

Mutagenic Activity, 78-6730

Phthalide, 4,5,6,7-Tetrabromo-3,3-bis(4-hydroxy-3-sulfophenyl)-

Contraceptives, Oral

Liver Function, Review, 78-6650

Piperazine, 2,5-Dimethyl-1,4-dinitroso-

Esterases

Plasma Levels, Mouse, 78-6743

Piperazine, 1,4-Dinitroso-

Esterases

Plasma Levels, Mouse, 78-6743

Piperazine, 1-Methyl-4-nitroso-

Esterases

Plasma Levels, Mouse, 78-6743

Piperidine, 2,6-Dimethyl-1-nitroso-

Esterases

Plasma Levels, Mouse, 78-6743

Pituitary Hormones

Mammary Neoplasms, Experimental

Estrus, 78-6778

Pituitary Neoplasms

Adenoma

Contraceptives, Oral, 78-7095

Prolactin, 78-7095

Contraceptives, Oral

Prolactin, 78-7095

Plant Agglutinins

Benz(a)anthracene

Aryl Hydrocarbon Hydroxylases, 78-6781

5,6-Benzoflavone

Aryl Hydrocarbon Hydroxylases, 78-6781

Bladder Neoplasms

Lymphocytes, 78-6735

Cholanthrene, 3-Methyl-

Aryl Hydrocarbon Hydroxylases, 78-6781

Leukemia

Immunosuppression, 78-7021

Leukemia, Myeloblastic

Colony Formation, 78-7150

Plant Agglutinins (cont'd)

- Lymphocytes
 - Cytotoxicity, 78-6988
- Mammary Neoplasms, Experimental
 - DNA Replication, 78-7004
 - Immunity, Cellular, 78-7004
 - Leukocyte Culture Test, Mixed, 78-7004
- Mycosis Fungoides
 - B-Lymphocytes, 78-7120
 - T-Lymphocytes, 78-7120
- Virus, Murine Leukemia
 - Lymphocytes, 78-6978
- Virus, Rauscher Murine Leukemia
 - Lymphocyte Transformation, 78-7017

Plasmacytoma

- Antigen-Antibody Reactions
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
- Antigens, Viral
 - Immune Response, Mouse, 78-7038
- Histocompatibility Antigens
 - Immune Response, Mouse, 78-7038
- Immunosuppression
 - Tumor Extract vs Culture Supernatant, 78-7003
- Kidney Neoplasms
 - Case Report, 78-7128
 - Ultrastructural Study, Giant Cells, 78-7128
- Macrophages
 - Cytotoxic Activity, 78-7002
- Radiation, Ionizing
 - Histocompatibility Antigens, 78-7038
- RNA Replication
 - Adenosine Triphosphate, 78-7190
 - α -Amanitine, 78-7190
 - Guanosine Triphosphate, 78-7190
 - RNA Polymerase, 78-7190
- RNA, Ribosomal
 - RNA Replication, 78-7190
- Silica
 - Histocompatibility Antigens, 78-7038
- Virus, Gross Murine Leukemia
 - Antigens, Viral, 78-7038

Plasmids

- see* Extrachromosomal Inheritance

Plasminogen Activators

- Carcinogen, Chemical
 - Two-Stage Carcinogenesis, Review, 78-6606
- Virus, SV40
 - Fibroblasts, Review, 78-6661

Platinum

- Ames Test
 - Carcinogenic Activity, Review, 78-6635

Plutonium

- Body Burden
 - Metabolism, Mouse, 78-6825
 - Specific Activity, Isotopes, 78-6825
- Lung Neoplasms
 - Risk Estimation, 78-6821
- Soil Pollutants
 - Inhalation Risk Factors, 78-6824
 - Plant Distribution, 78-6823
 - Size Fractionation Method, 78-6824

Plutonium Oxide

- Alpha Particles

Plutonium Oxide (cont'd)

- Dissolution, Water, 78-6818
- Body Burden
 - Inhalation Study, Rat, 78-6822
- Bone and Bones
 - Inhalation Study, Rat, 78-6834
- Lung Clearance
 - Hamster, Rat, 78-6819
 - Inhalation Study, Rat, 78-6834
- Lung Neoplasms
 - Inhalation Study, Rat, 78-6819, 78-6820
 - Radioisotopes, 78-6820
- Particle Size
 - Metabolism, Rat, 78-6822
- Uranium
 - Alpha Particles, 78-6818

Poly A

- Virus, Rauscher Murine Leukemia
 - RNA, Viral, 78-6886

Polyamines

- Kidney Neoplasms
 - Carboxy-Lyases, 78-6742
- Liver Neoplasms
 - Carboxy-Lyases, 78-6742

Polycyclic Hydrocarbons

- Air Pollution
 - Epidemiology, Review, 78-6664
 - Maximum Permissible Concentrations, Review 78-6664
- Brain Neoplasms
 - Species Difference, Review, 78-6639
- Carbonium Ion
 - Carcinogenic Metabolite, Review, 78-6623
- DNA Repair
 - Carcinogenic Activity, Review, 78-6622
- Hydro-Lyases
 - Metabolism, Review, 78-6623
- Microsomes, Liver
 - Ames Test, Review, 78-6635
- Oxygenases
 - Metabolism, Review, 78-6623
- Transferases
 - Metabolism, Review, 78-6623

Polyps

- Colitis, Granulomatous
 - Case Report, 78-7130
- Colitis, Ulcerative
 - Case Report, 78-7130
- Colon
 - Colitis, Granulomatous, 78-7130
 - Colitis, Ulcerative, 78-7130
- Enteritis, Regional
 - Case Report, 78-7130
- Intestinal Neoplasms
 - Precancerous Conditions, 78-7131
- Stomach
 - Enteritis, Regional, 78-7130
- Stomach Neoplasms
 - Achlorhydria, 78-7116
 - Epidemiology, 78-7116
 - Gastritis, 78-7116
 - Precancerous Conditions, 78-7116

Polyribonucleotides

- Leukemia
 - Drug Therapy, Review, 78-6669

polyribonucleotides (cont'd)

- Reverse Transcriptase
 - Enzymatic Activity, Review 78-6669
- Virus, Avian Sarcoma
 - RNA, Viral, 78-6972
- Virus, Leukemia
 - RNA, Viral, 78-6972
- Virus, Murine Sarcoma
 - RNA, Viral, 78-6972
- Virus, Rauscher Murine Leukemia
 - Ribonuclease, 78-6888
- Virus, RNA Tumor
 - Chromatography, 78-6972
 - RNA, Viral, 78-6972

polyribosomes

- Hepatoma
 - RNA, Messenger, 78-7189
- Virus, Herpes Simplex 1
 - RNA, Viral, 78-6942

polysaccharides

- Leukemia L1210
 - Immune Response, 78-7020

polyurethane Foam

- Respiratory Tract Neoplasms
 - Carcinogenic Potential, Rat, 78-6839

porphyria

- Hepatoma
 - Precancerous Conditions, 78-7123

precancerous Conditions

- Asbestos
 - Autoantibodies, 78-7061
 - Immunity, Cellular, 78-7061
 - B-Lymphocytes, 78-7061
 - T-Lymphocytes, 78-7061
- Bladder Neoplasms
 - 1-Butanol, 4-(Butylnitrosamino)-, 78-6735
 - Carcinoma In Situ, 78-7139
- Breast Neoplasms
 - Adenofibroma, 78-6700
 - Adipose Tissue, 78-7108
 - Contraceptives, Oral, 78-6648, 78-7107
 - Epidemiology, 78-6700
 - Hyperplasia, 78-7108
 - Mastitis, 78-6700
- Carcinogen, Chemical
 - Co-carcinogenic Activity, Review, 78-6611
- Carcinogen, Environmental
 - Co-carcinogenic Activity, Review, 78-6611
- Cervix Neoplasms
 - Contraceptives, Oral, 78-6648
 - Virus, Herpes Simplex 2, 78-6939, 78-7065
- Cholangioma
 - Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-, 78-6767
- Digestive System Neoplasms
 - Virus, Papilloma, 78-6899
- Esophageal Neoplasms
 - Carcinoma, Epidermoid, 78-7114
 - Carcinoma In Situ, 78-7114
 - Ethyl Alcohol, 78-7114
 - Smoking, 78-7114
- Gynecologic Neoplasms
 - 4,4'-Stilbenediol, α,α' -Diethyl-, 78-6652, 78-6789
- Head and Neck Neoplasms
 - Radiation, Ionizing, 78-6832

Precancerous Conditions (cont'd)

- Hepatoma
 - Acetamide, *N*-Fluoren-2-yl-, 78-6729
 - Aflatoxin B1, 78-6722
 - Australia Antigen, 78-6961
 - Diethylamine, *N*-Nitroso-, 78-6744
 - DNA Replication, 78-6744
 - Liver Cirrhosis, 78-7123
 - Porphyria, 78-7123
- Intestinal Neoplasms
 - Cell Differentiation, 78-7131
 - Polyps, 78-7131
 - Thymidine Incorporation, 78-7131
- Laryngeal Neoplasms
 - Enzymatic Activity, 78-7099
 - Neutrophils, 78-7099
- Leukemia, Lymphoblastic
 - Antigen-Antibody Reactions, 78-7026
 - Antigenic Determinants, 78-7026
 - Virus, Herpes, 78-6956
- Leukemia, Myeloblastic
 - Antigen-Antibody Reactions, 78-7026
 - Bronchitis, 78-7074
 - Case Report, 78-7074
 - Immunosuppressive Agents, 78-6695
- Liver Neoplasms
 - Glucosephosphate Dehydrogenase, 78-6787
 - Propane, 2-Nitro-, 78-6734
- Lung Neoplasms
 - Asbestos, 78-7061
- Lymphoma
 - Chromosome Aberrations, 78-6695
 - Immunosuppressive Agents, 78-6695
- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6774
 - 78-6776, 78-6780
 - Cell Differentiation, 78-6776
 - RNA, Viral, 78-6862
- Mouth Neoplasms
 - Areca, 78-6727
 - Basal Lamina, 78-7075
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-7075
 - Ultrastructural Study, Hamster, 78-7075
- Parathyroid Neoplasms
 - Transplantation, Autologous, 78-7103
- Prostatic Neoplasms
 - Reverse Transcriptase, 78-6976
 - Spermatozoa, 78-7142
- Sarcoma, Osteogenic
 - Strontium Radioisotopes, 78-6835
- Skin Neoplasms
 - Cholanthrene, 3-Methyl-, 78-6798
 - Collagen, 78-7087
 - Connective Tissue, 78-7087
 - Hyperplasia, 78-7087
- Stomach Neoplasms
 - Gastritis, 78-7116
 - Metaplasia, 78-7117
 - Polyps, 78-7116
 - Radiation, Ionizing, 78-6814
- Testicular Neoplasms
 - Spermatozoa, 78-7143
- Tracheal Neoplasms
 - Benz(a)anthracene, 78-6786
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6786
 - Benzo(a)pyrene, 78-6786
- Uterine Neoplasms

Precancerous Conditions (cont'd)

- Epidemiology, Review, 78-6642
- Virus, Epstein-Barr
- Antigen-Antibody Reactions, 78-6951

Progestational Hormones

- Liver Neoplasms
- Epidemiology, Review, 78-6650

Progesterone

- Breast Neoplasms
- Epidemiology, 78-7162
- Receptors, Hormone, Review, 78-6603
- Mammary Neoplasms, Experimental
- Virus, Murine Mammary Tumor, 78-6868

Prolactin

- Breast Neoplasms
- Alanine, 3-(3,4-Dihydroxyphenyl)-, 78-6728
- Epidemiology, 78-7162
- Melatonin, 78-6779
- Thyroid Dysfunction, Review, 78-6640
- Thyrotropin Releasing Hormone, 78-6640
- Mammary Neoplasms, Experimental
- Benz(a)anthracene, 7,12-Dimethyl-, 78-6778
- 78-6779
- Hypophyseal Isograft, 78-6868
- Thyroid Dysfunction, Review, 78-6640
- Virus, Murine Mammary Tumor, 78-6868
- Pituitary Neoplasms
- Adenoma, 78-7095
- Contraceptives, Oral, 78-7095

Proline, 4-Hydroxy-1-nitroso-

- Esterases
- Plasma Levels, Mouse, 78-6743

Propane, 2-Nitro-

- Liver Neoplasms
- Inhalation Study, Rat, 78-6734
- Precancerous Conditions, 78-6734

1-Propanol, 2,3-Dibromo-, Phosphate

- Occupational Hazard
- Textile Workers, Review, 78-6663

2-Propanol, 1,1'-Iminodi-N-nitroso-

- Brain Neoplasms
- Transplacental Carcinogenesis, 78-6749
- DNA Replication
- Brain, Rat, 78-6749

Prostaglandins

- Diterpene Esters
- Macrophages, 78-6719
- 12-O-Tetradecanoylphorbol-13-acetate
- Macrophages, 78-6719

Prostaglandins E

- Erythema
- Ultraviolet Rays, 78-6811
- Leukemia, Lymphoblastic
- Macrophages, 78-6993
- Macrophages
- Antitumor Activity, 78-6993
- Interferon, 78-6993
- Skin Neoplasms
- Cholanthrene, 3-Methyl-, 78-6798
- 12-O-Tetradecanoylphorbol-13-acetate
- Macrophages, 78-6782
- Ultraviolet Rays
- Indole-3-acetic Acid, 1-(p-Chlorobenzoyl)-5-

Prostaglandins E (cont'd)

- methoxy-2-methyl-, 78-6811
- Psoralen, 8-Methoxy-, 78-6811

Prostaglandins F

- Cell Division
- Embryo, Mouse, 78-7197
- Erythema
- Ultraviolet Rays, 78-6811
- Ultraviolet Rays
- Indole-3-acetic Acid, 1-(p-Chlorobenzoyl)-5-methoxy-2-methyl-, 78-6811
- Psoralen, 8-Methoxy-, 78-6811

Prostatic Hypertrophy

- Virus, Cytomegalo
- Antibodies, Viral, 78-7068
- Virus-Like Particles
- Isolation and Characterization, 78-6976

Prostatic Neoplasms

- Adenocarcinoma
- Reverse Transcriptase, 78-6976
- Epidemiology
- Iran, 78-7153
- Ethnic Groups
- Epidemiology, Hawaii, 78-7164
- Reverse Transcriptase
- Isolation and Characterization, 78-6976
- Precancerous Conditions, 78-6976
- Socioeconomic Factors
- Epidemiology, Review, 78-6698
- Spermatozoa
- Histological Study, Rat, 78-7142
- Precancerous Conditions, 78-7142
- Steroids
- Receptors, Hormone, Review, 78-6603
- Virus, Cytomegalo
- Antibodies, Viral, 78-7068
- Virus-Like Particles
- Reverse Transcriptase, 78-6976

Protein Kinase

- Virus, Rous Sarcoma
- Immunoglobulins, Heavy Chain, 78-6848
- Phosphoproteins, 78-6848

Proteins

- Acetic Acid, (Ethylenedinitrilo)tetra-
- Cell Transformation, Neoplastic, 78-6908
- Fibroblasts, 78-6908
- Cholanthrene, 3-Methyl-
- L Cells, 78-6908
- Heavy Chain Disease
- Amino Acids, 78-7060
- RNA, Messenger, 78-7060
- Trypsin
- Cell Transformation, Neoplastic, 78-6908
- Fibroblasts, 78-6908
- Urea
- Cell Transformation, Neoplastic, 78-6908
- Fibroblasts, 78-6908
- Virus, Avian Sarcoma
- Cell Membrane, 78-6841
- Cell Transformation, Neoplastic, 78-6841
- Isolation and Characterization, Chick Embryo Fibroblasts, 78-6841
- Virus, SV40
- Ethidium Bromide, 78-6915
- Fibroblasts, 78-6908

Endourea, 2-(2-Aminoethyl)-2-thio-

- Radiation, Ionizing
- Lymphocytes, 78-6979

Oralen, 8-Methoxy-

- Ames Test
- Mutagenic Activity, 78-6809
- Chromatids
- Hamster Ovary Cells, 78-6810
- Erythema
- Ultraviolet Rays, 78-6811

Escherichia coli

- Mutagenic Activity, 78-6809
- Hypoxanthine Phosphoribosyltransferase
- Mutagenic Activity, 78-6808

Mutagenic Activity

- Hamster V79 Cells, 78-6809
- Ultraviolet Rays
- Cell Transformation, Neoplastic, 78-6808
- Chromatids, 78-6810
- 5,8,11,14-Eicosatetraenoic Acid, 78-6811
- Mutagenic Activity, 78-6808, 78-6809
- Prostaglandins E, 78-6811
- Prostaglandins F, 78-6811

Oralen, 4,5'-8-Trimethyl-

- Virus, Herpes Simplex
- DNA, Viral, 78-6937

Oralen, 2-Amino-6-methoxy-

- Dimethylamine, *N*-Nitroso-
- DNA, Liver, 78-6625
- Enzymes, 78-6625

Hydrazine, 1,2-Dimethyl-

- DNA, Alkylation, 78-6755

Nitroso Compounds

- DNA, Liver, 78-6625
- Enzymes, 78-6625

Urea, Methyl Nitroso-

- Brain, Rat, 78-6757

azole

- Kidney Neoplasms
- Methanol, (Methyl-*ONN*-azoxy)-, 78-6760
- Skin Neoplasms
- Methanol, (Methyl-*ONN*-azoxy)-, 78-6760

ene

- Trachea
- Mucosal Toxicity, Rat, 78-6786
- Virus, Lymphocytic Choriomeningitis
- Cell Membrane, 78-6855
- Virus, Rous Sarcoma
- Cell Membrane, 78-6855

enebutyric Acid

- Virus, Lymphocytic Choriomeningitis
- Cell Membrane, 78-6855
- Virus, Rous Sarcoma
- Cell Membrane, 78-6855

Pyrrole-2,5-dione, 1-Ethyl-

- Radiation, Ionizing
- DNA Repair, 78-6812
- Ultraviolet Rays
- DNA Repair, 78-6812

macrine

- Leukemia L1210
- Ultraviolet Rays, 78-6807
- Ultraviolet Rays

Quinacrine (cont'd)

- DNA Repair, 78-6807

Quinoline

- Ames Test
- Mutagenic Activity, 78-6762
- Aryl Hydrocarbon Hydroxylases
- Mutagenic Activity, 78-6762
- Cholanthrene, 3-Methyl-
- Ames Test, 78-6762
- Microsomes, Liver
- Ames Test, 78-6762

Quinoline, 4-(Hydroxyamino)-, 1-Oxide

- Brain Neoplasms
- Transplacental Carcinogenesis, 78-6749
- DNA Replication
- Brain, Rat, 78-6749

Quinoline, 4-Nitro-, 1-Oxide

- Autoradiography
- Tissue Distribution, Mouse, 78-6763
- DNA Repair
- Carcinogenic Activity, Review, 78-6608
- Maternal-Fetal Exchange
- Tissue Distribution, Mouse, 78-6763

5-Quinolinol

- Ames Test
- Mutagenic Activity, 78-6762

8-Quinolinol

- Ames Test
- Mutagenic Activity, 78-6762

Radiation

- Abnormalities
- Teratogenic Effect, Review, 78-6616
- Ames Test
- Mutagenic Activity, 78-6792
- Chromosome Abnormalities
- Carcinogenic Activity, Review, 78-6610
- Hormones
- Co-carcinogenic Effect, Review, 78-6643
- Mutagenic Activity
- Micronucleus/Sperm Abnormality Assays, 78-6792

Radiation, Ionizing

- Fibrosarcoma
- Immunosuppression, Rat, 78-7067
- Leukemia
- Immunosuppression, Rat, 78-7067
- Mammary Neoplasms, Experimental
- Immunosuppression, Rat, 78-7067
- Adenoma
- Hyperthyroidism, 78-6831
- Ataxia Telangiectasia
- DNA Repair, Review, 78-6612
- Bone Marrow Cells
- Radiation-Protective Agents, 78-6979
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester
- DNA Repair, 78-6759
- Cell Transformation, Neoplastic
- Dose-Response Study, 78-6668
- Theoretical Model, 78-6665
- Corynebacterium parvum*
- Antibody Formation, 78-7057
- Antitumor Activity, 78-7057
- Phagocytosis, 78-7057
- DNA Repair
- Carcinogenic Activity, Review, 78-6619

Radiation, Ionizing (cont'd)

- Cell Transformation, Neoplastic, Review, 78-6613
- Cells, Cultured, 78-6812
- In Vivo Measurement Methods, Review, 78-6619
- Ear Neoplasms
 - Carcinogenic Potential, 78-6816
- Enzymatic Activity
 - Gastric Mucosa, Rat, 78-6814
- Eye Neoplasms
 - Carcinogenic Potential, 78-6816
- Fibrosarcoma
 - Cytotoxicity, Spleen Cells, 78-7053
 - Immunization, 78-7053
- Galactosidases
 - Enzymatic Activity, 78-6814
- Genetics
 - Epidemiology, Review, 78-6697
- Glucosidases
 - Enzymatic Activity, 78-6814
- Haemophilus influenzae*
 - DNA Repair, 78-6759
- Head and Neck Neoplasms
 - Carcinoma, Epidermoid, 78-6832
 - Precancerous Conditions, 78-6832
- Hearing Disorders
 - Epidemiology, 78-6816
- Hyperthyroidism
 - Case Report, 78-6831
- Leukemia
 - Case Report, 78-7072
 - Chromosome Abnormalities, 78-7072
 - Epidemiology, Review, 78-6667
 - Immunosuppression, 78-7021, 78-7022
 - Neoplasms, Multiple Primary, 78-7072
 - Occupational Hazard, 78-6667
- Leukemia L1210
 - Immunotherapy, 78-7020
- Leukemia, Myeloblastic
 - Case Report, 78-6815
- Leukemia, Myelocytic
 - Case Report, 78-6827
 - Chromosomes, Human, 21-22, 78-6827
- Leukemia, Radiation-Induced
 - DNA Nucleotidyltransferases, 78-6826
- Lymphocytes
 - Cysteine, 78-6979
 - Ethylamine, 2,2'-Dithiobis-, 78-6979
 - Glutathione, 78-6979
 - Pseudourea, 2-(2-Aminoethyl)-2-thio-, 78-6979
 - Radiation-Protective Agents, 78-6979
 - Serotonin, 78-6979
- Lymphoma
 - Immunity, Cellular, 78-7027
- Mammary Neoplasms, Experimental
 - Corynebacterium parvum*, 78-7057
 - Hormones, 78-6791
- Meningioma
 - Epidemiology, 78-7170
- Metaplasia
 - Gastric Mucosa, Rat, 78-6814
- Multiple Myeloma
 - Epidemiology, Review, 78-6667
 - Occupational Hazard, 78-6667
- Nuclear Reactors
 - Epidemiology, 78-6829
- Occupational Hazard
 - Dose-Response Study, Review, 78-6667

Radiation, Ionizing (cont'd)

- Epidemiology, Review, 78-6667
- Pancreatic Neoplasms
 - Epidemiology, Review, 78-6667
 - Occupational Hazard, 78-6667
- Pharyngeal Neoplasms
 - Carcinogenic Potential, 78-6816
- Plasmacytoma
 - Histocompatibility Antigens, 78-7038
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-
 - DNA Repair, 78-6812
- Saccharomyces cerevisiae*
 - DNA Repair, 78-6769
- Skin Neoplasms
 - Cholanthrene, 3-Methyl-, 78-7041
- Stomach Neoplasms
 - Precancerous Conditions, 78-6814
- Temporomandibular Joint Syndrome
 - Epidemiology, 78-6816
- Thymus Gland
 - Radiation-Protective Agents, 78-6979
- Thyroid Neoplasms
 - Adenocarcinoma, 78-7102
 - Adenoma, 78-7171
 - Case Report, 78-7102
 - Dose-Response Study, 78-7171
 - Epidemiology, 78-7171
 - Hyperthyroidism, 78-7102
- Trehalase
 - Enzymatic Activity, 78-6814
- Virus, Friend Murine Leukemia
 - Antigens, Viral, 78-6879
 - Virus Replication, 78-6878
- Virus, Marek's Disease Herpes
 - Immunity, Passive, 78-7023

Radioisotopes

- Leukemia
 - Thorium, 78-6830
- Lung Neoplasms
 - Americium Oxide, 78-6820
 - Curium Oxide, 78-6820
 - Plutonium Oxide, 78-6820
- Sarcoma, Osteogenic
 - Thorium, 78-6830

Radium

- Fibroma
 - Dose-Response Study, Mouse, 78-6830
- Hyperplasia
 - Spleen, Mouse, Rat, 78-6817
- Leukemia
 - Dose-Response Study, Mouse, 78-6830
- Sarcoma, Osteogenic
 - Dose-Response Study, Mouse, 78-6830

Receptors, Adrenergic

- Acetamide, *N*-Fluoren-2-yl-
 - Atria, Rat, Guinea Pig, 78-6732
 - Norepinephrine, 78-6732
- Diethylamine, *N*-Nitroso-
 - Atria, Rat, Guinea Pig, 78-6732

Rectal Neoplasms

- Water Supply
 - Epidemiology, 78-7175

Respiratory Tract Neoplasms

- Arsenic
 - Epidemiology, Review, 78-6634

Respiratory Tract Neoplasms (cont'd)

- Asbestos
 - Cummingtonite-Grunerite, 78-6838
 - Epidemiology, 78-6838
 - Epidemiology, Review, 78-6634
- Benzo(a)pyrene
 - 7-Acetic Acid, 5-Fluoro-12-methylbenzanthryl-78-7059
- Carcinogen, Chemical
 - Occupational Hazard, 78-6768
- Chromate
 - Epidemiology, Review, 78-6634
- Epidemiology
 - Iran, 78-7153
- Ethyl Alcohol
 - Epidemiology, Review, 78-6653
- Ethylene, Chloro-
 - Epidemiology, Review, 78-6634
- Nickel
 - Epidemiology, Review, 78-6634
- Occupational Hazard
 - Epidemiology, East Germany, 78-6768
 - Epidemiology, Review, 78-6634
 - Precancerous Conditions, Review, 78-6634
- Polyurethane Foam
 - Carcinogenic Potential, Rat, 78-6839
- Sex Factors
 - Epidemiology, England, Wales, 78-7157
- Smoking
 - Pulmonary Dysfunction, Review, 78-6658

Retinoblastoma

- Cell Differentiation
- Neoplasm Regression, Review, 78-6686

Retinol

- Benzo(a)pyrene
 - Carcinogenic Activity, Review, 78-6630
- Carcinogen, Chemical
 - Immune Response, Review, 78-6630
- Gastrointestinal Neoplasms
 - Mycotoxins, 78-6638
- Hepatoma
 - Carrier Proteins, 78-6796

Reverse Transcriptase

- Bladder Neoplasms
 - Carcinoma, Transitional Cell, 78-6973
 - Cells, Cultured, 78-6973
- Carcinoma, Epidermoid
 - Virus, Adeno 5, 78-6967
- Melanoma
 - Virus, Adeno 5, 78-6967
- Multiple Myeloma
 - Virus, C-Type RNA Tumor, 78-6897
- Myelofibrosis
 - Enzymatic Activity, Review, 78-6669
- Polyribonucleotides
 - Enzymatic Activity, Review, 78-6669
- Prostatic Neoplasms
 - Adenocarcinoma, 78-6976
 - Isolation and Characterization, 78-6976
 - Precancerous Conditions, 78-6976
 - Virus-Like Particles, 78-6976
- Sarcoma
 - Virus, Adeno 5, 78-6967
- Virus, Rauscher Murine Leukemia
 - Anti-Antibodies, 78-6888
 - Ribonuclease, 78-6885, 78-6888

Reverse Transcriptase (cont'd)

- Viral Proteins, 78-6887
- Virus, Sindbis, 78-6891
- Virus, RNA Tumor
 - Carcinogenic Activity, Review, 78-6669
- Virus, Rous Sarcoma
 - Endonucleases, 78-6854
 - RNA, Transfer, Binding, 78-6851
 - Temperature Sensitive Mutants, 78-6851

Rhabdomyosarcoma

- Virus-Like Particles
- Ultrastructural Study, 78-6977

Rhodium

- Ames Test
- Carcinogenic Activity, Review, 78-6635

Ribonuclease

- Bladder Neoplasms
 - Carcinoma, Transitional Cell, 78-6973
 - DNA Polymerase, 78-6973
- Pancreatic Neoplasms
 - Carcinoma, 78-7200
 - Carcinoma, Ductal, 78-7200
 - Islet Cell Tumor, 78-7200
 - Serum Levels, 78-7200
- Pancreatitis
 - Serum Levels, 78-7200
- Virus, Rauscher Murine Leukemia
 - Anti-Antibodies, 78-6888
 - Polyribonucleotides, 78-6888
 - Reverse Transcriptase, 78-6885, 78-6888
 - RNA, Viral, 78-6888
 - Temperature Sensitive Mutants, 78-6885
 - Thermolability, 78-6885
- Virus, Simian Sarcoma
 - RNA, Viral, 78-6888

Ribonucleotide Reductases

- Virus, Epstein-Barr
 - Urea, Hydroxy-, 78-6950
 - Uridine, 2'-Deoxy-5-iodo-, 78-6950
 - Virus Replication, 78-6950

Ribosomes

- Cholanthrene, 3-Methyl-
 - Proteins, Phosphorylation, 78-6969
- Virus, Adeno 12
 - Proteins, Phosphorylation, 78-6969
- Virus, Murine Sarcoma
 - Proteins, Phosphorylation, 78-6969

Rifampicin

- Virus, Polyoma
 - RNA Replication, 78-6901

RNA, Messenger

- Erythroleukemia
 - Globin, 78-7185
- Heavy Chain Disease
 - Immunoglobulins, 78-7060
 - Proteins, 78-7060
- Hepatoma
 - Nucleoproteins, 78-7189
 - Polyribosomes, 78-7189
- Nucleoproteins
 - Liver Regeneration, 78-7189
- Urea, Ethyl Nitroso-
 - Brain, Rat, 78-6757
- Virus, Adeno 2

RNA, Messenger (cont'd)

- Chromosomes, 78-6964
- DNA-RNA Hybridization, 78-6964
- Hybrid Cells, 78-6962
- Peptides, 78-6963
- RNA Replication, 78-6964
- Viral Proteins, 78-6964

Virus, Adeno 2 - SV40 Hybrid

- HeLa Cells, 78-6966
- Isolation and Characterization, 78-6966
- Nucleotide Sequence, 78-6966
- Viral Proteins, 78-6966

Virus, Avian Sarcoma

- Genes, Viral, 78-6840
- Viral Proteins, 78-6840
- Virus Replication, 78-6840

Virus, Polyoma

- Isolation and Characterization, 78-6904
- Viral Proteins, 78-6904

Virus, RNA Tumor

- Isolation and Characterization, Review, 78-6675
- Viral Proteins, Review, 78-6675

Virus, Rous Sarcoma

- Viral Proteins, 78-6852

Virus, SV40

- DNA-RNA Hybridization, 78-6920
- DNA, Viral, 78-6920
- RNA, Viral, 78-6921

RNA Polymerase

- Breast Neoplasms
 - Estrogens, 78-6603
- Plasmacytoma
 - RNA Replication, 78-7190
- Virus, Polyoma
 - DNA, Viral, 78-6901

RNA Replication

- Plasmacytoma
 - Adenosine Triphosphate, 78-7190
 - α -Amanitine, 78-7190
 - Guanosine Triphosphate, 78-7190
 - RNA Polymerase, 78-7190
 - RNA, Ribosomal, 78-7190
- Virus, Adeno 2
 - RNA, Messenger, 78-6964
- Virus, Murine Mammary Tumor
 - Dexamethasone, 78-6863
 - Xenopus laevis*, 78-6865
- Virus, Polio
 - L Cells, 78-6971
- Virus, Polyoma
 - DNA, Viral, 78-6901
 - Rifampicin, 78-6901

RNA, Ribosomal

- Acetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-
 - Liver, Rat, 78-6716
- 2-Butanone, 1,1-Dihydroxy-3-ethoxy-
 - Liver, Rat, 78-6716
- Plasmacytoma
 - RNA Replication, 78-7190
- Terbium
 - Fluorescence, 78-6716

RNA, Transfer

- Virus, Avian Sarcoma
 - Viral Proteins, 78-6840

RNA, Transfer, Methyltransferases

- Methionine, *S*-Adenosyl-
 - Ultraviolet Rays, 78-6621
- Uracil, 78-6621
- Uracil
 - Carcinoembryonic Antigen, 78-6621
 - Enzymatic Activity, 78-6621

RNA, Viral

- Mammary Neoplasms, Experimental
 - Hormones, 78-6868
 - Precancerous Conditions, 78-6862
 - Virus, Murine Mammary Tumor, 78-6868
- Virus, Abelson Murine Leukemia
 - DNA-RNA Hybridization, Review, 78-6678
- Virus, Avian Leukosis-Sarcoma
 - DNA-RNA Hybridization, 78-6843
- Virus, Avian Sarcoma
 - Cell-Free Translation, 78-6840
 - Polyribonucleotides, 78-6972
- Virus, Friend Spleen Focus-Forming
 - DNA-RNA Hybridization, Review, 78-6678
- Virus, Harvey Murine Sarcoma
 - Nucleotide Sequence, 78-6894
- Virus, Herpes Simplex 1
 - Isolation and Characterization, 78-6942
 - Polyribosomes, 78-6942
- Virus, Kirsten Murine Sarcoma
 - Nucleotide Sequence, 78-6894
- Virus, Leukemia
 - Polyribonucleotides, 78-6972
- Virus, Moloney Murine Leukemia
 - Nucleotide Sequence, 78-6882
 - Virus, Recombinant, 78-6882
- Virus, Murine Mammary Tumor
 - DNA-RNA Hybridization, 78-6864
 - Isolation and Characterization, 78-6864
 - Mammary Neoplasms, Experimental, 78-6862
 - Viral Proteins, 78-6865
- Virus, Murine Sarcoma
 - DNA-RNA Hybridization, Review, 78-6678
 - Polyribonucleotides, 78-6972
- Virus, Polyoma
 - Viral Proteins, 78-6904
- Virus, Rauscher Murine Leukemia
 - Isolation and Characterization, 78-6886
 - Poly A, 78-6886
 - Ribonuclease, 78-6888
- Virus, RNA Tumor
 - Polyribonucleotides, 78-6972
- Virus, Rous Sarcoma
 - Cell Transformation, Neoplastic, 78-6859
 - DNA-RNA Hybridization, 78-6859
 - Endonucleases, 78-6854
 - Nucleotide Sequence, 78-6859
 - Viral Proteins, 78-6852
- Virus, Simian Sarcoma
 - Ribonuclease, 78-6888
- Virus, SV40
 - Cytoplasm, Nucleus, 78-6921
 - Degradation, 78-6921
 - DNA-RNA Hybridization, 78-6921
 - RNA, Messenger, 78-6921

Ruthenium

- Ames Test
 - Carcinogenic Activity, Review, 78-6635

Saccharomyces cerevisiae

- Anthracycline
 - DNA Repair, 78-6769
- Antibiotics, Antineoplastic
 - DNA Repair, 78-6769
- Radiation, Ionizing
 - DNA Repair, 78-6769
- Sibiromycin
 - DNA Repair, 78-6769
- Tomaymycin
 - DNA Repair, 78-6769
- Ultraviolet Rays
 - DNA Repair, 78-6769

Salivary Gland Neoplasms

- Sjogren's Syndrome
 - Histological Study, 78-7082
 - Immunoglobulins, Light Chain, 78-7082

Salmonella typhimurium

- Acetohydroxamic Acid, *N*-Fluorenyl-2-yl-
 - Mutagenic Activity, 78-6730
- Mutagens
 - Azaguanine Resistance, 78-6761
 - Histidine Reversion, 78-6761

Sarcoma

- Acetic Anhydride
 - Transplantation Immunology, 78-7048
- Antigen-Antibody Complex
 - Complement Binding Assay, 78-7042
 - Transplantation Immunology, 78-7042
- Ascitic Fluid
 - Anti-Inflammatory Activity, 78-6981
- BCG*
 - Macrophages, 78-6995
 - Neoplasm Regression, 78-6995
- 2-Benzimidazolecarbamic Acid, 5-(2-Thienylcarbonyl)-
 - Methyl Ester
 - Cell Migration Inhibition, 78-7081
- Benzo(a)pyrene
 - Graft vs Host Reaction, 78-7045
 - B-Lymphocytes, 78-7045
 - T-Lymphocytes, 78-7045
- Bleomycin
 - Cell Migration Inhibition, 78-7081
- Carbodiimide, 1-Ethyl-3-(3-dimethylaminopropyl)-
 - Transplantation Immunology, 78-7048
- Cell Migration Inhibition
 - Cells, Cultured, 78-7081
- Cholanthrene, 3-Methyl-
 - Antigen-Antibody Complex, 78-7042
 - Histocompatibility Antigens, 78-7043
 - Macrophages, 78-6994
 - Neoplasm Metastasis, 78-7043
- Colchicine
 - Cell Migration Inhibition, 78-7081
- Colchicine, *N*-Deactyl-*N*-methyl-
 - Cell Migration Inhibition, 78-7081
- Cytosine, 1- β -*D*-Arabinofuranosyl-
 - Cell Migration Inhibition, 78-7081
- Epidemiology
 - Pediatric Patients, 78-7177
- Erythrocytes
 - Hemodynamics, 78-7078
 - Mitosis, 78-7078
- Glutaraldehyde
 - Transplantation Immunology, 78-7048
- Graft vs Host Reaction

Sarcoma (cont'd)

- B-Lymphocytes, 78-7045
 - T-Lymphocytes, 78-7045
 - Neoplasm Metastasis, 78-7045
 - Histocompatibility Antigens
 - Neoplasm Metastasis, 78-7043
 - Neoplasm Transplantation, 78-7043
 - Transplantation Immunology, 78-7044
 - Macrophages
 - Activation, Ascites Cells, 78-6994
 - Immune Response, 78-6994
 - Neoplasm Metastasis, 78-6994
 - Membrane Fraction
 - Immunogenetics, 78-7048
 - Methylamine
 - Transplantation Immunology, 78-7048
 - Neoplasm Metastasis
 - Animal Model, Mouse, 78-7043
 - Neoplasm Transplantation
 - Hemodynamics, 78-7078
 - Skin Neoplasms
 - Cholanthrene, 3-Methyl-, 78-7041
 - Sulfuric Acid, Dimethyl Ester
 - Transplantation Immunology, 78-7048
 - Uracil, 5-Fluoro
 - Cell Migration Inhibition, 78-7081
 - Vinblastine Sulfate
 - Cell Migration Inhibition, 78-7081
 - Vincristine Sulfate
 - Cell Migration Inhibition, 78-7081
 - Virus, Adeno 5
 - Reverse Transcriptase, 78-6967
 - Virus Activation, 78-6967
 - Virus Replication, 78-6967
 - Virus, C-Type RNA Tumor
 - Ultrastructural Study, Review, 78-6679
 - Virus, Marek's Disease Herpes
 - Antibodies, Viral, 78-6861
 - Antigens, Viral, 78-6861
 - Virus, Rous Sarcoma
 - Immunogenetics, 78-7048
 - Virus, SV40
 - Histocompatibility Antigens, 78-7044
 - Xanthine, 3-Hydroxy-
 - Carcinogenic Activity, Rat, 78-6718
- Sarcoma 180, Crocker**
- Lyases
 - Enzymatic Activity, Liver, 78-7187
 - Neoplasm Invasiveness
 - Mesenchyme, Chick Wing, 78-7079
 - Neoplasm Transplantation
 - Contact Inhibition, 78-7079
 - Neoplasm Invasiveness, 78-7079
- Sarcoma, Ewing's**
- Leukemia, Myeloblastic
 - Neoplasms, Multiple Primary, 78-6815
- Sarcoma, Mast Cell**
- Antigen-Antibody Reactions
 - Immunosuppression, 78-7003
 - Spleen Cells, 78-7003
 - Immunosuppression
 - Tumor Extract vs Culture Supernatant, 78-7003
- Sarcoma, Osteogenic**
- Bone Neoplasms
 - Neoplasm Recurrence, Local, 78-7147

- Sarcoma, Osteogenic (cont'd)**
 Strontium Radioisotopes, 78-6835
 Cells, Cultured
 Chromosomes, 78-7148
 Lactate Dehydrogenase Isoenzymes, 78-7148
 Radium
 Dose-Response Study, Mouse, 78-6830
 Strontium Radioisotopes
 Dose-Response Study, Rat, 78-6835
 Precancerous Conditions, 78-6835
 Thorium
 Dose-Response Study, Mouse, 78-6830
 Radioisotopes, 78-6830
 Yttrium Radioisotopes
 Dose-Response Study, Rat, 78-6835
- Sarcoma, Reticulum Cell**
 Chromosome Aberrations
 Case Report, 78-7080
 Chromosomes, Human, 13-15, 78-7080
 Lymph Nodes
 Karyotyping, 78-7080
 Virus, Epstein-Barr
 Antigens, Viral, 78-6948
 Case Report, 78-6948
 Virus-Like Particles, 78-6948
- Schistosoma**
 Neoplasms, Experimental
 Immune Response, Review, 78-6633
 Precancerous Conditions, Review, 78-6633
- Schistosoma haematobium**
 Bladder Neoplasms
 Carcinoma, Transitional Cell, 78-7140
 Histological Study, Monkey, 78-7140
 Neoplasm Transplantation, 78-7140
- Schistosoma mansoni**
 Silicic Acid, Magnesium Salt
 Immune Response, 78-6992
 Macrophage Activation, 78-6992
- Selenium**
 Breast Neoplasms
 Antitumor Activity, Review, 78-6696
 Epidemiology, Review, 78-6696
 Intestinal Neoplasms
 Adenocarcinoma, 78-6696
 Antitumor Activity, Review, 78-6696
 Epidemiology, Review, 78-6696
 Mammary Neoplasms, Experimental
 Antitumor Activity, Review, 78-6696
- Serotonin**
 Radiation, Ionizing
 Lymphocytes, 78-6979
- Serum Albumin, Bovine**
 Benzo(a)pyrene
 7-Acetic Acid, 5-Fluoro-12-methylbenzanthryl-
 78-7059
 Immunization, 78-7059
- Sex Chromosomes**
 Bladder Neoplasms
 Cell Line, Epithelioid, 78-7199
 Leukemia, Myelocytic
 Blast Crisis, Shay Tumor, 78-7069
 Neoplasm Transplantation, 78-7069
- Sibiromycin**
Saccharomyces cerevisiae
 DNA Repair, 78-6769
- Silica**
Corynebacterium parvum
 Antibody Formation, 78-7057
 Antitumor Activity, 78-7057
 Phagocytosis, 78-7057
 Lung Neoplasms
 Neoplasm Metastasis, 78-6998
 Phagocytosis, 78-6998
 Mammary Neoplasms, Experimental
Corynebacterium parvum, 78-7057
 Plasmacytoma
 Histocompatibility Antigens, 78-7038
- Silicic Acid, Magnesium Salt**
Listeria monocytogenes
 Immune Response, 78-6992
 Macrophage Activation, 78-6992
 Lung Neoplasms
 Immune Response, 78-6992
 Macrophage Activation, 78-6992
Schistosoma mansoni
 Immune Response, 78-6992
 Macrophage Activation, 78-6992
- Sjogren's Syndrome**
 Lymphoma
 Histological Study, 78-7082
 IgM, 78-7082
 Immunoglobulins, Light Chain, 78-7082
 B-Lymphocytes, 78-7082
 Salivary Gland Neoplasms
 Histological Study, 78-7082
 Immunoglobulins, Light Chain, 78-7082
- Skin Neoplasms**
 Areca
 Carcinogenic Potential, Ingredients, 78-6727
 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
 Croton Oil, 78-6660
 Carcinogen, Chemical
 Co-carcinogenic Activity, Review, 78-6611
 Carcinoma, Basal Cell
 Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
 Wounds and Injuries, 78-7092
 Carcinoma, Epidermoid
 Cholanthrene, 3-Methyl-, 78-7041
 Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
 Ultraviolet Rays, 78-6813
 Virus, Papilloma, 78-6684
 Xeroderma Pigmentosum, 78-7121
 Cholanthrene, 3-Methyl-
 Antilymphocyte Serum, 78-7041
 Calcitonin, 78-6798
 Cortisol, 78-6798
 Mouse, New Zealand Black, 78-7041
 Precancerous Conditions, 78-6798
 Prostaglandins E, 78-6798
 Radiation, Ionizing, 78-7041
 Thyroxine, 78-6798
 Collagen
 Precancerous Conditions, 78-7087
 Connective Tissue
 Mitosis, 78-7087
 Precancerous Conditions, 78-7087
 Diethylamine, 2,2'-Dichloro-*N*-methyl-

Skin Neoplasms (cont'd)

- Mycosis Fungoides, 78-6753
- Epidemiology
 - Iran, 78-7153
- Hair Dyes
 - Epidemiology, 78-7163
- Hyperplasia
 - Precancerous Conditions, 78-7087
- Keratoacanthoma
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
- Keratinosis
 - Ultraviolet Rays, 78-6813
- Mathematical Model
 - Age Factors, 78-6804
- Melanoma
 - Lymphocyte Transformation, Review, 78-6693
 - B-Lymphocytes, 78-6693
 - T-Lymphocytes, 78-6693
 - Precancerous Conditions, Review, 78-6693
- Methanol, (Methyl-*ONN*-azoxy)-
 - Pyrazole, 78-6760
- Mycosis Fungoides
 - Drug Therapy, 78-6753
- Neoplasms, Multiple Primary
 - Diethylamine, 2,2'-Dichloro-*N*-methyl-, 78-6753
 - Xeroderma Pigmentosum, 78-7121
- Papilloma
 - Methanol, (Methyl-*ONN*-azoxy)-, 78-6760
- Phosphonium, Tetrakis(hydroxymethyl)-, Chloride
 - Carcinogenic Activity, Mouse, 78-6725
 - Oxalate Salt, 78-6725
- Sarcoma
 - Cholanthrene, 3-Methyl-, 78-7041
- 12-*O*-Tetradecanoylphorbol-13-acetate
 - Two-Stage Carcinogenesis, Review, 78-6606
- Trace Elements
 - Epidemiology, Review, 78-6631
- Ultraviolet Rays
 - Age Factors, 78-6804
 - Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6806
 - p*-Cresol, 2,6-Di-*tert*-butyl-, 78-6806
 - Disulfide, Bis(diethylthiocarbamoyl)-, 78-6806
 - Enzymatic Activity, 78-6813
 - Epidemiology, 78-6804
 - Glucosephosphate Dehydrogenase, 78-6813
 - Glutathione, 78-6806
- Virus, Papilloma
 - Carcinogenic Activity, Review, 78-6686
 - Epidemiology, Review, 78-6684
 - Immune Response, Review, 78-6684
 - Immunity, Cellular, 78-7036
 - Isolation and Characterization, 78-7036
- Warts
 - Epidemiology, Review, 78-6684
 - Genetics, Review, 78-6684
 - Virus, Papilloma, 78-7036
- Xeroderma Pigmentosum
 - Case Report, 78-7121
 - DNA Repair, 78-7121
 - Genetics, 78-7121

Smallpox Vaccine

- Carcinoma, Basal Cell
- Cicatrix, 78-7089

Smoking

- Adenocarcinoma
 - Carcinoembryonic Antigen, 78-7118

Smoking (cont'd)

- Benzo(a)pyrene
 - Co-carcinogenic Activity, 78-6701
- Breast Neoplasms
 - Carcinoembryonic Antigen, 78-7066
- Bronchial Neoplasms
 - Epidemiology, Switzerland, 78-7151
- Carcinoembryonic Antigen
 - Genetics, 78-7118
- Cervix Neoplasms
 - Epidemiology, Yugoslavia, 78-6945
- Esophageal Neoplasms
 - Precancerous Conditions, 78-7114
- Ethyl Alcohol
 - Co-carcinogenic Activity, Review, 78-6653
- Ethylene, Chloro-
 - Chromosome Abnormalities, 78-6715
- Ethylene, Chloro- Polymer
 - Chromosome Abnormalities, 78-6715
- Lung Neoplasms
 - Epidemiology, 78-7156
- Respiratory Tract Neoplasms
 - Pulmonary Dysfunction, Review, 78-6658

Snuff

- Mouth Neoplasms
 - Epidemiology, 78-6720

Sodium Chloride

- Virus, Epstein-Barr
 - Anticomplement Immunofluorescence Test, 78-6953

Sodium Fluoride

- Benzenamine, *N,N*-Dimethyl-4-((3-methylphenyl)azo)-
 - Adenyl Cyclase, 78-6766

Soil Pollutants

- Plutonium
 - Inhalation Risk Factors, 78-6824
 - Plant Distribution, 78-6823
 - Size Fractionation Method, 78-6824

Solvents

- Leukemia
 - Case Report, 78-6765
 - Epidemiology, Child, 78-6765

Spermatozoa

- Prostatic Neoplasms
 - Histological Study, Rat, 78-7142
 - Precancerous Conditions, 78-7142
- Testicular Neoplasms
 - Histological Study, Rat, 78-7143
 - Precancerous Conditions, 78-7143

Spinal Cord Neoplasms

- Astrocytoma
 - Case Report, 78-7098
 - Neoplasms, Multiple Primary, 78-7098

Spinal Neoplasms

- Meningioma
 - Genetics, 78-7096

Stein-Leventhal Syndrome

- Uterine Neoplasms
 - Case Report, 78-7133

Sterigmatocystin

- Mesothelioma
 - Histological Study, Rat, 78-6800
- Mycotoxins

Sterigmatocystin (cont'd)

Carcinogenic Activity, Review, 78-6629

Peritoneal Neoplasms

Mesothelioma, 78-6800

Steroids**Breast Neoplasms**

Receptors, Hormone, Review, 78-6603

Prostatic Neoplasms

Receptors, Hormone, Review, 78-6603

4,4'-Stilbenediol, α,α' -Diethyl-**Antibody Formation**

IgG, 78-7011

Transplacental Effects, 78-7011

Antigens

Antibody Formation, 78-7011

Gynecologic Neoplasms

Epidemiology, Review, 78-6644

Histological Study, 78-6789

Precancerous Conditions, 78-6652, 78-6789

Transplacental Carcinogenesis, 78-6789

Transplacental Carcinogenesis, Review, 78-6652

Lipopolysaccharides

Antibody Formation, 78-7011

Lung Neoplasms

Phagocytosis, 78-6998

T-Lymphocytes

Immune Response, 78-7011

Testis

Abnormalities, Review, 78-6652

Stomach

Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester

Synthesis, Guinea Pig, Rat, 78-6754

Polyps

Enteritis, Regional, 78-7130

Stomach Neoplasms**Adenocarcinoma**

Histochemical Study, 78-7117

Transplantation, Heterologous, 78-7115

Asbestos

Occupational Hazard, 78-7172

Carcinoembryonic Antigen

Antigen-Antibody Reactions, 78-7058

Carcinoma, Mucinous

Histochemical Study, 78-7117

Diet

Epidemiology, Review, 78-6637

Gastritis

Precancerous Conditions, 78-7116

Metaplasia

Gastric Mucosa, 78-7117

Precancerous Conditions, 78-7117

Nitric Acid

Food Contamination, 78-6733

Nitrous Acid

Food Contamination, 78-6733

Occupational Hazard

Epidemiology, 78-7172

Latency Period, 78-7172

Polyps

Achlorhydria, 78-7116

Epidemiology, 78-7116

Gastritis, 78-7116

Precancerous Conditions, 78-7116

Radiation, Ionizing

Precancerous Conditions, 78-6814

Stomach Neoplasms (cont'd)**Transplantation, Heterologous**

Cell Differentiation, 78-7115

Mouse, Nude, 78-7115

Neoplasm Metastasis, 78-7115

Streptozotocin**Kidney Neoplasms**

Adenoma, 78-6799

Dietary Proteins, 78-6799

Strontium Radioisotopes**Bone Neoplasms**

Angiosarcoma, 78-6835

Fibrosarcoma, 78-6835

Sarcoma, Osteogenic, 78-6835

Hyperplasia

Spleen, Mouse, Rat, 78-6817

Sarcoma, Osteogenic

Dose-Response Study, Rat, 78-6835

Precancerous Conditions, 78-6835

Styrene**Leukemia, Lymphoblastic**

Case Report, 78-6765

Occupational Hazard

Printers, Review, 78-6663

Succinate Dehydrogenase**Ultraviolet Rays**

Enzymatic Activity, 78-6813

Succinic Acid

Aniline, *N,N*-Dimethyl-*p*-phenylazo-

Mitochondrial Swelling, 78-6764

Sulfide, Bis(2-chloroethyl)-**Carcinoma, Epidermoid**

Case Report, 78-7088

Neoplasm Recurrence, Local, 78-7088

Sulfobromophthalein

see Phthalide, 4,5,6,7-Tetrabromo-3,3-bis(4-hydroxy-3-sulfo)phenyl

Sulfuric Acid, Dimethyl Ester**Sarcoma**

Transplantation Immunology, 78-7048

Temporomandibular Joint Syndrome**Radiation, Ionizing**

Epidemiology, 78-6816

Teratoid Tumor**Cell Differentiation**

Neoplasm Regression, Review, 78-6686

Neoplasm Suppression, Review, 78-6705

Virus, Polyoma

Antigens, Viral, 78-6902

Hybrid Cells, 78-6902

Virus, SV40

Antigens, Viral, 78-6932

Terbium**Fluorescence**

DNA, 78-6716

Liver, Rat, 78-6716

Guanine

Fluorescence, 78-6716

RNA, Ribosomal

Fluorescence, 78-6716

Testicular Neoplasms

- Aneuploidy
 - Histological Study, Rat, 78-7143
- Chromosome Abnormalities
 - Histological Study, Rat, 78-7143
- Disgerminoma
 - Case Report, 78-7145
 - Chromosome Abnormalities, 78-7145
 - Histological Study, 78-7146
- Spermatozoa
 - Histological Study, Rat, 78-7143
 - Precancerous Conditions, 78-7143

Testis

- 4,4'-Stilbenediol, α, α' -Diethyl-
 - Abnormalities, Review, 78-6652

Testosterone

- Virus, Murine Leukemia
 - Viral Proteins, 78-6871

Testosterone, Propionate

- Neoplasms, Experimental
 - Virus, SV40, 78-6911

12-*O*-Tetradecanoylphorbol-13-acetate

- Benz(a)anthracene, 7,12-Dimethyl-
 - Two-Stage Carcinogenesis, Review, 78-6626
- Benzo(a)pyrene
 - Two-Stage Carcinogenesis, Review, 78-6626
- Cell Division
 - Embryo, Mouse, 78-7197
- Cells, Cultured
 - Mutagenic Activity, Review, 78-6606
- Fibroblasts
 - Two-Stage Carcinogenesis, Review, 78-6626
- Hepatoma
 - Carrier Proteins, 78-6796
- Macrophages
 - Antitumor Activity, 78-6782
 - Prostaglandins E, 78-6782
- Prostaglandins
 - Macrophages, 78-6719
- Skin Neoplasms
 - Two-Stage Carcinogenesis, Review, 78-6606
- Ultraviolet Rays
 - Mutagenic Activity, Review, 78-6606
 - Ouabain Resistance, 78-6606
 - Thioguanine Resistance, 78-6606

Thorium

- Leukemia,
 - Dose-Response Study, Mouse, 78-6830
 - Radioisotopes, 78-6830
- Sarcoma, Osteogenic
 - Dose-Response Study, Mouse, 78-6830
 - Radioisotopes, 78-6830

Thymidine

- Leukemia, Myeloblastic
 - Cell Cycle Kinetics, 78-7150

Thymidine Kinase

- Carcinoma, Transitional Cell
 - Virus, Equine Rhinopneumonitis, 78-6898
- Virus, Equine Rhinopneumonitis
 - Enzymatic Activity, 78-6898
- Fibroblasts, 78-6898
- Virus, Herpes Simplex 1
 - Mutagenic Activity, 78-6941

Thymine, 1- β -*D*-Arabinofuranosyl-

- Virus, Equine Rhinopneumonitis
 - Virus Replication, 78-6898

Thymoma

- Antibodies, Viral
 - Virus Replication, 78-6874
- Antigen-Antibody Complex
 - Transplantation Immunology, 78-7012
- T-Lymphocytes
 - Antigen-Antibody Complex, 78-7012
 - Immunoglobulins, Surface, 78-7012
- Myasthenia Gravis
 - Thymectomy, 78-7035
- Virus, Radiation Leukemia
 - Isolation and Characterization, 78-6874

Thymus Gland

- Leukemia, Radiation-Induced
 - DNA Nucleotidyltransferases, 78-6826
- Radiation, Ionizing
 - Radiation-Protective Agents, 78-6979

Thyroid Neoplasms

- Adenocarcinoma
 - Neoplasm Metastasis, 78-7102
 - Radiation, Ionizing, 78-7102
- Adenoma
 - Radiation, Ionizing, 78-7171
- Hair Dyes
 - Epidemiology, 78-7163
- Hyperthyroidism
 - Radiation, Ionizing, 78-7102
- Iodine Radioisotopes
 - Dose-Response Study, 78-6833
 - Mutagenic Activity, 78-6833
- Pheochromocytoma
 - Genetics, 78-7144
- Radiation, Ionizing
 - Case Report, 78-7102
 - Dose-Response Study, 78-7171
 - Epidemiology, 78-7171

Thyrotropin

- Breast Neoplasms
 - Thyroid Dysfunction, Review, 78-6640
 - Thyrotropin Releasing Hormone, 78-6640
- Mammary Neoplasms, Experimental
 - Benz(a)anthracene, 7,12-Dimethyl-, 78-6778
 - Thyroid Dysfunction, Review, 78-6640

Thyrotropin Releasing Hormone

- Breast Neoplasms
 - Prolactin, 78-6640
 - Thyrotropin, 78-6640

Thyroxine

- Skin Neoplasms
 - Cholanthrene, 3-Methyl-, 78-6798

Tomaymycin

- Saccharomyces cerevisiae*
 - DNA Repair, 78-6769

Tongue Neoplasms

- Carcinoma, Epidermoid
 - Case Report, 78-7098
 - Neoplasm Metastasis, 78-7098
- Urea, 1-(2-Chloroethyl)-3-cyclohexyl-1-nitroso-
 - 78-7098

Tooth Diseases

- Glioblastoma Multiforme
Bacteria, 78-7149
Epidemiology, 78-7149
- Meningioma
Epidemiology, 78-7149

Toxocara canis

- Granuloma
Connective Tissue, Mouse, 78-7085
Ultrastructural Study, Mouse, 78-7085

Trace Elements

- Gastrointestinal Neoplasms
Epidemiology, Review, 78-6631
- Lung Neoplasms
Epidemiology, Review, 78-6631
- Skin Neoplasms
Epidemiology, Review, 78-6631

Trachea

- Anthracene
Mucosal Toxicity, Rat, 78-6786
- Benz(a)anthracene
Mucosal Toxicity, Rat, 78-6786
- Benz(a)anthracene, 7,12-Dimethyl-
Mucosal Toxicity, Rat, 78-6786
- Benzo(a)pyrene
Mucosal Toxicity, Rat, 78-6786
- Benzo(b)triphenylene
Mucosal Toxicity, Rat, 78-6786
- Benzo(e)pyrene
Mucosal Toxicity, Rat, 78-6786
- Pyrene
Mucosal Toxicity, Rat, 78-6786

Tracheal Neoplasms

- Benz(a)anthracene
Precancerous Conditions, 78-6786
- Benz(a)anthracene, 7,12-Dimethyl-
Precancerous Conditions, 78-6786
- Benzo(a)pyrene
Precancerous Conditions, 78-6786

Transfection

- see Transformation, Genetic

Transferases

- Polycyclic Hydrocarbons
Metabolism, Review, 78-6623

Transformation, Genetic

- Oncogenic Viruses
Stress, Review, 78-6682
- Virus, Rous Sarcoma
DNA, Viral, 78-6847

Transplantation, Autologous

- Adenoma
Parathyroid Glands, 78-7103
- Parathyroid Neoplasms
Precancerous Conditions, 78-7103

Transplantation, Heterologous

- Brain Neoplasms
Nude Mouse, Review, 78-6639
- Burkitt's Lymphoma
Antilymphocyte Serum, 78-7086
Histological Study, Hamster, 78-7086
Neoplasm Invasiveness, 78-7086
- Fibroblasts
Cell Transformation, Neoplastic, 78-7079

Transplantation, Heterologous (cont'd)

- Mesenchyme, Chick Wing, 78-7079
- Leukemia
Burkitt's Lymphoma, 78-7086
- Neoplasms, Experimental
Neoplasm Invasiveness, 78-7079
- Stomach Neoplasms
Adenocarcinoma, 78-7115
Cell Differentiation, 78-7115
Mouse, Nude, 78-7115
Neoplasm Metastasis, 78-7115

Transplantation, Homologous

- Kidney Neoplasms
Immunosuppression, 78-6736
- Leukemia
Immunosuppression, 78-6736
- Liver Neoplasms
Immunosuppression, 78-6736
Precancerous Conditions, Review, 78-6706

Transplantation Immunology

- Carcinoma
BCG, 78-6996
- Fibrosarcoma
Benzo(a)pyrene, 78-7051
Corynebacterium parvum, 78-7055
T-Lymphocytes, 78-7051, 78-7054
Macrophages, 78-7051
- Hepatosoma
Antigen-Antibody Complex, 78-7042
- Leukemia, Lymphoblastic
Antibody Formation, 78-7013
Antigen-Antibody Complex, 78-7012
Antigens, Neoplasm, 78-7013
Immunoglobulins, 78-7013
T-Lymphocytes, 78-7013
- Lymph Nodes
Complement, 78-7044
- T-Lymphocytes
Antilymphocyte Serum, 78-7044
- Lymphosarcoma
Cells, Cultured, 78-6828
- Mammary Neoplasms, Experimental
Corynebacterium parvum, 78-7057
Macrophages, 78-7057
- Neoplasms, Experimental
Antigenic Determinants, 78-7031
Histocompatibility Antigens, 78-7044
Virus-Like Particles, 78-7031
Virus, SV40, 78-6911
- Sarcoma
Acetic Anhydride, 78-7048
Antigen-Antibody Complex, 78-7042
Carbodiimide, 1-Ethyl-3-(3-dimethylaminopropyl)-
78-7048
Glutaraldehyde, 78-7048
Histocompatibility Antigens, 78-7044
Methylamine, 78-7048
Sulfuric Acid, Dimethyl Ester, 78-7048
- Thymoma
Antigen-Antibody Complex, 78-7012
- Virus, Influenza
Histocompatibility Antigens, 78-6909
- Virus, Moloney Murine Sarcoma
Histocompatibility Antigens, 78-6909
- Virus, SV40
Freund's Adjuvant, 78-7005

Transplantation Immunology (cont'd)

- Hamster, 78-6985
- Histocompatibility Antigens, 78-6909

Trehalase

- Radiation, Ionizing
- Enzymatic Activity, 78-6814

Trypan Blue

- Lung Neoplasms
- Phagocytosis, 78-6998

Trypsin

- Proteins
- Cell Transformation, Neoplastic, 78-6908
- Fibroblasts, 78-6908
- Virus, Epstein-Barr
- Receptor Activity, 78-6949

Tryptophan

- Ultraviolet Rays
- DNA Repair, 78-6805
- Mutagenic Activity, 78-6805

Tuberculin

- Virus, Adeno 12
- BCG, 78-7008
- Virus, SV40
- BCG, 78-7008

Tunicamycin

- Glycoproteins
- Cell Membrane, 78-6906
- Virus, Polyoma
- Cell Adhesion, 78-6906
- Cell Transformation, Neoplastic, 78-6906
- Glycoproteins, 78-6906
- Virus, SV40
- Cell Adhesion, 78-6906
- Cell Transformation, Neoplastic, 78-6906
- Glycoproteins, 78-6906

Ultraviolet Rays

- 1,8,9-Anthracenetriol
- DNA Repair, 78-6807
- Benz(a)anthracene, 7,12-Dimethyl-
- Melanocytes, Mouse, 78-6777
- Caffeine
- Mutagenic Activity, Review, 78-6606
- Carbamic Acid, *N*-Methyl-*N*-nitroso-, 1-Naphthyl Ester
- DNA Repair, 78-6759
- Chromatids
- Hamster Ovary Cells, 78-6810
- Coumarin, 3-Benzyl-7-hydroxy-4-methyl-
- Occupational Hazard, 78-6663
- Textile Workers, Review, 78-6663
- DNA Polymerase
- Mutagenic Activity, Review, 78-6704
- Nucleotide Sequence, Review, 78-6704
- DNA Repair
- Carcinogenic Activity, Review, 78-6619
- Cell Transformation, Neoplastic, Review, 78-6613
- Cells, Cultured, 78-6812
- In Vivo Measurement Methods, Review, 78-6619
- Pyrimidine Dimers, Review, 78-6666
- Species Difference, Review, 78-6666
- Endonucleases
- DNA Repair, Review, 78-6620
- Erythema
- Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-, 78-6811

Ultraviolet Rays (cont'd)

- Prostaglandins E, 78-6811
- Prostaglandins F, 78-6811
- Psoralen, 8-Methoxy-, 78-6811
- Escherichia coli*
- DNA Repair, 78-6805
- Mutagenic Activity, 78-6805
- Fluorescent Dyes
- Occupational Hazard, 78-6663
- Textile Workers, Review, 78-6663
- Genetics
- Epidemiology, Review, 78-6697
- Glucosephosphate Dehydrogenase
- Enzymatic Activity, 78-6813
- Haemophilus influenzae*
- DNA Repair, 78-6759
- Hypoxanthine Phosphoribosyltransferase
- Mutagenic Activity, 78-6808
- Indole-3-acetic Acid, 1-(*p*-Chlorobenzoyl)-5-methoxy-2-methyl-
- 5,8,11,14-Eicosatetranoic Acid, 78-6811
- Prostaglandins E, 78-6811
- Prostaglandins F, 78-6811
- Keratosis
- DNA Repair, 78-7090, 78-7091
- Leukemia L1210
- 1,8,9-Anthracenetriol, 78-6807
- DNA Repair, 78-6807
- Quinacrine, 78-6807
- Leukocytes
- DNA Repair, 78-7091
- Lymphoma
- Antigens, Viral, 78-7018
- Melanoma
- Age Factors, 78-6804
- Epidemiology, 78-6803, 78-6804
- Precancerous Conditions, Review, 78-6693
- Sunspot Activity, 78-6803
- Methionine, *S*-Adenosyl-
- RNA, Transfer, Methyltransferases, 78-6621
- Psoralen, 8-Methoxy-
- Cell Transformation, Neoplastic, 78-6808
- Chromatids, 78-6810
- 5,8,11,14-Eicosatetranoic Acid, 78-6811
- Mutagenic Activity, 78-6808, 78-6809
- Prostaglandins E, 78-6811
- Prostaglandins F, 78-6811
- 1*H*-Pyrrole-2,5-dione, 1-Ethyl-
- DNA Repair, 78-6812
- Quinacrine
- DNA Repair, 78-6807
- Saccharomyces cerevisiae*
- DNA Repair, 78-6769
- Skin Neoplasms
- Age Factors, 78-6804
- Barbituric Acid, 5-Ethyl-5-phenyl-, 78-6806
- Carcinoma, Epidermoid, 78-6813
- p*-Cresol, 2,6-Di-*tert*-butyl-, 78-6806
- Disulfide, Bis(diethylthiocarbamoyl)-, 78-6806
- Enzymatic Activity, 78-6813
- Epidemiology, 78-6804
- Glucosephosphate Dehydrogenase, 78-6813
- Glutathione, 78-6806
- Keratosis, 78-6813
- Succinate Dehydrogenase
- Enzymatic Activity, 78-6813
- 12-*O*-Tetradecanoylphorbol-13-acetate

Ultraviolet Rays (cont'd)

- Mutagenic Activity, Review, 78-6606
- Ouabain Resistance, 78-6606
- Thioguanine Resistance, 78-6606

Tryptophan

- DNA Repair, 78-6805
- Mutagenic Activity, 78-6805

Virus, Epstein-Barr

- Lymphocyte Transformation, 78-6947
- Migration Inhibitory Factor, 78-6989

Virus, Herpes Simplex

- DNA, Viral, 78-6937

Virus, Herpes Simplex 1

- DNA Repair, 78-6941
- Mutagenic Activity, 78-6941

Virus, SV40

- DNA Repair, Review, 78-6620

Xeroderma Pigmentosum

- DNA Repair, 78-7121
- DNA Repair, Review, 78-6614, 78-6620, 78-6624

Uracil

- Methionine, S-Adenosyl-
RNA, Transfer, Methyltransferases, 78-6621

- RNA, Transfer, Methyltransferases
Carcinoembryonic Antigen, 78-6621
- Enzymatic Activity, 78-6621

Uracil, 5-Fluoro-

- 2-Benzimidazolecarbamic Acid, 5-(2-Thienylcarbonyl)-
Methyl Ester
Cell Migration Inhibition, 78-7081
- Sarcoma
Cell Migration Inhibition, 78-7081

Uranium

- Carcinoma, Basal Cell
Epidemiology, 78-7179
- Occupational Hazard, 78-7179
- Lung Neoplasms
Dose-Response Study, 78-7155
- Epidemiology, 78-7155
- Plutonium Oxide
Alpha Particles, 78-6818

Urea

- Proteins
Cell Transformation, Neoplastic, 78-6908
- Fibroblasts, 78-6908

Urea, 1-(2-Chloroethyl)-3-cyclohexyl-1-nitroso-

- Tongue Neoplasms
Carcinoma, Epidermoid, 78-7098

Urea, Ethyl Nitroso-

- Brain Neoplasms
Astrocytoma, 78-6749
- Glioma, 78-6749
- Oligodendroglioma, 78-6749
- Transplacental Carcinogenesis, 78-6749
- Chromosome Aberrations
Mutagenic Activity, Review, 78-6656
- DNA Replication
Brain, Rat, 78-6749
- Hepatoma
Neoplasm Metastasis, 78-7122
- Isoantigens
Transplacental Carcinogenesis, 78-6748
- Kidney Neoplasms
Transplacental Carcinogenesis, 78-6748

Urea, Ethyl Nitroso- (cont'd)

- Liver Neoplasms
Neoplasm Metastasis, 78-7122
- Nervous System Neoplasms
Carcinogenic Activity, Review, 78-6656
- Chronic Nerve Irritation, 78-6748
- Glycerolphosphate Dehydrogenase, 78-6757
- Neuroblastoma, 78-6756
- Transplacental Carcinogenesis, 78-6748, 78-6757
- RNA, Messenger
Brain, Rat, 78-6757
- Urea, Methyl Nitroso-
Transplacental Carcinogenesis, 78-6748

Urea, Hydroxy-

- Keratoses
DNA Repair, 78-7090
- Leukemia, Myeloblastic
Cell Cycle Kinetics, 78-7150
- Virus, Epstein-Barr
Ribonucleotide Reductases, 78-6950

Urea, Methyl Nitroso-

- 1,2-Benzisothiazolin-3-one, 1,1-Dioxide
Co-carcinogenic Effect, Review, 78-6660
- DNA Repair
Carcinogenic Activity, Review, 78-6622
- Leukemia L1210
Histones, Binding, 78-6752
- Nervous System Neoplasms
Transplacental Carcinogenesis, 78-6757
- Purine, 2-Amino-6-methoxy-
Brain, Rat, 78-6757
- Urea, Ethyl Nitroso-
Transplacental Carcinogenesis, 78-6748

Uridine, 2'-Deoxy-5-iodo-

- Virus, Epstein-Barr
Ribonucleotide Reductases, 78-6950
- Virus, Murine Leukemia
Isolation and Characterization, 78-6892

Urogenital Neoplasms

- Water Supply
Epidemiology, 78-7175

Uterine Neoplasms

- Adenocarcinoma
Genetics, 78-7118
- Laurence-Moon-Biedl Syndrome, 78-7133
- Digestive System Neoplasms
Case Report, 78-7119
- Peutz-Jeghers Syndrome, 78-7119
- Estrogens
Classification, 78-7181
- Epidemiology, 78-6647, 78-7181
- Epidemiology, Review, 78-6641, 78-6642, 78-6651
- Genetics, Review, 78-6647
- Menopause, Review, 78-6642, 78-6647, 78-6651
- Precancerous Conditions, Review, 78-6641
- Risk Factor Analysis, 78-7181
- Laurence-Moon-Biedl Syndrome
Case Report, 78-7133
- Precancerous Conditions
Epidemiology, Review, 78-6642
- Stein-Leventhal Syndrome
Case Report, 78-7133

Vaginal Neoplasms

- Adenocarcinoma

Vaginal Neoplasms (cont'd)

- Case Report, 78-7135
- Ultrastructural Study, 78-7135

Vinblastine Sulfate

- Cell Division
- Cells, Cultured, 78-7081
- Sarcoma
- Cell Migration Inhibition, 78-7081

Vincristine

- Leukemia, Myeloblastic
- Drug Therapy, 78-6815

Vincristine Sulfate

- Cell Division
- Cells, Cultured, 78-7081
- Sarcoma
- Cell Migration Inhibition, 78-7081

Viral Proteins

- Leukemia
 - Virus, Gross Murine Leukemia, 78-6880
- Virus, Adeno 2
 - RNA, Messenger, 78-6964
- Virus, Adeno 2 - SV40 Hybrid
 - RNA, Messenger, 78-6966
- Virus, Avian Myeloblastosis
 - Detergents, 78-6845
 - Electrolyte-Protein Complex, 78-6845
 - Isoelectric Focusing, 78-6845
 - Isolation and Characterization, 78-6845
- Virus, Avian Sarcoma
 - RNA, Messenger, 78-6840
 - RNA, Transfer, 78-6840
- Virus, Epstein-Barr
 - Cell Membrane, 78-6955
- Virus, Gross Murine Leukemia
 - Antigen-Antibody Reactions, 78-6880
 - Carbohydrates, 78-6880
 - Isolation and Characterization, 78-6880
 - Tryptic Peptides, 78-6880
- Virus, Harvey Murine Sarcoma
 - Isolation and Characterization, 78-6894
- Virus, Kirsten Murine Sarcoma
 - Isolation and Characterization, 78-6894
- Virus, Moloney Murine Leukemia
 - Antigenic Determinants, 78-6877
 - Virus, Recombinant, 78-6877
- Virus, Murine Leukemia
 - Antigenic Determinants, 78-6871, 78-6877
 - Host Range, 78-6872
 - Immune Serums, 78-6871
 - Mink Cell Focus-Inducing Isolate, 78-6877
 - Testosterone, 78-6871
 - Tissue Origin, 78-6871
 - Virus, Recombinant, 78-6872, 78-6877
- Virus, Murine Mammary Tumor
 - Antigen-Antibody Reactions, 78-6865
 - Hydrophobicity, 78-6870
 - Isolation and Characterization, 78-6865, 78-6870
 - RNA, Viral, 78-6865
- Virus, Polyoma
 - RNA, Messenger, 78-6904
 - RNA, Viral, 78-6904
- Virus, Radiation Leukemia
 - Antigenic Determinants, 78-6874
 - Isolation and Characterization, 78-6874
- Virus, Rauscher Murine Leukemia

Viral Proteins (cont'd)

- Antigenic Determinants, 78-6889
- Biosynthesis, 78-6887
- Hydrophobicity, 78-6870
- Immunoprecipitation, 78-6887
- Isolation and Characterization, 78-6870, 78-6887
- 78-6889
- Lymphocyte Transformation, 78-7017
- Proteolytic Enzyme Inhibitors, 78-6887
- Reverse Transcriptase, 78-6887
- Virus, Rous Sarcoma
 - Amino Acids, 78-6852
 - Antigen-Antibody Reactions, 78-6856
 - Antigenic Determinants, 78-6853
 - Cell Transformation, Neoplastic, 78-6856
 - Cytoplasm, 78-6856
 - Detergents, 78-6845
 - Electrolyte-Protein Complex, 78-6845
 - Isoelectric Focusing, 78-6845
 - Isolation and Characterization, 78-6845, 78-6852
 - 78-6856
 - RNA, Messenger, 78-6852
 - RNA, Viral, 78-6852
 - Temperature Sensitive Mutants, 78-6856
 - Tryptic Peptide Mapping, 78-6852
- Virus, Simian Adeno 7
 - Cytopathogenic Effect, Viral, 78-6968
- Virus, Simian Sarcoma
 - Antibody Specificity, 78-6936
 - Antigenic Determinants, 78-6936
 - Isolation and Characterization, 78-6936
- Virus, SV40
 - Antigens, Viral, 78-6924
 - Cell Transformation, Neoplastic, 78-6924
 - Temperature Sensitive Mutants, 78-6924
- Virus, Abelson Murine Leukemia
 - Cell Transformation, Neoplastic
 - Replication-Defective Virus, Review, 78-6678
 - Virus, Helper, Review, 78-6678
- Immunosuppression
 - Cell Transformation, Neoplastic, 78-7110
- Lymphoma
 - Neoplasm Metastasis, 78-6875
- RNA, Viral
 - DNA-RNA Hybridization, Review, 78-6678
- Virus, Adeno 2
 - Cell Transformation, Neoplastic
 - DNA-RNA Hybridization, 78-6962
- Chromosomes
 - Cell Transformation, Neoplastic, 78-6962
 - RNA, Messenger, 78-6964
- Cycloheximide
 - Peptides, 78-6963
- Cytosine, 1- β -D-Arabinofuranosyl-
 - Peptides, 78-6963
- DNA-RNA Hybridization
 - Ultrastructural Study, 78-6964
- DNA, Viral
 - Chromosomes, 78-6962
 - Hybrid Cells, 78-6962
- HeLa Cells
 - Virus Replication, 78-6965
- Hybrid Cells
 - DNA-RNA Hybridization, 78-6962
- Peptides
 - Isolation and Characterization, HeLa Cells, 78-6963

- Virus, Adeno 2 (cont'd)**
 RNA, Messenger, 78-6963
 RNA, Messenger
 DNA-RNA Hybridization, 78-6964
 Hybrid Cells, 78-6962
 RNA Replication, 78-6964
 Viral Proteins
 RNA, Messenger, 78-6964
 Virus Replication
 Cytoplasm, 78-6965
 Virus, SV40
 Virus, Helper, 78-6927
- Virus, Adeno 5**
 Carcinoma, Epidermoid
 Reverse Transcriptase, 78-6967
 Virus Activation, 78-6967
 Virus Replication, 78-6967
 Melanoma
 Reverse Transcriptase, 78-6967
 Virus Activation, 78-6967
 Virus Replication, 78-6967
 Sarcoma
 Reverse Transcriptase, 78-6967
 Virus Activation, 78-6967
 Virus Replication, 78-6967
 Virus, Simian Adeno 7
 Virus Replication, 78-6968
- Virus, Adeno 12**
 Antigens
 Immunity, Cellular, 78-7008
 Antigens, Viral
 Antigen-Antibody Reactions, 78-6970
 Cytosine, 1- β -D-Arabinofuranosyl-, 78-6970
 Isolation and Characterization, Chick Fibroblasts
 78-6970
 BCG
 Tuberculin, 78-7008
 Cell Transformation, Neoplastic
 Proteins, Phosphorylation, 78-6969
 Chromatin
 Proteins, Phosphorylation, 78-6969
 DNA, Viral
 Cell Transformation, Neoplastic, Review, 78-6670
 Immunity, Cellular
 Age Factors, 78-7008
 Animal Model, Hamster, 78-7008
 Lipopolysaccharides
 Antibody Formation, 78-7008
 T-Lymphocytes
 Cell Transformation, Neoplastic, 78-6984
 Rosette Formation, 78-6984
 Ribosomes
 Proteins, Phosphorylation, 78-6969
- Virus, Adeno 2 - SV40 Hybrid**
 HeLa Cells
 RNA, Messenger, 78-6966
 RNA, Messenger
 Isolation and Characterization, 78-6966
 Nucleotide Sequence, 78-6966
 Viral Proteins
 RNA, Messenger, 78-6966
- Virus, Avian Leukosis**
 Cell Transformation, Neoplastic
 Target Cells, Review, 78-6676
 Genes, Viral
- Virus, Avian Leukosis (cont'd)**
 Classification, Review, 78-6676
 Hematopoietic Stem Cells
 Target Cells, Review, 78-6676
- Virus, Avian Leukosis-Sarcoma**
 Genes, Viral
 Cell Transformation, Neoplastic, Review, 78-6677
 Nucleotide Sequence
 DNA-RNA Hybridization, 78-6843
 Genes, Viral, 78-6843
 Virus Subgroup, 78-6843
 RNA, Viral
 DNA-RNA Hybridization, 78-6843
 Virus, Helper
 Cell Transformation, Neoplastic, Review, 78-6677
- Virus, Avian Myeloblastosis**
 Glycoproteins
 Antigen-Antibody Reactions, 78-6844
 Cell Membrane, 78-6844
 Isolation and Characterization, 78-6844
 Osteopetrosis
 Antigen-Antibody Reactions, 78-6844
 Viral Proteins
 Detergents, 78-6845
 Electrolyte-Protein Complex, 78-6845
 Isoelectric Focusing, 78-6845
 Isolation and Characterization, 78-6845
- Virus, Avian Sarcoma**
 Cell Transformation, Neoplastic
 Target Cells, Review, 78-6676
 DNA, Viral
 Chromatin, Precipitation, 78-6842
 Purification Method, 78-6842
 Genes, Viral
 Classification, Review, 78-6676
 Proteins
 Cell Membrane, 78-6841
 Cell Transformation, Neoplastic, 78-6841
 Isolation and Characterization, Chick Embryo Fibroblasts, 78-6841
 RNA, Messenger
 Genes, Viral, 78-6840
 Virus Replication, 78-6840
 RNA, Viral
 Cell-Free Translation, 78-6840
 Polyribonucleotides, 78-6972
 Viral Proteins
 RNA, Messenger, 78-6840
 RNA, Transfer, 78-6840
- Virus, B77**
 BCG
 Cell Transformation, Neoplastic, 78-7006
- Virus, Baboon**
 Leukemia
 Isolation and Characterization, Review, 78-6672
 Virus, RD-114
 Antigenic Determinants, Review, 78-6674
- Virus, Bovine Leukemia**
 Leukemia, Lymphoblastic
 Epidemiology, 78-7169
 Multiple Myeloma
 Antibody Specificity, 78-6897
- Virus, C-Type RNA Tumor**
 Antibodies, Viral

Virus, C-Type RNA Tumor (cont'd)

- Radioimmunoassay, Tissue Samples, 78-6679
- Antigens, Viral
 - Radioimmunoassay, Tissue Samples, 78-6679
- Carcinoma
 - Ultrastructural Study, Review, 78-6679
- Leukemia
 - Ultrastructural Study, Review, 78-6679
 - Virus-Like Particles, Review, 78-6672
- Lupus Erythematosus, Systemic
 - Ultrastructural Study, Review, 78-6679
- Lymphoma
 - Ultrastructural Study, Review, 78-6679
- Multiple Myeloma
 - Isolation and Characterization, 78-6897
 - Reverse Transcriptase, 78-6897
 - Virus-Like Particles, 78-6897
- Sarcoma
 - Ultrastructural Study, Review, 78-6679
- Virus, Gross Murine Leukemia
 - Immunization, 78-7024
- Virus, Herpes Simplex
 - Virus Activation, 78-6937
- Virus, Rauscher Murine Leukemia
 - Immunization, 78-7024
- Xeroderma Pigmentosum
 - Ultrastructural Study, Review, 78-6679

Virus, Cytomegalo

- Bladder Neoplasms
 - Antibodies, Viral, 78-7068
 - Carcinoma, Transitional Cell, 78-7068
- Prostatic Hypertrophy
 - Antibodies, Viral, 78-7068
- Prostatic Neoplasms
 - Antibodies, Viral, 78-7068

Virus, DNA Tumor

- Chromosome Aberrations
 - Carcinogenic Activity, Review, 78-6670
- Genes, Viral
 - Cell Transformation, Neoplastic, Review, 78-6670
 - 78-6686

Virus, Epstein-Barr

- Antibodies, Viral
 - Immunosuppression, 78-6952
- Antigen-Antibody Reactions
 - Precancerous Conditions, 78-6951
- Antigenic Determinants
 - Cell Transformation, Neoplastic, 78-7015
 - Immunity, Cellular, 78-7015
- Antigens, Viral
 - Complement Fixation, 78-6953
 - DNA, Binding, 78-6953
 - Immunity, Cellular, 78-7015
 - Isolation and Characterization, 78-6953
 - Phosphonoacetic Acid, 78-6946
 - Raji Cells, 78-7015
- Burkitt's Lymphoma
 - Antigen-Antibody Reactions, 78-6951
 - Seroepidemiology, Uganda, 78-6951
- Cell Adhesion
 - Ultrastructural Study, Daudi Cells, 78-6955
- Cell Membrane
 - Raji Cells, 78-6949
 - Receptor Activity, 78-6949
- DNA Replication
 - Carcinogenic Activity, Review, 78-6683

Virus, Epstein-Barr (cont'd)

- DNA, Viral
 - DNA-DNA Hybridization, 78-6954
 - Endonucleases, 78-6954
 - Nucleotide Sequence, 78-6954
- Dwarfism
 - Lymphocyte Transformation, 78-6947
- Endonucleases
 - Nucleotide Sequence, 78-6954
- Glucose, 2-Deoxy-
 - Virus Replication, 78-6946
- Head and Neck Neoplasms
 - Antibodies, Viral, 78-7034
- Hodgkin's Disease
 - Migration Inhibitory Factor, 78-6989
- Immunosuppression
 - Case Report, 78-6952
- Infectious Mononucleosis
 - Migration Inhibitory Factor, 78-6989
- Leukemia, Myeloblastic
 - Antibodies, Viral, 78-7071
 - Case Report, 78-7071
 - Neoplasms, Multiple Primary, 78-7071
- Leukocytes
 - Migration Inhibitory Factor, 78-6989
- Lymphocytes
 - Interferon, 78-6982
- B-Lymphocytes
 - Antibody Specificity, 78-7015
 - Antigens, Viral, 78-7015
- T-Lymphocytes
 - Immunosuppression, 78-6952
- Lymphoma
 - Migration Inhibitory Factor, 78-6989
- Migration Inhibitory Factor
 - Immunity, Cellular, 78-6989
- Nasopharyngeal Neoplasms
 - Antibodies, Viral, 78-7034
 - Carcinoma, 78-7034
 - Cellular Inclusions, 78-7101
 - DNA, Viral, Review, 78-6632
 - Histological Study, Review, 78-6632
 - Lymphoma, 78-7034
 - Seroepidemiology, Review, 78-6632
- Nose Neoplasms
 - Case Report, 78-7071
 - Neoplasms, Multiple Primary, 78-7071
- Phosphonoacetic Acid
 - Virus Replication, 78-6946
- Ribonucleotide Reductases
 - Virus Replication, 78-6950
- Sarcoma, Reticulum Cell
 - Antigens, Viral, 78-6948
 - Case Report, 78-6948
 - Virus-Like Particles, 78-6948
- Sodium Chloride
 - Anticomplement Immunofluorescence Test, 78-6953
- Trypsin
 - Receptor Activity, 78-6949
- Ultraviolet Rays
 - Lymphocyte Transformation, 78-6947
 - Migration Inhibitory Factor, 78-6989
- Urea, Hydroxy-
 - Ribonucleotide Reductases, 78-6950
- Uridine, 2'-Deoxy-5-iodo-
 - Ribonucleotide Reductases, 78-6950
- Viral Proteins

- Virus, Epstein-Barr (cont'd)**
 Cell Membrane, 78-6955
 Virus, Marek's Disease Herpes
 Antibodies, Viral, 78-6861
 Xeroderma Pigmentosum
 Lymphocyte Transformation, 78-6947
- Virus, Equine Herpes 1**
 see Virus, Equine Rhinopneumonitis
- Virus, Equine Rhinopneumonitis**
 Carcinoma, Transitional Cell
 Thymidine Kinase, 78-6898
 Virus Replication, 78-6898
 Cells, Cultured
 Virus Replication, 78-6898
 Fibroblasts
 Thymidine Kinase, 78-6898
 Virus Replication, 78-6898
 Thymidine Kinase
 Enzymatic Activity, 78-6898
 Thymine, 1- β -D-Arabinofuranosyl-
 Virus Replication, 78-6898
- Virus, Feline Leukemia**
 Antigen-Antibody Reactions
 Seroepidemiology, Humans, 78-7165
 Antigens, Viral
 Isolation and Characterization, 78-6896
 Horizontal Transmission
 Seroepidemiology, Humans, 78-7165
 Leukemia
 Horizontal Transmission, 78-7165
 Lymphoma
 Horizontal Transmission, 78-7165
 Lymphosarcoma
 Antigen-Antibody Reactions, 78-6896
 Antigenic Determinants, 78-6896
 Occupational Hazard
 Antigen-Antibody Reactions, 78-7165
 Virus, RD-114
 Antigenic Determinants, Review, 78-6674
- Virus, Friend Murine Leukemia**
 Antigen-Antibody Reactions
 Immunosuppression, 78-7003
 Spleen Cells, 78-7003
 Antigens, Viral
 Anti-Antibodies, 78-6879
 Antibody-Dependent Cytotoxicity, 78-6879
 Cell Membrane, 78-6879
 Immune Response, 78-6879
 Erythroleukemia
 Antibody-Dependent Cytotoxicity, 78-6879
 Antigens, Viral, 78-6879
 Hematopoietic Stem Cells
 Colony Formation, 78-6878
 Virus Replication, 78-6878
 Immunity, Passive
 Antigens, Viral, 78-6879
 Immunosuppression
 Tumor Extract vs Culture Supernatant, 78-7003
 Radiation, Ionizing
 Antigens, Viral, 78-6879
 Virus Replication, 78-6878
 Virus, Moloney Murine Leukemia
 Virus, Recombinant, 78-6884
- Virus, Friend Spleen Focus-Forming**
 Cell Transformation, Neoplastic
- Virus, Friend Spleen Focus-Forming (cont'd)**
 Replication-Defective Virus, Review, 78-6678
 Virus, Helper, Review, 78-6678
 RNA, Viral
 DNA-RNA Hybridization, Review, 78-6678
- Virus, Gibbon Ape Lymphoma**
 Leukemia
 Isolation and Characterization, Review, 78-6672
- Virus, Gross Murine Leukemia**
 Formaldehyde
 Antibody Formation, 78-7024
 Immunity, Cellular, 78-7024
 Leukemia
 Viral Proteins, 78-6880
 Plasmacytoma
 Antigens, Viral, 78-7038
 Viral Proteins
 Antigen-Antibody Reactions, 78-6880
 Carbohydrates, 78-6880
 Isolation and Characterization, 78-6880
 Tryptic Peptides, 78-6880
 Virus, C-Type RNA Tumor
 Immunization, 78-7024
- Virus, Harvey Murine Sarcoma**
 RNA, Viral
 Nucleotide Sequence, 78-6894
 Viral Proteins
 Isolation and Characterization, 78-6894
 Virus, Kirsten Murine Sarcoma
 DNA-RNA Hybridization, 78-6894
- Virus, Helper**
 Virus, Avian Leukosis-Sarcoma
 Cell Transformation, Neoplastic, Review, 78-6677
 Virus, Murine Leukemia
 Cell Transformation, Neoplastic, Review, 78-6677
 Virus, Rat Leukemia
 Virus, Rat Sarcoma, 78-6895
 Virus, RNA Tumor
 Cell Transformation, Neoplastic, Review, 78-6686
 Virus, SV40
 Virus, Adeno 2, 78-6927
- Virus, Hepatitis**
 DNA Polymerase
 Isolation and Characterization, Woodchuck
 78-6959
 Hepatoma
 Australia Antigen, 78-6685
 Epidemiology, Vietnam, 78-6958
 Immune Serums, 78-6960
 Liver Cirrhosis, 78-6958
 Phosphodiesterases, 78-6960
 Seroepidemiology, Review, 78-6685
 Ultrastructural Study, Woodchuck, 78-6959
 Vertical Transmission, Review, 78-6685
 Virus-Like Particles, 78-6960
 Virus-Like Particles
 Isolation and Characterization, Woodchuck
 78-6959
- Virus, Herpes**
 DNA Replication
 Carcinogenic Activity, Review, 78-6683
 DNA, Viral
 Cell Transformation, Neoplastic, Review, 78-6673
 Isolation and Characterization, Review, 78-6673

Virus, Herpes (cont'd)

- Leukemia, Lymphoblastic
- Case Report, Child, 78-6956
- Precancerous Conditions, 78-6956

Virus, Herpes Saimiri

- Antigens, Viral
- Phosphonoacetic Acid, 78-6946
- Glucose, 2-Deoxy-
- Virus Replication, 78-6946
- Phosphonoacetic Acid
- Virus Replication, 78-6946

Virus, Herpes Simplex

- Psoralen, 4,5'-8-Trimethyl-
- DNA, Viral, 78-6937
- Ultraviolet Rays
- DNA, Viral, 78-6937
- Virus, C-Type RNA Tumor
- Virus Activation, 78-6937

Virus, Herpes Simplex 1

- Antigens, Viral
- Antibody Specificity, 78-6940
- Cell Transformation, Neoplastic, 78-6944
- Immune Serums, 78-6940
- Cervix Neoplasms
- Antibodies, Viral, 78-6939
- Lymphocyte Transformation, 78-7065
- DNA Replication
- Genes, Viral, 78-6943
- Temperature Sensitive Mutants, 78-6943
- DNA, Viral
- Antigens, Viral, 78-6944
- C-Type Virus Activation, Mouse, 78-6937
- Cell Transformation, Neoplastic, 78-6944
- Endonucleases
- Cell Transformation, Neoplastic, 78-6944
- Fibroblasts
- Cell Transformation, Neoplastic, 78-6944
- Ovarian Neoplasms
- Antibodies, Viral, 78-6939
- Peptides
- Physical Mapping, 78-6938
- Temperature Sensitive Mutants, 78-6943
- Virus, Recombinant, 78-6938
- RNA, Viral
- Isolation and Characterization, 78-6942
- Polyribosomes, 78-6942
- Thymidine Kinase
- Mutagenic Activity, 78-6941
- Ultraviolet Rays
- DNA Repair, 78-6941
- Mutagenic Activity, 78-6941

Virus, Herpes Simplex 2

- Antigens, Bacterial
- Antibodies, Viral, 78-7065
- Antigens, Viral
- Antibody Specificity, 78-6940
- Immune Serums, 78-6940
- Immunity, Cellular, 78-7065
- Cervix Neoplasms
- Antibodies, Viral, 78-6939, 78-7065
- Carcinogenic Activity, Review, 78-6683
- Carcinoma, 78-6939
- Carcinoma, Epidermoid, 78-6945
- Carcinoma In Situ, 78-6939
- Epidemiology, Yugoslavia, 78-6945

Virus, Herpes Simplex 2 (cont'd)

- Immunity, Cellular, 78-7065
- Lymphocyte Transformation, 78-7065
- Precancerous Conditions, 78-6939, 78-7065
- Seroepidemiology, 78-6939, 78-6945
- DNA Replication
- Carcinogenic Activity, Review, 78-6683
- DNA, Viral
- C-Type Virus Activation, Mouse, 78-6937
- Ovarian Neoplasms
- Antibodies, Viral, 78-6939
- Peptides
- Physical Mapping, 78-6938
- Virus, Recombinant, 78-6938

Virus, Influenza

- Cell Membrane
- Histocompatibility Antigens, 78-6909
- Histocompatibility Antigens
- Transplantation Immunology, 78-6909
- Virus, SV40
- Histocompatibility Antigens, 78-6909

Virus, Kirsten Murine Sarcoma

- Cell Transformation, Neoplastic
- Ultrastructural Study, Mink Lung Cells, 78-6893
- RNA, Viral
- Nucleotide Sequence, 78-6894
- Viral Proteins
- Isolation and Characterization, 78-6894
- Virus, Harvey Murine Sarcoma
- DNA-RNA Hybridization, 78-6894
- Virus, Murine Leukemia
- Isolation and Characterization, 78-6892
- Virus, Recombinant
- Isolation and Characterization, 78-6892

Virus, Leukemia

- Leukemia, Monoblastic
- Isolation and Characterization, 78-6975
- T-Lymphocytes, 78-6975
- Virus-Like Particles, 78-6975
- B-Lymphocytes
- Cells, Cultured, 78-6975
- Lymphocyte Transformation, 78-6975
- RNA, Viral
- Polyribonucleotides, 78-6972

Virus-Like Particles

- Cervix Neoplasms
- Condylomata Acuminata, 78-7136
- Condylomata Acuminata
- Ultrastructural Study, 78-7136
- Virus, Papilloma, 78-7136
- Virus, Papova, 78-7136
- Hepatitis
- Isolation and Characterization, Woodchuck
- 78-6959
- Hepatoma
- Isolation and Characterization, Woodchuck
- 78-6959
- Liver Cirrhosis, 78-6960
- Virus, Hepatitis, 78-6960
- Laryngeal Neoplasms
- Papilloma, 78-6900
- Letterer-Siwe Disease
- Ultrastructural Study, 78-6977
- Leukemia, Lymphoblastic
- Ultrastructural Study, 78-6977

Virus-Like Particles (cont'd)

- Leukemia, Monoblastic
 - Virus, Leukemia, 78-6975
- Mammary Neoplasms, Experimental
 - Virus, Murine Mammary Tumor, 78-7104
- Multiple Myeloma
 - Virus, C-Type RNA Tumor, 78-6897
- Neoplasms, Experimental
 - Transplantation Immunology, 78-7031
- Neuroblastoma
 - Ultrastructural Study, 78-6977
- Oncogenic Viruses
 - Ultrastructural Study, Hamster Lung Cells, 78-6974
- Prostatic Hypertrophy
 - Isolation and Characterization, 78-6976
- Prostatic Neoplasms
 - Reverse Transcriptase, 78-6976
- Rhabdomyosarcoma
 - Ultrastructural Study, 78-6977
- Sarcoma, Reticulum Cell
 - Virus, Epstein-Barr, 78-6948
- Virus, Hepatitis
 - Isolation and Characterization, Woodchuck 78-6959
- Virus, Murine Mammary Tumor
 - Cells, Cultured, 78-7104
- Virus, SV40
 - Histones, 78-6923
 - Mitomycin C, 78-6933
 - Ultrastructural Study, Monkey Cells, 78-6923

Virus, Lymphocytic Choriomeningitis

- Cell Membrane
 - Fluorescence, 78-6855
 - 1-Naphthylamine, *N*-Phenyl-, 78-6855
 - Pyrene, 78-6855
 - Pyrenebutyric Acid, 78-6855

Virus, Marek's Disease Herpes

- Antigens
 - Antigen-Antibody Reactions, 78-6860
 - Cell Membrane, 78-6860
 - Spleen Cells, 78-6860
- Breast Neoplasms
 - Antibodies, Viral, 78-6861
- Carcinoma
 - Antibodies, Viral, 78-6861
 - Antigens, Viral, 78-6861
- Colonic Neoplasms
 - Antibodies, Viral, 78-6861
- Concanavalin A
 - Immune Response, 78-7049
 - Lymphocyte Transformation, 78-7049
- Cyclophosphamide
 - Immunity, Passive, 78-7023
- Immunity, Cellular
 - Genetic Resistance, 78-7049
 - Strain Difference, 78-7050
- Immunity, Passive
 - Killer Cells, 78-7023
 - Spleen Cells, 78-7023
- Leukemia, Lymphoblastic
 - Immunity, Passive, 78-7023
- Liver Neoplasms
 - Antibodies, Viral, 78-6861
- Lung Neoplasms
 - Antibodies, Viral, 78-6861
- T-Lymphocytes

Virus, Marek's Disease Herpes (cont'd)

- Concanavalin A, 78-7050
- Lymphocyte Depletion, 78-7050
- Lymphoma
 - Antibodies, Viral, 78-6861
- Melanoma
 - Antibodies, Viral, 78-6861
 - Antigens, Viral, 78-6861
- Ovarian Neoplasms
 - Antibodies, Viral, 78-6861
- Radiation, Ionizing
 - Immunity, Passive, 78-7023
- Sarcoma
 - Antibodies, Viral, 78-6861
 - Antigens, Viral, 78-6861
- Virus, Epstein-Barr
 - Antibodies, Viral, 78-6861
- Virus, Turkey Herpes
 - Antigens, 78-6860

Virus, Mason-Pfizer Monkey

- Breast Neoplasms
 - DNA-RNA Hybridization, 78-6869

Virus, Moloney Murine Leukemia

- Antigenic Determinants
 - Virus, Recombinant, 78-6884
- Antigens, Viral
 - Ascitic Fluid, 78-7018
 - Cell Membrane, 78-7018
- Cell Membrane
 - Nucleotidases, 78-7018
- DNA, Viral
 - DNA-RNA Hybridization, 78-6881
 - Integration Sites, 78-6881
 - Virus Replication, 78-6881
- Lymphoma
 - Antigen-Antibody Reactions, 78-6883
 - Antigens, Viral, 78-7010, 78-7018
 - Cell Membrane, 78-7018
 - Histocompatibility Antigens, 78-6883
 - Immunity, Cellular, 78-6883, 78-7010
- RNA, Viral
 - Nucleotide Sequence, 78-6882
 - Virus, Recombinant, 78-6882
- Viral Proteins
 - Antigenic Determinants, 78-6877
 - Virus, Recombinant, 78-6877
- Virus, Friend Murine Leukemia
 - Virus, Recombinant, 78-6884
- Virus, Rauscher Murine Leukemia
 - Virus, Recombinant, 78-6884
- Virus, Recombinant
 - DNA-RNA Hybridization, 78-6884
 - Isolation and Characterization, 78-6882, 78-6884
 - Nucleotide Sequence, 78-6882

Virus, Moloney Murine Sarcoma

- Cell Membrane
 - Histocompatibility Antigens, 78-6909
- Histocompatibility Antigens
 - Transplantation Immunology, 78-6909
- Virus, SV40
 - Histocompatibility Antigens, 78-6909

Virus, Murine Leukemia

- Antibody Formation
 - Immune Response, 78-6876
- Antigenic Determinants

Virus, Murine Leukemia (cont'd)

- Genes, Viral, 78-6892
- Isolation and Characterization, 78-6892
- Antigens, Neoplasm
 - Immune Serums, 78-6978
 - Immunity, Cellular, 78-6978
- Antigens, Viral
 - Antigen-Antibody Reactions, 78-6876
 - Mouse, AKR, 78-6876
- Genes, Viral
 - Virus, Recombinant, 78-6675
- Immune Serums
 - Immunity, Cellular, 78-6978
- Leukemia
 - Antibody Formation, 78-7021, 78-7022
 - Immunosuppression, 78-7022
- Lymphocytes
 - Immune Serums, 78-6978
 - Plant Agglutinins, 78-6978
- T-Lymphocytes
 - Genes, Viral, 78-6677
 - Precancerous Conditions, Review, 78-6677
- Mammary Neoplasms, Experimental
 - Antigens, Neoplasm, 78-6978
- Testosterone
 - Viral Proteins, 78-6871
- Uridine, 2'-Deoxy-5-iodo-
 - Isolation and Characterization, 78-6892
- Viral Proteins
 - Antigenic Determinants, 78-6871, 78-6877
 - Host Range, 78-6872
 - Immune Serums, 78-6871
 - Mink Cell Focus-Inducing Isolate, 78-6877
 - Tissue Origin, 78-6871
 - Virus, Recombinant, 78-6872, 78-6877
- Virus, Helper
 - Cell Transformation, Neoplastic, Review, 78-6677
- Virus, Kirsten Murine Sarcoma
 - Isolation and Characterization, 78-6892
- Virus, Recombinant
 - Carcinogenic Activity, Review, 78-6675
 - Host Range, 78-6872
- Virus Replication
 - Mouse, AKR, 78-6876

Virus, Murine Mammary Tumor

- Antigens, Viral
 - Immune Response, 78-7016
 - Milk, Hybrid Mouse, 78-6866
 - Radioimmunoassay, Epididymis, Coagulating Gland 78-6867
 - Vertical Transmission, 78-6866
- Breast Neoplasms
 - DNA-RNA Hybridization, 78-6869
- Cells, Cultured
 - Virus Replication, 78-6863
- Dexamethasone
 - RNA Replication, 78-6863
- DNA, Viral
 - DNA-DNA Hybridization, 78-6863
 - Virus Replication, 78-6863
- Genetics
 - Cell Transformation, Neoplastic, 78-6866
- Leukemia
 - Antigens, Viral, 78-7016
- Liver, Mouse
 - DNA-RNA Hybridization, 78-6864
- Mammary Neoplasms, Experimental

Virus, Murine Mammary Tumor (cont'd)

- Antigen-Antibody Complex, 78-7058
- DNA-RNA Hybridization, 78-6864
- DNA, Viral, 78-6868
- Estrone, 78-6868
- Progesterone, 78-6868
- Prolactin, 78-6868
- RNA, Viral, 78-6862, 78-6868
- Virus-Like Particles, 78-7104
- RNA Replication
 - Xenopus laevis*, 78-6865
- RNA, Viral
 - DNA-RNA Hybridization, 78-6864
 - Isolation and Characterization, 78-6864
- Viral Proteins
 - Antigen-Antibody Reactions, 78-6865
 - Hydrophobicity, 78-6870
 - Isolation and Characterization, 78-6865, 78-6870
 - RNA, Viral, 78-6865
- Virus-Like Particles
 - Cells, Cultured, 78-7104
- Virus Replication
 - Mammary Epithelial Cells, 78-6863

Virus, Murine Sarcoma

- Cell Transformation, Neoplastic
 - Proteins, Phosphorylation, 78-6969
 - Replication-Defective Virus, Review, 78-6678
 - Virus, Helper, Review, 78-6678
- Chromatin
 - Proteins, Phosphorylation, 78-6969
- Lymphocytes
 - Interferon, 78-6982
- Ribosomes
 - Proteins, Phosphorylation, 78-6969
- RNA, Viral
 - DNA-RNA Hybridization, Review, 78-6678
 - Polyribonucleotides, 78-6972

Virus, Newcastle Disease

- Lymphocytes
 - Interferon, 78-6983

Virus, Papilloma

- Antigens, Viral
 - Carcinogenic Activity, Review, 78-6671
- Bracken Fern
 - Co-carcinogenic Effect, 78-6899
- Carcinoma, Epidermoid
 - Warts, 78-7036
- Condylomata Acuminata
 - Virus-Like Particles, 78-7136
- Digestive System Neoplasms
 - Carcinoma, Epidermoid, 78-6899
 - Epidemiology, Cattle, 78-6899
 - Precancerous Conditions, 78-6899
- Laryngeal Neoplasms
 - Isolation and Characterization, 78-6900
- Skin Neoplasms
 - Carcinogenic Activity, Review, 78-6686
 - Carcinoma, Epidermoid, 78-6684
 - Epidemiology, Review, 78-6684
 - Immune Response, Review, 78-6684
 - Immunity, Cellular, 78-7036
 - Isolation and Characterization, 78-7036
 - Warts, 78-7036
- Warts
 - Immunity, Cellular, 78-7036

- Virus, Papova**
 - Condylomata Acuminata
 - Virus-Like Particles, 78-7136
- Virus, Polio**
 - L Cells
 - RNA Replication, 78-6971
- Virus, Polyoma**
 - Antigens, Viral
 - Carcinogenic Activity, Review, 78-6671
 - Cell Differentiation, 78-6902
 - Hybrid Cells, 78-6902
 - Cell Transformation, Neoplastic
 - Antigenic Determinants, Review, 78-6671
 - Antigens, Review, 78-6671
 - DNA, Viral
 - Endonuclease-Resistant Mutants, 78-6903
 - RNA Polymerase, 78-6901
 - RNA Replication, 78-6901
 - Endonuclease-Resistant Mutants
 - Isolation and Characterization, 78-6903
 - Glycoproteins
 - Cell Transformation, Neoplastic, 78-6906
 - Rifampicin
 - RNA Replication, 78-6901
 - RNA, Messenger
 - Isolation and Characterization, 78-6904
 - Teratoid Tumor
 - Antigens, Viral, 78-6902
 - Hybrid Cells, 78-6902
 - Tunicamycin
 - Cell Adhesion, 78-6906
 - Cell Transformation, Neoplastic, 78-6906
 - Glycoproteins, 78-6906
 - Viral Proteins
 - RNA, Messenger, 78-6904
 - RNA, Viral, 78-6904
 - Virus Replication
 - Antigenic Determinants, Review, 78-6671
 - Endonuclease-Resistant Mutants, 78-6903
- Virus, Polyoma, BK**
 - Antigens, Viral
 - Epidemiology, 78-6905
 - Immune Serums
 - Antibodies, Viral, 78-6905
 - Neoplasms
 - Antigens, Viral, 78-6905
 - DNA-DNA Hybridization, 78-6905
- Virus, Radiation Leukemia**
 - Lymphoma
 - Antigenic Determinants, 78-7029
 - DNA Nucleotidyltransferases, 78-7029
 - Isoantigens, 78-7029
 - T-Lymphocytes, 78-7029
 - Ultrastructural Study, 78-7029
 - Thymoma
 - Isolation and Characterization, 78-6874
 - Viral Proteins
 - Antigenic Determinants, 78-6874
 - Isolation and Characterization, 78-6874
 - Virus Replication
 - Host Range, 78-6874
 - Mink. Cells, 78-6874
- Virus, Rat Leukemia**
 - Cell Transformation, Neoplastic
 - Carcinogenic Activity, 78-6895
- Virus, Rat Leukemia (cont'd)**
 - Cholanthrene, 3-Methyl-
 - Virus Activation, 78-6873
 - Fibrosarcoma
 - Antigenic Determinants, 78-6873
 - Cholanthrene, 3-Methyl-, 78-6873
 - Isolation and Characterization, 78-6873
 - Virus Cultivation
 - Carcinogenic Activity, 78-6895
 - Virus, Rat Sarcoma
 - Virus, Helper, 78-6895
- Virus, Rat Sarcoma**
 - Fibroblasts
 - Carcinogenic Activity, 78-6895
 - Virus Cultivation
 - Carcinogenic Activity, 78-6895
 - Virus, Rat Leukemia
 - Virus, Helper, 78-6895
- Virus, Rauscher Murine Leukemia**
 - Anti-Antibodies
 - Reverse Transcriptase, 78-6888
 - Ribonuclease, 78-6888
 - Antibody Formation
 - Cells, Cultured, 78-7021
 - Antigens, Viral
 - Immunization, 78-7024
 - Concanavalin A
 - Lymphocyte Transformation, 78-7017
 - DNA Polymerase
 - Temperature Sensitive Mutants, 78-6885
 - Thermolability, 78-6885
 - Formaldehyde
 - Antibody Formation, 78-7024
 - Immunity, Cellular, 78-7024
 - Glycoproteins
 - Cell Transformation, Neoplastic, 78-6890
 - Histocompatibility Antigens, 78-6890
 - T-Lymphocytes, 78-6890
 - Histocompatibility Antigens
 - Cell Transformation, Neoplastic, 78-6890
 - T-Lymphocytes, 78-6890
 - Leukemia
 - Antibody Formation, 78-7021
 - Lymphoma
 - Antigen-Antibody Reactions, 78-6883
 - Histocompatibility Antigens, 78-6883
 - Immunity, Cellular, 78-6883
 - Plant Agglutinins
 - Lymphocyte Transformation, 78-7017
 - Polyribonucleotides
 - Ribonuclease, 78-6888
 - Ribonuclease
 - Reverse Transcriptase, 78-6885, 78-6888
 - Temperature Sensitive Mutants, 78-6885
 - Thermolability, 78-6885
 - RNA, Viral
 - Isolation and Characterization, 78-6886
 - Poly A, 78-6886
 - Ribonuclease, 78-6888
 - Viral Proteins
 - Antigenic Determinants, 78-6889
 - Biosynthesis, 78-6887
 - Hydrophobicity, 78-6870
 - Immunoprecipitation, 78-6887
 - Isolation and Characterization, 78-6870, 78-6887
 - 78-6889

Virus, Rauscher Murine Leukemia (cont'd)

- Lymphocyte Transformation, 78-7017
- Proteolytic Enzyme Inhibitors, 78-6887
- Reverse Transcriptase, 78-6887

Virus, C-Type RNA Tumor

- Immunization, 78-7024

Virus, Moloney Murine Leukemia

- Virus, Recombinant, 78-6884

Virus, Sindbis

- Reverse Transcriptase, 78-6891
- Viral Interference, 78-6891
- Virus Replication, 78-6891

Virus, RD-114**Antigens, Viral**

- Isolation and Characterization, Review, 78-6674

Nucleic Acids

- Isolation and Characterization, Review, 78-6674

Virus, Baboon

- Antigenic Determinants, Review, 78-6674

Virus, Feline Leukemia

- Antigenic Determinants, Review, 78-6674

Virus Replication**Carcinoma, Epidermoid**

- Virus, Adeno 5, 78-6967

Carcinoma, Transitional Cell

- Virus, Equine Rhinopneumonitis, 78-6898

Lymphocytes

- Interferon, 78-6982

Melanoma

- Virus, Adeno 5, 78-6967

Sarcoma

- Virus, Adeno 5, 78-6967

Thymoma

- Antibodies, Viral, 78-6874

Virus, Adeno 2

- Cytoplasm, 78-6965
- HeLa Cells, 78-6965

Virus, Adeno 5

- Virus, Simian Adeno 7, 78-6968

Virus, Avian Sarcoma

- RNA, Messenger, 78-6840

Virus, Epstein-Barr

- Glucose, 2-Deoxy-, 78-6946
- Phosphonoacetic Acid, 78-6946
- Ribonucleotide Reductases, 78-6950

Virus, Equine Rhinopneumonitis

- Cells, Cultured, 78-6898
- Fibroblasts, 78-6898
- Thymine, 1- β -D-Arabinofuranosyl-, 78-6898

Virus, Friend Murine Leukemia

- Hematopoietic Stem Cells, 78-6878
- Radiation, Ionizing, 78-6878

Virus, Herpes Saimiri

- Glucose, 2-Deoxy-, 78-6946
- Phosphonoacetic Acid, 78-6946

Virus, Moloney Murine Leukemia

- DNA, Viral, 78-6881

Virus, Murine Leukemia

- Mouse, AKR, 78-6876

Virus, Murine Mammary Tumor

- Cells, Cultured, 78-6863
- DNA, Viral, 78-6863
- Mammary Epithelial Cells, 78-6863

Virus, Polyoma

- Antigenic Determinants, Review, 78-6671
- Endonuclease-Resistant Mutants, 78-6903

Virus Replication (cont'd)**Virus, Radiation Leukemia**

- Host Range, 78-6874
- Mink Cells, 78-6874

Virus, Rauscher Murine Leukemia

- Virus, Sindbis, 78-6891

Virus, Rous Sarcoma

- Chromatin Bodies, 78-6858
- Embryo Fibroblasts, Pigeon, 78-6850
- Tissue Culture, 78-6857

Virus, SV40

- Antigenic Determinants, Review, 78-6671
- Antigens, Viral, 78-6935
- Cell Nucleus, Microinjection, 78-6912
- Cells, Cultured, 78-6912
- Chromosomes, 78-6918
- DNA, Viral, 78-6927
- Endonucleases, 78-6927
- Nucleoproteins, 78-6918

Virus, RNA Tumor**Breast Neoplasms**

- Carcinoma, Ductal, 78-6869

Chromosome Aberrations

- Carcinogenic Activity, Review, 78-6670

Fibroblasts

- Cell Transformation, Neoplastic, 78-6928
- Growth Substances, 78-6928

Genes, Viral

- Cell Transformation, Neoplastic, Review, 78-6670

Polyribonucleotides

- Chromatography, 78-6972

Reverse Transcriptase

- Carcinogenic Activity, Review, 78-6669

RNA, Messenger

- Isolation and Characterization, Review, 78-6675
- Viral Proteins, Review, 78-6675

RNA, Viral

- Polyribonucleotides, 78-6972

Virus, Helper

- Cell Transformation, Neoplastic, Review, 78-6686

Virus, Rous Sarcoma**Actin**

- Cell Transformation, Neoplastic, 78-6846
- Immunofluorescence Microscopy, 78-6846
- Temperature Sensitive Mutants, 78-6846

Antigenic Determinants

- Cell Transformation, Neoplastic, 78-6853

Ascorbic Acid

- Cell Transformation, Neoplastic, 78-6849
- Glucose, 2-Deoxy-, 78-6849

BCG

- Cell Transformation, Neoplastic, 78-7006
- Immunity, Cellular, 78-7006
- Methanol Extraction Residue, 78-7006

Cell Membrane

- Cell Transformation, Neoplastic, 78-6855
- Fluorescence, 78-6855
- 1-Naphthylamine, *N*-Phenyl-, 78-6855
- Pyrene, 78-6855
- Pyrenebutyric Acid, 78-6855

Chromosomes

- Cell Transformation, Neoplastic, 78-6858
- Cells, Cultured, 78-6858
- Chromatin Bodies, 78-6858
- Metaphase, 78-6858

DNA Polymerase

Virus, Rous Sarcoma (cont'd)

- Temperature Sensitive Mutants, 78-6851
- DNA, Viral
 - Infectivity, Sheared Nucleic Acid, 78-6847
 - Transformation, Genetic, 78-6847
- Endonucleases
 - Escherichia coli*, 78-6854
 - Nucleotide Sequence, 78-6854
 - Reverse Transcriptase, 78-6854
- Genes, Viral
 - Cytoplasmic Effect, 78-6846
- Microtubules
 - Cell Transformation, Neoplastic, 78-6846
- Phosphoproteins
 - Cell Transformation, Neoplastic, 78-6848
 - Genes, Viral, 78-6848
 - Protein Kinase, 78-6848
 - Temperature Sensitive Mutants, 78-6848
- Pigeons
 - Host Range, 78-6850
- Protein Kinase
 - Immunoglobulins, Heavy Chain, 78-6848
- Reverse Transcriptase
 - RNA, Transfer, Binding, 78-6851
 - Temperature Sensitive Mutants, 78-6851
- RNA, Viral
 - Cell Transformation, Neoplastic, 78-6859
 - DNA-RNA Hybridization, 78-6859
 - Endonucleases, 78-6854
 - Nucleotide Sequence, 78-6859
 - Viral Proteins, 78-6852
- Sarcoma
 - Immunogenetics, 78-7048
- Viral Proteins
 - Amino Acids, 78-6852
 - Antigen-Antibody Reactions, 78-6856
 - Antigenic Determinants, 78-6853
 - Cell Transformation, Neoplastic, 78-6856
 - Cytoplasm, 78-6856
 - Detergents, 78-6845
 - Electrolyte-Protein Complex, 78-6845
 - Isoelectric Focusing, 78-6845
 - Isolation and Characterization, 78-6845, 78-6852, 78-6856
 - RNA, Messenger, 78-6852
 - Temperature Sensitive Mutants, 78-6856
 - Tryptic Peptide Mapping, 78-6852
- Virus Replication
 - Chromatin Bodies, 78-6858
 - Embryo Fibroblasts, Pigeon, 78-6850
 - Tissue Culture, 78-6857

Virus, Simian Adeno 7

- Viral Proteins
 - Cytopathogenic Effect, Viral, 78-6968
- Virus, Adeno 5
 - Virus Replication, 78-6968
- Virus, SV40
 - Viral Interference, 78-6968

Virus, Simian Sarcoma

- Leukemia
 - Isolation and Characterization, Review, 78-6672
- RNA, Viral
 - Ribonuclease, 78-6888
- Viral Proteins
 - Antibody Specificity, 78-6936
 - Antigenic Determinants, 78-6936

Virus, Simian Sarcoma (cont'd)

- Isolation and Characterization, 78-6936

Virus, Sindbis

- Virus, Rauscher Murine Leukemia
 - Reverse Transcriptase, 78-6891
 - Viral Interference, 78-6891
 - Virus Replication, 78-6891

Virus, SV40

- Aflatoxin B1
 - Virus Replication, Review, 78-6620
- Anemia, Aplastic
 - Twins, 78-6934
- Antigens
 - Antibody Formation, 78-7008
 - Immunity, Cellular, 78-7008
- Antigens, Viral
 - Carcinogenic Activity, Review, 78-6671
 - Cell Cycle Kinetics, 78-6932
 - Cell Transformation, Neoplastic, 78-6925, 78-6926, 78-6930, 78-6931, 78-6932
 - Chromosomes, Binding, 78-6916
 - Deletion Mutants, 78-6920, 78-6930
 - Endonucleases, 78-6925
 - Genetics, 78-6934
 - Hybrid Cells, 78-6914
 - Nucleotide Sequence, 78-6920
 - Temperature Sensitive Mutants, 78-6926
 - Viral Proteins, 78-6924
 - Virus Replication, 78-6935

BCG

- Tuberculin, 78-7008
- Blood Platelets
 - Growth Substances, 78-6928
- 2,3-Butanediol, 1,4-Dimercapto-
Chromosomes, 78-6907
- Cell Adhesion
 - Cell Transformation, Neoplastic, 78-6908
- Cell Fusion
 - Antigens, Viral, 78-6925
- Cell Membrane
 - Histocompatibility Antigens, 78-6909
- Cell Transformation, Neoplastic
 - Antigenic Determinants, Review, 78-6671
 - Antigens, Review, 78-6671
 - Cell Cycle Kinetics, 78-6910
 - Deletion Mutants, 78-6930
 - Fibroblasts, Rat, 78-6926
 - Fibroblasts, Review, 78-6661
 - A Gene Product, 78-6910
 - Genes, Viral, Review, 78-6661
 - Isolation and Characterization, 78-6924
 - Temperature Sensitive Mutants, 78-6926, 78-6931
- Chromosomes
 - Histones, 78-6907
 - Virus Replication, 78-6918
- Chromosomes, Human, 6-12
 - Hybrid Cells, 78-6919
- Chromosomes, Human, 16-18
 - Hybrid Cells, 78-6919
- Cytosol
 - DNA Replication, 78-6913
- DNA-DNA Hybridization
 - Ultrastructural Study, 78-6922
- DNA Replication
 - Cell Nucleus, 78-6913
 - Cell Transformation, Neoplastic, 78-6910

Virus, SV40 (cont'd)

- Chromosomes, 78-6913
- Deletion Mutants, 78-6917
- Nucleoproteins, 78-6913
- Nucleotide Sequence, 78-6917
- Temperature Sensitive Mutants, 78-6910
- DNA, Viral**
 - Antigens, Viral, 78-6920, 78-6927
 - C-Type Virus Activation, Mouse, 78-6937
 - Cell Transformation, Neoplastic, 78-6925, 78-6933
 - Cell Transformation, Neoplastic, Review, 78-6670
 - Chromosomes, Human, 6-12, 78-6919, 78-6929
 - Chromosomes, Human, 16-18, 78-6919
 - DNA-DNA Hybridization, 78-6922
 - DNA Replication, 78-6922
 - Endonucleases, 78-6922, 78-6925
 - Hybrid Cells, 78-6919, 78-6926
 - Mitomycin C, 78-6933
 - Purification Method, 78-6842
 - Virus Replication, 78-6927
- Endonucleases**
 - Chromosomes, 78-6907
 - Deletion Mutants, 78-6930
 - DNA Replication, 78-6927
 - Virus Replication, 78-6927
- Ethidium Bromide**
 - Cell Transformation, Neoplastic, 78-6915
 - DNA Replication, 78-6915
 - Drug-Resistant Mutants, 78-6915
 - Mitochondria, 78-6915
 - Proteins, 78-6915
- Fibroblasts**
 - Cell Transformation, Neoplastic, 78-6928
 - Genetics, 78-6934
 - Growth Substances, 78-6928
 - Proteins, 78-6908
- Freund's Adjuvant**
 - Immunity, Cellular, 78-7005
 - T-Lymphocytes, 78-7005
 - Transplantation Immunology, 78-7005
- Glycoproteins**
 - Cell Transformation, Neoplastic, 78-6906
- Histocompatibility Antigens**
 - Cell Transformation, Neoplastic, 78-6909
 - Transplantation Immunology, 78-6909
- Histones**
 - Cell Movement, 78-6907
 - Chromatin-DNA Hybrids, 78-6907
 - Chromosomes, 78-6923
- Hybrid Cells**
 - Antigens, Viral, 78-6929
 - Cell Transformation, Neoplastic, 78-6914, 78-6919
 - Chromosomes, 78-6914
 - Chromosomes, Human, 6-12, 78-6929
 - DNA, Viral, 78-6929
- Immune Serums**
 - Arming/Potentiating Activity, 78-6985
 - Lymphocytotoxicity, 78-6985
- Immunity, Cellular**
 - Age Factors, 78-7008
 - Animal Model, Hamster, 78-7008
 - Hamster, 78-6985
- Immunosuppression**
 - Cell Transformation, Neoplastic, 78-7110
- Leukemia**
 - Genetics, 78-6934
 - Twins, 78-6934

Virus, SV40 (cont'd)

- Leukemia, Lymphoblastic
 - Twins, 78-6934
- Leukemia, Subleukemic
 - Twins, 78-6934
- Lipopolysaccharides
 - Antibody Formation, 78-7008
- Lymphocytes
 - Immunity, Cellular, 78-6985
- T-Lymphocytes
 - Cell Transformation, Neoplastic, 78-6984
 - Rosette Formation, 78-6984
- Lymphoma
 - Genetics, 78-6934
 - Twins, 78-6934
- Methanesulfonic Acid, Ethyl Ester
 - Virus Replication, Review, 78-6620
- Methanesulfonic Acid, Methyl Ester
 - Virus Replication, Review, 78-6620
- Neoplasms, Experimental
 - Antilymphocyte Serum, 78-6911
 - Castration, 78-6911
 - Cortisone Acetate, 78-6911
 - Cyclophosphamide, 78-6911
 - Estrone, 78-6911
 - Testosterone, Propionate, 78-6911
 - Transplantation Immunology, 78-6911
- Nucleoproteins
 - Chromosomes, 78-6918
 - Isolation and Characterization, 78-6918
 - Virus Replication, 78-6918
- Plasminogen Activators
 - Fibroblasts, Review, 78-6661
- RNA, Messenger
 - DNA-RNA Hybridization, 78-6920
 - DNA, Viral, 78-6920
- RNA, Viral
 - Cytoplasm, Nucleus, 78-6921
 - Degradation, 78-6921
 - DNA-RNA Hybridization, 78-6921
 - RNA, Messenger, 78-6921
- Sarcoma
 - Histocompatibility Antigens, 78-7044
- Temperature Sensitive Mutants
 - Isolation and Characterization, 78-6924
- Teratoid Tumor
 - Antigens, Viral, 78-6932
- Transplantation Immunology
 - Hamster, 78-6985
- Tunicamycin
 - Cell Adhesion, 78-6906
 - Cell Transformation, Neoplastic, 78-6906
 - Glycoproteins, 78-6906
- Ultraviolet Rays
 - DNA Repair, Review, 78-6620
- Viral Proteins
 - Cell Transformation, Neoplastic, 78-6924
 - Temperature Sensitive Mutants, 78-6924
- Virus, Adeno 2
 - Virus, Helper, 78-6927
- Virus, Influenza
 - Histocompatibility Antigens, 78-6909
- Virus-Like Particles
 - Histones, 78-6923
 - Mitomycin C, 78-6933
 - Ultrastructural Study, Monkey Cells, 78-6923
- Virus, Moloney Murine Sarcoma

Virus, SV40 (cont'd)

Histocompatibility Antigens, 78-6909

Virus Replication

Antigenic Determinants, Review, 78-6671

Cell Nucleus, Microinjection, 78-6912

Cells, Cultured, 78-6912

Virus, Simian Adeno 7

Viral Interference, 78-6968

Virus, Turkey Herpes**Antigens**

Antigen-Antibody Reactions, 78-6860

Virus, Marek's Disease Herpes

Antigens, 78-6860

Virus, Vaccinia**DNA, Viral**

C-Type Virus Activation, Mouse, 78-6937

Lymphocytes

Interferon, 78-6983

Virus, Vesicular Stomatitis**Lymphocytes**

Interferon, 78-6983

Vitamin B**Areca**

Carcinogenic Potential, Ingredients, 78-6727

Mouth Neoplasms

Areca, 78-6727

Vitamin B12**Liver Neoplasms**

Carcinogen, Chemical, 78-6638

Vitamin EAcetohydroxamic Acid, *N*-Fluoren-2-yl-

Mutagenic Activity, 78-6730

Vulvar Neoplasms**Carcinoma**

Diagnosis, 78-7137

Epidemiology, 78-7137

Carcinoma, Epidermoid

Epidemiology, 78-7137

Carcinoma In Situ

Epidemiology, 78-7137

Warts**Burkitt's Lymphoma**

Genetics, Review, 78-6684

Carcinoma, Epidermoid

Genetics, 78-7036

Virus, Papilloma, 78-7036

Skin Neoplasms

Epidemiology, Review, 78-6684

Genetics, Review, 78-6684

Virus, Papilloma, 78-7036

Virus, Papilloma

Immunity, Cellular, 78-7036

Water Pollution**Nitrosamines**

Concentration Levels, 78-7183

Emission Sources, 78-7183

Wounds and Injuries**Carcinoma, Basal Cell**

Case Report, 78-7092

Meningioma

Epidemiology, 78-7170

Wounds and Injuries (cont'd)**Skin Neoplasms**

Carcinoma, Basal Cell, 78-7092

Xanthine, 1-Hydroxy-

Structure-Activity Relationship

Carcinogenic Activity, Rat, 78-6718

Xanthine, 3-Hydroxy-**Fibroma**

Carcinogenic Activity, Rat, 78-6718

Hepatoma

Carcinogenic Activity, Rat, 78-6718

Sarcoma

Carcinogenic Activity, Rat, 78-6718

Xanthine, 7-Hydroxy-**Liver Neoplasms**

Carcinogenic Activity, Rat, 78-6718

Xanthine, 9-Hydroxy-

Structure-Activity Relationship

Carcinogenic Activity, Rat, 78-6718

Xeroderma PigmentosumAcetamide, *N*-(Acetyloxy)-*N*-fluoren-2-yl-

DNA Repair, Review, 78-6624

Carcinogen, Chemical

DNA Repair, Review, 78-6614

Complementation Group

DNA Repair, Review, 78-6614

Genetics

Case Report, 78-7121

Skin Neoplasms

Carcinoma, Epidermoid, 78-7121

Case Report, 78-7121

DNA Repair, 78-7121

Genetics, 78-7121

Neoplasms, Multiple Primary, 78-7121

Ultraviolet Rays

DNA Repair, 78-7121

DNA Repair, Review, 78-6614, 78-6620, 78-6624

Virus, C-Type RNA Tumor

Ultrastructural Study, Review, 78-6679

Virus, Epstein-Barr

Lymphocyte Transformation, 78-6947

Xylene**Abnormalities**

Epidemiology, Child, 78-6765

Leukemia, Lymphoblastic

Case Report, 78-6765

In Utero Exposure, 78-6765

Yttrium Radioisotopes**Sarcoma, Osteogenic**

Dose-Response Study, Rat, 78-6835

Zinc Radioisotopes**Hyperplasia**

Spleen, Mouse, Rat, 78-6817

Zymosan**Leukemia L1210**

Immune Response, 78-7020

Chemical Abstracts Service Registry Number Index

- | | | |
|--|---|--|
| 35-25-6, 78-6766 | 56-49-5, 78-6762, 78-6773, 78-6781
78-6798, 78-6873, 78-6908
78-6969, 78-6994, 78-7007
78-7041, 78-7042, 78-7043
78-7044, 78-7053, 78-7110 | 64-17-5, 78-6653, 78-6699, 78-6945
78-7114 |
| 50-00-0, 78-6657, 78-7024 | | 64-86-8, 78-7081 |
| 50-02-2, 78-6863 | | 66-22-8, 78-6621 |
| 50-04-4, 78-6911, 78-6998 | | 66-27-3, 78-6620 |
| 50-06-6, 78-6607, 78-6729, 78-6742
78-6773, 78-6806 | 56-53-1, 78-6644, 78-6652, 78-6789
78-6998, 78-7011 | 66-81-9, 78-6963 |
| 50-07-7, 78-6759, 78-6933, 78-6988
78-7022 | 56-55-3, 78-6772, 78-6773, 78-6781
78-6786 | 67-21-0, 78-6737 |
| 50-18-0, 78-6815, 78-6911, 78-7023
78-7072 | 56-57-5, 78-6608, 78-6763 | 67-64-1, 78-6765 |
| 50-23-7, 78-6798, 78-6993, 78-7197
78-7198 | 56-65-5, 78-6737, 78-6764, 78-7190 | 67-68-5, 78-7185 |
| 50-28-2, 78-6794 | 56-87-1, 78-7039 | 68-19-9, 78-6638 |
| 50-29-3, 78-6659 | 57-13-6, 78-6908 | 68-22-4, 78-7126 |
| 50-32-8, 78-6623, 78-6626, 78-6657
78-6664, 78-6701, 78-6785
78-6786, 78-7043, 78-7046
78-7051, 78-7059, 78-7076
78-7122 | 57-14-7, 78-6743 | 68-26-8, 78-6630, 78-6638, 78-6796 |
| 50-67-9, 78-6979 | 57-22-7, 78-6815 | 69-05-6, 78-6807 |
| 50-76-0, 78-6815, 78-6973, 78-7072 | 57-30-7, 78-6758 | 70-18-8, 78-6806, 78-6979 |
| 50-81-7, 78-6730, 78-6849 | 57-63-6, 78-7125, 78-7126 | 70-25-7, 78-6608, 78-6612, 78-6622
78-6751, 78-6752, 78-6759
78-7110 |
| 50-89-5, 78-7150 | 57-83-0, 78-6603, 78-6868, 78-7162 | 70-34-8, 78-7025 |
| 51-21-8, 78-7081 | 57-85-2, 78-6911 | 71-43-2, 78-6627 |
| 51-31-0, 78-6766 | 57-88-5, 78-7180 | 72-33-3, 78-7125 |
| 51-41-2, 78-6732 | 57-97-6, 78-6626, 78-6745, 78-6772
78-6773, 78-6774, 78-6775
78-6776, 78-6777, 78-6778
78-6779, 78-6780, 78-6785
78-6786, 78-6984, 78-6991
78-7033, 78-7055, 78-7075 | 72-57-1, 78-6998 |
| 51-43-4, 78-7144 | | 73-22-3, 78-6805 |
| 51-48-9, 78-6798 | 58-08-2, 78-6606 | 73-31-4, 78-6779 |
| 51-75-2, 78-6753, 78-6759, 78-7072 | 58-22-0, 78-6871 | 73-40-5, 78-6716 |
| 51-79-6, 78-6743, 78-6792 | 58-82-2, 78-7001 | 74-89-5, 78-7048 |
| 51-85-4, 78-6730, 78-6979 | 59-02-9, 78-6730 | 74-90-8, 78-6657 |
| 52-90-4, 78-6802, 78-6979 | 59-51-8, 78-6638, 78-6730 | 75-01-4, 78-6634, 78-6636, 78-6712
78-6713, 78-6715 |
| 53-16-7, 78-6794, 78-6868, 78-6911 | 59-89-2, 78-6787 | 75-45-6, 78-6765 |
| 53-86-1, 78-6811 | 59-92-7, 78-6728 | 77-78-1, 78-7048 |
| 53-95-2, 78-6730, 78-6745 | 60-00-4, 78-6908 | 79-01-6, 78-6636, 78-6765, 78-7173 |
| 53-96-3, 78-6618, 78-6729, 78-6730
78-6731, 78-6732, 78-6745
78-6792 | 60-11-7, 78-6764, 78-7042, 78-7063
78-7064 | 79-46-9, 78-6734 |
| 54-12-6, 78-6805 | 60-23-1, 78-6802 | 81-07-2, 78-6660, 78-6738 |
| 54-42-2, 78-6892, 78-6950 | 60-24-2, 78-6988 | 83-89-6, 78-6807 |
| 55-17-4, 78-6737 | 60-57-1, 78-6659 | 84-65-1, 78-6663 |
| 55-18-5, 78-6732, 78-6742, 78-6743
78-6744, 78-6750, 78-7122 | 60-92-4, 78-6766 | 86-44-2, 78-6663 |
| 55-80-1, 78-6744, 78-6766, 78-6767 | 62-49-7, 78-6638 | 90-30-2, 78-6855 |
| 56-23-5, 78-6607, 78-6743, 78-6750 | 62-50-0, 78-6620, 78-6743, 78-6759 | 90-65-3, 78-6629 |
| | 62-75-9, 78-6625, 78-6739, 78-6740
78-6742, 78-6744, 78-6746
78-6802 | 91-22-5, 78-6762 |
| | 63-68-3, 78-6638, 78-6730 | 91-59-8, 78-6702 |
| | | 97-00-7, 78-7033, 78-7061 |
| | | 97-77-8, 78-6714, 78-6806 |
| | | 100-42-5, 78-6663, 78-6765 |

106-93-4, 78-6714	477-30-5, 78-7081	3902-71-4, 78-6937
107-02-8, 78-6657	480-22-8, 78-6807	4408-78-0, 78-6946
107-92-6, 78-7185	499-45-6, 78-7194	4615-63-8, 78-6772
108-24-7, 78-7048	505-60-2, 78-7088	4637-56-3, 78-6749
108-86-1, 78-6607	506-32-1, 78-6811	4803-27-4, 78-6769
109-89-7, 78-6747	531-85-1, 78-7122	5134-37-2, 78-6621
110-15-6, 78-6764	540-61-4, 78-6755	6051-87-2, 78-6781
111-30-8, 78-7048	540-73-8, 78-6755	6098-44-8, 78-6624, 78-6716, 78-6745
115-09-3, 78-6817	542-88-1, 78-6725	6416-25-7, 78-6663
118-00-3, 78-6725	551-11-1, 78-6811, 78-7197	7090-25-7, 78-6754, 78-6759
119-61-9, 78-6663	551-93-9, 78-6790	7439-89-6, 78-6654
120-12-7, 78-6786	569-64-2, 78-6663	7439-98-7, 78-6635
121-79-9, 78-6802	578-67-6, 78-6762	7440-02-0, 78-6634, 78-7156
124-09-4, 78-6817	578-76-7, 78-6755	7440-06-4, 78-6635
124-40-3, 78-6747	586-17-4, 78-6732	7440-07-5, 78-6821, 78-6823, 78-6824 78-6825
124-64-1, 78-6725	590-96-5, 78-6760	7440-16-6, 78-6635
126-72-7, 78-6663	604-59-1, 78-6730	7440-18-8, 78-6635
127-07-1, 78-6950, 78-7090, 78-7150	605-23-2, 78-6898	7440-27-9, 78-6716
128-37-0, 78-6730, 78-6743, 78-6806	621-64-7, 78-6743	7440-38-2, 78-6634
128-53-0, 78-6812	684-93-5, 78-6622, 78-6660, 78-6748 78-6752, 78-6757	7440-43-9, 78-6797
128-58-5, 78-6663	745-65-3, 78-6782, 78-6798, 78-6811 78-6993	7440-61-1, 78-6818, 78-7155, 78-7179
129-00-0, 78-6786, 78-6855	759-73-9, 78-6656, 78-6748, 78-6749 78-6756, 78-6757, 78-7122	7440-70-2, 78-6928
133-66-4, 78-6663	1162-65-8, 78-6620, 78-6629, 78-6721 78-6722	7553-56-2, 78-6640
134-03-2, 78-6802	1204-06-4, 78-6790	7631-86-9, 78-6998, 78-7038, 78-7057
140-79-4, 78-6743	1239-45-8, 78-6915	7632-00-0, 78-7170
143-67-9, 78-7081	1330-20-7, 78-6765	7647-14-5, 78-6953
147-94-4, 78-6963, 78-6970, 78-7081	1332-21-4, 78-6631, 78-6633, 78-6634 78-6654, 78-6697, 78-6836 78-6837, 78-6838, 78-7061 78-7113, 78-7172	7665-99-8, 78-6767
148-24-3, 78-6762	1385-95-1, 78-6721	7681-49-4, 78-6766
149-29-1, 78-6629	1421-85-8, 78-7110	7683-59-2, 78-6766
151-16-6, 78-6979	1892-57-5, 78-7048	7697-37-2, 78-6662, 78-6733
154-17-6, 78-6849, 78-6946	1932-15-6, 78-6718	7718-54-9, 78-6724
192-97-2, 78-6786	1937-37-7, 78-6663, 78-6771	7771-44-0, 78-6811
215-58-7, 78-6786	2068-78-2, 78-7081	7782-39-0, 78-6746
288-13-1, 78-6760	2481-94-9, 78-6764	7782-44-7, 78-6654
297-83-6, 78-6650	2602-46-2, 78-6663, 78-6771	7782-49-2, 78-6696
298-81-7, 78-6808, 78-6809, 78-6810 78-6811	2613-02-7, 78-6621	7782-77-6, 78-6655, 78-6662, 78-6733
305-03-3, 78-7072	2832-40-8, 78-6663	7787-47-5, 78-6817
306-60-5, 78-6741	3416-21-5, 78-7110	8001-28-3, 78-6626, 78-6660
309-00-2, 78-6659	3483-12-3, 78-6907	8015-95-0, 78-6801, 78-6981, 78-6997
311-45-5, 78-6730	3817-11-6, 78-6735	8063-94-3, 78-6715
362-74-3, 78-6993		9000-07-1, 78-6801, 78-6981, 78-6997
363-24-6, 78-6782, 78-6798, 78-6811 78-6993		9001-45-0, 78-7099
471-29-4, 78-6741, 78-6802		9001-48-3, 78-7077
		9001-78-9, 78-6713, 78-7099

9001-99-4, 78-6885, 78-6888, 78-6973
 78-7200
 9002-07-7, 78-6908, 78-6949
 9002-60-2, 78-6793
 9002-62-4, 78-6640, 78-6728, 78-6778
 78-6779, 78-6868, 78-7095
 78-7162
 9002-71-5, 78-6640, 78-6778
 9002-86-2, 78-6715
 9004-10-8, 78-7197
 9004-70-0, 78-6991
 9005-49-6, 78-7198
 9005-79-2, 78-6707
 9007-12-9, 78-6798
 9007-27-6, 78-7198
 9008-11-1, 78-6982, 78-6983, 78-6993
 9009-54-5, 78-6839
 9010-72-4, 78-7020
 9012-42-4, 78-6766, 78-7188
 9015-91-2, 78-6640
 9035-50-1, 78-6607
 10024-97-2, 78-6710
 10028-15-6, 78-6710
 10043-66-0, 78-6833
 10045-97-3, 78-6817
 10048-13-2, 78-6629, 78-6800
 10088-79-6, 78-6629
 10098-91-6, 78-6835
 10098-97-2, 78-6835
 10099-74-8, 78-6817
 10300-74-0, 78-6771
 11028-71-0, 78-6735, 78-6988, 78-7017
 78-7049, 78-7050
 11056-06-7, 78-7081
 11089-65-9, 78-6906
 11118-26-6, 78-6906

12001-28-4, 78-6631, 78-6633, 78-6634
 78-6654, 78-6697, 78-6836
 78-6837, 78-6838, 78-7061
 78-7113, 78-7172
 12001-29-5, 78-6631, 78-6633, 78-6634
 78-6654, 78-6697, 78-6836
 78-6837, 78-6838, 78-7061
 78-7113, 78-7172
 12001-76-2, 78-6727
 12005-67-3, 78-6817, 78-6820, 78-6834
 12016-67-0, 78-6820, 78-6834
 12059-95-9, 78-6818, 78-6819, 78-6820
 78-6822, 78-6834
 12172-73-5, 78-6631, 78-6633, 78-6634
 78-6654, 78-6697, 78-6836
 78-6837, 78-6838, 78-7061
 78-7113, 78-7172
 12268-14-3, 78-6634
 12587-46-1, 78-6818, 78-6821
 12684-33-2, 78-6769
 12736-98-0, 78-6817, 78-6820, 78-6834
 13010-47-4, 78-7098
 13073-35-3, 78-6737
 13233-32-4, 78-6830
 13292-46-1, 78-6901
 13479-29-3, 78-6718
 13907-45-4, 78-6634
 13907-47-6, 78-6634
 13981-16-3, 78-6822
 13982-39-3, 78-6817
 13982-63-3, 78-6817
 14046-83-4, 78-6772
 14158-27-1, 78-6817
 14987-04-3, 78-6992
 15623-47-9, 78-6830
 16339-07-4, 78-6743
 16561-29-8, 78-6606, 78-6626, 78-6719
 78-6782, 78-6796, 78-7197
 16757-86-1, 78-6783
 16757-87-2, 78-6783
 16757-89-4, 78-6783

16757-90-7, 78-6783
 16757-91-8, 78-6783
 16870-90-9, 78-6718
 16941-32-5, 78-6766
 17068-78-9, 78-6631, 78-6633, 78-6634
 78-6654, 78-6697, 78-6836
 78-6837, 78-6838, 78-7061
 78-7113, 78-7172
 17721-95-8, 78-6743
 18883-66-4, 78-6799
 20535-83-5, 78-6625, 78-6755, 78-6757
 21150-20-9, 78-7190
 21373-20-6, 78-7190
 21884-44-6, 78-6629
 22365-02-2, 78-6640
 22365-17-9, 78-6640
 23095-97-8, 78-6621
 23449-95-8, 78-7194
 24305-27-9, 78-6640
 24554-26-5, 78-7141
 24967-93-9, 78-7198
 27762-78-3, 78-6716
 28378-99-6, 78-6621
 29908-03-0, 78-6621
 30310-80-6, 78-6743
 31430-18-9, 78-7081
 33034-23-0, 78-7194
 34522-32-2, 78-7194
 35050-55-6, 78-6769
 39386-07-7, 78-6846
 39422-15-6, 78-6640
 39603-53-7, 78-6749
 39834-38-3, 78-6772
 51933-03-0, 78-6718
 55556-88-2, 78-6743
 59963-01-8, 78-6623, 78-6784

Wiswesser Line Notation Index

.AM..O2, 78-6817, 78-6820, 78-6834
 .AS, 78-6634
 .BE..G2 78-6817
 .CA, 78-6928
 .CD, 78-6797
 .CM..O2, 78-6820, 78-6834
 .CS, 78-6817
 .D, 78-6746
 .FE, 78-6654
 .I, 78-6640, 78-6833
 .MO, 78-6635
 .NA..F, 78-6766
 .NA..G, 78-6953
 .NA..N-O-Q, 78-7170
 .NI, 78-6634, 78-7156
 .NI..G2, 78-6724
 .N2..O, 78-6710
 .PB..N-O3*2, 78-6817
 .PT, 78-6635
 .PU, 78-6821, 78-6822, 78-6823, 78-6824, 78-6825
 .PU..O2, 78-6818, 78-6819, 78-6820, 78-6822, 78-6834
 .RA, 78-6817, 78-6830
 .RH, 78-6635
 .RU, 78-6635
 .SE, 78-6696
 .SI..O2, 78-6836, 78-6998, 78-7038, 78-7057
 .SR, 78-6817, 78-6835
 .TB, 78-6716
 .TH, 78-6830
 .UR, 78-6818, 78-7155, 78-7179
 .Y, 78-6835
 .ZN, 78-6817
 ER, 78-6607
 E1YE1O 3PO, 78-6663
 E2E, 78-6714
 GXGG YR DG&R DG, 78-6659
 GXGGG, 78-6607, 78-6743, 78-6750
 GYFF, 78-6765
 GYGU1G, 78-6636, 78-6765, 78-7173
 G1U1, 78-6634, 78-6636, 78-6712, 78-6713, 78-6715
 G101G, 78-6725
 G2N1&2G, 78-6753, 78-6759, 78-7072

G2S2G, 78-7088
 L B656 HHJ EMV1, 78-6618, 78-6729, 78-6730, 78-6731
 78-6732, 78-6745, 78-6792
 L B656 HHJ ENQV1, 78-6730, 78-6745
 L B677 MV&T&J CO1 DO1 EO1 JMV1 NO1, 78-7081
 L B677 MV&T&J CO1 DO1 EO1 JM1 NO1, 78-7081
 L C666 BV IVJ, 78-6663
 L C666J, 78-6786
 L D5 C555 A D- EU JUTJ AG AG BG IG JG KG, 78-6659
 L D6 B666 EUT&&&J GQ HQ, 78-6772
 L D6 B666 OU&&&TJ QQ RQ, 78-6772
 L D6 B666 QU &&&TJ OQ PQ, 78-6772
 L D6 B666&&&TJ OQ PQ, 78-6772
 L D6 B666J, 78-6772, 78-6773, 78-6781, 78-6786
 L D6 B666J C J, 78-6626, 78-6745, 78-6772, 78-6773
 78-6774, 78-6775, 78-6776, 78-6777, 78-6778
 78-6779, 78-6780, 78-6785, 78-6786, 78-6984
 78-6991, 78-7033, 78-7055, 78-7075
 L D6 B666T&&&J EQ FQ, 78-6772
 L D6 B6666 2AB TJ, 78-6623, 78-6626, 78-6657, 78-6664
 78-6701, 78-6785, 78-6786, 78-7045, 78-7046
 78-7051, 78-7059, 78-7076, 78-7122
 L D6 B6666 2AB TJ J O, 78-6783
 L D6 B6666 2AB TJ J Q, 78-6783
 L D6 B6666 2AB TJ L M, 78-6783
 L D6 B6666 2AB TJ O P, 78-6783
 L D6 B6666 2AB TJ O Q, 78-6783
 L D6 J6 C666J, 78-6786
 L D6666 B6 2AB TJ, 78-6786
 L E5 B666 FVTTT&J E OQ, 78-6794, 78-6868, 78-6911
 L E5 B666 LUTJ A E FY&3Y QQ -B&AEFO, 78-7180
 L E5 B666 OV MUTJ A CQ E FV1Q FQ -B&ACEF
 78-6798, 78-6993, 78-7197, 78-7198
 L E5 B666 OV MUTJ A E FQ -B&AEF, 78-6871
 L E5 B666 OV MUTJ A1 E1 FV1 -B&AEF, 78-6603
 78-6868, 78-7162
 L E5 B666TTT&J E FQ FIUU1 OQ, 78-7125, 78-7126
 L E5 B666TTT&J E FQ OQ, 78-6794
 L E6 D6656 1A T&&&T&J R, 78-6762, 78-6773, 78-6781
 78-6798, 78-6873, 78-6908, 78-6969, 78-6994
 78-7007, 78-7041, 78-7042, 78-7043, 78-7044
 78-7053, 78-7110
 L E6 K3 D6656 1A U LOT&TT&J S, 78-7110
 L6TJ AMVNNO&2G, 78-7098
 L6UTJ A BL/U1Y&U2/ 2Q C C -T, 78-6630, 78-6638
 78-6796

L66J BMR, 78-6855
 L66J CZ, 78-6702
 L666 B6 2AB PJ, 78-6786, 78-6855
 NCH, 78-6657
 ONN1&1, 78-6625, 78-6739, 78-6740, 78-6742, 78-6744
 78-6746, 78-6802
 ONN2&2, 78-6732, 78-6742, 78-6743, 78-6744, 78-6750
 78-7122
 ONN3&3, 78-6743
 OO, 78-6654
 OOO, 78-6710
 OS1&1, 78-7185
 QR BQ CQ EVO3, 78-6802
 QR BQ DYQ1MY -L, 78-6766
 QR BQ DYQ1M1 -LQR BQ DYQ1M1, 78-7144
 QR DY2& 2U, 78-6644, 78-6652, 78-6789, 78-6998, 78-7011
 QVYZ1R CQ DQ -L, 78-6728
 QVYZ2S1, 78-6638, 78-6730
 QVYZ2S2 -DL, 78-6737
 QV1N1VQ 22, 78-6908
 QV1UYO1&VYU1, 78-6629
 QV2VQ, 78-6764
 QV3, 78-7185
 QV3R DN2G2G, 78-7072
 QV4U3U3U3U6, 78-6811
 Q1 4P &G, 78-6725
 Q1NUNO&1, 78-6760
 Q2, 78-6653, 78-6699, 78-6945, 78-7114
 Q2K &Q, 78-6638
 Q4N4&NO, 78-6735
 R, 78-6627
 RVR, 78-6663
 SH2Q, 78-6988
 SH2Q1Q1SH, 78-6907
 T B666 HKJ EZ H2 IR& LZ &E &9/26, 78-6915
 T C566 DO LVOJ BO1, 78-6808, 78-6809, 78-6810, 78-6811
 T C6 B5665 2AB S BX IN QN NU JH&&TTTJ FO1 IVH
 KVO1 KQ LOV1 M2 E-NT F6 E596 A BN
 LM&&TTJ NVO1 RQ R2, 78-6815
 T C6 B65-24- A D E 2BC G& AV LO NO F&VM OU B&U
 D&U MH&&&TJ DQ GQ IQ J M QO1 R
 SOV1 T UQV WQ & E & E, 78-6901
 T C666 BNJ EG IMY&3N2&2 LO1, 78-6807
 T C666 BO EV INJ D FZ N G- K-/VM- OT5-16- AN FVN
 IVN LVO PVM SVTJ G J KY N RY 2
 78-6815, 78-6973, 78-7072

T D3 B556 BN EM JV MVTTT&J GO1 H1OVZ KZ L
 78-6759, 78-6933, 78-6988, 78-7022
 T E3 D5 C555 A D- FO KUTJ AG AG BG JG KG LG
 78-6659
 T E3 D6 B6666 2AB U FOTT&&&&J HQ IQ, 78-6623
 78-6784
 T E3 H6 D6 B666 FOT&T&&&J, 78-7110
 T E6 C6 B655 DO KV PO RO SU&&&TTJ IQ MO1
 78-6629, 78-6800
 T F5 C6 B655 DOV GV OO QO RUT&&TTJ LO1, 78-6620
 78-6629, 78-6721, 78-6722
 T 66 BNJ GQ, 78-6762
 T5MNJ, 78-6760
 T5OJ BNW E- ET5N CSJ BMVH, 78-7141
 T5OV EHJ CQ DQ EYQ1Q, 78-6730, 78-6849
 T5VNVJ B2, 78-6812
 T56 BM DN FMYMVJ GUM, 78-6716
 T56 BMJ DIYZVQ -L, 78-6805
 T56 BMJ D2Z GQ, 78-6979
 T56 BN DN FMYMVJ GUM D- BT5OTJ CQ DQ E1Q
 78-6725
 T56 BN DN FNVNVJ B F H, 78-6606
 T56 BOV GO IU&TJ FQ, 78-6629
 T56 BSVVMVJ, 78-6660, 78-6738
 T56 BVO DHJ D- D-/R DQ CSWQ& 2 FE HE IE, 78-6650
 T6MPOTJ BO BN2G2G, 78-6815, 78-6911, 78-7023, 78-7072
 T6MVMVJ EF, 78-7081
 T6N DNTJ ANO DNO, 78-6743
 T6N DNTJ A1 DNO, 78-6743
 T6N DOTJ ANO, 78-6787
 T6NVMVJ EI A- ET5OTJ B1Q CQ -A&C, 78-6892, 78-6950
 T6NVNJ DZ A- BT5OTJ CQ DQ E1Q, 78-6963, 78-6970
 78-7081
 T6VMVMV FHJ F2 FR, 78-6607, 78-6729, 78-6742, 78-6773
 78-6806
 T6VMVMV FHJ F2 FR &-NA-, 78-6758
 T6VMVTJ EIYQ- BL6VTJ D F, 78-6963
 T66 BNJ, 78-6762
 T66 BNJ BO EMQ, 78-6749
 T66 BNJ BO ENW, 78-6608, 78-6763
 T66 BNJ JQ, 78-6762
 VHH, 78-6657, 78-7024
 VH1U1, 78-6657
 VH1YQYQYQ1Q -BAA -D, 78-6849, 78-6946
 VH3VH, 78-7048
 WNMYUM&N1&NO, 78-6608, 78-6612, 78-6622, 78-6751
 78-6752, 78-6759, 78-7110

WNR BF ENW, 78-7025
WNR BG ENW, 78-7033, 78-7061
WNR DOPO&O2&O2, 78-6730
WNY, 78-6734
WS1&O1, 78-6620
WS1&O2, 78-6620, 78-6743, 78-6759
ZN1&1, 78-6743
ZR BV1, 78-6790
ZR DR DZ &GH &GH, 78-7122
ZVMQ, 78-6950, 78-7090, 78-7150
ZVN1&NO, 78-6622, 78-6660, 78-6748, 78-6752, 78-6757
ZVN2&NO, 78-6656, 78-6748, 78-6749, 78-6756, 78-6757
78-7122
ZVO2, 78-6743, 78-6792
ZVZ, 78-6908
ZV4YZVQ, 78-7039
Z1, 78-7048
Z1CN, 78-6755

Z1YQR CQ DQ -L, 78-6732
Z2SH, 78-6802
Z2SS2Z, 78-6730, 78-6979
Z6Z, 78-6817
1MM1, 78-6755
1M1, 78-6747
1N1&R DNUNR, 78-6764, 78-7042, 78-7063, 78-7064
1N1&R DNUNR C, 78-6744, 78-6766, 78-6767
1OSWO1, 78-7048
1R X, 78-6765
1U1R, 78-6663, 78-6765
1VOV1, 78-7048
1V1, 78-6765
1X&&R BQ E CX, 78-6730, 78-6743, 78-6806
2M2, 78-6747
2N2&YUS&S 2, 78-6714, 78-6806

RECEIVED JAN 11 1961



